

Exact solvability of an Ising-type model, and exact solvability of the 6-vertex, and 8-vertex, models

P. Rigas

Abstract

We compute the action-angle coordinates for an Ising type model whose L-operator has been previously studied in the literature by Bazhanov and Sergeev. In comparison to computations with such operators that have been examined previously by the author for the 4-vertex, 6-vertex, and 20-vertex, models, computations for asymptotically approximating a collection of sixteen identities with the Poisson bracket, which together constitute the Poisson structure of the Ising type model, exhibit dependencies upon nearest neighbor interactions. In spite of the fact that L-operators for the 20-vertex model are defined in terms of combinatorial, and algebraic, constituents unlike such operators for the 6-vertex model which are defined in terms of projectors and Pauli basis elements, L-operators for the Ising-type model can be used for concluding that a model which interpolates between the 6-vertex, and 8-vertex, models is exactly solvable. ^{1 2}

1 Introduction

1.1 Overview

Exactly solvable, and integrable, models of Statistical Physics have gathered significant attention, whether from the perspective of connections with the Bethe ansatz and correlations [12,20,26], continuum limits, [23], limit shapes, and associated asymptotic phenomena, [8,9,10,11,35,36,37,38,39,43], amongst several other closely related topics [1,2,4,6,17,18,19,22,24,25,27,28,29]. To make use of rapidly emerging developments for many models, and settings, at the intersection of Mathematical Physics, Statistical Physics, and Quantum Physics, [30,31,32,33,34], we propose a new perspective for applying the quantum inverse scattering method (QISM) to a model that has interactions which significantly differ from those of vertex models. From previous work of the author, [41,45], properties of the Poisson bracket - anticommutativity, Leibniz' rule, bilinearity, and the Jacobi identity - one can determine whether integrable, or exactly solvable, properties of vertex models exist. Besides classes of vertex models, which are but one type of many, models appearing in Statistical Physics, Ising type models are another. Such classes of models are interesting due to the fact that, from some perspectives, they greatly differ between the behavior of the Gaussian free field. Despite such differences, there can still be an adaptation of the QISM framework to such models, which not only relies upon similar steps of performing computations with an L-operator for asymptotically representing the transfer and quantum monodromy matrices, but also for formalizing notions of Poisson structures, which hold implications for accompanying Yang-Baxter algebras.

Further studying interactions between such themes, which span the fields of Discrete and Integrable Probability simultaneously, remains of interest. In determining whether integrable properties of models play roles in discrete structures, pursuing such a research program for vertex model, in comparison to Ising type models, predominantly relies upon the ice-rule. As a conversation rule for vertex models, in two dimensions the ice-rule states that there must be two incoming, and two outgoing arrows, surrounding each vertex of the square lattice \mathbf{Z}^2 , while in three dimensions it states that there must be three incoming, or three outgoing arrows, surrounding each vertex of the triangular lattice \mathbf{T} . For Ising type models, in the absence of the ice rule, one instead defines a probability measure over finite volume with a nearest neighbor interactions from a Hamiltonian, as well as with an inverse temperature. Notwithstanding of the differences in which vertex, and the Ising, models are defined, it is still possible to apply components of the QISM in each situation.

For vertex models, in comparison to Ising type models, asymptotic approximations for the transfer matrix, and quantum monodromy matrix, can be formulated from products of L-operators. That is, asymptotically in infinite volume, having a representation of the L-operator, whether in terms of Pauli

¹MSC: 34L25; 60K35

²Keywords: Statistical Physics, Mathematical Physics, eigenvalue problem, SOS model, 6-vertex model, 7-vertex model, 20-vertex model, 4-vertex model

basis elements and other closely related factors dictates which terms determine the approximation of each Poisson bracket, within the entire Poisson structure, in infinite volume. Moreover, whether a vertex model of Statistical Physics has *isotropic* parameters, namely a set of parameters that are all set to be equal, also plays a significant role in determining macroscopic properties of vertex configurations in weak finite limits that are taken towards infinite volume. Ising type models, albeit being introduced in a completely different way, and hence with a completely separate procedure for taking the finite volume limit for obtaining probability measures supported over the entirety of \mathbf{Z}^2 , can still be analyzed with similar computations from those provided by the author [41,45]. To this end, for some $a \in \mathbf{Z}$, introduce the L-operator for an Ising type model, [3],

$$\mathcal{L}_a^{a'} \equiv \left[\mathcal{L} \left[x, \frac{x'}{\xi}, y \right] \right]_a^{a'} \equiv \begin{bmatrix} \frac{xx'}{y^2} \delta_{a,a'+1} - \frac{y^2}{xx'} \delta_{a,a'-1} & q^a \left[\frac{x}{x'} \delta_{a,a'+1} - \frac{x'}{x} \delta_{a,a'-1} \right] \\ q^{-a} \left[\frac{x}{x'} \delta_{a,a'-1} - \frac{x'}{x} \delta_{a,a'+1} \right] & \frac{xx'}{y^2} \delta_{a,a'-1} - \frac{y^2}{xx'} \delta_{a,a'+1} \end{bmatrix},$$

where in the entries of the representation for the Ising type L-operator above, $x \neq x'$ denote two positions over the square lattice, while q parameters, which are widely studied in vertex models as "quantum" parameters, appear in other L-operators that have extensively been studied by the author in [45], which take the form,

$$\hat{L}(\xi) \equiv L_1^{3D} = \exp(\lambda_3(q^{-2}\xi^s)) \begin{bmatrix} q^{D_1} & q^{-2} a_1 q^{-D_1 - D_2} \xi^{s-s_1} & a_1 a_2 q^{-D_1 - 3D_2} \xi^{s-s_1-s_2} \\ a_1^\dagger q^{D_1} \xi^{s_1} & q^{-D_1 + D_2} q^{-2} q^{D_1 - D_2} \xi^s & -a_2 q^{D_1 - 3D_2} \xi^{s-s_2} \\ 0 & a_2^\dagger q^{D_2} \xi^{s_2} & q^{-D_2} \end{bmatrix},$$

given a spectral parameter ξ for the triangular lattice, and differential operators,

$$D_k^j \equiv (D \otimes \mathbf{1}) \mathbf{1}_{\{\mathbf{r} \equiv e_k\}}, \quad D_{k+1}^j \equiv (\mathbf{1} \otimes D) \mathbf{1}_{\{\mathbf{r} \equiv e_{k+1}\}}.$$

that was originally introduced in [6], given basis vectors e_k and e_{k+1} of \mathbf{T} . As a general observation, for adaptations of QISM for vertex models, one studies R-matrices, which satisfying the Yang Baxter equation. With several possible interpretations, such equations not only demonstrate the equality of partition functions for a vertex model in question, but also demonstrate how intertwining relations can be formalized.

1.2 Quantum inverse scattering type objects

For the Ising type model with L-operator \mathcal{L} introduced above, the fact that each entry of the two by two representation is the difference of two terms implies that it is equal to,

$$\begin{bmatrix} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{bmatrix} - \begin{bmatrix} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{bmatrix},$$

under the change of coordinates of $\mathcal{L}_a^{a'}$, which is given by,

$$\begin{bmatrix} x_i x_j \delta_{a,a'+1} - \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} - q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} - q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} - \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{bmatrix}.$$

To apply computations of the type that have previously been introduced by the author in [41,45], it suffices to consider cross terms from neighboring L-operators. In the form introduced above for sites x_i and x_j over the square lattice, one would consider products of the form,

$$\prod_{\substack{1 \leq i < j \leq N \\ x_i \neq x_j}} \left[\begin{bmatrix} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{bmatrix} - \begin{bmatrix} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{bmatrix} \right],$$

for some $N > 0$. As the weak finite volume limit is taken as $N \rightarrow +\infty$, products of the form above can also be taken over j such that,

$$\{j \in \mathbf{N} : j < i\},$$

with,

$$\prod_{\substack{1 \leq j < i \leq N \\ x_i \neq x_j}} \left[\begin{array}{cc} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{array} \right] - \left[\begin{array}{cc} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{array} \right],$$

Despite \mathcal{L} being introduced as an L-operator for an Ising type model, which the authors of [] observed as being a distant relative of the 6-vertex, and 8-vertex, models, several components parallel the structure of correlations, in addition to interactions, which are encapsulated by the ice rule of vertex models. Fundamentally, as a conservation rule, the ice rule significantly impacts the geometry of vertex configurations of the 6-vertex, and 20-vertex, models, [41,45], as well as characteristics of integrability, and exact solvability, that one would hope to prove about such models.

In the following, to obtain asymptotic approximations of the transfer matrix, and quantum monodromy matrix, for the Ising type model, we discuss behaviors of the expansion when $N \equiv 2$ first. Under this choice of N , the terms that emerge from the product,

$$\prod_{\substack{1 \leq i < j \leq 2 \\ x_i \neq x_j}} \left[\begin{array}{cc} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{array} \right] - \left[\begin{array}{cc} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{array} \right],$$

or from the product,

$$\prod_{\substack{1 \leq j < i \leq 2 \\ x_i \neq x_j}} \left[\begin{array}{cc} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{array} \right] - \left[\begin{array}{cc} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{array} \right],$$

include,

$$\begin{aligned} & \left[\begin{array}{cc} x_1 x_2 \delta_{a,a'+1} & q^a \frac{x_1}{x_2} \delta_{a,a'+1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} & \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \end{array} \right] \left[\begin{array}{cc} x_2 x_3 \delta_{a,a'+1} & q^a \frac{x_2}{x_3} \delta_{a,a'+1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'-1} & \frac{x_2 x_3}{y^2} \delta_{a,a'-1} \end{array} \right], \\ & - \left[\begin{array}{cc} \frac{y^2}{x_1 x_2} \delta_{a,a'-1} & q^a \frac{x_1}{x_2} \delta_{a,a'-1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'+1} & \frac{y^2}{x_1 x_2} \delta_{a,a'+1} \end{array} \right] \left[\begin{array}{cc} x_2 x_3 \delta_{a,a'+1} & q^a \frac{x_2}{x_3} \delta_{a,a'+1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'-1} & \frac{x_2 x_3}{y^2} \delta_{a,a'-1} \end{array} \right], \\ & - \left[\begin{array}{cc} x_1 x_2 \delta_{a,a'+1} & q^a \frac{x_1}{x_2} \delta_{a,a'+1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} & \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \end{array} \right] \left[\begin{array}{cc} \frac{y^2}{x_2 x_3} \delta_{a,a'-1} & q^a \frac{x_2}{x_3} \delta_{a,a'-1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'+1} & \frac{y^2}{x_2 x_3} \delta_{a,a'+1} \end{array} \right], \\ & \left[\begin{array}{cc} \frac{y^2}{x_1 x_2} \delta_{a,a'-1} & q^a \frac{x_1}{x_2} \delta_{a,a'-1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'+1} & \frac{y^2}{x_1 x_2} \delta_{a,a'+1} \end{array} \right] \left[\begin{array}{cc} \frac{y^2}{x_2 x_3} \delta_{a,a'-1} & q^a \frac{x_2}{x_3} \delta_{a,a'-1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'+1} & \frac{y^2}{x_2 x_3} \delta_{a,a'+1} \end{array} \right]. \end{aligned}$$

In each of the four terms above, minus signs accumulate in the middle two terms. In the finite weak volume limit as $N \rightarrow +\infty$, for either one of the the infinite products,

$$\begin{aligned} & \prod_{\substack{1 \leq i < j \leq +\infty \\ x_i \neq x_j}} \left[\begin{array}{cc} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{array} \right] - \left[\begin{array}{cc} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{array} \right], \\ & \prod_{\substack{1 \leq j < i \leq +\infty \\ x_i \neq x_j}} \left[\begin{array}{cc} x_i x_j \delta_{a,a'+1} & q^a \frac{x_i}{x_j} \delta_{a,a'+1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'-1} & \frac{x_i x_j}{y^2} \delta_{a,a'-1} \end{array} \right] - \left[\begin{array}{cc} \frac{y^2}{x_i x_j} \delta_{a,a'-1} & q^a \frac{x_i}{x_j} \delta_{a,a'-1} \\ q^{-a} \frac{x_i}{x_j} \delta_{a,a'+1} & \frac{y^2}{x_i x_j} \delta_{a,a'+1} \end{array} \right], \end{aligned}$$

of L-operators to asymptotically approximate the Ising type transfer matrix given by the mapping,

$$t^{\text{Ising Type}} : \mathbf{C}^2 \otimes (\mathbf{C}^2)^N \longrightarrow \mathbf{C}^2 \otimes (\mathbf{C}^2)^N \mapsto \prod_{1 \leq j \leq N} (t^{\text{Ising Type}})_j \propto \prod_{1 \leq j \leq N} \mathcal{L}_{a_j}^{a'_j},$$

where the points a'_j , and a_j , are contained within the finite volume $\mathcal{V}_i \subsetneq \mathbf{Z}^2$, for,

$$(t^{\text{Ising Type}})_j \cap j\mathbf{Z}^2 \neq \emptyset, \\ \text{support}(\mathcal{L}_{a_j}^{a'_j}) \subsetneq \text{support}(j\mathbf{Z}^2) \subsetneq \text{support}(\mathbf{Z}^2) \subsetneq \mathbf{Z}^2,$$

The quantum monodromy matrix given by the mapping,

$$T^{\text{Ising Type}} : \mathbf{C}^2 \otimes (\mathbf{C}^2)^N \longrightarrow \mathbf{C}^2 \otimes (\mathbf{C}^2)^N \mapsto \text{tr} \left[\prod_{1 \leq j \leq N} (t^{\text{Ising Type}})_j \right] \propto \text{tr} \left[\prod_{1 \leq j \leq N} \mathcal{L}_{a_j}^{a'_j} \right].$$

The two mappings above can be seen as lower dimensional objects in comparison to the three-dimensional quantum monodromy matrix for the 20-vertex model examined by the author in [45], which takes the form,

$$T_{a,b}^{3D}(\{u_i\}, \{v'_j\}, \{w''_k\}) : \mathbf{C}^3 \otimes (\mathbf{C}^3)^{\otimes(|N|+||M||_1)} \longrightarrow \mathbf{C}^3 \otimes (\mathbf{C}^3)^{\otimes(|N|+||M||_1)} \mapsto \prod_{j=0}^{-N} \prod_{k=0}^M \left[\text{diag}(\exp(\alpha(i, j, k))) \right. \\ \left. , \exp(\alpha(i, j, k)), \exp(\alpha(i, j, k)) \right) R_{ia,jb,kc}(u - u_i, u' - v'_j, w - w''_k) \Big],$$

where $R \equiv \mathcal{R}$ denotes the Universal R-matrix (see [6] for more details, including the factorization into contributions from the K-matrix). Beyond $N \equiv 3$ terms in the product of L-operators that has been previously discussed, a few of the terms that will be computed for the asymptotic expansion of the transfer matrix include,

$$- \begin{bmatrix} x_1 x_2 \delta_{a,a'+1} & q^a \frac{x_1}{x_2} \delta_{a,a'+1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} & \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \end{bmatrix} \begin{bmatrix} x_2 x_3 \delta_{a,a'+1} & q^a \frac{x_2}{x_3} \delta_{a,a'+1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'-1} & \frac{x_2 x_3}{y^2} \delta_{a,a'-1} \end{bmatrix} \begin{bmatrix} x_3 x_4 \delta_{a,a'+1} & q^a \frac{x_3}{x_4} \delta_{a,a'+1} \\ q^{-a} \frac{x_3}{x_4} \delta_{a,a'-1} & \frac{x_3 x_4}{y^2} \delta_{a,a'-1} \end{bmatrix}, \\ \begin{bmatrix} \frac{y^2}{x_1 x_2} \delta_{a,a'-1} & q^a \frac{x_1}{x_2} \delta_{a,a'-1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'+1} & \frac{y^2}{x_1 x_2} \delta_{a,a'+1} \end{bmatrix} \begin{bmatrix} x_2 x_3 \delta_{a,a'+1} & q^a \frac{x_2}{x_3} \delta_{a,a'+1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'-1} & \frac{x_2 x_3}{y^2} \delta_{a,a'-1} \end{bmatrix} \begin{bmatrix} x_3 x_4 \delta_{a,a'+1} & q^a \frac{x_3}{x_4} \delta_{a,a'+1} \\ q^{-a} \frac{x_3}{x_4} \delta_{a,a'-1} & \frac{x_3 x_4}{y^2} \delta_{a,a'-1} \end{bmatrix}, \\ \begin{bmatrix} \frac{y^2}{x_1 x_2} \delta_{a,a'-1} & q^a \frac{x_1}{x_2} \delta_{a,a'-1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'+1} & \frac{y^2}{x_1 x_2} \delta_{a,a'+1} \end{bmatrix} \begin{bmatrix} \frac{y^2}{x_2 x_3} \delta_{a,a'-1} & q^a \frac{x_2}{x_3} \delta_{a,a'-1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'+1} & \frac{y^2}{x_2 x_3} \delta_{a,a'+1} \end{bmatrix} \begin{bmatrix} \frac{y^2}{x_3 x_4} \delta_{a,a'-1} & q^a \frac{x_3}{x_4} \delta_{a,a'-1} \\ q^{-a} \frac{x_3}{x_4} \delta_{a,a'+1} & \frac{y^2}{x_3 x_4} \delta_{a,a'+1} \end{bmatrix}, \\ \begin{bmatrix} x_1 x_2 \delta_{a,a'+1} & q^a \frac{x_1}{x_2} \delta_{a,a'+1} \\ q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} & \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \end{bmatrix} \begin{bmatrix} \frac{y^2}{x_2 x_3} \delta_{a,a'-1} & q^a \frac{x_2}{x_3} \delta_{a,a'-1} \\ q^{-a} \frac{x_2}{x_3} \delta_{a,a'+1} & \frac{y^2}{x_2 x_3} \delta_{a,a'+1} \end{bmatrix} \begin{bmatrix} \frac{y^2}{x_3 x_4} \delta_{a,a'-1} & q^a \frac{x_3}{x_4} \delta_{a,a'-1} \\ q^{-a} \frac{x_3}{x_4} \delta_{a,a'+1} & \frac{y^2}{x_3 x_4} \delta_{a,a'+1} \end{bmatrix}.$$

Within the quantum inverse scattering framework, analyzing how interactions of either a vertex, or Ising type, model, are encoded through entries of the L-operator continues to remain of great interest. In L-operators for vertex models, one typically encounters Pauli basis elements, projectors, and possibly terms from a mapping into a unital associative algebra, whereas in L-operators for Ising type models, as indicated through the operator above, one encounters contributions from all possible pairwise contributions of two points within some finite volume. For an Ising type model, contributions from each entry of the representation for the L-operator differ from computations of L-operators for vertex models previously studied by the author in [41,45,46]. As one possible representative of an asymptotic expansion of the transfer matrix as the system size tends to that of \mathbf{Z}^2 , products of an L-operator can also be used for studying counterparts of the emptiness formation probability for the 20-vertex model, [47], from previous computations with contour-integral representations under domain-wall boundary conditions for the 6-vertex model [7]. As such, determining how the dimensionality of the underlying state space of a vertex model, whether in two or three dimensions, in addition to the dependency of the model on boundary

conditions, are of great significance. Furthermore, conjectures that have been raised with regards to the convergence of the scaling limit of the 6-vertex model to the Gaussian free field is also indicative of behaviors of other models in Statistical Physics - in which interactions interpolate between pointwise interactions such as those encoded through Hamiltonians of the Ising and Potts models, to interactions for the Gaussian free field which are not defined pointwise.

In the forthcoming sections, we provide computations for products of L-operators for the Ising type model. After having obtained a system of relations for each of the four entries of the product representation, as has already been investigated by the author for the 4-vertex, 6-vertex, and 20-vertex, models, we characterize integrability, and exact solvability, of the Ising type model.

1.3 Paper organization

To obtain the desired asymptotic expansions of the transfer, and quantum monodromy, matrices, we first compute all entries of the product representation for two L-operators. Beyond this first computation for products of L-operators, as in the case for adaptations of the quantum-inverse scattering method for the 4-vertex, 6-vertex, and 20-vertex, models, we demonstrate the existence of suitable action-angle variables for the Ising type model, $\Phi^{\text{Ising}} \equiv \Phi$, for which,

$$\{\Phi, \bar{\Phi}\} \equiv 0.$$

The existence of action-angle variables, such as the one provided above, not only implies that there exists a linearization of the model that sufficiently approximates nonlinear dynamics, and behaviors, of systems with a large system size, but also that integrability, and exact solvability, are expected to hold. For vertex models, as the dimensionality of the underlying state space increases to three dimensions, properties of integrability, and exact solvability, do not hold in the same manner due to the fact that Poisson brackets of action-angle coordinates do not necessarily vanish. In spite of the fact that difficulties exist in adapting the quantum inverse scattering method to show that integrability holds in the context of the 20-vertex model, one can still perform computations with the L-operator which can then be used to characterize Poisson structures in three dimensions, predominantly from the fact that asymptotic approximations for each entry of the product representation for the transfer matrix, and hence for the quantum monodromy matrix, lead to computations with the Poisson bracket.

For the Ising type model, obtaining an asymptotic representation for the transfer, and quantum monodromy, matrices sheds further light on integrability and exactly solvability of the model through the set of Poisson brackets that one must approximate, which together constitute the Poisson structure. Such a collection of brackets is obtained by exhausting all possible options for each of the two arguments of the Poisson bracket, which can further be analyzed to conclude that integrability for the model holds. Such observations, in addition to computations performed by the author in several previous works described at length near the end of the previous subsection above, can be used for analyzing asymptotic behaviors, integrability, and exact solvability, of the Ising type model. As a means of complementing, from previous characterizations of the transfer, and quantum monodromy, matrices in asymptotically large finite volume, To demonstrate the computations that one would expect to encounter with the Poisson bracket for demonstrating that integrability, and exact solvability, holds for the Ising type model within the quantum inverse scattering framework, consider the spanning set,

$$\begin{aligned} & \text{span}_{a,a' \in \mathbf{Z}^2} \left\{ x_1 x_3 \delta_{a,a'+1}, q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} q^a \frac{x_4}{x_1} \delta_{a,a'-1}, x_1 x_2 \delta_{a,a'+1} \frac{y^2}{x_1 x_3} \delta_{a,a'-1}, q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_3}{x_1} \delta_{a,a'-1} \right. \\ & \left. , q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{y^2}{x_1 x_4} \delta_{a,a'-1}, \frac{x_1 x_3}{y^2} \delta_{a,a'-1} q^a \frac{x_4}{x_1} \delta_{a,a'-1}, x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1}, q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1} \right\}, \end{aligned}$$

corresponding to the interactions between x_1 and x_3 . In comparison to L-operators for two-dimensional, and for three-dimensional, vertex models, additional possible interactions from L-operators for the Ising-type model are also dependent upon,

$$\begin{aligned} & \text{span}_{a,a' \in \mathbf{Z}^2} \left\{ x_1 x_2 \delta_{a,a'+1} \frac{y^2}{x_1 x_3} \delta_{a,a'-1}, q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1}, x_1 x_3 \delta_{a,a'+1} q^a \frac{x_4}{x_1}, q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} \frac{y^2}{x_1 x_4} \delta_{a,a'+1}, x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} \right. \\ & \left. , q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1}, q^a \frac{x_1}{x_3} \delta_{a,a'+1} q^{-a} \frac{x_4}{x_1} \delta_{a,a'+1}, \frac{x_1 x_3}{y^2} \delta_{a,a'-1} q^a \frac{x_4}{x_1} \delta_{a,a'-1} \right\}. \end{aligned}$$

Besides δ factors appearing next to the product, $x_1 x_3$, of two positions, or to the ratio, $\frac{x_1}{x_3}$, of two positions, q parameters parallel similar factors appearing in L-operators of vertex models,

$$\begin{aligned} & q^{D_k^j}, \\ & q^{-D_k^j - D_{k+1}^j}, \\ & q^{-D_k^j + D_{k+1}^j}, \\ & q^{-D_k^j}, \end{aligned}$$

in the second, first, and third, columns of the representation,

$$\begin{bmatrix} q^{D_k^j} & q^{-2} a_k^j q^{-D_k^j - D_{k+1}^j} \xi^{s-s_k^j} & a_k^j a_{k+1}^j q^{-D_k^j - 3D_{k+1}^j} \xi^{s-s_k^j - s_{k+1}^j} \\ \left(a_k^j \right)^\dagger q^{D_k^j} \xi^{s_k^j} & q^{-D_k^j + D_{k+1}^j} - q^{-2} q^{D_k^j - D_{k+1}^j} \xi^s & -a_k^j q^{D_k^j - 3D_{k+1}^j} \xi^{s-s_k^j} \\ 0 & a_k^\dagger q^{D_k^j} \xi^{s_k^j} & q^{-D_k^j} \end{bmatrix},$$

where the q parameters are raised to operators that are defined over the tensor product of two vector spaces, in addition to mappings ξ into a unital associative algebra. As $N \rightarrow +\infty$, asymptotically the Ising-type model transfer matrix is also dependent upon several additional spanning sets rather than the first two introduced above, one of which is,

$$\begin{aligned} \text{span}_{a,a' \in \mathbf{Z}^2} \left\{ x_1 x_3 \delta_{a,a'+1} x_1 x_4 \delta_{a,a'+1}, x_1 x_4 \delta_{a,a'+1} x_1 x_5 \delta_{a,a'+1}, q^{-a} \frac{x_1}{x_4} \delta_{a,a'-1} q^a \frac{x_1}{x_5} \delta_{a,a'+1}, x_1 x_3 \delta_{a,a'+1} x_1 x_6 \delta_{a,a'+1} \right. \\ \left. , q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} q^a \frac{x_1}{x_6} \delta_{a,a'+1}, x_1 x_3 \delta_{a,a'+1} q^{-a} \frac{x_1}{x_5} \delta_{a,a'-1}, \dots \right\}. \end{aligned}$$

In comparison to action-angle variables of the 20-vertex model, Φ^{20-V} , for which,

$$\{ \Phi^{20-V}, \Phi^{20\bar{0}-V} \} \neq 0,$$

stronger notions of Integrability follow from the vanishing Poisson bracket of the action-angle variables for the Ising type model with its complex conjugate. Terms appearing in the first, and second, spanning sets over elements of the Ising-type L-operator directly represent terms appearing in the Poisson bracket,

$$\{ \cdot, \cdot \},$$

with,

$$\begin{aligned} \left\{ \sum_{n \in \mathbf{Z}} \left[\frac{y^2}{x_1 x_{N-1}} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_{N-1}} \delta_{a,a'-1} + q^a \frac{X_{N-1}}{x_1} \delta_{a,a'+1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + \frac{y^2}{x_1 x_{N-2}} \delta_{a,a'-1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} \right. \right. \\ \left. \left. + q^a \frac{x_{N-2}}{x_1} \delta_{a,a'-1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} \right], \cdot \right\}. \end{aligned}$$

The second argument of the Poisson bracket, designed with \cdot , could equal,

$$\begin{aligned} & x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1}, \\ & q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1}, \\ & q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{y^2}{x_1 x_4} \delta_{a,a'-1}, \\ & q^a \frac{x_1}{x_2} \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1}, \\ & \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1}. \end{aligned}$$

Over the integers, additional contributions emerge by replacing the first entry of the Poisson bracket above with,

$$\sum_{N \in \mathbf{Z}} \left[q^{-a} \frac{x_1}{x_{N-2}} \delta_{a,a'-1} \frac{y^2}{x_1 x_N} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} q^{-a} \frac{x_N}{x_1} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} q^{-a} \frac{x_N}{x_1} \delta_{a,a'+1} + \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} q^a \frac{x_N}{x_1} \delta_{a,a'-1} \right].$$

Such computations parallel those of the 20-vertex model, in which the product of the vertex-type L-operator is proportional to,

$$(a_k^j)^\dagger q^{D_k^j} \xi^{s_k^j} q^{D_k^{j+1}} + q^{-D_k^j + D_{k+1}^j} (a_k^{j+1})^\dagger q^{D_k^{j+1}} \xi^{s_k^{j+1}} + (a_{k+1}^j)^\dagger q^{D_{k+1}^j} \xi^{s_{k+1}^j} (a_k^{j+1})^\dagger q^{D_k^{j+1}} \xi^{s_k^{j+1}}.$$

In the next section, we compute the first order approximation for products of L-operators from the Ising-type model. Albeit being defined pointwise, as more terms are taken in products of L-operators

2 Computations with the Ising type L-operator

Below, we obtain the entries of the two-dimensional representation for the transfer matrix. By making use of several observations from previous works of the author, [41,42,45,46,47,48], one may conclude that the Ising type model formulation of the 6-vertex model is exactly solvable.

Lemma 1 (*product representation of the transfer matrix from two, and three, L-operators of the Ising-type model*). Given the L-operator defined in the previous section for the Ising-type formulation of the 6-vertex model, in order to asymptotically approximate the transfer matrix in infinite volume, one has the following two representations, the first two of which are,

$$\begin{bmatrix} \mathcal{I}_1^1 & \mathcal{I}_1^2 \\ \mathcal{I}_1^3 & \mathcal{I}_1^4 \end{bmatrix}, \begin{bmatrix} \mathcal{I}_2^1 & \mathcal{I}_2^2 \\ \mathcal{I}_2^3 & \mathcal{I}_2^4 \end{bmatrix},$$

and the third of which is,

$$\begin{bmatrix} \mathcal{I}_1^1 \mathcal{I}_2^1 + \mathcal{I}_1^3 \mathcal{I}_2^3 & \mathcal{I}_1^2 \mathcal{I}_2^2 + \mathcal{I}_1^4 \mathcal{I}_2^4 \\ \mathcal{I}_1^1 \mathcal{I}_2^3 + \mathcal{I}_1^3 \mathcal{I}_2^4 & \mathcal{I}_1^2 \mathcal{I}_2^3 + \mathcal{I}_1^4 \mathcal{I}_2^4 \end{bmatrix}$$

where the entries of each representation equal,

$$\begin{aligned} & x_1 x_3 \delta_{a,a'+1} x_1 x_{N-1} \delta_{a,a'+1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + x_1 x_3 \delta_{a,a'+1} x_1 x_{N-1} \delta_{a,a'+1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} \\ & + q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1}, \\ & x_1 x_3 \delta_{a,a'+1} q^{-a} \frac{x_1}{x_{N-1}} \delta_{a,a'-1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + x_1 x_3 \delta_{a,a'+1} q^{-a} \frac{x_1}{x_{N-1}} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} \\ & + q^{-a} \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + q^{-a} \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1}, \\ & x_1 x_3 \delta_{a,a'+1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + x_1 x_3 \delta_{a,a'+1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} \\ & + q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} x_1 x_2 \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} + q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1}, \\ & x_1 x_3 \delta_{a,a'+1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_1}{x_N} \delta_{a,a'-1} + x_1 x_3 \delta_{a,a'+1} q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} \\ & + q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_1}{x_N} \delta_{a,a'-1} + q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1}. \end{aligned}$$

$$\begin{aligned}
& \left[q^a \frac{x_1}{x_2} \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} + \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1} \right] \left[q^a \frac{x_1}{x_3} \delta_{a,a'+1} q^{-a} \frac{x_4}{x_1} \delta_{a,a'+1} + \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \frac{y^2}{x_1 x_4} \delta_{a,a'+1} \right], \\
& \left[x_1 x_2 \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} + q^{-a} \frac{x_1}{x_2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1} \right] \left[q^a \frac{x_1}{x_3} \delta_{a,a'+1} \frac{y^2}{x_1 x_4} \delta_{a,a'-1} + \frac{x_1 x_3}{y^2} \delta_{a,a'-1} q^a \frac{x_4}{x_1} \delta_{a,a'-1} \right], \\
& \left[q^a \frac{x_1}{x_2} \delta_{a,a'+1} q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} + \frac{x_1 x_2}{y^2} \delta_{a,a'-1} \frac{y^2}{x_1 x_3} \delta_{a,a'+1} \right] \left[q^a \frac{x_1}{x_3} \delta_{a,a'+1} q^{-a} \frac{x_4}{x_1} \delta_{a,a'+1} + \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \frac{y^2}{x_1 x_4} \delta_{a,a'+1} \right], \\
& \left[\frac{y^2}{x_1 x_2} \delta_{a,a'-1} x_1 x_3 \delta_{a,a'+1} + q^{-a} \frac{x_2}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_3} \delta_{a,a'+1} \right] \left[\frac{y^2}{x_1 x_3} \delta_{a,a'-1} x_1 x_4 \delta_{a,a'+1} + q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_4} \delta_{a,a'+1} \right], \\
& \left[\frac{y^2}{x_1 x_3} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_4} \delta_{a,a'-1} + q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} \frac{x_1 x_4}{y^2} \delta_{a,a'-1} \right] \left[\frac{y^2}{x_1 x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_3} \delta_{a,a'+1} + q^a \frac{x_2}{x_1} \delta_{a,a'-1} \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \right], \\
& \left[\frac{y^2}{x_1 x_2} \delta_{a,a'-1} x_1 x_3 \delta_{a,a'+1} + q^{-a} \frac{x_2}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_3} \delta_{a,a'+1} \right] \left[\frac{y^2}{x_1 x_3} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_4} \delta_{a,a'-1} + q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} \frac{x_1 x_4}{y^2} \delta_{a,a'-1} \right], \\
& \left[\frac{y^2}{x_1 x_2} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_3} \delta_{a,a'-1} + q^{-a} \frac{x_2}{x_1} \delta_{a,a'+1} \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \right] \left[q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_4} \delta_{a,a'+1} + \frac{y^2}{x_1 x_4} \delta_{a,a'+1} \frac{x_1 x_4}{y^2} \delta_{a,a'-1} \right], \\
& \left[\frac{y^2}{x_1 x_2} \delta_{a,a'-1} q^a \frac{x_1}{x_3} \delta_{a,a'+1} + q^a \frac{x_2}{x_1} \delta_{a,a'-1} \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \right] \left[\frac{y^2}{x_1 x_3} \delta_{a,a'-1} q^{-a} \frac{x_1}{x_4} \delta_{a,a'-1} + q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} \frac{x_1 x_4}{y^2} \delta_{a,a'-1} \right], \\
& \left[q^{-a} \frac{x_2}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_3} \delta_{a,a'+1} + \frac{y^2}{x_1 x_3} \delta_{a,a'+1} \frac{x_1 x_3}{y^2} \delta_{a,a'-1} \right] \left[q^{-a} \frac{x_3}{x_1} \delta_{a,a'+1} q^a \frac{x_1}{x_4} \delta_{a,a'+1} + \frac{y^2}{x_1 x_4} \delta_{a,a'+1} \frac{x_1 x_4}{y^2} \delta_{a,a'-1} \right],
\end{aligned}$$

from which we conclude the argument. \square

Equipped with the two representations from the transfer matrix in the previous result, below we conclude that the Ising-type model is integrable. Following previous works of the author previously cited in other works, we demonstrate how arguments from each one of the Poisson bracket can be obtained, from which the existence of suitable action-angle variables can be inferred from the Poisson structure.

Lemma 1 (*action-angle variables for the Ising-type formulation of the 6-vertex model*). The Ising-type model is exactly solvable.

Proof of Lemma 1. It suffices to demonstrate that several computations with the Poisson bracket can be performed. As mentioned earlier in the previous section, the arguments that enter within each Poisson bracket take the form,

$$\begin{aligned}
& \sum_{N \in \mathbf{Z}} \left[x_1 x_{N-1} \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} \left[x_1 x_{N-1} \delta_{a,a'+1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + x_1 x_{N-2} \delta_{a,a'+1} \right. \right. \\
& \qquad \qquad \qquad \left. \left. \times x_1 x_N \delta_{a,a'+1} + q^{-a} \frac{x_1}{x_{N-2}} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'-1} \right] \right], \\
& \sum_{N \in \mathbf{Z}} \left[x_1 x_{N-1} \delta_{a,a'+1} x_1 x_N \delta_{a,a'+1} \left[x_1 x_{N-1} \delta_{a,a'+1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + x_1 x_{N-2} \delta_{a,a'+1} \right. \right. \\
& \qquad \qquad \qquad \left. \left. \times q^{-a} \frac{x_1}{x_N} \delta_{a,a'+1} + q^{-a} \frac{x_1}{x_{N-2}} \delta_{a,a'-1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} \right] \right],
\end{aligned}$$

for the next group of terms, and,

$$\begin{aligned} & \sum_{N \in \mathbf{Z}} \left[x_1 x_{N-2} \delta_{a,a'+1} q^{-a} \frac{x_1}{x_{N+1}} \delta_{a,a'-1} \left[q^{-a} \frac{x_1}{x_{N-1}} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} + \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + q^{-a} \frac{x_1}{x_{N-2}} \delta_{a,a'-1} \right. \right. \\ & \qquad \qquad \qquad \left. \left. \times q^a \frac{x_1}{x_N} \delta_{a,a'+1} + \frac{x_1 x_{N-2}}{y^2} \delta_{a,a'-1} \frac{x_1 x_{N+1}}{y^2} \delta_{a,a'-1} \right] \right], \\ & \sum_{N \in \mathbf{Z}} \left[q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} \frac{x_1 x_{N+1}}{y^2} \delta_{a,a'-1} \left[x_1 x_{N-1} \delta_{a,a'+1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N+1}} \delta_{a,a'+1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + x_1 x_{N-2} \delta_{a,a'+1} \right. \right. \\ & \qquad \qquad \qquad \left. \left. \times x_1 x_N \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-2}} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} \right] \right], \\ & \sum_{N \in \mathbf{Z}} \left[q^a \frac{x_1}{x_{N-1}} \delta_{a,a'+1} \frac{x_1 x_{N+1}}{y^2} \delta_{a,a'-1} \left[q^{-a} \frac{x_1}{x_{N-1}} \delta_{a,a'-1} q^a \frac{x_1}{x_N} \delta_{a,a'+1} + \frac{x_1 x_{N-1}}{y^2} \delta_{a,a'-1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} + x_1 x_{N-2} \delta_{a,a'+1} \right. \right. \\ & \qquad \qquad \qquad \left. \left. \times q^a \frac{x_1}{x_N} \delta_{a,a'+1} + q^a \frac{x_1}{x_{N-2}} \delta_{a,a'+1} \frac{x_1 x_N}{y^2} \delta_{a,a'-1} \right] \right], \end{aligned}$$

for the last group of terms. Straightforward computations with the Poisson bracket establish that the desired integrable, and exactly solvable, properties for the Ising type model hold, from which we conclude the argument. \square

3 References

- [1] Amico, L., Frahm, H., Osterloh, A., Ribeiro, G.A.P. Integrable spin-boson models descending from rational six-vertex models. *Nucl. Phys. B.* **787**: 283-300 (2007).
- [2] Alcaraz, F.C., Lazo, M.J. Exactly solvable interacting vertex models. *J. Stat. Mech.* P08008 (2007).
- [3] Bazhanov, V.V., Sergeev, S.S. An Ising-type formulation of the six-vertex model. *Nucl. Phys. B.* **986**, 1160055 (2023).
- [4] Baxter, R.J. Exactly Solved Models in Statistical Mechanics. Dover Publishing.
- [5] Boos, H., et al. Universal R-matrix and functional relations. *Rev. Math. Phys.* **26**: 143005 (2014).
- [6] Boos, H., Gohmann, F., Klumper, A., Nirov, K.S., Razumov, A.V.. Exercises with the universal R-matrix.
- [7] Colomo, F., Giulio, G.D., Pronko, A.G. Six-vertex model on a finite lattice: Integral representations for nonlocal correlation functions. *Nuclear Physics B* **972**: 115535 (2021).
- [8] Colomo, F., Pronko, A.G. The Arctic Circle Revisited. *Contemp. Math.* **458**: 361-376 (2008).
- [9] Deguchi, T. Introduction to solvable lattice models in statistical and mathematical physics. *CRC Press*: 9780429137891 (2003).
- [10] Duminil-Copin, H., Karrila, A., Manolescu, I., Oulamara, M. Delocalization of the height function of the six-vertex model. *J. Eur. Math. Soc.* (2024).
- [11] Duminil-Copin, H., Kozłowski, K.K., Krachun, D. et al. On the Six-Vertex Model's Free Energy. *Commun. Math. Phys.* 395, 1383–1430 (2022). <https://doi.org/>
- [12] de Vega, H.J. Bethe Ansatz and Quantum Groups. *LPTHE* **93/17** (1992).
- [13] de Vega, H.J. Boundary K-matrices for the XYZ, XXZ and XXX spin chains. *LPTHE-PAR* **93/29** (1993).
- [14] de Vega, H.J., Ruiz, A.G. Boundary K-matrices for the six vertex and the $n(2n - 1) A_{n-1}$ vertex models. *J. Phys. A: Math. Gen.* **26** (1993).

- [15] Di Francesco, Philippe. Twenty Vertex model and domino tilings of the Aztec triangle. *the electronic journal of combinatorics* **28**(4), (2021).
- [16] Faddeev, L.D., Takhtajan, L. A. *Hamiltonian Methods in the Theory*
- [17] Frahm, H., Seel, A. The staggered six-vertex model: Conformal invariance and corrections to scaling. *Nucl. Phys. B.* **879**: 382-406 (2014).
- [18] Garbali, A., de Gier, J., Mead, W., Wheeler, M. Symmetric functions from the six-vertex model in half-space. *arXiv: 2312.14348* (2023).
- [19] Glazman, A., Peled, R. On the transition between the disordered and antiferroelectric phases of the 6-vertex model. *Electron. J. Probab.* **28**: 1-53 (2023). DOI: 10.1214/23-EJP980.
- [20] Gohmann, F. Bethe ansatz. *arXiv: 2309.02008* (2023).
- [21] Gorbounov, V., Korff, C. Quantum integrability and generalised quantum Schubert calculus. *Adv. Math.* **313**: 282-356 (2017).
- [22] Gorbounov, V., Korff, C., Stroppel, C. Yang-Baxter Algebras as Convolution Algebras: The Grassmannian case. *Russian Mathematical Surveys* **75** (2020).
- [23] Ikhlef, Y., Jacobsen, J., Saleur, H. A staggered six-vertex model with non-compact continuum limit. *Nucl. Phys. B.* **789**(3): 483-524 (2008).
- [24] Keating, D., Reshetikhin, N., Sridhar, A. Integrability of Limit Shapes of the Inhomogeneous Six Vertex Model. *Commun. Math. Phys.* **391**, 1181–1207 (2022). [https : //doi.org/10.1007/s00220-022-04500-0](https://doi.org/10.1007/s00220-022-04500-0)
- [25] Kitanine, N., et al. Thermodynamic limit of particle-hole-form factors in the massless XXZ Heisenberg chain. *J. Stat. Mech.* *P05028* (2011).
- [26] Kitanine, N., et al. On correlation functions of integrable models associated with the six-vertex R-matrix. *J. Stat. Mech.* *P01022* (2007).
- [27] Kitanine, N., et al. On the spin-spin correlation functions of the XXZ spin- $\frac{1}{2}$ infinite chain. *J. Phys. A: Math. Gen.* **38** (2005).
- [28] Kozłowski, K.K. Riemann-Hilbert approach to the time-dependent generalized sine kernel. *Adv. Theor. Math. Phys.* **15**: 1655-1743 (2011).
- [29] Korff, C., McCoy, B.M. Loop symmetry of integrable vertex models at roots of unity. *Nucl. Phys. B.* **618**: 551-569 (2001).
- [30] Lieb, E. H. The Residual Entropy of Square Ice, *Phys Rev.* **162** 162-172 (1967).
- [31] Lamers, J. Introduction to quantum integrability. *10th Modave Summer School in Mathematical Physics* (2015).
- [32] Motegi, K. Symmetric functions and wavefunctions of the XXZ-type six-vertex models and elliptic Felderhof models by Izergin-Korepin analysis. *J. Math. Phys.* **59**: 053505 (2018).
- [33] Naprienko, S. Free fermionic Schur functions. *Adv. Math.* **436**: 109413 (2024).
- [34] Pauling, L. J. *Am. Chem. Soc.* **57**, 2680 (1935)
- [35] Reshetikhin, N., Sridhar, A. Integrability of limits shapes of the six-vertex model. *Comm. Math. Phys.* **356**: 535-565 (2017).
- [36] Rigas, P. From logarithmic delocalization of the six-vertex height function under sloped boundary conditions to weakened crossing probability estimates for the Ashkin-Teller, generalized random-cluster, and (q_σ, q_τ) -cubic models, *arXiv:2211.14934* (2022).

- [37] Rigas, P. Renormalization of crossing probabilities in the dilute Potts model. *arXiv: 2211.10979 v2* (2022).
- [38] Rigas, P. The phase transition for the Gaussian free field is sharp. *arXiv: 2307.12925* (2023).
- [39] Rigas, P. Phase transition of the long range Ising model in lower dimensions, for $d < \alpha \leq d + 1$, with a Peierls' argument. *arXiv: 2309.07943* (2023).
- [40] Rigas, P. Operator formalism for discreteley holomorphic parafermions of the two-color Ashkin-Teller, loop $O(1)$, staggered eight-vertex, odd eight-vertex, and abelian sandpile models. *arXiv: 2310.08212* (2023).
- [41] Rigas, P. Poisson structure and Integrability of a Hamiltonian flow for the inhomogeneous six-vertex model. *arXiv: 2310.15181* (2023).
- [42] Rigas, P. Open boundary conditions of the $D_3^{(2)}$ spin chain and sectors of conformal field theories. *arXiv: 2310.18499* (2023).
- [43] Rigas, P. Scaling limit of the triangular prudent walk. *arXiv: 2312.16236* (2023).
- [44] Rigas, P. Eigenvalue attraction in open quantum systems, biophysical systems, and Parity-time symmetric materials. *arXiv: 2309.07943* (2023).
- [45] Rigas, P. Quantum-inverse scattering for the 20-vertex model up to Dynkin automorphism: crossing probabilities, 3D Poisson structure, triangular height functions, weak integrability. *arXiv 2407.11066* (2024).
- [46] Rigas, P. Approximability of Poisson structures for the 4-vertex model, and higher-spin XXX chain, and Yang-Baxter algebras. *arXiv 2411.15188* (2024).
- [47] Rigas, P. The emptiness formation probability, and nonlocal representations of correlations, for the 20-vertex model. *arXiv 2409.05309* (2024).
- [48] Rigas, P. Intertwining vectors, and Boltzmann weight matrices, of a Solid-on-Solid model from the 20-vertex model. *arXiv 2412.04515* (2024).