

Vitalii Shpakivskyi

Vieta's Formulas for Quaternionic Polynomials

Abstract. The paper presents analogues of some Vieta formulas for quaternionic polynomials of the form $R_n(w) = \sum_{m=0}^n A_m x^m$. A criterion for the sphericity of the root (zero) of a polynomial $R_n(w)$ is established.

Key words. Quaternionic polynomials, Vieta's formulas, Basic polynomial, Spherical root.

Mathematics Subject Classification. Primary 30G35, Secondary 11R52.

1 Introduction

The results of this article were presented by the author in 2007 as a bachelor's thesis at Zhytomyr State University [1]. But have not been published until now.

The pioneer of the consideration of algebraic quaternion equations was Ivan Niven [2]. In the mentioned paper, I. Niven also proposed a method for computing the roots of one-side polynomials. In collaboration with S. Eilenberg, he was among the first to establish the fundamental theorem of algebra for quaternions [3]. They proved that every quaternion polynomial $R_n(w)$ has at least one zero. Later, B. Beck [4] proved that if the number of zeros of a polynomial of degree n is finite, then there are exactly n zeros counting possible repetition. Beck's result was re-proven in the paper [5] by A. Pogorui and M. Shapiro. In the paper [6], the authors proved the fundamental theorem of algebra for polynomials with coefficients in the skew field of quaternions and in the division algebra of octonions.

In [7], a new characterization of the types of zeros is provided, along with an algorithm for determining all zeros, including their types, without relying on an iterative process requiring convergence. In [8], a method for computing the zeros of a quaternionic polynomial $R_n(w)$ is developed, which is fundamentally based on Niven's algorithm.

In [9], a Weierstrass-like method was proposed for simultaneously finding all the zeros of unilateral quaternionic polynomials. The paper includes a convergence analysis and several numerical examples demonstrating the method's

performance. As established in [10] and [5] (cf. [11]), the zero set of a canonical quaternionic polynomial consists of p points and s two-dimensional spheres, where $p + 2s$ does not exceed the algebraic degree of the polynomial. The aim of [12] was to refine and extend the results on the roots of quaternionic polynomials initially presented in [10].

The thesis [13] explores recent work on finding the zeros of quaternionic polynomials and also the automorphisms of the zeros. The automorphisms of the zeros of quaternionic polynomials act on the zeros of the polynomial thereby preserving their algebraic properties. In [14], using hybrid Bézout matrices, necessary and sufficient conditions were established for a quaternionic polynomial to possess a complex root, a spherical root, or a complex isolated root. Work [15] is devoted to the study a method for approximating zeros of a quaternion polynomial based on the approximate complex zeros of a real polynomial. Moreover, it is showed how to derive bounds on the norm of zeros of a quaternion polynomial.

The paper [16] focuses on examining the properties of zeros in certain special quaternionic polynomials with restricted coefficients, where the real and imaginary components of the coefficients satisfy specific inequalities. Additionally, the articles [17], [18], and [19] explore various types of quadratic quaternionic equations.

Recently, two works staging the analogue of Vieta's formulas in Clifford geometric algebras have appeared. In the papers [20, 21], it was discussed a generalization of Vieta theorem (Vieta's formulas) to the case of Clifford geometric algebras. Also it was compared the generalized Vieta's formulas with the ordinary Vieta's formulas for characteristic polynomial containing eigenvalues.

In Section 3 of the present paper we establish analogues of some Vieta formulas for quaternionic polynomials of the form $R_n(w)$. In Section 4, a criterion for the sphericity of the root of a polynomial $R_n(w)$ is also established. The main research tool is the *basic polynomial* introduced in paper [5].

2 Preliminary

For a real quaternion we use the notations

$$\mathbb{H}(\mathbb{R}) = \{w = x_0 + x_1i + x_2j + x_3k : x_0, x_1, x_2, x_3 \in \mathbb{R}\},$$

where the following equalities are true for the imaginary units i, j, k :

$$i^2 = j^2 = k^2 = -1, \quad ij = -ji = k, \quad jk = -kj = i, \quad ki = -ik = j.$$

The number $\text{Sc}(w) := x_0$ is called the *scalar part* of the quaternion w ; $x_1i + x_2j + x_3k$ is called the *vector part* of w and is denoted by $\text{Vec}(w)$;

$$|w| = \sqrt{x_0^2 + x_1^2 + x_2^2 + x_3^2}$$

is called the *modulus* of w . Quaternion $\bar{w} := x_0 - x_1i - x_2j - x_3k$ is called the *conjugate* with respect to quaternion w . *Right (left) polynomial* [5] of quaternionic variable is called polynomial of the form

$$R_n(w) := \sum_{m=0}^n A_m w^m \quad \left(L_n(w) := \sum_{m=0}^n w^m A_m \right), \quad (1)$$

where $A_m, w \in \mathbb{H}(\mathbb{R})$.

The polynomial of the complex variable

$$\begin{aligned} F_{2n}^*(z) &:= \bar{A}_0 A_0 + (\bar{A}_0 A_1 + \bar{A}_1 A_0)z \\ &+ (\bar{A}_0 A_2 + \bar{A}_1 A_1 + \bar{A}_2 A_0)z^2 + \dots + \bar{A}_n A_n z^{2n} \end{aligned} \quad (2)$$

is called a *basic polynomial* for polynomial $R_n(x)$, and polynomial

$$G_{2n}^*(z) := A_0 \bar{A}_0 + (A_1 \bar{A}_0 + A_0 \bar{A}_1)z + (A_2 \bar{A}_0 + A_1 \bar{A}_1 + A_0 \bar{A}_2)z^2 + \dots + A_n \bar{A}_n z^{2n}$$

is called a *basic polynomial* for polynomial $L_n(x)$.

In the paper [5] it is proved that complex roots z of basic polynomial are related to the roots w of quaternionic polynomial by the following equalities

$$\text{Re } z = \text{Sc}(w), \quad |\text{Im } z| = |\text{Vec}(w)|. \quad (3)$$

The quaternion $w = x_0 + \text{Vec}(w)$ is called the *spherical root* of the polynomial (1) if any quaternion of the form $w_1 := x_0 + \text{Vec}(w_1)$ such that $|\text{Vec}(w)| = |\text{Vec}(w_1)|$ is also a root of this polynomial.

3 Analogues of some Viète formulas

Theorem 1. [5]. *The set of roots w of the quaternionic polynomial $R_n(w)$ consists of isolated points and spheres*

$$\{w : \text{Sc}(w) = x_0, \quad |\text{Vec}(w)| = r\}.$$

Moreover, the number of the isolated points together with the double number of the spheres is less or equal to n .

The multiplicity of a point root w of the polynomial $R_n(w)$ is called the multiplicity of the corresponding complex root z of the basic polynomial $F_{2n}^(z)$.*

The *multiplicity of a spherical root* is called the multiplicity of the corresponding complex root of the basic polynomial divided by two. Then the polynomial $R_n(w)$ has exactly n roots.

The *set of roots* of polynomial $R_n(w)$ is called the set $\{w_1, w_2, \dots, w_n\}$ which consists of all its isolated roots and taken any two representatives from each spherical root.

The *scalar product* of quaternions is denoted to similarly as the scalar product of vectors $(w, \vartheta) := \sum_{m=0}^3 x_m y_m$, where $w = x_0 + x_1 i + x_2 j + x_3 k$, $\vartheta = y_0 + y_1 i + y_2 j + y_3 k$.

Lemma 1. *For any $w, \vartheta \in \mathbb{H}(\mathbb{R})$ the following equality is true*

$$w\bar{\vartheta} + \vartheta\bar{w} = \bar{w}\vartheta + \bar{\vartheta}w = 2(w, \vartheta).$$

Proof. We first convert the expression

$$\begin{aligned} w\bar{\vartheta} + \vartheta\bar{w} &= (x_0 + \text{Vec}(w))(y_0 - \text{Vec}(\vartheta)) + (y_0 + \text{Vec}(\vartheta))(x_0 - \text{Vec}(w)) \\ &= x_0 y_0 - x_0 \text{Vec}(\vartheta) + y_0 \text{Vec}(w) - \text{Vec}(w) \text{Vec}(\vartheta) \\ &\quad + y_0 x_0 - y_0 \text{Vec}(\vartheta) + x_0 \text{Vec}(\vartheta) - \text{Vec}(\vartheta) \text{Vec}(w) \\ &= 2x_0 y_0 - \text{Vec}(w) \text{Vec}(\vartheta) - \text{Vec}(\vartheta) \text{Vec}(w) = \bar{w}\vartheta + \bar{\vartheta}w. \end{aligned}$$

Then, we consider the sum

$$\begin{aligned} \text{Vec}(w) \text{Vec}(\vartheta) + \text{Vec}(\vartheta) \text{Vec}(w) &= (-x_1 y_1 + x_1 y_2 k - x_1 y_3 j - x_2 y_1 k - x_2 y_2 \\ &\quad + x_2 y_3 i + x_3 y_1 j - x_3 y_2 i - x_3 y_3) + (-y_1 x_1 + y_1 a_2 k - y_1 x_3 j - y_2 x_1 k - x_2 y_2 \\ &\quad + y_2 x_3 i + y_3 x_1 j - y_3 x_2 i - y_3 x_3) = -2x_1 y_1 - 2x_2 y_2 - 2x_3 y_3. \end{aligned}$$

Using the two previous equalities, we obtain the statement of lemma.

Theorem 2. *For any polynomial $R_n(w)$ and for the set of its roots, taking into account their multiplicity, the following equality is true*

$$\prod_{m=1}^n |w_m| = \frac{|A_0|}{|A_n|}. \quad (4)$$

Proof. Let z_1, z_2, \dots, z_{2n} be the roots of basic polynomial. By definition (2)

$$F_{2n}^*(z) := |A_n|^2 z^{2n} + \dots + (\bar{A}_1 A_0 + \bar{A}_0 A_1) z + |A_0|^2.$$

Since the degree of the basic polynomial is $2n$, then it has exactly $2n$ roots. Let

$$z_2 = \bar{z}_1, \quad z_4 = \bar{z}_3, \quad z_{2n} = \bar{z}_{2n-1}.$$

Using the Vieta formula for the polynomial F_{2n}^* , we obtain

$$\prod_{m=1}^{2n} z_m = \frac{|A_0|^2}{|A_n|^2}.$$

Let

$$z_1 = \alpha_1 + i\beta_1, z_2 = \bar{z}_1 = \alpha_1 - i\beta_1, z_3 = \alpha_3 + i\beta_3, z_4 = \bar{z}_3 = \alpha_3 - i\beta_3, \dots, \\ z_{2n-1} = \alpha_{2n-1} + i\beta_{2n-1}, z_{2n} = \bar{z}_{2n-1} = \alpha_{2n-1} - i\beta_{2n-1}.$$

Then

$$(\alpha_1 + i\beta_1)(\alpha_1 - i\beta_1) \cdot \dots \cdot (\alpha_{2n-1} + i\beta_{2n-1})(\alpha_{2n-1} - i\beta_{2n-1}) = \frac{|A_0|^2}{|A_n|^2}, \quad (5)$$

whence

$$(\alpha_1^2 + \beta_1^2)(\alpha_3^2 + \beta_3^2) \cdot \dots \cdot (\alpha_{2n-1}^2 + \beta_{2n-1}^2) = \frac{|A_0|^2}{|A_n|^2}. \quad (6)$$

Using system (3), we obtain the equalities

$$\alpha^2 = \text{Sc}^2(w), \quad |\beta|^2 = |\text{Vec}(w)|^2.$$

Then equality (6) can be rewritten in the form

$$(\text{Sc}^2(w_1) + |\text{Vec}(w_1)|^2) (\text{Sc}^2(w_2) + |\text{Vec}(w_2)|^2) \\ \dots \times (\text{Sc}^2(w_n) + |\text{Vec}(w_n)|^2) = \frac{|A_0|^2}{|A_n|^2}$$

and

$$|w_1|^2 \cdot |w_2|^2 \cdot \dots \cdot |w_n|^2 = \frac{|A_0|^2}{|A_n|^2}.$$

The theorem is proved.

Theorem 3. *Let the conditions of Theorem 2 are satisfied. Then*

$$\sum_{m=1}^n \text{Sc}(w_m) = -\frac{(A_n, A_{n-1})}{|A_n|^2}.$$

Proof. Let us write the basic polynomial in the form

$$F_{2n}^*(z) = \sum_{m=0}^{2n} B_m z^m.$$

Let $z_2 = \bar{z}_1, z_4 = \bar{z}_3, \dots, z_{2n} = \bar{z}_{2n-1}$ be the roots of basic polynomial. Since $z + \bar{z} = 2\text{Re } z$, then using Viète formula for the polynomial F_{2n}^* , we obtain

$$\sum_{m=1}^n z_m = -\frac{B_{2n-1}}{B_{2n}}$$

hence

$$2(\operatorname{Re} z_1 + \operatorname{Re} z_3 + \dots + \operatorname{Re} z_{2n-1}) = -\frac{B_{2n-1}}{B_{2n}}. \quad (7)$$

Since for two complex conjugate roots of the basic polynomial corresponds one quaternionic root and equality (3) is true, equality (7) rewritten in the form

$$2 \sum_{m=1}^n \operatorname{Sc}(w_m) = -\frac{B_{2n-1}}{B_{2n}},$$

where $B_{2n-1} = \bar{A}_{n-1}A_n + \bar{A}_nA_{n-1}$, $B_{2n} = |A_n|^2$. By the Lemma 1

$$B_{2n-1} = \bar{A}_{n-1}A_n + \bar{A}_nA_{n-1} = 2(A_n, A_{n-1}).$$

Then

$$2 \sum_{m=1}^n \operatorname{Sc}(w_m) = -\frac{2(A_n, A_{n-1})}{|A_n|^2}.$$

The theorem is proved.

Lemma 2. *Let performed the conditions of Theorem 2 and $A_0 \neq 0$. Then none of the roots of the corresponding basic polynomial is not zero.*

Proof. Suppose the contrary, i. e. let exists a number $z = 0$ such that

$$F_{2n}^*(0) = |A_n|^2 \cdot 0^{2n} + \dots + (\bar{A}_1A_0 + \bar{A}_0A_1) \cdot 0 + |A_0|^2 = 0.$$

Since $A_0 \neq 0$, then $|A_0|^2 \neq 0$. This contradiction proves the Lemma.

Theorem 4. *Let performed the conditions of Theorem 2 and $A_0 \neq 0$. Then the following equation is true*

$$\sum_{m=1}^n \frac{\operatorname{Sc}(w_m)}{|w_m|^2} = -\frac{(A_1, A_0)}{|A_0|^2}.$$

Proof. Consider the Vieta equality

$$z_1 z_2 \cdot \dots \cdot z_{2n-1} + z_1 z_2 \cdot \dots \cdot z_{2n-2} z_{2n} + \dots + z_1 z_3 \cdot \dots \cdot z_{2n} + z_2 z_3 \cdot \dots \cdot z_{2n} = -\frac{B_1}{B_{2n}},$$

where B_1, B_{2n} are corresponding coefficients of the basic polynomial. Using equality (5) and relation $z \cdot \bar{z} = |z|^2$, we obtain

$$\begin{aligned} & |z_1|^2 \cdot |z_3|^2 \cdot \dots \cdot |z_{2n-3}|^2 (\alpha_{2n-1} + i\beta_{2n-1}) + |z_1|^2 \cdot |z_3|^2 \cdot \dots \cdot |z_{2n-3}|^2 (\alpha_{2n-1} - i\beta_{2n-1}) \\ & \dots + |z_3|^2 \cdot |z_5|^2 \cdot \dots \cdot |z_{2n-1}|^2 (\alpha_1 + i\beta_1) + |z_3|^2 \cdot |z_5|^2 \cdot \dots \cdot |z_{2n-1}|^2 (\alpha_1 - i\beta_1) = -\frac{B_1}{B_{2n}} \end{aligned}$$

or in equivalent form

$$2\alpha_{2n-1} |z_1|^2 \cdot |z_3|^2 \cdot \dots \cdot |z_{2n-3}|^2 + \dots + 2\alpha_1 |z_3|^2 \cdot |z_5|^2 \cdot \dots \cdot |z_{2n-1}|^2 = -\frac{B_1}{B_{2n}}.$$

Then, let us multiply and divide each term of the left-right side on the corresponding squares of modules:

$$\frac{2\alpha_{2n-1} \prod_{m=1}^n |z_{2m-1}|^2}{|z_{2n-1}|^2} + \dots + \frac{2\alpha_1 \prod_{m=1}^n |z_{2m-1}|^2}{|z_1|^2} = -\frac{B_1}{B_{2n}}$$

or

$$2 \prod_{m=1}^n |z_{2m-1}|^2 \left(\sum_{m=1}^n \frac{\alpha_{2m-1}}{|z_{2m-1}|^2} \right) = -\frac{B_1}{B_{2n}}.$$

Taking into account the equalities $B_{2n} = |A_n|^2$, $B_1 = \bar{A}_1 A_0 + \bar{A}_0 A_1 = 2(A_1, A_0)$, and relations (3), we obtain

$$2 \prod_{m=1}^n |w_m|^2 \left(\sum_{m=1}^n \frac{\text{Sc}(w_m)}{|w_m|^2} \right) = -\frac{2(A_1, A_0)}{|A_n|^2}.$$

Using Theorem 2, we have

$$2 \frac{|A_0|^2}{|A_n|^2} \left(\sum_{m=1}^n \frac{\text{Sc}(w_m)}{|w_m|^2} \right) = -\frac{2(A_1, A_0)}{|A_n|^2}.$$

Multiplying the previous equality on the $|A_n|^2$, we prove the theorem.

Corollary 1. *If all the roots of a quaternion polynomial $R_n(x)$ are pure vectors, then $(A_n, A_{n-1}) = 0$.*

Proof. By the condition of corollary we have $\sum_{m=1}^n \text{Sc}(w_m) = 0$. Therefore

$$-\frac{(A_n, A_{n-1})}{|A_n|^2} = 0.$$

Since $|A_n|^2 \neq 0$, then $(A_n, A_{n-1}) = 0$.

The following corollary is result of Theorem 4.

Corollary 2. *If all the roots of a quaternion polynomial $R_n(x)$ are pure vectors, then $(A_1, A_0) = 0$.*

4 Structure properties of roots of the polynomials

For a more detailed investigation of properties of the roots of quaternionic polynomials, we will find a general formula for calculate the degree of quaternion.

Consider the equalities:

$$w^0 = 1,$$

$$w^1 = w,$$

$$w^2 = 2x_0w - |w|^2,$$

$$\begin{aligned} w^3 &= w^2w = (2x_0w - |w|^2)w = 2x_0w^2 - |w|^2w = 2x_0(2x_0w - |w|^2) - |w|^2w = \\ &= 4x_0^2w - 2x_0^2|w|^2 - |w|^2w = (4x_0^2 - |w|^2)w - 2x_0|w|^2. \end{aligned}$$

Similarly

$$w^4 = (8x_0^3 - 4x_0|w|^2)w - (4x_0^2 - |w|^2)|w|^2,$$

$$w^5 = (16x_0^4 - 12x_0^2|w|^2 - |w|^4)w - (8x_0^3 - 4x_0|w|^2)|w|^2.$$

The coefficients near w and $|w|^2$ are real numbers, which we denote respectively by $Q_n(x_0, |w|^2)$ and $P_n(x_0, |w|^2)$. Then we get the equality

$$w^n = Q_n(x_0, |w|^2)w - P_n(x_0, |w|^2)|w|^2.$$

The functions Q_n and P_n are related by the recurrence formulae.

Property 1. *The following equalities are true*

$$1) \quad Q_n(w) = 2x_0Q_{n-1}(w) - P_{n-1}(w)|w|^2,$$

$$2) \quad P_n(w) = Q_{n-1}(w),$$

where $Q_0(w) := 0$, $P_0(w) := -\frac{1}{|w|^2}$, $Q_1(w) := 1$, $P_1(w) := 0$.

Proof. We will prove by mathematical induction. For $n = 1$ we have

$$Q_1(w) = 2x_0Q_0(w) - P_0(w)|w|^2 = 1,$$

$$P_1(w) = Q_0(w) = 0.$$

Suppose that formulas 1), 2) are true for all indexes less than n and prove this equalities for Q_n and P_n . Then we have the equality

$$w^{n-1} = Q_{n-1}(w)w - P_{n-1}(w)|w|^2.$$

Consider

$$\begin{aligned} w^n &= w^{n-1}w = (Q_{n-1}(w)w - P_{n-1}(w)|w|^2)w = \\ &= Q_{n-1}(w)(2x_0w - |w|^2) - P_{n-1}(w)|w|^2w = \\ &= (2x_0Q_{n-1}(w) - P_{n-1}(w)|w|^2)w - Q_{n-1}(w)|w|^2. \end{aligned}$$

From another hand

$$w^n = Q_n(w)w - P_n(w)|w|^2.$$

Equating the corresponding coefficients, we will get the desired equalities.

The next theorem is a *criterion of sphericity of root*.

Theorem 5. *The quaternion w_0 is a spherical root of polynomial $R_n(w)$ if and only if*

$$\sum_{m=1}^n A_m Q_m(w_0) = 0. \quad (8)$$

Proof. *Necessity.* If w_0 is a root of R_n , then $R_n(w_0) = 0$, i. e. $\sum_{m=0}^n A_m w_0^m = 0$.

Then

$$\begin{aligned} A_n (Q_n(w_0)w_0 - P_n(w_0)|w_0|^2) + A_{n-1} (Q_{n-1}(w_0)w_0 - P_{n-1}(w_0)|w_0|^2) \\ \dots + A_1 (Q_1(w_0)w_0 - P_1(w_0)|w_0|^2) + A_0 = 0, \end{aligned}$$

which implies

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) w_0 - |w_0|^2 \sum_{m=1}^n A_m P_m(w_0) + A_0 = 0. \quad (9)$$

Since w_0 is a spherical root, then any quaternion

$$w_1 = x_0 + \text{Vec}(w_1) \quad \text{with} \quad |\text{Vec}(w_0)| = |\text{Vec}(w_1)|$$

is also a root of this polynomial. Therefore

$$\begin{aligned} A_n (Q_n(w_1)w_1 - P_n(w_1)|w_1|^2) + A_{n-1} (Q_{n-1}(w_1)w_1 - P_{n-1}(w_1)|w_1|^2) \\ \dots + A_1 (Q_1(w_1)w_1 - P_1(w_1)|w_1|^2) + A_0 = 0. \end{aligned} \quad (10)$$

Denoted by

$$\begin{aligned} Q_n(w_0) &:= Q_n(x_0, |w_0|^2), & Q_n(w_1) &:= Q_n(x_0, |w_1|^2), \\ P_n(w_0) &:= P_n(x_0, |w_0|^2), & P_n(w_1) &:= P_n(x_0, |w_1|^2). \end{aligned}$$

Since $|w_0|^2 = |w_1|^2$, we obtain

$$Q_n(w_0) = Q_n(w_1), \quad P_n(w_0) = P_n(w_1). \quad (11)$$

Then equality (10) takes the form

$$\begin{aligned} A_n (Q_n(w_0)w_1 - P_n(w_0)|w_0|^2) + A_{n-1} (Q_{n-1}(w_0)w_1 - P_{n-1}(w_0)|w_0|^2) \\ \dots + A_1 (Q_1(w_0)w_1 - P_1(w_0)|w_0|^2) + A_0 = 0 \end{aligned}$$

or equivalent

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) w_1 - |w_0|^2 \sum_{m=1}^n A_m P_m(w_0) + A_0 = 0. \quad (12)$$

The equality (12) subtract from equality (9). Then

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) (w_0 - w_1) = 0,$$

whence

$$\sum_{m=1}^n A_m Q_m(w_0) = 0.$$

Sufficiency. Now we prove that any quaternion

$$w_1 = x_0 + \text{Vec}(w_1) \quad \text{such that} \quad |\text{Vec}(w_0)| = |\text{Vec}(w_1)|$$

is a root of polynomial $R_n(w)$. A consequence of equality (8) is identity

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) w_0 = 0. \quad (13)$$

The relations (11), (13) implies the equality

$$\left(\sum_{m=1}^n A_m Q_m(w_1) \right) w_1 = 0. \quad (14)$$

Now, from (13) and (14), we obtain

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) w_0 = \left(\sum_{m=1}^n A_m Q_m(w_1) \right) w_1. \quad (15)$$

Using equality (15), formula (9) takes the form

$$\left(\sum_{m=1}^n A_m Q_m(w_0) \right) w_0 - |w_0|^2 \sum_{m=1}^n A_m P_m(w_0) + A_0 = 0,$$

which is equivalent to the following condition

$$\begin{aligned} & A_n (Q_n(w_1)w_1 - P_n(w_1)|w_1|^2) + A_{n-1} (Q_{n-1}(w_1)w_1 - P_{n-1}(w_1)|w_1|^2) \\ & \dots + A_1 (Q_1(w_1)w_1 - P_1(w_1)|w_1|^2) + A_0 = 0, \end{aligned}$$

that is

$$\sum_{m=0}^n A_m w_1^m = 0.$$

Therefore, w_1 is a root of the polynomial $R_n(w)$. The theorem is proved.

The following statement follows from Theorem 5.

Corollary 3. *If w_0 and \bar{w}_0 are the roots of the polynomial $R_n(w)$ then this root is spherical.*

The following theorem establishes a relation between the roots of polynomials $R_n(w)$ and $L_n(w)$.

Theorem 6. *The following equality is true*

$$F_{2n}^*(w) = G_{2n}^*(w).$$

Proof. Let It is obviously that $\deg F_{2n}^*(w) = \deg G_{2n}^*(w)$. Now we prove the equality of the corresponding coefficients. For example, consider the second coefficients of basic polynomials corresponding for the right and left polynomials:

$$B_1^r = \bar{A}_0 A_1 + \bar{A}_1 A_0, \quad B_1^\ell = A_1 \bar{A}_0 + A_0 \bar{A}_1.$$

By Lemma 1 $B_1^r = B_1^\ell$. For the same reason all other coefficients also equal. The theorem is proved.

Remark. By Theorem 6, the all results for the right polynomials $R_n(w)$ also true for the left polynomials $L_n(w)$.

Acknowledgment

This work was supported by a grant from the Simons Foundation (1030291,V.S.Sh.).

References

- [1] V. S. Shpakivskyi, *Roots of polynomials of quaternionic variable* (in Ukrainian), Bachelor's thesis, Zhytomyr State University, 2007.
- [2] J. Niven, *Equations in quaternions*, Amer. Math. Monthly, (1941) 48: 654–661.
- [3] S. Eilenberg and I. Niven. *The "Fundamental Theorem of Algebra" for quaternions*, Bull. Amer. Math. Soc. (1944) 50: 246–248.
- [4] B. Beck, *Sur les huations polynomiales dans les quaternions*, L'Enseignement Mathématique XXV, 193–201, (1979).
- [5] A. Pogorui, M. Shapiro, *On the structure of the set of the zeros of quaternionic polynomials*, Complex Variables and Elliptic Equations, **49**, no. 6 (2004), 379–389.

- [6] G. Gentili, D. C. Struppa, F. Vlacc, *The fundamental theorem of algebra for Hamilton and Cayley numbers*, Math. Z. 259, 895–902 (2008).
- [7] D. Janovská, G. Opfer, *A note on the computation of all zeros of simple quaternionic polynomials*, SIAM J. Numer. Anal. 48, 244–256 (2010).
- [8] R. Serôdio, E. Pereira and J. Vitória, *Computing the zeroes of quaternion polynomials*, Comput., Math. Appl., (2001) 42(8–9), 1229–1237.
- [9] M. I. Falcão, F. Miranda, R. Severino, M. J. Soares, *Weierstrass method for quaternionic polynomial root-finding*, Mathematical Methods in the Applied Sciences, 41 (2018), 423–437.
- [10] N. Topuridze, *On the structure of the zero-set of a quaternionic polynomials*, Bull. Georgian Acad. Sci., **164**, 228–231 (2001).
- [11] G. Khimshiashvili, *Counting roots of quaternionic polynomials*, Bull. Georgian Acad. Sci., **165**, 465–468 (2002).
- [12] N. Topuridze, *On roots of quaternion polynomials*, J. Math. Sci. **160** (6), 843–855, (2009).
- [13] O. Ogunmefun, *Roots of Quaternionic Polynomials and Automorphisms of Roots*, (2023), Electronic Theses and Dissertations. Paper 4186. <https://dc.etsu.edu/etd/4186>
- [14] P. Dospra, D. Poulakis, *Complex Roots of Quaternion Polynomials*. In: Kotsireas, I., Martinez-Moro, E. (eds) Applications of Computer Algebra. ACA 2015. Springer Proceedings in Mathematics and Statistics, (2017), **198**, 45–58. Springer, Cham.
- [15] B. Kalantari, *Algorithms for quaternion polynomial root-finding*, Journal of Complexity, 2013, **29** (3–4), 302–322.
- [16] G. V. Milovanović, A. Mir, *Zeros of One Class of Quaternionic Polynomials*, Filomat 36:19 (2022), 6655–6667.
- [17] D. A. Mierzejewski, V. S. Szpakowski, *On solutions of some types of quaternionic quadratic equations*, Bull. Soc. Sci. Lettres Łódź **58**, Ser. Rech. Déform., 2008, **55**, 49–58.
- [18] V. S. Szpakowski, *Solution of general quadratic quaternionic equations*, Bull. Soc. Sci. Lettres Łódź **59**, Ser. Rech. Déform., 2009, **58**, 45–58.

- [19] D. Mierzejewski, *Investigation of quaternionic quadratic equations I. Factorization and passing to a system of real equations*, Bull. Soc. Sci. Lettres Łódź **58** Sér. Rech. Déform. **56** (2008), 17–26.
- [20] D. Shirokov, *On Noncommutative Vieta Theorem in Geometric Algebras*, In: Hitzer, E., Papagiannakis, G., Vasik, P. (eds) Empowering Novel Geometric Algebra for Graphics and Engineering. ENGAGE 2022. Lecture Notes in Computer Science, vol. 13862. Springer, Cham.
- [21] D. Shirokov, *Noncommutative Vieta theorem in Clifford geometric algebras*, Mathematical Methods in the Applied Sciences, 2024, **47** (14), 11305–11320.

Vitalii Shpakivskyi

Institute of Mathematics of the

National Academy of Sciences of Ukraine, Kyiv

e-mail: shpakivskyi86@gmail.com