# When is the Computation of a Feature Attribution Method Tractable?

**P.** Barcelo<sup>1,3,4</sup> and **R.** Cominetti<sup>1,2</sup> and **M. Morgado**<sup>1</sup>

<sup>1</sup> Institute for Mathematical and Computational Engineering, Universidad Católica de Chile

<sup>2</sup>Department of Industrial and Systems Engineering, Universidad Católica de Chile

<sup>3</sup>IMFD Chile and <sup>4</sup>CENIA Chile

pablo.barcelo, roberto.cominetti, mat.mw @uc.cl

#### Abstract

Feature attribution methods have become essential for explaining machine learning models. Many popular approaches, such as SHAP and Banzhaf values, are grounded in power indices from cooperative game theory, which measure the contribution of features to model predictions. This work studies the computational complexity of power indices beyond SHAP, addressing the conditions under which they can be computed efficiently. We identify a simple condition on power indices that ensures that computation is polynomially equivalent to evaluating expected values, extending known results for SHAP. We also introduce Bernoulli power indices, showing that their computation can be simplified to a constant number of expected value evaluations. Furthermore, we explore interaction power indices that quantify the importance of feature subsets, proving that their computation complexity mirrors that of individual features.

#### 1 Introduction

Feature attribution methods. Explainability of machine learning models has become one of the most pressing challenges associated with deploying this technology in critical applications. In fact, in many domains such as healthcare, finance, and autonomous systems, the ability to understand and trust model predictions is essential for ensuring safety, fairness, and accountability. As a result, research on providing robust and meaningful explanations for machine learning models has proliferated in recent years [\[Ribeiro](#page-6-0) *et al.*, 2016; [Doshi-Velez and Kim, 2017;](#page-6-1) [Lipton, 2018;](#page-6-2) [Marques-Silva and Ignatiev, 2022;](#page-6-3) [Darwiche, 2023\]](#page-6-4).

A popular approach to explaining a machine learning model's prediction is through *feature attribution methods*. These methods aim to quantify the contribution of each individual feature to the prediction  $F(\omega)$  made by a model F for a given input  $\omega$ . Several of the feature attribution methods used in practice are based on power indices from cooperative game theory, including the well-known *Shapley* and *Banzhaf* values [\[Lundberg and Lee, 2017;](#page-6-5) [Sundararajan and Najmi, 2020;](#page-6-6) [Karczmarz](#page-6-7) *et al.*, 2022; [Wang and Jia, 2023\]](#page-6-8). We provide a brief introduction to such indices below.

**Power indices.** Let  $Y = (Y_1, \ldots, Y_n)$  be a vector of features where each  $Y_i$  takes values in a finite set  $\Omega_i$ . We assume that the  $Y_i$ 's are independent random variables with corresponding product distribution

$$
\mathbb{P}(Y=\omega) = \prod_{i=1}^{n} \mathbb{P}(Y_i=\omega_i)
$$

over the set of possible outcomes  $\omega \in \Omega = \Omega_1 \times \ldots \times \Omega_n$ . We are also given a learned model  $F : \Omega \to \mathbb{R}$  that computes a prediction  $F(\omega)$ , for each  $\omega \in \Omega$ . Throughout the paper, we assume that  $F(\omega)$  can be computed in polynomial time in the number of features n.

Consider a fixed  $e \in \Omega$  and let  $N \triangleq \{1, \ldots, n\}$ . For each  $a \in N$ , we measure the relevance of the feature  $Y_a$  in explaining the outcome  $F(e)$  by using power indices from cooperative game theory. To do this, we first consider the conditional expectation of F on the event  ${Y<sub>S</sub> = e<sub>S</sub>}$ , that is

$$
\mathbb{E}[F|S] = \sum_{\omega \in \Omega} F(\omega) \mathbb{P}(Y = \omega | Y_S = e_S)
$$

$$
= \sum_{\omega \in \Omega} F(\omega) \prod_{i \in S} \delta_i(\omega_i) \prod_{i \notin S} \mathbb{P}(Y_i = \omega_i)
$$

with  $\delta_i$  defined as follows:

$$
\delta_i(w_i) = \begin{cases} 1 & \text{if } w_i = e_i, \\ 0 & \text{otherwise.} \end{cases}
$$

The *marginal contribution* of feature  $a \in N$  to a coalition  $S \subseteq N \setminus \{a\}$  is defined as

$$
m(a;S) \triangleq \mathbb{E}[F|S \cup \{a\}] - \mathbb{E}[F|S].
$$

Then, for a given probability distribution Q over the set of all subsets of  $N \setminus \{a\}$ , the *average contribution* of a according to  $Q$  is given by

$$
I(a; F) \triangleq \sum_{S \subseteq N \setminus \{a\}} Q(S) m(a; S).
$$

In cooperative games this corresponds to a *power index*, among which the Shapley and Banzhaf values are the most prominent examples:

- *Shapley value:* In this case,  $Q(S) = \frac{|S|!(n-|S|-1)!}{n!}$  for each  $S \subseteq N \setminus \{a\}.$
- *Banzhaf value:* This value considers a uniform probability distribution over the subsets of  $N \setminus \{a\}$ , i.e.,  $Q(S) = \frac{1}{2^{n-1}}$  for each  $S \subseteq N \setminus \{a\}.$

Throughout the paper, we will also consider some less common alternatives of power indices, such as the binomial indices, the Bernoulli indices, and others.

Previous results. The complexity of computing Shapley values, often referred to as SHAP scores in the context of ML models, has garnered significant attention in recent years [\[Van den Broeck](#page-6-9) *et al.*, 2022; [Arenas](#page-6-10) *et al.*, 2023; [Huang and Marques-Silva, 2024\]](#page-6-11). In particular, [Van den Broeck](#page-6-9) *et al.* have shown that the problem of computing SHAP scores over a model  $F$  is polynomially equivalent to computing expected values over F. Hence:

- The computation of SHAP scores is efficient for any class of models where expected values can be calculated in polynomial time. This means, e.g., that SHAP scores can be computed in polynomial time for linear regression models [\[Khosravi](#page-6-12) *et al.*, 2019], decision trees [\[Khosravi](#page-6-13) *et al.*, 2020], deterministic and decomposable Boolean circuits [\[Arenas](#page-6-10) *et al.*, 2023], and CNF formulas of bounded treewidth [\[Ferrara](#page-6-14) *et al.*, 2005].
- The computation of SHAP scores becomes intractable for model classes where computing expected values is itself intractable. Notably, this includes logistic regression models [\[Van den Broeck](#page-6-9) *et al.*, 2022] and (monotone) DNF formulas [\[Provan and Ball, 1983\]](#page-6-15).

Context. Although SHAP scores are the most widely studied feature attribution method, it has recently been argued that they are by no means a *silver bullet* for solving practical explainability challenges. In fact, it has been noted that, in some cases, other indices, such as the Banzhaf values, can provide more meaningful explainability results, depending on the class of models considered [\[Karczmarz](#page-6-7) *et al.*, 2022; [Wang and Jia, 2023\]](#page-6-8). Unfortunately, the computational complexity of power indices other than SHAP remains poorly understood, leading to the following natural question:

• Which power indices, as defined above, and for which classes of models, allow computation to be performed in polynomial time?

This question is critical for assessing the practical applicability of different power indices. Based on the previous observations, we rephrase this question as follows:

• Which power indices enable computation that is polynomially equivalent to evaluating expected values?

Results on simple power indices. We first focus on so called *simple* power indices, in which the value of  $Q(S)$  only depends on the cardinality of S, for each  $S \subseteq N \setminus \{a\}$ . In this case, we denote  $Q(S)$  as  $q_{|S|}$ . Notice that both Shapley and Banzhaf values are simple power indices.

• *Power index computation based on expected values:* We begin by observing that a straightforward extension of the results by [Van den Broeck](#page-6-9) *et al.* shows that the computation of any simple power index over a model  $F$  can be reduced, in polynomial time, to the computation of expected values over  $F$ . Specifically, the computation of an arbitrary simple power index over  $F$  can be carried out by evaluating the expected value of  $F$  over  $2n$ different probability distributions, where  $n$  is the number of features of  $F$ , and then performing polynomial interpolation on the resulting values.

• *Expected value computation based on power indices:* We then study the converse direction, i.e., whether the computation of expected values for  $F$  can be carried out in polynomial time if we are granted access to an oracle that computes power indices for  $F$  in polynomial time. We show that this is true for any simple power index that satisfies  $q_0 > 0$ , which includes many of the power indices studied in the literature (e.g., Shapley and Banzhaf values). While for Shapley values, this is a straightforward consequence of its *efficiency property*, for other indices that do not satisfy this property (including Banzhaf), the proof is more complicated and involves polynomial interpolation. We also observe that the condition  $q_0 > 0$  is necessary, in the sense that there are simple power indices with  $q_0 = 0$  and classes of models where the computation of the index is tractable, while the computation of expected values is intractable.

Results on Bernoulli power indices. As previously mentioned, simple power indices can be computed via polynomial interpolation over a polynomial number of expected values. We prove that this can be further simplified for what we term *Bernoulli* power indices. These indices are defined by probability distributions Q such that  $Q(S)$ , for any  $S \subseteq N \setminus \{a\}$ , is determined by independent Bernoulli trials. Notably, we show that for such power indices—which are not necessarily simple and include the important case of the Banzhaf value—computation can be reduced, in polynomial time, to calculating the difference between only *two* expected values.

Results on interaction power indices. A recent trend in feature attribution methods focuses on developing approaches that evaluate the importance not only of individual features, as exemplified by the previously introduced power indices, but also of feature sets of arbitrary size [\[Beliakov](#page-6-16) *et al.*, 2020; [Sundararajan](#page-6-17) *et al.*, 2020]. This shift aims to capture and quantify the significance of feature interactions. However, the computational complexity of interaction power indices across different classes of models remains unexplored. We show that for *simple* interaction power indices, the computational scenario mirrors that of simple power indices for individual features. Specifically, the computation of simple interaction power indices can be performed in polynomial time if and only if expected values can also be computed in polynomial time. Proving this result requires an extension of our previous techniques, as the computation of interaction power indices now relies on *bivariate* polynomial interpolation using expected values. We also investigate *Bernoulli* interaction power indices and show that, for feature sets of constant size, these indices can be computed with a fixed number of calls to an oracle for evaluating expected values.

Organization of the paper. Our results on simple power indices are presented in Section [2,](#page-2-0) those for Bernoulli power indices in Section [3,](#page-4-0) and those for interaction power indices in Section [4.](#page-4-1) Final remarks can be found in Section [5.](#page-6-18)

**Notations.** We let  $N = \{1, \ldots, n\}$  be the set of features. For  $A \subseteq N$  we denote  $|A|$  its cardinality,  $A^c = N \setminus A$  its complement,  $\mathcal{P}(A)$  the family of all subsets of A, and  $\mathcal{P}_k(A)$ the subsets of cardinality  $k$ .

# <span id="page-2-0"></span>2 Cardinality-based indices

We begin by considering the *simple power indices* where the probability  $Q(S) = q_{|S|}$  depends only on the cardinality of S, and hence  $q_0, \ldots, q_{n-1} \ge 0$  are such that  $\sum_{k=0}^{n-1} \binom{n-1}{k} q_k = 1$ .

#### 2.1 From expected values to power indices

In this section, we show that the computation of any simple power index for a model  $F$  can be reduced, in polynomial time, to evaluating expected values over  $F$ . This result builds on a straightforward extension of the techniques introduced by [Van den Broeck](#page-6-9) *et al.*, who established a similar reduction specifically for SHAP scores.

Denoting  $\mathcal{P}_k^a = \mathcal{P}_k(\lbrace a \rbrace^c)$  the set of coalitions  $S \subseteq \lbrace a \rbrace^c$ of size  $|S| = k$ , we have

$$
I(a;F) = \sum_{k=0}^{n-1} q_k m_k(a)
$$
 (1)

where

$$
m_k(a) = \sum_{S \in \mathcal{P}_k^a} m(a;S). \tag{2}
$$

.

Thus, in order to compute  $I(a; F)$  it suffices to calculate the sums  $m_k(a)$ . This can be done with the help of the following variant of [\[Van den Broeck](#page-6-9) *et al.*, 2022, Claim 2].

<span id="page-2-1"></span>**Lemma 1.** Let  $G : \Omega \to \mathbb{R}$  and  $c_k = \sum_{S \in \mathcal{P}_k} \mathbb{E}[G|S]$  with  $\mathcal{P}_k$  *the family of all subsets of* N *of size* k*.* For  $z \geq 0$  let  $Y^z = (Y_1^z, \ldots, Y_n^z)$  be independent random variables with

$$
\mathbb{P}(Y_i^z = \omega_i) = \frac{z \,\delta_i(\omega_i) + \mathbb{P}(Y_i = \omega_i)}{1 + z}
$$

*Then*  $\sum_{k=0}^{n} c_k z^k = (1+z)^n \mathbb{E}[G(Y^z)].$ 

*Proof.* By direct substitution we have

$$
\sum_{k=0}^{n} c_k z^k = \sum_{k=0}^{n} \sum_{S \in \mathcal{P}_k} \mathbb{E}[G|S] \cdot z^k
$$
  
= 
$$
\sum_{S \subseteq N} \mathbb{E}[G|S] \cdot z^{|S|}
$$
  
= 
$$
\sum_{S \subseteq N} z^{|S|} \sum_{\omega \in \Omega} G(\omega) \prod_{i \in S} \delta_i(\omega_i) \prod_{i \notin S} \mathbb{P}(Y_i = \omega_i).
$$

Exchanging the order of the sums we get

$$
\sum_{k=0}^{n} c_k z^k = \sum_{\omega \in \Omega} G(\omega) \sum_{S \subseteq N} \prod_{i \in S} z \, \delta_i(\omega_i) \times \prod_{i \notin S} \mathbb{P}(Y_i = \omega_i).
$$

Now, using the identity

<span id="page-2-3"></span>
$$
\sum_{S \subseteq N} \prod_{i \in S} u_i \times \prod_{i \in S^c} v_i = \prod_{i=1}^n (u_i + v_i) \tag{3}
$$

the inner sum above is exactly  $(1 + z)^n \mathbb{P}(Y^z = w)$  and then  $\sum_{k=0}^{n} c_k z^k = (1+z)^n \mathbb{E}[G(Y^z)].$  $\Box$ 

Using Lemma [1,](#page-2-1) we prove that the computation of simple power indices can be reduced to evaluating expected values in polynomial time. For the Shapley value—defined by  $q_k = \frac{k! (n-1-k)!}{(n-1)!}$ —this result was previously established in [\[Van den Broeck](#page-6-9) *et al.*, 2022].

<span id="page-2-2"></span>Theorem 2. *The computation of a simple power index*  $I(a; F)$ *, for any feature*  $a \in N$ *, can be reduced in polynomial time to evaluate* E[F(Z)] *for a family of* 2n *random vectors*  $Z = (Z_1, \ldots, Z_n)$ , each of them with a product distribution.

*Proof.* As mentioned earlier, computing  $I(a; F)$  boils down to calculate the coefficients  $c_k = m_k(a)$  for  $0 \leq k \leq n-1$ 1. Recall that we have been given a fixed partial instance  $e \in \Omega$ . Let  $\Omega^{-a} = \bigotimes_{i \neq a} \Omega_i$ . For  $\omega^{-a} \in \Omega^{-a}$ , we slightly abuse notation and write  $(e_a, \omega^{-a})$  for the element in  $\Omega$  that is obtained by extending  $\omega^{-a}$  with value  $e_a$  for feature a. Analogously, we write  $(Y_a, \omega^{-a})$  and  $(Y_a, (Y_i^z)_{i \neq a})$ , where  $Y_a$  is a random value that ranges over  $\Omega_a$ .

Consider the map  $G : \Omega^{-a} \to \mathbb{R}$  defined by

$$
G(\omega^{-a}) = F(e_a, \omega^{-a}) - \mathbb{E}[F(Y_a, \omega^{-a})]
$$

with a partial expectation of  $F(\cdot)$  with respect to  $Y_a$ . Then  $m(a; S) = \mathbb{E}[G|S]$  so that  $c_k = \sum_{S \in \mathcal{P}_k^a} \mathbb{E}[G|S]$  and Lemma [1](#page-2-1) gives the following:

$$
\sum_{k=0}^{n-1} c_k z^k = (1+z)^{n-1} \mathbb{E}[G((Y_i^z)_{i \neq a})].
$$

Evaluating at *n* different points  $z_0, z_1, \ldots, z_{n-1} \geq 0$ , we obtain a linear system for the coefficients  $c_0, \ldots, c_{n-1}$ . This system is defined by a Vandermonde matrix and can be solved uniquely. Thus, the computation of  $I(a; F)$  reduces to evaluate the expectations  $\mathbb{E}[G((Y_i^z)_{i\neq a})]$  for  $z = z_0, \dots, z_{n-1}$ . The proof is completed by noting that each such expectation for G amounts to compute the difference between the expectations of  $F$  for two product distributions, namely

$$
\mathbb{E}[G((Y_i^z)_{i\neq a})] = \mathbb{E}[F(Y^{1,z})] - \mathbb{E}[F(Y^{0,z})]
$$

with 
$$
Y^{1,z} = (e_a, (Y_i^z)_{i \neq a})
$$
 and  $Y^{0,z} = (Y_a, (Y_i^z)_{i \neq a})$ .  $\square$ 

#### 2.2 From power indices to expected values

Consider a simple power index  $I(a; F)$  with  $q_0 > 0$ . We show that the evaluation of the expected value for a model  $F$ reduces, in polynomial time, to the computation of  $I(a; F)$ for every feature  $a \in N$ . This property is well-known for the SHAP score, as it follows directly from the *efficiency* property of the Shapley value, which ensures that

$$
\sum_{a=1}^{n} I(a; F) = F(e) - \mathbb{E}[F].
$$

However, this identity does not hold for the Banzhaf value or other cardinality-based indices. Consequently, an alternative argument is required to establish the reduction in these cases.

Towards this end we first establish a technical Lemma, for which we introduce the following notations. Consider the random vectors  $Y^z$  as in Lemma [1,](#page-2-1) and denote  $\mathbb{E}_z[\cdot]$  the expectation for the corresponding product distribution, keeping  $\mathbb{E}[\cdot]$  for the original distribution of Y. Denote by  $m^2(a; S)$ and  $I^z(a;F)$  the marginal contributions and indices computed with  $Y^z$ . Finally, for  $k = 0, \ldots, n$  let

$$
\beta_k^n = k q_{k-1} - (n-k) q_k
$$

with  $q_{-1} = q_n = 0$ , and consider the polynomials

$$
P_{\ell}(z) \triangleq \sum_{k=0}^{\ell} {\ell \choose k} \beta_k^n (1+z)^k z^{\ell-k}.
$$

**Lemma 3.** Let  $\Theta(z) = \sum_{a=1}^{n} I^z(a;F)$  be the sum of all *indices computed for*  $Y^z$ *. Then, setting*  $c_\ell \triangleq \sum_{S \in \mathcal{P}_\ell} \mathbb{E}[F|S]$ *we have*  $\sum_{\ell=0}^{n} c_{\ell} P_{\ell}(z) = (1+z)^{n} \Theta(z)$ *.* 

*Proof.* By direct substitution we have

$$
\Theta(z) = \sum_{a=1}^{n} \sum_{k=0}^{n-1} \sum_{S \in \mathcal{P}_k^a} q_k m^z(a;S)
$$
  
= 
$$
\sum_{a=1}^{n} \sum_{k=0}^{n-1} \sum_{S \in \mathcal{P}_k^a} q_k \left( \mathbb{E}_z[F|S \cup \{a\}] - \mathbb{E}_z[F|S] \right).
$$

In this sum, each subset  $A \subseteq N$  of size  $|A| = k$  appears k times as a positive term  $q_{k-1} \mathbb{E}_z[F|S \cup \{a\}]$  for each  $a \in A$ , and  $n - k$  times as a negative term  $-q_k \mathbb{E}_{z}[F|S]$  for  $a \notin A$ , so that the sum can be expressed as

$$
\Theta(z) = \sum_{k=0}^{n} \sum_{A \in \mathcal{P}_k} \beta_k^n \, \mathbb{E}_z[F|A].
$$

Now, using Lemma [1](#page-2-1) and denoting  $P_j(A^c)$  the subsets of  $A^c$ of size j, we may compute  $\mathbb{E}_{z}[F|A]$  as

$$
\mathbb{E}_{z}[F|A] = \frac{1}{(1+z)^{n-k}} \sum_{j=0}^{n-k} z^j \sum_{B \in \mathcal{P}_j(A^c)} \mathbb{E}[F|A \cup B]
$$

which substituted above gives

$$
\Theta(z) = \sum_{k=0}^{n} \sum_{A \in \mathcal{P}_k} \sum_{j=0}^{n-k} \sum_{B \in \mathcal{P}_j(A^c)} \beta_k^n \frac{z^j}{(1+z)^{n-k}} \mathbb{E}[F|A \cup B].
$$

Next, we observe that every set  $S \subseteq N$  of size  $|S| = \ell$  admits  $\binom{\ell}{k}$  decompositions as  $S = A \cup B$  with  $A \in \mathcal{P}_k$  and  $B \in$  $\mathcal{P}_j(A^c)$ , where  $j = \ell - k$ , so that we may further rewrite

$$
\Theta(z) = \sum_{\ell=0}^{n} \sum_{S \in \mathcal{P}_{\ell}} \mathbb{E}[F|S] \sum_{k=0}^{\ell} {(\ell) \beta_k^n \over (1+z)^{n-k}}
$$

$$
= \frac{1}{(1+z)^n} \sum_{\ell=0}^{n} c_{\ell} P_{\ell}(z).
$$

Suppose we want to evaluate the expectation  $c_0 = \mathbb{E}[F]$ . Assuming that the simple power indices  $I^z(a;F)$  can be computed efficiently, we may evaluate  $(1+z)^n \Theta(z)$  at  $n+1$  different points  $z_0, \ldots, z_n$  in order to write a linear system  $Mc = b$ for the coefficients  $c = (c_0, \ldots, c_n)$  with  $M_{i,\ell} = P_{\ell}(z_i)$ and  $b_i = (1 + z_i)^n \Theta(z_i)$  for  $0 \leq i, \ell \leq n$ . Unfortunately the family  $\{P_\ell(\cdot): \ell = 0, \ldots, n\}$  is linearly dependent and the matrix  $M$  turns out to be singular (see below). However, since we have explicitly  $c_n = F(e)$ , we may write a reduced system in the unknowns  $(c_0, \ldots, c_{n-1})$ . We next establish sufficient conditions under which this reduced system is nonsingular, by exploiting the following simple fact.

REMARK. Let  $P_0(\cdot), \ldots, P_{n-1}(\cdot)$  be polynomials with  $deg(P_\ell) = \ell$ . Then, for any family  $z_0, \ldots, z_{n-1} \in \mathbb{R}$  of different reals, the following matrix is nonsingular

$$
M = \begin{pmatrix} P_0(z_0) & \cdots & P_{n-1}(z_0) \\ \vdots & & \vdots \\ P_0(z_{n-1}) & \cdots & P_{n-1}(z_{n-1}) \end{pmatrix}.
$$

Indeed, take  $\alpha \in \text{Ker}(M)$  so that  $M\alpha = 0$ . This implies that  $P(x) = \sum_{j=0}^{n-1} \alpha_j P_j(x)$  is a polynomial of degree  $n-1$  with *n* distinct roots  $z_0, \ldots, z_{n-1}$  and therefore  $P(\cdot) \equiv 0$ . Since  $deg(P_\ell) = \ell$ , the polynomials  $P_\ell(\cdot)$  are linearly independent so that  $\alpha = 0$  and therefore M is nonsingular. П

<span id="page-3-0"></span>Using this, we get the following converse of Theorem [2.](#page-2-2)

**Theorem 4.** *Consider a simple power index such that*  $q_0 > 0$ *. Then, computing* E[F] *reduces in polynomial time to evaluate the simple power indices*  $I^z(a;F)$  *for all*  $a = 1, \ldots, n$  *at*  $n$ *different positive reals*  $z = z_0, \ldots, z_{n-1} \in \mathbb{R}_+$ .

*Proof.* Recall that  $c_n = F(e)$  and consider the reduced linear system in the unknowns  $c = (c_0, \ldots, c_{n-1})$ :

$$
\sum_{\ell=0}^{n-1} c_{\ell} P_{\ell}(z_i) = (1+z_i)^n \Theta(z_i) - c_n P_n(z_i) \quad \forall \, 0 \leq i \leq n-1.
$$

The left hand side corresponds to  $Mc$  with  $M$  as in the remark above. We claim that since  $q_0 > 0$  each polynomial  $P_{\ell}(\cdot)$  has degree  $\ell$  for  $\ell = 0, \ldots, n - 1$ . Indeed, the leading coefficient of  $z^{\ell}$  in  $P_{\ell}(z)$  is  $a_{\ell} = \sum_{k=0}^{\ell} {\ell \choose k} \beta_k^n$ . Recalling that  $q_{-1} = 0$ and after some simple algebraic manipulations, we get

$$
a_{\ell} = \sum_{k=0}^{\ell} {\ell \choose k} (k q_{k-1} - (n-k) q_k)
$$
  
=  $(\ell - n) \sum_{k=0}^{\ell} {\ell \choose k} q_k.$ 

The latter is strictly negative for  $0 \leq \ell \leq n-1$  so that  $P_{\ell}(\cdot)$ has degree  $\ell$ . Invoking the previous remark it follows that  $M$ is nonsingular and then we can solve the linear system for c in order to recover  $c_0 = \mathbb{E}[F]$ . П

REMARK. Observe that for  $\ell = n$  the polynomial  $P_n(\cdot)$  has degree at most  $n-1$ , so that the family  $\{P_\ell(\cdot): \ell = 0, \ldots, n\}$ with  $P_n(\cdot)$  included is always linearly dependent. П

# Example 1.

(a) *Shapley value*: For  $q_k = \frac{k! (n-k-1)!}{n!}$  $\frac{-\kappa-1\mathrm{I}}{n!}$  a direct calculation gives  $P_{\ell}(z) = -z^{\ell}$  for  $\ell = 0, \ldots, n-1$  which are linearly independent with  $a_\ell = -1$ .

- (b) *Banzhaf value*: With  $q_k = 1/2^{n-1}$  for  $0 \leq k \leq n-1$ we have  $P_{\ell}(z) = \frac{1}{2^{n-1}} (\ell (1 + 2z)^{\ell-1} - (n - \ell)(1 + 2z)^{\ell})$ which has degree  $\ell$  with  $a_{\ell} = (\ell - n)2^{\ell - n + 1}$ .
- (c) *Binomial value*: Binomial indices correspond to the choice  $q_k = \theta^k (1-\theta)^{n-1-k}$  with  $\theta \in (0,1)$ . In this case  $P_{\ell}(z) = (1 - \theta)^{n - \ell - 1} (\ell(z + 1)(z + \theta)^{\ell - 1} - n(z + \theta)^{\ell})$ with  $a_{\ell} = (\ell - n)(1 - \theta)^{n - \ell - 1}$ .
- (d) *Dictatorial value*: This is given by  $q_0 = 1$  and  $q_k = 0$ for  $k = 1, \ldots, n - 1$ . Here  $p_{\ell}(z) = \ell z^{\ell-1} - (n - \ell)z^{\ell}$ and  $a_\ell = \ell - n$ .
- (e) *Marginal value*: In this case  $q_{n-1} = 1$  and  $q_k = 0$ for  $k = 0, \ldots, n - 2$ . Linear independence fails since  $P_{\ell}(z) \equiv 0$  for  $\ell = 0, \ldots, n-2$ . Moreover, even if  $\mathbb{E}[F]$  is difficult to compute, the marginal indices can be computed efficiently (assuming that  $F$  is) by using the explicit formula

$$
I(a;F) = \mathbb{E}[F|N] - \mathbb{E}[F|N \setminus \{a\}]
$$
  
=  $F(e) - \sum_{\omega_a \in \Omega_a} F(\omega_a, e_{-a}) \mathbb{P}(Y_a = \omega_a).$ 

This shows that in general there is no polynomial time reduction from the indices to the expectation, and some additional condition such as  $q_0 > 0$  is needed.  $\Box$ 

# <span id="page-4-0"></span>3 Bernoulli indices

As we show in this section, some power indices can be computed much more efficiently than through polynomial interpolation over a polynomial number of expected values. Let  $a \in N$  be a fixed feature and consider a power index in which the probability Q over  $\mathcal{P}(\{a\}^c)$  results from independent Bernoulli trials. Specifically, we consider a random subset  $S \subseteq \{a\}^c$  that includes each  $i \neq a$  with probabilities  $\theta_i \in [0, 1]$ , so that

$$
Q(S) = \prod_{i \in S} \theta_i \cdot \prod_{i \notin S \cup \{a\}} (1 - \theta_i).
$$

The corresponding index

$$
I(a;F) \triangleq \sum_{S \subseteq \{a\}^c} Q(S) (\mathbb{E}[F|S \cup \{a\}] - \mathbb{E}[F|S])
$$

is then called a *Bernoulli* power index.

REMARK. This class includes the binomial indices with  $\theta_i \equiv$  $\theta$  and the Banzhaf value with  $\theta_i \equiv \frac{1}{2}$ , both of which are also cardinality-based indices. However, apart from these cases, Bernoulli indices are not necessarily simple, as  $Q(S)$  does not need to depend solely on  $|S|$ . Observe that the SHAP score is not a Bernoulli index.  $\square$ 

Suppose that the feature  $a$  is also included with some probability  $\theta_a$  and let

$$
Q_{\theta}(S) \triangleq \prod_{i \in S} \theta_i \cdot \prod_{i \notin S} (1 - \theta_i)
$$

be the distribution over all subsets  $S \subseteq N$ . By considering  $\theta^1$  and  $\theta^0$  the vectors where we take respectively  $\theta_a = 1$  and  $\theta_a = 0$ , and denoting  $Q_1$  and  $Q_0$  the corresponding distributions over  $P(N)$ , the Bernoulli index can be expressed as

$$
I(a;F) = \sum_{S \subseteq N} Q_1(S) \mathbb{E}[F|S] - \sum_{S \subseteq N} Q_0(S) \mathbb{E}[F|S].
$$

We claim that this is a difference of expectations with respect to two product distributions, namely

<span id="page-4-2"></span>**Theorem 5.** Let  $Y^{\theta} = (Y_1^{\theta_1}, \dots, Y_n^{\theta_n})$  be a tuple of indepen*dent random variables with marginals given by the mixtures*  $\mathbb{P}(Y_i^{\theta_i} = \omega_i) = \theta_i \delta_i(\omega_i) + (1-\theta_i) \mathbb{P}(Y_i = \omega_i)$ . Then, denoting  $Y^{\theta,0}$  and  $Y^{\theta,1}$  the tuples  $Y^{\theta}$  with  $\theta_a = 0$  and  $\theta_a = 1$ , we have

$$
I(a;F) = \mathbb{E}\big[F(Y^{\theta,1})\big] - \mathbb{E}\big[F(Y^{\theta,0})\big].
$$

*Proof.* This follows from the general identity

$$
\sum_{S \subseteq N} Q_{\theta}(S) \mathbb{E}[F|S]
$$
\n
$$
= \sum_{S \subseteq N} Q_{\theta}(S) \sum_{\omega \in \Omega} F(\omega) \prod_{i \in S} \delta_i(\omega_i) \times \prod_{i \notin S} \mathbb{P}(Y_i = \omega_i)
$$
\n
$$
= \sum_{\omega \in \Omega} F(\omega) \sum_{S \subseteq N} \prod_{i \in S} \theta_i \delta_i(\omega_i) \times \prod_{i \notin S} (1 - \theta_i) \mathbb{P}(Y_i = \omega_i)
$$
\n
$$
= \sum_{\omega \in \Omega} F(\omega) \prod_{i \in N} (\theta_i \delta_i(\omega_i) + (1 - \theta_i) \mathbb{P}(Y_i = \omega_i))
$$
\n
$$
= \sum_{\omega \in \Omega} F(\omega) \mathbb{P}(Y^{\theta} = \omega)
$$
\n
$$
= \mathbb{E}[F(Y^{\theta})].
$$

REMARK. As mentioned earlier, the Banzhaf and binomial indices are Bernoulli indices, so by Theorem [5](#page-4-2) they can be calculated by computing just 2 expectations. This is more efficient than considering them as cardinality-based and using Theorem [2](#page-2-2) which requires 2n expectations.  $\Box$ 

# <span id="page-4-1"></span>4 Interaction Indices

*Interaction indices* aim not only to measure the relevance of a feature  $a \in N$  in isolation, but more generally to explain a subset of features  $A \subseteq N$ . An interaction index is defined as

$$
I(A;F) = \sum_{S \subseteq A^c} Q_A(S) m(A;S)
$$

where  $Q_A(\cdot)$  is a probability on  $\mathcal{P}(A^c)$ , and the marginal contribution of A to S is defined as in [\[Beliakov](#page-6-16) *et al.*, 2020; [Sundararajan and Najmi, 2020\]](#page-6-6) by

$$
m(A;S) = \sum_{B \subseteq A} (-1)^{|A \setminus B|} \mathbb{E}[F|S \cup B].
$$

Alternatively, we have

$$
I(A;F) = \sum_{B \subseteq A} (-1)^{|A \setminus B|} R(B),
$$

where  $R(B) = \sum_{S \subseteq A^c} Q_A(S) \mathbb{E}[F|S \cup B].$ 

#### 4.1 Cardinality-based interaction indices

An interaction index is said to be *simple* if it is characterized by a distribution  $Q_A(S) = q(|S|, |A|)$ , where the function q depends solely on the cardinalities of the sets  $S$  and  $A$ . We show that the computation of simple interaction indices retains the same properties as their single-feature counterparts:

- The computation of any interaction index for a model  $F$ can be reduced in polynomial time to the computation of expected values over F.
- Conversely, if  $q(1, 0) > 0$ , the computation of expected values over  $F$  can be reduced in polynomial time to the computation of interaction indices over F.

Note that it suffices to prove the first item, as the second follows directly from Theorem [4.](#page-3-0) This is because any index of the form  $I(a; F)$ , where  $a \in N$ , is also an interaction index. The proof of the first item (Theorem [6](#page-5-0) below) differs from that of Theorem [2](#page-2-2) due to the nature of interaction indices. Specifically, the computation now requires the use of *bivariate* polynomial interpolation to determine the indices based on a set of expected values.

<span id="page-5-0"></span>**Theorem 6.** Let  $A \subseteq N$  with  $|A| = m$ . Then a simple in*teraction index*  $I(A; F)$  *can be reduced to computing the expected values* E - F(Z) *for* (n−m+1)(m+1)*random vectors*  $Z = (Z_1, \ldots, Z_n)$  *with product distribution.* 

*Proof.* By splitting the sums according to the cardinality of the sets, and denoting

$$
c_{k,j} = \sum_{\substack{S \subseteq A^c \\ |S| = k}} \sum_{\substack{B \subseteq A \\ |B| = j}} \mathbb{E}\big[F|S \cup B\big]
$$

we have

$$
I(A;F) = \sum_{S \subseteq A^c} Q_A(S) \sum_{B \subseteq A} (-1)^{|A \setminus B|} \mathbb{E}[F|S \cup B]
$$
  
= 
$$
\sum_{k=0}^{n-m} \sum_{j=0}^{m} q(k,m) (-1)^{m-j} c_{k,j}
$$

so that  $I(A; F)$  reduces to computing the sums  $c_{k,j}$ .

Consider the bivariate polynomial

$$
P(z, y) = \sum_{k=0}^{n-m} \sum_{j=0}^{m} c_{k,j} z^k y^j.
$$

Proceeding as in the proof of Lemma [1,](#page-2-1) we have

$$
P(z, y) = \sum_{\omega \in \Omega} F(\omega) \sum_{S \subseteq A^c} \sum_{B \subseteq A} \pi_{S, B} \tag{4}
$$

with

$$
\pi_{S,B} = \prod_{i \in S} z \, \delta_i(\omega_i) \prod_{i \in B} y \, \delta_i(\omega_i) \prod_{i \notin S \cup B} \mathbb{P}(Y_i = \omega_i).
$$

The latter can be factorized as  $\pi_{S,B} = \pi_S \cdot \pi_B$  with

$$
\pi_S = \prod_{i \in S} z \, \delta_i(\omega_i) \prod_{i \in A^c \setminus S} \mathbb{P}(Y_i = \omega_i)
$$

$$
\pi_B = \prod_{i \in B} y \, \delta_i(\omega_i) \prod_{i \in A \setminus B} \mathbb{P}(Y_i = \omega_i)
$$

from which it follows that

$$
\sum_{S \subseteq A^c} \sum_{B \subseteq A} \pi_{S,B} = \left( \sum_{S \subseteq A^c} \pi_S \right) \left( \sum_{B \subseteq A} \pi_B \right).
$$

Now, using the identity [\(3\)](#page-2-3) (with  $A<sup>c</sup>$  instead of N) we get

$$
\sum_{S \subseteq A^c} \pi_S = \prod_{i \in A^c} \left( z \, \delta_i(\omega_i) + \mathbb{P}(Y_i = \omega_i) \right)
$$

$$
= (1+z)^{|A^c|} \prod_{i \in A^c} \mathbb{P}(Y_i^z = \omega_i)
$$

and similarly

$$
\sum_{B \subseteq A} \pi_B = (1+z)^{|A|} \prod_{i \in A} \mathbb{P}(Y_i^y = \omega_i).
$$

Combining these two expressions we deduce that

$$
\sum_{S \subseteq A^c} \sum_{B \subseteq A} \pi_{S,B} = (1+z)^n \mathbb{P}(Z=\omega)
$$

with  $Z = ((Y_i^y)_{i \in A}, (Y_i^z)_{i \in A^c})$ . Substituting this into [\(4\)](#page-5-1) it follows that computing  $P(z, y)$  reduces to evaluate an expected value  $P(z, y) = (1+z)^n \mathbb{E}[F(Z)]$  for a random vector Z with product distribution.

Now, evaluating  $P(z, y)$  over a rectangular grid

 $G = \{z_0, \ldots, z_{n-m}\} \times \{y_0, \ldots, y_m\}$ 

with all the  $z_i$ 's and  $y_j$ 's different, we obtain a linear system for the coefficients  $c_{k,j}$ . The matrix for this system is a bivariate Vandermonde matrix M with nonzero determinant (see [\[De Marchi and Usevich, 2014,](#page-6-19) page 23]):

$$
|\det(M)| = \left(\prod_{i < k} |z_i - z_k|\right)^{m+1} \times \left(\prod_{j < \ell} |y_j - y_\ell|\right)^{n-m+1} \neq 0
$$

where the first product is over all pairs  $i, k \in \{0, \ldots, n-m\}$ such that  $i < k$ , and the second over all  $j, \ell \in \{0, \ldots, m\}$ with  $j < l$ . Hence one can uniquely solve the system to find the coefficients  $c_{k,j}$ , and then compute  $I(A;F)$ . □

#### 4.2 Bernoulli interaction indices

Following with the ideas developed in Section [3,](#page-4-0) let us now consider a Bernoulli interaction index defined as follows for every  $A \subseteq N$ :

$$
Q_A(S) = \prod_{i \in S} \theta_i \times \prod_{i \in A^c \setminus S} (1 - \theta_i).
$$

<span id="page-5-1"></span>REMARK. Notice that for any specific set A we could choose different Bernoulli probabilities  $\theta_i^A$  for  $i \in A^c$ . Since here we consider a fixed A, we keep the simpler notation  $\theta_i$ . П

To conclude, we present the following straightforward extension of Theorem [5.](#page-4-2) This result establishes that if A has size m, then  $I(A; F)$  can be computed using an explicit formula involving  $2^m$  expected values over F. When  $m = 1$ , this reduces precisely to the statement of Theorem [5.](#page-4-2) Notice the difference with Theorem [6,](#page-5-0) as in this case the number of expected values one needs to compute does not depend on the number of features of the underlying model F.

Theorem 7. *A Bernoulli interaction index* I(A; F) *can be reduced to computing the expected values*  $\mathbb{E}[F(Z)]$  for  $2^m$ *random vectors*  $Z = (Z_1, \ldots, Z_n)$  *with product distribution.* 

*Proof.* Using  $I(A; F) = \sum_{B \subseteq A} (-1)^{|A \setminus B|} R(B)$ , it suffices to show that each  $R(B)$  is an expectation  $\mathbb{E}[F(Z)]$  for some Z with product distribution. Indeed, since

$$
R(B) = \sum_{S \subseteq A^c} \mathbb{E}\big[F|S \cup B\big] \prod_{i \in S} \theta_i \times \prod_{i \in A^c \setminus S} (1 - \theta_i)
$$

$$
\mathbb{E}\big[F|S \cup B\big] = \sum_{\omega \in \Omega} F(\omega) \prod_{i \in S \cup B} \delta_i(\omega_i) \times \prod_{i \notin S \cup B} \mathbb{P}(Y_i = \omega_i)
$$

after some straightforward algebraic manipulations we get

$$
R(B) = \sum_{\omega \in \Omega} F(\omega) \prod_{i \in B} \delta_i(\omega_i) \times \prod_{i \in A \setminus B} \mathbb{P}(Y_i = \omega_i) \times D(\omega)
$$

where

$$
D(\omega) = \sum_{S \subseteq A^c} \prod_{i \in S} \theta_i \delta_i(\omega_i) \times \prod_{i \in A^c \setminus S} (1 - \theta_i) \mathbb{P}(Y_i = \omega_i)
$$
  
= 
$$
\prod_{i \in A^c} (\theta_i \delta_i(\omega_i) + (1 - \theta_i) \mathbb{P}(Y_i = \omega_i)).
$$

Hence, denoting  $Y_i^{\theta}$  a random variable with distribution

$$
\mathbb{P}(Y_i^{\theta} = \omega_i) = \theta_i \delta_i(\omega_i) + (1 - \theta_i) \mathbb{P}(Y_i = \omega_i)
$$

it follows that  $R(B) = \mathbb{E}[F(Z)]$  where

$$
Z = ((e_i)_{i \in B}, (Y_i)_{i \in A \setminus B}, (Y_i^{\theta})_{i \in A^c}).
$$

□

# <span id="page-6-18"></span>5 Final remarks

We have provided a detailed analysis of the complexity of computing power indices, including scenarios where sets of features of arbitrary size are considered. As shown, the polynomial equivalence between computing the index and computing expected values is not unique to SHAP, but in some instances computing the index might actually be simpler than computing expected values. An open question remains regarding a precise characterization of indices that exhibit this property. While  $q_0 > 0$  guarantees this behavior, it is possible that certain indices with  $q_0 = 0$  might also satisfy it.

# References

- <span id="page-6-10"></span>[Arenas *et al.*, 2023] Marcelo Arenas, Pablo Barceló, Leopoldo E. Bertossi, and Mikaël Monet. On the complexity of shap-score-based explanations: Tractability via knowledge compilation and non-approximability results. *J. Mach. Learn. Res.*, 24:63:1–63:58, 2023.
- <span id="page-6-16"></span>[Beliakov *et al.*, 2020] Gleb Beliakov, Simon James, and J Wu. *Discrete fuzzy measures*. Springer, 2020.
- <span id="page-6-4"></span>[Darwiche, 2023] Adnan Darwiche. Logic for explainable AI. In *LICS*, pages 1–11, 2023.
- <span id="page-6-19"></span>[De Marchi and Usevich, 2014] Stefano De Marchi and Konstantin Usevich. On certain multivariate Vandermonde determinants whose variables separate. *Linear Algebra and its Applications*, 449:17–27, 2014.
- <span id="page-6-1"></span>[Doshi-Velez and Kim, 2017] Finale Doshi-Velez and Been Kim. Towards a rigorous science of interpretable machine learning. *arXiv preprint arXiv:1702.08608*, 2017.
- <span id="page-6-14"></span>[Ferrara *et al.*, 2005] Andrea Ferrara, Guoqiang Pan, and Moshe Y. Vardi. Treewidth in verification: Local vs. global. In *LPAR*, volume 3835 of *Lecture Notes in Computer Science*, pages 489–503. Springer, 2005.
- <span id="page-6-11"></span>[Huang and Marques-Silva, 2024] Xuanxiang Huang and João Marques-Silva. Updates on the complexity of SHAP scores. In *IJCAI*, pages 403–412. ijcai.org, 2024.
- <span id="page-6-7"></span>[Karczmarz *et al.*, 2022] Adam Karczmarz, Tomasz P. Michalak, Anish Mukherjee, Piotr Sankowski, and Piotr Wygocki. Improved feature importance computation for tree models based on the banzhaf value. In James Cussens and Kun Zhang, editors, *UAI*, volume 180, pages 969–979, 2022.
- <span id="page-6-12"></span>[Khosravi *et al.*, 2019] Pasha Khosravi, Yitao Liang, Yoo-Jung Choi, and Guy Van den Broeck. What to expect of classifiers? reasoning about logistic regression with missing features. In Sarit Kraus, editor, *IJCAI*, pages 2716– 2724, 2019.
- <span id="page-6-13"></span>[Khosravi *et al.*, 2020] Pasha Khosravi, Antonio Vergari, YooJung Choi, Yitao Liang, and Guy Van den Broeck. Handling missing data in decision trees: A probabilistic approach. *CoRR*, abs/2006.16341, 2020.
- <span id="page-6-2"></span>[Lipton, 2018] Zachary C Lipton. The mythos of model interpretability. *Queue*, 16(3):31–57, 2018.
- <span id="page-6-5"></span>[Lundberg and Lee, 2017] Scott M. Lundberg and Su-In Lee. A unified approach to interpreting model predictions. In *NeurIPS*, volume 30, pages 4765–4774, 2017.
- <span id="page-6-3"></span>[Marques-Silva and Ignatiev, 2022] João Marques-Silva and Alexey Ignatiev. Delivering trustworthy AI through formal XAI. In *AAAI*, pages 12342–12350, 2022.
- <span id="page-6-15"></span>[Provan and Ball, 1983] J. Scott Provan and Michael O. Ball. The complexity of counting cuts and of computing the probability that a graph is connected. *SIAM J. Comput.*, 12(4):777–788, 1983.
- <span id="page-6-0"></span>[Ribeiro *et al.*, 2016] Marco Tulio Ribeiro, Sameer Singh, and Carlos Guestrin. "why should i trust you?" explaining the predictions of any classifier. In *SIGKDD*, pages 1135–1144, 2016.
- <span id="page-6-6"></span>[Sundararajan and Najmi, 2020] Mukund Sundararajan and Amir Najmi. The many shapley values for model explanation. In *ICML*, volume 119, pages 9269–9278, 2020.
- <span id="page-6-17"></span>[Sundararajan *et al.*, 2020] Mukund Sundararajan, Kedar Dhamdhere, and Ashish Agarwal. The shapley taylor interaction index. In *ICML*, volume 119, pages 9259–9268, 2020.
- <span id="page-6-9"></span>[Van den Broeck *et al.*, 2022] Guy Van den Broeck, Anton Lykov, Maximilian Schleich, and Dan Suciu. On the tractability of SHAP explanations. *Journal of Artificial Intelligence Research*, 74:851–886, 2022.
- <span id="page-6-8"></span>[Wang and Jia, 2023] Jiachen T. Wang and Ruoxi Jia. Data banzhaf: A robust data valuation framework for machine

learning. In Francisco J. R. Ruiz, Jennifer G. Dy, and Jan-Willem van de Meent, editors, *AISTATS*, volume 206, pages 6388–6421, 2023.