Embedding integrable spin models in solvable vertex models on the square lattice.

M.J. Martins

Universidade Federal de São Carlos Departamento de Física C.P. 676, 13565-905, São Carlos (SP), Brazil

Abstract

Exploring a mapping among *n*-state spin and vertex models on the square lattice we argue that a given integrable spin model with edge weights satisfying the rapidity difference property can be formulated in the framework of an equivalent solvable vertex model. The Lax operator and the R-matrix associated to the vertex model are built in terms of the edge weights of the spin model and these operators are shown to satisfy the Yang-Baxter algebra. The unitarity of the R-matrix follows from an assumption that the vertical edge weights of the spin model satisfy certain local identity known as inversion relation. We apply this embedding to the scalar *n*-state Potts model and we argue that the corresponding R-matrix can be written in terms of the underlying Temperley-Lieb operators. We also consider our construction for the integrable Ashkin-Teller model and the respective R-matrix is expressed in terms of sixteen distinct weights parametrized by theta functions. We comment on the possible extention of our results to spin models whose edge weights are not expressible in terms of the difference of spectral parameters.

Keywords: Spin and vertex models, Integrability, Yang-Baxter equation

January 2025

1 Introduction

An important class of integrable two-dimensional lattice system of statistical mechanics are models in which the microstates variables or spins are located on the vertices of the lattice and the interactions occur along the lattice edges [1]. These systems are called spin or edge models and the simplest example is a two state model known as the Ising ferromagnet which was originally solved by Onsager in 1944 [2, 3]. Generalizations of Ising model are achieved considering that the spins take values on set of integer and the energy interactions are strictly short ranged depending only on the nearest neighbors spins states. On the square lattice the energy interactions among two neighboring spins (i, j) can be encoded in terms of local horizontal $W_h(i, j|x)$ and vertical $W_v(i, j|x)$ edge weights which here we assumed to be parametrized by the spectral variable x. The edge weights are schematically shown in Fig.(1) where the spin states are considered to take values on a discrete set $\{1, \ldots, n\}$.



Figure 1: Schematic representation of the horizontal $W_h(i, j|x)$ and the vertical $W_v(j, k|x)$ local Boltzmann weights of the spin models where (i, j, k) indicate spin states.

One of the most successful technique in solving two-dimensional lattice models turns out to be the method of commuting transfer matrices [1]. The use of transfer matrices to solve two-dimensional lattice models originates in the work of Kramers and Wannier [4] who have shown that the respective partition functions can be rewritten as trace of an ordered product of transfer matrices. For spin models on the square lattice a convenient transfer matrix operator is built by considering the product of the weights along a diagonal layer of the square lattice [5, 6]. The matrix elements of the diagonalto-diagonal transfer matrix for a lattice of size L with periodic boundary conditions are,

$$[T_{\text{dia}}(x)]_{a_1,\dots,a_L}^{b_1,\dots,b_L} = W_v(a_1,b_1|x)W_h(a_1,b_2|x)W_v(a_2,b_2|x)W_h(a_2,b_3|x)\dots W_v(a_L,b_L|x)W_h(a_L,b_1|x), \quad (1)$$

which have been schematically illustrated in Fig.(2).

It follows that the spin model partition function $Z_{\rm spin}(L)$ on the square lattice of size $L \times L$ can



Figure 2: Schematic representation of the diagonal-to-diagonal transfer matrix of spin models.

be written in terms of the following trace,

$$Z_{\rm spin}(L) = {\rm Tr}_{\mathcal{V}}\left[\left(T_{\rm dia}(x) \right)^L \right], \qquad (2)$$

where $\mathcal{V} = \prod_{j=1}^{L} \otimes \mathbb{C}^{n}$ is usually called the spin model quantum space.

Given a spin model with weights $W_{h,v}(i, j|x)$ there exist sufficient conditions for the existence of another spin model with distinct weights $W_{h,v}(i, j|y)$ such that their diagonal-to-diagonal transfer matrices commutes, $[T_{\text{dia}}(x), T_{\text{dia}}(y)] = 0$, for arbitrary lattice sizes. These are local relations among the horizontal and vertical weights originally discovered in the context of the Ising model [2, 7] which are often called star-triangle equations, see for instance refs.[8, 9, 10]. For spin models with weights depending on the difference of spectral parameters these set of relations are given by,

$$\sum_{d=1}^{n} W_{v}(d,c|y)W_{v}(b,d|x-y)W_{h}(a,d|x) = \mathcal{R}(x,y)W_{h}(a,b|y)W_{h}(a,c|x-y)W_{v}(b,c|x), \quad (3)$$

$$\sum_{d=1}^{n} W_{v}(c,d|y)W_{v}(d,b|x-y)W_{h}(d,a|x) = \mathcal{R}(x,y)W_{h}(b,a|y)W_{h}(c,a|x-y)W_{v}(c,b|x), \quad (4)$$

 $\mathcal{R}(x, y)$ denotes some factor independent of the spin states $a, b, c = 1, \ldots, n$. We remark that many solvable spin models indeed satisfy Eqs.(3,4) being notable examples the scalar Potts [11, 12, 13], the self-dual Ashkin-Teller [14, 15, 16, 17], the Fateev-Zamolodchikov [18], and the Kashiwara-Miwa [19, 20, 21] models.

It turns out that recently, we have argued that a *n*-state spin model can be viewed as a *n*-state vertex model in the sense that their partition functions coincide on the finite square lattice with periodic boundary conditions [22]. In particular, we have exhibited the expression of the Lax operator encoding the weights of the equivalent vertex model in terms of a special combination of the horizontal and vertical edge weights of the spin model. Therefore, assuming that the spin model is also integrable one expects that the same property should be extended to the equivalent vertex model. In

the case of a *n*-state vertex model, integrability is assured by exhibiting an invertible $n^2 \times n^2$ R-matrix which together with the Lax operator has to satisfy a quadratic algebra denominated Yang-Baxter algebra [23, 24]. We also remark that the determination of the R-matrix is an essential ingredient to formulate the exact solution of the vertex model in terms of the quantum inverse scattering method [24]. In this work, we argue that the matrix elements of the R-matrix can be constructed from the spin model edge weights and the respective Yang-Baxter algebra follows from the star-triangle equations (3,4). More precisely, we shall show that the expression of the underlying the R-matrix is,

$$R_{12}(x,y) = \sum_{i,j,k=1}^{n} \frac{W_h(j,i|x)W_v(j,k|x-y)}{W_h(k,i|y)} e_{ik} \otimes e_{ji},$$
(5)

where $e_{i,j}$ denotes the $n \times n$ matrix with only one non-vanishing entry with value 1 at row *i* and column *j*. We note that the R-matrix is not given solely in terms of the difference of the spectral parameters. This fact can be explored to generate generalizations of the one-dimensional quantum spin chains underlying the classical lattice spin models.

We have organized this paper as follows. In next section we elaborate on our previous results concerning the mentioned mapping among *n*-state spin and vertex models on the square lattice [22]. Here we note that Lax operator of the equivalent vertex model can be decomposed in terms of the product of two operators depending either on the horizontal or vertical edge weights. In section 3 we formulate the Yang-Baxter algebra and show that this algebra is satisfied as a consequence of the star-triangle relations for the spin model edge weights. We also discuss certain properties of the R-matrix such as unitarity relation and the Yang-Baxter equation. In section 4 we apply our results for one of the simplest integrable *n*-state spin model which is the scalar Potts model. We argue that the respective R-matrix of the equivalent vertex model can be rewritten in terms of Temperley-Lieb operators [26]. In section 5 we consider our construction for the integrable Ashkin-Teller model and show that the corresponding R-matrix elements can be expressed in terms of sixteen distinct weights. In section 6 we present our conclusion and comment on the possibility of extending our embedding to include integrable spin models whose edge weights are not parametrized in terms of the difference of spectral parameters. In Appendix A we present some technical details about the Yang-Baxter equation omitted in the main text.

2 The spin-vertex equivalence

In vertex models the state variables are assigned to the links of the square lattice and the Boltzmann weights depend on four spin variables i, j, k, l meeting together at the vertex. We express these weights by w(i, k; j, l|x) where the variable x represent the spectral parameter as illustrated in Fig.(3).



Figure 3: The local Boltzmann weights of vertex models with spin variables i, j, k, l taken values on the set $\{1, \ldots, n\}$.

For the vertex models with periodic boundary conditions it is convenient to built the transfer matrix considering the product of weights defined by two successive rows of spins states of the lattice. The matrix elements of the row-to-row transfer matrix are given by,

$$[T_{\rm ver}(x)]_{a_1,\dots,a_L}^{b_1,\dots,b_L} = \sum_{c_1,\dots,c_L}^n w(c_1,c_2;a_1,b_1|x)w(c_2,c_3;a_2,b_2|x)\dots w(c_L,c_1;a_L,b_L|x),\tag{6}$$

which is schematically illustrated in Fig.(4).



Figure 4: Schematic representation of the row-to-row transfer matrix of vertex models.

Once again the vertex model partition function $Z_{ver}(L)$ on the $L \times L$ square lattice is given by the trace of the *L*-th power of the row-to-row transfer matrix,

$$Z_{\rm ver}(L) = {\rm Tr}_{\mathcal{V}}\left[\left(T_{\rm ver}(x) \right)^L \right].$$
(7)

As argued in ref.[22] the partition functions of the *n*-state spin (2) and vertex (7) models coincide for a finite square lattice provide that the corresponding weights satisfy the following relation,

$$w(i,k;j,l|x) = W_v(j,k|x)W_h(j,i|x)\delta_{i,l}.$$
(8)

In fact, by substituting the vertex weights (8) in the row-to-row transfer matrix (6) and after summing over the Kronecker's delta symbols we obtain,

$$[T_{\text{ver}}(x)]_{a_1,\dots,a_L}^{b_1,\dots,b_L} = W_h(a_1,b_1|x)W_v(a_1,b_2|x)W_h(a_2,b_2|x)W_v(a_2,b_3|x)\dots W_h(a_L,b_L|x)W_v(a_L,b_1|x), \quad (9)$$

and we note that these matrix elements have the basic form of those of the diagonal-to-diagonal transfer matrix (1) except by the fact that the horizontal and vertical edge weights are interchanged. However, this exchange among the weights corresponds to rotation of the lattice by 90^{0} degrees and such operation does not affect the spin model partition function on the square lattice with periodic boundary conditions.

One interesting feature of the vertex models is that they have an inherent tensor structure. The Boltzmann weights can be encoded in terms of a local vertex operator usually called Lax operator which acts on the tensor product of two spaces associated to the horizontal and vertical spin variables. It turns out that the Lax operator can be written as follows,

$$\mathbb{L}_{12}(x) = \sum_{i,j,k,l=1}^{n} w(i,k;j,l|x)e_{ik} \otimes e_{jl} = \sum_{i,j,k=1}^{n} W_h(j,i|x)W_v(j,k|x)e_{ik} \otimes e_{ji},$$
(10)

and as result the respective row-to-row transfer matrix can be compactly expressed in terms of trace over an auxiliary space $\mathcal{A} \equiv C^n$, namely

$$T_{\rm ver}(x) = {\rm Tr}_{\mathcal{A}} \left[\mathbb{L}_{\mathcal{A}1}(x) \mathbb{L}_{\mathcal{A}2}(x) \dots \mathbb{L}_{\mathcal{A}L}(x) \right].$$
(11)

In our case we observe that the Lax operator can be decomposed in terms of product of two operators whose entries depend either on the horizontal or on the vertical spin edge weights. In fact, it is possible to the rewrite the Lax operator as $\mathbb{L}_{12}(x) = \mathbb{L}_{12}^{(h)}(x)\mathbb{L}_{12}^{(v)}(x)$ such that the components are given by,

$$\mathbb{L}_{12}^{(h)}(x) = \sum_{i,j=1}^{n} W_h(j,i|x) e_{ii} \otimes e_{jj}, \quad \mathbb{L}_{12}^{(v)}(x) = \sum_{i,j,k=1}^{n} W_v(j,k|x) e_{ik} \otimes e_{ji}, \tag{12}$$

where we note that the operator associated to the horizontal weights is a diagonal matrix.

We next would like to discuss the Hamiltonian limit associated to the *n*-state spin or vertex models. To this end we assume the existence of a particular spectral parameter point, say x = 0, in which the edge weights satisfy the following initial conditions,

$$W_h(i,j|0) = 1, \quad W_v(i,j|0) = \delta_{i,j},$$
(13)

implying that at x = 0 the Lax operator becomes the permutator on the tensor product $C^n \otimes C^n$,

$$\mathbb{L}_{12}(0) = P_{12} = \sum_{i,j=1}^{n} e_{ij} \otimes e_{ji}.$$
(14)

Considering the condition (14) the corresponding quantum spin chain is obtained by taking the logarithm derivative of the row-to-row transfer matrix (11) at the special point x = 0 [23]. This leads us to the following Hamiltonian,

$$H = \sum_{j=1}^{L} \left[\sum_{i,k=1}^{n} \frac{d}{dx} [W_h(i,k|x)] \Big|_{x=0} e_{ii}^{(j)} \otimes e_{kk}^{(j+1)} + \sum_{i,k,l=1}^{n} \frac{d}{dx} [W_v(i,k|x)] \Big|_{x=0} e_{ik}^{(j)} \otimes e_{ll}^{(j+1)} \right],$$
(15)

where $e_{ik}^{(j)}$ denotes the action of the matrix e_{ik} on the *j*-th site of a chain of size *L* and periodic boundary condition is imposed by defining $e_{ik}^{(L+1)} = e_{ik}^{(1)}$.

We would like to conclude by mentioning certain local identities satisfied by edge weights of solvable spin models such as the scalar Potts [11, 12, 13], the self-dual Ashkin-Teller [14, 15, 16, 17], the Fateev-Zamolodchikov [18], and the Kashiwara-Miwa [19, 20, 21] models. These identities are usually called inversion relations [1] and their expressions are given by,

$$W_h(i,j|x)W_h(i,j|-x) = \rho_1(x), \quad \sum_{k=1}^n W_v(i,k|x)W_v(k,j|-x) = \rho_2(x)\delta_{i,j}, \tag{16}$$

where $\rho_{1,2}(x)$ are arbitrary auxiliary normalization factors.

In next section, we argue that the inversion relation associated to the vertical edge weights is relevant to set up the unitarity property of the proposed R-matrix (5). This feature assures that the underlying R-matrix is invertible for almost all spectral parameters x and y.

3 The Yang-Baxter algebra

If the spin model is integrable it is natural to expect that this property will be also satisfied by the equivalent vertex model. An sufficient condition for commuting transfer matrices of vertex models was first introduced by Baxter in his analysis of the eight-vertex model [23] and afterwards elaborated in the context of the quantum inverse scattering approach [24]. This condition requires the existence an invertible R-matrix which together with the Lax operators satisfies the Yang-Baxter algebra. This algebra involves the action of these operators on the tensor product of three n-dimensional spaces, namely

$$R_{12}(x,y)\mathbb{L}_{13}(x)\mathbb{L}_{23}(y) = \mathbb{L}_{23}(y)\mathbb{L}_{13}(x)R_{12}(x,y),$$
(17)

where the lower indices indicate the spaces in which the Lax and the R-matrix operators acts nontrivially.

We observe that in the above relation the Lax operators are fixed by the edge weights of the spin model, see Eq.(10). In order to show that this algebra is satisfied by the R-matrix (5) we find convenient to rewrite the Yang-Baxter algebra (17) in terms of its components. To this end we express the R-matrix (5) as,

$$R_{12}(x,y) = \sum_{i,j,k,l} R_{i,j}^{k,l}(x,y) e_{ik} \otimes e_{jl}, \quad R_{i,j}^{k,l}(x,y) = \frac{W_h(j,i|x)W_v(j,k|x-y)}{W_h(k,i|y)} \delta_{i,l},$$
(18)

and by using the expression of the Lax operator (10) we find that the matrix elements of the Yang-Baxter algebra are,

$$\sum_{\gamma=1}^{n} \mathcal{R}_{a_{1},a_{2}}^{\gamma,b_{3}}(x,y) W_{h}(a_{3},\gamma|x) W_{v}(a_{3},b_{1}|x) W_{h}(\gamma,b_{3}|y) W_{v}(\gamma,b_{2}|y) = \delta_{a_{1},b_{3}} \sum_{\gamma,\gamma'=1}^{n} \mathcal{R}_{\gamma,\gamma'}^{b_{1},b_{2}}(x,y) W_{h}(a_{3},a_{2}|y) W_{v}(a_{3},\gamma'|y) W_{h}(a_{2},a_{1}|x) W_{v}(a_{2},\gamma|x).$$
(19)

In what follows we shall show that the Yang-Baxter algebra (19) is indeed fulfilled. We first substitute on the left hand side of Eq.(19) the structure constants $R_{a_1,a_2}^{\gamma,b_3}(x,y)$ and sum over the Kronecker's delta symbols. Further simplification can be carried out by using the first set of the star-triangle relations (3). As a result of these steps we obtain,

$$\sum_{\gamma=1}^{n} \mathcal{R}_{a_{1},a_{2}}^{\gamma,b_{3}}(x,y) W_{h}(a_{3},\gamma|x) W_{v}(a_{3},b_{1}|y) W_{h}(\gamma,b_{3}|y) W_{v}(\gamma,b_{2}|y) = \delta_{a_{1},b_{3}} W_{v}(a_{3},b_{1}|x) W_{h}(a_{2},a_{1}|x) \sum_{\gamma=1}^{n} W_{v}(\gamma,b_{2}|y) W_{h}(a_{2},\gamma|x-y) W_{h}(a_{3},\gamma|x) = \delta_{a_{1},b_{3}} W_{v}(a_{3},b_{1}|x) W_{h}(a_{2},a_{1}|x) \mathcal{R}(x,y) W_{h}(a_{3},a_{2}|y) W_{h}(a_{3},b_{2}|x-y) W_{v}(a_{2},b_{2}|x).$$
(20)

We now repeat the same procedure explained above for the right hand side of Eq.(19). Here we use the second set of the star-triangle relations (4) to simplify the sum of product of edge weights.

The steps are summarized below,

$$\delta_{a_1,b_3} \sum_{\gamma,\gamma'=1}^{n} \mathcal{R}_{\gamma,\gamma'}^{b_1,b_2}(x,y) W_h(a_3,a_2|y) W_v(a_3,\gamma'|y) W_h(a_2,a_1|x) W_v(a_2,\gamma|x) = \\ \delta_{a_1,b_3} \frac{W_h(a_2,a_1|x) W_h(a_3,a_2|y) W_v(a_2,b_2|x)}{W_h(b_1,b_2|y)} \sum_{\gamma'=1}^{n} W_v(a_3,\gamma'|y) W_h(\gamma',b_1|x-y) W_h(\gamma',b_2|x) = \\ \delta_{a_1,b_3} W_h(a_2,a_1|x) W_h(a_3,a_2|y) W_v(a_2,b_2|x) \mathcal{R}(x,y) W_h(a_3,b_2|x-y) W_v(a_3,b_1|x),$$
(21)

and by comparing Eqs.(20,21) we see that the left and the right hand sides of the Yang-Baxter algebra are the same.

The other important step for commuting transfer matrices is the assumption that the underlying R-matrix is invertible for almost all spectral parameters x and y. In our case this feature follows from the fact that the R-matrix (18) satisfy the unitarity property,

$$R_{12}(x,y)R_{21}(y,x) = \rho_2(x-y)I_n \otimes I_n,$$
(22)

where I_n denotes the $n \times n$ identity matrix. It turns out that this property can be derived from direct computation and with the help of the inversion relation for the vertical edge weights (16), namely

$$R_{12}(x,y)R_{21}(y,x) = \sum_{i,j,l=1}^{n} \frac{W_h(j,i|x)}{W_h(l,i|x)} \sum_{k=1}^{n} [W_v(j,k|x-y)W_v(k,l|y-x)] e_{ii} \otimes e_{j,l}$$

$$= \sum_{i,j,l=1}^{n} \frac{W_h(j,i|x)}{W_h(l,i|x)} \rho_2(x-y) \delta_{j,l} e_{ii} \otimes e_{j,l}$$

$$= \rho_2(x-y) \sum_{i,j=1}^{n} e_{ii} \otimes e_{j,j} = \rho_2(x-y) I_n \otimes I_n.$$
(23)

At this point we have discussed the basic ingredients showing that the equivalent *n*-state vertex model associated to an integrable *n*-state spin model indeed gives rise to a family of commuting rowto-row transfer matrices. In the context of vertex models is assumed that the Yang-Baxter algebra is an associative algebra when you reorder the product of three Lax operators with distinct rapidities x, y and z. It is well known that a sufficient condition for the associativity property is the celebrated Yang-Baxter equation,

$$R_{12}(x,y)R_{13}(x,z)R_{23}(y,z) = R_{23}(y,z)R_{13}(x,z)R_{12}(x,y),$$
(24)

where at z = 0 reduces to the Yang-Baxter algebra (17) since we have the identity $R_{ab}(x, 0) = L_{ab}(x)$ as a consequence of the initial condition (13). It is plausible to think that the Yang-Baxter equation (24) should follows from systematic applications of the star-triangle equations (3,4) combined with the help of the inversion relations (16). In this sense, we remark that it has been argued that from a particular combination of four edge weights of a given solvable spin model one is able to obtain the Boltzmann weights of a vertex model with commuting transfer matrices [10, 25]. In this construction, the number of spectral variables of the vertex model weights is duplicated due to the fact it is used two distinct sets of rapidities to parametrized of horizontal and vertical spin edge weights. We now follow the construction of refs.[10, 25] in the situation where the spin edge weights depend only on the difference of the spectral parameters. In our notation, the matrix elements of the Lax operator associated to such vertex model are,

$$\widetilde{\mathbb{L}}_{i,j}^{k,l}(x_1; x_2, y_1; y_2) = W_h(i, j | x_1 - y_1) W_v(j, k | x_1 - y_2) W_v(i, l | x_2 - y_1) W_h(l, k | x_2 - y_2)$$
(25)

where x_1, y_1 and x_2, y_2 denote two pairs of rapidities.

We next apply the expression (25) in a particular case of rapidities arrangements by choosing $x_2 = y_1$, $x_1 = y_1 + x$ and $y_2 = y_1 + y$ where x and y are arbitrary variables. By using the initial condition (13) for the vertical edge weights as well as the inversion relation (16) for the horizontal edge weights we obtain,

$$\widetilde{\mathbb{L}}_{i,j}^{k,l}(y_1 + x; y_1, y_1; y_1 + y) = W_h(i, j|x) W_v(j, k|x - y) W_h(l, k| - y) \delta_{i,l}
= \rho_1(y) \frac{W_h(i, j|x) W_v(j, k|x - y)}{W_h(l, k|y)} \delta_{i,l}$$
(26)

By comparing the above elements with the entries of the proposed R-matrix (18) we note that they are similar except by the fact that the spin states of the horizontal edge weights are exchanged. At this point we recall that the most known solvable spin models with weights parametrized by the difference of spectral parameters have the reflection symmetry $W_{h,v}(i, j|x) = W_{h,v}(j, i|x)$. Therefore, we expect that the R-matrices (18) associated to the scalar Potts, the self-dual Ashkin-Teller the Fateev-Zamolodchikov, and the Kashiwara-Miwa spin models should indeed satisfy the Yang-Baxter equation (24). In fact, we have confirmed this relation for the above mentioned spin models by using the explicit expressions of the corresponding edge weights. The technical details concerning this verification have been summarized in Appendix A.

The validity of the Yang-Baxter equation for a given R-matrix which is not of difference form has the following interesting consequence. We can formulated a generalized integrable n-state vertex model by replacing the Lax operator by the R-matrix in the row-to-row transfer matrix,

$$T(x, x_0) = \operatorname{Tr}_{\mathcal{A}} \left[\operatorname{R}_{\mathcal{A}1}(x, x_0) \operatorname{R}_{\mathcal{A}2}(x, x_0) \dots \operatorname{R}_{\mathcal{A}L}(x, x_0) \right],$$
(27)

where the second spectral parameter x_0 plays the role of an additional independent coupling of the model. We note that for $x_0 = 0$ we recover the transfer matrix of the *n*-state vertex model equivalent to a *n*-state spin model, see Eq.(11)

We next observe that for $x = x_0$ each operator $\mathbb{R}_{\mathcal{A}_j}(x, x_0)$ equals the permutator on the corresponding tensor product spaces $C^n \otimes C_j^n$. We can therefore construct a generalized quantum spin chain by taking the logarithmic derivative of the transfer matrix (27) at the point $x = x_0$. The expression of the respective Hamiltonian is,

$$H(x_{0}) = \sum_{j=1}^{L} \sum_{i,k=1}^{n} \left[\frac{1}{W_{h}(i,k|x_{0})} \frac{d}{dx} [W_{h}(i,k|x)] \Big|_{x=x_{0}} e_{ii}^{(j)} \otimes e_{kk}^{(j+1)} \right] \\ + \sum_{j=1}^{L} \left[\sum_{i,k,l=1}^{n} \frac{W_{h}(i,l|x_{0})}{W_{h}(k,l|x_{0})} \frac{d}{dx} [W_{v}(i,k|x-x_{0})] \Big|_{x=x_{0}} e_{ik}^{(j)} \otimes e_{ll}^{(j+1)} \right],$$
(28)

where periodic boundary condition is assumed. For $x_0 \neq 0$ the above Hamiltonian generalizes the quantum spin chain associated to the lattice *n*-state spin or vertex models, see Eq.(15).

In next sections we apply the above results in the cases of the solvable scalar Potts and the Ashkin-Teller spin models.

4 The scalar Potts model

The *n*-state scalar Potts model is a generalization of the Ising model when at each *i*-th site the spin variables σ_i can have $n \ge 2$ possible values. It is assumed that the interactions among the horizontal (vertical) adjacent spins variables have the same thermal energy $J_h(J_v)$ when the respective spins variables are alike and zero if they are different [11]. The total thermal energy of this spin model is given by,

$$\frac{E}{k_B T} = -J_v \sum_{\langle i,j \rangle \rangle} \delta(\sigma_i, \sigma_j) - J_v \sum_{\langle k,l \rangle} \delta(\sigma_k, \sigma_l),$$
(29)

where k_B is Boltzmann's constant and T is the temperature. The symbols $\langle i, j \rangle$ and $\langle k, l \rangle$ indicate the summations over all the horizontal and vertical edges of the square lattice, respectively.

This means that the edge weights have only two basic elements and they can be written as follows,

$$W_h(i,j) = \kappa_v \left[(\exp(J_h) - 1)\delta_{i,j} + 1 \right], \quad W_v(i,j) = \kappa_h \left[(\exp(J_v) - 1)\delta_{i,j} + 1 \right], \tag{30}$$

where $\kappa_{h,v}$ are normalization factors at our disposal. This model cannot be solved in general, but it is integrable when the couplings sit on the self-dual manifold [12],

$$(\exp(J_h) - 1)(\exp(J_v) - 1) = n.$$
(31)

It turns out that one possible parametrization of such solvable manifold is given as follows [12, 13],

$$W_h(i,j|x) = 1 + \sqrt{n} f_n(x) \delta_{i,j}, \quad W_v(i,j|x) = \frac{f_n(x)}{\sqrt{n}} + \delta_{i,j}, \tag{32}$$

where the function $f_n(x)$ is defined by,

and the normalization functions of the inversion relations are trivial, i.e $\rho_1(x) = \rho_2(x) = 1$.

The above parametrization can be traced back to the fact that the *n*-state Potts model can be seen as one of the possible representations of the Temperley-Lieb algebra [26]. In this context, the function $f_n(x)$ arises as a solution of certain functional equation associated to the Baxterization of a braid originated from the Temperley-Lieb monoid [27]. For the scalar Potts model the corresponding Temperley-Lieb generators can be expressed in terms the basic elements of the Z(n) algebra,

$$Z^n = X^n = 1, \quad ZX = \omega XZ, \tag{34}$$

where $\omega = \exp(2\pi i/n)$ and the entries of the $n \times n$ matrices Z and X are,

$$Z_{k,l} = \omega^{k-1} \delta_{k,l}, \quad X_{k,l} = \delta_{k,l+1} \pmod{n}.$$
(35)

By now the *n*-state scalar Potts representation of the Temperley-Lieb algebra is well known in the literature, see for instance [28]. In terms of the Z(n) operators such representation is given by,

$$E_{2j} = \frac{1}{\sqrt{n}} \sum_{k=0}^{n-1} \left(Z_j Z_{j+1}^{\dagger} \right)^k, \quad E_{2j-1} = \frac{1}{\sqrt{n}} \sum_{k=0}^{n-1} \left(X_j \right)^k, \tag{36}$$

where the operators Z_j and X_j acting at the *j*-th site of the chain obey the Z(n) algebra. These generators satisfy the following algebraic relations,

$$(E_j)^2 = \sqrt{n}E_j, \quad E_j E_{j\pm 1}E_j = E_j, \quad [E_j, E_k] = 0 \text{ for } |l-k| \ge 2.$$
 (37)

From the above discussion we see that for the scalar Potts model the most natural operators to use are the Temperley-Lieb generators (36). Therefore, one expects that both the Lax operator of the equivalent vertex model and the respective R-matrix satisfying the Yang-Baxter algebra should be rewritten in terms of the Temperley-Lieb generators. In fact, the expression for the Lax operator (10) is,

$$\mathbb{L}_{12}(x) = P_{12} \Big(\mathbf{I}_n \otimes \mathbf{I}_n + f_n(x) E_2 \Big) \Big(\mathbf{I}_n \otimes \mathbf{I}_n + f_n(x) E_1 \Big),$$
(38)

while the corresponding R-matrix (18) is given by,

$$R_{12}(x,y) = P_{12} \Big(I_n \otimes I_n + f_n(x-y) \left[E_1 + E_2 + f_n(x) E_2 E_1 + f_n(-y) E_1 E_2 \right] \Big).$$
(39)

By using trigonometric identities and the properties of the Temperley-Lieb operators one can verify that the R-matrix (39) indeed satisfy the Yang-Baxter equation. Interesting enough, this result tells us that a particular combination of two different Temperley-Lieb operators appears to be suitable for Baxterization once we consider that the R-matrix can not be parametrized only in terms of the difference of rapidities. As explained at the end of previous section we can use this R-matrix to generated a generalized Z(N) invariant quantum spin chain and as usual we write the respective Hamiltonian as,

$$H(x_0) = -J \Big[\sum_{j=1}^{L-1} H_{j,j+1}(x_0) + H_{L,1}(x_0) \Big],$$
(40)

where J is an overall normalization. The corresponding two-body Hamiltonian $H_{j,j+1}$, apart from an additive term, is given in terms of the Z(n) operators by the following expression¹,

$$H_{j,j+1}(x_0) = \sum_{k=1}^{n-1} (X_j)^k + \sum_{k=1}^{n-1} (Z_j Z_{j+1}^{\dagger})^k + g_n(x_0) \sum_{k,l=1}^{n-1} (Z_j Z_{j+1}^{\dagger})^k (X_j)^l + g_n(-x_0) \sum_{k,l=1}^{n-1} (X_j)^k (Z_j Z_{j+1}^{\dagger})^l,$$
(41)

where the function $g_n(x)$ is given by,

$$g_n(x) = \begin{cases} \frac{\sin(x)\sin(\gamma_n + x)}{\sin(2\gamma_n)\sin(\gamma_n)} & \text{for } n = 2, 3\\ \frac{x(1+x)}{2} & \text{for } n = 4\\ \frac{\sinh(x)\sinh(\gamma_n + x)}{\sinh(2\gamma_n)\sinh(\gamma_n)} & \text{for } n \ge 5. \end{cases}$$
(42)

The Hamiltonian defined by Eqs.(40,41,42) is an integrable Z(n) symmetric deformation of the quantum spin chain associated to the scalar Potts model. We note that this operator is Hermitian

¹We observe that for n = 2, 3 we recover the spin chains discussed in our previous work [22].

when x_0 is an imaginary number. The first two terms of the Hamiltonian correspond to the standard spin chain obtained within the so-called time-continuum limit of a certain transfer matrix formulation of the classical scalar Potts model [29, 30] while the last two terms are additional interactions among particular combinations of Z(n) generators. We recall that these type of extra interactions have appeared before in a quantum spin chain derived from the integrable higher-spin XXZ Heisenberg model for a specific choice of the quantum group deformation parameter [31]. We remark, however, that in the model of ref.[31] the extra interactions are weighted by suitable factors such that the underlying Hamiltonian is U(1)-invariant.

5 The Ashkin-Teller model

The Ashkin-Teller model [14] can be formulated in terms of two Ising models with thermal energies $J_{v,h}$ and $K_{v,h}$ which are coupled by a four spin interactions $L_{h,v}$ involving the product of the energy terms of the Ising models [32]. If we denote the Ising spins at a given *i*-th site by the variables σ_i and τ_i taking the values ± 1 the respective thermal energy can be written as follows,

$$\frac{E}{k_B T} = -\sum_{\langle i,j \rangle} \left(J_h \sigma_i \sigma_j + K_h \tau_i \tau_j + L_h \sigma_i \sigma_j \tau_i \tau_j \right) - \sum_{\langle k,l \rangle} \left(J_v \sigma_k \sigma_l + K_v \tau_k \tau_l + L_v \sigma_k \sigma_l \tau_k \tau_l \right),$$
(43)

where the sums $\langle i, j \rangle$ and $\langle k, l \rangle$ are over the horizontal and vertical edges of the square lattice.

This means that the Ashkin-Teller is a four-state spin model and the edge weights can be represented by the following 4×4 matrices,

$$W_{h} = \begin{pmatrix} a_{h} & b_{h} & c_{h} & d_{h} \\ b_{h} & a_{h} & d_{h} & c_{h} \\ c_{h} & d_{h} & a_{h} & b_{h} \\ d_{h} & c_{h} & b_{h} & a_{h} \end{pmatrix}, \quad W_{v} = \begin{pmatrix} a_{v} & b_{v} & c_{v} & d_{v} \\ b_{v} & a_{v} & d_{v} & c_{v} \\ c_{v} & d_{v} & a_{v} & b_{v} \\ d_{v} & c_{v} & b_{v} & a_{v} \end{pmatrix},$$
(44)

where the relation among the edge weights with the couplings are,

$$a_{h} = \kappa_{h} e^{(J_{h} + K_{h} + L_{h})}, \quad b_{h} = \kappa_{h} e^{(J_{h} - K_{h} - L_{h})}, \quad c_{h} = \kappa_{h} e^{(-J_{h} + K_{h} - L_{h})}, \quad d_{h} = \kappa_{h} e^{(-J_{h} - K_{h} + L_{h})},$$

$$a_{v} = \kappa_{v} e^{(J_{v} + K_{v} + L_{v})}, \quad b_{v} = \kappa_{v} e^{(J_{v} - K_{h} - L_{v})}, \quad c_{v} = \kappa_{v} e^{(-J_{v} + K_{v} - L_{v})} \quad d_{v} = \kappa_{v} e^{(-J_{v} - K_{v} + L_{v})}, \quad (45)$$

such that $\kappa_{h,v}$ are arbitrary normalizations factors.

It has been argued that the Ashkin-Teller spin model can be converted into a staggered eight vertex model on the square lattice [15] which becomes integrable when the vertex weights on the two sublattices are proportional [1]. If we denote by w_a , w_b , w_c and w_d the weights of the eight-vertex model it turns out that the edge weights of the Ashkin-Teller spin model are given by [16],

$$a_{h} = 1, \quad b_{h} = \frac{w_{a} - w_{d}}{w_{b} + w_{c}}, \quad c_{h} = \frac{w_{a} + w_{d}}{w_{b} + w_{c}}, \quad d_{h} = \frac{w_{c} - w_{b}}{w_{b} + w_{c}}$$
$$a_{v} = 1, \quad b_{v} = \frac{w_{b} - w_{d}}{w_{a} + w_{c}}, \quad c_{v} = \frac{w_{b} + w_{d}}{w_{a} + w_{c}}, \quad d_{v} = \frac{w_{c} - w_{a}}{w_{a} + w_{c}}, \tag{46}$$

where we have normalized the horizontal and vertical edge weights by a_h and a_v , respectively.

It is well known that weights of the eight-vertex model can be uniformized in terms of the theta elliptic functions [1]. By using this parametrization one finds that the spin edge weights can be expressed as follows [17],

$$a_{h}(x) = 1, \qquad a_{v}(x) = 1,$$

$$b_{h}(x) = \frac{\theta_{1}\left(\frac{\xi-x}{2},q\right)\theta_{3}\left(\frac{\xi+x}{2},q\right)}{\theta_{3}\left(\frac{\xi-x}{2},q\right)\theta_{1}\left(\frac{\xi+x}{2},q\right)}, \qquad b_{v}(x) = \frac{\theta_{1}\left(\frac{x}{2},q\right)\theta_{3}\left(\xi-\frac{x}{2},q\right)}{\theta_{3}\left(\frac{x}{2},q\right)\theta_{1}\left(\xi-\frac{x}{2},q\right)},$$

$$c_{h}(x) = \frac{\theta_{1}\left(\frac{\xi-x}{2},q\right)\theta_{4}\left(\frac{\xi+x}{2},q\right)}{\theta_{4}\left(\frac{\xi-x}{2},q\right)\theta_{1}\left(\frac{\xi+x}{2},q\right)}, \qquad c_{v}(x) = \frac{\theta_{1}\left(\frac{x}{2},q\right)\theta_{4}\left(\xi-\frac{x}{2},q\right)}{\theta_{4}\left(\frac{x}{2},q\right)\theta_{1}\left(\xi-\frac{x}{2},q\right)},$$

$$d_{h}(x) = \frac{\theta_{1}\left(\frac{\xi-x}{2},q\right)\theta_{1}\left(\frac{\xi+x}{2},q\right)}{\theta_{2}\left(\frac{\xi-x}{2},q\right)\theta_{1}\left(\frac{\xi+x}{2},q\right)}, \qquad d_{v}(x) = \frac{\theta_{1}\left(\frac{x}{2},q\right)\theta_{2}\left(\xi-\frac{x}{2},q\right)}{\theta_{2}\left(\frac{x}{2},q\right)\theta_{1}\left(\xi-\frac{x}{2},q\right)}, \qquad (47)$$

where ξ is an arbitrary parameter and $\theta_i(x,q)$ i = 1, ..., 4 are the four standard theta functions of nome q, with |q| < 1, defined by [33],

$$\theta_{1}(x,q) = 2q^{1/4}\sin(x)\prod_{k=1}^{\infty}(1-2q^{2k}\cos(2x)+q^{4k})(1-q^{2k})$$

$$\theta_{2}(x,q) = 2q^{1/4}\cos(x)\prod_{k=1}^{\infty}(1+2q^{2k}\cos(2x)+q^{4k})(1-q^{2k})$$

$$\theta_{3}(x,q) = \prod_{k=1}^{\infty}(1+2q^{2k-1}\cos(2x)+q^{4k-2})(1-q^{2k})$$

$$\theta_{4}(x,q) = \prod_{k=1}^{\infty}(1-2q^{2k-1}\cos(2x)+q^{4k-2})(1-q^{2k}),$$
(48)

while the normalizations entering the inversion relations are,

$$\rho_1(x) = 1, \quad \rho_2(x) = 4 \left[\frac{\theta_1(\frac{x}{2}, q)}{\theta_1(x, q)} \right]^2 \frac{\theta_1(\xi - x, q)\theta_1(\xi + x, q)}{\theta_1(\xi - \frac{x}{2}, q)\theta_1(\xi + \frac{x}{2}, q)}.$$
(49)

We now use the result (18) to built the R-matrix of the equivalent vertex model associated to the Ashkin-Teller model. We find that this operator has sixty four non-null vertex weights but many of them are the same due to the underlying $Z(2) \times Z(2)$ symmetry of the spin model. It turns out that we have only sixteen distinct weights and the explicit form of the 16 × 16 R-matrix is given by,

	$\begin{pmatrix} w_1 \end{pmatrix}$	0	0	0	w_2	0	0	0	w_3	0	0	0	w_4	0	0	0	١	
$\mathbf{R}_{12}(x,y) =$	w_5	0	0	0	w_6	0	0	0	w_7	0	0	0	w_8	0	0	0		
	w_9	0	0	0	w_{10}	0	0	0	w_{11}	0	0	0	w_{12}	0	0	0		
	w_{13}	0	0	0	w_{14}	0	0	0	w_{15}	0	0	0	w_{16}	0	0	0		
	0	w_6	0	0	0	w_5	0	0	0	w_8	0	0	0	w_7	0	0		
	0	w_2	0	0	0	w_1	0	0	0	w_4	0	0	0	w_3	0	0		
	0	w_{14}	0	0	0	w_{13}	0	0	0	w_{16}	0	0	0	w_{15}	0	0		
	0	w_{10}	0	0	0	w_9	0	0	0	w_{12}	0	0	0	w_{11}	0	0		(50)
	0	0	w_{11}	0	0	0	w_{12}	0	0	0	w_9	0	0	0	w_{10}	0	- , (30	(50)
	0	0	w_{15}	0	0	0	w_{16}	0	0	0	w_{13}	0	0	0	w_{14}	0		
	0	0	w_3	0	0	0	w_4	0	0	0	w_1	0	0	0	w_2	0		
	0	0	w_7	0	0	0	w_8	0	0	0	w_5	0	0	0	w_6	0		
	0	0	0	w_{16}	0	0	0	w_{15}	0	0	0	w_{14}	0	0	0	w_{13}		
	0	0	0	w_{12}	0	0	0	w_{11}	0	0	0	w_{10}	0	0	0	w_9		
	0	0	0	w_8	0	0	0	w_7	0	0	0	w_6	0	0	0	w_5		
	0	0	0	w_4	0	0	0	w_3	0	0	0	w_2	0	0	0	w_1	/	

where the vertex weights w_i are obtained in terms of the spin edge weights as follows,

$$w_{1} = 1, \quad w_{2} = \frac{b_{v}(x-y)}{b_{h}(y)}, \quad w_{3} = \frac{c_{v}(x-y)}{c_{h}(y)}, \quad w_{4} = \frac{d_{v}(x-y)}{d_{h}(y)}, \quad w_{5} = b_{h}(x)b_{v}(x-y),$$

$$w_{6} = \frac{b_{h}(x)}{b_{h}(y)}, \quad w_{7} = \frac{b_{h}(x)d_{v}(x-y)}{c_{h}(y)}, \quad w_{8} = \frac{b_{h}(x)c_{v}(x-y)}{d_{h}(y)}, \quad w_{9} = c_{h}(x)c_{v}(x-y),$$

$$w_{10} = \frac{c_{h}(x)d_{v}(x-y)}{b_{h}(y)}, \quad w_{11} = \frac{c_{h}(x)}{c_{h}(y)}, \quad w_{12} = \frac{b_{v}(x-y)c_{h}(x)}{d_{h}(y)}, \quad w_{13} = d_{h}(x)d_{v}(x-y),$$

$$w_{14} = \frac{c_{v}(x-y)d_{h}(x)}{b_{h}(y)}, \quad w_{15} = \frac{b_{v}(x-y)d_{h}(x)}{c_{h}(y)}, \quad w_{16} = \frac{d_{h}(x)}{d_{h}(y)}.$$
(51)

We have verified that the R-matrix (50,51) indeed satisfies the Yang-Baxter equation by using the explicit expressions of the edge weights (47). This can be done by using certain identities among theta functions and with the help of symbolic algebra packages.

We would like to conclude this section by discussing the Hamiltonian limit associated to the above R-matrix when the Ashkin-Teller spin model is layer isotropic. In this case the two independent Ising interactions are the same, i.e $J_h = K_h$ and $J_v = K_v$, and the Ashkin-Teller model becomes equivalent to a staggered six-vertex model with $w_d = 0$ [1, 34]. This corresponds to the limit $q \to 0$ in Eq.(47) and the respective weights are given in terms of trigonometric functions,

$$a_{h}(x) = 1, \quad b_{h}(x) = c_{h}(x) = \frac{\sin\left(\frac{\xi - x}{2}\right)}{\sin\left(\frac{\xi + x}{2}\right)}, \quad d_{h}(x) = \frac{\tan\left(\frac{\xi - x}{2}\right)}{\tan\left(\frac{\xi + x}{2}\right)},$$
$$a_{v}(x) = 1, \quad b_{v}(x) = c_{v}(x) = \frac{\sin\left(\frac{x}{2}\right)}{\sin\left(\xi - \frac{x}{2}\right)}, \quad d_{v}(x) = \frac{\tan\left(\frac{x}{2}\right)}{\tan\left(\xi - \frac{x}{2}\right)},$$
(52)

and from Eq.(51) we have now ten distinct weights w_i due to the identity $b_{h,v}(x) = c_{h,v}(x)$. The respective auxiliary functions associated to the inversion relations are,

$$\rho_1(x) = 1, \quad \rho_2(x) = \frac{\sin(\xi + x)\sin(\xi - x)}{\sin(\xi + \frac{x}{2})\sin(\xi - \frac{x}{2})\left[\cos\left(\frac{x}{2}\right)\right]^2}.$$
(53)

Inspired by the analysis in section 3 we write the Hamiltonian associated to the vertex model defined by the weights (51,52) as,

$$H(x_0) = -J \Big[\sum_{j=1}^{L-1} \Big(H_{j,j+1}^{(0)}(x_0) + H_{j,j+1}^{(1)}(x_0) \Big) + H_{L,1}^{(0)}(x_0) + H_{L,1}^{(1)}(x_0) \Big],$$
(54)

where the dynamics of the Hamiltonian will be described by the following two commuting sets of $\text{spin}-\frac{1}{2}$ Pauli matrices,

$$\sigma^{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \otimes \mathbf{I}_{2}, \quad \sigma^{z} = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \otimes \mathbf{I}_{2}, \quad \tau^{x} = \mathbf{I}_{2} \otimes \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \tau^{z} = \mathbf{I}_{2} \otimes \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
(55)

We find that the form of the two-body $H_{j,j+1}^{(0)}(x_0)$ term is similar to that obtained considering a particular time-continuous limit of the classical Ashkin-Teller model which preserves the self-dual property of such spin model [34], namely

$$H_{j,j+1}^{(0)}(x_0) = \sigma_j^z \sigma_{j+1}^z + \sigma_j^x + \tau_j^z \tau_{j+1}^z + \tau_j^x + \frac{\cos(\xi)}{\cos(x_0)} \left(\sigma_j^z \sigma_{j+1}^z \tau_j^z \tau_{j+1}^z, +\sigma_j^x \tau_j^x\right)$$
(56)

while the expression for the additional two-body $H_{j,j+1}^{(1)}(x_0)$ term is given by,

$$H_{j,j+1}^{(1)}(x_0) = -\frac{\cos(\xi)\sin(x_0)}{\cos(x_0)\sin(\xi)} \Big(\left(\sigma_j^x + \tau_j^x\right)\sigma_j^z\sigma_{j+1}^z\tau_j^z\tau_{j+1}^z - \left(\sigma_j^z\sigma_{j+1}^z + \tau_j^z\tau_{j+1}^z\right)\sigma_j^x\tau_j^x \Big) \\ + \frac{\cos(\xi)}{\cos(x_0)} \Big[\frac{\sin(x_0)}{\sin(\xi)}\Big]^2 \Big(\sigma_j^x\tau_j^z\tau_{j+1}^z + \tau_j^x\sigma_j^z\sigma_{j+1}^z + \sigma_j^x\tau_j^x\sigma_j^z\sigma_{j+1}^z\tau_{j+1}^z \Big) \\ - \frac{\sin(x_0)}{\sin(\xi)} \Big(\sigma_j^x\sigma_j^z\sigma_{j+1}^z + \tau_j^x\tau_j^z\tau_{j+1}^z \Big).$$
(57)

We finally note that the Hamiltonian defined by Eqs. (54, 56, 57) is a Hermitian operator when the parameter x_0 is imaginary.

6 Conclusions

The main purpose of this paper was to explore a recent correspondence among arbitrary n-state spin and vertex models on the square lattice [22] in the realm of exactly solvable two-dimensional systems. We have been able to formulate a given integrable classical spin model with edge weights depending on the difference of spectral parameters in the framework of an equivalent solvable vertex model on the square lattice. An immediate consequence of such embedding is that the exact solution of integrable spin models can in principle be considered within the quantum inverse scattering method [24].

In this sense, we have exhibited the expressions of Lax operator of the vertex model and the respective R-matrix in terms of the spin edge weights which are shown to satisfy the Yang-Baxter algebra. The unitarity of the R-matrix and therefore its invertibility is assured by assuming that the spin model edge weights satisfy certain inversion relation. It turns out that the R-matrix is not of difference form and this feature can be explored to construct deformed quantum spin chains with additional interactions generalizing those associated to the classical integrable spin models. We have applied this construction to the *n*-state scalar Potts [11] and the Ashkin-Teller [14] models and the expressions of the respective R-matrices are presented. In the case of the scalar Potts model we have argued that the R-matrix can be written in terms of the underlying Temperley-Lieb operators [26].

In principle, we can attempt to formulate the R-matrix associated to the equivalent vertex model without the need of any specific assumption about the parametrization of the edge weights of the spin model. To this end let us consider two spin models one of them with edge weights $W_{h,v}(i, j)$ and the other one with distinct prime edge weight $W'_{h,v}(i, j)$. The sufficient conditions for the commutation of the diagonal-to-diagonal transfer matrices of these spin models consist on the existence of double primed weights $W''_{h,v}(i, j)$ which are required to satisfy the following sets of star-triangle relations,

$$\sum_{d=1}^{n} W'_{v}(d,c)W''_{v}(b,d)W'_{h}(a,d) = \mathcal{R}W'_{h}(a,b)W''_{h}(a,c)W_{v}(b,c),$$

$$\sum_{d=1}^{n} W'_{v}(c,d)W''_{v}(d,b)W'_{h}(d,a) = \mathcal{R}W'_{h}(b,a)W''_{h}(c,a)W_{v}(c,b),$$
(58)

where again the factor \mathcal{R} is assumed to be independent of the spin variables $a, b, c = 1, \ldots n$.

The requirement (58) leads to $2n^3$ relations which in principle can be solved by eliminating auxiliary edge weights $W''_{h,v}(i,j)$ and the scalar factor \mathcal{R} and as result we end up with a number non-linear equations involving the edge weights $W_{h,v}(i,j)$ and $W'_{h,v}(i,j)$. We then assume that these non-linear equations can be solved in such way that both edge weights $W_{h,v}(i,j)$ and $W'_{h,v}(i,j)$ lie on the same algebraic variety. After performing the above tasks the auxiliary weights $W''_{h,v}(i,j)$ and the factor \mathcal{R} will be determined by polynomials whose variables are the edge weights $W_{h,v}(i,j)$ and $W'_{h,v}(i,j)$. Therefore, at this point we can shorten the notation and represent the set of edge weights $W_{h,v}(i,j)$ and $W'_{h,v}(i,j)$ by the symbols **w** and **w**', respectively. Now we can write the Yang-Baxter algebra associated to the equivalent vertex model as follows,

$$R_{12}(\mathbf{w}, \mathbf{w}') \mathbb{L}_{13}(\mathbf{w}) \mathbb{L}_{23}(\mathbf{w}') = \mathbb{L}_{23}(\mathbf{w}') \mathbb{L}_{13}(\mathbf{w}) R_{12}(\mathbf{w}, \mathbf{w}'),$$
(59)

where the respective Lax operator is given by Eq.(10), namely

$$\mathbb{L}_{12}(\mathbf{w}) = \sum_{i,j,k=1}^{n} W_h(j,i) W_v(j,k) e_{ik} \otimes e_{ji}.$$
(60)

Considering the reasoning of section 3 but now with the help of the star-triangle relations (58), it is possible to show that the following R-matrix,

$$\mathbf{R}_{12}(\mathbf{w}, \mathbf{w}') = \sum_{i,j,k=1}^{n} \frac{W_h(j,i) W_v''(j,k|\mathbf{w}, \mathbf{w}')}{W_h'(k,i)} e_{ik} \otimes e_{ji},$$
(61)

satisfy the Yang-Baxter algebra (59). Note that here we have emphasized the dependence of the auxiliary variables $W_{h,v}^{''}(i,j)$ on the spin edge weights $W_{h,v}(i,j)$ and $W_{h,v}^{'}(i,j)$.

The next step for the integrability is to assure that R-matrix has an inverse for most values of the spin edge weights \mathbf{w} and \mathbf{w}' . One way to guarantee this property is by imposing that the R-matrix satisfy the unitarity property,

$$R_{12}(\mathbf{w}, \mathbf{w}')R_{21}(\mathbf{w}', \mathbf{w}) = \rho_2(\mathbf{w}, \mathbf{w}')I_n \otimes I_n,$$
(62)

when we interchange the spin model edge weights, that is $\mathbf{w} \leftrightarrow \mathbf{w}'$. Considering the R-matrix expression (61) we find that the unitarity property is satisfied provided that the vertical auxiliary weights $W_v''(i, j)$ satisfy the following relation,

$$\sum_{k=1}^{n} W_{v}^{''}(i,k|\mathbf{w},\mathbf{w}')W_{v}^{''}(k,j|\mathbf{w}',\mathbf{w}) = \rho_{2}(\mathbf{w},\mathbf{w}')\delta_{i,j}$$
(63)

which is the analog of the second inversion relation given in Eq.(16).

We hope that the above abstract construction could be used to include spin models whose edge weights can not be presented in terms of the difference of two spectral parameters being the most known example the chiral Potts model [25]. However, it could be that such construction still need further adaptations to include the specific representation of the edge weights of the chiral Potts model in terms of two distinct points on an algebraic curve, in special attention to the constraint (63).

Acknowledgments

This work was supported in part by the Brazilian Research Council CNPq 305617/2021-4.

Appendix A: The Yang-Baxter equation

We start by rewriting the Yang-Baxter equation (24) in terms of its components,

$$\sum_{\gamma,\gamma',\gamma''=1}^{n} \mathcal{R}_{a_{1},a_{2}}^{\gamma,\gamma'}(x,y) \mathcal{R}_{\gamma,a_{3}}^{b_{1},\gamma''}(x,z) \mathcal{R}_{\gamma',\gamma''}^{b_{2},b_{3}}(y,z) = \sum_{\gamma,\gamma',\gamma''=1}^{n} \mathcal{R}_{a_{2},a_{3}}^{\gamma',\gamma''}(y,z) \mathcal{R}_{a_{1},\gamma''}^{\gamma,b_{3}}(x,z) \mathcal{R}_{\gamma,\gamma'}^{b_{1},b_{2}}(x,y).$$
(A.1)

By substituting the expression of the R-matrix (18) we observe that Yang-Baxter equation (A.1) is trivially satisfied for $b_3 \neq a_1$ since both sides of the equation are in fact zero. The situation is similar to that we have found for the Yang-Baxter algebra, see Eqs.(20,21). It turns out that for $b_3 = a_1$ the Yang-Baxter equation (A.1) becomes,

$$W_{v}(a_{3},b_{1}|x-z)\sum_{\gamma=1}^{n}\frac{W_{v}(a_{2},\gamma|x-y)W_{h}(a_{3},\gamma|x)W_{v}(\gamma,b_{2}|y-z)}{W_{h}(b_{1},\gamma|z)} = \frac{W_{h}(a_{3},a_{2}|y)}{W_{h}(b_{1},b_{2}|y)}W_{v}(a_{2},b_{2}|x-z)\sum_{\gamma'=1}^{n}\frac{W_{v}(a_{3},\gamma'|y-z)W_{h}(\gamma',b_{2}|x)W_{v}(\gamma',b_{1}|x-y)}{W_{h}(\gamma',a_{2}|z)}, \quad (A.2)$$

reducing the summations over three different labels into a sum over a single index.

We have checked that the relations (A.2) are satisfied by the edge weights of the scalar Potts, the self-dual Ashkin-Teller, the Fateev-Zamolodchikov, and the Kashiwara-Miwa spin models. This has been done by substituting the explicit expressions of the weights of the mentioned spin models in Eq.(A.2). The simplifications can be carried out by considering the addition properties between either trigonometric or elliptic theta functions and with help of symbolic algebra packages. In addition to that, we have used that the edge weights satisfy the property $W_{h,v}(i, j|x) = W_{h,v}(j, i|x)$ and in this case we note that Yang-Baxter relations (A.2) are trivially satisfied for the subset $b_1 = a_2$ and $b_2 = a_3$.

We now recall that the weights of the first two mentioned spin models have been already discussed in sections 4 and 5. Therefore, for sake of completeness, we provide below the edge weights of the Fateev-Zamolodchikov and Kashiwara-Miwa models in order to easy independent verifications for such spin models.

• The Fateev-Zamolodchikov model

The Fateev-Zamolodchikov model is *n*-state spin model with underlying Z(n) symmetry. The horizontal and vertical weights of this model are,

$$W_h(a,b|x) = \prod_{j=1}^{|a-b|} \frac{\sin\left((2j-1)\lambda - x\right)}{\sin\left((2j-1)\lambda + x\right)}, \quad W_v(a,b|x) = \prod_{j=1}^{|a-b|} \frac{\sin\left((2j-2)\lambda + x\right)}{\sin\left(2j\lambda - x\right)}, \tag{A.3}$$

where $\lambda = \frac{\pi}{2n}$.

For n = 2, 3 this spin model corresponds to the two and three states scalar Potts model. The normalizations of the inversion relations are given by,

$$\rho_1(x) = 1, \quad \rho_2(x) = n \prod_{j=1}^{[n/2]} \frac{\sin\left((2j-1)\lambda + x\right)\sin\left((2j-1)\lambda - x\right)}{\sin\left(2j\lambda + x\right)\sin\left(2j\lambda - x\right)}$$

$$\bullet \text{ The Kashiwara-Miwa model}$$
(A.4)

The Kashiwara-Miwa model is a generalization of the Fateev-Zamolodchikov model that breaks the Z(n) invariance but retains the weights reflection symmetry $W_{h,v}(i, j) = W_{h,v}(j, i)$ as well as the rapidity difference property. This model has also been investigated by Hasegawa and Yamada [20] and by Gaudin [21]. The corresponding edge weights are formulated in terms of the theta functions $\theta_1(x,q)$ and $\theta_4(x,q)$,

$$W_{h}(a,b|x) = [f(a)f(b)]^{-nx/\pi} \prod_{j=1}^{|a-b|} \frac{\theta_{1}((2j-1)\lambda - x,q)}{\theta_{1}((2j-1)\lambda + x,q)} \prod_{j=1}^{a+b} \frac{\theta_{4}((2j-1)\lambda - x,q)}{\theta_{4}((2j-1)\lambda + x,q)},$$

$$W_{v}(a,b|x) = [f(a)f(b)]^{n(x-\lambda)/\pi} \prod_{j=1}^{|a-b|} \frac{\theta_{1}((2j-2)\lambda + x,q)}{\theta_{1}(2j\lambda - x,q)} \prod_{j=1}^{a+b} \frac{\theta_{4}((2j-2)\lambda + x,q)}{\theta_{4}(2j\lambda - x,q)}, \quad (A.5)$$

where $f(a) = \frac{\theta_4(0,q)}{\theta_4(2\pi a/n,q)}$.

For n = 2 this spin model corresponds to Ising model solved originally by Onsager. The normalizations of the inversion relations are given by,

$$\rho_1(x) = 1, \quad \rho_2(x) = \frac{h(x)h(-x)}{[h(0)]^2},$$
(A.6)

where the function h(x) in terms of the theta functions is,

$$h(x) = \prod_{j=1}^{\lfloor n/2 \rfloor} \frac{\theta_1\left((2j-1)\lambda + x, q\right)\theta_4\left((2j-1)\lambda + x, q\right)}{\theta_1\left(2j\lambda + x, q\right)\theta_4\left(2j\lambda + x, q\right)}.$$
 (A.7)

References

- [1] R.J. Baxter, Exactly Solved Models in Statistical Mechanics, Academic Press, 1982
- [2] L. Onsager, *Phys.Rev.*, 65 (1944) 117
- [3] B.M. McCoy and T.T. Wu, The Two Dimensional Ising Model, Havard University Press, 1973
- [4] H.A. Kramers and G.H. Wannier, *Phys. Rev.* 60 (1941) 252
- [5] L. Onsager, Critical Phenomena in Alloys, Magnets and Supercondutors, eds. Mills, Ascher and Jaffee, McGraw-Hill, New York, 1971
- [6] M.J. Stephen and L. Mittag, J.Math. Phys. 13 (1972) 1944
- [7] G.H. Wannier, *Rev. Mod. Phys* 17 (1945) 50
- [8] H. Au-Yang and J.H.H Perk Advanced Studies in Pure Mathematics 19, (1989) 57
- [9] R.J. Baxter, Int.J.Mod.Phys.B 11 (1997) 27
- [10] J.H.H Perk and H. Au-Yang, Yang-Baxter Equations, Encyclopedia Vol.5, Elsevier, Amsterdam, 2006
- [11] R.B. Potts, Proc. Camb. Phil. Soc. 48 (1952) 106
- [12] R.J. Baxter, J.Phys.C 6 (1973) L445
- [13] V.S. Pokrovsky and Yu.A. Bashilov, Commun. Math. Phys. 84 (1982) 103
- [14] J. Ashkin and E. Teller, *Phys.Rev.* 64 (1943) 178
- [15] F.J. Wegner, J.Phys.C: Solid State Phys. 5 (1972) L131
- [16] P. Fendley and P. Ginsparg, Nucl. Phys. B 324 (1989) 549
- [17] P. Pearce and K.A. Seaton, J. Phys. A: Math. Gen. 23 (1990) 1191
- [18] V.A. Fateev and A.B. Zamolodchikov, Phys. Lett. A 92 (1982) 37
- [19] M. Kashiwara and T. Miwa, Nucl. Phys. B 275 (1986) 121

- [20] K. Hasegawa and Y. Yamada, *Phys.Lett.A* 146 (1990) 387
- [21] M. Gaudin, Journal de Physique I 1 (1991) 351
- [22] M.J. Martins, Nucl. Phys. B 1005 (2024) 116610
- [23] R.J. Baxter, Ann. Phys. 70 (1972) 193; Ann. Phys. 70 (1972) 323
- [24] L.A. Takhtadzhan and L.D. Faddev, Russ. Math. Surv. 34 (1979) 11
- [25] R.J. Baxter, J.H.H. Perk and H. Au-Yang Phys.Lett.A 128 (1988) 138
- [26] H.N.V. Temperley and E.H. Lieb, Proc. Roy. Soc. A 322 (1971) 251
- [27] V.F.R Jones, Int. J. Mod. Phys. B 4 (1990) 701
- [28] E. Berkan, Nucl. Phys. B 215 (1983) 68
- [29] L. Mittag and M.J. Stephen, J.Math.Phys. 12 (1971) 441
- [30] J. Solyom and P. Peuty, *Phys. Rev. B* 24 (1981) 218
- [31] E. Vernier, E. O'Brien and P. Fendley, J.Stat.Mech. (2019) 043107
- [32] C. Fan, *Phys.Lett.A* 39 (1972) 136
- [33] I.S. Gradshteyn and I.M. Ryzhik, Table of Integrals, Series and Products, Academic Press, New York, 1980
- [34] M. Kohmoto, M. Den Nijs and L.P. Kadanoff, Phys. Rev. B 24 (1981) 5229