POINT-WISE DOUBLING INDICES OF MEASURES AND ITS APPLICATION TO BI-LIPSCHITZ CLASSIFICATION OF BEDFORD-MCMULLEN CARPETS

HUI RAO, YAN-LI XU*, AND YUAN ZHANG

ABSTRACT. Doubling measure was introduced by Beurling and Ahlfors in 1956 and now it becomes a basic concept in analysis on metric space. In this paper, for a measure which is not doubling, we introduce a notion of point-wise doubling index, and calculate the point-wise doubling indices of uniform Bernoulli measures on Bedford-McMullen carpets. As an application, we show that, except a small class of Bedford-McMullen carpets, if two Bedford-McMullen carpets are bi-Lipschitz equivalent, then they have the same fiber sequence up to a permutation.

1. Introduction

A Borel probability measure μ on a metric space is said to be *doubling* if

$$\sup_{x \in \operatorname{supp}(\mu), \ r > 0} \frac{\mu(B_{2r}(x))}{\mu(B_r(x))} < \infty.$$

where $B_r(x)$ denotes the open ball with center x and radius r, and $\operatorname{supp}(\mu)$, the support of μ , consists of x such that $\mu(B_r(x)) > 0$ for every r > 0.

Doubling measure is introduced by Beurling and Ahlfors [4] in 1956. Since then, it is an important object in the analysis of metric spaces, see for instance [4,9,10,19] *etc.* Usually, a measure is considered to be 'nice' if it is doubling. But if a measure is not doubling, what can we say? The purpose of this paper is to study changing rate of a non-doubling measure point-wisely. Precisely, we introduce the following notion.

Definition 1.1. (Point-wise doubling index) Let μ be a measure on a metric space (X, d). Let $0 < \rho < 1$. Let $\varphi : (0, 1) \to \mathbb{R}^+$ be a strictly decreasing function satisfying $\lim_{r\to 0} \varphi(r) = \infty$, which we call a gauge function. For $z \in X$, the upper doubling index of μ at z (w.r.t. φ) is defined by

(1.1)
$$\overline{\delta}_{\varphi}(z;\mu,\rho) = \limsup_{r \to 0} \sup_{B_{\rho r}(z') \subset B_r(z)} \frac{\log[\mu(B_r(z))/\mu(B_{\rho r}(z'))]}{\varphi(r)};$$

Date: January 7, 2025.

The work is supported by NSFC No. 11971195.

2020 Mathematics Subject Classification: 28A80, 26A16.

Key words and phrases: Bedford-McMullen carpet, point-wise doubling index, run length. * Corresponding author.

the lower doubling index $\underline{\delta}_{\varphi}(z;\mu,\rho)$ is obtained by replacing lim sup by lim inf in (1.1). If the two values agree, the common value is called the *doubling index* and denoted by $\delta_{\varphi}(z; \mu, \rho)$.

Remark 1.1. It is seen that if μ is a doubling measure, then $\delta_{\varphi}(z; \mu, \rho) \equiv 0$ for all $z \in X$.

In this paper, we calculate doubling indices of uniform Bernoulli measures on Bedford-McMullen carpets, and we show that they are Lipschitz invariants. Thanks to these new invariants, we obtain some remarkable new results concerning the bi-Lipschitz classification of Bedford-McMullen carpets (see Theorem 1.3 and Theorem 1.4). Furthermore, we show that two Lipschitz equivalent Bedford-McMullen carpets have the same fiber sequence up to a permutation except a small class (see Theorem 1.5).

Let $n > m \ge 2$ be two integers and denote by diag(n,m) the 2 \times 2 diagonal matrix. Let $\mathcal{D} \subset \{0, 1, \dots, n-1\} \times \{0, 1, \dots, m-1\}$. For $\mathbf{d} \in \mathcal{D}$, set

(1.2)
$$S_{\mathbf{d}}(z) = \operatorname{diag}(n^{-1}, m^{-1})(z + \mathbf{d}),$$

then $\{S_{\mathbf{d}}\}_{\mathbf{d}\in\mathcal{D}}$ is an iterated function system (IFS), its invariant set $E = K(n, m, \mathcal{D})$, the unique non-empty compact set satisfying $E = \bigcup_{\mathbf{d} \in \mathcal{D}} S_{\mathbf{d}}(E)$, is called a *Bedford-McMullen carpet (abbreviated as BM-carpet)* [3, 16]. We denote by μ_E the unique Borel probability measure supported on E satisfying

$$\mu_E(\cdot) = \frac{1}{\#\mathcal{D}} \sum_{\mathbf{d}\in\mathcal{D}} \mu_E \circ S_{\mathbf{d}}^{-1}(\cdot),$$

and call μ_E the uniform Bernoulli measure of E, where #A denotes the cardinality of a set A.

Recall that two metric spaces (X, d_X) and (Y, d_Y) are said to be Lipschitz equivalent, denoted by $(X, d_X) \sim (Y, d_Y)$, if there exists a map $f : X \to Y$ which is bi-Lipschitz, that is, there is a constant C > 0 such that

$$C^{-1}d_X(x,y) \le d_Y(f(x), f(y)) \le Cd_X(x,y), \text{ for all } x, y \in X.$$

Recently, there are many works devoted to the Lipschitz classification of BMcarpets, see [2, 12, 14, 17, 18, 20, 21]. It turns out that the classification is much more complicated than the self-similar settings. Notably, there are a lot of Lipschitz invariants, which divide the BM-carpets into many sub-families invariant under bi-Lipschitz maps. The most well-known Lipschitz invariants are various dimensions, such as Hausdorff dimension, box dimension, Assouad dimension, etc. Fraser and Yu [8] showed that $\log m / \log n$ is a Lipschitz invariant, Rao, Yang and Zhang [18], and Banaji and Kolossváry [2] showed that the multifractal spectrum of the uniform Bernoulli measure on BM-carpet is a Lipschitz invariant, see also Huang, Rao, Wen and Xu [11].

Let $E = K(n, m, \mathcal{D})$ be a BM-carpet. We define

$$a_j = \#\{i: (i,j) \in \mathcal{D}\}, \quad j = 0, 1, \dots, m-1,$$

and call $(a_j)_{j=0}^{m-1}$ the fiber sequence of \mathcal{D} . Throughout the paper, we denote

 $\sigma = \log m / \log n.$

We say a BM-carpet is of *non-doubling type* if the associated uniform Bernoulli measure is not doubling. Li, Wei and Wen [15] characterized when a Bernoulli measure on a BM-carpet is doubling. According to their result, we have

Lemma 1.1. The uniform Bernoulli measure μ_E is not doubling if and only if $a_0a_{m-1} > 0$, $a_0 \neq a_{m-1}$, and $a_ja_{j+1} > 0$ for at least one $j \in \{0, 1, \ldots, m-2\}$.

For $\mathbf{i} = \mathbf{d}_1 \dots \mathbf{d}_k \in \mathcal{D}^k$, we define $S_{\mathbf{i}} = S_{\mathbf{d}_1} \circ \dots \circ S_{\mathbf{d}_k}$. Let $z \in E$, we call $(\mathbf{d}_j)_{j \ge 1}$ a *coding* of z if $\{z\} = \bigcap_{k \ge 1} S_{\mathbf{d}_1 \dots \mathbf{d}_k}([0, 1]^2)$. For convenience, we denote $\mathbf{d}_j = (x_j, y_j)$ and we also write a coding as $(\mathbf{x}, \mathbf{y}) = ((x_j)_{j \ge 1}, (y_j)_{j \ge 1})$.

Definition 1.2 ([14] [21]). We call z a *double vertical coding point* if z has two codings (\mathbf{x}, \mathbf{y}) and $(\mathbf{x}', \mathbf{y}')$ such that $\mathbf{y} \neq \mathbf{y}'$. Denote

(1.3) $V_E = \{z \in E; z \text{ has double vertical codings}\}.$

A BM-carpet E is said to satisfy the vertical separation condition (VSC) if $V_E = \emptyset$.

Li, Li and Miao [14] proved that if E = K(n, m, D) and F = K(n, m, D') are totally disconnected, satisfy the VSC and share the same fiber sequence, then $E \sim$ F. Yang and Zhang [21] strengthen this result by showing that if a BM-carpet is totally disconnected and satisfies the VSC, then it is Lipschitz equivalent to a certain symbolic space.

Now we state our main results. For the lower doubling index, we have

Theorem 1.1. Let E = K(n, m, D) be a BM-carpet of non-doubling type and assume that $a_0 > a_{m-1}$. Let $0 < \rho < n^{-3}$ and let φ be a gauge function satisfying

(1.4)
$$\lim_{k \to \infty} \frac{k}{\varphi(n^{-k})} = s \in [0, +\infty]$$

Then

$$\underline{\delta}_{\varphi}(z;\mu_E,\rho) = \begin{cases} s(1/\sigma - 1)\log(a_0/a_{m-1}), & \text{if } z \in V_E, \\ 0, & \text{otherwise.} \end{cases}$$

Remark 1.2. If the gauge function φ is too big such that the limit in (1.4) is zero, then $\underline{\delta}_{\varphi}(z; \mu_E, \rho)$ is constantly 0, and it is not interesting.

For
$$j \in \{0, 1, ..., m-1\}$$
, set $\Omega_j = \{(\mathbf{x}, \mathbf{y}) \in \mathcal{D}^{\infty}; \mathbf{y} \text{ ends with } j^{\infty}\}$. Clearly $V_E \subset \Omega_0 \cup \Omega_{m-1}$.

To compute the upper doubling index, we introduce a function $\beta(k; \boldsymbol{\omega})$, a kind of reverse run length function for a $\boldsymbol{\omega} \in \mathcal{D}^{\infty}$ at position k, in (3.7) of Section 3.

Theorem 1.2. Let E = K(n, m, D) be a BM-carpet of non-doubling type and assume that $a_0 > a_{m-1}$. Let φ be a gauge function satisfying

(1.5)
$$\lim_{k \to \infty} \frac{k}{\varphi(n^{-k})} = s \in (0, +\infty].$$

Let $0 < \rho < n^{-3}$. Let $z \in E$ and let $\boldsymbol{\omega}$ be a coding of z. Then (i) If $\boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}$, then

$$\overline{\delta}_{\varphi}(z;\mu_E,\rho) = \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \cdot \log(a_0/a_{m-1})$$

(ii) If $\boldsymbol{\omega} \in \Omega_0 \cup \Omega_{m-1}$, then $\overline{\delta}_{\varphi}(z; \mu_E, \rho) = s(1/\sigma - 1) \log(a_0/a_{m-1}) \cdot \mathbf{1}_{V_E}(z)$, where $\mathbf{1}_{V_E}$ is the indicator function of V_E .

Remark 1.3. From the above two theorems, we see that under the assumption (1.5), $\overline{\delta}_{\varphi}$ and $\underline{\delta}_{\varphi}$ are irrelevant with ρ . So from now on, we will use $\overline{\delta}_{\varphi}(z;\mu_E)$ and $\underline{\delta}_{\varphi}(z;\mu_E)$ instead of $\overline{\delta}_{\varphi}(z;\mu_E,\rho)$ and $\underline{\delta}_{\varphi}(z;\mu_E,\rho)$ respectively.

Example 1.1. Let $E = K(n, m, \mathcal{D})$ be a BM-carpet of non-doubling type and assume that $a_0 > a_{m-1}$. We are especially interested in the gauge functions $\varphi(r) =$ $-\log r$ and $\varphi(r) = \log |\log r|$.

(1) Let $\varphi(r) = -\log r$, and denote $\delta = \delta_{\varphi}$. Then

(1.6)
$$\overline{\delta}(z;\mu_E) = \begin{cases} \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{k} \cdot \log_n(a_0/a_{m-1}), & \text{if } \boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}, \\ \delta_{\max} \cdot \mathbf{1}_{V_E}(z), & \text{otherwise }, \end{cases}$$

where $\delta_{\max} = (\log n)^{-1} (1/\sigma - 1) \log(a_0/a_{m-1})$, and $\underline{\delta}(z; \mu_E) = \delta_{\max} \cdot \mathbf{1}_{V_E}(z)$. Moreover, we can show that for μ_E -a.e. $z \in E$, $\delta(z; \mu_E) = 0$, and the range of

 $\delta(z; \mu_E)$ is $[0, \delta_{\max}]$. (See Theorem 5.1.)

(2) Let $\varphi(r) = \log |\log r|$, and denote $\Delta = \delta_{\varphi}$. Then

(1.7)
$$\overline{\Delta}(z;\mu_E) = \begin{cases} \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\log k} \cdot \log(a_0/a_{m-1}), & \text{if } \boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}, \\ \infty \cdot \mathbf{1}_{V_E}(z), & \text{otherwise }, \end{cases}$$

and $\underline{\Delta}(z;\mu_E) = \infty \cdot \mathbf{1}_{V_E}(z).$

Moreover, we have that for μ_E -a.e. $z \in E$,

(1.8)
$$\overline{\Delta}(z;\mu_E) = -\log_{p_0}(a_0/a_{m-1}) := \Delta_{aver}$$

where $p_0 = a_0 / \# \mathcal{D}$, which tells us that $\varphi(r) = \log |\log r|$ is the gauge function for almost every $z \in E$. (See Theorem 5.2.)

In the following, we use the point-wise doubling indices to construct new Lipschitz invariants for classification of BM-carpets. We assume that $E = K(n, m, \mathcal{D}), F =$ $K(n, m, \mathcal{D}')$ are two BM-carpets. We use $(a_j)_{j=0}^{m-1}$ and $(a'_j)_{j=0}^{m-1}$ to denote the fiber sequences of \mathcal{D} and \mathcal{D}' , respectively. Denote by μ_E and μ_F the uniform Bernoulli measures of E and F respectively.

The following result shows that the VSC is a Lipschitz invariant.

Theorem 1.3. Let $E = K(n, m, \mathcal{D}), F = K(n, m, \mathcal{D}')$ be two BM-carpets of nondoubling type. Let $f: E \to F$ be a bi-Lipschitz map. Then for any $z \in E$,

(1.9)
$$\overline{\delta}_{\varphi}(z;\mu_E) = \overline{\delta}_{\varphi}(f(z);\mu_F), \quad \underline{\delta}_{\varphi}(z;\mu_E) = \underline{\delta}_{\varphi}(f(z);\mu_F).$$

Consequently,

 $V_F = f(V_E)$ and $\dim_H V_E = \dim_H V_F$.

Especially, E satisfies the VSC (that is, $V_E = \emptyset$) if and only if F does.

Using the Lipschitz invariants δ_{\max} , Δ_{aver} and γ_{\max} (which is defined in Section 6), we show that

Theorem 1.4. Let E = K(n, m, D), F = K(n, m, D') be two BM-carpets of nondoubling type. Suppose $E \sim F$. Then

(i) the fiber sequence of E is a permutation of that of F;

(ii) we have $a_0 = a'_0$ and $a_{m-1} = a'_{m-1}$ if we assume without loss of generality that $a_0 > a_{m-1}$ and $a'_0 > a'_{m-1}$.

We use $\mathcal{M}_{t,v,d,r}(n,m)$ to denote the class of BM-carpets which are totally disconnected (which is indicated by t) with expanding matrix diag(n,m), possess vacant rows (indicated by v), are of doubling type (indicated by d), and $\log m / \log n$ is rational (indicated by r). It is well-known that these four properties are all Lipschitz invariants.

Theorem 1.5. Let E = K(n, m, D), F = K(n, m, D') be two BM-carpets which are not in $\mathcal{M}_{t,v,d,r}(n,m)$. Then $E \sim F$ implies that the fiber sequence of E is a permutation of that of F.

Remark 1.4. Let $E = K(n, m, \mathcal{D})$ and $F = K(n, m, \mathcal{D}')$ be two BM-carpets. We remark that if they share the same fiber sequence up to a permutation, then μ_E and μ_F share the same multifractal spectrum, and this further implies that they share the same Hausdorff, box, Assouad and intermediate dimensions ([2,18]).

Remark 1.5. Set $H = \{i; (i, j), (i, j+1) \in \mathcal{D}\}$ and $I = \{i; (i, 0), (i, m-1) \in \mathcal{D}\}$. It is easy to show that if $H \neq \emptyset$ and $\#I \ge 2$, then $\dim_H V_E = \log(\#I)/\log n$; otherwise, V_E is either countable or empty.

Example 1.2. Let $E = K(8, 4, \mathcal{D})$ and $F = K(8, 4, \mathcal{D}')$ be two BM-carpets with digit sets indicated in Fig.1. Notice that the fiber sequence of \mathcal{D} is a permutation of that of \mathcal{D}' . They are indeed not Lipschitz equivalent, which can be obtained either by $\dim_H V_E = 0 \neq \dim_H V_F = \log 2/\log 8$, or by $a_0 \neq a'_0$.



FIGURE 1.

Remark 1.6. Recently, there are a lot of works devoted to the Assouad dimension of measures on metric spaces, see [6, 7, 13]. R. Anttila [1] introduced a notion of pointwise Assouad dimension of a measure, which provides another pointwise index of measures on metric spaces.

The paper is organized as follows. We recall some known results about approximate squares of BM-carpets in Section 2. In Section 3, we prove several important lemmas. Theorem 1.1 and Theorem 1.2 are proved in Section 4. We give some remarks on $\delta_{-\log r}$ and $\delta_{\log |\log r|}$ in Section 5. We discuss an alternative point-wise doubling index in Section 6. Theorems 1.3, 1.4 and 1.5 are proved in Section 7.

2. Estimates of measures of approximate squares

Let $E = K(n, m, \mathcal{D})$ be a BM-carpet. Throughout the paper, we denote $N = \#\mathcal{D}$ and use the notation $\ell(k) = \lfloor k/\sigma \rfloor$, where $\lfloor x \rfloor$ denotes the greatest integer no larger than x. Clearly,

$$m^{\ell(k)} < n^k < m^{\ell(k)+1}.$$

For $\mathbf{i} \in \mathcal{D}^k$, we call $E_{\mathbf{i}} = S_{\mathbf{i}}(E)$ a *cylinder* of E of rank k.

For two words \mathbf{i}, \mathbf{j} , we use $\mathbf{i} * \mathbf{j}$ to denote the concatenation of \mathbf{i} and \mathbf{j} , and use $\mathbf{i} \wedge \mathbf{j}$ to denote the maximal common prefix of \mathbf{i} and \mathbf{j} , and denote by $|\mathbf{i}|$ the length of \mathbf{i} . Let $\mathbf{w}|_q = w_1 * \cdots * w_q$ be the prefix of \mathbf{w} with length q. For $z, z' \in \mathbb{R}^2$, we use d(z, z') to denote the Euclidean distance between points z and z'. We define the coding map $\pi : \mathcal{D}^{\infty} \to E$ given by $\pi(\boldsymbol{\omega}) = \bigcap_{k \geq 1} S_{\boldsymbol{\omega}|_k}([0, 1]^2)$.

Let $(\mathbf{x}, \mathbf{y}) = ((x_i)_{i \ge 1}, (y_i)_{i \ge 1}) \in \mathcal{D}^{\infty}$. For $k \ge 1$, we set

0(1.)

(2.1)
$$Q_k(\mathbf{x}, \mathbf{y}) = \pi(\{(\mathbf{u}, \mathbf{v}) \in \mathcal{D}^{\infty}; |\mathbf{x} \wedge \mathbf{u}| \ge k \text{ and } |\mathbf{y} \wedge \mathbf{v}| \ge \ell(k)\}),$$

and call it the approximate square of E of rank k, or k-th approximate square of E.

Lemma 2.1 ([3,16]). If $Q_k(\mathbf{x}, \mathbf{y})$ is a k-th approximate square of E, then

$$\mu_E(Q_k(\mathbf{x}, \mathbf{y})) = \frac{\prod_{j=k+1}^{\ell(k)} a_{y_j}}{N^{\ell(k)}}, \text{ or } \mu_E(Q_k(\mathbf{x}, \mathbf{y})) = \frac{1}{N^k} \text{ if } \ell(k) = k.$$

Lemma 2.2. Let $\boldsymbol{\omega} \in \mathcal{D}^{\infty}$ and let $k \geq 1$, then

$$\frac{\mu_E(Q_{k+1}(\boldsymbol{\omega}))}{\mu_E(Q_k(\boldsymbol{\omega}))} \ge C_0^{-1}$$

where $C_0 = n N^{1+1/\sigma}$.

Proof. By Lemma 2.1, we have

$$\frac{\mu_E(Q_{k+1}(\boldsymbol{\omega}))}{\mu_E(Q_k(\boldsymbol{\omega}))} \ge \frac{1}{nN^{\ell(k+1)-\ell(k)}} \ge \frac{1}{nN^{1+1/\sigma}},$$

and the lemma is proved.

3. Lemmas

In this section, we always assume that E is a BM-carpet of non-doubling type and satisfies $a_0 > a_{m-1}$. Let μ be the uniform Bernoulli measure on E, let $0 < \rho < n^{-3}$ and let φ be a gauge function.

Let $z \in E$ and let $\boldsymbol{\omega} = (\mathbf{x}, \mathbf{y})$ be a coding of z. Let 0 < r < 1. Denote

(3.1)
$$U(z;r,\rho) = \sup_{z' \in E, B_{\rho r}(z') \subset B_r(z)} \frac{\mu(B_r(z))}{\mu(B_{\rho r}(z'))},$$

then $\overline{\delta}_{\varphi}(z;\mu,\rho) = \limsup_{r \to 0} \frac{\log U(z;r,\rho)}{\varphi(r)}$, so does $\underline{\delta}_{\varphi}(z;\mu,\rho)$.

In this section, we will always let k(r) be the integer such that

(3.2)
$$\frac{1}{n^{k(r)+2}} < r \le \frac{1}{n^{k(r)+1}}.$$

Let

(3.3)
$$\Xi_{k(r)}(z) = \{ Q_{k(r)}(\mathbf{u}, \mathbf{v}); \ Q_{k(r)}(\mathbf{u}, \mathbf{v}) \cap B_r(z) \neq \emptyset \}$$

Since $2r < n^{-k(r)}$, we conclude $\Xi_{k(r)}(z)$ contains at most four elements, and they locate in one row, or two adjacent rows of rank k(r).

Lemma 3.1. There is a constant $C_1 > 1$ such that

(3.4)
$$U(z; r, \rho) \le C_1 \frac{\max\{\mu(Q); \ Q \in \Xi_{k(r)}(z)\}}{\min\{\mu(Q); \ Q \in \Xi_{k(r)}(z)\}}$$

and

(3.5)
$$U(z; n^{3}r, \rho) \ge C_{1}^{-1} \frac{\max\{\mu(Q); \ Q \in \Xi_{k(r)}(z)\}}{\min\{\mu(Q); \ Q \in \Xi_{k(r)}(z)\}}.$$

Proof. For simplicity, we write k := k(r).

(i) Let k_0 be the integer such that $\frac{1}{n^{k_0+1}} < \rho \leq \frac{1}{n^{k_0}}$. Pick $z' \in E$ such that $B_{\rho r}(z') \subset B_r(z)$ and let ω' be a coding of z'. Since diam $(Q_{k+k_0+4}(\omega')) < \frac{\sqrt{m^2+1}}{n^{k+k_0+4}} < \rho r$, we have $Q_{k+k_0+4}(\omega') \subset B_{\rho r}(z')$, then by Lemma 2.2 we have (3.6)

$$\mu(B_{\rho r}(z')) \ge \mu(Q_{k+k_0+4}(\boldsymbol{\omega}')) \ge \frac{1}{C_0^{k_0+4}} \mu(Q_k(\boldsymbol{\omega}')) \ge \frac{1}{C_0^{k_0+4}} \min\{\mu(Q); \ Q \in \Xi_k(z)\},$$

where C_0 is the constant in Lemma 2.2 and the last inequality holds by $Q_k(\boldsymbol{\omega}') \in \Xi_k(z)$ since $B_{\rho r}(z') \subset B_r(z) \subset \bigcup_{Q \in \Xi_k(z)} Q$.

On the other hand, $\mu(B_r(z)) \leq 4 \max\{\mu(Q); Q \in \Xi_k(z)\}$, and the first assertion holds by setting $C_1 = 4C_0^{k_0+4}$.

(ii) Clearly, $B_{n^3r}(z)$ contains all elements of $\Xi_k(z)$ since $n^3r > r + \text{diam } Q$ for each $Q \in \Xi_k(z)$ (we remark that it holds no matter $\Xi_k(z)$ contains approximate squares of rank k of two adjacent rows or the same row), then

$$\mu(B_{n^{3}r}(z)) \ge \max\{\mu(Q); \ Q \in \Xi_{k}(z)\}.$$

Suppose $Q_k(\boldsymbol{\omega}')$ attains the minimal measure in $\Xi_k(z)$. Since $a_1 a_{m-1} \neq 0$, there exist $(i_1, 0), (i_2, m-1) \in \mathcal{D}$. Let z' be the point with the coding

$$(\boldsymbol{\omega}'|_{\ell(k)})\binom{i_1}{0}\binom{i_2}{m-1}^{\infty}.$$

Since $\rho n^3 r < n^{-(k+1)}$, we have that $B_{\rho n^3 r}(z') \cap E$ is covered by at most two approximate squares of rank k located in the same row as $Q_k(\omega')$, hence

$$\mu(B_{\rho n^{3}r}(z')) \leq 2\min\{\mu(Q); \ Q \in \Xi_{k}(z)\},\$$

and the second assertion holds.

Hereafter, we always assume that C_1 is the constant in Lemma 3.1.

3.1. Reverse run length function. Let $\mathcal{E} = \{j; a_j > 0\}$. Denote $a \lor b = \max\{a, b\}$. Now we define a kind of reverse run length function for a sequence over \mathcal{D} . Let $p \in \{0, m - 1\}$ and $\boldsymbol{\omega} = (\mathbf{x}, \mathbf{y}) \in \mathcal{D}^{\infty}$. For $k \ge 1$, let $h_p(k; \boldsymbol{\omega})$ be the maximal integer h such that $h \le \ell(k)$ and $y_h \ne p$. We define

$$\beta_0(k;\boldsymbol{\omega}) = \begin{cases} \ell(k) - (k \vee h_0(k;\boldsymbol{\omega})), & \text{if } y_{h_0(k;\boldsymbol{\omega})} \in \mathcal{E} + 1, \\ 0, & \text{otherwise (including } h_0(k;\boldsymbol{\omega}) = 0); \end{cases}$$
$$\beta_{m-1}(k;\boldsymbol{\omega}) = \begin{cases} \ell(k) - (k \vee h_{m-1}(k;\boldsymbol{\omega})), & \text{if } y_{h_{m-1}(k;\boldsymbol{\omega})} \in \mathcal{E} - 1, \\ 0, & \text{otherwise (including } h_{m-1}(k;\boldsymbol{\omega}) = 0). \end{cases}$$

Define

(3.7)
$$\beta(k;\boldsymbol{\omega}) = \beta_0(k;\boldsymbol{\omega}) \vee \beta_{m-1}(k;\boldsymbol{\omega}).$$

Lemma 3.2. There exists a constant $C_2 > 1$ such that for any $z = \pi(\boldsymbol{\omega}) \in E$ and $r \in (0, 1)$,

(3.8)
$$U(z;r,\rho) \le C_2 \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k(r);\boldsymbol{\omega})};$$

if $\beta(k(r); \boldsymbol{\omega}) \leq \ell(k(r)) - k(r) - 1$, then

(3.9)
$$U(z; n^6 r, \rho) \ge C_2^{-1} \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k(r);\omega)}$$

Proof. For simplicity, we write k := k(r).

We first prove (3.8). Clearly $Q_k(\boldsymbol{\omega}) = Q_k(\mathbf{x}, \mathbf{y}) \in \Xi_k(z)$. If all elements of $\Xi_k(z)$ are located in the same row as $Q_k(\mathbf{x}, \mathbf{y})$, then (3.8) holds by the first assertion of Lemma 3.1. So in the following we assume that there exists an approximate square $Q_k(\mathbf{u}, \mathbf{v}) \in \Xi_k(z)$ which locates in a row adjacent to the row of $Q_k(\mathbf{x}, \mathbf{y})$, in other words, $\mathbf{v}|_{\ell(k)}$ is adjacent to $\mathbf{y}|_{\ell(k)}$ in the lexicographical order.

Case 1. $y_{\ell(k)} \notin \{0, m-1\}.$

In this case $\beta(k; \boldsymbol{\omega}) = 0$. Since $\mathbf{v}|_{\ell(k)}$ and $\mathbf{y}|_{\ell(k)}$ are adjacent in the lexicographic order, we deduce that they differ only at the last letter. It follows that

(3.10)
$$\frac{1}{n} \le \frac{\mu(Q_k(\mathbf{x}, \mathbf{y}))}{\mu(Q_k(\mathbf{u}, \mathbf{v}))} \le n$$

By Lemma 3.1, we have $U(z; r, \rho) \leq C_1 n$. (3.8) holds in this case if we set $C_2 = C_1 n$.

Case 2. $y_{\ell(k)} = 0$ or m - 1.

Without loss of generality, we assume that $y_{\ell(k)} = 0$.

If $\mathbf{y} \prec \mathbf{v}$ in the lexicographical order, then $\mathbf{v}|_{\ell(k)} = \mathbf{y}|_{\ell(k)-1} * 1$, so (3.10) holds, which implies (3.8) holds as we did in Case 1.

If $\mathbf{v} \prec \mathbf{y}$, then there exist h and j such that

$$\mathbf{y}|_{\ell(k)} = y_1 \dots y_{h-1} * (j+1) * 0^{\ell(k)-h}, \quad \mathbf{v}|_{\ell(k)} = y_1 \dots y_{h-1} * j * (m-1)^{\ell(k)-h}.$$

Since $\beta(k; \boldsymbol{\omega}) = 0$ or $\ell(k) - k \vee h$, we deduce that

$$\frac{1}{n} \le \frac{\mu(Q_k(\mathbf{x}, \mathbf{y}))}{\mu(Q_k(\mathbf{u}, \mathbf{v}))} \le n \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k;\boldsymbol{\omega})}.$$

Therefore, by Lemma 3.1, no matter $\mu(Q_k(\mathbf{x}, \mathbf{y})) \ge \mu(Q_k(\mathbf{u}, \mathbf{v}))$ or not, we have

$$U(z;r,\rho) \le C_1 n \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k;\omega)}$$

and (3.8) also holds in this case if we set $C_2 = C_1 n$.

Next, we prove (3.9). Suppose $\beta(k; \boldsymbol{\omega}) \leq \ell(k) - k - 1$.

If $\beta(k; \boldsymbol{\omega}) \leq 4/\sigma$, then (3.9) holds if we set $C_2 \geq (a_0/a_{m-1})^{4/\sigma}$.

Hence, in the following we assume without loss of generality that $y_{\ell(k)} = 0$ and $\beta(k; \boldsymbol{\omega}) > 4/\sigma$. At this time $\beta_{m-1}(k; \boldsymbol{\omega}) = 0$ and $\beta_0(k; \boldsymbol{\omega}) = \beta(k; \boldsymbol{\omega}) \leq \ell(k) - (k+1)$, then $h := h_0(k; \boldsymbol{\omega}) \geq k + 1$. By the definition of $\beta(k; \boldsymbol{\omega})$, there exists i' such that $(i', y_h - 1) \in \mathcal{D}$. Since $a_{m-1} > 0$, there exists i'' such that $(i'', m-1) \in \mathcal{D}$. We define $\boldsymbol{\omega}' = (\mathbf{u}, \mathbf{v})$ as

$$\boldsymbol{\omega}' = \boldsymbol{\omega}|_{h-1} * \begin{pmatrix} i' \\ y_h - 1 \end{pmatrix} * \begin{pmatrix} i'' \\ m-1 \end{pmatrix}^{\infty} := \begin{pmatrix} \mathbf{x}' \\ \mathbf{y}' \end{pmatrix}.$$

Let $z'' = \pi(\boldsymbol{\omega}')$. Since z and z'' belong to a same cylinder of E of rank h-1 and $h-1 \ge k$, we obtain $d(z, z'') \le \sqrt{\frac{1}{n^{2k}} + \frac{m^2}{n^{2k}}} < \frac{1}{n^{k-1}}$, so $z'' \in B_{n^3r}(z)$.

We shall apply Lemma 3.1 to $B_{n^3r}(z)$ instead of $B_r(z)$. Notice that $k(n^3r) = k(r) - 3 = k - 3$, we define $\Xi'_{k-3}(z)$ to be the collection of approximates squares of rank (k-3) intersecting $B_{n^3r}(z)$. Clearly $Q_{k-3}(\omega), Q_{k-3}(\omega') \in \Xi'_{k-3}(z)$, moreover, $\beta(k;\omega) > 4/\sigma$ implies $h < \ell(k-3)$, then we have $y_{\ell(k-3)} = 0$ and $y'_{\ell(k-3)} = m - 1$, so $Q_{k-3}(\omega)$ and $Q_{k-3}(\omega')$ locate in different rows. By Lemma 2.1,

$$\frac{\mu(Q_{k-3}(\boldsymbol{\omega}))}{\mu(Q_{k-3}(\boldsymbol{\omega}'))} \ge \frac{1}{n} \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k-3;\boldsymbol{\omega})} \ge \frac{1}{n} \left(\frac{a_0}{a_{m-1}}\right)^{-3/\sigma-1} \cdot \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k;\boldsymbol{\omega})},$$

which together with the second assertion of Lemma 3.1 imply (3.9) if we set $C_2 = C_1 n (a_0/a_{m-1})^{3/\sigma+1}$.

The lemma is proved for all cases by setting $C_2 = C_1 n (a_0/a_{m-1})^{4/\sigma}$.

Corollary 3.1. It holds that

(3.11)
$$\overline{\delta}_{\varphi}(z;\mu,\rho) \leq \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \cdot \log(a_0/a_{m-1}).$$

Proof. Since k(r) runs over all integers as r runs over (0, 1), we have

$$\overline{\delta}_{\varphi}(z;\mu,\rho) = \limsup_{r \to 0} \frac{\log U(z;r,\rho)}{\varphi(r)} \\
\leq \limsup_{k \to \infty} \frac{\log C_2 + \beta(k;\boldsymbol{\omega}) \log\left(\frac{a_0}{a_{m-1}}\right)}{\varphi(n^{-k-1})} \\
\leq \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \cdot \log(a_0/a_{m-1}),$$

where the last inequality holds since φ is decreasing, and the corollary is proved. \Box

4. Proofs of Theorem 1.1 and Theorem 1.2

In this section, we assume the same assumptions on $E = K(n, m, \mathcal{D})$ as Section 3.

Lemma 4.1. Let φ be a gauge function satisfying

(4.1)
$$\lim_{k \to \infty} \frac{k}{\varphi(n^{-k})} = s \in (0, +\infty].$$

Let $z = \pi(\boldsymbol{\omega}) \in E$. If $\boldsymbol{\omega} \in \Omega_0 \cup \Omega_{m-1}$, then

$$\delta_{\varphi}(z;\mu,\rho) = s(1/\sigma - 1)\log(a_0/a_{m-1}) \cdot \mathbf{1}_{V_E}(z).$$

Proof. Case 1. $z \notin V_E$.

Without loss of generality, let us assume that y ends with 0^{∞} . Write y = $y_1 \cdots y_{n_0} 0^\infty$ where $y_{n_0} \neq 0$ for some $n_0 \in \mathbb{N}$.

Let L be the horizontal line containing z. Since z is not a double vertical coding point, any cylinder of E of rank n_0 below L does not contain z. Let K be the union of cylinders of E of rank n_0 below L, then dist(z, K) > 0.

Let $r < \operatorname{dist}(z, K)$. Let k = k(r) be defined in (3.2). Furthermore we choose r small so that $k > n_0$, then z locates at the bottom of $Q_k(\boldsymbol{\omega})$, so $\Xi_k(z)$ consists of at most two approximate squares of rank k in the same row, and one of them is $Q_k(\boldsymbol{\omega})$. It follows that $U(z; r, \rho) \leq C_1 n$ by Lemma 3.1. Therefore $\delta_{\varphi}(z; \mu, \rho) = 0$.

Case 2. $z \in V_E$.

Let $\omega' = (\mathbf{x}', \mathbf{y}')$ be another coding of z such that $\mathbf{y}' \neq \mathbf{y}$. Without loss of generality, let us assume that

$$\mathbf{y} = y_1 \dots y_{n_0} (j+1) 0^{\infty}, \quad \mathbf{y}' = y_1 \dots y_{n_0} j (m-1)^{\infty},$$

for some $n_0 \in \mathbb{N}$. Pick r > 0 and let k = k(r) be defined in (3.2). We choose r small so that $k > n_0$. Notice that $Q_k(\mathbf{x}, \mathbf{y}), Q_k(\mathbf{x}', \mathbf{y}') \in \Xi_k(z)$, and $\beta(k; \boldsymbol{\omega}) = \ell(k) - k$. By Lemma 3.1 we have

$$U(z; n^3 r, \rho) \ge C_1^{-1} \left(\frac{a_0}{a_{m-1}}\right)^{\beta(k;\omega)}$$

thus, using $\varphi(n^3 r) \leq \varphi(n^{-k})$, we have

$$\underline{\delta}_{\varphi}(z;\mu,\rho) \ge \lim_{k \to \infty} \frac{\ell(k) - k}{\varphi(n^{-k})} \log(a_0/a_{m-1}) = s(1/\sigma - 1) \log(a_0/a_{m-1}).$$

The other direction inequality is due to Corollary 3.1. The lemma is proved.

Proof of Theorem 1.1. Pick $z \in E$. Let $\boldsymbol{\omega} = (\mathbf{x}, \mathbf{y})$ be a coding of z. If $\boldsymbol{\omega} \in \mathbf{z}$ $\Omega_0 \cup \Omega_{m-1}$, the theorem holds by Lemma 4.1.

Now we assume that $\boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}$. Then **y** does not end in 0^{∞} and $(m-1)^{\infty}$. Let $(y_{j_t})_{t\geq 1}$ be a subsequence of $\mathbf{y} = (y_j)_{j\geq 1}$ satisfying that

$$(y_{j_t}, y_{j_{t+1}}) \neq (0, 0)$$
 and $(m - 1, m - 1), t \ge 1.$

Let k_t be the smallest integer such that $j_t + 1 \leq \ell(k_t)$, then $\beta(k_t; \boldsymbol{\omega}) \leq 1/\sigma + 2$.

Set $r_t = n^{-k_t-1}$. By the first assertion of Lemma 3.2, we have

$$\underline{\delta}_{\varphi}(z;\mu,\rho) \le \liminf_{t \to \infty} \frac{\log C_2 + (1/\sigma + 2)\log(a_0/a_{m-1})}{\varphi(n^{-k_t-1})} = 0.$$

The theorem is proved.

Proof of Theorem 1.2. Item (ii) is a consequence of Lemma 4.1. In the following, we prove Item (i). Moreover, by Corollary 3.1, we only need to show that

(4.2)
$$\overline{\delta}_{\varphi}(z;\mu,\rho) \geq \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \log(a_0/a_{m-1}), \text{ for } \boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}.$$

Let $z = \pi(\boldsymbol{\omega}) \in E$ with $\boldsymbol{\omega} \notin \Omega_0 \cup \Omega_{m-1}$.

Suppose $\beta(k; \boldsymbol{\omega}) = \ell(k) - k$ holds for infinitely many k. Let h be the largest integer such that $h \leq k-1$ and $y_{h+1} \neq 0$. Then $\beta(h; \boldsymbol{\omega}) = \ell(h) - h - 1$. Therefore, there exists a subsequence $(k_i)_{i\geq 1}$ of integers such that $\beta(k_i; \boldsymbol{\omega}) = \ell(k_i) - k_i - 1$. set $r_i = 1/n^{k_j+1}$. By the second assertion of Lemma 3.2, we have

$$\overline{\delta}_{\varphi}(z;\mu,\rho) \geq \limsup_{j \to \infty} \frac{\log U(z;n^6 r_j,\rho)}{\varphi(n^6 r_j)} \geq \limsup_{j \to \infty} \frac{\log C_2^{-1} + \beta(k_j;\boldsymbol{\omega}) \log(a_0/a_{m-1})}{\varphi(n^{-k_j+5})}$$
$$= \limsup_{j \to \infty} \frac{\ell(k_j) - k_j - 1}{\varphi(n^{-k_j+5})} \log(a_0/a_{m-1}) = s(1/\sigma - 1) \log(a_0/a_{m-1}),$$

so the lemma holds in this case since $\limsup_{k\to\infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \leq s(1/\sigma - 1).$

Suppose there exists $k_0 \in \mathbb{N}$ such that $\beta(k; \boldsymbol{\omega}) \leq \ell(k) - k - 1$ holds for $k \geq k_0$. Set $r_k = 1/n^{k+1}$. Similar as above, applying Lemma 3.2 to $k \ge k_0$ instead of k_j , we have

$$\overline{\delta}_{\varphi}(z;\mu,\rho) \geq \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k+5})} \log(a_0/a_{m-1}) \geq \limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{\varphi(n^{-k})} \log(a_0/a_{m-1}).$$

 _	

(The last equality holds since φ is decreasing.) This proves (4.2) as well as the item (i).

5. Remarks on $\delta_{-\log r}$ and $\delta_{\log |\log r|}$

Recall that $\delta = \delta_{-\log r}$ and $\Delta = \delta_{\log |\log r|}$ (See Example 1.1).

5.1. **Run length.** Let $\theta \in \{0, 1, \dots, m-1\}$. Set $\boldsymbol{\omega} = (\mathbf{x}, \mathbf{y}) \in \mathcal{D}^{\infty}$ with $\mathbf{y} = (y_j)_{j \ge 1}$, and let $l_{\theta}(k; \mathbf{y})$ be the run length of the letter θ in \mathbf{y} at the position k, *i.e.*, $l_{\theta}(k; \mathbf{y}) = t$ if $y_k = \dots = y_{k+t-1} = \theta$ and $y_{k+t} \neq \theta$. By convention we set $l_{\theta}(k; \mathbf{y}) = 0$ if $y_k \neq \theta$. It is well known that

Lemma 5.1 (Billingsley [5]). Let $\Sigma = \{0, 1, \ldots, m-1\}$. Let $\mathbf{p} = (p_0, \ldots, p_{m-1})$ be a probability weight which allows $p_j = 0$. Let $\nu_{\mathbf{p}}$ be the Bernoulli measure on Σ^{∞} associated with \mathbf{p} , then for all $p_{\theta} \neq 0$,

$$\nu_{\mathbf{p}}\left(\left\{\mathbf{y}\in\Sigma^{\infty};\ \limsup_{k\to\infty}\frac{l_{\theta}(k;\mathbf{y})}{-\log_{p_{\theta}}k}=1\right\}\right)=1.$$

Definition 5.1 (Modified run length). Let $\theta_0 \in \Sigma$ be a fixed letter. Define $l_{\theta_0,\theta}(k, \mathbf{y}) = t$ if $y_{k-1} = \theta_0$, $y_k = \cdots = y_{k+t-1} = \theta$ and $y_{k+t} \neq \theta$; otherwise, set $l_{\theta_0,\theta}(k, \mathbf{y}) = 0$.

By the same argument as the proof of Lemma 5.1, one can show that

Lemma 5.2. Let $\Sigma = \{0, 1, \dots, m-1\}$. Let $\mathbf{p} = (p_0, \dots, p_{m-1})$ be a probability weight which allows $p_j = 0$. Let $\nu_{\mathbf{p}}$ be the Bernoulli measure on Σ^{∞} associated with \mathbf{p} . If $p_{\theta_0} \neq 0$ and $p_{\theta} \neq 0$, then

$$\nu_{\mathbf{p}}\left(\left\{\mathbf{y}\in\Sigma^{\infty};\ \limsup_{k\to\infty}\frac{l_{\theta_{0},\theta}(k;\mathbf{y})}{-\log_{p_{\theta}}k}=1\right\}\right)=1.$$

Let \mathbb{P} be the uniform Bernoulli measure on \mathcal{D}^{∞} .

Lemma 5.3. Let E = K(n, m, D) be a BM-carpet of non-doubling type. Assume that $a_0 > a_{m-1}$. Then

$$\mathbb{P}\left(\left\{\boldsymbol{\omega}\in\mathcal{D}^{\infty};\ \limsup_{k\to\infty}\frac{\beta(k;\boldsymbol{\omega})}{-\log_{p_0}k/\sigma}=1\right\}\right)=1,$$

where $p_0 = a_0 / \# \mathcal{D}$.

Proof. Pick θ_0 such that $\theta_0, \theta_0 - 1 \in \mathcal{E}$. Let $L \ge 1$ be an integer and set $\epsilon = 1/L$. Denote

$$H_{L,1} = \left[(\mathbf{x}, \mathbf{y}) \in \mathcal{D}^{\infty} : l_0(k; \mathbf{y}) + l_{m-1}(k; \mathbf{y}) < (1+\epsilon) \log k / (-\log p_0) \text{ eventually} \right],$$

 $H_{L,2} = [(\mathbf{x}, \mathbf{y}) \in \mathcal{D}^{\infty} : l_{\theta_0,0}(k; \mathbf{y}) \ge (1-\epsilon) \log k / (-\log p_0) \text{ infinitely often}].$

Since $p_0 > p_{m-1}$, by Lemma 5.1 we have $\mathbb{P}(H_{L,1}) = 1$. On the other hand, we have $\mathbb{P}(H_{L,2}) = 1$ by Lemma 5.2. Let $H_L = H_{L,1} \cap H_{L,2}$, then $\mathbb{P}(H_L) = 1$.

Fix $\boldsymbol{\omega} = (\mathbf{x}, \mathbf{y}) \in H_L$. For k large enough, we have

$$\beta(k;\boldsymbol{\omega}) \le l_0(\ell(k) - \beta(k;\boldsymbol{\omega}) + 1; \mathbf{y}) + l_{m-1}(\ell(k) - \beta(k;\boldsymbol{\omega}) + 1; \mathbf{y}) \le (1+\epsilon)\log\ell(k)/(-\log p_0)$$

eventually since $\boldsymbol{\omega} \in H_{L,1}$. It follows that

(5.1)
$$\limsup_{k \to \infty} \frac{\beta(k; \boldsymbol{\omega})}{-\log_{p_0} k/\sigma} \le 1 + \epsilon$$

On the other hand, by $\boldsymbol{\omega} \in H_{L,2}$, there exists infinite many h such that

$$l_{\theta_0,0}(h; \mathbf{y}) \ge (1-\epsilon) \log h / (-\log p_0).$$

Set $k = \sigma(h-1+(1-\epsilon)(-\log_{p_0} h))$, we obtain $\beta(k; \boldsymbol{\omega}) \ge (1-\epsilon)\log h/(-\log p_0))-1$. It follows that

(5.2)
$$\limsup_{k \to \infty} \frac{\beta(k; \boldsymbol{\omega})}{-\log_{p_0} k/\sigma} \ge 1 - \epsilon.$$

Set $H = \bigcap_{L>1} H_L$, then H has full measure, and each $\omega \in H$,

$$\limsup_{k \to \infty} \frac{\beta(k; \boldsymbol{\omega})}{-\log_{p_0} k / \sigma} = 1$$

The lemma is proved.

5.2. Remarks on δ and Δ .

Theorem 5.1. Let E = K(n, m, D) be a BM-carpet of non-doubling type. Assume that $a_0 > a_{m-1}$. Then for μ_E -a.e. $z \in E$, $\delta(z; \mu_E) = 0$, and the range of $\overline{\delta}(z; \mu_E)$ is $[0, \delta_{\max}]$.

Proof. By Lemma 5.3,

$$\limsup_{k \to \infty} \frac{\beta(k; \boldsymbol{\omega})}{k} = 0$$

for \mathbb{P} -a.e. $\boldsymbol{\omega} \in \mathcal{D}^{\infty}$, hence $\delta(z; \mu_E) = 0$ for μ_E -a.e. $z \in E$ by (1.6) in Example 1.1 and Theorem 1.1.

Next, we prove that $\{\overline{\delta}(z;\mu_E); z \in E\} = [0, \delta_{\max}]$. Pick $t \in [0, \delta_{\max}]$, let $t' = t/\log_n(a_0/a_{m-1})$. Since E is of non-doubling type, there exist $(i', j), (i, j + 1) \in \mathcal{D}$ and $(i_0, 0) \in \mathcal{D}$.

Take an integer $p_1 > \sigma/(1-\sigma)$, and set $p_{k+1} = \ell(p_k)$ for $k \ge 1$. Then $p_k \to \infty$ as $k \to \infty$. Set

(5.3)
$$\boldsymbol{\omega} = \begin{pmatrix} i_0 \\ 0 \end{pmatrix}^{p_1} \prod_{k=1}^{\infty} \left[\begin{pmatrix} i \\ j+1 \end{pmatrix}^{\ell(p_k) - p_k - \lfloor t' p_k \rfloor} \begin{pmatrix} i_0 \\ 0 \end{pmatrix}^{\lfloor t' p_k \rfloor} \right],$$

and let $z = \pi(\boldsymbol{\omega})$. Then for $k \ge p_1$,

$$\beta(k; \boldsymbol{\omega}) \leq t'k$$
 and $\beta(p_k; \boldsymbol{\omega}) = \lfloor t'p_k \rfloor.$

So by (1.6), we have

(5.4)
$$\bar{\delta}(z;\mu_E) = \left(\limsup_{k \to \infty} \frac{\beta(k;\boldsymbol{\omega})}{k}\right) \log_n(a_0/a_{m-1}) = t' \log_n(a_0/a_{m-1}) = t.$$

Г		٦	
-	-		

r		1

Theorem 5.2. Let $E = K(n, m, \mathcal{D})$ be a BM-carpet of non-doubling type. Assume that $a_0 > a_{m-1}$. Then for μ_E -a.e. $z \in E$,

(5.5)
$$\overline{\Delta}(z;\mu_E,\rho) = -\log_{p_0}(a_0/a_{m-1})$$

where $p_0 = a_0 / \# \mathcal{D}$.

Proof. Clearly $\mu_E(\{z \in E; z = \pi(\boldsymbol{\omega}) \text{ with } \boldsymbol{\omega} \in \Omega_0 \cup \Omega_{m-1}\}) = 0$. So the theorem is a direct consequence of Lemma 5.3 and (1.7).

6. A point-wise doubling index γ

In this section, we define an alternative point-wise doubling index.

Let μ be a measure on a metric space X. Fix $0 < \rho < 1$. For $z \in X$, the upper doubling index of μ at z is defined by

$$\overline{\gamma}(z;\mu,\rho) = \limsup_{r \to 0} \sup_{B_{\rho r}(z') \subset B_r(z)} \frac{\log \mu(B_{\rho r}(z'))}{\log \mu(B_r(z))};$$

similarly, we define $\gamma(z; \mu, \rho)$. If the two values agree, we denote it by $\gamma(z; \mu, \rho)$.

Let $E = K(n, m, \mathcal{D})$ be a BM-carpet, and μ be the uniform Bernoulli measure on E. Denote $N = #\mathcal{D}$, and denote

$$\gamma_{\max} = \frac{\log N - (1 - \sigma) \log a_{m-1}}{\log N - (1 - \sigma) \log a_0}.$$

Theorem 6.1. Let E = K(n, m, D) be a BM-carpet of non-doubling type. Assume that $a_0 > a_{m-1}$. Let $0 < \rho < n^{-3}$. Then $\overline{\gamma}(z; \mu, \rho) \in [1, \gamma_{\max}]$ and γ_{\max} is attainable.

Proof. Let $z \in E$ and let $\boldsymbol{\omega}$ be a coding of z. Pick 0 < r < 1. Let k = k(r) be the integer such that $\frac{1}{n^{k(r)+2}} < r \leq \frac{1}{n^{k(r)+1}}$ and let

(6.1)
$$\Xi_k(z) := \{ Q_k(\mathbf{u}, \mathbf{v}); \ Q_k(\mathbf{u}, \mathbf{v}) \cap B_r(z) \neq \emptyset \}$$

as in Section 3.

Let $z' \in E$ and $B_{\rho r}(z') \subset B_r(z)$. By the first assertion of Lemma 3.2, we have

(6.2)
$$\frac{\log \mu(B_{\rho r}(z'))}{\log \mu(B_r(z))} \le \frac{-\log U(z; r, \rho)}{\log \mu(B_r(z))} + 1 \le \frac{-\log C_2 - \beta(k; \boldsymbol{\omega}) \log(a_0/a_{m-1})}{\log(4 \cdot \max\{\mu(Q); Q \in \Xi_k(z)\})} + 1$$

where the last inequality holds since $\Xi_k(z)$ is a cover of $B_r(z)$.

Let $Q_k(\mathbf{u}, \mathbf{v})$ be an approximate square located in a row adjacent to the row of $Q_k(\boldsymbol{\omega})$. By analysing the three cases that $\mathbf{v}|_{\ell(k)}$ equals to $\mathbf{y}|_{\ell(k)-1} * v_{\ell(k)}, \mathbf{y}|_{h-1} * v_h *$ $0^{\ell(k)-h}$, or $\mathbf{y}|_{h-1} * v_h * (m-1)^{\ell(k)-h}$, we have

$$\mu(Q_k(\mathbf{u}, \mathbf{v})) \le \frac{n(a^*)^{\ell(k)-k-\beta(k;\boldsymbol{\omega})}a_0^{\beta(k;\boldsymbol{\omega})}}{N^{\ell(k)}},$$

where $a^* = \max_{j \in \mathcal{E}} a_j$. It follows that

(6.3)
$$\frac{\log \mu(B_{\rho r}(z'))}{\log \mu(B_{r}(z))} \le \frac{-\log C_{2} + \log \left(\frac{a_{m-1}}{a_{0}}\right)^{\beta(k;\boldsymbol{\omega})}}{\log(4n) + \log \frac{(a^{*})^{\ell(k)-k-\beta(k;\boldsymbol{\omega})}a_{0}^{\beta(k;\boldsymbol{\omega})}}{N^{\ell(k)}}} + 1$$

It is easy to show that after removing the unimportant constants $-\log C_2$ and $\log(4n)$, the right hand side of the above formula attains maximum when $\beta(k; \boldsymbol{\omega}) = \ell(k) - k$. Hence

(6.4)
$$\overline{\gamma}(z;\mu,\rho) \le \limsup_{k \to \infty} \left[\log \left(\frac{a_{m-1}}{a_0} \right)^{\ell(k)-k} \left(\log \frac{a_0^{\ell(k)-k}}{N^{\ell(k)}} \right)^{-1} + 1 \right] = \gamma_{\max}.$$

By the second assertion of Lemma 3.2, if $\beta(k; \boldsymbol{\omega}) \leq \ell(k) - k - 1$, we have

(6.5)
$$\sup_{z' \in E, B_{\rho n^6 r}(z') \subset B_{n^6 r}(z)} \frac{\log \mu(B_{\rho n^6 r}(z'))}{\log \mu(B_{n^6 r}(z))} = \frac{-\log U(z; n^6 r, \rho)}{\log \mu(B_{n^6 r}(z))} + 1$$
$$\geq \frac{\log C_2 - \beta(k; \boldsymbol{\omega}) \log(a_0/a_{m-1})}{\log(\max\{\mu(Q); \ Q \in \Xi_k(z)\})} + 1$$

where the last inequality holds since $B_{n^3r}(z)$ covers every element of $\Xi_k(z)$.

Let $\boldsymbol{\omega}$ be a point in \mathcal{D}^{∞} such that $\beta(k; \boldsymbol{\omega}) = \ell(k) - k - 1$ for infinitely many k. For example, the $\boldsymbol{\omega}$ in (5.3) will do with a slight adjustment for $t' = 1/\sigma - 1$. Let $z = \pi(\boldsymbol{\omega})$. By (6.5) we have

(6.6)
$$\overline{\gamma}(z;\mu,\rho) \ge \lim_{j \to \infty} \left(\frac{\log C_2 + \log \left(\frac{a_{m-1}}{a_0}\right)^{\ell(k_j) - k_j - 1}}{\log(a^*) + \log \frac{a_0^{\ell(k_j) - k_j - 1}}{N^{\ell(k_j)}}} + 1 \right) = \gamma_{\max}.$$

The theorem is proved.

Remark 6.1. Under the assumption of Theorem 6.1, one can show that

- (i) If $\boldsymbol{\omega} \in \Omega_0 \cup \Omega_{m-1}$, then $\gamma(z; \mu, \rho) = \max\{\gamma_{\max} \cdot \mathbf{1}_{V_E}(z), 1\}$. (ii) $\gamma(z; \mu, \rho) = \max\{\gamma_{\max} \cdot \mathbf{1}_{V_E}(z), 1\}$.
- (iii) For μ -a.e. $z \in E, \overline{\gamma}(z; \mu, \rho) = 1$.

7. Proofs of Theorems 1.3, 1.4 and 1.5

In this section, we always assume that $E = K(n, m, \mathcal{D}), F = K(n, m, \mathcal{D}')$ are two BM-carpets and $f : E \to F$ is a bi-Lipschitz map with Lipschitz constant c_0 . We use $(a_j)_{j=0}^{m-1}$ and $(a'_j)_{j=0}^{m-1}$ to denote the fiber sequences of \mathcal{D} and \mathcal{D}' , respectively. Denote $N = \#\mathcal{D}$ and $N' = \#\mathcal{D}'$. Denote $s = \#\{j; a_j > 0\}$ and $s' = \#\{j; a'_j > 0\}$. Let

(7.1)
$$a_1^* > a_2^* > \dots > a_{\tilde{p}}^*$$

be the distinct non-zero terms of $(a_j)_{j=0}^{m-1}$, and let M_i be the occurrence of a_i^* , let

(7.2)
$$b_1^* > b_2^* > \dots > b_{\tilde{q}}^*$$

be the distinct non-zero terms of $(a'_j)_{j=0}^{m-1}$, and let M'_i be the occurrence of b^*_i .

Denote by μ_E and μ_F the uniform Bernoulli measures of E and F respectively. Huang *et al.* [11] obtained the following Theorem, see also Rao *et al.* [18].

Theorem 7.1 ([11, 18]). Let E and F be two BM-carpets. If $f : E \to F$ is a bi-Lipschitz map, then $\mu_F \circ f$ is equivalent to μ_E , namely, there exists $\zeta > 0$ such that

(7.3)
$$\zeta^{-1}\mu_E(A) \le \mu_F(f(A)) \le \zeta\mu_E(A)$$

for any Borel set $A \subset E$. Consequently, μ_E and μ_F have the same multifractal spectrum.

By the above result, Rao *et al.* [18] characterized when μ_E and μ_F have the same multifractal spectrum, see also Banaji *et al.* [2].

Theorem 7.2 ([2, 18]). Let E = K(n, m, D) and F = K(n, m, D') be two BMcarpets. Then μ_E and μ_F have the same multifractal spectrum if and only if

(7.4)
$$\tilde{p} = \tilde{q} \quad and \quad \frac{a_i^*}{b_i^*} = \left(\frac{M_i'}{M_i}\right)^{1/\sigma} = \left(\frac{s'}{s}\right)^{1/\sigma} = \left(\frac{N}{N'}\right)^{1/(1-\sigma)}, \text{ for } i = 1, \dots, \tilde{p}.$$

Proof of Theorem 1.3. Without loss of generality, we assume that $a_0 > a_{m-1} > 0$ and $a'_0 > a'_{m-1} > 0$. Let ζ be the constant in Theorem 7.1 with respect to E and F. Let $f: E \to F$ be a bi-Lipschitz map with a Lipschitz constant c_0 .

Firstly, we prove (1.9). Pick any $z \in E$. Let 0 < r < 1. Then

$$f(B_{c_0^{-1}r}(z)) \subset B_r(f(z)) \subset f(B_{c_0r}(z)).$$

By Theorem 7.1, we have

(7.5)
$$\zeta^{-1}\mu_E(B_{c_0^{-1}r}(z)) \le \mu_F(B_r(f(z))) \le \zeta\mu_E(B_{c_0r}(z)).$$

Let $0 < \rho < n^{-3}c_0^{-2}$. Pick any $\omega' \in F$ satisfying $B_{\rho r}(\omega') \subset B_r(f(z))$. Similar as (7.5), we have

(7.6)
$$\zeta^{-1}\mu_E(B_{c_0^{-1}\rho r}(f^{-1}(\omega'))) \le \mu_F(B_{\rho r}(\omega')) \le \zeta\mu_E(B_{c_0\rho r}(f^{-1}(\omega'))).$$

By (7.5) and (7.6), we have

$$\log\left(\frac{\mu_{F}(B_{r}(f(z)))}{\mu_{F}(B_{\rho r}(\omega'))}\right) \le \log\zeta^{2} + \log\left(\frac{\mu_{E}(B_{c_{0}r}(z))}{\mu_{E}(B_{c_{0}}^{-1}\rho r}(f^{-1}(\omega')))}\right).$$

Since $B_{\rho r}(\omega') \subset B_r(f(z))$, we have $B_{c_0^{-1}\rho r}(f^{-1}(\omega')) \subset B_{c_0 r}(z)$, so

$$\overline{\delta}_{\varphi}(f(z);\mu_F,\rho) \le \overline{\delta}_{\varphi}(z;\mu_E,c_0^{-2}\rho);$$

by symmetry, we have $\overline{\delta}_{\varphi}(z; \mu_E, \rho c_0^2) \leq \overline{\delta}_{\varphi}(f(z); \mu_F, \rho)$. Finally, since $\overline{\delta}_{\varphi}$ is irrelevant with ρ , we obtain

$$\overline{\delta}_{\varphi}(z;\mu_E) = \overline{\delta}_{\varphi}(f(z);\mu_F).$$

Clearly, the above equality also holds if we replace $\overline{\delta}_{\varphi}$ by $\underline{\delta}_{\varphi}$. (1.9) is proved.

Let $\delta = \delta_{-\log r}$, then $\underline{\delta}(z; \mu_E) = \delta_{\max} \cdot \mathbf{1}_{V_E}(z)$. Hence, $z \in V_E$ if and only if $\underline{\delta}(z; \mu_E) > 0$, if and only if $\underline{\delta}(f(z); \mu_F) > 0$, if and only if $f(z) \in V_F$. This proves

 $V_F = f(V_E)$. In particular, $V_E = \emptyset$ if and only if $V_F = \emptyset$, that is, E satisfies the VSC if and only if F does.

Remark 7.1. Under the assumption of Theorem 1.3, we also have

(7.7)
$$\overline{\gamma}(z;\mu_E) = \overline{\gamma}(f(z);\mu_F).$$

Proof of Theorem 1.4. Without loss of generality, we assume that $a_0 > a_{m-1} > 0$ and $a'_0 > a'_{m-1} > 0$. By Theorem 5.1 and Theorem 1.3, we have $\delta_{E,\max} = \delta_{F,\max}$, then

(7.8)
$$\frac{a_0}{a_{m-1}} = \frac{a'_0}{a'_{m-1}}.$$

Similarly, by Theorem 6.1 and Remark 7.1 we have $\gamma_{E,\max} = \gamma_{F,\max}$, thus

$$\frac{(1-\sigma)\log(a_0/a_{m-1})}{\log N - (1-\sigma)\log a_0} = \frac{(1-\sigma)\log(a_0'/a_{m-1}')}{\log N' - (1-\sigma)\log a_0'}$$

by using $\gamma_{E,\max} - 1 = \gamma_{F,\max} - 1$, summing up with (7.8) implies that

(7.9)
$$\frac{a_0}{a'_0} = \frac{a_{m-1}}{a'_{m-1}} = \left(\frac{N}{N'}\right)^{\frac{1}{1-\sigma}}$$

By Theorem 5.2, we have

$$\overline{\Delta}(z;\mu_E,\rho) = -\log_{p_0}(a_0/a_{m-1}) \text{ for } \mu_E\text{-a.e. } z \in E,$$

$$\overline{\Delta}(f(z);\mu_F,\rho) = -\log_{p_0'}(a_0'/a_{m-1}') \text{ for } \mu_F\text{-a.e. } f(z) \in F,$$

where $p'_0 = a'_0/N'$. So $\log_{p_0}(a_0/a_{m-1}) = \log_{p'_0}(a'_0/a'_{m-1})$ by (1.9) with $\overline{\Delta} = \overline{\delta}_{\log|\log r|}$ and it follows that

$$\frac{a_0}{N} = p_0 = p'_0 = \frac{a'_0}{N'},$$

which together with (7.9) imply N = N'. By (7.4) and (7.9), the theorem is proved.

Proof of Theorem 1.5. Let E and F be two BM-carpets and $E \sim F$.

If E and F are of non-doubling type, the theorem holds by Theorem 1.4. If E and F do not possess vacant rows, then s = s' = m, thus N = N' by (7.4) in Theorem 7.2, the theorem holds in this case. Hence, in the following, we always assume that $E, F \in \mathcal{M}_{v,d}$.

If E and F are not totally disconnected, since both E and F possess vacant rows, all connected components of E and F are horizontal line segments of length 1, then $a_1^* = (a_1')^* = n$, thus N = N' by (7.4) in Theorem 7.2, and the theorem holds in this case. This proves that the theorem holds provided $E, F \notin \mathcal{M}_{t.v.d.}$

Finally, by [18, Theorem 1.4], the theorem holds in the case that $E, F \in \mathcal{M}_{t,v,d}$ and $\log m / \log n$ is irrational. The theorem is proved.

References

- R. Anttila, Pointwise Assouad dimension for measures, Proc. Roy. Soc. Edinburgh Sect. A, 153 (2023), no. 6, 2053-2078.
- [2] A. Banaji and I. Kolossváry, Intermediate dimensions of Bedford-McMullen carpets with applications to Lipschitz equivalence, Adv. Math., 449 (2024), Paper No. 109735, 69 pp.
- [3] T. Bedford, Crinkly curves, Markov partitions and dimensions, PhD Thesis, University of Warwick, (1984).
- [4] A. Beurling and L. Ahlfors, The boundary correspondence under quasiconformal mappings, Acta Math., 96 (1956), 125-142.
- [5] P. Billingsley, Probability and Measure, John Wiley & Sons, New York, 1995.
- [6] J.M. Fraser, Assouad dimension and fractal geometry, Cambridge Tracts in Mathematics (Cambridge University Press, 2020).
- [7] J.M. Fraser and D.C. Howroyd, On the upper regularity dimensions of measures, Indiana Univ. Math. J., 69 (2020), 685-712.
- [8] J.M. Fraser and H. Yu, Assouad-type spectra for some fractal families, Indiana Univ. Math. J., 67 (2018), 2005-2043.
- [9] P. Hajłasz and P. Koskela, Sobolev met Poincaré, Mem. Amer. Math. Soc., 145 (2000), x+101.
- [10] J. Heinonen, Lectures on analysis on metric spaces, Universitext (Springer-Verlag, New York, 2001).
- [11] L.Y. Huang, H. Rao, Z.Y. Wen and Y.L. Xu, Box-counting measures of metric spaces and its applications, Preprint (2022), (arXiv:2111.00752[math.GT]).
- [12] L.Y. Huang and Y. Zhang, Lipschitz classification of Bedford-McMullen carpets with locally uniform horizontal fibers, Topology Appl., 350 (2024), Paper No. 108906, 15 pp.
- [13] A. Käenmäki, J. Lehrbäck and M. Vuorinen, Dimensions, Whitney covers, and tubular neighborhoods, Indiana Univ. Math. J., 62 (2013), 1861-1889.
- [14] B.M. Li, W.X. Li and J.J. Miao, Lipschitz equivalence of McMullen sets, Fractals, 21(3 & 4) (2013), 1350022, 11 pages.
- [15] H. Li, C. Wei and S.Y. Wen, Doubling property of self-affine measures on carpets of Bedford and McMullen. Indiana Univ. Math. J., 65 (2016), 833-865.
- [16] C. McMullen, The Hausdorff dimension of general Sierpiński carpets, Nagoya Math. J., 96 (1984), 1-9.
- [17] J.J. Miao, L.F. Xi and Y. Xiong, Gap sequences of McMullen sets, Proc. Amer. Math. Soc., 145 (2017), 1629-1637.
- [18] H. Rao, Y.M. Yang and Y. Zhang, Invariance of multifractal spectrum of uniform self-affine measures and its applications. Preprint (2020), (arXiv:2005.07451 [math.DS]).
- [19] E. M. Stein, Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals, Princeton Mathematical Series, 43 (Princeton University Press, Princeton, NJ, 1993).
- [20] Y.M. Yang and Y. Zhang, Lipschitz classification of Bedford-McMullen carpets with uniform horizontal fibers, J. Math. Anal. Appl., 495 (2021), 124742.
- [21] Y.M. Yang and Y. Zhang, Constructing bi-Lipschitz maps between Bedford-McMullen carpets via symbolic spaces, J. Math. Anal. Appl., 528 (2023), 127514.

Department of Mathematics and Statistics, Central China Normal University, Wuhan, 430079, China

Email address: hrao@mail.ccnu.edu.cn

DEPARTMENT OF MATHEMATICS AND STATISTICS, CENTRAL CHINA NORMAL UNIVERSITY, WUHAN, 430079, CHINA

Email address: xu_yl@mails.ccnu.edu.cn

Department of Mathematics and Statistics, Central China Normal University, Wuhan, 430079, China

Email address: yzhang@mail.ccnu.edu.cn