

ON THE JUMP OF THE COVER TIME IN RANDOM GEOMETRIC GRAPHS

CARLOS MARTINEZ AND DIETER MITSCHKE

ABSTRACT. In this paper we study the cover time of the simple random walk of the giant component of supercritical d -dimensional random geometric graphs on n vertices. We show that the cover time undergoes a jump at the connectivity threshold radius r_c : with r_g denoting the threshold for having a giant component, we show that if the radius r satisfies $r_c < r \leq (1 - \varepsilon)r_g$ for any $\varepsilon > 0$, the cover time of the giant component is asymptotically almost surely $\Theta(n \log^2 n)$. On the other hand, we show that for $r \geq (1 + \varepsilon)r_c$, the cover time of the graph is asymptotically almost surely $\Theta(n \log n)$ (which was known for $d = 2$ only for a radius larger by a constant factor).

Keywords: cover time, random walk, random geometric graphs

MSC Class: 05C80, 60D05, 05C81

1. INTRODUCTION

We denote by $\mathcal{G}(n, r, d)$ the by now classic model of random geometric graphs defined as follows: let d and n be positive integers, and let r be a positive real number, possibly depending on n . The vertices of the graph are the points of a Poisson Point Process (PPP) of intensity 1 in $\Lambda_n = \left[-\frac{n^{1/d}}{2}, \frac{n^{1/d}}{2}\right]^d$, that is, the d -dimensional axis-parallel cube of volume n centered at the origin, and two vertices are connected by an edge if their Euclidean distance is at most r . Random geometric graphs were originally introduced by Gilbert [24] as a model for telecommunication networks, and since then they received a lot of attention both from a theoretical point of view (see for example the monograph of Penrose [35], or also [36]) as well as from an applied point of view (see for example [3, 33, 23]).

The simple random walk on the vertices of a (throughout the paper finite and connected) graph $G = (V, E)$ is a discrete-time stochastic process with random variables X_0, X_1, \dots such that $X_0 = v$ for some $v \in V$, and for $i \geq 0$, X_{i+1} is a vertex chosen uniformly at random from the neighbors of X_i . The *cover time* of the simple random walk is then defined as

$$\min\{t \geq 0 : \bigcup_{s=0}^t \{X_s\} = V\}.$$

The cover time in general depends on the starting vertex, and it therefore makes sense to consider

$$\max_{v \in V, X_0=v} \min\{t \geq 0 : \bigcup_{s=0}^t \{X_s\} = V\}.$$

Also, since the cover time is a random variable, we are interested in its expected value (starting from a worst-case vertex). We thus define for $G(V, E)$,

$$\tau_{cov}(G) = \mathbb{E}\left[\max_{v \in V, X_0=v} \min\{t \geq 0 : \bigcup_{s=0}^t \{X_s\} = V\}\right].$$

Dieter Mitsche has been partially supported by grant GrHyDy ANR-20-CE40-0002 and by Fondecyt grant 1220174.

1.1. Main results. In order to state our results, we need a bit of additional notation. For a sequence $\{\mathcal{A}_n\}_n$ of events, we say that \mathcal{A}_n holds *asymptotically almost surely (a.a.s.)*, if $\Pr[\mathcal{A}_n] \rightarrow 1$, as $n \rightarrow \infty$. For $G \in \mathcal{G}(n, r, d)$ (identifying G with its vertex set), define $r_c = r_c(n, d)$ as the smallest radius such that a.a.s. a random geometric graph with such a radius is connected. It is well known (see [25]) that $r_c = (\log n/V_d)^{1/d}$, where V_d is the volume of the unit ball in d dimensions. Define also $r_g = r_g(n, d)$ as the smallest radius such that a.a.s. a random geometric graph with such a radius has a giant component, that is, a component of size linear in n . It is well known that $r_g = \Theta(1)$ (the existence of such a threshold is well known, the exact value is only conjectured, even in 2 dimensions). For rigorous lower and upper bounds on r_g see [35].

The cover time of $\mathcal{G}(n, r, d)$ was, as far as we know, only studied in two papers: first, Avin and Ercal (see [5]) showed that for $G \in \mathcal{G}(n, r, 2)$ with $r^2 > 8 \log n$, that is, a constant factor above the connectivity threshold, $\tau_{cov}(G) = \Theta(n \log n)$ a.a.s. Second, Cooper and Frieze [12] showed that for $d \geq 3$ and $G \in \mathcal{G}(n, r, d)$ with for $r = (c \log n/V_d)^{1/d}$ with $c > 1$ being arbitrary and V_d as before being the volume of the d -dimensional unit ball, a.a.s. $\tau_{cov} = c \log\left(\frac{c}{c-1}\right) n \log n$ (their result does not apply for $d = 2$). That is, these two papers study random geometric graphs above the connectivity threshold, and in the case $d = 2$, only by a constant factor above the connectivity threshold. In this paper, we look at sparser graphs not dealt with by these papers: for $G \in \mathcal{G}(n, r, d)$ and $r \geq r_g$, we denote by $L_1(G)$ its largest component (that is, the connected graph induced by the vertex set of this component). Our first main result is the following:

Theorem 1.1. *Let $\varepsilon > 0$ be arbitrarily small. Let $d \geq 2$. Let $G \in \mathcal{G}(n, r, d)$ be a random geometric graph with $r_g < r \leq (1 - \varepsilon)r_c$. Then, a.a.s., $\tau_{cov}(L_1(G)) = \Theta(n \log^2 n)$.*

Our second main result complements the previous result, by considering a random geometric graph right above the connectivity threshold (which was left open in the case $d = 2$):

Theorem 1.2. *Let $\varepsilon > 0$ be an arbitrarily small constant. Let $d = 2$. Let $G \in \mathcal{G}(n, r, d)$ be a random geometric graph with $r \geq (1 + \varepsilon)r_c$. Then, a.a.s. $\tau_{cov}(G) = \Theta(n \log n)$.*

Theorem 1.1 and 1.2 together show that for $G \in \mathcal{G}(n, r, d)$, τ_{cov} undergoes a jump at r_c , decreasing from $\Theta(n \log^2 n)$ to $\Theta(n \log n)$. Our proofs shed some light into the behavior of τ_{cov} right at r_c , and we comment on this in the concluding remarks (we do not know the exact behavior, though).

1.2. Further related work. Apart from the aforementioned papers, the cover time is a classic parameter in the study of random walks that has been analyzed in many contexts and on different graphs: on the one hand, for general graphs, Feige showed in [19, 20] that for any connected graph on n vertices, the cover time is a.a.s. at least $(1 - o(1))n \log n$ and at most $(4/27)n^3 + o(n^3)$, and he also gave graphs attaining the asymptotic bounds.

On the one hand, from an algorithmic point of view, for general graphs, Kahn, Kim, Lóvasz and Vu in [26] first established a $(\log \log n)^2$ -polynomial time approximation algorithm for the cover time. Later this was improved by Ding, Lee and Peres, who in a breakthrough result in [18] found a polynomial-time approximation algorithm approximating the cover time up to a constant factor (later this result was refined by Ding [17] and Zhai [38]).

Again in another context, on the random graph $\mathcal{G}(n, p)$, a series of papers of Cooper and Frieze analyzed the cover time of the largest component of $\mathcal{G}(n, p)$ for different values of p (summarized in Theorem 1 of [22]). Moreover, in the aforementioned paper [22], Frieze, Pegden and Tkocz obtained more precise estimates for the cover time of the emerging giant component, improving on previous results of Barlow, Ding, Nachmias and Peres [6] and Cooper, Frieze, Lubetzky [13].

Finally, in a geometric setup that is the most similar to the current work, in a series of papers, the cover time of the d -dimensional toroidal grid of side length n (and thus with n^d vertices) has been studied, and remarkably precise results are known by now. For $d = 2$, an upper bound of the cover time of the correct order was first found by Aldous [4], and a non-matching first order lower bound by Lawler [29]. Finally, the correct first order was obtained in the landmark paper

of Dembo, Peres, Rosen and Zeitouni [15]: they showed that the cover time of the 2-dimensional torus satisfies

$$\tau_{cov}/(n^2 \log^2 n) \rightarrow 4/\pi,$$

as $n \rightarrow \infty$, where the convergence is in probability. Ding [16] improved the result by showing that $\sqrt{\tau_{cov}/n^2} - \frac{2}{\sqrt{\pi}} \log n$ is of order $\log \log n$ with probability tending to 1 as $n \rightarrow \infty$. Abe [2] then found the second order term: he showed that there exists $c \in (0, 1)$, such that

$$2 \log n - \log \log n - (\log \log n)^c \leq \frac{\tau_{cov}}{\frac{2}{\pi} n^2 \log n} \leq 2 \log n - \log \log n + (\log \log n)^c$$

with probability tending to 1 as $n \rightarrow \infty$, similar to a previous bound in the continuous setup of Brownian motion by Belius and Kistler [8]. Recently, using a comparison with the extremal landscape of the discrete Gaussian free field, Louidor and Saggiatti [32] went even further: they showed that the square root of the cover time normalized by the size of the subset is tight around $\frac{1}{\sqrt{\pi}} \log n - \frac{1}{4\sqrt{\pi}} \log \log n$. The latter approach was inspired by an analysis of the cover time on the binary tree of depth n : in this case, Cortines, Louidor and Saggiatti showed in [14] that the cover time, normalized by $2^{n+1}n$, and then centered by $(\log 2)n - \log n$, admits a weak limit as $n \rightarrow \infty$. The limiting distribution in this case obtained was a Gumbel random variable with rate one, shifted randomly by the logarithm of the sum of the limits of the derivative martingales associated with two negatively correlated discrete Gaussian free fields on the infinite version of the tree. For $d \geq 3$, Belius in [7] showed that, normalized by the volume of the torus and then centered by a constant times $\log n^d$, the cover time converges in law to a Gumbel distribution.

In the context of a percolated grid (of side length n), some results are known as well: whereas the cover time of (unpercolated) \mathbb{Z}^2 is a.a.s. of the order $n^2 \log^2 n$ and of (unpercolated) \mathbb{Z}^d for $d \geq 3$ of the order $n^d \log n$, Abe in [1] showed that for any $d \geq 2$, the cover time of the giant component of supercritical \mathbb{Z}^d is a.a.s. of the order $n^2 \log^2 n$, thereby showing a quantitative difference of unpercolated and percolated \mathbb{Z}^d for $d \geq 3$.

In the context of random hyperbolic graphs, recently, Kiwi, Schepers and Sylvester studied the cover time of the giant component of random hyperbolic graphs on n vertices. They showed that the cover time of the giant component of these graphs is $\Theta(n \log^2 n)$ (see [28]).

1.3. Outline of proofs. The lower bound of the proof of Theorem 1.1 is comparably simpler: depending on the regime of the radius, either we find pending paths of order $\log n/r^d$ (as in the paper of Abe [1]), or we will find many vertices of degree 1 inside the giant component. The upper bound is also inspired by Abe's result in [1] on τ_{cov} of a supercritical percolation cluster on \mathbb{Z}^d : we first mimic Abe's backbone construction from [1] over Bernoulli percolation now using a suitable renormalization scheme on the random geometric graph for fixed radius r . We then use this structure to build a flow that dissipates energy in a convenient way: we have to be careful with the construction as there are additional obstacles that do not appear in Bernoulli percolation, such as the fact that crossings could be arbitrarily close to one another or that we may have to extract induced paths from paths in densely populated areas. We then extend this result to larger radii below the connectivity threshold by a simple coupling. We remark that if we were interested in showing Theorem 1.1 only for $r = \Theta(1)$ and $r \geq Cr_g$ for C being a large constant, the theorem would directly follow as a direct application of [1], using a tessellation of cubes of side length equal to $r/\sqrt{5}$ (for $d = 2$, and changing the constant for other dimensions) and a coupling with Bernoulli percolation. However, since we show it for $r > r_g$, we need the already mentioned renormalization scheme for the upper bound.

The proof of Theorem 1.2 elaborates on the idea of the upper bound of Theorem 1.1 by additional ingredients: on the one hand, since the typical degree is high, one extra splitting of the flow is introduced in order to be able to bound the effective resistance between any two vertices of a certain backbone (which is large and well spread out) by $O(1/\log n)$. On the other hand, we then show that other vertices outside this backbone cannot be separated from the backbone by taking out too few vertices: we will find many short internally vertex-disjoint paths between any vertex outside and some vertex inside the backbone, on which we can split the flow again evenly.

Further notation. For $j \in \mathbb{Z}^d$, we denote by j_i the i -th coordinate. We denote by e_i the i -th standard basis vector in d dimensions. For a set A , we denote its cardinality by $|A|$. For $G = (V, E)$ and $x, y \in V$, we write $x \sim y$ for $\{x, y\} \in E$. We denote by $d_G(x, y)$ the graph distance between x and y , that is, the number of edges on a shortest path between x and y . If x and y are identified with points, we also denote by $d_E(x, y)$ the Euclidean distance between x and y . We say that two paths of vertices $u_0 := x, u_1, \dots, u_m, u_{m+1} := y$ and $v_0 := x, v_1, \dots, v_n, v_{n+1} := y$ between x and y are internally vertex-disjoint, if $\{u_0, \dots, u_{m+1}\} \cap \{v_0, \dots, v_{n+1}\} = \{x, y\}$. A path of vertices v_1, \dots, v_k in $G(V, E)$ is induced, if $\{v_i, v_j\} \in E$ implies $j = i + 1$ for some $i = 1, \dots, k - 1$. Note that in an induced path of $G \in \mathcal{G}(n, r, d)$ for every vertex of the path except for the first and last there are exactly two other vertices in the path at Euclidean distance at most r , and for the first and last vertex there is only one other vertex in the path at Euclidean distance at most r . Observe also, by taking out vertices if needed, one may obtain an induced path from a path that is not, and that has the same endpoints as the original path.

Plan of paper. In Section 2 we recall basic facts on random walks, in particular on the resistance, as well as facts on random geometric graphs and Bernoulli percolation. In Section 3 we then introduce a renormalization scheme that will turn out to be very useful in our proofs. Section 4 is then dedicated to the proof of the lower bound of Theorem 1.1. In Section 5 we then show the corresponding upper bound of Theorem 1.1, first for $r = \Theta(1)$, and later we show the small modifications needed for $r = \omega(1)$. Finally, in Section 6 we prove Theorem 1.2.

2. PRELIMINARIES

In this section we recall basics on different topics, and we also prove some easy preliminary results.

2.1. Markov chains and random walks. A network is a connected graph $G = (V, E)$ together with a set of positive real values $(\tau(e))_{e \in E}$, the so called *resistances* of the edges. We denote the network simply as G and make no distinction with the underlying graph. We define the conductance \mathfrak{c} of the edge e as $\mathfrak{c}(e) = \frac{1}{\tau(e)}$. Given a network, we define the associated transition probabilities by

$$P(x, y) = \frac{\mathfrak{c}(x, y)}{\mathfrak{c}(x)}$$

where $\mathfrak{c}(x) = \sum_{y \sim x} \mathfrak{c}(x, y)$. For $\{x, y\} \in E$, denote by $\vec{e} = \overrightarrow{xy}$ an oriented edge, oriented from x to y , and by $-\vec{e} = \overrightarrow{yx}$ the oriented edge, oriented from y to x . For $x, y \in V$, a *unit flow* from x to y is a function θ on oriented edges such that $\theta(\vec{xy}) = -\theta(\overrightarrow{yx})$, $\sum_{x \sim a} \theta(\overrightarrow{xa}) = 1$, $\sum_{a \sim y} \theta(\overrightarrow{ay}) = 1$ and

$$\sum_{z \sim a} \theta(\overrightarrow{za}) = 0 \text{ for all } z \in V \setminus \{x, y\}.$$

Next, we have a central definition:

Definition 2.1. The *energy* of a unit flow θ is given by

$$E(\theta) = \sum_{e \in E} \theta^2(e) \tau(e).$$

The *effective resistance* between $x, y \in V$ is then defined as

$$\mathcal{R}(x \longleftrightarrow y, G) = \inf\{E(\theta) : \theta \text{ is a unit flow from } x \text{ to } y\}$$

Remark 2.2. By definition, we may get upper bounds on the effective resistance by specifying a unit flow between x and y . A trivial upper bound for $\mathcal{R}(x \longleftrightarrow y, G)$ is therefore given by the graph distance between the vertices (define the unit flow over the shortest path with all edges having flow equal to 1).

The effective resistance can be interpreted as the value of the resistance that appears when replacing the entire network with only one edge between x and y that mimics the resistance between them in the original network. A *sub-network* G' of G is a connected sub-graph of G with the same

values of resistances on the remaining edges. We now collect a few results that will turn very useful for us. First, effective resistance can not decrease when we delete edges:

Theorem 2.3 (Rayleigh, Theorem 9.12 of [30]). *Let G' be a sub-network of G . Then, for $x, y \in G'$ we have*

$$\mathcal{R}(x \longleftrightarrow y, G) \leq \mathcal{R}(x \longleftrightarrow y, G').$$

When there is no room for confusion on the underlying network G we simply denote the effective resistance by $\mathcal{R}(x \longleftrightarrow y)$. Next, we state that the triangle inequality holds for the effective resistance:

Lemma 2.4 (Corollary 10.8 of [30]). *Let $G = (V, E)$ and let $x, y, z \in V$. Then*

$$\mathcal{R}(x \longleftrightarrow z) \leq \mathcal{R}(x \longleftrightarrow y) + \mathcal{R}(y \longleftrightarrow z)$$

For $G = (V, E)$ and $x, y \in V$, we say that $\Pi \subseteq E$ is an *edge-cutset* separating x from y if every path from x to y contains at least one edge in Π . In other words, if we remove Π from E then x is isolated from y . We will use the following Nash-Williams inequality to obtain lower bounds for effective resistances.

Lemma 2.5 (Proposition 9.15 of [30]). *Let $\{\Pi_k\}_k$ be a pairwise edge-disjoint family of edge-cutsets that separate x from y . Then*

$$\mathcal{R}(x \longleftrightarrow y) \geq \sum_k \left(\sum_{e \in \Pi_k} \mathfrak{c}(e) \right)^{-1}.$$

Note that for $G = (V, E)$, the simple random walk admits a natural network representation when the set of resistances is given by $\mathfrak{r}(e) = 1$ for all $e \in E$. We next relate τ_{cov} to the effective resistance:

Theorem 2.6 (Theorem 2.4 of [10]). *Let $G = (V, E)$ be a connected graph. Then there exists a universal constant $\gamma_1 > 0$ such that*

$$\tau_{cov} \leq \gamma_1 |E| \log(|V|) \left(\max_{x, y \in V} \mathcal{R}(x \longleftrightarrow y) \right).$$

Theorem 2.7 (Theorem 1.3 of [26]). *Let $G = (V, E)$ be a connected graph. Given $V' \subseteq V$, there exists a universal constant γ_2 such that*

$$\tau_{cov} \geq \gamma_2 |E| \log(|V'|) \left(\min_{\substack{x \neq y \\ x, y \in V'}} \mathcal{R}(x \longleftrightarrow y) \right).$$

2.2. Random geometric graphs. We next state and prove a few useful facts about random geometric graphs. We start with a large deviation bound for the number of edges (the result is much more general, we only state a simplified form that is sufficient to deduce the corollary needed for our purposes): We recall that a sequence of random variables X_n satisfies an upper tail large deviation principle with speed $s(n)$ and rate function $I(x)$, if for any closed set $F \subset [0, \infty)$,

$$\limsup_{n \rightarrow \infty} \frac{1}{s(n)} \log \mathbb{P} \left(\frac{X_n - \mathbb{E}[X_n]}{\mathbb{E}[X_n]} \in F \right) \leq - \inf_{x \in F} I(x),$$

and for any open set $G \subset [0, \infty)$,

$$\liminf_{n \rightarrow \infty} \frac{1}{s(n)} \log \mathbb{P} \left(\frac{X_n - \mathbb{E}[X_n]}{\mathbb{E}[X_n]} \in G \right) \geq - \inf_{x \in G} I(x).$$

Theorem 2.8 (Theorem 2.2 of [11]). *Denote by V_d the volume of the d -dimensional unit ball. Let $G = (V, E) \in \mathcal{G}(n, r, d)$ with $r = \Theta(1)$, so that $\mu_n := \mathbb{E}[|E|] = V_d n r^d / 2$. Let $I(x) := (\frac{1}{2}) \sqrt{2x}$. Then $|E|$ satisfies an upper tail large deviation principle with speed $\sqrt{\mu_n} \log n$ and rate function $I(x)$.*

We obtain the following corollary:

Corollary 2.9. *Let $d \geq 2$, and let $G \in \mathcal{G}(n, r, d)$ with $r = \Theta(1)$. A.a.s., every axis-parallelepiped with integer coordinates (inside Λ_n) of side lengths $Mr \log n$ for some constant $M > 0$ in 2 dimensions and Mr in the remaining $d - 2$ dimensions contains $O(\log^2 n)$ vertices and has $O(\log^2 n)$ edges with at least endpoint in it.*

Proof. First note that by standard Chernoff bounds for Poisson point processes, together with a union bound over all $O(n)$ parallelepipeds, each such parallelepiped contains a.a.s. $O(\log^2 n)$ vertices, showing the first part of the corollary. Condition on this below. Fix then any parallelepiped P and denote by X_v the degree of vertex v . Observe that the expected degree of each vertex v then satisfies $\mathbb{E}[X_v] = \Theta(1)$, and the total expected degree of vertices $X = \sum_{v \in P} X_v$ in P satisfies $\mathbb{E}[X] = O(\log^2 n)$. Thus, the same asymptotic bound holds for the number of edges $|E|$ with at least one endpoint in P . By Theorem 2.8, for any constant $c > 0$, $\mathbb{P}(|E| - \mathbb{E}[|E|]) \geq c\mathbb{E}[|E|] \leq e^{-I(c)\sqrt{\mu_n} \log n}$. For every M , since $I(c)$ is increasing in c , we may thus choose $c = c(M)$ large enough, so that $I(c)\sqrt{\mu_n} \geq 1.1$. Thus, with probability at least $1 - n^{-1.1}$, inside P the number of edges is $O(\log^2 n)$, and by a union bound over all $O(n)$ parallelepipeds, the corollary follows. \square

The following standard result will be used several times in the sequel. Since we were unable to find a reference, we give a proof for completeness.

Lemma 2.10. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$, and let $r > r_g$. A.a.s., the number of edges of $L_1(G)$ is $\Theta(nr^d)$.*

Proof. The upper bound on the number of edges follows trivially, as a.a.s. the total number of edges in G is $\Theta(nr^d)$. For the lower bound, first sprinkle a Poisson point process of intensity $1 - \varepsilon$ where $\varepsilon > 0$ is small enough so that the resulting graph has a giant component of size $\Theta(n)$ a.a.s. Now, each vertex of this giant has in expectation $\Theta(r^d)$ neighbors among vertices of the second PPP of intensity ε , and so the total number of edges of edges emanating from the giant component of the first PPP is $\Theta(nr^d)$. By a standard second moment method (we may assume $r = O(\log n)$), concentration follows. \square

We will make use of the following proposition that relates the Euclidean distance of two vertices of $L_1(G)$ to the graph distance. By Theorem 3 and Remark 5 of [21], we have:

Theorem 2.11. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$ and let $r > r_g$ and also $r = O(1)$. A.a.s. the following is true: there exist positive constants $C = C(r, d)$ and $C' = C'(r, d)$, such that for any two vertices $x, y \in L_1(G)$ with $d_E(x, y) \geq C \log n$, we have $d_G(x, y) \leq C' d_E(x, y)$.*

We get the following immediate corollary:

Corollary 2.12. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$ and let $r > r_g$ and also $r = O(1)$. Let $V' \subseteq L_1(G)$ be such that for any vertex $x \in L_1(G)$ we have that $d_E(x, V') < C \log n$ for the same constant C as in Theorem 2.11. Then, a.a.s., there exists a constant $c' > 0$ such that $d_G(x, V) \leq c' \log n$ for any vertex $x \in L_1(G)$.*

Proof. On the one hand, for every $x \in L_1(G)$ there exists $y \in V'$ with $d_E(x, y) < C \log n$, but on the other hand, since $x \in L_1(G)$, a.a.s. there exists $v \in L_1(G)$ with $d_E(x, v) \geq n^\varepsilon$ for some $\varepsilon > 0$. Thus, there exists $z \in L_1(G)$ such that $d_E(x, z) \geq C \log n$ and $d_E(x, z) \leq 2C \log n$: indeed, consider a shortest path $x = v_0, v_1, \dots, v_k = v$ for some $k \in \mathbb{N}$: since the Euclidean distance between two consecutive vertices is at most r , there exist vertices v_i and v_{i+1} on this path such that $d_E(x, v_i) < C \log n$ and $d_E(x, v_{i+1}) \geq C \log n$, but at the same time $d_E(x, v_{i+1}) \leq C \log n + r \leq 2C \log n$. Therefore, by Theorem 2.11, $d_G(x, v_{i+1}) \leq c' \log n$ for some $c' > 0$ and since $v_{i+1} \in V'$, the statement follows. \square

Next, we also use the following result (see for example [35], Theorem 10.18):

Theorem 2.13. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$ and let $r > r_g$. A.a.s., there exists $c > 0$ such that the second largest connected component of G is of size at most $c \log^2 n$.*

2.3. Bernoulli percolation. We also need results from Bernoulli percolation. Consider an i.i.d. family of independent Bernoulli variables with mean p , indexed by the edges of \mathbb{Z}^d (typically called bonds). We say that the bond e is open if its associated variable is equal to 1. For a given realization, we generate a subgraph of \mathbb{Z}^d with edges given by the open bonds, this random graph model is known as Bernoulli bond percolation. For $d \geq 2$, it is known that this model undergoes a non-trivial phase transition: there exists $0 < p_c^{\text{bond}} < 1$ such that if $p > p_c^{\text{bond}}$ then a.s. there is a unique (maximal) infinite connected component \mathcal{C} (see for example [9]), which we call the infinite cluster. If one considers variables indexed by the vertices of \mathbb{Z}^d instead of bonds we have Bernoulli site percolation. We say that a vertex is open if its associated variable is 1. We say that a path of vertices is open if all its vertices are open. As in bond percolation, for $d \geq 2$ there exists $0 < p_c^{\text{site}} < 1$ such that if $p > p_c^{\text{site}}$ then there is a.s. a unique infinite cluster.

We denote by $\mathbb{P}_p^{\text{site}}$ the law of Bernoulli site percolation with parameter p , and by $\mathbb{P}_p^{\text{bond}}$ the law of Bernoulli bond percolation with parameter p , respectively. We denote by L_1^{bond} the largest connected component of Bernoulli bond percolation inside the box $[-n^{1/d}/2, n^{1/d}/2]^d$ if it exists. Equipping L_1^{bond} with the network structure induced by the simple random walk, we have the following sharp estimate for the effective resistance of this network:

Theorem 2.14 (Theorem 1.1 of [1]). *Let $d \geq 2$. For bond percolation on \mathbb{Z}^d and $p \in (p_c^{\text{bond}}, 1)$ there are constants $c_1, c_2 > 0$ such that a.a.s., it is true that*

$$c_1 \log n \leq \max_{x, y \in L_1^{\text{bond}}} \mathcal{R}(x \longleftrightarrow y, L_1^{\text{bond}}) \leq c_2 \log n.$$

We also make use of the following percolation estimations of crossings in boxes for 2-dimensional site percolation. Given $[0, m] \times [0, n] \subseteq \mathbb{Z}^2$, a horizontal crossing is an open and self-avoiding path entirely contained within the box such that the first vertex of the path has its first coordinate equal to 0 and the final vertex has its first coordinate equal to m (path going from left to right). We define a vertical crossing in a similar way. A long crossing of a rectangle $[a, b] \times [c, d]$ is an open self-avoiding path along the longest dimension of the rectangle (ties broken arbitrarily). We will use the following result:

Lemma 2.15 (Theorem 11.1 of [27]). *Consider site percolation on \mathbb{Z}^2 with parameter $p > p_c^{\text{site}}$. There exist positive constants c_1, c_2, c_3 such that*

$$\mathbb{P}_p^{\text{site}}(\text{There exist at least } c_1 n \text{ disjoint horizontal crossings} \\ \text{in } ([0, m] \times [0, n]) \cap \mathbb{Z}^2) \geq 1 - c_2 m e^{-c_3 n}.$$

Finally, we need a result on the chemical distance for Bernoulli site percolation (see Lemma 2.3 of [1]):

Theorem 2.16. *Let $d \geq 2$ and $p > p_c^{\text{site}}$. There exist constants κ_0, c such that a.a.s. the following holds: for any $x, y \in L_1^{\text{bond}}$, and any $\kappa > \kappa_0$, if $d_E(x, y) \leq \kappa \log n$, then $d_G(x, y) \leq c\kappa \log n$.*

3. RENORMALIZATION SCHEME

In this section we explain the renormalization scheme that we use in order to obtain the upper bound of Theorem 1.1. The objective of the renormalization used by Abe in [1] is similar, here it is used to prove that there is a backbone of crossings of $L_1(G)$. We encounter several technical challenges proper of random geometric graphs that we have to consider. Fix throughout this section $M > 0$ large enough. For each $i \in \mathbb{Z}^d$ we define the cube

$$C_i = Mri + [-Mr/2, Mr/2]^d.$$

That is, C_i is an axis-parallel cube with center Mri and side length Mr . We say it has label i . We say that two cubes C_i and C_j are neighbors if $i \sim j$ (on the renormalized lattice resulting from tessellation of \mathbb{Z}^d into cubes of sidelength Mr). Denote by \mathcal{K}_n the set of cubes that are totally contained within $\Lambda_n = \left[-\frac{n^{1/d}}{2}, \frac{n^{1/d}}{2}\right]^d$. The choice of M depends on the section: in Section 4 the choice of M depends on another renormalization scheme that is introduced there. In the other sections, we may assume that by an appropriate choice M , all cubes C_i are either completely contained in Λ_n or they share at most some faces with Λ_n .

Now, let $\mu = \mu(d)$ be largest constant so that any two vertices in adjacent cubes (that is, sharing a $(d-1)$ -dimensional face) of side length r/μ are connected by an edge (for example, in $d=2$, $\mu = \sqrt{5}$). We will consider a tessellation using such cubes Q_i later on as well, for now we use it to define crossings. Let $U = [a_1, a_2] \times [b_1, b_2] \times U_{d-2}$ be a parallelepiped in \mathbb{R}^d , with U_{d-2} a parallelepiped in \mathbb{R}^{d-2} . We say that the parallelepiped U has a *crossing component*, if there exists a connected component (of the underlying $G \in \mathcal{G}(n, r, d)$) within the parallelepiped that contains vertices at Euclidean distance at most $\frac{r}{\mu}$ from each of the faces of the parallelepiped. Similarly as we did for \mathbb{Z}^d , we define also crossings: a *horizontal crossing* of $G \in \mathcal{G}(n, r, d)$ in U is a self-avoiding path of G such that the vertices of the path are totally contained in U , the first vertex of the path has first coordinate in the interval $(a_1, a_1 + r/\mu)$, and the last vertex of the path has first coordinate in the interval $(a_2 - r/\mu, a_2)$. A vertical crossing is defined in the same way, exchanging the first coordinate by the second coordinate and a_1, a_2 by b_1, b_2 , and as before, we use also the definition of a *long crossing* for a crossing in the longer dimension. We remark that while a horizontal (or vertical) crossing is a self-avoiding path, a crossing component is in general not a path (but it contains such a path).

We then define the indicator event U_i related to cube C_i to be true iff the following two conditions hold:

- (a) $C_i \cup C_j$ contains a crossing component for each $j \sim i$
- (b) C_i has only one connected component with diameter larger than $Mr/5$.




FIGURE 1. The central cube C_i and its surrounding paths. The colored lines represent vertex paths of $G \in \mathcal{G}(n, r, d)$.




FIGURE 2. A 2-dimensional slice \mathcal{K}_n^2 , with a red α -logarithmic strip.

By a straight-forward modification of a result of Penrose and Pisztora (Theorem 2 of [37]) (in the original definition of Penrose and Pisztora, the crossing component is at distance at most r from all the faces of the parallelepiped, but the proof works in exactly the same way for r/μ) we may state the following lemma:

Lemma 3.1. *Let $d \geq 2$, let M be a large enough constant, let $G \in \mathcal{G}(n, r, d)$ and let $r > r_g$. Fix $i \sim j \in \mathbb{Z}^d$. There exists a constant $\gamma > 0$ such that*

$$\mathbb{P}(C_i \cup C_j \text{ has a crossing component}) \geq 1 - e^{-\gamma M r^d},$$

$$\mathbb{P}(C_i \text{ has only one connected component with diameter larger than } Mr/5) \geq 1 - e^{-\gamma M r^d}.$$

Therefore, for M large enough there is a constant $\gamma' > 0$ such that for any $i \in \mathbb{Z}^d$ we have $\mathbb{P}(U_i = 1) \geq 1 - e^{-\gamma' M r^d}$. If $U_i = 1$ we say that the cube C_i is *good* (see Figure 1). Note that the indicator events U_i and U_j are independent if i, j are at graph distance (in the renormalized lattice graph) of at least 2. By Theorem 0.0 of [31], for M large enough, $(U_i)_{i \in \mathbb{Z}^d}$ stochastically dominates a family $(Y_i)_{i \in \mathbb{Z}^d}$ of i.i.d. Bernoulli variables with parameter p , where $p \rightarrow 1$ as $M \rightarrow \infty$. In particular, seeing the good cubes as open sites of Bernoulli site percolation, the set of good cubes thus has a giant component. Moreover, if a cube C_i is good and C_i is in the giant component of good cubes, then any vertex $x \in C_i$ belonging to the largest component of G restricted to C_i belongs to $L_1(G)$: indeed, by repeated application of Lemma 3.1, the giant component of the renormalized lattice contains, since the component with large diameter is unique, a.s. a long crossing of order $\Omega(n^{1/d}/r) \geq n^\varepsilon$ for some $\varepsilon > 0$ (we may assume $r = O(\log n)$ throughout the paper), which by Theorem 2.13 a.s. is the giant component of G .

A 2-dimensional slice of \mathcal{K}_n is a subset of cubes with indices i such that all but 2 coordinates are fixed (see Figure 2 for an example). We denote such a slice by \mathcal{K}_n^2 . Since every 2-dimensional slice is isomorphic to \mathbb{Z}^2 restricted to a box of a certain side length, by abuse of notation we can thus also define both vertical and horizontal orientations for each slice. We remark also, that, as before, the good cubes of \mathcal{K}_n^2 dominate supercritical 2-dimensional site percolation restricted to a large box.

Now, divide a 2-dimensional slice \mathcal{K}_n^2 into vertical disjoint strips of cubes C_i of width $\lceil \alpha \log n \rceil$. The total Euclidean width of the strip is $\lceil \alpha \log n \rceil M r$ (as each cube has side length $M r$), whereas its Euclidean diameter (that is, the largest Euclidean distance between any two points in such a strip) is $\Theta(n^{1/d})$, see Figure 2 (the strips that are at the boundary of the slice have a smaller width). We call them α -logarithmic vertical strips (or simply strips, in case the orientation is not important), and analogously we define α -logarithmic horizontal strips.

We prove the following result regarding long crossings of good cubes inside of Λ_n :

Lemma 3.2. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$ and assume $r_g < r < r_c$ and assume M large enough. Let α be a large enough positive constant. Then, a.s. there exists $c > 0$ such that each α -logarithmic strip that is not in the boundary of Λ_n of every 2-dimensional slice of Λ_n has at least $c \log n$ disjoint long crossings of good cubes.*

Proof. As noted after Lemma 3.1, for M large enough, the good cubes of the tessellation stochastically dominate a super-critical Bernoulli site percolation process. Observe that the number of slices of Λ_n is $O(n^{1/d})$, and also the number of α -logarithmic strips inside a slice is $O(n^{1/d})$. Hence, by Lemma 2.15 applied to every α -logarithmic strip, together with a union bound, the probability that at least one strip does not have $c \log n$ long crossings of good cubes is at most $O(n^{1/d})O(n^{1/d}c_2n^{1/d}e^{-c_3\alpha \log n}) = o(1)$, where we assumed α large enough. \square

4. LOWER BOUND ON THE COVER TIME

Recall that $\mu = \mu(d)$ is the largest constant so that any two vertices in adjacent cubes (that is, two cubes sharing a $d - 1$ -dimensional face) of sidelength r/μ are connected by an edge. We are going to use the tessellation into cubes C_i as described in Section 3 of side length $M r$ and introduce moreover a new finer tessellation of Λ_n into cubes Q_i , of the already mentioned sidelength $\frac{r}{\mu}$. The cube with index $i \in \mathbb{Z}^d$ is given by:

$$Q_i = i \frac{r}{\mu} + \left[0, \frac{r}{\mu} \right]^d$$

Let $\phi(d) = \phi$ be the smallest integer such that any pair of vertices in $G \in \mathcal{G}(n, r, d)$, contained in cubes Q_i and Q_j , is not connected by an edge if $|i - j|_1 > \phi$. For example, $\phi(2) = 4$. Let $x \in \Lambda_n$ be a vertex of $G \in \mathcal{G}(n, r, d)$ with $x \in Q_j$. We say that x is *m-special* if the following two conditions hold:

- Each of the cubes Q_k , with $k = j + e_1 s$ and $1 \leq s \leq m$, has exactly one point of the PPP. We call the set of cubes $\{Q_k\}$ by slight abuse of notation *m-pending path* with root at x (note that the graph induced by a pending path is not necessarily a path, since vertices in two cubes at distance 2 in the cube metric might also be connected by an edge).
- All cubes at distance less or equal to ϕ to the pending path in the $|\cdot|_1$ norm (where the cubes are seen as points of \mathbb{Z}^d) are empty (see Figure 3 for an example).

Tessellate Λ_n into cubes of side length Mr as in Section 3, such that Mr is large enough and also an integer multiple of r/μ (in this way cubes Q are either totally contained in cubes C or they share faces or they are disjoint). Note that such a choice of M only depends on the dimension. We only consider cubes of sidelength Mr fully contained in Λ_n ; it might happen that close to the boundary some part of Λ_n remains uncovered by the tessellation, but this causes no problem with the proofs of this section.

The following lemma tells us that for $c_1 > 0$ small enough, there are relatively many $\lfloor c_1 \frac{\log n}{r^d} \rfloor$ -pending paths, whose root is attached to the giant of $G \in \mathcal{G}(n, r, d)$.

Lemma 4.1. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$. There exist small constants c_1, ξ, γ (ξ depends only on d, c_1, γ), such that if $r_g < r < \gamma r_c$, the event*

$$\left\{ \left| x \in L_1(G) : x \text{ is } \left(\left\lfloor c_1 \frac{\log n}{r^d} \right\rfloor \right) \text{-special} \right| \geq n^\xi \right\}$$

occurs a.a.s.

Proof. In order to make sure that a $\lfloor c_1 \frac{\log n}{r^d} \rfloor$ -pending path completely is contained in Λ_n , we would like its root to be inside Λ_{m_n} , where

$$m_n = n^{1/d} - \frac{r}{\mu} \left\lfloor c_1 \frac{\log n}{r^d} \right\rfloor$$

Observe that by Lemma 2.13, a.a.s. all vertices that are in a horizontal crossing of Λ_{m_n} belong to $L_1(G)$, as a crossing in Λ_{m_n} is a connected component of size $\Omega(n^{1/d})$. Since we are interested in a result that holds a.a.s., we may condition on this.

Let $\mathcal{K}_{m_n}^2$ be the 2-dimensional slice where the first two coordinates are not fixed and all others are equal to 0 (see Figure 2). By Lemma 3.2, a.a.s., there are at least $c_2 m_n$ horizontal disjoint crossings of good cubes inside $\mathcal{K}_{m_n}^2$ for some $c_2 > 0$, and as before, we may condition on this as well. Each horizontal crossing of good cubes contains a crossing of G inside Λ_{m_n} (recall that the right endpoints of the crossing in G are at distance less than r/μ from the right boundary of the slice $\mathcal{K}_{m_n}^2$).

Let $x \in Q_i$ and $y \in Q_j$ be vertices of $G \in \mathcal{G}(n, r, d)$, with $i_1 = j_1$ (that is, with first coordinates being equal). Note that if $|i - j| > 2\phi$ (see the definition of ϕ above) then the events of the vertices x, y being $\left(\left\lfloor c_1 \frac{\log n}{r^d} \right\rfloor \right)$ -special are independent. Also, note that the configuration inside \mathcal{K}_{m_n} is independent of what happens in $\Lambda_n \setminus \mathcal{K}_{m_n}$. In particular, the event that a vertex x that is inside of $\mathcal{K}_{m_n}^2$, is at distance less than r/μ from the right boundary of $\mathcal{K}_{m_n}^2$ and belongs to $L_1(G)$ does not have an effect on the probability to have a $\lfloor c_1 \frac{\log n}{r^d} \rfloor$ -pending path in $\Lambda_n \setminus \mathcal{K}_{m_n}^2$ attached to it. Using the fact that $\lfloor c_1 \log n / r^d \rfloor \geq 1$, the probability for the latter event to happen is equal to

$$e^{-(r/\mu)^d (r/\mu)^d} \lfloor c_1 \frac{\log n}{r^d} \rfloor (e^{-(r/\mu)^d})^{c'} \lfloor c_1 \frac{\log n}{r^d} \rfloor (e^{-(r/\mu)^d})^{c''} \geq n^{-\rho},$$

where c', c'' are constants that depend only on the dimension and ρ is a constant that depends on $c_1, d, \varepsilon, \gamma$. Observe that we can choose c_1 small enough such that $\rho < 1/d$, and then we can

choose γ small enough such that $c_1 \frac{\log n}{r^d}$ is always greater than 1. Hence, using the fact that a.a.s. there are at least $c_3 n^{1/d}$ right endpoints of horizontal crossings of $L_1(G)$ in Λ_{m_n} (taken from long crossings of good cubes, and that are at distance less than r/μ of the right boundary of the slice), and that at least a constant fraction of all of them give rise to independent events of being $\lfloor c_1 \log n / r^d \rfloor$ -special, we have

$$\mathbb{P}(\{|\{x \in L_1(G) : x \text{ is } (\lfloor c_1 \log n / r^d \rfloor)\text{-special}\}| < n^\xi\}) \leq \mathbb{P}\left(\text{Bin}\left(c_4 n^{1/d}, n^{-\rho}\right) < n^\xi\right).$$

By choosing $\xi = (1/d - \rho)/2$ and applying Chernoff bounds, the theorem follows. \square

Theorem 4.2. *Let ε be a positive constant. Let $r_g < r < (1 - \varepsilon)r_c$, then a.a.s.,*

$$\tau_{\text{cov}}(L_1(G)) \geq \delta n \log^2 n,$$

where δ is a constant that only depends on ε .

Proof. Case 1: $r < \gamma' r_c$, where $\gamma' \leq \gamma$ (γ is the same constant from Lemma 4.1) is a small enough constant to be fixed later. Define as before

$$m_n = n^{1/d} - \frac{r}{\mu} \left\lfloor \frac{c_1 \log n}{r^d} \right\rfloor.$$

Let \mathcal{P} be the set of $\lfloor \frac{c_1 \log n}{r^d} \rfloor$ -special vertices of Λ_{m_n} that are contained within $L_1(G)$ (restricted to Λ_{m_n}). Denote by \mathcal{F} the set of vertices of $L_1(G)$ that are in the rightmost cube of a pending path with root at a vertex in \mathcal{P} (see Figure 3):

$$\mathcal{F} = \{y \in L_1(G) : y \in C_j, j = i + \lfloor c_1 \log n / r^d \rfloor, x \in Q_i \text{ and } x \in \mathcal{P}\}$$

Note that $|\mathcal{P}| = |\mathcal{F}|$. By Lemma 4.1, $|\mathcal{F}| \geq n^\xi$ a.a.s., and by construction, $\mathcal{F} \subseteq L_1(G)$. Now, we give a lower bound for the effective resistance between any pair of vertices of \mathcal{F} . Take $x, y \in \mathcal{F}$, and let us construct edge-disjoint edge-cutsets separating x from y . Let $x \in Q_i$. Let $\sigma(d) = \sigma$ be the minimum integer such that given σ consecutive cubes of a pending path, any path of G that connects any two vertices in different components of the pending path when taking out vertices in these σ cubes must have at least one edge with both endpoints inside the σ many consecutive cubes. For $1 \leq k \leq \lfloor \frac{1}{\sigma} \lfloor \frac{c_1 \log n}{r^d} \rfloor \rfloor - 1$ (the -1 is to avoid boundary cases that require separate calculations), let Π_k be the set of edges of $L_1(G)$ with both endpoints are inside the cube Q_j with $j = i - e_1 s$ and $\sigma(k-1) \leq s < \sigma k$. Choose γ' small enough such that $\lfloor \frac{1}{\sigma} \lfloor \frac{c_1 \log n}{r^d} \rfloor \rfloor$ is greater or equal to 2. Note that $|\Pi_k| \leq \binom{\sigma}{2}$, as there is only one vertex per cube Q_j . Moreover, observe that $\Pi := \{\Pi_k\}_k$ is a family of disjoint edge-cutsets between x, y , as any path between both vertices has to contain an edge belonging to Π_k (by definition of σ), for any k . By Lemma 2.5 we then have

$$\mathcal{R}(x \longleftrightarrow y, L_1(G)) \geq \sum_{k=1}^{\lfloor \frac{1}{\sigma} \lfloor \frac{c_1 \log n}{r^d} \rfloor \rfloor - 1} \frac{1}{\binom{\sigma}{2}} \geq \delta_1 \frac{\log n}{r^d},$$

for some $\delta_1 > 0$. Using Lemma 2.10 together with the formula of Theorem 2.7 we obtain the desired result for this case.

Case 2: $\gamma' r_c \leq r \leq (1 - \varepsilon)r_c$. In this case we will use a different strategy: we will show that a.a.s. the number of vertices of L_1 with degree 1 is $\Omega(n^{\xi'})$ for some $\xi' > 0$. First, note that the expected number of vertices $v \in G$ having degree exactly 1, and having the only neighbor at radial distance at least $r/2$ is at least $(1 + o(1))n^{1-\xi'}$ for some $0 < \xi' < 1$: indeed, for each vertex, by our assumption on r , the probability to have such only neighbor is $e^{-c_1(r/2)^d} c_2 r^d e^{-c_3 r^d} = n^{-\xi'}$ for some dimension-dependent $c_1, c_2, c_3 > 0$ and $0 < \xi' < 1$, and the statement follows since a.a.s. there are $n(1 + o(1))$ vertices. Moreover, by a standard second moment method, there are a.a.s. $(1 + o(1))n^{1-\xi'}$ such vertices. The same holds when restricting to vertices at distance at least say r^2 (the value of r^2 is arbitrary, only to make sure that paths below remain inside Λ_n) from all boundaries of Λ_n , and we will assume this in the sequel.

We need to show that for each vertex there is at least a constant chance to be part of L_1 . Denote first by C_d be the smallest dimension-dependent integer such that any cube of a given tessellation has at most C_d adjacent cubes in the tessellation (that is, cubes sharing a lower-dimensional face). We will now show that every component different from L_1 , by our choice of r , a.a.s. has Euclidean diameter at most $O(r)$ in G , or equivalently, Euclidean diameter $O(1)$ on the renormalized lattice: indeed, It is a standard topological fact that in the tessellation of side length r/μ , such component has to be separated by a diagonally connected set of cells from L_1 , all being empty. The expected number of such components with more than L cells is at most $n(C_d e^{-V_d(r/\mu)^d})^L$. Plugging in our assumption on r and sufficiently large $L = L(d)$, we see that this expectation is $o(1)$, and hence, by Markov's inequality there is a.a.s. no such set. Therefore, a.a.s. every other component has Euclidean diameter at most Lr , and we condition on this as well.

Next, having explored the neighborhood of one vertex $v = v_0$ and having found exactly one vertex v_1 at Euclidean distance at least $r/2$, we see that the ball of radius r around v_1 that was not yet explored is of area at least $c''r^d$ for some $c'' > 0$. Assume w.l.o.g. that the x -coordinate of v_1 is at least $r/2$ larger than the one of v_0 . Conditional on this, the probability that v_1 has at least one neighbor with x -coordinate at least $r/2$ larger than the one of v_1 is again $1 - n^{-\xi''}$ for some $\xi'' > 0$, since this area had not been explored before and the unexplored area is $\Theta(\log n)$. Conditional under having found such a vertex, say v_2 , we proceed iteratively: having found v_i in this way, we look again for at least one neighbor v_{i+1} with x -coordinate at least $r/2$ larger than v_i . By the same argument, independently of the found vertices, the probability to find such a vertex is at least $1 - n^{-\xi''}$. Hence, with L from before, by a union bound over all steps, with probability at least $1 - L^2 n^{-\xi''} = 1 - o(1)$ we find at least L^2 consecutive vertices in the same component as v_0 . We advance at least $L^2 r/2 > Lr$ in the x -direction, and conditional under the statements before, v_0 must therefore belong to L_1 . Hence, with probability $1 - o(1)$, every vertex of degree 1, whose neighbor is at distance at least $r/2$, belongs to L_1 . Thus, the expected number of vertices of L_1 having degree 1 is at least $(1 - o(1))n^{1-\xi'}$. Since for every vertex the L^2 -neighborhood a.a.s. contains a polylogarithmic number of vertices, another second moment method shows that this also holds a.a.s.

To conclude, let V' then be the set of vertices of L_1 of degree exactly 1. By the above, a.a.s. $|V'| = \Omega(n^{1-\xi'})$ for some $0 < \xi' < 1$. For any different $x, y \in V'$, taking out the only edge incident to say x , we obtain an edge-cutset of size 1 between x and y , and thus, by Lemma 2.5,

$$\mathcal{R}(x \longleftrightarrow y, L_1(G)) \geq 1.$$

Since this holds for any different $x, y \in V'$, using Lemma 2.10, by Theorem 2.7, a.a.s. there exist some $\gamma_1, \gamma_2 > 0$, such that

$$\tau_{cov} \geq \delta_2 n r^d \log n \geq \delta_3 n \log^2 n,$$

and this case is finished as well, and by choosing $\delta = \min\{\delta_1, \delta_3\}$ the proof of the theorem is finished. \square

5. UPPER BOUND ON THE COVER TIME

To prove the upper bound, we first introduce a structure of random paths between good cubes of a 2-dimensional slice. This construction was first introduced by Abe [1]. Our first step will then be to show that for any vertices x, y belonging to corresponding good cubes C_i and C_j as well as to the largest components of these cubes, and therefore in particular belonging to $L_1(G)$, we can suitably bound from above the resistance between x and y . We thus focus in the next subsection on good cubes only. Throughout this section, we consider the tessellation of cubes C_i introduced in Section 3. In this section we will allow M to change slightly with n so that a cube C_i is either totally contained within Λ_n or shares at most some faces with Λ_n , and so Λ_n is completely covered by the tessellation. Given an integer M , we can find for each n , the closest value to it that satisfies this is at most $O(1/n)$ away, which will not play any role below.




FIGURE 3. A pending path in 2 dimensions with root at x (x is thus m -special for some m); all cubes surrounding the path are empty; y is the vertex associated to x that belongs to \mathcal{R} . Observe that a vertex in the path could be connected by an edge to more than 2 vertices (the maximum being 5 vertices).

5.1. Random path of good cubes. We assume r fixed for the rest of this subsection.

Given two self-avoiding paths \mathcal{L}_1 and \mathcal{L}_2 of cubes (with direction given by the ordering of the cubes in the path) and $C \in \mathcal{L}_1 \cap \mathcal{L}_2$, the concatenation of both paths at C is the path obtained from starting at \mathcal{L}_1 until hitting C and following \mathcal{L}_2 after that. Note that if C is the last cube of intersection between the two paths (seen from \mathcal{L}_2) the concatenation is self-avoiding.

In the same way we define the concatenation of two self-avoiding paths \mathcal{J}_1 and \mathcal{J}_2 of $G \in \mathcal{G}(n, r, d)$, given at a vertex x that belongs to both paths, as the path that results from starting at \mathcal{J}_1 until hitting x and after that continuing with \mathcal{J}_2 .

Recall that by Lemma 3.2, for α and M large enough, a.a.s. there are at least $\lfloor c \log n \rfloor$ long crossings of good cubes in each α -logarithmic strip (that is not in the boundary) of every 2-dimensional slice of Λ_n . In what follows we condition on this event.

Without loss of generality suppose in the following that \mathcal{K}_n^2 is the 2-dimensional slice made out of cubes C_i with only the first two coordinates not equal to 0. Divide \mathcal{K}_n^2 into disjoint strips of α -logarithmic size (see Figure 2). Recall that for a cube C_j , we denote by j_1, \dots, j_d its d coordinates. For $s \in \mathbb{Z}$, denote the s -th α -logarithmic horizontal strip by

$$H_s = \{C_j \in \mathcal{K}_n^2 : \lceil \alpha \log n \rceil - s \leq j_2 < s + \lceil \alpha \log n \rceil\},$$

and similarly denote the s -th α -logarithmic vertical strip by

$$V_s = \{C_j \in \mathcal{K}_n^2 : \lceil \alpha \log n \rceil - s \leq j_1 < s + \lceil \alpha \log n \rceil\}$$

Recall also that strips at the boundary of the slice might have a smaller width. Select $\lfloor c \log n \rfloor$ disjoint long crossings of good cubes for each strip H_s, V_s that is not on the boundary. We enumerate the vertical crossings of good cubes of the vertical strip V_s , counting from right to left and denote them by V_s^ℓ for $1 \leq \ell \leq \lfloor c \log n \rfloor$ (this way of enumerating the vertical crossings is going to be important later). Similarly, we enumerate the horizontal crossings of H_s , this time counting from bottom to top and denoting them by H_s^ℓ , for $1 \leq \ell \leq \lfloor c \log n \rfloor$.

For $r, s \in \mathbb{Z}$, denote by $T_{s,r}$ the smallest *parallelepiped* that contains all cubes C_i from $V_s \cap H_r$. Note that the parallelepiped $T_{s,r}$ that is not at the boundary of the 2-dimensional slice has Euclidean side length $Mr \lceil \alpha \log n \rceil$ in the first 2 dimensions and Mr in the others, as the side length of a cube C_i is Mr (we consider in the construction that follows only parallelepipeds that are not at the boundary, this will be sufficient for our purposes). Also note that any vertical crossing of good cubes V_s^ℓ of V_s and any horizontal crossing of good cubes H_r^ℓ of H_r intersect in at least one good cube C that lies inside of $T_{s,r}$, as the slice is a 2-dimensional object. Let now \mathcal{L} be a self-avoiding path of parallelepipeds T between T_{s_1, r_1} and T_{s_2, r_2} (that is, a sequence of adjacent

parallelepipeds of the same slice such that no parallelepiped appears more than once). We then construct an ordered set $(T_{a_1, b_1}, T_{a_2, b_2}, \dots, T_{a_k, b_k})$ of parallelepipeds that belong to \mathcal{L} and that we use to identify the change of its direction in the following way: set $T_{a_1, b_1} = T_{s_1, r_1}$. Let $T_{a_l, b_l} \in \mathcal{L}$, and suppose without loss of generality that the next step of the path \mathcal{L} is to the right. $T_{a_{l+1}, b_{l+1}}$ is then defined as the last parallelepiped of \mathcal{L} that is visited by going only to the right of T_{a_l, b_l} . The construction is analogous if the path \mathcal{L} after T_{a_l, b_l} goes to the left, up or down, respectively. We finish at $T_{a_k, b_k} = T_{s_2, r_2}$. If $(a_{i+1}, b_{i+1}) - (a_i, b_i) = me_1$ for some m , we say that \mathcal{L} has horizontal orientation at step (a_i, b_i) , and analogously, we say that it has vertical orientation if instead of e_1 we have e_2 . See Figure 4 for an example of the construction.




FIGURE 4. Path \mathcal{L} of parallelepipeds T . The path changes direction when visiting the parallelepipeds T_{a_i, b_i}

Assume that $r_1 = r_2$ for the parallelepipeds T_{s_1, r_1} and T_{s_2, r_2} , and assume that $s_2 > s_1$, so that T_{s_2, r_2} is to the right of T_{s_1, r_1} . We will also throughout this subsection require $s_2 - s_1$ to be odd, so that the parallelepipeds T_{s_1, r_1} and T_{s_2, r_2} are separated by an even number of parallelepipeds: we will later lift these restrictions. We now construct a self-avoiding random path \mathcal{L}_X (based on the straight path \mathcal{L}) of parallelepipeds between T_{s_1, r_1} and T_{s_2, r_2} : let X be a continuous random variable chosen uniformly in $[0, \frac{s_2 - s_1 + 1}{2}]$, and let $y_X \in \Lambda_n$ be the point that results from summing $(0, Mr[\alpha \log n]X, 0, \dots, 0)$ to the left bottom corner of $T_{r_1, \frac{s_2 + s_1 + 1}{2}}$ (the x -coordinate of this point is the midpoint of T_{s_1, r_1} and T_{s_2, r_2} , see Figure 5). Note that we assume here that y_X is inside of Λ_n ; if this were not the case we subtract $(0, Mr[\alpha \log n], 0, \dots, 0)$ from the midpoint of T_{s_1, r_1} and T_{s_2, r_2} , and the construction that follows is reflected vertically.

We denote by \mathcal{L}_X the a.s. unique random path of parallelepipeds intersecting the segment between the bottom left corner of T_{s_1, r_1} and the point y_X , and the segment between the bottom right corner of T_{s_2, r_2} and y_X (see Figure 6). Let $C_i \in T_{s_1, r_1}$ and $C_j \in T_{s_2, r_2}$ be any good cubes such that both of them belong to some vertical crossings of the strips V_{s_1} and V_{s_2} , respectively (see the leftmost part of Figure 7 for an example of such crossings for T_{s_1, r_1}). Either such crossings have length $O(\log n)$, or otherwise, by Theorem 2.16, a.a.s. we may select crossings with this property. We denote the selected vertical crossings by $V(C_i)$ and $V(C_j)$. From the vertical crossing of good cubes $V(C_i)$ that contains C_i we then extract a bottom-to-top crossing of T_{s_1, r_1} made from good cubes which we denote by $V'(C_i)$ (see yellow cubes in the middle picture of Figure 7). Similarly, we extract a bottom-to-top-crossing of T_{s_2, r_2} made from good cubes, from the vertical crossing $V(C_j)$ that contains C_j , and we denote it by $V'(C_j)$. Given $1 \leq \ell \leq \lfloor c \log n \rfloor$, we construct a self-avoiding path of good cubes between C_i and C_j that is entirely contained within \mathcal{L}_X and has index ℓ .

Since $V'(C_i)$ is self-avoiding, we may split it into two paths of good cubes with intersection only at C_i . The paths in question are the ones originated from starting at C_i and going up or down




FIGURE 5. Position of y_X relative to the parallelepipeds




FIGURE 6. The red cubes form the path \mathcal{L}_X

following $V'(C_i)$, see the rightmost picture of Figure 7. Denote the paths by $V'_1(C_i)$ and $V'_2(C_i)$ and do the same for $V'(C_j)$. At least one of these paths of good cubes that start at C_i intersects $H_{r_1}^\ell$ as the bottom-to-top crossing (of T_{s_1, r_1}) $V'(C_i)$ made from good cubes intersects the horizontal crossing of good cubes (of the strip H_{r_1}) $H_{r_1}^\ell$ in at least one good cube (this is true as the slice is 2-dimensional). Suppose without loss of generality that $V'_1(C_i)$ is intersecting this horizontal crossing. Concatenate then $V'_1(C_i)$ with $H_{r_1}^\ell$ at the last good cube of $H_{r_1}^\ell$ that intersects $V'_1(C_i)$; we denote the resulting path by \mathcal{Q}_1 and note that it is non-intersecting (see Figure 8, the bottom picture shows the final \mathcal{Q}_1). Now, observe that $(a_3, b_3) - (a_2, b_2)$ must be vertically oriented. We concatenate \mathcal{Q}_1 to the vertical crossing $V_{a_3}^\ell$ (seen as path that goes from bottom to top) at the last cube of $V_{a_3}^\ell$ that intersects \mathcal{Q}_1 ; we denote this path by \mathcal{Q}_2 .




FIGURE 7. The path of smaller cubes is $V(C_i)$. The path in yellow is $V'(C_i)$. In green and blue the two paths $V'_1(C_i)$ and $V'_2(C_i)$, respectively, obtained from $V'(C_i)$ that start from C_i .

FIGURE 8. The gray path is $H_{r_1}^\ell$. \mathcal{Q}_1 is the path that is left after concatenation in the picture below, adding the green and gray parts.

We continue in this way until we cross the midpoint (that is, we cross y_X). Here we continue interpreting the horizontal crossings as paths from left to right but change the orientation of the




FIGURE 9. The red path is $\mathcal{L}_X^{\ell_1}$, and the blue path is $\mathcal{L}_X^{\ell_2}$, they are disjoint as a consequence of the enumeration introduced for the long crossings. The green part (subset of $V'(C_i) \cup V'(C_j)$) allows us to reach C_i and C_j .

vertical crossings of good cubes to be from top to bottom. Furthermore, now instead of taking the ℓ -th vertical crossing of good cubes we take the $(\lceil c \log n \rceil - \ell)$ -th vertical crossing of good cubes to create the concatenations (the way of enumerating here is different so that that the concatenations are disjoint for different ℓ if we delete the good cubes from $V'(C_i) \cup V'(C_j)$, see Figure 9). To finish the construction, once we arrive at T_{s_2, r_2} , what is left is the concatenation to the vertical crossing of good cubes $V'(C_j)$. Here we proceed just as we did for C_i . Select one path that starts from C_j and is made from good cubes of $V'(C_j)$ and intersects \mathcal{Q}_{k-1} (the last concatenation available and that passes through T_{s_2, r_2} ; there exists at least one such path of good cubes). Concatenate \mathcal{Q}_{k-1} to the last path selected to obtain \mathcal{Q}_k . Note that the resulting path is self-avoiding by construction, and also note that for different ℓ , the paths are disjoint if we delete the initial and final good cubes that come from $V'(C_i)$ and $V'(C_j)$. We denote \mathcal{Q}_k by \mathcal{Q}_X^ℓ . Delete from \mathcal{Q}_k the good cubes that belong to $V'(C_i)$ and $V'(C_j)$, except for the cubes where the concatenations with $V'(C_i)$ and $V'(C_j)$ were made at the beginning and the end, and denote the resulting random path by \mathcal{L}_X^ℓ (see Figure 9). In particular, note that the path \mathcal{L}_X^ℓ is a sub-path of \mathcal{Q}_X^ℓ since the latter has C_i and C_j as endpoints (in Figure 9, \mathcal{Q}_X^ℓ corresponds to the blue path together with the green paths in Figure 9 while the path \mathcal{L}_X^ℓ is only the blue path). We point out that the construction of \mathcal{Q}_X^ℓ will be used only in the proof of the upper bound of the cover time above the connectivity threshold in Section 6.

5.2. Proof of the upper bound on the cover time with fixed radius.

Theorem 5.1. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$, and let $\varepsilon > 0$ be arbitrarily small. Assume that $r > (1 + \varepsilon)r_g$ and also assume $r = O(1)$. A.a.s., there is a constant $\gamma(d, r) > 0$ such that*

$$\tau_{\text{cov}}(L_1) \leq \gamma n \log^2 n.$$

Proof. Since we use the construction given in Subsection 5.1, we recall our initial assumptions made on the good cubes C_i and C_j : both cubes belong to the same strip and they both belong to parallelepipeds not at the boundary of the slice and separated by an even number of parallelepipeds. We will lift these assumptions later.

Consider any two vertices x, y that belong to the good cubes C_i and C_j respectively, and both belong to the largest connected component of the renormalized graph of good cubes. As the good cubes C_i and C_j belong to long crossings of good cubes (in the renormalized graph) which at the

same time contain long crossings of G of length $\Theta(n^{1/d})$, by Theorem 2.13 x, y then also belong to $L_1(G)$ a.a.s.

Select an induced path of $G \in \mathcal{G}(n, r, d)$ that starts at x , is contained in $V'_1(C_i)$ and has its final vertex at the last cube of $V'_1(C_i)$. Denote this path by \mathcal{V}_x^1 (note that this selection does not depend on the random variable X). We do the same with $V'_2(C_j)$ and denote the path by \mathcal{V}_y^2 . We proceed in the same way for the vertex y .

Without loss of generality suppose that $V'_1(C_i)$ intersects \mathcal{L}_X^ℓ . Denote by C^1 the only cube of \mathcal{L}_X^ℓ that intersects $V'_1(C_i)$ (note that conditioned on X the selection of C^1 depends on ℓ). Also suppose that $V'_1(C_j)$ intersects \mathcal{L}_X^ℓ in some good cube, and denote by C^2 the first good cube of \mathcal{L}_X^ℓ that intersects $V'_1(C_j)$.

As \mathcal{L}_X^ℓ is a path of good cubes that begins at C^1 and ends at C^2 , we may obtain an induced path (totally contained within \mathcal{L}_X^ℓ) of $G \in \mathcal{G}(n, r, d)$ that starts at some vertex of \mathcal{V}_x^1 inside C^1 and finishes at some vertex of \mathcal{V}_y^1 inside C^2 (all cubes are good, and so we can use the giant components of the good cubes), and that intersects \mathcal{V}_x^1 and \mathcal{V}_y^1 only at its initial and final point, respectively. We denote this path by J_X^ℓ , see Figure 10. We concatenate \mathcal{V}_x^1 , J_X^ℓ and \mathcal{V}_y^1 in the obvious way (as Figure 10 suggests for the first part of the concatenation of blue and red). From this path, we may then extract an induced path with the same endpoints.




FIGURE 10. \mathcal{V}_x^1 and J_X^ℓ have exactly one vertex in common. We use this vertex to concatenate both paths.

We collect some properties of \mathcal{J}_X^ℓ :

- Let $\ell_1 \neq \ell_2$. The set of vertices of $\mathcal{J}_X^{\ell_1}$ that are within good cubes that belong to $\mathcal{L}_X^{\ell_1}$ is disjoint from the set of vertices of $\mathcal{J}_X^{\ell_2}$ that are within the good cubes that belong to $\mathcal{L}_X^{\ell_2}$. If $\mathcal{J}_X^{\ell_1}$ and $\mathcal{J}_X^{\ell_2}$ have common vertices, these are inside of $V'(C_i)$ or $V'(C_j)$.
- The number of edges that have some endpoint inside of $V'(C_i) \cup V'(C_j)$ and belong to some path \mathcal{J}_X^ℓ is $\Theta(\log n)$.

For point (a) recall that if $\ell_1 \neq \ell_2$, then $\mathcal{L}_X^{\ell_1}$ does not intersect $\mathcal{L}_X^{\ell_2}$, and as \mathcal{J}_X^ℓ is totally contained within $\mathcal{L}_X^\ell \cup V'(C_i) \cup V'(C_j)$, part (a) follows.

For (b), recall that $V'(C_i)$ has logarithmic length (in the renormalized lattice where we see cubes as points of \mathbb{Z}^d). Since the path \mathcal{V}_x^1 is chosen to be an induced path it has a logarithmic number

of edges. Note that the other edges of \mathcal{J}_X^ℓ with at least one endpoint that is inside of $V'(C_i)$ must have this endpoint inside of C^1 . Since the path is induced and the cube C^1 has constant size, the number of edges from some realization \mathcal{J}_X^ℓ and with at least one endpoint that is inside of C^1 is $O(1)$. Finally, since there is a logarithmic number of such cubes C^1 (one for each ℓ) the number of such vertices is logarithmic. The same is true for $V'(C_j)$ and we obtain property (b).

Now, let Y be a discrete uniform random variable with values in $\{1, \dots, \lfloor c \log n \rfloor\}$ and independent from X . Define the random path \mathcal{J}_X^Y and the function

$$\psi(u, v) = \begin{cases} 1, & \text{if } \{u, v\} \in \mathcal{J}_X^Y \text{ and the path visits } u \text{ first,} \\ -1, & \text{if } \{u, v\} \in \mathcal{J}_X^Y \text{ and the path visits } v \text{ first,} \\ 0, & \text{otherwise.} \end{cases}$$

We define a unit flow θ between x and y by

$$\theta(u, v) = \mathbb{E}(\psi(u, v))$$

for $u, v \in G$, where the expectation is with respect to X, Y . Note that

$$|\theta(u, v)| \leq \mathbb{P}_{X, Y}(\{u, v\} \in \mathcal{J}_X^Y) =: p(u, v)$$

We next group the parallelepipeds T that contain some realization of \mathcal{J}_X^Y according to their level t as follows (see Figure 11):

$$D_t = \begin{cases} \{T_{i,j} : i = s_1 + t, r_1 \leq j \leq r_1 + t\}, & \text{if } 0 \leq t \leq \frac{s_2 - s_1 + 1}{2}, \\ \{T_{i,j} : i = s_1 + t, r_1 \leq j \leq r_1 + s_2 - s_1 - t - 1\}, & \text{if } \frac{s_2 - s_1 + 1}{2} + 1 \leq t \leq s_2 - s_1 + 1. \end{cases}$$

We are now ready to bound the effective resistance between x, y . Recall that $\tau(e) = 1$, and so,




FIGURE 11. Parallelepipeds at level t (i.e., their projections onto the first 2 dimensions). See the projected probability over the right line of D_t inside the blue dotted lines.




FIGURE 12. Using triangle inequality for vertices in different strips and in the same 2-dimensional slice.

since $\theta^2(u, v) \leq p^2(u, v)$, by Definition 2.1, in order to bound $\mathcal{R}(x \longleftrightarrow y)$, it suffices to bound $\sum_{e \in E} p^2(e)$. More precisely, we may split the edge set according to levels of parallelepipeds and get:

$$\mathcal{R}(x \longleftrightarrow y) \leq \sum_{u \in V'(C_i) \cup V'(C_j)} \sum_{\substack{v \in L_1 \\ v \sim u}} p^2(u, v) + \sum_t \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(C_i) \cup V'(C_j)}} \sum_{\substack{v \in L_1 \\ v \sim u}} p^2(u, v). \quad (5.1)$$

The first double sum on the right hand side can be bounded by $O(\log n)$ as the number of edges that have positive probability of belonging to the path is $\Theta(\log n)$ (see properties of \mathcal{J}_X^ℓ above).

Consider then the remaining term. Let $u \in T_{i,j}$ with $T_{i,j} \in D_t$ and $\{u, v\} \in E$ and $u \notin V'(C_i) \cup V'(C_j)$. We now claim that the probability that $T_{i,j} \in D_t$, with $1 \leq t \leq \frac{s_2-s_1+1}{2}$, is part of the path of parallelepipeds \mathcal{L}_X is bounded above by $\frac{c'}{t}$ for some constant c' that can be chosen independent of n and t (for $t = 0$ we bound this probability by 1). To show this, denote by y_X^t be the intersection of the segment between the left bottom corner point of T_{s_1, r_1} and y_X and the left line of level t (see Figure 11). By similarity of triangles, y_X^t is uniform over the left line of level t (see for example the triangles $Ay_X^t B$ and $Ay_X C$ in Figure 11). Now, observe that in order for $T_{i,j} \in D_t$ to be part of \mathcal{L}_X the random point y_X^t has to belong to the left face of $T_{i,j}$ or to $T_{i,j-1}$ (the left face of the parallelepiped immediately below, in case such parallelepiped exists): indeed, if y_X^t were above $T_{i,j}$, then the parallelepiped does not belong to the path, and if y_X^t were below $T_{i,j-1}$, as the angle of the red line is at most $\pi/4$, the red line in Figure 11 does not intersect $T_{i,j}$. Therefore, the probability that $T_{i,j} \in D_t$ belongs to \mathcal{L}_X is bounded from above by the probability that y_X^t belongs to the left faces of two cubes at level t . Since y_X^t is uniform in its range, then this probability is at most $\frac{2}{t}$. By symmetry, the same is true for $\frac{s_1-s_1+1}{2} + 1 \leq t \leq s_2 - s_1$ (for $t = s_2 - s_1 + 1$ we bound this probability by 1); in this case we obtain an upper bound of $\frac{2}{s_2-s_1-t-1}$ for the probability that $T_{i,j} \in D_t$.

Consider a fixed realization of \mathcal{L}_X , and suppose first that $\{u, v\} \in E$ such that u belongs to some parallelepiped $T_{i,j}$ of \mathcal{L}_X . We may also assume that the vertex u does not belong to the good cubes of $V'(C_i) \cup V'(C_j)$ (otherwise the contribution is already counted in the first double sum). Then, recall that $\{u, v\}$ belongs to at most one path of the form \mathcal{J}_X^Y (by property (a) of the path \mathcal{J}_X^Y above). By independence of X, Y and as $\{u, v\}$ belongs to one unique path \mathcal{J}_X^Y (with X fixed), and as the path is chosen uniformly from a set of $\lfloor c \log n \rfloor$ possible realizations, we have

$$\mathbb{P}_{X,Y}(\{u, v\} \in \mathcal{J}_X^Y | X) \leq \frac{1}{\lfloor c \log n \rfloor}.$$

Furthermore, recall that u belongs to the parallelepiped of $T_{i,j} \in D_t$. Thus, by the above, since the choice of the parallelepiped in D_t (with $1 \leq t \leq \frac{s_2-s_1+1}{2}$) and the choice of the crossing are independent, $p(u, v) \leq \frac{c''}{t \log n}$ for some $c'' > 0$. The same occurs by symmetry for the other levels.

Fix then a parallelepiped T that belongs to some level t . Recall that the volume of the parallelepiped T is $\Theta(\log^2 n)$, as in 2 dimensions its side length is $Mr \lceil \alpha \log n \rceil$ and it is Mr in the other dimensions (recall that r is assumed to be constant in this section). Hence, the number of edges that have at least one endpoint inside of T is by Corollary 2.9 a.a.s. $O(\log^2 n)$. We substitute these observations into the right hand expression of (5.1) and obtain

$$\begin{aligned} \sum_{1 \leq t \leq \frac{s_2-s_1+1}{2}} \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(C_i) \cup V'(C_j)}} \sum_{v \in L_1} p^2(u, v) &\leq \sum_{1 \leq t \leq \frac{s_2-s_1+1}{2}} \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(C_i) \cup V'(C_j)}} \sum_{v \in L_1} \frac{c'''}{t^2 \log^2 n} \\ &\leq \sum_{1 \leq t \leq \frac{s_2-s_1+1}{2}} \sum_{T_{i,j} \in D_t} \log^2 n \frac{c'''}{t^2 \log^2 n} \\ &\leq \sum_{1 \leq t \leq \frac{s_2-s_1+1}{2}} (t+1) \log^2 n \frac{c'''}{t^2 \log^2 n} \\ &= \sum_{1 \leq t \leq \frac{s_2-s_1+1}{2}} O\left(\frac{1}{t}\right) = O(\log n). \end{aligned}$$

It is clear that the contribution of D_0 to the inequality is $O(1)$. By symmetry around y_X , one obtains the same bound for the remaining levels $\frac{s_2-s_1+1}{2} < t \leq s_2 - s_1 + 1$. Then $\mathcal{R}(x \longleftrightarrow y) = O(\log n)$.

We now lift the assumptions on the parallelepipeds: consider first two vertices x, y of L_1 that are in the same strip of a 2-dimensional slice, but now separated by an odd number of parallelepipeds that are not at the boundary. Let $x \in T_{s'_1, r_1}$ and $y \in T_{s'_2, r_1}$ with $s'_2 - s'_1$ even such that neither

parallelepiped is at the boundary. By the construction given above there exists $x' \in L_1(G)$ with $x' \in T_{s'_1+1, r_1}$ such that $\mathcal{R}(x' \longleftrightarrow y) = O(\log n)$. Note that $d_E(x, x') = O(\log n)$, since x and x' are in adjacent parallelepipeds, and any parallelepiped has logarithmic side length in the first 2 dimensions and fixed side length in the other dimensions. By Lemma 2.12, a.a.s., the graph distance between x and x' is then also $O(\log n)$, and thus a.a.s. $\mathcal{R}(x \longleftrightarrow x') = O(\log n)$. Then, by application of the triangle inequality (see Lemma 2.4), also a.a.s. $\mathcal{R}(x \longleftrightarrow y) = O(\log n)$. Also, since any two vertices in the same parallelepiped (not necessarily in good cubes, but still in $L_1(G)$) satisfy that their graph distance is $O(\log n)$ (see Lemma 2.12, the largest side length of a parallelepiped is $O(\log n)$), by another application of the triangle inequality, also a.a.s. the effective resistance between them is $O(\log n)$. Hence, for any $x', y' \in L_1(G)$ belonging to the largest connected component of the renormalized graph inside the same strip of a 2-dimensional slice not at the boundary, we have

$$\mathcal{R}(x \longleftrightarrow y) = O(\log n).$$

By the same idea, we may extend our bounds to vertices in parallelepipeds of a strip on the boundary of the 2-dimensional slice, as for these vertices there exists a vertex at graph distance $O(\log n)$ in a parallelepiped not at the boundary.

We now extend the result to any two vertices x, y of $L_1(G)$ in the same 2-dimensional slice: to do so, suppose that $x, y \in \mathcal{K}_n^2$. Let $x \in H_r$ (x is in the r -th horizontal strip) and let $y \in V_s$ (y in the s -th vertical strip). Note that this election is always possible as any vertex belongs to one vertical strip and one horizontal strip. Pick $z \in L_1$ such that $z \in V_s \cap H_r = T_{s,r}$ (see Figure 12). Note that such z must exist as $T_{s,r}$ is the intersection of two strips that contain long crossings of good cubes, which at the same time contain long crossings of L_1 . As x and y are in the same strip, by the above, a.a.s. $\mathcal{R}(x \longleftrightarrow z) = O(\log n)$. Similarly, a.a.s. $\mathcal{R}(y \longleftrightarrow z) = O(\log n)$, and applying again the triangle inequality the result holds as well.

Now, in general, suppose that there is no 2-dimensional slice containing x, y (while always a plane containing x, y exists, we cannot ensure the existence of a slice, which by our definition is assumed to be axis-parallel). Note that in case two different slices have non-empty intersection, this intersection is a line of cubes such that only one coordinate is not fixed. In this case, as before, there exists at least a vertex $z \in G \in \mathcal{G}(n, r, d)$ that belongs to L_1 and that is in the intersection of both slices. Then, for any two vertices x', y' such that x' is in the first slice and y' is in the second slice, by the above, a.a.s. $\mathcal{R}(x' \longleftrightarrow z) = O(\log n)$ and $\mathcal{R}(y' \longleftrightarrow z) = O(\log n)$, and thus, by the triangle inequality, also $\mathcal{R}(x' \longleftrightarrow y') = O(\log n)$. To conclude, observe that for any two vertices $x, y \in G$, there exists a sequence of slices $(\mathcal{K}_n^2)_1, \dots, (\mathcal{K}_n^2)_s$, with $s \leq d-1$, such that for any $1 \leq i \leq s-1$, $(\mathcal{K}_n^2)_s$ has non-empty intersection with $(\mathcal{K}_n^2)_{s+1}$, and moreover, $x \in (\mathcal{K}_n^2)_1$, and $y \in (\mathcal{K}_n^2)_s$. By applying the triangle inequality iteratively, we then have a.a.s. $\mathcal{R}(x \longleftrightarrow y) = O(\log n)$.

By this result, together with Lemma 2.10 we then immediately conclude that a.a.s.

$$\tau_{cov} \leq \gamma n \log^2 n,$$

for some $\gamma(d, r) > 0$, and the theorem follows. \square

We now extend this result to larger radii, but still (arbitrarily) smaller than the connectivity threshold:

Theorem 5.2. *Let $d \geq 2$, let $G \in \mathcal{G}(n, r, d)$, and let ε be an arbitrarily small positive constant. Assume furthermore $r_g < r \leq (1 - \varepsilon)r_c$. Then, a.a.s.,*

$$\tau_{cov}(L_1(G)) \leq \delta' n \log^2 n$$

where δ' is a constant that only depends on ε .

Proof. Given any tessellation, let C_d be the dimension-dependent constant so that every d -dimensional cube is adjacent (sharing any lower-dimensional face) to at most C_d other cubes, and let $r_0 = r_0(d)$ be the smallest radius r so that $C_d e^{-V_d r^d} \leq 0.99$. Note that $r_0 = \Theta(1)$. In the

proof of Theorem 5.1 we showed that the effective resistance between any two vertices of the giant component of $G \in \mathcal{G}(n, r, d)$ is a.a.s. $O(\log n)$ for every $r \leq r_0$. Consider thus $r > r_0$. Observe that the graph $G \in \mathcal{G}(n, r_0, d)$ can be naturally coupled with the graph $G' \in \mathcal{G}(n, r, d)$ such that the giant of G is a subgraph of the giant of G' . Hence, by Theorem 2.3, for vertices x, y in both giants, the effective resistance between any two such vertices is also in G' a.a.s. bounded by $O(\log n)$.

By the triangle inequality for the effective resistance (Lemma 2.4), it thus remains to show that a.a.s. for any vertex x that is in the giant of G' but not in the giant of G , there exists x' in the giant of G , such that $d_G(x, x') = O(\log n)$.

To see this, note that in G , there is a.a.s. no diagonally connected set of cells of side length r/μ with integer left bottom corner inside Λ_n of size $C \log n$ for C large enough, with the property that all its cells are empty: indeed, letting $c := C_d e^{-V_d r_g^d} \leq 0.99$, the expected number of such sets is at most $nc^{C \log n}$, since there are at most n choices for the starting cell, at most C_d choices for continuing at each step, and $e^{-V_d r_g^d}$ is the probability that the selected cell is indeed empty. Note that for $C = C(c)$ large enough, $nc^{C \log n} = o(1)$, and hence, by Markov's inequality, a.a.s. such a set does not exist. Since every connected component different from the giant in G has to have an empty diagonally connected outer boundary (if needed, we restrict to the outer boundary's largest connected piece inside Λ_n , which is still of the same logarithmic order, only by a constant only depending on the dimension smaller than the total boundary), each such component thus has to have Euclidean boundary at most $O(\log n)$, and thus, for every x' in the giant of G' , there must exist a vertex x in the giant of G , such that $d_E(x, x') = O(\log n)$. Then, by Corollary 2.12, a.a.s. also $d_G(x, x') = O(\log n)$, and the theorem follows. \square

Proof of Theorem 1.1. By combining Theorem 4.2 and Theorem 5.2, Theorem 1.1 follows immediately. \square

6. UPPER BOUND RIGHT ABOVE THE THRESHOLD FOR CONNECTIVITY

Perhaps surprisingly, for $\mathcal{G}(n, r, 2)$, no upper bound of $O(n \log n)$ on the cover time of the simple random walk had been known right above the connectivity threshold. For $d \geq 3$, in the already mentioned paper of Cooper and Frieze [12] the result was obtained, but their result does not apply for $d = 2$. As mentioned in the introduction, for $d = 2$, in [5] the authors found the asymptotic order of the cover time only for a radius that was by a constant factor larger than the connectivity threshold. In this section, we close the gap left by the papers for $d = 2$. We note that it is enough to prove the upper bound, as the lower bound is immediate by the general lower bound on cover times for connected graphs by Feige [19]. Our main result will therefore be the following theorem (since our result is only new for 2 dimensions, we state it only for 2 dimensions, although it holds more generally):

Theorem 6.1. *Let $\varepsilon > 0$ arbitrarily small and consider $G \in \mathcal{G}(n, r, 2)$. Suppose that $r^2 = (1 + \varepsilon) \frac{\log n}{\pi}$. Then, a.a.s, for any two vertices $x, y \in G \in \mathcal{G}(n, r, 2)$,*

$$\mathcal{R}(x \longleftrightarrow y) = O\left(\frac{1}{\log n}\right).$$

Proof of Theorem 1.2 assuming Theorem 6.1: By Lemma 2.10, a.a.s., $|E| = \Theta(nr^d) = \Theta(n \log n)$. Theorem 1.2 then follows immediately by Theorem 2.6. \square

It thus remains to prove Theorem 6.1. The approach of the proof is similar to the one in Section 5: first we prove that the upper bound on the effective resistance is true for any pair of vertices inside a *backbone*, and in a second step we then prove that the resistance between any two vertices is of the desired order. We now proceed to the details:

Proof of Theorem 6.1: Recall the definition of μ introduced in the tessellation of cubes (Q_i) in Section 4 of cubes of side length r/μ (in fact cubes here are cells, but we keep the notation of the

previous section). In this section, we choose $\mu(n) = \mu + O(1/n)$ to be the smallest real number larger or equal to μ such that any cube Q is either completely contained in Λ_n or it shares at most some faces, so that we avoid boundary issues. With abuse of notation, we denote below $\mu(n) = \mu$. Let $\beta = \beta(\varepsilon)$ be a constant chosen sufficiently small (to be fixed below). We say that Q_i is *dense* if it has at least $\beta \log n$ vertices of $G \in \mathcal{G}(n, r, 2)$, otherwise we say that it is *non-dense*.

See cubes Q as sites of \mathbb{Z}^2 . The probability that a cube is non-dense is at most $n^{-\kappa}$ for some $\kappa = \kappa(\beta) > 0$, since the cube Q has area $\Theta(\log n)$. Considering dense cubes as occupied sites of a Bernoulli site percolation model, we are thus in the super-critical phase of Bernoulli site percolation, i.e., there exists a giant component corresponding to dense cubes. Moreover, as before, it is a standard topological fact that every (maximal) connected component of cubes, which might contain dense and non-dense cubes and that has empty intersection with the giant component of dense cubes, has a diagonally connected outer boundary of non-dense cubes. As there are at most 8^L diagonally connected paths of length L when the starting vertex is fixed, the probability that there exists a diagonally connected path of non-dense cubes of length greater or equal than L is, by a union bound over all starting vertices, at most

$$n^{1-L\beta}8^L,$$

which tends to 0 for $L = L(\beta)$ sufficiently large. Therefore, since every connected component (in the renormalized lattice of cubes Q) of area at least CL^2 (for C large enough), by a standard isoperimetric inequality has to have an outer boundary of length at least L , a.a.s. every component different from the giant of the associated Bernoulli site percolation has at most constant size CL^2 , and Euclidean diameter at most L . In other words, for every site, at distance L one finds a site belonging to the giant component (of the renormalized lattice), and for any vertex of the underlying random geometric graph, at Euclidean distance at most $Lr/\sqrt{5}$ we find a vertex y belonging to a cube of the giant component of the Bernoulli site percolation.

Step 1: Upper bound on the effective resistance between two vertices inside the giant component of dense cubes

We first prove that the effective resistance between any two vertices inside the giant component of dense cubes is $O\left(\frac{1}{\log n}\right)$. Divide Λ_n into disjoint α -logarithmic strips of cubes Q_i . We use the structure introduced in Subsection 5.1, with the difference that we consider cubes Q_i instead of cubes C_i and dense cubes playing the role of good cubes.

Let \mathcal{Q}_X^Y (here Y is a uniform discrete random variable independent from X with values in $\{1, \dots, \lfloor c \log n \rfloor\}$, see the construction of Subsection 5.1 for the definition of c and Theorem 5.1 for details of the construction of \mathcal{Q}_X^ℓ with a fixed value ℓ of Y) be the random path of dense cubes between two fixed dense cubes Q_i and Q_j that both belong to the giant component of dense cubes, and as in Subsection 5.1, we may assume that the picked dense cubes belong to some parallelepipeds $Q_i \in \mathbb{T}_{s_1, r_1}$ and $Q_j \in \mathbb{T}_{s_2, r_2}$ such that $r_1 = r_2$, $s_2 > s_1$, $s_2 - s_1$ is odd, and that both parallelepipeds are not at the boundary of the 2-dimensional slice that contains them. Pick vertices x, y of $G \in \mathcal{G}(n, r, 2)$ such that $x \in Q_i$ and $y \in Q_j$. For each dense cube that is part of some realization of \mathcal{Q}_X^Y select $\lfloor \beta \log n \rfloor$ vertices, except for Q_i and Q_j where we may choose $\lfloor \beta \log n \rfloor - 1$ vertices distinct from x, y (or $\lfloor \beta \log n \rfloor - 2$ in case $Q_i = Q_j$).

Conditional on X and Y , we construct a random path of $G \in \mathcal{G}(n, r, 2)$ that starts at x , finishes at y , and is entirely contained within the path of dense cubes \mathcal{Q}_X^Y as follows. First, select uniformly at random and independently from X, Y one of the $\lfloor \beta \log n \rfloor - 1$ vertices that belong to the dense cube Q_i (one out of the $\lfloor \beta \log n \rfloor - 1$ vertices previously selected), and connect it by an edge to the vertex x : this is the first edge of our path. For subsequent steps we do something similar: suppose that the last vertex that we chose belongs to the dense cube Q_s . We then choose uniformly and independently at random from the $\lfloor \beta \log n \rfloor$ vertices that belong to the dense cube that follows in the path \mathcal{L}_X^Y . We proceed until the last selected vertex belongs to the cube just before Q_j (the dense cube that contains y). In this case we select uniformly at random between the $\lfloor \beta \log n \rfloor - 1$ selected vertices in Q_j and finally connect all these last vertices by an edge to y (if we had $Q_i = Q_j$, then we would pick $\lfloor \beta \log n \rfloor - 2$ vertices inside the cube and form paths of length 2 from x to y

through these vertices). This process gives rise to $\lfloor \beta \log n \rfloor - 2$ internally vertex-disjoint random paths of $G \in \mathcal{G}(n, r, 2)$ with initial vertex x and final vertex y . We denote this random path of $G \in \mathcal{G}(n, r, 2)$ by $\mathcal{J}_{X,Z}^Y$ and denote its law by $\mathbb{P}_{X,Y,Z}$, where Z denotes the randomness that comes from the path selection conditioned on X, Y . Observe that the number of edges that have some endpoint inside of $V'(Q_i) \cup V'(Q_j)$ is $\Theta(\log^3 n)$, since $V'(Q_i)$ (see also Subsection 5.1) is made of $O(\log n)$ dense cubes, each of which has $O(\log n)$ selected vertices and between any two of such dense cubes there are $O(\log^2 n)$ edges. Since between any two consecutive dense cubes there are a.a.s. $\Theta(\log^2 n)$ edges, and we choose uniformly at random one of them, we obtain in the case $u \neq x, u \neq y, v \neq x, v \neq y$,

$$\mathbb{P}_{X,Y,Z}(\{u, v\} \in \mathcal{J}_Z^Y | X, Y) = O\left(\frac{1}{\log^2 n}\right),$$

whereas in the case $|\{u, v\} \cap \{x, y\}| = 1$, we have

$$\mathbb{P}_{X,Y,Z}(\{u, v\} \in \mathcal{J}_{X,Z}^Y | X, Y) = O\left(\frac{1}{\log n}\right).$$

As before, denote by $p(u, v)$ the probability that the edge $\{u, v\}$ belongs to the random path $\mathcal{J}_{X,Z}^Y$. Using the random path $\mathcal{J}_{X,Y}^Y$, we define a flow in the same way as in the proof of Theorem 5.1, and we obtain

$$\begin{aligned} \mathcal{R}(x \longleftrightarrow y) &\leq \sum_{\substack{v \in L_1 \\ y \sim v}} p^2(y, v) + \sum_{\substack{v \in L_1 \\ x \sim v}} p^2(x, v) \\ &\quad + \sum_{\substack{u \in V'(Q_i) \cup V'(Q_j) \\ u \neq x, u \neq y}} \sum_{\substack{v \in L_1 \\ v \neq x, v \neq y \\ u \sim v}} p^2(u, v) \\ &\quad + \sum_t \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(Q_i) \cup V'(Q_j)}} \sum_{\substack{v \in L_1 \\ u \sim v}} p^2(u, v). \end{aligned}$$

To bound the first two sums, since $p(x, v) = O\left(\frac{1}{\log n}\right)$, and since x and y send flow to different $O(\log n)$ vertices, we have $\sum_{v \in L_1} (p^2(y, v) + p^2(x, v)) = O(1/\log n)$. For the sum in the second row, since the number of edges with at least one endpoint inside of $V'(Q_i) \cup V'(Q_j)$ is $O(\log^3 n)$, and since $p(u, v) = O(1/\log^2 n)$,

$$\sum_{\substack{u \in V'(Q_i) \cup V'(Q_j) \\ u \neq x, u \neq y}} \sum_{\substack{v \in L_1 \\ v \neq x, v \neq y \\ u \sim v}} p^2(u, v) = O\left(\frac{1}{\log n}\right).$$

To bound the term in the last row, recall that, as in the proof of Theorem 5.1 in Section 5, the probability that a dense cube inside of some parallelepiped $T \in D_t$ belongs to \mathcal{Q}_X^Y is $O\left(\frac{1}{t \log n}\right)$ for $1 \leq t \leq \frac{s_2 - s_1 + 1}{2}$ and $O\left(\frac{1}{\log n}\right)$ for $t = 0$, since there are $\Omega(t)$ parallelepipeds at level t , and we choose one of the $\Omega(\log n)$ crossings of dense cubes, uniformly at random. Now, observe that we have the additional randomness coming from the fact for any pair of consecutive dense cubes, we choose u.a.r. a vertex inside each cube, giving another factor $O(1/\log^2 n)$, as mentioned above. All three choices are independent of each other, and thus, for edges $\{u, v\}$ appearing in the last term, we have

$$p(u, v) = O\left(\frac{1}{t \log^3 n}\right).$$

The number of edges with positive probability p and with at least one endpoint inside of a parallelepiped T is a.a.s. $O(\log^4 n)$, as there are $O(\log^2 n)$ dense cubes inside T (the volume of T is $\Theta(\log^3 n)$, and each cube has area $\Theta(\log n)$), each dense cube contains $O(\log n)$ vertices, and the

selected vertices are connected by an edge to another $O(\log n)$ selected vertices by construction. Thus

$$\begin{aligned}
\sum_{1 \leq t \leq \frac{s_2 - s_1 + 1}{2}} \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(Q_i) \cup V'(Q_j)}} \sum_{\substack{v \in L_1 \\ u \sim v}} p^2(u, v) &\leq \sum_{1 \leq t \leq \frac{s_2 - s_1 + 1}{2}} \sum_{T_{i,j} \in D_t} \sum_{\substack{u \in T_{i,j} \\ u \notin V'(Q_i) \cup V'(Q_j)}} \sum_{\substack{v \in L_1 \\ u \sim v}} \frac{c'''}{t^2 \log^6 n} \\
&\leq \sum_{1 \leq t \leq \frac{s_2 - s_1 + 1}{2}} \sum_{T_{i,j} \in D_t} \log^4 n \frac{c'''}{t^2 \log^6 n} \\
&\leq \sum_{1 \leq t \leq \frac{s_2 - s_1 + 1}{2}} (t+1) \frac{c'''}{t^2 \log^2 n} \\
&= O\left(\frac{1}{\log^2 n}\right) \sum_{1 \leq t \leq \frac{s_2 - s_1 + 1}{2}} O\left(\frac{1}{t}\right) \\
&= O\left(\frac{1}{\log n}\right).
\end{aligned}$$

it is clear that the contribution of D_0 the contribution to the formula is $O\left(\frac{1}{\log^2 n}\right)$. By symmetry, the contribution of D_t with $\frac{s_2 - s_1}{2} < t \leq s_2 - s_1 + 1$ is also $O\left(\frac{1}{t \log^2 n}\right)$, and thus

$$\mathcal{R}(x \longleftrightarrow y) = O\left(\frac{1}{\log n}\right).$$

Recall that we assumed that x, y belong to some dense cubes Q_i and Q_j , that at the same time are inside of some parallelepipeds in the same horizontal strip and separated by an even number of parallelepipeds. Before we lift these conditions, we first provide a useful construction: we claim that if $x', y' \in L_1(G)$ with $x' \in Q_i'$ and $y' \in Q_j'$ with both cubes being dense, and such that Q_i' and Q_j' are connected by a path \mathcal{L} of dense cubes of length $O(\log n)$, then $\mathcal{R}(x' \longleftrightarrow y') = O\left(\frac{1}{\log n}\right)$.

To see this, let x' send an amount of $\frac{1}{\lfloor \beta \log n \rfloor - 1}$ flow to the $\lfloor \beta \log n \rfloor - 1$ previously selected vertices of Q_i' . From there on, let the previously selected vertices to send a uniformly distributed flow to the selected $\lfloor \beta \log n \rfloor$ vertices that belong to the dense cube that follows in the path that connects Q_i' and Q_j' . We continue distributing in a uniform way the flow until we arrive until Q_j' . Once the flow reached the $\lfloor \beta \log n \rfloor - 1$ selected vertices of Q_j' , send all the flow back to y' . Denote this flow by θ . We then have

$$\mathcal{R}(x' \longleftrightarrow y') \leq \sum_{\substack{z \in Q_i' \\ x' \sim z}} \theta^2(x', z) + \sum_{\substack{z \in Q_j' \\ y' \sim z}} \theta^2(y', z) + \sum_{Q \in \mathcal{L}} \sum_{\substack{u \in Q \\ v \in L_1 \\ u \sim v}} \theta^2(u, v)$$

As before, the first two sums contribute $O(\log n)O(1/\log^2 n) = O(1/\log n)$. For the third sum, note that there are $O(\log n)$ dense cubes in \mathcal{L} , and between two such cubes there are $O(\log^2 n)$ edges with flow $O\left(\frac{1}{\log^2 n}\right)$ on each of them. Thus

$$\mathcal{R}(x' \longleftrightarrow y') = O\left(\frac{1}{\log n}\right) + \sum_{|\mathcal{L}|} O(\log^2 n) O\left(\frac{1}{\log^4 n}\right) = O\left(\frac{1}{\log n}\right),$$

and the claim is proved.

We now show how this claim allows us to prove the bound on the effective resistance for any $x \in C_i' \in T_{s_1, r_1}$ and $y \in C_j' \in T_{s_2, r_2}$ in the same strip, first assuming that both parallelepipeds do not belong to the boundary. First, if x and y are separated by an odd number of parallelepipeds (that is, $s_2 - s_1$ even), we find x' in a parallelepiped adjacent to x , at Euclidean distance (in the renormalized lattice of cubes) $O(\log n)$ from x . Thus, by the claim, $\mathcal{R}(x \longleftrightarrow x') = O\left(\frac{1}{\log n}\right)$, and since x' and y are separated by an even number of parallelepipeds, by the above, $\mathcal{R}(x' \longleftrightarrow y) =$

$O\left(\frac{1}{\log n}\right)$, and finally by Lemma 2.4, also $\mathcal{R}(x \longleftrightarrow y) = O\left(\frac{1}{\log n}\right)$. As before, by the same idea, we may extend our bounds also to arbitrary vertices inside the same parallelepiped (in cubes not satisfying all restrictions) and to vertices in parallelepipeds of a strip on the boundary.

To generalize the inequality between any pair of strips we use the same idea as in Section 5 from Figure 12: for any pair of strips, we find another vertex z in the intersection of the strips that contain x and y and that intersect at some parallelepiped T that contains the vertex z such that $\mathcal{R}(x \longleftrightarrow z)$ and $\mathcal{R}(y \longleftrightarrow z)$ are $O\left(\frac{1}{\log n}\right)$ (such z must exist, as there is a logarithmic amount of crossings of dense cubes in each strip). Together with the triangle inequality (Lemma 2.4) we thus finished proving that a.a.s. the effective resistance between any two vertices that belong to dense cubes that belong to the giant component of dense cubes is $O\left(\frac{1}{\log n}\right)$.

Step 2: Upper bound for the effective resistance between a vertex not in the giant and another in the giant

Now we prove that the effective resistance between any vertex $x \in G$ that is in a cube, being dense or non-dense, that is not in the giant component of the auxiliary Bernoulli site process, and some vertex $y \in G$, belonging to a dense cube in the giant component of the auxiliary Bernoulli site process is at most $O\left(\frac{1}{\log n}\right)$: by the triangle inequality (Lemma 2.4) together with Step 1, this will then show that for any pair of vertices $x, y \in G$, we have $\mathcal{R}(x \longleftrightarrow y) = O\left(\frac{1}{\log n}\right)$.

In order to show this bound on the effective resistance, it suffices to show that for every vertex x we find y belonging to the giant of the auxiliary Bernoulli site process together with a set $\mathcal{P}_{x,y}$ of $\Omega(\log n)$ internal vertex-disjoint paths between x and y of constant length. Indeed, in this case one can split evenly the unit flow emanating from x between all these paths, such that the flow on each such edge of each path is $O(1/\log n)$, and the resistance, by Definition 2.1 thus satisfies

$$\mathcal{R}(x \longleftrightarrow y) = \sum_{e \in \mathcal{P}_{x,y}} O(1/\log^2 n) = O\left(\frac{1}{\log n}\right).$$

We will follow this plan in detail now. We first assume that x is at distance at least $3C\sqrt{\log n}$ from all boundaries of Λ_n , where C is a sufficiently large constant. Recall that by the beginning of the proof, for every $x \in G$ there exists $y \in G$ at distance $rL/\sqrt{5} = O(\log n)$, with y belonging to a dense cube of the giant component of the auxiliary Bernoulli process (and therefore also $y \in L_1(G)$). We thus may find, an axis-parallel square S of side length $3C\sqrt{\log n}$ containing both x and y , such that both x and y are at distance at least say $C\sqrt{\log n}$ from the boundary of S , and such that the left bottom corner of S is an integer coordinate. We now show that we cannot separate x from y by removing less than $\varepsilon' \log n$ many vertices inside S (and therefore we cannot separate x and y in G either), for some $0 < \varepsilon' < \varepsilon$. Since this will hold a.a.s. for all x, y simultaneously (with S changing according to positions of x, y), we conclude that the graph is $\varepsilon' \log n$ -connected. By Menger's theorem, for every pair of vertices there exist $\varepsilon' \log n$ internally vertex-disjoint paths inside S between any pair of vertices, and we will show then that they can be chosen of constant length.

To do so, we adapt ideas from a classic k -connectivity argument on random geometric graphs (it is less precise than the proof of Theorem 1.1, in particular Proposition 5.2 in Penrose's paper [34], but some ideas are adapted), applied with $k = \varepsilon' \log n$. First, we show that after removing less than $\varepsilon' \log n$ many vertices, G a.a.s. does not have two components \mathcal{C}_1 and \mathcal{C}_2 both of Euclidean diameter larger than $C\sqrt{\log n}/5$ (C the same as before). In fact, we show that this cannot happen inside S . Indeed, if this were the case, then consider the cells of the tessellation with cells of sidelength $r/\sqrt{20}$ forming the exterior boundary of the smaller of the two components, say \mathcal{C}_1 (formally, this exterior boundary is defined as the exterior boundary of the component which contains cells that either contain a vertex of \mathcal{C}_1 or contain an edge of \mathcal{C}_1 , where an edge is drawn as a straight line). It is a standard topological fact that in \mathbb{R}^2 the outer boundary is diagonally connected, and its restriction to Λ_n contains at least a quarter of this boundary. The number of cells of this outer

boundary inside Λ_n is least $(C\sqrt{\log n}/20)/(r/\sqrt{20}) \geq C/4$. In fact, by choice of S , this lower bound holds also for the longest connected part of the exterior boundary inside S . Observe that the maximal Euclidean distance between two vertices in any two diagonally adjacent cells is at most $\sqrt{3r^2/20} \leq r/2$, and hence, if there were any vertex in a cell of the exterior boundary, such a vertex would be connected by an edge to a vertex in \mathcal{C}_1 (indeed, if the neighboring square was hitting only an edge of \mathcal{C}_1 , its distance to the edge were at most $r/2$, and from there to the closer endpoint of the edge at most $r/2$, so in total at most r). Now, denote by X the number of vertices in a fixed set of $C/4$ cells of radius $r/\sqrt{20}$. We have by standard Chernoff bounds for Poisson random variables

$$\Pr(X \leq \varepsilon' \log n) = \Pr(\text{Poi}(Cr^2/80) \leq \varepsilon' \log n) \leq (eC/(\pi\varepsilon'80))^{\varepsilon' \log n} e^{-Cr^2/80} \leq e^{-Cr^2/200},$$

assuming C large enough and $\varepsilon' < \varepsilon$. The number of such lattice animals is at most $O(n8^{C/4})$, and thus, by Chernoff bounds together with a union bound over all such animals, the probability of having in one of them less than $\varepsilon' \log n$ many vertices is at most $O(n8^{C/4}e^{-r^2C/200}) = O(n8^{C/4}e^{-(1+\varepsilon)C \log n/(200\pi)}) = o(n^{-2})$, where the equality follows by choosing C large enough. Hence, for C large, by taking a union bound over all pairs of vertices, a.a.s. such two components cannot exist.

Next, if after removing less than $\varepsilon' \log n$ vertices there would exist a connected component \mathcal{C}_1 of Euclidean diameter at most $C\sqrt{\log n}/5$ inside S , this component a.a.s. has $O(\log n)$ vertices. If the Euclidean diameter of such a component \mathcal{C}_1 were less than $\varepsilon''\sqrt{\log n}$ (for $\varepsilon'' < \varepsilon' < \varepsilon$ small enough; we define the Euclidean diameter of an isolated vertex to be zero and include this in this case), then note that the probability to have at most $\varepsilon' \log n$ points in the annulus obtained by taking the difference of the ball of outer radius r and inner radius $\varepsilon''\sqrt{\log n}/\pi$ centered around any of the vertices in \mathcal{C}_1 , is, once again by Chernoff bounds for Poisson variables, at most

$$O(\log n) \Pr(\text{Poi}((1+\varepsilon-\varepsilon'') \log n) \leq \varepsilon' \log n) = (e(1+\varepsilon-\varepsilon'')/\varepsilon')^{\varepsilon' \log n} e^{-(1+\varepsilon-\varepsilon''+o(1)) \log n} \leq n^{-1-\varepsilon'''},$$

for some $\varepsilon''' > 0$, where we assumed $\varepsilon'' < \varepsilon'$. By a union bound over all at most $n(1+o(1))$ components such a component a.a.s. does not exist.

Otherwise, if the Euclidean diameter were at least $\varepsilon''\sqrt{\log n}/\pi$ (but still at most $C\sqrt{\log n}/5$, and therefore still completely inside S), then observe that the smallest axis-parallel rectangle R containing all vertices of such a component has to have without loss of generality horizontal length at least $(\varepsilon''\sqrt{\log n}/\pi)/2$. A.s., such rectangle (of area $O(\log n)$) is defined by at most 4 points, and hence, for every fixed S , there are at most $O(\log^4 n)$ choices for the rectangle. Therefore, the halfcircle of radius r to the left of the leftmost point of R as well as the halfcircle to the right of the rightmost point of R have to contain at most $\varepsilon' \log n$ points (otherwise by removing less than $\varepsilon' \log n$ vertices \mathcal{C}_1 is not disconnected). The probability for this to happen is, by a union bound over all $O(n)$ choices for S and thus in total over all at most a.a.s. $O(n \log^4 n)$ choices of rectangles, by another Chernoff bound for Poisson random variables, at most

$$O(n \log^4 n) \Pr(\text{Poi}((1+\varepsilon) \log n) \leq \varepsilon' \log n) \leq (e(1+\varepsilon)/\varepsilon')^{\varepsilon' \log n} n^{-\varepsilon+o(1)} = o(1),$$

where we again assumed $\varepsilon' < \varepsilon$. Hence, a.a.s. such component also cannot exist, and for any $x \in G$ we find inside S the desired vertex y .

We now sketch the modifications when x is close to the boundary: suppose first that x is at distance at most $3C\sqrt{\log n}$ from exactly one boundary of Λ_n . In case of having two components \mathcal{C}_1 and \mathcal{C}_2 of diameter larger than $C\sqrt{\log n}/5$ after removing less than $\varepsilon' \log n$ vertices (one of them containing x), replacing S from before by $S \cap \Lambda_n$, the same argument as before shows that this cannot happen a.a.s. Similarly, a component \mathcal{C}_1 containing x of diameter less than $\varepsilon''\sqrt{\log n}$ cannot happen, as the semiannulus around x obtained by the difference of the ball of radius r around x and the one of radius $\varepsilon''\sqrt{\log n}$ cannot be empty a.a.s. (the union bound again is only over all $n^{1/2+o(1)}$ many vertices). Also, if the Euclidean diameter is of size at most $C\sqrt{\log n}/5$, but larger than $\varepsilon''\sqrt{\log n}$, a semicircle of radius r plus an extra area of size $\Theta(r)$ would have to be empty, and a.a.s. by a union bound over all $n^{1/2+o(1)}$ vertices this cannot happen.

In case x is at distance at most $3C\sqrt{\log n}$ from two boundaries of Λ_n , note that there are only $n^{o(1)}$ such vertices a.a.s., and the adaptations of the other arguments are also analogous.

Hence, as a conclusion, we see that G restricted to $S \cap \Lambda_n$ is a.a.s. $\varepsilon' \log n$ -connected. By Menger's theorem, for any x, y there exist thus ε' internally vertex-disjoint paths between x and y inside S . Moreover, since these paths are inside S , one may shorten these to make them paths of constant length by making it an induced path: for each path from x to y , we shorten it by maintaining only the first and last vertex of the path passing through any cube Q_i inside S (the path in between is replaced by a direct edge between these vertices), and hence its length is at most constant. The theorem follows.

7. CONCLUDING REMARKS AND FUTURE WORK

We showed that the cover time undergoes a jump at the threshold of connectivity, decreasing from $\Theta(n \log^2 n)$ to $\Theta(n \log n)$ a.a.s. It is therefore natural to ask what happens right at the threshold of connectivity: assume then that $r^d = (\log n + f(n))/V_d$, where $f(n)$ may be positive or negative, and $|f(n)| = o(\log n)$. If $f(n)$ is negative and $|f(n)| = \omega(1)$, the same argument as in Case 2 of Section 4, can be used to show that the number of vertices of degree 1 inside the giant component is a.a.s. $\Omega(nr^d V_d e^{-r^d V_d}) = \Omega(e^{-f(n)})$ (which is positive since $f(n)$ is negative). Hence, we obtain a set V' of those vertices of degree 1 vertices x, y for which $\log(|V'|) = -f(n)$, and any pair of vertices of $x, y \in V'$ satisfies $\mathcal{R}(x \longleftrightarrow y) \geq 1$. Thus, by Theorem 2.7, a.a.s. $\tau_{\text{cov}} \geq \gamma n \log n |f(n)|$. We do not know for which values of $f(n)$ this lower bound is tight: for some values of $f(n)$ a corresponding upper bound could perhaps be obtained by splitting the cubes into certain good cubes where the flow can be nicely dissipated and bad cubes where not, and combining them in an optimal way. We do not know what happens if $f(n)$ is positive.

More generally, in the same spirit as the very precise works on the cover time for the random walk on \mathbb{Z}^d , one could ask for more precise estimates on the cover time for any $r \leq r_c$. Since we do not have very precise estimates on the Green's function (this being essential for the calculations there) and we neither know of an isomorphism translating $\mathcal{G}(n, r, 2)$ into a problem on Gaussian free fields, this seems to be out of reach with the current techniques.

REFERENCES

- [1] Y. Abe. Effective resistances for supercritical percolation clusters in boxes. *Annales de l'IHP Probabilités et Statistiques*, 51(3):935–946, 2015.
- [2] Y. Abe. Second-order term of cover time for planar simple random walk. *Journal of Theoretical Probability*, 34:1689–1747, 2021.
- [3] I. F. Akyildiz, W. Su, Y. Sankarasubramaniam, and E. Cayirci. Wireless sensor networks: a survey. *Comput. Netw.*, 38(4):393–422, 2002.
- [4] D. Aldous. Threshold limits for cover times. *Journal of Theoretical Probability*, 4:197–211, 1991.
- [5] C. Avin and G. Ercal. On the cover time and mixing time of random geometric graphs. *Theoretical Computer Science*, 380(1-2):2–22, 2007.
- [6] M. Barlow, J. Ding, A. Nachmias, and Y. Peres. The evolution of the cover time. *Combin. Probab. Comput.*, 20:331–345, 2011.
- [7] D. Belius. Gumbel fluctuations for cover times in the discrete torus. *Probability Theory and Related Fields*, 157:635–689, 2013.
- [8] D. Belius and N. Kistler. The subleading order of two dimensional cover times. *Probability Theory and Related Fields*, 167(1-2):461–552, 2017.
- [9] B. Bollobás and O. Riordan. *Percolation*. Cambridge University Press, 2006.
- [10] A. K. Chandra, P. Raghavan, W. L. Ruzzo, and R. Smolensky. The electrical resistance of a graph captures its commute and cover times. In *Proceedings of the twenty-first annual ACM Symposium on Theory of Computing*, pages 574–586, 1989.
- [11] S. Chatterjee and M. Harel. Localization in random geometric graphs with too many edges. *Annals of Probability*, 48(2):574–621, 2020.
- [12] C. Cooper and A. Frieze. The cover time of random geometric graphs. *Random Structures and Algorithms*, 38(3):324–349, 2011.
- [13] C. Cooper, A. Frieze, and E. Lubetzky. Cover time of a random graph with given degree sequence ii: Allowing vertices of degree two. *Random Structures and Algorithms*, 45:627–674, 2014.

- [14] A. Cortines, O. Loidor, and S. Saglietti. A scaling limit for the cover time of the binary tree. *Advances in Mathematics*, 391:107974, 2021.
- [15] A. Dembo, Y. Peres, J. Rosen, and O. Zeitouni. Cover times for Brownian motion and random walks in two dimensions. *Annals of Mathematics*, 160:433–464, 2004.
- [16] J. Ding. On cover times for 2d lattices. *Electronic Journal of Probability*, 17:1–18, 2012.
- [17] J. Ding. Asymptotic of cover times via Gaussian Free Fields: Bounded degree graphs and general trees. *Ann. Probab.*, 42:464–496, 2014.
- [18] J. Ding, J. R. Lee, and Y. Peres. Cover times, blanket times, and majorizing measures. *Annals of Mathematics*, 175:1409–1471, 2012.
- [19] U. Feige. A tight lower bound on the cover time for random walks on graphs. *Random Structures and Algorithms*, 6(4):433–438, 1995.
- [20] U. Feige. A tight upper bound on the cover time for random walks on graphs. *Random Structures and Algorithms*, 6(1):51–54, 1995.
- [21] T. Friedrich, T. Sauerwald, and A. Stauffer. Diameter and broadcast time of random geometric graphs in arbitrary dimensions. *Algorithmica*, 67:65–88, 2013.
- [22] A. Frieze, W. Pegden, and T. Tkocz. On the cover time of the emerging giant. *SIAM Journal on Discrete Mathematics*, 36(3):10.1137/21M1441468, 2022.
- [23] D. Funke, S. Lamm, U. Meyer, M. Penschuck, P. Sanders, C. Schulz, D. Strash, and M. von Looz. Communication-free massively distributed graph generation. *J. Parallel Distrib. Comput.*, 131:200–217, 2019.
- [24] E. N. Gilbert. Random plane networks. *J. Soc. Ind. Appl. Math.*, 9:533–543, 1961.
- [25] P. Gupta and P. Kumar. Critical power for asymptotic connectivity in wireless networks. In: *Stochastic Analysis, Control, Optimization and Applications: A Volume in Honor of W. H. Fleming*, pages 547–566, 1998.
- [26] J. Kahn, J. H. Kim, L. Lovász, and V. H. Vu. The cover time, the blanket time, and the Matthews bound. In *Proceedings 41st Annual Symposium on Foundations of Computer Science*, pages 467–475. IEEE, 2000.
- [27] H. Kesten. *Percolation theory for mathematicians (Vol. 2)*. Boston: Birkhäuser, 1982.
- [28] M. Kiwi, M. Schepers, and J. Sylvester. Cover and hitting times of hyperbolic random graphs. *Random Structures and Algorithms*, 65(4):915–978, 2024.
- [29] G. Lawler. On the covering time of a disc by simple random walk in two dimensions. *Seminar on Stochastic Processes*, pages 189–207, 1992.
- [30] D. A. Levin and Y. Peres. *Markov chains and mixing times*, volume 107. American Mathematical Society, 2017.
- [31] T. M. Liggett, R. H. Schonmann, and A. M. Stacey. Domination by product measures. *The Annals of Probability*, 25(1):71–95, 1997.
- [32] O. Loidor and S. Saglietti. Tightness for the cover time of wired planar domains. *Preprint available at <https://arxiv.org/pdf/2406.11034>*, 2024.
- [33] M. Nekovee. Worm epidemics in wireless ad-hoc networks. *New J. Phys.*, 9:189, 2007.
- [34] M. Penrose. On k-connectivity for a geometric random graph. *Random Structures and Algorithms*, 15(2):145–164, 1999.
- [35] M. Penrose. *Random geometric graphs*, volume 5 of *Oxf. Stud. Probab.* Oxford: Oxford University Press, 2003.
- [36] M. Penrose. Lectures on random geometric graphs. In N. Fountoulakis and D. Hefetz, editors, *Random graphs, geometry and asymptotic structure*, pages 67–101. Cambridge: Cambridge University Press, 2016.
- [37] M. D. Penrose and A. Pisztora. Large deviations for discrete and continuous percolation. *Advances in Applied Probability*, 28(1):29–52, 1996.
- [38] A. Zhai. Exponential concentration of cover times. *Electron. J. Probab.*, 23:1–22, 2018.

(Carlos Martinez) IMCA, UNIVERSIDAD NACIONAL DE INGENIERÍA, LOS BIÓLOGOS 245, LA MOLINA, LIMA, PERU
 Email address: `carlos.martinez@imca.edu.pe`

(Dieter Mitsche) IMC, PONTIFICIA UNIVERSIDAD CATÓLICA, AVDA. VICUÑA MACKENNA 4860, SANTIAGO, CHILE
 Email address: `dieter.mitsche@uc.cl`