q-analogues of sums of consecutive powers of natural numbers and extended Carlitz q-Bernoulli numbers and polynomials

BAKIR FARHI

National Higher School of Mathematics P.O.Box 75, Mahelma 16093, Sidi Abdellah (Algiers)

Algeria

bakir.farhi@nhsm.edu.dz

http://farhi.bakir.free.fr/

Abstract

In this paper, we investigate a specific class of q-polynomial sequences that serve as a q-analogue of the classical Appell sequences. This framework offers an elegant approach to revisiting classical results by Carlitz and, more interestingly, to establishing an important extension of the Carlitz q-Bernoulli polynomials and numbers. In addition, we establish explicit series representations for our extended Carlitz q-Bernoulli numbers and express them in terms of q-Stirling numbers of the second kind. This leads to a novel formula that explicitly connects the Carlitz q-Bernoulli numbers with the q-Stirling numbers of the second kind.

MSC 2020: Primary 05A30; Secondary 11B68, 11B73, 05A40.

Keywords: Carlitz *q*-Bernoulli numbers and polynomials, *q*-analogue of Appell sequences, *q*-analogues of sums of consecutive powers of natural numbers, *q*-Stirling numbers of the second kind, umbral calculus.

1 Introduction and Notation

Throughout this paper, we let \mathbb{N} and \mathbb{N}_0 respectively denote the set of positive integers and the set of nonnegative integers. For $n \in \mathbb{N}_0$, we let $(X)_n$ denote the *falling factorial* of X to depth n, defined as

$$(X)_n := X(X-1)\cdots(X-n+1).$$

The Stirling numbers of the second kind, denoted S(n,k) $(n,k \in \mathbb{N}_0, n \ge k)$, are then defined through the polynomial identity:

$$X^{n} = \sum_{k=0}^{n} S(n,k)(X)_{k} \qquad (\forall n \in \mathbb{N}_{0}).$$

The Bernoulli polynomials are denoted, as usual, by $B_n(X)$ and the Bernoulli numbers by B_n $(n \in \mathbb{N}_0)$. A modern definition of the Bernoulli polynomials and numbers employs their exponential generating functions, given by

$$\frac{t}{e^t - 1}e^{Xt} = \sum_{n=0}^{\infty} B_n(X)\frac{t^n}{n!} \quad , \quad \frac{t}{e^t - 1} = \sum_{n=0}^{\infty} B_n\frac{t^n}{n!}$$

(so $B_n = B_n(0)$ for all $n \in \mathbb{N}_0$). As is well-known, the Bernoulli polynomials and numbers play a central role in various branches of mathematics, including number theory, mathematical analysis, and algebraic geometry. For a modern and exhaustive perspective, the reader is referred to the paper by Kouba [7]. An interesting and well-known formula expressing the Bernoulli numbers in terms of the Stirling numbers of the second kind is given by:

$$B_n = \sum_{k=0}^n (-1)^k \frac{k!}{k+1} S(n,k) \qquad (\forall n \in \mathbb{N}_0)$$
(1.1)

(see e.g., [4, Corollary 2.8]).

It should be noted that the Bernoulli polynomial sequence $(B_n(X))_{n \in \mathbb{N}_0}$ belongs to the broader class of Appell sequences (see [3, 9]). These are sequences of polynomials $(P_n(X))_{n \in \mathbb{N}_0}$ characterized by the following properties: $P_0(X)$ is a nonzero constant polynomial, and for all $n \in \mathbb{N}$, the derivatives satisfy the relation $P'_n(X) = nP_{n-1}(X)$. Using the Taylor expansion, an Appell polynomial sequence $(P_n(X))_n$ can also be characterized by its general term, which takes the form

$$P_n(X) = \sum_{k=0}^n \binom{n}{k} p_k X^{n-k} \qquad (\forall n \in \mathbb{N}_0)$$

where $(p_k)_{k \in \mathbb{N}_0}$ is a sequence of scalars with $p_0 \neq 0$.

Further, let q be a positive real parameter. The q-analogue of X, whether treated as an indeterminate or a number, is defined by:

$$[X] := \frac{q^X - 1}{q - 1}$$

For $k \in \mathbb{N}_0$, the q-analogue $[X]_k$ of $(X)_k$ is defined as

$$[X]_k := [X] \cdot [X-1] \cdots [X-k+1].$$

For $n \in \mathbb{N}_0$, the q-analogue of n! is given by

$$[n]! := [n]_n = [n] \cdot [n-1] \cdots [1].$$

Furthermore, for $n, k \in \mathbb{N}_0$ with $n \ge k$, the q-analogue $\begin{bmatrix} n \\ k \end{bmatrix}$ of the binomial coefficient $\binom{n}{k}$ is defined as

$$\begin{bmatrix} n \\ k \end{bmatrix} := \frac{[n]!}{[k]![n-k]!} = \frac{[n]_k}{[k]!}.$$

These numbers appear, in particular, in the famous Gauss binomial formula:

$$\prod_{k=0}^{n-1} \left(x + q^k y \right) = \sum_{k=0}^n q^{\frac{k(k-1)}{2}} {n \brack k} x^{n-k} y^k \qquad (\forall n \in \mathbb{N}, \forall x, y \in \mathbb{R}),$$
(1.2)

and, just like the classical binomial coefficients, they also have a combinatorial interpretation (see, e.g., [5]).

On the other hand, we let Δ denote the forward difference operator which acts linearly on $\mathbb{R}[X]$ by the formula:

$$(\Delta P)(X) := P(X+1) - P(X) \qquad (\forall P \in \mathbb{R}[X]).$$

It is well known and easy to verify that the *n*-fold composition of Δ is given by the formula:

$$(\Delta^n P)(X) = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} P(X+k) \qquad (\forall P \in \mathbb{R}[X], \forall n \in \mathbb{N}_0).$$
(1.3)

In this paper, we call a *q*-polynomial any polynomial in [X] with real coefficients. Equivalently, a *q*-polynomial is a polynomial in q^X with real coefficients. The degree of a *q*-polynomial refers to its degree as a polynomial in [X] (or equivalently in q^X). We denote by \mathscr{E} the \mathbb{R} vector space of all *q*-polynomials and by \mathscr{E}_d ($d \in \mathbb{N}_0$) the \mathbb{R} -vector subspace of \mathscr{E} consisting of *q*-polynomials of degree $\leq d$. An important example of *q*-polynomials arises naturally in the evaluation of sums of the form

$$S_{n,r}(N) := \sum_{k=0}^{N-1} q^{rk} [k]^n$$

 $(N, n \in \mathbb{N}_0, r \in \mathbb{N})$, which are q-analogues of the sum of powers $\sum_{k=0}^{N-1} k^n$. More interestingly, when expressed as a linear combination of the q-polynomials $q^{kN}[N]^{n-k}$ ($0 \le k \le n$), $S_{n,r}(N)$ leads to a q-analogue example of Appell polynomial sequences (see Theorem 2.6). Perhaps it was by exploring this fact that Carlitz [1] obtained important q-analogues of Bernoulli numbers and polynomials. By means of the symbolic calculus, Carlitz [1] defined two real sequences $(\eta_n)_{n\in\mathbb{N}_0}$ and $(\beta_n)_{n\in\mathbb{N}_0}$ (depending on q) by:

$$\begin{cases} \eta_0 = 1, \eta_1 = 0, \\ (q\eta + 1)^n = \eta^n \quad (\forall n \ge 2) \end{cases},$$
(1.4)

$$\begin{cases} \beta_0 = 1, \\ q(q\beta + 1)^n - \beta^n = \delta_{n,1} \quad (\forall n \in \mathbb{N}) \end{cases},$$
(1.5)

where $\delta_{i,j}$ denotes the Kronecker delta. Furthermore, it is noted that for all $n \in \mathbb{N}_0$:

$$\beta_n = \eta_n + (q-1)\eta_{n+1}.$$
 (1.6)

Although both sequences are intriguing, the sequence $(\beta_n)_n$ is particularly noteworthy because its terms are well-defined at q = 1 (unlike $(\eta_n)_n$, for which only the first two terms are defined at q = 1). Moreover, specializing q = 1 in β_n recovers the n^{th} Bernoulli number B_n . These facts are far from trivial. To prove them, Carlitz was led to introduce q-analogues of the Stirling numbers of the second kind, $S_q(n,k)(n,k \in \mathbb{N}_0, n \ge k)$, which he defined via the following identity of q-polynomials:

$$[X]^{n} = \sum_{k=0}^{n} q^{\frac{1}{2}k(k-1)} S_{q}(n,k) [X]_{k}$$
(1.7)

(for all $n \in \mathbb{N}_0$). Specializing q = 1 in $S_q(n, k)$ $(n, k \in \mathbb{N}_0, n \ge k)$ recovers the classical Stirling numbers of the second kind S(n, k). Using the $S_q(n, k)$'s, Carlitz [1] established the following formula for β_n $(n \in \mathbb{N}_0)$:

$$\beta_n = \sum_{k=0}^n (-1)^k \frac{[k]!}{[k+1]} S_q(n,k), \qquad (1.8)$$

which confirms that β_n is always well-defined at q = 1, with value equal to the Bernoulli number B_n (according to (1.1)).

Carlitz also associated to the sequences $(\eta_n)_n$ and $(\beta_n)_n$ the q-polynomial sequences $(\eta_n(X))_n$ and $(\beta_n(X))_n$, given by:

$$\eta_n(X) = \sum_{k=0}^n \binom{n}{k} \eta_k q^{kX} [X]^{n-k},$$
(1.9)

$$\beta_n(X) = \sum_{k=0}^n \binom{n}{k} \beta_k q^{kX} [X]^{n-k}$$
(1.10)

(for all $n \in \mathbb{N}_0$). The original sequences $(\eta_n)_n$ and $(\beta_n)_n$ are then respectively obtained as the values of $(\eta_n(X))_n$ and $(\beta_n(X))_n$ at X = 0. Especially, $(\beta_n(X))_n$ constitutes a q-analogue of the classical Bernoulli polynomial sequence $(B_n(X))_n$. That said, alternative definitions of the Carlitz q-Bernoulli numbers and polynomials exist, such as those based on generating functions or p-adic q-integrals on \mathbb{Z}_p . For example, Koblitz [6] constructed q-analogues of the p-adic Dirichlet L-series that interpolate the Carlitz q-Bernoulli numbers.

In this work, we first incorporate the Carlitz q-polynomial sequences $(\eta_n(X))_n$ and $(\beta_n(X))_n$ into a broader class of q-polynomial sequences, which we call "Carlitz-type q-polynomial sequences". We establish a fundamental theorem showing that this class of q-polynomial sequences serves as a q-analogue of the classical Appell polynomial sequences. Next, We verify that the closed forms of the q-analogues of power sums of consecutive natural numbers can essentially be expressed in terms of Carlitz-type q-polynomial sequences. Furthermore, we show that a specific subclass of Carlitz-type q-polynomial sequences (including $(\eta_n(X))_n$ and $(\beta_n(X))_n)$ can be generated using simple symbolic formulas, such as those established by Carlitz in [1]. We then establish several properties of Carlitz-type q-polynomial sequences and leverage these properties to extend $(\eta_n(X))_n$ and $(\beta_n(X))_n$ into $(\beta_n^{(r)}(X))_n$ $(r \in \mathbb{N}_0)$, which provide important q-analogues of the classical Bernoulli polynomials for $r \geq 1$. Additionally, we represent the numbers $\beta_n^{(r)} := \beta_n^{(r)}(0)$ as series and express them in terms of the *q*-Stirling numbers of the second kind. Finally, we derive a novel formula connecting the Carlitz *q*-Bernoulli numbers with the *q*-Stirling numbers of the second kind.

2 The results and the proofs

2.1 Carlitz-type *q*-polynomial sequences

We first introduce the following definition:

Definition 2.1. Let $(T_n(X))_{n \in \mathbb{N}_0}$ be a *q*-polynomial sequence. We say that $(T_n(X))_n$ is *Carlitz-type* if its general term $T_n(X)$ $(n \in \mathbb{N}_0)$ has the form:

$$T_n(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k q^{kX},$$

where $(a_k)_{k\in\mathbb{N}_0}$ is a real sequence. In this case, $(a_k)_k$ is called the associated sequence of the Carlitz-type q-polynomial sequence $(T_n(X))_n$.

An important characterization of the Carlitz-type q-polynomial sequences is given by the following fundamental theorem:

Theorem 2.2. Let $(T_n(X))_{n \in \mathbb{N}_0}$ be a q-polynomial sequence, and for all $n \in \mathbb{N}_0$, set $t_n := T_n(0)$. Then $(T_n(X))_n$ is Carlitz-type if and only if we have for all $n \in \mathbb{N}_0$:

$$T_n(X) = \sum_{k=0}^n \binom{n}{k} t_k q^{kX} [X]^{n-k},$$

that is (symbolically):

$$T_n(X) = \left(q^X t + [X]\right)^n.$$

To prove this theorem, we need to use the inversion formula from the following lemma:

Lemma 2.3 (see e.g, [2, 8]). Let $(u_n)_{n \in \mathbb{N}_0}$ and $(v_n)_{n \in \mathbb{N}_0}$ be two real sequences. Then the two following identities are equivalent:

$$u_n = \sum_{k=0}^n \binom{n}{k} v_k \qquad (\forall n \in \mathbb{N}_0),$$
$$v_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} u_k \qquad (\forall n \in \mathbb{N}_0).$$

Proof of Theorem 2.2. Suppose that $(T_n(X))_n$ is Carlitz-type. So there exists a real sequence $(a_n)_{n \in \mathbb{N}_0}$ for which we have for all $n \in \mathbb{N}_0$:

$$T_n(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k q^{kX}.$$

In particular, we have for all $n \in \mathbb{N}_0$:

$$t_n := T_n(0) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k,$$

that is

$$(q-1)^n t_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k.$$

By inverting this last formula (using Lemma 2.3), we derive that for all $n \in \mathbb{N}_0$:

$$a_n = \sum_{k=0}^n \binom{n}{k} (q-1)^k t_k$$

Then, by inserting this into the expression of $T_n(X)$, we obtain that for all $n \in \mathbb{N}_0$:

$$T_n(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \left(\sum_{i=0}^k \binom{k}{i} (q-1)^i t_i \right) q^{kX}$$
$$= (q-1)^{-n} \sum_{k=0}^n \sum_{i=0}^k (-1)^{n-k} \binom{n}{k} \binom{k}{i} (q-1)^i t_i q^{kX}.$$

By inverting the summations and remarking that $\binom{n}{k}\binom{k}{i} = \binom{n}{i}\binom{n-i}{k-i}$ (for all $n \in \mathbb{N}_0, k \in \{0, 1, \dots, n\}$, and $i \in \{0, 1, \dots, k\}$), we derive that for all $n \in \mathbb{N}_0$:

$$T_n(X) = (q-1)^{-n} \sum_{0 \le i \le n} \sum_{i \le k \le n} (-1)^{n-k} \binom{n}{i} \binom{n-i}{k-i} (q-1)^i t_i q^{kX}$$

= $(q-1)^{-n} \sum_{0 \le i \le n} \sum_{0 \le j \le n-i} (-1)^{n-i-j} \binom{n}{i} \binom{n-i}{j} (q-1)^i t_i q^{(i+j)X}$ (where we set $j = k-i$)
= $(q-1)^{-n} \sum_{0 \le i \le n} \binom{n}{i} (q-1)^i t_i q^{iX} \sum_{0 \le j \le n-i} \binom{n-i}{j} q^{jX} (-1)^{n-i-j}.$

But according to the binomial formula, we have for all $n \in \mathbb{N}_0$ and all $i \in \{0, 1, \dots, n\}$:

$$\sum_{0 \le j \le n-i} \binom{n-i}{j} q^{jX} (-1)^{n-i-j} = \left(q^X - 1\right)^{n-i} = (q-1)^{n-i} [X]^{n-i}.$$

By inserting this into the last obtained expression for $T_n(X)$, we finally derive that for all $n \in \mathbb{N}_0$:

$$T_n(X) = \sum_{0 \le i \le n} \binom{n}{i} t_i q^{iX} [X]^{n-i},$$

as required.

Conversely, suppose that we have for all $n \in \mathbb{N}_0$:

$$T_n(X) = \sum_{k=0}^n \binom{n}{k} t_k q^{kX} [X]^{n-k},$$

and define for all $n \in \mathbb{N}_0$:

$$a_n := \sum_{k=0}^n \binom{n}{k} (q-1)^k t_k$$
(2.1)

and

$$S_n(X) := (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k q^{kX}.$$

Equivalently, $(S_n(X))_{n \in \mathbb{N}_0}$ is a Carlitz-type q-polynomial sequence, with associated sequence $(a_n)_{n \in \mathbb{N}_0}$. So, according to the first part of this proof, we have for all $n \in \mathbb{N}_0$:

$$S_n(X) = \sum_{k=0}^n \binom{n}{k} s_k q^{kX} [X]^{n-k},$$
(2.2)

where

$$s_n := S_n(0) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k.$$
(2.3)

But the inversion of Formula (2.1) (using Lemma 2.3) gives for all $n \in \mathbb{N}_0$:

$$(q-1)^n t_n = \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k,$$

that is

$$t_n = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k = s_n$$

(according to (2.3)). Hence we have for all $n \in \mathbb{N}_0$:

$$T_n(X) = \sum_{k=0}^n \binom{n}{k} t_k q^{kX} [X]^{n-k}$$
$$= \sum_{k=0}^n \binom{n}{k} s_k q^{kX} [X]^{n-k}$$
$$= S_n(X) \qquad (according to (2.2))$$

Consequently $(T_n(X))_n = (S_n(X))_n$ is Carlitz-type, as required. This complete this proof. *Remark* 2.4. From the perspective of Theorem 2.2, the Carlitz-type q-polynomial sequences can be regarded as a q-analogue of the Appell polynomial sequences.

Examples 2.5. In [1], Carlitz introduced two important q-polynomial sequences, which are $(\eta_n(X))_{n \in \mathbb{N}_0}$ and $(\beta_n(X))_{n \in \mathbb{N}_0}$, defined for all $n \in \mathbb{N}_0$ by:

$$\eta_n(X) := (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{k}{[k]} q^{kX}, \qquad (2.4)$$

$$\beta_n(X) := (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{k+1}{[k+1]} q^{kX}$$
(2.5)

(where we take by convention $\frac{k}{[k]} = 1$ for k = 0). From their definitions, it is clear that $(\eta_n(X))_n$ and $(\beta_n(X))_n$ are both Carlitz-type with associated real sequences $(\frac{k}{[k]})_k$ and $(\frac{k+1}{[k+1]})_k$

respectively. In addition, Carlitz [1] explored the real sequences constituted of the values of $(\eta_n(X))_n$ and $(\beta_n(X))_n$ at X = 0; that is the sequences $(\eta_n)_{n \in \mathbb{N}_0}$ and $(\beta_n)_{n \in \mathbb{N}_0}$, defined by:

$$\eta_n := \eta_n(0) \text{ and } \beta_n := \beta_n(0) \quad (\forall n \in \mathbb{N}_0).$$

So, according to Theorem 2.2, we have important alternative expressions for $\eta_n(X)$ and $\beta_n(X)$ $(n \in \mathbb{N}_0)$, which are

$$\eta_n(X) = \sum_{k=0}^n \binom{n}{k} \eta_k q^{kX} [X]^{n-k},$$
(2.6)

$$\beta_n(X) = \sum_{k=0}^n \binom{n}{k} \beta_k q^{kX} [X]^{n-k}.$$
 (2.7)

Using the symbolic calculus, Formulas (2.6) and (2.7) can be written more simply as:

$$\eta_n(X) = \left(q^X \eta + [X]\right)^n, \qquad (2.8)$$

$$\beta_n(X) = \left(q^X \beta + [X]\right)^n \tag{2.9}$$

(for all $n \in \mathbb{N}_0$).

We now see that Carlitz-type q-polynomial sequences naturally emerge when deriving closed-form expressions for q-analogues of power sums of consecutive positive integers. We begin with the following theorem:

Theorem 2.6. Let r be a positive integer. For all nonnegative integer n and all positive integer N, we have

$$\sum_{k=0}^{N-1} q^{rk} [k]^n = q^{rN} S_n(N) - S_n(0),$$

where $(S_k(X))_{k \in \mathbb{N}_0}$ is the Carlitz-type q-polynomial sequence associated to the real sequence of general term $a_k = \frac{1}{q-1} \frac{1}{[k+r]}$ ($\forall k \in \mathbb{N}_0$).

Proof. Let $n \in \mathbb{N}_0$ and $N \in \mathbb{N}$ be fixed. We have

$$\begin{split} \sum_{k=0}^{N-1} q^{rk} [k]^n &= \sum_{k=0}^{N-1} q^{rk} \left(\frac{q^k - 1}{q - 1} \right)^n \\ &= (q - 1)^{-n} \sum_{k=0}^{N-1} q^{rk} \left(q^k - 1 \right)^n \\ &= (q - 1)^{-n} \sum_{k=0}^{N-1} q^{rk} \sum_{i=0}^n \binom{n}{i} q^{ki} (-1)^{n-i} \qquad \text{(by the binomial formula)} \\ &= (q - 1)^{-n} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \sum_{k=0}^{N-1} q^{(i+r)k} \\ &= (q - 1)^{-n} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \frac{q^{(i+r)N} - 1}{q^{i+r} - 1} \\ &= (q - 1)^{-n} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \frac{1}{(q - 1)[i + r]} \left(q^{(i+r)N} - 1 \right). \end{split}$$

Setting for all $n \in \mathbb{N}_0$:

$$S_n(X) := (q-1)^{-n} \sum_{i=0}^n (-1)^{n-i} \binom{n}{i} \frac{1}{(q-1)[i+r]} q^{iX}$$

 $(so (S_n(X))_n$ is the Carlitz-type q-polynomial sequence associated to the real sequence of general term $a_i = \frac{1}{(q-1)[i+r]}$, we conclude that:

$$\sum_{n=0}^{N-1} q^{rk} [k]^n = q^{rN} S_n(N) - S_n(0),$$

as required.

The following generalization of Theorem 2.6 aims to provide a larger class of real sequences, for which the associated Carlitz-type q-polynomial sequences naturally arise when evaluating q-analogues of power sums of consecutive positive integers.

Theorem 2.7. Let r be a positive integer and d be a nonnegative integer such that $r \ge d$. For all integer $n \ge d$ and all positive integer N, we have

$$\sum_{k=0}^{N-1} q^{rk} [k]^{n-d} = \frac{q^{(r-d)N} T_n(N) - T_n(0)}{n(n-1)\cdots(n-d+1)},$$

where $(T_k(X))_{k \in \mathbb{N}_0}$ is the Carlitz-type q-polynomial sequence associated to the real sequence of general term

$$b_k = (q-1)^{d-1} \frac{k(k-1)\cdots(k-d+1)}{[k+r-d]} \qquad (\forall k \in \mathbb{N}_0).$$

Proof. Let n and N be fixed integers such that $n \ge d$ and $N \ge 1$. According to Theorem 2.6, we have that

$$\sum_{k=0}^{N-1} q^{rk} [k]^{n-d} = q^{rN} S_{n-d}(N) - S_{n-d}(0),$$

where $(S_k(X))_{k \in \mathbb{N}_0}$ is the Carlitz-type *q*-polynomial sequence associated to the real sequence of general term $a_k = \frac{1}{(q-1)[k+r]}$ ($\forall k \in \mathbb{N}_0$). By introducing the *q*-polynomial sequence $(T_k(X))_{k \in \mathbb{N}_0}$ defined by:

$$T_k(X) = \begin{cases} 0 & \text{if } k < d\\ k(k-1)\cdots(k-d+1)q^{dX}S_{k-d}(X) & \text{else} \end{cases} \quad (\forall k \in \mathbb{N}_0)$$

the closed form of the sum of q-powers above is transformed to

$$\sum_{k=0}^{N-1} q^{rk} [k]^{n-d} = \frac{q^{(r-d)N} T_n(N) - T_n(0)}{n(n-1)\cdots(n-d+1)}.$$

So it remains to verify that $(T_k(X))_k$ is Carlitz-type with the associated sequence that given by the theorem. For an integer $k \ge d$, we have

$$T_k(X) := k(k-1)\cdots(k-d+1)q^{dX}S_{k-d}(X)$$

9

$$= k(k-1)\cdots(k-d+1)q^{dX}\cdot(q-1)^{-k+d}\sum_{i=0}^{k-d}(-1)^{k-d-i}\binom{k-d}{i}\frac{1}{(q-1)[i+r]}q^{iX}$$
$$= k(k-1)\cdots(k-d+1)q^{dX}(q-1)^{-k+d}\sum_{j=d}^{k}(-1)^{k-j}\binom{k-d}{j-d}\frac{1}{(q-1)[j+r-d]}q^{(j-d)X}$$

(by setting j = i + d). But since $k(k-1)\cdots(k-d+1)\binom{k-d}{j-d} = j(j-1)\cdots(j-d+1)\binom{k}{j}$ (for all integers $k, j \ge d$), it follows that for all integer $k \ge d$:

$$T_k(X) = (q-1)^{-k} \sum_{j=d}^k (-1)^{k-j} \binom{k}{j} (q-1)^{d-1} \frac{j(j-1)\cdots(j-d+1)}{[j+r-d]} q^{jX}$$
$$= (q-1)^{-k} \sum_{j=0}^k (-1)^{k-j} \binom{k}{j} (q-1)^{d-1} \frac{j(j-1)\cdots(j-d+1)}{[j+r-d]} q^{jX}.$$

By remarking that the last formula for $T_k(X)$ is obviously valid for $0 \le k < d$, we conclude that $(T_k(X))_k$ is Carlitz-type and its associated real sequence has the general term $b_j = (q-1)^{d-1} \frac{j(j-1)\cdots(j-d+1)}{[j+r-d]} \quad (\forall j \in \mathbb{N}_0).$ The theorem is thus proved.

Remark 2.8. It is important to note that the general term of the real sequence $(b_k)_{k \in \mathbb{N}_0}$ associated to the *q*-polynomial sequence in Theorem 2.7 has the form $\frac{P(k)}{[k+s]}$, with *P* is a real polynomial and *s* is a nonnegative integer. This fact will be used below for establishing some important symbolic formulas.

In the following theorem, we will establish symbolic formulas generating the values at 0 of some particular Carlitz-type q-polynomial sequences.

Theorem 2.9. Let r and d be two nonnegative integers and P be a real polynomial of degree d. Let also $(T_n(X))_{n\in\mathbb{N}_0}$ be the Carlitz-type q-polynomial sequence whose associated real sequence is $\left(\frac{P(k)}{[k+r]}\right)_{k\in\mathbb{N}_0}$ and $(t_n)_{n\in\mathbb{N}_0}$ be the real sequence defined by $t_n := T_n(0)$ ($\forall n \in \mathbb{N}_0$). Then $(t_n)_n$ satisfies the symbolic formula:

$$q^r \left(qt+1\right)^n = t^n \qquad (\forall n > d).$$

Proof. Let n > d be a fixed integer. From the definition of $T_n(X)$, we have that

$$T_n(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{P(k)}{[k+r]} q^{kX}.$$

Thus

$$q^{r}T_{n}(1) - T_{n}(0) = (q-1)^{-n} \sum_{k=0}^{n} (-1)^{n-k} {n \choose k} \frac{P(k)}{[k+r]} (q^{k+r} - 1)$$

= $(q-1)^{-n+1} \sum_{k=0}^{n} (-1)^{n-k} {n \choose k} P(k)$ (since $q^{k+r} - 1 = (q-1)[k+r]$)
= $(q-1)^{-n+1} (\Delta^{n} P) (0)$ (according to (1.3))
= 0 (since $n > \deg P = d$).

Consequently, we have

$$q^{r}T_{n}(1) = T_{n}(0) = t_{n}.$$
(2.10)

On the other hand, we have (according to Theorem 2.2) the symbolic formula:

$$T_n(X) = \left(q^X t + [X]\right)^n$$

In particular, we have symbolically

$$T_n(1) = \left(qt+1\right)^n.$$

By inserting this into (2.10), it finally follows that we have symbolically

$$q^r \left(qt+1\right)^n = t^n,$$

as required.

Examples 2.10.

1. Since the general term of the associated real sequences of the Carlitz-type q-polynomial sequences $(\eta_n(X))_{n \in \mathbb{N}_0}$ and $(\beta_n(X))_{n \in \mathbb{N}_0}$ are respectively $\frac{k}{[k]}$ and $\frac{k+1}{[k+1]}$ then (according to Theorem 2.9) the real sequences $(\eta_n)_{n \in \mathbb{N}_0}$ and $(\beta_n)_{n \in \mathbb{N}_0}$ satisfy the symbolic formulas:

$$(q\eta + 1)^{n} = \eta^{n}$$

$$q (q\beta + 1)^{n} = \beta^{n}$$

 $(\forall n > 1)$

2. Let us put ourselves in the situation of Theorem 2.7 and set $t_k := T_k(0)$ (for all $k \in \mathbb{N}_0$). Since the general term of the associated real sequence of the Carlitz-type q-polynomial sequence $(T_k(X))_k$ has the form $\frac{P(k)}{[k+r-d]}$, where P is a real polynomial of degree d, then (according to Theorem 2.9) the sequence $(t_k)_k$ satisfy the symbolic formula:

$$q^{r-d} \left(qt+1\right)^k = t^k \qquad (\forall k > d).$$

In the following proposition, we provide some important properties of Carlitz-type qpolynomial sequences. For simplicity, given a real or functional sequence $(f_n)_{n \in \mathbb{N}_0}$, we denote (with a slight abuse of notation) by $(nf_{n-1})_{n \in \mathbb{N}_0}$ the sequence whose terms are defined as

$$\begin{cases} 0 & \text{if } n = 0, \\ n f_{n-1} & \text{otherwise} \end{cases}.$$

Proposition 2.11. Let $(T_n(X))_{n \in \mathbb{N}_0}$ be a Carlitz-type q-polynomial sequence and $(a_k)_{k \in \mathbb{N}_0}$ be its associated real sequence. Then the following properties hold.

1. The q-polynomial sequence $(q^X n T_{n-1}(X))_{n \in \mathbb{N}_0}$ is Carlitz-type with associated real sequence $((q-1)ka_{k-1})_{k \in \mathbb{N}_0}$.

- 2. If $T_0(X) = 0_{\mathscr{E}}$ (i.e., $a_0 = 0$) then the q-polynomial sequence $\left(\frac{q^{-X}T_{n+1}(X)}{n+1}\right)_{n \in \mathbb{N}_0}$ is Carlitz-type with associated real sequence $\left(\frac{1}{q-1}\frac{a_{k+1}}{k+1}\right)_{k \in \mathbb{N}_0}$.
- 3. The sequence $(q^{-X}(T_n(X) + (q-1)T_{n+1}(X)))_{n \in \mathbb{N}_0}$ is a Carlitz-type q-polynomial sequence with associated real sequence $(a_{k+1})_{k \in \mathbb{N}_0}$.
- 4. The q-polynomial sequence $\left(n\left(T_{n-1}(X)+(q-1)T_n(X)\right)\right)_{n\in\mathbb{N}_0}$ is Carlitz-type with associated real sequence $\left((q-1)ka_k\right)_{k\in\mathbb{N}_0}$.

Proof. By definition, we have for all $n \in \mathbb{N}_0$:

$$T_n(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k q^{kX}.$$
(2.11)

• Let us show the 1st item of the proposition. Using (2.11), we have for all $n \in \mathbb{N}_0$:

$$q^{X}nT_{n-1}(X) = (q-1)^{-n+1} \sum_{k=0}^{n-1} (-1)^{n-1-k} n\binom{n-1}{k} a_{k} q^{(k+1)X}$$
$$= (q-1)^{-n} \sum_{k=1}^{n} (-1)^{n-k} n\binom{n-1}{k-1} (q-1) a_{k-1} q^{kX}.$$

Remarking that $n\binom{n-1}{k-1} = k\binom{n}{k}$ (for all $1 \le k \le n$), it follows that for all $n \in \mathbb{N}_0$:

$$q^{X}nT_{n-1}(X) = (q-1)^{-n} \sum_{k=0}^{n} (-1)^{n-k} \binom{n}{k} \left((q-1)ka_{k-1} \right) q^{kX},$$

showing that the q-polynomial sequence $(q^X n T_{n-1}(X))_{n \in \mathbb{N}_0}$ is Carlitz-type with associated real sequence $((q-1)ka_{k-1})_{k \in \mathbb{N}_0}$. The first item of the proposition is proved.

• Next, let us show the 2nd item of the proposition. So, suppose that $T_0(X) = 0_{\mathscr{E}}$, that is $a_0 = 0$. Using (2.11), we have for all $n \in \mathbb{N}_0$:

$$q^{-X} \frac{T_{n+1}(X)}{n+1} = (q-1)^{-n-1} \sum_{k=1}^{n+1} (-1)^{n+1-k} \frac{1}{n+1} \binom{n+1}{k} a_k q^{(k-1)X}$$
$$= (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \frac{1}{n+1} \binom{n+1}{k+1} \frac{1}{q-1} a_{k+1} q^{kX}.$$

Remarking that $\frac{1}{n+1} \binom{n+1}{k+1} = \frac{1}{k+1} \binom{n}{k}$ (for all $0 \le k \le n$), it follows that for all $n \in \mathbb{N}_0$:

$$q^{-X}\frac{T_{n+1}(X)}{n+1} = (q-1)^{-n}\sum_{k=0}^{n} (-1)^{n-k} \binom{n}{k} \left(\frac{1}{q-1}\frac{a_{k+1}}{k+1}\right) q^{kX},$$

showing that the q-polynomial sequence $\left(q^{-X}\frac{T_{n+1}(X)}{n+1}\right)_{n\in\mathbb{N}_0}$ is Carlitz-type with associated real sequence $\left(\frac{1}{q-1}\frac{a_{k+1}}{k+1}\right)_{k\in\mathbb{N}_0}$. The second item of the proposition is proved.

• Now, let us show the 3rd item of the proposition. Using (2.11), we have for all $n \in \mathbb{N}_0$:

$$q^{-X} \left(T_n(X) + (q-1)T_{n+1}(X) \right) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_k q^{(k-1)X} + (q-1)^{-n} \sum_{k=0}^{n+1} (-1)^{n+1-k} \binom{n+1}{k} a_k q^{(k-1)X} = (q-1)^{-n} \sum_{k=0}^{n+1} (-1)^{n-k+1} \left(\binom{n+1}{k} - \binom{n}{k} \right) a_k q^{(k-1)X}.$$

But since $\binom{n+1}{k} - \binom{n}{k} = \begin{cases} 0 & \text{if } k = 0\\ \binom{n}{k-1} & \text{if } k \ge 1 \end{cases}$ (for all $0 \le k \le n+1$), it follows that:

$$q^{-X} \left(T_n(X) + (q-1)T_{n+1}(X) \right) = (q-1)^{-n} \sum_{k=1}^{n+1} (-1)^{n-k+1} \binom{n}{k-1} a_k q^{(k-1)X}$$
$$= (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} a_{k+1} q^{kX},$$

showing that the sequence $(q^{-X}(T_n(X) + (q-1)T_{n+1}(X)))_{n \in \mathbb{N}_0}$ is q-polynomial of Carlitz-type with associated real sequence $(a_{k+1})_{k \in \mathbb{N}_0}$. The third item of the proposition is proved.

• Finally, the 4^{th} item of the proposition follows by successively applying its 3^{rd} and 1^{st} items, which have already been proven. This completes the proof.

Example 2.12. Since the real sequence associated to $(\eta_n(X))_{n\in\mathbb{N}_0}$ (which is a Carlitz-type q-polynomial sequence) is $\left(\frac{k}{[k]}\right)_{k\in\mathbb{N}_0}$ then (according to Item 3 of Proposition 2.11) the sequence $\left(q^{-X}\left(\eta_n(X)+(q-1)\eta_{n+1}(X)\right)\right)_{n\in\mathbb{N}_0}$ is a Carlitz-type q-polynomial sequence with associated real sequence $\left(\frac{k+1}{[k+1]}\right)_{k\in\mathbb{N}_0}$. Hence $\left(q^{-X}\left(\eta_n(X)+(q-1)\eta_{n+1}(X)\right)\right)$ coincides with $(\beta_n(X))_n$; that is

$$\beta_n(X) = q^{-X} \left(\eta_n(X) + (q-1)\eta_{n+1}(X) \right) \qquad (\forall n \in \mathbb{N}_0).$$
(2.12)

We now derive from Theorem 2.7 formulas for q-analogues of sums of powers of positive integers, which involve the q-polynomial sequences $(\eta_n(X))_{n \in \mathbb{N}_0}$ and $(\beta_n(X))_{n \in \mathbb{N}_0}$ introduced by Carlitz in [1]. The following corollaries are obtained:

Corollary 2.13. For all positive integers n and N, we have

$$\sum_{k=0}^{N-1} q^k [k]^{n-1} = \frac{\eta_n(N) - \eta_n}{n}$$

Proof. Take in Theorem 2.7: r = d = 1.

Corollary 2.14. For all positive integers n and N, we have

$$n\sum_{k=0}^{N-1} q^{2k} [k]^{n-1} + (q-1)\sum_{k=0}^{N-1} q^k [k]^n = q^N \beta_n(N) - \beta_n.$$

Proof. Let n and N be fixed positive integers. By applying Theorem 2.7 for the couples (r, d) = (2, 1) and (r, d) = (1, 0), we respectively obtain the two following formulas:

$$\sum_{k=0}^{N-1} q^{2k} [k]^{n-1} = \frac{q^N T_n(N) - T_n(0)}{n},$$
$$\sum_{k=0}^{N-1} q^k [k]^n = q^N S_n(N) - S_n(0),$$

where $(T_k(X))_{k \in \mathbb{N}_0}$ and $(S_k(X))_{k \in \mathbb{N}_0}$ represent the Carlitz-type *q*-polynomial sequences, associated respectively to the real sequences $\left(\frac{k}{[k+1]}\right)_{k \in \mathbb{N}_0}$ and $\left(\frac{1}{q-1}\frac{1}{[k+1]}\right)_{k \in \mathbb{N}_0}$. By combining the two above formulas, we derive that:

$$n\sum_{k=0}^{N-1} q^{2k} [k]^{n-1} + (q-1)\sum_{k=0}^{N-1} q^k [k]^n = q^N U_n(N) - U_n(0),$$

where

$$U_k(X) := T_k(X) + (q-1)S_k(X) \qquad (\forall k \in \mathbb{N}_0).$$

Next, from the fact that $(T_k(X))_k$ and $(S_k(X))_k$ are Carlitz-type *q*-polynomial sequences whose associated real sequences are respectively $\left(\frac{k}{[k+1]}\right)_k$ and $\left(\frac{1}{q-1}\frac{1}{[k+1]}\right)_k$, we derive that $(U_k(X))_k$ is also a Carlitz-type *q*-polynomial sequence, with an associated real sequence having the general term:

$$\frac{k}{[k+1]} + (q-1)\left(\frac{1}{q-1}\frac{1}{[k+1]}\right) = \frac{k+1}{[k+1]} \qquad (\forall k \in \mathbb{N}_0)$$

Consequently, $(U_k(X))_k$ is nothing else than the *q*-polynomial sequence $(\beta_k(X))_k$ of Carlitz *q*-Bernoulli polynomials. This completes the proof.

2.2 Extended Carlitz *q*-Bernoulli numbers and polynomials

2.2.1 Definitions and basic results

We begin by introducing the following definition:

Definition 2.15. We define $\left(\beta_n^{(r)}(X)\right)_{r,n\in\mathbb{N}_0}$ recursively by:

$$\beta_n^{(0)}(X) := \eta_n(X) \qquad (\forall n \in \mathbb{N}_0), \qquad (2.13)$$

$$\beta_n^{(r+1)}(X) := q^{-X} \left(\beta_n^{(r)}(X) + (q-1)\beta_{n+1}^{(r)}(X) \right) \qquad (\forall r, n \in \mathbb{N}_0).$$
(2.14)

Besides, we define $\left(\beta_n^{(r)}\right)_{r,n\in\mathbb{N}_0}$ by:

$$\beta_n^{(r)} := \beta_n^{(r)}(0) \qquad (\forall r, n \in \mathbb{N}_0).$$

$$(2.15)$$

So, it follows from (2.13) and (2.14) that:

$$\beta_n^{(0)} = \eta_n \qquad (\forall n \in \mathbb{N}_0), \qquad (2.16)$$

$$\beta_n^{(r+1)} = \beta_n^{(r)} + (q-1)\beta_{n+1}^{(r)} \qquad (\forall r, n \in \mathbb{N}_0).$$
(2.17)

Since $\left(\beta_n^{(0)}(X)\right)_{n\in\mathbb{N}_0} = (\eta_n(X))_{n\in\mathbb{N}_0}$ is a Carlitz-type *q*-polynomial sequence (see Examples 2.5) then (according to Item 3 of Proposition 2.11 and a simple induction on *r*) all the sequences $\left(\beta_n^{(r)}(X)\right)_{n\in\mathbb{N}_0}$ ($r\in\mathbb{N}_0$) are Carlitz-type *q*-polynomial sequences.

Definition 2.16. Giving $r \in \mathbb{N}_0$, the q-polynomials $\beta_n^{(r)}(X)$ $(n \in \mathbb{N}_0)$ are called the extended Carlitz q-Bernoulli polynomials of order r, and the numbers $\beta_n^{(r)}$ $(n \in \mathbb{N}_0)$ are called the extended Carlitz q-Bernoulli numbers of order r.

Taking r = 0 in (2.14), we find in particular that for all $n \in \mathbb{N}_0$:

$$\beta_n^{(1)}(X) = q^{-X} \left(\eta_n(X) + (q-1)\eta_{n+1}(X) \right) = \beta_n(X)$$

(according to (2.12)). So, the Carlitz q-Bernoulli polynomials of order 1 are nothing else than the Carlitz q-Bernoulli polynomials $\beta_n(X)$ (introduced by Carlitz in [1]). Consequently, the Carlitz q-Bernoulli numbers of order 1 are nothing else than the Carlitz q-Bernoulli numbers β_n of [1]. Furthermore, Carlitz [1] showed that the q-polynomials $\beta_n(X) = \beta_n^{(1)}(X)$ ($n \in \mathbb{N}_0$) are well-defined for q = 1 and coincide with the Bernoulli polynomials when specializing q to 1; that is $\beta_n^{(1)}(X)|_{q=1} = B_n(X)$ for all $n \in \mathbb{N}_0$. Consequently, in view of (2.14), the q-polynomials $\beta_n^{(r)}(X)$ ($r \in \mathbb{N}, n \in \mathbb{N}_0$) are also well-defined for q = 1, and for all $r \in \mathbb{N}$ and $n \in \mathbb{N}_0$, we have:

$$\beta_n^{(r+1)}(X)|_{q=1} = \beta_n^{(r)}(X)|_{q=1}.$$

By induction on r, it follows that for all $r \in \mathbb{N}$ and $n \in \mathbb{N}_0$:

$$\beta_n^{(r)}(X)|_{q=1} = \beta_n^{(1)}(X)|_{q=1} = B_n(X).$$

This demonstrates that for all $r \in \mathbb{N}$, the *q*-polynomials $\beta_n^{(r)}(X)$ $(n \in \mathbb{N}_0)$ can serve as a *q*-analogue of the classical Bernoulli polynomials. In view of (2.15), it follows that for all $r \in \mathbb{N}$, the numbers $\beta_n^{(r)}$ $(n \in \mathbb{N}_0)$ can also serve as a *q*-analogue of the classical Bernoulli numbers. This justify the terminology introduced in Definition 2.16.

The following proposition gathers several formulas for the Carlitz q-Bernoulli polynomials and numbers of a given order, which can be directly derived from the results of §2.1.

Proposition 2.17. For all $r, n \in \mathbb{N}_0$, we have that:

$$\beta_n^{(r)}(X) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{k+r}{[k+r]} q^{kX}, \qquad (2.18)$$

$$\beta_n^{(r)} = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{k+r}{[k+r]},$$
(2.19)

$$\beta_n^{(r)}(X) = \sum_{k=0}^n \binom{n}{k} \beta_k^{(r)} q^{kX} [X]^{n-k}.$$
(2.20)

In addition, for a given $r \in \mathbb{N}_0$, the Carlitz q-Bernoulli polynomials and numbers of order r satisfy the following symbolic formulas:

$$q^{r} (q\beta^{(r)} + 1)^{n} = (\beta^{(r)})^{n} \qquad (\forall n \ge 2),$$
 (2.21)

$$\beta_n^{(r)}(X) = \left(q^X \beta^{(r)} + [X]\right)^n \qquad (\forall n \in \mathbb{N}_0).$$
(2.22)

Proof. Using Item 3 of Proposition 2.11 together with Formula (2.14), we immediately show by induction that for all $r \in \mathbb{N}_0$, the associated real sequence of the Carlitz-type q-polynomial sequence $\left(\beta_n^{(r)}(X)\right)_{n\in\mathbb{N}_0}$ is $\left(\frac{k+r}{[k+r]}\right)_{k\in\mathbb{N}_0}$. Formula (2.18) of the proposition then follows. Next, Formula (2.19) of the proposition follows by taking X = 0 in Formula (2.18), which we have just proven. Formulas (2.20) and (2.22) of the proposition immediately follow from Theorem 2.2. Finally, Formula (2.21) of the proposition immediately follows from Theorem 2.9. This completes this proof.

We now see, for a general $r \in \mathbb{N}_0$, how the *q*-polynomials $\beta_n^{(r)}(X)$ arise in *q*-analogues of sums of powers of consecutive positive integers. We have the following theorem providing a generalization of Corollaries 2.13 and 2.14.

Theorem 2.18. Let r be a nonnegative integer. Then we have for all positive integers n and N:

$$n\sum_{k=0}^{N-1} q^{(r+1)k} [k]^{n-1} + (q-1)r\sum_{k=0}^{N-1} q^{rk} [k]^n = q^{rN} \beta_n^{(r)}(N) - \beta_n^{(r)}.$$

Proof. Let n and N be fixed positive integers. By applying Theorem 2.7 for the couples (r, 0) and (r + 1, 1) (instead of (r, d)), we respectively obtain the two following formulas:

$$\sum_{k=0}^{N-1} q^{rk} [k]^n = q^{rN} T_n(N) - T_n(0),$$
$$\sum_{k=0}^{N-1} q^{(r+1)k} [k]^{n-1} = \frac{q^{rN} S_n(N) - S_n(0)}{n},$$

where $(T_k(X))_{k\in\mathbb{N}_0}$ and $(S_k(X))_{k\in\mathbb{N}_0}$ are the Carlitz-type *q*-polynomial sequences associated respectively to the real sequences of general term:

$$a_k = \frac{1}{q-1} \frac{1}{[k+r]}$$
 and $b_k = \frac{k}{[k+r]}$ $(\forall k \in \mathbb{N}_0).$

By combining the above formulas, we derive that:

$$n\sum_{k=0}^{N-1} q^{(r+1)k}[k]^{n-1} + (q-1)r\sum_{k=0}^{N-1} q^{rk}[k]^n = q^{rN}U_n(N) - U_n(0),$$

where

$$U_k(X) := S_k(X) + (q-1)rT_k(X) \qquad (\forall k \in \mathbb{N}_0).$$

On the other hand, from the fact that $(T_k(X))_k$ and $(S_k(X))_k$ are Carlitz-type *q*-polynomial sequences whose associated real sequences are respectively $\left(\frac{1}{q-1}\frac{1}{[k+r]}\right)_k$ and $\left(\frac{k}{[k+r]}\right)_k$, we derive that $(U_k(X))_k$ is also a Carlitz-type *q*-polynomial sequence, with an associated real sequence having the general term:

$$\frac{k}{[k+r]} + (q-1)r\left(\frac{1}{q-1}\frac{1}{[k+r]}\right) = \frac{k+r}{[k+r]} \qquad (\forall k \in \mathbb{N}_0).$$

Hence $(U_k(X))_k$ coincides with $(\beta_k^{(r)}(X))_k$, concluding to the required result.

2.2.2 Representation of the $\beta_n^{(r)}$'s as series

We now turn our attention to the representation of the extended Carlitz q-Bernoulli numbers $\beta_n^{(r)}$ as numerical series. We have the following theorem:

Theorem 2.19. Suppose that |q| < 1. Then we have for all $r \in \mathbb{N}_0$ and $n \in \mathbb{N}$:

$$\beta_n^{(r)} = \begin{cases} -n \sum_{\substack{k=0 \\ +\infty}}^{+\infty} q^k [k]^{n-1} + (1-q)^{-n} & \text{if } r = 0\\ -n \sum_{\substack{k=0 \\ k=0}}^{+\infty} q^{(r+1)k} [k]^{n-1} + r(1-q) \sum_{\substack{k=0 \\ k=0}}^{+\infty} q^{rk} [k]^n & \text{else} \end{cases}$$

Proof. The result follows by letting $N \to +\infty$ in the formula of Theorem 2.18. However, before doing so, we must evaluate $\lim_{N\to+\infty} q^{rN}\beta_n^{(r)}(N)$ (for given $r \in \mathbb{N}_0$, $n \in \mathbb{N}$). To achieve this, we use Formula (2.18) of Proposition 2.17. Let $r \in \mathbb{N}_0$ and $n, N \in \mathbb{N}$ be fixed. According to Formula (2.18) of Proposition 2.17, we have that:

$$\beta_n^{(r)}(N) = (q-1)^{-n} \sum_{k=0}^n (-1)^{n-k} \binom{n}{k} \frac{k+r}{[k+r]} q^{kN}$$
$$= (q-1)^{-n} \left[(-1)^n \frac{r}{[r]} + \sum_{k=1}^n (-1)^{n-k} \binom{n}{k} \frac{k+r}{[k+r]} q^{kN} \right]$$

Since |q| < 1 by hypothesis, we derive that:

$$\lim_{N \to +\infty} \beta_n^{(r)}(N) = (1-q)^{-n} \frac{r}{[r]}.$$

Thus

$$\lim_{N \to +\infty} q^{rN} \beta_n^{(r)}(N) = \begin{cases} \lim_{N \to +\infty} \beta_n^{(r)}(N) = (1-q)^{-n} & \text{if } r = 0\\ 0 & \text{else} \end{cases}$$

Finally, by letting $N \to +\infty$ in the formula of Theorem 2.18 and incorporating the result we just obtained, we get

$$n\sum_{k=0}^{+\infty} q^{(r+1)k} [k]^{n-1} + (q-1)r\sum_{k=0}^{+\infty} q^{rk} [k]^n = \begin{cases} (1-q)^{-n} - \beta_n^{(r)} & \text{if } r = 0\\ -\beta_n^{(r)} & \text{else} \end{cases}$$

implying the desired result of the theorem and completes the proof.

Remark 2.20. For $r \in \mathbb{N}$, the formula of Theorem 2.19 can be written more simply as:

$$\beta_n^{(r)} = \sum_{k=0}^{+\infty} \left(rq^{rk} - (n+r)q^{(r+1)k} \right) [k]^{n-1}.$$
(2.23)

Taking in particular r = 1, we derive the following important representation of the Carlitz q-Bernoulli numbers as series:

$$\beta_n = \sum_{k=0}^{+\infty} \left(q^k - (n+1)q^{2k} \right) \left[k \right]^{n-1}$$
(2.24)

(valid for |q| < 1 and $n \in \mathbb{N}$).

From Remark 2.20 and the fact that the extended Carlitz q-Bernoulli numbers of any positive order are q-analogues of the classical Bernoulli numbers, we immediately derive the following corollary:

Corollary 2.21. For all $r, n \in \mathbb{N}$, we have that

$$B_n = \lim_{\substack{q \to 1 \\ q \to 1}} \sum_{k=0}^{+\infty} \left(rq^{rk} - (n+r)q^{(r+1)k} \right) [k]^{n-1}.$$

Remark 2.22. If, in the formula of Corollary 2.21, we allow the limit to be taken inside the sum and also permit the manipulation of divergent series, we obtain, for all $n \in \mathbb{N}$:

$$B_n = \sum_{k=0}^{+\infty} -nk^{n-1} = -n\sum_{k=1}^{+\infty} k^{n-1} = -n\zeta(1-n).$$

This provides a well-known and valid formula for the classical Bernoulli numbers, where ζ denotes the Riemann zeta function.

2.2.3 Expression of the $\beta_n^{(r)}$'s in terms of q-Stirling numbers of the second kind

The following theorem provides a generalization of Formula (1.8) of Carlitz, which express the Carlitz *q*-Bernoulli numbers in terms of *q*-Stirling numbers of the second kind.

Theorem 2.23. For all $r \in \mathbb{N}$ and $n \in \mathbb{N}_0$, we have

$$\beta_n^{(r)} = \sum_{k=0}^n \varphi_r(q,k) S_q(n,k),$$

where

$$\varphi_r(q,k) = (-1)^k [k]! \sum_{i=0}^{r-1} q^{ik} \frac{[r-1]_i}{[k+r]_{i+1}}$$
$$= (-1)^k [k]! [r-1]! q^{-r+1} \sum_{i=0}^{r-1} \frac{(q-1)^i q^{\frac{1}{2}i(i+1)-i(r-1)}}{[r-1-i]!} \frac{1}{[k+i+1]}$$

(for all $k \in \mathbb{N}$).

The proof of this theorem needs some preparations which are presented below. We begin with the following proposition:

Proposition 2.24. For all $r, N \in \mathbb{N}$ and $k \in \mathbb{N}_0$, we have

$$\sum_{i=0}^{N-1} q^{ir}[i]_k = \left(\sum_{\ell=0}^{r-1} q^{(r-\ell-1)N+(\ell+1)k} \frac{[r-1]_\ell}{[k+r]_{\ell+1}}\right) [N]_{k+1}.$$

Proof. Let $k \in \mathbb{N}_0$ and $N \in \mathbb{N}$ be fixed. For all $r \in \mathbb{N}$, we have

$$\begin{split} \Delta \left(q^{(r-1)X}[X]_{k+1} \right) &= q^{(r-1)(X+1)}[X+1]_{k+1} - q^{(r-1)X}[X]_{k+1} \\ &= q^{(r-1)(X+1)}[X+1][X] \cdots [X-k+1] - q^{(r-1)X}[X][X-1] \cdots [X-k] \\ &= \left(q^{(r-1)(X+1)}[X+1] - q^{(r-1)X}[X-k] \right) [X]_k. \end{split}$$

But since

$$\begin{split} q^{(r-1)(X+1)}[X+1] - q^{(r-1)X}[X-k] &= q^{(r-1)(X+1)} \frac{q^{X+1} - 1}{q-1} - q^{(r-1)X} \frac{q^{X-k} - 1}{q-1} \\ &= \frac{q^{r(X+1)} - q^{(r-1)(X+1)} - q^{rX-k} + q^{(r-1)X}}{q-1} \\ &= \frac{q^{rX-k} \left(q^{k+r} - 1\right) - q^{(r-1)X} \left(q^{r-1} - 1\right)}{q-1} \\ &= q^{rX-k} [k+r] - q^{(r-1)X} [r-1], \end{split}$$

it follows that:

$$\Delta\left(q^{(r-1)X}[X]_{k+1}\right) = \left(q^{rX-k}[k+r] - q^{(r-1)X}[r-1]\right)[X]_k.$$

By multiplying both sides of this last equality by $\frac{q^k}{[k+r]}$ and then summing both sides of the resulting equality from X = 0 to N - 1, we get (after simplifying and rearranging)

$$\sum_{X=0}^{N-1} q^{rX} [X]_k = \frac{q^{(r-1)N+k}}{[k+r]} [N]_{k+1} + q^k \frac{[r-1]}{[k+r]} \sum_{X=0}^{N-1} q^{(r-1)X} [X]_k.$$

By reiterating this last formula r times, we get

$$\begin{split} \sum_{X=0}^{N-1} q^{rX} [X]_k &= q^{(r-1)N+k} \frac{1}{[k+r]} [N]_{k+1} + q^{(r-2)N+2k} \frac{[r-1]}{[k+r][k+r-1]} [N]_{k+1} \\ &+ q^{(r-3)N+3k} \frac{[r-1]_2}{[k+r]_3} [N]_{k+1} + \dots + q^{rk} \frac{[r-1]_{r-1}}{[k+r]_r} [N]_{k+1} + q^{rk} \frac{[r-1]_r}{[k+r]_r} \sum_{X=0}^{N-1} [X]_k \\ &= \left(\sum_{\ell=0}^{r-1} q^{(r-\ell-1)N+(\ell+1)k} \frac{[r-1]_\ell}{[k+r]_{\ell+1}} \right) [N]_{k+1} \qquad (\text{since } [r-1]_r = 0). \end{split}$$

The proposition is proved.

In what follows, for $r, N \in \mathbb{N}$ and $k \in \mathbb{N}_0$, we set

$$f_r(q,k,N) := q^{\frac{k(k-1)}{2}} \sum_{i=0}^{N-1} q^{ir}[i]_k,$$

which is q-polynomial in N (according to Proposition 2.24). We let also $f_r(q, k, X)$ denote the unique q-polynomial interpolating $f_r(q, k, N)$ at the positive integers X = N, and $g_r(q, k)$ denote the limit:

$$g_r(q,k) := \lim_{X \to 0} \frac{1}{[X]} f_r(q,k,X).$$

A first useful expression for $g_r(q, k)$ is directly derived from Proposition 2.24. We have the following corollary:

Corollary 2.25. For all $r \in \mathbb{N}$ and $k \in \mathbb{N}_0$, we have

$$g_r(q,k) = (-1)^k [k]! \left(\sum_{i=0}^{r-1} q^{ik} \frac{[r-1]_i}{[k+r]_{i+1}} \right).$$

Proof. Let $r \in \mathbb{N}$ and $k \in \mathbb{N}_0$ be fixed. According to Proposition 2.24, we have for all $N \in \mathbb{N}$:

$$f_r(q,k,N) = q^{\frac{k(k-1)}{2}} \left(\sum_{\ell=0}^{r-1} q^{(r-\ell-1)N+(\ell+1)k} \frac{[r-1]_{\ell}}{[k+r]_{\ell+1}} \right) [N]_{k+1}.$$

Hence

$$f_r(q,k,X) = q^{\frac{k(k-1)}{2}} \left(\sum_{\ell=0}^{r-1} q^{(r-\ell-1)X+(\ell+1)k} \frac{[r-1]_{\ell}}{[k+r]_{\ell+1}} \right) [X]_{k+1}.$$

Consequently, we have

$$g_r(q,k) := \lim_{X \to 0} \frac{1}{[X]} f_r(q,k,X)$$

$$= \lim_{X \to 0} q^{\frac{k(k-1)}{2}} \left(\sum_{\ell=0}^{r-1} q^{(r-\ell-1)X+(\ell+1)k} \frac{[r-1]_{\ell}}{[k+r]_{\ell+1}} \right) [X-1]_k$$

$$= q^{\frac{k(k+1)}{2}} \left(\sum_{\ell=0}^{r-1} q^{\ell k} \frac{[r-1]_{\ell}}{[k+r]_{\ell+1}} \right) [-1]_k$$

$$= (-1)^k [k]! \sum_{\ell=0}^{r-1} q^{\ell k} \frac{[r-1]_{\ell}}{[k+r]_{\ell+1}}$$

(since $[-1]_k = (-1)^k q^{-\frac{k(k+1)}{2}}[k]!$). The corollary is proved.

An equally interesting formula for $g_r(q, k)$ $(r \in \mathbb{N}, k \in \mathbb{N}_0)$ can be derived by decomposing the q-rational function $\frac{q^{iX}}{[X]_{i+1}}$ $(0 \le i \le r-1)$ into q-partial fractions. This decomposition is given by the following proposition:

Proposition 2.26. For all $k \in \mathbb{N}_0$, we have

$$\frac{q^{kX}}{[X]_{k+1}} = \frac{1}{[k]!} \sum_{i=0}^{k} \left((-1)^{k-i} {k \brack i} q^{\frac{k^2+k}{2} + \frac{i^2-i}{2}} \frac{1}{[X-i]} \right).$$

Proof. Let $k \in \mathbb{N}_0$ be fixed. We will first establish the existence and uniqueness of real numbers $\alpha_{k,i}(q)$ $(0 \le i \le k)$ such that the following holds:

$$\frac{q^{kX}}{[X]_{k+1}} = \sum_{i=0}^{k} \frac{\alpha_{k,i}(q)}{[X-i]}.$$
(2.25)

We have:

$$(2.25) \iff q^{kX} = \sum_{i=0}^{k} \alpha_{k,i}(q) \frac{[X]_{k+1}}{[X-i]}$$
$$\iff q^{kX} = \sum_{i=0}^{k} \alpha_{k,i}(q) \left(\prod_{\substack{0 \le j \le k \\ j \ne i}} [X-j]\right).$$
(2.26)

Consider the family \mathscr{B} of (k+1) q-polynomials of degree k given by:

$$\mathscr{B} := \left\{ \prod_{\substack{0 \le j \le k \\ j \ne i}} [X - j] \right\}_{0 \le i \le k}$$

It can be observed that any k arbitrary q-polynomials from \mathscr{B} always share a common root that is not a root of the remaining q-polynomial in \mathscr{B} . Therefore, no q-polynomial in \mathscr{B} can be expressed as a linear combination of the others, which shows that \mathscr{B} is a linearly independent family in \mathscr{E}_k . Consequently, since $\operatorname{Card}\mathscr{B} = \dim \mathscr{E}_k = k + 1$, \mathscr{B} is a basis of \mathscr{E}_k . It follows that the q-polynomial q^{kX} (belonging to \mathscr{E}_k) can be uniquely expressed as a linear combination of the q-polynomials in \mathscr{B} . This establishes the existence and uniqueness of real numbers $\alpha_{k,i}(q)$ $(0 \le i \le k)$ satisfying (2.26) (equivalently (2.25)).

Let us now determine the real numbers $\alpha_{k,i}(q)$ $(0 \le i \le k)$. For a given $\ell \in \{0, 1, \ldots, k\}$, by multiplying both sides of (2.25) by $[X - \ell]$ and then substituting $X = \ell$ into the resulting identity, we obtain:

$$\frac{q^{k\ell}}{[\ell][\ell-1]\cdots[1]\cdot[-1][-2]\cdots[\ell-k]} = \alpha_{k,\ell}(q).$$

But since $[\ell][\ell-1]\cdots[1] = [\ell]!$ and $[-1][-2]\cdots[\ell-k] = (-1)^{k-\ell}q^{-\frac{(k-\ell)(k-\ell+1)}{2}}[k-\ell]!$ (using the property $[-X] = -q^{-X}[X]$), it follows that:

$$\alpha_{k,\ell}(q) = \frac{(-1)^{k-\ell} q^{\frac{(k-\ell)(k-\ell+1)}{2}+k\ell}}{[\ell]![k-\ell]!}$$
$$= \frac{1}{[k]!} (-1)^{k-\ell} {k \brack \ell} q^{\frac{k^2+k}{2}+\frac{\ell^2-\ell}{2}}$$

as required. This completes the proof.

Using the formula from Proposition 2.26, the formula of Corollary 2.25 for $g_r(q, k)$ can be transformed into another, more interesting form for studying the limiting case as $q \to 1$. We have the following proposition:

Proposition 2.27. For all $r \in \mathbb{N}$ and $k \in \mathbb{N}_0$, we have

$$g_r(q,k) = (-1)^k [k]! [r-1]! q^{-r+1} \sum_{i=0}^{r-1} \frac{(q-1)^i q^{\frac{1}{2}i(i+1)-i(r-1)}}{[r-1-i]!} \frac{1}{[k+i+1]!}$$

Proof. Let $r \in \mathbb{N}$ and $k \in \mathbb{N}_0$ be fixed. By successively applying Corollary 2.25 and Proposition 2.26, we obtain

$$g_{r}(q,k) = (-1)^{k}[k]! \sum_{i=0}^{r-1} q^{-ir}[r-1]_{i} \left\{ \frac{q^{i(k+r)}}{[k+r]_{i+1}} \right\}$$
$$= (-1)^{k}[k]! \sum_{i=0}^{r-1} q^{-ir}[r-1]_{i} \frac{1}{[i]!} \sum_{j=0}^{i} (-1)^{i-j} {i \brack j} q^{\frac{i^{2}+i}{2} + \frac{j^{2}-j}{2}} \frac{1}{[k+r-j]}$$
$$= (-1)^{k}[k]! \sum_{\substack{0 \le i \le r-1\\0 \le j \le i}} q^{-ir} \frac{[r-1]_{i}}{[i]!} {i \brack j} (-1)^{i-j} q^{\frac{i^{2}+i}{2} + \frac{j^{2}-j}{2}} \frac{1}{[k+r-j]}.$$

By inverting the summations over i and j and noting that $\frac{[r-1]_i}{[i]!} {i \brack j} = {r-1 \brack j} {r-1-j \brack i-j}$ and that $q^{-ir} \cdot q^{\frac{i^2+i}{2} + \frac{j^2-j}{2}} = q^{\frac{(i-j)^2+(i-j)}{2} - i(r-j)}$ (for all $0 \le i \le r-1$ and $0 \le j \le i$), we derive that:

$$g_r(q,k) = (-1)^k [k]! \sum_{0 \le j \le r-1} \left(\sum_{i=j}^{r-1} {r-1-j \brack i-j} (-1)^{i-j} q^{\frac{(i-j)^2 + (i-j)}{2} - i(r-j)} \right) {r-1 \brack j} \frac{1}{[k+r-j]}.$$

Then, performing the change of index $\ell = i - j$ in the inner summation gives

$$g_r(q,k) = (-1)^k [k]! \sum_{0 \le j \le r-1} \left(\sum_{\ell=0}^{r-1-j} q^{\frac{\ell(\ell-1)}{2}} {r-1-j \choose \ell} \left(-q^{1-r+j} \right)^\ell \right) q^{j^2-jr} {r-1 \choose j} \frac{1}{[k+r-j]}$$

But according to the Gauss binomial formula (1.2), we have for all $j \in \{0, 1, ..., r-1\}$:

$$\sum_{\ell=0}^{r-1-j} q^{\frac{\ell(\ell-1)}{2}} {r-1-j \choose \ell} \left(-q^{1-r+j}\right)^{\ell} = \left(1-q^{1-r+j}\right) \left(1-q^{2-r+j}\right) \cdots \left(1-q^{-1}\right)$$
$$= q^{1-r+j} \left(q^{r-1-j}-1\right) \cdot q^{2-r+j} \left(q^{r-2-j}-1\right) \cdots q^{-1}(q-1)$$
$$= q^{-\frac{(r-1-j)(r-j)}{2}} (q-1)^{r-1-j} [r-1-j]!.$$

Substituting this into the last obtained expression for $g_r(q, k)$, we get

$$g_r(q,k) = (-1)^k [k]! \sum_{j=0}^{r-1} q^{-\frac{(r-1-j)(r-j)}{2}} (q-1)^{r-1-j} [r-1-j]! q^{j^2-jr} {r-1 \brack j} \frac{1}{[k+r-j]}.$$

Finally, changing the index i = r - 1 - j and rearranging yields

$$g_r(q,k) = (-1)^k [k]! [r-1]! q^{-r+1} \sum_{i=0}^{r-1} \frac{(q-1)^i q^{\frac{1}{2}i(i+1)-i(r-1)}}{[r-1-i]!} \frac{1}{[k+i+1]},$$

as required.

We are now ready to prove Theorem 2.23.

Proof of Theorem 2.23. Let $r \in \mathbb{N}$ and $n \in \mathbb{N}_0$ be fixed. In view of Corollary 2.25 and Proposition 2.27, we need to show that:

$$\beta_n^{(r)} = \sum_{k=0}^n g_r(q,k) S_q(n,k).$$

To achieve this, consider $N \in \mathbb{N}$ and evaluate the sum

$$\sum_{i=0}^{N-1} q^{ri} [i]^n$$

in two different ways. On the one hand, from Theorem 2.6, we have

$$\sum_{i=0}^{N-1} q^{ri}[i]^n = q^{rN} S_n(N) - S_n(0),$$

_	

where $(S_k(X))_{k\in\mathbb{N}_0}$ is the Carlitz-type q-polynomial sequence associated to the real sequence of general term $a_k := \frac{1}{q-1} \frac{1}{[k+r]}$ ($\forall k \in \mathbb{N}_0$). So, putting $s_k := S_k(0)$ ($\forall k \in \mathbb{N}_0$) and using the formula of Theorem 2.2 for $S_k(X)$, we derive that:

$$\sum_{i=0}^{N-1} q^{ri}[i]^n = q^{rN} \sum_{k=0}^n \binom{n}{k} s_k q^{kN} [N]^{n-k} - s_n$$
$$= \sum_{k=0}^{n-1} \binom{n}{k} s_k q^{(k+r)N} [N]^{n-k} + (q^{(n+r)N} - 1) s_n.$$
(2.27)

On the other hand, by using Formula (1.7), we have that:

$$\sum_{i=0}^{N-1} q^{ri}[i]^n = \sum_{i=0}^{N-1} q^{ri} \left(\sum_{k=0}^n q^{\frac{1}{2}k(k-1)} S_q(n,k)[i]_k \right)$$
$$= \sum_{k=0}^n S_q(n,k) \left(q^{\frac{1}{2}k(k-1)} \sum_{i=0}^{N-1} q^{ri}[i]_k \right)$$
$$= \sum_{k=0}^n S_q(n,k) f_r(q,k,N).$$
(2.28)

Comparing (2.27) and (2.28), we derive the identity:

$$\sum_{k=0}^{n-1} \binom{n}{k} s_k q^{(k+r)N} [N]^{n-k} + \left(q^{(n+r)N} - 1\right) s_n = \sum_{k=0}^n S_q(n,k) f_r(q,k,N).$$

Since this identity consists of q-polynomials in N and it is valid for all $N \in \mathbb{N}$ then it extends as a q-polynomial identity. Namely, we have

$$\sum_{k=0}^{n-1} \binom{n}{k} s_k q^{(k+r)X} [X]^{n-k} + \left(q^{(n+r)X} - 1\right) s_n = \sum_{k=0}^n S_q(n,k) f_r(q,k,X).$$

Dividing through by [X] and letting $X \to 0$, we derive

$$ns_{n-1} + (q-1)(n+r)s_n = \sum_{k=0}^n g_r(q,k)S_q(n,k).$$
(2.29)

Now, setting for all $k \in \mathbb{N}_0$:

$$T_k(X) := kS_{k-1}(X) + (q-1)(k+r)S_k(X)$$

= $(q-1)rS_k(X) + k(S_{k-1}(X) + (q-1)S_k(X)),$

it is clear that $ns_{n-1} + (q-1)(n+r)s_n = T_n(0)$. Next, the q-polynomial sequence $((q-1)rS_k(X))_{k\in\mathbb{N}_0}$ is clearly Carlitz-type with associated real sequence $((q-1)ra_k)_k = \left(\frac{r}{[k+r]}\right)_k$. Besides, according to Item 4 of Proposition 2.11, the q-polynomial sequence $\left(k\left(S_{k-1}(X) + (q-1)S_k(X)\right)\right)_{k\in\mathbb{N}_0}$ is also Carlitz-type with associated real sequence $((q-1)ka_k)_k = \left(\frac{k}{[k+r]}\right)_k$. Consequently, the q-polynomial sequence $(T_k(X))_{k\in\mathbb{N}_0}$ is Carlitz-type with associated real sequence $\left(\frac{r}{[k+r]}\right)_k + \left(\frac{k}{[k+r]}\right)_k = \left(\frac{k+r}{[k+r]}\right)_k$. Thus $(T_k(X))_k$ coincides with

 $(\beta_k^{(r)}(X))_k$, that is $T_k(X) = \beta_k^{(r)}(X)$ ($\forall k \in \mathbb{N}_0$). In particular, we have $ns_{n-1} + (q-1)(n+r)s_n = T_n(0) = \beta_n^{(r)}(0) = \beta_n^{(r)}$. Substituting this into (2.29) yields the required formula. This completes the proof.

From Theorem 2.23, we derive the following curious corollary:

Corollary 2.28. For all $n \in \mathbb{N}_0$, we have

$$\beta_n + q\beta_{n+1} = \sum_{k=0}^n (-1)^k \frac{[k]!}{[k+2]} S_q(n,k).$$

Proof. Let $n \in \mathbb{N}_0$ be fixed. According to Formula (2.17), we have

$$\beta_n^{(2)} = \beta_n^{(1)} + (q-1)\beta_{n+1}^{(1)} = \beta_n + (q-1)\beta_{n+1}$$

which gives $\beta_{n+1} = \frac{\beta_n^{(2)} - \beta_n}{q-1}$, and then

$$\beta_n + q\beta_{n+1} = \beta_n + q\left(\frac{\beta_n^{(2)} - \beta_n}{q - 1}\right) = \frac{q\beta_n^{(2)} - \beta_n^{(1)}}{q - 1}$$

It follows by using Theorem 2.23 that

$$\beta_n + q\beta_{n+1} = \sum_{k=0}^n \frac{q\varphi_2(q,k) - \varphi_1(q,k)}{q-1} S_q(n,k).$$
(2.30)

Further, from the second expression of $\varphi_r(q,k)$ in Theorem 2.23, we have that

$$\varphi_1(q,k) = (-1)^k \frac{[k]!}{[k+1]}$$
 and $\varphi_2(q,k) = (-1)^k [k]! q^{-1} \left(\frac{1}{[k+1]} + \frac{q-1}{[k+2]}\right)$.

Thus

$$\frac{q\varphi_2(q,k) - \varphi_1(q,k)}{q-1} = (-1)^k \frac{[k]!}{[k+2]!}$$

Substituting this into (2.30) yields the required formula. This achieves the proof.

Letting $q \rightarrow 1$ in Corollary 2.28 gives the following result on the classical Bernoulli numbers, which is very recently pointed out by the author [4, Page 12, Formula (2.3)]:

Corollary 2.29. For all $n \in \mathbb{N}_0$, we have

$$B_n + B_{n+1} = \sum_{k=0}^n (-1)^k \frac{k!}{k+2} S(n,k).$$

References

 L. CARLITZ. q-Bernoulli numbers and polynomials, Duke Math. J., 15, n°4 (1948), p. 987-1000.

- [2] L. COMTET. Advanced Combinatorics. The Art of Finite and Infinite Expansions, revised and enlarged ed., D. Reidel Publ. Co., Dordrecht, 1974.
- F. A. COSTABILE. Modern umbral calculus: an elementary introduction with applications to linear interpolation and operator approximation theory, 72, Walter de Gruyter GmbH & Co KG, 2019.
- B. FARHI. New formulas involving Bernoulli and Stirling numbers of both kinds, preprint, 2024. Available at https://arxiv.org/pdf/2410.05207v2.
- [5] V. KAC & P. CHEUNG. Quantum Calculus, Universitext, Springer-Verlag, 2002.
- [6] N. KOBLITZ. On Carlitz's q-Bernoulli numbers, J. Number Theory, 14 (1982), p. 332-339.
- [7] O. KOUBA. Lecture notes, Bernoulli polynomials and numbers, preprint, 2016. Available at https://arxiv.org/pdf/1309.7560v2.
- [8] J. RIORDAN. Combinatorial Identities, John Wiley and Sons Inc., New York, 1968.
- [9] S. ROMAN. The umbral calculus, Academic Press, New York, 1984.