

THE DENSITY OF ZEROS OF RANDOM POWER SERIES WITH STATIONARY COMPLEX GAUSSIAN COEFFICIENTS

TOMOYUKI SHIRAI

ABSTRACT. We study the zeros of random power series with stationary complex Gaussian coefficients, whose spectral measure is absolutely continuous. We analyze the precise asymptotic behavior of the radial density of zeros near the boundary of the circle of convergence. The dependence of the coefficients generally reduces the density of zeros compared with that of the hyperbolic Gaussian analytic function (the i.i.d. coefficients case), where the spectral density and its zeros plays a crucial role in this reduction. We also show the relationship between the support of the spectral measure and the analytic continuation at the boundary of the circle of convergence.

1. INTRODUCTION

Gaussian analytic functions have been the subject of extensive research from various points of view over the years (cf. [11, 21, 22, 19, 14, 9, 17, 16, 20, 5, 7, 3, 6, 1, 12, 2, 8] and references therein). The case of random power series with i.i.d. coefficients has been extensively studied. In this paper, we focus on Gaussian power series with dependent coefficients under the following setting. Let $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$ be a stationary, centered, complex Gaussian process with covariance

$$\gamma(k - l) = \mathbb{E}[\xi_k \bar{\xi}_l],$$

where we always assume that $\gamma(0) = 1$, i.e., the distribution of each marginal ξ_k is complex standard normal. We consider the Gaussian power series with dependent coefficients $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$

$$X_\Xi(z) := \sum_{k=0}^{\infty} \xi_k z^k$$

and study its random zeros. The radius of convergence of $X_\Xi(z)$ is equal to 1 a.s. since $\lim_{k \rightarrow \infty} |\xi_k|^{1/k} = 1$ a.s. as long as $\gamma(0) = \mathbb{E}[|\xi_k|^2] = 1$ (cf. Lemma 6.1.) The zeros of $X_\Xi(z)$ form a point process $\mathcal{Z}_{X_\Xi} := \sum_{z \in \mathbb{D}: X_\Xi(z)=0} \delta_z$ on $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$. When $\gamma(k) = \delta_{k0}$, i.e., $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$ is an i.i.d. sequence, the Gaussian analytic function (GAF) is called the hyperbolic GAF, and its zero process is known to be the determinantal point process associated with the Bergman kernel $K_{\text{Berg}}(z, w) = (1 - z\bar{w})^{-2}$ and the background measure $\pi^{-1}m(dz)$ [19]. From this fact, it is easily seen that the expected number of zeros of the hyperbolic GAF within $\mathbb{D}_r = \{z \in \mathbb{C} : |z| < r\}$ is exactly equal to $r^2(1 - r^2)^{-1}$. In [15] the authors treated the case where the coefficient Gaussian process Ξ corresponds to a fractional Gaussian noise with Hurst index $0 \leq H < 1$ and gave an estimate for $\mathbb{E}[\mathcal{Z}_{X_\Xi}(\mathbb{D}_r)]$, showing that it is small by $O((1 - r^2)^{-1/2})$ compared with the hyperbolic GAF

Date: 2025/1/7.

2020 Mathematics Subject Classification. 30B20; 30C15; 60G15; 60G55.

case. In [18], the case where Ξ is finitely dependent, i.e., $\gamma(k)$ is a trigonometric polynomial, was considered, and the precise asymptotics of the expected number of zeros were analyzed. It is also shown that the dependence of the coefficients always reduces the expected number of zeros. Furthermore, the spectral density and its zeros of the coefficient Gaussian process Ξ , in particular, the degeneracy of zeros, plays a crucial role in determining the extent of this reduction. With the results of [18] in mind, this paper discusses the precise asymptotic behavior of the radial density of zeros as it approaches the boundary of the circle of convergence, i.e., $\partial\mathbb{D}$. Moreover, we discuss the analytic continuation of $X_\Xi(z)$ in relation to the strength of the dependence among the coefficients in Ξ . Intuitively, we will see that the stronger the dependence of Ξ , the easier the analytic continuation becomes. These observations provide deeper insights into the relationship between the spectral measure of the coefficients and the distributions of zeros.

The rest of the paper is organized as follows: In Section 2, we present the main results of the paper. Section 3 discusses the spectral representation of the coefficient Gaussian process Ξ and provides a spectral representation of the density of zeros of $X_\Xi(z)$. In Section 4, we refine the asymptotic behavior of the Poisson integral and its variant near the boundary of the unit disk. Section 5 contains the proof of the main Theorem 2.1. Finally, in Section 6, we discuss the analytic continuation of $X_\Xi(z)$ in terms of the spectral measure of Ξ .

2. MAIN RESULTS

The variance of $X_\Xi(z)$ in terms of the spectral measure $F(dt)$ of the coefficient Gaussian process Ξ is given by

$$K(z, z) := \mathbb{E}[|X_\Xi(z)|^2] = \frac{1}{1-r^2} \int_{-\pi}^{\pi} P_r(\varphi-t) F(dt),$$

where $z = re^{i\varphi}$ and $P_r(s)$ is the Poisson kernel

$$P_r(s) = \frac{1-r^2}{1-2r\cos s+r^2} \quad \text{for } s \in [-\pi, \pi] \text{ and } 0 \leq r < 1.$$

The covariance kernel of $X_\Xi(z)$ will be given in (3.3). We often identify $e^{i\varphi}$ with $\varphi \in [-\pi, \pi]$ and we write $f(\varphi)$ for $f(e^{i\varphi})$. If $F(dt)$ is absolutely continuous, i.e., $F(dt) = f(t)dt$, then we have the following representation:

$$K(z, z) = \frac{1}{1-|z|^2} \mathbb{E}_z[f(B_\tau)],$$

where B_t is the complex Brownian motion starting at $z \in \mathbb{D}$ and τ is the first hitting time to the boundary $\partial\mathbb{D}$. Thus, as z approaches the boundary $e^{i\varphi}$, i.e., $|z|$ approaches 1, the variance $K(z, z)$ diverges and $X_\Xi(z)$ oscillates very fast (and it tends to have more zeros) unless $f(e^{i\varphi})$ vanishes. For example, it is well-known that a Gaussian process $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$ is *purely non-deterministic* if and only if its spectral measure $F(ds)$ is absolutely continuous, i.e., $F(ds) = f(s)ds$ such that $\int_{-\pi}^{\pi} \log f(s)ds > -\infty$ [10, p.112]. In this case, $f(s) > 0$ a.e. on $\partial\mathbb{D}$, and hence $X_\Xi(z)$ oscillates infinitely often near the boundary $\partial\mathbb{D}$, and the boundary $\partial\mathbb{D}$ becomes the natural boundary (see Section 6).

In order to state the main result, we introduce some notations. The symmetrization and anti-symmetrization of h are defined by

$$\hat{h}(s) = \frac{h(s) + h(-s)}{2}, \quad \check{h}(s) = \frac{h(s) - h(-s)}{2}. \quad (2.1)$$

We define the operator T acting on symmetric functions smooth at $s = 0$ by

$$Th(s) := \begin{cases} \frac{h(s) - h(0)}{1 - \cos s} & \text{for } s \neq 0, \\ h''(0) & \text{for } s = 0. \end{cases}$$

and \mathcal{I} is the integral operator defined by

$$\mathcal{I}(h) = \frac{1}{2\pi} \int_{-\pi}^{\pi} h(s) ds.$$

The following is the first main theorem of this paper.

Theorem 2.1. *Let $F_{\varphi}(ds) = F(d(s+\varphi))$ on $(-\pi, \pi]$ modulo 2π . Suppose $F_{\varphi}(ds) = f_{\varphi}(s)ds$ and $f_{\varphi}(s)$ is smooth at $s = 0$.*

(i) *When $f_{\varphi}(0) > 0$,*

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)^2} - \frac{1}{4f_{\varphi}(0)^2} \left\{ f'_{\varphi}(0)^2 + \mathcal{I}(T\hat{f}_{\varphi})^2 \right\} + O(1-r^2).$$

(ii) *When $f_{\varphi}(0) = 0$,*

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)} \frac{f''_{\varphi}(0)}{2\mathcal{I}(T\hat{f}_{\varphi})} + O(1)$$

(iii) *When $f_{\varphi}(0) = f''_{\varphi}(0) = 0$. Then,*

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi} \frac{\mathcal{I}(T^2\hat{f}_{\varphi})^2 - \mathcal{I}(T^2(\hat{f}_{\varphi} \cos s))^2 - \mathcal{I}(T^2(\check{f}_{\varphi} \sin s))^2}{4\mathcal{I}(T\hat{f}_{\varphi})^2} + O(1-r^2).$$

When the spectral density vanishes at some points on the unit circle, it is observed that the order of the density of zeros in the direction approaching those points decreases according to the degree of the degeneracy. This is related to the phenomenon analyzed in [18], where the expected number of zeros is further reduced when the spectral density has zeros.

Remark 2.2. Since $\hat{f}_{\varphi}(s)$ is of $O(s^2)$ at $s = 0$ when $f_{\varphi}(0) = 0$, we have

$$\mathcal{I}(T\hat{f}_{\varphi}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{\hat{f}_{\varphi}(s)}{1 - \cos s} ds.$$

Note that $\hat{f}_{\varphi}(s)$, $\hat{f}_{\varphi}(s) \cos s$, $\check{f}_{\varphi}(s) \sin s$ are of $O(s^4)$ when $f_{\varphi}(0) = f''_{\varphi}(0) = 0$. For such a function g of $O(s^4)$ at $s = 0$, we have

$$\mathcal{I}(T^2g) = \frac{1}{2\pi} \int_{-\pi}^{\pi} \frac{g(s)}{(1 - \cos s)^2} ds.$$

Then, the numerator of the first term in (iii) can be expressed as

$$\frac{1}{4} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \frac{G_{\varphi}(s, t) + G_{\varphi}(s, -t)}{(1 - \cos s)^2 (1 - \cos t)^2} ds dt,$$

where $G_{\varphi}(s, t) := \{f_{\varphi}(s)f_{\varphi}(t) + f_{\varphi}(-s)f_{\varphi}(-t)\}\{1 - \cos(s-t)\}$, and it is obviously positive.

Example 2.3 (1-dependent case). We consider the case where the coefficient Gaussian process Ξ is 1-dependent, i.e., the covariance function is given by

$$\gamma(k) = \begin{cases} 1 & k = 0, \\ a & |k| = 1, \\ 0 & |k| \geq 2 \end{cases}$$

with $|a| \leq 1/2$. In this case, $f_\varphi(s) = 1 + 2a \cos(s + \varphi)$. Then,

$$f_\varphi(0) = 1 + 2a \cos \varphi, \quad f'_\varphi(0) = -2a \sin \varphi, \quad f''_\varphi(0) = -2a \cos \varphi,$$

and

$$\hat{f}_\varphi(s) = 1 + 2a \cos \varphi \cos s, \quad T\hat{f}_\varphi(s) = \frac{\hat{f}_\varphi(s) - \hat{f}_\varphi(0)}{1 - \cos s} = -2a \cos \varphi.$$

Note that $f_\varphi(0) = 0$ only if $(a, \varphi) = (1/2, \pi)$ or $(a, \varphi) = (-1/2, 0)$.

(i) When $|a| \leq 1/2$ and $(a, \varphi) \neq (1/2, \pi), (-1/2, 0)$, since $\mathcal{I}(T\hat{f}_\varphi) = -2a \cos \varphi$, we have

$$R(f_\varphi) := \frac{1}{4f_\varphi(0)^2} \left(f'_\varphi(0)^2 + \mathcal{I}(T\hat{f}_\varphi)^2 \right) = \frac{a^2}{(1 + 2a \cos \varphi)^2},$$

and then

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)^2} - \frac{a^2}{(1+2a \cos \varphi)^2} + O(1-r^2).$$

(ii) When $(a, \varphi) = (1/2, \pi)$ or $(a, \varphi) = (-1/2, 0)$, we have

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)^2} \frac{f''_\varphi(0)}{2\mathcal{I}(T\hat{f}_\varphi)} + O(1) = \frac{1}{2\pi(1-r^2)^2} + O(1).$$

Example 2.4. We consider the case $F(ds) = I_{[-\pi/2, \pi/2]}(s)ds$ and let $\{\xi_k\}_{k \in \mathbb{Z}}$ be the corresponding Gaussian process. The right panel of Fig. 1 shows the zeros of the approximate polynomial of $X_F(z) = \sum_{k=0}^{\infty} \xi_k z^k$. The zeros in the right half are distributed almost the same as in the left panel, while the zeros in the left half are neatly aligned near the unit circle. These correspond to the zeros that might disappear in the limit as $N \rightarrow \infty$, which is consistent with the fact that the density of the zeros on the left is $O(1)$.

Suppose $\pi/2 < \varphi < \pi$ and

$$f_\varphi(s) = \begin{cases} 1 & -\pi < s < -(\varphi - \pi/2), \\ 0 & -(\varphi - \pi/2) < s < 3\pi/2 - \varphi, \\ 1 & 3\pi/2 - \varphi < s < \pi. \end{cases}$$

Then, we see that

$$\hat{f}_\varphi(s) = \begin{cases} 0 & |s| < \varphi - \pi/2, \\ 1/2 & \varphi - \pi/2 < |s| < 3\pi/2 - \varphi, \\ 1 & 3\pi/2 - \varphi < |s| < \pi, \end{cases}$$

and

$$\check{f}_\varphi(s) = \begin{cases} 0 & |s| < \varphi - \pi/2, \\ -\text{sgn}(s)/2 & \varphi - \pi/2 < |s| < 3\pi/2 - \varphi, \\ 0 & 3\pi/2 - \varphi < |s| < \pi. \end{cases}$$




FIGURE 1. Left: The zeros of the approximate polynomial $X_{\text{hyp}}^{(400)}(z) = \sum_{k=0}^{400} \xi_k z^k$ for $F(dt) = dt/(2\pi)$, the i.i.d. case. Red points indicate zeros inside the unit disk, while blue points indicate zeros outside the unit disk. Right: The zeros of an approximate polynomial of $X_F(z)$ for $F(dt) = \mathbf{1}_{[-\pi/2, \pi/2]}(t)dt/\pi$.




FIGURE 2. The case $f(s) = I_{[-\pi/2, \pi/2]}(s)$ and $\varphi = 3\pi/4$. From left, $f_\varphi(s)$, $\hat{f}_\varphi(s)$, and $\check{f}_\varphi(s)$.

In this case, $f_\varphi(0) = f_\varphi''(0) = 0$. After some direct calculations for (iii) in Theorem 2.1 using Remark 2.2, we obtain the following: as $r \rightarrow 1$,

$$\rho_1(re^{i\varphi}) = \frac{1}{12\pi \cos^2 \varphi} + O(1-r^2) \quad \varphi \in (\pi/2, \pi].$$

This shows the behavior of zeros on the left-half plane.

Finally, we discuss analytic continuation of X_Ξ . For a power series $h = h(z)$, a point $re^{i\theta}$ on the circle of convergence of h is *singular* if there does not exist any analytic continuation of h across an arc of $\{|z| = r\}$ containing $re^{i\theta}$. A point on $\{|z| = r\}$ is called *regular* if it is not singular. We call the totality of the regular points the *regular set* of X_Ξ . Suppose F is absolutely continuous with $F(dt) = f(t)dt$. From Theorem 2.1 (iii), if $f(s) = 0$ on a closed arc I , then the number of zeros of X_Ξ near I is of $O(1)$. This suggests that $X_\Xi(z)$ could be analytically continued across I . This is indeed the case, and the support of the spectral measure remains crucial even when it is not absolutely continuous.

Theorem 2.5. *Let $\text{supp } F$ be the support of the spectral measure $F(dt)$ of Ξ . Then $(\text{supp } F)^c$ is the regular set of X_Ξ , i.e., $X_\Xi(z)$ can only be analytically extended outside the disk of convergence across $(\text{supp } F)^c$.*

Conceptually, stronger dependence of Ξ makes analytic continuation more feasible. The dependence of Ξ is closely related to the spectral measure and its degeneracy. (i) If Ξ is purely non-deterministic (see Introduction for the condition), then the circle of convergence is the natural boundary for X_Ξ . (ii) In Example 2.4, $X_\Xi(z)$ can be analytically continued across the unit circle in the left-half plane. (iii) If $\Xi = (\xi_k)_{k \in \mathbb{Z}}$ is periodic in k with period q , the spectral measure is atomic. In this case, X_Ξ has the meromorphic extension with poles at $\{e^{2\pi ik/q}\}_{k=0,1,\dots,q-1}$.

3. EDELMAN-KOSLTAN'S FORMULA AND SPECTRAL MEASURES

We consider a centered, stationary Gaussian process $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$ with covariance function $\{\gamma(k)\}_{k \in \mathbb{Z}}$. Suppose $\gamma(0) = 1$. Then, by Herglotz's theorem, there exists a probability measure $F(dt)$ on $(-\pi, \pi]$ such that

$$\gamma(k) = \int_{-\pi}^{\pi} e^{-ikt} F(dt).$$

The measure $F(dt)$ is called the spectral measure of Gaussian process Ξ . The Gaussian process Ξ with spectral measure $F(dt)$ admits the following spectral representation (cf. [10]):

$$\xi_k = \int_{-\pi}^{\pi} e^{-ikt} dZ(t), \quad (3.1)$$

where $Z(t)$ is the complex Gaussian process with independent increments satisfying

$$\mathbb{E}[Z(A)\overline{Z(B)}] = F(A \cap B).$$

Every element η of the span $H(\Xi) := \text{span}\{\xi_k, k \in \mathbb{Z}\} \subset L^2(\mathbb{P})$ of $\Xi = \{\xi_k\}_{k \in \mathbb{Z}}$ is represented as

$$\eta = \int_{-\pi}^{\pi} u(t) dZ(t).$$

for some $u \in L^2(F)$. Here $L^2(F)$ is the space of square-integrable functions with respect to the inner product

$$\langle f, g \rangle_F := \int_{-\pi}^{\pi} f(t)\overline{g(t)} F(dt).$$

The correspondence between $\eta \in H(\Xi)$ and $u \in L^2(F)$ is a unitary isomorphism.

From (3.1), we see that

$$X(z) = \sum_{k=0}^{\infty} \left(\int_{-\pi}^{\pi} e^{-ikt} dZ(t) \right) z^k = \int_{-\pi}^{\pi} \frac{1}{1 - ze^{-it}} dZ(t). \quad (3.2)$$

We have an integral representation of the covariance kernel of the GAF X_Ξ from this expression as follows:

$$K(z, w) = \int_{-\pi}^{\pi} \frac{1}{1 - ze^{-it}} \overline{\frac{1}{1 - we^{-it}}} F(dt) \quad (3.3)$$

In particular, for $z = re^{i\varphi}$,

$$K(z, z) = \int_{-\pi}^{\pi} \frac{1}{|1 - ze^{-it}|^2} F(dt) = \frac{1}{1 - r^2} \tilde{F}(re^{i\varphi}) \quad (3.4)$$

where \tilde{F} is the harmonic extension of $F(dt)$ defined as the Poisson integral

$$\tilde{F}(re^{i\varphi}) = \int_{-\pi}^{\pi} P_r(\varphi - t)F(dt) = \int_{-\pi}^{\pi} P_r(s)F_\varphi(ds)$$

with $F_\varphi(ds) = F(d(s + \varphi))$ on $(-\pi, \pi]$ modulo 2π . Here we give a spectral representation of the 1-intensity $\rho_1(z)$.

Proposition 3.1. *The 1-intensity of zeros of $X_\Xi(z)$ is given by the formula*

$$\rho_1(z) = \frac{1}{\pi(1-r^2)^2} \frac{\int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \{1 - \cos(t-s)\} P_r(s)^2 P_r(t)^2 F_\varphi(ds) F_\varphi(dt)}{\left(\int_{-\pi}^{\pi} P_r(s) F_\varphi(ds) \right)^2},$$

where $z = re^{i\varphi}$ and $F_\varphi(ds) = F(d(s + \varphi))$.

Proof. Edelman-Kostlan's formula [4] shows that

$$\rho_1(z) = \frac{1}{\pi} \partial_z \partial_{\bar{z}} \log K(z, z) = \frac{1}{\pi} \frac{\partial_z \partial_{\bar{z}} K(z, z) \cdot K(z, z) - \partial_z K(z, z) \cdot \partial_{\bar{z}} K(z, z)}{K(z, z)^2}.$$

From (3.4),

$$K(z, z)^2 = \frac{1}{(1-r^2)^2} \tilde{F}(re^{i\varphi})^2.$$

Also we see that

$$\begin{aligned} & \partial_z \partial_{\bar{z}} K(z, z) \cdot K(z, z) - \partial_z K(z, z) \cdot \partial_{\bar{z}} K(z, z) \\ &= \frac{1}{2} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \frac{|e^{it} - e^{is}|^2}{|1 - ze^{-it}|^4 |1 - ze^{-is}|^4} F(ds) F(dt) \\ &= \frac{1}{(1-r^2)^4} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \{1 - \cos(t-s)\} P_r(\varphi - t)^2 P_r(\varphi - s)^2 F(ds) F(dt) \\ &= \frac{1}{(1-r^2)^4} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \{1 - \cos(v-u)\} P_r(u)^2 P_r(v)^2 F_\varphi(du) F_\varphi(dv). \end{aligned}$$

By change of variables $s = \varphi + u$ and $t = \varphi + v$, we obtain the assertion. \square

For an integrable function h , we introduce

$$\mathcal{P}_r(h) := \frac{1}{2\pi} \int_{-\pi}^{\pi} h(s) P_r(s) ds, \quad \mathcal{Q}_r(h) := \frac{1}{2\pi} \int_{-\pi}^{\pi} h(s) P_r(s)^2 ds. \quad (3.5)$$

Suppose $F_\varphi(ds) = f_\varphi(s) ds$. Since $P_r(s)$ is symmetric, we have $\mathcal{Q}_r(f_\varphi) = \mathcal{Q}_r(\hat{f}_\varphi)$, $\mathcal{Q}_r(f_\varphi \cos s) = \mathcal{Q}_r(\hat{f}_\varphi \cos s)$, and $\mathcal{Q}_r(f_\varphi \sin s) = \mathcal{Q}_r(\check{f}_\varphi \sin s)$, where \hat{f}_φ and \check{f}_φ are symmetrization and anti-symmetrization of f_φ defined in (2.1). Therefore, when $F_\varphi(ds) = f_\varphi(s) ds$, we can restate Proposition 3.1 as

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)^2} \frac{\mathcal{S}_r(f_\varphi)}{\mathcal{P}_r(\hat{f}_\varphi)^2}. \quad (3.6)$$

where

$$\mathcal{S}_r(f_\varphi) := \mathcal{Q}_r(\hat{f}_\varphi)^2 - \mathcal{Q}_r(\hat{f}_\varphi \cos s)^2 - \mathcal{Q}_r(\check{f}_\varphi \sin s)^2. \quad (3.7)$$

In the next section, we will give asymptotic expansions for $\mathcal{P}_r(h)$ and $\mathcal{Q}_r(h)$ as $r \rightarrow 1$.

4. PRECISE ASYMPTOTICS OF POISSON INTEGRAL AND ITS RELATIVE

The following notation will be used in this section and hereafter.

$$x = x(s) = 1 - \cos s, \quad y = y(r) = 1 - r^2. \quad (4.1)$$

Let $\mathcal{H} := \{h : (-\pi, \pi] \rightarrow \mathbb{R} : h \text{ is smooth around } s = 0, \text{ and symmetric, i.e. } h(-s) = h(s)\}$ and $\mathcal{H}_b := \{h \in \mathcal{H} : h \text{ is bounded}\}$. For $h \in \mathcal{H}$, we define

$$Th(s) := \frac{h(s) - h(0)}{x(s)}, \quad Th(0) = h''(0).$$

Then, $T : \mathcal{H} \rightarrow \mathcal{H}$. If, in addition, h is bounded, so is Th ; T sends \mathcal{H}_b to itself. We also remark that

$$T^2h(0) = (Th)''(0) = \frac{1}{6}(h''(0) + h^{(4)}(0)). \quad (4.2)$$

4.1. Asymptotic expansion I. Fatou's theorem states that if h is integrable on $(-\pi, \pi]$ and continuous at $s = 0$ then $\mathcal{P}_r(h)$ converges to $h(0)$ (cf. [13]). The following is a precise version of Fatou's theorem.

Proposition 4.1. *Suppose $h \in \mathcal{H}_b$. Then, as $r \rightarrow 1$,*

$$\begin{aligned} \mathcal{P}_r(h) &= h(0) + \frac{\mathcal{I}(Th)}{2}y + \frac{1}{4} \left\{ \mathcal{I}(Th) - \frac{h''(0)}{2} \right\} y^2 \\ &\quad + \frac{1}{8} \left\{ \frac{3}{2}\mathcal{I}(Th) - h''(0) - \frac{1}{2}\mathcal{I}(T^2h) \right\} y^3 + O(y^4), \end{aligned} \quad (4.3)$$

where $y = 1 - r^2$.

Proof. We use Lemma 4.2 twice to obtain

$$\begin{aligned} \mathcal{P}_r(h) &= h(0) + \frac{y}{2r}\mathcal{I}(Th) - \frac{y^2}{2r(1+r)^2}Th(0) - \frac{y^3}{4r^2(1+r)^2}\mathcal{I}(T^2h) \\ &\quad + \frac{y^4}{4r^2(1+r)^4}\mathcal{P}_r(T^2h). \end{aligned}$$

By expanding the functions of r in y , we obtain the assertion. \square

We observe a recursion relation of the Poisson integral.

Lemma 4.2. *Suppose $h \in \mathcal{H}_b$. Then,*

$$\mathcal{P}_r(h) = h(0) + \frac{y}{2r}\mathcal{I}(Th) - \frac{y^2}{2r(1+r)^2}\mathcal{P}_r(Th).$$

Proof. We also define the operator \mathcal{U}_r by

$$\mathcal{U}_r(f) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(s) \frac{2x}{(1-r)^2 + 2rx} ds.$$

Then,

$$\mathcal{P}_r(h) = h(0) + \frac{1-r^2}{2}\mathcal{U}_r(Th).$$

On the other hand, the following identity

$$r \frac{2x}{(1-r)^2 + 2rx} = 1 - \frac{1-r^2}{(1+r)^2}P_r(s)$$

yields

$$r\mathcal{U}_r(f) = \mathcal{I}(f) - \frac{1-r^2}{(1+r)^2}\mathcal{P}_r(f).$$

Hence we obtain the equality. \square

4.2. Asymptotic expansion II. In this subsection, we give the following asymptotic expansion of $Q_r(h)$ as $r \rightarrow 1$. Since $P_r(s)ds$ is close to $\delta_0(ds)$, $P_r(s)^2ds$ is close to $\delta_0(ds)^2$, which gives a diverging term.

Proposition 4.3. *Suppose $h \in \mathcal{H}_b$. Then, as $r \rightarrow 1$,*

$$\begin{aligned} \mathcal{Q}_r(h) &= 2h(0)y^{-1} - h(0) + \frac{1}{4}h''(0)y + \left(\frac{1}{4}\mathcal{I}(T^2h) + \frac{1}{8}h''(0)\right)y^2 \\ &\quad + \left(\frac{1}{4}\mathcal{I}(T^2h) + \frac{1}{16}h''(0) - \frac{1}{64}h^{(4)}(0)\right)y^3 \\ &\quad + \left(\frac{1}{4}\mathcal{I}(T^2h) - \frac{1}{16}\mathcal{I}(T^3h) + \frac{1}{32}h''(0) - \frac{3}{128}h^{(4)}(0)\right)y^4 \\ &\quad + O(y^5), \end{aligned}$$

where $y = 1 - r^2$.

Proof. Before proving this expansion formula, we first observe that

$$\mathcal{Q}_r(h) = h(0)\mathcal{Q}_r(1) + Th(0)\mathcal{Q}_r(x) + \mathcal{Q}_r(T^2h \cdot x^2) \quad (4.4)$$

since $h(s) = h(0) + Th(0)x + T^2h(s)x^2$. To expand the third term, we introduce two more operators:

$$\begin{aligned} \mathcal{V}_r(h) &:= \frac{1}{2\pi} \int_{-\pi}^{\pi} h(s) \left\{ \frac{2x}{(1-r)^2 + 2rx} \right\}^2 ds, \\ \mathcal{K}_r(h) &:= \frac{1}{2\pi} \int_{-\pi}^{\pi} h(s) \frac{\{2x - (1-r)\}\{(1-r)^2 + 2(1+r)x\}}{\{(1-r)^2 + 2rx\}^2} ds. \end{aligned}$$

We note that

$$\mathcal{V}_r(h) = \mathcal{I}(h) + (1-r)\mathcal{K}_r(h). \quad (4.5)$$

Since $Q_r(h \cdot x^2) = \frac{y^2}{4}\mathcal{V}_r(h)$, from (4.4) and (4.5), we have

$$\mathcal{Q}_r(h) = h(0)\mathcal{Q}_r(1) + h''(0)\mathcal{Q}_r(x) + \frac{y^2}{4}\mathcal{I}(T^2h) + \frac{y^3}{4(1+r)}\mathcal{K}_r(T^2h). \quad (4.6)$$

It is easy to see that

$$\begin{aligned} \mathcal{Q}_r(1) &= \frac{1+r^2}{1-r^2} = \frac{2}{y} - 1, \\ \mathcal{Q}_r(x) &= \frac{1-r}{1+r} = \frac{1}{4}y + \frac{1}{8}y^2 + \frac{5}{64}y^3 + \frac{7}{128}y^4 + O(y^5). \end{aligned} \quad (4.7)$$

By (4.2), (4.8) below, and $(1+r)^{-1} = \frac{1}{2} + \frac{1}{8}y + O(y^2)$, we obtain the assertion. \square

We have a recursion formula for \mathcal{K}_r , from which we obtain an expansion formula.

Lemma 4.4. *Suppose $h \in \mathcal{H}_b$. As $r \rightarrow 1$,*

$$\mathcal{K}_r(h) = 2\mathcal{I}(h) - \frac{3}{4}h(0) + \left(-\frac{15}{16}h(0) + \frac{3}{2}\mathcal{I}(h) - \frac{1}{2}\mathcal{I}(Th)\right)y + O(y^2). \quad (4.8)$$

Proof. From the identity

$$\begin{aligned} & \frac{\{2x - (1-r)\}\{(1-r)^2 + 2(1+r)x\}}{\{(1-r)^2 + 2rx\}^2} \\ &= 2 + (1-r) \left\{ (1+2r) \left(\frac{2x}{(1-r)^2 + 2rx} \right)^2 + (2r-3) \frac{1}{(1-r)^2 + 2rx} + r(2r-3) \frac{2x}{\{(1-r)^2 + 2rx\}^2} \right\}, \end{aligned}$$

we see that

$$\mathcal{K}_r(h) = 2\mathcal{I}(h) + \frac{1+2r}{1+r} \mathcal{V}_r(h)y + \frac{2r-3}{1+r} \mathcal{P}_r(h) + (1-r)r(2r-3) \left\{ h(0) \frac{2}{(1+r)^3} \frac{1}{1-r} + \frac{1}{2} \mathcal{V}_r(Th) \right\}.$$

For the fourth term, we used $h(s) = h(0) + xTh(s)$ and (4.7). By using Lemma 4.2 and (4.5), we obtain the following recursion equation:

$$\begin{aligned} \mathcal{K}_r(h) &= 2\mathcal{I}(h) + \frac{(2r-3)(1+4r+r^2)}{(1+r)^3} h(0) + \left\{ \frac{1+2r}{1+r} \mathcal{I}(h) + \frac{(2r-3)(1+r^2)}{2r(1+r)} \mathcal{I}(Th) \right\} y \\ &+ \left\{ \frac{1+2r}{(1+r)^2} \mathcal{K}_r(h) - \frac{2r-3}{2r(1+r)^3} \mathcal{P}_r(Th) + \frac{r(2r-3)}{2(1+r)^2} \mathcal{K}_r(Th) \right\} y^2. \quad (4.9) \end{aligned}$$

From (4.9) together with the Taylor expansions of the coefficients at $r = 1$, we see that

$$\begin{aligned} \mathcal{K}_r(h) &= 2\mathcal{I}(h) + \left(-\frac{3}{4} - \frac{15}{16}y + O(y^2) \right) h(0) \\ &+ \left\{ \left(\frac{3}{2} - \frac{1}{8}y + O(y^2) \right) \mathcal{I}(h) + \left(-\frac{1}{2} - \frac{5}{8}y + O(y^2) \right) \mathcal{I}(Th) \right\} y + O(y^2), \end{aligned}$$

from which we obtain the assertion. \square

5. PROOF OF THEOREM 2.1

Suppose $F_\varphi(ds) = f_\varphi(s)ds$ and $f_\varphi(s) > 0$. From (3.6) and (3.7), we recall

$$\rho_1(re^{i\varphi}) = \frac{1}{\pi(1-r^2)^2} \frac{\{\mathcal{Q}_r(\hat{f}_\varphi)\}^2 - \{\mathcal{Q}_r(\hat{f}_\varphi \cos s)\}^2 - \{\mathcal{Q}_r(\check{f}_\varphi \sin s)\}^2}{\{\mathcal{P}_r(\hat{f}_\varphi)\}^2},$$

where \hat{f}_φ and \check{f}_φ are symmetrization and anti-symmetrization of f_φ , respectively, as defined in (2.1).

Suppose $h \in \mathcal{H}_b$. From Proposition 4.3, setting $y = 1 - r^2$, we have

$$\mathcal{Q}_r(h) =: \sum_{k=-1}^4 Q_k(h)y^k + O(y^5),$$

where

$$\begin{aligned} Q_{-1}(h) &= 2h(0), \quad Q_0(h) = -h(0), \quad Q_1(h) = \frac{1}{4}h''(0), \\ Q_2(h) &= \frac{1}{4} \left\{ \mathcal{I}(T^2h) + \frac{h''(0)}{2} \right\}, \\ Q_3(h) &= \frac{1}{4} \left\{ \mathcal{I}(T^2h) + \frac{1}{4}h''(0) - \frac{1}{16}h^{(4)}(0) \right\} \\ Q_4(h) &= \frac{1}{4} \left\{ \mathcal{I}(T^2h) - \frac{1}{4}\mathcal{I}(T^3h) + \frac{1}{8}h''(0) - \frac{3}{32}h^{(4)}(0) \right\}. \end{aligned}$$

Then, we have

$$\begin{aligned}\mathcal{R}_r(h) &:= \mathcal{Q}_r(h)^2 = \sum_{p=-2}^3 \left(\sum_{k+l=p} Q_k(h)Q_l(h) \right) y^p + O(y^4) \\ &=: \sum_{k=-2}^3 R_k(h)y^k + O(y^4).\end{aligned}$$

A direct computation shows that

$$\begin{aligned}R_{-2}(h) &= 4h(0)^2, \quad R_{-1}(h) = -4h(0)^2, \\ R_0(h) &= h(0)\{h(0) + h''(0)\}, \quad R_1(h) = h(0)\mathcal{I}(T^2h) \\ R_2(h) &= h(0) \left\{ \frac{1}{2}\mathcal{I}(T^2h) - \frac{1}{16}h^{(4)}(0) \right\} + \frac{1}{16}\{h''(0)\}^2 \\ &= \frac{1}{2}R_1(h) + \frac{1}{16}\{h''(0)^2 - h(0)h^{(4)}(0)\} \\ R_3(h) &= \frac{1}{2} \left\{ h(0) + \frac{1}{4}h''(0) \right\} \mathcal{I}(T^2h) - \frac{1}{4}h(0)\mathcal{I}(T^3h) + \frac{1}{16}\{h''(0)^2 - h(0)h^{(4)}(0)\} \\ &= R_2(h) + \frac{1}{8}h''(0)\mathcal{I}(T^2h) - \frac{1}{4}h(0)\mathcal{I}(T^3h).\end{aligned}$$

For g (instead of f_φ) which is not necessarily symmetric, we now compute (3.7), i.e.,

$$S_r(g) := \mathcal{R}_r(\hat{g}) - \mathcal{R}_r(\hat{g} \cos s) - \mathcal{R}_r(\check{g} \sin s) = \sum_{k=-2}^3 S_k(g)y^k + O(y^4), \quad (5.1)$$

which is equivalent to

$$S_k(g) = R_k(\hat{g}) - R_k(\hat{g} \cos s) - R_k(\check{g} \sin s).$$

We note that if $h \in \mathcal{H}$, then

$$(h \cos s)''(0) = h''(0) - h(0), \quad h^{(4)}(0) - (h \cos s)^{(4)}(0) = -h(0) + 6h''(0).$$

and

$$Th - T(h \cos s) = h$$

Then we can compute $S_k(g)$ as follows:

$$\begin{aligned}S_{-2}(g) &= S_{-1}(g) = 0, \quad S_0(g) = \hat{g}(0)^2 \\ S_1(g) &= \hat{g}(0)\{\mathcal{I}(T^2\hat{g} - T^2(\hat{g} \cos s))\} = \hat{g}(0)\mathcal{I}(T\hat{g}) \\ S_2(g) &= \frac{1}{2}\hat{g}(0)\mathcal{I}(T\hat{g}) + \frac{1}{16}\{\hat{g}''(0)^2 - (\hat{g} \cos s)''(0)^2 - (\check{g} \sin s)''(0)^2\} \\ &\quad - \frac{1}{16}\hat{g}(0)\{\hat{g}^{(4)}(0) - (\hat{g} \cos s)^{(4)}(0)\} \\ &= \frac{1}{2}\hat{g}(0)\mathcal{I}(T\hat{g}) - \frac{1}{4}\left(\hat{g}(0)\hat{g}''(0) + \check{g}'(0)^2\right) \\ S_3(g) &= S_2(g) + \frac{1}{8}\{\hat{g}''(0)\mathcal{I}(T\hat{g}) + \hat{g}(0)\mathcal{I}(T^2\hat{g} \cos s) - 2\check{g}'(0)\mathcal{I}(T^2\check{g} \sin s)\} - \frac{1}{4}\hat{g}(0)\mathcal{I}(T^2\hat{g}).\end{aligned}$$

When $g(0) > 0$,

$$\mathcal{S}_r(g) = g(0)^2 + g(0)\mathcal{I}(T\hat{g})y + \left\{ \frac{1}{2}g(0)\mathcal{I}(T\hat{g}) - \frac{1}{4}\left(g(0)g''(0) + g'(0)^2\right) \right\} y^2 + O(y^3). \quad (5.2)$$

When $g(0) = 0$, $g'(0) = 0$, we have

$$\mathcal{S}_r(g) = \frac{1}{8}g''(0)\mathcal{I}(T\hat{g})y^3 + O(y^4). \quad (5.3)$$

5.1. The case $f_\varphi(0) > 0$. Suppose the spectral density at φ is strictly positive, i.e., $f_\varphi(0) > 0$. From (3.6), (4.3), and (5.2), we see that

$$\begin{aligned} \rho_1(re^{i\varphi}) &= \frac{1}{\pi(1-r^2)^2} \frac{\mathcal{S}_r(f_\varphi)}{\mathcal{P}_r(\hat{f}_\varphi)^2} \\ &= \frac{1}{\pi(1-r^2)^2} \frac{f_\varphi(0)^2 + f_\varphi(0)\mathcal{I}(T\hat{f}_\varphi)y + b_\varphi y^2 + O(y^3)}{f_\varphi(0)^2 + f_\varphi(0)\mathcal{I}(T\hat{f}_\varphi)y + c_\varphi y^2 + O(y^3)} \\ &= \frac{1}{\pi(1-r^2)^2} \left\{ 1 + \frac{b_\varphi - c_\varphi}{f_\varphi(0)^2} y^2 + O(y^3) \right\} \\ &= \frac{1}{\pi(1-r^2)^2} \left\{ 1 - \frac{1}{4f_\varphi(0)^2} \{f'_\varphi(0)^2 + \{\mathcal{I}(T\hat{f}_\varphi)\}^2\} y^2 + O(y^3) \right\}. \end{aligned}$$

We used the expansion formula

$$\frac{p + au + bu^2 + O(u^3)}{p + au + cu^2 + O(u^3)} = 1 + \frac{b-c}{p} u^2 + O(u^3) \quad \text{as } u \rightarrow 0.$$

5.2. The case $f_\varphi(0) = 0$. First, note that when $f_\varphi(0) = 0$, we have $f'_\varphi(0) = 0$ since $f_\varphi(s) \geq 0$. We have the following asymptotics from Propositions 4.1 and 4.3

$$\mathcal{P}_r(\hat{f}_\varphi) = \frac{\mathcal{I}(T\hat{f}_\varphi)}{2}y + \frac{1}{4} \left(\mathcal{I}(T\hat{f}_\varphi) - \frac{f''_\varphi(0)}{2} \right) y^2 + O(y^3),$$

and thus

$$\mathcal{P}_r(\hat{f}_\varphi)^2 = \frac{1}{4}\mathcal{I}(T\hat{f}_\varphi)^2 y^2 + \frac{1}{4}\mathcal{I}(T\hat{f}_\varphi) \left(\mathcal{I}(T\hat{f}_\varphi) - \frac{f''_\varphi(0)}{2} \right) y^3 + O(y^4). \quad (5.4)$$

On the other hand, we see that

$$\mathcal{S}_r(f_\varphi) = \frac{1}{8}f''_\varphi(0)\mathcal{I}(T\hat{f}_\varphi)y^3 + O(y^4). \quad (5.5)$$

Therefore, from (5.4) and (5.5), we obtain

$$\frac{\mathcal{S}_r(f_\varphi)}{\mathcal{P}_r(f_\varphi)^2} = \frac{f''_\varphi(0)}{2\mathcal{I}(T\hat{f}_\varphi)}y + O(y^2).$$

This implies Theorem 2.1(ii).

5.3. **The case** $f_\varphi(0) = f_\varphi''(0) = 0$. When $f_\varphi(0) = f_\varphi''(0) = 0$ (and thus $f_\varphi'(0) = 0$), from Proposition 4.1, we have

$$\mathcal{P}_r(\hat{f}_\varphi) = \frac{1}{2}\mathcal{I}(T\hat{f}_\varphi)y + \frac{1}{4}\mathcal{I}(T\hat{f}_\varphi)^2y^2 + O(y^3),$$

and hence

$$\mathcal{P}_r(\hat{f}_\varphi)^2 = \frac{1}{4}\mathcal{I}(T\hat{f}_\varphi)^2y^2 + O(y^3). \quad (5.6)$$

From Proposition 4.3, we have

$$\mathcal{Q}_r(\hat{f}_\varphi) = \frac{1}{4}\mathcal{I}(T^2\hat{f}_\varphi)y^2 + O(y^3),$$

and thus

$$\begin{aligned} \mathcal{S}_r(f_\varphi) &= \mathcal{Q}_r(\hat{f}_\varphi)^2 - \mathcal{Q}_r(\hat{f}_\varphi \cos s)^2 - \mathcal{Q}_r(\check{f}_\varphi \sin s)^2 \\ &= \frac{1}{16} \left\{ \mathcal{I}(T^2\hat{f}_\varphi)^2 - \mathcal{I}(T^2\hat{f}_\varphi \cos s)^2 - \mathcal{I}(T^2\check{f}_\varphi \sin s)^2 \right\} y^4 + O(y^5) \end{aligned} \quad (5.7)$$

Therefore, from (5.6) and (5.7), we obtain

$$\begin{aligned} \rho_1(re^{i\varphi}) &= \frac{1}{\pi(1-r^2)^2} \frac{\mathcal{S}_r(f_\varphi)}{\mathcal{P}_r(f_\varphi)^2} \\ &= \frac{1}{\pi} \frac{\mathcal{I}(T^2\hat{f}_\varphi)^2 - \mathcal{I}(T^2(\hat{f}_\varphi \cos s))^2 - \mathcal{I}(T^2(\check{f}_\varphi \sin s))^2}{4\mathcal{I}(T\hat{f}_\varphi)^2} + O(y). \end{aligned}$$

Remark 5.1. The numerator of the first term above can be expressed as

$$\frac{1}{4} \int_{-\pi}^{\pi} \int_{-\pi}^{\pi} \frac{G_\varphi(s, t) + G_\varphi(s, -t)}{(1 - \cos s)^2(1 - \cos t)^2} ds dt,$$

where $G_\varphi(s, t) := \{f_\varphi(s)f_\varphi(t) + f_\varphi(-s)f_\varphi(-t)\}\{1 - \cos(s-t)\}$.

6. ANALYTIC CONTINUATION

First we give a proof of the following basic fact.

Lemma 6.1. *Let $\{\zeta_n\}_{n=0}^\infty$ be a sequence of standard complex normal random variables that are not necessarily independent. Then,*

$$\lim_{n \rightarrow \infty} |\zeta_n|^{1/n} = 1 \text{ a.s.}$$

Proof. Since $\mathbb{P}(|\zeta| \geq r) = e^{-r^2}$ for $\zeta \sim N_{\mathbb{C}}(0, 1)$, we have

$$\sum_{n=1}^{\infty} \mathbb{P}(|\zeta_n| \geq \sqrt{2 \log n}) \leq \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty$$

and

$$\sum_{n=1}^{\infty} \mathbb{P}(|\zeta_n| < \frac{1}{n}) = \sum_{n=1}^{\infty} (1 - e^{-\frac{1}{n^2}}) \leq \sum_{n=1}^{\infty} \frac{1}{n^2} < \infty,$$

which imply

$$\mathbb{P}\left(\frac{1}{n} \leq |\zeta_n| \leq \sqrt{2 \log n} \text{ f.e.}\right) = 1.$$

Therefore, we obtain the assertion. \square

Suppose $h(z)$ is a power series of convergence radius $\rho \in (0, \infty)$. We recall that a point b on $|z| = \rho$ is singular (or a singular point) if there does not exist any analytic continuation of $h(z)$ across an arc of $\{|z| = \rho\}$ containing b ; otherwise b is called regular (or a regular point). A subset of an arc $|z| = \rho$ is regular if it consists of regular points. If there exist $0 < r < 1$ and $0 < \eta < \pi$ such that

$$\rho(r) := \left(\limsup_{k \rightarrow \infty} \left| \frac{1}{k!} X_{\Xi}^{(k)}(r) \right|^{1/k} \right)^{-1} > |e^{i\eta} - r| = (1 - 2r \cos \eta + r^2)^{1/2}, \quad (6.1)$$

then the arc $I_\eta = \{e^{is} : |s| \leq \eta\} \subset \partial\mathbb{D}$ is regular (cf. [11, p.39]) since $\rho(r)$ is the radius of convergence centered at r and the disk with radius $\rho(r)$ centered at r contains the arc I_η . Using this fact, we provide a proof of Theorem 2.5.

Proof of Theorem 2.5. Since the open set $(\text{supp } F)^c$ can be expressed as at most a countable union of open arcs, we regard each open arc as $I_\delta^\circ := \{e^{is} : s \in (-\delta, \delta)\}$ for some $\delta > 0$ by applying an appropriate rotation if needed. We check (6.1) to show the open arc I_δ° is regular. From the spectral representation (3.2), we see that

$$\alpha_k = \alpha_k(z) := \frac{1}{k!} X_{\Xi}^{(k)}(z) = \int_{-\pi}^{\pi} \frac{e^{-ikt}}{(1 - ze^{-it})^{k+1}} dZ(t).$$

Therefore, for $z = r \in (0, 1)$,

$$\begin{aligned} \text{var}(\alpha_k) &= \mathbb{E} \left[\left| \frac{1}{k!} X_{\Xi}^{(k)}(z) \right|^2 \right] \\ &= \int_{-\pi}^{\pi} \frac{1}{|1 - re^{-it}|^{2(k+1)}} F(dt) \\ &= \int_{-\pi}^{\pi} \frac{1}{(1 - 2r \cos t + r^2)^{k+1}} F(dt). \end{aligned}$$

Since $\{\alpha_k\}$ is a sequence of centered Gaussian random variables and $\alpha_k = \text{var}(\alpha_k)^{1/2} \zeta_k$ with $\zeta_k := \alpha_k / \text{var}(\alpha_k)^{1/2} \sim N_{\mathbb{C}}(0, 1)$, by Lemma 6.1, we have

$$\rho(r) = \left(\limsup_{k \rightarrow \infty} |\alpha_k|^{1/k} \right)^{-1} = \limsup_{k \rightarrow \infty} \text{var}(\alpha_k)^{-1/2k} \quad a.s.$$

Since the function $1 - 2r \cos t + r^2$ is increasing in $t \in (0, \pi)$ for $r \in (0, 1)$, and $\text{supp } F \subset (I_\delta^\circ)^c$, we have

$$\begin{aligned} \rho(r) &= \limsup_{k \rightarrow \infty} \left(\int_{-\pi}^{\pi} \frac{1}{(1 - 2r \cos t + r^2)^{k+1}} F(dt) \right)^{-1/2k} \\ &\geq (1 - 2r \cos \delta + r^2)^{1/2}. \end{aligned}$$

From (6.1), the arc I_η is regular for any $\eta \in (0, \delta)$, and thus I_δ° is regular.

On the other hand, when $0 \in \text{supp } F$, for every $\epsilon \in (0, \pi)$, we see that

$$\begin{aligned} \rho(r) &\leq \limsup_{k \rightarrow \infty} \left(\int_{-\epsilon}^{\epsilon} \frac{1}{(1 - 2r \cos t + r^2)^{k+1}} F(dt) \right)^{-1/2k} \\ &\leq \limsup_{k \rightarrow \infty} \left(F([- \epsilon, \epsilon]) \cdot (1 - 2r \cos \epsilon + r^2)^{-(k+1)} \right)^{-1/2k} \\ &= (1 - 2r \cos \epsilon + r^2)^{1/2}. \end{aligned}$$

Since $\epsilon > 0$ is arbitrary, $\rho(r) \leq 1 - r$ for every $r \in (0, 1)$, which implies that $X_{\Xi}(z)$ cannot be analytically extended across $z = 1$, i.e., $z = 1$ is a singular point. Therefore, we can conclude that $(\text{supp } F)^c$ is the regular set of X_{Ξ} . \square

Acknowledgment. This work was supported by JSPS KAKENHI Grant Numbers JP22H05105, JP23H01077 and JP23K25774, and also supported in part JP21H04432 and JP24KK0060.

REFERENCES

- [1] Rémi Bardenet and Adrien Hardy. Time-frequency transforms of white noises and Gaussian analytic functions. *Appl. Comput. Harmon. Anal.*, 50:73–104, 2021.
- [2] Jeremiah Buckley and Alon Nishry. Gaussian complex zeroes are not always normal: limit theorems on the disc. *Probab. Math. Phys.*, 3(3):675–706, 2022.
- [3] Jeremiah Buckley, Alon Nishry, Ron Peled, and Mikhail Sodin. Hole probability for zeroes of Gaussian Taylor series with finite radii of convergence. *Probab. Theory Related Fields*, 171(1-2):377–430, 2018.
- [4] Alan Edelman and Eric Kostlan. How many zeros of a random polynomial are real? *Bull. Amer. Math. Soc. (N.S.)*, 32(1):1–37, 1995.
- [5] Naomi D. Feldheim. Zeroes of gaussian analytic functions with translation-invariant distribution. *Israel Journal of Mathematics*, 195(1):317–345, 2013.
- [6] Naomi Dvora Feldheim. Variance of the number of zeroes of shift-invariant Gaussian analytic functions. *Israel J. Math.*, 227(2):753–792, 2018.
- [7] Subhroshekhar Ghosh and Yuval Peres. Rigidity and tolerance in point processes: Gaussian zeros and Ginibre eigenvalues. *Duke Math. J.*, 166(10):1789–1858, 2017.
- [8] Antti Haimi, Günther Koliander, and José Luis Romero. Zeros of Gaussian Weyl-Heisenberg functions and hyperuniformity of charge. *J. Stat. Phys.*, 187(3):Paper No. 22, 41, 2022.
- [9] J. Ben Hough, Manjunath Krishnapur, Yuval Peres, and Bálint Virág. *Zeros of Gaussian analytic functions and determinantal point processes*, volume 51 of *University Lecture Series*. American Mathematical Society, Providence, RI, 2009.
- [10] I. A. Ibragimov and Y. A. Rozanov. *Gaussian random processes*, volume 9 of *Applications of Mathematics*. Springer-Verlag, New York-Berlin, 1978. Translated from the Russian by A. B. Aries.
- [11] Jean-Pierre Kahane. *Some random series of functions*, volume 5 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1985.
- [12] Avner Kiro and Alon Nishry. Fluctuations for zeros of Gaussian Taylor series. *J. Lond. Math. Soc. (2)*, 104(3):1172–1203, 2021.
- [13] Paul Koosis. *Introduction to H_p spaces*, volume 115 of *Cambridge Tracts in Mathematics*. Cambridge University Press, Cambridge, second edition, 1998. With two appendices by V. P. Havin [Viktor Petrovich Khavin].
- [14] Manjunath Krishnapur. From random matrices to random analytic functions. *Ann. Probab.*, 37(1):314–346, 2009.
- [15] Safari Mukeru, Mmboniseni P. Mulaudzi, Joseph Nzabanita, and Marc M. Mpanda. Zeros of Gaussian power series with dependent random variables. *Illinois J. Math.*, 64(4):569–582, 2020.
- [16] F. Nazarov and M. Sodin. Correlation functions for random complex zeroes: strong clustering and local universality. *Comm. Math. Phys.*, 310(1):75–98, 2012.
- [17] Alon Nishry. Hole probability for entire functions represented by Gaussian Taylor series. *J. Anal. Math.*, 118(2):493–507, 2012.
- [18] Kohei Noda and Tomoyuki Shirai. Expected number of zeros of random power series with finitely dependent Gaussian coefficients. *J. Theoret. Probab.*, 36(3):1534–1554, 2023.
- [19] Yuval Peres and Bálint Virág. Zeros of the i.i.d. Gaussian power series: a conformally invariant determinantal process. *Acta Math.*, 194(1):1–35, 2005.
- [20] Tomoyuki Shirai. Limit theorems for random analytic functions and their zeros. In *Functions in number theory and their probabilistic aspects*, volume B34 of *RIMS Kôkyûroku Bessatsu*, pages 335–359. Res. Inst. Math. Sci. (RIMS), Kyoto, 2012.
- [21] M. Sodin. Zeros of Gaussian analytic functions. *Math. Res. Lett.*, 7(4):371–381, 2000.

- [22] Mikhail Sodin and Boris Tsirelson. Random complex zeroes. I. Asymptotic normality. *Israel J. Math.*, 144:125–149, 2004.

INSTITUTE OF MATHEMATICS FOR INDUSTRY, KYUSHU UNIVERSITY, 744 MOTOOKA, NISHI-KU,
FUKUOKA 819-0001, JAPAN

Email address: shirai@imi.kyushu-u.ac.jp