

BESICOVITCH-WEIGHTED ERGODIC THEOREMS WITH CONTINUOUS TIME

SEMYON LITVINOV

ABSTRACT. Given $1 \leq p < \infty$, we show that ergodic flows in the L^p -space over a σ -finite measure space generated by strongly continuous semigroups of Dunford-Schwartz operators and modulated by bounded Besicovitch almost periodic functions converge almost uniformly (in Egorov's sense). The corresponding local ergodic theorem is proved with identification of the limit. Then we extend these results to arbitrary fully symmetric spaces, including Orlicz, Lorentz, and Marcinkiewicz spaces.

1. INTRODUCTION

Since the groundbreaking work [21], pointwise convergence of Besicovitch-weighted ergodic averages has been examined quite extensively; see, for example, [2, 3, 16, 10, 8]. However, all these studies were concerned with discrete time, thus, there is a natural question what happens in the realm of continuous time. In this case, of course, one must replace Besicovitch sequences by the well-known Besicovitch almost periodic functions [1].

Initially, this problem was studied in the non-commutative setting—for actions of strongly continuous semigroups of *positive* Dunford-Schwartz operators—in [19] for $t \rightarrow 0^+$ and in [7] for $t \rightarrow 0^+$ and for $t \rightarrow \infty$. In both articles, non-commutative counterparts of the almost uniform convergence (in Egorov's sense) were employed.

Back to the commutative case with infinite measure, it is worth asking whether *almost uniform convergence*—which is generally stronger than almost everywhere convergence—of Besicovitch-weighted ergodic averages with continuous time will hold for actions of strongly continuous semigroups of Dunford-Schwartz operators that are *not presumed positive*. Besides, as the measure is not finite, there is another question: what happens within the the universal $(L^1 + L^\infty)$ -space *outside the L^1 -space*. These are the three questions we aim to address in this article.

Let (Ω, μ) be a σ -finite measure space. Given $1 \leq p \leq \infty$, denote $\mathcal{L}^p = \mathcal{L}^p(\Omega, \mu)$. Let $\mathcal{T} = \{T_s\}_{s \geq 0}$ be a semigroup of Dunford-Schwartz operators in $\mathcal{L}^1 + \mathcal{L}^\infty$ which is strongly continuous in \mathcal{L}^1 . In Section 4 of the article, we show that the corresponding ergodic averages $\{M_t(\mathcal{T})(f)\}_{t > 0}$ converge almost uniformly as $t \rightarrow \infty$ and as $t \rightarrow 0^+$ for every $f \in \mathcal{L}^p$, where $1 \leq p < \infty$. Note that the pointwise, that is almost everywhere, convergence of these averages was studied in [23] and later in [18] and [22].

Then we proceed in Section 5 by showing that almost uniform convergence in the space \mathcal{L}^p , where $1 \leq p < \infty$, holds when the above ergodic averages are modulated

Date: January 5, 2025.

2010 Mathematics Subject Classification. 47A35(primary), 46L52(secondary).

Key words and phrases. Sigma-finite measure, Dunford-Schwartz operator, continuous semigroup, bounded Besicovitch function, almost uniform convergence, fully symmetric space.

by the bounded Besicovitch (when $t \rightarrow \infty$) or locally Besicovitch (when $t \rightarrow 0^+$) almost periodic functions. In the case $t \rightarrow 0^+$, the limits are identified.

In Section 6, we extend these results to arbitrary fully symmetric spaces. Such include the well-known Orlicz, Lorentz, and Marcinkiewicz spaces.

2. PRELIMINARIES

Let $(\Omega, \mathcal{A}, \mu)$ be a σ -finite infinite measure space and let $\mathcal{L}^0 = \mathcal{L}^0(\Omega, \mu)$ be the $*$ -algebra of equivalence classes of almost everywhere (a.e.) finite complex-valued measurable functions on Ω . Given $1 \leq p \leq \infty$, let $\mathcal{L}^p \subset \mathcal{L}^0$ be the \mathcal{L}^p -space on Ω equipped with the standard Banach norm $\|\cdot\|_p$.

A net $\{f_\alpha\} \subset \mathcal{L}^0$ is said to converge *almost uniformly (a.u.)* to $f \in \mathcal{L}^0$ (in Egorov's sense) if for every $\varepsilon > 0$ there exists a set $G \subset \Omega$ such that $\mu(\Omega \setminus G) \leq \varepsilon$ and $\|(f - f_\alpha)\chi_G\|_\infty \rightarrow 0$, where χ_G is the characteristic function of set G . It is clear that every a.u. convergent net converges almost everywhere (a.e.) and that the converse is not true in general.

Note that a.u. convergence may not agree with any topology in \mathcal{L}^0 – see, for example, [20]. A proof of the following statement can be found in [17, Theorem 3.2]. Although it is given there in the non-commutative setting, one only needs to adjust notation and then verbatim repeat the argument.

Proposition 2.1. *The space \mathcal{L}^0 is complete with respect to a.u. convergence*

Traditionally, a linear operator $T : \mathcal{L}^1 \rightarrow \mathcal{L}^1$ is called a *Dunford-Schwartz operator* (in short, $T \in DS$) if

$$\|T(f)\|_1 \leq \|f\|_1 \quad \forall f \in \mathcal{L}^1 \quad \text{and} \quad \|T(f)\|_\infty \leq \|f\|_\infty \quad \forall f \in \mathcal{L}^\infty \cap \mathcal{L}^1;$$

see [9, Ch. VIII, § 6], [12], [14, §§ 4.1, 4.2].

However, as it was proved in [8, Theorem 3.3] (see also [22, p. 42]), one can assume without loss of generality that $T \in DS$ contracts the entire space \mathcal{L}^∞ :

Theorem 2.1. *For any Dunford-Schwartz operator T there exists a unique linear operator $\tilde{T} : \mathcal{L}^1 + \mathcal{L}^\infty \rightarrow \mathcal{L}^1 + \mathcal{L}^\infty$ such that*

$$\tilde{T}(f) = T(f) \quad \forall f \in \mathcal{L}^1, \quad \|\tilde{T}(f)\|_\infty \leq \|f\|_\infty \quad \forall f \in \mathcal{L}^\infty,$$

and $\tilde{T}|_{\mathcal{L}^\infty}$ is $\sigma(\mathcal{L}^\infty, \mathcal{L}^1)$ -continuous.

This observation is of the essence for us as we aim to extend the results initially developed for \mathcal{L}^1 to arbitrary fully symmetric spaces $E \subset \mathcal{L}^1 + \mathcal{L}^\infty$.

If a Dunford-Schwartz operator T is positive, that is, $T(f) \geq 0$ whenever $f \geq 0$, we will write $T \in DS^+$.

Let X be a Banach space, and let $M_\alpha : X \rightarrow \mathcal{L}^0$, $\alpha \in \Lambda$, be a family of linear maps. The function on X given by

$$M^*(f) = \sup_{\alpha \in \Lambda} |M_\alpha(f)|, \quad f \in X,$$

is called the *maximal function* of the family $\{M_\alpha\}_{\alpha \in \Lambda}$. If $M^*(f) \in \mathcal{L}^0$ for all $f \in X$, then the map $M^* : X \rightarrow \mathcal{L}^0$ is called the *maximal operator* of $\{M_\alpha\}_{\alpha \in \Lambda}$.

Given $f \in \mathcal{L}^1 + \mathcal{L}^\infty$ and $T \in DS$, denote

$$M_n(f) = \frac{1}{n} \sum_{k=0}^{n-1} T^k(f), \quad n = 1, 2, \dots \quad \text{and} \quad M(T)^*(f) = \sup_n |M_n(f)|.$$

For a proof of the following *maximal ergodic inequality* we refer the reader to [6, Theorem 3.3].

Theorem 2.2. *If $T \in DS$ and $1 \leq p < \infty$, then*

$$\mu\{M(T)^*(|f|) > \lambda\} \leq \left(2 \frac{\|f\|_p}{\lambda}\right)^p \quad \forall f \in \mathcal{L}^p, \lambda > 0.$$

In what follows, $\mathcal{T} = \{T_s\}_{s \geq 0} \subset DS$ is a semigroup, *strongly continuous in \mathcal{L}^1* , that is,

$$\|T_s(f) - T_{s_0}(f)\|_1 \rightarrow 0 \quad \text{whenever } s \rightarrow s_0 \text{ for all } f \in \mathcal{L}^1.$$

It is well-known that if a linear operator contracts both \mathcal{L}^1 and \mathcal{L}^∞ , it also contracts each space \mathcal{L}^p for $1 < p < \infty$. In addition, utilizing the idea of the proof of Proposition 2.1 in [5], we establish the following fact:

Proposition 2.2. *If a semigroup $\{T_s\}_{s \geq 0} \subset DS$ is strongly continuous in \mathcal{L}^1 , then it is strongly continuous in each \mathcal{L}^p for $1 < p < \infty$.*

Proof. Clearly, it is sufficient to verify that

$$\|T_{s_n}(f) - T_{s_0}(f)\|_p \rightarrow 0 \quad \text{as } s_n \rightarrow s_0 \quad \forall f \in \mathcal{L}^p.$$

So, let $f \in \mathcal{L}^p$ and, given $m \in \mathbb{N}$, denote

$$f_m = f \chi_{\{|f| \leq m^{-1}\}} + f \chi_{\{|f| \geq m\}} \quad \text{and} \quad g_m = f - f_m.$$

Fix $\varepsilon > 0$ and define m_0 to be such that $\|f_{m_0}\|_p \leq \varepsilon/4$. As T_s is a contraction in \mathcal{L}^p for each $s \geq 0$, we have

$$\|T_{s_n}(f_{m_0}) - T_{s_0}(f_{m_0})\|_p \leq \frac{\varepsilon}{2} \quad \forall n \in \mathbb{N}.$$

Next, note that $g_{m_0} \in \mathcal{L}^1 \cap \mathcal{L}^\infty$. Since we can assume that $f \neq 0$, we can also assume, without loss of generality, that $g_{m_0} \neq 0$. Then, if we let

$$h_n = \frac{1}{2} \|g_{m_0}\|_\infty^{-1} (T_{s_n}(g_{m_0}) - T_{s_0}(g_{m_0})),$$

it follows, by the assumption, that $\|h_n\|_1 \rightarrow 0$ as $n \rightarrow \infty$. Besides, since T_s is a contraction in \mathcal{L}^∞ for each $s \geq 0$, we have $-\mathbf{1} \leq h_n \leq \mathbf{1}$, hence $|h_n|^p \leq |h_n|$, for each n , implying that

$$\|h_n\|_p^p = \int_\Omega |h_n|^p d\mu \leq \int_\Omega |h_n| d\mu = \|h_n\|_1 \rightarrow 0 \quad \text{as } n \rightarrow \infty.$$

Thus, there exists $n_0 \in \mathbb{N}$ such that

$$\|T_{s_n}(g_{m_0}) - T_{s_0}(g_{m_0})\|_p \leq \frac{\varepsilon}{2} \quad \forall n \geq n_0.$$

Therefore, if $n \geq n_0$, it follows that

$$\|T_{s_n}(f) - T_{s_0}(f)\|_p \leq \|T_{s_n}(f_{m_0}) - T_{s_0}(f_{m_0})\|_p + \|T_{s_n}(g_{m_0}) - T_{s_0}(g_{m_0})\|_p \leq \varepsilon,$$

and the proof is complete. \square

The following proposition is to justify the existence of ergodic averages we are going to examine in this article. Note that the averages $M_t(\mathcal{T})(f)$ below were actually introduced in [18] and [22] but in a less straightforward manner, based on a non-trivial application of Fubini's theorem to the product measure space $\mathbb{R}_+ \otimes \Omega$.

Proposition 2.3. *Let ν be Lebesgue measure on \mathbb{R}_+ , and let $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ be a ν -measurable function with $\|\beta\|_\infty < \infty$. Given $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, the function $\beta(s)T_s(f)$ acting on the measure space (\mathcal{R}_+, ν) with the values in the Banach space $(\mathcal{L}^p, \|\cdot\|_p)$ is Bochner ν -integrable on $[0, t]$ for every $t > 0$. Therefore, the ergodic averages*

$$M_t(\mathcal{T}, \beta)(f) := \frac{1}{t} \int_0^t \beta(s)T_s(f)ds \in \mathcal{L}^p.$$

are defined for all $f \in \mathcal{L}^p$ and $t > 0$.

In particular, the averages

$$M_t(\mathcal{T})(f) := \frac{1}{t} \int_0^t T_s(f)ds \in \mathcal{L}^p$$

are also defined for all $f \in \mathcal{L}^p$ and $t > 0$.

Proof. Fix $f \in \mathcal{L}^p$ and let q be such that $p^{-1} + q^{-1} = 1$. Then, by virtue of Proposition 2.2 and Hölder's inequality, given $g \in \mathcal{L}^q$, the function

$$\mathbb{R}_+ \ni s \mapsto \int_\Omega T_s(f)gd\mu$$

is continuous on \mathbb{R}_+ , implying that the map

$$S_f : \mathbb{R}_+ \rightarrow \mathcal{L}^p \text{ given by } S_f(s) = T_s(f)$$

is *weakly ν -measurable* as of [24, Ch. V, § 4]. Since $S_f(\mathcal{R}_+)$ is a separable subset of the Banach space $(\mathcal{L}^p, \|\cdot\|_p)$, Pettis theorem [24, Ch. V, § 4] implies that S_f is *strongly ν -measurable*, that is, there exists a sequence $U_n : \mathbb{R}_+ \rightarrow \mathcal{L}^p$ of simple maps, as in [24, Ch. V, § 4], such that

$$\lim_{n \rightarrow \infty} \|T_s(f) - U_n(s)\|_p = 0, \text{ hence } \|T_s(f)\|_p = \lim_{n \rightarrow \infty} \|U_n(s)\|_p, \nu\text{-a.e.}$$

Therefore, as for any n the function $\mathbb{R}_+ \ni s \mapsto \|U_n(s)\|_p$ is trivially ν -measurable, it follows that the function

$$\mathbb{R}_+ \ni s \mapsto \|T_s(f)\|_p$$

is also ν -measurable. Since, in addition,

$$\|T_s(f)\|_p \leq \|f\|_p \quad \forall s \in \mathbb{R}_+,$$

the function $\|T_s(f)\|_p$, as well as the function

$$\|\beta(s)T_s(f)\|_p = |\beta(s)| \|T_s(f)\|_p,$$

is ν -integrable on $[0, t]$, for each $t > 0$.

Therefore, by [24, Ch. V, § 5, Theorem 1], the function $\beta(s)T_s(f)$ is Bochner ν -integrable on $[0, t]$ for any given $t > 0$, which completes the proof. \square

Given $1 \leq p < \infty$, we can now define the maximal function of the family $\{M_t(\mathcal{T})\}_{t>0}$ on \mathcal{L}^p as

$$M(\mathcal{T})^*(f) = \sup_{t>0} |M_t(\mathcal{T})(f)|, \quad f \in \mathcal{L}^p.$$

We will need the following, stronger version, of Riesz convergence theorem:

Proposition 2.4. *If $f_n \rightarrow f$ in measure, then there exists a subsequence $\{n_k\}$ such that $f_{n_k} \rightarrow f$ a.u.*

Proof. Fix $\varepsilon > 0$, Since, for every given $k \in \mathbb{N}$, we have

$$\mu\{|f_n - f| > 1/k\} \rightarrow 0,$$

there exists n_k such that

$$\mu\{|f_{n_k} - f| > 1/k\} \leq \frac{\varepsilon}{2^k}.$$

Therefore, if we define

$$\Omega_k = \{|f_{n_k} - f| \leq 1/k\} \text{ and } \Omega_\varepsilon = \bigcap_{k=1}^{\infty} \Omega_k,$$

then $\mu(\Omega \setminus \Omega_k) \leq \frac{\varepsilon}{2^k}$ for each $k \in \mathbb{N}$ implies that $\mu(\Omega \setminus \Omega_\varepsilon) \leq \varepsilon$ and

$$\| |f_{n_k} - f| \chi_{\Omega_\varepsilon} \|_\infty \leq 1/k, \text{ hence } \| |f_{n_k} - f| \chi_{\Omega_\varepsilon} \|_\infty \rightarrow 0,$$

that is, $f_{n_k} \rightarrow f$ a.u. □

We shall also notice the following known fact, together with a corollary, that will be used repeatedly in the sequel. A short proof is provided, due to their special importance in this article.

Proposition 2.5. *Given $1 \leq p < \infty$, the unit ball in \mathcal{L}^p is sequentially closed with respect to a.e. convergence.*

Proof. Let $\|f_n\|_p \leq 1$ for each n and $f_n \rightarrow f$ a.e. Then $|f_n|^p \rightarrow |f|^p$ a.e. also, implying, by Fatou's lemma, that

$$\int_{\Omega} |f|^p d\mu \leq \liminf \int_{\Omega} |f_n|^p d\mu \leq 1,$$

hence $\|f\|_p \leq 1$. □

Since a.u. convergence implies a.e. convergence, and in view of Proposition 2.4, we also have this:

Corollary 2.1. *For every $1 \leq p < \infty$, the unit ball in \mathcal{L}^p is sequentially closed with respect to a.u. convergence and convergence in measure.*

3. MAXIMAL ERGODIC INEQUALITY FOR THE FAMILY $\{M_t(\mathcal{T}, \beta)\}_{t>0}$

Let $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ is a Lebesgue measurable function with $\|\beta\|_\infty \leq C < \infty$. Denote

$$M(\mathcal{T}, \beta)^*(f) = \sup_{t>0} |M_t(\mathcal{T}, \beta)(f)|,$$

the maximal function of the family $\{M_t(\mathcal{T}, \beta)\}_{t>0}$.

Theorem 3.1. *Let $1 \leq p < \infty$. If $\mathcal{T} = \{T_s\}_{s \geq 0}$ and $\beta(s)$ are as above, then*

$$(1) \quad \mu \left\{ M(\mathcal{T}, \beta)^*(f) > \lambda \right\} \leq \left(\frac{4C \|f\|_p}{\lambda} \right)^p$$

and, in particular,

$$(2) \quad \mu \left\{ M(\mathcal{T})^*(f) > \lambda \right\} \leq \left(\frac{4 \|f\|_p}{\lambda} \right)^p$$

for all $f \in \mathcal{L}^p$ and $\lambda > 0$.

Proof. Fix $f \in \mathcal{L}^p$, $\lambda > 0$, $\frac{n}{m} \in \mathbb{Q}$, where $n, m \in \mathbb{N}$, and denote

$$g = \int_0^1 |T_{s/m}(f)| ds.$$

Let $|T|$ be the linear modulus of $T \in DS$ – see, for example, [14, §4.1]. Then $|T| \in DS^+$ and

$$|T^k(f)| \leq |T|^k(|f|) \quad \forall f \in \mathcal{L}^p,$$

thus, we can write

$$\begin{aligned} |M_{n/m}(\mathcal{T}, \beta)(f)| &= \left| \frac{m}{n} \int_0^{\frac{n}{m}} \beta(s) T_s(f) ds \right| \leq 2 \frac{m}{n} \int_0^{\frac{n}{m}} |\beta(s) T_s(f)| ds \\ &= 2 \frac{m}{n} \int_0^{\frac{n}{m}} |\beta(s)| |T_s(f)| ds \leq 2C \frac{m}{n} \int_0^{\frac{n}{m}} |T_s(f)| ds \\ &= \frac{2C}{n} \int_0^n |T_{s/m}(f)| ds = \frac{2C}{n} \sum_{k=0}^{n-1} \int_k^{k+1} |T_{s/m}(f)| ds \\ &= \frac{2C}{n} \sum_{k=0}^{n-1} \int_0^1 |T_{(s+k)/m}(f)| ds = \frac{2C}{n} \sum_{k=0}^{n-1} \int_0^1 |T_{1/m}^k(T_{s/m}(f))| ds \\ &\leq \frac{2C}{n} \sum_{k=0}^{n-1} \int_0^1 |T_{1/m}|^k (|T_{s/m}(f)|) ds \leq 2C \frac{1}{n} \sum_{k=0}^{n-1} |T_{1/m}|^k(g) \\ &= 2C M_n(|T_{1/m}|)(g), \end{aligned}$$

which, by Theorem 2.2 applied to the operator $|T_{1/m}| \in DS^+$, entails, as $g \geq 0$,

$$\begin{aligned} \mu \{ |M_{n/m}(\mathcal{T}, \beta)(f)| > \lambda \} &\leq \mu \left\{ M_n(|T_{1/m}|)(g) > \frac{\lambda}{2C} \right\} \\ &\leq \left(\frac{4C \|g\|_p}{\lambda} \right)^p \leq \left(\frac{4C \|f\|_p}{\lambda} \right)^p \end{aligned}$$

for all $m, n \in \mathbb{N}$, hence

$$\mu \left\{ \sup_{0 < r \in \mathbb{Q}} |M_r(\mathcal{T}, \beta)(f)| > \lambda \right\} \leq \left(\frac{4C \|f\|_p}{\lambda} \right)^p.$$

To finish the argument, fix $f \in \mathcal{L}^p$ and $\lambda > 0$ and denote

$$f_t = |M_t(\mathcal{T}, \beta)(f)|, \quad E = \left\{ \sup_{0 < r \in \mathbb{Q}} f_r > \lambda \right\}.$$

Then we have

$$\mu(E) \leq \left(\frac{4C \|f\|_p}{\lambda} \right)^p \quad \text{and} \quad \|f_r \chi_{\Omega \setminus E}\|_\infty \leq \lambda \quad \forall 0 < r \in \mathbb{Q}.$$

If $t > 0$ and $0 < r_n \rightarrow t$ with $\{r_n\} \subset \mathbb{Q}$, then we have $f_{r_n} \rightarrow f_t$ in \mathcal{L}^p , hence in measure. Then, by Proposition 2.4, there exists a subsequence $\{r_{n_k}\}$ such that $f_{r_{n_k}} \rightarrow f$ a.u., implying that

$$\|f_t \chi_{\Omega \setminus E}\|_\infty \leq \lambda,$$

hence

$$\mu \{ f_t > \lambda \} \leq \mu(E) \leq \left(\frac{4C \|f\|_p}{\lambda} \right)^p \quad \forall t > 0,$$

which entails the maximal inequality (1), hence (2), for all $f \in \mathcal{L}^p$ and $\lambda > 0$. \square

Now, as a corollary of Theorem 3.1, we will show that the a.u.-convergence set of the net $\{M_t(\mathcal{T}, \beta)\}_{t>0}$ and, in particular, of $\{M_t(\mathcal{T})\}_{t>0}$, as $t \rightarrow \infty$ (or as $t \rightarrow 0$) is closed in the Banach space $(\mathcal{L}^p, \|\cdot\|_p)$ for each $1 \leq p < \infty$. For this we will utilize the notion of *measure topology* in \mathcal{L}^0 , which is given by the following system of neighborhoods of zero:

$$\mathcal{N}(\varepsilon, \delta) = \{f \in \mathcal{L}^0 : \mu\{|f| > \delta\} \leq \varepsilon\}, \quad \varepsilon > 0, \delta > 0.$$

Proposition 3.1. *Let $(X, \|\cdot\|)$ be a Banach space, and let $M_t : X \rightarrow \mathcal{L}^0$, $t > 0$, be a family of linear operators. If the maximal operator $M^* : (X, \|\cdot\|) \rightarrow (\mathcal{L}^0, t_\mu)$ is continuous at $0 \in X$, then the set*

$$X_c = \{f \in X : \{M_t(f)\} \text{ converges a.u. as } t \rightarrow \infty \text{ (or as } t \rightarrow 0)\}$$

is closed in X .

Proof. Assume that $t \rightarrow \infty$ in the definition of X_c . Fix $\varepsilon > 0$ and $\delta > 0$ and let a sequence $\{f_k\} \subset X_c$ and $f \in X$ be such that $\|f - f_k\| \rightarrow 0$. Then the continuity of M^* at $0 \in X$ implies there exist f_{k_0} and $G_0 \subset \Omega$ satisfying conditions

$$\mu(\Omega \setminus G_0) \leq \frac{\varepsilon}{2} \quad \text{and} \quad \left\| \sup_{t>0} M_t(f - f_{k_0}) \chi_{G_0} \right\|_\infty \leq \frac{\delta}{3}.$$

Next, since the net $\{M_t(f_{k_0})\}$ converges a.u. as $t \rightarrow \infty$, there exist $G_1 \subset \Omega$ and $t_0 > 0$ such that

$$\mu(\Omega \setminus G_1) \leq \frac{\varepsilon}{2} \quad \text{and} \quad \|(M_{t'}(f_{k_0}) - M_{t''}(f_{k_0})) \chi_{G_1}\|_\infty \leq \frac{\delta}{3} \quad \forall t', t'' \geq t_0.$$

Now, if we define $G = G_0 \cap G_1$, then $\mu(\Omega \setminus G) \leq \varepsilon$ and

$$\begin{aligned} \|(M_{t'}(f) - M_{t''}(f)) \chi_G\|_\infty &\leq \|M_{t'}(f - f_{k_0}) \chi_G\|_\infty + \|M_{t''}(f - f_{k_0}) \chi_G\|_\infty \\ &\quad + \|(M_{t'}(f_{k_0}) - M_{t''}(f_{k_0})) \chi_G\|_\infty \leq \delta \quad \forall t', t'' \geq t_0, \end{aligned}$$

thus, the net $\{M_t(f)\}_{t>0}$ is Cauchy as $t \rightarrow \infty$. This, in view of Proposition 2.1, implies that this net converges a.u. as $t \rightarrow \infty$ in \mathcal{L}^0 , hence $f \in X_c$.

The case $t \rightarrow 0$ is dealt with in the same manner. \square

Corollary 3.1. *Given $1 \leq p < \infty$, the a.u.-convergence set of the net $\{M_t(\mathcal{T}, \beta)\}_{t>0}$, hence of $\{M_t(\mathcal{T})\}_{t>0}$, as $t \rightarrow \infty$ (or as $t \rightarrow 0$) is closed in \mathcal{L}^p .*

Proof. Without loss of generality, assume that $C > 0$. Then, given $\varepsilon > 0$ and $\delta > 0$, condition $\|f\|_p \leq \frac{\varepsilon^{1/p} \delta}{4C}$ implies, by Theorem 3.1, that

$$\mu\{M(\mathcal{T}, \beta)^*(f) > \delta\} \leq \varepsilon,$$

hence $M(\mathcal{T}, \beta)^*(f) \in \mathcal{N}(\varepsilon, \delta)$. Therefore $M(\mathcal{T}, \beta)^* : (\mathcal{L}^p, \|\cdot\|_p) \rightarrow (\mathcal{L}^0, t_\mu)$ is continuous at zero in \mathcal{L}^p , and the statement follows from Proposition 3.1. \square

4. ALMOST UNIFORM CONVERGENCE OF NON-WEIGHTED AVERAGES IN $\mathcal{L}^p(\Omega, \mu)$

In this section we aim to establish a.u. convergence of non-weighted ergodic averages with continuous time in \mathcal{L}^p . For a.e. convergence of such averages, see, for example, [22].

We begin with the case of a *flow*, that is, when $t \rightarrow \infty$:

Theorem 4.1. *Given $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, the averages $M_t(\mathcal{T})(f)$ converge a.u. to some $\widehat{f} \in \mathcal{L}^p$ as $t \rightarrow \infty$.*

Proof. Pick $h \in \mathcal{L}^1 \cap \mathcal{L}^\infty$ and denote $g = \int_0^1 T_s(h) ds$. As in the proof of Theorem 3.1, one can see that

$$M_n(\mathcal{T})(h) = \sum_{k=0}^{n-1} T_1^k(g) = M_n(T_1)(g), \quad n = 1, 2, \dots$$

Therefore, by [6, Theorem 3.5], we conclude that $M_n(\mathcal{T})(h) \rightarrow \widehat{h}$ a.u. for some $\widehat{h} \in \mathcal{L}^p$. Thus, given $\varepsilon > 0$, there exists $G \subset \Omega$ such that $\mu(\Omega \setminus G) \leq \varepsilon$ and

$$\|(M_n(\mathcal{T})(h) - \widehat{h})\chi_G\|_\infty \rightarrow 0.$$

Next, given $t > 1$, we have

$$\begin{aligned} \|M_t(\mathcal{T})(h) - M_{[t]}(\mathcal{T})(h)\|_\infty &= \left\| \frac{1}{t} \int_0^t T_s(h) ds - \frac{1}{[t]} \int_0^{[t]} T_s(h) ds \right\|_\infty \\ &= \left\| \frac{1}{t} \int_{[t]}^t T_s(h) ds - \left(\frac{1}{t} - \frac{1}{[t]} \right) \int_0^{[t]} T_s(h) ds \right\|_\infty \\ &\leq 2 \left(1 - \frac{[t]}{t} \right) \|h\|_\infty \rightarrow 0 \quad \text{as } t \rightarrow \infty. \end{aligned}$$

Since $\{M_{[t]}(\mathcal{T})(h)\}_{t>1} = \{M_n(\mathcal{T})(h)\}_{n=1}^\infty$ as sequences, it follows that

$$\begin{aligned} \|(M_t(\mathcal{T})(h) - \widehat{h})\chi_G\|_\infty &= \|M_t(\mathcal{T})(h) - M_{[t]}(\mathcal{T})(h)\|_\infty \\ &\quad + \|(M_{[t]}(\mathcal{T})(h) - \widehat{h})\chi_G\|_\infty \rightarrow 0 \quad \text{as } t \rightarrow \infty, \end{aligned}$$

and we conclude $M_t(\mathcal{T})(h) \rightarrow \widehat{h}$ a.u. for all $h \in \mathcal{L}^1 \cap \mathcal{L}^\infty$.

Thus, since the set $\mathcal{L}^1 \cap \mathcal{L}^\infty$ is dense in \mathcal{L}^p , an application of Corollary 3.1 yields that for every $f \in \mathcal{L}^p$ there exists $\widehat{f} \in \mathcal{L}^0$ such that $M_t(\mathcal{T})(f) \rightarrow \widehat{f}$ a.u. Then, there is a sequence $t_n \rightarrow \infty$ for which $M_{t_n}(\mathcal{T})(f) \rightarrow \widehat{f}$ a.u. as $n \rightarrow \infty$. Hence, as the sequence $\{M_{t_n}(\mathcal{T})(f)\}_{n=1}^\infty$ is bounded in \mathcal{L}^p , Corollary 2.1 implies that $\widehat{f} \in \mathcal{L}^p$. \square

Next, we shall establish a.u. convergence in the *local ergodic theorem*, that is, when $t \rightarrow 0^+$:

Theorem 4.2. *Given $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, the averages $M_t(\mathcal{T})(f)$ converge to f as $t \rightarrow 0$.*

To prove Theorem 4.2, we will need a local mean ergodic theorem. Let us begin with a lemma:

Lemma 4.1. *If $t_0 > 0$, $g \in \mathcal{L}^p$, and $f = M_{t_0}(g)$, then $\|M_t(\mathcal{T})(f) - f\|_p \rightarrow 0$ as $t \rightarrow 0$. In addition, if $g \in \mathcal{L}^1 \cap \mathcal{L}^\infty$, then $\|M_t(\mathcal{T})(f) - f\|_\infty \rightarrow 0$ as $t \rightarrow 0$.*

Proof. Let $0 < t < t_0$. Then we have

$$\begin{aligned}
 \|M_t(\mathcal{T})(f) - f\|_p &= \left\| \frac{1}{t} \int_0^t (T_s(f) - f) ds \right\|_p \leq \|T_s(f) - f\|_p \\
 &= \frac{1}{t_0} \left\| \int_0^{t_0} T_{s+t}(g) ds - \int_0^{t_0} T_s(g) ds \right\|_p \\
 &= \frac{1}{t_0} \left\| \int_t^{t_0+t} T_s(g) ds - \int_0^{t_0} T_s(g) ds \right\|_p \\
 &= \frac{1}{t_0} \left\| \int_t^{t_0} T_s(g) ds + \int_{t_0}^{t_0+t} T_s(g) ds - \int_0^t T_s(g) ds - \int_t^{t_0} T_s(g) ds \right\|_p \\
 &= \frac{1}{t_0} \left\| \int_{t_0}^{t_0+t} T_s(g) ds - \int_0^t T_s(g) ds \right\|_p \leq \frac{2t}{t_0} \|g\|_p \rightarrow 0 \quad \text{as } t \rightarrow 0.
 \end{aligned}$$

Proof of the second statement is carried out exactly the same way by simply replacing L^1 -norm by L^∞ -norm. \square

Here is the local mean ergodic theorem in \mathcal{L}^p :

Theorem 4.3. *If $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, then $\|M_t(\mathcal{T})(f) - f\|_p \rightarrow 0$ as $t \rightarrow 0$.*

Proof. Since, by Proposition 2.2, the semigroup $\{T_s\}_{t \geq 0}$ is strongly continuous in \mathcal{L}^p , it follows that

$$\|T_s(f) - f\|_p \rightarrow 0 \quad \text{as } s \rightarrow 0.$$

Therefore, for every $n \in \mathbb{N}$ there exists $t_n > 0$ such that

$$\|T_s(f) - f\|_p < \frac{1}{n} \quad \forall 0 < s \leq t_n$$

hence, denoting $f_n = M_{t_n}(f)$, we have

$$\|f_n - f\|_p = \frac{1}{t_n} \left\| \int_0^{t_n} (T_s(f) - f) ds \right\|_p < \frac{1}{n}.$$

Fix $\varepsilon > 0$ and choose n_0 to be such that

$$\|f_{n_0} - f\|_p < \frac{\varepsilon}{4}.$$

Further, by Lemma 4.1,

$$\|M_t(\mathcal{T})(f_{n_0}) - f_{n_0}\|_p \rightarrow 0 \quad \text{as } t \rightarrow 0,$$

thus, there exists $t_0 > 0$ such that

$$\|M_t(\mathcal{T})(f_{n_0}) - f_{n_0}\|_p < \frac{\varepsilon}{2} \quad \forall 0 < t < t_0.$$

Now, we have

$$\begin{aligned}
 \|M_t(\mathcal{T})(f) - f\|_p &\leq \|M_t(\mathcal{T})(f) - M_t(\mathcal{T})(f_{n_0})\|_p + \|M_t(\mathcal{T})(f_{n_0}) - f_{n_0}\|_p \\
 &\quad + \|f_{n_0} - f\|_p \leq 2\|f_{n_0} - f\|_p + \frac{\varepsilon}{2} < \varepsilon
 \end{aligned}$$

whenever $0 < t < t_0$, thus completing the argument. \square

Proof of Theorem 4.2. Let a sequence of real numbers $t_n \rightarrow 0$. Then, by Theorem 4.3, the set

$$\mathcal{D} = \{M_{t_n}(\mathcal{T})(h) : n \in \mathbb{N}, h \in \mathcal{L}^1 \cap \mathcal{L}^\infty\}$$

is dense in $\mathcal{L}^1 \cap \mathcal{L}^\infty$, hence in \mathcal{L}^p . Besides, by Lemma 4.1,

$$\|M_{t_n}(g) - g\|_\infty \rightarrow 0, \text{ hence } M_{t_n}(\mathcal{T})(g) \rightarrow g \text{ a.u., } \forall g \in \mathcal{D}.$$

Now, as Corollary 3.1 implies that the a.u.-convergence set of the averages $\{M_{t_n}(\mathcal{T})\}$ is closed in \mathcal{L}^p , we conclude that the sequence $\{M_{t_n}(\mathcal{T})(f)\}$ converges a.u. for all $f \in \mathcal{L}^p$. Further, given $f \in \mathcal{L}^p$, Theorem 4.3 yields that $M_{t_n}(\mathcal{T})(f) \rightarrow f$ in \mathcal{L}^p , hence in measure. Therefore, by Proposition 2.4, $M_{t_{n_k}}(\mathcal{T})(f) \rightarrow f$ a.u. for some subsequence $\{t_{n_k}\}$, and we conclude that

$$M_{t_n}(\mathcal{T})(f) \rightarrow f \text{ a.u. as } t_n \rightarrow 0 \forall f \in \mathcal{L}^p.$$

Next, assume that $f \in \mathcal{L}^1 \cap \mathcal{L}^\infty$. Then we have $M_{1/n}(\mathcal{T})(f) \rightarrow f$ a.u., in particular,

$$M_{[1/t]^{-1}}(f) \rightarrow f \text{ a.u. as } t \rightarrow 0.$$

So, for a fix $\varepsilon > 0$, there exists $G \subset \Omega$ such that $\mu(\Omega \setminus G) \leq \varepsilon$ and

$$\|(M_{[1/t]^{-1}}(f) - f)\chi_G\|_\infty \rightarrow 0 \text{ as } t \rightarrow 0.$$

Further,

$$\begin{aligned} \|M_t(\mathcal{T})(f) - M_{[1/t]^{-1}}(\mathcal{T})(f)\|_\infty &= \left\| \frac{1}{t} \int_0^t T_s(f) ds - \left[\frac{1}{t} \right] \int_0^{[1/t]^{-1}} T_s(f) ds \right\|_\infty \\ &= \left\| \left(\frac{1}{t} - \left[\frac{1}{t} \right] \right) \int_0^t T_s(f) ds - \left[\frac{1}{t} \right] \int_t^{[1/t]^{-1}} T_s(f) ds \right\|_\infty \\ &\leq 2 \left(1 - \left[\frac{1}{t} \right] t \right) \|f\|_\infty \rightarrow 0 \text{ as } t \rightarrow 0. \end{aligned}$$

Therefore, we can write

$$\begin{aligned} \|(M_t(\mathcal{T})(f) - f)\chi_G\|_\infty &\leq \|M_t(\mathcal{T})(f) - M_{[1/t]^{-1}}(\mathcal{T})(f)\|_\infty \\ &\quad + \|(M_{[1/t]^{-1}}(f) - f)\chi_G\|_\infty \rightarrow 0 \text{ as } t \rightarrow 0. \end{aligned}$$

Thus, the averages $\{M_t(\mathcal{T})\}_{t>0}$ converge a.u. on the set $\mathcal{L}^1 \cap \mathcal{L}^\infty$, which is dense in \mathcal{L}^p , implying, in view of Corollary 3.1, that the net $\{M_t(\mathcal{T})(f)\}_{t>0}$ converges a.u. as $t \rightarrow 0$ for all $f \in \mathcal{L}^p$. This, in turn, entails that $M_t(\mathcal{T})(f) \rightarrow f$ a.u. as $t \rightarrow 0$, in the same way as in the first part of this proof, as we now know that, given $f \in \mathcal{L}^p$, there is a sequence $t_n \rightarrow 0$ such that $\{M_{t_n}(\mathcal{T})(f)\}$ converges a.u. \square

5. A.U. CONVERGENCE OF THE WEIGHTED AVERAGES IN $\mathcal{L}^p(\Omega, \mu)$

Let \mathbb{C} be the field of complex numbers, and let $\mathbb{C}_1 = \{z \in \mathbb{C} : |z| = 1\}$. A function $p : \mathbb{R}_+ \rightarrow \mathbb{C}$ is called a *trigonometric polynomial* if $p(s) = \sum_{j=1}^n w_j \lambda_j^s$, where $n \in \mathbb{N}$, $\{w_j\}_1^n \subset \mathbb{C}$, and $\{\lambda_j\}_1^n \subset \mathbb{C}_1$.

A Lebesgue measurable function $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ will be called *bounded Besicovitch (locally Besicovitch) function* if $\|\beta\|_\infty < \infty$, and for every $\varepsilon > 0$ there is a trigonometric polynomial p_ε such that

$$(3) \quad \limsup_{t \rightarrow \infty} \frac{1}{t} \int_0^t |\beta(s) - p_\varepsilon(s)| ds < \varepsilon$$

(respectively,

$$(4) \quad \limsup_{t \rightarrow 0^+} \frac{1}{t} \int_0^t |\beta(s) - p_\varepsilon(s)| ds < \varepsilon).$$

Let ν stand for Lebesgue measure in \mathbb{C}_1 and consider the product measure space

$$(\mathbb{C}_1 \otimes \Omega, \nu \otimes \mu) = (\mathbb{C}_1, \nu) \otimes (\Omega, \mu).$$

As in [8, Lemma 4.7]), we obtain the following.

Lemma 5.1. *Let (X, ν) and (Y, μ) be σ -finite measure spaces. If a net $\{g_t\}_{t \geq 0} \subset \mathcal{L}^0(X \otimes Y)$ is such that $g_t \rightarrow g$ a.u. in $X \otimes Y$ as $t \rightarrow \infty$ or $t \rightarrow 0^+$, then $g_t(x, y) \rightarrow g(x, y)$ a.u. in Y as $t \rightarrow \infty$ or $t \rightarrow 0^+$ for almost all $x \in X$.*

Recall that a semigroup $\{T_t\}_{t \geq 0} \subset DS$ is presumed to be strongly continuous in $\mathcal{L}^1 = \mathcal{L}^1(\Omega)$. Define

$$f_z(\omega) = \tilde{f}(z, \omega) \quad \text{whenever } \tilde{f} \in \mathcal{L}^1(\mathbb{C}_1 \otimes \Omega) + \mathcal{L}^\infty(\mathbb{C}_1 \otimes \Omega),$$

and note that if $\tilde{f} \in \mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)$, then, by Fubini's theorem, $f_z \in \mathcal{L}^1 = \mathcal{L}^1(\Omega)$ for ν -almost all $z \in \mathbb{C}_1$.

Next, given $\lambda \in \mathbb{C}_1$, define

$$T_s^\lambda(\tilde{f})(z, \omega) = T_s(f_{\lambda^s z})(\omega) \quad \text{for } s \geq 0 \quad \text{and } \tilde{f} \in \mathcal{L}^1(\mathbb{C}_1 \otimes \Omega) + \mathcal{L}^\infty(\mathbb{C}_1 \otimes \Omega).$$

Then it is easily verified that $\{T_s^{(\lambda)}\}_{s \geq 0} \subset DS$ on $\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega) + \mathcal{L}^\infty(\mathbb{C}_1 \otimes \Omega)$ for each $\lambda \in \mathbb{C}_1$. Besides, given $r, s \geq 0$, we have

$$\begin{aligned} T_{r+s}^{(\lambda)}(\tilde{f})(z, \omega) &= T_{r+s}(f_{\lambda^{r+s} z})(\omega) = T_r(T_s(f_{\lambda^s \lambda^r z}))(\omega) = T_r(T_s^{(\lambda)}(f_{\lambda^r z}))(\omega) \\ &= T_s^{(\lambda)}(T_r(f_{\lambda^r z}))(\omega) = T_r^{(\lambda)} T_s^{(\lambda)}(\tilde{f})(z, \omega), \end{aligned}$$

that is, $\{T_t^{(\lambda)}\}_{t \geq 0}$ is a semigroup.

Proposition 5.1. *The semigroup $\{T_s^{(\lambda)}\}_{s \geq 0}$ is strongly continuous in $\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)$ for each $\lambda \in \mathbb{C}_1$.*

Proof. Clearly, it suffices to show that, given $\tilde{f} \in \mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)$ and $\lambda \in \mathbb{C}_1$, we have

$$\|T_{s_n}^{(\lambda)}(\tilde{f}) - T_s^{(\lambda)}(\tilde{f})\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)} \rightarrow 0 \quad \text{as } s_n \searrow s \quad \text{and as } s \nearrow s.$$

If $s_n \searrow s$ ($s_n \nearrow s$), then, as

$$\|T_{s_n}^{(\lambda)}(\tilde{f}) - T_s^{(\lambda)}(\tilde{f})\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)} \leq \|T_{s_n-s}^{(\lambda)}(\tilde{f}) - \tilde{f}\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)}$$

(respectively, as

$$\|T_{s_n}^{(\lambda)}(\tilde{f}) - T_s^{(\lambda)}(\tilde{f})\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)} \leq \|T_{s-s_n}^{(\lambda)}(\tilde{f}) - \tilde{f}\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)},$$

we conclude that it is sufficient to verify that

$$\|T_{s_n}^{(\lambda)}(\tilde{f}) - \tilde{f}\|_{\mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)} = \int_{\mathbb{C}_1} \|T_{s_n}(\tilde{f}(\lambda^{s_n} z, \cdot)) - \tilde{f}(z, \cdot)\|_1 d\nu(z) \rightarrow 0 \quad \text{as } s_n \rightarrow 0^+.$$

We have

$$\begin{aligned} \|T_{s_n}(\tilde{f}(\lambda^{s_n} z, \cdot)) - \tilde{f}(z, \cdot)\|_1 &\leq \|T_{s_n}(\tilde{f}(\lambda^{s_n} z, \cdot)) - T_{s_n}(\tilde{f}(z, \cdot))\|_1 \\ &\quad + \|T_{s_n}(\tilde{f}(z, \cdot)) - \tilde{f}(z, \cdot)\|_1 \\ &\leq \|\tilde{f}(\lambda^{s_n} z, \cdot) - \tilde{f}(z, \cdot)\|_1 + \|T_{s_n}(\tilde{f}(z, \cdot)) - \tilde{f}(z, \cdot)\|_1. \end{aligned}$$

Since $\{T_s\}_{s \geq 0}$ is strongly continuous in \mathcal{L}^1 , it follows that

$$\|T_{s_n}(\tilde{f}(z, \cdot)) - \tilde{f}(z, \cdot)\|_1 \text{ for } \nu\text{-almost all } z \in \mathbb{C}_1,$$

which, together with $\|T_{s_n}(\tilde{f}(z, \cdot)) - \tilde{f}(z, \cdot)\|_1 \leq 2\|\tilde{f}(z, \cdot)\|_1$ ν -a.e. on \mathbb{C}_1 , implies that

$$\int_{\mathbb{C}_1} \|T_{s_n}(\tilde{f}(z, \cdot)) - \tilde{f}(z, \cdot)\|_1 d\nu(z) \rightarrow 0 \text{ as } s_n \rightarrow 0^+.$$

Thus, it remains to verify that

$$(5) \quad \int_{\mathbb{C}_1} \|\tilde{f}(\lambda^{s_n} z, \cdot) - \tilde{f}(z, \cdot)\|_1 d\nu(z) \rightarrow 0 \text{ as } s_n \rightarrow 0^+.$$

To that end, let

$$(6) \quad \tilde{h}(z, \omega) = \sum_{i=1}^m h_i \chi_{[\lambda_1, \lambda_{i+1}]}(z), \quad z \in \mathbb{C}_1, \quad h_i \in \mathcal{L}^1 \text{ for } i = 1, \dots, m$$

be a simple Bochner-measurable function, where $\{\lambda_i\}_{i=1}^{m+1}$, with $\lambda_{m+1} = \lambda_1$, is a partition of \mathbb{C}_1 . Then, given $s \in \mathbb{R}$, we have

$$\tilde{h}(\lambda^s z, \omega) = \sum_{i=1}^m h_i \chi_{[\lambda^s \lambda_1, \lambda^s \lambda_{i+1}]}(z),$$

where $\{\lambda^s \lambda_i\}_{i=1}^{m+1}$ is another partition of \mathbb{C}_1 . Since $\lambda^{s_n} \lambda_i \rightarrow \lambda_i$ as $s_n \rightarrow 0$ for each i , it follows that

$$\int_{\mathbb{C}_1} \|\tilde{h}(\lambda^{s_n} z, \cdot) - \tilde{h}(z, \cdot)\|_1 d\nu(z) \leq \sum_{i=1}^m |\chi_{[\lambda^{s_n} \lambda_i, \lambda^{s_n} \lambda_{i+1}]} - \chi_{[\lambda_i, \lambda_{i+1}]}| \|h_i\|_1 \rightarrow 0$$

as $s_n \rightarrow 0^+$.

Now, as $\tilde{f} \in \mathcal{L}^1(\mathbb{C}_1 \otimes \Omega)$, one can justify that there exists a sequence $\{\tilde{h}_k(z, \omega)\}$ of functions of the form (6) for which

$$\int_{\mathbb{C}_1} \|\tilde{f}(z, \cdot) - \tilde{h}_k(z, \cdot)\|_1 d\nu(z) \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Then we have

$$\begin{aligned} \int_{\mathbb{C}_1} \|\tilde{f}(\lambda^{s_n} z, \cdot) - \tilde{f}(z, \cdot)\|_1 d\nu(z) &\leq \int_{\mathbb{C}_1} \|\tilde{f}(\lambda^{s_n} z, \cdot) - \tilde{h}_k(\lambda^{s_n} z, \cdot)\|_1 d\nu(z) \\ &\quad + \int_{\mathbb{C}_1} \|\tilde{h}_k(\lambda^{s_n} z, \cdot) - \tilde{h}_k(z, \cdot)\|_1 d\nu(z) \\ &\quad + \int_{\mathbb{C}_1} \|\tilde{h}_k(z, \cdot) - \tilde{f}(z, \cdot)\|_1 d\nu(z) = 2 \int_{\mathbb{C}_1} \|\tilde{h}_k(z, \cdot) - \tilde{f}(z, \cdot)\|_1 d\nu(z) \\ &\quad + \int_{\mathbb{C}_1} \|\tilde{h}_k(\lambda^{s_n} z, \cdot) - \tilde{h}_k(z, \cdot)\|_1 d\nu(z) \rightarrow 0 \text{ as } s_n \rightarrow 0^+ \quad \forall k, \end{aligned}$$

and (5) follows, completing the proof. \square

Lemma 5.2. *Let a semigroup $\mathcal{T} = \{T_s\}_{s \geq 0} \subset DS$ be strongly continuous in \mathcal{L}^1 , and let $p(s) = \sum_{j=1}^n w_j \lambda_j^s$ be a trigonometric polynomial. If $1 \leq p < \infty$, $f \in \mathcal{L}^p$, and*

$$(7) \quad P_t(\mathcal{T})(f) = \frac{1}{t} \int_0^t p(s) T_s(f) ds,$$

then

- (i) the net $P_t(\mathcal{T})(f)$ converges a.u. as $t \rightarrow \infty$;
- (ii) the net $P_t(\mathcal{T})(f)$ converges a.u. to $p(0)f$ as $t \rightarrow 0$.

Proof. (i) Fix $\lambda \in \mathbb{C}_1$. In view of Propositions 2.2 and 5.1, $\{T_s^{(\lambda)}\}_{s \geq 0} \subset DS$ is a semigroup strongly continuous in $\mathcal{L}^p(\mathbb{C}_1 \otimes \Omega)$. Then, by Theorem 4.1, given $\tilde{f} \in \mathcal{L}^p(\mathbb{C}_1 \otimes \Omega)$, the net

$$(8) \quad \frac{1}{t} \int_0^t T_s^{(\lambda)}(\tilde{f}) ds$$

converges a.u. as $t \rightarrow \infty$. Therefore, by Lemma 5.1, the net

$$\frac{1}{t} \int_0^t T_s^{(\lambda)}(\tilde{f})(z, \omega) ds = \frac{1}{t} \int_0^t T_s(\tilde{f}(\lambda^s z, \omega)) ds$$

converges a.u. as $t \rightarrow \infty$ ν -a.e. on \mathbb{C}_1 . In particular, letting $\tilde{f}(z, \omega) = zf(\omega)$, we conclude that the net

$$z \frac{1}{t} \int_0^t \lambda^s T_s(f) ds$$

converges a.u. as $t \rightarrow \infty$ for some $0 \neq z \in \mathbb{C}_1$, implying that the net

$$\frac{1}{t} \int_0^t \lambda^s T_s(x) ds$$

converges a.u. as $t \rightarrow \infty$. Therefore, by linearity, the net $P_t(\mathcal{T})(x)$ converges a.u. as $t \rightarrow \infty$.

(ii) Now, by Theorem 4.3, given $\tilde{f} \in \mathcal{L}^p(\mathbb{C}_1 \otimes \Omega)$, it follows that the net (8) converges a.u. to \tilde{f} as $t \rightarrow 0$. Then, letting $f(z, \omega) = zf(\omega)$, we see as above that

$$\frac{1}{t} \int_0^t \lambda^s T_s(f) ds \rightarrow f \quad \text{a.u.}$$

as $t \rightarrow 0$, and the result follows by linearity. \square

Now we are ready to present the main results of this section.

Theorem 5.1. *Let a semigroup $\{T_s\}_{s \geq 0} \subset DS$ be strongly continuous in \mathcal{L}^1 , and let $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ be a bounded Besicovitch function with $\|\beta\|_\infty < C < \infty$. Then, given $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, the net $\{M_t(\mathcal{T}, \beta)(f)\}_{t > 0}$ converges a.u. as $t \rightarrow \infty$ to some $\hat{f} \in \mathcal{L}^p$.*

Proof. Assume that $f \in \mathcal{L}^1 \cap \mathcal{L}^\infty$. Fix $\varepsilon > 0$ and choose a trigonometric polynomial $p = p_\varepsilon$ to satisfy condition (3). Let $\{P_t(x)\}_{t > 0}$ be given by (7). Then we have

$$(9) \quad \|M_t(\mathcal{T}, \beta)(f) - P_t(\mathcal{T})(f)\|_\infty \leq \frac{1}{t} \int_0^t |\beta(s) - p(s)| ds \|f\|_\infty < \varepsilon \|f\|_\infty$$

for all big enough values of t . Since, by Lemma 5.2, the averages $P_t(\mathcal{T})(f)$ converge a.u., it follows that the net $\{M_t(\mathcal{T}, \beta)(f)\}_{t > 0}$ is a.u. Cauchy as $t \rightarrow \infty$. Therefore, by Proposition 2.1, this net converges to some $\hat{f} \in \mathcal{L}^0$, and as in the end of proof of Theorem 4.1, it follows that $\hat{f} \in \mathcal{L}^p$. \square

Theorem 5.2. *Let a semigroup $\{T_t\}_{t \geq 0} \subset DS$ be strongly continuous in \mathcal{L}^1 , and let $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ be a bounded locally Besicovitch function with $\|\beta\|_\infty < C < \infty$. Then, given $1 \leq p < \infty$ and $f \in \mathcal{L}^p$, the averages $\{M_t(\mathcal{T}, \beta)(f)\}_{t > 0}$ converge a.u. to $\lambda(f)f$ as $t \rightarrow 0$ for some $\lambda(f) \in \mathbb{C}$.*

Proof. Assume that $f \in \mathcal{L}^1 \cap \mathcal{L}^\infty$. Fix $\varepsilon > 0$ and choose a trigonometric polynomial $p = p_\varepsilon$ to satisfy condition (4). If the net $\{P_t(\mathcal{T})(f)\}_{t>0}$ is given by (7), then (9) holds for all small enough positive values of t . Since, by Lemma 5.2, the averages $P_t(\mathcal{T})(f)$ converge a.u. as $t \rightarrow 0$, it follows that the net $\{M_t(\mathcal{T}, \beta)(f)\}_{t>0}$ is a.u. Cauchy as $t \rightarrow 0$. Then, as in Theorem 5.1, we conclude that for any $f \in \mathcal{L}^p$ the averages $\{M_t(\mathcal{T}, \beta)(f)\}_{t>0}$ converge a.u. to some $\widehat{f} \in \mathcal{L}^p$ as $t \rightarrow 0$.

Let p_n be a trigonometric polynomial to satisfy condition (4) with $\varepsilon = 1/n$, and let $\{P_t^{(n)}(\mathcal{T})(f)\}_{t>0}$ be the corresponding averages given by (7). Then there exists $t_n > 0$ such that

$$(10) \quad \|M_t(\mathcal{T}, \beta)(f) - P_t^{(n)}(\mathcal{T})(f)\|_p \leq \frac{1}{t} \int_0^t |\beta(s) - p_n(s)| ds \|f\|_p < \frac{\|f\|_p}{n}$$

holds for all $0 < t < t_n$.

By Lemma 5.2, $P_t^{(n)}(\mathcal{T})(f) \rightarrow p_n(0) f$ a.u. as $t \rightarrow 0$, implying that there exists a sequence $t_m \rightarrow 0$ such that

$$M_{t_m}(\mathcal{T}, \beta)(f) - P_{t_m}^{(n)}(\mathcal{T})(f) \rightarrow \widehat{f} - p_n(0) f \quad \text{a.u. as } m \rightarrow \infty.$$

Since, by Corollary 2.1, the unit ball in \mathcal{L}^p is sequentially closed with respect to a.u. convergence, (10) entails that

$$\|\widehat{f} - p_n(0) f\|_p \leq \frac{\|f\|_p}{n},$$

hence $\widehat{f} = \|\cdot\|_p - \lim_{n \rightarrow \infty} p_n(0) f$, and we conclude that $\widehat{f} = \lambda(f) f$ for some $\lambda(f) \in \mathbb{C}$, which completes the argument. \square

6. EXTENSION TO FULLY SYMMETRIC SPACES

Now we aim to extend the results of Section 5 to the fully symmetric spaces $E \subset \mathcal{L}^1 + \mathcal{L}^\infty$ with $\mathbf{1} = \chi_\Omega \notin E$.

As before, a semigroup $\mathcal{T} = \{T_s\}_{s>0} \subset DS$ is assumed to be strongly continuous in \mathcal{L}^1 . Then, by Proposition 2.3, if $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ is a Lebesgue measurable function such that $\|\beta\|_\infty < C < \infty$, the averages $M_t(\mathcal{T}, \beta)(f)$ are defined for every $f \in \mathcal{L}^1$ and $t > 0$. Furthermore, as the operators $C^{-1} M_t(\mathcal{T}, \beta)$ contract both \mathcal{L}^1 and $\mathcal{L}^1 \cap \mathcal{L}^\infty$, we may assume, in view of Theorem 2.1, that the averages $M_t(\mathcal{T}, \beta)(f)$ are defined for all $f \in \mathcal{L}^1 + \mathcal{L}^\infty$. Note that then $C^{-1} M_t(\mathcal{T}, \beta) \in DS$ for all $t > 0$.

Denote

$$\mathcal{R}_\mu = \{f \in \mathcal{L}^1 + \mathcal{L}^\infty : \mu\{|f| > \lambda\} < \infty \text{ for all } \lambda > 0\},$$

the Fava's space – see [11, Section (2.5.5)]. (One can observe that $\bigcup_{1 \leq p < \infty} \mathcal{L}^p \subsetneq \mathcal{R}_\mu$.)

As it was shown – see [4, 6, 15] – that the pointwise Dunford-Schwartz ergodic theorem holds for $f \in \mathcal{L}^1 + \mathcal{L}^\infty$ if and only if $f \in \mathcal{R}_\mu$, we will only be interested in the functions that belong to the Fava's space \mathcal{R}_μ .

We will utilize the following characterization of \mathcal{R}_μ – see [8, Proposition 2.1]:

Proposition 6.1. *Let $f \in \mathcal{L}^1 + \mathcal{L}^\infty$. Then $f \in \mathcal{R}_\mu$ if and only if for each $\varepsilon > 0$ there exist $g_\varepsilon \in \mathcal{L}^1$ and $h_\varepsilon \in \mathcal{L}^\infty$ such that*

$$f = g_\varepsilon + h_\varepsilon \quad \text{and} \quad \|h_\varepsilon\|_\infty \leq \varepsilon.$$

Corollary 6.1. *If $T \in DS$, then $T(\mathcal{R}_\mu) \subset \mathcal{R}_\mu$, hence $M_t(\mathcal{T}, \beta)(\mathcal{R}_\mu) \subset \mathcal{R}_\mu$ for every Lebesgue measurable function $\beta : \mathbb{R}_+ \rightarrow \mathbb{C}$ such that $\|\beta\|_\infty < \infty$ and $t > 0$.*

Let us also note the following – see [17, Theorem 2.3], where the proof can be I expect it to be a challenging task to solve, say, 3 problems in 10 minutes without making sure that you know how to do them before coming to the class, i.e., by thoroughly accomplishing your homework. straightforwardly adapted to the current commutative setting:

Proposition 6.2. *The space \mathcal{R}_μ is complete with respect to a.u. convergence.*

Theorem 6.1. *If a semigroup $\mathcal{T} = \{T_s\}_{s \geq 0} \subset DS$ is strongly continuous in \mathcal{L}^1 and β is a bounded Besicovitch (locally Besicovitch) function, then for every $f \in \mathcal{R}_\mu$ the averages $M_t(\mathcal{T}, \beta)(f)$ converge to some $\hat{f} \in \mathcal{R}_\mu$ as $t \rightarrow \infty$ (respectively, as $t \rightarrow 0$).*

Proof. Assume that β is a bounded Besicovitch function with $\|\beta\|_\infty < C < \infty$. Fix $\varepsilon > 0$ and $\delta > 0$. By Proposition 6.1, there exist $g \in \mathcal{L}^1$ and $h \in \mathcal{L}^\infty$ such that

$$f = g + h \quad \text{and} \quad \|h\|_\infty \leq \frac{\delta}{3C}.$$

Since $g \in \mathcal{L}^1$, Theorem 5.1 implies that there exists $G \subset \Omega$ and $t_0 > 0$ such that

$$\mu(\Omega \setminus G) \leq \varepsilon \quad \text{and} \quad \|(M_{t'}(\mathcal{T}, \beta)(g) - M_{t''}(\mathcal{T}, \beta)(g))\chi_G\|_\infty \leq \frac{\delta}{3} \quad \forall t', t'' \geq t_0.$$

Then, given $t', t'' \geq t_0$, we have

$$\begin{aligned} \|(M_{t'}(\mathcal{T}, \beta)(f) - M_{t''}(\mathcal{T}, \beta)(f))\chi_G\|_\infty &\leq \|(M_{t'}(\mathcal{T}, \beta)(g) - M_{t''}(\mathcal{T}, \beta)(g))\chi_G\|_\infty \\ &\quad + \|(M_{t'}(\mathcal{T}, \beta)(h) - M_{t''}(\mathcal{T}, \beta)(h))\chi_G\|_\infty \\ &\leq \frac{\delta}{3} + \|M_{t'}(\mathcal{T}, \beta)(h)\|_\infty + \|M_{t''}(\mathcal{T}, \beta)(h)\|_\infty \leq \frac{\delta}{3} + 2C\|h\|_\infty \leq \delta. \end{aligned}$$

Therefore, the net $\{M_t(\mathcal{T}, \beta)(f)\}_{t > 0} \subset \mathcal{R}_\mu$ is a.u. Cauchy, implying, by Proposition 6.2, that it converges a.u. to some $\hat{f} \in \mathcal{R}_\mu$.

Next, let us assume that β is a bounded locally Besicovitch function. Then, repeating the above argument with Theorem 5.2 instead of Theorem 5.1, we conclude that the net $\{M_t(\mathcal{T}, \beta)(f)\}_{t > 0}$ converges a.u. to some $\hat{f} \in \mathcal{R}_\mu$ as $t \rightarrow 0$. \square

Define

$$\mathcal{L}_\mu^0 = \{f \in \mathcal{L}^0 : \mu\{|f| > \lambda\} < \infty \text{ for some } \lambda > 0\}.$$

Given $f \in \mathcal{L}_\mu^0$, the *non-increasing rearrangement* of f is defined as

$$f^*(t) = \inf \{\lambda > 0 : \mu\{|f| > \lambda\} < t\}, \quad t > 0.$$

A non-zero linear subspace $E \subset \mathcal{L}_\mu^0$ with a Banach norm $\|\cdot\|_E$ is called *symmetric* (fully symmetric) if $g \in E$, $f \in \mathcal{L}_\mu^0$, and

$$f^*(t) \leq g^*(t) \quad (\text{respectively, } \int_0^s f^*(t)dt \leq \int_0^s g^*(t)dt \text{ (in short, } f \prec\prec g)) \quad \forall s > 0$$

imply that $f \in E$ and $\|f\|_E \leq \|g\|_E$.

Immediate examples of fully symmetric spaces are the spaces $\mathcal{L}^p = \mathcal{L}^p(\Omega, \mu)$, $1 \leq p \leq \infty$, with the standard norms $\|\cdot\|_p$. Further examples include the well-known Orlicz, Lorentz, and Marcinkiewicz spaces – for details, see, for example, [6, Sec. 5], [8, Sec. 6].

Here is a criterion for a symmetric space to belong to the space \mathcal{R}_μ – see [8, Proposition 6.1]:

Proposition 6.3. *Assume that $\mu(\Omega) = \infty$. Then a symmetric space E is contained in \mathcal{R}_μ if and only if $\mathbf{1} \notin E$.*

Let us now present an application of Theorem 6.1 to fully symmetric spaces:

Theorem 6.2. *Let E be a fully symmetric space such that $\mathbf{1} \notin E$, and let \mathcal{T} and $\beta(s)$ be as in Theorem 6.1. Then, given $f \in E$, the averages $M_t(\mathcal{T}, \beta)(f)$ converge a.u. to some $\hat{f} \in E$ as $t \rightarrow \infty$ (respectively, as $t \rightarrow 0$).*

Proof. Since $\mathbf{1} \notin E$, it follows that $E \subset \mathcal{R}_\tau$. Therefore, by Theorem 6.1, the averages $\{M_t(\mathcal{T}, \beta)(f)\}_{t>0}$ converge a.u. to some $\hat{f} \in \mathcal{R}_\tau$ as $t \rightarrow \infty$ (respectively, as $t \rightarrow 0$), so, it remains to show that $\hat{f} \in E$, for which we will follow the argument of that of [8, Theorem 6.2],

Let $\{t_n\}$ be a sequence such that the averages $\{M_{t_n}(\mathcal{T}, \beta)(f)\}_{n=1}^\infty$ converge a.u. when $t_n \rightarrow \infty$ (respectively, $t_n \rightarrow 0$). If C is such that $\|\beta\|_\infty < C$, then $C^{-1}M_{t_n}(\mathcal{T}, \beta) \in DS$ for every n , implying, by [13, Ch. II, § 3, Sec. 4], that

$$(11) \quad C^{-1}M_{t_n}(\mathcal{T}, \beta)(f) \prec\prec f \quad \forall n.$$

Next, as $M_{t_n}(\mathcal{T}, \beta)(f) \rightarrow \hat{f}$ also in measure, we observe, by [13, Ch. II, § 2, 11⁰], that

$$(C^{-1}M_{t_n}(\mathcal{T}, \beta)(f))^* \rightarrow (C^{-1}\hat{f})^* \quad \text{a.e. on } (0, s) \text{ as } n \rightarrow \infty \quad \forall s > 0.$$

Therefore, it follows from Fatou's lemma and (11) that

$$\int_0^s (C^{-1}\hat{f})^* dt \leq \liminf_n \int_0^s (C^{-1}M_{t_n}(\mathcal{T}, \beta)(f))^* dt \leq \int_0^s f^* dt \quad \forall s > 0,$$

hence $(\hat{f})^* \prec\prec C f^*$. Since E is a fully symmetric space and $f \in E$, we conclude that $\hat{f} \in E$. \square

Remark 6.1. One may observe that all the above arguments remain applicable if the net $\{t\}_{t>0}$ is replaced by an arbitrary net $\{t_\alpha\}_{\alpha \in \Lambda} \subset (0, \infty)$ with the directed set Λ such that $\lim_{\alpha \in \Lambda} t_\alpha = \infty$, or $\lim_{\alpha \in \Lambda} t_\alpha = 0$.

Acknowledgment. The author would like to express his gratitude to Professor Vladimir Chilin for invaluable learning experiences he had generously created over the many years of our collaboration. This work was motivated by these experiences and partially supported by the Pennsylvania State University traveling program.

REFERENCES

- [1] A. S. Besicovitch, *Almost Periodic Functions*, Cambridge Univ. Press, 1932.
- [2] J. R. Baxter, J. H. Olsen, Weighted and subsequential ergodic theorems, *Canad. J. Math.* **35** (1983) 145-166.
- [3] A. Bellow, V. Losert, The weighted pointwise ergodic theorem and the individual ergodic theorem along sequences, *Trans. Amer. Math. Soc.* **288** (1) (1985) 307-345.
- [4] V. Chilin, S. Litvinov, The validity space of Dunford–Schwartz pointwise ergodic theorem, *J. Math. Anal. Appl.* **461** (2018) 234–247.
- [5] V. Chilin, S. Litvinov, Local ergodic theorems in symmetric spaces of measurable operators, *Integr. Equ. Oper. Theory* (2019) 91:15.
- [6] V. Chilin, S. Litvinov, Almost uniform and strong convergences in ergodic theorems for symmetric spaces, *Acta Math. Hungar.* **157** (1) (2019) 229-253.

- [7] V. Chilin, S. Litvinov, Noncommutative weighted individual ergodic theorems with continuous time, *Inf. Dimen. Anal. Quant. Prob. Rel. Topics* **23** (2) (2020) (20 pages).
- [8] V. Chilin, D. Çömez, and S. Litvinov, Individual ergodic theorems for infinite measure, *Colloq. Math.* **167** (2) (2022) 219-238.
- [9] N. Dunford, J. T. Schwartz, *Linear Operators, Part I: General Theory*, Wiley, 1988.
- [10] C. Demeter, R. L. Jones, Besicovitch weights and the necessity of duality restrictions in the weighted ergodic theorem, Chapel Hill Ergodic Theory Workshops, Contemp. Math. **356**, Amer. Math. Soc., Providence, RI, (2004), pp. 127-135.
- [11] G. A. Edgar, L. Sucheston, *Stopping Times and Directed Processes* (Cambridge University Press, 1992).
- [12] A. M. Garsia, *Topics in Almost Everywhere Convergence*, Markham, 1970.
- [13] S. G. Krein, Ju. I. Petunin, and E. M. Semenov, *Interpolation of Linear Operators* (Translations of Mathematical Monographs, Amer. Math. Soc., **54**, 1982).
- [14] U. Krengel, *Ergodic Theorems*, de Gruyter, 1985.
- [15] D. Kunszenti-Kovács, Counter-examples to the Dunford–Schwartz pointwise ergodic theorem on $L^1 + L^\infty$, *Arch. Math. (Basel)* **112** (2019) 205–212.
- [16] M. Lin, J. Olsen, and A. Tempelman, On modulated ergodic theorems for Dunford-Schwartz operators, *Illinois J. Math.* **43** (3) (1999) 542-567.
- [17] S. Litvinov, Notes on noncommutative ergodic theorems, *Proc. Amer. Math. Sci.* **150** (8) (2024) 3381-3391.
- [18] S. A. McGrath, Local ergodic theorems for noncommuting semigroups, *Proc. Amer. Math. Sci.* **79** (2) (1980) 212-216.
- [19] F. Mukhamedov and A. Karimov, On noncommutative weighted local ergodic theorems on L^p -spaces, *Period. Math. Hungar.* **55** (2007) 223–235.
- [20] E. T. Ordman, Classroom Notes: Convergence almost everywhere is not topological, *Amer. Math. Monthly* **73** (2) (1966) 182–183.
- [21] C. Ryll-Nardzewski, Topics in ergodic theory, *Lecture Notes in Math.*, **472** (1975), Springer-Verlag, 131-156.
- [22] R. Sato, Ergodic theorems for d -parameter semigroups of Dunford-Schwartz operators, *Math. J. Okayama Univ.* **23** (1981) 41-57.
- [23] T. R. Terrell, The local ergodic theorem and semigroups of non-positive operators, *J. Funct. Anal.* **10** (1972) 424-429.
- [24] K. Yosida, *Functional Analysis* (Springer Verlag, Berlin-Göttingen-Heidelberg, 1965).

PENNSYLVANIA STATE UNIVERSITY, 76 UNIVERSITY DRIVE, HAZLETON, PA 18202, USA
 Email address: sn12@psu.edu