Parsings of Stationary Processes, Stopping Times and the Fundamental Pointwise Convergence Theorems of Ergodic Theory

Matan Tal

The Hebrew University of Jerusalem

Abstract

The idea of a parsing of a stationary process according to a collection of words is introduced, and the basic framework required for the asymptotic analysis of these parsings is presented. We demonstrate how the pointwise ergodic theorem and the Shannon-McMillan-Breiman theorem can be deduced from their respective weaker convergence in probability versions combined with our observations regarding parsings, where the parsings are done according to collections that originate in stopping times tailored for that purpose.

1 Introduction

For a set Λ , let us denote by σ the shift map of $\Lambda^{\mathbb{Z}}$.

Notation: For any $x \in \Lambda^{\mathbb{Z}}$ we shall denote the word $x_r x_{r+1} \dots x_{r+d}$ by x_r^{r+d} , and by $\left[x_r^{r+d}\right]$ the cylinder sub-set of $\Lambda^{\mathbb{Z}}$ determined by the word x_r^{r+d} beginning in the zero coordinate.

Let V be a collection of words in Λ such that no one word is a prefix of another. Given any $x \in \Lambda^{\mathbb{Z}}$ and a coordinate we choose to start from, a parsing of x accroding to V is obtained by moving from the starting position in the positive direction until we approach a coordinate in which an element of V begins, and then continuing to progress in the

same manner moving in the positive direction from the first coordinate after the end of that element. Notice that this parsing depends on the choice of the starting position and that it may contain gaps.

Observation: If in a portion of $1 - \varepsilon$ of the coordinates of x_1^M begin elements of V (that also end not after coordinate M) then in the parsing of x_1^M starting at the first coordinate the gaps occupy less than a portion of ε of the coordinates $1, \ldots, M$ (because in these coordinates of elements of V do not begin).

This trivial observation together with the mean ergodic theorem suffices to derive a very short proof of the pointwise ergodic theorem for bounded functions. This is what will be done in the next section.

After that we continue to study parsings. Now we assume Λ to be finite and equip $\Lambda^{\mathbb{Z}}$ with μ - a σ -invariant ergodic probability measure on $\Lambda^{\mathbb{Z}}$ - and analyze the behaviour of the parsings of outcomes of this stationary process. In particular, we deduce the following theorem. For its formulation we denote by \tilde{V} the collection obtained by adding to V all elements of Λ (treated as single letter words) - the ones that not already appear in it.

Theorem 1: Let $(\Lambda^{\mathbb{Z}}, \sigma, \mu)$ be an ergodic system (Λ a finite set). Given a finite prefixfree collection V, there exists for every $\alpha \in \tilde{V}$ (as defined above) an $0 \leq \eta_{\alpha} \leq \mu([\alpha])$ such that for an $x \in \Lambda^{\mathbb{Z}}$ μ -almost-surely exists a parsing for which every $\alpha \in \tilde{V}$ occupies asymptotically $|\alpha| \eta_{\alpha}$ of the coordinates of x. In particular, $p_{\alpha} := \frac{\eta_{\alpha}}{\sum_{\beta \in \tilde{V}} \eta_{\beta}}$ is the asymptotic frequency of appearances of each $\alpha \in \tilde{V}$ among the elements of \tilde{V} in such a parsing.

With Theorem 1 in our hands we will be ready to prove the Shannon-McMillan-Breiman theorem. The proof is rather combinatorial in its nature. Both this proof and the proof of the pointwise ergodic theorem presented utilize stopping time arguments, i.e. appropriate decompositions of the system into parts of different scales, in order to construct the collection V.

Remark: Ornstein and Weiss' proof of their recurrence-time theorem (published in

[5] and also appearing in [6]) as well as their proof of Theorem 3 stated here in section 4, are also based on a parsing of a stationary process like that is dealt with here (and the proof of the recurrence time theorem also uses a stopping time to establish the collection V). However, They exploited the special character of their problem and thus were not in need of our general analysis of parsings. In the closing argument of [1] there also is use of parsings according to collections (though quite implicitly) and also the above observation is applied. Probably these parsings appear in many more works in ergodic theory.

Acknowledgements: This work follows in spirit the book "Single Orbit Dynamics" by B. Weiss [6].

2 A Proof of the Pointwise Ergodic Theorem

In this section we prove the pointwise ergodic theorem for a bounded function by our methods that avoid use of the maximal ergodic theorem (the well-known reduction from merely an integrable function to this case does require the maximal ergodic theorem).

Proof: Assume that (X, T, ν) is an ergodic system (ν is a probability measure) and that $f: X \to [-L, L]$ is measurable.

By ergodicity, there exist constants $-L \le c' \le c \le L$ so that

 $\limsup_{N \to \infty} \frac{1}{N} \sum_{n=0}^{N-1} T^i(f(x)) = c \text{ and } \liminf_{N \to \infty} \frac{1}{N} \sum_{n=0}^{N-1} T^i(f(x)) = c' \text{ ν-almost-surely. Let us assume to the contrary that } c' < c, \text{ and assume without loss of generalitys}$ that $c > \int\limits_X f \, d\nu$ (if this is not the case then $c' < \int\limits_X f \, d\nu$ and we could work with -f).

We define \tilde{f} , a finite-valued version of f: $\tilde{f}(x) = \left| \frac{2f(x)}{c - \int_{X} f \, d\nu} \right| \frac{c - \int_{X} f \, d\nu}{2}$.

 $\limsup_{N\to\infty} \frac{1}{N} \sum_{n=0}^{N-1} T^i\left(\tilde{f}\left(x\right)\right) \text{ equals a constant } \tilde{c} \text{ ν-almost-everywhere and } \tilde{c} + \frac{c-\int f \, d\nu}{\frac{X}{2}} > c. \text{ Because also } c > \int\limits_X f \, d\nu + \frac{c-\int f \, d\nu}{\frac{X}{2}} \ge \int\limits_X \tilde{f} \, d\nu + \frac{c-\int f \, d\nu}{\frac{X}{2}} \text{ we conclude that } \tilde{c} > \int\limits_V \tilde{f} \, d\nu.$

Denoting by A the finite set $\left\{k\frac{c-\int f\,d\nu}{X}:\left[-\frac{2L}{c-\int f\,d\nu}\right] \le k \le \left[\frac{2L}{c-\int f\,d\nu}\right]\right\}$ we define $\varphi:X\to A^{\mathbb{Z}}$ by

 $\left(\varphi \left(x\right) \right) _{i}=\widetilde{f}\left(T^{i}\left(x\right) \right)$ and push-forward the measure ν to obtain a finite-valued stationary process.

For $y \in A^{\mathbb{Z}}$, we define a stopping time N_y to be the minimal $N \geq 0$ for which $\frac{1}{N} \sum_{n=0}^{N-1} y_n > \frac{\tilde{c} + \int \tilde{f} \, d\nu}{2}$. We denote by V_k the words of length k in the alphabet A that stop this stopping time. For a small $0 < \varepsilon < \frac{\tilde{c} - \int \tilde{f} \, d\nu}{3}$ to be chosen at the end, we define \tilde{K} to be the minimal K for which $\sum_{k=1}^{K} \varphi_* \nu\left(V_k\right) > 1 - \varepsilon$ (by $\varphi_* \nu\left(V_k\right)$) we mean the sum of $\varphi_* \nu$ -probabilities of all the words in V_k), and define V to be $\bigcup_{k=1}^{\tilde{K}} V_k$.

By the mean ergodic theorem, there exists an M'>0 such that for every $M\geq M'$ there is a set $E\subseteq A^{\mathbb{Z}}$ of $\varphi_*\nu$ -measure greater than $\frac{1}{2}$ such that for every $y\in E$ $\left|\frac{1}{M}\sum_{n=0}^{M-1}y_n-\int_X \tilde{f}\,d\nu\right|<\varepsilon$ and in a portion greater than $1-\varepsilon$ of the coordinates of y_0^{M-1} words of V begin (and are contained in y_0^{M-1}). $\frac{1}{2}$ was arbitrarily chosen here, any other positive number would do, for we only need to know that E is of positive measure and thus non-empty.

For any $y \in E$, the words of V in the parsing starting at the beginning of y_0^{M-1} occupy a portion greater than $1-\varepsilon$ of it (by the observation in the introduction) and the average of all the values of y_0^{M-1} in these occupied places is greater than $\frac{\tilde{c}+\int \tilde{f} \, d\nu}{\frac{X}{2}}$. Thus we obtain $\frac{1}{M}\sum_{n=0}^{M-1}y_n \geq (1-\varepsilon)\frac{\tilde{c}+\int \tilde{f} \, d\nu}{\frac{X}{2}} - \varepsilon L$. For ε small enough we get $\frac{1}{M}\sum_{n=0}^{M-1}y_n \geq \int \tilde{f} \, d\nu + \frac{\tilde{c}-\int \tilde{f} \, d\nu}{\frac{X}{3}} > \int \tilde{f} \, d\nu + \varepsilon$. A contradiction.

3 The Parsings Space

We consider again the system $(\Lambda^{\mathbb{Z}}, \sigma, \mu)$ (Λ a finite set, σ the shift map and μ a σ -invariant ergodic probability measure).

A finite collection V of words in the Λ -alphabet shall be called prefix-free if no word in it is a prefix of another. By abuse of language, we will continue to refer to the μ -

probability of the union of the cylinders representing the words of V beginning in the zero coordinate as the μ -probability of the collection and even dare to denote it by $\mu(V)$, i.e. $\mu(V) := \sum_{\alpha \in V} \mu([\alpha])$ (the prefix-free property means exactly that these cylinders are disjoint).

Fixing a coordinate to start from we parse uniquely any point in $\Lambda^{\mathbb{Z}}$ starting from that coordinate until we meet an element in V and continue in the same manner after it indefinetly in the positive direction (only that direction is parsed, and the parsing is with gaps). Notice that it is not always possible to extend the parsing into the negative direction, for a word in V overlapping the starting position may be met before reaching a last letter of another word in V.

If a parsing starts at a certain coordinate and another starts before it, we shall identify the two if the latter agrees with the former in the former's domain of definition. Under this identification, the number of parsings of any $x \in \Lambda^{\mathbb{Z}}$ can still very well be greater than 1 because the parsing depends on the coordinate we choose to start from, however, it is not greater than $2|V|\max\{|\alpha|\}_{\alpha\in V}$ (this bound is not optimal in general and even less so to most specific collections V). To see this, consider a word appearing in some fixed parsing, its length is of course not greater than $\max\{|\alpha|\}_{\alpha\in V}$, and other parsings with a starting position prior to this word need to have a word that overlaps with it. Hence, together with this fixed one, there surely are at most $2|V|\max\{|\alpha|\}_{\alpha\in V}$ such parsings (choose from the collection containing this word and all those words that overlap it a word of maximal length and count all words in V as either ending or beginig in one of its letters). Now, If there were more than $2|V|\max\{|\alpha|\}_{\alpha\in V}$ parsings, then there is a parsing and a specific word that appears in some place of it such that at least $2|V|\max\{|\alpha|\}_{\alpha\in V}$ other parsings have a starting position prior to it, and this is a contradicion.

Up to now, we ignored the measure μ . Putting it into play, it is natural to ask - in the spirit of the pointwise ergodic theorem - whether the asymptotic frequency of appearances of every word in V appearing in such a parsing is generically the same as the μ -probability of the cylinders they represent. The answer to this question is negative, as shown by the following example.

Example: $\Lambda = \{A, B, C\}$ and we define a Markov chain with transition probabilities

$$p_{AB}=1,\,p_{AC}=0,\,p_{BA}=0,\,p_{BC}=1,\,p_{CA}=rac{1}{2},\,p_{CB}=rac{1}{2}.$$

Its unique stationary distribution is $p_A = \frac{1}{5}$, $p_B = \frac{2}{5}$, $p_C = \frac{2}{5}$, and thus an ergodic stationary process is defined. The collection $\{ABC, BC, C\}$ is a prefix-free collection (of total probability 1), and for generic points the parsing according to it is unique (under the above identification), with no gaps and the word C appears with asymmtotic frequency 0.

The above example shows that these asymptotic frequencies need not equal the respective μ -probabilities. However, one may ask if even μ -almost-surely parsings of a point in $\Lambda^{\mathbb{Z}}$ posses asymptotic word frequencies.

We denote by \tilde{V} the collection obtained by adding to V all elements of Λ (treated as single letter words). We consider the coding map $\varphi: \tilde{V}^{\mathbb{Z}} \to \Lambda^{\mathbb{Z}}$ that codes consecutively each element of \tilde{V} by just the word it is in the alphabet Λ where the word appearing in the zero coordinate of a point in $\tilde{V}^{\mathbb{Z}}$ is coded such that it begins in the zero coordinate of the point in $\Lambda^{\mathbb{Z}}$. φ does not commute with the shift maps. To fix this, we create a new space $X' = \{(z, n): z \in \tilde{V}^{\mathbb{Z}}, n \in \mathbb{Z}, 0 \leq n < |z_1|\}$ (where $|z_1|$ is the length of the word z) and define on it:

$$\tilde{\sigma}((z,n)) = \begin{cases} (z, n+1) & n \neq n-1 \\ (\sigma(z), 0) & n = n-1 \end{cases}$$

(in more fancy words in the theory of dynamical systems, the system thus obtained is the discrete suspension of $(\tilde{V}^{\mathbb{Z}}, \sigma)$ formed by the ceiling function $z \mapsto |z_1|$). And also define a map $\tilde{\varphi}: X' \to \Lambda^{\mathbb{Z}}$ by $\tilde{\varphi}((z,n)) = \sigma^n(\varphi(z))$. Now $\tilde{\varphi}$ is an equivariant map from the system $(X', \tilde{\sigma})$ onto the system $(\Lambda^{\mathbb{Z}}, \sigma)$. We equip X' with the topology inherited from $\tilde{V}^{\mathbb{Z}} \times \mathbb{Z}_{\geq 0}$, and with this topology it is compact and $\tilde{\varphi}$ is a continuous factor map. However we want to restrict out attention to its closed invariant sub-set X of which consecutive elements in $\tilde{V} \setminus V$ that form together a word in V are forbidden - for that cannot happen in the parsing process. So X is the space of all the possible parsings according to V, and $\tilde{\varphi}$ restricted to X is already onto.

Thus there exists a probability measure η on X which is ergodic relative to $\tilde{\sigma}$ for which $\tilde{\varphi}$ is a factor map from $(X, \tilde{\sigma}, \eta)$ to $(\Lambda^{\mathbb{Z}}, \sigma, \mu)$ as measure-preserving systems (η need not be a unique such measure).

Open Question: If V was infinite then the space X need not be compact. Is such an η still guaranteed to exist?

For every $\alpha \in V$, let us denote by $[(\alpha,0)]$ the set of all points $(z,n) \in X$ with $z=\alpha$ and n=0. Now the pointwise ergodic theorem tells us that η -almost-surely the sum $\frac{1}{N} \sum_{i=1}^{N} \chi_{[(\alpha,0)]} \left(\tilde{\sigma} \left((z,n) \right) \right)$ (where $\chi_{[(\alpha,0)]}$ is the indicator function of the set $[(\alpha,0)]$) converges to $\eta \left([(\alpha,0)] \right)$. Notice that $[(\alpha,0)]$ is contained in the inverse image under $\tilde{\varphi}$ of $[\alpha]$ and thus $\eta \left([(\alpha,0)] \right) \leq \mu \left([\alpha] \right)$.

Proof of Theorem 1: η_{α} corresponds to $\eta([(\alpha,0)])$ in the preceding discussion.

Open Question: Do μ -almost-surely all parsings of $x \in \Lambda^{\mathbb{Z}}$ admit such limits?

Continuing in the notation of Theorem 1, notice that

$$\sum_{\alpha \in V} |\alpha| \, \eta_{\alpha} = 1 - \sum_{\alpha \in \tilde{V} \setminus V} \eta_{\alpha} \ge 1 - (1 - \mu(V)) = \mu(V) \,,$$

where the inequality is based on the observation in the introduction.

Theorem 1 is all we need for the proof of the Shannon-McMillan-Breiman theorem presented in the next section. However, this preliminary treatment of the parsings space X will not be complete without addressing its entropy and writing down the elementary bounds on p_{α} . The latter can be thus bounded from above:

$$p_{\alpha} = \frac{\eta_{\alpha}}{\sum\limits_{\beta \in \tilde{V}} \eta_{\beta}} = \frac{\max\left\{|\beta|\right\}_{\beta \in V} \eta_{\alpha}}{\max\left\{|\beta|\right\}_{\beta \in V} \sum\limits_{\beta \in \tilde{V}} \eta_{\beta}} \leq \frac{\max\left\{|\beta|\right\}_{\beta \in V} \eta_{\alpha}}{\sum\limits_{\beta \in \tilde{V}} |\beta| \eta_{\beta}} = \max\left\{|\beta|\right\}_{\beta \in V} \eta_{\alpha}.$$

In the other direction one similarly obtains $p_{\alpha} = \frac{\eta_{\alpha}}{\sum\limits_{\beta \in V} \eta_{\beta}} \geq \min\{|\beta|\}_{\beta \in V} \eta_{\alpha}$. All words

in V being of the same length and $\mu(V) = 1$ is a sufficient condition to guarantee equality in these two inequalities. However, they are not always helpful.

Example: We return to the previous example. The probability of transitioning in a parsing from ABC to BC is $\frac{1}{2}$ and so is the transition in the opposite direction. This is again a markov process with stationary probability $\frac{1}{2}$ to each state of these two. The upper bound on the frequency above gives $\frac{3}{5}$ to $\alpha = ABC$, $\frac{6}{5}$ to $\alpha = BC$ and $\frac{6}{5}$ to $\alpha = C$ (although that frequency is zero!). $\frac{6}{5} \geq 1$ so it is not of any help.

Regarding the entropy of the system $(X, \tilde{\sigma}, \eta)$, let us denote by ξ the alphabet partition of $\Lambda^{\mathbb{Z}}$ relative to the zero coordinate and the partition of X relative to its zero coordinate by ζ . The important fact is that for every $x \in \Lambda^{\mathbb{Z}}$ and every $n \in \mathbb{N}$ the inverse image through $\tilde{\varphi}$ of the atom $\xi_0^n(x)$ is a union of not more than $2|V|\max\{|\alpha|\}_{\alpha\in V}$ atoms of the partition ζ_0^n (where, as usual, the subscipt and superscript indexes refer to refinements of the partitions by the maps of the systems). In particular this impies that the entropy of $(X, \tilde{\sigma}, \eta)$ is equal to that of $(\Lambda^{\mathbb{Z}}, \sigma, \mu)$.

Remark: In an information theoretic point of view, parsing is often applied to encodings of stationary processes, and those encoded processes need not be themselves stationary, so our analysis does not suit this case. Furthermore, it seems that there is no natural probabilistic scenario that constructs a stationary probability measure on the symbolic space of the the encoded process out of the non-stationary measure it possesses. However, putting encoded processes aside, a choice of a prefix-free collection of words in the alphabet of a stationary process is equivalent to defining a stopping time on it, and as we see in this paper, questions about parsing of the process according to such a collection may become relevant.

4 The Shannon-McMillan-Breiman Theorem

For the present section and the next one, we continue with the ergodic system $(\Lambda^{\mathbb{Z}}, \sigma, \mu)$ (Λ being a finite alphabet) and denote its entropy by h.

Theorem 2' (Shannon-McMillan): For every $\varepsilon > 0$ there exists some N_{ε} such that for every $n > N_{\varepsilon}$ the μ -probability of $\{x : e^{-(h+\varepsilon)n} < \mu([x_1^n]) < e^{-(h-\varepsilon)n}\}$ is greater than $1 - \varepsilon$.

This is how the Shannon-McMillan theorem is probably most usually formulated. However, our point of departure will be the following, also very well known, equivalent formulation.

Theorem 2 (Shannon-McMillan): For every $\varepsilon > 0$ there exists some N_{ε} such that for every $n > N_{\varepsilon}$ the following is satisfied:

- (i) There exists a collection of less than $e^{(h+\varepsilon)n}$ length n words that has a μ -probability greater than $1-\varepsilon$.
- (ii) Every collection of less than $e^{(h-\varepsilon)n}$ length n words has a μ -probability smaller than ε .

While Theorem 1 deals with statistics in space, D. Ornstein and B. Weiss proved an analogous version for statistics in time (the latter is stronger than the former) - cf. [4] or chapter 9 in [6].

Theorem 3 (Ornstein-Weiss): For $x \in \Lambda^{\mathbb{Z}}$ μ -almost-surely, given an $\varepsilon > 0$ there exists some $N_{x,\varepsilon} \in \mathbb{N}$ such that for every $N \geq N_{x,\varepsilon}$ the following is satisfied:

- (i) For every $M \in \mathbb{N}$, there exists a collection C_N of length N words of cardinality smaller than $e^{(h+\varepsilon)N}$ such that its elements cover a fraction greater than $1-\varepsilon$ of x_1^M .
- (ii) For every $M \geq e^{hN}$, the elements of any collection of length N words of cardinality smaller than $e^{(h-\varepsilon)N}$ cover less than an ε fraction of x_1^M .

Theorem 2 is not applied in the sequel, but is stated here as its first - and easier to prove - part was the inspiration for Lemma 5 in the proof of the Shannon-McMillan-Breiman theorem (cf. [2, 3]) presented in this paper.

Theorem 4 (Shannon-McMillan-Breiman): $\lim_{n\to\infty} \frac{-\log \mu([x_1^n])}{n} = h$ for $x \in \Lambda^{\mathbb{Z}}$ μ -almost-surely.

Remark: For an infinite alphabet and an ergodic stationary measure of finite entropy the same claim follows easily from the above result (formulated for finite Λ) by refining finite alphabet partitions. Alternatively, one can carry out our proof directly for an infinite alphabet without much additional sophistication involved (Section 3 need not be generalized for an infinite alphabet for that purpose).

5 A Proof of the Shannon-McMillan-Breiman Theorem

Before proving Theorem 4 we need to prove a corollary of Theorem 2 (its proof is trivial but the statement is inspired by theorem 3 (i)).

Lemma 6: For every $\varepsilon > 0$ there exists an $N_{\varepsilon} \in \mathbb{N}$ and collections C_N for all $N \geq N_{\varepsilon}$, each of cardinality smaller than $e^{(h+\varepsilon)N}$, such that for every $x \in \Lambda^{\mathbb{Z}}$ which is μ -generic relative to σ , the following is satisfied:

for every $N \geq N_{\varepsilon}$ the elements of the collection C_N can cover a fraction greater than $1 - \varepsilon$ of x_1^M with no overlaps for a sufficiently large M.

Proof: We claim that the N_{ε} and the collections C_N of Theorem 2 do the job. Given an $N \geq N_{\varepsilon}$, by the pointwise ergodic theorem for any μ -generic $x \in \Lambda^{\mathbb{Z}}$ the elements of C_N begin in a fraction greater than $1 - \varepsilon$ of the coordinates of x_1^M for a sufficiently large M (where the coordinates $M - N + 2, \ldots, M$ are counted as coordinates in which these elements do not begin). Parse x_1^M starting from the first coordinate (in the sense of the preceding section), and apply the observation from the introduction.

We will also need this elementary lemma.

Lemma 7: Given an
$$x \in \Lambda^{\mathbb{Z}}$$
 and $s \in \mathbb{N}$: $\liminf_{n \to \infty} \left(-\frac{\log \mu([x_1^n])}{n} \right) = \frac{1}{s} \liminf_{n \to \infty} \left(-\frac{\log \mu([x_1^{ns}])}{n} \right)$.

Proof: Surely " \leq " holds. To see the other direction notice that, given $0 \leq r < s$ we

have $\mu\left(\left[x_1^{ns+r}\right]\right) \leq \mu\left(\left[x_1^{ns}\right]\right)$, and so $-\log\mu\left(\left[x_1^{ns+r}\right]\right) \geq -\log\mu\left(\left[x_1^{ns}\right]\right)$. Thus

$$\frac{ns+r}{ns} \cdot \frac{-\log \mu\left(\left[x_1^{ns+r}\right]\right)}{ns+r} \geq \frac{-\log \mu\left(\left[x_1^{ns}\right]\right)}{ns}. \blacksquare$$

Proof of Theorem 4: To simplify matters, let us first notice that there are constants $0 \le c_2 \le c_1$ for which μ -almost-surely the \liminf and \limsup of the \liminf in question equal respectively. This follows from ergodicity of the invariant measure μ and the fact that

$$-\frac{\log\mu\left(\left[\left(\sigma x\right)_{1}^{n}\right]\right)}{n} \leq -\frac{n+1}{n} \cdot \frac{\log\mu\left(\left[x_{1}^{n+1}\right]\right)}{n+1}.$$

Assume to the contrary that $c_1 > h$. Given a large number $a \in \mathbb{N}$ that we will later choose, for μ -almost-every $x \in \Lambda^{\mathbb{Z}}$ we define a stopping time T_x to be the minimal T > a for which $\frac{-\log \mu\left(\left[x_1^T\right]\right)}{T} > \frac{h+c_1}{2}$ (for the zero measure set of points with no such a T it remains undefined). Denote by V_k the collection of positive μ -probability words of the form $x_1^{T_x}$ with $T_x = k$. The union $\bigcup_{k=a+1}^{\infty} V_k$ is a prefix-free collection. For a 0 < q < 1 very close to 1 that we will later choose, we restrict our attention to the finite sub-collection $V := \bigcup_{k=a+1}^{\tilde{T}} V_k$, where \tilde{T} is the minimal integer T > a for which $\mu\left(\bigcup_{k=a+1}^{\tilde{T}} V_k\right) = \sum_{k=a+1}^{T} \mu\left(\left\{x \in \Lambda^{\mathbb{Z}} : T_x = k\right\}\right) > q$.

We fix some $y \in \Lambda^{\mathbb{Z}}$ for which both Lemma 6 and Theorem 1 (with V) apply. For a small $\varepsilon > 0$ to be chosen later, we obtain N_{ε} (following the notation of Lemma 6) and fix some large $N \geq N_{\varepsilon}$ whose value we will later choose. So there exists a collection C_N of cardinality smaller than $e^{(h+\varepsilon)N}$ whose elements cover with no overlaps a fraction greater than $1 - \varepsilon$ of of y_1^M for any sufficiently large M.

By Theorem 1, applied with our V, there exists a parsing of y according to V with \tilde{V} , η_{α} and p_{α} as in the theorem. For a small $\delta > 0$ to be also chosen later, we choose an $M \in \mathbb{N}$ so large that the requirement involving C_N applies to it and also that y_1^M

contains any $\alpha \in \tilde{V}$ with a frequency greater than $(1 - \delta) p_{\alpha}$ and smaller than $(1 + \delta) p_{\alpha}$ in that parsing. The idea now is to show that the presence of these words contradicts the existence of C_N .

By the observation from the introduction

$$\sum_{\alpha \in V} |\alpha| \, \eta_{\alpha} \ge \mu(V) > q,$$

implies that for $L \in \mathbb{N}$ to be chosen later

$$\sum_{k \in \left\{k : a < k \le \tilde{T}, \sum_{\alpha \in V_k} k \eta_\alpha > \frac{q}{L} \frac{6}{\pi^2} \frac{1}{k^2}\right\}} \sum_{\alpha \in V_k} k \eta_\alpha > q - \frac{q}{L}.$$

So, if δ is chosen to be small enough, $k \in \left\{k : a < k \leq \tilde{T}, \sum_{\alpha \in V_k} k \eta_\alpha > \frac{q}{L} \frac{6}{\sigma^2} \frac{1}{k^2}\right\}$ parsing a fraction greater than $q - \frac{q}{L}$ of y_1^M . Thus a covering with no overlaps by elements of C_N that covers a fraction greater than $1 - \varepsilon$ of y_1^M must cover a fraction greater than $q - \frac{q}{L} - \varepsilon$ of the part of y_1^M composed in the parsing of words in the collections V_k for $k \in \left\{k : a < k \leq \tilde{T}, \sum_{\alpha \in V_k} k \eta_\alpha > \frac{q}{L} \frac{6}{\sigma^2} \frac{1}{k^2}\right\}$. This means that there exists some $k_0 \in \left\{k : a < k \leq \tilde{T}, \sum_{\alpha \in V_k} k \eta_\alpha > \frac{q}{L} \frac{6}{\sigma^2} \frac{1}{k^2}\right\}$ so that C_N can cover with its elements with no overlaps a fraction greater than $q - \frac{q}{L} - \varepsilon$ of the part of y_1^M composed of words in V_{k_0} in the parsing. And for a small r > 0 to be chosen later we denote by V'_{k_0} the sub-collection of V_{k_0} that is composed of words of which the above covering covers a fraction greater than $(1-r)\left(q - \frac{q}{L} - \varepsilon\right)$ of the total appearances of each of them in the parsing of y_1^M . So for a $\alpha \in V_{k_0} \setminus V'_{k_0}$, a fraction of at most $(1-r)\left(q - \frac{q}{L} - \varepsilon\right)$ of its appearances in the parsing of y_1^M is coverd. The calculation

$$(1-r)\left(q-\frac{q}{L}-\varepsilon\right)\left(1-\frac{\sum\limits_{\alpha\in V_{k_0}'}p_\alpha}{\sum\limits_{\alpha\in V_{k_0}}p_\alpha}\right)+\frac{\sum\limits_{\alpha\in V_{k_0}'}p_\alpha}{\sum\limits_{\alpha\in V_{k_0}}p_\alpha}>q-\frac{q}{L}-\varepsilon$$

$$\left(1-(1-r)\left(q-\frac{q}{L}-\varepsilon\right)\right)\frac{\sum\limits_{\alpha\in V_{k_0}}p_\alpha}{\sum\limits_{\alpha\in V_{k_0}}p_\alpha} > r\left(q-\frac{q}{L}-\varepsilon\right)$$

$$\frac{\sum\limits_{\alpha\in V_{k_0}}p_\alpha}{\sum\limits_{\alpha\in V_{k_0}}p_\alpha} > \frac{r\left(q-\frac{q}{L}-\varepsilon\right)}{1-(1-r)\left(q-\frac{q}{L}-\varepsilon\right)}$$
 shows that
$$\frac{\sum\limits_{\alpha\in V_{k_0}}p_\alpha}{\sum\limits_{\alpha\in V_{k_0}}p_\alpha} > \frac{r\left(q-\frac{q}{L}-\varepsilon\right)}{1-(1-r)\left(q-\frac{q}{L}-\varepsilon\right)} = \frac{r\left(q-\frac{q}{L}-\varepsilon\right)}{1-\left(q-\frac{q}{L}-\varepsilon\right)+r\left(q-\frac{q}{L}-\varepsilon\right)}.$$

Now if $\alpha \in V_{k_0}$ then $p_{\alpha} = \frac{\eta_{\alpha}}{\sum\limits_{\beta \in \tilde{V}} \eta_{\beta}} \leq \frac{\mu([\alpha])}{\sum\limits_{\beta \in \tilde{V}} \eta_{\beta}} < \frac{e^{-\frac{(h+c_1)}{2}k_0}}{\sum\limits_{\beta \in \tilde{V}} \eta_{\beta}}$ (the first inequality was explained in the previous section and the second follows from the definition of the stopping time). This means that

$$\begin{split} \left|V_{k_0}'\right| &\geq \left\lceil\frac{\sum\limits_{\alpha \in V_{k_0}'} p_{\alpha}}{\max\left\{p_{\alpha}\right\}_{\alpha \in V_{k_0}'}}\right\rceil > \left\lceil\sum\limits_{\beta \in \tilde{V}} \eta_{\beta} \cdot e^{\frac{(h+c_1)}{2}k_0} \sum\limits_{\alpha \in V_{k_0}'} p_{\alpha}\right\rceil = \left\lceil\sum\limits_{\beta \in \tilde{V}} \eta_{\beta} \cdot e^{\frac{(h+c_1)}{2}k_0} \frac{\sum\limits_{\alpha \in V_{k_0}} p_{\alpha}}{\sum\limits_{\alpha \in V_{k_0}} p_{\alpha}} \sum\limits_{\alpha \in V_{k_0}} p_{\alpha}\right\rceil \\ &= \left|\sum\limits_{\beta \in \tilde{V}} \eta_{\beta} \cdot e^{\frac{(h+c_1)}{2}k_0} \frac{r\left(q - \frac{q}{L} - \varepsilon\right)}{1 - \left(q - \frac{q}{L} - \varepsilon\right) + r\left(q - \frac{q}{L} - \varepsilon\right)} \sum\limits_{\alpha \in V_{k_0}} \sum\limits_{\beta \in \tilde{V}} \eta_{\beta}\right| \\ &> \left[e^{\frac{(h+c_1)}{2}k_0} \frac{r\left(q - \frac{q}{L} - \varepsilon\right)}{1 - \left(q - \frac{q}{L} - \varepsilon\right) + r\left(q - \frac{q}{L} - \varepsilon\right)} \frac{q}{L} \frac{6}{\pi^2} \frac{1}{k_0^3}\right]. \end{split}$$

We know that a fraction greater than $(1-r)\left(q-\frac{q}{L}-\varepsilon\right)-2\frac{N}{k_0}$ of each of the words in V'_{k_0} can be covered by no overplapping words in C_N (the subtrahend is needed since when now referring to covering of individual words by elements in C_N we do not permit only parts of such elements being used in the two boundaries of the word). But this means we can bound from above the number of words in V'_{k_0} :

$$\begin{aligned} |V'_{k_0}| &\leq |\Lambda|^{\left(1 - (1 - r)\left(q - \frac{q}{L} - \varepsilon\right) + 2\frac{N}{k_0}\right)k_0} \sum_{i=1}^{\left\lfloor \frac{k_0}{N} \right\rfloor} \binom{k_0}{i} e^{(h + \varepsilon)Ni} \\ &\leq |\Lambda|^{\left(1 - (1 - r)\left(q - \frac{q}{L} - \varepsilon\right) + 2\frac{N}{k_0}\right)k_0} e^{(h + \varepsilon)k_0} e^{\left(-\frac{1}{N}\log\frac{1}{N} - \left(1 - \frac{1}{N}\right)\log\left(1 - \frac{1}{N}\right)\right)k_0} \end{aligned}$$

Notice that k_0 depends on our choices of a, q, L, M, N. In particular $k_0 > a$ and thus

for every choice of q, L, M, N we can choose a to make k_0 as large as we wish. Thus it is clear that we can choose a, L, M, N large, q close to 1 and ε, r close to 0 to make this upper bound on $\left|V_{k_0}'\right|$ smaller than the lower bound $\left[e^{\frac{(h+c_1)}{2}k_0}\frac{r\left(q-\frac{q}{L}-\varepsilon\right)}{1-\left(q-\frac{q}{L}-\varepsilon\right)+r\left(q-\frac{q}{L}-\varepsilon\right)}\frac{q}{L}\frac{6}{\pi^2}\frac{1}{k_0^3}\right]$ and arrive to a contradiction. Thus $c_1 \leq h$. //

If h=0 the proof is already done. So we may deal in the sequel only with the case h>0.

Now let us assume to the contrary that $c_2 < h$.

Given a small enough $\varepsilon > 0$, we are going to contradict the second part of Theorem 2. In fact, we will show that for every N large enough there is a collection of cardinality smaller than $e^{(h-\varepsilon)N}$ of probability greater than $1-\varepsilon$.

We are entitled to suppose without loss of generality that h, the entropy of (X, μ, σ) , satisfies the inequality $\frac{h+c_2}{2} + 2 < h - \varepsilon$ (which will turn out to be useful). To see this we notice that, by lemma 7, for every $s \in \mathbb{N}$, the system of σ^s perceived as acting on $(\Lambda^s)^{\mathbb{Z}}$ has an in its ergodic decomposition a component for which the difference between its entropy and liminf in question not smaller than $s(h-c_2)$. And, for a large enough s, we take the system of this ergodic component as our new system (X, μ, σ) (so our new Λ will be the old Λ^s etc).

For a large enough a, we define (almost-surely) a stopping time S_x as the minimal S > a satisfying

$$\frac{-\log\mu\left(\left[x_1^S\right]\right)}{S} < \frac{h+c_2}{2}. \text{ We denote by } \tilde{S} \text{ the minimal } S \text{ for which }$$

$$\sum_{k=a+1}^S \mu\left(\left\{x\in\Lambda^\mathbb{Z}\,:\, S_x=k\right\}\right) \,>\, 1\,-\,\varepsilon, \text{ and for every } a\,<\,k\,\leq\,\tilde{S} \text{ by } W_k \text{ the collection of length } k \text{ words that stop this stopping time.} \quad \text{Each } |W_k| \text{ is smaller than }$$

$$\mu\left(\left\{x\in\Lambda^\mathbb{Z}\,:\, S_x=k\right\}\right)e^{\frac{h+c_2}{2}k}.$$

For a small $\delta > 0$, by Theorem 1, applied with the collection $W = \bigcup_{k=a+1}^{\tilde{S}} W_k$, there exists for every $\alpha \in \tilde{W}$ a $p_{\alpha} \geq 0$ and some $N\left(\varepsilon,a\right)$ such that for any $N > N\left(\varepsilon,a\right)$ there is

a probability greater than $1-\varepsilon$ that for $x\in\Lambda^{\mathbb{Z}}$ there exists a a parsing of x_1^N according to \tilde{W} such that each $\alpha\in\tilde{W}$ appears in it in a frequency between $(1-\delta)\,p_\alpha$ and $(1+\delta)\,p_\alpha$. Hence, by the observation from the introduction, a parsing of x_1^N according to W covers a fraction greater than $1-\varepsilon$ of x_1^N (since $\mu(W)>1-\varepsilon$) and in such a covering the average length of appearances of the words belonging to these collections in x_1^N is greater than $\sum_{k=a+1}^{\tilde{S}} k^{\frac{1-\delta}{1+\delta}} \frac{\sum_{\alpha\in W_k}^{p_\alpha}}{\sum_{\alpha\in W}^{p_\alpha}}.$ Hence there are at most $\frac{N}{\sum_{k=a+1}^{\tilde{S}} k^{\frac{1-\delta}{1+\delta}} \frac{\sum_{\alpha\in W_k}^{p_\alpha}}{\sum_{\alpha\in W}^{p_\alpha}}}$ appearances of words in

W in this parsing. Let us denote the collection of all x_1^N of that sort by U. Then |U| can be estimated from above by

$$\begin{pmatrix} N \\ \lfloor \frac{N}{a} \rfloor \end{pmatrix} e^{\frac{N}{\sum\limits_{k=a+1}^{\tilde{S}} k \frac{1-\delta}{1+\delta} \frac{\alpha \in W_k}{\alpha \in W}^{p_{\alpha}}}} \begin{pmatrix} \sum\limits_{k=a+1}^{\tilde{S}} -\frac{\sum\limits_{\alpha \in W_k}^{p_{\alpha}} p_{\alpha}}{\sum\limits_{\alpha \in W}^{p_{\alpha}}} \log \begin{pmatrix} \sum\limits_{\alpha \in W_k}^{p_{\alpha}} p_{\alpha} \\ \sum\limits_{\alpha \in W}^{p_{\alpha}} p_{\alpha} \end{pmatrix} + \delta \end{pmatrix}$$

$$\cdot \left(\prod_{k=a+1}^{\tilde{S}} |W_k|^{\frac{1-\delta}{\alpha \in W_k} p_\alpha \frac{N}{\sum\limits_{j=a+1}^{\tilde{S}} j \frac{1-\delta}{1+\delta} \frac{\alpha \in W_j}{\sum\limits_{\alpha \in W}^{p_\alpha}}}} \right) \left(N \atop \lfloor \varepsilon N \rfloor \right) |\Lambda|^{\varepsilon N}$$

Also,

$$\prod_{k=a+1}^{\tilde{S}} |W_k| \sum_{\alpha \in W_k} p_\alpha \frac{N}{\sum\limits_{j=a+1}^{\tilde{S}} j \frac{1-\delta}{1+\delta} \frac{\alpha \in W_j}{\alpha \in W}^{p_\alpha}}} \leq \prod_{k=a+1}^{\frac{(h+c_2)k}{2}} (1+\delta)^{\frac{k}{\alpha} \frac{\sum\limits_{\alpha \in W_k} p_\alpha}{k}} \frac{N}{\sum\limits_{j=a+1}^{\tilde{S}} j \frac{1-\delta}{1+\delta} \frac{\alpha \in W_j}{\alpha \in W}^{p_\alpha}}}$$

$$=\prod_{k=a+1}^{\tilde{S}} e^{\frac{\frac{h+c_2}{2}\frac{(1+\delta)^2}{1-\delta}\frac{\sum\limits_{\alpha\in W_k}^{k}p_{\alpha}}{\sum\limits_{j=a+1}^{\tilde{S}}\sum\limits_{\alpha\in W_k}^{j}p_{\alpha}}\sum\limits_{\alpha\in W}^{k}p_{\alpha}N} = e^{\frac{h+c_2}{2}\frac{(1+\delta)^2}{1-\delta}\sum\limits_{\alpha\in W}^{k}p_{\alpha}N} \leq e^{\frac{h+c_2}{2}\frac{(1+\delta)^2}{1-\delta}N}.$$

Thus

$$|U| < e^{H\frac{1}{a}} e^{\sum_{k=a+1}^{\tilde{\Sigma}} -\frac{\sum_{\alpha \in W_k}^{p_{\alpha}} \log \left(\frac{\sum_{\alpha \in W_k}^{p_{\alpha}}}{\sum_{\alpha \in W}^{p_{\alpha}}}\right) + \delta}{\sum_{k=a+1}^{\tilde{\Sigma}} k\frac{1-\delta}{1+\delta} \sum_{\alpha \in W_k}^{p_{\alpha}} \sum_{\alpha \in W}^{p_{\alpha}} N} e^{\frac{h+c_2}{2} \frac{(1+\delta)^2}{1-\delta} N} e^{H_{\varepsilon} N} e^{\log|\Lambda| \varepsilon N}$$

where $H_p := -p \log p - (1-p) \log (1-p)$. And for a, N large enough and ε, δ small

enough, this is smaller than

$$e^{\frac{\sum\limits_{k=a+1}^{\tilde{S}}-\frac{\sum\limits_{\alpha\in W_k}^{p_{\alpha}}\log\left(\sum\limits_{\alpha\in W_k}^{p_{\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}k\frac{\sum\limits_{\alpha\in W_k}^{p_{\alpha}}}{\sum\limits_{\alpha\in W_k}^{p_{\alpha}}}}e^{\frac{\sum\limits_{\alpha\in W_k}^{p_{\alpha}}}{\sum\limits_{\alpha\in W_k}^{p_{\alpha}}}}e^{\frac{h+c_2}{2}N}.$$

We are left to deal with $\frac{\sum\limits_{k=a+1}^{\tilde{S}}-\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}\frac{k}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}}. \text{ Luckily, } \sum\limits_{k=a+1}^{\tilde{S}}k\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\text{ can be seen}$ as an expectation of the length of a coding for an alphabet indexed by $k\in\left\{a+1,a+2,\ldots,\tilde{S}\right\}$ with probabilites $\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\text{ such that the k-th symbol has a code of length k (there are many such codes). But then we know that <math display="block">\sum\limits_{k=a+1}^{\tilde{S}}k\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\text{ is not smaller than the entropy of that distribution which is just }\sum\limits_{k=a+1}^{\tilde{S}}-\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\log\left(\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\right)$ Hence $\frac{\sum\limits_{k=a+1}^{\tilde{S}}-\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W}^{p_\alpha}}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}k\frac{\sum\limits_{\alpha\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W}^{p_\alpha}}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{\alpha\in W_k}^{\tilde{S}}k^{\frac{\tilde{S}}{\alpha}}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}k\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}k\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}}\log\left(\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\sum\limits_{\alpha\in W_k}^{p_\alpha}}\right)}{\sum\limits_{k=a+1}^{\tilde{S}}\frac{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}{\alpha\overset{\sum}{\in W_k}^{p_\alpha}}}}}$

Where the last inequality follows from the assumption $\frac{h+c_2}{2}+2 < h-\varepsilon$. This contradicts the second part of Theorem 2'.

Remark: The only pointwise result we have essentially applied was the use of the pointwise ergodic theorem in the proof of Lemma 6 (A weaker version of Theorem 1 without the pointwise ergodic theorem but satisfactory for our needs can easily be proved). In fact, instead of Lemma 6 we could similarly prove with only the mean ergodic theorem the following result and it would have been sufficient for our needs (thus gaining a pointwise theorem - Theorem 4 - without using any previous pointwise result).

Theorem: Given an $\varepsilon > 0$ there exists some $N_{\varepsilon} \in \mathbb{N}$ such that for every $N \geq N_{\varepsilon}$ there

exists a collection C_N of length N words of cardinality smaller than $e^{(h+\varepsilon)N}$ such that with μ -probability greater than $1-\varepsilon$ a point $x\in\Lambda^{\mathbb{Z}}$ satisfies that for every large enough M the collection C_N can cover a fraction greater than $1-\varepsilon$ of x_1^M .

Thus this proof of the Shannon-McMillan-Breiman Theorem need not use any previous pointwise result.

References

- J. Bourgain, H. Furstenberg, Y. Katznelson, D.S. Ornstein (1989). Return Times of Dynamical Systems. Publications Mathématiques de l'Institut des Hautes Études Scientifiques, 69, 42-45.
- [2] L. Breiman (1957). The Individual Ergodic Theorem of Information Theory. The Annals of Mathematical Statistics, 28(3), 809-811.
- [3] L. Breiman (1960). Correction Notes: Correction to "The Individual Ergodic Theorem of Information Theory". The Annals of Mathematical Statistics, 31(3), 809-810.
- [4] D. S. Ornstein, B. Weiss (1990). How Sampling Reveals a Process. The Annals of Probability, 18(3), 905-930.
- [5] D. S. Ornstein, B. Weiss (1993). Entropy and Data Compression Schemes. IEEE Transactions on information theory, 39(1), 78-83.
- [6] B. Weiss (2000). Single Orbit Dynamics. CBMS Regional Conference Series in Mathematics, 95. American Mathematical Society, Providence, RI.

Einstein Institute of Mathematics, Edmond J. Safra campus, The Hebrew University of Jerusalem, Israel.

matan.tal@mail.huji.ac.il