ON THE CATEGORIFIED WRAPPING NUMBER CONJECTURE

BENJAMIN DANIELS AND MELISSA ZHANG

ABSTRACT. We prove the Categorified Wrapping Number Conjecture for large classes of annular links, including alternating annular links and tangle closures exhibiting plumbed link phenomena. We do so by characterizing when a resolution is sufficient to produce a nonzero homology class in k-grading wrap(L) on its own. This characterization primarily concerns the type of crossing resolutions abutting trivial circles.

CONTENTS

| 1. Introduction | 1 |
|--|----|
| 1.1. Acknowledgements | 4 |
| 2. Annular Khovanov homology | 4 |
| 2.1. Cube of resolutions | 4 |
| 2.2. The annular Khovanov chain complex | 5 |
| 2.3. Wrapping number of diagrams and resolutions | 8 |
| 3. Perfectly wrapped and uniform resolutions | 8 |
| 3.1. Terminology and notation | 8 |
| 3.2. Type 0 and type 1 trivial circles | 10 |
| 3.3. Proof of the main result | 11 |
| 3.4. An expanded class of diagrams admitting perfectly wrapped uniform resolutions | 13 |
| 4. Classes of links admitting perfectly wrapped uniform resolutions | 16 |
| 4.1. Alternating links | 17 |
| 4.2. Other classes and modifications of links | 17 |
| References | 20 |

1. INTRODUCTION

In 2000, Khovanov categorified the Jones polynomial link invariant [Kho00], sparking a revolution in the development of link homologies. In the following years, his theory has been adapted and generalized in numerous ways (see [BN05], [Lee05], and [Ras10] for some highly impactful early constructions).

One such generalization is Asaeda–Przytycki–Sikora's extension of Khovanov homology to links in Ibundles over surfaces [APS04], where I is the unit interval. This homology theory is sometimes referred to as APS homology in reference to the authors' names. The most popular version of APS homology is annular Khovanov homology, which provides an invariant for annular links, i.e. links in the thickened annulus.

Let $A = \mathbb{R}^2 - \{0\}$, and let $L \subset A \times I$ be an annular link. The annular Khovanov homology of L, denoted by AKh(L), is a triply-graded homology-type invariant of L computed from an annular diagram for L, i.e. a link diagram that misses the origin, which we will denote by an asterisk (*) in the plane. In addition to the homological and quantum gradings inherited from Khovanov homology, AKh has an additional 'k-grading' determined by the location of the *: this endows the usual Khovanov complex with a k-filtration grading, and AKh is the homology of the associated graded object.

Annular Khovanov homology has attracted considerable interest over the past decades. For example, AKh admits spectral sequences to gauge and Floer theories (see [Rob13] and [Xie21]), enjoys an $\mathfrak{sl}_2(\mathbb{C})$ action (see [GLW18] and [Kim21]), and has been extended to equivariant settings (see [BPW19] and [Akh23]).

As a categorification of the Kauffman bracket skein module of the annulus KBSM(A) [Prz91], annular Khovanov homology inherits long-open questions from skein theory, including the conjecture at the center of

Date: January 7, 2025.

the present article. The wrapping number of an annular link L, denoted by wrap(L), is the minimal geometric intersection between a link L and a meridional disk in $A \times I$. This quantity is related to the annular winding number grading, or k-grading, in both KBSM(A) and AKh, which keeps track of the algebraic wrapping of each Kauffman state around the *, in a sense. In particular, AKh(L) is supported in k-gradings with absolute value $\leq \operatorname{wrap}(L)$ [GLW18].

This naturally leads to the following conjecture of Grigsby from 2010^{-1} :

Conjecture 1. [Grigsby, MSRI 2010] (Categorified Wrapping Number Conjecture) The annular Khovanov homology of an annular link L is nontrivial in k-grading wrap(L).

We direct the reader to [Kim21] for a summary of present bounds and attempts to prove Conjecture 1. Grigsby's conjecture is motivated by the following older conjecture, first posed in a seminal paper by Hoste and Przytycki in 1995:

Conjecture 2. [HP95] (Wrapping Number Conjecture) The Kauffman bracket skein module of an annular link L is nonzero in annular degree wrap(L).

In support of their conjecture, Hoste–Przytycki proved that Conjecture 2 holds for *adequately wrapped* annular links. An adequately wrapped annular link enjoys diagrams with particular resolutions, ones which guarantee that the link satisfies Conjecture 2.

In this paper, we generalize Hoste–Przytycki's results in the categorified setting by identifying a broader class of resolutions which may be associated to a nonvanishing annular Khovanov homology class. We refer to these resolutions as *perfectly wrapped* and *uniform*. The property of being perfectly wrapped limits the type of cobordisms entering and exiting the resolution. Uniformity is a condition on trivial circles; it requires each trivial circle to abut only 0-resolutions of crossings, or only 1-resolutions of crossings.

An abridged version of our main theorem, Theorem 30, is the following:

Theorem 3. Let L be an annular link with a diagram D admitting a perfectly wrapped uniform resolution D_u . Then AKh(L) is nontrivial in k-grading wrap(L).

In particular, every perfectly wrapped uniform resolution D_u produces at least one distinguished generator $x \in \text{CAKh}(D_u)$ such that

generates a nonzero annular Khovanov homology class in wrap(L) k-grading.

Theorem 3 follows from the data of a single resolution and the cobordisms entering and exiting it. Namely, we look at the distinguished generators which are always sent to zero by $\operatorname{wrap}(L)$ k-graded differential components, and the distinguished generators which are never mapped to by $\operatorname{wrap}(L)$ k-graded differential components.

Theorem 3 encompasses Hoste–Przytycki's adequately wrapped links, along with a number of other annular link classes, chief among them alternating annular links.

The proof for alternating annular links requires a transformation between another type of resolution and perfectly wrapped uniform resolutions. This former type is called *almost uniform* as opposed to uniform. It has an alternate condition on its trivial circles: Each trivial circle in an almost uniform resolution abuts a singular kind of crossing resolution on its exterior, and the opposite kind on its interior.

The definition is motivated entirely by Lemma 40, which gives a perfectly wrapped almost uniform resolution for every alternating diagram for an annular link. The following theorem and corollary, adapted from Theorem 38 and Corollary 42, completes the proof. They are proven entirely in the setting of resolutions.

Theorem 4. Let D be a diagram for an annular link L, with no nugatory crossings (understood in the annular setting). Suppose that D admits a perfectly wrapped almost uniform resolution D_u . Then D admits a perfectly wrapped uniform resolution D_v .

Corollary 5. The Categorified Wrapping Number Conjecture holds for every alternating annular link.

Beyond alternating annular links, we examine phenomena associated with plumbed links. The following takes from Corollaries 43 and 44:

¹See the anecdote under Conjecture 1.2 on page 2 of [Mar23].





(a) A 4-braid with inserted horizontal 2-braids ('chains') inserted in two locations.

(b) A braid wearing unknotted belts.

FIGURE 1. Annular closures of these tangles are examples of the two classes of annular links from Corollary 6.

Corollary 6. The annular closures of the following classes of tangles satisfy the conditions of Theorem 3, and therefore Conjecture 1:

- (1) A braid (drawn vertically) with some crossings replaced by horizontal 2-braids; in other words, the result of surgering a braid along some twisted horizontal bands. See Figure 1a for an example.
- (2) A composition of braids and identity braids wearing belts. See Figure 1b for an example.

The proof of Corollary 6 relies on work of Grigsby–Ni, who showed that AKh over \mathbb{C} coefficients is 1dimensional at k-grading wrap(L) for braid closures [GN14]. We note the similarities between these types and work of Martin; she found additional classes of adequately wrapped annular links, proving both Conjecture 1 and Conjecture 2 for these classes [Mar23].

Hoste–Przytycki used the fact that the blackboard-framed n-cabling of an adequately wrapped diagram for an annular link is also adequately wrapped [HP95]. We obtain an analogous result, with an additional observation about adding unlink components to diagrams. The following is adapted from Corollaries 45 and 46.

Corollary 7. Suppose that D is a diagram for an annular link L that admits a perfectly wrapped uniform resolution. Then

- (1) The blackboard-framed n-cabling of D admits a perfectly wrapped uniform resolution (see Figure 2a for an example), and
- (2) The addition of an arbitrary number of disjoint loops ('earrings'), each wrapping around a single strand, to D produces a diagram with a perfectly wrapped uniform resolution (see Figure 2b for an example).

It remains to be seen how far this approach can be taken. In one direction, Theorem 3 could likely be upgraded to prove Conjecture 2 for certain annular links (perhaps without additional conditions). In another direction, we are motivated to ask about the extent to which links admit perfectly wrapped uniform resolutions.

Question 8. Does every annular link admit a diagram with a perfectly wrapped uniform resolution?

If true, Question 8 would imply Conjecture 1 by Theorem 3. It would also imply that the rank of the annular Khovanov homology in k-grading wrap(L) (over \mathbb{Z} coefficients) is nonzero in general.

In light of the above, it is feasible that some annular link will not admit a diagram with a perfectly wrapped uniform resolution; we may have a link whose annular Khovanov homology in k-grading wrap(L) is a direct sum of torsion groups.

We can also ask about the maximal number of perfectly wrapped uniform resolutions an annular link may admit for a given diagram. Denote this quantity by U(L) in the manner of Corollary 31.



(a) A blackboard-framed 2-cabling of a square knot in the thickened annulus.



FIGURE 2. Examples of operations from Corollary 7.

Question 9. Can we identify, for any annular link L, a class of diagrams D for L such that the number of perfectly wrapped uniform resolutions of D is equal to U(L)?

The authors believe that this equality holds when D is a minimal crossing diagram.

1.1. Acknowledgements. Special thanks to Jozef Przytycki for providing the authors access to his manuscript. We also thank Wren Burrill and Nathan Singh for useful conversations.

2. ANNULAR KHOVANOV HOMOLOGY

Throughout the article, we work over \mathbb{Z} coefficients. We will indicate when we use \mathbb{C} coefficients as appropriate, e.g. in reference to the $\mathfrak{sl}_2(\mathbb{C})$ action on AKh over \mathbb{C} .

2.1. Cube of resolutions. We recount the method of computation for annular Khovanov homology. Let $L \subseteq A \times I$ be an annular link, presented via an annular diagram D with n crossings: n_+ positive and n_- negative. We now form the *cube of resolutions* for D.

Start with a directed graph Γ , where each vertex is some binary string $u \in \{0,1\}^n$. Let |u| denote the sum of each digit in u; i.e., the number of 1's in u. Then construct an edge $u \to v$ if and only if |u| = |v| - 1, written $u \prec_1 v$. An example for n = 3 is given below:



Number the crossings in D. Then we assign to any given binary string u in Γ a complete resolution of D, where the *i*th crossing is resolved according to the *i*th digit of u. The two types of resolutions are as follows:

$$\Big\rangle\Big\langle \xleftarrow{0} \\ \xleftarrow{1} \\ \xrightarrow{1} \\ \ddots \\ \cdot$$

We then refer to the complete resolution of D as D_u , or the complete resolution of D corresponding to the binary string u. In general, D_u will consist of a collection of planar circles of two types: *trivial* and *nontrivial*. The former class of circles are those which do not wind around the *, while the latter class are those which do.

There is now the question of what topological data the edges should encode. As each edge presently represents a bit flip, the resolution at the source of an edge will differ from the resolution at the target by a single crossing resolution swap, from type 0 to type 1 as above. Enacting this swap can affect the planar circles in the source resolution in six different ways, as shown in Figure 3.

Each merge or split is a cobordism between the collections of planar circles. For instance, a merge of two circles into one can be locally represented as a "pair of pants" cobordism, shown in Figure 4. Throughout,



FIGURE 3. Left column, from top to bottom: merges $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}, \mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$, and $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$, respectively. Right column, from top to bottom: splits $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}, \mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$, and $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$.



FIGURE 4. A pair of pants cobordism, which represents a merging of two circles when read from top to bottom. The red arc is the surgery arc on the 0-resolution; the blue arc is the dual surgery arc on the 1-resolution.

we use red arcs on 0-resolutions to indicate the surgery arc, so that surgering along this red arc results in the 1-resolution of the crossing. Dually, we use blue arcs on 1-resolutions to indicate the dual surgery arc.

Remark 10. Throughout, we use the term 'arc' to refer to 'surgery arc', as explained above. Unfortunately, components of knot diagrams between crossings are also sometimes referred to as arcs; we will not be using this terminology.

2.2. The annular Khovanov chain complex. With the combinatorics of crossing resolution switches handled, we are prepared to transform the cube of resolutions from a graph consisting of topological data to a chain complex consisting of algebraic data.

Let D_u be a given complete resolution corresponding to some binary string u. Begin by numbering each circle in D_u .

Let $\mathbb{V} = \mathbb{Z}v_+ \oplus \mathbb{Z}v_-$ and $\mathbb{W} = \mathbb{Z}w_+ \oplus \mathbb{Z}w_-$; these are bi-graded modules with quantum grading gr_q and k-grading gr_k determined by the bi-gradings on their generators:

- $(gr_a, gr_k)(v_{\pm}) = (\pm 1, \pm 1)$
- $(\operatorname{gr}_{a}, \operatorname{gr}_{k})(w_{\pm}) = (\pm 1, 0).$

Remark 11. In terms of the $\mathfrak{sl}_2(\mathbb{C})$ action, $\mathbb{V} \otimes_{\mathbb{Z}} \mathbb{C}$ is the fundamental representation for $\mathfrak{sl}_2(\mathbb{C})$, and $\mathbb{W} \otimes_{\mathbb{Z}} \mathbb{C}$ is two copies of the trivial representation. The annular k-grading is the $\mathfrak{sl}_2(\mathbb{C})$ weight-space grading.

Define the module $\operatorname{CAKh}(D_u)$ as

$$CAKh(D_u) := \bigotimes_i \begin{cases} \mathbb{V} & \text{if the } i\text{th circle is nontrivial} \\ \mathbb{W} & \text{if the } i\text{th circle is trivial} \end{cases}.$$

Up to reordering, we may number our circles such that the module $CAKh(D_u)$ can be written as

 $\mathbb{V}^{\otimes \#}$ of nontrivial circles in $D_u \otimes \mathbb{W}^{\otimes \#}$ of trivial circles in D_u .

The pure tensors that generate $CAKh(D_u)$ are called *distinguished generators*. Observe that these are homogeneous in bidegree, and are easily computed by the formulas

$$\begin{split} \mathrm{gr}_q &= |u| + \# v_+ + \# w_+ - \# v_- - \# w_- \\ \mathrm{gr}_k &= \# v_+ - \# v_-. \end{split}$$

Remark 12. We can interpret a distinguished generator x in a summand $\operatorname{CAKh}(D_u)$ as a choice of orientation for each trivial and nontrivial circle in D_u , with v_+, w_+ associated with a counterclockwise (CCW) orientation, and v_-, w_- associated with a clockwise (CW) orientation. Denote this oriented 1-manifold embedded in the annulus by (D_u, x) .

The reader may check that the gr_k of distinguished generator corresponds to the winding number of the oriented 1-manifold (D_u, x) around the *: this is the algebraic intersection number of (D_u, x) and a generic curve γ connecting * with ∞ .

The next step is to associate to every edge a module map. Along an edge $D_u \to D_v$, a circle in D_u or D_v is called *active* if it abuts the surgery arc or the dual surgery arc. Otherwise, it is *passive*. Below, we give the explicit definition of the module map $d_{u,v}$ associated to the cobordism $D_u \to D_v$.

Definition 13. The annular Khovanov differentials are explicitly defined as follows, extended by identity on passive circles. (Refer to Figure 3 for the associated cobordisms.)

• A merge of two trivial circles $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$:

$$\begin{array}{l} d_{u,v}: \mathbb{W} \otimes \mathbb{W} \to \mathbb{W} \\ w_+ \otimes w_+ \mapsto w_+ \\ w_- \otimes w_+ \mapsto w_- \\ w_+ \otimes w_- \mapsto w_- \\ w_- \otimes w_- \mapsto 0 \end{array}$$

• A merge of a nontrivial circle and a trivial circle $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$:

$$\begin{array}{l} d_{u,v}: \mathbb{V}\otimes\mathbb{W}\rightarrow\mathbb{V}\\ v_+\otimes w_+\mapsto v_+\\ v_+\otimes w_-\mapsto 0\\ v_-\otimes w_+\mapsto v_-\\ v_-\otimes w_-\mapsto 0 \end{array}$$

• A merge of two nontrivial circles $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$:

$$\begin{array}{l} d_{u,v}: \mathbb{V} \otimes \mathbb{V} \to \mathbb{W} \\ v_+ \otimes v_+ \mapsto 0 \\ v_- \otimes v_+ \mapsto w_- \\ v_+ \otimes v_- \mapsto w_- \\ v_- \otimes v_- \mapsto 0 \end{array}$$



FIGURE 5. A dots-and-arrows depiction of a \mathbb{Z} -module merge map $m : \mathbb{V} \otimes \mathbb{V} \to \mathbb{W}$. There are two arrows a, b with sources $v_+ \otimes v_-$ and $v_- \otimes v_+$, respectively. The target of both arrows is w_- .

• A split of one trivial circle into two trivial circles $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}$:

$$d_{u,v}: \mathbb{W} \to \mathbb{W} \otimes \mathbb{W}$$
$$w_+ \mapsto w_- \otimes w_+ + w_+ \otimes w_-$$
$$w_- \mapsto w_- \otimes w_-$$

• A split of one nontrivial circle into a nontrivial circle and a trivial circle $\mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$:

$$d_{u,v}: \mathbb{V} \to \mathbb{V} \otimes \mathbb{W}$$
$$v_+ \mapsto v_+ \otimes w_-$$
$$v_- \mapsto v_- \otimes w_-$$

• A split of one trivial circle into two nontrivial circles $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$:

$$\begin{aligned} d_{u,v} &: \mathbb{W} \to \mathbb{V} \otimes \mathbb{V} \\ w_+ &\mapsto v_- \otimes v_+ + v_+ \otimes v_- \\ w_- &\mapsto 0 \end{aligned}$$

The reader may wish to verify that these maps preserve the (gr_q, gr_k) bi-grading.

Notation 14. We will also take a more combinatorial perspective on these modules and maps. Any chain $\mathbf{x} \in \text{CAKh}(D)$ can be written as a finite sum of distinguished generators $\sum_i c_i g_i$. Let $\langle x, g_i \rangle$ denote the coefficient of the distinguished generator g_i in \mathbf{x} , using the distinguished basis.

Let g, g' be distinguished generators, and observe that

$$\langle d(g), g' \rangle \in \{-1, 0, 1\}.$$

This special property of Khovanov differentials allows us to interpret the chain complex (CAKh(D), d) using "dots and arrows", as follows and as depicted in Figure 5.

- Each distinguished generator is represented by a dot.
- The differential is a collection of arrows; there is an arrow from g to g' if and only if $\langle d(g), g' \rangle = \pm 1$. (We may decorate the arrow if we wish to keep track of signs.)

Suppose there is an arrow a from g to g'. Then we call g the source of the arrow a, and g' the target of the arrow a.

We now have the data of a collection of modules, each on the vertices of our graph Γ , along with module maps corresponding to the edges in Γ .

To combine these into a chain complex, we first assign signs $\epsilon_{u,v} \in \{\pm 1\}$ to each edge $u \prec_1 v$ in the cube, so that each square in Γ has an odd number of -1's. Then, we define the (unshifted; see Remark 15) annular Khovanov chain complex at (co)homological² grading *i* to be

$$\operatorname{CAKh}^{i}(D) = \bigoplus_{|u|=i} \operatorname{CAKh}(D_{u})$$

 $^{^{2}}$ Khovanov homology was defined with cohomological grading conventions; the differential increases homological grading by 1.

and the differential $d_i : \operatorname{CAKh}^i(D) \to \operatorname{CAKh}^{i+1}(D)$ by

$$d_i = \bigoplus_{|u|=i} \sum_{u \prec_1 v} \epsilon_{u,v} d_{u,v}$$

To check that $d^2 = 0$, it suffices to check that any face of the cube



commutes, and that the additional signs make the faces anti-commute.

The homology of this complex is denoted AKh(D). Let AKh(L;k) = AKh(D;k) denote the annular Khovanov homology of L at $gr_k = k$.

Remark 15. The expert reader may notice that our definition of AKh is not quite the usual definition, because we omit the global homological and quantum shifts that make annular Khovanov homology diagramindependent. The reason for this is that we are focused on the k-grading, which does not have a global shift, and it is easier to work directly with the unshifted complex.

For a diagram D of an annular link L, with n_{\pm} crossings with sign \pm , respectively, our unshifted AKh(D) is related to the usual annular Khovanov homology AKh(L), as it appears in the literature, by

$$AKh^{i,j,k}(L) = AKh^{i-n_-,j+n_+-2n_k}(D).$$

In particular, if we forget the homological and quantum gradings, we are justified in writing AKh(L;k), as this is indeed independent of the choice of diagram D.

2.3. Wrapping number of diagrams and resolutions. Let D be a diagram for an annular link L. We define the wrapping number of D, denoted wrap(D), to be the minimal geometric intersection between D and a meridional arc in A. Then the wrapping number of L is

$$\operatorname{wrap}(L) = \min_{\operatorname{diagrams} D} \operatorname{wrap}(D).$$

For a resolution D_u of D, we can similarly define wrap (D_u) ; wrap (D_u) can also be computed as the number of nontrivial circles in D_u .

If we suppose that D satisfies wrap(L) = wrap(D), then every resolution of D will have wrapping number less than or equal to wrap(L). Consequently, computing the annular Khovanov homology from our cube of resolutions for D will produce generators x_i with $|\text{gr}_k(x_i)| \leq \text{wrap}(L)$. Therefore AKh(L) is nontrivial only within these k-gradings as well.

The reader may now verify that every annular Khovanov differential preserves the parity of nontrivial circles. We conclude that AKh(L) must have support contained in the set of k-gradings

$$\{-wrap(L), -wrap(L) + 2, ..., wrap(L) - 2, wrap(L)\},\$$

as [GLW18] found.

3. Perfectly wrapped and uniform resolutions

In this section, we identify instances in which the existence of a single type of resolution suffices to verify Conjecture 1, in the style of [HP95] and [Mar23].

3.1. Terminology and notation. We first establish terminology and notation for the coming sections. For a given resolution D_u , we let $|D_u| = |\pi_0(D_u)|$, i.e. the number of components.

We denote a cobordism merging two trivial circles into one by $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$, a cobordism merging a trivial circle a nontrivial circle into a nontrivial circle by $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$, etc.

For an annular link L, we are only concerned with the component of CAKh(L) at $\text{gr}_k = \text{wrap}(L)$. As the annular differential respects k-grading, we may analyze this component in isolation. We initially restrict our search for nonvanishing classes to resolutions with wrapping number equivalent to wrap(L).

Definition 16. Let D be a diagram for an annular link L. We say that a resolution D_u of D is *exactly wrapped* if the number of nontrivial circles in D_u is equal to wrap(L).

To justify our frequent usage of these resolutions, we prove that they exist for arbitrary diagrams. We will show a couple of lemmas to aid in this proof.

Lemma 17. For any annular diagram D for an annular link L, wrap $(D) \equiv wrap(L) \mod 2$.

Proof. The diagram $D \hookrightarrow A$ represents some $\mathbb{Z}/2$ homology class $[D] \in H_1(A; \mathbb{Z}/2) \cong \mathbb{Z}/2$. This is invariant under annular isotopy and annular Reidemeister moves, so wrap $(D) \equiv \operatorname{wrap}(D') \mod 2$ for any diagrams D, D' of L. In particular, wrap $(D) \equiv \operatorname{wrap}(L) \mod 2$, because wrap $(L) \mod 2 = [L] \in H_1(A \times I; \mathbb{Z}/2) \cong \mathbb{Z}/2$. \Box

Lemma 18. Given any connected diagram D for L, there are resolutions D_u for each wrapping number between wrap $(D) \mod 2$ and wrap(D) (of the correct parity).

Proof. Take a resolution D_u of D such that $\operatorname{wrap}(D_u) = \operatorname{wrap}(D)$. In particular, $\operatorname{wrap}(D_u)$ is maximal out of all resolutions for D. As D is connected, we may use a series of cobordisms of the type $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$ and $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$ to obtain a resolution D_v of D where $\operatorname{wrap}(D_v) = \operatorname{wrap}(D)$ and every circle in D_v is nontrivial. Using the fact that D is connected once again, each nontrivial circle in D_v must abut some other nontrivial circle in D_v by a crossing resolution. We can then obtain the desired resolutions by repeated applications of merges $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$, which preserve the parity of the number of nontrivial circles in a resolution. \Box

Let D, D' be annular link diagrams, viewed as diagrams in $S^1 \times [1,2]$ and $S^1 \times [2,3]$, respectively. The horizontal composition of D and D' is the annular diagram $D \sqcup D'$ in $S^1 \times [1,3]$.

Proposition 19. Let D be a diagram for an annular link L. Then D has an exactly wrapped resolution.

Proof. From Lemma 17, we know that $\operatorname{wrap}(D) \equiv \operatorname{wrap}(L) \mod 2$. Moreover, $\operatorname{wrap}(D) \geq \operatorname{wrap}(L)$ by definition. Lastly, for a diagram $D = D_1 \sqcup \ldots \sqcup D_n$ consisting of the horizontal composition of n connected diagrams, we have

$$wrap(D) = wrap(D_1) + \dots + wrap(D_n)$$

Our proposition follows immediately from Lemma 18 applied to $D_1, ..., D_n$.

We will invoke Proposition 19 implicitly throughout the coming sections.

Within a collection of resolutions, we may select especially useful ones with restrictions on their incoming and outgoing cobordisms.

Definition 20. Let D be a diagram for an annular link L. A resolution D_u is *insulated* if

- Every cobordism with source D_u is of the form $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}, \mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}, \mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}, \text{ or } \mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}.$
- Every cobordism with target D_u is of the form $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}, \mathcal{V} \to \mathcal{V} \sqcup \mathcal{V}, \text{ or } \mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}.$

In other words, the resolutions adjacent to D_u either have strictly less trivial circles than D_u , or have a different number of nontrivial circles. This definition is similar to the condition on 0- and 1-smoothings of links in the classical definition of adequacy.

Definition 21. A resolution D_u is *perfectly wrapped* if D_u is both exactly wrapped and insulated.

We now show that perfectly wrapped resolutions exist as a corollary of Proposition 19.

Lemma 22. Every diagram D for an annular link L admits a perfectly wrapped resolution.

Proof. Let D_u be an exactly wrapped resolution of D such that, for any other exactly wrapped resolution D_v ,

of trivial circles in $D_u \ge \#$ of trivial circles in D_v .

Suppose that D_v is an exactly wrapped resolution for which there is a cobordism $D_v \to D_u$. Then this cobordism must be of the form $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}$ or $\mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$ by the above condition. If D_w is an exactly wrapped resolution for which there is a cobordism $D_u \to D_w$, then this cobordism must be of the form $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$ or $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{W}$ in a similar fashion. This ensures that every incoming and outgoing cobordism is as desired.

From the perspective of a single resolution, we wish to analyze the data of its incident cobordisms.

Definition 23. We define the *null set of* D_u , written $\mathcal{K}(D_u)$, to be the set of all distinguished generators which are not the source of any arrow at $\operatorname{gr}_k = \operatorname{wrap}(L)$. We define the *target set of* D_u , written $\mathcal{I}(D_u)$, to be the set of all distinguished generators that are targets for some arrow at $\operatorname{gr}_k = \operatorname{wrap}(L)$.

Usually, $\mathcal{I}(D_u)$ is not a subset of $\mathcal{K}(D_u)$ in absence of a cocycle condition. However, these objects still may allow us to verify Conjecture 1 in the case that $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$, as a distinguished generator $x \in \mathcal{K}(D_u) \setminus \mathcal{I}(D_u)$ represents a nonvanishing homology class in AKh. We can also obtain information about D_u .

Lemma 24. If $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$, then D_u is insulated. Furthermore, $AKh(L; wrap(L)) \cong 0$.

Proof. For the first statement, we will prove the contrapositive. A cobordism with source D_u of the form $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}$ or $\mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$ induces an always nonzero map in wrap(L) grading, implying that $\mathcal{K}(D_u) = \emptyset$. A cobordism with target D_u of the form $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$ or $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$ induces a map whose image spans $\operatorname{CAKh}(D_u)$, meaning that $\mathcal{K}(D_u) \subseteq \mathcal{I}(D_u)$, and our claim holds.

For the second statement, let $x \in \mathcal{K}(D_u) \setminus \mathcal{I}(D_u)$. Then the element

$$(0, ..., 0, x, 0, ..., 0) \in \operatorname{CAKh}^{|u|}(D)$$

is in the kernel of $d_{|u|}$, as $x \in \mathcal{K}(D_u)$. As $x \notin \mathcal{I}(D_u)$, x is not the target of a single arrow by definition. Consequently, $d_{|u|-1}$ does not contain (0, ..., 0, x, 0, ..., 0) in its image. Therefore $AKh(L; wrap(L)) \ncong 0$. \Box

Our goal is to discern exactly which resolutions satisfy the conditions of Lemma 24, so that we may verify Conjecture 1 for the link we are resolving.

3.2. Type 0 and type 1 trivial circles. We now define the qualities of trivial circles which we seek to investigate.

Definition 25. Let D_u be a resolution. We say a trivial circle W in D_u is type 0 if it only abuts 0-resolutions of crossings (i.e. red arcs). Similarly, we say a trivial circle W in D_u is type 1 if it only abuts 1-resolutions of crossings (i.e. blue arcs).

In the lemmas below, we use the shorthand notation w_{\pm} to mean 'either w_{\pm} or w_{\pm} '.

Lemma 26. Let D_u be perfectly wrapped. Let n_1 denote the number of type 1 trivial circles in D_u . Then, for a distinguished generator $x \in \text{CAKh}(D_u)$, we have

$$x \in \mathcal{K}(D_u)$$

if and only if x is of the form

(1)
$$\underbrace{v_{+}^{\otimes \operatorname{wrap}(L)}}_{nontrivial} \otimes \underbrace{w_{\pm}^{\otimes n_{1}}}_{type 1} \otimes \underbrace{w_{-}^{\otimes (|D_{u}| - \operatorname{wrap}(L) - n_{1})}}_{not type 1},$$

where every trivial circle that is **not** type 1 is labeled w_{-} .

Proof. Recall that $\mathcal{K}(D_u)$ only contains distinguished generators with k-grading equal to wrap(L), by definition. Therefore, for a distinguished generator to be in $\mathcal{K}(D_u)$, every nontrivial circle must be labeled v_+ .

For the matter of trivial circles, observe that a trivial circle W in D_u is active for some cobordism with source D_u if and only if W is not type 1.

We show that if a distinguished generator is not of the form above, then it is not in the kernel. Consider a distinguished generator

$$y = \underbrace{v_{\pm}^{\otimes \operatorname{wrap}(L)}}_{\operatorname{nontrivial}} \otimes \underbrace{w_{\pm}^{\otimes n_1}}_{\operatorname{type 1}} \otimes \underbrace{w_{\pm}}_{\operatorname{not type 1}} \otimes \underbrace{w_{\pm}^{\otimes (|D_u| - \operatorname{wrap}(L) - n_1 - 1)}}_{\operatorname{not type 1}},$$

where some trivial circle W that is not type 1 is labeled w_+ . As W is not type 1, W is active for some cobordism $D_u \to D_v$. The map induced by this cobordism will be either $\mathbb{W} \otimes \mathbb{W} \to \mathbb{W}$, $\mathbb{V} \otimes \mathbb{W} \to \mathbb{W}$, or $\mathbb{W} \to \mathbb{V} \otimes \mathbb{V}$. By inspection of the AKh differentials (Definition 13), one may check that y is not in the kernel of any of these maps. Hence, $y \notin \mathcal{K}(D_u)$.

On the other hand, let x be a distinguished generator of the form shown in (1), where every trivial circle that is not type 1 is labeled w_- . As D_u is perfectly wrapped, there are four possible cobordisms with source D_u , namely $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}, \mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}, \mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$, and $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$.

- (1) Consider the map $m_{W,W'} : \mathbb{W} \otimes \mathbb{W} \to \mathbb{W}$ corresponding to the cobordism $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$, which merges trivial circles W and W'. As both trivial circles are active for this cobordism, they are connected by a 0 crossing resolution, meaning that neither are type 1. Therefore, they are both labeled w_- . Hence, $m_{W,W'}(x) = 0$.
- (2) For the map $m_{V,W} : \mathbb{V} \otimes \mathbb{W} \to \mathbb{V}$ corresponding to the cobordism $\mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}$, merging a nontrivial circle V and a trivial circle W, we know that W is not type 1 as it is active for this cobordism. Hence, it is labeled w_{-} , implying $m_{V,W}(x) = 0$.
- (3) Next, we consider the map $m_{V,V'} : \mathbb{V} \otimes \mathbb{V} \to \mathbb{W}$ corresponding to the cobordism $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$, which merges nontrivial circles V and V'. This map is trivial at the wrap(L) k-grading, so $m_{V,V'}(x) = 0$.
- (4) Finally, we have the map $\Delta_{V,V'} : \mathbb{W} \to \mathbb{V} \otimes \mathbb{V}$ corresponding to the cobordism $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$, which splits a trivial circle W into two nontrivial circles V, V'. Given that W is active for this cobordism, W is not type 1, and is thus labeled by w_- . Therefore, $\Delta_{V,V'}(x) = 0$.

Therefore, along each edge of the cube of resolutions with source D_u , the induced map sends x to 0, so $x \in \mathcal{K}(D_u)$.

The following lemma is, in a sense, dual to the previous lemma.

Lemma 27. Let D_u be perfectly wrapped. Let n_0 denote the number of type 0 trivial circles in D_u . Then, for a distinguished generator $x \in \text{CAKh}(D_u)$, we have

$$x \notin \mathcal{I}(D_u)$$

if and only if x is of the form

(2)
$$\underbrace{v_{+}^{\otimes \operatorname{wrap}(L)}}_{nontrivial} \otimes \underbrace{w_{\pm}^{\otimes n_{0}}}_{type \ 0} \otimes \underbrace{w_{+}^{\otimes (|D_{u}| - \operatorname{wrap}(L) - n_{0})}}_{not \ type \ 0}$$

where every trivial circle that is not type 0 is labeled w_+ .

Proof. By the same reasoning as in Lemma 26, we must label our nontrivial circles by v_+ .

A trivial circle W in D_u is active for a cobordism $D_v \to D_u$ if and only if W is **not** type 0. Consider a distinguished generator

$$y = \underbrace{v_{+}^{\otimes \operatorname{wrap}(L)}}_{\operatorname{nontrivial}} \otimes \underbrace{w_{\pm}^{\otimes n_{0}}}_{\operatorname{type } 0} \otimes \underbrace{w_{-}}_{\operatorname{not type } 0} \otimes \underbrace{w_{\pm}^{\otimes (|D_{u}| - \operatorname{wrap}(L) - n_{0} - 1)}}_{\operatorname{not type } 0},$$

where some trivial circle W that is not type 0 is labeled w_- . By the previous remark, there is a cobordism $D_v \to D_u$ affecting W. This cobordism will induce a map $\mathbb{W} \to \mathbb{W} \otimes \mathbb{W}$, $\mathbb{V} \otimes \mathbb{V} \to \mathbb{W}$, or $\mathbb{V} \to \mathbb{V} \otimes \mathbb{W}$. We may then check that y is in the image of each of these maps, so that $y \in \mathcal{I}(D_u)$.

Let x be of the form shown in (2), where every trivial circle that is not type 0 is labeled w_+ . As D_u is perfectly wrapped, there are four possible cobordisms with target D_u : (1) $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}$, (2) $\mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$, (3) $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$, and (4) $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$.

Case (3) is immediate, as the induced map is trivial in wrap(L) k-grading. Cases (1) and (2) are both splits, where the active trivial circles in D_u are all labeled +; therefore, we may check that x is not in the image of either of these induced maps. For case (4), the active trivial circle is labeled +. Once again, the induced map does not contain x in its image. We conclude that $x \notin \mathcal{I}(D_u)$ as desired.

3.3. Proof of the main result. We will now introduce a key definition, one which will allow us to accomplish our goal of classifying resolutions D_u for which Lemma 24 applies.

Definition 28. We say a resolution D_u of a diagram D for an annular link L is *uniform* if every trivial circle in D_u is of type 0 or type 1.

Note that we allow for the possibility of split nullhomologous unknot components of D in the annulus; these are trivial circles that are both type 0 and type 1.

Lemma 29. If $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$, then D_u is uniform.

Proof. We will prove the contrapositive. Assume that D_u is not uniform. First note that if D_u is not insulated, then Lemma 24 implies that $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) = \emptyset$, and we are done. So, we may assume that D_u is insulated.

As D_u is not uniform, there exists some trivial circle W in D_u which abuts both a 0-resolution and a 1-resolution.

As W abuts a 0-resolution, it is active for a cobordism $D_u \to D_v$. If y is a distinguished generator in $\mathcal{K}(D_u)$, we may check by cases on the induced maps (as we do in Lemma 26) that W must be labeled by w_- in y, keeping in mind that D_u is insulated.

As W also abuts a 1-resolution, it is active for a cobordism $D_w \to D_u$ of the form $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}, \mathcal{V} \to \mathcal{V} \sqcup \mathcal{W}$, or $\mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$. Recalling that W is labeled w_{-} in y, we may again check that each of these cases places y in $\mathcal{I}(D_u)$.

Thus $\mathcal{K}(D_u) \subseteq \mathcal{I}(D_u)$, and so $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) = \emptyset$ as desired. \Box

We are now ready to prove the main theorem.

Theorem 30. Let D be a diagram for an annular link L. Then a resolution D_u of D is perfectly wrapped and uniform if and only if $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$. In this case, let n_0, n_1, n_2 respectively denote the number of trivial circles which are type 0 and not type 1, type 1 and not type 0, and both type 0 and type 1. Then, up to renumbering of trivial circles, $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u)$ contains exactly those distinguished generators x of the form

(3)
$$v_{+}^{\otimes \operatorname{wrap}(L)} \otimes \underbrace{w_{-}^{\otimes n_{0}}}_{type \ 0} \otimes \underbrace{w_{\pm}^{\otimes n_{2}}}_{both} \otimes \underbrace{w_{+}^{\otimes n_{1}}}_{type \ 1}.$$

In particular, L satisfies Conjecture 1.

Proof. (\Longrightarrow) Suppose that D_u is perfectly wrapped and uniform. Up to renumbering of trivial circles, we may use Lemma 26 to conclude that $\mathcal{K}(D_u)$ contains exactly those distinguished generators of the form

$$v_{+}^{\otimes \operatorname{wrap}(L)} \otimes \underbrace{w_{-}^{\otimes n_{0}}}_{\operatorname{type 0}} \otimes \underbrace{w_{\pm}^{\otimes n_{2}} \otimes w_{\pm}^{\otimes n_{1}}}_{\operatorname{type 1}},$$

in which each type 0 trivial circle is labeled w_{-} . Similarly, Lemma 27 tells us that $\mathcal{I}(D_u)$ exactly excludes those distinguished generators of the form

$$v_{+}^{\otimes \operatorname{wrap}(L)} \otimes \underbrace{w_{\pm}^{\otimes n_{0}} \otimes w_{\pm}^{\otimes n_{2}}}_{\operatorname{type 0}} \otimes \underbrace{w_{\pm}^{\otimes n_{1}}}_{\operatorname{type 1}}$$

in which each type 1 trivial circle is labeled w_+ .

Now define

(4)
$$x = v_{+}^{\otimes \operatorname{wrap}(L)} \otimes \underbrace{w_{-}^{\otimes n_{0}}}_{\operatorname{type 0}} \otimes \underbrace{w_{\pm}^{\otimes n_{2}}}_{\operatorname{both}} \otimes \underbrace{w_{+}^{\otimes n_{1}}}_{\operatorname{type 1}}.$$

By inspection, $x \in \mathcal{K}(D_u) \setminus \mathcal{I}(D_u)$, so $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$.

(\Leftarrow) Assume that $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u) \neq \emptyset$.

By Lemmas 24 and 29, D_u is insulated and uniform. Moreover, wrap $(D_u) \ge \text{wrap}(L)$ as $\mathcal{K}(D_u)$ is nonempty (recall that $\mathcal{K}(D_u)$ contains only generators at k-grading wrap(L)). If wrap $(D_u) > \text{wrap}(L)$, then we may consider the distinguished generator

(5)
$$z = v_{+}^{\otimes \operatorname{wrap}(D_{u})} \otimes \underbrace{w_{-}^{\otimes n_{0}}}_{\operatorname{type 0}} \otimes \underbrace{w_{\pm}^{\otimes n_{2}}}_{\operatorname{both}} \otimes \underbrace{w_{+}^{\otimes n_{1}}}_{\operatorname{type 1}}.$$

For every cobordism with source D_u , the induced map takes z to 0. Moreover, for every cobordism with target D_u , the induced map does not contain z in its image. This means that

$$(0, ..., 0, z, 0, ..., 0) \in CAKh^{|u|}(D)$$

generates a non-zero homology class in $AKh(L; wrap(D_u))$, contradicting the fact that AKh(L) is supported in k-gradings less than or equal to $wrap(D_u)$. Therefore, $wrap(D_u) = wrap(L)$ must hold, implying that D_u is perfectly wrapped and uniform by definition.

The final claims are fairly immediate; only distinguished generators in $\text{CAKh}(D_u)$ of the form in (4) are both in the null set of D_u and not in the target set of D_u . Hence, $\mathcal{K}(D_u) \setminus \mathcal{I}(D_u)$ contains exactly the generators of the form (4). An application of Lemma 24 completes the proof, showing Conjecture 1 for L. Corollary 31. Let L be an annular link. Define

$$U(L) = \max_{\text{diagrams } D} \#(\text{perfectly wrapped uniform resolutions of } D).$$

Then

rank
$$AKh(L; wrap(L)) \ge U(L)$$

Proof. From Theorem 30 and Lemma 24, every perfectly wrapped uniform resolution corresponds to at least one distinct homology class in AKh(L; wrap(L)), generating \mathbb{Z} .

Corollary 32. If $AKh(L; wrap(L))/Tors \cong 0$, then U(L) = 0. Namely, L does not admit a perfectly wrapped uniform resolution.

3.4. An expanded class of diagrams admitting perfectly wrapped uniform resolutions. We will now work to identify a general case in which a perfectly wrapped resolution can be converted into a perfectly wrapped uniform resolution; the motivation is alternating links, which we will handle in Section 4.

Recalling our description of arcs from Subsection 2.1, we will be specific as to the location of these arcs.

Definition 33. Fix a trivial circle W in D_u . By the Jordan curve theorem, $A \setminus W$ consists of two connected components. We say a crossing resolution, or equivalently, an arc, a is W-exterior if it is in the same connected component as the *, and W-interior otherwise. Additionally, we refer to a as W-adjacent-exterior (resp. W-adjacent-interior) if a is W-exterior (resp. W-interior) and abuts W.

A different set of desirable trivial circles can now be defined.

Definition 34. We say a trivial circle W is of type (0, 1) if every W-adjacent-interior arc is red, and every W-adjacent-exterior arc is blue. Similarly, W is of type (1, 0) if every W-adjacent-interior arc is blue, and every W-adjacent-exterior arc is red.

Definition 35. A resolution D_u of a diagram D for an annular link L is almost uniform if every trivial circle in D_u is of type (0,1) or type (1,0).

Note that an almost uniform trivial circle is uniform if and only if it has no adjacent-interior arcs or no adjacent-exterior arcs.

Similar to Definition 33, we want a notion of trivial circles on the interior and exterior of W.

Definition 36. Let W be a trivial circle in a resolution D_u . We say a trivial circle W' is an *exterior trivial* circle of W if W' is contained in the same connected component of $A \setminus W$ as the *. Otherwise, W' is an *interior trivial circle* of W.

The matter of finding a perfectly wrapped uniform resolution from a perfectly wrapped almost uniform resolution comes down to analyzing interior trivial circles. To make these notions precise, we establish an intuitive idea of depth.

Definition 37. Let D_u be a resolution of a diagram D, and let $W \subset D_u$ be a trivial circle. Let a be a point just outside W. The *depth* of W in D_u , denoted by Dep(W), is the minimal geometric intersection of a curve γ connecting a to ∞ and the set of trivial circles in D_u .

In other words, the depth of a trivial circle W is the number of trivial circles containing W, not counting W.

Finally Figure 6 illustrates the difference between two types of nugatory crossings (for diagrams in the plane) in the presence of the marked point *. The terms *removable nugatory crossing* and *essential nugatory crossing* come from Page 10 of [BKS23].

Theorem 38. Suppose that L admits a diagram D without removable nugatory crossings, where D has a perfectly wrapped almost uniform resolution D_u . Then D admits a perfectly wrapped uniform resolution D_v , and in particular, L satisfies Conjecture 1.

Proof. Let $W_1, ..., W_k$ denote the set of trivial circles in D_u such that $Dep(W_i) = 0$. To construct the resolution D_v , we will build a series of perfectly wrapped resolutions $D_{u_1}, ..., D_{u_k}$, each of which is progressively more uniform.

We start with $W_1 \subseteq D_u$. Suppose that W_1 is type (0,1). Let D_{u_1} be the resolution where every W_1 exterior arc is unchanged and every W_1 -interior red arc is switched to a blue arc. The resulting resolution is



FIGURE 6. A removable nugatory crossing can be eliminated via Reidemeister moves representing an isotopy supported within the solid torus.



FIGURE 7. p' is a crossing resolution in D_{u_1} abutting X alone.

similar to D_u , except that W_1 and each of its interior trivial circles have been replaced by a collection of type 1 circles; since the * is outside W_1 , these new circles are indeed trivial. Let $\mathbb{W}D_{u_1}$ be the local 'subresolution' consisting of this new collection of type 1 trivial circles, along with all arcs that were W_1 -interior in D_u .

Switching the color of a W_1 -interior arc must induce a cobordism $\mathcal{W} \to \mathcal{W} \sqcup \mathcal{W}$ or $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}$, since these arcs are, by definition, on the interior of a trivial circle. Hence, D_{u_1} is exactly wrapped, as it is related to D_u by cobordisms which preserve the number of nontrivial circles.

We claim that D_{u_1} is in fact perfectly wrapped. As D_u is insulated, every crossing resolution in D_u is of one of the forms below:

- (I-0) If c is a 0-resolution, then switching c to a 1-resolution induces a cobordism of the form $\mathcal{W} \sqcup \mathcal{W} \to \mathcal{W}, \mathcal{V} \sqcup \mathcal{W} \to \mathcal{V}, \mathcal{V} \sqcup \mathcal{V} \to \mathcal{W}$, or $\mathcal{W} \to \mathcal{V} \sqcup \mathcal{V}$.
- (I-1) If c is a 1-resolution, then switching c to a 0-resolution induces a cobordism of the form $\mathcal{W} \leftarrow \mathcal{W} \sqcup \mathcal{W}, \mathcal{V} \leftarrow \mathcal{V} \sqcup \mathcal{W}, \mathcal{W} \leftarrow \mathcal{V} \sqcup \mathcal{V}$, or $\mathcal{V} \sqcup \mathcal{V} \leftarrow \mathcal{W}$.

If this is the case for each crossing resolution in D_{u_1} , then D_{u_1} is insulated, and thus perfectly wrapped by definition. Note that a crossing resolution that was W_1 -exterior in D_u remains the same color in D_{u_1} and still satisfies either (I-0) or (I-1); in particular, the W_1 -adjacent-exterior crossing resolutions are still merge maps. Therefore, it suffices to show that WD_{u_1} is insulated. This will cover the case of all remaining arcs.

Each arc c' in $\mathbb{W}D_{u_1}$ is blue and abuts only trivial circles. If c' abuts two different trivial circles, then switching this resolution induces a cobordism $\mathcal{W} \leftarrow \mathcal{W} \sqcup \mathcal{W}$. Thus, for D_{u_1} to be insulated, it remains to show that a given c' may not abut only one trivial circle.

Suppose for contradiction that there exists a resolution p' in D_{u_1} , corresponding to some W_1 -interior resolution p in D_u , such that p' abuts only one type 1 trivial circle X in $\mathbb{W}D_{u_1}$. This phenomenon is detailed in Figure 7.

For the crossing resolutions q'_i and r'_i in D_{u_1} , let q_i and r_i denote the respective crossing resolutions in D_u . We now have two cases: p is blue, or p is red. Figure 8 pictures the first case.

The arc p is W_1 -interior, and thus switching p cannot induce a cobordism $\mathcal{V} \sqcup \mathcal{V} \leftarrow \mathcal{W}$. Therefore, as D_u is perfectly wrapped, it must be the case that X_1 and X_2 are different trivial circles.



FIGURE 8. p is a blue arc in D_u



FIGURE 9. q_1 and r_i are on opposite sides of a trivial circle, but are colored the same, so this trivial circle can be neither type (0, 1) nor type (1, 0)

- (B-1) If m = 0 and $n \ge 0$, then p is the resolution of a removable nugatory crossing, contradicting our original assumption.
- (B-2) If m > 0 and n = 0, then q_1 must abut the trivial circle X_1 . By almost uniformity of D_u , we must either have (1) q_1 is blue and on the same side of X_1 as p or (2) q_1 is red and on the opposite side of X_1 as p. In either case, we see that q'_1 is blue and on the same side of X_1 as p, contradicting our definition of the q_i arcs.
- (B-3) If m > 0 and n > 0, then we move clockwise around X in D_{u_1} . If q_1 immediately succeeds p in clockwise order, then we observe the same contradiction as in Case (B-2). Suppose otherwise that $r_1, ..., r_i$ succeed p and precede q_1 . If r_i is blue, then we obtain the same contradiction as Case (B-2). Therefore, r_i is red. But then, as D_u is almost uniform, q_1 is red as well. In Figure 9, we observe that this as a contradiction.

It follows p must be red, as depicted in Figure 10. As before, X_1 and X_2 are distinct trivial circles; we think of X_1 as a trivial circle that abuts some q_i or r_i .

We proceed in an similar fashion.

- (R-1) If m = 0 and $n \ge 0$, then p represents the resolution of a removable nugatory crossing.
- (R-2) If m > 0 and n = 0, then q_1 must be red. We see that this conflicts with the fact that D_u is almost uniform, as p and q_1 are on opposite sides of a trivial circle but are both red.



FIGURE 10. p is red

(R-3) If m > 0 and n > 0, then we proceed as in Case (B-3): we consider crossings in a clockwise fashion. If q_1 immediately proceeds p in clockwise order, then the argument of Case (R-2) may be used. Suppose otherwise that $r_1, ..., r_j$ succeed p and precede q_1 . If $r_1, ..., r_j$ are all blue arcs, then q_1 must be red (otherwise q_1 and p are on the same side of X_1 , but oppositely colored). However, this means that q_1 and r_j are on the same side of X_1 , but are oppositely colored.

Hence, some r_l must be red; we may let r_l be the red arc in D_u which is most advanced in clockwise order out of $r_1, ..., r_j$. Then $r_l, ..., r_j$ are blue arcs; the contradiction follows as in Case (R-2) once again.

We conclude that D_{u_1} is perfectly wrapped. In summary, our process has resulted in a perfectly wrapped resolution, identical to D_u aside from the insertion of $\mathbb{W}D_{u_1}$ in place of W_1 and its interior trivial circles.

Next, consider W_2 . W_2 is a trivial circle in D_u with $\text{Dep}(W_2) = 0$. The algorithm we performed in D_u has been conducted away from W_2 , so W_2 corresponds to a trivial circle in D_{u_1} (which we will also call W_2); furthermore, this trivial circle is not a circle in $\mathbb{W}D_{u_1}$ by the same reasoning. Therefore, W_2 is either type (0, 1) or type (1, 0).

If W_2 is type (0, 1), we repeat our previous process. If W_2 is type (1, 0), we can use a completely analogous argument (leave W_2 -exterior arcs the same and switch W_2 -interior blue arcs to red arcs) to replace W_2 and its interior trivial circles by a collection of type 0 trivial circles. We obtain a perfectly wrapped diagram D_{u_2} with a replacement for W_2 .

We now repeat this procedure. With only finitely many trivial circles to replace, it terminates in a perfectly wrapped resolution D_v , where every trivial circle is contained in some collection of type 1 or type 0 trivial circles. Thus, D_v is perfectly wrapped and uniform, as desired.

The following corollary is implicit in the previous argument, but we state and prove it for emphasis.

Corollary 39. Suppose that L admits a diagram D without removable nugatory crossings, along with a perfectly wrapped resolution D_u . For depth 0 trivial circles in $W_1, ..., W_n$, suppose that there exist a collection of disks $U_1, ..., U_n$ in the annulus such that

- (1) U_i contains W_i ,
- (2) $U_i \cap U_j = \emptyset$ for $i \neq j$, and
- (3) Within each U_i , D_u is either almost uniform or uniform.

Then there exists a perfectly wrapped uniform resolution of D.

Proof. Condition (1) implies that every trivial circle in D_u is contained in some disk. Moreover, if some W_i abuts some W_j , they must have the same color of exterior arcs by condition (3). Finally, by condition (2), we can perform the algorithm of Theorem 38 within every U_i for which D_u is almost uniform. The resulting resolution is perfectly wrapped and uniform.

4. Classes of links admitting perfectly wrapped uniform resolutions

We may now investigate certain links which adhere to the conditions of Theorem 30, and thereby satisfy Conjecture 1.



FIGURE 11. d_i is red and exterior



FIGURE 12. d_i is blue and interior

4.1. Alternating links. As mentioned before, Subsection 3.4, and specifically Definition 35, was motivated by alternating links. With Lemma 40, we see how type (0, 1) and (1, 0) circles arise naturally as a corollary of alternating phenomena.

Lemma 40. Let D be an alternating diagram for an annular link L. Let D_u be a resolution of D. Then any trivial circle W in D_u is either of type (0,1) or type (1,0).

Proof. W will have a collection $d_1, ..., d_n$ of arcs around it, which we may label in clockwise order (as in d_{i+1} is clockwise from d_i around W).

We now look at any d_j abutting W, and check that d_{j+1} is either of identical color to d_j if it is on the same side of W as d_j , or is of opposite color to d_j if it is on the opposite side of W as d_j . Suppose that if d_j is exterior, it is red, and if d_j is interior, it is blue; other cases follow completely analogously. The alternating condition then forces almost uniformity, as we see in Figures 11 and 12. This concludes the proof.

Definition 41 (Page 10, [BKS23]). We say a diagram *D* for an annular link *L* is a *reduced alternating diagram* if *D* is alternating and has no removable nugatory crossings.

Corollary 42. Let L be an alternating link. Then L satisfies Conjecture 1.

Proof. In S^3 , every nugatory crossing is removable, and it is a classical result that every alternating link in S^3 admits a reduced alternating diagram. From the S^3 case, we can conclude that an alternating annular link must admit a reduced alternating annular diagram.

We take D to be such a diagram for L. By Lemma 22, there is a perfectly wrapped resolution D_u of D, which by Lemma 40 is almost uniform. Then an application of Theorem 38 to the diagram D gives a perfectly wrapped uniform resolution. Theorem 30 concludes the proof.

4.2. Other classes and modifications of links. Beyond alternating links, there are immediate candidates for the existence of perfectly wrapped uniform resolutions which may exhibit arbitrarily non-alternating behavior. In particular, we are motivated by plumbed links, arborescent links, and other such classes; these display some of the phenomena we discuss below.

Corollary 43. Suppose that T is a tangle which is braided outside of disjoint disks $U_1, ..., U_n$, each of which contain a positively or negatively oriented "chain" (or 2-braid) and no other crossings. Then the annular closure of T, \hat{T} , admits a perfectly wrapped uniform resolution.



FIGURE 13. A partial resolution of \hat{T} outside of $\cup_i U_i$



FIGURE 14. The braidlike resolution of a negative chain



FIGURE 15. The chainlike resolution of a negative chain

Proof. To construct a perfectly wrapped uniform resolution D_u of \hat{T} , our first step is to take the braidlike resolution (or, in other words, the oriented resolution of \hat{T} with respect to the braidlike orientation) of \hat{T} outside of $\cup_i U_i$. The corresponding partial resolution is displayed in Figure 13.

For our complete resolution of T, we work by cases on the orientation of the chain C_i in U_i . If C_i is in line with the braid, then we can consider it as part of the braid, and we take the braidlike resolution of C_i as shown in Figure 14. If C_i is not in line with the braid, then we resolve chainlike, as in Figure 15.

The resulting complete resolution D_u is exactly wrapped by construction. Moreover, every trivial circle in D_u arises as a result of some chainlike resolution since \hat{T} is resolved braidlike outside of the U_i disks. It follows by inspection that D_u is perfectly wrapped and uniform.

Corollary 44. Consider an (n, n)-tangle T_k of the type displayed in Figure 16, where the unlink component encircles any k strands on the trivial n-braid in the pictured manner.

Let L be the link obtained by vertically stacking $T_{k_1}, ..., T_{k_m}$ along with n-braids $B_1, ..., B_l$ in any order, and taking the annular closure. Then L admits at least 2^m perfectly wrapped uniform resolutions.

Proof. Resolve each B_i braidlike, and take any one of the resolutions of each T_{k_j} displayed in Figures 17 and 18.

We see that the resulting exactly wrapped resolution D_u is perfectly wrapped and uniform, as every trivial circle in D_u is of the form displayed in either Figure 17 or Figure 18.



FIGURE 16. The (n, n)-tangle T_k



FIGURE 17. The 0-type perfectly wrapped uniform resolution



FIGURE 18. The 1-type perfectly wrapped uniform resolution



FIGURE 19. Cabling resolution applied to a 3-cabling

In the spirit of the previous class, we can examine the operation of cabling in the setting of perfectly wrapped uniform resolutions.

Corollary 45. Let D be a diagram for an annular link L which admits a perfectly wrapped uniform resolution D_u . Let D^n be the blackboard-framed n-cabling of D, representing an annular link L^n . Then D^n admits a perfectly wrapped uniform resolution D_v^n .

Proof. At any given crossing c in D, we see a grid pattern of crossings in D^n . We then resolve each crossing in this grid exactly as we did for c. Figure 19 demonstrates this algorithm with a 0-resolution of c in the case n = 3.



FIGURE 20. A linked loop, or earring



FIGURE 21. Two resolutions of a linked loop

 D_v^n is the blackboard-framed *n*-cabling of D_u . If a trivial circle in D_u is abutted by all 0-resolutions, then each of the *n* corresponding trivial circles in D_v^n are abutted by all 0-resolutions. The same holds for 1-resolutions. Hence, D_u is uniform.

Moreover,

$$\operatorname{wrap}(D_v^n) = n \cdot \operatorname{wrap}(D_u) = n \cdot \operatorname{wrap}(L) = \operatorname{wrap}(L^n).$$

Hence, D_v^n is exactly wrapped.

Finally, if switching a crossing in D_u enacts some type of cobordism, then switching any crossing in the corresponding grid in D_v^n either enacts that same type of cobordism or results in a merge of two adjacent circles. Therefore, as D_u is insulated, D_v^n is insulated as well.

Corollary 46. Let L be a link with a diagram D that admits a perfectly wrapped uniform resolution D_u . Suppose that D' is a diagram (for a link L'), which is identical to D outside of disjoint disks $U_1, ..., U_n$. Each U_i contains a single linked loop, or "earring," as shown in Figure 20, and no other crossings. Then D' admits a perfectly wrapped uniform resolution.

Proof. We construct a perfectly wrapped uniform resolution D'_v by first considering a partial resolution of D'. Outside of $\bigcup_i U_i$, we resolve each crossing in D' as in D_u . Now within a given U_i , consider two resolutions of the linked loop, displayed in Figure 21. We note that for each linked loop O_i in U_i , there is a choice of whether to produce a type 0 or type 1 trivial circle.

In our partial resolution of D', the arc passing through a loop O_i may be, thus far, part of some nontrivial circle. In this case, we can use either choice of pictured resolution for O_i . Alternatively, if the arc passing through O_i is thus far part of some trivial circle, this circle will abut only 0-resolutions or only 1-resolutions in the partial resolution. Hence, we resolve O_i in the manner that allows for compatible type.

Our resulting complete resolution D'_v is functionally identical to D_u , but within each U_i we insert a trivial circle of the kind in Figure 21. These trivial circles will be of compatible type with the circles they abut. Hence, D'_v is uniform. Moreover, we see immediately that D'_v is perfectly wrapped within each U_i . Given that D'_v is identical to D_u outside of $\bigcup_i U_i$, it follows that D'_v is perfectly wrapped. \Box

References

- [Akh23] Rostislav Akhmechet. Equivariant annular Khovanov homology. J. Knot Theory Ramifications, 32(2):Paper No. 2350002, 31, 2023.
- [APS04] Marta M. Asaeda, Józef H. Przytycki, and Adam S. Sikora. Categorification of the Kauffman bracket skein module of *I*-bundles over surfaces. *Algebr. Geom. Topol.*, 4:1177–1210, 2004.

- [BKS23] Hans U. Boden, Homayun Karimi, and Adam S. Sikora. Adequate links in thickened surfaces and the generalized Tait conjectures. Algebr. Geom. Topol., 23(5):2271–2308, 2023.
- [BN05] Dror Bar-Natan. Khovanov's homology for tangles and cobordisms. Geom. Topol., 9:1443-1499, 2005.
- [BPW19] Anna Beliakova, Krzysztof K. Putyra, and Stephan M. Wehrli. Quantum link homology via trace functor I. Invent. Math., 215(2):383–492, 2019.
- [GLW18] J. Elisenda Grigsby, Anthony M. Licata, and Stephan M. Wehrli. Annular Khovanov homology and knotted Schur-Weyl representations. *Compos. Math.*, 154(3):459–502, 2018.
- [GN14] J. Elisenda Grigsby and Yi Ni. Sutured Khovanov homology distinguishes braids from other tangles. Math. Res. Lett., 21(6):1263–1275, 2014.
- [HP95] Jim Hoste and Józef H. Przytycki. The (2,∞)-skein module of Whitehead manifolds. J. Knot Theory Ramifications, 4(3):411–427, 1995.
- [Kho00] Mikhail Khovanov. A categorification of the Jones polynomial. Duke Math. J., 101(3):359–426, 2000.
- [Kim21] Juhyun Kim. Annular Links with Sl₂-Irreducible Annular Khovanov Homology. ProQuest LLC, Ann Arbor, MI, 2021. Thesis (Ph.D.)–California Institute of Technology.
- [Lee05] Eun Soo Lee. An endomorphism of the Khovanov invariant. Adv. Math., 197(2):554–586, 2005.
- [Mar23] Gage Martin. Annular Khovanov homology and meridional disks. J. Knot Theory Ramifications, 32(2):Paper No. 2250088, 14, 2023.
- [Prz91] Józef H. Przytycki. Skein modules of 3-manifolds. Bull. Polish Acad. Sci. Math., 39(1-2):91–100, 1991.
- [Ras10] Jacob Rasmussen. Khovanov homology and the slice genus. Invent. Math., 182(2):419-447, 2010.
- [Rob13] Lawrence P. Roberts. On knot Floer homology in double branched covers. Geom. Topol., 17(1):413-467, 2013.
- [Xie21] Yi Xie. Instantons and annular Khovanov homology. Adv. Math., 388:Paper No. 107864, 51, 2021.

DEPARTMENT OF MATHEMATICS, UC DAVIS, ONE SHIELDS AVE., DAVIS, CA 95616-8633, U.S.A. *Email address:* bkdaniels@ucdavis.edu

DEPARTMENT OF MATHEMATICS, UC DAVIS, ONE SHIELDS AVE., DAVIS, CA 95616-8633, U.S.A. *Email address:* mlzhang@ucdavis.edu