Major-minor mean field games: common noise helps

Fr[a](#page-0-0)nçois Delarue^a and Chenchen Mou^{[b](#page-0-1)}

^a*Université Côte d'Azur, CNRS, Laboratoire J.A.Dieudonné, Parc Valrose, France-06108 NICE Cedex 2, francois.delarue@univ-cotedazur.fr* ^b*City University of Hong Kong, Department of Mathematics, China-Hong Kong SAR, chencmou@cityu.edu.hk*

Abstract. The objective of this work is to study the existence, uniqueness, and stability of equilibria in mean field games involving a major player and a continuum of minor players over finite intervals of arbitrary length. Following earlier articles addressing similar questions in the context of classical mean field games, the cost functions for the minor players are assumed to satisfy the Lasry-Lions monotonicity condition. In this contribution, we demonstrate that if, in addition to the monotonicity condition, the intensity of the (Brownian) noise driving the major player is sufficiently high, then—under further mild regularity assumptions on the coefficients—existence, uniqueness, and stability of equilibria are guaranteed. A key challenge is to show that the threshold (beyond which the noise intensity must be taken) can be chosen independently of the length of the time interval over which the game is defined. Building on the stability properties thus established, we further show that the associated system of master equations admits a unique classical solution. To the best of our knowledge, this is the first result of its kind for major-minor mean field games defined over intervals of arbitrary length.

[MSC2020 subject classifications:](https://mathscinet.ams.org/mathscinet/msc/msc2020.html) Primary 49N80, 91A16; secondary 35R60, 60H10, 60H15 *Keywords:* Major-minor mean field games, Master equation, BMO martingale

1. Introduction

Mean field games were introduced simultaneously by Lasry and Lions [\[44](#page-87-0)[–46\]](#page-87-1), and by Huang, Caines, and Malhamé [\[35](#page-86-0), [36\]](#page-86-1), with the aim of providing an asymptotic description of differential games for large populations of weakly interacting agents. While the theory for the most popular form of mean field games has reached a certain level of maturity (as evidenced by Lions' lectures at the Collège de France, [\[50\]](#page-87-2), as well as –among other references– the books by Cardaliaguet et al. [\[14\]](#page-86-2), Carmona and Delarue [\[19,](#page-86-3) [20\]](#page-86-4) and Gomes et al. [\[33\]](#page-86-5), along with the lecture notes by Cardaliaguet and Porretta [\[1](#page-85-0)] and Delarue [\[27\]](#page-86-6), and the surveys by Caines et al. [\[10\]](#page-86-7) and Cardaliaguet and Delarue [\[15](#page-86-8)]), many questions remain open. These arise either in regimes where the assumptions are too weak to be covered by the existing literature or in situations where the very structure of the game is significantly more complex and thus escapes known results.

In this regard, one example currently generating considerable interest is the case of mean field games subject to common noise (see, for some of the earlier works on the subject, the books [\[14,](#page-86-2) [20](#page-86-4)] and the papers by Bertucci, Lasry and Lions [\[6\]](#page-86-9), Carmona, Delarue and Lacker [\[21](#page-86-10)], Carmona, Fouque and Sun [\[22\]](#page-86-11) and Lacker and Zariphopoulou [\[43](#page-87-3)]). By common noise, we mean a noise that impacts all the agents in the population in a common manner, thereby leading to a randomization of the population state. This contrasts with the traditional version of mean field games, where the population state—obtained by taking the limit over a population of asymptotically identically distributed and independent players—is deterministic. In the presence of common noise, the mathematical challenges are numerous: they may involve adapting existing concepts (see for instance the works by Bertucci [\[5\]](#page-86-12), Bertucci, Lasry and Lions [\[7\]](#page-86-13), Bertucci and Maynard [\[8\]](#page-86-14), Cardaliaguet and Souganidis [\[17\]](#page-86-15), Gangbo et al. [\[32\]](#page-86-16), and Mou and Zhang [\[51\]](#page-87-4) for works in connection with the master equation, now a cornerstone of the field; see also Cardaliaguet and Souganidis [\[18\]](#page-86-17) for the analysis of the mean field game system itself) or exploring additional effects induced solely by the noise (see Bayraktar et al. [\[2](#page-86-18)] and Foguen Tchuendom [\[31\]](#page-86-19)).

In our work, we consider a more "advanced" version of mean field games with common noise: games involving a major player and minor players. In the regime we study, the major player is subject to a Brownian diffusion, thereby forcing the entire population of minor players to become random. However, mean field games with a major player introduce additional difficulties not present in games involving only minor players (even with common noise). In games with a 2

major player, the population of minor agents competes with the major player. In addition to the standard mean field game played among the minor agents once the major player has determined her strategy, there is also an equilibrium condition stemming from the ability of the major player to react to the strategies of the minor players. This entanglement makes the formalization of major-minor games quite subtle. For this reason, different concepts of solution have been proposed in the literature, each reflecting a particular form of equilibrium among the various actors.

The first article introducing the concept of a major player is due to Huang [\[34\]](#page-86-20). Initially formulated in infinite horizon and for a linear-quadratic structure, the paper was later adapted by Nourian and Huang [\[52](#page-87-5)] to the finite horizon case, and by Nourian and Caines [\[53\]](#page-87-6) to the nonlinear case. In these works, the complexity inherent to the presence of two types of players is resolved by imposing a sequential structure on the equilibria, where the major player's response is computed for a fixed state of the population. An alternative sequential approach is proposed in the paper of Bensoussan, Chau, and Yam [\[3\]](#page-86-21), where the major player anticipates the rational behaviour of the minor players. For a review of other types of equilibria considered in the literature, we refer to the works of Cardaliaguet, Cirant, and Porretta [\[12](#page-86-22)] and Bergault et al. [\[4](#page-86-23)].

In the current paper, we focus on full-fledged Nash equilibria, which implicitly require updating the state of the minor player population when changing the major player's strategy. In this regard, two regimes are possible, depending on whether the major player employs open-loop or closed-loop strategies and leading to possibly distinct solutions. The open-loop case has been studied by Carmona and Zhu [\[24\]](#page-86-24), as well as Carmona and Wang [\[23\]](#page-86-25) and Huang and Tang [\[37\]](#page-86-26), primarily using the stochastic Pontryagin principle. The closed-loop case has led to several articles by Lasry and Lions [\[47](#page-87-7)], and Cardaliaguet, Cirant, and Porretta [\[12,](#page-86-22) [13\]](#page-86-27). This is precisely the framework we consider in our contribution. In this context, the primary work to which we compare our results is that of Cardaliaguet, Cirant, and Porretta [\[13\]](#page-86-27), where the authors address the existence of classical solutions to the Nash system over short time horizons. This system is expressed as two partial differential equations (called master equations) posed on the space of probability measures.

In comparison, the objective of our work is to study stability of the Nash equilibria and then existence of classical solutions to the Nash system, for games set over finite but arbitrarily long time horizons. The main idea to prevent the emergence, over intervals of arbitrary length, of multiple Nash equilibria or, equivalently, of singular solutions to the Nash system, is twofold: (i) we assume that the costs to the minor players satisfy the Lasry-Lions monotonicity condition; this provides stability for the system governing the minor players when the major player is fixed—this is well understood in the case of common noise models (see, for example $[14]$)–; (ii) we assume that the major player's dynamics is subjected to a non-degenerate Brownian noise, which is referred to as a common noise in the rest of the text; the diffusion structure induces a regularizing effect that is expected to contribute to the uniqueness and stability of equilibria. In fact, we show that this regularizing effect indeed exists if the intensity of the common noise exceeds a certain threshold. Furthermore, we provide conditions under which this threshold is independent of the duration T of the game. A concise version of this result, requiring detailed mild regularity assumptions outlined in Section [2,](#page-2-0) is presented in Theorem [2.14.](#page-10-0) As a corollary, Theorem [2.16](#page-16-0) establishes the existence and uniqueness of classical solutions to the corresponding Nash system. All the results are stated for minor players evolving in the torus (of dimension d). This simplification is already present in [\[14\]](#page-86-2). Here (and similarly to what occurs in the long-term study of classical mean field games, see Cardaliaguet et al. [\[11\]](#page-86-28) and Cardaliaguet and Porretta [\[16](#page-86-29)]), the periodic structure allows for easier control of certain key quantities, independently of the length T .

The proof structure relies on a natural representation of the major and minor players' values in the form of two coupled forward-backward stochastic equations. The forward-backward system representing the minor players can be interpreted as a mean field game system embedded in the random environment induced by the major player. It is made of a forward Kolmogorov equation with random coefficients and a backward stochastic Hamilton-Jacobi-Bellman equation, accounting respectively for the evolution and the value of the minor player population. We thus call this system a Forward-Backward Stochastic Partial Differential Equation (FBSPDE). As for the major player, the forward-backward equation is a standard finite-dimensional Forward-Backward Stochastic Differential Equation (FBSDE) with random coefficients depending on the state of the minor player population. Stochastic control theory says that the major player's strategy is represented by the martingale component of the corresponding forward-backward equation. Our approach then combines stability methods for each of the two systems. On the one hand, the Lasry-Lions condition is crucial to apply to the FBSPDE the techniques introduced in [\[14](#page-86-2)] for solving mean field game systems with common noise. On the other hand, we use tools from stochastic analysis and the theory of backward stochastic differential equations (BSDEs) to address the FBSDE.

In fact, the key challenge is to study the stability properties of the FBSDE, which has here the peculiarity to be strongly coupled with the FBSPDE. As is often the case in stochastic analysis, Girsanov's transformation simplifies the forward equation of the FBSDE, reducing it to a standard Brownian motion and thus decoupling it from the FBSPDE system for the minor players. Quantitatively, a major step is to estimate the impact of this Girsanov transformation on the overall stability properties of the system comprising both the FBSDE and the FBSPDE. For this, we make systematic use of Bounded Mean Oscillation (BMO) martingale theory introduced in [\[40](#page-86-30)]. In short, these BMO bounds decrease as the intensity of the common noise increases, which explains the condition on the intensity of the noise.

Major Minor MFGs 3

This quantitative analysis ultimately yields *a priori* bounds on the spatial gradient of the game's value functions. Once these bounds are established, existence and uniqueness of equilibria are obtained by iterating contraction principles applied in small time intervals. Remarkably, the resulting solutions are strong, meaning that they are adapted to the filtration generated by the common noise. The resolution of the master equations is also achieved iteratively by demonstrating that, over short time intervals, the system of forward-backward equations mentioned earlier serves as the characteristic system for the master equations. The derivation of classical solutions from this characteristic system follows a linearization procedure similar to that developed by [\[14\]](#page-86-2). In this regard, our approach differs (even in short time) from the splitting method proposed in the work [\[13\]](#page-86-27).

The paper is organized as follows. In Section [2,](#page-2-0) we introduce the problem and the equilibrium representation in the form of two coupled forward-backward equations, the aforementioned FBSDE and the FBSPDE. We also state the main results. In Section [3,](#page-18-0) we derive a priori estimates for solutions to the FBSDE and FBSPDE, including the essential BMO bounds necessary for measure changes. Section [4](#page-30-0) is devoted to obtaining a priori bounds on the gradients of the value functions by linearizing the forward-backward equations. Section [5](#page-50-0) contains several short-time results required to conclude, though their novelty is of lesser significance.

2. Set-up and main results

2.1. A brief presentation of the model

We introduce the Major/Minor MFG through a probabilistic formulation that comprises two filtered probability spaces $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ and $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ equipped with two Brownian motions $(B_t^0)_{0 \le t \le T}$ and $(B_t)_{0 \le t \le T}$ with values in \mathbb{R}^d . In brief, the space labelled with a '0' carries the 'common noise' underpinning the major player and the other spaces carries the 'idiosyncratic noise' underpinning one tagged minor player.

We then consider random variables constructed on Ω^0 , Ω or $\Omega^0 \times \Omega$. When dealing with a random variable X defined on the product space $\Omega^0 \times \Omega$, the expectation of X (if well-defined, and possibly with values in \mathbb{R}^n for some $n \ge 1$) is denoted $\mathbb{E}^0 \mathbb{E}(X)$ and the conditional law of X given \mathcal{F}^0 is denoted $\mathcal{L}^0(X)$ (i.e., for $\omega^0 \in \Omega^0$, $\mathcal{L}^0(X)(\omega^0)$ is the law of the random variable $\omega \in \Omega \mapsto X(\omega^0, \omega)$).

Given an \mathbb{F}^0 -adapted continuous random flow of measures $\mu = (\mu_t)_{0 \le t \le T}$ with values in $\mathcal{P}(\mathbb{T}^d)$, the game is equipped with two kinds of players:

1. **Major player**: For a (fixed) \mathcal{F}_0^0 -measurable initial condition $X_0^0 : \Omega^0 \to \mathbb{R}^d$ and for a (square-integrable) \mathbb{F}^0 progressively measurable control process $\alpha^0 = (\alpha^0_t)_{0 \le t \le T}$ with values in \mathbb{R}^d , the major player has dynamics

$$
dX_t^0 = \alpha_t^0 dt + \sigma_0 dB_t^0, \quad t \in [0, T],
$$

where α^0 is a velocity field from \mathbb{R}^d into itself and σ_0 is a (strictly) positive diffusion coefficient, and has cost functional

$$
J^{0}(\alpha^{0}; \mu) = \mathbb{E}^{0}\bigg[g^{0}(X_{T}^{0}, \mu_{T}) + \int_{0}^{T}\bigg(f_{t}^{0}(X_{t}^{0}, \mu_{t}) + L^{0}(X_{t}^{0}, \alpha_{t}^{0})\bigg)\mathrm{d}t\bigg].
$$

Here, f^0 (resp. g^0) is a cost coefficient from $[0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ (resp. $\mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$) to $\mathbb R$ and L^0 is a Lagrangian from $\mathbb{R}^d \times \mathbb{R}^d$ to \mathbb{R} .

2. **Minor player**: For a (fixed) \mathcal{F}_0 -measurable initial condition $X_0: \Omega \to \mathbb{T}^d$, and for a (square-integrable) $\mathbb{F}^0 \otimes \mathbb{F}$ progressively measurable control process $\alpha = (\alpha_t)_{0 \le t \le T}$ with values in \mathbb{R}^d , the minor player has dynamics

$$
dX_t = \alpha_t dt + dB_t, \quad t \in [0, T],
$$

and has cost functional

$$
J(\boldsymbol{\alpha};\boldsymbol{\alpha}^0,\boldsymbol{\mu}) = \mathbb{E}\bigg[g(X_T^0,X_T,\mu_T) + \int_0^T \Big(f_t(X_t^0,X_t,\mu_t) + L(X_t,\alpha_t)\Big)dt\bigg].
$$

Here, f (resp. g) is a cost coefficients from $[0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$ (resp. $\mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$) to $\mathbb R$ and L is a Lagrangian from $\mathbb{T}^d \times \mathbb{R}^d$ to \mathbb{R} .

As far as the major player is concerned, we look for strategies in Markov feedback form $\alpha^0 : [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$, namely α^0 reads in the form $\alpha_t^0 = \alpha^0(t, X_t^0, \mu_t)$, $t \in [0, T]$. This implicitly puts some constraints on the state equation for $(X_t^0)_{0 \le t \le T}$, which are addressed in Subsection [2.4.](#page-6-0) As for the minor player, we also look for strategies in Markov

feedback form $\alpha : [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$, namely $\alpha_t = \alpha(t, X_t^0, X_t, \mu_t)$, $t \in [0, T]$, which also puts some constraints on the state equation for $(X_t)_{0 \le t \le T}$. The goal is to find a Nash equilibrium in the sense of [\[12,](#page-86-22) [13,](#page-86-27) [47](#page-87-7)]. A clean definition (though in a weaker setting) is given in Subsection [2.4](#page-6-0) below. In a nutshell, an equilibrium has the following three features: (i) the process μ solves the Fokker-Planck equation associated with the random velocity field $x \mapsto \alpha(t, X_t^0, x, \mu_t)$; (ii) with μ as in (i), the strategy α minimizes the cost J when α^0 and μ are fixed; (iii) the strategy α^0 minimizes the cost J^0 with the peculiarity that μ therein depends on α^0 itself through the state X^0 of the major player. The latter point makes Major/Minor MFGs more difficult to solve than MFGs with common noise.

2.2. General notations

We first state with several notations that are necessary in our analysis.

Functional and distributional spaces on \mathbb{T}^d . For an index $s \in \mathbb{N}$, we call $\mathcal{C}^s(\mathbb{T}^d)$ the space of functions from \mathbb{T}^d into R that have $\lfloor s \rfloor$ continuous derivatives. We equip the space $\mathcal{C}^s(\mathbb{T}^d)$ with the norm

$$
||f||_s = \sup_{k=0,\cdots, \lfloor s \rfloor} \sup_{x \in \mathbb{T}^d} |\nabla^k f(x)|.
$$

In particular, $||f||_0$ coincides with the L^{∞} norm, sometimes denoted $||f||_{L^{\infty}}$, of f. When $s > 0$ and $s \notin \mathbb{N}$, we call $\mathcal{C}^{s}(\mathbb{T}^{d})$ the space of functions from \mathbb{T}^d into $\mathbb R$ that have $\lfloor s \rfloor$ derivatives and such that the derivative of order $\lfloor s \rfloor$ is $s - \lfloor s \rfloor$ Hölder continuous. We equip the space $\mathcal{C}^s(\mathbb{T}^d)$ with the norm

$$
||f||_{s} = \sup_{k=0,\dots, \lfloor s \rfloor} \sup_{x \in \mathbb{T}^d} |\nabla^k f(x)| + \sup_{x,x' \in \mathbb{T}^d: x \neq x'} \frac{|\nabla^{\lfloor s \rfloor} f(x) - \nabla^{\lfloor s \rfloor} f(x')|}{|x - x'|^{s - \lfloor s \rfloor}}.
$$

We also make use of the (topological) dual space of $C^s(\mathbb{T}^d)$, which we denote $C^{-s}(\mathbb{T}^d)$ and which we equip with the dual norm:

$$
||q||_{-s} = \sup_{||f||_s \le 1} (f, q)_{s, -s},
$$

where $(\cdot, \cdot)_{s,-s}$ is here used to denote the duality product between elements of $C^{-s}(\mathbb{T}^d)$ and $C^{s}(\mathbb{T}^d)$. Quite often the precise index s in the duality $(f, q)_{s,-s}$ between a function $f \in C^{s}(\mathbb{T}^{d})$ and a distribution $q \in C^{-s}(\mathbb{T}^{d})$ is implicitly understood and thus omitted, and the duality product between f and q is merely written (f, q) .

We next introduce the Sobolev space $\mathcal{H}^s(\mathbb{T}^d)$. We denote $(e_k)_{k\in\mathbb{Z}^d}$ the standard (complex valued) Fourier basis of \mathbb{T}^d and $(\cdot, \cdot)_{0,2}$ the standard inner product in $L^2(\mathbb{T}^d)$ For $s > 0$, we call $\mathcal{H}^s(\mathbb{T}^d)$ the space of functions $f \in L^2(\mathbb{T}^d)$ such that $||f||_{s,2}^2 := \sum_{k \in \mathbb{Z}^d} (1+|k|^2)^s |(f, e_k)_{0,2}|^2 < \infty$. The $\mathcal{H}^s(\mathbb{T}^d)$ -norm is $||\cdot||_{s,2}$. See for instance [\[28](#page-86-31)].

Distances on the space $\mathcal{P}(\mathbb{T}^d)$. For two probability measures $\mu, \nu \in \mathcal{P}(\mathbb{T}^d)$, $\|\mu - \nu\|_{-1}$ coincides with $\mathbb{W}_1(\mu, \nu)$, where we recall that, for any $p \geq 1$,

$$
\mathbb{W}_p(\mu,\nu) := \inf_{\pi} \left[\int_{\mathbb{T}^d \times \mathbb{T}^d} |x - y|^p \mathrm{d}\pi(x,y) \right]^{1/p},
$$

with the infimum being taken over 'couplings' π , i.e. over elements $\pi \in \mathcal{P}(\mathbb{T}^d \times \mathbb{T}^d)$, whose images by the first and second projection mappings from $\mathbb{T}^d \times \mathbb{T}^d$ into \mathbb{T}^d are respectively μ and ν .

Similarly, $\|\mu - \nu\|_{-0}$ coincides with $d_{TV}(\mu, \nu)$, where we recall that

$$
d_{\text{TV}}(\mu, \nu) := \sup_{\|\ell\|_{\infty} \le 1} \left| \int_{\mathbb{T}^d} \ell(x) \mathrm{d}\big(\mu - \nu\big)(x) \right|,
$$

with the supremum in the right-hand side being taken over functions $\ell : \mathbb{T}^d \to \mathbb{R}$ that are bounded by 1.

Derivatives on $\mathcal{P}(\mathbb{T}^d)$. Throughout, we use two standard notions of derivatives on $\mathcal{P}(\mathbb{T}^d)$. We refer the reader to [\[19,](#page-86-3) Chapter 5] and [\[14\]](#page-86-2). Briefly, we say that a function $\ell : \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ is continuously differentiable in the flat sense if there exists a jointly continuous function $\delta_{\mu} \ell : \mathcal{P}(\mathbb{T}^d) \times \mathbb{R}^d \to \mathbb{R}$ such that, for any two $\mu, \nu \in \mathcal{P}(\mathbb{T}^d)$,

$$
\ell(\nu) - \ell(\mu) = \int_0^1 \delta_\mu \ell(r\nu + (1-r)\mu, y) d(\nu - \mu)(y).
$$

Because the flat derivative is just defined up to an additive constant, we require (by convention) that

$$
\forall \mu \in \mathcal{P}(\mathbb{T}^d), \quad \int_{\mathbb{T}^d} \delta_\mu \ell(\mu, y) \mathrm{d}\mu(y) = 0. \tag{2.1}
$$

When the function $(\mu, y) \mapsto \delta_{\mu} \ell(\mu, y)$ is differentiable with respect to y, we let

$$
\partial_{\mu}\ell(\mu, y) = \nabla_{y} (\delta_{\mu}\ell)(\mu, y), \quad (\mu, y) \in \mathcal{P}(\mathbb{T}^{d}) \times \mathbb{T}^{d},
$$

which we sometimes refer to as the 'Wasserstein' derivative of ℓ . It is standard to observe that, if $\partial_{\mu}\ell$ is bounded (with respect to μ and y), then the function ℓ is Lipschitz continuous with respect to the 1-Wasserstein distance \mathbb{W}_1 .

Stochastic processes. On a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, we introduce the following spaces. For a Euclidean space $(E, |\cdot|)$ and an exponent $p \in [1, \infty]$, we call $\mathcal{S}^p(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P}; E)$ (or just $\mathcal{S}^p(E)$) the collection of **F**-adapted processes with continuous E-valued trajectories $(S_t)_{0 \leq t \leq T}$ such that

$$
\sup_{0\leq t\leq T}|S_t|\in L^p(\Omega,\mathcal{F},\mathbb{P};\mathbb{R}),
$$

and, for $p \in [1,\infty)$, we call $\mathcal{H}^p(\Omega,\mathcal{F},\mathbb{F},\mathbb{P};E)$ (or just $\mathcal{H}^p(E)$) the collection of \mathbb{F} -progressively-measurable E-valued processes $(H_t)_{0 \leq t \leq T}$ such that

$$
\mathbb{E}\bigg[\bigg(\int_0^T |H_t|^2 \mathrm{d}t\bigg)^{p/2}\bigg] < \infty.
$$

For a **F**-continuous local martingale $M = (M_t)_{0 \le t \le T}$ with values in **R**, we call $\langle M \rangle = (\langle M \rangle_t)_{0 \le t \le T}$ the standard bracket of M and $(\mathscr{E}_t(M)_{0 \leq t \leq T})$ the Doléans-Dade exponential

$$
\mathcal{E}_t(\boldsymbol{M}) := \exp\Big(M_t - \frac{1}{2} \langle M \rangle_t\Big), \quad t \in [0, T].
$$

If in addition, M is uniformly integrable, we let

$$
\|{\boldsymbol{M}}\|^2_{\text{BMO}} := \sup_\tau \bigl\| \mathbb{E}\bigl[|M_T - M_\tau|^2 | \mathcal{F}_\tau \bigr] \bigr\|_{L^\infty(\Omega)},
$$

with the supremum being taken over the collection of **F**-stopping times τ and with $\|\cdot\|_{L^{\infty}(\Omega)}$ here denoting the L^{∞} norm on $(\Omega, \mathcal{F}, \mathbb{P})$. When $M \in \mathcal{S}^2(\mathbb{R})$, the above is the same as

$$
||M||_{\rm BMO}^2 = \sup_{\tau} \left\| \mathbb{E} \left[\langle M \rangle_T - \langle M \rangle_\tau | \mathcal{F}_\tau \right] \right\|_{L^\infty(\Omega)}.
$$

We refer to [\[40](#page-86-30)] for more details on BMO martingales.

2.3. Assumptions on the coefficients

In order to state the assumptions, we introduce a set of four generic conditions, which will be also very useful in the rest of the paper. For any three (strictly) positive reals Λ , Λ and Δ , we let:

Condition $\mathscr{C}^0(\Lambda, \mathfrak{r})$. We say that a function $h^0: (x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto h^0(x_0, \mu) \in \mathbb{R}$ satisfies $\mathscr{C}^0(\Lambda, \mathfrak{r})$ if

(i) h^0 is continuously differentiable with respect to x_0 and μ ;

(ii) for any $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the function $\delta_{\mu}h^0(x_0, \mu, \cdot)$: $y \in \mathbb{T}^d \mapsto \delta_{\mu}h^0(x_0, \mu, y) \in \mathbb{R}$ belongs to $\mathcal{C}^{\mathcal{P}}(\mathbb{T}^d)$;

(iii) the functions $\nabla_{x_0} h^0$ and $\delta_\mu h^0$ satisfy

$$
\forall x_0 \in \mathbb{R}^d, \ \mu \in \mathcal{P}(\mathbb{T}^d), \quad \left| h^0(x_0, \mu) \right| + \left| \nabla_{x_0} h^0(x_0, \mu) \right| \leq \Lambda, \quad \|\delta_{\mu} h^0(x_0, \mu, \cdot)\|_{\mathbb{T}} \leq \Lambda.
$$

Condition $\mathscr{C}(\Lambda, \mathcal{P}, \mathcal{P})$. We say that a function $h : (x_0, x, \mu) \in \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto h(x_0, x, \mu) \in \mathbb{R}$ satisfies $\mathscr{C}(\Lambda, \mathcal{P}, \mathcal{P})$ if

(i) h is continuously differentiable with respect to x_0 , x and μ ;

- (ii) for any $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the function $\nabla_{x_0} h(x_0, \cdot, \mu) : x \mapsto \nabla_{x_0} h(x_0, x, \mu) \in \mathbb{R}^d$ belongs to $\mathcal{C}^{\delta}(\mathbb{T}^d)$;
- (iii) the function $\delta_\mu h(x_0, \cdot, \mu, \cdot)$: $(x, y) \in \mathbb{R}^d \times \mathbb{T}^d \mapsto \delta_\mu h(x_0, \cdot, \mu, \cdot)$ has continuous joint derivatives up to the order $\lfloor \mathfrak{n} \rfloor$ in y and $|\triangle|$ in x;

(iv) the functions $\nabla_{x_0} h$ and $\delta_\mu h$ satisfy

$$
\forall x_0 \in \mathbb{R}^d, \ \mu \in \mathcal{P}(\mathbb{T}^d), \quad \left\|h(x_0, \cdot, \mu)\right\|_{\mathfrak{b}} + \left\|\nabla_{x_0}h(x_0, \cdot, \mu)\right\|_{\mathfrak{b}} \leq \Lambda, \quad \sup_{l=0, \cdots, \lfloor n \rfloor} \sup_{y \in \mathbb{T}^d} \left\|\nabla_y^l \delta_\mu h(x_0, \cdot, \mu, y)\right\|_{\mathfrak{b}} \leq \Lambda;
$$

$$
\forall y, y' \in \mathbb{T}^d, \quad \left\|\nabla_y^{\lfloor n \rfloor} \delta_\mu h(x_0, \cdot, \mu, y') - \nabla_y^{\lfloor n \rfloor} \delta_\mu h(x_0, \cdot, \mu, y)\right\|_{\mathfrak{b}} \leq \Lambda |y' - y|^{n - \lfloor n \rfloor}.
$$

Of course, the very last line above can be removed if π is an integer.

Notice that condition (iv) right above is equivalent to

$$
\forall x_0 \in \mathbb{R}^d, \ \mu \in \mathcal{P}(\mathbb{T}^d), \quad \left\|h(x_0, \cdot, \mu)\right\|_{\mathfrak{b}} + \left\|\nabla_{x_0}h(x_0, \cdot, \mu)\right\|_{\mathfrak{b}} \leq \Lambda, \quad \sup_{k=0, \cdots, \lfloor \mathfrak{b} \rfloor} \sup_{x \in \mathbb{T}^d} \|\nabla_x^k \delta_\mu h(x_0, x, \mu, \cdot)\|_{\mathfrak{b}} \leq \Lambda;
$$

$$
\forall x, x' \in \mathbb{T}^d, \quad \left\|\nabla_x^{\lfloor \mathfrak{b} \rfloor} \delta_\mu h(x_0, x', \mu, \cdot) - \nabla_x^{\lfloor \mathfrak{b} \rfloor} \delta_\mu h(x_0, x, \mu, \cdot)\right\|_{\mathfrak{b}} \leq \Lambda |x' - x|^{\delta - \lfloor \mathfrak{b} \rfloor}.
$$

Condition $\mathscr{D}^0(\Lambda, \mathfrak{n})$. We say that a function $h^0: (x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto h^0(x_0, \mu) \in \mathbb{R}$ satisfies $\mathscr{D}^0(\Lambda, \mathfrak{n})$ if it satisfies the first two conditions in $\mathscr{C}^0(\Lambda,\infty)$ and, for all $l \in \{0, \dots, \lfloor \tau \rfloor\}, x_0, x'_0 \in \mathbb{R}^d, \mu, \mu' \in \mathcal{P}(\mathbb{T}^d)$ and $y', y \in \mathbb{T}^d$,

$$
\begin{aligned} \left| \nabla_{x_0} h^0(x'_0, \mu') - \nabla_{x_0} h^0(x_0, \mu) \right| &\leq \kappa \Big(|x'_0 - x_0| + \mathbb{W}_1(\mu', \mu) \Big), \\ \left| \nabla_y^l \delta_\mu h^0(x'_0, \mu', y') - \nabla_y^l \delta_\mu h^0(x_0, \mu, y) \right| &\leq \kappa \Big(|x'_0 - x_0| + \mathbb{W}_1(\mu', \mu) + |y' - y|^{n - \lfloor n \rfloor} \Big). \end{aligned}
$$

Condition $\mathscr{D}(\Lambda, \mathcal{P}, \mathcal{P})$. We say that a function $h : (x_0, x, \mu) \in \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto h(x_0, x, \mu) \in \mathbb{R}$ satisfies $\mathscr{D}(\Lambda, \mathcal{P}, \mathcal{P})$ it satisfies the first two conditions in $\mathscr{C}(\Lambda,\mathfrak{r},\mathfrak{a})$ and, for all $l \in \{0, \dots, \lfloor \mathfrak{r}_k \rfloor\}$, $x_0, x'_0 \in \mathbb{R}^d$, $\mu, \mu' \in \mathcal{P}(\mathbb{T}^d)$ and $y, y' \in \mathbb{T}^d$,

$$
\begin{aligned}\n\left\| \nabla_{x_0} h(x'_0, \cdot, \mu') - \nabla_{x_0} h(x_0, \cdot, \mu) \right\|_{\mathfrak{b}} &\leq \Lambda \Big(|x'_0 - x_0| + \mathbb{W}_1(\mu', \mu) \Big), \\
\left\| \nabla_y^l \delta_\mu h(x'_0, \cdot, \mu', y') - \nabla_y^l \delta_\mu h(x_0, \cdot, \mu, y) \right\|_{\mathfrak{b}} &\leq \Lambda \Big(|x'_0 - x_0| + \mathbb{W}_1(\mu', \mu) + |y' - y|^{n - \lfloor n \rfloor} \Big). \n\end{aligned}
$$

We now introduce the two sets of assumptions.

Assumption A. There exist three reals $\lambda > 0$, $\kappa > 0$ and $\lambda > d/2 + 5$, $\lambda \notin \mathbb{N}$, such that

(A1) The functions $(f_t^0)_{0 \le t \le T}$ and g^0 satisfy $\mathscr{D}^0(\kappa, \lfloor \Delta \rfloor - (d/2 + 1))$; the functions $(f_t)_{0 \le t \le T}$ and g satisfy $\mathscr{D}(\kappa, \lfloor \Delta \rfloor (d/2+1), \Delta$).

(A2) The Lagrangians L^0 and L are λ -strictly convex in the variables α_0 and α respectively, uniformly with respect to the other variables. Moreover, L^0 has continuous joint derivatives up to the order 2 in x_0 and $\lfloor \lambda \rfloor + 1$ in α_0 , and L has continuous joint derivatives up to the order $\lfloor \Delta \rfloor + 1$ in (x, α) . The quantity $L^0(x_0, \alpha_0)$ is bounded by $\kappa (1 + |\alpha_0|^2)$ and the quantity $L(x, \alpha)$ is bounded by $\kappa(1+|\alpha|^2)$. The gradient $\nabla_{\alpha_0}L^0(x_0, \alpha_0)$ is bounded by $\kappa(1+|\alpha_0|)$ and the gradient $\nabla_{\alpha} L(x_0, x, \alpha)$ is bounded by $\kappa(1 + |\alpha|)$. All the other existing derivatives are bounded by κ .

(A3) The functions f and q satisfy the following Lasry-Lions monotonicity conditions

$$
\int_{\mathbb{R}^d} (f_t(x_0, x, \mu') - f_t(x_0, x, \mu)) d(\mu' - \mu)(x) \ge 0, \quad \int_{\mathbb{R}^d} (g(x_0, x, \mu') - g(x_0, x, \mu)) d(\mu' - \mu)(x) \ge 0,
$$

for all $t \in [0, T]$, $x_0 \in \mathbb{R}^d$, and $\mu, \mu' \in \mathcal{P}(\mathbb{T}^d)$.

(A4) The coefficient σ_0 is greater than 1.

Assumption B. On top of Assumption (A) and for the same parameters as therein, there exists two functions $F^0 : \mathbb{R}^d \to \mathbb{R}$ and $F: \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ such that

(B1) The function F^0 is continuously differentiable and bounded by κ , and its gradient is also bounded by κ and is κ -Lipschitz continuous; the function F, seen as a function of (x_0, x, μ) that would be constant in the variable x_0 , satisfies $\mathscr{D}(\kappa, |\lambda| - (d/2 + 1), \lambda).$

(B2) The following two bounds hold true:

$$
\int_0^T \sup_{x_0 \in \mathbb{T}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} |f_t^0(x_0, \mu) - F^0(x_0)| \, \mathrm{d}t \le \kappa, \quad \int_0^T \sup_{x_0 \in \mathbb{T}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|f_t(x_0, x, \mu) - F(x, \mu)\|_{\mathbf{A}} \, \mathrm{d}t \le \kappa.
$$

(B3) The functions $(f_t^0)_{0 \le t \le T}$ satisfy

$$
\int_0^T \sup_{x_0 \in \mathbb{T}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|\delta_{\mu} f_t^0(x_0, \mu, \cdot)\|_1 dt \leq \kappa.
$$

Remark 2.1. The following remarks are in order:

- 1. Assumption (B) has a simple interpretation. In long time, the running costs associated with the major and minor players become independent of the state of the other player. As made clear below, this assumption is very important to obtain a lower bound independent of T for the intensity σ_0 beyond which the Major/Minor MFG has the desired solvability properties.
- 2. Construction of examples satisfying (B) is quite simple. It suffices to start from given coefficients F^0 and F and to add perturbations $f_t^0 - F^0$ and $f_t - F$ that decay sufficiently fast as t tends to ∞ .
- 3. The form of Assumption (B) explains (up to some extent) our choice to restrict the analysis to Lagrangians L^0 and L that depend on the state of one player only (and not on the state of the other player). If one of the two Lagrangians were depending on both states, we would need a convenient form of condition (**B2**).

Hamiltonians. With the two Lagrangians L^0 and L, we associate the following two Hamiltonians:

$$
H^{0}(x_{0}, p) := \sup_{\alpha \in \mathbb{R}^{d}} [-p \cdot \alpha - L^{0}(x, \alpha)],
$$

\n
$$
H(x, p) = \sup_{\alpha \in \mathbb{R}^{d}} [-p \cdot \alpha - L(x, \alpha)].
$$
\n(2.2)

Under the standing standing assumptions on the Lagrangian L , we have the following representation formula for H :

$$
\nabla_p H(x,p) = (\nabla_\alpha L)^{\circ - 1}(x,p), \quad H(x,p) = p \cdot \nabla_p H(x,p) - L(x, -\nabla_p H(x,p)),\tag{2.3}
$$

and similarly for H^0 . We easily deduce that there exist two (strictly) positive constants λ' and κ' such that (notice that (A5) below is not an assumption but a consequence of Assumption (A)):

(A5) The Hamiltonians H^0 and H are λ' -strictly convex in the variables p_0 and p respectively, uniformly with respect to the other variables. Moreover, H^0 has continuous joint derivatives up to the order 2 in x_0 and $|z| + 1$ in p, and H has continuous joint derivatives up to the order $\lfloor \Delta \rfloor + 1$ in (x, p) . The quantity $H^0(x_0, p)$ is bounded by $\kappa'(1 + |p|^2)$ and the quantity $H(x, p)$ is bounded by $\kappa'(1+|p|^2)$. The gradient $\nabla_p H^0(x_0, p)$ is bounded by $\kappa'(1+|p|)$ and the gradient $\nabla_p H(x, p)$ is bounded by $\kappa'(1+|p|)$. All the other existing derivatives are bounded by κ' .

2.4. Weak formulation of the Major/Minor MFG

Below, we address a weak formulation of the game in which the spaces Ω^0 and Ω are taken as the canonical spaces $\Omega_{\text{canonical}}^0 := \mathcal{C}([0,T],\mathbb{R}^d) \times \mathcal{C}([0,T],\mathcal{P}(\mathbb{T}^d)) \times \mathcal{C}([0,T],\mathbb{R}^d)$ and $\Omega_{\text{canonical}} := \mathcal{C}([0,T],\mathbb{R}^d) \times \mathcal{C}([0,T],\mathbb{R}^d)$, equipped with their respective Borel σ -fields $\mathcal{B}(\Omega_{\text{canon}}^0)$ and $\mathcal{B}(\Omega_{\text{canon}})$. For simplicity, the canonical processes on Ω_{canon}^0 and Ω_{canon} are still denoted $(X^0, \mu, B^0) = (X_t^0, \mu_t, B_t^0)_{0 \le t \le T}$ and $(X, B) = (X_t, B_t)_{0 \le t \le T}$, and the canonical filtrations are still denoted $\mathbb{F}^0 = (\mathcal{F}^0_t)_{0 \le t \le T}$ and $\mathbb{F} = (\mathcal{F}_t)_{0 \le t \le T}$.

Definition 2.2. For fixed initial conditions $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, a pair (α^0, α) , with $\alpha^0 : [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ and $\alpha: [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ is said to be admissible if α^0 and α are measurable, α is bounded and there exists a unique probability measure \mathbb{P}^0 on $\Omega_{\text{canonical}}^0$ such that: (i) under \mathbb{P}^0 , the process B^0 is an- \mathbb{F}^0 Brownian motion starting from 0; (ii) the pair (X^0, μ) satisfies $\mathbb{P}^0(\{(X^0_0, \mu_0) = (x_0, \mu)\}) = 1$; (iii) the two equations

$$
dX_t^0 = \alpha^0(t, X_t^0, \mu_t)dt + dB_t^0,
$$

\n
$$
\partial_t \mu_t = \frac{1}{2} \Delta_x \mu_t - \text{div}_x (\alpha(t, X_t^0, \cdot, \mu_t) \mu_t), \quad t \in [0, T],
$$
\n(2.4)

are satisfied under \mathbb{P}^0 ; (iv) the BMO condition

$$
\left\| \int_0^{\cdot} \alpha^0(t, X_t^0, \mu_t) \cdot \mathrm{d}B_t^0 \right\|_{\text{BMO}} < \infty \tag{2.5}
$$

holds true. When needed to clarify the set-up, we put an additional index in \mathbb{P}^0 and write $\mathbb{P}^0_{(\alpha^0,\alpha)}$.

Remark 2.3. In [\(2.4\)](#page-6-1), the Kolmogorov equation is understood in a weak sense, namely, for \mathbb{P}^0 -almost every $\omega^0 \in \Omega_{\text{canon}}^0$, for any function $\phi: [0, T] \times \mathbb{T}^d \to \mathbb{R}^d$ in the space $\mathcal{W}_{1,2}^{d+1}$ of functions with first and second order x-derivatives in space and first order *t*-derivative in $L^{d+1}(\mathbb{T}^d)$,

$$
(\phi_t, \mu_t) = (\phi_0, \mu_0) + \int_0^t \left(\partial_t \phi_s(\cdot) + \frac{1}{2} \Delta_x \phi_s(\cdot) + \alpha(s, X_s^0, \cdot, \mu_s) \cdot \nabla_x \phi_s(\cdot), \mu_s\right) \mathrm{d}s, \quad t \in [0, T].\tag{2.6}
$$

By an obvious separability argument, the above is equivalent to having the same expansion for any $\phi \in \mathcal{W}_{1,2}^{d+1}$ and for \mathbb{P}^0 -almost every $ω^0$ ∈ Ω 0 _{canon}.

The definition of the costs in the weak formulation relies on the following lemma (we recall that X^0 and μ are part of the canonical process on Ω_{canon}^0):

Lemma 2.4. Let $\mu \in \mathcal{P}(\mathbb{T}^d)$ and $\alpha : [0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ be a bounded and mesurable function as in Definition *[2.2.](#page-6-2) Then, there exists a measurable mapping*

$$
\Omega_{\text{canon}}^0 \ni \omega^0 \mapsto \mathbb{P}_{\omega^0} \in \mathcal{P}(\Omega_{\text{canon}}),
$$

such that, for any $\omega^0 \in \Omega_{\text{canon}}^0$, \mathbb{P}_{ω^0} is the unique probability measure on Ω_{canon} satisfying the following three items: (i) *under* \mathbb{P}_{ω^0} , the process B is an- $\mathbb F$ Brownian motion starting from 0; (ii) the pair (X, B) satisfies $\mathbb{P}_{\omega^0} \circ X_0^{-1} = \mu$; (iii) *the equation*

$$
dX_t = \alpha(t, X_t^0, X_t, \mu_t)dt + dB_t, \quad t \in [0, T],
$$
\n(2.7)

is satisfied under \mathbb{P}_{ω^0} *. When needed to clarify the set-up, we put an additional index in* \mathbb{P}_{ω^0} *and write* $\mathbb{P}_{\omega^0,\alpha}$ *.*

Remark 2.5. Using Itô-Krylov formula, it is easy to check, for $\omega^0 \in \Omega^0$ and for any function $\phi : [0, T] \times \mathbb{T}^d \to \mathbb{R}^d$ in the space $W_{1,2}^{d+1}$, the flow of probability measures $\nu := (\nu_t = \mathbb{P}_{\omega^0} \circ X_t^{-1})_{0 \le t \le T}$ satisfies the Fokker-Planck equation

$$
\partial_t \nu_t = \frac{1}{2} \Delta_x \nu_t - \text{div}_x \big(\alpha(t, X_t^0, \cdot, \mu_t) \nu_t \big), \quad t \in [0, T]; \quad \nu_0 = \mu,
$$
\n
$$
(2.8)
$$

in the same weak sense as in Remark [2.3.](#page-7-0) As such, ν is \mathbb{F}^0 -adapted, because for any bounded test function $\phi : \mathbb{T}^d \to \mathbb{R}$, the process $(\int_{\mathbb{T}^d} \phi(x) d\nu_t(x) = \mathbb{E}_{\omega^0}[\phi(X_t)])_{0 \le t \le T}$ is \mathbb{F}^0 -progressively measurable.

In fact, this is the unique weak solution to [\(2.8\)](#page-7-1). Indeed, any weak solution, say $\tilde{\nu} = (\tilde{\nu}_t)_{0 \leq t \leq T}$, to the Fokker-Planck equation (understood in the same weak sense as (2.6) in Remark [2.3\)](#page-7-0) is necessarily equal to the flow of marginal laws of the solution to the SDE [\(2.33\)](#page-17-0). This follows from a standard duality argument that consists in solving the parabolic equation $\partial_t \phi_t + \frac{1}{2} \Delta_x \phi_t + \alpha(t, X_t(\omega^0), \cdot, \mu_t) \cdot \nabla_x \phi_t = 0$, with $\phi_T = \ell$ for a prescribed smooth function $\ell : \mathbb{T}^d \to \mathbb{R}$, and then in expanding $((\phi_t, \tilde{\nu}_t))_{0 \le t \le T}$. In this way, we get $(\ell, \tilde{\nu}_T) = (\phi_T, \tilde{\nu}_T) = (\phi_0, \mu)$ for any smooth ℓ , which suffices to identify $\tilde{\nu}_T$. Replacing T by t, we can proceed in a similar manner and identify $\tilde{\nu}_t$ for any $t \in [0, T]$.

The proof of Lemma [2.4](#page-7-3) is postponed to the end of the section, see Subsection [2.7.](#page-16-1) For the time being, the statement makes it possible to let:

Definition 2.6. For a fixed initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and with the same notation as in Definitions [2.2](#page-6-2) and Lemma [2.4,](#page-7-3) we let:

1. Cost to the major: Let (α^0, α) be an admissible pair and $\mathbb{P}^0_{(\alpha^0, \alpha)}$ be the probability associated with it by Definition [2.2.](#page-6-2) Then, the cost to the major player is defined by

$$
J_{\mathbf{w}}^0(\alpha^0, \alpha) = \mathbb{E}^0_{(\alpha^0, \alpha)} \bigg[g^0(X_T^0, \mu_T) + \int_0^T \bigg(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, \alpha_t^0) \bigg) dt \bigg];
$$

2. Cost to the minor: Let $\alpha : [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ be a bounded and mesurable function and \mathbb{P}^0 be a probability measure on $\Omega_{\text{canonical}}^0$. Then, the cost to the minor player in the environment \mathbb{P}^0 is defined by

$$
J_{\mathbf{w}}(\alpha; \mathbb{P}^0) = \int_{\Omega^0} \mathbb{E}_{\omega^0, \alpha} \bigg[g(X_T^0, X_T, \mu_T) + \int_0^T \Big(f_t(X_t^0, X_t, \mu_t) + L(X_t, \alpha_t) \Big) dt \bigg] d\mathbb{P}^0(\omega^0).
$$

The following definition is inspired from [\[12,](#page-86-22) [13,](#page-86-27) [47\]](#page-87-7):

Definition 2.7. Let $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ be a fixed initial condition and (α^0, α) be an admissible pair in the sense of Definition [2.2.](#page-6-2) Recalling the notation (X^0, μ, B^0) for the canonical process on $\Omega_{\text{canonical}}^0$ and the notation (X, B) for the canonical process on Ω_{canon} , the pair (α^0, α) is said to be a mean field equilibrium if

1. **Deviation of the minor**: For any bounded and measurable function β : $[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$,

$$
J_{\mathbf{w}}\big(\alpha; \mathbb{P}^0_{(\alpha^0, \alpha)}\big) \leq J_{\mathbf{w}}\big(\beta; \mathbb{P}^0_{(\alpha^0, \alpha)}\big);
$$

2. **Deviation of the major**: For any feedback function β^0 : $[0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ such that the pair (β^0, α) is admissible,

$$
J_{\mathbf{w}}^0(\alpha^0,\alpha) \le J_{\mathbf{w}}^0(\beta^0,\alpha).
$$

Remark 2.8. It is worth emphasizing that the nature of the equilibrium would be different if α^0 and α were required to be in open loop form. The analysis of equilibria over controls in Markov feedback form is in fact more difficult.

As far as the control β is concerned in item 1 right above, the nature of it (open versus closed) does not make any difference. Intuitively, this comes from the fact that any deviation of the minor player has no influence on the state of the population (encoded through μ) nor on the state of the major player (encoded through X^0). This fact is standard in MFG theory.

We now give conditions under which a pair (α^0, α) is admissible in the sense of Definition [2.2:](#page-6-2)

Lemma 2.9. Let $\alpha^0 : [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ and $\alpha : [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d$ be two Borel measurable *functions such that* α *is bounded and satisfies*

$$
\sup_{t\in[0,T]}\sup_{x_0\in\mathbb{R}^d\mu,\mu'\in\mathcal{P}(\mathbb{T}^d):\mu\neq\mu'}\sup_{x\in\mathbb{T}^d}\frac{|\alpha(t,x_0,x,\mu')-\alpha(t,x_0,x,\mu)|}{\mathbb{W}_1(\mu,\mu')}<\infty.
$$
\n(2.9)

Then:

(a) For any $(x_0,\mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, there exists at most one probability measure \mathbb{P}^0 on $\Omega_{\text{canonical}}^0$ such that items (i), (ii), *(iii) and (iv) in Definition [2.2](#page-6-2) are satisfied.*

 (a) For any $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, there exists a unique probability measure $\overline{\mathbb{P}}^0$ on $\Omega_{\text{canonical}}^0$ such that items (i), (ii) and (*iii*) in Definition [2.2](#page-6-2) are satisfied when $\alpha^0 \equiv 0$.

(c) If, for a given $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the BMO condition

$$
\left\| \int_0^{\cdot} \alpha^0(t, X_t^0, \mu_t) \cdot \mathrm{d}B_t^0 \right\|_{\text{BMO}} < \infty \tag{2.10}
$$

is satisfied under $\bar{\mathbb{P}}^0$, then there exists a (hence unique) probability measure \mathbb{P}^0 on $\Omega_{\rm canon}^0$ such that items (i), (ii), (iii) and (iv) are satisfied under Definition [2.2.](#page-6-2) Conversely, if there exists a probability measure \mathbb{P}^0 on Ω_{canon}^0 such that items (i), (*ii*), (*iii*) and (*iv*) are satisfied, then the BMO condition [\(2.10\)](#page-8-0) is satisfied under $\bar{\mathbb{P}}^0$.

Similar to Lemma [2.4,](#page-7-3) Lemma [2.9](#page-8-1) is proven in Subsection [2.7.](#page-16-1)

2.5. Forward-backward characterization

Our analysis below relies on a stochastic forward-backward system, which plays the same role as the MFG system in the standard setting. This system reads in the form of two coupled stochastic forward-backward equations, which are understood in a strong sense and thus posed on any arbitrary filtered probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ satisfying the usual conditions and equipped with a Brownian motion $(B_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d . In particular, we NO longer regard $(\bm{X}^0,\bm{\mu},\bm{B}^0)=(X_t^0,\mu_t,B_t^0)_{0\leq t\leq T}$ and $(\bm{X},\bm{B})=(X_t,B_t)_{0\leq t\leq T}$ as canonical processes in this subsection.

The first equation provides a Lagrangian description of the state of the major player (at equilibrium):

$$
dX_t^0 = -\nabla_p H^0(X_t^0, Z_t^0) dt + \sigma_0 dB_t^0, \quad t \in [0, T],
$$

\n
$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0))\right) dt + \sigma_0 Z_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$
\n
$$
X_0^0 = x_0, \quad Y_T^0 = g^0(X_T^0, \mu_T).
$$
\n(2.11)

Above, the measure argument $(\mu_t)_{0 \le t \le T}$ in the dynamics of $(Y_t^0)_{0 \le t \le T}$ corresponds to the statistical law of the minor player, whose evolution (at equilibrium) is described by the stochastic system (which is the second of the two aforementioned forward-backward equation):

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t) \mu_t \right) = 0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mathrm{d}_t u_t(x) = \left(-\frac{1}{2} \Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t) \right) \mathrm{d}t + \sigma_0 v_t^0(x) \cdot \mathrm{d}B_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mu_0 = \mu, \quad u_T(x) = g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d.
$$
 (2.12)

Definition 2.10. Solutions to the above two systems are understood in the following sense:

- 1. For a given initial condition $x_0 \in \mathbb{R}^d$ and an \mathbb{F}^0 -adapted continuous process $\mu = (\mu_t)_{0 \le t \le T}$ with values in $\mathcal{P}(\mathbb{T}^d)$, we call solution to (2.11) any \mathbb{F}^0 -progressively measurable process $(\mathbf{X}^0, \mathbf{Y}^0, \mathbf{Z}^0)$ with values in $\mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$, such that: (i) X^0 and Y^0 have continuous trajectories; (ii) $\sup_{0 \le t \le T} |X_t^0| \in L^2(\Omega^0, \mathbb{P}^0)$ and $\sup_{0 \le t \le T} |Y_t^0| \in$ $L^{\infty}(\Omega^0, \mathbb{P}^0)$; (iii) $\sup_{\tau} ||\mathbb{E}^0[\int_{[\tau, T]} |Z_t^0|^2 dt | \mathcal{F}^0_{\tau}]] ||_{L^{\infty}(\Omega^0, \mathbb{P}^0)} < \infty$, the supremum being taken over the collection of stopping times τ with respect to the usual augmentation of the filtration generated by (X^0, Y^0, Z^0, B^0) ; (iv) the system (2.11) is satisfied \mathbb{P}^0 - almost surely.
- 2. For a given initial condition $\mu \in \mathcal{P}(\mathbb{T}^d)$ and an \mathbb{F}^0 -adapted continuous process $\mathbf{X}^0 = (X_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d such that $\sup_{0 \le t \le T} |X_t^0| \in L^2(\Omega^0, \mathbb{P}^0)$, we call solution to [\(2.12\)](#page-9-0) any \mathbb{P}^0 -progressively measurable process (μ, u, v^0) with values in $\mathcal{P}(\mathbb{T}^d) \times \mathcal{C}^{\lambda}(\mathbb{T}^d) \times \mathcal{C}^{\lambda-1}(\mathbb{T}^d)$, such that: (i) μ and u have continuous trajectories in $\mathcal{P}(\mathbb{T}^d) \times \mathcal{C}^{\gamma}(\mathbb{T}^d)$ for any $\gamma < \delta$; (ii) $\sup_{0 \le t \le T} ||u_t||_{\delta} \in L^{\infty}(\Omega^0, \mathbb{P}^0)$; (iii) $\mathbb{E}^0 \int_{[0,T]} ||v_t^0||_{\lfloor \delta \rfloor - d/2 - 1}^2 ds < \infty$; (iv) the forward equation [\(2.12\)](#page-9-0) is satisfied **P** 0 almost surely in the same weak sense as in Remark [2.3;](#page-7-0) (v) the backward equation is satisfied **P** 0 -almost surely (in the classical sense).
- 3. For a given initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, we call a solution to the coupled systems [\(2.11\)](#page-8-2)–[\(2.12\)](#page-9-0) a pair $((X^0, Y^0, Z^0), (\mu, u, v^0))$ satisfying items 1 and 2 right above.

Remark 2.11. The following remarks are in order:

- 1. The process $(Y_t^0)_{0 \le t \le T}$ is expected to describe the evolution of the equilibrium cost to the major player. Intuitively, $(-\nabla_p H^0(X_t^0, Z_t^0))_{0 \le t \le T}$ is the corresponding equilibrium feedback.
- 2. In the stochastic forward-backward system [\(2.12\)](#page-9-0), the (random) function $(t, x) \mapsto u_t(x)$ is the equilibrium value of the minor player. The optimal feedback is $(t, x) \mapsto -\nabla_p H(x, \nabla_x u_t(x))$. Obviously, $(u_t)_{0 \le t \le T}$ is random under the presence of the noise B^0 acting on the major player. The term $(v_t^0)_{0 \le t \le T}$ is here to ensure that $(u_t)_{0 \le t \le T}$ is indeed **F** 0 -adapted.

It is worth mentioning that, in comparison with the two forward and backward equations (4.2) and (4.3) introduced in [\[14,](#page-86-2) Chapter 4] in the analysis of MFGs with a common noise, our own system is simpler because the dynamics of the minor player are NOT forced by the common noise B^0 . In particular, the forward equation is not a stochastic Fokker-Planck equation (like [\[14,](#page-86-2) (4.2)]) but a Fokker-Planck equation with random coefficients. For the same reason, the backward equation does not contain any Itô-Wentzell correction comparable to the one appearing in $[14, (4.3)]$ $[14, (4.3)]$.

3. Below, the analysis of the backward SPDE in [\(2.12\)](#page-9-0) is inspired by the study carried out in the monograph [\[14\]](#page-86-2), in which a similar equation is treated within the framework of mean field games with common noise (see Chapter 4 therein). However, we have slightly changed the spaces in which solutions are taken: we feel clearer to see them as random processes with values in non-integer Hölder spaces, whilst they are regarded as random process with values in integer Hölder spaces in [\[14](#page-86-2)]. This requires some care because, for $\lambda \notin \mathbb{N}$, the space $C^{\lambda}(\mathbb{T}^d)$ is not separable. Working with (non-separable) Banach-valued random variables is indeed an issue, see for instance [\[48\]](#page-87-8). One standard way to overcome the lack of separability is to strengthen the notion of measurability and to work with *Bochner* measurable random variables. As explained in [\[49\]](#page-87-9) (see also [\[39\]](#page-86-32) for an overview), a random variable with values in a Banach space E (measurability being understood with respect to the standard Borel σ -field on E) is *Bochner* measurable if it takes values in a separable subspace of E. Any such *Bochner* measurable random variable has a tight distribution and can be approximated by simple random variables, which makes its manipulation easier. In the specific framework of [\(2.12\)](#page-9-0), one can typically choose $C^{b'}(\mathbb{T}^d)$, for $b' > b$, as separable subspace of $C^b(\mathbb{T}^d)$. Indeed, Schauder estimates make it possible to gain some extra regularity on u_t , for $t < T$, and then to regard the latter as an element of $C^{S'}(\mathbb{T}^d)$, for $S' > S$. At time $t = T$, *Bochner* measurability can be checked directly thanks to the properties of g . Combined with the continuity properties stated in Definition [2.10](#page-9-1) and an interpolation inequality in Hölder spaces, this argument says even more: u has continuous trajectories from $[0, T - \varepsilon]$ to $\mathcal{C}^{\Delta'}$, for any $\varepsilon > 0$ and some $s' > s$. As such, u can be written as the almost everywhere limit in $[0, T] \times \Omega$ of simple processes of the form $\sum_{i=0}^{n-1} X_{t_i} \mathbf{1}_{(t_i,t_{i+1}]}$, with $n \ge 1$, $t_0 = 0 < t_1 < \cdots < t_n = T$ and X_{t_i} an \mathcal{F}_{t_i} *Bochner* measurable $\mathcal{C}^{\Delta}(\mathbb{T}^d)$ valued random variable for each $i = 0, \dots, n - 1$. This extends the notion of Bochner measurability to processes. We will come back to these measurability questions when needed, but the message is clear: measurability properties stated in Definition [2.10](#page-9-1) are in fact understood in the *Bochner* sense. Of course, there is no similar difficulty with the forward component of the system (2.12) because $\mathcal{P}(\mathbb{T}^d)$ is compact when equipped with any standard distance metricizing the weak topology.

4. In the same vein as above, notice that, in the second item of Definition [2.10,](#page-9-1) $\sup_{0 \le t \le T} ||u_t||_{\delta}$ is necessarily measurable if u has continuous trajectories in $\mathcal{C}^{\pi}(\mathbb{T}^d)$ for any $\pi < \infty$. Indeed, for $\lfloor \Delta \rfloor < \pi < \infty$, the norm $\|\cdot\|_{\infty}$ is lower semi-continuous on $\mathcal{C}^n(\mathbb{T}^d)$, which shows that $\sup_{0 \le t \le T} ||u_t||_{\delta}$ is then equal to $\sup_{t \in [0,T] \cap \mathbb{Q}} ||u_t||_{\delta}$.

The following statement clarifies the connection with Definition [2.10:](#page-9-1)

Proposition 2.12. Assume that, on any filtered probability space $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$ equipped with an \mathbb{F}^0 -Brownian *motion* $\mathbf{B}^0 = (B_t^0)_{0 \le t \le T}$ *with values in* \mathbb{R}^d , for any initial condition $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the system (2.11) – (2.12) has a unique solution (in the sense of Definition [2.10\)](#page-9-1), denoted $(X_s^{0,t,x_0,\mu}, Y_s^{0,t,x_0,\mu}, Z_s^{0,t,x_0,\mu})_{t\leq s\leq T}$ and $(\mu_s^{t,x_0,\mu}, u_s^{t,x_0,\mu}, v_s^{0,t,x_0,\mu})_{t\leq s\leq T}$, and satisfying

$$
\sup_{t \in [0,T]} \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu,\mu' \in \mathcal{P}(\mathbb{T}^d): \mu \neq \mu'} \frac{\|u_t^{t,x_0,\mu}(\cdot) - u_t^{t,x_0,\mu'}(\cdot)\|_1}{\mathbb{W}_1(\mu,\mu')} < \infty.
$$
\n(2.13)

Then, there exists a pair (α^0,α) satisfying the Definition [2.7](#page-7-4) of a mean field equilibrium such that, for any initial condition $(x_0,\mu)\in\mathbb{R}^d\times\mathcal{P}(\mathbb{T}^d)$ to the Major/Minor MFG at time 0, the law of the forward path $(X_s^{0,0,x_0,\mu},\mu_s^{0,x_0,\mu},B_s^0)_{0\leq s\leq T}$ *coincides with the measure* $P_{(\alpha^0, \alpha)}$ *defined in Definition [2.7.](#page-7-4)*

Moreover, assume that for a fixed $(x_0,\mu)\in\mathbb{R}^d\times\mathcal{P}(\mathbb{T}^d)$, there exists another mean field equilibrium $(\tilde{\alpha}^0,\tilde{\alpha})$ to the *Major/Minor initialized at* (x_0, μ) *at time* 0 *such that: (i)* $\tilde{\alpha}$ *satisfies*

$$
\sup_{t \in [0,T]} \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu,\mu' \in \mathcal{P}(\mathbb{T}^d): \mu \neq \mu'} \sup_{x \in \mathbb{T}^d} \frac{|\tilde{\alpha}(t,x_0,x,\mu') - \tilde{\alpha}(t,x_0,x,\mu)|}{\mathbb{W}_1(\mu,\mu')} < \infty; \tag{2.14}
$$

and (ii) the state equation [\(2.4\)](#page-6-1) driven by $(\tilde{\alpha}^0, \tilde{\alpha})$ and defined on the canonical space $\Omega_{\rm canon}^0$ has, for any starting point $(t, \tilde{x}_0, \tilde{\mu}) \in [0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, a solution that is adapted with respect to the (augmentation of the) filtration generated by $(B_s^0 - B_t^0)_{t \le s \le T}$ *(which is here the third component of the canonical process). Then,* $\mathbb{P}_{(\tilde{\alpha}^0, \tilde{\alpha})} = \mathbb{P}_{(\alpha^0, \alpha)}$.

Remark 2.13. Our statement may seem rather complicated at first sight. In fact, the main idea is to limit the analysis to the existence and uniqueness of Nash equilibria that are adapted to the common noise. By analogy with the terminology used in the theory of SDEs, those equilibria should be called "strong". As for the existence of such strong equilibria, the key point here is that the system (2.11) – (2.12) is assumed to be uniquely strongly solvable. This forces the solutions (to the system) to be adapted to the common noise and also implies the existence of a feedback function, see the first step of the proof below. As for uniqueness, the main assumption consists of the two items (i) and (ii) in the second part of the statement. As shown in the third step of the proof, the combination of both forces the equation [\(2.4\)](#page-6-1) (when driven by driven by $(\tilde{\alpha}^0, \tilde{\alpha})$ to be uniquely strongly solvable.

Here is now the main statement of our article:

Theorem 2.14. *Under Assumption* (A), for any $T > 0$, there exists a threshold $\sigma_0^*(T) \in (0, +\infty)$ *such that, for* $\sigma_0 \geq \sigma_0^*(T)$, for any initial condition $(x_0,\mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the system (2.11) – (2.12) has a unique solution in the $\lim_{n \to \infty} \frac{\partial f}{\partial n}$ *(on any filtered probability space* $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$ *equipped with an* \mathbb{F}^0 -*Brownian motion* $B^0 = (B_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d). Moreover, [\(2.13\)](#page-10-1) holds true.

If in addition, Assumption (**B**) is also in force, then we can choose $\sigma_0^*(T)$ independently of T. Namely, we can find a *threshold* $\sigma_0^* \in (0, +\infty)$ such that existence and uniqueness hold true on any interval $[0, T]$, $T > 0$, and for any $\sigma_0 \ge \sigma_0^*$.

Notice that [\(2.13\)](#page-10-1) implicitly requires to solve [\(2.11\)](#page-8-2)–[\(2.12\)](#page-9-0) when the initial condition is fixed at any time $t \in [0, T]$. Observe in particular that, combined with Proposition [2.12,](#page-10-2) Theorem [2.14](#page-10-0) says the the Major/Minor MFG has a unique "strong" equilibrium when $\sigma_0 \ge \sigma_0^*(T)$ (and (A) holds true) or $\sigma_0 \ge \sigma_0^*$ (and (B) holds true).

The proof of Theorem [2.14](#page-10-0) is deferred to Sections [3](#page-18-0) and [4,](#page-30-0) see in particular Theorem [4.11](#page-41-0) for a refined version of it. The proof of the latter makes explicit use of the condition $s > d/2 + 5$. Notice in fact that it suffices to prove the second part of Theorem [2.14,](#page-10-0) namely the claim under Assumption (B). Indeed, once the conclusion has been proved to hold true under Assumption (B), one can easily the derive the first part of the statement by modifying the constant κ in (A) in such 12

a way that (B2) and (B3) hold true. This is possible to do so by replacing κ by a new constant that is allowed to depend on T .

We now turn to the proof of Proposition [2.12](#page-10-2) (the reader may skip it on a first reading):

Proof of Proposition [2.12.](#page-10-2) The proof is divided into three steps.

First Step. The first step is to prove that, for a fixed initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ at time 0 and under the standing unique solvability property of the system (2.11) – (2.12) , the latter induces a Nash equilibrium to the Major/Minor MFG. To do so, we consider a probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ equipped with a Brownian motion $\mathbf{B}^0 = (B_t^0)_{0 \le t \le T}$. We assume that \mathbb{F}^0 is the \mathbb{P}^0 -completion of the filtration.generated by B^0 , from which we deduce that solutions to [\(2.11\)](#page-8-2)–[\(2.12\)](#page-9-0), when constructed on this space, are necessarily adapted to the (completion) of the filtration generated by B^0 .

Following $[20,$ Proposition 1.31], strong uniqueness of the solution to (2.11) – (2.12) implies that the mapping that sends the initial condition (t, x_0, μ) onto the law of $(X^{t, x_0, \mu}_{s \lor t}, \mu^{t, x_0, \mu}_{s \lor t})_{0 \le s \le T}$ on $\mathcal{C}([0, T], \mathbb{R}^d) \times \mathcal{C}([0, T], \mathcal{P}(\mathbb{T}^d))$ is measurable, from which we deduce that the solution to (2.11) – (2.12) forms a strong Markov process. Then, following [\[38,](#page-86-33) Proposition 3.2 & Theorem 3.4] (which relies on Theorem 6.27 in [\[25](#page-86-34)]), we can find a Borel function $\psi^0 : [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \rightarrow$ $\mathbb{R}^d \otimes \mathbb{R}^d$ such that, for any initial time $t \in [0, T]$ and any time $s \in [t, T]$,

$$
\mathbb{P}^0\Big(\big\{Z^{0,t,x_0,\mu}_s=\psi^0(s,X^{0,t,x_0,\mu}_s,\mu^{0,t,x_0,\mu}_s)\big\}\Big)=1,
$$

which shows the existence of a Markov feedback function for the process $X^{0,t,x_0,\mu}$ (understood below as the state of the major player).

Also, by observing from the unique strong solvability of (2.11) – (2.12) that the process $u^{t,x_0,\mu}$ is, for any $(t,x_0,\mu) \in$ $[0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, adapted to the (completion of the) filtration generated by $(B_s^0 - B_t^0)_{t \le s \le T}$, we can define the value to the minor player as

$$
V(t, x_0, x, \mu) = u_t^{t, x_0, \mu}(x), \quad x \in \mathbb{T}^d, \quad (t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d).
$$

Since the process $u^{t,x_0,\mu}$ takes values in $\mathcal{C}^{\delta}(\mathbb{T}^d)$, we deduce that V is differentiable in x. The gradient $\nabla_x V$ induces a measurable mapping from $[0, T] \times \mathbb{R}^d \times T^d \times \mathcal{P}(T^d)$ to \mathbb{R}^d . By unique strong solvability of [\(2.11\)](#page-8-2)–[\(2.12\)](#page-9-0), we know that, for any $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and any $s \in [t, T]$,

$$
\mathbb{P}^0\Big(\big\{\forall x \in \mathbb{T}^d, \ u_s^{0,t,x_0,\mu}(x) = V(s,X^{0,t,x_0,\mu}_s,x,\mu_s^{0,t,x_0,\mu})\big\}\Big) = 1,
$$

and then,

$$
\mathbb{P}^0\Big(\big\{\forall x\in \mathbb{T}^d, \; \nabla_x u_s^{0,t,x_0,\mu}(x)=\nabla_x V\big(s,X_s^{0,t,x_0,\mu},x,\mu_s^{0,t,x_0,\mu}\big)\big\}\Big)=1,
$$

which shows the existence of a Markov feedback function for the minor player. Moreover, under the assumption [\(2.13\)](#page-10-1), the partial derivative $\nabla_x V$ appearing in the above event is Lipschitz continuous in the measure argument, uniformly with respect to the other parameters. This supplies us with the following two feedback functions:

$$
\alpha^{0} : [0, T] \times \mathbb{R}^{d} \times \mathcal{P}(\mathbb{T}^{d}) \ni (x_{0}, \mu) \mapsto -\nabla_{p} H^{0}(x_{0}, \psi^{0}(t, x_{0}, \mu)),\alpha : [0, T] \times \mathbb{R}^{d} \times \mathbb{T}^{d} \times \mathcal{P}(\mathbb{T}^{d}) \ni (x_{0}, x, \mu) \mapsto -\nabla_{p} H(x, \nabla_{x} V(t, x_{0}, x, \mu)).
$$
\n(2.15)

The next point is to show that, for any fixed initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ (choose, to simplify, 0 as initial time), the pair (α^0, α) is admissible in the sense of Definition [2.2.](#page-6-2) By transferring the law of $(\mathbf{X}^{0,0,x_0,\mu}, \boldsymbol{\mu}^{0,x_0,\mu}, \boldsymbol{B}^0)$ onto the canonical space Ω_{canon}^0 , we get the existence of a probability measure $\mathbb{P}^0_{(\alpha^0,\alpha)}$ under which the system [\(2.4\)](#page-6-1) (with (α^0,α) given by [\(2.15\)](#page-11-0)) is satisfied. Denoting here by (X^0, μ, B^0) the canonical process on Ω_{canon}^0 , the pair (X^0, μ) is adapted to the completion under $\mathbb{P}_{(\alpha^0,\alpha)}^0$ of the filtration generated by \mathcal{B}^0 . Hence, under the completion of $\mathbb{P}_{(\alpha^0,\alpha)}^0$ (still denoted $\mathbb{P}^0_{(\alpha^0,\alpha)}$), one can solve the backward equation in [\(2.11\)](#page-8-2) and then denote the solution by $(\mathbf{Y}^0,\mathbf{Z}^0)=(Y_t^0,Z_t^0)_{0\leq t\leq T}$, see [\[41](#page-87-10)] for standard solvability results for quadratic BSDEs. Then, observing that the law of the solution to the backward equation in [\(2.11\)](#page-8-2) is uniquely determined by the law of the input $(\mathbf{X}^0, \boldsymbol{\mu}, \mathbf{B}^0)$, one necessarily has, for Leb $\times \mathbb{P}_{(\alpha^0, \alpha)}$ almost every $(t, \omega^0) \in [0, T] \times \Omega_{\text{canon}}^0$, $Z_t^0 = \psi^0(t, X_t^0, \mu_t)$, which proves that $(\mathbf{X}^0, \mathbf{Y}^0, \mathbf{Z}^0)$ solves the forward-backward system [\(2.11\)](#page-8-2) under $\mathbb{P}^0_{(\alpha^0,\alpha)}$. The BMO condition [\(2.5\)](#page-6-3) is then established by means of standard results for backward SDEs, see e.g. [\[57](#page-87-11)].

Uniqueness of this probability measure is a consequence of Lemma [2.9,](#page-8-1) which says that the law of the solution to [\(2.4\)](#page-6-1) (with (α^0, α) given by [\(2.15\)](#page-11-0)) is uniquely determined. This shows that the pair (α^0, α) fits the requirements of Definition [2.2.](#page-6-2)

Second Step. We now prove that the pair (α^0, α) in [\(2.15\)](#page-11-0) defines an equilibrium in the sense of Definition [2.7.](#page-7-4) As in the first step, we do so for a fixed initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ at time 0.

We first check item 1 in Definition [2.7.](#page-7-4) For β as in item 1 and for $(\mathbb{P}_{\omega^0,\beta})_{\omega^0\in\Omega_{\text{cmon}}^0}$ as in Lemma [2.4,](#page-7-3) and with $(\mathbf{X}^0, \mu, \mathbf{B}^0) = (X_t^0, \mu_t, B_t^0)_{0 \le t \le T}$ and $(\mathbf{X}, \mathbf{B}) = (X_t, B_t)_{0 \le t \le T}$ denoting the canonical processes on Ω_{canon}^0 and Ω_{canon} , we know that, for any $\omega^0 \in \Omega^0$, the following equation holds true under $\mathbb{P}^0_{\omega^0,\beta}$:

$$
dX_t = \beta(t, X_t^0, X_t, \mu_t)dt + dB_t, \quad t \in [0, T],
$$

with $\mathbb{P}^0_{\omega^0,\beta} \circ X_0^{-1} = \mu$. We then introduce, on $\Omega_{\text{canon}}^0 \times \Omega_{\text{canon}}$, the probability measure $\mathbb{P}^0_{(\alpha^0,\alpha)} \otimes \mathbb{P}_{\cdot,\beta}$ defined by

$$
\mathbb{P}^0_{(\alpha^0,\alpha)}\otimes \mathbb{P}_{\cdot,\beta}\big(A^0\times A\big)=\int_{\Omega^0_{\text{canon}}}\mathbf{1}_{A^0}(\omega^0)\mathbb{P}_{\omega^0,\beta}(A)\mathrm{d}\mathbb{P}^0_{(\alpha^0,\alpha)}(\omega^0),\quad A^0\in \mathcal{B}(\Omega^0_{\text{canon}}),\ A\in \mathcal{B}(\Omega_{\text{canon}}).
$$

We then expand $(u_t(X_t))_{0\leq t\leq T}$ by using Itô-Wentzell formula (which does not raise any difficulty in this setting because $(u_t)_{0 \le t \le T}$ is independent of $(B_t)_{0 \le t \le T}$, see^{[1](#page-12-0)}). We obtain, under $\mathbb{P}^0_{(\alpha^0,\alpha)} \otimes \mathbb{P}_{\cdot,\beta}$, for all $t \in [0,T]$,

$$
u_T(X_T) - u_t(X_t) + \int_t^T \left[f_s(X_s^0, X_s, \mu_s) + L(X_s, \beta_s) \right] ds
$$
\n
$$
\geq \int_t^T \left[H\left(X_s, \nabla_x u_s(X_s)\right) + \beta_s \cdot \nabla_x u_s(X_s) + L(X_s, \beta_s) \right] ds + \int_t^T \nabla_x u_s(X_s) \cdot dB_s + \int_t^T v_s^0(X_s) \cdot dB_s^0,
$$
\n(2.16)

with the short-hand notation $(\beta_s := \beta(s, X_s^0, X_s, \mu_s))_{0 \le s \le T}$.

By construction, see (2.2) , the integrand in the ds integral is non-negative. Taking expectation (which is licit thanks to the properties of u stated in Definition [2.10\)](#page-9-1), we deduce that $J_{\rm w}(\beta; \mathbb{P}_{(\alpha^0,\alpha)}^0) \geq \mathbb{E}_{(\alpha^0,\alpha)}^0 \otimes \mathbb{E}_{\cdot,\beta}[u_0(X_0)] =$ $\mathbb{E}^0_{(\alpha^0,\alpha)}[(u_0,\mu_0)]$, with the inequality becoming an equality when β is equal to α (see [\(2.15\)](#page-11-0) for the definition of the latter), i.e., $J_{\mathbf{w}}(\beta; \mathbb{P}_{(\alpha^0, \alpha)}^0) \geq J_{\mathbf{w}}(\alpha; \mathbb{P}_{(\alpha^0, \alpha)}^0)$.

Next, we prove item 2 in Definition [2.7.](#page-7-4) For this, we consider a new Markov feedback function β^0 such that the pair (β^0, α) is admissible. Then, we rewrite the forward-backward system [\(2.11\)](#page-8-2) solved by $(X^0, \mu, B^0, Y^0, Z^0)$ on $(\Omega_{\text{canon}}^0, \mathbb{P}_{(\alpha^0, \alpha)}^0)$ (see the first step) in the form

$$
dX_t^0 = \beta^0(t, X_t^0, \mu_t)dt + \sigma_0 d\tilde{B}_t^0
$$

\n
$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0))\right)dt + \left(\nabla_p H^0(X_t^0, Z_t^0) + \beta^0(t, X_t^0, \mu_t)\right) \cdot Z_t^0 dt
$$
\n
$$
+ \sigma_0 Z_t^0 \cdot d\tilde{B}_t^0,
$$
\n(2.17)

where

$$
\tilde{B}_t^0 := B_t^0 - \sigma_0^{-1} \left(\nabla_p H^0(X_t^0, Z_t^0) + \beta^0(t, X_t^0, \mu_t) \right) dt, \quad t \in [0, T]. \tag{2.18}
$$

¹The proof is as follows. For a time $h > 0$, we write $u_{t+h}(X_{t+h}) - u_t(X_t) = u_{t+h}(X_{t+h}) - u_{t+h}(X_t) + u_{t+h}(X_t) - u_t(X_t)$. And then, by standard Itô's formula, we get on the one hand, for any fixed $\omega^0 \in \Omega^0$, under $\mathbb{P}_{\omega^0,\beta}$,

$$
u_{t+h}(X_{t+h}) - u_{t+h}(X_t) = \frac{1}{2} \int_t^{t+h} \Delta_x u_{t+h}(X_s) ds + \int_t^{t+h} \nabla_x u_{t+h}(X_s) \cdot \left(\beta(s, X_s^0, X_s, \mu_s) ds + dB_s\right), \quad t \in [0, T-h].
$$

On the other hand, using the fact that X_t is independent of $(B_s^0 - B_t^0)_{t \le s \le t+h}$, we can formally replace x by X_t in the backward SPDE of $(u_s)_{t\leq s\leq t+h}$ and then get, with probability 1 under $\mathbb{P}^0_{(\alpha^0,\alpha)}\otimes\mathbb{P}_{\cdot,\beta}$,

$$
u_{t+h}(X_t) - u_t(X_t) = -\frac{1}{2} \int_t^{t+h} \Delta_x u_s(X_t) ds + \int_t^{t+h} H\big(X_t, \nabla_x u_s(X_t)\big) ds - \int_t^{t+h} f_s(X_s^0, X_t, \mu_s) ds + \sigma_0 \int_t^{t+h} v_s^0(X_t) \cdot dB_s^0.
$$

We then sum the last two displays over a mesh of stepsize h. We handle the Lebesgue integrals by using the fact that $\sup_{0 \leq t \leq T} ||u_t||_{\delta} \in L^{\infty}(\Omega^0, \mathbb{P}^0)$. In fact, the most difficult term to handle is the stochastic integral. We have, for any deterministic exponent $\eta \in (0,1)$, with $\eta < |\lambda| - d/2 - 1$,

$$
\mathbb{E}^{0} \int_{t}^{t+h} |v_{s}^{0}(X_{t}) - v_{s}^{0}(X_{s})|^{2} ds \leq \mathbb{E}^{0} \bigg[\Big(1 \wedge \sup_{|s-r| \leq h} |X_{s} - X_{r}|^{2\eta} \Big) \int_{t}^{t+h} ||v_{s}^{0}||_{\eta}^{2} ds \bigg].
$$

Using the fact that β satisfies [\(2.5\)](#page-6-3) together with the bound $\mathbb{E}^0 \int_{[0,T]} \|v_s^0\|_{\lfloor\delta\rfloor - d/2 - 1}^2 ds < \infty$, we can easily sum the above right-hand side over a mesh of stepsize h and let h tend to 0. We obtain [\(2.16\)](#page-12-1).

We justify in the fourth step below that we can apply Girsanov theorem to prove that \tilde{B}^0 is a Brownian motion under the probability measure

$$
\frac{\mathrm{d}\tilde{\mathbb{P}}_{(\alpha^0,\alpha)}^0}{\mathrm{d}\mathbb{P}_{(\alpha^0,\alpha)}^0} = \mathscr{E}_T\left(\sigma_0^{-1} \int_0^\cdot \left(\nabla_p H^0(X_t^0, Z_t^0) + \beta^0(t, X_t^0, \mu_t)\right) \cdot \mathrm{d}B_t^0\right). \tag{2.19}
$$

And then, we observe that the dt term in the backward equation appearing in [\(2.17\)](#page-12-2) is greater than $-[f_t^0(X_t^0, \mu_t) +$ $L^0(X_t^0, \beta^0(t, X_t^0, \mu_t))]$, which follows from the convexity of L^0 :

$$
L^{0}(X_{t}^{0}, \beta^{0}(t, X_{t}^{0}, \mu_{t}))
$$

\n
$$
\geq L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0})) + (\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0}) + \beta^{0}(t, X_{t}^{0}, \mu_{t})) \cdot \nabla_{\alpha}L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0}))
$$

\n
$$
= L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0})) - (\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0}) + \beta^{0}(t, X_{t}^{0}, \mu_{t})) \cdot Z_{t}^{0}.
$$

Observe now that the stochastic integral in [\(2.17\)](#page-12-2) has zero expectation under $\tilde{P}^0_{(\alpha^0,\alpha)}$ because $\mathbb{E}^0[|\int_{[0,T]}|Z_t^0|^2dt|^p]<\infty$ for any $p \ge 1$ and the change of measure in [\(2.19\)](#page-13-0) has a finite exponential moment, see [\[40,](#page-86-30) Theorem 2.2] and the fourth step below for the proof of the related BMO property). This suffices to say that

$$
Y_0^0 \le \tilde{\mathbb{E}}^0_{(\alpha^0,\alpha)} \bigg[g^0(X_T^0, \mu_T) + \int_0^T \Big(f^0(X_t^0, \mu_t) + L^0(X_t^0, \beta^0(t, X_t^0, \mu_t)) \Big) dt \bigg].
$$

Observing that the law of (X^0, μ) under $\tilde{P}^0_{(\alpha^0, \alpha)}$ is the same as the law of (X^0, μ) under $P^0_{(\beta^0, \alpha)}$ (because (β^0, α) is admissible, see Definition [2.2\)](#page-6-2), this shows that $J_w^0(\alpha^0, \alpha) = Y_0^0 \leq J_w^0(\beta^0, \alpha)$, as required.

Third Step. The last step is to prove uniqueness. To do so, we take an admissible pair (α^0, α) (different from the one constructed right above, but denoted in the same manner) satisfying the requirements of Definition [2.7.](#page-7-4) By item (ii) in the statement and with (X^0, μ, B^0) denoting again the canonical process on Ω_{canon}^0 , we know that, under the completion of $\mathbb{P}^0_{(\alpha^0,\alpha)}$ (still denoted $\mathbb{P}^0_{(\alpha^0,\alpha)}$), the process $(\bm{X}^0,\bm{\mu})$ is adapted to the augmentation of the filtration generated by $\bm{B}^0.$ This makes it possible to solve the backward stochastic HJB equation within the same class as in Definition [2.10](#page-9-1) (solvability of this equation is explained in Subsection [3.1\)](#page-18-1):

$$
d_t u_t(x) = \left(-\frac{1}{2}\Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t)\right)dt + \sigma_0 v_t^0(x) \cdot dB_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
u_T(x) = g(X_T^0, x, \mu_T).
$$
 (2.20)

Then, by expanding the duality product $((u_t, \mu_t))_{0 \le t \le T}$ (or by expanding $(u_t(X_t))_{0 \le t \le T}$ under $\mathbb{P}^0_{(\alpha^0, \alpha)} \otimes \mathbb{P}_{\cdot, \alpha}$), we can reproduce the verification argument used in the second step to show that necessarily, for Leb \times $\mathbb{P}^0_{(\alpha^0,\alpha)}$ -almost every $(t, \omega^0) \in [0, T] \times \Omega_{\text{canon}}^0$

$$
\alpha(t, X_t^0, x, \mu_t) = -\nabla_p H(X_t^0, \nabla_x u_t(x)), \quad x \in \mathbb{T}^d
$$

Combined with the second equation in [\(2.4\)](#page-6-1), this suffices to show that the pair (X^0, μ) together with (u, v^0) introduced in [\(2.20\)](#page-13-1) solves [\(2.12\)](#page-9-0) under $\mathbb{P}^0_{(\alpha^0,\alpha)}$.

We now handle the major player. As above, we can solve, under $\mathbb{P}^0_{(\alpha^0,\alpha)}$, the BSDE

$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, \alpha^0(t, X_t^0, \mu_t))\right)dt + \sigma_0 Z_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$

\n
$$
Y_T^0 = g^0(X_T^0, \mu_T).
$$
\n(2.21)

.

Under the standing assumption (see in particular (2.14) in item (i) of the statement) and by Lemma [2.9,](#page-8-1) the state equation [\(2.4\)](#page-6-1) has at most one weak solution, for any starting point in $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. By item (ii) in the statement, we know that equation [\(2.4\)](#page-6-1) is already assumed to have at least one strong solution, and this for any starting point in $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. By a straightforward modification of the Yamada-Watanabe theorem, we deduce that the state equation [\(2.4\)](#page-6-1) is uniquely strongly solvable for any starting point in $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. And then, by adapting the proof of [\[20,](#page-86-4) Proposition 1.31] (which is itself inspired from the remark below [\[56,](#page-87-12) Theorem 6.2.2]), strong uniqueness implies that the solution to [\(2.4\)](#page-6-1) forms a strong Markov process. Then, following once again [\[38,](#page-86-33)

Proposition 3.2 & Theorem 3.4], we can find a new Borel function $\varphi^0 : [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}^d \otimes \mathbb{R}^d$ such that, for any $s \in [0, T]$,

$$
\mathbb{P}^0\Big(\big\{Z_s^0 = \varphi^0(s, X_s^0, \mu_s)\big\}\Big) = 1.
$$
\n(2.22)

Recalling the assumption [\(2.5\)](#page-6-3), it is quite easy to prove that $\sup_{0 \le t \le T} |Y_t^0| \in L^{\infty}(\Omega^0, \mathbb{P}^0)$. And then, expanding $(|Y_t^0|^2)_{0 \le t \le T}$ by means of Itô's formula, we obtain $||\int_0^T Z_t^0 dB_t^0||_{BMO} < \infty$, and then $||\int_0^T \nabla_p H^0(X_t^0, Z_t^0) \cdot dB_t^0||_{BMO} <$ ∞.

We then define the tilted noise

$$
\tilde{B}_t^0 := B_t^0 + \sigma_0^{-1} \left(\nabla_p H^0(X_t^0, Z_t^0) + \alpha^0(t, X_t^0, \mu_t) \right) dt, \quad t \in [0, T]. \tag{2.23}
$$

We justify in the fourth step below that we can apply Girsanov theorem to prove that \tilde{B}^0 is a Brownian motion under the probability measure

$$
\frac{\mathrm{d}\tilde{\mathbb{P}}_{(\alpha^0,\alpha)}^0}{\mathrm{d}\mathbb{P}_{(\alpha^0,\alpha)}^0} = \mathscr{E}_T\bigg(-\sigma_0^{-1}\int_0^\cdot \left(\nabla_p H^0(X_t^0, Z_t^0) + \alpha^0(t, X_t^0, \mu_t)\right) \cdot \mathrm{d}B_t^0\bigg). \tag{2.24}
$$

And we rewrite the BSDE in (2.21) as

$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, \alpha^0(t, X_t^0, \mu_t))\right)dt - \left(\nabla_p H^0(X_t^0, Z_t^0) + \alpha^0(t, X_t^0, \mu_t)\right) \cdot Z_t^0 dt
$$

+ $\sigma_0 Z_t^0 \cdot d\tilde{B}_t^0, \quad t \in [0, T].$ (2.25)

As before, we observe from the (strict) convexity property of L^0 that

$$
L^{0}(X_{t}^{0}, \alpha^{0}(t, X_{t}^{0}, \mu_{t})) + (\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0}) + \alpha^{0}(t, X_{t}^{0}, \mu_{t})) \cdot Z_{t}^{0} \geq L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0})),
$$

with equality if and only if $\nabla_p H^0(X_t^0, Z_t^0) + \alpha^0(t, X_t^0, \mu_t) = 0$. We then have (the BMO condition, which can be transferred from the original probability measure $\mathbb{P}^0_{(\alpha^0,\alpha)}$ to the new probability measure $\tilde{\mathbb{P}}^0_{(\alpha^0,\alpha)}$ -see Theorem [\[40,](#page-86-30) Theorems 2.3 and 3.6]–, makes it possible to take expectation in (2.25))

$$
J_{\mathbf{w}}^{0}(\alpha^{0}, \alpha) = Y_{0}^{0} \ge \tilde{\mathbb{E}}_{(\alpha^{0}, \alpha)}^{0} \bigg[g^{0}(X_{T}^{0}, \mu_{T}) + \int_{0}^{T} \bigg(f_{t}(X_{t}^{0}, \mu_{t}) + L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0})) \bigg) \mathrm{d}t \bigg]. \tag{2.26}
$$

Under $\tilde{\mathbb{P}}^0_{(\alpha^0, \alpha)}, (X^0, \mu, \tilde{B}^0) = (X^0_t, \mu_t, \tilde{B}^0_t)_{0 \leq t \leq T}$ solves the system

$$
dX_t^0 = -\nabla_p H^0(X_t^0, \varphi^0(s, X_s^0, \mu_s))dt + d\tilde{B}_t^0,
$$

\n
$$
\partial_t \mu_t = \frac{1}{2}\Delta_{x_0}\mu_t - \text{div}_{x_0}(\alpha(t, X_t^0, \cdot, \mu_t)\mu_t), \quad t \in [0, T].
$$

Here, we observe that $\| \int_0^{\cdot} \nabla_p H^0(X_t^0, -\nabla_p H^0(X_t^0, \varphi^0(s, X_s^0, \mu_s)) \cdot d\tilde{B}_t^0 \|\text{BMO} = \| \int_0^{\cdot} \nabla_p H^0(X_t^0, Z_t^0) \cdot d\tilde{B}_t^0 \|\text{BMO} < \infty$ (again, this follows from Theorem $[40,$ Theorem 3.6). This provides one weak solution to the state equation (2.4) , driven by $\beta^{0}(t, x_{0}, \mu) := -\nabla_{p}H^{0}(x_{0}, \varphi^{0}(t, x_{0}, \mu))$, that satisfies the BMO condition [\(2.5\)](#page-6-3). By Lemma [2.9,](#page-8-1) this weak solution is necessarily unique and, therefore, the pair (β^0, α) is admissible in the sense of Definition [2.2.](#page-6-2) Then, the right-hand side in [\(2.26\)](#page-14-1) coincides with $J_w^0(\beta^0, \alpha)$,

By item 2 in Definition [2.7,](#page-7-4) the inequality in [\(2.26\)](#page-14-1) must become an equality and then, for almost every $(t, \omega^0) \in$ $[0, T] \times \Omega^0$ under the measure Leb $\times \mathbb{P}^0$, it holds $\nabla_p H^0(X_t^0, Z_t^0) + \alpha^0(t, X_t^0, \mu_t) = 0$, which proves that (X^0, μ) coincides with the solution of (2.11) – (2.12) .

Fourth Step. We now justify the application of Girsanov theorem in (2.17) , (2.18) and (2.19) . We start from (2.4) for an admissible pair (α^0, α) as in the statement of Definition [2.2,](#page-6-2) with α satisfying [\(2.9\)](#page-8-3) (which is the case in [\(2.15\)](#page-11-0) because of (2.13)). Then, Lemma [2.9](#page-8-1) says that the BMO condition (2.10) is satisfied under \overline{P}^0 . This observation applies here to both α^0 as in [\(2.15\)](#page-11-0) and $\alpha^0 \equiv \beta^0$ (with β^0 as in [\(2.17\)](#page-12-2)). In particular, choosing now α^0 as in [\(2.15\)](#page-11-0), the process $(\int_{[0,t]}[\beta^0(s,X_s^0,\mu_s)-\alpha^0(s,X_s^0,\mu_s)]\cdot dB_s^0)_{0\leq t\leq T}$ is BMO under $\bar{\mathbb{P}}^0$. Let now

$$
\check{\mathbb{P}}^0 := \mathscr{E}_T \biggl(\int_0^\cdot \alpha^0 \bigl(s,X_s^0,\mu_s\bigr) \cdot \mathrm{d} B_s^0 \biggr) \cdot \overline{\mathbb{P}}^0.
$$

16

By [\[40,](#page-86-30) Theorem 3.6], we know that $(\int_{[0,t]} [\beta^0(s, X_s^0, \mu_s) - \alpha^0(s, X_s^0, \mu_s)] \cdot d\check{B}_s^0)_{0 \le t \le T}$ is BMO under $\check{\mathbb{P}}^0$, where $\check{B}^0 =$ $(\check{B}_t^0 := B_t^0 - \int_0^t \alpha^0(s,X_s^0,\mu_s) \mathrm{d}s)_{0 \le t \le T}$ is a Brownian motion under $\check{\mathbb{P}}^0$. Since $\check{\mathbb{P}}^0 \circ (\mathbf{X}^0,\mu, \check{\mathbf{B}}^0)^{-1} = \mathbb{P}^0_{(\alpha^0,\alpha)},$ this shows that $(\int_{[0,t]} [\beta^0(s,X_s^0,\mu_s) - \alpha^0(s,X_s^0,\mu_s)] \cdot dB_s^0)_{0 \le t \le T}$ is BMO under $\mathbb{P}^0_{(\alpha^0,\alpha)}$, which suffices to apply Girsanov theorem in (2.17) , (2.18) and (2.19) .

We now proceed in a similar manner to justify the Girsanov transformation in (2.23) and (2.24) . In fact, it suffices to apply the same argument as above but with α^0 as in the third step, see [\(2.21\)](#page-13-2), and with $\beta^0(t, x_0, \mu)$ = $-\nabla_p H^0(x_0, \varphi^0(t, x_0, \mu))$, see [\(2.22\)](#page-14-4). 口

2.6. System of master equations

Following the analysis performed in [\[13](#page-86-27)], we associate with the Major/Minor MFG a system of master equations. Formally, it reads as a pair of two equations for the value V^0 to the major player and the value V to the minor player.

The equation for V^0 reads

$$
\partial_t V^0(t, x_0, \mu) + \frac{1}{2} \sigma_0^2 \Delta_{x_0} V^0(t, x_0, \mu) - H^0(x_0, \nabla_{x_0} V^0(t, x_0, \mu)) + f_t^0(x_0, \mu) \n+ \int_{\mathbb{T}^d} \left\{ \frac{1}{2} \text{div}_y (\partial_\mu V^0(t, x_0, \mu, y)) - \partial_\mu V^0(t, x_0, \mu, y) \cdot \nabla_p H(y, \nabla_x V(t, x_0, y, \mu)) \right\} d\mu(y) = 0, \tag{2.27}
$$
\n
$$
V^0(T, x_0, \mu) = g^0(x_0, \mu),
$$

for $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. The equation for V is

$$
\partial_t V(t, x_0, x, \mu) + \frac{1}{2} \Delta_x V(t, x_0, x, \mu) + \frac{1}{2} \sigma_0^2 \Delta_{x_0} V(t, x_0, x, \mu) - H(x_0, x, \nabla_x V(t, x_0, x, \mu)) + f_t(x_0, x, \mu) \n- \nabla_p H^0(x_0, \nabla_{x_0} V^0(t, x_0, \mu)) \cdot \nabla_{x_0} V(t, x_0, x, \mu) \n+ \int_{\mathbb{T}^d} \left\{ \frac{1}{2} \text{div}_y(\partial_\mu V(t, x_0, x, \mu, y)) - \partial_\mu V(t, x_0, x, \mu, y) \cdot \nabla_p H(y, \nabla_x V(t, x_0, y, \mu)) \right\} d\mu(y) = 0, \tag{2.28}
$$
\n
$$
V(T, x_0, x, \mu) = g(x_0, x, \mu), \tag{2.29}
$$

for $(t, x_0, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$.

The following statement clarifies the connection between (2.27) – (2.28) and (2.11) – (2.12) .

Proposition 2.15. Assume that the master equation [\(2.27\)](#page-15-0)–[\(2.28\)](#page-15-1) has a classical solution (V^0, V) in the sense that

- $I.$ $(t, x_0, \mu) \mapsto (\partial_t V^0(t, x_0, \mu), \nabla_{x_0} V^0(t, x_0, \mu), \nabla_{x_0}^2 V^0(t, x_0, \mu))$ *is continuous on* $[0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ (with the *latter factor being equipped with any distance metricizing weak convergence on* $P(\mathbb{T}^d)$, for instance \mathbb{W}_1); $(t, x_0, \mu, y) \mapsto (\partial_\mu V^0(t, x_0, \mu, y), \nabla_y \partial_\mu V^0(t, x_0, \mu, y))$ *is continuous on* $[0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$;
- 2. $(t, x_0, x, \mu) \mapsto (\partial_t V(t, x_0, x, \mu), \nabla_{x_0} V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu), \nabla_{x_0}^2 V(t, x_0, x, \mu), \nabla_x^2 V(t, x_0, x, \mu))$ *is con-* \mathcal{L} *tinuous on* $[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$; $(t, x_0, x, \mu, y) \mapsto (\partial_\mu V(t, x_0, x, \mu, y), \nabla_y \partial_\mu V(t, x_0, x, \mu, y))$ *is continuous on* $[0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$.

Assume also that $(t, x_0, \mu) \mapsto \nabla_{x_0} V^0(t, x_0, \mu)$ and $(t, x_0, x, \mu) \mapsto (\nabla_{x_0} V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu))$ are Lipschitz *continuous with respect to* (x_0, μ) and (x_0, x, μ) *respectively (using the distance* \mathbb{W}_1 to handle the argument μ) and that the initial condition X_0^0 (in Subsection [2.1\)](#page-2-1) is square-integrable. Then, the triplets $(X_t^0, Y_t^0, Z_t^0)_{0 \le t \le T}$ and $(\mu_t, u_t, v_t^0)_{0 \le t \le T}$ *obtained by solving, on a product structure comprising two filtered probability spaces* $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$ and $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ equipped with two Brownian motions $(B_t^0)_{0 \le t \le T}$ and $(B_t)_{0 \le t \le T}$ with values in \mathbb{R}^d , the (coupled for*ward) equations*

$$
dX_t^0 = -\nabla_p H^0(X_t^0, \nabla_{x_0} V^0(t, X_t^0, \mu_t))dt + \sigma_0 dB_t^0, \quad t \in [0, T],
$$

\n
$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x (\nabla_p H(x, \nabla_x V(t, X_t^0, x, \mu_t)) \mu_t) = 0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$
\n(2.29)

and then by letting

$$
Y_t^0 := V^0(t, X_t^0, \mu_t), \quad Z_t^0 := \nabla_{x_0} V^0(t, X_t^0, \mu_t), \quad t \in [0, T],
$$

$$
u_t(x) := V(t, X_t^0, x, \mu_t), \quad v_t^0(x) := \nabla_{x_0} V(t, X_t^0, x, \mu_t), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$
 (2.30)

are solutions of [\(2.11\)](#page-8-2)*–*[\(2.12\)](#page-9-0)*.*

As the proof shows (see (2.32)), the solutions to (2.11) – (2.12) that we obtain in this manner just supply us with 'true' martingales in the two equations. Notice that, to simplify, we do not prove that the the triplets $(X_t^0, Y_t^0, Z_t^0)_{0 \le t \le T}$ and $(\mu_t, u_t, v_t^0)_{0 \le t \le T}$ satisfy all the conditions in Definition [2.10,](#page-9-1) as this would be useless at this stage of the paper. Obviously, this would require further assumptions on V^0 and V.

Our main statement regarding the solvability of the master equation is

Theorem 2.16. *In addition to Assumption* (A)*, assume that*

- i. the coefficient $(t, x_0, \mu) \mapsto f_t^0(x_0, \mu)$ is Hölder continuous in time, uniformly in (x_0, μ) ; the coefficient $(t, x_0, x, \mu) \mapsto$ $f_t(x_0, x, \mu)$ *is Hölder continuous in time, uniformly in* (x_0, x, μ) *;*
- ii. $x_0 \mapsto g^0(x_0,\mu)$ has Hölder continuous second-order derivatives, uniformly in μ ; $x_0 \mapsto g(x_0,x,\mu)$ has Hölder con*tinuous second-order derivatives, uniformly in* (x, μ) *.*

Then, for any $T > 0$ and for $\sigma_0 \ge \sigma_0^*(T) \in (0, +\infty)$ *(with the latter being defined as in the statement of Theorem* [2.14\)](#page-10-0), the system (2.27) – (2.28) admits a unique solution in the class of functions $(V⁰, V)$ that satisfy items (1) and (2) in the statement of Proposition [2.15](#page-15-2) and such that: (3) $(t, x_0, \mu) \mapsto \nabla_{x_0} V^0(t, x_0, \mu)$ and $(t, x_0, x, \mu) \mapsto$ $(\nabla_{x_0} V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu))$ are Lipschitz continuous with respect to (x_0, μ) and (x_0, x, μ) respectively (us*ing the distance* \mathbb{W}_1 *to handle the argument* μ *);* (4) $(t, x_0, \mu) \mapsto (V^0(t, x_0, \mu), \nabla_{x_0} V^0(t, x_0, \mu))$ *is globally bounded,* $(t, x_0, \mu) \mapsto (\|V(t, x_0, \cdot, \mu)\|_{\infty}, \|\nabla_{x_0} V(t, x_0, \cdot, \mu)\|_{\infty})$ *is globally bounded for any* $\pi \in [1, |\Lambda| - (d/2 + 1)] \setminus \mathbb{N}$.

If Assumption (B) *is also in force, then existence and uniqueness hold true on any interval* [0, T]*,* $T > 0$ *, and for any* $\sigma_0 \geq \sigma_0^*$ (with the latter being defined as in the statement of Theorem [2.14\)](#page-10-0).

The proof of Theorem [2.16](#page-16-0) is deferred to Section [4,](#page-30-0) see Theorem [4.12](#page-44-0) for a refined version. We now turn to the proof of Proposition [2.15:](#page-15-2)

Proof of Proposition [2.15.](#page-15-2) To establish the solvability of [\(2.29\)](#page-15-3), we consider the conditional McKean-Vlasov equation

$$
dX_t^0 = -\nabla_p H^0\Big(X_t^0, \nabla_{x_0} V^0(t, X_t^0, \mathcal{L}^0(X_t))\Big)dt + \sigma_0 dB_t^0,
$$

$$
dX_t = -\nabla_p H\Big(X_t^0, X_t, \nabla_x V\big(t, X_t^0, X_t, \mathcal{L}^0(X_t)\big)\Big)dt + dB_t, \quad t \in [0, T],
$$

with the same initial conditions X_0^0 and X_0 as in Subsection [2.1.](#page-2-1)

Because $\nabla_{x_0} V^0$ and $\nabla_x V$ are jointly continuous in all the arguments and Lipschitz continuous with respect to (x_0, μ) and (x_0, x, μ) respectively (Lipschitz continuity in the argument μ holding true with respect to \mathbb{W}_1), the above system has a unique solution, see [\[20](#page-86-4), Proposition 2.8]. It satisfies

$$
\mathbb{E}^{0}\Big[\sup_{0\leq t\leq T}|X_{t}^{0}|^{2}\Big]<\infty.\tag{2.31}
$$

(A similar bound holds true for $\sup_{0 \le t \le T} |X_t|^2$ under $\mathbb{E}^0 \mathbb{E}$ when X is implicitly regarded as being \mathbb{R}^d -valued, but this bound has little interest since X is regarded as being \mathbb{T}^d -valued.) Letting $(\mu_t := \mathcal{L}^0(X_t))_{0 \le t \le T}$, this makes it possible to define $(Y_t^0, Z_t^0)_{0 \le t \le T}$ and $(u_t, v_t^0)_{0 \le t \le T}$ as in [\(2.30\)](#page-15-4). Combining the integrability condition [\(2.31\)](#page-16-3) with the regularity properties of $\nabla_{x_0} V^{\overline{0}}, \overline{\nabla}_{x_0} V$ and $\nabla_x V$, we get

$$
\mathbb{E}^{0}\Big[\sup_{0\leq t\leq T}\big(|Z_{t}^{0}|^{2}+\sup_{x\in\mathbb{T}^{d}}|\nabla_{x}V(t,X_{t}^{0},x,\mu_{t})|^{2}+\sup_{x\in\mathbb{T}^{d}}|v_{t}^{0}(x)|^{2}\big)\Big]<\infty,
$$
\n(2.32)

The derivation of the forward equation in (2.12) is straightforward. As for the two backward equations in (2.11) and [\(2.12\)](#page-9-0), they are obtained by combining the chain rule proved in Appendix, see Proposition [6.1,](#page-83-0) with the two PDEs [\(2.27\)](#page-15-0) and (2.28) . The bound (2.32) shows that the martingale terms are 'true' martingales. \Box

2.7. About Fokker-Planck equations with random coefficients

The purpose of this subsection is to prove Lemmas [2.4](#page-7-3) and [2.9.](#page-8-1)

Proof of Lemma [2.4.](#page-7-3) We call \tilde{P} the (completion of the) probability measure on Ω_{canon} under which the canonical process (X, B) satisfies: (i) $\tilde{P} \circ X_0^{-1} = \mu$; (ii) B is an \mathbb{F} -Brownian motion starting from 0; (iii) $X_t - X_0 = B_t$ for all $t \in [0, T]$. For $\omega \in \Omega^0$, we then let

$$
\tilde{\mathbb{P}}_{\omega^0} := \mathscr{E}_T \bigg(\int_0^{\cdot} \alpha(t, X_t^0(\omega^0), X_t, \mu_t(\omega^0)) \cdot \mathrm{d}B_t \bigg) \cdot \tilde{\mathbb{P}}.
$$

Since α is bounded, $\tilde{\mathbb{P}}_{\omega^0}$ is a probability measure. It is standard to check that, under $\tilde{\mathbb{P}}_{\omega^0}$, the process $(\tilde{B}_t := B_t - \int_{\alpha}^t \alpha(s, X^0(\omega^0), X_s, \mu_s(\omega^0))ds)_{0 \le t \le T}$ is an F-Brownian motion startin $\int_0^t \alpha(s, X_s^0(\omega^0), X_s, \mu_s(\omega^0))ds)_{0 \le t \le T}$ is an **F**-Brownian motion starting from 0, X_0 is distributed according to μ and, for any $t \in [0, T]$,

$$
dX_t = \alpha(t, X_t^0(\omega^0), X_t, \mu_t(\omega^0))dt + d\tilde{B}_t, \quad t \in [0, T].
$$
\n(2.33)

We then define

$$
\mathbb{P}_{\omega^0} := \tilde{\mathbb{P}}_{\omega^0} \circ \left(\mathbf{X}, \tilde{\mathbf{B}} \right)^{-1}.
$$

Using Fubini's theorem, it is easy to prove that, for any event C of $\mathcal{B}(\Omega_{\text{canon}})$, the mapping $\omega^0 \mapsto \mathbb{P}_{\omega^0}(C)$ is measurable, which proves the measurability of the mapping $\Omega_{\text{canon}}^0 \ni \omega^0 \mapsto \mathbb{P}_{\omega^0}$.

Using the fact that the SDE (2.7) (with a prescribed initial condition) is uniquely strongly (and thus weakly) solvable for any $\omega^0 \in \Omega_{\text{canonical}}^0$ (because α is bounded), we get that \mathbb{P}_{ω^0} is unique. 口

Proof of Lemma [2.9.](#page-8-1) *First Step.* With $x^0 := (x_t^0)_{0 \le t \le T}$ an element of $C([0,T], \mathbb{R}^d)$, we associate the Fokker-Planck equation

$$
\partial_t \tilde{\mu}_t^{\mathbf{x}^0} = \tfrac{1}{2} \Delta_x \tilde{\mu}_t^{\mathbf{x}^0} - \mathrm{div}_x \Big(\alpha \big(t, x_t^0, \cdot, \tilde{\mu}_t^{\mathbf{x}^0} \big) \tilde{\mu}_t^{\mathbf{x}^0} \Big), \quad t \in [0, T],
$$

with μ_0 as initial condition (at time 0).

By [\[42](#page-87-13)] and Remark [2.5,](#page-7-6) we know that, for any $x^0 \in C([0, T], \mathbb{R}^d)$, the Fokker-Planck equation right above admits a unique solution in the same weak sense as in Remark [2.3,](#page-7-0) which can be obtained by iterating the mapping, denoted $\Phi(\mathbf{x}^0, \cdot)$, that sends an element $\boldsymbol{\nu} = (\nu_t)_{0 \le t \le T} \in C([0, T], \mathcal{P}(\mathbb{T}^d))$ onto the solution $\boldsymbol{\mu} = (\mu_t)_{0 \le t \le T}$ of the equation

$$
\partial_t \mu_t = \frac{1}{2} \Delta_x \mu_t - \text{div}_x \big(\alpha(t, x_t^0, \cdot, \nu_t) \mu_t \big), \quad t \in [0, T],
$$

with μ_0 as initial condition. By Lemma [2.4,](#page-7-3) $(\mu_t)_{0 \le t \le T}$ is the flow of marginal laws of the process X under the tilted measure

$$
\mathbb{P}_{\mathbf{x}^0}:=\mathscr{E}_T\biggl(\int_0^\cdot\alpha\bigl(t,x_t^0,X_0+B_t,\nu_t\bigr)\cdot\mathrm{d} B_t\biggr)\cdot\tilde{\mathbb{P}},
$$

where \tilde{P} is the same probability measure on Ω_{canon} as in the proof of Lemma [2.4.](#page-7-3) Proceeding as in the latter proof, we easily deduce that μ , seen as an element of $\mathcal{C}([0,T], \mathcal{P}(\mathbb{T}^d))$, is the image by a measurable function Φ of the pair (x^0, ν) , seen as an element of $C([0,T], \mathbb{R}^d) \times C([0,T], \mathcal{P}(\mathbb{T}^d))$, i.e., $\boldsymbol{\mu} = \Phi(\boldsymbol{x}^0, \boldsymbol{\nu})$. And then, writing

$$
\boldsymbol{\mu}^{\boldsymbol{x}^0} = \lim_{n \to \infty} \left[\Phi(\boldsymbol{x}^0, \cdot) \right]^{\circ n} (\boldsymbol{\nu}^0),
$$

for an arbitrarily fixed element $v^0 \in C([0,T], \mathcal{P}(\mathbb{T}^d))$, we deduce that the mapping $\mathcal{C}([0,T], \mathbb{R}^d) \ni x^0 \mapsto \mu^{x^0} \in$ $\mathcal{C}([0,T], \mathcal{P}(\mathbb{T}^d))$ is measurable.

Considering on an arbitrary probability space a d-dimensional Brownian motion $(\bar{B}_t^0)_{0 \le t \le T}$ and replacing x^0 by $(x_0 + \bar{B}_t^0)_{0 \le t \le T}$, we get the existence of a (measurable) solution to

$$
\begin{aligned} \mathrm{d}\bar{X}_t^0 &= \mathrm{d}\bar{B}_t^0, \\ \partial_t \bar{\mu}_t &= \tfrac{1}{2} \Delta_x \bar{\mu}_t - \mathrm{div}_x \big(\alpha(t, \bar{X}_t^0, \cdot, \bar{\mu}_t) \bar{\mu}_t \big), \quad t \in [0, T], \end{aligned}
$$

with (x_0, μ) as initial condition (at time 0). In fact, the solution is (replacing the interval [0, T] by the interval [0, S], for S running between 0 and T in the above measurability argument) progressively-measurable with respect to the filtration generated by \bar{B}^0 . Also, it is pathwise unique. Below, we call $\bar{\mathbb{P}}^0$ the law of $(\bar{X}^0,\bar{\mu},\bar{B}^0)$ on Ω_canon^0 . This proves (b) in the statement.

Second Step. Assume now that the BMO condition [\(2.10\)](#page-8-0) is satisfied under \bar{P}^0 . Under the latter probability, the canonical process $(\bm{X}^0,\bm{\mu},\bm{B}^0)$ on Ω_{canon}^0 satisfies

$$
dX_t^0 = dB_t^0,
$$

\n
$$
\partial_t \mu_t = \frac{1}{2} \Delta_x \mu_t - \text{div}_x \big(\alpha(t, X_t^0, \cdot, \mu_t) \mu_t \big), \quad t \in [0, T],
$$

with (x_0, μ) as initial condition (at time 0). We introduce the tilted measure

$$
\tilde{\mathbb{P}}^0 := \mathscr{E}_T \bigg(\int_0^\cdot \alpha^0(t,X_t^0,\mu_t) \cdot \mathrm{d}B_t^0 \bigg) \cdot \bar{\mathbb{P}}^0,
$$

which is a probability measure thanks to the BMO condition [\(2.5\)](#page-6-3). Letting $(\tilde{B}_t^0 := B_t^0 - \int_0^t \alpha^0(s, X_s^0, \mu_s) ds)_{0 \le t \le T}$, we have

$$
dX_t^0 = \alpha^0(t, X_t^0, \mu_t)dt + d\tilde{B}_t^0,
$$

\n
$$
\partial_t \mu_t = \frac{1}{2} \Delta_{x_0} \mu_t - \text{div}_{x_0} (\alpha(t, X_t^0, \cdot, \mu_t) \mu_t), \quad t \in [0, T],
$$

with (x_0, μ) as initial condition (at time 0), and \tilde{B}^0 is a Brownian motion under $\tilde{\mathbb{P}}^0$. It then remains to let $\mathbb{P}^0 := \tilde{\mathbb{P}}^0$ \circ $(X^0, \mu, \tilde{B}^0)^{-1}$. It is a probability measure on Ω_{canon}^0 and it satisfies the requirements of Lemma [2.9.](#page-8-1) Notice in particular that item (iv) in Definition [2.2](#page-6-2) follows from [\[40,](#page-86-30) Theorems 2.3 and 3.3]. This proves the existence part in item (c) in the statement.

Third Step. Uniqueness is proven in a similar manner. Assuming that we are given a probability measure (still denoted) P⁰ satisfying items (i), (ii), (iii) and (iv) in Definition [2.2](#page-6-2) and considering without any loss of generality its completion, we introduce the tilted measure

$$
\tilde{\mathbb{P}}^0 := \mathscr{E}_T \bigg(- \int_0^\cdot \alpha^0(t,X^0_t,\mu_t) \cdot \mathrm{d} B^0_t \bigg) \cdot \mathbb{P}^0,
$$

which a probability measure thanks to the BMO condition (2.5) under \mathbb{P}^0 .

Letting $(\tilde{B}_t^0 := X_t^0 - x_0)_{0 \le t \le T}$, we have

$$
dX_t^0 = d\tilde{B}_t^0,
$$

\n
$$
\partial_t \mu_t = \frac{1}{2} \Delta_{x_0} \mu_t - \text{div}_{x_0} \left(\alpha(t, X_t^0, \cdot, \mu_t) \mu_t \right), \quad t \in [0, T],
$$

and \tilde{B}^0 is a Brownian motion under \tilde{P}^0 . And by the uniqueness result established in the first step, we see that the law of (X^0, μ, \tilde{B}^0) under $\tilde{\mathbb{P}}^0$ is equal to the probability $\tilde{\mathbb{P}}^0$ constructed in the second step. By [\[40,](#page-86-30) Theorems 2.3 and 3.3] again, the BMO condition (2.10) is satisfied under \bar{P}^0 . Also, we have

$$
\mathbb{P}^0 = \mathscr{E}_T \left(\int_0^{\cdot} \alpha^0(t, X_t^0, \mu_t) \cdot d\tilde{B}_t^0 \right) \cdot \tilde{\mathbb{P}}^0.
$$

In the end, \mathbb{P}^0 , which is tautologically equal to the law of (X^0, μ, B^0) under \mathbb{P}^0 , can be regarded as the law of $(X^0, \mu, (\tilde{B}_t^0 - \int_0^{\cdot} \alpha^0(s, X_s^0, \mu_s) ds)_{0 \le t \le T}$ under $\mathscr{E}_T (\int_0^{\cdot} \alpha^0(t, X_t^0, \mu_t) \cdot d\tilde{B}_t^0) \cdot \tilde{\mathbb{P}}^0$. Since $\tilde{\mathbb{P}}^0 \circ (X^0, \mu, \tilde{B}^0)^{-1} =$ $\overline{P}^0 \circ (\mathbf{X}^0, \mu, \mathbf{B}^0)^{-1}$, we deduce that $\overline{P^0}$ coincides with the law of $(\mathbf{X}^0, \mu, (B_t^0 - \int_0^{\cdot} \alpha^0(s, X_s^0, \mu_s) ds)_{0 \le t \le T})$ under $\mathscr{E}_T\left(\int_0^{\cdot} \alpha^0(t, X_t^0, \mu_t) \cdot dB_t^0\right) \cdot \bar{\mathbb{P}}^0$. This is exactly the construction achieved in the previous step. \Box

3. A priori estimates for the forward-backward system

The objective of this section is to obtain a series of *a priori* estimates for the solution(s) to the forward-backward system (2.11) – (2.12) , when posed on an arbitrary filtered probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}^0)$ satisfying the usual conditions and equipped with a Brownian motion $(B_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d , with \mathbb{F}^0 being generated by \mathcal{F}_0^0 and \mathbf{B}^0 .

3.1. HJB equation for the minor player

The main result of this subsection concerns the regularity of the solution to the stochastic HJB equation in [\(2.12\)](#page-9-0). We proceed very much as in the monograph [\[14](#page-86-2)]. We also refer to [\[29](#page-86-35)] for a related analysis but in Sobolev (instead of Hölder) spaces.

Throughout this subsection, we fix an \mathbb{F}^0 -adapted continuous path $X^0 = (X_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d (not neces-sarily solving the forward equation in [\(2.11\)](#page-8-2)) and an \mathbb{F}^0 -adapted continuous path $\mu = (\mu_t)_{0 \le t \le T}$ with values in $\mathcal{P}(\mathbb{T}^d)$ (not necessarily solving the forward equation in (2.12)). With the two of them, we associate the (backward) stochastic

$$
d_t u_t(x) = \left(-\frac{1}{2}\Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t)\right)dt + dm_t(x), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
u_T(x) = g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d,
$$
 (3.1)

where $(m_t(x))_{0 \le t \le T}$ is an \mathbb{F}^0 -martingale for any $x \in \mathbb{T}^d$.

The class within which the equation (3.1) is solved is clarified in the following statement, which is taken from [\[14,](#page-86-2) Proposition 4.3.8]:

Lemma 3.1. *Under Assumption* (A) *and within the framework described above, the equation* [\(3.1\)](#page-19-0) *has a unique solution* $(\mathbf{u},\mathbf{m})=(u_t,m_t)_{0\leq t\leq T}$, such that

- *1.* $(u_t)_{0 \le t \le T}$ is an \mathbb{F}^0 -adapted process with values in $\mathcal{C}^{\lambda}(\mathbb{T}^d)$, with continuous paths in $\mathcal{C}^{\lambda}(\mathbb{T}^d)$ for any $\lambda < \lambda$ satis*fying* $\sup_{t \in [0,T]} ||u_t||_{\delta} \in L^{\infty}(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ *.*
- *2.* $(m_t)_{0 \le t \le T}$ is an \mathbb{F}^0 -adapted process with values in $\mathcal{C}^{3-2}(\mathbb{T}^d)$, with continuous paths in $\mathcal{C}^n(\mathbb{T}^d)$ for any $n < 3-2$, $satisfying \sup_{t\in[0,T]}\|m_t\|_{\infty-2}\in L^{\infty}(\Omega^0,\mathcal{F}^0,\mathbb{P}^0)$, with $m_0\equiv 0$, and with $(m_t(x))_{0\leq t\leq T}$ being an \mathbb{F}^0 -martingale *for any* $x \in \mathbb{T}^d$.

In fact, uniqueness holds in a wider class of solutions (u, m) *for which the above holds true with respect to some* $s' > 2$ *in lieu of* δ *.*

Proof. The result mainly follows from [\[14](#page-86-2), Proposition 4.3.8], with two main differences: (i) one must here handle a quadratic HJB equation, whilst the nonlinearity is of linear growth in $[14]$; (ii) one here claims that continuity of u holds in $\mathcal{C}^{\pi}(\mathbb{T}^d)$ for any $\pi < \infty$, whilst continuity in [\[14\]](#page-86-2) is obtained up to the order [Δ]; (iii) as explained in the third item of Remark [2.11,](#page-9-2) the pair (u,m) is directly seen as a random variable with values in non-integer Hölder spaces, which are not separable. Generally speaking, the argument to treat (ii) is as follows: using the fact that solutions are proven to satisfy $\sup_{t\in[0,T]}\|u_t\|_{\delta}\in L^{\infty}(\Omega^0,\mathcal{F}^0,\mathbb{P}^0)$, continuity with values in any $\mathcal{C}^{\gamma}(\mathbb{T}^d)$, with $\gamma<\delta$, follows from the fact that the embedding from $\mathcal{C}^{\wedge}(\mathbb{T}^d)$ to $\mathcal{C}^{\wedge}(\mathbb{T}^d)$ is compact. As for (iii), we follow the outline given in Remark [2.11.](#page-9-2) Once *u* is known to be an \mathbb{F}^0 -adapted (continuous) process with values in $\mathcal{C}^{\lfloor \Delta \rfloor - 1}(\mathbb{T}^d)$, the same argument based on Schauder's estimates as the one used in [\[14\]](#page-86-2) permits to show that, for any $t \in [0, T)$, u_t takes values in $\mathcal{C}^{\delta}(\mathbb{T}^d)$ and m_t in $C^{s'-2}(\mathbb{T}^d)$, for a certain $s' > s$. Then, using the same argument as in the proof of [\[49,](#page-87-9) Proposition II.2, (1b)], we deduce that, for any $t \in [0, T)$, u_t and m_t are \mathcal{F}_t^0 -Bochner measurable with values in $\mathcal{C}^{\delta}(\mathbb{T}^d)$ and $\mathcal{C}^{\delta-2}(\mathbb{T}^d)$ respectively. At time $t = T$, the field $x \mapsto g(X_T^0, x, \mu_T)$ can be proven to be *Bochner* measurable: X_T^0 and μ_T are the almost sure limits of simple random variables with values in \mathbb{R}^d and $\mathcal{P}(\mathbb{T}^d)$ respectively. Using the fact that the function $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto g(x_0, \cdot, \mu) \in \mathcal{C}^{\delta}(\mathbb{T}^d)$ is continuous, we deduce that u_T is the almost sure limit of simple random variables with values in $C^{\delta}(\mathbb{T}^d)$, which shows *Bochner* measurability. It remains to prove that m_T is also *Bochner* measurable with values in $\mathcal{C}^{\lambda-2}(\mathbb{T}^d)$. In fact, it suffices to prove that the integral from 0 to T of the driver in the backward equation of [\(3.1\)](#page-19-0) is *Bochner* measurable with values in $\mathcal{C}^{b-2}(\mathbb{T}^d)$. By [\[49,](#page-87-9) Proposition II.2, (2)], it suffices to prove the same property but for the integral from 0 to $T - \varepsilon$, and this for any $\varepsilon \in (0, T)$. The latter is a mere consequence of the fact that the integral from 0 to $T - \varepsilon$ takes in fact values in $\mathcal{C}^{s'-2}(\mathbb{T}^d)$ for a certain $s' > s$.

In order to tackle (i), we proceed by considering first a truncated version of the Hamiltonian, namely we consider a function H^R on \mathbb{R}^d such that

$$
H^{R}(x, p) = H(x, p), \quad \text{if } |p| \leq R \, ; \quad \sup_{x \in \mathbb{T}^d} \sup_{p \in \mathbb{R}^d} |\nabla_p H^{R}(x, p)| < \infty,
$$
\n
$$
\sup_{x \in \mathbb{T}^d} \sup_{p \in \mathbb{R}^d} |\nabla_x H^{R}(x, p)| \leq \sup_{x \in \mathbb{T}^d} \sup_{p \in \mathbb{R}^d} |\nabla_x H(x, p)| + 1.
$$

Then, [\[14,](#page-86-2) Proposition 4.3.8] applies to the following equation:

$$
d_t u_t^R(x) = \left(-\frac{1}{2}\Delta_x u_t^R(x) + H^R(x, \nabla_x u_t^R(x)) - f_t(X_t^0, x, \mu_t)\right) dt + dm_t^R(x), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
u_T^R(x) = g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d,
$$
\n(3.2)

and supplies us with the existence of a unique solution (u^R, m^R) satisfying the prescriptions of items 1 and 2 in the statement of Lemma [3.1.](#page-19-1)

In order to pass from the equation [\(3.2\)](#page-19-2) to the original equation [\(3.1\)](#page-19-0), it suffices to show that $|\nabla_x u_t^R|$ can be bounded by a deterministic constant independent of R , which can be done by the classical Bernstein argument. Differentiating with respect to x (which is licit from the results in $[14,$ Proposition 4.3.8]), we have

$$
d_t \partial_{x_i} u_t^R(x) = \left(-\frac{1}{2} \Delta_x \partial_{x_i} u_t^R(x) + \nabla_p H^R(x, \nabla_x u_t^R(x)) \cdot \partial_{x_i} \nabla_x u_t^R(x) + \partial_{x_i} H^R(x, \nabla_x u_t^R(x))\right)
$$

$$
- \partial_{x_i} f_t(X_t^0, x, \mu_t)\right) dt + d\partial_{x_i} m_t^R(x), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
\partial_{x_i} u_T^R(x) = \partial_{x_i} g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d,
$$
 (3.3)

for any coordinate $i \in \{1, \dots, d\}$. Then, we can interpret the equation as a backward stochastic transport diffusion equation, for which we have a maximum principle. In turn, we get that

$$
|\partial_{x_i} u_t^R(x)| \leq \|\nabla_x g\|_{L^\infty} + T \|\nabla_x f\|_{L^\infty} + T \left(\|\nabla_x H\|_{L^\infty} + 1 \right).
$$

We hence get a bound C for the left-hand side. Obviously, this bound C is deterministic and independent of R, which allows us to retrieve the original equation by choosing R larger than C . Uniqueness of the hence constructed solution is obvious because we restricted our analysis to solutions that are bounded in $C^{\delta}(\mathbb{T}^d)$ by a deterministic constant. As mentioned in the statement, we can even state uniqueness in a wider class. This follows from the fact that the parameter n in [\[14](#page-86-2), Proposition 4.3.8] can be taken equal to 2. This completes the proof. \Box

Since \mathbb{F}^0 is generated by \mathcal{F}_0^0 and \mathbf{B}^0 , we can represent, for any $x \in \mathbb{T}^d$, the martingale $(m_t(x))_{0 \le t \le T}$ in the form

$$
m_t(x) = \sigma_0 \int_0^t v_s^0(x) \cdot \mathrm{d}B_s^0, \quad 0 \le t \le T,
$$
\n(3.4)

where $(v_t^0(x))_{0 \le t \le T}$ is an \mathbb{R}^d -valued \mathbb{F}^0 -progressively measurable process. The regularity of v^0 is given by the following statement:

Lemma 3.2. The representation term in [\(3.4\)](#page-20-0) induces a process $(v_t^0)_{0 \le t \le T}$ with values in $\mathcal{H}^{\lfloor \Delta \rfloor}(\mathbb{T}^d)$ (which is embedded \int *in* $\mathcal{C}^{\mathcal{S}-d/2-1}(\mathbb{T}^d)$ *).* Moreover,

$$
\mathbb{E}^{0} \int_{0}^{T} \|v_{t}^{0}\|_{\lambda - d/2 - 1}^{2} \mathrm{d}s < \infty,\tag{3.5}
$$

and, for any $k \in \{0, \dots, \lfloor \Delta - d/2 \rfloor - 1\}$, \mathbb{P}^0 -almost surely, for all $x \in \mathbb{T}^d$,

$$
\nabla_x^k m_t(x) = \sigma_0 \int_0^t \nabla_x^k v_s^0(x) \cdot \mathrm{d}B_s^0, \quad t \in [0, T]. \tag{3.6}
$$

Notice that $\mathcal{H}^{[\lambda]}(\mathbb{T}^d)$ (equipped with $\|\cdot\|_{\lambda-d/2-1}$) is separable: for any $t \in [0,T]$, v_t^0 , when regarded as a random variable with values in $C^{\Delta-d/2-1}(\mathbb{T}^d)$ is obviously *Bochner* measurable.

Proof. Because $\sup_{t\in[0,T]}||u_t||_{\infty} \in L^{\infty}(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$, we can see the equation [\(3.1\)](#page-19-0) as a linear equation with a source term in $\mathcal{C}^{b-1}(\mathbb{T}^d)$. This makes it possible to apply [\[30](#page-86-36), Theorem 2.3] (with a modicum of care because the latter result is stated on the Euclidean space, but the adaptation to the periodic setting is straightforward). We deduce that the representation term $(v_t^0)_{0 \le t \le T}$ in [\(3.4\)](#page-20-0) takes values in $\mathcal{H}^{\lfloor \Delta \rfloor}(\mathbb{T}^d)$. Moreover, it satisfies

$$
\mathbb{E}^0 \int_0^T \|v_t^0\|_{\lfloor \lambda \rfloor,2}^2 \mathrm{d} s < \infty.
$$

By Sobolev embedding, we obtain [\(3.5\)](#page-20-1) (because $|\triangle| - d/2 > \triangle - d/2 - 1$). Then, [\(3.6\)](#page-20-2) can be easily established by induction on $k \in \{1, \dots, |\lambda - d/2| - 1\}$, by passing to the limit (as ε tends to 0) in the representation formula

$$
\frac{1}{\varepsilon} \left(\nabla_x^{k-1} m_t(x + \varepsilon \xi) - \nabla_x^{k-1} m_t(x) \right) = \sigma_0 \int_0^t \left[\frac{1}{\varepsilon} \left(\nabla_x^{k-1} v_s^0(x + \varepsilon \xi) - \nabla_x^{k-1} v_s^0(x) \right) \right] \cdot \mathrm{d}B_s,
$$

for any $x \in \mathbb{T}^d$ and unitary $\xi \in \mathbb{R}^d$. This proves that, for all $x \in \mathbb{T}^d$, \mathbb{P}^0 -almost surely, for all $t \in [0, T]$, [\(3.6\)](#page-20-2) holds true. Using the regularity properties of the left-hand side in [\(3.6\)](#page-20-2), we easily deduce that, for any integer $p \ge 1$, there exists a constant C_p such that, for any $k \in \{0, \dots, \lfloor \Delta - d/2 \rfloor - 1\}$ and any $x, x' \in \mathbb{T}^d$,

$$
\mathbb{E}\bigg[\sup_{0\leq t\leq T}\bigg|\int_0^t\Big(\nabla^k_xv^0_s(x')-\nabla^k_xv^0_s(x)\Big)\cdot\mathrm{d}B_s^0\bigg|^p\bigg]=\mathbb{E}\bigg[\sup_{0\leq t\leq T}|\nabla^k m_t(x')-\nabla^k m_t(x)|^p\bigg]\leq C_p|x'-x|^p,
$$

which proves, by Kolmogorov continuity theorem, that we can find, for each $x \in \mathbb{T}^d$, a version of the stochastic process $(\int_0^t \nabla_x^k v_s^0(x) \cdot dB_s)_{0 \le t \le T}$ that is continuous in x. This permits to exchange the quantifiers in [\(3.6\)](#page-20-2) and then get the formula \mathbb{P}^0 -almost surely, for all $(t, x) \in [0, T] \times \mathbb{T}^d$.

3.2. Uniform in time estimates for the HJB equation

Within the same framework as in Subsection [3.1,](#page-18-1) we now provide a uniform in time estimate of the gradient of the solution to (3.1) . This bound is key in our analysis.

Proposition 3.3. *Under Assumption* (A), *there exists a constant* R_0 *, only depending on the parameters in* (A) *except* (σ_0, T) *(in particular,* R_0 *is independent of* μ *), such that*

$$
\mathbb{P}^0\left(\sup_{0\leq t\leq T} \|\nabla_x u_t\|_{L^\infty} \leq R_0\right) = 1.
$$
\n(3.7)

Proof. The proof of the bound [\(3.7\)](#page-21-0) is inspired from the proof of Lemma 1.5 in [\[16\]](#page-86-29), but adapted to the SPDE setting.

The very first step of the proof is to consider, for a given $\xi \in \mathbb{R}^d$ with $|\xi| = 1$, the equation satisfied by $(D_{\xi\xi}^2 u_t(x)) =$ $(\nabla^2_{xx} u_t(x)\xi) \cdot \xi)_{0 \le t \le T, x \in \mathbb{T}^d}$. By a straightforward adaptation of the computations as in [\[16\]](#page-86-29), we get that $D^2_{\xi\xi} u$ solves the following equation:

$$
d_t\left(D_{\xi\xi}^2u_t(x)\right) = \left[-\frac{1}{2}\Delta_x D_{\xi\xi}^2u_t(x) + \left(\nabla_{xx}^2 H\left(x,\nabla_x u_t(x)\right)\xi\right)\cdot\xi + 2\left(\nabla_{px}^2 H\left(x,\nabla_x u_t(x)\right)\xi\right)\cdot\nabla_x D_{\xi}u_t(x) + \left(\nabla_{pp}^2 H\left(x,\nabla_x u_t(x)\right)\nabla_x D_{\xi}u_t(x)\right)\cdot\nabla_x D_{\xi}u_t(x) + \nabla_p H\left(x,\nabla_x u_t(x)\right)\cdot\nabla_x D_{\xi\xi}^2u_t(x) - D_{\xi\xi}^2 f_t(X_t^0,x,\mu_t)\right]dt
$$

+ d\left(D_{\xi\xi}^2 m_t\right)(x),

where the notations D_{ξ} and $D_{\xi\xi}^2$ are extended in an obvious manner to the first and second order derivatives in the direction ξ of (possibly random) functions different from $u_t(\cdot)$. Above, $(D_{\xi\xi}^2 m_t(x))_{0 \le t \le T}$ is a martingale.

The key point in the proof is to observe that there exists a constant $C > 0$, only depending on the parameters in (A5), such that, for $x \in \mathbb{T}^d$, $p, q \in \mathbb{R}^d$,

$$
\left(\nabla_{xx}^2 H(x,p)\xi\right) \cdot \xi + 2\left(\nabla_{px}^2 H(x,p)\xi\right) \cdot q + \left(\nabla_{pp}^2 H(x,p)q\right) \cdot q - D_{\xi\xi}^2 f_t(X_t^0, x, \mu_t)
$$
\n
$$
\geq C^{-1}|q|^2 - C. \tag{3.8}
$$

We next consider the same Brownian motion $(B_t)_{0 \le t \le T}$ as in the dynamics of the minor player. In particular, $(B_t)_{0 \leq t \leq T}$ is independent of the common noise $(B_t^0)_{0 \leq t \leq T}$. We then consider the SDE

$$
dX_t = -\nabla_p H(X_t, \nabla_x u_t(X_t))dt + dB_t, \quad t \in [t_0, T],
$$

for a given $t_0 \in [0, T]$ and for an initial condition $X_{t_0} = x$. Notice that the drift is random, but the solution is uniquely defined thanks to the regularity properties granted by Lemma [3.1.](#page-19-1) We also let

$$
U_t := D_{\xi\xi}^2 u_t(X_t).
$$

By applying Itô-Wentzell formula to $(U_t)_{0\leq t\leq T}$ and by invoking Lemma [3.2](#page-20-3) to guarantee that $\mathbb{E}^0 \int_{[0,T]} \|D^2_{\xi\xi}v^0_t\|_1^2 \mathrm{d}t < \infty$ (thanks to the condition $\lambda > d/2 + 5$), we get (see footnote [1](#page-12-0) for the proof)

$$
dU_t = \left[\left(\nabla_{xx}^2 H(X_t, \nabla_x u_t(X_t)) \xi \right) \cdot \xi + 2 \left(\nabla_{px}^2 H(X_t, \nabla_x u_t(X_t)) \xi \right) \cdot \nabla_x D_{\xi} u_t(X_t) + \left(\nabla_{pp}^2 H(X_t, \nabla_x u_t(X_t)) \nabla_x D_{\xi} u_t(X_t) \right) \cdot \nabla_x D_{\xi} u_t(X_t) - D_{\xi\xi}^2 f_t(X_t^0, X_t, \mu_t) \right] dt + dn_t,
$$

where $(n_t)_{0 \le t \le T}$ is a new generic martingale term, but now independent of the entry x. By [\(3.8\)](#page-21-1) and thanks to the lower bound $|\nabla_x D_{\xi} u_t(X_t)|^2 \ge |\nabla_x D_{\xi} u_t(X_t) \cdot \xi|^2 = |D^2_{\xi \xi} u_t(X_t)|^2 = |U_t|^2$, we get

$$
dU_t \ge (C^{-1}|U_t|^2 - C)dt + dn_t.
$$

= $C^{-1}(U_t - C)(U_t + C)dt + dn_t,$ (3.9)

which gives

$$
d\Big[\exp\bigg(-C^{-1}\int_{t_0}^t (U_s + C)ds\bigg)\big(U_t - C\big)\Big] \ge dh_t.
$$

In particular, assuming without any loss of generality that $||D^2_{\xi\xi}g||_{\infty} \leq C$ (i.e. $C \geq \kappa$, with the notations used in Assumption (A)), the boundary condition at time T in the above left-hand side has a non-positive value, from which we deduce (by conditioning on $\mathcal{F}_{t_0}^0 \otimes \mathcal{F}_{t_0}$) that $U_{t_0} \leq C$. By initializing X from any $x \in \mathbb{T}^d$ at time t_0 , this shows that, \mathbb{P}^0 -almost surely,

$$
\sup_{x \in \mathbb{T}^d} \sup_{\xi \in \mathbb{R}^d : |\xi| = 1} \nabla_x^2 u_{t_0}(x) \xi \cdot \xi \le C.
$$

By (15) in [\[16](#page-86-29)], which says that, on the torus, the Lipschitz constant of a smooth function can be controlled by its semiconcavity constant, we deduce that $\mathbb{P}^0(\{\|\nabla_x u_{t_0}\|_{L^\infty} \leq R_0\}) = 1$, with $R_0 := C\sqrt{d}$. By continuity in time of the process $(\Vert u_t \Vert_2)_{0 \le t \le T}$, we complete the proof. \Box

We eventually extend the result of Proposition [3.3](#page-21-2) to higher-derivatives:

Proposition 3.4. *Under Assumption* (A) *and within the context described in the beginning of Subsection [3.1,](#page-18-1) there exists a* constant $R_{\lambda-1}$, only depending on the parameters in (A) except (σ_0 , T) (in particular, R is independent of μ), such that

$$
\mathbb{P}^0\Big(\sup_{0\leq t\leq T} \|\nabla_x u_t\|_{\lambda-1} \leq R_{\lambda-1}\Big) = 1.
$$
\n(3.10)

Proof. When t is close to T (say $T - t \le 1$), the result follows from [\[14](#page-86-2), Proposition 4.3.8] (which applies since the gradient in the Hamiltonian is known to be bounded by means of Proposition [3.3\)](#page-21-2).

In particular, we can assume that $T - t \geq 1$. The only difficulty is to get a bound that is independent of T. By taking one derivative in the equation (3.1) as done in (3.3) and then by using standard heat kernel estimates, we claim that, for any integer $k \in \{1, \dots, |\Delta| - 1\}$, any $t, S \in [0, T]$, with $S - 1 \le t < S$,

$$
\begin{split} \|\nabla_x u_t\|_{\delta-k} &\leq \frac{C}{\sqrt{S-t}} \mathbb{E}^0 \big[\|\nabla_x u_S\|_{\delta-(k+1)} \, | \, \mathcal{F}_t^0 \big] \\ &+ C \mathbb{E}^0 \bigg[\int_t^S \frac{1}{\sqrt{r-t}} \sup_{i=1,\cdots,d} \left\| \nabla_p H(\cdot,\nabla_x u_r(\cdot)) \cdot \partial_{x_i} \nabla_x u_r \right\|_{\delta-(k+1)} \mathrm{d} r \, | \, \mathcal{F}_t^0 \bigg] \\ &+ C \mathbb{E}^0 \bigg[\int_t^S \frac{1}{\sqrt{r-t}} \Big(\left\| \nabla_x H(\cdot,\nabla_x u_r(\cdot)) \right\|_{s-(k+1)} + \left\| \nabla_x f_r(X_r^0,\cdot,\mu_r) \right\|_{s-(k+1)} \Big) \mathrm{d} r \, | \, \mathcal{F}_t^0 \bigg]. \end{split}
$$

Assuming that we have uniform (in time and in $\omega \in \Omega^0$) bounds for $\|\nabla_x u_t\|_{\omega-(k+1)}$, we deduce that there exists a constant C_k such that

$$
\|\nabla_x u_t\|_{\delta-k} \le \frac{C_k}{\sqrt{S-t}} + C_k \mathbb{E}^0 \left[\int_t^S \frac{1}{\sqrt{r-t}} \|\nabla_x u_r\|_{\delta-k} \mathrm{d}r \, |\, \mathcal{F}_t^0 \right]. \tag{3.11}
$$

And then, for any $\delta \in (0, S)$, with $\delta \leq 1$, (the mapping $t \mapsto \text{essup}_{\omega \in \Omega^0} ||\nabla_x u_t||_{\delta-k}$ is measurable as limit of $t \mapsto$ $\mathbb{E}^0[\|\nabla_x u_t\|_{\Delta-k}^p]^{1/p}$ as p tends to $\infty)$

$$
\begin{split} & \int_{S-\delta}^{S} \frac{1}{\sqrt{t-(S-\delta)}} \mathrm{essup}_{\omega \in \Omega^{0}} \|\nabla_{x} u_{t}\|_{\mathbf{\mathbf{\mathbf{b}}-k}} \mathrm{d}t \\ & \leq C_{k} B(\tfrac{1}{2},\tfrac{1}{2}) \ + C_{k} \int_{S-\delta}^{S} \frac{1}{\sqrt{t-(S-\delta)}} \bigg(\int_{t}^{S} \frac{1}{\sqrt{r-t}} \mathrm{essup}_{\omega \in \Omega^{0}} \|\nabla_{x} u_{r}\|_{\mathbf{\mathbf{\mathbf{b}}-k}} \mathrm{d}r \bigg) \mathrm{d}t \\ & = C_{k} B(\tfrac{1}{2},\tfrac{1}{2}) \ + C_{k} \int_{S-\delta}^{S} \mathrm{essup}_{\omega \in \Omega^{0}} \|\nabla_{x} u_{r}\|_{\mathbf{\mathbf{\mathbf{b}}-k}} \bigg(\int_{S-\delta}^{r} \frac{1}{\sqrt{r-t}} \frac{1}{\sqrt{t-(S-\delta)}} \mathrm{d}t \bigg) \mathrm{d}r \\ & = C_{k} B(\tfrac{1}{2},\tfrac{1}{2}) \ + C_{k} B(\tfrac{1}{2},\tfrac{1}{2}) \int_{S-\delta}^{S} \mathrm{essup}_{\omega \in \Omega^{0}} \|\nabla_{x} u_{r}\|_{\mathbf{\mathbf{\mathbf{b}}-k}} \mathrm{d}r \\ & \leq C_{k} B(\tfrac{1}{2},\tfrac{1}{2}) \ + C_{k} \sqrt{\delta} B(\tfrac{1}{2},\tfrac{1}{2}) \int_{S-\delta}^{S} \frac{1}{\sqrt{t-(S-\delta)}} \mathrm{essup}_{\omega \in \Omega^{0}} \|\nabla_{x} u_{r}\|_{\mathbf{\mathbf{\mathbf{b}}-k}} \mathrm{d}r, \end{split}
$$

where $B(\cdot, \cdot)$ is the Euler Bêta function. And then, we get a bound in the left hand-side for $\sqrt{\delta}B(1/2, 1/2)$ = $\min(1/2, 1/2C_k)$. Choosing $S = t + \delta$ in [\(3.11\)](#page-22-0), we get, for $t \leq T - 1$, a bound for essup_{$\omega \in \Omega$ 0 $||\nabla_x u_t||_{\delta - k}$.} \Box

3.3. BMO estimates for the minor MFG system

We now fix an \mathbb{F}^0 -adapted continuous path $X^0 = (X_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d (not necessarily solving the forward equation in [\(2.11\)](#page-8-2)). With it, we associate the stochastic MFG system [\(2.12\)](#page-9-0). Since f and g are assumed to be monotone in Lasry Lions sense, and since we have an *a priori* estimate for the gradient of the backward component, we can easily do as if the Hamiltonian were at most of linear growth and hence follow the proof of [\[14](#page-86-2), Theorem 4.3.1] to get a unique solution to the MFG system. Importantly, the backward component satisfies the conclusion of Lemma [3.1](#page-19-1) and Proposition [3.3.](#page-21-2) Our first step is to strengthen the estimate provided by Proposition [3.4.](#page-22-1)

Throughout the subsection, we consider a fixed \mathbb{F}^0 -stopping time τ with values in $[0,T]$ and we consider the auxiliary MFG system

$$
\partial_t \tilde{\mu}_t - \frac{1}{2} \Delta_x \tilde{\mu}_t - \text{div}_x \left(\nabla_p H \left(\cdot, \nabla_x \tilde{u}_t \right) \tilde{\mu}_t \right) = 0, \quad (t, x) \in [\tau, T] \times \mathbb{T}^d,
$$

\n
$$
\partial_t \tilde{u}_t(x) = -\frac{1}{2} \Delta_x \tilde{u}_t(x) + H \left(x, \nabla_x \tilde{u}_t(x) \right) - F(x, \tilde{\mu}_t), \quad (t, x) \in [\tau, T] \times \mathbb{T}^d,
$$

\n
$$
\tilde{u}_T(x) = 0, \quad x \in \mathbb{T}^d, \quad \tilde{\mu}_T = \mu_\tau,
$$
\n(3.12)

where F is as in Assumption (B). Except for the fact that the initial time is random, this MFG system is deterministic. Since F satisfies the Lasry-Lions monotonicity condition, the system (3.12) has a unique solution.

Proposition 3.5. *Under Assumption* (B) *and with the the same notation as above, there exists a constant* C*, only depending on the parameters in* (B) *except* (σ_0, T) *(in particular, C is also independent of* τ *), such that*

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\int_{\tau}^T\bigg(w_1(\mu_t,\tilde{\mu}_t)^2 + \Big\|(u_t-\tilde{u}_t) - \int_{\mathbb{T}^d}(u_t-\tilde{u}_t)(x)\,\mathrm{d}x\Big\|_{\mathbb{A}}^2\bigg)\,\mathrm{d}t\,|\,\mathcal{F}^0_{\tau}\bigg] \leq C\bigg\}\bigg) = 1.
$$

Proof. For simplicity, we assume in the first three steps of the proof that $\tau = 0$. We then explain in the very last step how the proof has to be changed when τ is general.

First Step. By monotonicity of the coefficient F and from the standard duality method of mean field games (see [\[14,](#page-86-2) Lemma 3.1.2]), we have, for any $t \in [0, T]$,

$$
-\langle u_t - \tilde{u}_t, \mu_t - \tilde{\mu}_t \rangle + c \mathbb{E}^0 \bigg[\int_t^T \big(\mu_r + \tilde{\mu}_r, |\nabla_x (u_r - \tilde{u}_r)|^2 \big) dr \, | \mathcal{F}_t^0 \bigg]
$$

$$
\leq -\mathbb{E}^0 \bigg[\int_t^T \big(f_r(X_r^0, \cdot, \mu_r) - F(\cdot, \mu_r), \mu_r - \tilde{\mu}_r \big) dr + (u_T - \tilde{u}_T, \mu_T - \tilde{\mu}_T) \, | \mathcal{F}_t^0 \bigg],
$$

for a constant $c > 0$ only depending on the parameters in (**B**) except (σ_0, T) .

By boundedness of u_T and \tilde{u}_T (but the latter is null), we obtain

$$
-(u_t - \tilde{u}_t, \mu_t - \tilde{\mu}_t) + c \mathbb{E}^0 \bigg[\int_t^T (\mu_r + \tilde{\mu}_r, |\nabla_x (u_r - \tilde{u}_r)|^2) dr \, | \, \mathcal{F}_t^0 \bigg]
$$

$$
\leq C \mathbb{E}^0 \bigg[1 + \int_t^T \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} || f_r(x_0, \cdot, \mu) - F(\cdot, \mu) ||_{L^\infty} dr \, | \, \mathcal{F}_t^0 \bigg].
$$
 (3.13)

We give below two applications of (3.13) .

Second Step. First, we study the equation for $(\mu_t - \tilde{\mu}_t)_{0 \le t \le T}$. We write

$$
\partial_t (\mu_t - \tilde{\mu}_t) = \frac{1}{2} \Delta_x (\mu_t - \tilde{\mu}_t) + \text{div}_x \Big(\nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot)) (\mu_t - \tilde{\mu}_t) \Big) + \text{div}_x \Big(\big[\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot)) \big] \mu_t \Big).
$$

Following Lemma [6.2](#page-84-0) and noticing that \tilde{u} satisfies the same bounds as u in Proposition [3.4,](#page-22-1) consider now the backward PDE

$$
\partial_t \varphi_t + \frac{1}{2} \Delta_x \varphi_t - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot)) \cdot \nabla_x \varphi_t = 0, \quad t \in [0, S]; \quad \varphi_S = \phi,
$$

for a smooth terminal function ϕ and for some $S \in [0, T]$ (since \tilde{u} is deterministic, so is the above PDE and there is no need to add an additional martingale term as it would be the case if we were considering the same equation but driven by u). By duality, we obtain

$$
\frac{\mathrm{d}}{\mathrm{d}t}(\varphi_t, \mu_t - \tilde{\mu}_t) = -\Big(\big[\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot))\big]\mu_t, \nabla_x \varphi_t\Big).
$$

Therefore, using the fact that $\mu_0 - \tilde{\mu}_0 = 0$, we obtain

$$
(\phi, \mu_S - \tilde{\mu}_S) \le \int_0^S \|\nabla_x \varphi_t\|_{L^\infty} (\mu_t, |\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot))|^2)^{1/2} dt
$$

$$
\le ||\phi||_1 \int_0^S \exp(-\gamma (S - t)) (\mu_t, |\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot))|^2)^{1/2} dt,
$$

with the exponential decay following from Lemma [6.2,](#page-84-0) and for γ only depending on the parameters in (B) but not on (σ_0, T) . And then,

$$
\|\mu_S - \tilde{\mu}_S\|_{-1} \leq \int_0^S \exp\left(-\gamma(S-t)\right) \left(\mu_t, |\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot))|^2\right)^{1/2} dt
$$

$$
\leq \frac{1}{\gamma^{1/2}} \left(\int_0^S \exp\left(-\gamma(S-t)\right) \left(\mu_t, |\nabla_p H(\cdot, \nabla_x u_t(\cdot)) - \nabla_p H(\cdot, \nabla_x \tilde{u}_t(\cdot))|^2\right) dt\right)^{1/2},
$$

which yields (replacing S by t in the left-hand side and t by r in the right-hand side)

$$
\mathbb{E}^{0}\left[\int_{0}^{T} \|\mu_{t}-\tilde{\mu}_{t}\|_{-1}^{2} dt\right] \leq \frac{1}{\gamma} \mathbb{E}^{0} \int_{0}^{T} \left(\int_{0}^{t} \exp\left(-\gamma(t-r)\right) \left(\mu_{r}, |\nabla_{p} H(\cdot, \nabla_{x} u_{r}(\cdot)) - \nabla_{p} H(\cdot, \nabla_{x} \tilde{u}_{r}(\cdot))|^{2}\right) dr\right) dt
$$

\n
$$
= \frac{1}{\gamma} \mathbb{E}^{0} \int_{0}^{T} \left(\mu_{r}, |\nabla_{p} H(\cdot, \nabla_{x} u_{r}(\cdot)) - \nabla_{p} H(\cdot, \nabla_{x} \tilde{u}_{r}(\cdot))|^{2}\right) \left(\int_{r}^{T} \exp\left(-\gamma(t-r)\right) dt\right) dr
$$

\n
$$
\leq \frac{1}{\gamma^{2}} \mathbb{E}^{0} \int_{0}^{T} \left(\mu_{r}, |\nabla_{p} H(\cdot, \nabla_{x} u_{r}(\cdot)) - \nabla_{p} H(\cdot, \nabla_{x} \tilde{u}_{r}(\cdot))|^{2}\right) dr.
$$

We now insert [\(3.13\)](#page-24-0) (with $t = 0$, recalling that $\mu_0 = \tilde{\mu}_0$) into the above bound. We obtain (recalling that $\|\mu - \tilde{\mu}\|_{-1}$ is the same as $\mathbb{W}_1(\mu, \tilde{\mu})$

$$
\mathbb{E}^{0}\bigg[\int_{0}^{T} \mathbb{W}_{1}(\mu_{t},\tilde{\mu}_{t})^{2}dt\bigg] \leq C\bigg(1+\mathbb{E}^{0}\int_{0}^{T} \sup_{x_{0}\in\mathbb{R}^{d}} \sup_{\mu\in\mathcal{P}(\mathbb{T}^{d})} \|f_{r}(x_{0},\cdot,\mu)-F(\cdot,\mu)\|_{\infty} dr\bigg). \tag{3.14}
$$

Third Step. We provide a similar estimate for the difference of the two backward components $u - \tilde{u}$. We have

$$
d_t(u_t - \tilde{u}_t)(x) = \left[-\frac{1}{2}\Delta_x(u_t - \tilde{u}_t)(x) + H(x, \nabla_x u_t(x)) - H(x, \nabla_x \tilde{u}_t(x)) - (f_t(X_t^0, x, \mu_t) - F(x, \tilde{\mu}_t)) \right] dt
$$

+ $\sigma_0 v_t^0(x) \cdot dB_t^0$.

Rewriting the difference of the two Hamiltonians as

$$
H(x, \nabla_x u_t(x)) - H(x, \nabla_x \tilde{u}_t(x))
$$

=
$$
\left(\int_0^1 \nabla_p H(x, \theta \nabla_x u_t(x) + (1 - \theta) \nabla_x \tilde{u}_t(x)) d\theta \right) \cdot (\nabla_x u_t(x) - \nabla_x \tilde{u}_t(x)),
$$

we apply Lemma [6.3](#page-85-1) with

$$
b_t(x) = -\left(\int_0^1 \nabla_p H\big(x, \theta \nabla_x u_t(x) + (1 - \theta) \nabla_x \tilde{u}_t(x)\big) d\theta\right).
$$
 (3.15)

For a smooth initial condition q satisfying $\int_{\mathbb{T}^d} q(x) dx = 0$ and for an initial time $t_0 \in [0, T]$, we solve the Fokker-Planck equation

$$
\partial_t q_t - \frac{1}{2} \Delta_x q_t + \text{div}_x (b_t(\cdot) q_t) = 0, \quad t \in [t_0, T]; \quad q_{t_0} = q.
$$

We observe that this equation is a random conservative equation, due to the random nature of $(b_t)_{0 \le t \le T}$. By decomposing q in positive and negative parts, it suffices to construct solutions for q a probability density. Following the proof of Lemma [2.4](#page-7-3) and the first step of the proof of Lemma [2.9,](#page-8-1) we deduce that the solution is measurable with respect to ω^0 : $(q_t)_{0 \le t \le T}$ can be regarded as an \mathbb{F}^0 -progressively measurable process with values in the space of finite signed measures on \mathbb{T}^d . Then, by duality, we expand (see [\[14](#page-86-2), Lemma 4.3.11])

$$
d_t(u_t - \tilde{u}_t, q_t) = -(f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t), q_t)dt + dn_t, \quad t \in [t_0, T],
$$
\n(3.16)

where $(n_t)_{t_0 \le t \le T}$ is a generic martingale term. We obtain, for a constant C depending on the parameters in (B) except (σ_0, T) ,

$$
(u_{t_0} - \tilde{u}_{t_0}, q) \leq \mathbb{E}^0 \bigg[\|q_T\|_{-\delta} \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|g(x_0, \cdot, \mu)\|_{\delta} + C \int_{t_0}^T \|q_t\|_{-\delta} \mathbb{W}_1(\mu_t, \tilde{\mu}_t) dt + \int_{t_0}^T \|q_t\|_{-\delta} \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|f_t(x_0, \cdot, \mu) - F(\cdot, \mu)\|_{\delta} dt \, | \mathcal{F}_{t_0}^0 \bigg].
$$

By Lemma 6.3 (which is stated for deterministic conservative equations but which applies here because b is bounded by a deterministic constant), we can find new values of C and γ such that $||q_t||_{-\delta} \leq C \exp(-\gamma(t - t_0))||q||_{-\delta}$. Therefore, choosing q as $q(\cdot) = (-1)^l \partial^l \rho(\cdot - x)$ and then as $q(\cdot) = (-1)^l [\partial^l \rho(\cdot - x) - \partial^l \rho(\cdot - x')]$, for $x, x' \in \mathbb{T}^d$ and ρ a smooth density on \mathbb{R}^d , where $\partial^l \rho$ denotes the derivative of ρ along l (arbitrary) directions of \mathbb{R}^d (with possible repetitions), and then letting ρ tend to the Delta mass in 0, we get

$$
\left\| u_{t_0} - \tilde{u}_{t_0} - \int_{\mathbb{T}^d} \left(u_{t_0} - \tilde{u}_{t_0} \right)(x) dx \right\|_{\mathfrak{d}} \leq C \exp(-\gamma(T - t_0)) + C \mathbb{E}^0 \bigg[\int_{t_0}^T \exp(-\gamma(t - t_0)) \mathbb{W}_1(\mu_t, \tilde{\mu}_t) dt \, | \, \mathcal{F}_{t_0}^0 \bigg] + C \int_{t_0}^T \exp(-\gamma(t - t_0)) \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \| f_t(x_0, \cdot, \mu) - F(\cdot, \mu) \|_{\mathfrak{d}} dt.
$$

Squaring as done in the second step, we obtain (allowing the constant C to vary from line to line as long as it only depends on the various parameters in (**B**) except (σ_0, T))

$$
\|u_{t_0} - \tilde{u}_{t_0} - \int_{\mathbb{T}^d} (u_{t_0} - \tilde{u}_{t_0})(x) dx\|_{\Delta}^2 \leq C \exp(-\gamma(T - t_0)) + C \mathbb{E}^0 \left[\int_{t_0}^T \exp(-\gamma(t - t_0)) \mathbb{W}_1(\mu_t, \tilde{\mu}_t)^2 dt \, | \, \mathcal{F}_{t_0}^0 \right]
$$

$$
+ C \int_{t_0}^T \exp(-\gamma(t - t_0)) \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|f_t(x_0, \cdot, \mu) - F(\cdot, \mu)\|_{\Delta} dt,
$$

where we used the fact that f_t and F are uniformly bounded. Above, γ is implicitly hidden in the multiplicative constant C . Hence, by integrating in t_0 and using the same Fubini argument as before, we deduce that

$$
\mathbb{E}^{0}\bigg[\int_{0}^{T}\Big\|\big(u_{t}-\tilde{u}_{t}\big)-\int_{\mathbb{T}^{d}}\big(u_{t}-\tilde{u}_{t}\big)(x)\,\mathrm{d}x\Big\|_{\mathcal{S}}^{2}\mathrm{d}t\bigg] \leq C + C\mathbb{E}^{0}\bigg[\int_{0}^{T}\mathbb{W}_{1}(\mu_{t},\tilde{\mu}_{t})^{2}\mathrm{d}t\bigg] + C\int_{0}^{T}\sup_{x_{0}\in\mathbb{T}^{d}}\sup_{\mu\in\mathcal{P}(\mathbb{T}^{d})}\|f_{t}(x_{0},\cdot,\mu)-F(\cdot,\mu)\|_{\mathcal{S}}\mathrm{d}t.
$$

Substituting the last term on the first line by the upper bound in (3.14) , we obtain

$$
\mathbb{E}^{0}\bigg[\int_{0}^{T}\bigg\|\big(u_{t}-\tilde{u}_{t}\big)-\int_{\mathbb{T}^{d}}\big(u_{t}-\tilde{u}_{t}\big)(x)\,\mathrm{d}x\bigg\|_{\mathbb{A}}^{2}\mathrm{d}t\bigg]\leq C+C\int_{0}^{T}\sup_{x_{0}\in\mathbb{T}^{d}}\sup_{\mu\in\mathcal{P}(\mathbb{T}^{d})}\|f_{t}(x_{0},\cdot,\mu)-F(\cdot,\mu)\|_{\mathbb{A}}\mathrm{d}t.
$$

The above right-hand side is bounded thanks to (**B2**).

Fourth Step. Replacing the initial time 0 by a more general stopping time τ with values in [0, T] (as given in the assumption of the lemma), we get in a very similar manner (using conditional expectation given \mathcal{F}^0_τ instead of expectation) that there exists a constant C, independent of σ_0 , τ and T, such that

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\int_{\tau}^T \Big\|\big(u_t-\tilde{u}_t\big)-\int_{\mathbb{T}^d} \big(u_t-\tilde{u}_t\big)(x)\,\mathrm{d} x\Big\|_{\mathcal{A}}^2\mathrm{d} t\,\big|\,\mathcal{F}^0_{\tau}\bigg]\leq C\bigg\}\bigg)=1.
$$

Reformulating [\(3.14\)](#page-24-1) in a similar manner, we complete the proof.

An important application is (recall (3.4) for the definition of the term v^0 below)

Proposition 3.6. *Under Assumption* (B), *there exist an exponent* $\gamma > 0$ *and a constant* C, *only depending on the parameters in* (**B**) *except* (σ_0, T) *, such that, for any* \mathbb{F}^0 -stopping time τ *with values in* $[0, T]$ *,*

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\exp\bigg(\gamma\sigma_0^2\int_{\tau}^T\bigg\|v_r^0(\cdot)-\int_{\mathbb{T}^d}v_r^0(x)\,\mathrm{d}x\bigg\|_{\lfloor\lambda\rfloor-(d/2+1)}^2\,\mathrm{d}r\bigg)\,|\,\mathcal{F}_\tau^0\bigg]\leq C\bigg\}\bigg)=1.
$$

In the sequel, we let

$$
\bar{v}_t^0(x) := v_t^0(x) - \int_{\mathbb{T}^d} v_t^0(y) \, dy, \quad (t, x) \in [0, T] \times \mathbb{T}^d. \tag{3.17}
$$

Proof. The proof relies on BMO estimates similar to those used in Section [2,](#page-2-0) but there is a subtlety due to the fact that the argument in the exponential involves a functional norm.

We proceed by Sobolev embeddings. We recall that $C^{\delta}(\mathbb{T}^d)$ is included in the Sobolev space $\mathcal{H}^{\delta}(\mathbb{T}^d)$. Also, because $H^{\delta-1}(\mathbb{T}^d)$ embeds continuously in $\mathcal{C}^{[\delta]-(d/2+1)}(\mathbb{T}^d)$, we obtain

$$
\left\|\bar{v}^0_r\right\|_{\lfloor\delta\rfloor-(d/2+1)}^2 \leq C\left\|\bar{v}^0_r\right\|_{H^{\delta-1}(\mathbb{T}^d)}^2 = C\left\|\bar{v}^0_r\right\|_{\delta-1,2}^2 = C\sum_{k\in\mathbb{Z}^d\backslash\{0\}}\left(1+|k|^2\right)^{\delta-1}\lvert (v^0_r,e_k)\rvert^2,
$$

where we recall $(e_k)_{k\in\mathbb{Z}^d}$ is the standard (complex valued) Fourier basis of \mathbb{T}^d and (\cdot, \cdot) denotes here the standard inner product in $L^2(\mathbb{T}^d)$ (understood coordinate-wise since v^0 takes values in \mathbb{R}^d). Therefore, for τ as in the statement,

$$
\mathbb{E}^{0}\bigg[\int_{\tau}^{T} \left\|\bar{v}_{r}^{0}\right\|_{\lfloor\delta\rfloor - (d/2+1)}^{2} \mathrm{d}r \,|\,\mathcal{F}_{\tau}^{0}\bigg] \leq C \sum_{k \neq 0} \left(1 + |k|^{2}\right)^{\delta - 1} \mathbb{E}^{0}\bigg[\int_{\tau}^{T} |(v_{r}^{0}, e_{k})|^{2} \mathrm{d}r \,|\,\mathcal{F}_{\tau}^{0}\bigg].\tag{3.18}
$$

Next, we compute the dynamics of the Fourier coefficients. For $k \in \mathbb{Z}^d$,

$$
d_t(u_t - \tilde{u}_t, e_k) = 2\pi^2 |k|^2 (u_t - \tilde{u}_t, e_k) dt + (e_k, H(\cdot, \nabla_x u_t(\cdot)) - H(\cdot, \nabla_x \tilde{u}_t(\cdot))) dt - (e_k, f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t)) dt + \sigma_0(v_t^0, e_k) dB_t^0, \quad t \in [0, T].
$$

Taking squared norm, we obtain

$$
d_t \vert (u_t - \tilde{u}_t, e_k) \vert^2 \geq 2(u_t - \tilde{u}_t, e_k) (e_k, H(\cdot, \nabla_x u_t(\cdot)) - H(\cdot, \nabla_x \tilde{u}_t(\cdot))) dt - 2(u_t - \tilde{u}_t, e_k) (e_k, f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t)) dt + \sigma_0^2 \vert (v_t^0, e_k) \vert^2 dt + dn_t,
$$

 \Box

where $(n_t)_{0 \leq t \leq T}$ is a generic martingale term. Therefore,

$$
\sigma_0^2 \mathbb{E}^0 \bigg[\int_{\tau}^T \sum_{k \neq 0} (1+|k|^2)^{\delta-1} |(v_t^0, e_k)|^2 dt | \mathcal{F}_{\tau}^0 \bigg]
$$

\n
$$
\leq \mathbb{E}^0 \bigg[\sum_{k \neq 0} (1+|k|^2)^{\delta-1} |(u_T, e_k)|^2 | \mathcal{F}_{\tau}^0 \bigg]
$$

\n
$$
- 2 \mathbb{E}^0 \bigg[\int_{\tau}^T \sum_{k \neq 0} (1+|k|^2)^{\delta-1} (u_t - \tilde{u}_t, e_k) (e_k, f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t)) dt | \mathcal{F}_{\tau}^0 \bigg]
$$

\n
$$
+ 2 \mathbb{E}^0 \bigg[\int_{\tau}^T \sum_{k \neq 0} (1+|k|^2)^{\delta-1} (u_t - \tilde{u}_t, e_k) (e_k, H(\cdot, \nabla u_t(\cdot)) - H(\cdot, \nabla \tilde{u}_t(\cdot))) dt | \mathcal{F}_{\tau}^0 \bigg].
$$

And then, by (3.18) and for a constant C as in the statement,

$$
\sigma_0^2 \mathbb{E}^0 \Biggl[\int_{\tau}^{T} ||\bar{v}_r^0||_{\lfloor \lambda \rfloor - (d/2+1)}^2 dr | \mathcal{F}_r^0 \Biggr] \n\leq C \mathbb{E}^0 \Biggl[||u_T||_{\lambda - 1, 2}^2 | \mathcal{F}_r^0 \Biggr] \n+ C \mathbb{E}^0 \Biggl[\int_{\tau}^{T} \Biggl\| (u_t - \tilde{u}_t) - \int_{\mathbb{T}^d} (u_t - \tilde{u}_t)(x) dx \Biggr\|_{\lambda - 1, 2} ||f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t)||_{\lambda - 1, 2} dt | \mathcal{F}_r^0 \Biggr] \n+ C \mathbb{E}^0 \Biggl[\int_{\tau}^{T} \Biggl\| (u_t - \tilde{u}_t) - \int_{\mathbb{T}^d} (u_t - \tilde{u}_t)(x) dx \Biggr\|_{\lambda - 1, 2} \Biggr\| H(\cdot, \nabla_x u_t(\cdot)) - H(\cdot, \nabla_x \tilde{u}_t(\cdot)) \Biggr\|_{\lambda - 1, 2} dt | \mathcal{F}_r^0 \Biggr] \n\leq C \mathbb{E}^0 \Biggl[||u_T||_{\lambda - 1}^2 | \mathcal{F}_r^0 \Biggr] + C \mathbb{E}^0 \Biggl[\int_{\tau}^{T} \Biggl\| f_t(X_t^0, \cdot, \mu_t) - F(\cdot, \tilde{\mu}_t) \Biggr\|_{\lambda - 1}^2 dt | \mathcal{F}_r^0 \Biggr] \n+ C \mathbb{E}^0 \Biggl[\int_{\tau}^{T} \Biggl\| (u_t - \tilde{u}_t) - \int_{\mathbb{T}^d} (u_t - \tilde{u}_t)(x) dx \Biggr\|_{\lambda - 1}^2 dt | \mathcal{F}_r^0 \Biggr] \n+ C \mathbb{E}^0 \Biggl[\int_{\tau}^{T} \Biggl\| H(\cdot, \nabla_x u_t(\cdot)) - H(\cdot, \nabla_x \tilde{u}_t(\cdot)) \Biggr\|_{\lambda - 1}^2 dt | \mathcal{F}_r^0 \Biggr].
$$

The key point to treat the very last term is to observe that

$$
H(\cdot, \nabla_x u_t(\cdot)) - H(\cdot, \nabla_x \tilde{u}_t(\cdot)) = -b_t(\cdot) \cdot (\nabla_x u_t - \nabla_x \tilde{u}_t)(\cdot),
$$

with $b_t(\cdot)$ as in [\(3.15\)](#page-25-0). By Proposition [3.4](#page-22-1) (with a similar result holding true for the derivatives of \tilde{u}), we have a bound for $||b_t||_{\delta-1}$, independently of (σ_0, T) . Thanks to this,

$$
\left\|H(\cdot,\nabla_x u_t(\cdot)) - H(\cdot,\nabla_x \tilde{u}_t(\cdot))\right\|_{\delta-1} \leq C\left\|\nabla_x (u_t - \tilde{u}_t)\right\|_{\delta-1} \leq C\left\|(u_t - \tilde{u}_t) - \int_{\mathbb{T}^d} (u_t - \tilde{u}_t)(x) \,dx\right\|_{\delta}.
$$

And then, using the Lipschitz property of F w.r.t. μ in \mathcal{C}^{δ} , we obtain

$$
\sigma_0^2 \mathbb{E}^0 \bigg[\int_{\tau}^T \left\| \bar{v}^0(r, \cdot) \right\|_{\lfloor \lambda \rfloor - (d/2 + 1)}^2 \mathrm{d}r \, |\mathcal{F}_{\tau}^0 \bigg]
$$

\n
$$
\leq C \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|g(x_0, \cdot, \mu)\|_{\lambda} + C \int_{\tau}^T \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|f_t(x_0, \cdot, \mu) - F(\cdot, \mu)\|_{\lambda} \mathrm{d}t
$$

\n
$$
+ C \mathbb{E}^0 \bigg[\int_{\tau}^T \mathbb{W}_1(\mu_t, \tilde{\mu}_t)^2 \mathrm{d}t \, |\mathcal{F}_{\tau}^0 \bigg] + C \mathbb{E}^0 \bigg[\int_{\tau}^T \left\| \left(u_t - \tilde{u}_t \right) - \int_{\mathbb{T}^d} \left(u_t - \tilde{u}_t \right)(x) \, \mathrm{d}x \right\|_{\lambda}^2 \mathrm{d}t \, |\mathcal{F}_{\tau}^0 \bigg],
$$

where we removed the square in the $f_t - F$ term by using boundedness of the latter. Thanks to (B2). and Proposition [3.5,](#page-23-1) we can bound the left-hand side by C (for a possibly new value of C). The conclusion follows from the theory of BMO martingale, see [\[40](#page-86-30), Theorem 2.2]. \Box **Remark 3.7.** The same proof shows that there exist an exponent $\gamma_T > 0$ and a constant C_T , possibly depending on T, such that, for any \mathbb{F}^0 -stopping time τ with values in $[0, T]$,

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\exp\bigg(\gamma_T\sigma_0^2\int_{\tau}^T\big\|v_r^0\big\|_{\lfloor\delta\rfloor-(d/2+1)}^2\,\mathrm{d} r\bigg)\,|\,\mathcal{F}_\tau^0\bigg]\leq C_T\bigg\}\bigg)=1.
$$

The only difference is that, without the additional centring, we loose the exponential decay in the second and third steps of the proof of Proposition [3.5.](#page-23-1)

3.4. Forward-backward SDE for the major player

We here provide similar results for the system (2.11) satisfied by the major player, using now the following auxiliary deterministic HJB equation:

$$
\partial_t w^0(t, x_0) + \frac{1}{2}\sigma_0^2 \Delta_{x_0} w^0(t, x_0) + F^0(x_0) - H^0(x_0, \nabla_{x_0} w^0(t, x_0)) = 0, \quad (t, x_0) \in [0, T] \times \mathbb{R}^d,
$$

\n
$$
w^0(T, x_0) = 0.
$$
\n(3.19)

Existence and uniqueness of a (classical) solution is standard. The key point is that, by [\[58](#page-87-14), Theorem 1.20], we have a bound for $\sup_{(t,x_0)\in[0,T]\times\mathbb{R}^d} |\nabla_{x_0} w^0(t,x_0)|$ that only depends on the parameters in (**B**), except T. Using the fact that $F⁰$ has bounded first-order derivatives, we deduce from standard regularization properties of the heat kernel that $\sup_{(t,x_0)\in[0,T]\times\mathbb{R}^d} |\nabla_{x_0}^2 w^0(t,x_0)|$ is bounded with a bound that only depends on the parameters in (**B**), except T. Notice that when $F^0 \equiv 0$ and $H^0(\cdot, 0) \equiv 0$, then $w^0 \equiv 0$.

Lemma 3.8. *Under Assumption* (**B**), for any \mathbb{F}^0 -adapted continuous path $(\mu_t)_{0 \le t \le T}$ with values in $\mathcal{P}(\mathbb{T}^d)$ and any **F** 0 *-stopping time* τ *with values in* [0, T]*, any solution to the forward-backward SDE* [\(2.11\)](#page-8-2) *(in the sense of item 1 in Definition [2.10\)](#page-9-1) satisfies*

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\int_\tau^T \sigma_0^2 |Z_r^0 - \nabla_{x_0} w^0(X_t^0)|^2 dr \,|\,\mathcal{F}_\tau^0\bigg] \le C\bigg\}\bigg) = 1,\tag{3.20}
$$

for a constant C *only depending on the parameters in* (**B**) *except the parameters* (σ_0, T) *.*

Proof. Using smoothness of w^0 , we can expand $(Y_t^0 - w^0(t, X_t^0))_{0 \le t \le T}$ by means of Itô's formula. We get

$$
d[Y_t^0 - w^0(t, X_t^0)] = -(f_t^0(X_t^0, \mu_t) - F^0(X_t^0))dt
$$

$$
- (L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0)) + H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) - \nabla_p H^0(X_t^0, Z_t^0) \cdot \nabla_{x_0} w^0(t, X_t^0)) dt
$$

$$
+ \sigma_0(Z_t^0 - \nabla_{x_0} w^0(t, X_t^0)) \cdot dB_t^0, \quad t \in [0, T].
$$

We recall the standard formula (see (2.3))

$$
H^{0}(x_{0}, p_{0}) = p_{0} \cdot \nabla_{p} H^{0}(x_{0}, p_{0}) - L^{0}(x_{0}, -\nabla_{p} H^{0}(x_{0}, p_{0})), \qquad (3.21)
$$

from which we obtain (using $p_0 = Z_t^0$)

$$
d[Y_t^0 - w^0(t, X_t^0)] = -(f_t^0(X_t^0, \mu_t) - F^0(X_t^0))dt
$$

$$
-(-H^0(X_t^0, Z_t^0) + H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) - \nabla_p H^0(X_t^0, Z_t^0) \cdot [\nabla_{x_0} w^0(t, X_t^0) - Z_t^0] dt
$$

$$
+ \sigma_0(Z_t^0 - \nabla_{x_0} w^0(t, X_t^0)) \cdot dB_t^0, \quad t \in [0, T].
$$

Here, we write

$$
H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0})) - H^{0}(X_{t}^{0}, Z_{t}^{0}) - \nabla_{p} H^{0}(X_{t}^{0}, Z_{t}^{0}) \cdot [\nabla_{x_{0}} w^{0}(t, X_{t}^{0}) - Z_{t}^{0}]
$$

\n
$$
= \int_{0}^{1} [\nabla_{p} H^{0}(X_{t}^{0}, \theta \nabla_{x_{0}} w^{0}(t, X_{t}^{0}) + (1 - \theta)Z_{t}^{0}) - \nabla_{p} H^{0}(X_{t}^{0}, Z_{t}^{0})] \cdot (\nabla_{x_{0}} w^{0}(t, X_{t}^{0}) - Z_{t}^{0}) d\theta
$$

\n
$$
= [A_{t}^{0}(\nabla_{x_{0}} w^{0}(t, X_{t}^{0}) - Z_{t}^{0})] \cdot (\nabla_{x_{0}} w^{0}(t, X_{t}^{0}) - Z_{t}^{0}),
$$

where we have let

$$
A_t^0 := \int_0^1 \int_0^1 \nabla_{pp}^2 H^0(X_t^0, \theta \varphi \nabla_{x_0} w^0(t, X_t^0) + (1 - \varphi \theta) Z_t^0) d\theta d\varphi.
$$

So, with the notations

$$
P_t^0 := Y_t^0 - w^0(t, X_t^0), \quad Q_t^0 := Z_t^0 - \nabla_{x_0} w^0(t, X_t^0),
$$

we obtain the following BSDE:

$$
dP_t^0 = \left[-\left(f_t^0(X_t^0, \mu_t) - F^0(X_t^0)\right) - (A_t^0 Q_t^0) \cdot Q_t^0 \right] dt + \sigma_0 Q_t^0 \cdot dB_t^0, \quad t \in [0, T].
$$

Using Girsanov transformation, it is standard (see [\[9](#page-86-37)] and the references therein about quadratic BSDEs) to deduce that

$$
|P_t^0| \le \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} |g^0(x_0, \mu)| + \int_t^T \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} |f_r^0(x_0, \mu) - F^0(\mu)| dr \le C,
$$
\n(3.22)

with C independent of (σ_0, T) , see (B2).

We now let $E^0 := (E_t^0 := \exp(\nu P_t^0))_{0 \le t \le T}$, for some parameter $\nu > 0$. Notice from [\(3.22\)](#page-29-0) that we have a global bound for both P^0 and E^0 . This bound is independent of σ_0 and T (and of τ). We then expand

$$
dE_t^0 = \nu E_t^0 dP_t^0 + \frac{1}{2} \nu^2 E_t^0 d \langle P^0 \rangle_t
$$

= $\nu E_t^0 \Big[- \big(f_t^0 (X_t^0, \mu_t) - F^0 (X_t^0) \big) - (A_t^0 Q_t^0) \cdot Q_t^0 \Big] dt + \frac{1}{2} \nu^2 \sigma_0^2 E_t^0 |Q_t^0|^2 dt + \nu \sigma_0 E_t^0 Q_t^0 \cdot dB_t^0.$

Recall that $\sigma_0 \ge 1$, choose $\nu \ge 2(\|\nabla^2_{pp} H^0\|_{\infty} + 1)$ and deduce that, for any \mathbb{F}^0 -stopping time τ with values in $[0, T]$,

$$
\sigma_0^2 \mathbb{E}^0 \bigg[\int_{\tau}^T E_t^0 |Q_t^0|^2 dt \, | \, \mathcal{F}_{\tau}^0 \bigg] \leq \mathbb{E}^0 \bigg[E_T^0 + \int_{\tau}^T E_t^0 \big| f_t^0(X_t^0, \mu_t) - F^0(X_t^0) \big| dt \, | \, \mathcal{F}_{\tau}^0 \bigg].
$$

Thanks to the bounds for P^0 and E^0 and to (B2) again, we obtain

$$
\sigma_0^2 \mathbb{E}^0 \bigg[\int_{\tau}^{T} E_t^0 |Q_t^0|^2 dt \, | \, \mathcal{F}_{\tau}^0 \bigg] \leq C,
$$

with the constant C being independent of σ_0 and T.

Bound [\(3.20\)](#page-28-0) is a BMO bound. It ensures that the Doléans-Dade exponential

$$
\mathcal{E}_t^0 := \mathscr{E}_t \bigg(\int_0^{\cdot} \sigma_0^{-1} \big(\nabla_p H^0(X_r^0, Z_r^0) - \nabla_p H^0(X_r^0, \nabla_{x_0} w^0(r, X_r^0)) \big) \cdot \mathrm{d}B_r^0 \bigg), \quad t \in [0, T],
$$

is a martingale with respect to \mathbb{F}^0 , see [\[40,](#page-86-30) Theorem 2.3]. In particular, the measure

$$
\widetilde{\mathbb{P}}^0 := \mathcal{E}_T^0 \cdot \mathbb{P}^0 \tag{3.23}
$$

is a probability measure. In fact, the very benefit of Lemma [3.8](#page-28-1) is to provide an L^{1+} bound for \mathcal{E}^0 , independently of σ_0 and T. Indeed, from [\[40,](#page-86-30) Theorem 3.1] (together with the fact that $\sigma_0 \ge 1$), we claim

Lemma 3.9. *Under Assumption* (**B**)*, there exist two constants* $\gamma_0 > 0$ *and* $C_0 \ge 0$ *, only depending on the parameters in* **(B)** *except* (σ_0, T) *such that, for any* \mathbb{F}^0 -*stopping time* τ *with values in* $[0, T]$ *,*

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\Big[\big(\mathcal{E}^0_T(\mathcal{E}^0_\tau)^{-1}\big)^{1+\gamma_0}|\,\mathcal{F}^0_\tau\Big]\leq C_0\bigg\}\bigg)=1,
$$

and

$$
\mathbb{P}^0\bigg(\bigg\{\mathbb{E}^0\bigg[\exp\bigg(\gamma_0\sigma_0^2\int_{\tau}^T|Z_r^0-\nabla_{x_0}w^0(r,X_r^0)|^2\mathrm{d}r\bigg)\,|\,\mathcal{F}_\tau^0\bigg]\leq C_0\bigg\}\bigg)=1.
$$

 \Box

4. Weak formulation of the major-minor MFG

4.1. Girsanov transformation of the coupled major-minor forward-backward system

The preliminary estimates we obtained in the previous section allow us to make a change of measure that leads to a new formulation of the major-minor stochastic MFG coupled system, which we call weak formulation (precisely because it is formulated on a new probability space depending on the solution itself). On an arbitrary filtered probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ satisfying the usual conditions and equipped with a Brownian motion $(B_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d , we introduce the tilted Brownian motion

$$
\widetilde{B}_t^0 := B_t^0 - \sigma_0^{-1} \int_0^t \left[\nabla_p H^0(X_r^0, Z_r^0) - \nabla_p H^0(X_r^0, \nabla_{x_0} w(r, X_r^0)) \right] dr, \quad t \in [0, T].
$$

Thanks to Lemma [3.8,](#page-28-1) we know that $\mathbf{B}^0 = (B_t^0)_{0 \le t \le T}$ is a Brownian motion under \mathbb{P}^0 defined in [\(3.23\)](#page-29-1), and then (2.11) – (2.12) become

$$
dX_t^0 = -\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0)) dt + \sigma_0 d\widetilde{B}_t^0,
$$

\n
$$
dY_t^0 = \left[-f_t^0(X_t^0, \mu_t) - L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0)) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0)) \right] dt + \sigma_0 Z_t^0 \cdot d\widetilde{B}_t^0, \quad t \in [0, T],
$$

\n
$$
Y_T^0 = g^0(X_T^0, \mu_T),
$$
\n(4.1)

and

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \mu_t \right) = 0, \quad \text{on } (0, T) \times \mathbb{T}^d,
$$
\n
$$
\begin{aligned}\n\mathbf{d}_t u_t(x) &= \left(-\frac{1}{2} \Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t) \right) \mathbf{d}t \\
&\quad + \left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0)) \right) \cdot v_t^0(x) \mathbf{d}t + \sigma_0 v_t^0(x) \cdot \mathbf{d} \widetilde{B}_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d, \\
u_T(x) &= g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d.\n\end{aligned}
$$
\n
$$
(4.2)
$$

This is the system that we use below to get estimates on the values of the Major/Minor MFG. The very interest is that the forward equation of the major player is now independent of the remaining three equations.

It is important to observe that the Girsanov transformation (3.23) does not impact the conclusion of Proposition [3.6:](#page-26-1) provided we change γ and C in the statement, the bound therein remains true under the tilted measure. This follows from the fact that, with γ_0 as in Lemma [3.9,](#page-29-2) for any \mathbb{F}^0 -stopping time τ with values in $[0, T]$,

$$
\tilde{\mathbb{E}}^{0}\left[\exp\left(\gamma \frac{\gamma_{0}}{1+\gamma_{0}}\sigma_{0}^{2}\int_{\tau}^{T}\left\|\bar{v}_{r}^{0}\right\|_{\lfloor\delta\rfloor-(d/2+1)}^{2}\mathrm{d}r\right)|\mathcal{F}_{\tau}^{0}\right]
$$
\n
$$
=\mathbb{E}^{0}\left[\mathcal{E}_{T}\mathcal{E}_{\tau}^{-1}\exp\left(\gamma \frac{\gamma_{0}}{1+\gamma_{0}}\sigma_{0}^{2}\int_{\tau}^{T}\left\|\bar{v}_{r}^{0}\right\|_{\lfloor\delta\rfloor-(d/2+1)}^{2}\mathrm{d}r\right)|\mathcal{F}_{\tau}^{0}\right]
$$
\n
$$
\leq\mathbb{E}^{0}\left[\left(\mathcal{E}_{T}\mathcal{E}_{\tau}^{-1}\right)^{1+\gamma_{0}}|\mathcal{F}_{\tau}^{0}\right]^{1/(1+\gamma_{0})}\mathbb{E}^{0}\left[\exp\left(\gamma \sigma_{0}^{2}\int_{\tau}^{T}\left\|\bar{v}_{r}^{0}\right\|_{\lfloor\delta\rfloor-(d/2+1)}^{2}\mathrm{d}r\right)|\mathcal{F}_{\tau}^{0}\right]^{\gamma_{0}/(1+\gamma_{0})}
$$
\n
$$
\leq C_{0}^{1/(1+\gamma_{0})}C^{\gamma_{0}/(1+\gamma_{0})},
$$

with C_0 as in Lemma [3.9.](#page-29-2)

Letting

$$
\tilde{\gamma}_0:=\gamma_0 \frac{\gamma_0}{1+\gamma_0}, \quad \tilde{\gamma}:=\gamma \frac{\gamma_0}{1+\gamma_0}, \quad \tilde{C}:=C_0^{1/(1+\gamma_0)} C^{\gamma_0/(1+\gamma_0)},
$$

$$
\widetilde{\mathbb{E}}^{0}\bigg[\exp\bigg(\tilde{\gamma}\sigma_0^2 \int_{\tau}^{T} \left\|\bar{v}_r^0\right\|_{\lfloor\lambda\rfloor - (d/2+1)}^2 dr\bigg) \,|\,\mathcal{F}_\tau^0\bigg] \leq \tilde{C},
$$
\n
$$
\widetilde{\mathbb{E}}^{0}\bigg[\exp\bigg(\tilde{\gamma}_0\sigma_0^2 \int_{\tau}^{T} |Z_r^0 - \nabla_{x_0} w^0(X_r^0)|^2 dr\bigg) \,|\,\mathcal{F}_\tau^0\bigg] \leq \tilde{C}_0.
$$
\n(4.3)

4.2. Strong formulation of the tilted system and related linearized system

We now come to the heart of the paper. We study the tilted system when posed in the strong form, meaning on the original probability space, with the given B^0 as driving Brownian motion. In clear, we directly address

$$
dX_t^0 = -\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0)) dt + \sigma_0 dB_t^0,
$$

\n
$$
dY_t^0 = \left[-f_t^0(X_t^0, \mu_t) - L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0)) \right.
$$

\n
$$
+ Z_t^0 \cdot (\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0))) \right] dt + \sigma_0 Z_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$

\n
$$
Y_T^0 = g^0(X_T^0, \mu_T),
$$
\n(4.4)

and

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \mu_t \right) = 0, \quad \text{on } (0, T) \times \mathbb{T}^d,
$$

\n
$$
\begin{aligned}\n\mathbf{d}_t u_t(x) &= \left(-\frac{1}{2} \Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t) \right) \mathbf{d}t \\
&\quad + \left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(X_t^0)) \right) \cdot v_t^0(x) \mathbf{d}t + \sigma_0 v_t^0(x) \cdot \mathbf{d}B_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d, \\
u_T(x) &= g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d,\n\end{aligned}
$$
\n(4.5)

on the space $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$.

The following statement clarifies the passage from strong existence and uniqueness for (2.11) – (2.12) to strong existence and uniqueness for (4.4) – (4.5) :

Proposition 4.1. *Under Assumption* (**B**), assume further that, on any arbitrary probabilistic set-up $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$ *and for a fixed initial condition* $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the system (2.11) – (2.12) has a unique solution in the sense of *Definition* [2.10.](#page-9-1) Assume also that there exist two bounded and measurable mappings $\mathcal{U}^0: [0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ and $\mathcal{U}:[0,T]\times\mathbb{R}^d\times\mathbb{T}^d\times\mathcal{P}(\mathbb{T}^d)\to\mathbb{R}$, with \mathcal{U}^0 and \mathcal{U} being both differentiable in the \mathbb{R}^d -variable, with \mathcal{U} being differentiable in the \mathbb{T}^d -variable and with \mathcal{U}^0 and $\nabla_x\mathcal{U}$ being Lipschitz continuous in the \mathbb{R}^d and $\mathcal{P}(\mathbb{T}^d)$ -variables uniformly in the other *variables, such that,* **P** ⁰ *almost surely,*

$$
Y_t^0 = \mathcal{U}^0(t, X_t^0, \mu_t), \ Z_t^0 = \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t), \quad t \in [0, T],
$$

\n
$$
u_t(x) = \mathcal{U}(t, X_t^0, x, \mu_t), \ v_t^0(x) = \nabla_{x_0} \mathcal{U}(t, X_t^0, x, \mu_t), \quad (t, x) \in [0, T] \times \mathbb{T}^d.
$$
\n(4.6)

Then, on the same (and thus on any) probabilistic set-up $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}^0)$ *and for the same initial condition* $(x_0, \mu) \in$ $\mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the system [\(4.4\)](#page-31-0)–[\(4.5\)](#page-31-1) has a unique (hence strong) solution satisfying the requirements of Definition [2.10](#page-9-1) *together with*

$$
\sup_{\tau} \left\| \mathbb{E}^0 \left[\int_{\tau}^T \|v^0_r\|_{\lfloor \lambda \rfloor - (d/2 + 1)}^2 \mathrm{d}r \, \big| \, \mathcal{F}_\tau \right] \right\|_{L^\infty(\Omega^0, \mathbb{P}^0)} < \infty. \tag{4.7}
$$

Remark 4.2. The fact that existence and uniqueness hold true on any $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$ (equipped with a Brownian motion B^0) implies that any solution constructed on a sub-filtration of \mathbb{F}^0 (still carrying B^0) must coincide with the solution constructed on \mathbb{F}^0 . In particular, without any loss of generality, \mathbb{F}^0 can always be taken as the filtration generated by \mathcal{F}_0^0 and B^0 . In this way, we can recover the setting of Section [3.](#page-18-0)

Major Minor MFGs 33

Proof. Existence of a (weak) solution to (4.4) – (4.5) follows from the aforementioned Girsanov argument (as before, the word 'weak' means that the solution is driven by another Brownian motion, different from the original B^0). This weak solution satisfies the integrability conditions in Definition [2.10](#page-9-1) because (ii) and (iii) in item 1 in Definition 2.10 are preserved by BMO changes of measure (see [\[40](#page-86-30), Theorem 3.3]) and (ii) in item 2 is also preserved. The bound (4.7) is a consequence of Remark [3.7.](#page-28-2) In fact, the Lipschitz property of \mathcal{U}^0 gives here an L^{∞} bound on \mathbf{Z}^0 that is even stronger than the BMO condition stated in (iii) in item 1 in Definition [2.10.](#page-9-1)

Thanks to (iii) in item 1 of Definition 2.10 (but for (4.4) – (4.5)), the Girsanov transformation can be reverted, hence proving weak uniqueness.

Strong uniqueness is proved by means of (4.6) . By weak uniqueness, any weak solution to (4.4) – (4.5) is in fact obtained by changing the Brownian motion in the original (2.11) – (2.12) . As such, it must satisfy [\(4.6\)](#page-31-3). Inserting the representation [\(4.6\)](#page-31-3) into the forward equation of [\(4.5\)](#page-31-1) (which is the same as the forward equation of [\(2.12\)](#page-9-0)) and using the fact that $\nabla_x U$ is Lipschitz in the $\mathcal{P}(\mathbb{T}^d)$ -argument, one gets (following the proof of Lemma [2.9\)](#page-8-1) that the pair $(\mathbf{X}^0, \boldsymbol{\mu})$ is necessarily progressively-adapted to the (usual augmentation of the) filtration generated by B^0 and pathwise unique. By [\(4.6\)](#page-31-3), we deduce that solutions to [\(4.4\)](#page-31-0)–[\(4.5\)](#page-31-1) are progressively-adapted to the filtration generated by B^0 and pathwise unique.

Existence and uniqueness of a solution to the system (2.11) – (2.12) (which is the main purpose of this section) will be eventually established in Subsection [4.6](#page-41-1) (using auxiliary results proven in Section [5\)](#page-50-0). At this stage, we are given a flow of solutions

$$
\big((\bm{X}^{0,x_0,\mu},\bm{Y}^{0,x_0,\mu},\bm{Z}^{0,x_0,\mu}),(\bm{\mu}^{x_0,\mu},\bm{u}^{x_0,\mu},\bm{v}^{0,x_0,\mu})\big)_{x_0,\mu}
$$

to the systems [\(4.4\)](#page-31-0) and [\(4.5\)](#page-31-1), parametrized by $(x_0, \mu) \in \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$ (x_0 should be understood as $X_0^{0,x_0,\mu}$ and μ as $\mu_0^{x_0,\mu}$). Implicitly, these solutions are required to satisfy the same constraints as in Definition [2.10.](#page-9-1)

Throughout, we fix $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and $(x'_0, \mu') \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, and we consider the system satisfied by the following derivatives, for $t \in [0, T]$:

$$
\left(\delta X_t^0, \delta Y_t^0, \delta Z^0\right) = \frac{\mathrm{d}}{\mathrm{d}\varepsilon}_{|\varepsilon=0} \left(X_t^{0, x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}, Y_t^{0, x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}, Z_t^{0, x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}\right)
$$

$$
\left(\delta \mu_t, \delta u_t, \delta v_t^0\right) = \frac{\mathrm{d}}{\mathrm{d}\varepsilon}_{|\varepsilon=0} \left(\mu_t^{x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}, u_t^{x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}, v_t^{0, x_0 + \varepsilon(x_0' - x_0), \mu + \varepsilon(\mu' - \mu)}\right).
$$

The sense given to the above derivatives will be clarified below (see in particular the proof of Proposition [4.4](#page-34-0) in Subsection [4.7\)](#page-45-0). What matters now is the form of the system that is satisfied by the two triples $(\delta X^0, \delta Y^0, \delta Z^0)$ and $(\delta \mu, \delta u, \delta v^0)$, which we call 'linearized' processes. Just say that the first triple takes values in $\mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$ whilst the second one takes values in a functional space, namely ν is Schwartz-distributional valued and u and v^0 are functional valued. Throughout, we use the following notation. For a generic (possibly random and say real-valued) function Φ of (some of) the inputs (X_t^0, Z_t^0, μ_t) , we denote by $\delta[\Phi(X_t^0, Z_t^0, \mu_t)]$ the term

$$
\delta[\Phi(X_t^0, Z_t^0, \mu_t)] := \nabla_x \Phi(X_t^0, Z_t^0, \mu_t) \cdot \delta X_t^0 + \nabla_z \Phi(X_t^0, Z_t^0, \mu_t) \cdot \delta Z_t^0 + (\delta_\mu \Phi(X_t^0, Z_t^0, \mu_t)(\cdot), \delta \mu_t).
$$

Then, using the relationship (see (2.3))

$$
- L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0})) + Z_{t}^{0} \cdot (\nabla_{p}H^{0}(X_{t}^{0}, Z_{t}^{0}) - \nabla_{p}H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0})))
$$

\n
$$
= H^{0}(X_{t}^{0}, Z_{t}^{0}) - (Z_{t}^{0} - \nabla_{x_{0}} w^{0}(t, X_{t}^{0})) \cdot \nabla_{p}H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0}))
$$

\n
$$
- \nabla_{x_{0}} w^{0}(t, X_{t}^{0}) \cdot \nabla_{p}H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0}))
$$

\n
$$
= (H^{0}(X_{t}^{0}, Z_{t}^{0}) - H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0}))) - (Z_{t}^{0} - \nabla_{x_{0}} w^{0}(t, X_{t}^{0})) \cdot \nabla_{p}H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0}))
$$

\n
$$
- L^{0}(X_{t}^{0}, -\nabla_{p}H^{0}(X_{t}^{0}, \nabla_{x_{0}} w^{0}(t, X_{t}^{0}))),
$$

and noticing that

$$
\delta \left[-H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) + \nabla_{x_0} w^0(t, X_t^0) \cdot \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right]
$$

= $-\nabla_{x_0} H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \cdot \delta X_t^0 + \nabla_{x_0} w^0(t, X_t^0) \cdot \delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right],$

we obtained, as linearized system for the major player,

$$
d\delta X_t^0 = -\delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] dt,
$$

34

$$
d\delta Y_t^0 = -\delta \left[f_t^0(X_t^0, \mu_t) \right] dt + \left[-\left(Z_t^0 - \nabla_{x_0} w^0(t, X_t^0) \right) \cdot \delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] - \delta \left[L^0(X_t^0, -\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] \right] dt + \left(\nabla_{x_0} H^0(X_t^0, Z_t^0) - \nabla_{x_0} H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \delta X_t^0 dt + \left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \delta Z_t^0 dt + \sigma_0 \delta Z_t^0 \cdot dB_t^0, \quad t \in [0, T], \delta Y_T^0 = \delta \left[g^0(X_T^0, \mu_T) \right],
$$
\n(4.8)

with the initial condition $\delta X_0 = x'_0 - x_0$ and where (X_t^0, Y_t^0, Z_t^0) is a shorter notation for $(X_t^{0,x_0,\mu}, Y_t^{0,x_0,\mu}, Z_t^{0,x_0,\mu})$. Similarly, the linearized system of the minor player is

$$
\partial_t \delta \mu_t - \frac{1}{2} \Delta_x \delta \mu_t - \text{div}_x \Big(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \delta \mu_t \Big) - \text{div}_x \Big(\delta \big[\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \big] \mu_t \Big) = 0,
$$

\n
$$
\mathrm{d}_t \delta u_t(x) = \Big(-\frac{1}{2} \Delta_x \delta u_t(x) + \nabla_p H(x, \nabla_x u_t(x)) \cdot \nabla_x \delta u_t(x) - \delta \big[f_t(X_t^0, x, \mu_t) \big] \Big) \mathrm{d}t
$$

\n
$$
+ \Big(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \Big) \cdot \delta v_t^0(x) \mathrm{d}t
$$

\n
$$
+ \delta \Big[\Big(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \Big) \Big] \cdot v_t^0(x) \mathrm{d}t + \sigma_0 \delta v_t^0(x) \cdot \mathrm{d}B_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\delta u_T(x) = \delta \big[g(X_T^0, x, \mu_T) \big], \quad x \in \mathbb{T}^d,
$$

with the initial condition $\delta\mu_0 = \mu' - \mu$ and with the same convention as before that, for a (say smooth real-valued) function $\Phi(\nabla_x u_t(x))$ of $\nabla_x u_t(x)$,

$$
\delta\big[\Phi(\nabla_x u_t(x))\big] = \nabla\Phi(\nabla_x u_t(x))\cdot\nabla_x \delta u_t(x).
$$

Solutions to the two systems (4.8) – (4.9) are understood in the following sense:

Definition 4.3. Let $((X^0, Y^0, Z^0), (\mu, u, v^0))$ be a solution to [\(4.4\)](#page-31-0)–[\(4.5\)](#page-31-1) in the sense of Definition [2.10,](#page-9-1) for an initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. Then, for another $(x'_0, \mu') \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$,

- 1. Given the initial condition $x'_0 x_0$ and an \mathbb{F}^0 -adapted process $\delta \mu = (\delta \mu_t)_{0 \le t \le T}$ with continuous trajectories in $\mathcal{C}^{-\lfloor \Delta \rfloor + (d/2+1)}(\mathbb{T}^d)$, we call solution to [\(4.8\)](#page-33-0) any \mathbb{F}^0 -progressively measurable process $(\delta \mathbf{X}^0, \delta \mathbf{Y}^0, \delta \mathbf{Z}^0)$ with values in $\mathbb{R}^d \times \mathbb{R} \times \mathbb{R}^d$, such that: (i) $\delta \mathbf{X}^0$ and $\delta \mathbf{Y}^0$ have continuous trajectories; (ii) $\sup_{0 \le t \le T} |\delta X_t^0| \in L^\infty(\Omega^0, \mathbb{P}^0)$, $\sup_{0 \leq t \leq T} |\delta Y_t^0| \in L^{\infty}(\Omega^0, \mathbb{P}^0);$ (iii) $\sup_{\tau} \| \mathbb{E}^0[\int_{[\tau, T]} |\delta Z_t^0|^2 \mathrm{d}t | \mathcal{F}^0_{\tau}]\|_{L^{\infty}(\Omega^0, \mathbb{P}^0)} < \infty$, the supremum being taken over all stopping times τ ; (iv) the system [\(4.8\)](#page-33-0) is satisfied \mathbb{P}^0 -almost surely.
- 2. Given the initial condition $\mu' \mu$, an \mathbb{F}^0 -adapted process $\delta \mathbf{X}^0 = (\delta X_t^0)_{0 \le t \le T}$ with $\sup_{0 \le t \le T} |\delta X_t^0| \in L^\infty(\Omega^0, \mathbb{P}^0)$ and continuous trajectories in \mathbb{R}^d , and an \mathbb{F}^0 -progressively measurable process $\delta \mathbf{Z}^0 = (\delta Z_t^0)_{0 \le t \le T}$ with $\sup_{\tau} \|\mathbb{E}^0[\int_{[\tau,T]} |\delta Z_t^{\hat{0}}|^2 dt | \mathcal{F}_{\tau}^0] \|_{L^{\infty}(\Omega^0,\mathbb{P}^0)} < \infty$, we call solution to [\(4.9\)](#page-33-1) any *Bochner* \mathbb{F}^0 -progressively measurable process $(\delta \mu, \delta u, \delta v^0)$ with values in $C^{-1}(\mathbb{T}^d) \times C^{3+\epsilon}(\mathbb{T}^d) \times C^{3+\epsilon}(\mathbb{T}^d)$ for some $\epsilon > 0$, such that: (i) $(\delta \mu, \delta u)$ has continuous trajectories in $C^{-1}(\mathbb{T}^d) \times C^{3+\epsilon}(\mathbb{T}^d)$; (ii) $\sup_{0 \le t \le T} \|\delta u_t\|_{3+\epsilon} \in L^{\infty}(\Omega^0, \mathbb{P}^0)$, $\sup_{0 \le t \le T} \|\delta \mu_t\|_{-1} \in$ $L^{\infty}(\Omega^0, \mathbb{P}^0)$; (iii) $\mathbb{E}^0[\int_0^T \|\delta v_t^0\|_{3+\epsilon}^2 dt] < \infty$; (iv) the forward equation in [\(4.9\)](#page-33-1) is satisfied \mathbb{P}^0 -almost surely in the weak sense; (v) the backward equation in [\(4.9\)](#page-33-1) is satisfied \mathbb{P}^0 -almost surely in the classical sense.
- 3. For a given initial condition $(x'_0 x_0, \mu' \mu)$, we call a solution to the coupled systems [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1) a pair $((\delta \mathbf{X}^{0}, \delta \mathbf{Y}^{0}, \delta \mathbf{Z}^{0}), (\delta \boldsymbol{\mu}, \delta \boldsymbol{u}, \delta \boldsymbol{v}^{0}))$ satisfying items 1 and 2 right above.

Regarding (iv) in item 2, $\delta\mu$ is said to satisfy [\(4.9\)](#page-33-1) \mathbb{P}^0 -almost surely in the weak sense if, \mathbb{P}^0 , for any test function function $\varphi : [0, T] \times \mathbb{T}^d \to \mathbb{R}$ in the (separable) space of functions that are once differentiable in t and three times differentiable in space, with jointly continuous derivatives, it holds, for any $t \in [0, T]$,

$$
\left(\varphi(t,\cdot),\delta\mu_t\right) - \left(\varphi(0,\cdot),\delta\mu_0\right) = \int_0^t \left(\partial_t\varphi(s,\cdot),\delta\mu_s\right)ds + \frac{1}{2}\int_0^t \left(\Delta_x\varphi(s,\cdot),\delta\mu_s\right)ds + \int_0^t \left(\nabla_x\varphi(s,\cdot)\cdot\nabla_p H(\cdot,\nabla_x u_s(\cdot)),\delta\mu_s\right)ds + \int_0^t \left(\nabla_x\varphi(s,\cdot)\cdot\delta\left[\nabla_p H(\cdot,\nabla_x u_s(\cdot))\right],\mu_s\right)ds.
$$
\n(4.10)

Regarding the solvability of the system [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1), the following statement is proven in Subsection [4.7:](#page-45-0)

Proposition 4.4. *Let Assumption* (B) *and the assumptions of Proposition [4.1](#page-31-4) hold true (for any initial condition of the system* [\(2.11\)](#page-8-2)–[\(2.12\)](#page-9-0)). Assume also that there exist two positive reals α and L such that, for all $t \in [0, T]$, $\mathcal{U}^0(t, \cdot, \cdot) \in$ $\mathscr{D}^0(L,1)$ and $\mathcal{U}(t,\cdot,\cdot,\cdot) \in \mathscr{D}(L,1,3+\alpha)$, and all $(t,x_0,\mu) \in [0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$,

$$
\lim_{h \to 0} \left(\left\| \nabla_{x_0} \mathcal{U}(t+h, x_0, \cdot, \mu) - \nabla_{x_0} \mathcal{U}(t, x_0, \cdot, \mu) \right\|_{3+\alpha} \right)
$$
\n
$$
+ \sup_{l \in \{0, 1\}} \sup_{y \in \mathbb{T}^d} \left\| \nabla_y^l \delta_\mu \mathcal{U}(t+h, x_0, \cdot, \mu, \cdot) - \nabla_y^l \delta_\mu \mathcal{U}(t, x_0, \cdot, \mu, y) \right\|_{3+\alpha} = 0.
$$
\n(4.11)

Let $(x_0, \mu), (x'_0, \mu') \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. Then, on any arbitrary probabilistic set-up $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$, and for the given *initial condition* $(x'_0 - x_0, \mu' - \mu)$ *, the system* [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1) *has a (hence strong) solution* $((\delta X^0, \delta Y^0, \delta Z^0), (\delta \mu, \delta u, \delta v^0))$ *satisfying the requirements of Definition [4.2](#page-33-1) and,* **P** ⁰ *almost surely,*

$$
\delta Y_t^0 = \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t) \cdot \delta X_t^0 + \left(\delta_\mu \mathcal{U}^0(t, X_t^0, \mu_t), \delta \mu_t\right), \quad t \in [0, T],
$$

\n
$$
\delta u_t(x) = \nabla_{x_0} \mathcal{U}(t, X_t^0, x, \mu_t) \cdot \delta X_t^0 + \left(\delta_\mu \mathcal{U}(t, X_t^0, x, \mu_t), \delta \mu_t\right), \quad (t, x) \in [0, T] \times \mathbb{T}^d.
$$
\n(4.12)

In the following Subsections [4.3,](#page-34-1) [4.4](#page-35-0) and [4.5,](#page-37-0) the assumptions of Propositions [4.1](#page-31-4) and [4.4](#page-34-0) are assumed to hold **true (in addition to Assumption (B)).** Namely, there exist two positive reals $\kappa, \alpha > 0$ and two mappings \mathcal{U}^0 : $[0, T] \times$ $\mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ and $\mathcal{U}: [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ in $\mathscr{D}^0(L, 3 + \alpha)$ and $\mathscr{D}(L, 1, 3 + \alpha)$ respectively, satisfying the continuity property [\(4.11\)](#page-34-2), such that the (hence unique) solution to (2.11) – (2.12) satisfies the representation formula [\(4.6\)](#page-31-3).

4.3. Tilting the linearized system

The objective is to provide (*a priori*) estimates, for the linearized systems [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1), that are independent of T . This is the key point in the study of the solvability of the (double) forward-backward system (4.4) – (4.5) . Our strategy relies on a new change of measure which we explain now. Indeed, we know (under the standing assumption) that the solution satisfies the analogue of [\(4.3\)](#page-31-5) but under **P** 0 . Therefore, we can apply a new Girsanov transformation (see [\[40,](#page-86-30) Theorem 2.3]). Letting

$$
\overline{\mathcal{E}}_t := \mathcal{E}_t \left(-\sigma_0^{-1} \int_0^{\cdot} \left(\nabla_p H^0(X_r^0, Z_r^0) - \nabla_p H^0(X_r^0, \nabla_{x_0} w^0(r, X_r^0)) \right) \cdot \mathrm{d}B_r^0 \right), \quad t \in [0, T], \tag{4.13}
$$

we have that $(\overline{\mathcal{E}}_t)_{0 \le t \le T}$ is an \mathbb{F}^0 -martingale (under \mathbb{P}^0) and we can define the probability measure

$$
\overline{\mathbb{P}}^0 := \overline{\mathcal{E}}_T \cdot \mathbb{P}^0. \tag{4.14}
$$

Then, following the proof of Lemma 3.9 , and then proceeding as in the derivation of (4.3) , we deduce

Lemma 4.5. *Under Assumption* (**B**)*, there exist constants* $\overline{\gamma}$, $\overline{\gamma}$ ₀ > 0 *and* \overline{C} , \overline{C} ₀ ≥ 0*, only depending on the parameters in* **(B)** expect (σ_0, T) , such that, for any \mathbb{F}^0 -stopping time τ with values in $[0, T]$,

$$
\overline{\mathbb{E}}^{0}\left[\exp\left(\overline{\gamma}\sigma_{0}^{2}\int_{\tau}^{T}\left\|\overline{v}_{r}^{0}\right\|_{\lfloor\delta\rfloor-(d/2+1)}^{2}\mathrm{d}r\right)\right|\mathcal{F}_{\tau}^{0}\right] \leq \overline{C},
$$
\n
$$
\overline{\mathbb{E}}^{0}\left[\exp\left(\overline{\gamma}_{0}\sigma_{0}^{2}\int_{\tau}^{T}|Z_{r}^{0}-\nabla_{x_{0}}w^{0}(r,X_{r}^{0})|^{2}\mathrm{d}r\right)|\mathcal{F}_{\tau}^{0}\right] \leq \overline{C}_{0}.
$$
\n(4.15)

Obviously, under the new measure \overline{P}^0 , the process

$$
\overline{B}_t^0 := B_t^0 + \sigma_0^{-1} \int_0^t \left(\nabla_p H^0(X_r^0, Z_r^0) - \nabla_p H^0(X_r^0, \nabla_{x_0} w^0(r, X_r^0)) \right) dr, \quad t \in [0, T],
$$

is an \mathbb{F}^0 -Brownian motion and the forward-backward system (4.8) – (4.9) writes

 $d\delta X_t^0 = -\delta \big[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \big] dt,$

$$
d\delta Y_t^0 = -\delta \left[f_t^0(X_t^0, \mu_t) \right] dt + \left[-\left(Z_t^0 - \nabla_{x_0} w^0(t, X_t^0) \right) \cdot \delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] - \delta \left[L^0(X_t^0, -\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] \right] dt + \left(\nabla_{x_0} H^0(X_t^0, Z_t^0) - \nabla_{x_0} H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \delta X_t^0 dt + \sigma_0 \delta Z_t^0 \cdot d\overline{B}_t^0, \quad t \in [0, T],
$$
\n(4.16)
\n
$$
\delta Y_T^0 = \nabla_{x_0} g^0(X_T^0, \mu_T) \delta X_T^0 + (\delta_\mu g^0(X_T^0, \mu_T), \delta \mu_T),
$$

and

$$
\partial_t \delta \mu_t - \frac{1}{2} \Delta_x \delta \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \delta \mu_t \right) - \text{div}_x \left(\delta \left[\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \right] \mu_t \right) = 0,
$$

\n
$$
\mathrm{d}_t \delta u_t(x) = \left(-\frac{1}{2} \Delta_x \delta u_t(x) + \nabla_p H(x, \nabla_x u_t(x)) \cdot \nabla_x \delta u_t(x) - \delta \left[f_t(X_t^0, x, \mu_t) \right] \right) \mathrm{d}t
$$

\n
$$
+ \delta \left[\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \right] \cdot v_t^0(x) \mathrm{d}t + \sigma_0 \delta v_t^0(x) \cdot \mathrm{d} \overline{B}_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\delta u_T(x) = \nabla_{x_0} g(X_T^0, x, \mu_T) \delta X_T^0 + (\delta_\mu g(X_T^0, x, \mu_T), \delta \mu_T),
$$
\n(4.17)

where $((X^0, Y^0, Z^0), (\mu, u, v^0))$ solves (under the mesure $\overline{\mathbb{P}}^0$)

$$
dX_t^0 = -\nabla_p H^0(X_t^0, Z_t^0) dt + \sigma_0 d\overline{B}_t^0, \quad t \in [0, T],
$$

\n
$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0))\right) dt + \sigma_0 Z_t^0 \cdot d\overline{B}_t^0, \quad t \in [0, T],
$$
\n
$$
X_0^0 = x_0, \quad Y_T^0 = g^0(X_T^0, \mu_T).
$$
\n(4.18)

and

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t) \mu_t \right) = 0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mathbf{d}_t u_t(x) = \left(-\frac{1}{2} \Delta_x u_t(x) + H(x, \nabla_x u_t(x)) - f_t(X_t^0, x, \mu_t) \right) \text{d}t + \sigma_0 v_t^0(x) \cdot \text{d}\overline{B}_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mu_0 = \mu, \quad u_T(x) = g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d.
$$
 (4.19)

4.4. A priori estimates for the linearized systems

We now state several standard *a priori* estimates for the linearized systems [\(4.16\)](#page-35-1) and [\(4.17\)](#page-35-2). All of them are borrowed from the book [\[14](#page-86-2)]. As a preliminary observation, we notice from the regularity of w^0 that

Lemma 4.6. *Under Assumption* (B)*, there exists a constant* C_T *, only depending on the parameters in* (B)*, such that*

$$
\sup_{0 \le t \le T} |\delta X_t^0| \le C_T |x_0 - x'_0|.
$$

Proof. This is a straightforward consequence of Gronwall's lemma.

We continue with the analysis of the backward equation.

Lemma 4.7. *Under Assumption* (B)*, there exist two constants* C_T *and* C *, only depending on the parameters in* (B) *but with* C being independent of σ_0 and T, such that, with probability 1, for any $t \in [0,T]$, and for $\pi \in (0, \lfloor \Delta \rfloor - (d/2+1)] \setminus \mathbb{N}$,

$$
\|\delta u_t\|_{\mathcal{R}} \leq C_T |x_0 - x'_0| + C \overline{\mathbb{E}}^0 \bigg[\|\delta \mu_T\|_{-\mathcal{R}} + \int_t^T \bigg(\|\delta \mu_r\|_{-\mathcal{R}} + |\delta Z_r^0| \|\bar{v}_r^0\|_{\mathcal{R}} \bigg) \, \mathrm{d} r \, |\mathcal{F}_t^0 \bigg],
$$

where we recall the notation $\bar{v}_t^0 = v_t^0 - \int_{\mathbb{T}^d} v_t^0(x) dx$, see [\(3.17\)](#page-26-2).

Proof. The strategy is inspired from a duality argument developed in the proof of Lemma 4.3.2 in [\[14\]](#page-86-2). It relies on Lemma [6.3](#page-85-1) in Appendix (which we already used in the previous section, see [\(3.15\)](#page-25-0) and [\(3.16\)](#page-25-1)). For $t_0 \in [0, T]$ and for q a smooth (deterministic) function with $||q||_{\neg r} \leq 1$, we consider the conservation equation

$$
\partial_t q_t - \frac{1}{2} \Delta_x q_t - \text{div}_x (\nabla_p H(\cdot, \nabla_x u_t(\cdot)) q_t) = 0, \quad t \in [t_0, t]; \quad q_{t_0} = q,
$$

 \Box
we compute

$$
d(\delta u_t, q_t) = -\left(\delta \left[f_t(X_t^0, x, \mu_t)\right], q_t\right)dt + \delta \left[\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0))\right)\right] \cdot (q_t, v_t^0)dt
$$

+ $\sigma_0(q_t, \delta v_t^0) \cdot d\overline{B}_t^0, \quad t \in [t_0, T].$

And then, using the regularity properties of g and f together with Lemmas [4.6](#page-35-0) and [6.3,](#page-85-0) we obtain

$$
(\delta u_{t_0}, q) = \overline{\mathbb{E}}^0 \left[(\delta u_T, q_T) + \int_{t_0}^T \left(\delta \big[f_t(X_t^0, x, \mu_t) \big], q_t \right) dt - \int_{t_0}^T \delta \Big[\Big(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \Big) \Big] \cdot (q_t, v_t^0) dt \, | \, \mathcal{F}_{t_0}^0 \right] \n\leq C_T |x_0 - x_0'| + C \overline{\mathbb{E}}^0 \left[\| \delta \mu_T \|_{-\tau_0} + \int_{t_0}^T \Big(\| \delta \mu_t \|_{-\tau_0} + \Big(| \delta X_t^0| + | \delta Z_t^0| \Big) \, \| \bar{v}_t^0 \|_{\tau_0} \Big) dt \, | \, \mathcal{F}_{t_0}^0 \right].
$$

By [\(4.15\)](#page-34-0) and Lemma [4.6](#page-35-0) again, the above bound can be rewritten (for a new value of the constant C_T)

$$
(\delta u_{t_0}, q) \leq C_T |x_0 - x'_0| + C \overline{\mathbb{E}}^0 \bigg[\|\delta \mu_T\|_{-\tau_1} + \int_{t_0}^T \bigg(\|\delta \mu_t\|_{-\tau_1} + |\delta Z_t^0| \, \|\bar{v}_t^0\|_{\tau_1} \bigg) \, \mathrm{d}t \, |\, \mathcal{F}_{t_0}^0 \bigg].
$$

For $l \in \{0, \dots, \lfloor n \rfloor\}$, x and h two elements of \mathbb{R}^d and ρ a smooth density on \mathbb{R}^d , we observe that, for $q(\cdot) = (-1)^l \partial^l \rho(\cdot - 1)^{l-1}$ x) (with $\partial^l \rho$ denoting the derivative of ρ along l arbitrary directions of \mathbb{R}^d with possible repetitions) and any $\varphi \in C^{\infty}(\mathbb{T}^d)$ satisfying $\|\varphi\|_{r} \leq 1$,

$$
|(\varphi, q)| \le \left| \nabla^l \varphi * \rho(x) \right| \le 1.
$$

Similarly, for $q(\cdot) = (-1)^{\lfloor n \rfloor} |h|^{-n+\lfloor n \rfloor} [\partial^{\lfloor n \rfloor} \rho(\cdot - x) - \partial^{\lfloor n \rfloor} \rho(\cdot - (x+h)],$

$$
\left|(\varphi,q)\right|\le |h|^{-\mathrm{Tr}+\lfloor \mathrm{Tr}\rfloor}\left|\nabla^{\lfloor \mathrm{Tr}\rfloor}\varphi*\rho(x)-\nabla^{\lfloor \mathrm{Tr}\rfloor}\varphi*\rho(x+h)\right|\le 1.
$$

And then, denoting by $(\rho_n)_{n\geq 1}$ a sequence of mollifiers on \mathbb{R}^d , we deduce that, $\overline{\mathbb{P}}^0$ -almost surely, for any $l \in$ $\{0, \dots, \lfloor n \rfloor\}$, any $n \ge 1$ and any $x, h \in \mathbb{Q}^d$,

$$
\left|\nabla^{l}\left(\delta u_{t_{0}} * \rho_{n}\right)(x)\right| \leq C_{T}|x_{0} - x'_{0}| + C\overline{\mathbb{E}}^{0}\bigg[\|\delta\mu_{T}\|_{-\tau_{1}} + \int_{t_{0}}^{T}\left(\|\delta\mu_{t}\|_{-\tau_{1}} + |\delta Z_{t}^{0}| \|\bar{v}_{t}^{0}\|_{\tau_{1}}\right)dt \,|\,\mathcal{F}_{t_{0}}^{0}\bigg],
$$

and

|h| −˚rffl+⌊˚rffl⌋ ∇ ⌊˚rffl⌋ δu^t⁰ ∗ ρⁿ (x) − ∇⌊˚rffl⌋ δu^t⁰ ∗ ρⁿ (x + h) ≤ C^T |x⁰ − x ′ 0 | + C**E** 0 kδµ^T k−˚rffl + Z ^T t0 ^kδµtk−˚rffl ⁺ [|]δZ⁰ t | kv¯ 0 t k˚rffl ^dt| F⁰ t0 .

Obviously, this holds true almost surely, for any $l \in \{0, \dots, \lfloor n \rfloor\}$, any $n \ge 1$ and any $x, h \in \mathbb{R}^d$. And since δu_{t_0} is already known to be in $C^{3+\epsilon}(\mathbb{T}^d)$ (see Proposition [4.4\)](#page-34-1), we deduce that it belongs to $C^{r}(\mathbb{T}^d)$ with

$$
\|\delta u_{t_0}\|_{\mathcal{R}} \leq C_T |x_0 - x'_0| + C \overline{\mathbb{E}}^0 \bigg[\|\delta \mu_T\|_{-\mathcal{R}} + \int_{t_0}^T \bigg(\|\delta \mu_t\|_{-\mathcal{R}} + |\delta Z_t^0| \|v_t^0\|_{\mathcal{R}} \bigg) dt \, | \, \mathcal{F}_{t_0}^0 \bigg].
$$

Here, we also recall that δu has continuous values in $C^{3+\epsilon}(\mathbb{T}^d)$. We deduce that the above is true, almost surely, for any \Box $t_0 \in [0, T]$.

Lemma 4.8. *Under Assumption* (B)*, there exist an exponent* γ > 0 *and a constant* C*, only depending on the parameters in* (**B**) *except* σ_0 *and* T *, such that, with probability 1, for any* $t \in [0, T]$ *, for any* $\tau \in [1, |\mathcal{A}| - (d/2 + 1)] \setminus \mathbb{N}$ *,*

$$
\|\delta\mu_t\|_{-\tau} \le C\bigg[\|\delta\mu_0\|_{-\tau} + \int_0^t \exp(-\gamma(t-r))\bigg\|\nabla_x \delta u_r \,\mu_r\bigg\|_{-\tau+1} \mathrm{d}r\bigg].
$$

In particular,

$$
\|\delta \mu_t\|_{-\tau}^2 \le C \bigg[\|\delta \mu_0\|_{-\tau}^2 + \int_0^t \exp(-\gamma (t-r)) \big(|\nabla_x \delta u_r|^2, \mu_r \big) dr \bigg],
$$

for a possibly new value of C*.*

Proof. We fix $t_0 \in [0, T]$. We use again a duality argument by computing $d_t(\delta \mu_t, \varphi_t)$, where $(\varphi_t)_{0 \le t \le t_0}$ solves the (random) backward equation

$$
\partial_t \varphi_t = -\frac{1}{2}\Delta \varphi_t + \nabla_p H(\cdot, \nabla_x u_t(\cdot)) \cdot \nabla \varphi_t, \quad t \in [0, t_0]; \quad \varphi_{t_0} = \phi,
$$

where ϕ is a deterministic function in $\mathcal{C}^{\pi}(\mathbb{T}^d)$. Observe that $(\varphi_t)_{0 \le t \le t_0}$ may be anticipative. By Lemma [6.2](#page-84-0) together with the fact that $\nabla_x u$ takes values in $\mathcal{C}^{3+\epsilon}(\mathbb{T}^d)$ for some $\epsilon > 0$, we deduce that $(\varphi_t)_{0 \le t \le t_0}$ takes values in $\mathcal{C}^{3+\epsilon}(\mathbb{T}^d)$. In particular, $(\partial_t \varphi_t)_{0\leq t\leq t_0}$ takes values in $\mathcal{C}^{1+\epsilon}(\mathbb{T}^d)$. Because $\delta\mu$ takes values in $\mathcal{C}^{-1}(\mathbb{T}^d)$, this makes it possible to expand the duality product $(\delta \mu_t, \varphi_t)_{0 \leq t \leq t_0}$. We obtain

$$
\mathrm{d}_t(\delta \mu_t, \varphi_t) = -\Big(\nabla_x \varphi_t, \delta\big[\nabla_p H(\cdot, \nabla_x u_t(\cdot))\big]\mu_t\Big) \mathrm{d}t.
$$

Then,

$$
(\delta\mu_{t_0}, \phi) \leq (\delta\mu_0, \varphi_0) + C \int_0^{t_0} \left\| \nabla_x \varphi_t \right\|_{\mathcal{R}-1} \left\| \delta \left[\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \right] \mu_t \right\|_{-(\mathcal{R}-1)} dt
$$

$$
\leq C \|\phi\|_{\mathcal{R}} \left[\|\delta\mu_0\|_{-\mathcal{R}} + \int_0^{t_0} \exp(-\gamma(t_0 - t)) \left\| \nabla_x \delta u_t \mu_t \right\|_{-(\mathcal{R}-1)} dt \right],
$$

where, to get the last line, we used the identity $\delta \left[\nabla_p H(\cdot, \nabla_x u_t(\cdot))\right] = \nabla^2_{pp} H(\cdot, \nabla_x u_t(\cdot)) \nabla_x \delta u_t$ together with the bound $\|\nabla^2_{pp}H(\cdot,\nabla_x u_t(\cdot))\|_{\mathcal{R}-1}\leq C$, which follows from Proposition [3.4.](#page-22-0) Taking the supremum over ϕ in the unit ball of $\mathcal{C}^{\pi}(\mathbb{T}^d)$, we complete the proof.

4.5. A key functional

We now introduce a key object. For a parameter $A > 0$, we call $(e_t)_{0 \le t \le T}$ the solution to the backward SDE

$$
d_t e_t = \frac{\sigma_0^2}{A} e_t \|\bar{v}_t^0\|_{\lfloor \delta \rfloor - (d/2 + 1)}^2 dt + \ell_t d\overline{B}_t^0, \quad t \in [0, T]; \quad e_T = 1,
$$
\n(4.20)

where we recall the notation $\bar{v}_t^0 = v_t^0 - \int_{\mathbb{T}^d} v_t^0(x) dx$, see [\(3.17\)](#page-26-0).

The solution is given by the following lemma, which is a straightforward consequence of [\(4.15\)](#page-34-0):

Lemma 4.9. *Under Assumption* (B)*, the BSDE* [\(4.20\)](#page-37-0) *has a unique solution, which is given by*

$$
e_t = \overline{\mathbb{E}}^0 \left[\exp \left(-\frac{\sigma_0^2}{A} \int_t^T \|\bar{v}_r^0\|_{\lfloor \lambda \rfloor - (d/2 + 1)}^2 \, \mathrm{d}r \right) | \, \mathcal{F}_t^0 \right], \quad t \in [0, T].
$$

In particular, there exists a constant $c \in (0,1)$ *, only depending on the parameters in* (**B**) *except* σ_0 *and* T *, such that, for* A large enough (independently of σ_0 and T), with probability 1, for all $t \in [0, T]$,

$$
c < e_t \le 1. \tag{4.21}
$$

Proof. For any solution $(e_t)_{0 \le t \le T}$ as in [\(4.20\)](#page-37-0),

$$
d_t \left[e_t \exp \left(-\frac{\sigma_0^2}{A} \int_0^t ||\bar{v}_r^0||^2_{\lfloor \lambda \rfloor - (d/2 + 1)} dr \right) \right] = e_t \exp \left(-\frac{\sigma_0^2}{A} \int_0^t ||\bar{v}_r^0||^2_{\lfloor \lambda \rfloor - (d/2 + 1)} dr \right) \ell_t d\overline{B}_t^0.
$$

Under the standard conditions that

$$
\overline{\mathbb{E}}^0 \bigg[\sup_{0 \le t \le T} |e_t|^2 + \int_0^T |\ell_t|^2 dt \bigg] < \infty,
$$

we see that the right-hand side in the latter expansion yields a martingale. We deduce that

$$
e_t = \overline{\mathbb{E}}^0 \left[\exp \left(-\frac{\sigma_0^2}{A} \int_t^T \|\bar{v}_r^0\|_{\lfloor \delta \rfloor - (d/2 + 1)}^2 \mathrm{d}r \right) | \mathcal{F}_t^0 \right], \quad t \in [0, T],
$$

and there is no difficulty for proving that the above right-hand side induces a solution. In order to prove the estimate (4.21) (which is the key point in the statement), we recall that for any positive random variable ξ with positive values

$$
\overline{\mathbb{E}}^0 \left[\xi \, | \, \mathcal{F}_t^0 \right] \ge \overline{\mathbb{E}}^0 \left[\xi^{-1} \, | \, \mathcal{F}_t^0 \right]^{-1}.
$$

Choosing

$$
\xi = \exp\bigg(-\frac{\sigma_0^2}{A} \int_t^T \|\bar{v}_r^0\|_{\lfloor \delta \rfloor - (d/2 + 1)}^2 \mathrm{d}r\bigg),\,
$$

we complete the proof of the lower bound by using (4.15) . Importantly, we notice that the lower bound c does not depend on A (as soon as the latter one satisfies $1/A \leq \overline{\gamma}$). The upper bound is obvious. 口

Here is the way we use the process $(e_t)_{0 \le t \le T}$. To clarify the dependence on the various parameters in (B), we let $\kappa_{H^0} := \|\nabla_{pp}^2 H^0\|_{\infty}/2$ and we call λ_H the largest real such that $\nabla_{pp}^2 H \ge \lambda_H I_d$. We also introduce

$$
\theta_t := \sup_{x_0 \in \mathbb{R}^d} \sup_{\mu \in \mathcal{P}(\mathbb{T}^d)} \|\delta_\mu f_t^0(x_0, \mu)(\cdot)\|_{\mathbb{T}} + \min(1, \frac{1}{T}), \quad \Theta_t = \int_0^t \theta_s \, ds, \quad t \in [0, T].
$$

By assumption (see (B3)), $\Theta_T = \int_0^T \theta_t dt$ is bounded by a constant independent of T (and of course of σ_0). And by construction, $\theta_t^{-1} \le \max(1,T)$. Then, for three (positive) parameters ε_1 , ε_2 and ϱ , we next compute

$$
\mathrm{d}_t \bigg[\varepsilon_1 \exp(\Theta_t) |\delta Y_t^0|^2 - (\delta \mu_t, \delta u_t) + \varepsilon_2 e_t \bigg(\|\delta \mu_0\|_{-\tau}^2 + \int_0^t \left(|\nabla_x \delta u_r|^2, \mu_r \right) \mathrm{d}r \bigg) \bigg],
$$

for a real $\pi \in [1, \lfloor \Delta \rfloor - (d/2 + 1)] \setminus \mathbb{N}$, where we recall $\delta \mu_0 = \mu - \mu'$.

We start with the expansion of $d_t[\varepsilon_1 \exp(\Theta_t) | \delta Y_t^0|^2]$. Back to [\(4.16\)](#page-35-1), we write

$$
d_t \left[\varepsilon_1 \exp(\Theta_t) |\delta Y_t^0|^2 \right] = \varepsilon_1 \exp(\Theta_t) \left[\sigma_0^2 |\delta Z_t^0|^2 + \theta_t |\delta Y_t^0|^2 + \delta Y_t^0 |\left(\mathcal{O}_{\mathbb{R}^d}(1) + \mathcal{O}_{\mathbb{R}^d}(|Z_t^0 - \nabla_{x_0} w^0(t, X_t^0)|) \right) \cdot \delta X_t^0 \right] + \delta Y_t^0 \left(\mathcal{O}_{\mathcal{C}^{\wedge}(\mathbb{T}^d)}(\theta_t), \delta \mu_t \right) \right] dt + dn_t,
$$

where $(n_t)_{t>0}$ is a generic martingale term, whose precise value may vary from line to line. For a normed space $(E, \|\cdot\|)$ and for a possibly random real $\Delta \geq 0$, $\mathcal{O}_E(\Delta)$ stands for an E-valued term satisfying $\|\mathcal{O}_E(\Delta)\| \leq C|\Delta|$, for a constant C that only depends on the various parameters in (B) except σ_0 and T.

Next, we compute $d_t(\delta \mu_t, \delta u_t)$. By duality, we obtain from [\(4.17\)](#page-35-2):

$$
d_t(\delta \mu_t, \delta u_t)
$$

= $-\left(\delta[\nabla_x u_t], \delta[\nabla_p H(\cdot, \nabla_x u_t(\cdot))] \mu_t\right) dt - \left(\delta \mu_t, \delta[f_t(X_t^0, x, \mu_t)]\right) dt$
+ $\delta\left[\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0))\right)\right] \cdot (v_t^0, \delta \mu_t) dt + dn_t$
= $\left[-\left((\nabla_x \delta u_t)^\top \nabla_{pp}^2 H(\cdot, \nabla_x u_t(\cdot)) \nabla_x \delta u_t, \mu_t\right) - \left((\delta_\mu f_t(X_t^0, \cdot, \mu_t, \cdot), \delta \mu_t(\cdot)), \delta \mu_t(\cdot)\right) + \mathcal{O}_{\mathbb{R}}\left(\frac{|\delta X_t^0|}{|\delta \mu_t|_{-r}}\right) + \mathcal{O}_{\mathbb{R}^d}\left(\frac{|\delta X_t^0|}{|\delta \mu_t|_{-r}}\right) \cdot (v_t^0, \delta \mu_t) + \left(\nabla_{pp}^2 H^0(X_t^0, Z_t^0) \delta Z_t^0\right) \cdot (v_t^0, \delta \mu_t)\right] dt + dn_t.$

Above, we notice that $(v_t^0, \delta \mu_t) = (\bar{v}_t^0, \delta \mu_t)$ because the duality bracket $(\delta \mu_t, 1)$ (with 1 standing for the constant function, equal to 1) is equal to 0.

Lastly, we handle $d_t[\varepsilon_2 e_t(\|\delta\mu_0\|_{-\infty}^2 + \int_0^t (\nabla_x \delta u_r|^2, \mu_r) dr)]$. We have, for all $t \in [0, T]$,

$$
d_{t} \left[\varepsilon_{2} e_{t} \left(\|\delta \mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} \left(\left| \nabla_{x} \delta u_{r} \right|^{2}, \mu_{r} \right) dr \right) \right]
$$

= $\left[\varepsilon_{2} e_{t} \left(\left| \nabla_{x} \delta u_{t} \right|^{2}, \mu_{t} \right) + \frac{\varepsilon_{2} \sigma_{0}^{2}}{A} e_{t} \|\bar{v}_{t}^{0}\|_{\lfloor \delta \rfloor - (d/2+1)}^{2} \left(\|\delta \mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} \left(\left| \nabla_{x} \delta u_{r} \right|^{2}, \mu_{r} \right) dr \right) \right] dt + dn_{t}.$

By combining the last three displays, we get

$$
d_{t}\left[\varepsilon_{1} \exp(\Theta_{t})|\delta Y_{t}^{0}|^{2} - (\delta\mu_{t}, \delta u_{t}) + \varepsilon_{2} e_{t}\left(\|\delta\mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t}(|\nabla_{x}\delta u_{r}|^{2}, \mu_{r})dr\right)\right]
$$

\n
$$
= \varepsilon_{1} \exp(\Theta_{t})\left[\sigma_{0}^{2}|\delta Z_{t}^{0}|^{2} + \theta_{t}|\delta Y_{t}^{0}|^{2}\right]dt
$$

\n
$$
+ \varepsilon_{1} \exp(\Theta_{t})\left[\delta Y_{t}^{0}\left[\left(\mathcal{O}_{\mathbb{R}^{d}}(1) + \mathcal{O}_{\mathbb{R}^{d}}(|Z_{t}^{0} - \nabla_{x_{0}}w^{0}(t, X_{t}^{0})|)\right) \cdot \delta X_{t}^{0}\right] + \delta Y_{t}^{0}\left(\mathcal{O}_{\mathcal{C}^{\tau_{1}}(\mathbb{T}^{d})}(\theta_{t}), \delta\mu_{t}\right)\right]dt
$$

\n
$$
+ \left[\left((\nabla_{x}\delta u_{t})^{\top}\nabla_{pp}^{2}H(\cdot, \nabla_{x}u_{t}(\cdot))\nabla_{x}\delta u_{t}, \mu_{t}\right) + \left((\delta_{\mu}f_{t}(X_{t}^{0}, \cdot, \mu_{t}, \cdot), \delta\mu_{t}(\cdot)), \delta\mu_{t}(\cdot)\right)\right]dt
$$

\n
$$
+ \left[\mathcal{O}_{\mathbb{R}}(|\delta X_{t}^{0}|||\delta\mu_{t}||_{-\tau_{1}}) + \mathcal{O}_{\mathbb{R}^{d}}(|\delta X_{t}^{0}|) \cdot (\bar{v}_{t}^{0}, \delta\mu_{t}) + (\nabla_{pp}^{2}H^{0}(X_{t}^{0}, Z_{t}^{0})\delta Z_{t}^{0}) \cdot (\bar{v}_{t}^{0}, \delta\mu_{t})\right]dt
$$

\n
$$
+ \left[\varepsilon_{2}e_{t}(|\nabla_{x}\delta u_{t}|^{2}, \mu_{t}) + \frac{\varepsilon_{2}\sigma_{0}^{2}}{A}e_{t}|\bar{v}_{t}^{0}||_{\lfloor\delta_{1}-(d/2+1)}\left(\|\delta
$$

where $(n_t)_{0 \le t \le T}$ stands for a generic martingale term. All the terms on the second line of the right-hand side are handled by means of Young's inequality, in such a way that the corresponding term δY_t^0 is 'absorbed' by $\theta_t |\delta Y_t^0|^2$ (which is greater than $\min(1, T^{-1})|\delta Y_t^0|^2$ on the first line of the right-hand side. The two terms on the third line of the right-hand side are non-negative: for the first one, this follows from the convexity of H in the variable p ; for the second one, this follows from the monotonicity of f in the variables (x, μ) . The first two terms on the fourth line of the right-hand side are also handled by means of Young's inequality: among the resulting two terms, one of them is $|\delta X_t^0|^2$ multiplied by a possibly large constant. The last term on the fourth line of the right-hand side is handled by a new application of Young's inequality, with the term δZ_t^0 being now 'absorbed' by $\varepsilon_1 \sigma_0^2 |\delta Z_t^0|^2$ on the first line.

Then, using Lemma [4.9](#page-37-2) together with the fact that $\theta_t^{-1} \le \max(1,T)$, $\theta_t \le \kappa + 1$ and $\Theta_T \le C$, we observe that the above expansion is greater than

$$
d_{t}\left[\varepsilon_{1} \exp(\Theta_{t})|\delta Y_{t}^{0}|^{2} - (\delta\mu_{t}, \delta u_{t}) + \varepsilon_{2} e_{t}\left(\|\delta\mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} (|\nabla_{x}\delta u_{r}|^{2}, \mu_{r}) dr\right)\right]
$$

\n
$$
\geq \left[\frac{\varepsilon_{1}\sigma_{0}^{2}}{2}|\delta Z_{t}^{0}|^{2} + \frac{c\varepsilon_{2}\sigma_{0}^{2}}{2A}\|\bar{v}_{t}^{0}\|_{\lfloor\delta\rfloor-(d/2+1)}^{2} \left(\|\delta\mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} (|\nabla_{x}\delta u_{r}|^{2}, \mu_{r}) dr\right) - \frac{\kappa_{H^{0}}}{\varepsilon_{1}\sigma_{0}^{2}}\|\delta\mu_{t}\|_{-\tau_{1}}^{2}\|\bar{v}_{t}^{0}\|_{\lfloor\delta\rfloor-(d/2+1)}^{2} + \lambda_{H} (|\nabla_{x}\delta u_{t}|^{2}, \mu_{t}) - C_{\varepsilon_{1},\sigma_{0},T}|\delta X_{t}^{0}|^{2} - C_{\varepsilon_{1},\sigma_{0},T}|Z_{t}^{0} - \nabla_{x_{0}}w^{0}(X_{t}^{0})|^{2}|\delta X_{t}^{0}|^{2} - C\varepsilon_{1}\|\delta\mu_{t}\|_{-\tau_{1}}^{2}\right]dt + dn_{t},
$$

for a constant C only depending on the parameters in (B) except σ_0 and T and for a constant $C_{\epsilon_1,\sigma_0,T}$ only depending on ε_1 and the parameters in (B) (including σ_0 and T). We now recall the upper bound for $\|\delta\mu_t\|_{-\infty}^2$ (see Lemma [4.8\)](#page-36-0). Allowing the value of C to vary from line to line, we obtain

$$
d_{t}\left[\varepsilon_{1} \exp(\theta_{t})|\delta Y_{t}^{0}|^{2} - (\delta\mu_{t}, \delta u_{t}) + \varepsilon_{2} e_{t}\left(\|\delta\mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} (|\nabla_{x}\delta u_{r}|^{2}, \mu_{r}) dr\right)\right]
$$

\n
$$
\geq \left[\frac{\varepsilon_{1}\sigma_{0}^{2}}{2}|\delta Z_{t}^{0}|^{2} + \lambda_{H}(|\nabla_{x}\delta u_{t}|^{2}, \mu_{t}) + \left\{\left(\frac{c\varepsilon_{2}\sigma_{0}^{2}}{2A} - \frac{C\kappa_{H^{0}}}{\varepsilon_{1}\sigma_{0}^{2}}\right) \|\bar{v}_{t}^{0}\|_{\infty}^{2} - (d/2+1) - C\varepsilon_{1}\right\} \left(\|\delta\mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{t} \exp(-\gamma(t-r))\left(\left|\nabla_{x}\delta u_{r}\right|^{2}, \mu_{r}\right) dr\right)
$$
\n
$$
- C_{\varepsilon_{1},\sigma_{0},T} |\delta X_{t}^{0}|^{2} - C_{\varepsilon_{1},\sigma_{0},T} |Z_{t}^{0} - \nabla_{x_{0}} w^{0}(X_{t}^{0})|^{2} |\delta X_{t}^{0}|^{2}\right] dt + dn_{t},
$$
\n(4.22)

We deduce the following main inequality:

Proposition 4.10. *Under Assumption* (B)*, there exist a constant* $C \ge 1$ *only depending on the parameters in* (B) *except* (σ_0, T) and a constant $C_{\varepsilon_1,\sigma_0,T}$ only depending on ε_1 and the parameters in (**B**) (including (σ_0, T)) and non-decreasing with *T*, such that, with the two notations $\kappa_{H^0} := \|\nabla^2_{pp} H^0\|_{\infty}/2$ and $\lambda_H := \sup\{\theta : \nabla^2_{pp} H \ge \theta I_d\}$, for *A* and *c* as in *Lemma [4.9,](#page-37-2) and under the following two conditions:*

i.
$$
\varepsilon_1 + \varepsilon_2 < \frac{\lambda_H}{C}
$$
,
ii. $\frac{c\varepsilon_2 \sigma_0^2}{2A} > \frac{C\kappa_{H^0}}{\varepsilon_1 \sigma_0^2}$

it holds, for any $\lceil (n+1)/2 \rceil - (d/2 + 1) \rceil \setminus \mathbb{N}$,

,

$$
|\delta Y_0^0|^2 + \|\delta u_0\|_{\mathcal{R}}^2 \le C_{\varepsilon_1, \sigma_0, T} \left(|x_0 - x'_0|^2 + \|\mu - \mu'\|_{-\mathcal{R}}^2 \right).
$$
 (4.23)

It is worth pointing out that the constant C in the statement may depend on λ_H and κ_{H^0} themselves. This makes the two conditions *i* and *ii* less explicit than might be assumed at first sight. However, we think this formulation is useful for the proof.

Proof. Integrating [\(4.22\)](#page-40-0) from 0 to T, taking expectation and using condition *ii* in the statement, we obtain (for a constant C only depending on the parameters in (**B**) except (σ_0, T))

$$
\begin{split}\n\overline{\mathbb{E}}^{0} \Big[&\varepsilon_{1} \exp(\Theta_{T}) |\delta Y_{T}^{0}|^{2} - (\delta \mu_{T}, \delta u_{T}) + \varepsilon_{2} e_{T} \Big(\|\delta \mu_{0}\|_{-\tau_{1}}^{2} + \int_{0}^{T} \big(|\nabla_{x} \delta u_{r}|^{2}, \mu_{r} \big) dr \Big) \Big] \\
&\geq \overline{\mathbb{E}}^{0} \Big\{ \varepsilon_{1} |\delta Y_{0}^{0}|^{2} - (\delta \mu_{0}, \delta u_{0}) + \varepsilon_{2} e_{0} \|\delta \mu_{0}\|_{-\tau_{1}}^{2} \\
&+ \int_{0}^{T} \Big[\frac{\varepsilon_{1} \sigma_{0}^{2}}{2} |\delta Z_{t}^{0}|^{2} + \lambda_{H} \big(|\nabla_{x} \delta u_{t}|^{2}, \mu_{t} \big) \Big] dt - C \varepsilon_{1} \int_{0}^{T} \Big[\big(|\nabla_{x} \delta u_{t}|^{2}, \mu_{t} \big) \int_{t}^{T} \exp(-\gamma (r - t)) dr \Big] dt \\
&- C_{\varepsilon_{1}, \sigma_{0}, T} \int_{0}^{T} \Big[|\delta X_{t}^{0}|^{2} + |Z_{t}^{0} - \nabla_{x_{0}} w^{0}(X_{t}^{0})|^{2} |\delta X_{t}^{0}|^{2} \Big] dt - C_{\varepsilon_{1}, \sigma_{0}, T} \|\delta \mu_{0}\|_{-\tau_{1}}^{2} \Big\}.\n\end{split} \tag{4.24}
$$

Applying [\(4.15\)](#page-34-0) and Lemma [4.6,](#page-35-0) we get that the whole term on the last line is less $C_{\epsilon_1,\sigma_0,T} |x_0 - x'_0|^2$, for a constant $C_{\epsilon_1,\sigma_0,T}$ depending only on the parameters in (B) (including σ_0 and T) and on ϵ_1 . Moreover, by the monotonicity condition (A3) and Lemma [4.8,](#page-36-0) we have (for possibly new values of $C_{\varepsilon_1,\sigma_0,T}$ and C)

$$
\begin{split}\n\overline{\mathbb{E}}^{0} \Big[& (\delta \mu_{T}, \delta u_{T}) - \varepsilon_{1} \exp(\Theta_{T}) |\delta Y_{T}^{0}|^{2} \Big] \\
&= \overline{\mathbb{E}}^{0} \Big[\delta X_{T}^{0} \cdot (\nabla_{x_{0}} g(X_{T}^{0}, \cdot, \mu_{T}), \delta \mu_{T}) + \Big((\delta_{\mu} g(X_{T}^{0}, \cdot, \mu_{T}, \cdot), \delta \mu_{T}(\cdot)), \delta \mu_{T}(\cdot) \Big) \\
&- \varepsilon_{1} \exp(\Theta_{T}) \Big(\nabla_{x_{0}} g^{0}(X_{T}^{0}, \mu_{T}) \cdot \delta X_{T}^{0} + (\delta_{\mu} g^{0}(X_{T}^{0}, \mu_{T}), \delta \mu_{T}) \Big)^{2} \Big] \\
&\geq -C_{\varepsilon_{1}, \sigma_{0}, T} \overline{\mathbb{E}}^{0} \Big[|\delta X_{T}^{0}|^{2} \Big] - C \varepsilon_{1} \exp(\Theta_{T}) ||\delta \mu_{T}||_{-r_{1}}^{2} \\
&\geq -C_{\varepsilon_{1}, \sigma_{0}, T} |x_{0} - x_{0}'|^{2} - C \varepsilon_{1} \Big(||\delta \mu_{0}||_{-r_{1}}^{2} + \overline{\mathbb{E}}^{0} \Big[\int_{0}^{T} \Big(|\nabla_{x} \delta u_{r}|^{2}, \mu_{r} \Big) dr \Big] \Big). \n\end{split} \tag{4.25}
$$

Therefore, by adding [\(4.24\)](#page-40-1) and [\(4.25\)](#page-41-0), and by dropping the positive term $\varepsilon_2 e_0 \|\delta \mu_0\|_{-\infty}^2$, we obtain (for a new value of C)

$$
\overline{\mathbb{E}}^{0}\left[\left(\lambda_{H}-C(\varepsilon_{1}+\varepsilon_{2})\right)\int_{0}^{T}\left(\left|\nabla_{x}\delta u_{r}\right|^{2},\mu_{r}\right)dr+\frac{\varepsilon_{1}\sigma_{0}^{2}}{2}\int_{0}^{T}\left|\delta Z_{r}^{0}\right|^{2}dr\right]
$$
\n
$$
\leq\left[C_{\varepsilon_{1},\sigma_{0},T}\left(|x_{0}-x_{0}'|^{2}+\left\|\delta\mu_{0}\right\|_{-\infty}^{2}\right)-\varepsilon_{1}\left|\delta Y_{0}^{0}\right|^{2}+\left(\delta u_{0},\delta\mu_{0}\right)\right].
$$
\n(4.26)

Next, we recall Lemmas [4.7](#page-35-3) and [4.8,](#page-36-0) and use [\(4.15\)](#page-34-0) to derive that

$$
\|\delta u_0\|_{\mathcal{R}} \leq C_T |x_0 - x'_0| + C \overline{\mathbb{E}}^0 \bigg[\|\delta \mu_T\|_{-\mathcal{R}} + \int_0^T \Big(\|\delta \mu_r\|_{-\mathcal{R}} + |\delta Z_r^0| \|\bar{v}_r^0\|_{\mathcal{L}^1} - (d/2+1) \Big) dr \bigg] \leq C_T (|x_0 - x'_0| + \|\delta \mu_0\|_{-\mathcal{R}}) + C \overline{\mathbb{E}}^0 \bigg[\int_0^T \big(|\nabla_x \delta u_r|^2, \mu_r \big) dr \bigg]^{1/2} + C \overline{\mathbb{E}}^0 \bigg[\int_0^T |\delta Z_r^0|^2 dr \bigg]^{1/2},
$$

Squaring it, and applying assumption i in the statement together with (4.26) , we obtain

$$
\begin{split} \|\delta u_{0}\|_{\mathcal{H}}^{2} &\leq C_{T} \left(|x_{0}-x_{0}'|^{2}+\|\delta\mu_{0}\|_{-\mathcal{H}}^{2} \right) + C \overline{\mathbb{E}}^{0} \bigg[\int_{0}^{T} \left(\left| \nabla_{x}\delta u_{r} \right|^{2}, \mu_{r} \right) \mathrm{d}r \bigg] + C \overline{\mathbb{E}}^{0} \bigg[\int_{0}^{T} |\delta Z_{r}^{0}|^{2} \mathrm{d}r \bigg] \\ &\leq C_{\varepsilon_{1},\sigma_{0},T} \left(|x_{0}-x_{0}'|^{2}+\|\delta\mu_{0}\|_{-\mathcal{H}}^{2}+\left|(\delta u_{0},\delta\mu_{0})\right| \right) - \frac{1}{C_{\varepsilon_{1},\sigma_{0},T}} |\delta Y_{0}^{0}|^{2}, \end{split}
$$

where we assumed without any loss of generality that $C_{\epsilon_1,\sigma_0,T} \geq 1$. Rearranging it (and handling the duality bracket with a Young's inequality), we complete the proof. a Young's inequality), we complete the proof.

4.6. Application to existence and uniqueness

In this subsection, we NO longer take for granted the assumptions of Propositions [4.1](#page-31-0) and [4.4.](#page-34-1)

The main objective is to prove the following statement, which subsumes Theorem [2.14:](#page-10-0)

Theorem 4.11. *Under Assumption* (B)*, there exist a constant* $C \geq 1$ *, only depending on the parameters in* (B) *except* (σ_0, T) , and a constant $C_{\epsilon_1,\sigma_0,T}$, only depending on ϵ_1 and the parameters in (**B**) (including (σ_0, T)) and non-decreasing with *T*, such that, with the two notations $\kappa_{H^0} := \|\nabla^2_{pp} H^0\|_{\infty}/2$ and $\lambda_H := \sup\{\theta : \nabla^2_{pp} H \ge \theta I_d\}$, for *A* and *c* as in *Lemma [4.9,](#page-37-2) and under the following two conditions:*

i.
$$
\varepsilon_1 + \varepsilon_2 < \frac{\lambda_H}{C}
$$
,
\n*ii.* $\frac{c\varepsilon_2 \sigma_0^2}{2A} > \frac{C\kappa_{H^0}}{\varepsilon_1 \sigma_0^2}$,

the following holds true:

a. for all $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, the system (2.11) – (2.12) has a unique solution;

- b. there exist two continuous mappings \mathcal{U}^0 : $[0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ and \mathcal{U} : $[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ such that, f or any $r \in [1, \lfloor \mathcal{B} \rfloor - (d/2 + 1)] \setminus \mathbb{N}$, $\mathcal{U}^0(t, \cdot, \cdot)$ and $\mathcal{U}(t, \cdot, \cdot, \cdot)$ *belong to* $\mathscr{C}^0(C_{\varepsilon_1, \sigma_0, T}, \mathcal{D})$ and $\mathscr{C}(C_{\varepsilon_1, \sigma_0, T}, \mathcal{D}, \mathcal{D})$ for all $t \in [0, T]$ *and the representation formula* [\(4.6\)](#page-31-1) *holds true*;
- *c.* for a certain $\alpha > 0$, the continuity condition [\(4.11\)](#page-34-2) is satisfied and $\mathcal{U}^0(t,\cdot,\cdot)$ and $\mathcal{U}(t,\cdot,\cdot,\cdot)$ belong to $\mathscr{D}^0(C_{\varepsilon_1,\sigma_0,T},1)$ *and* $\mathscr{D}(C_{\varepsilon_1,\sigma_0,T},1,3+\alpha)$ *respectively.*

Proof. The proof relies on a standard induction argument, used first in [\[26\]](#page-86-0).

First Step. We prove in Section [5,](#page-50-0) see Proposition [5.19](#page-80-0) with $R_0 := 2 \max(C_{\varepsilon_1, \sigma_0, T}, \kappa)$ and $s := \lfloor \Delta \rfloor - (d/2 + 1) > 3$ (since $s > d/2 + 5$) that, under Assumptions (A1), (A2) and (A4) (but not (A3)), there exists a constant $\delta > 0$ (corresponding to $\mathfrak C$ in Proposition [5.19\)](#page-80-0), depending on the parameters in Assumption (A) but only depending on g^0 and g through any pair of (strictly) positive reals $(\tilde{L}, \tilde{\alpha})$ such that $g^0 \in \mathscr{C}^0(\tilde{L}, 3 + \tilde{\alpha})$ and $g \in \mathscr{C}(\tilde{L}, 3 + \tilde{\alpha}, 3 + \tilde{\alpha})$ (the roles of \tilde{L} and $\tilde{\alpha}$ in the statement of Proposition [5.19](#page-80-0) are respectively played by \mathfrak{L} and $s - |s|$) and satisfying the following variant of *a*, *b* and *c*:

- *a*. [see part *iii* in Proposition [5.19](#page-80-0) together with the definition of **r** in Theorem [5.1\]](#page-50-1) *for all* $(t_0, x_0, \mu) \in [T \delta, T] \times \mathbb{R}^d \times$ $\mathcal{P}(\mathbb{T}^d)$, the system [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0) has a unique solution when: 1. the BMO borm of $(\int_0^t Z_s^0 \cdot dB_s^0)_{0 \leq t \leq T}$ is required to be bounded by R_0 ; 2. u is regarded as a continuous process with values in $C^{3+\beta}(\mathbb{T}^d)$ *, for a certain* $\beta \in (0,\tilde{\alpha})$ *, and* the gradient $(\nabla_x u_t)_{0\leq t\leq T}$ is also required to be bounded by R_0 ; 3. the stochastic integral $(\int_{t_0}^t v_s^0(\cdot)\cdot dB_s^0)_{t_0\leq t\leq T}$ is *just regarded as a martingale* $(m_t)_{t_0 \le t \le T}$ *with values in* $C^{1+\beta}(\mathbb{T}^d)$ *, for the same value of* β *;*
- *b'.* [see part *i* in Proposition [5.19](#page-80-0) together with the representation formulas in [\(5.26\)](#page-59-0) and Proposition [5.13\]](#page-71-0) *there exist two positive reals* L *and* α *(corresponding to* C *and* (**s** + ⌊**s**⌋)/2 *in Proposition [5.19\)](#page-80-0) and two continuous mappings* $\mathcal{U}^0:[T-\delta,T]\times\mathbb{R}^d\times\mathcal{P}(\mathbb{T}^d)\to\mathbb{R}$ and $\mathcal{U}:[T-\delta,T]\times\mathbb{R}^d\times\mathbb{T}^d\times\mathcal{P}(\mathbb{T}^d)\to\mathbb{R}$ such that $\mathcal{U}^0(t,\cdot,\cdot)$ and $\mathcal{U}(t,\cdot,\cdot,\cdot)$ *belong to* $\mathscr{D}^0(L,1)$ and $\mathscr{D}(L,1,3+\alpha)$ for all $t \in [T-\delta,T]$, and the representation formula [\(4.6\)](#page-31-1) (restricted to the *interval* $[T - \delta, T]$ *and without the representation of* v^0 *which has not been defined yet) holds true;*
- *c'.* [see part *ii* in Proposition [5.19\]](#page-80-0) *the continuity condition* [\(4.11\)](#page-34-2) *is satisfied.*

We now fix $(t_0, x_0, \mu) \in [T - \delta, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$. By Lemma [3.1](#page-19-0) (which one can invoke with μ given by item *a*' above), the pair (u, m) takes values in $C^{\lambda}(\mathbb{T}^d) \times C^{\lambda-2}(\mathbb{T}^d)$. We can also apply Lemma [3.2:](#page-20-0) it permits to represent the martingale as a stochastic integral (with v^0 as integrand). The combination of the latter two lemmas says that the solution given above fits the requirements of Definition [2.10.](#page-9-1) By Proposition [3.4](#page-22-0) and Lemma [3.8,](#page-28-0) we get bounds for $|Y_{t_0}^0|$ Solution given above fits the requirements of Definition 2.10. By Proposition 3.4 and Lemma 3.8, we get bounds for $[Y_{t_0}^{\perp}]$
and $||u_{t_0}||_{\rho}$ from which we deduce that, for a possibly new value of $C_{\epsilon_1,\sigma_0,T}$, $|\$ bounded by $C_{\epsilon_1,\sigma_0,T}$. By Lemma [5.14](#page-71-1) (applied to $(\mathbf{U}_0^0,\mathbf{V}^0)=(u_t(x),v_t(x))_{t_0\leq t\leq T}$ and $\mathcal{V}^0=\mathcal{U}(\cdot,\cdot,x,\cdot)$ for any fixed $x \in \mathbb{T}^d$), we get the representation formula [\(4.6\)](#page-31-1) for $v_t^0(x)$ (observing that v_t^0 and $\nabla_{x_0} \mathcal{U}$ are continuous in x, we get the formula [\(4.6\)](#page-31-1) for almost every $(t, \omega^0) \in [0, T] \times \Omega^0$, for all $x \in T^d$). By Lemma [3.8](#page-28-0) again, we also deduce that, whatever the solution $(\mathbf{X}^{0,\prime}, \mathbf{Y}^{0,\prime}, \mathbf{Z}^{0,\prime}, \boldsymbol{\mu}', \boldsymbol{u}', \boldsymbol{m}')$ to [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0), the BMO norm of $(\int_0^t Z_s^{0,\prime} \cdot dB_s^0)_{0 \le t \le T}$ is bounded by R_0 (again, this may require to modify $C_{\varepsilon_1,\sigma_0,T}$), which proves from *a*' that $(X^0, Y^0, Z^0, \mu, u, m)$ is the unique solution to [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0). Furthermore, the assumptions of Propositions [4.1](#page-31-0) and [4.4](#page-34-1) are satisfied over the interval $[T - \delta, T]$. In turn, we can invoke Proposition [4.10](#page-40-2) to deduce that [\(4.23\)](#page-40-3) (at least, the analogue of it, but on the interval $[T - \delta, T]$) holds true. We explain below how to use this bound.

We recall that the initial conditions to [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1) (which are now initialized from t_0) are $\delta X_0 = x'_0 - x_0$ and $\delta \mu_0 =$ $\mu' - \mu$. And then, by the representation formula [\(4.12\)](#page-34-3), we have, for $\pi \in [3, \lfloor \Delta \rfloor - (d/2 + 1)] \setminus \mathbb{N}$ (which is possible because $\Delta > d/2 + 5$ and then $|\Delta| > d/2 + 4 = (d/2 + 1) + 3$,

$$
\left| \nabla_{x_0} \mathcal{U}^0(t_0, x_0, \mu) \cdot (x'_0 - x_0) + \left(\delta_\mu \mathcal{U}^0(t_0, x_0, \mu), \mu' - \mu \right) \right| \leq C_{\varepsilon_1, \sigma_0, T} \left(|x'_0 - x_0| + ||\mu' - \mu||_{-\varepsilon} \right),
$$
\n
$$
\left\| \nabla_{x_0} \mathcal{U}(t_0, x_0, \cdot, \mu) \cdot (x'_0 - x_0) + (\delta_\mu \mathcal{U}(t_0, x_0, \cdot, \mu), \mu' - \mu) \right\|_{\varepsilon} \leq C_{\varepsilon_1, \sigma_0, T} \left(|x'_0 - x_0| + ||\mu' - \mu||_{-\varepsilon} \right).
$$
\n(4.27)

Next, we focus on $\delta_\mu U$ (which is the most difficult term). Choosing $x'_0 = x_0$, we rewrite the last inequality as

$$
\left\| \int_{\mathbb{T}^d} \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y) \mathrm{d} \big(\mu' - \mu \big)(y) \right\|_{\mathcal{D}} \leq C_{\varepsilon_1, \sigma_0, T} \| \mu' - \mu \|_{-\mathcal{D}}.
$$
\n(4.28)

Assume for a while that for some constant $\gamma > 0$, μ is bounded from below by γ Leb_T^d</sub> (which we denote $\mu \geq \gamma$ Leb_{T^{d}}).</sub> For a given smooth even density ρ and for $l \in \{1, \dots, \lfloor n \rfloor\}$, consider the *l*th partial derivative $\partial^l \rho$ of ρ along *l* (arbitrary) directions of \mathbb{R}^d (with possible repetitions). Then, for $\epsilon \in \mathbb{R}$ small enough and $y_0 \in \mathbb{T}^d$ fixed, $\mu + \epsilon \partial^l \rho(y_0 - \cdot) \cdot \text{Leb}_{\mathbb{T}^d}$ is a probability measure and the above yields:

$$
\left\| \nabla_{y_0}^l \left(\int_{\mathbb{R}^d} \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y) \rho(y_0 - y) \mathrm{d}y \right) \right\|_{\mathcal{D}} \leq C_{\varepsilon_1, \sigma_0, T} \|\nabla^l \rho \cdot \mathbf{Leb}_{\mathbb{T}^d} \|_{-\mathcal{D}} \leq C_{\varepsilon_1, \sigma_0, T}.\tag{4.29}
$$

The case $l = 0$ can be easily included in the left-hand side by returning back to [\(4.28\)](#page-42-0), choosing therein $\mu' = \epsilon \delta_{y_0} + (1 - \epsilon \delta_{y_0})$ ϵ) μ and then invoking [\(2.1\)](#page-4-0).

Denoting the integral in the right-hand side of [\(4.29\)](#page-42-1) by $[\delta_\mu U(t_0, x_0, \mu, \cdot, \cdot) * \rho](y_0)$ (with the convolution acting implicitly on the last argument), we now address the Hölder regularity of $\nabla_y^{[r]}[\delta_\mu\mathcal{U}(t_0, x_0, \mu, \cdot, \cdot) * \rho]$ in the variable y by a similar argument. For a vector $z \in \mathbb{R}^d$, we apply [\(4.28\)](#page-42-0) to μ (still satisfying $\mu \ge \gamma \text{Leb}_{\mathbb{T}^d}$) and $\mu' = \mu + \epsilon [\partial^l \rho(\cdot - (y_0 + \cdot))$ (z)) – $\partial^l \rho(\cdot - y_0)$] (for ϵ small enough and with $l = \lfloor r \rfloor$). We obtain

$$
\left\| \nabla_{y_0}^{\lfloor r_1 \rfloor} \left(\int_{\mathbb{R}^d} \left[\delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y_0 + z - y) - \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y_0 - y) \right] \rho(y) dy \right) \right\|_{r_1}
$$
\n
$$
\leq C_{\varepsilon_1, \sigma_0, T} \left\| \nabla^{\lfloor r_1 \rfloor} \left[\rho(\cdot + z) - \rho \right] \cdot \text{Leb}_{\mathbb{T}^d} \right\|_{-r_1} \leq C_{\varepsilon_1, \sigma_0, T} |z|^{r_1 - \lfloor r_1 \rfloor} . \tag{4.30}
$$

The two inequalities [\(4.29\)](#page-42-1) and [\(4.30\)](#page-43-0) are true for any fixed (smooth) density ρ . Considering a sequence of mollifiers $(\rho_n)_{n\geq 1}$, this shows that, for any $l_x, l_y \in \{0, \dots, \lfloor r \rfloor\}$, the functions $(\nabla_x^{l_x} \nabla_y^{l_y} [\delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, \cdot) * \rho_n])_{n\geq 1}$ are relatively compact for the topology of uniform convergence on $\mathbb{T}^d \times \mathbb{T}^d$. Since $\mathcal{U}(t, \cdot, \cdot, \cdot)$ belongs to $\mathscr{D}(L, 1, 3 + \alpha)$ for all $t \in$ $[T - \delta, T]$, we already know that $\delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, \cdot) * \rho_n$ converges to $\delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, \cdot)$ for the topology of uniform convergence on $\mathbb{T}^d \times \mathbb{T}^d$. We deduce that, for any $l_x, l_y \in \{0, \dots, \lfloor r \rfloor\}$, the derivatives $\nabla_x^{l_x} \nabla_y^{l_y} \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, \cdot)$ exist and satisfy

$$
\sup_{l=0,\cdots,[\n\pi]} \left\| \nabla_{y_0}^l \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y) \right\|_{\mathcal{R}} \leq C_{\varepsilon_1, \sigma_0, T},
$$
\n
$$
\left\| \nabla_{y_0}^{|\mathcal{R}|} \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y_0 + z) - \nabla_{y_0}^{|\mathcal{R}|} \delta_\mu \mathcal{U}(t_0, x_0, \mu, \cdot, y_0) \right\|_{\mathcal{R}} \leq C_{\varepsilon_1, \sigma_0, T} |z|^{\mathcal{R}| |\mathcal{R}|}.
$$
\n(4.31)

The above holds true under the condition $\mu \ge \gamma \text{Leb}_{\mathbb{T}^d}$ for some $\gamma > 0$. Also, thanks to the continuity of $\delta_u \mathcal{U}$ in μ (given again by the fact that $U(t, \cdot, \cdot, \cdot)$ belongs to $\mathscr{D}(L, 1, 3 + \alpha)$ for all $t \in [T - \delta, T]$), we can prove that [\(4.31\)](#page-43-1) is true for any $\mu \in \mathcal{P}(\mathbb{T}^d)$ by a new approximation argument. Indeed, for any $\mu \in \mathcal{P}(\mathbb{T}^d)$ and $\epsilon \in (0,1)$, we can consider $(1 - \epsilon)\mu + \mu \text{Leb}_{\mathbb{T}^d}$ as new probability measure: [\(4.31\)](#page-43-1) is true for all $\epsilon \in (0, 1)$, and we can argue by a new compactness argument. Together with [\(4.27\)](#page-42-2), this shows that, for any $t_0 \in [T - \delta, T]$, the function $\mathcal{U}(t_0, \cdot, \cdot, \cdot)$ satisfies the condition $\mathscr{C}(C_{\varepsilon_1,\sigma_0,T},\pi,\pi)$ By the same argument, $\mathcal{U}^0(t_0,\cdot,\cdot)$ satisfies the condition $\mathscr{C}^0(C_{\varepsilon_1,\sigma_0,T},\pi)$.

Second Step. The next step is to iterate, considering now the problem (2.11) – (2.12) with $\mathcal{U}^0(T-\delta,\cdot,\cdot)$ and $\mathcal{U}(T-\delta,\cdot,\cdot,\cdot)$ as new terminal conditions at time $T - \delta$ (in place of g^0 and g at time T). By the first step (which does not require the terminal condition g to satisfy (A3)) and thanks to the lower bound $r > 3$ (without any loss of generality, we can assume $r > 3 + \tilde{\alpha}$ for the same $\tilde{\alpha}$ as the one used in the first step), we can find a new constant $\delta' > 0$ only depending on the parameters in (B) and on the constant $C_{\varepsilon_1,\sigma_0,T}$ (which also depends on the parameters in (B)) such that the items *a*', *b*' and *c*' above are satisfied but on $[T - (\delta + \delta'), T - \delta]$ in place of $[T - \delta, T]$. This makes it possible to extend the fields U^0 and U to the interval $[T - (\delta + \delta'), T]$. By combining the regularity properties given by item *b*' on the intervals $[T - (\delta + \delta'), T - \delta]$ and $[T - \delta, T]$ respectively, we deduce that *b*' holds true on the entire $[T - (\delta + \delta'), T]$. Similarly, we can easily check that *c*' is satisfied on the entire $[T - (\delta + \delta'), T]$.

Now, we observe that any solution given by *a*' on the interval $[T - (\delta + \delta'), T - \delta]$ can be extended to the interval $[T-\delta,T]$ by solving the problem addressed in the first step but restarting from the random initial condition $(X^0_{T-\delta}, \mu_{T-\delta})$. Solvability of the problem (2.11) – (2.12) with a random initial condition is not an issue. The analysis of forward-backward systems with random initial conditions has been well documented in other examples: see for instance [\[26\]](#page-86-0) for forwardbackward stochastic differential equations in finite dimension and [\[14,](#page-86-1) Subsection 5.1] for a similar analysis in the mean field case. Here, one can use the collection of regular conditional probabilities $(\mathbb{P}_{T-\delta}^0(\omega^0,\cdot))_{\omega^0\in\Omega^0}$ of \mathbb{P}^0 given the σ field generated by $(B_s^0)_{T-(\delta+\delta')\leq s\leq T-\delta}$ in order to reduce the system (initialized from a random initial condition) to a problem with a deterministic initial condition. Namely, for an initial condition $(X_{T-\delta}^0, \mu_{T-\delta})$ as above, one can solve the pair of two forward equations:

$$
dX_t^0 = -\nabla_p H^0(X_t^0, \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t^0)) dt + \sigma_0 dB_t^0,
$$

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x \mathcal{U}(t, X_t^0, \cdot, \mu_t^0)) \mu_t \right) = 0, \quad t \in [T - \delta, T].
$$

Since $\mathcal{U}^0 \in \mathcal{D}^0(L, 3 + \alpha)$ and $\mathcal{U} \in \mathcal{D}(L, 1, 3 + \alpha)$, existence and uniqueness are standard (it suffices to solve a conditional McKean-Vlasov equation, see for instance [\[20](#page-86-2), Chapter 2]). Then, defining $(Y_t^0, Z_t^0)_{T-\delta \le t \le T}$ and $(u_t, v_t^0)_{T-\delta \le t \le T}$ as suggested by (4.6) , namely letting

$$
Y_t^0 := \mathcal{U}^0(t, X_t^0, \mu_t), \ Z_t^0 := \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t), \quad t \in [T - \delta, T],
$$

$$
u_t(x) := \mathcal{U}(t, X_t^0, x, \mu_t), \ v_t^0(x) := \nabla_{x_0} \mathcal{U}(t, X_t^0, x, \mu_t), \quad (t, x) \in [T - \delta, T] \times \mathbb{T}^d
$$

we know from Proposition [4.1](#page-31-0) that, ω^0 - \mathbb{P}^0 -almost surely, under the disintegrated probability $\mathbb{P}^0_{T-\delta}(\omega^0, \cdot)$, the two triplets $(X_t^0, Y_t^0, Z_t^0)_{T-\delta \le t \le T}$ and $(\mu_t, u_t, v_t^0)_{T-\delta \le t \le T}$ that have been hence defined solve the two forward-backward equations in [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0) (with the realization $(X_{T-\delta}^{0}(\omega^{0}), \mu_{T-\delta}(\omega^{0}))$ as deterministic initial condition). And then, by construction of the regular conditional probability, they solve the systems $(2.11)-(2.12)$ $(2.11)-(2.12)$ $(2.11)-(2.12)$ under the original probability \mathbb{P}^0 , with $(X_{T-\delta}^0, \mu_{T-\delta})$ as (now random) initial condition.

By combining the solution originally given by *a*' on $[T - (\delta + \delta'), T - \delta]$ with its extension to $[T - \delta, T]$, we get a solution to (2.11) – (2.12) in the sense of *a*', starting from (t_0, x_0, μ) with $t_0 \in [T - (\delta + \delta'), T]$. This solution is necessarily unique. Indeed, if $(\tilde{X}_t^0, \tilde{Y}_t^0, \tilde{Z}_t^0)_{T-(\delta+\delta')\leq t\leq T}$ and $(\tilde{\mu}_t, \tilde{u}_t, \tilde{v}_t^0)_{T-(\delta+\delta')\leq t\leq T}$ form another solution with the same initial condition, then it satisfies the system [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0) on $[T - \delta, T]$ with $(\tilde{X}_{T-\delta}^0, \tilde{\mu}_{T-\delta})$ as (random) initial condition. In particular, for \mathbb{P}^0 -almost every ω^0 , it satisfies the system (2.1 with the realization $(\tilde{X}_{T-\delta}^0(\omega^0), \tilde{\mu}_{T-\delta}(\omega^0))$ as deterministic initial condition. And, then, by the uniqueness property on $[T - \delta, T]$, this new solution can be represented as in [\(4.6\)](#page-31-1) on the interval $[T - \delta, T]$. In particular, $\tilde{Y}_{T-\delta}^0 = \mathcal{U}^0(T - \delta, T)$ δ , $\widetilde{X}_{T-\delta}^0$, $\widetilde{\mu}_{T-\delta}$ and $\widetilde{u}_{T-\delta} = \mathcal{U}(T-\delta, \widetilde{X}_{T-\delta}^0; \cdot, \widetilde{\mu}_{T-\delta})$. This says that the triplets $(\widetilde{X}_t^0, \widetilde{Y}_t^0, \widetilde{Z}_t^0)_{T-(\delta+\delta')} \leq t \leq T-\delta$ and $(\widetilde{X}_t^0, \widetilde{X}_t^0, \widetilde{Z}_t^0)_{T-(\delta+\$ $(\widetilde{\mu}_t, \widetilde{u}_t, \widetilde{v}_t^0)_{T-(\delta+\delta') \leq t \leq T-\delta}$ satisfy the same equations as $(X_t^0, Y_t^0, Z_t^0)_{T-(\delta+\delta') \leq t \leq T-\delta}$ and $(\mu_t, u_t, v_t^0)_{T-(\delta+\delta') \leq t \leq T-\delta}$ on $[T - (\delta + \delta'), T - \overline{\delta}]$. By (short time) uniqueness on $[T - (\delta + \delta'), T - \delta]$, the latter two processes coincide. Then, they restart from the same states at time $T - \delta$ and eventually coincide on the entire $[T - (\delta + \delta'), T]$.

In the end, this proves that the three items *a*, *b* and *c* listed in the statement now hold true but on the wider interval $[T-(\delta+\delta'),T].$

Third Step. The proof now follows the scheme introduced in [\[26](#page-86-0)]. We can restart from the first step, but replacing $[T \delta$, T] by $[T - (\delta + \delta'), T]$. We get that, for any $t_0 \in [T - (\delta + \delta'), T]$, $\mathcal{U}^0(t_0, \cdot, \cdot)$ and $\mathcal{U}(t_0, \cdot, \cdot, \cdot)$ satisfy the conditions $\mathscr{C}^0(C_{\varepsilon_1,\sigma_0,T},\hbar)$ and $\mathscr{C}(C_{\varepsilon_1,\sigma_0,T},\hbar,\hbar)$ respectively. And then, we can repeat the second step and extend the maps \mathcal{U}^0 and U to $[T - (\delta + 2\delta'), T - (\delta + \delta')]$ for the same δ' as in the second step. By iterating the argument, we can cover the entire interval to $[0, T]$. Details are left to the reader. \Box

We are now in position to prove Theorem [2.16:](#page-16-0)

Theorem 4.12. *In addition to Assumption* (B)*, assume that*

- i. the coefficient $(t, x_0, \mu) \mapsto f_t^0(x_0, \mu)$ is Hölder continuous in time, uniformly in (x_0, μ) ; the coefficient $(t, x_0, x, \mu) \mapsto$ $f_t(x_0, x, \mu)$ *is Hölder continuous in time, uniformly in* (x_0, x, μ) *;*
- ii. $x_0 \mapsto g^0(x_0,\mu)$ has Hölder continuous second-order derivatives, uniformly in μ ; $x_0 \mapsto g(x_0,x,\mu)$ has Hölder con*tinuous second-order derivatives, uniformly in* (x, μ) *.*

Then, under the same conditions (i) and (ii) as in the statement of Theorem [4.11,](#page-41-2) the functions U ⁰ *and* ^U *are the unique* solutions to the master equation (2.27) – (2.28) in the class of pairs $(V⁰, V)$ such that

- $1.$ $(t, x_0, \mu) \mapsto (\partial_t V^0(t, x_0, \mu), \nabla_{x_0} V^0(t, x_0, \mu), \nabla_{x_0}^2 V^0(t, x_0, \mu))$ is continuous on $[0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$; $(t, x_0, \mu, y) \mapsto$ $(\partial_{\mu}V^{0}(t,x_{0},\mu,y),\nabla_{y}\partial_{\mu}V^{0}(t,x_{0},\mu,y))$ is continuous on $[0,T]\times\mathbb{R}^{d}\times\mathcal{P}(\mathbb{T}^{d})\times\mathbb{T}^{d}$;
- $2. (t, x_0, x, \mu) \mapsto (\partial_t V(t, x_0, x, \mu), \nabla_{x_0} V(t, x_0, x, \mu), \nabla_{x_0}^2 V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu), \nabla_x^2 V(t, x_0, x, \mu))$ is con*tinuous on* $[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$; $(t, x_0, x, \mu, y) \mapsto (\partial_\mu V(t, x_0, x, \mu, y), \nabla_y \partial_\mu V(t, x_0, x, \mu, y))$ *is continuous* $\mathcal{O}[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$
- 3. $(t, x_0, \mu) \mapsto \nabla_{x_0} V^0(t, x_0, \mu)$ and $(t, x_0, x, \mu) \mapsto (\nabla_{x_0} V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu))$ are Lipschitz continuous *with respect to* (x_0, μ) *and* (x_0, x, μ) *respectively (using the distance* \mathbb{W}_1 *to handle the argument* μ *), uniformly in* t*;*
- $4.$ $(t, x_0, \mu) \mapsto (V^0(t, x_0, \mu), \nabla_{x_0} V^0(t, x_0, \mu))$ and $(t, x_0, \mu) \mapsto (||V(t, x_0, \cdot, \mu)||_{\tau}$, $||\nabla_{x_0} V(t, x_0, \cdot, \mu)||_{\tau}$ are glob*ally bounded, for any* $r \in [1, |\lambda| - (d/2 + 1)] \setminus \mathbb{N}$ *.*

In particular, U ⁰ *and* ^U *have first-order derivative in* ^t *and second-order derivatives in* ^x0*, which are jointly continuous with respect to all their arguments.*

Proof. The proof relies on the iteration principle introduced in the proof of Theorem [4.11.](#page-41-2) We use in particular the same notation as therein.

,

On the interval $[T - \delta, T]$, the result follows from Corollary [5.17](#page-78-0) and Proposition [5.19](#page-80-0) (item iv). It says in particular that, at time $T - \delta$, the function $x_0 \mapsto \mathcal{U}^0(T - \delta, x_0, \mu)$ has Hölder continuous second-order derivatives, uniformly in μ , and the function $x_0 \mapsto \mathcal{U}(T - \delta, x_0, x, \mu)$ has Hölder continuous second-order derivatives, uniformly in (x, μ) . This makes it possible to reapply Corollary [5.17](#page-78-0) and Proposition [5.19](#page-80-0) (item iv) but on the interval $[T - (\delta + \delta'), T - \delta]$. By continuity properties of U^0 and U (and of their derivatives), we get a solution on $[T - (\delta + \delta'), T]$. Again, Corollary [5.17](#page-78-0) and Proposition [5.19](#page-80-0) (item iv) say that the functions $x_0 \mapsto \mathcal{U}^0(T - (\delta + \delta'), x_0, \mu)$ has Hölder continuous secondorder derivatives, uniformly in μ , and the function $x_0 \mapsto \mathcal{U}(T - (\delta + \delta'), x_0, x, \mu)$ has Hölder continuous second-order derivatives, uniformly in (x, μ) . By iterating, we get that \mathcal{U}^0 and \mathcal{U} solve the two master equations on the entire $[0, T]$.

We now turn to uniqueness. By Proposition [2.15,](#page-15-2) we know that any solution (V^0, V) to the master equation, as given in the statement, induces a solution to the forward-backward system (2.11) – (2.12) (for a given initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$). The bounds on (V^0, V) required in item (4) of the statement say that the resulting solution (u, v^0) to [\(2.12\)](#page-9-0) takes values in $C^n(\mathbb{T}^d) \times C^n(\mathbb{T}^d)$, for any $n \in [1, \lfloor \Delta \rfloor - (d/2 + 1)] \setminus \mathbb{N}$. Then, Lemmas [3.1](#page-19-0) and [3.2](#page-20-0) make it possible to gain extra regularity and guarantee that (ii) and (iii) in item 2 of Definition [2.10](#page-9-1) are satisfied. This shows that, for any fixed initial condition $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, (V^0, V) induces a solution to (2.11) – (2.12) in the sense of Definition [2.10.](#page-9-1) Uniqueness to [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0), as given by Theorem [4.11](#page-41-2) under the initial condition (x_0, μ) , together with the representation formula [\(4.6\)](#page-31-1) imply that (V^0, V) is necessarily equal to $(\mathcal{U}^0, \mathcal{U})$. \square

4.7. Proof of Proposition [4.4](#page-34-1)

Proof. Recall that $((X^0, Y^0, Z^0), (\mu, u, v^0))$ is the unique solution of [\(4.4\)](#page-31-2)–[\(4.5\)](#page-31-3) and has the representation [\(4.6\)](#page-31-1). For any $\varepsilon > 0$, let $((X^{0,\varepsilon}, Y^{0,\varepsilon}, Z^{0,\varepsilon}), (\mu^{\varepsilon}, u^{\varepsilon}, v^{0,\varepsilon}))$ be the unique solution of [\(4.4\)](#page-31-2)–[\(4.5\)](#page-31-3) corresponding to the initial data $(x_0 + \varepsilon (x'_0 - x_0), \mu + \varepsilon (\mu' - \mu))$. By Proposition [4.1,](#page-31-0) we have, \mathbb{P}^0 -almost surely,

$$
Y_t^{0,\varepsilon} = \mathcal{U}^0(t, X_t^{0,\varepsilon}, \mu_t^{\varepsilon}), Z_t^{0,\varepsilon} = \nabla_{x_0} \mathcal{U}^0(t, X_t^{0,\varepsilon}, \mu_t^{\varepsilon}), \quad t \in [0, T],
$$

\n
$$
u_t^{\varepsilon}(x) = \mathcal{U}(t, X_t^{0,\varepsilon}, x, \mu_t^{\varepsilon}), v_t^{0,\varepsilon}(x) = \nabla_{x_0} \mathcal{U}(t, X_t^{0,\varepsilon}, x, \mu_t^{\varepsilon}), \quad (t, x) \in [0, T] \times \mathbb{T}^d.
$$
\n
$$
(4.32)
$$

We let

$$
\begin{aligned} &\big((\triangle X^{0,\varepsilon},\triangle Y^{0,\varepsilon},\triangle Z^{0,\varepsilon}),(\triangle\mu^{\varepsilon},\triangle u^{\varepsilon},\triangle v^{0,\varepsilon})\big)\\ &:=\bigg(\Big(\frac{X^{0,\varepsilon}-X^0}{\varepsilon},\frac{Y^{0,\varepsilon}-Y^0}{\varepsilon},\frac{Z^{0,\varepsilon}-Z^0}{\varepsilon}\Big),\Big(\frac{\mu^{\varepsilon}-\mu}{\varepsilon},\frac{u^{\varepsilon}-u}{\varepsilon},\frac{v^{0,\varepsilon}-v^0}{\varepsilon}\Big)\bigg)\,. \end{aligned}
$$

First Step. Let us first solve the following forward system

$$
d\delta X_t^0 = -\delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] dt,
$$

\n
$$
\partial_t \delta \mu_t - \frac{1}{2} \Delta_x \delta \mu_t - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \delta \mu_t \right) - \text{div}_x \left(\delta \left[\nabla_p H(\cdot, \nabla_x \mathcal{U}(t, X_t^0, \cdot, \mu_t)) \right] \mu_t \right) = 0,
$$
\n(4.33)

for $t \in [0, T]$, with the two initial conditions $\delta X_0^0 = x_0' - x_0$ and $\delta \mu_0 = \mu' - \mu$. Above, the second equation is formally obtained from the forward equation in [\(4.9\)](#page-33-1) by replacing the term $\text{div}_x(\delta[\nabla_pH(\cdot,\nabla_xu_t(\cdot))] \mu_t)$ by $\text{div}_x(\delta[\nabla_p H(\cdot, \nabla_x \mathcal{U}(t, X_t^0, \cdot, \mu_t))] \mu_t)$. Since $\nabla_{x_0} w^0$ is x_0 -continuously differentiable with a bounded derivative (see [\(3.19\)](#page-28-1)), we easily deduce from the theory of ODEs that the first equation in [\(4.33\)](#page-45-0) admits a unique solution δX^0 in the sense of item 1 in Definition [4.2.](#page-33-1) It satisfies $\sup_{t\in[0,T]}|\delta X_t^0|\leq C_T|x_0'-x_0|$. Moreover, \mathbb{P}^0 -almost surely,

$$
\sup_{t \in [0,T]} |\Delta X_t^{0,\varepsilon}| \le C_T |x'_0 - x_0|, \quad \lim_{\varepsilon \to 0} \sup_{t \in [0,T]} \left| X_t^{0,\varepsilon} - X_t^0 \right| = 0 \quad \text{and} \quad \lim_{\varepsilon \to 0} \sup_{t \in [0,T]} \left| \Delta X_t^{0,\varepsilon} - \delta X_t^0 \right| = 0. \tag{4.34}
$$

Now let us turn to the second equation in [\(4.33\)](#page-45-0), which we prove to be solvable by linearizing the forward equation in [\(4.5\)](#page-31-3). To do so, we represent the solutions by means on an SDE, very much in the spirit of Lemma [2.4.](#page-7-0) Let $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ be a filtered probability space (different from $(\Omega^0, \mathcal{F}^0, \mathbb{F}^0, \mathbb{P}^0)$) equipped with an \mathbb{R}^d -valued \mathbb{F} -Brownian $(B_t)_{0 \le t \le T}$ and two \mathbb{R}^d -valued random variables ξ and ξ^ε satisfying $\mathbb{P} \circ \xi^{-1} = \mu_0 = \mu$, $\mathbb{P} \circ (\xi^\varepsilon)^{-1} = \mu_0^\varepsilon = \mu + \varepsilon (\mu' - \mu)$ and $\mathbb{E}[|\xi^{\varepsilon} - \xi|] = \mathbb{W}_1(\mu_0^{\varepsilon}, \mu_0)$. Let also $(X_t^{\varepsilon})_{0 \le t \le T}$, $(X_t)_{0 \le t \le T}$ be the solutions to

$$
X_t^{\varepsilon} = \xi^{\varepsilon} - \int_0^t \nabla_p H\left(X_s^{\varepsilon}, \nabla_x \mathcal{U}(s, X_s^{0,\varepsilon}, X_s^{\varepsilon}, \mu_s^{\varepsilon})\right) ds + B_t, \quad t \in [0, T],\tag{4.35}
$$

$$
X_t = \xi - \int_0^t \nabla_p H\left(X_s, \nabla_x \mathcal{U}(s, X_s^0, X_s, \mu_s)\right) \mathrm{d}s + B_t, \quad t \in [0, T].\tag{4.36}
$$

Then $\mathbb{P} \circ (X_t^{\varepsilon})^{-1} = \mu_t^{\varepsilon}$ and $\mathbb{P} \circ X_t^{-1} = \mu_t$ for any $t \in [0, T]$, \mathbb{P}^0 -almost surely. We have

$$
\mathbb{E}\left[|X_t^{\varepsilon}-X_t|\right]\leq \mathbb{E}\left[|\xi^{\varepsilon}-\xi|\right]+\mathbb{E}\bigg[\int_0^t \Big|\nabla_p H\left(X_s^{\varepsilon},\nabla_x \mathcal{U}(s,X_s^{0,\varepsilon},X_s^{\varepsilon},\mu_s^{\varepsilon})\right)-\nabla_p H\left(X_s,\nabla_x \mathcal{U}(s,X_s^{0},X_s,\mu_s)\right)\Big|ds\bigg].
$$

Since $U(t, \cdot, \cdot, \cdot)$ belongs to $\mathscr{C}(\kappa, 1, 2)$ for every $t \in [0, T]$ and for some $\kappa > 0$, the gradient $\nabla_x U$ is Lipschitz in the variable (x_0, x, μ) (with $\mathcal{P}(\mathbb{T}^d)$ being equipped with \mathbb{W}_1), uniformly in $t \in [0, T]$. By [\(4.34\)](#page-45-1) and Gronwall's lemma, we deduce that, **P** 0 -almost surely,

$$
\lim_{\varepsilon \to 0} \sup_{t \in [0,T]} \mathbb{W}_1(\mu_t^{\varepsilon}, \mu_t) = 0. \tag{4.37}
$$

In fact, we claim that the result also holds in total variation. When $\mu_0^{\epsilon} = \mu$ (i.e., $\mu' = \mu$), this follows from [\[42](#page-87-0), (3.1) and Proof of Theorem 2.4] (which rely on Pinsker's inequality and Girsanov theorem) together with [\(4.34\)](#page-45-1) and [\(4.37\)](#page-46-0). When $\mu' \neq \mu$, one must introduce the solution $\hat{\mathbf{X}}^{\varepsilon}$ to the SDE [\(4.35\)](#page-45-2) when initialized from ξ at time 0. Then, [\[42,](#page-87-0) (3.1) and Proof of Theorem 2.4] say that, \mathbb{P}^0 -almost surely,

$$
\lim_{\varepsilon \to 0} \sup_{t \in [0,T]} d_{\mathrm{TV}} \left(\mathbb{P} \circ (\hat{X}_t^{\varepsilon})^{-1}, \mu_t \right) = 0. \tag{4.38}
$$

Also, since $d_{\text{TV}}(\mu_0^{\varepsilon}, \mu_0) = d_{\text{TV}}((1 - \varepsilon)\mu + \varepsilon\mu', \mu) \leq 4\varepsilon$, we can now choose ξ^{ε} in [\(4.35\)](#page-45-2) such that $\mathbb{P}(\{\xi = \xi^{\varepsilon}\}) \geq 1 - 4\varepsilon$. Then, it is easy to see that $\mathbb{P}(\{\hat{X}_{t}^{\varepsilon}=X_{t}^{\varepsilon}\}) \geq 1-4\varepsilon$, for all $t \in [0,T]$. Since $\mathbb{P} \circ (\xi^{\varepsilon})^{-1}$ is still required to be equal to μ_{0}^{ε} , this does not change the flow $(\mathbb{P} \circ (X_t^{\varepsilon})^{-1})_{0 \le t \le T}$, which is still equal to $(\mu_t^{\varepsilon})_{0 \le t \le T}$. We deduce that, \mathbb{P}^0 -almost surely,

$$
\lim_{\varepsilon \to 0} \sup_{t \in [0,T]} d_{\text{TV}}(\mu_t^{\varepsilon}, \mu_t) = 0. \tag{4.39}
$$

We note that $\Delta \mu^{\epsilon} = (\Delta \mu^{\epsilon}_t)_{0 \leq t \leq T}$ satisfies (in a weak sense similar to [\(4.10\)](#page-33-2))

$$
\partial_t \left(\Delta \mu_t^{\varepsilon} \right) - \frac{1}{2} \Delta_x \left(\Delta \mu_t^{\varepsilon} \right) \n- \operatorname{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \Delta \mu_t^{\varepsilon} \right) - \operatorname{div}_x \left(\mu_t^{\varepsilon} \left[\Gamma_t^{1,\varepsilon}(\cdot) \Delta X_t^{0,\varepsilon} + \left(\Gamma_t^{2,\varepsilon}(\cdot, \cdot), \Delta \mu_t^{\varepsilon} \right) \right] \right) = 0,
$$
\n(4.40)

where

$$
\Gamma_t^{1,\varepsilon}(x) := \left(\int_0^1 \nabla_{pp}^2 H\Big(x, \theta \nabla_x \mathcal{U}(t, X_t^{0,\varepsilon}, x, \mu_t^{\varepsilon}) + (1 - \theta) \nabla_x \mathcal{U}(t, X_t^{0}, x, \mu_t) \Big) d\theta \right) \cdot \left(\int_0^1 \nabla_{xx_0}^2 \mathcal{U}\Big(t, \theta X_t^{0,\varepsilon} + (1 - \theta) X_t^{0}, x, \theta \mu_t^{\varepsilon} + (1 - \theta) \mu_t \Big) d\theta \right), \n\Gamma_t^{2,\varepsilon}(x, y) := \left(\int_0^1 \nabla_{pp}^2 H\Big(x, \theta \nabla_x \mathcal{U}(t, X_t^{0,\varepsilon}, x, \mu_t^{\varepsilon}) + (1 - \theta) \nabla_x \mathcal{U}(t, X_t^{0}, x, \mu_t) \Big) d\theta \right) \cdot \left(\int_0^1 \nabla_x \delta_\mu \mathcal{U}\Big(t, \theta X_t^{0,\varepsilon} + (1 - \theta) X_t^{0}, x, \theta \mu_t^{\varepsilon} + (1 - \theta) \mu_t, y \Big) d\theta \right),
$$

the dots right above being understood as matricial products and the duality bracket between $\triangle \mu_t^{\varepsilon}$ and $\Gamma_t^{2,\varepsilon}(\cdot,\cdot)$ in [\(4.40\)](#page-46-1) being acting on the variable y of the latter. Following the proof of Lemma [4.8](#page-36-0) and using the fact that $\mathcal{U}(t, \cdot, \cdot, \cdot) \in$ $\mathscr{C}(\kappa, 1, 2)$, we get, \mathbb{P}^0 -almost surely, for any $t \in [0, T]$,

$$
\|\Delta \mu_t^{\varepsilon}\|_{-1} \le C \bigg[\|\mu' - \mu\|_{-1} + \int_0^t \Big[\|\Gamma_s^{1,\varepsilon}(\cdot)\|_{L^\infty} \big|\Delta X_s^{0,\varepsilon}\big| + \Big\| \Big(\Gamma_s^{2,\varepsilon}(\cdot,\cdot),\Delta \mu_s^{\varepsilon}\Big) \Big\|_{L^\infty} \Big] ds \bigg] \le C \bigg[\|\mu' - \mu\|_{-1} + \int_0^t \Big(|\Delta X_s^{0,\varepsilon}| + \|\Delta \mu_s^{\varepsilon}\|_{-1} \Big) ds \bigg].
$$

By [\(4.34\)](#page-45-1) and Gronwall's lemma, we obtain, with probability 1 under \mathbb{P}^0 ,

$$
\sup_{t \in [0,T]} \|\Delta \mu_t^{\varepsilon}\|_{-1} \le C \bigg[|x'_0 - x_0| + \|\mu' - \mu\|_{-1} \bigg]. \tag{4.41}
$$

Now, for any sequence of reals $(\varepsilon_k)_{k\geq 1}$ converging to 0, we let

$$
\begin{aligned}\n\triangle^2 \mu_t^{k',k} &:= \triangle \mu_t^{\varepsilon_{k'}} - \triangle \mu_t^{\varepsilon_k}, \quad \triangle^2 X_t^{0,k',k} &:= \triangle X_t^{0,\varepsilon_{k'}} - \triangle X_t^{0,\varepsilon_k}, \\
\triangle \Gamma_t^{1,k',k} &:= \Gamma_t^{1,\varepsilon_{k'}} - \Gamma_t^{1,\varepsilon_k}, \quad \triangle \Gamma_t^{2,k',k} &:= \Gamma_t^{2,\varepsilon_{k'}} - \Gamma_t^{2,\varepsilon_k}, \quad t \in [0,T].\n\end{aligned}
$$

Then, we can verify that $\triangle^2 \mu^{k',k} = (\triangle^2 \mu^{k',k}_t)_{0 \le t \le T}$ satisfies

$$
\partial_t \left(\Delta^2 \mu_t^{k',k} \right) - \frac{1}{2} \Delta_x \left(\Delta^2 \mu_t^{k',k} \right) - \text{div}_x \left(\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \Delta^2 \mu_t^{k',k} \right) \n- \text{div}_x \left(\mu_t^{\varepsilon_k} \left[\Gamma_t^{1,\varepsilon_k}(\cdot) \Delta^2 X_t^{0,k',k} + \left(\Gamma_t^{2,\varepsilon_k}(\cdot,\cdot), \Delta^2 \mu_t^{k',k} \right) \right] \right) \n- \text{div}_x \left(\mu_t^{\varepsilon_k} \left[\Delta \Gamma_t^{1,k',k}(\cdot) \Delta X_t^{0,\varepsilon_{k'}} + \left(\Delta \Gamma_t^{2,k',k}(\cdot,\cdot), \Delta \mu_t^{\varepsilon_{k'}} \right) \right] \right) \n- \text{div}_x \left(\left(\mu_t^{\varepsilon_{k'}} - \mu_t^{\varepsilon_k} \right) \left[\Gamma_t^{1,\varepsilon_{k'}}(\cdot) \Delta X_t^{0,\varepsilon_{k'}} + \left(\Gamma_t^{2,\varepsilon_{k'}}(\cdot,\cdot), \Delta \mu_t^{\varepsilon_{k'}} \right) \right] \right) = 0, \quad t \in [0,T].
$$

Again, following the proof of Lemma [4.8,](#page-36-0) we can show that, \mathbb{P}^0 -almost surely, for all $t \in [0, T]$,

$$
\label{eq:4.13} \begin{split} \|\triangle^{2}\mu_{t}^{k',k}\|_{-1} & \leq C\int_{0}^{t}\biggl[\left\|\Gamma_{s}^{1,\varepsilon_{k}}(\cdot)\right\|_{L^{\infty}}\bigl|\triangle^{2}X_{t}^{0,k',k}\bigr| + \bigl\|\triangle\Gamma_{s}^{1,k',k}(\cdot)\bigr\|_{L^{\infty}}\bigl|\triangle X_{t}^{0,\varepsilon_{k'}}\bigr| \\ & + \Bigl\|\Bigl(\Gamma_{s}^{2,\varepsilon_{k}}(\cdot,\cdot),\triangle^{2}\mu_{s}^{k',k}\Bigr)\Bigr\|_{L^{\infty}} + \Bigl\|\Bigl(\triangle\Gamma_{s}^{2,k',k}(\cdot,\cdot),\triangle\mu_{s}^{\varepsilon_{k'}}\Bigr)\Bigr\|_{L^{\infty}} \\ & + d_{\text{TV}}\bigl(\mu_{s}^{\varepsilon_{k'}},\mu_{s}^{\varepsilon_{k}}\bigr)\Bigl(\bigl\|\Gamma_{s}^{1,\varepsilon_{k'}}(\cdot)\bigr\|_{L^{\infty}}\bigl|\triangle X_{s}^{0,\varepsilon_{k'}}\bigr| + \Bigl\|\Bigl(\Gamma_{s}^{2,\varepsilon_{k'}}(\cdot,\cdot),\triangle\mu_{s}^{\varepsilon_{k'}}\Bigr)\Bigr\|_{L^{\infty}}\Bigr] \mathrm{d}s. \end{split}
$$

Using the fact that $U(t, \cdot, \cdot)$ belongs to $\mathcal{D}(\kappa, 1, 2)$ for all $t \in [0, T]$, we deduce from [\(4.34\)](#page-45-1) and [\(4.37\)](#page-46-0) that, \mathbb{P}^0 -almost surely,

$$
\lim_{n\to\infty}\sup_{k,k'\geq n}\sup_{s\in[0,T]}\Bigl(\bigl\|\triangle\Gamma^{1,k',k}_s(\cdot)\bigr\|_{L^\infty}+\sup_{x\in\mathbb{T}^d}\bigl\|\triangle\Gamma^{2,k',k}_s(x,\cdot)\bigr\|_1\Bigr)=0.
$$

Moreover, by [\(4.34\)](#page-45-1) again, \mathbb{P}^0 -almost surely,

$$
\lim_{n \to \infty} \sup_{k,k' \ge n} \sup_{s \in [0,T]} \left| \triangle^2 X_s^{0,k',k} \right| = 0.
$$

By [\(4.39\)](#page-46-2) and [\(4.41\)](#page-46-3), we deduce that, \mathbb{P}^0 -almost surely, there exists a positive sequence $(\delta_n)_{n\geq 1}$, converging to 0, such that, for all $k, k' \geq n$ and for all $t \in [0, T]$,

$$
\|\triangle^{2}\mu_{t}^{k',k}\|_{-1} \leq \delta_{n} + C \int_{0}^{t} \|\triangle^{2}\mu_{s}^{k',k}\|_{-1} \mathrm{d}s.
$$

And then, by Gronwall's lemma, \mathbb{P}^0 -almost surely, $(\Delta \mu^{\varepsilon_k})_{k\geq 1}$ is a Cauchy sequence in $\mathcal{C}([0,T], \mathcal{C}^{-1}(\mathbb{T}^d))$. Therefore, there exists a continuous \mathbb{F}^0 -progressively measurable process $\delta \mu = (\delta \mu_t)_{0 \le t \le T}$ with values in $C^{-1}(\mathbb{T}^d)$ such that, \mathbb{P}^0 almost surely,

$$
\lim_{k \to \infty} \sup_{t \in [0,T]} ||\triangle \mu_t^k - \delta \mu_t||_{-1} = 0.
$$

Passing to the limit in [\(4.40\)](#page-46-1), we deduce that $\delta \mu$ satisfies (in the same sense as [\(4.10\)](#page-33-2))

$$
\partial_t (\delta \mu_t) - \frac{1}{2} \Delta_x (\delta \mu_t) - \text{div}_x (\nabla_p H(\cdot, \nabla_x u_t(\cdot)) \delta \mu_t)
$$

$$
- \text{div}_x \Big(\mu_t \Big[\nabla_{pp}^2 H(\cdot, \nabla_x u_t(\cdot)) \nabla_{xx_0}^2 \mathcal{U}(t, X_t^0, x, \mu_t) \delta X_t^0 + \Big(\nabla_{pp}^2 H(\cdot, \nabla_x u_t(\cdot)) \nabla_x \delta_\mu \mathcal{U}(t, \cdot, \cdot), \delta \mu_t \Big) \Big] \Big) \qquad (4.42)
$$

= 0, $t \in [0, T].$

Moreover, passing to the limit in [\(4.41\)](#page-46-3), we obtain

$$
\sup_{t \in [0,T]} \|\delta \mu_t\|_{-1} \le C \bigg[|x'_0 - x_0| + \|\mu' - \mu\|_{-1} \bigg]. \tag{4.43}
$$

This proves that $\delta \mu$ solves the second equation in [\(4.33\)](#page-45-0) in the sense of item 2 in Definition [4.2.](#page-33-1) *Second Step.* We now address the linearized backward equations

$$
d\delta Y_t^0 = -\delta \left[f_t^0(X_t^0, \mu_t) \right] dt + \left[-\left(Z_t^0 - \nabla_{x_0} w^0(t, X_t^0) \right) \cdot \delta \left[\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] - \delta \left[L^0(X_t^0, -\nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right] \right] dt + \left(\nabla_{x_0} H^0(X_t^0, Z_t^0) - \nabla_{x_0} H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \delta X_t^0 dt + \left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \delta Z_t^0 dt + \sigma_0 \delta Z_t^0 \cdot dB_t^0, \quad t \in [0, T], \delta Y_T^0 = \delta \left[g^0(X_T^0, \mu_T) \right],
$$
\n(4.44)

and

$$
d_t \delta u_t(x) = \left(-\frac{1}{2}\Delta_x \delta u_t(x) + \nabla_p H(x, \nabla_x u_t(x)) \cdot \nabla_x \delta u_t(x) - \delta \left[f_t(X_t^0, x, \mu_t)\right]\right) dt,
$$

+
$$
\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0))\right) \cdot \delta v_t^0(x) dt
$$

+
$$
\delta \left[\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0))\right)\right] \cdot v_t^0(x) dt
$$

+
$$
\sigma_0 \delta v_t^0(x) \cdot dB_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
\delta u_T(x) = \delta \left[g(X_T^0, x, \mu_T)\right], \quad x \in \mathbb{T}^d.
$$
 (4.45)

We first address [\(4.44\)](#page-48-0). Recalling that $Z_t^0 = \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t)$ is bounded, it follows from the standard BSDE theory that [\(4.44\)](#page-48-0) admits a solution $(\delta Y^0, \delta Z^0) \in \mathscr{S}^2(\mathbb{R}) \times \mathscr{H}^2(\mathbb{R}^d)$. Moreover, following [\[54\]](#page-87-1), we have

$$
\lim_{\varepsilon \to 0} \mathbb{E}^0 \bigg[\sup_{t \in [0,T]} |\Delta Y_t^{0,\varepsilon} - \delta Y_t^0|^2 + \int_0^T |\Delta Z_t^{0,\varepsilon} - \delta Z_t^0|^2 dt \bigg] = 0,\tag{4.46}
$$

which, together with the first step and the fact that $\mathcal{U}^0(t, \cdot, \cdot) \in \mathcal{D}^0(\kappa, 1)$ for all $t \in [0, T]$, implies

$$
\delta Y_t^0 = \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t) \cdot \delta X_t^0 + \left(\delta_\mu \mathcal{U}^0(t, X_t^0, \mu_t, \cdot), \delta \mu_t \right), \quad t \in [0, T].
$$

Because $\mathcal{U}^0(t,\cdot,\cdot)\in\mathscr{C}^0(\kappa,1)$, for all $t\in[0,T]$, and $\|\delta X^0\|$ and $\|\delta\mu\|_{-1}$ are bounded, we have

$$
\Big\|\sup_{0\leq t\leq T}|\delta Y^0_t|\Big\|_{L^\infty(\Omega^0,{\mathbb P}^0)}<\infty.
$$

Then we can follow the proof of Lemma [3.8](#page-28-0) to deduce that

$$
\sup_{\tau}\left\|\mathbb{E}^0\Big[\int_{[\tau,T]}|\delta Z_t^0|^2\mathrm{d} t |\mathcal{F}^0_{\tau}\Big]\right\|_{L^\infty(\Omega^0,\mathbb{P}^0)}<\infty,
$$

where the supremum is taken over all stopping times τ . This justifies that $(\delta Y^0, \delta Z^0)$ solves [\(4.44\)](#page-48-0) in the sense of item 1 in Definition [4.2.](#page-33-1)

The well-posedness of the equation [\(4.45\)](#page-48-1) is obtained via [\[30,](#page-86-3) Theorem 2.2], with solutions in the set of progressivelymeasurable processes $(\delta u, \delta v^0)$ with values in (the separable space) $\mathcal{H}^1(\mathbb{T}^d) \times [\mathcal{H}^1(\mathbb{T}^d)]^d$, such that δu has continuous trajectories and for which the following bound holds true:

$$
\mathbb{E}^{0}\left[\sup_{0\leq t\leq T}\|\delta u_{t}\|^{2}_{1,2}+\int^{T}_{0}\|\delta v^0_{t}\|^{2}_{1,2}\mathrm{d}t\right]<\infty.
$$

Note that

$$
d_t(\Delta u_t^{\varepsilon}) = \left[-\frac{1}{2}\Delta_x \left(\Delta u_t^{\varepsilon} \right) + \Upsilon_t^{\varepsilon}(x) \cdot \nabla_x \Delta u_t^{\varepsilon} - \nabla_x f_t^{\varepsilon}(x) \cdot \Delta X_t^{0,\varepsilon} - (\delta_\mu f_t^{\varepsilon}(x, \cdot), \Delta \mu_t^{\varepsilon}) \right] dt
$$

+
$$
\left(\nabla_p H^0(X_t^0, Z_t^0) - \nabla_p H^0(X_t^0, \nabla_{x_0} w^0(t, X_t^0)) \right) \cdot \Delta v_t^{0,\varepsilon} dt
$$

+
$$
\left[(j_t^{1,\varepsilon} - j_t^{3,\varepsilon}) \Delta X_t^{0,\varepsilon} + j_t^{2,\varepsilon} \Delta Z_t^{0,\varepsilon} \right] \cdot v_t^{0,\varepsilon}(x) dt + \sigma_0 \Delta v_t^{0,\varepsilon} \cdot dB_t^0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

$$
\Delta u_T^{\varepsilon}(x) = \nabla_x g^{\varepsilon}(x) \cdot \Delta X_T^{0,\varepsilon} + (\delta_\mu g^{\varepsilon}(x, \cdot), \Delta \mu_T^{\varepsilon}),
$$

where

$$
\begin{split} &\Upsilon^\varepsilon_t(x):=\int_0^1\nabla_p H\big(x,\theta\nabla_x u^\varepsilon_t(x)+(1-\theta)\nabla_x u_t(x)\big)\text{d}\theta,\\ &\nabla_x f^\varepsilon_t(x):=\int_0^1\nabla_x f_t\big(\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t,x,\theta\mu^\varepsilon_t+(1-\theta)\mu_t\big)\text{d}\theta,\quad\text{and similarly for }\nabla_x g^\varepsilon(x)\\ &\delta_\mu f^\varepsilon_t(x,y):=\int_0^1\delta_\mu f_t\big(\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t,x,\theta\mu^\varepsilon_t+(1-\theta)\mu_t,y\big)\text{d}\theta,\quad\text{and similarly for }\partial_\mu g^\varepsilon(x,y)\\ &j^{1,\varepsilon}_t:=\int_0^1\nabla_{x_0p}^2H^0\big(\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t,\theta Z^{0,\varepsilon}_t+(1-\theta)Z^0_t\big)\text{d}\theta,\\ &j^{2,\varepsilon}_t:=\int_0^1\nabla_{p}^2H^0\big(\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t,\theta Z^{0,\varepsilon}_t+(1-\theta)Z^0_t\big)\text{d}\theta,\\ &j^{3,\varepsilon}_t:=\int_0^1\big(\nabla_{x_0p}^2H^0+\nabla_{p p}^2H^0\nabla_{x_0x_0}^2w^0\big)\big(\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t,\nabla_{x_0}w^0(t,\theta X^{0,\varepsilon}_t+(1-\theta)X^0_t)\big)\text{d}\theta. \end{split}
$$

Using the first step together with the representation formulas [\(4.32\)](#page-45-3), there is no difficulty in proving that

$$
\lim_{\varepsilon \to 0} \mathbb{E}^{0} \int_{0}^{T} \left[\left\| \left(\Upsilon_{t}^{\varepsilon} - \Upsilon_{t}^{0} \right) \cdot \nabla_{x} \Delta u_{t}^{\varepsilon} \right\|_{0}^{2} \right] dt = 0,
$$
\n
$$
\lim_{\varepsilon \to 0} \mathbb{E}^{0} \int_{0}^{T} \left[\left\| \nabla_{x} f_{t}^{\varepsilon} \cdot \Delta X_{t}^{0,\varepsilon} - \nabla_{x} f_{t} (X_{t}^{0}, \cdot, \mu_{t}) \cdot \delta X_{t}^{0} \right\|_{0}^{2} \right] dt = 0,
$$
\n
$$
\lim_{\varepsilon \to 0} \mathbb{E}^{0} \int_{0}^{T} \left[\sup_{x \in \mathbb{T}^{d}} \left| \left(\delta_{\mu} f_{t}^{\varepsilon}, \Delta \mu_{t}^{\varepsilon} \right) - \left(\nabla_{\mu} f_{t} (X_{t}^{0}, x, \mu_{t}, \cdot), \delta \mu_{t} \right) \right|^{2} \right] dt = 0,
$$
\n
$$
\lim_{\varepsilon \to 0} \mathbb{E}^{0} \int_{0}^{T} \left\| \left[\left(j_{t}^{1,\varepsilon} - j_{t}^{3,\varepsilon} \right) \Delta X_{t}^{0,\varepsilon} + j_{t}^{2,\varepsilon} \Delta Z_{t}^{0,\varepsilon} \right] \cdot v_{t}^{0,\varepsilon} - \left[\left(j_{t}^{1,0} - j_{t}^{3,0} \right) \delta X_{t}^{0} + j_{t}^{2,0} \delta Z_{t}^{0} \right] \cdot v_{t}^{0} \right\|_{0}^{2} dt = 0.
$$

Proceeding in the same manner for the terminal condition, we can invoke [\[30](#page-86-3), Theorem 2.2] again to obtain

$$
\lim_{\varepsilon\to 0}\mathbb{E}^0\left[\sup_{0\leq t\leq T}\|\triangle u_t^\varepsilon-\delta u_t\|_{1,2}^2+\int_0^T\|\triangle v_t^{0,\varepsilon}-\delta v_t^0\|_{1,2}^2\mathrm{d} t\right]=0.
$$

Since $\mathcal{U}(t, \cdot, \cdot, \cdot) \in \mathcal{D}(\kappa, 1, 3 + \alpha)$ for all $t \in [0, T]$, we can derive that

$$
\delta u_t(x) = \nabla_{x_0} \mathcal{U}(t, X_t^0, x, \mu_t) \cdot \delta X_t^0 + \left(\delta_\mu \mathcal{U}(t, X_t^0, x, \mu_t, \cdot), \delta \mu_t \right),
$$

for any $t \in [0, T]$ with probability 1 under \mathbb{P}^0 , which implies that $\sup_{0 \le t \le T} ||\delta u_t||_{3+\alpha} \in L^{\infty}(\Omega^0, \mathbb{P}^0)$. Further applying [\(4.11\)](#page-34-2), we can verify that δu has continuous trajectories in $C^{3+\alpha}(\mathbb{T}^d)$. Inserting the above identity in [\(4.42\)](#page-47-0), we obtain $(4.9).$ $(4.9).$

Moreover, from the representation formula [\(4.32\)](#page-45-3) together with [\(4.34\)](#page-45-1) and [\(4.41\)](#page-46-3), and the fact that $\mathcal{U}(t,\cdot,\cdot,\cdot)$ belongs to $\mathscr{D}(\kappa, 1, 3 + \alpha)$, we deduce that

$$
\|\triangle v_t^{0,\varepsilon}\|_{3+\alpha} \le C\bigg[|x'_0 - x_0| + \|\mu' - \mu\|_{-1}\bigg],
$$

which proves by passing to the limit that δv_t^0 takes values in a compact subset of $C^{3+\epsilon}(\mathbb{T}^d)$, for any $\epsilon \in (0,\alpha)$. Therefore, δv^0 , regarded as a process taking values in $\mathcal{C}^{3+\epsilon}(\mathbb{T}^d)$, is progressively-measurable (in the *Bochner* sense).

In the end, we obtain $(\delta u, \delta v^0)$ solves the second equation of the backward system [\(4.9\)](#page-33-1) in the sense of item 2 in Definition [4.2.](#page-33-1) We finally get $((\delta \mathbf{X}^{0}, \delta \mathbf{Y}^{0}, \delta \mathbf{Z}^{0}), (\delta \mu, \delta \mathbf{u}, \delta \mathbf{v}^{0}))$ is a solution to the system [\(4.8\)](#page-33-0)–[\(4.9\)](#page-33-1) in the sense of Definition [4.2.](#page-33-1) \Box

5. Short time analysis

In this section we study the local well-posedness of the forward-backward system (2.11) – (2.12) (and then of the master equations $(2.27)-(2.28)$ $(2.27)-(2.28)$ $(2.27)-(2.28)$). One of the difficulties to do so is that the coefficients of $(2.11)-(2.12)$ $(2.11)-(2.12)$ are not Lipschitz continuous (due to the quadratic growth of the two Lagrangians L^0 and L in the control variable α). Conceptually, this does not cause any major trouble because we know in the end (from the analysis carried out in the previous sections, see in particular [\(4.23\)](#page-40-3)) that $|Z_t^0|$ and $|\nabla_x u_t|$ are bounded (by deterministic constants). However, these bounds cannot be directly used in the small time analysis.

To circumvent this difficulty, we introduce two auxiliary functions \hat{L}^0 : $\mathbb{R}^d \times \mathbb{R}^d \to \mathbb{R}$ and \hat{H} : $\mathbb{T}^d \times \mathbb{R}^d \to \mathbb{R}$ that are globally Lipschitz continuous. The function $(x_0, p) \mapsto \hat{L}^0(x_0, p)$ is understood as an approximation of the function $(x_0, p) \mapsto L^0(x_0, -\nabla_p H^0(x_0, p))$ and the function $(x, p) \in \mathbb{T}^d \times \mathbb{R}^d \mapsto \hat{H}(x, p)$ is understood as an approximation of H.

In a first time, we thus focus on the systems of two forward-backward equations (set on an arbitrary filtered probability space $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ satisfying the usual conditions and equipped with a Brownian motion $(B_t^0)_{0 \le t \le T}$ with values in \mathbb{R}^d):

$$
dX_t^0 = -\nabla_p H^0(X_t^0, Z_t^0) dt + \sigma_0 dB_t^0, \quad t \in [0, T],
$$

\n
$$
dY_t^0 = -\left(f_t^0(X_t^0, \mu_t) + \hat{L}^0(X_t^0, Z_t^0)\right) dt + \sigma_0 Z_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$

\n
$$
X_0^0 = x_0, \quad Y_T^0 = g^0(X_T^0, \mu_T),
$$
\n(5.1)

and

$$
\partial_t \mu_t - \frac{1}{2} \Delta_x \mu_t - \text{div}_x \left(\nabla_p \hat{H} \left(\cdot, \nabla_x u_t \right) \mu_t \right) = 0, \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mathrm{d}_t u_t(x) = \left(-\frac{1}{2} \Delta_x u_t(x) + \hat{H} \left(x, \nabla_x u_t(x) \right) - f_t(X_t^0, x, \mu_t) \right) \mathrm{d}t + \mathrm{d}m_t(x), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\mu_0 = \mu, \quad u_T(x) = g(X_T^0, x, \mu_T), \quad x \in \mathbb{T}^d.
$$
 (5.2)

This is only in the end of the section that we return back to the original equations (2.11) and (2.12) , see Subsection [5.8.](#page-79-0) Notice in particular that, differently from (2.12) , the martingale part in the backward equation of (5.2) is not represented as a stochastic integral (as there is no need at this stage).

We thus introduce the following variant of Assumption (A):

Assumption \hat{A} . There exist four reals $\mathcal{L} > 0$, $\kappa > 0$, $\hat{\kappa} > 0$ and $s > 3$ ($s \notin \mathbb{N}$) such that

 $(A1)$ The function $(f_t^0)_{0 \le t \le T}$ satisfies $\mathscr{D}^0(\kappa, \mathbf{s})$ and the function $(f_t)_{0 \le t \le T}$ satisfies $\mathscr{D}(\kappa, \mathbf{s}, \mathbf{s})$.

 $(\hat{A2})$ The function g^0 satisfies $\mathscr{C}^0(\mathfrak{L}, s)$ and $\mathscr{D}^0(\hat{\kappa}, s)$, and the function g satisfies $\mathscr{C}(\mathfrak{L}, s, s)$ and $\mathscr{D}(\hat{\kappa}, s, s)$.

(Â3) The functions H^0 and \hat{L}^0 have continuous joint derivatives up to the order 2 in x_0 and $\lfloor s \rfloor + 1$ in p, and the function \hat{H} has continuous joint derivatives up to the order $|s| + 1$ in (x, p) . All the existing derivatives (of order greater than 1) of \hat{L}^0 and \hat{H} are bounded by κ and the quantity $\hat{L}^0(x_0, p)$ is bounded by $\kappa(1+|p|)$. The quantity $H^0(x_0, p)$ is bounded by $\kappa(1+|p|^2)$, the gradient $\nabla_p H^0(x_0, p)$ is bounded by $\kappa(1+|p|)$. All the other existing derivatives of H^0 are bounded by $κ$.

(Â4) The function \hat{H} satisfies the coercivity condition $\nabla^2_{pp}\hat{H}(x,p) > 0$ (in the sense of positive definite matrices), for all $(x, p) \in \mathbb{T}^d \times \mathbb{R}^d$. For any $r > 0$, we let

$$
\zeta(r) := \inf_{(x,p)\in \mathbb{T}^d\times\mathbb{R}^d:|p|\leq r} \inf_{\rho\in\mathbb{R}^d:|\rho|=1} \rho \cdot (\nabla^2_{pp}\hat{H}(x,p)\rho) > 0.
$$

5.1. Local well-posedness

Theorem 5.1. *Under Assumption* (\hat{A})*, there exists a constant* $\mathfrak{C} > 0$ *, only depending on* \mathfrak{L} *,* κ *,* σ_0 *and* σ *s such that for* $T \leq \mathfrak{C}$ *,* the forward-backward system [\(5.1\)](#page-50-3)-[\(5.2\)](#page-50-2) admits a unique solution (X^0,Y^0,Z^0,μ,u,m) in the sense of Definition [2.10,](#page-9-1) i.e., (X^0,Y^0,Z^0) is required to be \mathbb{F}^0 -progressively measurable, with the trajectories of X^0 and Y^0 being continuous, such that $\sup_{0 \le t \le T} |X_t^0| \in L^2(\Omega^0, \mathbb{P}^0)$ and

$$
\Big\| \sup_{t \in [0,T]} |Y_t^0| \Big\|_{L^\infty(\Omega^0, \mathbb{P}^0)} + \sup_{\tau} \Big\| \mathbb{E}^0 \bigg[\int_{\tau}^T |Z_t^0|^2 dt \, | \mathcal{F}_\tau^0 \bigg] \Big\|_{L^\infty(\Omega^0, \mathbb{P}^0)} < \infty,
$$
\n
$$
(5.3)
$$

the second supremum being taken over \mathbb{F}^0 -stopping times τ ; and (μ, u, m) is required to be an \mathbb{F}^0 -progressively mea s urable process with values in $\mathcal{P}(\mathbb{T}^d)\times\mathcal{C}^s(\mathbb{T}^d)\times\mathcal{C}^{s-2}(\mathbb{T}^d)$, with continuous trajectories in $\mathcal{P}(\mathbb{T}^d)\times\mathcal{C}^r(\mathbb{T}^d)\times\mathcal{C}^{r-2}(\mathbb{T}^d)$ *for any* $r < s$ *, such that*

$$
\Big\| \sup_{t \in [0,T]} \Big[\|u_t\|_{s} \Big] \Big\|_{L^{\infty}(\Omega^0, \mathbb{P}^0)} + \Big\| \sup_{t \in [0,T]} \Big[\|m_t\|_{s-2} \Big] \Big\|_{L^{\infty}(\Omega^0, \mathbb{P}^0)} < \infty,
$$
\n(5.4)

and, for any $x \in \mathbb{T}^d$, $(m_t(x))_{0 \le t \le T}$ *is an* \mathbb{F}^0 -martingale satisfying $m_0(x) \equiv 0$.

Moreover, for any pair of initial conditions $((x_0^i, \mu^i))_{i=1,2} \in [\mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)]^2$, the corresponding pair of solutions $((\mathbf{X}^{0,i}, \mathbf{Y}^{0,i}, \mathbf{Z}^{0,i}, \boldsymbol{\mu}^i, \mathbf{u}^i, \mathbf{m}^i))_{i=1,2}$ to [\(5.1\)](#page-50-3)-[\(5.2\)](#page-50-2) satisfy, for any $p \in [1, 8]$,

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[\|u_{t}^{1}-u_{t}^{2}\|_{s}^{2p}+\|m_{t}^{1}-m_{t}^{2}\|_{s-2}^{2p}+\mathbb{W}_{1}(\mu_{t}^{1},\mu_{t}^{2})^{2p}\right]\right] \n+\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[|X_{t}^{0,1}-X_{t}^{0,2}|^{2p}+|Y_{t}^{0,1}-Y_{t}^{0,2}|^{2p}\right]+\left(\int_{0}^{T}|Z_{t}^{0,1}-Z_{t}^{0,2}|^{2}\mathrm{d}t\right)^{p}\right] \n\le C\left[|x_{0}^{1}-x_{0}^{2}|^{2p}+\mathbb{W}_{1}(\mu^{1},\mu^{2})^{2p}\right],
$$
\n(5.5)

for a constant C *only depending on* d*,* L*,* κ*,* σ⁰ *and* **s***. For the same* C*, we can assume that the left-hand sides of* [\(5.3\)](#page-51-0) *and* [\(5.4\)](#page-51-1) *are bounded by* C*.*

Proof. The proof relies on a suitable application of the Banach fixed point theorem. Throughout, T is less than 1.

First Step. Assuming that \mathbb{F}^0 is generated by \mathcal{F}^0_0 and \mathbf{B}^0 , we first construct the mapping to which we will eventually apply the fixed point theorem. Inputs X^0 are taken in the set $\mathscr{S}^2(\mathbb{R}^d)$ of \mathbb{F}^0 -progressively measurable continuous processes with values in \mathbb{R}^d such that $\sup_{t\in[0,T]}|X_t^0|\in L^2(\Omega^0,\mathbb{P}^0)$. For $X^0\in\mathscr{S}^2(\mathbb{R}^d)$, we can follow the proof of [\[14,](#page-86-1) Theorem 4.3.1] to obtain, for a small time horizon, a unique solution to

$$
\partial_t \tilde{\mu}_t - \frac{1}{2} \Delta_x \tilde{\mu}_t - \text{div}_x \left(\nabla_p \hat{H} (\cdot, \nabla_x \tilde{u}_t) \tilde{\mu}_t \right) = 0,
$$

\n
$$
\mathrm{d}_t \tilde{u}_t(x) = \left(-\frac{1}{2} \Delta_x \tilde{u}_t(x) + \hat{H}(x, \nabla_x \tilde{u}_t(x)) - f_t(X_t^0, x, \tilde{\mu}_t) \right) \mathrm{d}t + \mathrm{d}m_t(x), \quad (t, x) \in [0, T] \times \mathbb{T}^d,
$$

\n
$$
\tilde{\mu}_0 = \mu, \quad \tilde{u}_T(x) = g(X_T^0, x, \tilde{\mu}_T), \quad x \in \mathbb{T}^d,
$$
\n(5.6)

where $(\tilde{\boldsymbol{\mu}}, \tilde{\boldsymbol{u}}, \tilde{\boldsymbol{m}})$ is an \mathbb{F}^0 -progressively measurable processes with values in $\mathcal{P}(\mathbb{T}^d) \times \mathcal{C}^s(\mathbb{T}^d) \times \mathcal{C}^{s-2}(\mathbb{T}^d)$, with continuous trajectories in $\mathcal{P}(\mathbb{T}^d) \times \mathcal{C}^{\mathbb{T}}(\mathbb{T}^d) \times \mathcal{C}^{\mathbb{T}^{-2}}(\mathbb{T}^d)$ for any $\mathbb{T} < \mathbb{s}$, such that

$$
\sup_{t \in [0,T]} \{ \|\tilde{u}_t\|_{s} + \|m_t\|_{s-2} \} \in L^{\infty}(\Omega^0, \mathcal{F}^0, \mathbb{P}^0), \tag{5.7}
$$

and $(m_t(x))_{0 \le t \le T}$ is an \mathbb{F}^0 -martingale starting from 0 for any $x \in \mathbb{T}^d$.

Actually, the framework is not (exactly) the same as in [\[14,](#page-86-1) Theorem 4.3.1]. First, X^0 plays the role of the common noise, even though it is not acting additively (as B^0 does in [\[14\]](#page-86-1)). Second (and this is the main difference), [\[14,](#page-86-1) Theorem 4.3.1] is proven under the Lasry-Lions monotonicity condition, which we have not assumed here. However, the proof of [\[14,](#page-86-1) Theorem 4.3.1] (which relies on the method of continuation) remains conceptually the same in our context. Essentially, the parameter ϖ in [\[14](#page-86-1), Subsection 4.3.5] is here taken as 0 and the role of ϵ (which is assumed to be small in [\[14\]](#page-86-1)) is now played by T. Although the correspondence between ϵ and T is not immediate, the similarity between the two parameters can be well noticed in the proof of [\[14](#page-86-1), Lemma 4.3.9]: the Cauchy-Schwarz inequality that permits to pass from [\[14,](#page-86-1) (4.30)] to [\[14](#page-86-1), (4.31)] makes an additional \sqrt{T} appear in front of the parameter ϑ in [\[14,](#page-86-1) (4.30)]. This additional \sqrt{T} then plays the role of ϵ in the second step of [\[14,](#page-86-1) Subsection 4.3.5]. With this analogy in mind, one notices

from (c) in the second step of [\[14,](#page-86-1) Subsection 4.3.5] that the threshold $\mathfrak{C} > 0$ below which T must be taken only depends on q through $\mathfrak L$ (and not $\hat \kappa$).

To be complete, one mentions another subtlety: in [\[14,](#page-86-1) Theorem 4.3.1], continuities of \tilde{u} and \tilde{m} are stated in $\mathcal{C}^{\lfloor s \rfloor}(\mathbb{T}^d)$ and $C^{\lfloor s \rfloor - 2}(\mathbb{T}^d)$ respectively, but there is no difficulty in replacing $\lfloor s \rfloor$ by any $\mathbb{r} \in (\lfloor s \rfloor, s)$ (for instance by combining the above bound with an interpolation argument). In this regard, we emphasize that *Bochner* measurability (with values in $C^{s}(\mathbb{T}^{d})$ and $C^{s-2}(\mathbb{T}^{d})$) can be obtained as explained in Remark [2.11,](#page-9-2) by noticing that, for $t < T$, u_t and m_t takes values in $C^{5'}(\mathbb{T}^d)$ and $C^{5'-2}(\mathbb{T}^d)$ for some $s' > s$. This follows from the smoothing properties of the heat kernel: we refer to [\[14](#page-86-1), (4.16)] for the way it can be used here. Following Proposition [3.4](#page-22-0) (and [\[14,](#page-86-1) Proposition 4.3.8]), we notice that $\sup_{t\in[0,T]}\{\|\tilde{u}_t\|_{s}+\|m_t\|_{s-2}\}$ can be bounded by constants only depending on the parameters $\mathfrak{L}, \kappa, \sigma_0$, \mathfrak{s} and T (but not $\hat{\kappa}$) in $(\hat{\mathbf{A}})$.

Next we define $(\tilde{X}^0, \tilde{Y}^0, \tilde{Z}^0)$ as the strong solution to the following FBSDE system (see [\[26\]](#page-86-0)):

$$
d\tilde{X}_t^0 = -\nabla_p H^0(\tilde{X}_t^0, \tilde{Z}_t^0) dt + \sigma_0 dB_t^0, \quad t \in [0, T],
$$

\n
$$
d\tilde{Y}_t^0 = -\left(f_t^0(\tilde{X}_t^0, \tilde{\mu}_t) + \hat{L}^0(\tilde{X}_t^0, \tilde{Z}_t^0)\right) dt + \sigma_0 \tilde{Z}_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$
\n
$$
\tilde{X}_0^0 = x_0, \quad \tilde{Y}_T^0 = g^0(\tilde{X}_T^0, \tilde{\mu}_T).
$$
\n(5.8)

Setting $\mathfrak{T}(\mathbf{X}^0) := \tilde{\mathbf{X}}^0$, we define a map $\mathfrak{T} : \mathscr{S}^2(\mathbb{R}^d) \to \mathscr{S}^2(\mathbb{R}^d)$. In the rest of the proof, we show that \mathfrak{T} is a contraction mapping from $\mathscr{S}^2(\mathbb{R}^d)$ into itself (for T small enough).

Second Step. Denote the solutions to [\(5.6\)](#page-51-2) with two different inputs $X^{0,1}$ and $X^{0,2}$ with $X_0^{0,1} = X_0^{0,2} = x_0$ by $(\tilde{\mu}^1, \tilde{u}^1, \tilde{m}^1)$ and $(\tilde{\mu}^2, \tilde{u}^2, \tilde{m}^2)$, and the corresponding solutions to [\(5.8\)](#page-52-0) by $(\tilde{X}^{0,1}, \tilde{Y}^{0,1}, \tilde{Z}^{0,1})$ and $(\tilde{X}^{0,2}, \tilde{Y}^{0,2}, \tilde{Z}^{0,2})$. By the standard duality method of mean field games (without using the monotonicity condition of f_t and g , see for instance $[14, (4.29)]$ $[14, (4.29)]$) and thanks to [\(5.7\)](#page-51-3) (which, together with $(A4)$, says that we are working on a subdomain of the space on which the Hessian matrix $\nabla^2_{pp}H$ is bounded from below by a strictly positive constant), we have

$$
\mathbb{E}^{0}\bigg[\int_{0}^{T}\Big(|\nabla_{x}(\tilde{u}_{t}^{1}-\tilde{u}_{t}^{2})|^{2},\tilde{\mu}_{t}^{1}+\tilde{\mu}_{t}^{2}\Big)dt\bigg] \leq C\mathbb{E}^{0}\bigg[\sup_{t\in[0,T]}|X_{t}^{0,1}-X_{t}^{0,1}|^{2}+\sup_{t\in[0,T]}|\mathcal{N}_{1}(\tilde{\mu}_{t}^{1},\tilde{\mu}_{t}^{2})^{2}\bigg].\tag{5.9}
$$

As the maps $x \mapsto \nabla_p H(x, \nabla_x \tilde{u}^1(t, x))$ and $p \mapsto \nabla_p H(x, p)$ are Lipschitz continuous (with Lipschitz constants only depending on the parameters $\mathfrak{L}, \kappa, \sigma_0$, s and T, but not $\hat{\kappa}$), we use [\(5.7\)](#page-51-3) and [\(5.9\)](#page-52-1) to derive (the reader may compare with the derivation of $[14, (3.7)]$

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[\mathbb{W}_{1}(\tilde{\mu}^{1}_{t},\tilde{\mu}^{2}_{t})^{2}\right]\right] \leq C \mathbb{E}^{0}\left[\left(\int_{0}^{T}\left(|\nabla_{x}(\tilde{u}^{1}_{t}-\tilde{u}^{2}_{t})|^{2},\tilde{\mu}^{1}_{t}+\tilde{\mu}^{2}_{t}\right)^{\frac{1}{2}}\mathrm{d}t\right)^{2}\right] \n\leq C T \mathbb{E}^{0}\left[\int_{0}^{T}\left(|\nabla_{x}(\tilde{u}^{1}_{t}-\tilde{u}^{2}_{t})|^{2},\tilde{\mu}^{1}_{t}+\tilde{\mu}^{2}_{t}\right)\mathrm{d}t\right] \n\leq C T \mathbb{E}^{0}\left[\sup_{t\in[0,T]}|X^{0,1}_{t}-X^{0,1}_{t}|^{2}+\sup_{t\in[0,T]} \mathbb{W}_{1}(\tilde{\mu}^{1}_{t},\tilde{\mu}^{2}_{t})^{2}\right],
$$

where C only depends on $\mathfrak{L}, \kappa, \sigma_0$, s. And then, for $CT \leq 1/2$, we obtain

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[\mathbb{W}_{1}(\tilde{\mu}_{t}^{1},\tilde{\mu}_{t}^{2})^{2}\right]\right] \leq 2CT\mathbb{E}^{0}\left[\sup_{t\in[0,T]}|X_{t}^{0,1}-X_{t}^{0,1}|^{2}\right].
$$
\n(5.10)

By standard short time FBSDE estimates (see [\[26\]](#page-86-0)), we can derive, still under the condition $CT \leq 1/2$ (but for a possibly new value of C that only depends on \mathfrak{L} , κ , σ_0 and s),

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[|\tilde{X}_{t}^{0,1}-\tilde{X}_{t}^{0,2}|^{2}\right]\leq C\sup_{t\in[0,T]}\mathbb{E}^{0}\left[\mathbb{W}_{1}(\tilde{\mu}_{t}^{1},\tilde{\mu}_{t}^{2})^{2}\right]\leq CT\mathbb{E}^{0}\left[\sup_{t\in[0,T]}|X_{t}^{0,1}-X_{t}^{0,1}|^{2}\right].
$$
 (5.11)

Since $CT \leq 1/2$, we get

$$
\mathbb{E}^{0}\bigg[\underset{t\in[0,T]}{\sup}|\tilde{X}^{0,1}_t-\tilde{X}^{0,2}_t|^2\bigg] \leq \frac{1}{2}\mathbb{E}^{0}\bigg[\underset{t\in[0,T]}{\sup}|X^{0,1}_t-{X}^{0,2}_t|^2\bigg],
$$

which implies $\mathfrak T$ is a contraction mapping from $\mathscr{S}^2(\mathbb{R}^d)$ to $\mathscr{S}^2(\mathbb{R}^d)$. Therefore, there exists a unique fixed point $X^0 \in \mathscr{S}^2(\mathbb{R}^d)$ with $X_0^0 = x_0$. For this choice of X^0 , the FBSPDE (μ, u, m) . Since X^0 is the fixed point of $\mathfrak T$, the FBSDE system [\(5.1\)](#page-50-3) admits a unique strong solution (X^0, Y^0, Z^0) . In this way, we get existence and uniqueness of a solution to the pair (5.1) – (5.2) . The bound (5.4) follows from (5.7) and [\(5.3\)](#page-51-0) follows from the argument used in the proof of Lemma [3.8.](#page-28-0)

Third Step. We now prove the stability property, removing the assumption that \mathbb{F}^0 is generated by \mathcal{F}^0_0 and \mathbf{B}^0 (which will Final step. We now prove the stability property, removing the assumption that \mathbb{F} is generated by \mathcal{F}_0 and \mathbf{B} (which will imply in particular uniqueness of the solution constructed above, but on the larger conditions $(x_0^1, \mu^1), (x_0^2, \mu^2) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, let $(X^{0,1}, Y^{0,1}, Z^{0,1}, \mu^1, u^1, m^1)$ and $(X^{0,2}, Y^{0,2}, Z^{0,2}, \mu^2, u^2, m^2)$ be the solutions of the forward-backward system (5.1) - (5.2) with (respective) initial data (x_0^1, μ^1) and (x_0^2, μ^2) .

Note that

$$
d_t \left(u_t^1(x) - u_t^2(x) \right)
$$

= $\left[-\frac{1}{2} \Delta_x \left(u_t^1(x) - u_t^2(x) \right) - \left(f_t(X_t^{0,1}, x, \mu_t^1) - f_t(X_t^{0,2}, x, \mu_t^2) \right) \right]$
+ $\left\{ \int_0^1 \nabla_p \hat{H} \left(x, r \nabla_x u_t^1(x) + (1 - r) \nabla_x u^2(x) \right) dr \right\} \cdot \nabla_x \left(u_t^1(x) - u_t^2(x) \right) \right] dt + d \left(m_t^1(x) - m_t^2(x) \right).$ (5.12)

and

$$
u_T^1(x) - u_T^2(x) = g(X_T^{0,1}, x, \mu_T^1) - g(X_T^{0,2}, x, \mu_T^2).
$$

Following [\[14,](#page-86-1) (4.14)–(4.16)] and recalling $T \le 1$, we have, for all $t \in [0, T]$,

$$
\|u_t^1 - u_t^2\|_{s}
$$

\n
$$
\leq C\mathbb{E}^0 \Big[\sup_{r \in [0,T]} \left\|f_r(X_r^{0,1}, \cdot, \mu_r^1) - f_r(X_r^{0,2}, \cdot, \mu_r^2)\right\|_{s-1} + \left\|g(X_T^{0,1}, \cdot, \mu_T^1) - g(X_T^{0,2}, \cdot, \mu_T^2)\right\|_{s} |\mathcal{F}_t\Big]
$$

\n
$$
\leq C\mathbb{E}^0 \Big[\sup_{r \in [0,T]} |X_r^{0,1} - X_r^{0,2}| + \sup_{r \in [0,T]} \mathbb{W}_1(\mu_r^1, \mu_r^2) |\mathcal{F}_t\Big],
$$

for a constant C that only depends on $\mathfrak{L}, \kappa, \sigma_0$ and s.

Taking power $2p$, for $p \in [1, 8]$, and then expectation (under \mathbb{P}^0) on both sides, we obtain, by means of Doob's inequality,

$$
\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|u_{t}^{1}-u_{t}^{2}\|_{s}^{2p}\Big]\leq C\mathbb{E}\Big[\sup_{r\in[0,T]}|X_{r}^{0,1}-X_{r}^{0,2}|^{2p}+\sup_{r\in[0,T]}\mathbb{W}_{1}(\mu_{r}^{1},\mu_{r}^{2})^{2p}\Big],\tag{5.13}
$$

where we allowed the constant C to vary from line to line as long as it only depends on $\mathfrak{L}, \kappa, \sigma_0, \mathfrak{s}$.

Moreover, similar to the derivation of (5.10) (but with a power $2p$ in the computations, which does not change the argument), we can obtain (from (5.13))

$$
\mathbb{E}^{0}\Big[\sup_{t\in[0,T]} \mathbb{W}_{1}(\mu_{t}^{1},\mu_{t}^{2})^{2p}\Big] \leq C\Big\{\mathbb{W}_{1}(\mu^{1},\mu^{2})^{2p} + \mathbb{E}^{0}\Big[\Big(\int_{0}^{T} \big(|\nabla_{x}(u_{t}^{1}-u_{t}^{2})|^{2},\mu_{t}^{1}+\mu_{t}^{2}\big)\mathrm{d}t\Big)^{2p}\Big]\Big\}\leq C\Big\{\mathbb{W}_{1}^{2}(\mu^{1},\mu^{2}) + T^{2p}\mathbb{E}^{0}\Big[\sup_{t\in[0,T]} \big\|\nabla_{x}(u_{t}^{1}-u_{t}^{2})\big\|_{L^{\infty}}^{2p}\Big]\Big\}\leq C\Big\{\mathbb{W}_{1}^{2}(\mu^{1},\mu^{2}) + T^{2p}\mathbb{E}^{0}\Big[\sup_{t\in[0,T]} |X_{t}^{0,1} - X_{t}^{0,2}|^{2p} + \sup_{t\in[0,T]} \mathbb{W}_{1}(\mu_{t}^{1},\mu_{t}^{2})^{2p}\Big]\Big\}.
$$

For $CT \leq 1/2$ (and since $T \leq 1$), we obtain

$$
\mathbb{E}^{0}\Big[\sup_{t\in[0,T]} \mathbb{W}_1(\mu_t^1, \mu_t^2)^{2p}\Big] \le 2C\bigg\{\mathbb{W}_1(\mu^1, \mu^2)^{2p} + T\mathbb{E}^{0}\Big[\sup_{t\in[0,T]} |X_t^{0,1} - X_t^{0,2}|^{2p}\Big]\bigg\}.\tag{5.14}
$$

By standard short time FBSDE estimates (see again [\[26\]](#page-86-0)), we obtain (still under a condition of the form $CT \leq 1/2$)

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left[|X_t^{0,1}-X_t^{0,2}|^{2p}+|Y_t^{0,1}-Y_t^{0,2}|^{2p}\right]+\left(\int_0^T|Z_t^{0,1}-Z_t^{0,2}|^2\mathrm{d}t\right)^p\right] \leq C\left(|x_0^1-x_0^2|^{2p}+\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\mathbb{W}_1(\mu_t^1,\mu_t^2)^{2p}\right]\right) \leq \max(C,2C^2)\left(|x_0^1-x_0^2|^{2p}+\mathbb{W}_1(\mu^1,\mu^2)^{2p}+T\mathbb{E}^{0}\left[\sup_{t\in[0,T]}|X_t^{0,1}-X_t^{0,2}|^{2p}\right]\right).
$$

If $\max(C, 2C^2)T \leq 1/2$ (and $T \leq 1$), then

$$
\mathbb{E}^{0}\bigg[\sup_{t\in[0,T]} \left[|X_t^{0,1} - X_t^{0,2}|^{2p} + |Y_t^{0,1} - Y_t^{0,2}|^{2p} \right] + \left(\int_0^T |Z_t^{0,1} - Z_t^{0,2}|^2 dt \right)^p \bigg]
$$

$$
\leq 2 \max(C, 2C^2) \left(|x_0^1 - x_0^2|^{2p} + \mathbb{W}_1(\mu^1, \mu^2)^{2p} \right).
$$

Plugging the above inequality into (5.13) and (5.14) , we get (for a new value of C)

$$
\mathbb{E}^0\Big[\sup_{t\in[0,T]}\|u^1_t-u^2_t\|^{2p}_{\rm s}+\sup_{t\in[0,T]}\mathbb{W}_1(\mu^1_t,\mu^2_t)^{2p}\Big]\leq C\Big[|x^1_0-x^2_0|^{2p}+\mathbb{W}_1(\mu^1,\mu^2)^{2p}\Big].
$$

Recalling [\(5.12\)](#page-53-2) and inserting the above bound, we deduce that

$$
\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|m_{t}^{1}-m_{t}^{2}\|_{\mathbb{S}-2}^{2p}\Big]\leq C\Big[|x_{0}^{1}-x_{0}^{2}|^{2p}+\mathbb{W}_{1}(\mu^{1},\mu^{2})^{2p}\Big]
$$

.

In summary, we obtain (5.5) .

5.2. Linearized system

We now address the following general linearized forward-backward system, set on $[0, T]$:

$$
d\delta X_t^0 = \left[-p_t \delta X_t^0 - q_t \delta Z_t^0 + a_t \right] dt,
$$

\n
$$
d\delta Y_t^0 = \left[l_t \cdot \delta X_t^0 + o_t \cdot \delta Z_t^0 - \nabla_{x_0} f_t^0(X_t^0, \mu_t) \cdot \delta X_t^0 - \left(\delta_\mu f_t^0(X_t^0, \mu_t), \delta \mu_t \right) + b_t \right] dt + \sigma_0 \delta Z_t^0 \cdot dB_t^0,
$$

\n
$$
\delta X_0^0 = \triangle x_0, \quad \delta Y_T^0 = \nabla_{x_0} g^0(X_T^0, \mu_T) \cdot \delta X_T^0 + \left(\delta_\mu g^0(X_T^0, \mu_T), \delta \mu_T \right) + c_T;
$$

\n
$$
\partial_t \delta \mu_t - \frac{1}{2} \Delta_x \delta \mu_t - \text{div}_x (\Upsilon_t(x) \delta \mu_t + \mu_t \Gamma_t(x) \nabla_x \delta u_t(x)) - \text{div}_x (d_t) = 0, \quad \text{on } \mathbb{T}^d,
$$

\n
$$
d_t \delta u_t(x) = \left[-\frac{1}{2} \Delta_x \delta u_t(x) + \Upsilon_t(x) \cdot \nabla_x \delta u_t(x) - \nabla_{x_0} f_t(X_t^0, x, \mu_t) \cdot \delta X_t^0 - \left(\delta_\mu f_t(X_t^0, x, \mu_t), \delta \mu_t \right) + j_t(x) \right] dt
$$

\n
$$
+ d_t \delta m_t(x), \quad x \in \mathbb{T}^d,
$$

\n
$$
\delta \mu_0 = \triangle \mu, \quad \delta u_T(x) = \nabla_{x_0} g(X_T^0, x, \mu_T) \cdot \delta X_T^0 + \left(\delta_\mu g(X_T^0, x, \mu_T), \delta \mu_T \right) + k_T(x) \quad x \in \mathbb{T}^d,
$$

\n(5.15)

which should be interpreted as a generalized version of the system satisfied by the derivative of the flow induced by the solution to (5.1) - (5.2) . On top of Assumption (\hat{A}) , the coefficients driving (5.15) are required to satisfy the set of conditions below:

Assumption C. For a real $r \in (\lfloor s \rfloor, s)$ and for another real $\mathfrak{K} > 1$,

- (i) The initial conditions $\triangle \mu$ and $\triangle x_0$ are deterministic, taking values in C^{-r+1} and \mathbb{R}^d respectively;
- (ii) The random variable k_T is \mathcal{F}_T^0 -measurable with values in $\mathcal{C}^s(\mathbb{T}^d)$, and satisfies $\exp_{\omega^0 \in \Omega^0} ||k_T||_{\mathbb{S}} < +\infty$;
- (iii) The process $(j_t)_{0 \le t \le T}$ is \mathbb{F}^0 -adapted with continuous paths from $[0, T]$ to $\mathcal{C}^{s'}(\mathbb{T}^d)$, for any $s' \in (0, s 1)$, and satisfies the bound $\overline{\text{essup}}_{\omega^0 \in \Omega^0} \sup_{t \in [0,T]} \|j_t\|_{s-1} < \infty$; the process $(d_t)_{0 \le t \le T}$ is \mathbb{F}^0 -adapted with continuous paths from $[0, T]$ to $[\mathcal{C}^{-\mathbf{r}' + 2}(\mathbb{T}^d)]^d$, for any $\mathbf{r}' > \mathbf{r}$ and satisfies $\operatorname{essup}_{\omega^0 \in \Omega^0} \operatorname{sup}_{t \in [0, T]} ||d_t||_{-\mathbf{r} + 2} < +\infty;$

 \Box

- (iv) The process $(\Upsilon_t)_{0 \le t \le T}$ is \mathbb{F}^0 -adapted with continuous paths from $[0, T]$ to $\mathcal{C}^{\mathfrak{s}'}(\mathbb{T}^d, \mathbb{R}^d) \simeq [\mathcal{C}^{\mathfrak{s}'}(\mathbb{T}^d)]^d$, for any $\mathfrak{s}' \in$ $(0, s - 1)$, and satisfies essup_{ω0∈Ω0} sup_{t∈[0,T]} $||\Upsilon_t||_{s-1} \leq \mathfrak{K}$;
- (v) The process $(\Gamma_t)_{0 \le t \le T}$ is \mathbb{F}^0 -adapted with continuous paths from $[0, T]$ to $\mathcal{C}^1(\mathbb{T}^d, \mathbb{R}^{d \times d}) \simeq [\mathcal{C}^1(\mathbb{T}^d)]^{d \times d}$; it satisfies $\text{essup}_{\omega^0 \in \Omega^0} \text{sup}_{t \in [0,T]} ||\Gamma_t||_1 \leq \mathfrak{K}$ and, \mathbb{P}^0 -almost surely,

$$
\forall (t, x) \in [0, T] \times \mathbb{T}^d, \quad \mathfrak{K}^{-1} I_d \le \Gamma_t(x) \le \mathfrak{K} I_d;
$$

(vi) The processes $(a_t)_{0 \le t \le T}$ and $(b_t)_{0 \le t \le T}$ belong to $\mathcal{H}^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}^0; \mathbb{R}^d)$ and $\mathcal{H}^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0; \mathbb{R})$ respectively; the random variable c_T is in $L^2(\Omega^0, \mathcal{F}_T^0, \mathbb{P}^0; \mathbb{R})$; the processes $(p_t)_{0 \le t \le T}$, $(q_t)_{0 \le t \le T}$, $(l_t)_{0 \le t \le T}$ and $(o_t)_{0 \le t \le T}$ are \mathbb{F}^0 -adapted, with respective values in \mathbb{R}^d , $\mathbb{R}^{d \times d}$, \mathbb{R}^d and \mathbb{R}^d , and satisfies essup_{$\omega^0 \in \Omega^0$} sup $t \in [0,T]$ $[|p_t| +$ $|q_t| + |l_t| + |o_t| \leq \mathfrak{K}.$

Within the above framework and under the additional condition that $(d_t)_{0 \le t \le T}$ takes values in $C^{-\tau'+2}(\mathbb{T}^d)$ for some $\mathbb{r}' < \mathbb{r}$, we let, for any $p \geq 1$,

$$
M_p := |\triangle x_0|^{2p} + ||\triangle \mu||_{-\mathbb{r}+1}^{2p} + \mathbb{E}^0 \Big[\sup_{t \in [0,T]} \Big(||d_t||_{-\mathbb{r}+2}^{2p} + ||j_t||_{\mathbb{S}-1}^{2p} \Big) + \Big(\int_0^T (|a_t| + |b_t|) dt \Big)^{2p} + |c_T|^{2p} + ||k_T||_{\mathbb{S}}^{2p} \Big].
$$
 (5.16)

Theorem 5.2. *Under Assumptions* (\hat{A}) *and* (C)*, there exists a real* $\mathfrak{C} > 0$ *, only depending on d,* κ *,* \hat{A} *,* \mathfrak{L} *,* σ_0 *<i>and* ($\mathfrak{r}, \mathfrak{s}$)*, such that for* $T \leq \mathfrak{C}$, the forward-backward system [\(5.15\)](#page-54-0) admits a unique solution $(\delta \mu, \delta u, \delta m, \delta X^0, \delta Y^0, \delta Z^0)$, adapted with respect to the filtration \mathbb{F}^0 and with values in $C^{-r+1}(\mathbb{T}^d)\times C^s(\mathbb{T}^d)\times C^{s-2}(\mathbb{T}^d)\times\mathbb{R}^d\times\mathbb{R}\times\mathbb{R}^d$, satisfying

 (a) $(\delta\mu, \delta u, \delta m)$ has continuous trajectories in $C^{-u+1}(\mathbb{T}^d) \times C^u(\mathbb{T}^d) \times C^{u-2}(\mathbb{T}^d)$ for any $u \in (\mathbb{r}, \mathbb{s})$, and

$$
\text{essup}_{\omega \in \Omega^0} \sup_{t \in [0,T]} \left(\|\delta u_t\|_{s} + \|\delta m_t\|_{s-2} + \|\delta \mu_t\|_{-r+1} \right) < +\infty; \tag{5.17}
$$

- (*b*) for any $x \in \mathbb{T}^d$, $(\delta m_t(x))_{0 \le t \le T}$ *is an* \mathbb{F}^0 -martingale;
- (c) $(\delta X^0, \delta Y^0, \delta Z^0)$ belongs to $\mathscr{S}^2(\mathbb{R}^d) \times \mathscr{S}^2(\mathbb{R}) \times \mathscr{H}^2(\mathbb{R}^d)$.

Moreover, if $(d_t)_{t \in (0,T]}$ takes values in $[\mathcal{C}^{-\mathfrak{r}'+2}(\mathbb{T}^d)]^d$ for some $\mathfrak{r}' < \mathfrak{r}$ *, there exists a constant* C*, only depending on d,* κ *,* \mathfrak{K} *,* \mathfrak{L} *,* σ_0 *and* $(\mathbb{r}, \mathfrak{s})$ *, such that, for any* $p \in [1, 8]$ *,*

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]} \left(\|\delta\mu_t\|_{-\mathbb{r}+1}^{2p} + \|\delta u_t\|_{\mathbb{S}}^{2p} + \|\delta m_t\|_{\mathbb{S}-2}^{2p} \right) + \sup_{t\in[0,T]} \left(|\delta X_t^0|^{2p} + |\delta Y_t^0|^{2p} \right) + \left(\int_0^T |\delta Z_t^0|^2 dt \right)^p \right] \tag{5.18}
$$
\n
$$
\leq C_p M_p.
$$

Remark 5.3. 1. In the statement of Theorem [5.2,](#page-55-0) measurability and progressive-measurability are understood in the same *Bochner* sense as in the third item of Remark [2.11.](#page-9-2) In comparison, the novelty here is that the forward component of the FBSPDE is also impacted by issues of measurability. The difficulty is the same as in Definition [2.10:](#page-9-1) the dual space $C^{-r+1}(\mathbb{T}^d)$ of $C^{r-1}(\mathbb{T}^d)$ is not separable. The remedy is the same: we can easily prove (see for instance [\[14,](#page-86-1) Lemma 3.3.1]) that, for $t \in (0,T]$, $\delta \mu_t$ is in fact in the subspace $C^{-\mathbf{r}'+1}(\mathbb{T}^d)$, for some $\mathbf{r}' < \mathbf{r}$, which is separable when equipped with $\|\cdot\|_{-\mathbb{r}+1}$.

Actually, some care is also needed in the formulation of Assumption (C). In item (ii) therein, measurability is also understood in *Bochner* sense. In item (iii), each j_t is regarded as a random variable with values in $C^{s'}(\mathbb{T}^d)$, for any s' ∈ (0, s − 1). Since j_t takes almost surely values in $\mathcal{C}^{s-1}(\mathbb{T}^d)$, this says that j_t is in fact *Bochner* measurable with values in $C^{s'}(\mathbb{T}^d)$, for any $s' < s - 1$. The same argument applies to $(d_t)_{0 \le t \le T}$ and $(\Psi_t)_{0 \le t \le T}$ in items (iii) and (iv) in Assumption (C) .

2. In [\(5.17\)](#page-55-1), $\sup_{t\in[0,T]} \|\delta\mu_t\|_{-\tau+1}$ can be shown to be measurable by the same argument as above. For any $\varepsilon > 0$, we have in fact a (deterministic) bound for $\sup_{\varepsilon \le t \le T} \|\delta \mu_t\|_{-\pi'+1}$ for some $\pi' < \pi$, from which we deduce that $\delta \mu$ has continuous trajectories from $[\varepsilon, T]$ to $C^{-r+1}(\mathbb{T}^d)$. This proves that $\sup_{t \in [\varepsilon, T]} \|\delta \mu_t\|_{-\varepsilon+1}$ is measurable. Letting ε tend to 0, we deduce that $\sup_{t\in(0,T]}\|\delta\mu_t\|_{-\varepsilon+1}$ and then $\sup_{t\in[0,T]}\|\delta\mu_t\|_{-\varepsilon+1}$ are measurable. That said, it must be clear that there is actually no need of measurability of the supremum to write down the result. Instead, we can just say that there exists $C > 0$ such that $\mathbb{P}^0(\{\forall t \in [0, T], \|\delta \mu_t\|_{-\mathbb{r}+1} \le C\}) = 1$, which is licit since \mathbb{P}^0 is complete. In particular, this is exactly how the condition $\text{essup}_{\omega^0 \in \Omega^0} \text{sup}_{t \in [0,T]} ||d_t||_{-\tau+2} < +\infty$ in item (iii) of (C) should be understood: formally, we cannot prove that $\sup_{t\in[0,T]} ||dt||_{-r+2}$ is measurable, but the condition still makes sense. (Notice that, in comparison, sup_{t∈[0,T]} $||j_t||_{s-1}$ – also in item (iii) of Assumption (C)— is a random variable, see Remark [2.11.](#page-9-2) Here, the same argument does not hold for processes taking values in the dual space $\mathcal{C}^{-\mathbb{r}+2}(\mathbb{T}^d)$ because the norm $\|\cdot\|_{-\mathbb{r}+2}$ is not lower-semicontinuous on $\mathcal{C}^{-\mathbb{r}'+2}(\mathbb{T}^d)$ for $\mathbb{r}' > \mathbb{r}$.)

3. The fact that $\sup_{t\in[0,T]}||d_t||_{-\tau+2}$ may not be a random variable explains why we need another condition to define M in [\(5.16\)](#page-55-2) (in order to guarantee that the *sup* is in fact measurable). Indeed, if d_t takes values in $\left[\mathcal{C}^{-r'+2}(\mathbb{T}^d) \right]^{d}$ for some $\mathbb{r}' < \mathbb{r}$, then (for any $\omega^0 \in \Omega^0$)

$$
||d_t||_{-\mathbb{r}+2} = \sup_{g \in \mathcal{C}^{\infty}(\mathbb{T}^d): ||g||_{\mathbb{r}-2}} (d_t, g).
$$

And then, using the fact that the process d has continuous trajectories with values in $[\mathcal{C}^{-\mathfrak{r}''+2}(\mathbb{T}^d)]^d$ for any $\mathfrak{r}'' > \mathfrak{r}$ (see item (ii) in Assumption (C)), we deduce have that, for any $g \in C^{\infty}(\mathbb{T}^d)$,

$$
(d_t, g) = \lim_{s \to t, s \in \mathbb{Q}} (d_s, g) \le \liminf_{s \to t, s \in \mathbb{Q}} ||d_s||_{-\mathbb{r}+2},
$$

and then

$$
\sup_{t \in [0,T]} \|d_t\|_{-\mathbb{r}+2} = \sup_{t \in [0,T] \cap \mathbb{Q}} \|d_t\|_{-\mathbb{r}+2}.
$$

It then remains to prove that each d_t is a *Bochner* random variable with values in $[\mathcal{C}^{-r+2}(\mathbb{T}^d)]^d$, but this follows from the fact that it is a random variable with values in $\left[\mathcal{C}^{-\mathbf{r}''+2}(\mathbb{T}^d)\right]^d$ for any \mathbf{r}'' $\geq \mathbf{r}$ and that it takes values in $\mathcal{C}^{-\mathbb{r}'+2}(\mathbb{T}^d)$ for some $\mathbb{r}' < \mathbb{r}$.

Proof. Small time unique solvability of [\(5.15\)](#page-54-0) is proven by means of a suitable contraction argument, which is explained in the first two steps below. In the third step, we establish [\(5.18\)](#page-55-3). Proceeding as in the proof of Theorem [5.1,](#page-50-1) one can assume that \mathbb{F}^0 is generated by \mathcal{F}^0_0 and \mathbf{B}^0 .

First Step. The first of the two steps underpinning the contraction argument relies on the following idea. For a given input $\delta X^0 := (\delta X_t^0)_{0 \le t \le T} \in \mathscr{S}^2(\mathbb{R}^d)$ satisfying $\delta X_0^0 = \Delta x_0$, we want to solve the FBSPDE, on $[t, T]$,

$$
\partial_t \delta \tilde{\mu}_t - \frac{1}{2} \Delta_x \delta \tilde{\mu}_t - \text{div}_x \Big(\Upsilon_t(x) \delta \tilde{\mu}_t + \mu_t \Gamma_t(x) \nabla_x \delta \tilde{u}_t(x) \Big) - \text{div}_x (d_t) = 0, \quad \text{on } \mathbb{T}^d,
$$

\n
$$
\mathrm{d}_t \delta \tilde{u}_t(x) = \Big[-\frac{1}{2} \Delta_x \delta \tilde{u}_t(x) + \Upsilon_t(x) \cdot \nabla_x \delta \tilde{u}_t(x) - \nabla_{x_0} f_t(X_t^0, x, \tilde{\mu}_t) \cdot \delta X_t^0 - \Big(\delta_\mu f_t(X_t^0, x, \mu_t), \delta \tilde{\mu}_t \Big) + j_t(x) \Big] \text{d}t
$$

\n
$$
+ \mathrm{d}_t \delta \tilde{m}_t(x), \quad x \in \mathbb{T}^d,
$$

\n
$$
\delta \tilde{\mu}_0 = \Delta \mu, \quad \delta \tilde{u}_T(x) = \nabla_{x_0} g(X_T^0, x, \mu_T) \cdot \delta X_T^0 + \Big(\delta_\mu g(X_T^0, x, \mu_T), \delta \tilde{\mu}_T \Big) + k_T(x), \quad x \in \mathbb{T}^d,
$$

\n(5.19)

with paths in the space $C^0([0,T], C^{-\nu+1}(\mathbb{T}^d) \times C^{\nu}(\mathbb{T}^d) \times C^{\nu-2}(\mathbb{T}^d))$ and with

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]}\left\{\|\delta\tilde{\mu}_{t}\|_{-\omega+1}^{2}+\|\delta\tilde{u}_{t}\|_{\omega}^{2}+\|\delta\tilde{m}_{t}\|_{\omega-2}^{2}\right\}\right]<+\infty.
$$
\n(5.20)

Basically, we would like to apply [\[14,](#page-86-1) Theorem 4.4.2] in order to guarantee that there is a unique solution $(\delta \tilde{\mu}, \delta \tilde{\mu}, \delta \tilde{m})$, adapted with respect to the filtration \mathbb{F}^0 , to (5.19) – (5.20) . Notice that if we can indeed apply [\[14](#page-86-1), Theorem 4.4.2], then *Bochner* measurability with values in any $C^{-u+1}(\mathbb{T}^d) \times C^u(\mathbb{T}^d) \times C^{u-2}(\mathbb{T}^d)$ is obvious because [\[14,](#page-86-1) Theorem 4.4.2] then implies that the solution is also in $C^{-\mathbf{u}''+1}(\mathbb{T}^d) \times C^{\mathbf{u}'}(\mathbb{T}^d) \times C^{\mathbf{u}'-2}(\mathbb{T}^d)$ for $\mathbf{r} < \mathbf{u}'' < \mathbf{u} < \mathbf{u}' < \mathbf{s}$.

Actually, the difficulty is that the current setting does not exactly fit the assumption of [\[14](#page-86-1), Theorem 4.4.2]. One first difficulty is that [\[14](#page-86-1), Theorem 4.4.2] is stated in the monotone framework. This is the same issue as the one mentioned in the proof of Theorem [5.1](#page-50-1) and the remedy is very similar. Here is an overview of it. Because of the lack of monotonicity, there is an additional term in [\[14,](#page-86-1) (4.44)] that writes (with the notations from [\[14\]](#page-86-1)) $\mathbb{E}[\sup_{t\in[0,T]}\|\tilde{\rho}_t-\tilde{\rho}_t'\|^2_{-(n+\alpha')}]$ (in our system of notations, $\tilde{\rho}_t$ corresponds to $\delta \tilde{\mu}_t$, $\tilde{\rho}'_t$ to another (forward) solution $\delta \tilde{\mu}'_t$ and $n + \alpha'$ to $u - 1$, for $u \in (\mathbb{r}, \mathbb{s})$). If T is small enough, then Cauchy-Schwarz inequality yields an additional factor \sqrt{T} in [\[14,](#page-86-1) (4.47)] so that [\[14](#page-86-1), (4.49)] remains small enough, then Cauchy-Schwarz inequality yields an additional factor \sqrt{T} in [14, (4.47)] true, with an additional factor T in front of $\sup_{t\in[0,T]}\|\tilde{z}_t-\tilde{z}_t'\|^2_{n+1+\alpha}$ (again with the notations from [\[14](#page-86-1)], \tilde{z}_t being here understood as $\delta \tilde{u}_t$). This makes it possible to apply the same contraction argument as in the proof of [\[14,](#page-86-1) Proposition 4.4.7], the role of ε (which is assumed to be small therein) being now played by \sqrt{T} .

Another difference is that the process δX^0 (which appears here both in the driver and in the boundary condition of the backward equation) is not bounded (in ω^0). This creates another difficulty since the perturbations \tilde{f}_t^0 and \tilde{g}_T^0 in [\[14](#page-86-1), (4.37)] are assumed to be bounded. In order to construct a solution, we thus apply [\[14,](#page-86-1) Theorem 4.4.2] to an approximation of [\(5.19\)](#page-56-0), in which $(\delta X_t^0)_{0 \le t \le T}$ is replaced by $(\varphi_R(\delta X_t^0))_{0 \le t \le T}$ for a real $R > 0$ and a function $\varphi_R : \mathbb{R}^d \to \mathbb{R}^d$ that is bounded by 2R and equal to the identity mapping (from \mathbb{R}^d into itself) on the d-dimensional ball of center 0 and radius R. This makes it possible to combine [\[14,](#page-86-1) Theorem 4.4.2] and [\[14](#page-86-1), Proposition 4.4.5]: The first statement shows that there is a solution for each R (and for a fixed initial condition), denoted $(\delta \tilde{\mu}^R, \delta \tilde{u}^R, \delta \tilde{m}^R)$, and the second one shows (by comparing the solutions for two different values of R) that the family of (hence constructed) solutions is Cauchy as R tends to ∞ . Here, the Cauchy property is understood in the same L^2 sense as in [\(5.20\)](#page-56-1). The limit of the Cauchy sequence is a solution of (5.19) . Uniqueness is slightly more challenging. We take one solution to (5.19) satisfying [\(5.20\)](#page-56-1). We then introduce the stopping time $\tau_R := \inf\{t \in [0,T] : |\delta X_t^0| + ||\delta \tilde{u}_t||_{\mathfrak{s}} \ge R\}$. Writing the system solved by $(\delta\tilde{\mu}_{t\wedge\tau_R}, \delta\tilde{\mu}_{t\wedge\tau_R}, \delta\tilde{\mu}_{t\wedge\tau_R})_{0\leq t\leq T}$ as a forward-backward system on $[0, T]$, with

$$
\delta\tilde{u}_{\tau_R}(x) = \left[\nabla_{x_0} g(X_T^0, x, \mu_T) \cdot \delta X_T^0 + \left(\delta_\mu g(X_T^0, x, \mu_T), \delta\tilde{\mu}_T\right) + k_T(x)\right] \mathbf{1}_{\{\tau_R = T\}} + \delta\tilde{u}_{\tau_R}(x) \mathbf{1}_{\{\tau_R < T\}}, \quad x \in \mathbb{T}^d,
$$

as terminal condition, we can compare $(\delta\tilde{\boldsymbol{\mu}}_{t\wedge\tau_R},\delta\tilde{\boldsymbol{u}}_{t\wedge\tau_R},\delta\tilde{\boldsymbol{v}}_{t\wedge\tau_R}^0)_{0\leq t\leq T}$ with $(\delta\tilde{\boldsymbol{\mu}}^R,\delta\tilde{\boldsymbol{u}}^R,\delta\tilde{\boldsymbol{v}}^{0,R})$ by means of [\[14,](#page-86-1) Proposition 4.4.5]. Thanks to the above decomposition, we observe that, on the event $\{\tau_R = T\}$, the terminal condition of the forward-backward system obtained by making the difference of the two solutions just writes $(\delta_{\mu}g(X_T^0,x,\mu_T), \delta\tilde{\mu}_T \delta \tilde{\mu}_T^R$) and is thus purely linear (equivalently, the intercept is zero). On the event $\{\tau_R < T\}$, the terminal condition writes $[\delta \tilde{u}_{\tau_R}(x) - \delta \tilde{u}_T^R(x)] \mathbf{1}_{\{\tau_R < T\}}$, which tends, by a standard uniform integrability argument, to 0 in L^2 as R tends to 0. By [\[14](#page-86-1), Proposition 4.4.5], this proves uniqueness. Also, this makes it possible to compare any two solutions satisfying [\(5.19\)](#page-56-0) and [\(5.20\)](#page-56-1) by means of [\[14,](#page-86-1) Proposition 4.4.5] again. All these claims hold true for $T \leq \mathfrak{C}$, where $\mathfrak{C} > 0$ only depends on d, κ , \mathfrak{K} , \mathfrak{L} , σ_0 and $(\mathfrak{r},\mathfrak{s})$. In particular, \mathfrak{C} is independent of the initial condition.

Next we define $(\delta \tilde{\mathbf{X}}^0, \delta \tilde{\mathbf{Y}}^0, \delta \tilde{\mathbf{Z}}^0)$ as the strong solution to the following FBSDE system (on $[0,T]$)

$$
d\delta \tilde{X}_t^0 = \left[-p_t \delta \tilde{X}_t^0 - q_t \delta \tilde{Z}_t^0 + a_t \right] dt,
$$

\n
$$
d\delta \tilde{Y}_t^0 = \left[l_t \cdot \delta \tilde{X}_t^0 + o_t \delta \tilde{Z}_t^0 - \left(\delta_\mu f_t^0(X_t^0, \mu_t), \delta \tilde{\mu}_t \right) + b_t \right] dt + \sigma_0 \delta \tilde{Z}_t^0 \cdot dB_t^0, \quad t \in [0, T],
$$

\n
$$
\delta \tilde{X}_0^0 = \Delta x_0, \quad \delta \tilde{Y}_T^0 = \nabla_{x_0} g^0(X_T^0, \mu_T) \cdot \delta \tilde{X}_T^0 + \left(\delta_\mu g^0(X_T^0, \mu_T), \delta \tilde{\mu}_T \right) + c_T.
$$
\n(5.21)

Here, solutions are required to satisfy

$$
\mathbb{E}^{0}\left[\sup_{0\leq t\leq T}\left(|\delta\tilde{X}_{t}^{0}|^{2}+|\delta\tilde{Y}_{t}^{0}|^{2}\right)+\int_{0}^{T}|\delta\tilde{Z}_{t}^{0}|^{2}\mathrm{d}t\right]<\infty.\tag{5.22}
$$

By [\[26](#page-86-0)], existence and uniqueness of a solution hold true for $T \leq \mathfrak{C}$, for a possibly new value of the constant \mathfrak{C} . Next, we implicitly require T to be less than \mathfrak{C} .

Setting $\delta \mathfrak{T}(\delta X^0) := \delta \tilde{X}^0$, this defines a map $\delta \mathfrak{T} : \mathscr{S}^2(\mathbb{R}^d) \to \mathscr{S}^2(\mathbb{R}^d)$.

Second Step. In this step, we show that, for $T \leq \mathfrak{C}$ (again for a possibly new value of \mathfrak{C} , depending on d, κ , \mathfrak{K} , \mathfrak{L} and (\mathbb{r}, \mathbb{s})), $\delta \mathfrak{T}$ is a contraction mapping from $\mathscr{S}^2(\mathbb{R}^d)$ to $\mathscr{S}^2(\mathbb{R}^d)$.

For two inputs $\delta X^{0,1}$ and $\delta X^{0,2}$ with $\delta X_0^{0,1} = \delta X_0^{0,2} = \Delta x_0$ as initial conditions, we denote by $(\delta \tilde{\mu}^1, \delta \tilde{u}^1, \delta \tilde{m}^1)$ and $(\delta \tilde{\mu}^2, \delta \tilde{u}^2, \delta \tilde{m}^2)$ the corresponding solutions to [\(5.19\)](#page-56-0), and next by $(\delta \tilde{X}^{0,1}, \delta \tilde{Y}^{0,1}, \delta \tilde{Z}^{0,1})$ and $(\delta \tilde{X}^{0,2}, \delta \tilde{Y}^{0,2}, \delta \tilde{Z}^{0,2})$ the corresponding solutions to [\(5.21\)](#page-57-0). Using [\[14,](#page-86-1) Proposition 4.4.5] (which is licit thanks to the analysis performed in the first step, and with the following notations therein: the primed solution is null, $\vartheta = 1$, $\tilde{m}_t = \mu_t$, $\tilde{z}_t(x) = \delta \tilde{u}_t^1(x)$, $\tilde{z}'_t(x) = \delta \tilde{v}_t^2(x)$ $\delta \tilde{u}_t^2(x), \tilde{f}_t^0(x) = j_t(x) - \nabla_{x_0} f_t(X_t^0, x, \tilde{\mu}_t) \cdot \delta X_t^{0,1}, \tilde{f}_t^{0,\tilde{\ell}}(x) = j_t(x) - \nabla_{x_0} f_t(X_t^0, x, \tilde{\mu}_t) \cdot \delta X_t^{0,2}, \tilde{V}_t(x) = \tilde{V}_t^{\ell}(x) = \Upsilon_t(x),$ $\tilde{b}_t^0(x) = \tilde{b}_t^{0,1}(x) = d_t(x)$, we get (thanks to Remark [5.3,](#page-55-4) the term inside the expectation symbol in the left-hand side below is a random variable):

$$
\mathbb{E}^{0}\bigg[\sup_{t\in[0,T]} \|\delta\tilde{\mu}^1_t-\delta\tilde{\mu}^2_t\|^2_{-\mathrm{r}+1}\bigg] \leq C\mathbb{E}^{0}\bigg[\sup_{t\in[0,T]} |\delta X_t^{0,1}-\delta X_t^{0,2}|^2\bigg],
$$

where C only depends on d, κ , $\hat{\mathcal{R}}$, $\hat{\mathcal{L}}$, and (Γ , $\hat{\mathcal{S}}$). By standard short time stability estimates for FBSDEs, see [\[26,](#page-86-0) Theorem 1.3], we can derive (for $T \leq \mathfrak{C}$, allowing the value of $\mathfrak C$ to vary from line to line as long as it just depends on the same parameters as before)

$$
\mathbb{E}^{0}\left[\int_{0}^{T} |\delta \tilde{Z}_{t}^{0,1} - \delta \tilde{Z}_{t}^{0,2}|^{2} dt\right]
$$
\n
$$
\leq C \mathbb{E}^{0}\left[\left(\int_{0}^{T} \left(\delta_{\mu} f_{t}^{0}(X_{t}^{0}, \mu_{t}), \delta \tilde{\mu}_{t}^{1} - \delta \tilde{\mu}_{t}^{2}\right) dt\right)^{2} + \left(\delta_{\mu} g^{0}(X_{T}^{0}, \mu_{T}), \delta \tilde{\mu}_{T}^{1} - \delta \tilde{\mu}_{T}^{2}\right)^{2}\right]
$$
\n
$$
\leq C \mathbb{E}^{0}\left[\sup_{t \in [0,T]} \|\delta \tilde{\mu}_{t}^{1} - \delta \tilde{\mu}_{t}^{2}\|_{-\kappa+1}^{2}\right] \leq C \mathbb{E}^{0}\left[\sup_{t \in [0,T]} |X_{t}^{0,1} - X_{t}^{0,2}|^{2}\right].
$$
\n(5.23)

Inserting the above estimate in the forward equation of [\(5.21\)](#page-57-0), we deduce that, for $T \leq \mathfrak{C}$,

$$
\mathbb{E}^{0}\biggl[\sup_{t\in[0,T]}|\delta\tilde{X}^{0,1}_t-\delta\tilde{X}^{0,2}_t|^2\biggr]\leq \frac{1}{2}\mathbb{E}^{0}\biggl[\sup_{t\in[0,T]}|\delta X^{0,1}_t-\delta X^{0,2}_t|^2\biggr],
$$

which implies $\delta \mathfrak{T}$ is a contraction mapping from $\mathscr{S}^2(\mathbb{R}^d)$ to $\mathscr{S}^2(\mathbb{R}^d)$. Therefore, there exists a unique fixed point $\delta \mathbf{X}^0 \in \mathbb{R}^d$ $\mathscr{S}^2(\mathbb{R}^d)$ with $\delta X_0^0 = \triangle x_0$.

Then, the coupled system comprising both the (finite-dimensional) FBSDE and the FBSPDE in [\(5.15\)](#page-54-0) admits a unique solution, which is adapted with respect to the filtration \mathbb{F}^0 . The solution to the FBSPDE system is denoted $(\delta\mu,\delta u,\delta v^0)$. It satisfies (a) and (b) in the statement. The solution to the finite-dimensional FBSDE system in [\(5.15\)](#page-54-0) admits a unique strong solution $(\delta X^0, \delta Y^0, \delta Z^0)$. It satisfies (c) in the statement.

Third Step. We now estimate the 6-tuple $(\delta\mu, \delta u, \delta m, \delta X^0, \delta Y^0, \delta Z^0)$, solution to the forward-backward system [\(5.15\)](#page-54-0). The proof is similar to that one of [\[14,](#page-86-1) Corollary 4.4.6]. The point is to compare the solution $(\delta\mu, \delta u, \delta m)$ of the FBSPDE system in [\(5.15\)](#page-54-0) with the trivial system driven by null coefficients and thus having a null solution. Comparison is obtained by means of [\[14](#page-86-1), Proposition 4.4.5]. One of the key point in the proof is to fix one instant $t \in [0, T]$ and then to apply [\[14](#page-86-1), Proposition 4.4.5] to the restriction of the triple $(\delta\mu, \delta u, \delta m)$ to the interval $[t, T]$, seen as a solution of the FBSPDE system in [\(5.15\)](#page-54-0) on [t, T], with $\delta\mu_t$ as initial solution and under the conditional probability distribution of \mathbb{P}^0 given the σ -field \mathcal{F}_t^0 .

We get, with probability 1 under \mathbb{P}^0 ,

$$
\|\delta u_t\|_{\text{s}}^2\leq C\|\delta\mu_t\|_{-\text{r}+1}^2+C\mathbb{E}^0\Big[\sup_{r\in[t,T]}\|d_r\|_{-\text{r}+2}^2+\sup_{r\in[t,T]}\|j_r\|_{\text{s}-1}^2+\sup_{r\in[t,T]}\left|\delta X_r^0\right|^2+\|k_T\|_{\text{s}}^2\,|\,\mathcal{F}_t^0\Big],
$$

where C only depends on d, κ , $\hat{\mathcal{R}}$, $\hat{\mathcal{L}}$ and $(\mathbb{r}, \mathfrak{s})$ (recalling that $T \leq \mathfrak{C}$) and is allowed to vary from line to line. Measurability of the arguments in the above right-hand side follows from the third item in Remark [5.3.](#page-55-4)

Taking power $p \in (1, 8]$ and then expectation (under \mathbb{P}^0) on both sides, we obtain by means of Doob's inequality,

$$
\label{eq:2.10} \begin{split} &\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|\delta u_t\|^{2p}_\mathbb{S}\Big]\\ &\leq C\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|\delta\mu_t\|^{2p}_{-r+1}+\sup_{t\in[0,T]}\|d_t\|^{2p}_{-r+2}+\sup_{t\in[0,T]}\|j_t\|^{2p}_{\mathbb{S}-1}+\sup_{t\in[0,T]}\|\delta X^0_t|^{2p}+\|k_T\|^{2p}_\mathbb{S}\Big]. \end{split}
$$

Now, by an obvious adaptation of Lemma [6.3](#page-85-0) (see for instance [\[14,](#page-86-1) Lemma 3.3.1]), we have, \mathbb{P}^0 -almost surely,

$$
\sup_{t\in[0,T]}\|\delta\mu_t\|_{-\mathbb{r}+1}\leq C\|\Delta\mu\|_{-\mathbb{r}+1}+CT\Big[\sup_{t\in[0,T]}\|\delta u_t\|_{\mathbb{S}-1}+\sup_{t\in[0,T]}\|d_t\|_{-\mathbb{r}+2}\Big].
$$

And, then, assuming $CT \leq 1/2$,

$$
\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|\delta\mu_{t}\|^{2p}_{-\mathbb{r}+1}\Big]\leq C\Big(\|\triangle\mu\|^{2p}_{-\mathbb{r}+1}+\mathbb{E}^{0}\Big[\sup_{t\in[0,T]}\|d_{t}\|^{2p}_{-\mathbb{r}+2}+\sup_{t\in[0,T]}\|j_{t}\|^{2p}_{\mathbb{s}-1}+\sup_{t\in[0,T]}\|\delta X_{t}^{0}|^{2p}+\|k_{T}\|^{2p}_{\mathbb{s}}\Big]\Big).
$$

And then, by [\[26,](#page-86-0) Theorem A.5],

$$
\mathbb{E}^{0}\left[\left(\int_{0}^{T}|\delta Z_{t}^{0}|^{2}dt\right)^{p}\right]
$$
\n
$$
\leq C\left(|\Delta x_{0}|^{2p} + \mathbb{E}^{0}\left[\sup_{t\in[0,T]}||\delta\mu_{t}||^{2p}_{-\mathbb{r}+1} + \left(\int_{0}^{T}(|a_{t}|+|b_{t}|)dt\right)^{2p} + |c_{T}|^{2p}\right]\right)
$$
\n
$$
\leq C\left(|\Delta x_{0}|^{2p} + ||\Delta\mu||^{2p}_{-\mathbb{r}+1} + ||\Delta\mu||^{2p}_{-\mathbb{r}+1} + ||\Delta\mu||^{2p}_{-\mathbb{r}+2} + ||j_{t}||^{2p}_{s-1} + |\delta X_{t}^{0}|^{2p}\right) + \left(\int_{0}^{T}(|a_{t}|+|b_{t}|)dt\right)^{2p} + |c_{T}|^{2p} + ||k_{T}||_{s}^{2p}\right).
$$

Inserting the above bound into the equation for δX^0 , we deduce that, for $CT \leq 1/2$ (and $T \leq 1$),

$$
\mathbb{E}^{0}\left[\sup_{0\leq t\leq T}|\delta X_{t}^{0}|^{2p}\right]
$$
\n
$$
\leq C\left(|\Delta x_{0}|^{2p} + ||\Delta \mu||_{-r+1}^{2p} + ||j_{t}||_{s-1}^{2p}\right) + \left(\int_{0}^{T}\left(|a_{t}| + |b_{t}|\right)dt\right)^{2p} + |c_{T}|^{2p} + ||k_{T}||_{s}^{2p}\right)
$$
\n
$$
= CM_{p},
$$

and the proof is easily completed by recombining all the intermediary steps.

$$
\Box
$$

5.3. First order derivatives of the master fields

For simplicity, the results in the last two subsections are stated with $t = 0$ as initial time. However, they remain true for any initial time $t \in [0, T]$ (provided T is small enough). In particular, for any initial condition $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times$ $\mathcal{P}(\mathbb{T}^d)$, the system [\(5.1\)](#page-50-3)–[\(5.2\)](#page-50-2) has a unique solution, denoted $(X^{0,t,x_0,\mu}, Y^{0,t,x_0,\mu}, Z^{0,t,x_0,\mu}, \mu^{t,x_0,\mu}, u^{t,x_0,\mu}, m^{t,x_0,\mu})$, on $[t, T]$ with T given in Theorem [5.1,](#page-50-1) which makes it possible to let

$$
\mathcal{U}(t, x_0, x, \mu) := u^{t, x_0, \mu}(t, x), \quad x \in \mathbb{T}^d,
$$
\n(5.24)

and

$$
\mathcal{U}^{0}(t,x_{0},\mu) := Y_{t}^{0,t,x_{0},\mu}.
$$
\n(5.25)

Elaborating on the proof of $[14, \text{Lemma 5.1.1}]$ and using the uniqueness result of (5.1) – (5.2) together with the stability property [\(5.5\)](#page-51-4), we have, with probability 1 under \mathbb{P}^0 ,

$$
u^{t,x_0,\mu}(r,x) = \mathcal{U}(r, X_r^{0,t,x_0,\mu}, x, \mu_r^{t,x_0,\mu}) \quad \text{and} \quad Y_r^{0,t,x_0,\mu} = \mathcal{U}^0(r, X_r^{0,t,x_0,\mu}, \mu_r^{t,x_0,\mu}), \quad \text{for any } r \in [t, T]. \tag{5.26}
$$

Thanks to [\(5.3\)](#page-51-0) and [\(5.4\)](#page-51-1), \mathcal{U}^0 and \mathcal{U} are bounded. In fact, by [\(5.5\)](#page-51-4) again, we also have

$$
\mathcal{U}: [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \ni (t, x_0, \mu) \mapsto \mathcal{U}(t, x_0, \cdot, \mu) \in \mathcal{C}^s(\mathbb{T}^d) \text{ is uniformly Lipschitz continuous in } x_0, \mu,
$$

$$
\mathcal{U}^0: [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \ni (t, x_0, \mu) \mapsto \mathcal{U}^0(t, x_0, \mu) \in \mathbb{R} \text{ is uniformly Lipschitz continuous in } x_0, \mu,
$$
 (5.27)

where the Lipschitz continuity in μ is measured by the \mathbb{W}_1 metric. Continuity in time is ensured by the following lemma:

Lemma 5.4. *Under Assumption* (\hat{A}) *, there exists a constant* \mathfrak{C} *only depending on d,* \mathfrak{L} *,* κ *,* σ_0 *and s, such that the following holds true for* $T \leq \mathfrak{C}$ *. For any* $\mathfrak{r} \in (\lfloor \mathfrak{s} \rfloor, \mathfrak{s})$ *, we can find a constant* C*, only depending on* d*,* \mathfrak{L} *,* κ *,* σ_0 *,* \mathfrak{r} *and* \mathfrak{s} *, such that, for any* $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ *and* $r \in [t, T]$ *,*

$$
\|\mathcal{U}(r, x_0, \cdot, \mu) - \mathcal{U}(t, x_0, \cdot, \mu)\|_{\mathbb{F}} \le C(r - t)^{(s - r)/2},\tag{5.28}
$$

Major Minor MFGs 61

and

$$
|\mathcal{U}^{0}(r,x_{0},\mu)-\mathcal{U}^{0}(t,x_{0},\mu)| \leq C(r-t)^{1/2}.
$$
\n(5.29)

Proof. *First Step.* We first show (5.28) . Following [\[14,](#page-86-1) (4.12)], note that, with probability 1 under \mathbb{P}^0 ,

$$
u_t(\cdot) = \mathbb{E}^0\bigg[P_{r-t}u_r(\cdot) - \int_t^r P_{s-t}\Big(\hat{H}(\cdot,\nabla_x u_s(\cdot)) - f_s(X_s^0,\cdot,\mu_s)\Big)ds\bigg], \quad \text{for any } r \in [t,T],
$$

where $(P_t)_{t\in[0,T]}$ stands here for the standard heat kernel on the torus \mathbb{T}^d . The expectation in the right-hand side should be understood as follows: here and below, we write $\mathbb{E}^0[h(\cdot)]$ for the mapping $x \in \mathbb{T}^d \mapsto \mathbb{E}^0[h(x)]$ when $h: \Omega^0 \times \mathbb{T}^d \to \mathbb{R}$ is a (measurable) random field such that $\mathbb{E}^0[[h(x)]]$ is finite for any $x \in \mathbb{T}^d$. Of course, Lebesgue's theorem makes it possible to address the regularity of $\mathbb{E}^0[h(\cdot)]$ when h itself is regular in \mathbb{T}^d . Also, we recall from [\(5.4\)](#page-51-1) that there exists a constant C, only depending on the parameters in (\hat{A}) , such that $\mathbb{P}^0(\{\sup_{t\leq s\leq T} ||\nabla_x u_s||_{s-1} \leq C\}) = 1$. And then (with id denoting the identity mapping),

$$
u_t(\cdot) - \mathbb{E}^0[u_r(\cdot)] = \mathbb{E}^0\bigg[\Big(P_{r-t} - \mathrm{id}\Big)u_r(\cdot) - \int_t^r P_{s-t}\Big(\hat{H}(\cdot,\nabla_x u_s(\cdot)) - f_s(X_s^0,\cdot,\mu_s)\Big)\mathrm{d} s\bigg],
$$

which further implies

$$
||u_t - \mathbb{E}^0[u_r]||_{\mathbf{r}} \le \mathbb{E}^0\bigg[||(P_{r-t} - id)u_r||_{\mathbf{r}} + C \int_t^r (s-t)^{-1/2} ||\hat{H}(\cdot, \nabla_x u_s(\cdot)) - f_s(X_s^0, \cdot, \mu_s)||_{\mathbf{r}-1}ds\bigg]
$$

$$
\le \mathbb{E}^0\bigg[||(P_{r-t} - id)u_r||_{\mathbf{r}}\bigg] + C(r-t)^{1/2},
$$

with the constant C only depends on the parameters in Assumption (\hat{A}) . It is standard to have

$$
\mathbb{E}^{0}\Big[\big\|(P_{r-t}-\mathrm{id})u_r\big\|_{\mathbb{F}}\Big]\leq C(r-t)^{(s-r)/2}\mathbb{E}^{0}\big[\|u_r\|_{\mathbb{S}}\big]\leq C(r-t)^{(s-r)/2}.
$$

Therefore, allowing the constant C to vary from line to line,

$$
\left\|u_t - \mathbb{E}^0[u_r]\right\|_{\mathbb{P}} \leq C(r-t)^{(\mathbb{s}-\mathbb{P})/2}.
$$

By (5.24) , we can derive

$$
||u_t - \mathcal{U}(r, x_0, \cdot, \mu)||_r \le ||u_t - \mathbb{E}^0[u_r]||_r + ||\mathbb{E}^0[u_r] - \mathcal{U}(r, x_0, \cdot, \mu)||_r
$$

\n
$$
\le C(r - t)^{(s - r)/2} + ||\mathbb{E}^0[\mathcal{U}(r, X_r^{0, t, x_0, \mu}, \cdot, \mu_r^{t, x_0, \mu}) - \mathcal{U}(r, x_0, \cdot, \mu)]||_r
$$

\n
$$
\le C\Big\{(r - t)^{(s - r)/2} + \mathbb{E}^0[\mathbb{W}_1(\mu_r^{t, x_0, \mu}, \mu) + |X_r^{0, t, x_0, \mu} - x_0|]\Big\}
$$

\n
$$
\le C(r - t)^{(s - r)/2}.
$$

Second Step. We now show [\(5.29\)](#page-60-0). We first have

$$
\begin{split}\n& \left| \mathcal{U}^{0}(r,x_{0},\mu) - \mathcal{U}^{0}(t,x_{0},\mu) \right| \\
&\leq \left| \mathcal{U}^{0}(r,x_{0},\mu) - \mathbb{E}^{0}\left[\mathcal{U}^{0}(r,X_{r}^{0,t,x_{0},\mu},\mu_{r}^{t,x_{0},\mu}) \right] \right| + \left| \mathbb{E}^{0}\left[\mathcal{U}^{0}(r,X_{r}^{0,t,x_{0},\mu},\mu_{r}^{t,x_{0},\mu}) - \mathcal{U}^{0}(t,x_{0},\mu) \right] \right| \\
&\leq C \Big(\mathbb{E}^{0}\big[\mathbb{W}_{1}(\mu_{r}^{t,x_{0},\mu},\mu) + |X_{r}^{0,t,x_{0},\mu} - x_{0}| \big] + \left| \mathbb{E}^{0}\big[Y_{r}^{0,t,x_{0},\mu} - Y_{t}^{0,t,x_{0},\mu} \big] \right| \Big) \\
&\leq C \Big(|r-t|^{1/2} + \left| \mathbb{E}^{0}\big[Y_{r}^{0,t,x_{0},\mu} - Y_{t}^{0,t,x_{0},\mu} \big] \right| \Big). \end{split} \tag{5.30}
$$

By means of [\(5.3\)](#page-51-0), we have a bound for $\mathbb{E}^0 \int_t^r |Z_s^{0,t,x_0,\mu}|^2 ds$, from which we obtain

$$
\mathbb{E}^{0}[|Y_r^{0,t,x_0,\mu} - Y_t^{0,t,x_0,\mu}|] \le C|r - t|^{1/2}.
$$
\n(5.31)

Combining (5.30) and (5.31) , we can derive

$$
|\mathcal{U}^{0}(r,x_{0},\mu)-\mathcal{U}^{0}(t,x_{0},\mu)|\leq C(r-t)^{1/2},
$$

which completes the proof.

We now would like to establish the continuous differentiability of U and U_0 in x_0 and μ . For any initial condition $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and $(\Delta x_0, \Delta \mu) \in \mathbb{R}^d \times \mathcal{C}^{-\mathbb{T}+1}(\mathbb{T}^d)$, for some $\mathbb{T} \in (\lfloor s \rfloor, s)$, we consider the solution $(\delta X^0, \delta Y^0, \delta Z^0, \delta \mu, \delta u, \delta m)$ (its dependence on the initial condition is omitted for simplicity) to the linearized forwardbackward system, set on $[t, T]$,

$$
d\delta X_r^0 = \left[-\nabla_{px_0}^2 H^0(X_r^0, Z_r^0) \delta X_r^0 - \nabla_{pp}^2 H^0(X_r^0, Z_r^0) \delta Z_r^0 \right] dr,\nd\delta Y_r^0 = \left[\nabla_{x_0} \hat{L}^0(X_r^0, Z_r^0) \cdot \delta X_r^0 + \nabla_p \hat{L}^0(X_r^0, Z_r^0) \cdot \delta Z_r^0 \right. \n- \nabla_{x_0} f_r^0(X_r^0, \mu_r) \cdot \delta X_r^0 - \left(\delta_\mu f_r^0(X_r^0, \mu_r), \delta \mu_r \right) \right] dr + \sigma_0 \delta Z_r^0 \cdot dB_r^0,\n\delta X_t^0 = \triangle x_0, \quad \delta Y_T^0 = \nabla_{x_0} g^0(X_T^0, \mu_T) \cdot \delta X_T^0 + \left(\delta_\mu g^0(X_T^0, \mu_T), \delta \mu_T \right);\n\partial_r \delta \mu_r - \frac{1}{2} \Delta_x \delta \mu_r - \text{div}_x (\nabla_p \hat{H}(x, \nabla_x u_r(x)) \delta \mu_r + \mu_r \nabla_{pp}^2 \hat{H}(x, \nabla_x u_r(x)) \nabla_x \delta u_r(x)) = 0,\nd_r \delta u_r(x) = \left[-\frac{1}{2} \Delta_x \delta u_r(x) + \nabla_p \hat{H}(x, \nabla_x u_r(x)) \cdot \nabla_x \delta u_r(x) - \nabla_{x_0} f_r(X_r^0, x, \mu_r) \cdot \delta X_r^0 \right. \n- \left(\delta_\mu f_r(X_r^0, x, \mu_r), \delta \mu_r \right) \right] dr + d_r \delta m_r(x), \quad x \in \mathbb{T}^d,
$$

\n $\delta \mu_t = \triangle \mu, \quad \delta u_T(x) = \nabla_{x_0} g(X_T^0, x, \mu_T) \cdot \delta X_T^0 + \left(\delta_\mu g(X_T^0, x, \mu_T), \delta \mu_T \right) \quad \text{on } \mathbb{T}^d,$

where $(X^0, Y^0, Z^0, \mu, u, m)$ is the unique solution to the system [\(5.1\)](#page-50-3)–[\(5.2\)](#page-50-2) corresponding to the initial condition (t, x_0, μ) . Applying Theorem [5.2](#page-55-0) to [\(5.32\)](#page-61-0), we have the following theorem.

Theorem 5.5. Let $\mathbb{r} \in (\vert \mathbf{s} \vert, \mathbf{s})$. Under Assumption ($\hat{\mathbf{A}}$), there exist a constant c only depending on d, κ , \mathfrak{L} and \mathbf{s} and *a constant* $\mathfrak C$ *only depending on* d, κ , $\mathfrak L$, $\zeta(c)$, σ_0 *and* ($\mathfrak r$, $\mathfrak s$) *such that, for* $T \leq \mathfrak C$, the forward-backward system [\(5.15\)](#page-54-0) $admits a unique solution $(\delta\mu, \delta u, \delta m, \delta X^0, \delta Y^0, \delta Z^0)$, adapted with respect to the filtration \mathbb{F}^0 and with values in$ $\mathcal{C}^{-\mathbb{r}+1}(\mathbb{T}^d)\times\mathcal{C}^{\mathfrak{s}}(\mathbb{T}^d)\times\mathbb{C}^{s-2}(\mathbb{T}^d)\times\mathbb{R}^d\times\mathbb{R}\times\mathbb{R}^d$, and satisfying items (a), (b) and (c) in the statement of Theorem [5.2.](#page-55-0)

Moreover, there exists a constant C *only depending on the parameters in Assumption* (Â) *and on* **r** *such that, for any* $p \in [1, 8]$,

$$
\mathbb{E}^{0}\left[\sup_{t\in[0,T]} \left(\|\delta\mu_t\|_{-\mathbf{r}+1}^{2p} + \|\delta u_t\|_{\mathbf{s}}^{2p} + \|\delta m_t\|_{\mathbf{s}-2}^{2p}\right) + \sup_{t\in[0,T]} \left(|\delta X_t^0|^{2p} + |\delta Y_t^0|^{2p}\right) + \left(\int_0^T |\delta Z_t^0|^2 dt\right)^p\right] \tag{5.33}
$$

$$
\leq C\left(\|\delta\mu\|_{-\mathbf{r}+1}^{2p} + |\Delta x_0|^{2p}\right).
$$

Proof. The only subtlety in the application of Theorem [5.2](#page-55-0) lies in the verification of (v) in Assumption (C). In fact, by Theorem [5.1,](#page-50-1) we have (for $T \leq \mathfrak{C}$) a bound for $||u_t||_{s}$. And then, we can insert this bound in the function ζ in Assumption (\hat{A} 4) The rest of the proof does not raise any difficulty (Â4). The rest of the proof does not raise any difficulty.

Given $(t, x_0, y, \mu) \in [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d), i \in \{1, \cdots, d\}$ and a d -tuple $l \in \{0, \cdots, \lfloor s \rfloor - 1\}^d$ with $|l| := \sum_{j=1}^d l_j \leq$ \lfloor s \rfloor − 1, we first denote by $(\delta\mu^i, \delta u^i, \delta m^i, \delta X^{0,i}, \delta Y^{0,i}, \delta Z^{0,i})$ the solution to the system [\(5.32\)](#page-61-0) on [t, T] with $\triangle \mu = 0$ and $\Delta x^0 = e_i$, and then denote by $(\delta \mu^{l,y}, \delta u^{l,y}, \delta m^{l,y}, \delta X^{0,l,y}, \delta Y^{0,l,y}, \delta Z^{0,l,y})$ the solution to the system [\(5.32\)](#page-61-0) on $[t, T]$ with $\triangle \mu = (-1)^{|l|} \nabla^{l} \delta_y \in C^{-(r-1)}(\mathbb{T}^d)$ (δ_y denoting the Dirac mass at point y) for any $r \in (\lfloor s \rfloor, s)$, and $\triangle x^0 = 0$. Define

$$
K_i^{x_0}(t, x_0, x, \mu) := \delta u_t^i(x), \ K_i^{0, x_0}(t, x_0, \mu) := \delta Y_t^{0, i},
$$

\n
$$
K_l^{\mu}(t, x_0, x, \mu, y) := \delta u_t^{l, y}(x), \ K_l^{0, \mu}(t, x_0, \mu, y) := \delta Y_t^{0, l, y}.
$$
\n(5.34)

Then, choosing for instance $\mathbf{r} = (\lfloor s \rfloor + s)/2$, we can apply Theorem [5.5](#page-61-1) to obtain (for $T \leq \mathfrak{C}$, for some $\mathfrak{C} > 0$ only depending on the parameters d, κ , \mathfrak{L} , ζ and σ in Assumption (\hat{A}))

$$
||K_i^{x_0}(t, x_0, \cdot, \mu)||_{\mathfrak{s}} + ||K_l^{\mu}(t, x_0, \cdot, \mu, y)||_{\mathfrak{s}} + |K_i^{0, x_0}(t, x_0, \mu)| + |K_l^{0, \mu}(t, x_0, \mu, y)| \leq C,
$$
\n(5.35)

 \Box

Major Minor MFGs 63

for a constant C only depending on the parameters in Assumption (\hat{A}). In what follows, we just write $K^{x_0}(t, x_0, \cdot, \mu)$ and $K^{0,x_0}(t, x_0, \mu)$ for the d-tuples $(K_i^{x_0}(t, x_0, \cdot, \mu))_{i=1,\dots,d}$ and $(K_i^{0,x_0}(t, x_0, \mu))_{i=1,\dots,d}$.

The following statement is the analogue of $[14, \text{Lemma } 5.2.2]$:

Lemma 5.6. *Given* $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and $p \in [1, 8]$, we have under the same generic condition $T \leq \mathfrak{C}$ as *before, for any* d -tuple $l \in \{0, \dots, \lfloor \mathfrak{s} \rfloor - 1\}^d$ with $|l| := \sum_{i=1}^d l_i \leq \lfloor \mathfrak{s} \rfloor - 1$ and any $y \in \mathbb{T}^d$,

$$
\lim_{h \in \mathbb{R}^d, h \to 0} \mathbb{E}^0 \Big[\sup_{r \in [t,T]} \left\| \delta u_r^{l, y+h} - \delta u_r^{l, y} \right\|_{s}^{2p} + \sup_{r \in [t,T]} \left\| \delta \mu_r^{l, y+h} - \delta \mu_r^{l, y} \right\|_{-s+1}^{2p} + \sup_{r \in [t,T]} \left| \delta Y_r^{0, l, y+h} - \delta Y_r^{0, l, y} \right|^{2p} \Big] = 0.
$$
\n(5.36)

Moreover, for any $l \in \{0, \dots, \lfloor s \rfloor - 2\}^d$ *with* $|l| \leq \lfloor s \rfloor - 2$ *, and any* $i \in \{1, \dots, d\}$ *,*

$$
\lim_{h \in \mathbb{R}, h \to 0} \mathbb{E}^{0} \Big[\sup_{r \in [t,T]} \Big\| \frac{1}{h} (\delta u_r^{l, y + he_i} - \delta u_r^{l, y}) - \delta u_r^{l+e_i, y} \Big\|_{\mathfrak{s}}^{2p} + \sup_{r \in [t,T]} \Big\| \frac{1}{h} (\delta \mu_r^{l, y + he_i} - \delta \mu_r^{l, y}) - \delta \mu_r^{l+e_i, y} \Big\|_{-\mathfrak{s}+1}^{2p} \n+ \sup_{r \in [t,T]} \Big| \frac{1}{h} (\delta Y_r^{0, l, y + he_i} - \delta Y_r^{0, l, y}) - \delta Y_r^{0, l+e_i, y} \Big\|_{\mathfrak{s}}^{2p} \Big] = 0.
$$

In particular, the function $(x, y) \in \mathbb{T}^d \times \mathbb{T}^d \mapsto K_0^{\mu}(t, x_0, x, \mu, y)$ and $y \in \mathbb{T}^d \mapsto K_0^{0,\mu}(t, x_0, \mu, y)$ are $(\lfloor s \rfloor - 1)$ -times continuously differentiable with respect to y, and for any $l \in \{0, \dots, \lfloor s \rfloor - 1\}$ with $|l| \leq \lfloor s \rfloor - 1$, the derivative $y \in \mathbb{T}^d \mapsto \nabla_y^l K_0^\mu(t, x_0, \cdot, \mu, y) \in C^s(\mathbb{T}^d)$ is continuous. It holds, for any $(x, y) \in \math$

$$
\nabla_y^l K_0^{\mu}(t, x_0, x, \mu, y) = K_l^{\mu}(t, x_0, x, \mu, y) \quad \text{and} \quad \nabla_y^l K_0^{0, \mu}(t, x_0, \mu, y) = K_l^{0, \mu}(t, x_0, \mu, y).
$$

Moreover, there exists a constant C*, only depending on the parameters in Assumption* (Â)*, such that*

$$
\sup_{(t,x_0,\mu,y)\in[0,T]\times\mathbb{R}^d\times\mathcal{P}(\mathbb{T}^d)\times\mathbb{T}^d}\left\|K_l^\mu(t,x_0,\cdot,\mu,y)\right\|_{\text{s}}\leq C,
$$

and

$$
\sup_{(t,x_0,\mu,y)\in[0,T]\times\mathbb{R}^d\times\mathcal{P}(\mathbb{T}^d)\times\mathbb{T}^d} \left|K_l^{0,\mu}(t,x_0,\mu,y)\right| \leq C.
$$

Proof. By Theorem [5.5,](#page-61-1) for any fixed $r \in (\lfloor s \rfloor, s)$ and any $p \in [1, 8]$, we can find a constant C_p such that, for all $(t, x, \mu, y) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$ and $l \in \{1, \cdots, \lfloor s \rfloor - 1\}$ with $|l| \leq \lfloor s \rfloor - 1$,

$$
\mathbb{E}^{0}\Big[\sup_{r\in[t,T]}\big\|\delta u^{l,y}_r\big\|^{2p}_{\text{s}}+\sup_{r\in[t,T]}\big\|\delta\mu^{l,y}_r\big\|^{2p}_{-\textbf{r}+1}+\sup_{r\in[t,T]}\big|\delta Y^{0,l,y}_r\big|^{2p}\Big]\leq C_p.
$$

In particular,

$$
||K_l^{\mu}(t, x_0, \cdot, \mu, y)||_{\epsilon} + |K_l^{0,\mu}(t, x_0, \mu, y)| \leq C.
$$

Notice that

$$
\lim_{h\in\mathbb{R}^d,h\to 0}\left\|\nabla_y^l\delta_{y+h}-\nabla_y^l\delta_{y}\right\|_{-\mathbb{r}+1}=0.
$$

Therefore, Theorem [5.5](#page-61-1) gives [\(5.36\)](#page-62-0). This yields

$$
\lim_{h \in \mathbb{R}^d, h \to 0} \left(\left\| K_l^{\mu}(t, x_0, \cdot, \mu, y + h) - K_l^{\mu}(t, x_0, \cdot, \mu, y) \right\|_{\mathfrak{s}} + \left| K_l^{0,\mu}(t, x_0, \mu, y + h) - K_l^{0,\mu}(t, x_0, \mu, y) \right| \right) = 0,
$$

which implies that the mappings $y \in \mathbb{T}^d \mapsto K_l^{\mu}(t, x_0, \cdot, \mu, y) \in C^s(\mathbb{T}^d)$ and $y \in \mathbb{T}^d \mapsto K_l^{0,\mu}(t, x_0, \mu, y) \in \mathbb{R}$ are continuous. Similarly, for any $l \in \{0, \dots, \lfloor s \rfloor - 2\}^d$ with $|l| \leq \lfloor s \rfloor - 2$, and any $i \in \{1, \dots, d\}$,

$$
\lim_{h \in \mathbb{R}, h \to 0} \left\| \frac{1}{h} \left(\nabla^l \delta_{y+he_i} - \nabla^l \delta_y \right) + \nabla^{l+e_i} \delta_y \right\|_{-\tau+1} = 0.
$$

Therefore, again by Theorem [5.5,](#page-61-1) we get

$$
\lim_{h \in \mathbb{R}, h \to 0} \left(\left\| \frac{1}{h} \left(K_l^{\mu}(t, x_0, \cdot, \mu, y + he_i) - K_l^{\mu}(t, x_0, \cdot, \mu, y) \right) - K_{l+e_i}^{\mu}(t, x_0, \cdot, \mu, y) \right\|_{\mathfrak{s}} + \left| \frac{1}{h} \left(K_l^{0,\mu}(t, x_0, \mu, y + he_i) - K_l^{0,\mu}(t, x_0, \mu, y) \right) - K_{l+e_i}^{0,\mu}(t, x_0, \mu, y) \right| \right) = 0,
$$

which implies, by induction, that

$$
\nabla_y^l K_0^{\mu}(t, x_0, x, \mu, y) = K_l^{\mu}(t, x_0, x, \mu, y) \quad \text{and} \quad \nabla_y^l K_0^{0,\mu}(t, x_0, \mu, y) = K_l^{0,\mu}(t, x_0, \mu, y).
$$

The proof is completed.

The following statement is the analogue of [\[14](#page-86-1), Lemma 5.2.3]:

Lemma 5.7. Given $\triangle x_0 \in \mathbb{R}^d$ and a finite signed measure $\triangle \mu$ on \mathbb{T}^d , the solution $(\delta \mu, \delta u, \delta m, \delta X^0, \delta Y^0, \delta Z^0)$ to the *system* [\(5.32\)](#page-61-0) *has the following representation formulas*

$$
\delta u_t(x) = K^{x_0}(t, x_0, x, \mu) \cdot \Delta x_0 + \int_{\mathbb{T}^d} K_0^{\mu}(t, x_0, x, \mu, y) \Delta \mu(\mathrm{d}y), \tag{5.37}
$$

and

$$
\delta Y_t^0 = K^{0,x_0}(t, x_0, \mu) \cdot \Delta x_0 + \int_{\mathbb{T}^d} K_0^{0,\mu}(t, x_0, \mu, y) \Delta \mu(\mathrm{d}y). \tag{5.38}
$$

Proof. By compactness of the torus, we can find, for a given $\varepsilon > 0$, a covering $\{U_i\}_{1 \le i \le N}$ of \mathbb{T}^d , made of disjoint Borel subsets with diameter less than ε . Choosing, for each $i \in \{1, \dots, N\}$, $y_i \in U_i$, we then let

$$
\triangle \mu^{\varepsilon} := \sum_{i=1}^N \triangle \mu(U_i) \delta_{y_i}.
$$

Then,

$$
\|\Delta \mu - \Delta \mu^{\varepsilon}\|_{-1} = \sup_{\|\phi\|_{1} \le 1} \left| \int_{\mathbb{T}^{d}} \phi(y) (\Delta \mu(\mathrm{d}y) - \Delta \mu^{\varepsilon}(\mathrm{d}y)) \right|
$$

$$
= \sup_{\|\phi\|_{1} \le 1} \left| \sum_{i=1}^{N} \int_{U_{i}} (\phi(y) - \phi(y_{i})) \Delta \mu(\mathrm{d}y) \right| \le C |\Delta \mu|_{\mathrm{TV}} \varepsilon,
$$

where $|\triangle \mu|_{TV}$ is the total mass of $\triangle \mu$. Consider now [\(5.15\)](#page-54-0) with $(\triangle x_0, \triangle \mu^{\epsilon})$ as initial condition at time t and denote the corresponding solution by $(\delta\mu^{\varepsilon}, \delta u^{\varepsilon}, \delta m^{\varepsilon}, \delta X^{0,\varepsilon}, \delta Y^{0,\varepsilon}, \delta Z^{0,\varepsilon})$. By linearity of the system [\(5.32\)](#page-61-0), we observe that

$$
\delta Y_t^{0,\varepsilon} = K^{0,x_0}(t, x_0, \mu) \cdot \Delta x_0 + \sum_{i=1}^N K_0^{0,\mu}(t, x_0, \mu, y_i) \Delta \mu(U_i)
$$

$$
= K^{0,x_0}(t, x_0, \mu) \cdot \Delta x_0 + \sum_{i=1}^N \int_{U_i} K_0^{0,\mu}(t, x_0, \mu, y_i) \Delta \mu(\mathrm{d}y),
$$

$$
\delta u_t^{\varepsilon}(x) = K^{x_0}(t, x_0, x, \mu) \cdot \Delta x_0 + \sum_{i=1}^N K_0^{\mu}(t, x_0, x, \mu, y_i) \Delta \mu(U_i)
$$

$$
= K^{x_0}(t, x_0, x, \mu) \cdot \Delta x_0 + \sum_{i=1}^N \int_{U_i} K_0^{\mu}(t, x_0, x, \mu, y_i) \Delta \mu(\mathrm{d}y), \quad x \in \mathbb{T}^d.
$$

 \Box

And then, by invoking Theorem [5.5](#page-61-1) in order to compare $(\delta u, \delta Y^0)$ and $(\delta u^{\varepsilon}, \delta Y^{0,\varepsilon})$, we obtain

$$
\left\|\delta u_t(\cdot) - K^{x_0}(t, x_0, \cdot, \mu) \cdot \Delta x_0 - \sum_{i=1}^N \int_{U_i} K_0^{\mu}(t, x_0, \cdot, \mu, y_i) \Delta \mu(\mathrm{d}y) \right\|_{\mathfrak{s}} + \left|\delta Y_t^0 - K^{0, x_0}(t, x_0, \mu) \cdot \Delta x_0 - \sum_{i=1}^N \int_{U_i} K_0^{0, \mu}(t, x_0, \mu, y_i) \Delta \mu(\mathrm{d}y) \right| \leq C \|\Delta \mu - \Delta \mu^{\varepsilon}\|_{-\mathfrak{s}+1} \leq C \|\Delta \mu - \Delta \mu^{\varepsilon}\|_{-1} \leq C \|\Delta \mu|_{\mathrm{TV}} \varepsilon.
$$

By the smoothness of K_0^{μ} and $K_0^{0,\mu}$ derived in Lemma [5.6,](#page-62-1) we can easily derive that

$$
\left\|\delta u_t(\cdot) - K^{x_0}(t, x_0, \cdot, \mu) \cdot \Delta x_0 - \sum_{i=1}^N \int_{U_i} K_0^{\mu}(t, x_0, \cdot, \mu, y) \Delta \mu(\mathrm{d}y) \right\|_{\mathfrak{s}} + \left|\delta Y_t^0 - K^{0, x_0}(t, x_0, \mu) \cdot \Delta x_0 - \sum_{i=1}^N \int_{U_i} K_0^{0, \mu}(t, x_0, \mu, y) \Delta \mu(\mathrm{d}y) \right| \leq C |\Delta \mu|_{\text{TV}} \varepsilon.
$$
\n(5.39)

Letting $\varepsilon \to 0$ in [\(5.39\)](#page-64-0), we obtain [\(5.37\)](#page-63-0) and [\(5.38\)](#page-63-1).

5.4. Differentiability of the fields \mathcal{U}^0 and \mathcal{U} in the variables x_0 and μ

The following statement provides a second-order expansion of the fields \mathcal{U}^0 and \mathcal{U} :

Proposition 5.8. *Under Assumption* (\hat{A}), *consider two initial conditions* (t, x_0, μ) , $(t, \hat{x}_0, \hat{\mu}) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and call $(\mu, u, m, X^0, Y^0, Z^0)$ and $(\hat{\mu}, \hat{u}, \hat{m}, \hat{X}^0, \hat{Y}^0, \hat{Z}^0)$ the solutions of the system [\(5.1\)](#page-50-3)–[\(5.2\)](#page-50-2) with (t, x_0, μ) and $(t, \hat{x}_0, \hat{\mu})$ as respective initial conditions and $(\delta\mu, \delta\mathbf{u}, \delta\mathbf{m}, \delta\mathbf{X}^0, \delta\mathbf{Y}^0, \delta\mathbf{Z}^0)$ the solution of the system [\(5.32\)](#page-61-0) with $(t, \hat{x}_0 - x_0, \hat{\mu} - \mu)$ *as initial condition, so that we can let*

$$
\Delta^2 \mathbf{X}^0 := \hat{\mathbf{X}}^0 - \mathbf{X}^0 - \delta \mathbf{X}^0, \quad \Delta^2 \mathbf{Y}^0 := \hat{\mathbf{Y}}^0 - \mathbf{Y}^0 - \delta \mathbf{Y}^0, \quad \Delta^2 \mathbf{Z}^0 := \hat{\mathbf{Z}}^0 - \mathbf{Z}^0 - \delta \mathbf{Z}^0,
$$

$$
\Delta^2 \mu := \hat{\mu} - \mu - \delta \mu, \quad \Delta^2 \mathbf{u} := \hat{\mathbf{u}} - \mathbf{u} - \delta \mathbf{u}, \quad \Delta^2 \mathbf{m} := \hat{\mathbf{m}} - \mathbf{m} - \delta \mathbf{m}.
$$

Let $\Gamma \in (|\mathfrak{s}|, \mathfrak{s})$ *. Then, for* $T \leq \mathfrak{C}$ *, for some* $\mathfrak{C} > 0$ *only depending on the parameters* d*,* κ *,* \mathfrak{L} *,* ζ *,* σ_0 *and* \mathfrak{s} *in Assumption* (Â) *and on* **r***, we can find a constant* C*, only depending on the parameters in Assumption* (Â) *and on* **r***, such that, for any* $p \in [1, 8]$ *,*

$$
\mathbb{E}^{0}\bigg[\sup_{r\in[t,T]}\Big(\|\triangle^{2}\mu_{r}\|_{-\mathbf{r}+1}^{p}+\|\triangle^{2}u_{r}\|_{\mathbf{s}}^{p}+\|\triangle^{2}m_{r}\|_{\mathbf{s}-2}^{p}\Big)\bigg]
$$
\n
$$
+\mathbb{E}^{0}\bigg[\sup_{r\in[t,T]}\|\triangle^{2}X_{r}^{0}|^{p}+\sup_{r\in[t,T]}\|\triangle^{2}Y_{t}^{0}|^{p}+\int_{t}^{T}\|\triangle^{2}Z_{r}^{0}|^{p}\mathrm{d}r\bigg]\leq C_{p}\big(\mathbb{W}_{1}(\hat{\mu},\mu)^{2p}+|\hat{x}_{0}-x_{0}|^{2p}\big).
$$
\n(5.40)

In particular, by choosing $p = 1$ *, we get (with* $C := C_1$ *)*

$$
\left\| \mathcal{U}(t, \hat{x}_0, \cdot, \hat{\mu}) - \mathcal{U}(t, x_0, \cdot, \mu) - K^{x_0}(t, x_0, \cdot, \mu) \cdot (\hat{x}_0 - x_0) - \int_{\mathbb{T}^d} K_0^{\mu}(t, x_0, \cdot, \mu, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)) \right\|_{\mathfrak{s}} \n\leq C \Big(\mathbb{W}_1(\hat{\mu}, \mu)^2 + |\hat{x}_0 - x_0|^2 \Big),
$$
\n
$$
\left| \mathcal{U}^0(t, \hat{x}_0, \hat{\mu}) - \mathcal{U}^0(t, x_0, \mu) - K^{0, x_0}(t, x_0, \mu) \cdot (\hat{x}_0 - x_0) - \int_{\mathbb{T}^d} K_0^{0, \mu}(t, x_0, \mu, y) (\hat{\mu} - \mu)(\mathrm{d}y) \right| \n\leq C \Big(\mathbb{W}_1(\hat{\mu}, \mu)^2 + |\hat{x}_0 - x_0|^2 \Big),
$$

 \Box

and, thus, for any $(t, x) \in [0, T] \times \mathbb{T}^d$ *, the mappings*

$$
(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto \mathcal{U}(t, x_0, x, \mu)
$$
 and $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \mapsto \mathcal{U}^0(t, x_0, \mu)$

are differentiable with respect to x_0 *and* μ *and the derivatives read, for any* $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$,

$$
\nabla_{x_0} \mathcal{U}(t, x_0, x, \mu) = K^{x_0}(t, x_0, x, \mu), \quad \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu) = K^{0, x_0}(t, x_0, \mu),
$$

and

$$
\delta_{\mu}\mathcal{U}(t,x_0,x,\mu,y) = K_0^{\mu}(t,x_0,x,\mu,y), \quad \delta_{\mu}\mathcal{U}^0(t,x_0,\mu,y) = K_0^{0,\mu}(t,x_0,\mu,y), \quad \text{for any } y \in \mathbb{T}^d.
$$

Proof. We first note that $(\triangle^2 \mathbf{X}^0, \triangle^2 \mathbf{Y}^0, \triangle^2 \mathbf{Z}^0)$ solves, on $[t, T]$,

$$
d_r \Delta^2 X_r^0 = \left(-\nabla_{px_0} H^0(X_r^0, Z_r^0) \Delta^2 X_r^0 - \nabla_{pp} H^0(X_r^0, Z_r^0) \Delta^2 Z_r^0 + a_r \right) dr,
$$

\n
$$
d_r \Delta^2 Y_r^0 = \left[-\nabla_{x_0} f_r^0(X_r^0, \mu_r) \cdot \Delta^2 X_r^0 - \left(\delta_\mu f_r^0(X_r^0, \mu_r, \cdot), \Delta^2 \mu_r \right) \right.
$$

\n
$$
+ \nabla_{x_0} \hat{L}^0(X_r^0, Z_r^0) \Delta^2 X_r^0 + \nabla_p \hat{L}^0(X_r^0, Z_r^0) \Delta^2 Z_r^0 + b_r \right] dr + \sigma_0 \Delta^2 Z_r^0 \cdot dB_r^0, \text{ on } [t, T],
$$

\n
$$
\Delta^2 X_t^0 = 0, \quad \Delta^2 Y_T^0 = \nabla_{x_0} g^0(X_T^0, \mu_T) \cdot \Delta^2 X_T^0 + \left(\delta_\mu g^0(X_T^0, \mu_T, \cdot), \Delta^2 \mu_T \right) + c_T,
$$

where

$$
a_r = \left(\nabla^2_{px_0} H^0(X_r^0, Z_r^0) - \int_0^1 \nabla^2_{px_0} H^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{Z}_r^0 + (1 - \theta)Z_r^0)d\theta\right)(\hat{X}_r^0 - X_r^0)
$$

+
$$
\left(\nabla^2_{pp} H^0(X_r^0, Z_r^0) - \int_0^1 \nabla^2_{pp} H^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{Z}_r^0 + (1 - \theta)Z_r^0)d\theta\right)(\hat{Z}_r^0 - Z_r^0),
$$

$$
b_r = \left(\nabla_{x_0} f^0(X_r^0, \mu_r) - \int_0^1 \nabla_{x_0} f^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{\mu}_r + (1 - \theta)\mu_r)d\theta\right) \cdot (\hat{X}_r^0 - X_r^0)
$$

+
$$
\left(\delta_{\mu} f^0(X_r^0, \mu_r, \cdot) - \int_0^1 \delta_{\mu} f^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{\mu}_r + (1 - \theta)\mu_r, \cdot)d\theta, \hat{\mu}_r - \mu_r\right)
$$

-
$$
\left(\nabla_{x_0} \hat{L}^0(X_r^0, Z_r^0) - \int_0^1 \nabla_{x_0} \hat{L}^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{Z}_r^0 + (1 - \theta)Z_r^0)d\theta\right) \cdot (\hat{X}_r^0 - X_r^0)
$$

-
$$
\left(\nabla_p \hat{L}^0(X_r^0, Z_r^0) - \int_0^1 \nabla_p \hat{L}^0(\theta \hat{X}_r^0 + (1 - \theta)X_r^0, \theta \hat{Z}_r^0 + (1 - \theta)Z_r^0)d\theta\right) \cdot (\hat{Z}_r^0 - Z_r^0)
$$

and

$$
c_T = -\left(\nabla_{x_0} g^0(X_T^0, \mu_T) - \int_0^1 \nabla_{x_0} g^0(\theta \hat{X}_T^0 + (1 - \theta) X_T^0, \theta \hat{\mu}_T + (1 - \theta) \mu_T) d\theta\right) \cdot (\hat{X}_T^0 - X_T^0) - \left(\delta_\mu g^0(X_T^0, \mu_T, \cdot) - \int_0^1 \delta_\mu g^0(\theta X_T^0 + (1 - \theta) X_T^0, \theta \hat{\mu}_T + (1 - \theta) \mu_T, \cdot), \hat{\mu}_T - \mu_T\right).
$$

We then notice that $(\Delta^2 \mu, \Delta^2 u, \Delta^2 m)$ satisfies, on $[t, T]$,

$$
\partial_r \triangle^2 \mu_r - \frac{1}{2} \Delta_x \triangle^2 \mu_r - \text{div}_x (\nabla_p \hat{H}(x, \nabla_x u_r(x)) \triangle^2 \mu_r + \mu_r \nabla^2_{pp} \hat{H}(x, \nabla_x u_r(x)) \nabla_x \delta^2 u_r(x))
$$

\n
$$
- \text{div}_x (d_r(x)) = 0, \quad \text{on } \mathbb{T}^d,
$$

\n
$$
\mathrm{d}_r \delta^2 u_r(x) = \left[-\frac{1}{2} \Delta_x \delta^2 u_r(x) + \nabla_p \hat{H}(x, \nabla_x u_r(x)) \cdot \nabla_x \delta^2 u_r(x) - \nabla_{x_0} f_r(X_r^0, x, \mu_r) \cdot \delta^2 X_r^0 - \left(\delta_\mu f_r(X_r^0, x, \mu_r, \cdot), \delta^2 \mu_r \right) + j_r(x) \right] \mathrm{d}r + \mathrm{d}_r \delta^2 m_r(x), \quad x \in \mathbb{T}^d,
$$

\n
$$
\delta \mu_t = 0, \quad \delta u_T(x) = \nabla_{x_0} g(X_T^0, x, \mu_T) \cdot \delta^2 X_T^0 + \left(\delta_\mu g(X_T^0, x, \mu_T, \cdot), \delta^2 \mu_T \right) + k_T, \quad x \in \mathbb{T}^d,
$$

where

$$
d_r(x) = (\hat{\mu}_r - \mu_r) \left(\int_0^1 \nabla_{pp}^2 \hat{H}(x, \theta \nabla_x \hat{u}_r(x) + (1 - \theta) \nabla_x u_r(x)) d\theta \right) (\nabla_x \hat{u}_r(x) - \nabla_x u_r(x))
$$

+
$$
\mu_r \int_0^1 \int_0^1 \theta \nabla_{ppp}^3 \hat{H}(x, \nabla_x u_r(x) + \tilde{\theta} \theta (\nabla_x \hat{u}_r(x) - \nabla_x u_r(x))) (\nabla_x \hat{u}_r(x) - \nabla_x u_r(x))^{\otimes 2} d\tilde{\theta} d\theta,
$$

$$
j_r(x) = -\left(\nabla_p \hat{H}(x, \nabla_x u_r(x)) - \int_0^1 \nabla_p \hat{H}(x, \theta \nabla_x \hat{u}_r(x) + (1 - \theta) \nabla_x u_r(x)) d\theta \right) \cdot (\nabla_x \hat{u}_r(x) - \nabla_x u_r(x))
$$

+
$$
\left(\nabla_{x_0} f_r(X_r^0, x, \mu_r) - \int_0^1 \nabla_{x_0} f_r(\theta \hat{X}_r^0 + (1 - \theta) X_r^0, x, \theta \hat{\mu}_r + (1 - \theta) \mu_r) d\theta \right) \cdot (\hat{X}_r^0 - X_r^0)
$$

+
$$
\left(\delta_\mu f_r(X_r^0, x, \mu_r, \cdot) - \int_0^1 \delta_\mu f_r(\theta \hat{X}_r^0 + (1 - \theta) X_r^0, x, \theta \hat{\mu}_r + (1 - \theta) \mu_r, \cdot) d\theta, \hat{\mu}_r - \mu_r \right),
$$

and

$$
k_T(x) = -\left(\nabla_{x_0} g(X_T^0, x, \mu_T) - \int_0^1 \nabla_{x_0} g(\theta \hat{X}_T^0 + (1 - \theta) X_T^0, x, \theta \hat{\mu}_T + (1 - \theta) \mu_T) d\theta\right) \cdot (\hat{X}_T^0 - X_T^0) - \left(\delta_{\mu} g(X_T^0, x, \mu_T) - \int_0^1 \delta_{\mu} g(\theta X_T^0 + (1 - \theta) X_T^0, x, \theta \hat{\mu}_T + (1 - \theta) \mu_T, \cdot), \hat{\mu}_T - \mu_T\right).
$$

Next, we recall μ and $\hat{\mu}$ take values in the space $\mathcal{P}(\mathbb{T}^d)$. We deduce that

$$
||d_r||_{-1} \leq C(|\hat{u}_r - u_r||_2 \mathbb{W}_1(\hat{\mu}_t, \mu_t) + ||\hat{u}_r - u_r||_1^2), \quad r \in [t, T]
$$

\n
$$
||k_T||_{\mathfrak{s}} \leq C(|\hat{X}_T^0 - X_T^0|^2 + \mathbb{W}_1(\hat{\mu}_T, \mu_T)^2),
$$

\n
$$
||j_r||_{\mathfrak{s}-1} \leq C(|\hat{u}_r - u_r||_{\mathfrak{s}}^2 + |\hat{X}_r^0 - X_r^0|^2 + \mathbb{W}_1(\hat{\mu}_r, \mu_r)^2),
$$

\n
$$
\int_t^T |a_r| dr \leq C\left[\sup_{r \in [t, T]} |\hat{X}_r^0 - X_r^0|^2 + \int_t^T |\hat{Z}_r^0 - Z_r^0|^2 dr\right]
$$

\n
$$
\int_t^T |b_r| dr \leq C\left[\sup_{r \in [t, T]} |\hat{X}_r^0 - X_r^0|^2 + \int_t^T \mathbb{W}_1(\hat{\mu}_r, \mu_r)^2 dr + \int_t^T |\hat{Z}_r^0 - Z_r^0|^2 dr\right],
$$

\n
$$
|c_T| \leq C(|\hat{X}_T^0 - X_T^0|^2 + \mathbb{W}_1^2(\hat{\mu}_T, \mu_T)).
$$

We further apply [\(5.5\)](#page-51-4) to obtain, for any $p \ge 1$,

$$
\mathbb{E}\Big[\sup_{r\in[t,T]}||d_r||_{-\mathbb{r}+2}^p\Big] \leq C_p \big(\mathbb{W}_1^{2p}(\hat{\mu},\mu) + |\hat{x}_0 - x_0|^{2p}\big),
$$
\n
$$
\mathbb{E}\Big[\|k_T\|_{s}^p\Big] \leq C \big(\mathbb{W}_1^{2p}(\hat{\mu},\mu) + |\hat{x}_0 - x_0|^{2p}\big),
$$
\n
$$
\mathbb{E}\Big[\sup_{r\in[t,T]}||j_r||_{s-1}^p\Big] \leq C \big(\mathbb{W}_1^{2p}(\hat{\mu},\mu) + |\hat{x}_0 - x_0|^{2p}\big),
$$
\n
$$
\mathbb{E}\Big[\bigg(\int_t^T |a_r|dr\bigg)^p + \bigg(\int_t^T |b_r|dr\bigg)^p + c_T^2\Bigg] \leq C \big(\mathbb{W}_1^{2p}(\hat{\mu},\mu) + |\hat{x}_0 - x_0|^{2p}\big).
$$

Therefore, by Theorem [5.2,](#page-55-0) we have [\(5.40\)](#page-64-1).

Corollary 5.9. *There exist a constant* $\mathfrak{C} > 0$ *, only depending on the parameters d,* κ *,* \mathfrak{L} *,* ζ *,* σ_0 *and s in Assumption* (\hat{A})*,* and a constant C, only depending on the parameters in Assumption ($\hat{\bf A}$), such that, for $T\leq{\frak C}$, for all $t\in[0,T]$, ${\cal U}^{0}(t,\cdot,\cdot)$ satisfies (ii) and (iii) in the definition of $\mathscr{C}^0(C, [\mathfrak{s}]-1)$ and $\mathcal{U}(t, \cdot, \cdot, \cdot)$ satisfies (ii), (iii) and (iv) in the definition of $\mathscr{C}(C, \lfloor \mathfrak{s} \rfloor - 1, \mathfrak{s}).$

$$
\Box
$$

Proof. The bounds for $|\mathcal{U}^0(t, x_0, \mu)|$ and $||\mathcal{U}(t, x_0, \cdot, \mu)||_s$ follow from [\(5.3\)](#page-51-0) and [\(5.4\)](#page-51-1). The bounds for $|\nabla_{x_0} \mathcal{U}^0(t, x_0, \mu)|$ and $\|\nabla_{x_0} \mathcal{U}(t, x_0, \cdot, \mu)\|_{\mathfrak{s}}$ follow from the inequality [\(5.35\)](#page-61-2). Moreover, the bounds for $\|\delta_\mu \mathcal{U}^0(t, x_0, \mu, \cdot)\|_{\lfloor \mathfrak{s} \rfloor - 1}$ and $\max_{l=0,\cdots,\lfloor s\rfloor-1} \|\nabla_y^l \delta_\mu \mathcal{U}^0(t,x_0,\cdot,\mu,\cdot)\|_s$ follow from Lemma [5.6.](#page-62-1) Continuity of the derivatives as stated in the definition of \mathscr{C}^0 and \mathscr{C} also follows from Lemma [5.6.](#page-62-1)

In the following statement, we address the spatial regularity of the derivatives of \mathcal{U} and \mathcal{U}^0 :

Proposition 5.10. Given two 4-tuples $(t, x_0, \mu, y), (t, \hat{x}_0, \hat{\mu}, \hat{y}) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$, a unitary vector $e \in \mathbb{R}^d$, *two scalars* $\vartheta_0, \vartheta \in [0,1]$, a fixed $\mathbb{r} \in (\lfloor \mathfrak{s} \rfloor, \mathfrak{s})$ and a d-tuple $l \in \{0, \cdots, \lfloor \mathfrak{s} \rfloor - 1\}^d$ with $|l| := \sum_{i=1}^d l_i \leq \lfloor \mathfrak{s} \rfloor - 1$, call $(\mu, u, m, X^0, Y^0, Z^0)$ and $(\hat{\mu}, \hat{u}, \hat{m}, \hat{X}^0, \hat{Y}^0, \hat{Z}^0)$ the solutions of the system [\(5.1\)](#page-50-3)–[\(5.2\)](#page-50-2) with (t, x_0, μ) and $(t, \hat{x}_0, \hat{\mu})$ as respective initial conditions and then $(\delta\mu, \delta u, \delta m, \delta X^0, \delta Y^0, \delta Z^0)$ and $(\delta\hat\mu, \delta\hat u, \delta\hat m, \delta\hat X^0, \delta\hat Y^0, \delta\hat Z^0)$ the solutions *of the linearized system* [\(5.32\)](#page-61-0) *with* $(t, \vartheta_0 e, \vartheta(-1)^{|l|} \nabla^l \delta_y)$ *and* $(t, \vartheta_0 e, \vartheta(-1)^{|l|} \nabla^l \delta_y)$ *as respective initial conditions.*

Then, for $T \leq \mathfrak{C}$ *, for some* $\mathfrak{C} > 0$ *only depending on the parameters* d, κ *,* \mathfrak{L} *,* ζ *,* σ ₀ and s in Assumption (\hat{A}) and on Γ *, we can find a constant* C, only depending on the parameters in Assumption (\hat{A}) and on τ , such that, for any $p \in [1, 4]$,

$$
\mathbb{E}^{0}\left[\sup_{r\in[t,T]}\left(\|\delta\hat{\mu}_{r}-\delta\mu_{r}\|_{-r+1}^{2p}+\|\delta\hat{u}_{r}-\delta u_{r}\|_{s}^{2p}+\|\delta\hat{m}_{r}-\delta m_{r}\|_{s-2}^{2p}\right)\right] \n+\mathbb{E}^{0}\left[\sup_{r\in[t,T]}\left(|\delta\hat{X}_{r}^{0}-\delta X_{r}^{0}|^{2p}+|\delta\hat{Y}_{t}^{0}-\delta Y_{t}^{0}|^{2p}\right)+\left(\int_{t}^{T}|\delta\hat{Z}_{r}^{0}-\delta Z_{r}^{0}|^{2}\mathrm{d}r\right)^{p}\right] \n\leq C\left(|\hat{x}_{0}-x_{0}|^{2p}+\mathbb{W}_{1}^{2p}(\hat{\mu},\mu)+|\hat{y}-y|^{2p(r-\lfloor s\rfloor)}\right).
$$
\n(5.41)

In particular, for any $y, \hat{y} \in \mathbb{T}^d$, *for any* $k \in \{0, \dots, \lfloor \mathfrak{s} \rfloor - 1\}$ *,*

$$
\begin{split}\n\left\| \nabla_{x_0} \mathcal{U}(t, x_0, \cdot, \mu) - \nabla_{x_0} \mathcal{U}(t, \hat{x}_0, \cdot, \hat{\mu}) \right\|_{\mathfrak{s}} + \left| \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu) - \nabla_{x_0} \mathcal{U}^0(t, \hat{x}_0, \hat{\mu}) \right| &\leq C \Big(\left| \hat{x}_0 - x_0 \right| + \mathbb{W}_1(\hat{\mu}, \mu) \Big), \\
\left\| \nabla_y^k \delta_\mu \mathcal{U}(t, x_0, \cdot, \mu, y) - \nabla_y^k \delta_\mu \mathcal{U}(t, \hat{x}_0, \cdot, \hat{\mu}, \hat{y}) \right\|_{\mathfrak{s}} + \left| \nabla_y^k \delta_\mu \mathcal{U}^0(t, x_0, \mu, y) - \nabla_y^k \delta_\mu \mathcal{U}^0(t, \hat{x}_0, \hat{\mu}, \hat{y}) \right| \\
&\leq C \Big(\left| \hat{x}_0 - x_0 \right| + \mathbb{W}_1(\hat{\mu}, \mu) + \left| \hat{y} - y \right|^{r - \lfloor \mathfrak{s} \rfloor} \Big),\n\end{split} \tag{5.42}
$$

and, for any $t \in [0, T]$, $\mathcal{U}^0(t, \cdot, \cdot) \in \mathcal{D}^0(C, \mathbb{r} - 1)$ *and* $\mathcal{U}(t, \cdot, \cdot, \cdot) \in \mathcal{D}(C, \mathbb{r} - 1, \mathbb{s})$ *.*

Proof. Applying Theorem [5.1,](#page-50-1) we obtain (for $T \leq \mathfrak{C}$)

$$
\mathbb{E}^{0}\left[\sup_{r\in[t,T]}\left(\|\hat{u}_{r}-u_{r}\|_{s}^{4p}+\mathbb{W}_{1}^{4p}(\hat{\mu}_{r},\mu_{r})+|\hat{X}_{r}^{0}-X_{r}^{0}|^{4p}+|\hat{Y}_{r}^{0}-Y_{r}^{0}|^{4p}\right)+\left(\int_{t}^{T}|\hat{Z}_{r}^{0}-Z_{r}^{0}|^{2}\mathrm{d}r\right)^{2p}\right]
$$
\n
$$
\leq C\left[|\hat{x}_{0}-x_{0}|^{4p}+\mathbb{W}_{1}^{4p}(\hat{\mu},\mu)\right].
$$
\n(5.43)

By Theorem [5.5,](#page-61-1)

$$
\mathbb{E}^{0}\left[\sup_{r\in[t,T]}\left(\|\delta\mu_{r}\|_{-r+1}^{4p}+\|\delta\hat{\mu}_{r}\|_{-r+1}^{4p}+\|\delta u_{r}\|_{s}^{4p}+\|\delta\hat{u}_{r}\|_{s}^{4p}+|\delta X_{r}^{0}|^{4p}+|\delta\hat{X}_{r}^{0}|^{4p}+|\delta Y_{r}^{0}|^{4p}+|\delta\hat{Y}_{r}^{0}|^{4p}\right)\right] + \left(\int_{0}^{T} \left(|\delta Z_{t}^{0}|^{2}+|\delta\hat{Z}_{t}^{0}|^{2}\right)dt\right)^{2p}\right]
$$
\n
$$
\leq C.
$$
\n(5.44)

It remains to see that, since $|l| \leq |s| - 1$, we have

$$
\|\nabla^l \delta_{\hat{y}} - \nabla^l \delta_{y}\|_{-\mathbb{r}+1} \leq |\hat{y} - y|^{\mathbb{r} - \lfloor \mathfrak{s} \rfloor}.
$$

In order to conclude, we follow the proof of Proposition [5.8.](#page-64-2) The idea is to write $(\delta \hat{\mu} - \delta \mu, \delta \hat{u} - \delta u, \delta \hat{m} - \delta m, \delta \hat{X}^0$ δX^0 , $\delta \hat{Y}^0 - \delta Y^0$, $\delta \hat{Z}^0 - \delta Z^0$) as a solution of a linear system of the form [\(5.15\)](#page-54-0), with suitable coefficients $(a_t)_{0 \le t \le T}$, $(b_t)_{0 \le t \le T}$, $(c_T, (d_t)_{0 \le t \le T}$, $(j_t)_{0 \le t \le T}$ and k_T . Briefly, all these terms can be bounded from above by products of two terms: one of the norms appearing in (5.43) and one of the norms of (5.44) . Cauchy-Schwarz inequality permits to handle the $2p$ moment of any of these products. By (5.43) and (5.44) , we obtain (5.41) .

Then, we get the first line in [\(5.42\)](#page-67-3) by choosing $p = 1$, $\vartheta = 1$, $\vartheta = 0$ and $y = \hat{y}$. The second inequality is obtained by choosing $p = 1$, $\vartheta_0 = 0$ and $\vartheta = 1$. \square

5.5. Regularity in time of the derivatives

We now address the time continuity of the derivatives. We start with the following statement:

Proposition 5.11. *Let* $\mathbf{r} \in (\mathbf{s} \mid \mathbf{s})$ *and let Assumption* ($\hat{\mathbf{A}}$) *be in force. Then, there exists a constant* $\mathfrak{C} > 0$ *, only depending* on d, κ , \mathfrak{L} , ζ , σ_0 and s in Assumption (\hat{A}) and on \mathbb{r} , such that, for $T \leq \mathfrak{C}$, for any $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, it *holds*

$$
\lim_{h \to 0} \left(\left\| \nabla_{x_0} \mathcal{U}(t+h, x_0, \cdot, \mu) - \nabla_{x_0} \mathcal{U}(t, x_0, \cdot, \mu) \right\|_{\mathbb{F}} + \left\| \delta_\mu \mathcal{U}(t+h, x_0, \cdot, \mu, \cdot) - \delta_\mu \mathcal{U}(t, x_0, \cdot, \mu, \cdot) \right\|_{\mathbb{F}, \mathbb{F} - 1} \right) = 0, \quad (5.45)
$$

where $\| \cdot \|_{\mathbb{r}, \mathbb{r}-1}$ *denotes the standard Hölder norm^{[2](#page-68-0)} on the space* $C^{\mathbb{r}, \mathbb{r}-1}(\mathbb{T}^d \times \mathbb{T}^d)$ *of functions* $h : (x, y) \in \mathbb{T}^d \times \mathbb{T}^d \mapsto$ $h(x, y)$ *that have crossed derivatives in* (x, y) *up the order* $|\mathbf{r}|$ *in* x and $|\mathbf{r}| - 1$ *in* y, all the derivatives being $\mathbf{r} - |\mathbf{s}|$ *Hölder continuous in* (x, y) *. Similarly,*

$$
\lim_{h \to 0} (|\nabla_{x_0} \mathcal{U}^0(t + h, x_0, \mu) - \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu)| + ||\delta_\mu \mathcal{U}^0(t + h, x_0, \mu, \cdot) - \delta_\mu \mathcal{U}^0(t, x_0, \mu, \cdot)||_{\mathbb{F}_{-1}}) = 0.
$$
 (5.46)

Proof. Given (x_0, μ) , $(\hat{x}_0, \hat{\mu}) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$, we can derive from Proposition [5.8](#page-64-2) that, for $T \leq \mathfrak{C}$ and for any $t \in [0, T]$,

$$
\mathcal{U}(t, \hat{x}_0, \cdot, \hat{\mu}) - \mathcal{U}(t, x_0, \cdot, \mu)
$$

= $\nabla_{x_0} \mathcal{U}(t, x_0, \cdot, \mu) \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \delta_\mu \mathcal{U}(t, x_0, \cdot, \mu, y) (\hat{\mu}(dy) - \mu(dy)) + O(|\hat{x}_0 - x_0|^2 + \mathbb{W}_1^2(\hat{\mu}, \mu)),$ (5.47)

where the equality holds true in $C^{s}(\mathbb{T}^d)$ and the Landau notation $O(\cdot)$ is uniform in (t, x_0, μ) , and

$$
\mathcal{U}^{0}(t, \hat{x}_{0}, \hat{\mu}) - \mathcal{U}^{0}(t, x_{0}, \mu)
$$
\n
$$
= \nabla_{x_{0}} \mathcal{U}^{0}(t, x_{0}, \mu) \cdot (\hat{x}_{0} - x_{0}) + \int_{\mathbb{T}^{d}} \delta_{\mu} \mathcal{U}^{0}(t, x_{0}, \mu, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)) + O(|\hat{x}_{0} - x_{0}|^{2} + \mathbb{W}_{1}^{2}(\hat{\mu}, \mu)). \tag{5.48}
$$

We now fix $(x_0, \mu) \in \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and let $\mathbf{u} := (\mathbf{r} + \mathbf{s})/2$. Recall [\(5.35\)](#page-61-2) and the fact that $\nabla_{x_0} \mathcal{U}^0(t, x_0, \mu) =$ $K^{0,x_0}(t, x_0, \mu)$. Therefore, the collection $\{\nabla_{x_0} \mathcal{U}^0(t, x_0, \mu), t \in [0, T]\}$ is bounded and thus relatively compact in \mathbb{R}^d . By Proposition [5.10](#page-67-4) (applied with the indices (u, s) in lieu of (r, s)) and under the same generic condition $T \leq \mathfrak{C}$ as therein, the collection of functions $\{x \in \mathbb{T}^d \mapsto \nabla_{x_0} \mathcal{U}(t, x_0, x, \mu), t \in [0, T]\}$ is relatively compact in $\mathcal{C}^r(\mathbb{T}^d)$, the collection of functions $\{y \in \mathbb{T}^d \mapsto \delta_\mu U^0(t, x_0, \mu, y), t \in [0, T]\}$ is relatively compact in $\mathcal{C}^{r-1}(\mathbb{T}^d)$, and the collection of functions $\{(x, y) \in \mathbb{T}^d \times \mathbb{T}^d \mapsto \delta_\mu \mathcal{U}(t, x_0, x, \mu, y), t \in [0, T]\}$ is relatively compact in $\mathcal{C}^{\mathbb{F}, \mathbb{F}-1}(\mathbb{T}^d \times \mathbb{T}^d)$. Any limits $\Phi^{0, x_0} \in \mathbb{R}^d$, $\Phi^{x_0}, \Phi^{0,\mu}: \mathbb{T}^d \mapsto \mathbb{R}$ and $\Phi^{\mu}: T^d \times \mathbb{T}^d \mapsto \mathbb{R}$ obtained by letting t tend to some $t_0 \in [0, T]$ in [\(5.47\)](#page-68-1) and [\(5.48\)](#page-68-2) must satisfy

$$
\mathcal{U}(t_0, \hat{x}_0, \cdot, \hat{\mu}) - \mathcal{U}(t_0, x_0, \cdot, \mu)
$$

= $\Phi^{x_0}(\cdot) \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \Phi^{\mu}(\cdot, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)) + O(|\hat{x}_0 - x_0|^2 + \mathbb{W}_1^2(\hat{\mu}, \mu)),$

² Let $r, s > 0$ with $r, s \notin \mathbb{N}$. If a function $h : (x, y) \in \mathbb{T}^d \times \mathbb{T}^d \mapsto h(x, y)$ has mixed derivatives up to the order $\lfloor r \rfloor$ in x and $\lfloor s \rfloor$ in y, we set

$$
\|g\|_{\lfloor r \rfloor, \lfloor s \rfloor} := \sup_{k=0, \cdots, \lfloor r \rfloor} \sup_{l=0, \cdots \lfloor s \rfloor} \sup_{x, y \in \mathbb{T}^d} |\nabla_x^k \nabla_y^l g(x, y)|,
$$

and, if $\nabla_x^{\lfloor r \rfloor} \nabla_y^{\lfloor s \rfloor} h$ is $(r - \lfloor r \rfloor, s - \lfloor s \rfloor)$ -Hölder continuous, we also set

$$
\|g\|_{(r,s)} := \|g\|_{\lfloor r \rfloor, \lfloor s \rfloor} + \sup_{(x,y),(x',y') \in \mathbb{T}^d \times \mathbb{T}^d : (x,y) \neq (x',y')} \frac{|\nabla_x^{\lfloor r \rfloor} \nabla_y^{\lfloor s \rfloor} g(x,y) - \nabla_x^{\lfloor r \rfloor} \nabla_y^{\lfloor s \rfloor} g(x',y')|}{|x - x'|^{r - \lfloor r \rfloor} + |y - y'|^{s - \lfloor s \rfloor}}.
$$

where the equality holds true in $C^{r}(\mathbb{T}^{d})$, and

$$
\mathcal{U}^{0}(t_{0}, \hat{x}_{0}, \hat{\mu}) - \mathcal{U}^{0}(t_{0}, x_{0}, \mu)
$$

= $\Phi^{0,x_{0}} \cdot (\hat{x}_{0} - x_{0}) + \int_{\mathbb{T}^{d}} \Phi^{0,\mu}(y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)) + O(|\hat{x}_{0} - x_{0}|^{2} + \mathbb{W}_{1}^{2}(\hat{\mu}, \mu)).$

The last two displays show that

$$
\nabla_{x_0} \mathcal{U}(t_0, x_0, x, \mu) \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \delta_\mu \mathcal{U}(t_0, x_0, x, \mu, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y))
$$

= $\Phi^{x_0}(x) \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \Phi^\mu(x, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)), \quad x \in \mathbb{T}^d,$

and

$$
\nabla_{x_0} \mathcal{U}^0(t, x_0, \mu) \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \delta_\mu \mathcal{U}^0(t, x_0, \mu, y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y))
$$

= $\Phi^{0, x_0} \cdot (\hat{x}_0 - x_0) + \int_{\mathbb{T}^d} \Phi^{0, \mu}(y) (\hat{\mu}(\mathrm{d}y) - \mu(\mathrm{d}y)),$

which implies that

$$
\nabla_{x_0} \mathcal{U}(t_0, x_0, x, \mu) = \Phi^{x_0}(x), \quad \delta_\mu \mathcal{U}(t_0, x_0, x, \mu, y) = \Phi^\mu(x, y),
$$

and

$$
\nabla_{x_0} \mathcal{U}^0(t_0, x_0, \mu) = \Phi^{0, x_0}, \quad \delta_{\mu} \mathcal{U}^0(t_0, x_0, \mu, y) = \Phi^{0, \mu}(y).
$$

This proves (5.45) and (5.46) .

We now refine Proposition [5.11.](#page-68-5) (using the same notation as therein for the norm $\|\cdot\|_{r,r-1}$, see footnote [2\)](#page-68-0):

Proposition 5.12. Let $r \in (\lfloor s \rfloor, s)$. Under Assumption (\hat{A}), there exist a constant $\mathfrak{C} > 0$, only depending on the parameters d, \mathfrak{L} , κ , ζ , σ_0 and $\sin A$ assumption (\hat{A}) and on \mathfrak{r} , and a constant $C > 0$, only depending on the parameters in Assumption (A) and on \mathbb{r} , such that, for $T \leq \mathfrak{C}$, for all $(t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ and $h > 0$ such that $t + h \leq T$,

$$
\|\nabla_{x_0}\mathcal{U}(t+h,x_0,\cdot,\mu)-\nabla_{x_0}\mathcal{U}(t,x_0,\cdot,\mu)\|_{\mathbb{F}}+\left\|\delta_\mu\mathcal{U}(t+h,x_0,\cdot,\mu,\cdot)-\delta_\mu\mathcal{U}(t,x_0,\cdot,\mu,\cdot)\right\|_{\mathbb{F},\mathbb{F}-1}
$$
\n
$$
\leq Ch^{1/2},\tag{5.49}
$$

and

$$
\left|\nabla_{x_0}\mathcal{U}^0(t+h,x_0,\mu) - \nabla_{x_0}\mathcal{U}^0(t,x_0,\mu)\right| + \left\|\delta_\mu\mathcal{U}^0(t+h,x_0,\mu,\cdot) - \delta_\mu\mathcal{U}^0(t,x_0,\mu,\cdot)\right\|_{r-1} \le Ch^{1/2}.\tag{5.50}
$$

In fact, one could have directly established Proposition [5.12,](#page-69-0) but we felt better to start with Proposition [5.11,](#page-68-5) because we use repeatedly the two properties (5.45) and (5.46) in the rest of the analysis.

Proof. We start with the proof of the second inequality in (5.49) . To do so, we recall from Lemma [5.6](#page-62-1) and Proposition [5.8](#page-64-2) that $\nabla_y^l \delta_\mu \mathcal{U}(t, x_0, x, \mu, y) = K_l^{\mu}(t, x_0, x, \mu, y)$, for $y \in \mathbb{T}^d$ and for $l \in \{0, \dots, \lfloor s \rfloor - 1\}^d$ with $|l| \leq \lfloor s \rfloor - 1$, with K^{x_0} and K_l^{μ} being defined in [\(5.34\)](#page-61-3). By the latter, we have (with the same notation as therein and under the generic notation $T \leq \mathfrak{C}$

$$
K_l^{\mu}(t, x_0, x, \mu, y) - K_l^{\mu}(t + h, x_0, x, \mu, y) = \delta u_t^{l, y}(x) - \nabla_y^l \delta_{\mu} \mathcal{U}(t + h, x_0, x, \mu, y),
$$

 \Box

and then, by recalling [\(5.32\)](#page-61-0),

$$
K_{l}^{\mu}(t, x_{0}, x, \mu, y) - K_{l}^{\mu}(t + h, x_{0}, x, \mu, y)
$$

\n
$$
= \mathbb{E}^{0} [\delta u_{t+h}^{l,y}(x)] - \nabla_{y}^{l} \delta_{\mu} \mathcal{U}(t + h, x_{0}, x, \mu, y)
$$

\n
$$
+ \mathbb{E}^{0} \int_{t}^{t+h} \left[-\frac{1}{2} \Delta_{x} \delta u_{r}^{l,y}(x) + \nabla_{p} \hat{H}(x, \nabla_{x} u_{r}(x)) \cdot \nabla_{x} \delta u_{r}^{l,y}(x) - \nabla_{x_{0}} f_{r}(X_{r}^{0}, x, \mu_{r}) \cdot \delta X_{r}^{0,l,y} \right. \tag{5.51}
$$

\n
$$
- \left(\delta_{\mu} f_{r}(X_{r}^{0}, x, \mu_{r}), \delta \mu_{r}^{l,y} \right) \Big] dr.
$$

Again, it is worth stressing that the pair of triples $(\delta X^{0,l,y}, \delta Y^{0,l,y}, \delta Z^{0,l,y})$ and $(\delta \mu^{l,y}, \delta u^{l,y}, \delta m^{l,y})$ solve the linearized system [\(5.32\)](#page-61-0) with $\Delta \mu = (-1)^{|l|} \nabla^l \delta_y \in C^{1-\mathbb{r}}(\mathbb{T}^d)$ (for the same \mathbb{r} as in the statement) and $\Delta x^0 = 0$ as initial conditions at time t, when X^0 starts from x_0 at time t and μ from μ . Moreover, using [\(5.33\)](#page-61-4), we can bound the $L^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0)$ norms of all the terms entering the integrand of [\(5.51\)](#page-70-0). As a result,

$$
\begin{aligned} &\left|\nabla_{y}^{l}\delta_{\mu}\mathcal{U}(t,x_{0},x,\mu,y)-\nabla_{y}^{l}\delta_{\mu}\mathcal{U}(t+h,x_{0},x,\mu,y)\right| \\ &\leq \mathbb{E}^{0}\Big[\Big|\nabla_{y}^{l}\delta_{\mu}\mathcal{U}(t+h,X_{t+h}^{0},x,\mu_{t+h},y)-\nabla_{y}^{l}\delta_{\mu}\mathcal{U}(t+h,x_{0},x,\mu,y)\Big|\Big] + Ch. \end{aligned} \tag{5.52}
$$

Using the forward equation in (5.1) , we have

$$
\mathbb{E}^{0}[|X_{t+h}^{0}-x_{0}|^{2}]=\mathbb{E}^{0}[|X_{t+h}^{0}-X_{t}^{0}|^{2}]\leq Ch\mathbb{E}^{0}\int_{t}^{t+h}(1+|Z_{s}^{0}|^{2})ds.
$$

By (5.3) , the right-hand side above is bounded by Ch for a possibly new value of C.

Also, using the forward equation in [\(5.2\)](#page-50-2) together with the bound [\(5.4\)](#page-51-1), we have $\mathbb{W}_1(\mu_{t+h}, \mu_t) \leq Ch$. And then, using the Lipschitz regularity of $\nabla_y^l \delta_\mu \mathcal{U}$ in the variables x_0 and μ (see [\(5.42\)](#page-67-3)), we obtain

$$
\left|\nabla_y^l \delta_\mu \mathcal{U}(t, x_0, x, \mu, y) - \nabla_y^l \delta_\mu \mathcal{U}(t + h, x_0, x, \mu, y)\right| \le Ch^{1/2},\tag{5.53}
$$

which proves the second inequality in (5.49) .

We prove the second inequality in a similar manner, using the representation $\nabla_y^l \delta_\mu U^0(t, x_0, \mu, y) = K_l^{0, \mu}(t, x_0, \mu, y) =$ $\delta Y_t^{0,l,y}$ (with the same definition as above for $\delta Y^{0,l,y}$). We have

$$
\nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t, x_{0}, \mu, y) - \nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t + h, x_{0}, \mu, y)
$$

\n
$$
= K_{l}^{0,\mu}(t, x_{0}, \mu, y) - K_{l}^{0,\mu}(t + h, x_{0}, \mu, y)
$$

\n
$$
= \mathbb{E}^{0} \left[\delta Y_{t+h}^{0,l,y} \right] - K_{l}^{0,\mu}(t + h, x_{0}, \mu, y)
$$

\n
$$
- \mathbb{E}^{0} \int_{t}^{t+h} \left[\nabla_{x_{0}} \hat{L}^{0}(X_{r}^{0}, Z_{r}^{0}) \cdot \delta X_{r}^{0} + \nabla_{p} \hat{L}^{0}(X_{r}^{0}, Z_{r}^{0}) \cdot \delta Z_{r}^{0} \right. \left. - \nabla_{x_{0}} f_{r}^{0}(X_{r}^{0}, \mu_{r}) \cdot \delta X_{r}^{0} - \left(\delta_{\mu} f_{r}^{0}(X_{r}^{0}, \mu_{r}), \delta \mu_{r} \right) \right] dt.
$$

This is the same as before except for the fact that the bound for the term δZ_r^0 in [\(5.33\)](#page-61-4) is weaker than the bound satisfied by the other terms therein. Here is how we remedy this difficulty: it suffices to say (thanks to Cauchy-Schwarz inequality) that $\mathbb{E} \int_{t}^{t+h} |\delta Z_{r}^{0}| dr \leq C h^{1/2}$. Proceeding as in the derivation of [\(5.52\)](#page-70-1), we deduce

$$
\begin{split} & \left| \nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t, x_{0}, \mu, y) - \nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t+h, x_{0}, \mu, y) \right| \\ & \leq \mathbb{E}^{0} \left[\left| \nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}, y) - \nabla_{y}^{l} \delta_{\mu} \mathcal{U}^{0}(t+h, x_{0}, \mu, y) \right| \right] + C h^{1/2}. \end{split} \tag{5.54}
$$

And then, following the proof of (5.53) , we get

$$
\left|\nabla_y^l \delta_\mu \mathcal{U}^0(t, x_0, \mu, y) - \nabla_y^l \delta_\mu \mathcal{U}^0(t + h, x_0, \mu, y)\right| \le Ch^{1/2}.
$$
\n(5.55)

This completes the proof.

 \Box

5.6. Identification of the martingale representation term

The next step is to identify the process Z^0 in [\(2.11\)](#page-8-0):

Proposition 5.13. *Under Assumption* (\hat{A})*, there exists a constant* $\mathfrak{C} > 0$ *, only depending on the parameters d,* \mathfrak{L} *,* κ *,* ζ *,* σ_0 and $\sin A$ ssumption (\hat{A}), such that, for $T \leq \mathfrak{C}$, the process Z^0 in [\(5.1\)](#page-50-3) admits the following representation:

$$
\forall t \in [0, T], \quad \mathbb{P}\left(\left\{Z_t^0 = \nabla_{x_0} \mathcal{U}^0(t, X_t^0, \mu_t)\right\}\right) = 1,
$$

and this, for any choice of initial condition (x_0, μ) *in* [\(5.1\)](#page-50-3)*.*

The result is in fact a consequence of the more general representation property (the reader can skip the demonstration on first reading):

Lemma 5.14. *Under Assumption* (\hat{A}) *, consider* $\mathfrak{C} > 0$ *as in Theorem [5.1.](#page-50-1) Then, for* $T \leq \mathfrak{C}$ *and*

- *1. for any initial condition* (x_0, μ) *in* [\(5.1\)](#page-50-3)*,*
- 2. for any continuous function V^0 : $[0,T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \to \mathbb{R}$ with the property that, for some $C_0 > 0$ and some \mathbf{r} ∈ ($\lfloor \mathbf{s} \rfloor$, \mathbf{s})*,* $\mathcal{V}^0(t, \cdot, \cdot) \in \mathscr{D}^0(C_0, \mathbf{r} - 1)$ *for all* $t \in [0, T]$ *, and*

$$
\forall (t, x_0, \mu) \in [0, T) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d),
$$

$$
\lim_{h \to 0} \left(\left| \nabla_{x_0} \mathcal{V}^0(t+h, x_0, \mu) - \nabla_{x_0} \mathcal{V}^0(t, x_0, \mu) \right| + \left\| \delta_\mu \mathcal{U}^0(t+h, x_0, \mu, \cdot) - \delta_\mu \mathcal{U}^0(t, x_0, \mu, \cdot) \right\|_{\mathbb{F}^{-1}} \right) = 0,
$$

3. for any triplet $(\mathbf{U}^0, \mathbf{V}^0, \ell^0)$ in $\mathscr{S}^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0; \mathbb{R}) \times \mathscr{H}^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0; \mathbb{R}) \times \mathscr{H}^2(\Omega^0, \mathcal{F}^0, \mathbb{P}^0; \mathbb{R})$ satisfy*ing*

$$
dU_t^0 = \ell_t^0 dt + \sigma_0 V_t^0 \cdot dB_t^0, \quad U_T^0 = \mathcal{V}^0(T, X_T^0, \mu_T), \tag{5.56}
$$

and, \mathbb{P}^0 -*almost surely,* $U_t^0 = \mathcal{V}^0(t, X_t^0, \mu_t)$,

the process V ⁰ *admits the following representation:*

$$
\forall t \in [0, T], \quad \mathbb{P}\left(\{V_t^0 = \nabla_{x_0} \mathcal{V}^0(t, X_t^0, \mu_t)\}\right) = 1.
$$

Proof. The proof is standard. Here, we take it from [\[19,](#page-86-4) Lemma 4.11]. We consider a uniform subdivision $\tau := (t_i)_{i=0,\dots,n}$ with $0 =: t_0 < t_1 < \dots < t_n := T$ of stepsize h together with a simple \mathbb{R}^d -valued adapted process $\boldsymbol{\pi} = (\pi_t)_{0 \leq t \leq T}$ of the form

$$
\pi_t = \sum_{i=0}^{n-1} \pi^{(i)} \mathbf{1}_{(t_i, t_{i+1}]}(t), \quad t \in [0, T], \tag{5.57}
$$

where, for some $K \ge 0$ and for each $i = 0, \dots, n - 1, \pi^{(i)}$ belongs to $L^{\infty}(\Omega, \mathcal{F}_{t_i}^0, \mathbb{P}^0; \mathbb{R}^d)$ with $\mathbb{P}^0(\{|\pi^{(i)}| \le K\}) = 1$. For $i = 0, \dots, n - 1$, we then have

$$
\mathbb{E}^{0}\bigg[\bigg(\int_{t_i}^{t_{i+1}} \pi_t \cdot dB_t^0\bigg) \times U_{t_{i+1}}^0\bigg] = \mathbb{E}^{0}\bigg[\big(\pi^{(i)} \cdot (B_{t_{i+1}}^0 - B_{t_i}^0)\big) \times U_{t_{i+1}}^0\bigg].
$$

First Step. Using the backward equation [\(5.56\)](#page-71-2), it is standard to prove that there exists a constant C (only depending on π through K) such that, for all $i = 0, \dots, n - 1$,

$$
\begin{split}\n&\left|\mathbb{E}^{0}\left[\left(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot \mathrm{d}B_{t}^{0}\right) \times U_{t_{i+1}}^{0}\right] - \sigma_{0} \mathbb{E}^{0}\left[\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot V_{t}^{0} \mathrm{d}t\right]\right| \\
&= \left|\mathbb{E}^{0}\left[\left(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot \mathrm{d}B_{t}^{0}\right) \times \int_{t_{i}}^{t_{i+1}} \ell_{t}^{0} \mathrm{d}t\right]\right| \\
&\leq (t_{i+1} - t_{i})^{1/2} \mathbb{E}^{0}\left[\int_{t_{i}}^{t_{i+1}} |\pi_{t}|^{2} \mathrm{d}t\right]^{1/2} \mathbb{E}^{0}\left[\int_{t_{i}}^{t_{i+1}} |\ell_{t}^{0}|^{2} \mathrm{d}t\right]^{1/2} \\
&\leq C(t_{i+1} - t_{i}) \left\{\int_{t_{i}}^{t_{i+1}} \mathbb{E}^{0}\left[|\ell_{t}^{0}|^{2}\right] \mathrm{d}t\right\}^{1/2} \leq Ch\epsilon(h),\n\end{split} \tag{5.58}
$$
where we have let

$$
\epsilon(h) := \left\{ h + \sup_{0 \le s < t < T: |t - s| \le h} \int_s^t \mathbb{E}^0 \left[|Z_r^0|^2 + |\ell_r^0|^2 \right] dr \right\}^{1/2}, \quad h > 0. \tag{5.59}
$$

Because $\int_0^T \mathbb{E}^0[|Z_r^0|^2 + |\ell_r^0|^2]dr < \infty$, we have $\lim_{h \to 0} \epsilon(h) = 0$.

Now, we observe that the left-hand side in the top line of [\(5.58\)](#page-71-0) is equal to

$$
\mathbb{E}^{0}\Bigg[\Bigg(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot dB_{t}^{0}\Bigg) \times U_{t_{i+1}}^{0}\Bigg] \n= \mathbb{E}^{0}\Bigg[\Bigg(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot dB_{t}^{0}\Bigg) \times \Big(\mathcal{V}^{0}(t_{i+1}, X_{t_{i+1}}^{0}, \mu_{t_{i+1}}) - \mathcal{V}^{0}(t_{i+1}, X_{t_{i}}^{0}, \mu_{t_{i}}) \Bigg] \n= \mathbb{E}^{0}\Bigg[\Bigg(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot dB_{t}^{0}\Bigg) \times \Bigg(\int_{0}^{1} \Bigg[\nabla_{x_{0}} \mathcal{V}^{0}\Big(t_{i+1}, rX_{t_{i+1}}^{0} + (1-r)X_{t_{i}}^{0}, r\mu_{t_{i+1}} + (1-r)\mu_{t_{i}}\Big) \cdot \Big(X_{t_{i+1}}^{0} - X_{t_{i}}^{0}\Big) \n+ \int_{\mathbb{T}^{d}} \delta_{\mu} \mathcal{V}^{0}\Big(t_{i+1}, rX_{t_{i+1}}^{0} + (1-r)X_{t_{i}}^{0}, r\mu_{t_{i+1}} + (1-r)\mu_{t_{i}}, y\Big) d\big(\mu_{t_{i+1}} - \mu_{t_{i}}\big)(y)\Bigg] dr \Bigg)\Bigg].
$$
\n(5.60)

Second Step. We first handle the term on the last line of [\(5.60\)](#page-72-0). Using the forward equation in [\(5.2\)](#page-50-0), we have, for any $r \in [0, 1],$

$$
\int_{\mathbb{T}^d} \delta_\mu \mathcal{V}^0(t_{i+1}, rX_{t_{i+1}}^0 + (1-r)X_{t_i}^0, r\mu_{t_{i+1}} + (1-r)\mu_{t_i}, y) d(\mu_{t_{i+1}} - \mu_{t_i})(y)
$$
\n
$$
= \int_{t_i}^{t_{i+1}} \int_{\mathbb{T}^d} \frac{1}{2} \Delta_y \delta_\mu \mathcal{V}^0(t_{i+1}, rX_{t_{i+1}}^0 + (1-r)X_{t_i}^0, r\mu_{t_{i+1}} + (1-r)\mu_{t_i}, y) d\mu_s(y) ds
$$
\n
$$
- \int_{t_i}^{t_{i+1}} \int_{\mathbb{T}^d} \nabla_y \delta_\mu \mathcal{V}^0(t_{i+1}, rX_{t_{i+1}}^0 + (1-r)X_{t_i}^0, r\mu_{t_{i+1}} + (1-r)\mu_{t_i}, y) \cdot \nabla_p \hat{H}(y, \nabla_x u_s(y)) d\mu_s(y) ds.
$$

By item 2 in the statement, the two functions $\Delta_y \delta_\mu V^0$ and $\nabla_y \delta_\mu V^0$ are bounded. Therefore, we can modify the value of C such that, for all $i = 0, \dots, n - 1$, with probability 1 under \mathbb{P}^0 ,

$$
\left| \int_{\mathbb{T}^d} \delta_\mu \mathcal{V}^0(t_{i+1}, rX_{t_{i+1}}^0 + (1-r)X_{t_i}^0, r\mu_{t_{i+1}} + (1-r)\mu_{t_i}, y) d(\mu_{t_{i+1}} - \mu_{t_i})(y) \right| \le C(t_{i+1} - t_i). \tag{5.61}
$$

And then, by Cauchy-Schwarz' inequality (and recalling [\(5.59\)](#page-72-1)),

$$
\left| \mathbb{E}^{0} \left[\left(\int_{t_{i}}^{t_{i+1}} \pi_{t} \cdot d B_{t}^{0} \right) \times \left(\int_{0}^{1} \left[\int_{\mathbb{T}^{d}} \delta_{\mu} \mathcal{V}^{0} \left(t_{i+1}, r X_{t_{i+1}}^{0} + (1-r) X_{t_{i}}^{0}, r \mu_{t_{i+1}} + (1-r) \mu_{t_{i}}, y \right) d(\mu_{t_{i+1}} - \mu_{t_{i}})(y) \right] dr \right) \right] \qquad (5.62)
$$

$$
\leq C (t_{i+1} - t_{i})^{3/2} \leq C h \epsilon(h).
$$

Third Step. We now address the term on the third line of [\(5.60\)](#page-72-0). Proceeding as in the derivation of [\(5.61\)](#page-72-2), we have, for all $i = 0, \dots, n - 1$, with probability 1 under \mathbb{P}^0 ,

$$
\mathbb{W}_1(\mu_{t_i}, \mu_{t_{i+1}}) \leq C (t_{i+1} - t_i)^{1/2}.
$$

Therefore, thanks to the boundedness and Lipschitz regularity of $\nabla_{x_0} \mathcal{V}^0$ (see item 2 in the statement), we get (recalling again [\(5.59\)](#page-72-1))

$$
\left| \mathbb{E}^{0} \left[\int_{0}^{1} \left[\nabla_{x_{0}} \mathcal{V}^{0} \left(t_{i+1}, r X_{t_{i+1}}^{0} + (1-r) X_{t_{i}}^{0}, r \mu_{t_{i+1}} + (1-r) \mu_{t_{i}} \right) \cdot \left(X_{t_{i+1}}^{0} - X_{t_{i}}^{0} \right) \right] dr \right. \\
\left. - \sigma_{0} \nabla_{x_{0}} \mathcal{V}^{0} \left(t_{i+1}, X_{t_{i}}^{0}, \mu_{t_{i}} \right) \cdot \left(B_{t_{i+1}}^{0} - B_{t_{i}}^{0} \right) \right|^{2} \right]^{1/2} \\
\leq C (t_{i+1} - t_{i})^{1/2} \epsilon (t_{i+1} - t_{i}) \\
+ \sigma_{0} \mathbb{E}^{0} \left[\left| \int_{0}^{1} \left[\nabla_{x_{0}} \mathcal{V}^{0} \left(t_{i+1}, r X_{t_{i+1}}^{0} + (1-r) X_{t_{i}}^{0}, r \mu_{t_{i+1}} + (1-r) \mu_{t_{i}} \right) \cdot \left(B_{t_{i+1}}^{0} - B_{t_{i}}^{0} \right) \right] dr \right. \\
\left. - \nabla_{x_{0}} \mathcal{V}^{0} \left(t_{i+1}, X_{t_{i}}^{0}, \mu_{t_{i}} \right) \cdot \left(B_{t_{i+1}}^{0} - B_{t_{i}}^{0} \right) \right|^{2} \right]^{1/2} \\
\leq C (t_{i+1} - t_{i})^{1/2} \epsilon (t_{i+1} - t_{i}) \tag{5.63}
$$

$$
\leq C(t_{i+1} - t_i)^{1/2} \epsilon(t_{i+1} - t_i)
$$

+ $C\mathbb{E}^0 \Biggl[\Biggl\{ 1 \wedge \Bigl(\big| X_{t_{i+1}}^0 - X_{t_i}^0 \big|^2 + \mathbb{W}_1 \bigl(\mu_{t_i}, \mu_{t_{i+1}} \bigr)^2 \Bigr) \Biggr\} |B_{t_{i+1}}^0 - B_{t_i}^0|^2 \Biggr]^{1/2}$

$$
\leq C(t_{i+1} - t_i)^{1/2} \epsilon(t_{i+1} - t_i) + C(t_{i+1} - t_i)^{1/2} \mathbb{E}^0 \Biggl[1 \wedge \Bigl((t_{i+1} - t_i) \int_{t_i}^{t_{i+1}} |Z_r^0|^2 dr \Bigr)^2 \Biggr]^{1/4}
$$

$$
\leq C(t_{i+1} - t_i)^{1/2} \epsilon(t_{i+1} - t_i) + C(t_{i+1} - t_i)^{1/2} \mathbb{E}^0 \Biggl[1 \wedge \Bigl((t_{i+1} - t_i) \int_{t_i}^{t_{i+1}} |Z_r^0|^2 dr \Bigr) \Biggr]^{1/4}
$$

$$
\leq C(t_{i+1} - t_i)^{1/2} \epsilon(t_{i+1} - t_i) \leq C h^{1/2} \epsilon(h),
$$

with the last line following from Young's inequality.

Fourth Step. Finally, by combining [\(5.58\)](#page-71-0),[\(5.60\)](#page-72-0), [\(5.62\)](#page-72-3) and [\(5.63\)](#page-73-0).

$$
\left|\mathbb{E}^0\bigg[\int_{t_i}^{t_{i+1}} \pi_s \cdot \left(\sigma_0 V_s^0\right) \mathrm{d}s - \int_{t_i}^{t_{i+1}} \pi_s \cdot \left(\sigma_0 \nabla_{x_0} \mathcal{V}^0(t_{i+1}, X_{t_i}^0, \mu_{t_i})\right) \mathrm{d}s\bigg]\right| \leq Ch \epsilon(h),
$$

which yields

$$
\lim_{h \to 0} \left| \mathbb{E}^{0} \left[\int_{0}^{T} \pi_{s} \cdot \left(\sigma_{0} V_{s}^{0} \right) \mathrm{d}s - \int_{0}^{T} \pi_{s} \cdot \left(\sigma_{0} \nabla_{x_{0}} \mathcal{V}^{0} \left(t^{+}(\tau, s), X_{t^{-}(\tau, s)}^{0}, \mu_{t^{-}(\tau, s)} \right) \right) \mathrm{d}s \right] \right| \leq C \epsilon(h), \tag{5.64}
$$

with

$$
t^+(\tau,s) := \sum_{i=0}^{n-1} t_{i+1} \mathbf{1}_{(t_i,t_{i+1}]}(s), \quad t^-(\tau,s) := \sum_{i=0}^{n-1} t_i \mathbf{1}_{[t_i,t_{i+1}]}(s), \quad s \in [0,T].
$$

We observe that, as the stepsize h of τ tends to $0, t^+(\tau, s)$ and $t^-(\tau, s)$ tend to s.

By boundedness and continuity of ∇_{x_0} \mathcal{V}^0 (see item 2) and by Lebesgue's dominated convergence theorem, it is straightforward to prove that

$$
\lim_{h \to 0} \mathbb{E}^{0} \left[\int_{0}^{T} \left| \nabla_{x_{0}} \mathcal{V}^{0}(s, X_{s}^{0}, \mu_{s}) - \nabla_{x_{0}} \mathcal{V}^{0}(t^{+}(\tau, s), X_{t^{-}(\tau, s)}^{0}, \mu_{t^{-}(\tau, s)}) \right| ds \right] = 0.
$$
\n(5.65)

It remains to choose π as a bounded simple process (as in [\(5.57\)](#page-71-1)) but with respect to a fixed partition τ^0 := $(t_i^0)_{i=0,\dots,n_0}$, with $0 =: t_0^0 < t_1^0 < \dots < t_{n_0} := T$. The stepsize of τ^0 is denoted h_0 and is kept fixed. We then refine the partition τ^0 considering another partition $\tau \supset \tau^0$. The stepsize of τ is denoted h and is sent to zero in the next lines. The main observation is that π is also a bounded simple process with respect to the partition τ . This makes it possible to apply (5.64) and (5.65) , from which we deduce (sending h to 0) that

$$
\mathbb{E}^{0}\bigg[\int_{0}^{T} \pi_{t} \cdot \Big(\sigma_{0} \Big[V_{t}^{0} - \nabla_{x_{0}} \mathcal{V}^{0} \big(t, X_{t}, \mu_{t} \big) \Big] \Big) dt \bigg] = 0.
$$

The above is true for any bounded simple process π . By a density argument, it is true for any bounded progressively measurable process, from which we deduce that, for Leb $\times \mathbb{P}^0$ -almost every (t, ω^0) ,

$$
\sigma_0 \Big[V_t^0 - \nabla_{x_0} \mathcal{V}^0(t, X_t, \mu_t) \Big] = 0.
$$

This completes the proof.

5.7. Derivation of the master equations

We now address the two PDEs (called master equations) that are satisfied by \mathcal{U}^0 and \mathcal{U} . The derivation of the master equation for \mathcal{U}^0 relies on the following formula (which is very much in the spirit of a Kolmogorov formula):

Proposition 5.15. *Under Assumption* (\hat{A})*, there exists a constant* $\mathfrak{C} > 0$ *, only depending on the parameters* d*,* κ *,* \mathfrak{L} *,* ζ , σ_0 and s in Assumption (\hat{A}) such that, for $T \leq \mathfrak{C}$, we have the following representation of $\mathcal{U}^0(t_0, x_0, \mu)$, for any $(t_0, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$:

$$
\mathcal{U}^{0}(t_{0},x_{0},\mu) = \mathbb{E}^{0}\left[g^{0}(\tilde{X}_{T}^{0},\mu)\right] - \mathbb{E}^{0}\left[\frac{1}{2}\int_{t_{0}}^{T}\int_{\mathbb{T}^{d}}\Delta_{y}\delta_{\mu}\mathcal{U}^{0}(s,\tilde{X}_{s}^{0},\mu,y)d\mu(y)ds\right] \n+ \mathbb{E}^{0}\left[\int_{t_{0}}^{T}\int_{\mathbb{T}^{d}}\nabla_{y}\delta_{\mu}\mathcal{U}^{0}(s,\tilde{X}_{s}^{0},\mu,y)\cdot\nabla_{p}\hat{H}(y,\nabla_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},y,\mu))d\mu(y)ds\right] \n- \mathbb{E}^{0}\left[\int_{t_{0}}^{T}\left(f_{s}^{0}(\tilde{X}_{s}^{0},\mu)+\hat{L}^{0}(\tilde{X}_{s}^{0},\nabla_{x_{0}}\mathcal{U}^{0}(s,\tilde{X}_{s}^{0},\mu))\right)ds\right],
$$
\n(5.66)

where $\tilde{\boldsymbol{X}}^0=(\tilde{X}^0_t)_{t_0\leq t\leq T}$ solves the SDE

$$
d\tilde{X}_t^0 = -\nabla_p H^0(\tilde{X}_t^0, \nabla_{x_0} \mathcal{U}^0(t, \tilde{X}_t^0, \mu))dt + dB_t^0, \quad t \in [t_0, T]; \quad \tilde{X}_{t_0}^0 = x_0.
$$
\n(5.67)

Pay attention that μ is fixed in [\(5.67\)](#page-74-0) (equivalently, (5.67) is not a McKean-Vlasov equation).

Proof. Throughout the proof, T is implicitly taken less than $\mathfrak C$ so that all the results already proved in this section can be freely applied.

For simplicity, we establish the representation at time $t_0 = 0$. Thanks to the regularity of $\nabla_{x_0} \mathcal{U}^0$ (see Proposition [5.10\)](#page-67-0) the equation (5.67) has a unique solution.

First Step. We fix $t \in [0, T]$. With \tilde{X}^0 as in the statement, we consider the solution $(X^0, \mu) = (X^0_s, \mu_s)_{t \le s \le T}$ to the two forward equations in [\(5.1\)](#page-50-1) and [\(5.2\)](#page-50-0) with $X_t^0 = \tilde{X}_t^0$ and $\mu_t = \mu$ as initial condition at time t. By Proposition [5.13,](#page-71-2) the equation for X^0 can be rewritten as

$$
dX_s^0 = -\nabla_p H^0(X_s^0, \nabla_{x_0} \mathcal{U}^0(s, X_s^0, \mu_s)) ds + dB_s^0, \quad s \in [t, T]; \quad X_t^0 = \tilde{X}_t^0.
$$
\n(5.68)

In particular, by Lipschitz regularity of $\nabla_{x_0} U^0$ and from the identity $\mu_t = \mu$, we obtain, for a fixed (small) $h > 0$,

$$
\sup_{t \le s \le t+h} |\tilde{X}_s^0 - X_s^0| \le C \int_t^{t+h} \mathbb{W}_1(\mu_s, \mu_t) ds \le C h^{3/2}.
$$
\n(5.69)

Now, by [\(5.69\)](#page-74-1),

$$
\mathbb{E}^{0}[U^{0}(t+h, \tilde{X}_{t+h}^{0}, \mu) | \mathcal{F}_{t}^{0}] = \mathbb{E}^{0}[U^{0}(t+h, \tilde{X}_{t+h}^{0}, \mu_{t}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
= \mathbb{E}^{0}[U^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
+ \mathbb{E}^{0}[U^{0}(t+h, \tilde{X}_{t+h}^{0}, \mu_{t}) - U^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
= \mathbb{E}^{0}[U^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
+ \mathbb{E}^{0}[U^{0}(t+h, X_{t+h}^{0}, \mu_{t}) - U^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) | \mathcal{F}_{t}^{0}] + O(h^{3/2}),
$$
\n(5.70)

 \Box

where $|O(h^{3/2})| \leq C h^{3/2}$.

Second Step. We handle the term on the last line of (5.70) . Using the regularity of U^0 in the third argument together with the equation for μ in [\(5.2\)](#page-50-0), we have (the reader may compare the expansion below with [\[19,](#page-86-0) Theorem 5.99]):

$$
\mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) - \mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{t})
$$
\n
$$
= \frac{1}{2} \int_{t}^{t+h} \int_{\mathbb{T}^{d}} \Delta_{y} \delta_{\mu} \mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{s}, y) d\mu_{s}(y) ds
$$
\n
$$
- \int_{t}^{t+h} \int_{\mathbb{T}^{d}} \nabla_{y} \delta_{\mu} \mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{s}, y) \cdot \nabla_{p} \hat{H}(y, \nabla_{x} \mathcal{U}(s, X_{s}^{0}, y, \mu_{s})) d\mu_{s}(y) ds.
$$

And then, using [\(5.69\)](#page-74-1) together with the Lipschitz regularity of $\Delta_y \delta_\mu U^0$, $\nabla_y \delta_\mu U^0$ and $\nabla_x U$ in the variables x_0 and μ and the Hölder regularity of the same functions in y (see (5.42)), we deduce that

$$
\mathbb{E}^{0}\left[\mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) - \mathcal{U}^{0}(t+h, X_{t+h}^{0}, \mu_{t}) | \mathcal{F}_{t}^{0}\right]
$$
\n
$$
= \mathbb{E}^{0}\left[\frac{1}{2} \int_{t}^{t+h} \int_{\mathbb{T}^{d}} \Delta_{y} \delta_{\mu} \mathcal{U}^{0}(t+h, \tilde{X}_{s}^{0}, \mu_{t}, y) d\mu_{t}(y) ds - \int_{t}^{t+h} \int_{\mathbb{T}^{d}} \nabla_{y} \delta_{\mu} \mathcal{U}^{0}(t+h, \tilde{X}_{s}^{0}, \mu_{t}, y) \cdot \nabla_{p} \hat{H}\left(y, \nabla_{x} \mathcal{U}(s, \tilde{X}_{s}^{0}, y, \mu_{t})\right) d\mu_{t}(y) ds | \mathcal{F}_{t}^{0}\right] + O\left(h^{1 + (\mathbb{r} - \lfloor s \rfloor)/2}\right).
$$
\n(5.71)

We then handle the term on the penultimate line of (5.70) . Using the backward equation in (5.1) , we have

$$
\mathbb{E}^{0}[U^{0}(t+h, X_{t+h}^{0}, \mu_{t+h}) | \mathcal{F}_{t}^{0}] = \mathbb{E}^{0}[Y_{t+h}^{0} | \mathcal{F}_{t}^{0}]
$$
\n
$$
= Y_{t}^{0} - \mathbb{E}^{0}\bigg[\int_{t}^{t+h} \Big(f_{s}^{0}(X_{s}^{0}, \mu_{s}) + \hat{L}^{0}(X_{s}^{0}, \nabla_{x_{0}}U^{0}(s, X_{s}^{0}, \mu_{s}))\Big)ds | \mathcal{F}_{t}^{0}\bigg]
$$
\n
$$
= U^{0}(t, X_{t}^{0}, \mu_{t}) - \mathbb{E}^{0}\bigg[\int_{t}^{t+h} \Big(f_{s}^{0}(\tilde{X}_{s}^{0}, \mu_{t}) + \hat{L}^{0}(\tilde{X}_{s}^{0}, \nabla_{x_{0}}U^{0}(s, \tilde{X}_{s}^{0}, \mu_{t}))\Big)ds | \mathcal{F}_{t}^{0}\bigg] + O(h^{3/2}).
$$
\n(5.72)

Third Step. We now insert [\(5.71\)](#page-75-0) and [\(5.72\)](#page-75-1) into [\(5.70\)](#page-74-2). Recalling $X_t^0 = \tilde{X}_t^0$ and $\mu_t = \mu$, we obtain

$$
\mathbb{E}^{0}\left[\mathcal{U}^{0}(t+h,\tilde{X}_{t+h}^{0},\mu)\,|\,\mathcal{F}_{t}^{0}\right] = \mathcal{U}^{0}(t,\tilde{X}_{t}^{0},\mu)
$$
\n
$$
-\mathbb{E}^{0}\left[\frac{1}{2}\int_{t}^{t+h}\int_{\mathbb{T}^{d}}\Delta_{y}\delta_{\mu}\mathcal{U}^{0}(t+h,\tilde{X}_{s}^{0},\mu,y)d\mu(y)ds\,|\,\mathcal{F}_{t}^{0}\right]
$$
\n
$$
+\mathbb{E}^{0}\left[\int_{t}^{t+h}\int_{\mathbb{T}^{d}}\nabla_{y}\delta_{\mu}\mathcal{U}^{0}(t+h,\tilde{X}_{s}^{0},\mu,y)\cdot\nabla_{p}\hat{H}\left(y,\nabla_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},y,\mu)\right)d\mu(y)ds\,|\,\mathcal{F}_{t}^{0}\right]
$$
\n
$$
-\mathbb{E}^{0}\left[\int_{t}^{t+h}\left(f_{s}^{0}(\tilde{X}_{s}^{0},\mu)+\hat{L}^{0}(\tilde{X}_{s}^{0},\nabla_{x_{0}}\mathcal{U}^{0}(s,\tilde{X}_{s}^{0},\mu))\right)ds\,|\,\mathcal{F}_{t}^{0}\right] + O\left(h^{1+(r-\lfloor s\rfloor)/2}\right).
$$
\n(5.73)

We are now given a subdivision $\tau = (t_i)_{i=0,\dots,n}$ of $[0,T]$, with $0 = t_0 < t_1 < \dots < t_n = T$ and with stepsize h. Then, [\(5.73\)](#page-75-2) says

$$
\mathbb{E}^{0}[{\mathcal{U}}^{0}(t_{i+1}, \tilde{X}_{t_{i+1}}^{0}, \mu)] = \mathbb{E}^{0}[{\mathcal{U}}^{0}(t_{i}, \tilde{X}_{t_{i}}^{0}, \mu)] \n- \mathbb{E}^{0}\bigg[\frac{1}{2}\int_{t_{i}}^{t_{i+1}} \int_{\mathbb{T}^{d}} \Delta_{y}\delta_{\mu}{\mathcal{U}}^{0}(t_{i+1}, \tilde{X}_{s}^{0}, \mu, y) d\mu(y) ds\bigg] \n+ \mathbb{E}^{0}\bigg[\int_{t_{i}}^{t_{i+1}} \int_{\mathbb{T}^{d}} \nabla_{y}\delta_{\mu}{\mathcal{U}}^{0}(t_{i+1}, \tilde{X}_{s}^{0}, \mu, y) \cdot \nabla_{p}\hat{H}\big(y, \nabla_{x}{\mathcal{U}}(s, \tilde{X}_{s}^{0}, y, \mu)\big) d\mu(y) ds\bigg] \n- \mathbb{E}^{0}\bigg[\int_{t_{i}}^{t_{i+1}} \Big(f_{s}^{0}(\tilde{X}_{s}^{0}, \mu) + \hat{L}^{0}(\tilde{X}_{s}^{0}, \nabla_{x_{0}}{\mathcal{U}}^{0}(s, \tilde{X}_{s}^{0}, \mu))\Big) ds\bigg] + O\big(h^{1 + (r - \lfloor s \rfloor)/2}\big).
$$

Major Minor MFGs 77

Letting $t(\tau, s) := t_{i+1}$ if $s \in (t_i, t_{i+1}]$, for $i = 0, \ldots, n-1$, and summing over i the above expansion, we get

$$
\mathcal{U}^{0}(0,x_{0},\mu) = \mathbb{E}^{0}\left[g^{0}(\tilde{X}_{T}^{0},\mu)\right]
$$

\n
$$
-\mathbb{E}^{0}\left[\frac{1}{2}\int_{0}^{T}\int_{\mathbb{T}^{d}}\Delta_{y}\delta_{\mu}\mathcal{U}^{0}(t(\tau,s),\tilde{X}_{s}^{0},\mu,y)d\mu(y)ds\right]
$$

\n
$$
+\mathbb{E}^{0}\left[\int_{0}^{T}\int_{\mathbb{T}^{d}}\nabla_{y}\delta_{\mu}\mathcal{U}^{0}(t(\tau,s),\tilde{X}_{s}^{0},\mu,y)\cdot\nabla_{p}\hat{H}(y,\nabla_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},y,\mu))d\mu(y)ds\right]
$$

\n
$$
-\mathbb{E}^{0}\left[\int_{0}^{T}\left(f_{s}^{0}(\tilde{X}_{s}^{0},\mu)+\hat{L}^{0}(\tilde{X}_{s}^{0},\nabla_{x_{0}}\mathcal{U}^{0}(s,\tilde{X}_{s}^{0},\mu))\right)ds\right] + O\left(h^{(\mathbb{r}-\lfloor s\rfloor)/2}\right).
$$

Using the boundedness and time continuity of $\Delta_y \delta_\mu U^0$ and $\nabla_y \delta_\mu U^0$ (recall in particular Proposition [5.12\)](#page-69-0) and observing that $\lim_{h\to 0} t(\tau, s) = s$, we can easily let h tend to 0 in the formula. By Lebesgue's dominated convergence theorem, we get the expected formula. \Box

We have a similar result for U :

Proposition 5.16. *Under Assumption* (\hat{A})*, there exists a constant* $\mathfrak{C} > 0$ *, only depending on the parameters d,* κ *,* \mathfrak{L} *,* ζ , σ_0 and s in Assumption (\hat{A}) such that, for $T \leq \mathfrak{C}$, the following representation of $\mathcal{U}(t_0, x_0, \mu, x)$ holds true, for any $(t_0, x_0, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$:

$$
\mathcal{U}(t_0, x_0, x, \mu) = \mathbb{E}^0 \Big[g(\tilde{X}_T^0, x, \mu) \Big] - \mathbb{E}^0 \Big[\frac{1}{2} \int_{t_0}^T \int_{\mathbb{T}^d} \Delta_y \delta_\mu \mathcal{U}(s, \tilde{X}_s^0, x, \mu, y) d\mu(y) ds \Big] \n+ \mathbb{E}^0 \Big[\int_{t_0}^T \int_{\mathbb{T}^d} \nabla_y \delta_\mu \mathcal{U}(s, \tilde{X}_s^0, x, \mu, y) \cdot \nabla_p \hat{H}(y, \nabla_x \mathcal{U}(s, \tilde{X}_s^0, y, \mu)) d\mu(y) ds \Big] \n+ \mathbb{E}^0 \Big[\int_0^T \Big(-\frac{1}{2} \Delta_x \mathcal{U}(s, \tilde{X}_s^0, x, \mu) + \hat{H}(x, \nabla_x \mathcal{U}(s, \tilde{X}_s^0, x, \mu)) - f_s(\tilde{X}_s^0, x, \mu) \Big) ds \Big],
$$
\n(5.74)

where $\tilde{\mathbf{X}}^0 = (\tilde{X}_t^0)_{t_0 \leq t \leq T}$ solves the SDE [\(5.67\)](#page-74-0).

Proof. The proof is similar to the proof of Proposition [5.15.](#page-74-3) In particular, we just establish the representation at time $t_0 = 0.$

First Step. For a fixed $t \in [0, T]$, we reuse the notation [\(5.68\)](#page-74-4) and the bound [\(5.69\)](#page-74-1). Following [\(5.70\)](#page-74-2), we deduce from the Lipschitz regularity of U (see [\(5.27\)](#page-59-0)) that

$$
\mathbb{E}^{0}[U(t+h, \tilde{X}_{t+h}^{0}, x, \mu) | \mathcal{F}_{t}^{0}] = \mathbb{E}^{0}[U(t+h, \tilde{X}_{t+h}^{0}, x, \mu_{t}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
= \mathbb{E}^{0}[U(t+h, X_{t+h}^{0}, x, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
+ \mathbb{E}^{0}[U(t+h, \tilde{X}_{t+h}^{0}, x, \mu_{t}) - U(t+h, X_{t+h}^{0}, x, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
= \mathbb{E}^{0}[U(t+h, X_{t+h}^{0}, x, \mu_{t+h}) | \mathcal{F}_{t}^{0}]
$$
\n
$$
+ \mathbb{E}^{0}[U(t+h, X_{t+h}^{0}, x, \mu_{t}) - U(t+h, X_{t+h}^{0}, x, \mu_{t+h}) | \mathcal{F}_{t}^{0}] + O(h^{3/2}),
$$
\n(5.75)

where $|O(h^{3/2})| \leq C h^{3/2}$.

-

Second Step. We follow the second step in the proof of Proposition [5.15](#page-74-3) and address the term on the last line of [\(5.75\)](#page-76-0). Using the regularity of U in the third argument (see Proposition [5.10\)](#page-67-0) together with the equation for μ in [\(5.2\)](#page-50-0), we have

$$
\mathcal{U}(t+h, X_{t+h}^0, x, \mu_{t+h}) - \mathcal{U}(t+h, X_{t+h}^0, x, \mu_t)
$$

= $\frac{1}{2} \int_t^{t+h} \int_{\mathbb{T}^d} \Delta_y \delta_\mu \mathcal{U}^0(t+h, X_{t+h}^0, x, \mu_s, y) d\mu_s(y) ds$

$$
- \int_t^{t+h} \int_{\mathbb{T}^d} \nabla_y \delta_\mu \mathcal{U}^0(t+h, X_{t+h}^0, x, \mu_s, y) \cdot \nabla_p \hat{H}(y, \nabla_x \mathcal{U}(s, X_s^0, y, \mu_s)) d\mu_s(y) ds.
$$

And then, using [\(5.69\)](#page-74-1) together with the Lipschitz regularity of $\Delta_y \delta_\mu \mathcal{U}$, $\nabla_y \delta_\mu \mathcal{U}$ and $\nabla_x \mathcal{U}$ in the variables x_0 and μ and their Hölder regularity in y , we deduce that

$$
\mathbb{E}^{0}\left[\mathcal{U}(t+h, X_{t+h}^{0}, x, \mu_{t+h}) - \mathcal{U}(t+h, X_{t+h}^{0}, x, \mu_{t}) | \mathcal{F}_{t}^{0}\right]
$$
\n
$$
= \mathbb{E}^{0}\left[\frac{1}{2}\int_{t}^{t+h}\int_{\mathbb{T}^{d}}\Delta_{y}\delta_{\mu}\mathcal{U}(t+h, \tilde{X}_{s}^{0}, x, \mu_{t}, y)d\mu_{t}(y)ds - \int_{t}^{t+h}\int_{\mathbb{T}^{d}}\nabla_{y}\delta_{\mu}\mathcal{U}(t+h, \tilde{X}_{s}^{0}, x, \mu_{t}, y) \cdot \nabla_{p}\hat{H}(y, \nabla_{x}\mathcal{U}(s, \tilde{X}_{s}^{0}, y, \mu_{t}))d\mu_{t}(y)ds | \mathcal{F}_{t}^{0}\right] + O\left(h^{1 + (r - \lfloor s \rfloor)/2}\right).
$$
\n(5.76)

We then handle the term on the penultimate line of (5.75) . Using the backward equation in (5.2) (with the prescribed initial conditions for the two forward equations in (5.1) and (5.2)) and (5.27) again, we have

$$
\mathbb{E}^{0}[U(t+h, X_{t+h}^{0}, x, \mu_{t+h}) | \mathcal{F}_{t}^{0}] = \mathbb{E}^{0}[u_{t+h}(x) | \mathcal{F}_{t}^{0}]
$$
\n
$$
= u_{t}(x) + \mathbb{E}^{0}\left[\int_{t}^{t+h} \left(-\frac{1}{2}\Delta_{x}u_{s}(x) + \hat{H}(x, \nabla_{x}u_{s}(x)) - f_{s}(X_{s}^{0}, x, \mu_{s})\right)ds | \mathcal{F}_{t}^{0}\right]
$$
\n
$$
= U(t, X_{t}^{0}, x, \mu_{t})
$$
\n
$$
+ \mathbb{E}^{0}\left[\int_{t}^{t+h} \left(-\frac{1}{2}\Delta_{x}U(s, X_{s}^{0}, x, \mu_{s}) + \hat{H}(x, \nabla_{x}U(s, X_{s}^{0}, x, \mu_{s})) - f_{s}(X_{s}^{0}, x, \mu_{s})\right)ds | \mathcal{F}_{t}^{0}\right]
$$
\n
$$
= U(t, X_{t}^{0}, x, \mu_{t})
$$
\n
$$
+ \mathbb{E}^{0}\left[\int_{t}^{t+h} \left(-\frac{1}{2}\Delta_{x}U(s, \tilde{X}_{s}^{0}, x, \mu_{t}) + \hat{H}(x, \nabla_{x}U(s, \tilde{X}_{s}^{0}, x, \mu_{t})) - f_{s}(\tilde{X}_{s}^{0}, x, \mu_{t})\right)ds | \mathcal{F}_{t}^{0}\right] + O(h^{3/2}).
$$
\n(5.77)

Third Step. We now insert [\(5.76\)](#page-77-0) and [\(5.77\)](#page-77-1) into [\(5.75\)](#page-76-0). We obtain

$$
\mathbb{E}^{0}\left[\mathcal{U}(t+h,\tilde{X}_{t+h}^{0},x,\mu_{t})\,|\,\mathcal{F}_{t}^{0}\right] = \mathcal{U}^{0}(t,\tilde{X}_{t}^{0},x,\mu)
$$
\n
$$
-\mathbb{E}^{0}\left[\frac{1}{2}\int_{t}^{t+h}\int_{\mathbb{T}^{d}}\Delta_{y}\delta_{\mu}\mathcal{U}(t+h,\tilde{X}_{s}^{0},x,\mu,y)d\mu(y)ds\,|\,\mathcal{F}_{t}^{0}\right]
$$
\n
$$
+\mathbb{E}^{0}\left[\int_{t}^{t+h}\int_{\mathbb{T}^{d}}\nabla_{y}\delta_{\mu}\mathcal{U}(t+h,\tilde{X}_{s}^{0},x,\mu,y)\cdot\nabla_{p}\hat{H}(y,\nabla_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},y,\mu))d\mu(y)ds\,|\,\mathcal{F}_{t}^{0}\right]
$$
\n
$$
+\mathbb{E}^{0}\left[\int_{t}^{t+h}\left(-\frac{1}{2}\Delta_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},x,\mu)+\hat{H}(x,\nabla_{x}\mathcal{U}(s,\tilde{X}_{s}^{0},x,\mu))-f_{s}(\tilde{X}_{s}^{0},x,\mu)\right)ds\,|\,\mathcal{F}_{t}^{0}\right]
$$
\n
$$
+O\left(h^{1+(\mathbf{r}-\lfloor s\rfloor)/2}\right).
$$
\n(5.78)

We are now given a subdivision $\tau = (t_i)_{i=0,\dots,n}$ of $[0,T]$, with $0 = t_0 < t_1 < \dots < t_n = T$ and with stepsize h. Then, [\(5.78\)](#page-77-2) says

$$
\mathbb{E}^{0}\left[\mathcal{U}(t_{i+1}, \tilde{X}_{t_{i+1}}^{0}, x, \mu)\right] = \mathbb{E}^{0}\left[\mathcal{U}(t_{i}, \tilde{X}_{t_{i}}^{0}, x, \mu)\right] \n- \mathbb{E}^{0}\left[\frac{1}{2} \int_{t_{i}}^{t_{i+1}} \int_{\mathbb{T}^{d}} \Delta_{y} \delta_{\mu} \mathcal{U}(t_{i+1}, \tilde{X}_{s}^{0}, x, \mu, y) d\mu(y) ds\right] \n+ \mathbb{E}^{0}\left[\int_{t_{i}}^{t_{i+1}} \int_{\mathbb{T}^{d}} \nabla_{y} \delta_{\mu} \mathcal{U}(t_{i+1}, \tilde{X}_{s}^{0}, x, \mu, y) \cdot \nabla_{p} \hat{H}\left(y, \nabla_{y} \mathcal{U}(s, \tilde{X}_{s}^{0}, y, \mu)\right) d\mu(y) ds\right] \n+ \mathbb{E}^{0}\left[\int_{t_{i}}^{t_{i+1}} \left(-\frac{1}{2} \Delta_{x} \mathcal{U}(s, \tilde{X}_{s}^{0}, x, \mu) + \hat{H}\left(x, \nabla_{x} \mathcal{U}(s, \tilde{X}_{s}^{0}, x, \mu)\right) - f_{s}(\tilde{X}_{s}^{0}, x, \mu)\right) ds\right] + O\left(h^{1+(\mathbb{r}-\lfloor s \rfloor)/2}\right).
$$

Major Minor MFGs 79

Letting $t(\tau, s) := t_{i+1}$ if $s \in (t_i, t_{i+1}]$, for $i = 0, \ldots, n-1$, and summing over i the above expansion, we get

$$
\mathcal{U}(0, x_0, x, \mu) = \mathbb{E}^0 \left[g(\tilde{X}_T^0, x, \mu) \right]
$$

\n
$$
- \mathbb{E}^0 \left[\frac{1}{2} \int_0^T \int_{\mathbb{T}^d} \Delta_y \delta_\mu \mathcal{U}(t(\tau, s), \tilde{X}_s^0, x, \mu, y) d\mu(y) ds \right]
$$

\n
$$
+ \mathbb{E}^0 \left[\int_0^T \int_{\mathbb{T}^d} \nabla_y \delta_\mu \mathcal{U}(t(\tau, s), \tilde{X}_s^0, x, \mu, y) \cdot \nabla_p \hat{H}(y, \nabla_y \mathcal{U}(s, \tilde{X}_s^0, y, \mu)) d\mu(y) ds \right]
$$

\n
$$
+ \mathbb{E}^0 \left[\int_0^T \left(-\frac{1}{2} \Delta_x \mathcal{U}(s, \tilde{X}_s^0, x, \mu) + \hat{H}(x, \nabla_x \mathcal{U}(s, \tilde{X}_s^0, x, \mu)) - f_s(\tilde{X}_s^0, x, \mu) \right) ds \right] + O(h^{(r - \lfloor s \rfloor)/2}).
$$

Using the boundedness and time continuity of all the derivatives appearing above (see Lemma [5.4](#page-59-1) and Proposition [5.12\)](#page-69-0) and observing that $\lim_{h\to 0} t(\tau, s) = s$, we can easily let h tend to 0 in the formula. By Lebesgue's dominated convergence theorem, we get the expected formula. \Box

The following result makes the connection between Propositions [5.15](#page-74-3) and [5.16](#page-76-1) and the master equation.

Corollary 5.17. *Let Assumption* (Â) *be in force. Assume also that:*

- i. the coefficient $(t, x_0, \mu) \mapsto f^0_t(x_0, \mu)$ is Hölder continuous in time, uniformly in (x_0, μ) ; the coefficient $(t, x_0, x, \mu) \mapsto$ $f_t(x_0, x, \mu)$ *is Hölder continuous in time, uniformly in* (x_0, x, μ) *;*
- ii. $x_0 \mapsto g^0(x_0,\mu)$ has Hölder continuous second-order derivatives, uniformly in μ ; $x_0 \mapsto g(x_0,x,\mu)$ has Hölder con*tinuous second-order derivatives, uniformly in* (x, μ) *.*

Then, there exists a constant $\mathfrak{C} > 0$ *, only depending on the parameters* d, \mathfrak{L} , κ , ζ *,* σ_0 *and* \sin *Assumption* (\hat{A}) *such that, for* $T \leq \mathfrak{C}$ *, the system of two master equations*

$$
\partial_t V^0(t, x_0, \mu) + \frac{1}{2} \sigma_0^2 \Delta_{x_0} V^0(t, x_0, \mu) - \nabla_p H^0(x_0, \nabla_{x_0} V^0(t, x_0, \mu)) \cdot \nabla_{x_0} V^0(t, x_0, \mu) \n+ \hat{L}^0(x_0, \nabla_{x_0} V^0(t, x_0, \mu)) + f_t^0(x_0, \mu) \n+ \int_{\mathbb{T}^d} \left\{ \frac{1}{2} \text{div}_y(\partial_\mu V^0(t, x_0, \mu, y)) - \partial_\mu V^0(t, x_0, \mu, y) \cdot \nabla_p \hat{H}(y, \nabla_x V(t, x_0, y, \mu)) \right\} d\mu(y) = 0, \tag{5.79}
$$
\n
$$
V^0(T, x_0, \mu) = g^0(x_0, \mu), \tag{5.79}
$$

 $for (t, x_0, \mu) \in [0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$ *, and*

$$
\partial_t V(t, x_0, x, \mu) + \frac{1}{2} \Delta_x V(t, x_0, x, \mu) + \frac{1}{2} \sigma_0^2 \Delta_{x_0} V(t, x_0, x, \mu) - \hat{H}(x_0, x, \nabla_x V(t, x_0, x, \mu)) + f_t(x_0, x, \mu) \n- \nabla_p H^0(x_0, \nabla_{x_0} V^0(t, x_0, \mu)) \cdot \nabla_{x_0} V(t, x_0, x, \mu) \n+ \int_{\mathbb{T}^d} \left\{ \frac{1}{2} \text{div}_y(\partial_\mu V(t, x_0, x, \mu, y)) - \partial_\mu V(t, x_0, x, \mu, y) \cdot \nabla_p \hat{H}(y, \nabla_x V(t, x_0, y, \mu)) \right\} d\mu(y) = 0, \tag{5.80}
$$
\n
$$
V(T, x_0, x, \mu) = g(x_0, x, \mu), \tag{5.81}
$$

for $(t, x_0, x, \mu) \in [0, T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$, has a classical solution (V^0, V) in the sense that

- $1.$ $(t, x_0, \mu) \mapsto (\partial_t V^0(t, x_0, \mu), \nabla_{x_0} V^0(t, x_0, \mu), \nabla_{x_0}^2 V^0(t, x_0, \mu))$ is continuous on $[0, T] \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$; $(t, x_0, \mu, y) \mapsto$ $(\partial_{\mu}V^{0}(t, x_{0}, \mu, y), \nabla_{y}\partial_{\mu}V^{0}(t, x_{0}, \mu, y))$ *is continuous on* $[0, T] \times \mathbb{R}^{d} \times \mathcal{P}(\mathbb{T}^{d}) \times \mathbb{T}^{d}$;
- 2. $(t, x_0, x, \mu) \mapsto (\partial_t V(t, x_0, x, \mu), \nabla_{x_0} V(t, x_0, x, \mu), \nabla_{x_0}^2 V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu), \nabla_x^2 V(t, x_0, x, \mu))$ is con- \mathcal{L} *tinuous on* $[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d)$; $(t, x_0, x, \mu, y) \mapsto (\partial_\mu V(t, x_0, x, \mu, y), \nabla_y \partial_\mu V(t, x_0, x, \mu, y))$ *is continuous* $\mathcal{O}[0,T] \times \mathbb{R}^d \times \mathbb{T}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d$
- 3. $(t, x_0, \mu) \mapsto \nabla_{x_0} V^0(t, x_0, \mu)$ and $(t, x_0, x, \mu) \mapsto (\nabla_{x_0} V(t, x_0, x, \mu), \nabla_x V(t, x_0, x, \mu))$ are Lipschitz continuous *with respect to* (x_0, μ) *and* (x_0, x, μ) *respectively (using the distance* \mathbb{W}_1 *to handle the argument* μ *), uniformly in* t*.*

One solution is given by the master fields U^0 and U introduced in [\(5.25\)](#page-59-2) and [\(5.24\)](#page-59-3). In particular, U^0 and U have *first-order derivative in* t *and second-order derivatives in* x⁰ *and these derivatives are jointly continuous (with respect*

to all their arguments). Moreover, for any $t \in [0,T]$, the mapping $x_0 \mapsto \mathcal{U}^0(t,x_0,\mu)$ has Hölder continuous secondorder derivatives, uniformly in μ , and the mapping $x_0\mapsto\mathcal{U}^0(t,x_0,x,\mu)$ has Hölder continuous second-order derivatives, *uniformly in* (x, μ) *.*

Proof. We use Propositions [5.15](#page-74-3) and [5.16](#page-76-1) to prove that U^0 and U solve equations [\(5.79\)](#page-78-0) and [\(5.80\)](#page-78-1) respectively.

We start with [\(5.79\)](#page-78-0). We claim that all the terms entering the time integrals in the right-hand side of [\(5.66\)](#page-74-5) are Hölder continuous in the pair (s, \tilde{X}_s^0) , uniformly in the other parameters. For the term on the first line of [\(5.66\)](#page-74-5), this follows from the fact that, for all $t \in [0,T]$, $\mathcal{U}^0(t,\cdot,\cdot)$ belongs to $\mathcal{D}^0(C,2)$ for some $C \ge 0$ (see Proposition [5.10\)](#page-67-0) and that its derivatives are Hölder continuous in time (see Proposition [5.12\)](#page-69-0). For the second line of [\(5.66\)](#page-74-5), the regularity of the term $\nabla_y \delta_\mu \mathcal{U}^0(s, \tilde{X}_s^0, \mu, y)$ follows from the same argument. The regularity of the term $\nabla_x \mathcal{U}(s, \tilde{X}_s^0, y, \mu)$ also follows from the fact that $U(t, \cdot, \cdot, \cdot)$ belongs to $\mathscr{D}(C, 1, 2)$ for all $t \in [0, T]$, and that its derivatives in x are Hölder continuous in time (see Lemma [5.4\)](#page-59-1). The argument is the same for the terms on the third line of [\(5.66\)](#page-74-5).

Moreover, we observe that the drift of the SDE [\(5.67\)](#page-74-0) is also Hölder continuous in the arguments (t, \tilde{X}_t^0) .

And then we can interpret [\(5.66\)](#page-74-5) as the Kolmogorov formula for representing $U^0(t_0, x_0, \mu)$ as the solution to a secondorder PDE, denoted (E^{μ}) , in the variables $(t_0, x_0) \in [0, T] \times \mathbb{R}^d$. This second-order PDE (E^{μ}) is driven by the generator of [\(5.67\)](#page-74-0). It has $g^0(\cdot,\mu)$ as terminal condition and the remaining three terms in the right-hand side of [\(5.66\)](#page-74-5) as source terms. By standard Schauder's theory for uniformly parabolic PDEs with Hölder continuous coefficients, the PDE (E^{μ}) has a classical solution with second Hölder derivatives that are Hölder continuous in time and space. This says two things. First, for a fixed μ , $(t_0, x_0) \mapsto \mathcal{U}^0(t_0, x_0, \mu)$ solves the PDE (E^{μ}) . By expanding all the terms, one easily recovers the master equation [\(5.79\)](#page-78-0). In particular, U^0 has a first-order derivative in time and second-order derivatives in x_0 . Second, those derivatives are jointly continuous in (t, x_0, μ) (the proof is given a few lines below). In particular, \mathcal{U}^0 satisfies items 1 and 3 in the statement (properties of the first order derivatives of \mathcal{U}^0 in x_0 and μ follow directly from Propositions [5.10](#page-67-0) and [5.12\)](#page-69-0). Joint regularity of $\partial_t \mathcal{U}^0$ and $\nabla_{x_0}^2 \mathcal{U}^0$ in (t, x_0, μ) follows from Schauder's theory again: the (t, x_0) -Hölder norms of $\partial_t \mathcal{U}^0$ and $\nabla_{x_0}^2 \mathcal{U}^0$ can be bounded independently of μ ; therefore, the latter two derivatives are locally compact (in the space of Hölder functions) when μ varies; using continuity of the coefficients of (E^{μ}) with respect to the argument μ , one easily deduces that $\partial_t \mathcal{U}^0$ and $\nabla_{x_0}^2 \mathcal{U}^0$ are indeed jointly continuous (using in particular the fact that $\mathcal{U}^0(t, \cdot, \cdot) \in \mathscr{D}^0(C, 2 + (\mathbb{S} - \lfloor \mathbb{S} \rfloor)/2)$ for some $C \ge 0$).

Thanks to Proposition [5.16,](#page-76-1) one can proceed in a similar way to prove that U solves the master equation [\(5.80\)](#page-78-1) and satisfies the properties claimed in the statement.

5.8. Back to the original coefficients

We now want to come back to the original system (2.11) – (2.12) . To do this, we use the following lemma:

Lemma 5.18. Assume that L^0 and L are as in Assumption (A) and define the corresponding Hamiltonians H^0 and H. *Then, for any given* $R > 0$, there exist two coefficients \hat{L}^0 and \hat{H} satisfying Assumption (\hat{A}) with respect to a new constant κ *and* a function ζ *only depending on the parameters* κ *and* λ *in* (**A**)*, such that, for any* $(x, x_0, p) \in \mathbb{T}^d \times \mathbb{R}^d \times \mathbb{R}^d$ *, with* $|p| \leq R$,

$$
\hat{H}(x,p) = H(x,p), \quad \hat{L}^0(x_0,p) = L^0(x_0, -\nabla_p H^0(x_0,p)).
$$
\n(5.81)

Proof. Without any loss of generality, we can assume $R > 2$. We then introduce two auxiliary functions. We call ζ a smooth cut-off function from \mathbb{R}^d to $[0,1]$ such that $\zeta(p) = 1$ for $|p| \le R$, $\zeta(p) = 0$ for $|p| \ge R^2$, and $|\nabla_p \zeta(p)| \le c/R^2$. and $|\nabla^2_{pp}\zeta(p)| \le c/R^4$ for any p, where $c > 0$ is a universal constant. We call φ an even smooth function from $\mathbb R$ to $\mathbb R$ such that φ'' is non-decreasing on $[0, R^2]$ and non-increasing on $[R^2, +\infty)$ with $\varphi''(r) = 1$ for $r \in [0, R]$, $\varphi''(r) = r/R$ for $r \in [2R, R^2]$, and $\varphi''(r) = 2R^5/r^2$ for $r \geq 2R^2$. Assuming that $\varphi(0) = 0$ and $\varphi'(0) = 0$, we have $\varphi(r) = r^2/2$ for $r \in [-R, R], \varphi'(r) \ge r^2/(2R)$ for $r \in [2R, R^2]$ and $\varphi'(r) \in (0, 4R^3]$ for all $r \in \mathbb{R} \setminus \{0\}$. We then write $\Phi(p)$ for $\varphi(|p|)$. The function Φ is smooth on \mathbb{R}^d and

$$
\nabla_{pp}^2 \Phi(p) = \frac{\varphi'(|p|)}{|p|} \Big[1 - \frac{p}{|p|} \otimes \frac{p}{|p|} \Big] + \varphi''(|p|) \frac{p}{|p|} \otimes \frac{p}{|p|}, \quad p \in \mathbb{R}^d \setminus \{0\}. \tag{5.82}
$$

In particular, $\nabla^2_{pp}\Phi(p) = I_d$ for $|p| \le R$. For $|p| \in [R, R^2]$, $\nabla^2_{pp}\Phi(p) \ge [|p|/(2R)]I_d$ (in the sense of comparison of symmetric matrices) and, for $|p| > R^2$, $\nabla^2_{pp} \Phi(p)$ is positive definite.

With λ' as in (A5), we consider

$$
\hat{H}(x,p) := \left(H(x,p) - \frac{\lambda'}{4}|p|^2\right)\zeta(p) + \frac{\lambda'}{2}\Phi(p).
$$

 \Box

For $|p| \leq R$,

$$
\hat{H}(x,p) = H(x,p) - \frac{\lambda'}{4}|p|^2 + \frac{\lambda'}{4}|p|^2 = H(x,p).
$$

Moreover,

$$
\nabla_{pp}^{2} \hat{H}(x,p) = \left(\nabla_{pp}^{2} H(x,p) - \frac{\lambda'}{2} I_d\right) \zeta(p) + \left(\nabla_{p} H(x,p) - \frac{\lambda'}{2} p\right) \otimes \nabla \zeta(p) + \nabla \zeta(p) \otimes \left(\nabla_{p} H(x,p) - \frac{\lambda'}{2} p\right) + \left(H(x,p) - \frac{\lambda'}{4}|p|^{2}\right) \nabla_{pp}^{2} \zeta(p) + \frac{\lambda'}{2} \nabla_{pp}^{2} \Phi(p).
$$

And then, using (A5), we know that (in the sense of comparison of symmetric matrices)

$$
\nabla_{pp}^2 \hat{H}(x,p) \geq -C\Big(|p| \times \frac{1}{R^2} + |p|^2 \times \frac{1}{R^4}\Big) {\bf 1}_{\{|p| \in [R,R^2]\}} I_d + \frac{\lambda'}{2} \nabla_{pp}^2 \Phi(p),
$$

for a constant C only depending on the properties of H. And then, by [\(5.82\)](#page-79-0), we get, for $|p| \in [R, R^2]$, $\nabla^2_{pp}\hat{H}(x, p) \ge$ $\left[\lambda'/p\right]/(4R) - C(|p|/R^2 + |p|^2/R^4)]I_d = (|p|/R)\left[\lambda'/4 - C/R - C|p|/R^3\right]I_d$. For R large enough, the right-hand side is positive. And, then, it is straightforward to deduce that H satisfies (\hat{A} 4). Property (\hat{A} 3) follows from the fact that φ has bounded derivatives (of order greater than 1) and thus Φ also has bounded derivatives (of order greater than 1). \Box

As for the construction of $\hat{L}^0(x_0, p)$, it suffices to consider $L^0(x_0, -\zeta(p)\nabla_p H^0(x_0, p))$.

Proposition 5.19. Assume that $(f_t^0)_{0 \le t \le T}$, $(f_t)_{0 \le t \le T}$, L^0 and L are as in Assumptions (A1), (A2) and (A4) and define the corresponding Hamiltonians H^0 and H . Consider g^0 and g as in ($\hat{A}2$), for a certain $s \in (3, \lfloor s \rfloor - (d/2 + 1)]$. Given $R_0 \geq 2\mathfrak{L}$ (with \mathfrak{L} *as in* ($\hat{A}2$)), there exists a constant $\mathfrak{C} > 0$, only depending on d, κ , \mathfrak{L} , λ , R_0 , σ_0 and ϵ , such that, for $T \leq \mathfrak{C}$, the system [\(5.1\)](#page-50-1)–[\(5.2\)](#page-50-0) with \hat{H} and \hat{L}^0 given by Lemma [5.18](#page-79-1) is uniquely solvable (in the sense of Theorem [5.1\)](#page-50-2) and *satisfies the following conclusions for* $r := (\lfloor s \rfloor + s)/2$ *and for another constant* C, *only depending on the parameters in* (A)*:*

i. For all $t \in [0, T]$, $\mathcal{U}^0(t, \cdot, \cdot) \in \mathcal{D}^0(C, \mathbb{r} - 1)$ and $\mathcal{U}(t, \cdot, \cdot, \cdot) \in \mathcal{D}(C, \mathbb{r} - 1, \mathbb{s})$;

ii. With the same notation for the norm $\|\cdot\|_{\mathbb{F},\mathbb{F}-1}$ as in footnote [2,](#page-68-0)

$$
\lim_{h\to 0}\left(\left\|\nabla_{x_0}\mathcal{U}(t+h,x_0,\cdot,\mu)-\nabla_{x_0}\mathcal{U}(t,x_0,\cdot,\mu)\right\|_{\mathbb{F}}+\left\|\delta_\mu\mathcal{U}(t+h,x_0,\cdot,\mu,\cdot)-\delta_\mu\mathcal{U}(t,x_0,\cdot,\mu,\cdot)\right\|_{\mathbb{F},\mathbb{F}-1}\right)=0,
$$

and

$$
\lim_{h \to 0} \left(\left| \nabla_{x_0} \mathcal{U}^0(t+h, x_0, \mu) - \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu) \right| + \left\| \delta_\mu \mathcal{U}^0(t+h, x_0, \mu, \cdot) - \delta_\mu \mathcal{U}^0(t, x_0, \mu, \cdot) \right\|_{\mathbb{R}^{-1}} \right) = 0;
$$

- *iii.* The gradients $(t, x_0, \mu) \mapsto \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu)$ and $(t, x_0, x, \mu) \mapsto \nabla_x \mathcal{U}(t, x_0, x, \mu)$ are bounded by R_0 . In particu*lar, the solution to* [\(5.1\)](#page-50-1)*–*[\(5.2\)](#page-50-0) *is also a solution to* [\(2.11\)](#page-8-0)*–*[\(2.12\)](#page-9-0) *and any other solution to* [\(2.11\)](#page-8-0)*–*[\(2.12\)](#page-9-0) *satisfying* $|| \int_0^{\cdot} Z_s^0 \cdot dB_s^0 ||_{\text{BMO}} \leq R_0$ and $\sup_{t \in [0,T]} |\nabla_x u_t| \leq R_0$ *coincides with the unique solution of* [\(5.1\)](#page-50-1)–[\(5.2\)](#page-50-0);
- *iv.* If the coefficients f^0 , f, g^0 and g satisfy the additional conditions of Corollary [5.17,](#page-78-2) then (U^0, U) solves the master *equations* [\(2.27\)](#page-15-0)*–*[\(2.28\)](#page-15-1)*.*

Proof. We let $s := (\lfloor \Delta \rfloor - (d/2 + 1) + 3)/2$. For $R \ge R_0$ and $T \le \mathfrak{C}$, with \mathfrak{C} depending on d, \mathfrak{L} , λ , κ , R , σ_0 and s , the system [\(5.1\)](#page-50-1)–[\(5.2\)](#page-50-0) with \hat{H} and \hat{L}^0 given by Lemma [5.18](#page-79-1) has a unique solution. Item (i) follows from Proposition [5.10.](#page-67-0) Item (ii) follows from (5.45) and (5.46) .

The main challenge is to prove item (iii). We first prove the bound for $(t, x_0, x, \mu) \mapsto \nabla_{x_0} \mathcal{U}(t, x_0, x, \mu)$. To do so, we come back to the proof of the bound [\(5.4\)](#page-51-0). In fact, now that u has been found, we can regard the term $H(x, \nabla_x u_t(x))$ as part of the source term in (5.2) , with an explicit bound (depending on R). The idea is that the influence of the source term is very small as t gets close to T . In particular, we can use any standard estimates for linear backward SPDEs, see for instance [\[14](#page-86-1), Lemma 4.3.7], to a get a bound for $|\nabla_x u_t|$ of the form $\mathfrak{L} + C_R(T - t)$, where C_R depends on R. Briefly, $\mathfrak L$ comes from the boundary condition g and C_R comes from the source term. For $T - t$ small enough, $C_R(T - t)$ can be bounded by L. Equivalently, we can decrease the value of C so that $(t, x_0, x, \mu) \mapsto \nabla_x \mathcal{U}(t, x_0, x, \mu)$ is bounded by R_0 . The proof of the bound for $(t, x_0, \mu) \mapsto \nabla_{x_0} \mathcal{U}^0(t, x_0, \mu)$ is very similar, even though slightly more difficult. The first step is to use the representation of $\nabla_{x_0} \mathcal{U}^0$ in terms of K^{0,x_0} , see Proposition [5.8,](#page-64-0) and then to return to the interpretation of the latter in terms of δY^0 , see [\(5.34\)](#page-61-0). The point is thus to come back to the equation [\(5.32\)](#page-61-1) for δY^0 . At this stage, we already have a bound for δX^0 and δZ^0 . Therefore, we can also obtain a bound for $|\delta Y_t^0|$ of the form $\mathfrak{L} + C_R(T-t)$ and then complete the proof as done for $|\nabla_x u|$.

We now turn to the rest of the claim (iii). The fact that the solution to (5.1) – (5.2) is also a solution to (2.11) – (2.12) is easily checked and follows from (5.81) (by the way, the argument is similar for item (iv)). It thus remains to prove that uniqueness to [\(2.11\)](#page-8-0)–[\(2.12\)](#page-9-0) holds true within the class of solutions satisfying $\|\int_0^{\infty} Z_s^0 \cdot dB_s^0\|_{\text{BMO}} \le R_0$ (which condition is obviously satisfied by the solution to (5.1) – (5.2) since $T \le 1$) and $\sup_{t \in [0,T]} |\nabla_x u_t| \le R_0$. Under the latter bound, we can replace $H(x, \nabla_x u_t)$ by $\hat{H}(x, \nabla_x u_t)$ in the system [\(2.12\)](#page-9-0). This makes it possible to identify the two systems [\(2.12\)](#page-9-0) and [\(5.2\)](#page-50-0) (i.e., the systems for the minor player). The difficulty comes from the fact that we cannot do the same for the major player and identify the systems [\(2.11\)](#page-8-0) and [\(5.1\)](#page-50-1): indeed, we cannot replace $L^0(X_t^0, -\nabla_p H^0(X_t^0, Z_t^0))$ by $\hat{L}^0(X_t^0, Z_t^0)$ because the BMO bound on Z^0 is too weak to do so. To overcome this issue, we must revisit the proof $\mathcal{L}^{\{X_t, Z_t\}}$ because the BMO bound on $\mathbb{Z}^{\{X_t\}}$ is too weak to do so. To overcome this issue, we must revisit the proof of Theorem [5.1](#page-50-2) with $\hat{L}^0(x_0, p)$ being replaced by $L^0(x_0, -\nabla_p H^0(x_0, p))$ (but with difficulty only comes from the new 'major' system (5.1) , the only point is to prove that (5.11) remains true in this new setting with the additional assumption that $\tilde{X}^{0,i} = X^{0,i}$ for $i = 1, 2$ (since we are dealing with uniqueness, we can restrict the analysis to fixed points of the mapping $\mathfrak T$, with the latter being defined right below [\(5.8\)](#page-52-1)). Here is the way we get the analogue of (5.11) . Removing the 'tilde' in the second step of the proof of Theorem [5.1,](#page-50-2) we first observe from the Lipschitz property of $\nabla_p H^0$ and Gronwall's lemma that

$$
\sup_{t \in [0,T]} |X_t^{0,1} - X_t^{0,2}| \le C \int_0^T |Z_t^{0,1} - Z_t^{0,2}| \mathrm{d}t,\tag{5.83}
$$

for a constant C only depending on d, κ , \mathfrak{L} , λ , σ_0 and s, where here and below we assume without any loss of generality that $T \le 1$ (equivalently $\mathfrak{C} \le 1$). As for the backward equation, we notice that there exist two progressively measurable processes $(\theta_t)_{0 \le t \le T}$ and $(\theta_t)_{0 \le t \le T}$, with values in \mathbb{R}^d , such that $|\theta_t| \le C(1 + |Z_t^{0,1}| + |Z_t^{0,2}|)$ and $|\theta_t| \le C$ for all $t \in [0, T]$ and a possibly new value of C, such that

$$
d(Y_t^{0,1} - Y_t^{0,2}) = -\left[f_t(X_t^{0,1}, \mu_t^1) - f_t(X_t^{0,2}, \mu_t^2) + \vartheta_t \cdot (X_t^{0,1} - X_t^{0,2})\right]dt
$$

$$
- \theta_t \cdot (Z_t^{0,1} - Z_t^{0,2})dt + \sigma_0(Z_t^{0,1} - Z_t^{0,2}) \cdot dB_t^0, \quad t \in [0, T].
$$

We then let

$$
\mathcal{E}_t := \mathcal{E}_t \bigg(\sigma_0^{-1} \int_0^{\cdot} \theta_s \cdot \mathrm{d}B_s^0 \bigg), \quad t \in [0, T].
$$

And then, by the BMO properties of $Z^{0,1}$ and $Z^{0,2}$, we can apply Girsanov theorem and then obtain

$$
Y_t^{0,1} - Y_t^{0,2} = \mathcal{E}_t^{-1} \mathbb{E}^0 \bigg\{ \mathcal{E}_T \bigg[g^0(X_T^{0,1}, \mu_T^1) - g^0(X_T^{0,2}, \mu_T^2) + \int_t^T \big[f_s^0(X_s^{0,1}, \mu_s^1) - f_s^0(X_s^{0,2}, \mu_s^2) \big] ds + \int_t^T \vartheta_s \cdot (X_s^{0,1} - X_s^{0,2}) ds \bigg] \, | \, \mathcal{F}_t^0 \bigg\}, \quad t \in [0, T].
$$

In fact, the BMO norm of θ is bounded by a known constant. We deduce from [\[40,](#page-86-2) Theorem 3.1] that there exist two conjugate exponents $p, q > 1$, only depending on d, κ , \mathfrak{L} , λ , σ_0 and s, such that

$$
\mathbb{E}^{0}\Big[\Big(\mathcal{E}_t^{-1}\mathcal{E}_T\Big)^q|\,\mathcal{F}_t^0\Big]\leq C,
$$

the value of C being allowed to vary from line to line. Then, by Hölder inequality,

$$
|Y_t^{0,1} - Y_t^{0,2}| \le C \mathbb{E}^0 \left[\sup_{t \le s \le T} |X_s^{0,1} - X_s^{0,2}|^p + \sup_{t \le s \le T} \mathbb{W}_1(\mu_s^1, \mu_s^2)^p | \mathcal{F}_t^0 \right]^{1/p}
$$

$$
\le C \mathbb{E}^0 \left[\sup_{t \le s \le T} |X_s^{0,1} - X_s^{0,2}|^{2p} + \sup_{t \le s \le T} \mathbb{W}_1(\mu_s^1, \mu_s^2)^{2p} | \mathcal{F}_t^0 \right]^{1/(2p)}.
$$
 (5.84)

Also, by squaring the difference $Y^{0,1} - Y^{0,2}$ (in [\(2.11\)](#page-8-0)), we obtain the following variant of [\[55,](#page-87-0) Proposition 2.2]: \mathbb{P}^0 almost surely, for all $t \in [0, T]$,

$$
\label{eq:Y01} \begin{split} &|Y_t^{0,1}-Y_t^{0,2}|^2+\tfrac{1}{2}\sigma_0^2\mathbb{E}^0\biggl[\int_t^T|Z_s^{0,1}-Z_s^{0,2}|^2\mathrm{d}s\,|\,\mathcal{F}_t^0\biggr] \\ &\leq C\mathbb{E}\biggl[\sup_{s\in[0,T]}|X_s^{0,1}-X_s^{0,2}|^2+\sup_{s\in[0,T]}|\mathbb{V}_t^{0,1}\mu_s^2)^2+\sup_{s\in[t,T]}|Y_s^{0,1}-Y_s^{0,2}|^2\int_t^T\bigl(1+|Z_s^{0,1}|+|Z_s^{0,2}|\bigr)^2\mathrm{d}s\,|\,\mathcal{F}_t^0\biggr]. \end{split}
$$

Here, using again the BMO property together with [\[40](#page-86-2), Theorem 2.2], we can assume without any loss of generality that

$$
\mathbb{E}\bigg[\bigg(\int_t^T \big(1+|Z^{0,1}_s|+|Z^{0,2}_s|\big)^2\mathrm{d} s\bigg)^q\,|\,\mathcal{F}^0_t\bigg]\leq C.
$$

And then, by a new application of Hölder inequality (assuming without any loss of generality that $p \in (2, +\infty)$ and $q \in (1, 2)$,

$$
\label{eq:Y01} \begin{aligned} &|Y_t^{0,1}-Y_t^{0,2}|^2+\tfrac{1}{2}\sigma_0^2\mathbb{E}^0\bigg[\int_t^T|Z_s^{0,1}-Z_s^{0,2}|^2\mathrm{d}s\,|\,\mathcal{F}_t^0\bigg] \\ &\leq C\mathbb{E}\bigg[\sup_{s\in[0,T]}|X_s^{0,1}-X_s^{0,2}|^2+\sup_{s\in[0,T]}\mathbb{W}_1(\mu^1_s,\mu^2_s)^2\,|\,\mathcal{F}_t^0\bigg]+C\mathbb{E}\bigg[\sup_{s\in[t,T]}|Y_s^{0,1}-Y_s^{0,2}|^{2p}\,|\,\mathcal{F}_t^0\bigg]^{1/p}. \end{aligned}
$$

Inserting [\(5.84\)](#page-81-0) and using Doob's inequality, we get

$$
\begin{aligned} &|Y_t^{0,1} - Y_t^{0,2}|^2 + \tfrac{1}{2}\sigma_0^2 \mathbb{E}^0 \bigg[\int_t^T |Z_s^{0,1} - Z_s^{0,2}|^2 \mathrm{d}s \, |\, \mathcal{F}_t^0 \bigg] \\ &\leq C \mathbb{E} \bigg[\sup_{s \in [0,T]} |X_s^{0,1} - X_s^{0,2}|^{2p} + \sup_{s \in [0,T]} \mathbb{W}_1(\mu_s^1, \mu_s^2)^{2p} \, |\, \mathcal{F}_t^0 \bigg]^{1/p} \end{aligned}
$$

.

Taking the power p and then the supremum over $t \in [0, T]$, we get

$$
\label{eq:21} \begin{split} &\mathbb{E}^{0}\bigg[\sup_{t\in[0,T]}\mathbb{E}^{0}\bigg[\int_{t}^{T}|Z_{s}^{0,1}-Z_{s}^{0,2}|^{2}\textrm{d} s\,|\,\mathcal{F}_{t}^{0}\bigg]^{p}\,|\,\mathcal{F}_{0}^{0}\bigg]\\ &\leq C T^{p}\mathbb{E}^{0}\bigg[\bigg(\int_{0}^{T}|Z_{s}^{0,1}-Z_{s}^{0,2}|^{2}\textrm{d} s\bigg)^{p}\,|\,\mathcal{F}_{0}^{0}\bigg]+C\mathbb{E}^{0}\bigg[\sup_{s\in[0,T]}\mathbb{W}_{1}(\mu_{s}^{1},\mu_{s}^{2})^{2p}\,|\,\mathcal{F}_{0}^{0}\bigg], \end{split}
$$

with the second line following from [\(5.83\)](#page-81-1). Bounding from below the left-hand side by the value at $t = 0$, we get (for a possibly new value of $\mathfrak{C} \leq 1$:

$$
\mathbb{E}^{0}\Big[\sup_{s\in[0,T]}|X_{s}^{0,1}-X_{s}^{0,2}|^{2p}\Big]\leq CT\mathbb{E}^{0}\Big[\sup_{s\in[0,T]}\mathbb{W}_{1}(\mu_{s}^{1},\mu_{s}^{2})^{2p}\Big].
$$

The fact that we use here $2p$ as exponent (and not 2) makes a difference with [\(5.11\)](#page-52-0). However, we can easily complete the proof of uniqueness by inserting the above bound in (5.14) (with a possibly new value of p). \Box

6. Appendix

6.1. Convenient form of the chain rule over $\mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d)$

In this subsection, we consider two filtered probability spaces $(\Omega^0, \mathcal{F}^0, \mathbb{P}^0, \mathbb{P}^0)$ and $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ equipped with two Brownian motions $(B_t^0)_{0 \le t \le T}$ and $(B_t)_{0 \le t \le T}$ with values in \mathbb{R}^d . We are also given two \mathbb{R}^d -valued Itô processes

$$
dX_t^0 = b_t^0 dt + \varsigma_t^0 dB_t^0,
$$

$$
dX_t = b_t dt + \varsigma_t dB_t, \quad t \ge 0,
$$

with square-integrable conditions X_0^0 and X_0 , respectively \mathcal{F}_0^0 and \mathcal{F}_0 -measurable. The processes $(b_t^0)_{t\geq0}$ and $(\varsigma_t^0)_{t\geq0}$ are constructed on Ω^0 and are \mathbb{F}^0 -progressively measurable (with values in \mathbb{R}^d and $\mathbb{R}^d \otimes \mathbb{R}^d$ respectively), and the processes $(b_t)_{t\geq0}$ and $(\varsigma_t)_{t\geq0}$ are constructed on $\Omega^0 \times \Omega$ and are $\mathbb{F}^0 \otimes \mathbb{F}$ -progressively measurable (also with values in \mathbb{R}^d and $\mathbb{R}^d \otimes \mathbb{R}^d$ respectively).

We also assume that, for any $T > 0$, the processes $(\zeta_t^0)_{0 \le t \le T}$ and $(\zeta_t)_{0 \le t \le T}$ are bounded by a deterministic constant. And, we assume that

$$
\mathbb{E}^{0}\Bigl[\sup_{0\leq t\leq T}|b_{t}^{0}|^{2}\Bigr]+\mathbb{E}^{0}\mathbb{E}\Bigl[\sup_{0\leq t\leq T}|b_{t}|^{2}\Bigr]<\infty.
$$

Lastly, for any $t \geq 0$, we let

$$
\mu_t(\omega^0) := \mathcal{L}^0(X_t)(\omega^0), \quad \omega^0 \in \Omega^0,
$$

when $(X_t)_{t\geq 0}$ is regarded as a random variable with values in \mathbb{T}^d . Equivalently, $\mu_t(\omega^0)$ is seen as an element of $\mathcal{P}(\mathbb{T}^d)$.

Proposition 6.1. Let $\ell : [0, +\infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \ni (t, x_0, \mu) \mapsto \ell(t, x_0, \mu) \in \mathbb{R}$ be differentiable with respect to t, x_0 and µ *such that*

- *1.* The functions $(t, x_0, \mu) \mapsto \partial_t \ell(t, x_0, \mu)$ and $(t, x_0, \mu) \mapsto \nabla_{x_0} \ell(t, x_0, \mu)$ are jointly continuous (with respect to \mathbb{W}_1 *in the argument* μ *);*
- *2. The function* $[0, +\infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d \ni (t, x_0, \mu, y) \mapsto \partial_\mu \ell(t, x_0, \mu, y)$ *is jointly continuous (with respect to* \mathbb{W}_1 *in the argument* μ *) and is differentiable with respect to y, the derivative* $[0, +\infty) \times \mathbb{R}^d \times \mathcal{P}(\mathbb{T}^d) \times \mathbb{T}^d \ni$ $(t, x_0, \mu, y) \mapsto \nabla_y \partial_\mu \ell(t, x_0, \mu, y)$ *being also jointly continuous.*

Then, \mathbb{P}^0 -almost surely, for any $t \geq 0$,

$$
d_t[\ell(t, X_t^0, \mu_t)] = \left[\partial_t \ell(t, X_t^0, \mu_t) + \mathbb{E}\left[b_t \cdot \partial_\mu \ell(t, X_t^0, \mu_t, X_t)\right] + \frac{1}{2} \mathbb{E}\left[\text{Tr}\left(\varsigma_t \varsigma_t^\dagger \nabla_y \partial_\mu \ell(t, X_t^0, \mu_t, X_t)\right)\right] \right. \\ \left. + b_t^0 \cdot \nabla_{x_0} \ell(t, X_t^0, \mu_t) + \frac{1}{2} \text{Tr}\left(\varsigma_t^0 \left(\varsigma_t^0\right)^\dagger \nabla_{x_0}^2 \ell(t, X_t^0, \mu_t)\right)\right] dt + \left(\varsigma_t^0\right)^\dagger \nabla_{x_0} \ell(t, X_t^0, \mu_t) \cdot d_t^0.
$$

Proof. Fix $T > 0$. It suffices to prove the formula for $t \in [0, T]$. For a mesh $0 = t_0 < t_1 < \cdots < t_n = t$ of the interval $[0, t]$, we have

$$
\ell(t, X_t^0, \mu_t) - \ell(0, X_0^0, \mu_0)
$$

=
$$
\sum_{i=1}^n \Big[\{ \ell(t_i, X_{t_i}^0, \mu_{t_i}) - \ell(t_{i-1}, X_{t_i}^0, \mu_{t_{i-1}}) \} + \{ \ell(t_{i-1}, X_{t_i}^0, \mu_{t_{i-1}}) - \ell(t_{i-1}, X_{t_{i-1}}^0, \mu_{t_{i-1}}) \} \Big].
$$

By freezing ω^0 , we can expand $\ell(t_i, X_{t_i}^0, \mu_{t_i}) - \ell(t_{i-1}, X_{t_i}^0, \mu_{t_{i-1}})$ by means of the standard chain rule on $[0, +\infty) \times$ $\mathcal{P}(\mathbb{T}^d)$, see [\[19,](#page-86-0) Theorem 5.99]. For $i \in \{1, \dots, n\}$, we obtain

$$
\ell(t_{i}, X_{t_{i}}^{0}, \mu_{t_{i}}) - \ell(t_{i-1}, X_{t_{i}}^{0}, \mu_{t_{i-1}})
$$
\n
$$
= \int_{t_{i-1}}^{t_{i}} \partial_{t} \ell(s, X_{t_{i}}^{0}, \mu_{s}) ds + \int_{t_{i-1}}^{t_{i}} \mathbb{E}[b_{s} \cdot \partial_{\mu} \ell(s, X_{t_{i}}^{0}, \mu_{s}, X_{s})] ds + \frac{1}{2} \int_{t_{i-1}}^{t_{i}} \mathbb{E}[Tr(\varsigma_{s} \varsigma_{s}^{\dagger} \nabla_{y} \partial_{\mu} \ell(s, X_{t_{i}}^{0}, \mu_{s}, X_{s}))] ds
$$
\n
$$
= \int_{t_{i-1}}^{t_{i}} \partial_{t} \ell(s, X_{s}^{0}, \mu_{s}) ds + \int_{t_{i-1}}^{t_{i}} \mathbb{E}[b_{s} \cdot \partial_{\mu} \ell(s, X_{s}^{0}, \mu_{s}, X_{s})] ds + \frac{1}{2} \int_{t_{i-1}}^{t_{i}} \mathbb{E}[Tr(\varsigma_{s} \varsigma_{s}^{\dagger} \nabla_{y} \partial_{\mu} \ell(s, X_{s}^{0}, \mu_{s}, X_{s}))] ds
$$
\n
$$
+ \varpi_{t_{i-1}, t_{i}}^{0},
$$

where $(\varpi_{r,s}^0)_{0 \le r \le s \le T}$ is a collection of \mathcal{F}_T^0 -measurable random variables satisfying $\sum_{i=1}^n |\varpi_{t_{i-1},t_i}^0| \to 0$ in \mathbb{P}^0 probability as n tends to ∞ (and the step size of the mesh tends to 0). The derivation of the above identity relies on the fact that the path $(X_s^0)_{0 \le s \le T}$ is continuous and the derivatives $(s, \xi, \mu, y) \mapsto \partial_\mu \ell(s, \xi, \mu, y)$ and $(s, \xi, \mu, y) \mapsto$ $\nabla_y \partial_\mu \ell(s, \xi, \mu, y)$ are continuous (and thus) bounded on (compact) sets of the form $[0, T] \times {\{\xi \in \mathbb{R}^d : |\xi| \le a\}} \times \mathcal{P}(\mathbb{T}^d) \times$ \mathbb{T}^d , for any $a > 0$ (and similarly for the derivative $\partial_t \ell$). Moreover, by standard Itô formula, we have in a similar manner:

$$
\ell(t_{i-1}, X_{t_i}^0, \mu_{t_{i-1}}) - \ell(t_{i-1}, X_{t_{i-1}}^0, \mu_{t_{i-1}})
$$
\n
$$
= \int_{t_{i-1}}^{t_i} b_s^0 \cdot \nabla_{x_0} \ell(t_{i-1}, X_s^0, \mu_{t_{i-1}}) ds + \frac{1}{2} \int_{t_{i-1}}^{t_i} \text{Tr} \Big(\zeta_s^0 (\zeta_s^0)^\dagger \nabla_{x_0}^2 \ell(t_{i-1}, X_s^0, \mu_{t_{i-1}}) \Big) ds
$$
\n
$$
+ \int_{t_{i-1}}^{t_i} (\zeta_s^0)^\dagger \nabla_{x_0} \ell(t_{i-1}, X_s^0, \mu_{t_{i-1}}) \cdot dB_s^0
$$
\n
$$
= \int_{t_{i-1}}^{t_i} b_s^0 \cdot \nabla_{x_0} \ell(s, X_s^0, \mu_s) ds + \frac{1}{2} \int_{t_{i-1}}^{t_i} \text{Tr} \Big(\zeta_s^0 (\zeta_s^0)^\dagger \nabla_{x_0}^2 \ell(s, X_s^0, \mu_s) \Big) ds
$$
\n
$$
+ \int_{t_{i-1}}^{t_i} (\zeta_s^0)^\dagger \nabla_{x_0} \ell(s, X_s^0, \mu_s) \cdot dB_s^0 + \varpi_{t_{i-1}, t_i}^0,
$$

for a possible new choice of $(\varpi_{r,s}^0)_{0\leq r < s \leq T}$. The above identity follows from the continuity of the path $(\mu_s)_{0\leq s \leq T}$ (with respect to \mathbb{W}_1).

Combining the last two displays, summing over i and letting the step size of the mesh tend to 0, we complete the proof. \Box

6.2. Estimates for transport-diffusion equations

We here collect several results regarding the long-time behaviour of transport-diffusion equation on the torus. We start with the following first lemma:

Lemma 6.2. Let $T > 0$, $r \ge 1$ and $b : [0, \infty) \times \mathbb{T}^d \to \mathbb{R}^d$ be a measurable function such that $\sup_{0 \le t \le T} ||b(t, \cdot)||_{r-1} < \infty$. *Then, there exist two constants* C *and* $\gamma > 0$ *, depending on the quantity* sup_{0≤t≤T} $||b(t, \cdot)||_{r-1}$ *but not on* T, such that the solution to the transport-diffusion equation (set on $[0,T]\times\mathbb{T}^d$)

$$
\partial_t \varphi_t + \frac{1}{2} \Delta \varphi_t + b(t, \cdot) \cdot \nabla \varphi_t = 0, \quad t \in [0, T]; \quad \varphi_T = \phi,
$$

for $\phi \in C^{\mathbb{r}}(\mathbb{T}^d)$ *, satisfies*

$$
\|\varphi_t - \bar{\varphi}_t\|_{\mathbb{r}} \le C \exp\left(-\gamma(T-t)\right) \|\phi\|_{\mathbb{r}},\tag{6.1}
$$

where $\bar{\varphi}_t = \int_{\mathbb{T}^d} \varphi_t(x) dx$.

Proof. Notice that the result is standard for t close to T (say $T - t \le 1$). It just provides a control of the $C^{\mathbb{r}}$ -norm of the solution in term of the C^{Γ} -norm of the terminal condition and the $C^{\Gamma-1}$ -norm of the velocity field.

In order to get the exponential decay when t gets away from T , we recall from [\[11,](#page-86-3) Lemma 7.4] that

$$
\|\varphi_t - \overline{\varphi}_t\|_{L^\infty} \le C \exp\bigl(-\gamma(T-t)\bigr) \|\phi\|_{L^\infty}, \quad t \in [0, T], \tag{6.2}
$$

with C and γ as in the statement. Take now $\delta \in (0,1)$. For $t \in [0, T - \delta]$, we write

$$
\varphi_t = P_{\delta} \varphi_{t+\delta} + \int_t^{t+\delta} P_{r-t} \big[b(r,\cdot) \cdot \nabla \varphi_r \big] dr,
$$

where $(P_s)_{s\geq 0}$ stands here for the standard heat kernel on the torus \mathbb{T}^d , namely $(P_s)_{s\geq 0}$ is the semi-group generated by the operator $\frac{1}{2}\Delta$. And then,

$$
\varphi_t - \bar{\varphi}_{t+\delta} = P_{\delta} \big(\varphi_{t+\delta} - \bar{\varphi}_{t+\delta} \big) + \int_t^{t+\delta} P_{r-t} \big[b(r,\cdot) \cdot \nabla \varphi_r \big] dr.
$$

In particular, for any integer $k \in \{0, \dots, \lfloor \mathbf{r} \rfloor - 1\}$ and any real $\eta \in (0, 1)$,

$$
\|\nabla^{k+1}\varphi_t\|_{\eta} = \left\|\nabla^{k+1}\left(\varphi_t - \bar{\varphi}_{t+\delta}\right)\right\|_{\eta} \leq C_{\delta} \|\varphi_{t+\delta} - \bar{\varphi}_{t+\delta}\|_{k} + C \int_{t}^{t+\delta} (r-t)^{-(1+\eta)/2} \|b(r,\cdot)\cdot\nabla\varphi_r\|_{k} dr,
$$

where C_{δ} depends on δ .

Assuming that [\(6.2\)](#page-84-0) holds true with respect to $\|\cdot\|_k$ instead of $\|\cdot\|_{\mathbb{F}}$, we can bound $\|b(r, \cdot)\cdot\nabla\varphi_r\|_k = \|b(r, \cdot)\cdot\nabla(\varphi_r - \varphi_r)\|_k$ $\|\bar{\varphi}_r\|_k$ by $C_k(\exp(-\gamma(T-r))\|\phi\|_{\mathbb{F}} + \|\nabla^{k+1}\varphi_r\|_n)$. We obtain

$$
\|\nabla^{k+1}\varphi_t\|_{\eta} \leq C_{k,\delta} \exp\bigl(-\gamma(T-t)\bigr) \|\phi\|_{\mathbb{F}} + C_k \int_t^{t+\delta} (r-t)^{-(1+\eta)/2} \|\nabla^{k+1}\varphi_r\|_{\eta} dr,
$$

and then, for $t + \delta \leq T$,

$$
\exp(\gamma(T-t))\|\nabla^{k+1}\varphi_t\|_{\eta} \leq C_{k,\delta}\|\phi\|_{\mathbb{F}} + C_k\delta^{(1-\eta)/2}\exp(\gamma\delta)\sup_{t\leq r\leq t+\delta}\Bigl[\exp\bigl(\gamma(T-r)\bigr)\|\nabla^{k+1}\varphi_r\|_{\eta}\Bigr].
$$

Choosing δ small enough and then taking the supremum over $t \leq T - \delta$, we get a bound for the left-hand side. We then get the result by iterating on the value of k. get the result by iterating on the value of k .

Lemma 6.3. Let $T > 0$, $r \ge 1$ and $b : [0, \infty) \times \mathbb{T}^d \to \mathbb{R}^d$ be a measurable function such that $\sup_{0 \le t \le T} ||b(t, \cdot)||_r < \infty$. *Then, there exist two constants* C *and* $\gamma > 0$ *, depending on* $\sup_{0 \le t \le T} ||b(t, \cdot)||_{r} < \infty$ *but not on* T, *such that the solution to the conservation equation (set on* $[0,T] \times \mathbb{T}^d$ *)*

$$
\partial_t q_t - \frac{1}{2} \Delta q_t + \text{div}_x (b(t, \cdot) q_t) = 0, \quad t \in [0, T]; \quad q_0 = q,
$$
\n(6.3)

for a smooth initial condition $q: \mathbb{T}^d \to \mathbb{R}$ *with* $\int_{\mathbb{T}^d} q(x) dx = 0$ *<i>, satisfies*

$$
||q_t||_{-\mathbb{r}} \leq C \exp(-\gamma t) ||q||_{-\mathbb{r}}, \quad t \in [0, T].
$$

When $\bar{q} := \int_{\mathbb{T}^d} q(x) dx \neq 0$, we deduce from the conservative structure that

$$
||q_t - \bar{q}||_{-\Gamma} \le C||q - \bar{q}||_{-\Gamma}, \quad \int_{\mathbb{T}^d} q_t(x) dx = \bar{q}, \quad t \in [0, T].
$$

Proof. The proof is done by duality. For ϕ and $(\varphi_t)_{0 \le t \le T}$ as in the statement of Lemma [6.2,](#page-84-1) we compute (by decomposing q in positive and negative parts, it suffices to derive the identity below when q is a probability measure, in which case the result follows on Itô-Krylov formula for Itô processes with a non-degenerate diffusion coefficient and abounded drift)

$$
d_t(\varphi_t, q_t) = 0, \quad t \in [0, T],
$$

where (\cdot, \cdot) is here understood as the duality bracket between $C^r(\mathbb{T}^d)$ and $C^{-r}(\mathbb{T}^d)$, so that

$$
(\phi,q_T)=(\varphi_0,q)\leq C\exp\bigl(-\gamma T\bigr)\|q\|_{-\mathbb{r}}\|\phi\|_{\mathbb{r}}.
$$

The result follows by maximizing over $\phi \in C^{\mathbb{r}}(\mathbb{T}^d)$ satisfying $\|\phi\|_{\mathbb{r}} \leq 1$.

Acknowledgment

François Delarue acknowledges the financial support of the European Research Council (ERC) under the European Union's Horizon Europe research and innovation programme (AdG ELISA project, Grant Agreement No. 101054746). Views and opinions expressed are however those of the authors only and do not necessarily reflect those of the European Union or the European Research Council Executive Agency. Neither the European Union nor the granting authority can be held responsible for them.

Chenchen Mou acknowledges the financial support of the Hong Kong Research Grants Council (RGC) under Grants No. GRF 11311422 and GRF 11303223.

References

[1] Y. Achdou, P. Cardaliaguet, F. c. Delarue, A. Porretta, and F. Santambrogio, *Mean field games*, Lecture Notes in Mathematics, vol. 2281, Springer, Cham; Centro Internazionale Matematico Estivo (C.I.M.E.), Florence, 2020, Notes from the CIME School held in Cetraro, June 2019, Edited by Cardaliaguet and Porretta, Fondazione CIME/CIME Foundation Subseries.

 \Box

- [2] E. Bayraktar, A. Cecchin, A. Cohen, and F. Delarue, *Finite state mean field games with Wright-Fisher common noise*, J. Math. Pures Appl. (9) 147 (2021), 98–162.
- [3] A. Bensoussan, M. H. M. Chau, and S. C. P. Yam, *Mean field games with a dominating player*, Appl. Math. Optim. 74 (2016), no. 1, 91–128.
- [4] P. Bergault, P. Cardaliaguet, and C. Rainer, *Mean field games in a Stackelberg problem with an informed major player*, SIAM J. Control Optim. 62 (2024), no. 3, 1737–1765.
- [5] C. Bertucci, *Monotone solutions for mean field games master equations: continuous state space and common noise*, Comm. Partial Differential Equations 48 (2023), no. 10-12, 1245–1285.
- [6] C. Bertucci, J.-M. Lasry, and P.-L. Lions, *Some remarks on mean field games*, Comm. Partial Differential Equations 44 (2019), no. 3, 205–227.
- [7] , *On Lipschitz solutions of mean field games master equations*, J. Funct. Anal. 287 (2024), no. 5, Paper No. 110486, 42.
- [8] C. Bertucci and C. Meynard, *Noise through an additional variable for mean field games master equation on finite state space*, (2024).
- [9] P. Briand and R. Elie, *A simple constructive approach to quadratic bsdes with or without delay*, Stochastic Processes and their Applications 123 (2013), no. 8, 2921–2939.
- [10] P. E. Caines, M. Huang, and R. P. Malhamé, *Mean field games*, Handbook of dynamic game theory, Springer, Cham, 2018, pp. 345–372.
- [11] P. Cardaliaguet, J.-M. Lasry, P.-L. Lions, and A. Porretta, *Long time average of mean field games with a nonlocal coupling*, SIAM Journal on Control and Optimization 51 (2013), no. 5, 3558–3591.
- [12] P. Cardaliaguet, M. Cirant, and A. Porretta, *Remarks on Nash equilibria in mean field game models with a major player*, Proceedings of the American Mathematical Society 148 (2020), no. 10, 4241–4255.
- [13] , *Splitting methods and short time existence for the master equations in mean field games*, J. Eur. Math. Soc. (JEMS) 25 (2023), no. 5, 1823–1918.
- [14] P. Cardaliaguet, F. Delarue, J.-M. Lasry, and P.-L. Lions, *The master equation and the convergence problem in mean field games*, Annals of Mathematics Studies, vol. 201, Princeton University Press, Princeton, NJ, 2019.
- [15] P. Cardaliaguet and F. Delarue, *Selected topics in mean field games*, ICM—International Congress of Mathematicians. Vol. 5. Sections 9–11, EMS Press, Berlin, 2023, pp. 3660–3703.
- [16] P. Cardaliaguet and A. Porretta, *Long time behavior of the master equation in mean-field game theory*, Analysis & PDE 12 (2019), no. 6, 1397– 1453.
- [17] P. Cardaliaguet and P. Souganidis, *Monotone solutions of the master equation for mean field games with idiosyncratic noise*, SIAM J. Math. Anal. 54 (2022), no. 4, 4198–4237.
- [18] P. Cardaliaguet and P. E. Souganidis, *On first order mean field game systems with a common noise*, Ann. Appl. Probab. 32 (2022), no. 3, 2289– 2326.
- [19] R. Carmona and F. Delarue, *Probabilistic theory of mean field games with applications. I*, Probability Theory and Stochastic Modelling, vol. 83, Springer, Cham, 2018, Mean field FBSDEs, control, and games.
- [20] , *Probabilistic theory of mean field games with applications. II*, Probability Theory and Stochastic Modelling, vol. 84, Springer, Cham, 2018, Mean field games with common noise and master equations.
- [21] R. Carmona, F. Delarue, and D. Lacker, *Mean field games with common noise*, Annals of Probability 44 (2016), 3740–3803.
- [22] R. Carmona, J.-P. Fouque, and L.-H. Sun, *Mean field games and systemic risk*, Commun. Math. Sci. 13 (2015), no. 4, 911–933.
- [23] R. Carmona and P. Wang, *An alternative approach to mean field game with major and minor players, and applications to herders impacts*, Appl. Math. Optim. 76 (2017), no. 1, 5–27.
- [24] R. Carmona and X. Zhu, *A probabilistic approach to mean field games with major and minor players*, Ann. Appl. Probab. 26 (2016), no. 3, 1535–1580.
- [25] E. Çinlar, J. Jacod, P. Protter, and M. J. Sharpe, *Semimartingales and Markov processes*, Z. Wahrsch. Verw. Gebiete 54 (1980), no. 2, 161–219.
- [26] F. Delarue, *On the existence and uniqueness of solutions to FBSDEs in a non-degenerate case*, Stochastic Processes and their Applications 99 (2002) , no. 2, $209 - 286$.
- [27] F. Delarue (ed.), *Mean field games*, Proceedings of Symposia in Applied Mathematics, vol. 78, American Mathematical Society, Providence, RI, 2021, AMS Short Course, Mean Field Games: Agent Based Models to Nash Equilibria, January 13–14, 2020, Denver, Colorado.
- [28] E. Di Nezza, G. Palatucci, and E. Valdinoci, *Hitchhiker's guide to the fractional Sobolev spaces*, Bull. Sci. Math. 136 (2012), no. 5, 521–573.
- [29] K. Du and S. Chen, *Backward stochastic partial differential equations with quadratic growth*, Journal of Mathematical Analysis and Applications 419 (2014), no. 1, 447–468.
- [30] K. Du and Q. Meng, *A revisit to w2n-theory of super-parabolic backward stochastic partial differential equations in* r^d *, Stochastic Processes and* their Applications 120 (2010), no. 10, 1996–2015.
- [31] R. Foguen Tchuendom, *Uniqueness for linear-quadratic mean field games with common noise*, Dyn. Games Appl. 8 (2018), no. 1, 199–210.
- [32] W. Gangbo, A. R. Mészáros, C. Mou, and J. Zhang, *Mean field games master equations with nonseparable Hamiltonians and displacement monotonicity*, Ann. Probab. 50 (2022), no. 6, 2178–2217.
- [33] D. A. Gomes, E. A. Pimentel, and V. Voskanyan, *Regularity theory for mean-field game systems*, SpringerBriefs in Mathematics, Springer, Cham, 2016.
- [34] M. Huang, *Large-population LQG games involving a major player: the Nash equivalence principle*, SIAM Journal on Control and Optimization 48 (2010), 3318–3353.
- [35] M. Huang, P. Caines, and R. Malhamé, *Individual and mass behavior in large population stochastic wireless power control problems: centralized and Nash equilibrium solutions*, (2003), 98 – 103.
- [36] M. Huang, R. P. Malhamé, and P. E. Caines, *Large population stochastic dynamic games: Closed-loop McKean-Vlasov systems and the Nash certainty equivalence principle*, Commun. Inf. Syst. 6 (2006), no. 3, 221–251.
- [37] Z. Huang and S. Tang, *Mean field games with major and minor agents: the limiting problem and nash equilibrium*, Stochastics 0 (2024), no. 0, 1–34.
- [38] P. Imkeller, A. Réveillac, and A. Richter, *Differentiability of quadratic BSDEs generated by continuous martingales*, Ann. Appl. Probab. 22 (2012), no. 1, 285–336 (English).
- [39] S. Janson and S. Kaijser, *Higher moments of Banach space valued random variables*, Mem. Amer. Math. Soc. 238 (2015), no. 1127, vii+110.
- [40] N. Kazamaki, *Continuous exponential martingales and BMO*, Lecture Notes in Mathematics, vol. 1579, Springer-Verlag, Berlin, 1994.
- 88
- [41] M. Kobylanski, *Backward stochastic differential equations and partial differential equations with quadratic growth*, Ann. Probab. 28 (2000), no. 2, 558–602.
- [42] D. Lacker, *On a strong form of propagation of chaos for McKean-Vlasov equations*, Electronic Communications in Probability 23 (2018), no. none, $1 - 11$.
- [43] D. Lacker and T. Zariphopoulou, *Mean field and* n*-agent games for optimal investment under relative performance criteria*, Math. Finance 29 (2019), no. 4, 1003–1038.
- [44] J.-M. Lasry and P.-L. Lions, *Jeux à champ moyen. I. Le cas stationnaire*, C. R. Math. Acad. Sci. Paris 343 (2006), no. 9, 619–625.
- [45] , *Jeux à champ moyen. II. Horizon fini et contrôle optimal*, C. R. Math. Acad. Sci. Paris 343 (2006), no. 10, 679–684.
- [46] , *Mean field games*, Jpn. J. Math. 2 (2007), no. 1, 229–260.
- [47] , *Mean-field games with a major player*, C. R. Math. Acad. Sci. Paris 356 (2018), no. 8, 886–890.
- [48] M. Ledoux and M. Talagrand, *Probability in Banach spaces*, Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)], vol. 23, Springer-Verlag, Berlin, 1991, Isoperimetry and processes.
- [49] D. Li and H. Queffélec, *Introduction to Banach spaces: analysis and probability. Vol. 2*, Cambridge Studies in Advanced Mathematics, vol. 167, Cambridge University Press, Cambridge, 2018, Translated from the French.
- [50] P.-L. Lions, *Cours au Collège de France, Equations aux dérivées partielles et applications*, https://www.college-de-france.fr/site/pierre-louislions/course.htm.
- [51] C. Mou and J. Zhang, *Wellposedness of Second Order Master Equations for Mean Field Games with Nonsmooth Data*, Mem. Amer. Math. Soc. 302 (2024), no. 1515.
- [52] S. Nguyen and M. Huang, *Linear-quadratic-Gaussian mixed games with continuum-parametrized minor players*, SIAM Journal on Control and Optimization (2012).
- [53] M. Nourian and P. Caines, ǫ*-Nash mean field game theory for nonlinear stochastic dynamical systems with major and minor agents*, SIAM Journal on Control and Optimization 50 (2013), 2907–2937.
- [54] E. Pardoux and S. Peng, *Backward stochastic differential equations and quasilinear parabolic partial differential equations*, Stochastic partial differential equations and their applications (Charlotte, NC, 1991), Lect. Notes Control Inf. Sci., vol. 176, Springer, Berlin, 1992, pp. 200-217.
- [55] E. Pardoux and S. Peng, *Adapted solution of a backward stochastic differential equation*, Systems & Control Letters 14 (1990), no. 1, 55–61.
- [56] D. W. Stroock and S. R. S. Varadhan, *Multidimensional diffusion processes*, Classics in Mathematics, Springer-Verlag, Berlin, 2006, Reprint of the 1997 edition.
- [57] R. Tevzadze, *Solvability of backward stochastic differential equations with quadratic growth*, Stochastic Process. Appl. 118 (2008), no. 3, 503– 515.
- [58] H. V. Tran, *Hamilton-Jacobi equations—theory and applications*, Graduate Studies in Mathematics, vol. 213, American Mathematical Society, Providence, RI, 2021.