

On the components of random geometric graphs in the dense limit

Mathew D. Penrose^{1,2} and Xiaochuan Yang^{1,3}

University of Bath and Banque Internationale á Luxembourg S.A.

January 7, 2025

Abstract

Consider the geometric graph on n independent uniform random points in a connected compact region A of \mathbb{R}^d , $d \geq 2$ with smooth boundary, or in the unit square, with distance parameter r_n . Let K_n be the number of components of this graph, and R_n the number of vertices not in the giant component. Let S_n be the number of isolated vertices. We show that if r_n is chosen so that nr_n^d tends to infinity but slowly enough that $\mathbb{E}[S_n]$ also tends to infinity, then K_n , R_n and S_n are all asymptotic to μ_n in probability as $n \rightarrow \infty$ where (with $|A|$, θ_d and $|\partial A|$ denoting the volume of A , the volume of the unit d -ball, and the perimeter of A respectively) $\mu_n := ne^{-\pi nr_n^d/|A|}$ if $d = 2$ and $\mu_n := ne^{-\theta_d nr_n^d/|A|} + \theta_{d-1}^{-1} |\partial A| r_n^{1-d} e^{-\theta_d nr_n^d/(2|A|)}$ if $d \geq 3$. We also give variance asymptotics and central limit theorems for K_n and R_n in this limiting regime when $d \geq 3$, and for Poisson input with $d \geq 2$. We extend these results (substituting $\mathbb{E}[S_n]$ for μ_n) to a class of non-uniform distributions on A .

Contents

1	Introduction	2
2	Statement of results	5
2.1	Results for general f	6
2.2	Results for the uniform case	8
2.3	Overview of proofs	9

¹Supported by EPSRC grant EP/T028653/1

²Corresponding author: Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK: m.d.penrose@bath.ac.uk

³Banque Internationale à Luxembourg S.A., 69 Route d'Esch, L-2953 Luxembourg: xiaochuan.j.yang@gmail.com

3	Preliminaries	10
3.1	Geometrical tools	11
3.2	Probabilistic tools	12
3.3	Percolation type estimates	13
4	The number of isolated vertices	18
4.1	Mean and variance of the number of isolated vertices	19
4.2	Asymptotic distribution of the singleton count	22
4.3	Asymptotics for I_n in the uniform case	22
5	Proof of first-order asymptotics	29
5.1	Asymptotics of means	30
5.2	Proof of first order limit theorems	35
6	Asymptotics of variances	38
6.1	Variances for small components: Poisson input	39
6.2	Variances for small components: binomial input	41
6.3	Variance estimates for medium components	45
6.4	Variance estimates for large components	48
6.5	Variance estimates: conclusion	50
6.6	Proof of convergence in distribution results	51

1 Introduction

Given a compact set A with a nice boundary in Euclidean space \mathbb{R}^d , $d \geq 2$, the *random geometric graph* (RGG) based on a random point set $\mathcal{X} \subset A$ is the graph $G(\mathcal{X}, r)$ with vertex set \mathcal{X} and edges between each pair of points distant at most r apart, in the Euclidean metric, for a specified distance parameter $r > 0$. Such graphs are important in a variety of applications (see [7]), including modern topological data analysis (TDA), where the topological properties of the graph are used to help understand the topology of A .

In this paper we consider the *number of components* of the graph G , denoted $K(G)$, where $G = G(\mathcal{X}, r)$ with \mathcal{X} a random sample of n points in A (denoted \mathcal{X}_n) or the corresponding Poisson process (denoted \mathcal{P}_n , and defined more formally later). In particular, we investigate asymptotic properties for large n with $r = r(n)$ specified and decaying to zero according to a certain limiting regime (see (1.1), (1.2) below). Our results add significantly to the existing literature about the limit theory of Betti numbers, an area that has received intensive recent attention in TDA. Indeed, the number of components of $G(\mathcal{X}, r)$ is the 0-th Betti number of the occupied Boolean set $\cup_{x \in \mathcal{X}} B_{r/2}(x)$, where $B_r(x)$ or $B(x, r)$ denotes the closed Euclidean ball of radius r centred on x . Given the sample \mathcal{X} , keeping track of $K(G(\mathcal{X}, r))$ while varying r corresponds to the 0-th persistent homology, which leads to sparse topological descriptors in a 2D persistence diagram. See [1] for related geometric models of TDA.

We are also concerned with the *giant component* - the component of $G(\mathcal{X}, r)$ with the largest order. For the graphs we consider, most of the vertices lie in the giant component, so for more detailed information we consider the total number of vertices $R(G(\mathcal{X}, r))$ that are *not* in the giant component of $G(\mathcal{X}, r)$. To be precise, given a finite graph G of order n , list the orders of its components in decreasing order as $L_1(G), L_2(G), \dots, L_{K(G)}(G)$. Set $R(G) := n - L_1(G)$.

We shall consider the limiting behaviour of $K_n := K(G(\mathcal{X}_n, r_n))$, $K'_n := K(G(\mathcal{P}_n, r_n))$, $R_n := R(G(\mathcal{X}_n, r))$ and $R'_n := R(G(\mathcal{P}_n, r_n))$ as $n \rightarrow \infty$ with r_n specified for all $n \geq 1$. Let θ_d (or θ for short) denote the volume of the unit radius ball in \mathbb{R}^d , i.e. $\theta_d := \pi^{d/2}/\Gamma(1+d/2)$. For points uniformly distributed in A (which we call the *uniform case*), the main limiting regime for r_n that we consider here is to assume that as $n \rightarrow \infty$,

$$nr_n^d \rightarrow +\infty; \quad (1.1)$$

$$\gamma_n := n(\theta/\lambda(A))r_n^d - (2 - 2/d)(\log n - \mathbf{1}_{\{d \geq 3\}} \log \log n) \rightarrow -\infty, \quad (1.2)$$

where λ denotes the Lebesgue measure on \mathbb{R}^d . We call this the *intermediate* or *mildly dense* regime because the average vertex degree is of order $\Theta(nr_n^d)$ and therefore grows to infinity as n becomes large, but only slowly in this regime.

Other limiting regimes of r are better understood. In the *thermodynamic regime* where $nr_n^d \rightarrow a\lambda(A)$ with $a \in (0, \infty)$ as $n \rightarrow \infty$, it holds as $n \rightarrow \infty$ that

$$\frac{K_n}{n} \xrightarrow{\mathbb{P}} c(a); \quad \frac{R_n}{n} \xrightarrow{\mathbb{P}} c'(a), \quad (1.3)$$

where $c(a) \in (0, 1)$ is given explicitly in [7, Theorem 13.25], and $c'(a) \in (0, 1]$ is given less explicitly in [7, Theorem 11.9]. If a lies below a certain percolation threshold $a_c := a_c(d) \in (0, \infty)$ then $c'(a) = 1$. Central limit theorems for K_n and for R_n in this regime are also proved in [7] (these results hold for K'_n and R'_n as well as for K_n and R_n).

In the *sparse regime* where $nr_n^d \rightarrow 0$, the average vertex degree goes to 0 and we still have (1.3) with $c(0) = c'(0) = 1$. This can be deduced from the fact that $c(a) \rightarrow 1$ as $a \rightarrow 0$ (which can be deduced from the formula in [7]), along with coupling arguments.

On the other hand, if $\gamma_n \rightarrow +\infty$, and ∂A is smooth or A is a convex polygon, it follows from [11, Theorem 1.1] that with probability tending to 1 as $n \rightarrow \infty$, $G(\mathcal{X}_n, r_n)$ is fully connected so that $K_n = 1$ and $R_n = 0$. We here call this limiting regime the *connectivity regime* (in [7] this terminology was used slightly differently).

As well as the mildly dense regime (1.1), (1.2), in this paper we also consider the case where γ_n is bounded away from $-\infty$ and $+\infty$ as $n \rightarrow \infty$; we call this the *critical regime for connectivity*. Thus we consider the whole range of possible limiting behaviours for r in between the thermodynamic and connectivity regimes.

In TDA one is interested in understanding (for a fixed sample \mathcal{X}_n) the number of components of $G(\mathcal{X}_n, r)$ in the whole range of values from $r = 0$, right up to the connectivity threshold (i.e. the smallest r such that $G(\mathcal{X}_n, r)$ is connected). Therefore it seems well worth trying to understand K_n in the mildly dense regime, as well as in other regimes. Likewise, studying R_n in this regime helps us understand the rate at which the giant component swallows up the whole vertex set as r approaches the connectivity threshold.

Our main results for the uniform case refer to constants μ_n defined by

$$\mu_n := ne^{-n\theta r_n^d/\lambda(A)} + \theta_{d-1}^{-1}|\partial A|r_n^{1-d}e^{-n\theta r_n^d/(2\lambda(A))}\mathbf{1}\{d \geq 3\}, \quad (1.4)$$

where ∂A denotes the topological boundary of A and $|\partial A|$ denotes the $(d-1)$ -dimensional Hausdorff measure of ∂A . We say A has a C^2 boundary (for short, $\partial A \in C^2$) if for each $x \in \partial A$ there exists a neighbourhood U of x and a real-valued function ϕ that is defined on an open set in \mathbb{R}^{d-1} and twice continuously differentiable, such that $\partial A \cap U$, after a rotation, is the graph of the function ϕ .

We can now present our main results for the uniform case. In all of our results we assume either that $d = 2$ and $A = [0, 1]^2$, or that $d \geq 2$ and $A \subset \mathbb{R}^d$ is compact and connected, and A is the closure of its interior (for short: $A = \overline{A^\circ}$), with $\partial A \in C^2$. Also we assume $r_n \in (0, \infty)$ is given for all $n \geq 1$. Let $N(0, 1)$ denote a standard normal random variable, and for $t \in (0, \infty)$ let Z_t be a Poisson random variable with mean t . Let $\xrightarrow{\mathcal{D}}$, respectively $\xrightarrow{L^1}$, denote convergence in distribution, respectively in the L^1 norm. Define $\sigma_A := |\partial A|/(\lambda(A)^{1-1/d})$; this is sometimes called the *isoperimetric ratio* of A .

Theorem 1.1 (Basic results for the uniform case). *Let ξ_n denote either $K_n - 1$ or R_n , and let ξ'_n denote either $K'_n - 1$ or R'_n .*

(a) *Suppose $(r_n)_{n \geq 1}$ satisfy (1.1) and (1.2). Then in the uniform case, as $n \rightarrow \infty$ we have the convergence results: $\mu_n \rightarrow \infty$, and $(\xi_n/\mu_n) \xrightarrow{L^1} 1$, and $(\xi'_n/\mu_n) \xrightarrow{L^1} 1$. Also $\mu_n^{-1}\text{Var}[\xi'_n] \rightarrow 1$, and $\mu_n^{-1/2}(\xi'_n - \mathbb{E}[\xi'_n]) \xrightarrow{\mathcal{D}} N(0, 1)$. If $d \geq 3$ then $\mu_n^{-1}\text{Var}[\xi_n] \rightarrow 1$, and $\mu_n^{-1/2}(\xi_n - \mathbb{E}[\xi_n]) \xrightarrow{\mathcal{D}} N(0, 1)$.*

(b) *Suppose instead that $\gamma_n \rightarrow \gamma \in \mathbb{R}$ as $n \rightarrow \infty$. Set $c_{d,A} := \theta_{d-1}^{-1}(\theta/(2-2/d))^{1-1/d}\sigma_A$. Then as $n \rightarrow \infty$, $\xi_n \xrightarrow{\mathcal{D}} Z_{e^{-\gamma}}$ if $d = 2$ and $\xi_n \xrightarrow{\mathcal{D}} Z_{c_{d,A}e^{-\gamma/2}}$ if $d \geq 3$, and likewise for ξ'_n .*

In Section 2 we shall provide a more detailed version of Theorem 1.1: we shall give estimates on the rates of convergence, and also generalize to allow for non-uniformly distributed points in A .

To the best of our knowledge, the only previous results on K_n and R_n in the mildly dense regime are by Ganesan [3] in the special case of $d = 2$ and $A = [0, 1]^2$, where he proved that there exists a constant $c > 0$ such that as $n \rightarrow \infty$,

$$\mathbb{P}[K_n \leq ne^{-cnr_n^2}] \rightarrow 1; \quad \mathbb{P}[R_n \leq ne^{-cnr_n^2}] \rightarrow 1. \quad (1.5)$$

In other words, the proportionate number of components and the proportionate number of vertices not in the largest component decay exponentially in nr_n^2 but the exact exponent is not identified; Ganesan's proof, while ingenious, does not provide much of a clue as to the optimal value of c satisfying (1.5), or whether this optimal value is the same for K_n and for R_n . Moreover, his proof of the second part of (1.5) does not appear to generalize to higher dimensions.

One possible reason why the mildly dense limiting regime was not previously well understood is an apparently strong dependence between contributions from different regions of space; one has to look a long way from a given vertex to tell whether it lies in the giant component. A second reason is the importance of boundary effects in this regime and the

necessity of dealing with the curved boundary of A quantitatively; note the factor of $|\partial A|$ in the definition of μ_n at (1.4). Another reason is that in contrast to the thermodynamic regime, it seems not to be possible to re-scale space to obtain a limiting Poisson process to work with, as was often done in previous works on these kinds of limit theorems, for example [9]. In Section 2.3 we shall provide an overview of the methods we develop to deal with these issues.

Our results show that the phenomenon of exponential decay is common to all dimensions and more general sets A , and we identify the optimal value of c in (1.5). Furthermore, we prove a central limit theorem (CLT) for the fluctuations of K'_n and R'_n (for all $d \geq 2$) and for K_n, R_n (for $d \geq 3$). Our CLT is ‘weakly quantitative’ in the sense that we provide bounds on the rate of convergence to normal, although our bounds might not be optimal.

We expect that our approach can shed some light on the limiting behaviour of higher dimensional homology, and higher Betti numbers, of random geometric complexes in the mildly dense regime for which the correct scaling is so far not well understood; see the last paragraph of [1, Section 2.4.1]. This is beyond the scope of this paper and we leave it for future work.

2 Statement of results

We now describe our setup more precisely. Let $d \geq 2$ and let $A \subset \mathbb{R}^d$ be a compact connected set, such that $A = \overline{A^\circ}$. Assume either that $\partial A \in C^2$ or that $d = 2$ and $A = [0, 1]^2$. Let $f : \mathbb{R}^d \rightarrow [0, \infty)$ be a probability density function with support A . We refer to the special case where f is constant on A as the *uniform case* but in general we allow possibly non-constant f . Set $f_0 := \inf_A f(x)$, $f_1 := \inf_{\partial A} f$, and $f_{\max} := \sup_A f(x)$. We assume throughout that $f_0 > 0$ and that f is continuous on A (so in particular $f_{\max} < \infty$). We use ν to denote the measure with density f , i.e. $\nu(dx) = f(x)dx$. Clearly in the uniform case $f_0 = \lambda(A)^{-1}$.

Let (X_1, X_2, \dots) be a sequence of independent random vectors in \mathbb{R}^d with common density f , and for $n \in \mathbb{N}$ set $\mathcal{X}_n := \{X_1, \dots, X_n\}$, which is a binomial point process. Also let $(Z_t)_{t>0}$ be a unit intensity Poisson counting process, independent of (X_1, X_2, \dots) , so that for $n \in [1, \infty)$, Z_n is a Poisson random variable with mean n , and set $\mathcal{P}_n := \mathcal{X}_{Z_n}$. Then \mathcal{P}_n is a Poisson point process with intensity measure $n\nu$. We use n to denote both the number of points in \mathcal{X}_n and the average number of points in a Poisson sample \mathcal{P}_n with the convention that $n \in [1, \infty)$ in the latter case.

We are concerned with the quantities $K_n := K(G(\mathcal{X}_n, r_n))$ and $R_n := R(G(\mathcal{X}_n, r_n))$ and their Poisson counterparts $K'_n := K(G(\mathcal{P}_n, r_n))$ and $R'_n := R(G(\mathcal{P}_n, r_n))$, with $r_n \in (0, \infty)$ specified for each n .

Given $g : (0, \infty) \rightarrow \mathbb{R}$, and $h : (0, \infty) \rightarrow (0, \infty)$, we write $g(x) = O(h(x))$ if $\limsup |g(x)|/h(x) < \infty$, and write $g(x) = o(h(x))$ if $\limsup |g(x)|/h(x) = 0$, $g(x) = \Omega(h(x))$ if $\liminf (g(x)/h(x)) > 0$. We write $g(x) = \Theta(h(x))$ if both $g(x) = O(h(x))$ and $g(x) = \Omega(h(x))$, and $g(x) \sim h(x)$ if $\lim(g(x)/h(x)) = 1$. Here, the limit is taken either as $x \rightarrow 0$ or $x \rightarrow \infty$ to be specified in each appearance.

To present quantitative CLTs, we recall that for random variables X, Y , the Kolmogorov distance d_K and the total variation distance d_{TV} between them are defined re-

spectively by

$$d_K(X, Y) := \sup_{z \in \mathbb{R}} |\mathbb{P}[X \leq z] - \mathbb{P}[Y \leq z]|; \quad d_{TV}(X, Y) := \sup_{A \in \mathcal{B}(\mathbb{R})} |\mathbb{P}[X \in A] - \mathbb{P}[Y \in A]|,$$

where the second supremum is taken over all Borel measurable subsets of \mathbb{R} . Note that convergence in the Kolmogorov distance implies convergence in distribution.

2.1 Results for general f

We now give our results for the component count and the number of vertices not in the giant component in the general case with f not assumed necessarily to be constant on A . For general f , instead of μ_n defined at (1.4), our results refer to constants I_n defined by

$$I_n := n \int_A \exp(-n\nu(B_{r_n}(x)))\nu(dx). \quad (2.1)$$

We define the *critical regime for connectivity* to be when r_n is chosen so that $I_n = \Theta(1)$ as $n \rightarrow \infty$, and the *mildly dense regime* to be when r_n is chosen so that (1.1) holds but also $I_n \rightarrow \infty$ as $n \rightarrow \infty$. As discussed later in Remark 2.9, The latter condition turns out to be equivalent to (1.2) in the uniform case.

Theorem 2.1 (First order moment asymptotics for general f). *Suppose f is continuous on A with $f_0 > 0$, and that r_n satisfies (1.1) and also $I_n \rightarrow \infty$ as $n \rightarrow \infty$. Let ξ_n denote any of $K_n - 1$, R_n , $K'_n - 1$ or R'_n , and let ζ_n be either ξ_n or $\xi_n + 1$. Then as $n \rightarrow \infty$ we have*

$$\mathbb{E}[\xi_n] = I_n(1 + O((nr_n^d)^{1-d})); \quad (2.2)$$

$$\mathbb{E}[|(\zeta_n/I_n) - 1|] = O((nr_n^d)^{1-d} + I_n^{-1/2}). \quad (2.3)$$

In particular $(\zeta_n/I_n) \xrightarrow{L^1} 1$.

We can use the L^1 convergence in Theorem 2.1, together with an asymptotic analysis of I_n , to determine the optimal exponent c in Ganesan's result (1.5). First we introduce some further notation. Given $(r_n)_{n \geq 1}$ we define

$$b^+ := \limsup_{n \rightarrow \infty} (n\theta r_n^d / \log n); \quad b^- := \liminf_{n \rightarrow \infty} (n\theta r_n^d / \log n). \quad (2.4)$$

and $b := b^+ = b^-$ whenever $b^+ = b^-$. Loosely speaking, b is the logarithmic growth rate of the degree of a typical vertex, at least in the uniform case with $\lambda(A) = 1$. We identify two critical values for b , namely

$$b_c := \max\left(\frac{1}{f_0}, \frac{2 - 2/d}{f_1}\right); \quad b'_c := \begin{cases} (d(f_0 - f_1/2))^{-1} & \text{if } f_0 > f_1/2; \\ +\infty & \text{if } f_0 \leq f_1/2 \end{cases} \quad (2.5)$$

(so in the uniform case $b_c = (2 - 2/d)/f_0$ and $b'_c = 2/(df_0)$, and hence $b'_c < b_c$ if $d \geq 3$). The following result shows b_c is the critical value of the logarithmic growth rate b above which $I_n \rightarrow 0$, and below which $I_n \rightarrow \infty$.

Proposition 2.2. *If $b^+ < b_c$ then $I_n \rightarrow \infty$ as $n \rightarrow \infty$. Conversely, if $b^- > b_c$ then $I_n \rightarrow 0$ as $n \rightarrow \infty$, and if $\liminf_{n \rightarrow \infty} I_n > 0$ then $b^+ \leq b_c$.*

The next result arises from the fact that for $b < b'_c$ the main contribution to I_n , and hence to K_n or R_n , comes from the interior of A , while for $b'_c < b < b_c$ the main contribution to I_n comes from near the boundary of A . Given random variables $(Y_n)_{n \geq 1}$ we write $Y_n = o_{\mathbb{P}}(1)$ to mean $Y_n \rightarrow 0$ in probability as $n \rightarrow \infty$.

Theorem 2.3. *Under the conditions of Theorem 2.1, as $n \rightarrow \infty$ we have*

$$\zeta_n = n \exp(-\theta f_0 n r_n^d (1 + o_{\mathbb{P}}(1))) \quad \text{if } b^+ \leq b'_c; \quad (2.6)$$

$$\zeta_n = n^{1-1/d} \exp\left(-\theta f_1 n r_n^d \left(\frac{1}{2} + o_{\mathbb{P}}(1)\right)\right) \quad \text{if } b^- \geq b'_c. \quad (2.7)$$

In particular, if $b^+ = b^- = b$ then $\zeta_n = n^{1-\min(f_0 b, (1/d) + f_1 b/2) + o_{\mathbb{P}}(1)}$.

If $d = 2$ then since $f_1 \geq f_0$ we have $d(f_0 - f_1/2) \leq f_0$ and $b'_c \geq f_0^{-1} = b_c$. Thus, if $n r_n^2 \rightarrow \infty$ and $I_n \rightarrow \infty$, then (2.6) applies and Ganesan's result (1.5) for $A = [0, 1]^2$ holds whenever $c < \pi f_0$, and fails whenever $c > \pi f_0$ (in the latter case the probabilities in (1.5) tend to zero). A similar remark holds when $d \geq 3$, provided also $b^+ < b'_c$.

Next we give distributional results. The first one says that in the critical regime for connectivity, both $K_n - 1$ and R_n (along with $K'_n - 1$ and R'_n) are asymptotically Poisson.

Theorem 2.4 (Poisson convergence in the connectivity regime for general f). *Suppose f is continuous on A with $f_0 > 0$, and suppose that $I_n = \Theta(1)$ as $n \rightarrow \infty$. Let ξ_n denote any of $K_n - 1$, R_n , $K'_n - 1$ or R'_n . Then $d_{\text{TV}}(\xi_n, Z_{I_n}) = O((\log n)^{1-d})$ as $n \rightarrow \infty$. In particular, if $\lim_{n \rightarrow \infty} I_n = c$ for some $c \in (0, \infty)$, then $\xi_n \xrightarrow{D} Z_c$ as $n \rightarrow \infty$.*

Our next result demonstrates asymptotic normality of K'_n and of R'_n for $d \geq 2$, and of K_n and R_n for $d \geq 3$, in the whole of the mildly dense limiting regime for r .

Theorem 2.5 (Variance asymptotics and CLT for general f). *Suppose f is continuous on A with $f_0 > 0$, and that r_n satisfies (1.1) and also $I_n \rightarrow \infty$ as $n \rightarrow \infty$. Let ξ_n denote either $K_n - 1$ or R_n ; let ξ'_n denote either $K'_n - 1$ or R'_n . Then as $n \rightarrow \infty$ we have*

$$\text{Var}[\xi'_n] = I_n (1 + O((n r_n^d)^{(1-d)/2})); \quad (2.8)$$

$$d_K(I_n^{-1/2}(\xi'_n - \mathbb{E}[\xi'_n]), N(0, 1)) = O((n r_n^d)^{(1-d)/3} + I_n^{-1/2}). \quad (2.9)$$

If $d \geq 3$ then also

$$\text{Var}[\xi_n] = I_n (1 + O(n r_n^d)^{1-d/2}); \quad (2.10)$$

$$d_K(I_n^{-1/2}(\xi_n - \mathbb{E}[\xi_n]), N(0, 1)) = O((n r_n^d)^{(2-d)/3} + I_n^{-1/2}). \quad (2.11)$$

Remark 2.6. 1. In view of Theorem 2.4, our limiting regime for r_n for Theorems 2.1, 2.3 and 2.5 (namely, $n r_n^d \rightarrow \infty$ and $I_n \rightarrow \infty$) covers the whole range of limiting regimes between the thermodynamic regime and the critical regime for connectivity.

2. We can often simplify the expression (1.4) for μ_n depending on the logarithmic growth rate of $n r_n^d$. Indeed, if $d = 2$ or $b^+ f_0 < 2/d$ then $\mu_n \sim n e^{-n \theta f_0 r_n^d}$, and if $d \geq 3$ and $b^- f_0 > 2/d$ then $\mu_n \sim \theta_{d-1}^{-1} |\partial A| r_n^{1-d} e^{-n \theta f_0 r_n^d / 2}$.

2.2 Results for the uniform case

In the uniform case, we can replace I_n with the quantity μ_n defined at (1.4). Indeed, in Propositions 4.7 and 4.8 we shall show that in the uniform case, $I_n = \mu_n(1 + O((nr_n^2)^{-1/2}))$ as $n \rightarrow \infty$ if $d = 2$ and $I_n = \mu_n\left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right)$ as $n \rightarrow \infty$ if $d \geq 3$. Therefore in the convergence results arising from Theorems 2.1, 2.4 and 2.5 we can replace I_n with μ_n ; this gives us Theorem 1.1. The more quantitative versions of the results in Theorem 1.1, where we keep track of rates of convergence, go as follows.

Theorem 2.7 (First order results for the uniform case). *Suppose $f \equiv f_0 1_A$ with $f_0 = \lambda(A)^{-1}$. Let ξ_n denote any of $K_n - 1$, R_n , $K'_n - 1$ or R'_n , define γ_n as at (1.2) and define μ_n by (1.4).*

(a) *If $(r_n)_{n \geq 1}$ satisfies $|\gamma_n| = O(1)$ as $n \rightarrow \infty$, then $d_{\text{TV}}(\xi_n, Z_{\mu_n}) = O((\log n)^{-1/2})$ if $d = 2$ and $d_{\text{TV}}(\xi_n, Z_{\mu_n}) = O\left(\left(\frac{\log \log n}{\log n}\right)^2\right)$ if $d \geq 3$.*

(b) *If $\gamma_n \rightarrow \gamma \in \mathbb{R}$ as $n \rightarrow \infty$, then as $n \rightarrow \infty$, $\xi_n \xrightarrow{\mathcal{D}} Z_{e^{-\gamma}}$ if $d = 2$ and $\xi_n \xrightarrow{\mathcal{D}} Z_{c_{d,A} e^{-\gamma/2}}$ if $d \geq 3$, where we set $c_{d,A} := \theta_{d-1}^{-1}(\theta/(2 - 2/d))^{1-1/d} \sigma_A$ and $\sigma_A := |\partial A|/(\lambda(A)^{1-1/d})$.*

(c) *Suppose (1.1) and (1.2) hold. If $d = 2$ then as $n \rightarrow \infty$:*

$$\mathbb{E}[\xi_n] = \mu_n(1 + O((nr_n^2)^{-1/2})); \quad (2.12)$$

$$\mathbb{E}\left[\left|\frac{\xi_n}{\mu_n} - 1\right|\right] = O((nr_n^2)^{-1/2} + \mu_n^{-1/2}), \quad (2.13)$$

while if $d \geq 3$ then as $n \rightarrow \infty$:

$$\mathbb{E}[\xi_n] = \mu_n\left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right); \quad (2.14)$$

$$\mathbb{E}\left[\left|\frac{\xi_n}{\mu_n} - 1\right|\right] = O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2 + \mu_n^{-1/2}\right). \quad (2.15)$$

Theorem 2.8 (Variance asymptotics and CLT for the uniform case). *Suppose $f \equiv f_0 1_A$ with $f_0 = \lambda(A)^{-1}$. Suppose r_n satisfies (1.1) and (1.2), and define μ_n by (1.4). Let ξ_n denote either K_n or R_n , and let ξ'_n denote either K'_n or R'_n . If $d = 2$ then as $n \rightarrow \infty$:*

$$\text{Var}[\xi'_n] = \mu_n(1 + O((nr_n^2)^{-1/2})); \quad (2.16)$$

$$d_K(\mu_n^{-1/2}(\xi'_n - \mathbb{E}[\xi'_n]), N(0, 1)) = O((nr_n^2)^{-1/3} + \mu_n^{-1/2}). \quad (2.17)$$

If $d \geq 3$ then as $n \rightarrow \infty$:

$$\text{Var}[\xi'_n] = \mu_n \left(1 + O \left((nr_n^d)^{(1-d)/2} + \left(\frac{\log(nr_n^d)}{nr_n^d} \right)^2 \right) \right); \quad (2.18)$$

$$\text{Var}[\xi_n] = \mu_n \left(1 + O \left((nr_n^d)^{1-d/2} + \left(\frac{\log(nr_n^d)}{nr_n^d} \right)^2 \right) \right); \quad (2.19)$$

$$d_K(\mu_n^{-1/2}(\xi'_n - \mathbb{E}[\xi'_n]), N(0, 1)) = O \left((nr_n^d)^{(1-d)/3} + \left(\frac{\log(nr_n^d)}{nr_n^d} \right)^{4/3} + \mu_n^{-1/2} \right); \quad (2.20)$$

$$d_K(\mu_n^{-1/2}(\xi_n - \mathbb{E}[\xi_n]), N(0, 1)) = O \left((nr_n^d)^{(2-d)/3} + \left(\frac{\log(nr_n^d)}{nr_n^d} \right)^{4/3} + \mu_n^{-1/2} \right). \quad (2.21)$$

Remark 2.9. 1. In the uniform case, we have $f_0 = f_1$ and $b_c = (2 - 2/d)/f_0$.

2. From Theorem 2.7 we see for the uniform case that in the whole of the mildly dense regime both K_n and R_n scale like μ_n (and if $d = 2$ or $f_0 b^+ < 2/d$, like $n \exp(-nf_0 \theta r_n^d)$) in probability, rather than like a constant times n as given by (1.3) in the thermodynamic regime.
3. In the uniform case, if $nr_n^d \rightarrow \infty$ and $\limsup \gamma_n < \infty$, then by Propositions 4.7 and 4.8, $I_n \sim \mu_n$ as $n \rightarrow \infty$. Hence by (5.25), if $\gamma_n \rightarrow \gamma \in \mathbb{R}$ as $n \rightarrow \infty$, then $I_n \rightarrow e^{-\gamma}$ if $d = 2$ and $I_n \rightarrow c_{d,A} e^{-\gamma/2}$ if $d \geq 3$. Using this and the fact that I_n is decreasing in r_n while γ_n is increasing in r_n we can deduce that $\gamma_n \rightarrow -\infty$ if and only if $I_n \rightarrow \infty$, as claimed earlier.

2.3 Overview of proofs

The main insight behind our results is that the dominant contribution both for $K_n - 1$ and for R_n comes from the singletons, i.e. the isolated vertices. Let S_n , respectively S'_n , denote the number of singletons of $G(\mathcal{X}_n, r)$, resp. $G(\mathcal{P}_n, r_n)$. Our starting point for a proof of Theorem 2.5 is a similar collection of results for S_n and S'_n , of interest in their own right, which go as follows:

Proposition 2.10 (Results on singletons). *Suppose f is continuous on A with $f_0 > 0$, and that r_n satisfies (1.1) and also $I_n \rightarrow \infty$ as $n \rightarrow \infty$. Let ζ_n be either S_n or S'_n . Then there exists $\delta > 0$ such that as $n \rightarrow \infty$ we have $\mathbb{E}[\zeta_n] = I_n(1 + O(e^{-\delta nr_n^d}))$, and $\text{Var}[\zeta_n] = I_n(1 + O(e^{-\delta nr_n^d}))$, and also*

$$d_K(I_n^{-1/2}(S'_n - I_n), N(0, 1)) = O(e^{-\delta nr_n^d} + I_n^{-1/2}); \quad (2.22)$$

$$\text{if } \partial A \in C^2, \quad d_K(\tilde{I}_n^{-1/2}(S_n - \mathbb{E}[S_n]), N(0, 1)) = O(e^{-\delta nr_n^d} + I_n^{-1/2}). \quad (2.23)$$

Proposition 2.10 extends results in [12], where the same conclusions are derived under the extra condition $b^+ < 1/\max(f_0, d(f_0 - f_1/2))$ rather than the weaker condition $I_n \rightarrow \infty$ that we consider here.

To get from Proposition 2.10 to Theorem 2.5, let ξ_n be either $K_n - 1$ or R_n and ξ'_n be either K'_n or R'_n . We show that both the mean and the variance of both $\xi_n - S_n$ and

$\xi'_n - S'_n$, are asymptotically negligible relative to I_n . To do this we deal separately with the contribution to $\xi_n - S_n$ or $\xi'_n - S'_n$ from components with Euclidean diameters that categorized as ‘small’, ‘medium’ or ‘large’ compared to r_n , using different arguments for the three different categories. This requires us to deal with a lot of different cases, as a result of which Section 6, containing the second moment estimates, is quite long (the proofs of the first order results can be read without referring to that section). Once we have the moment estimates, we can derive the ‘quantitative’ CLT for ξ_n or ξ'_n from the one for S_n or S'_n by using a quantitative version of Slutsky’s theorem.

Our argument for small components has geometrical ingredients (presented in Section 3.1) and takes boundary effects into account. The argument for large components involves discretization and path-counting arguments seen in continuum percolation theory. The argument for medium-sized components involves both geometry and discretization.

To derive our results with more explicit constants in the uniform case (Theorems 2.7 and 2.8) we need to demonstrate asymptotic equivalence of I_n and μ_n . This can be done by adapting the method of polytopal approximation that was recently used for related problems in [4], [10] and [11]. We do this in Section 4.3.

The rest of the paper is organised as follows. After some preliminary lemmas in Section 3, in Section 4 we give an asymptotic analysis of S_n and S'_n , and of I_n , in particular proving Propositions 2.2 and 2.10.

In Section 5 we give estimates of $\mathbb{E}[\zeta_n - S_n]$ and $\mathbb{E}[\zeta'_n - S'_n]$, where ζ_n is either $K_n - 1$ or R_n , and ζ'_n is either $K'_n - 1$ or R'_n . where ζ_n is either $K_n - 1$, and then conclude the proof of Theorems 2.1, 2.3 and 2.4. In Section 6 we complete the proof of Theorems 2.5 and 2.8.

3 Preliminaries

Throughout the rest of this paper we assume that the density f of the measure ν is supported by a compact set $A \subset \mathbb{R}^d$ with $f_0 := \inf_A f > 0$, and that f is continuous on A , so that $f_{\max} := \sup_A f$ is finite. Moreover we assume either that $d = 2$ and $A = [0, 1]^2$, or that $d \geq 2$ and $A \subset \mathbb{R}^d$ is compact and connected with $A = \overline{A^\circ}$ and $\partial A \in C^2$. Also we assume $r_n \in (0, \infty)$ is given for all $n \geq 1$.

Given $D, D' \subset \mathbb{R}^d$, we set $D \oplus D' := \{x + y : x \in D, y \in D'\}$, the Minkowski sum of D and D' . Also let D° denote the interior of D . Let o denote the origin in \mathbb{R}^d . Let $\|\cdot\|$ denote the Euclidean norm on \mathbb{R}^d . Given $a > 0$ we set $aD := \{ax : x \in D\}$. Also we set $D^{(-a)} := \{x \in D : B(x, a) \subset D\}$. Given $n \in \mathbb{N}$, we write $[n]$ for the set $\{1, \dots, n\}$.

We introduce an ordering \prec on A : for $x, y \in A$, if $\partial A \in C^2$ we say $x \prec y$ if either $\text{dist}(x, \partial A) < \text{dist}(y, \partial A)$ (using the Euclidean distance) or $\text{dist}(x, \partial A) = \text{dist}(y, \partial A)$ and x precedes y (strictly) in the lexicographic ordering. If $A = [0, 1]^2$ we say $x \prec y$ if the ℓ_1 distance from x to the nearest corner of A is less than that of y , or if these two distances are equal and x precedes y lexicographically. In either case, given $x \in A$ we write $A_x := \{y \in A : x \prec y\}$.

For non-empty $U \subset \mathbb{R}^d$ set $\text{diam}(U) := \sup_{x, y \in U} \{\|x - y\|\}$, and let $\#(U)$ denote the number of elements of U .

3.1 Geometrical tools

Lemma 3.1. *There exist $\delta_1 > 0$ and $s_0 > 0$ depending on d and A such that if $s \in (0, s_0)$ and $x, y \in A$ with $x \prec y$, then*

$$\lambda(A \cap B_s(y) \setminus B_s(x)) \geq 2\delta_1 s^d \quad \text{if } \|y - x\| \geq s; \quad (3.1)$$

$$\lambda(A \cap B_s(y) \setminus B_s(x)) \geq 2\delta_1 s^{d-1} \|y - x\| \quad \text{if } \|y - x\| \leq 3s, \quad (3.2)$$

and if $\partial A \in C^2$ then (3.1) still holds if we drop the condition $x \prec y$.

Note that when $A = [0, 1]^2$ we do require $x \prec y$ for (3.1); otherwise y could be ‘jammed into a corner’ of A , for example when x is near $(2^{-1/2}s, 2^{-1/2}s)$.

Proof. Note first that it suffices to prove the second inequality (3.2) for $\|y - x\| \leq s$, since it can be proved in the case $s \leq \|x - y\| \leq 3s$ by using the first inequality (3.1) and changing δ_1 .

In the case with $\partial A \in C^2$, (3.1) comes from [12, Lemma 5.9], which does not require the condition $x \prec y$, while (3.2) comes from [4, Lemma 3.2].

Now suppose $A = [0, 1]^2$. Without loss of generality, the nearest corner of A to x is the origin. Writing $x = (x_1, x_2)$ and $y = (y_1, y_2)$, assume also without loss of generality that $x_2 \leq y_2$. Also assume $y_1 \leq x_1$ (otherwise (3.1) and (3.2) are easy to see).

If $\|y - x\| \geq s$ then $y_2 \geq x_2 + 2^{-1/2}s$ (otherwise the condition $x \prec y$ fails). Then the ball of radius $0.05s$ centred on $(y_1 + 0.05s, y_2 + 0.8s)$ is contained in $A \cap B_s(y) \setminus B_s(x)$, and (3.1) follows for this case.

For (3.2), we assume without loss of generality that $\|y - x\| \leq s$. Consider the segment S of $B(x, s)$ that is cut off from $B(x, s)$ by the line parallel to $[x, y]$ and at a distance $2^{-1/2}s$ from x , away from the origin. Then as illustrated in Figure 1, $S \oplus \{y - x\} \subset A$, and by Fubini’s theorem there is a constant $\delta_1 > 0$ such that $\lambda((S \oplus \{y - x\}) \setminus S) \geq 2\delta_1 s \|y - x\|$. \square

Lemma 3.2. *Let $0 < \varepsilon < K < \infty$. Then there exists $\delta_2 = \delta_2(\varepsilon, K)$, $s_0 > 0$ depending on ε, K, d and A , such that for all $s \in (0, s_0)$ and all compact $B \subset A$ with $\text{diam } B \in [\varepsilon s, Ks]$ and $x_0 \in B$ with $x_0 \prec y$ for all $y \in B$, we have*

$$\lambda((B \oplus B_s(o)) \cap A) \geq \lambda(B) + \lambda(B_s(x_0) \cap A) + 2\delta_2 s^d. \quad (3.3)$$

Proof. In the case with $\partial A \in C^2$, we can use [11, Lemma 2.5].

If instead $d = 2$ and $A = [0, 1]^2$, we can argue similarly for x not close to any corner of A . In the other case we can use [7, Proposition 5.15]. \square

Lemma 3.3. *Let $\varepsilon > 0$. Suppose $\partial A \in C^2$. There exists $s_0 > 0$ depending on d, A and ε such that if $s \in (0, s_0)$ and $y \in A, z \in \partial A$, then*

$$\lambda(A \cap B_s(y)) \geq ((1/2) - \varepsilon)\theta s^d; \quad (3.4)$$

$$\lambda(A \cap B_s(z)) \leq ((1/2) + \varepsilon)\theta s^d. \quad (3.5)$$

If instead $d = 2$ and $A = [0, 1]^2$ there exists $s_0 > 0$ such that if $y \in A, s \in (0, s_0)$ then $\lambda(A \cap B_s(y)) \geq (\pi/4)s^2$.




Figure 1: The shaded region is $(S \oplus \{y - x\}) \setminus S$, as described in the proof of Lemma 3.1.

Proof. The first inequality (3.4) comes from [7, Lemma 5.6] or [4, Lemma 3.2].

The second inequality (3.5) comes from [7, Lemma 5.6]; alternatively it can be proved by repeatedly using [11, Lemma 2.1(ii)].

The third inequality is obvious. □

3.2 Probabilistic tools

Lemma 3.4 (Chernoff bounds). *Suppose $n \in \mathbb{N}$, $p \in (0, 1)$, $t > 0$ and $0 < k < n$.*

(i) *If $k \geq e^2 np$ then $\mathbb{P}[\text{Bin}(n, p) \geq k] \leq \exp(-(k/2) \log(k/(np))) \leq e^{-k}$.*

(ii) *For all t large, $\mathbb{P}[Z_t \geq t + t^{3/4}] \leq \exp(-\sqrt{t}/9)$ and $\mathbb{P}[Z_t \leq t - t^{3/4}] \leq \exp(-\sqrt{t}/9)$.*

(iii) *If $k \geq e^2 t$ then $\mathbb{P}[Z_t \geq k] \leq e^{-k}$.*

Proof. See e.g. [7, Lemmas 1.1, 1.2 and 1.4]. □

Let $\mathbf{N}(\mathbb{R}^d)$ be the space of all finite subsets of \mathbb{R}^d , equipped with the smallest σ -algebra $\mathcal{S}(\mathbb{R}^d)$ containing the sets $\{\mathcal{X} \in \mathbf{N}(\mathbb{R}^d) : |\mathcal{X} \cap B| = m\}$ for all Borel $B \subset \mathbb{R}^d$ and all $m \in \mathbb{N} \cup \{0\}$. Given $F : \mathbf{N}(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $x \in \mathbb{R}^d$, define the *add-one cost* $D_x F(\mathcal{X}) := F(\mathcal{X} \cup \{x\}) - F(\mathcal{X})$ for all $\mathcal{X} \in \mathbf{N}(\mathbb{R}^d)$. Also define $D^+ F(\mathcal{X}) := \max(D_x F(\mathcal{X}), 0)$ and $D^- F(\mathcal{X}) := \max(-D_x F(\mathcal{X}), 0)$, the positive and negative parts of $D_x F(\mathcal{X})$.

Lemma 3.5 (Poincaré and Efron-Stein inequalities). *Suppose $F : \mathbf{N}(\mathbb{R}^d) \rightarrow \mathbb{R}$ is measurable and $n > 0$. If $\mathbb{E}[F(\mathcal{P}_n)^2] < \infty$ then*

$$\text{Var}[F(\mathcal{P}_n)] \leq n \int_A \mathbb{E}[|D_x F(\mathcal{P}_n)|^2] \nu(dx). \quad (3.6)$$

Also, if $n \in \mathbb{N}$ and $\mathbb{E}[F(\mathcal{X}_n)^2] < \infty$ then

$$\text{Var}[F(\mathcal{X}_n)] \leq n \int_A \mathbb{E}[|D_x F(\mathcal{X}_{n-1})|^2] \nu(dx). \quad (3.7)$$

Proof. The first assertion (3.6) is the Poincaré inequality [5, Theorem 18.7]. For the second assertion (3.7), we use Efron and Stein's jackknife estimate for the variance of functions of iid random variables. Let $\tilde{F}_n : \mathbb{R}^{dn} \rightarrow \mathbb{R}$ be given by $\tilde{F}_n((x_1, \dots, x_n)) = F(\{x_1, \dots, x_n\})$ for all $x_1, \dots, x_n \in \mathbb{R}$. Then \tilde{F}_n is measurable. The Efron-Stein inequality (see e.g. [2]) says that

$$\mathrm{Var}[\tilde{F}_n(\mathbf{X}_n)] \leq \frac{1}{2} \sum_{i=1}^n \mathbb{E}[(\tilde{F}_n(\mathbf{X}_n) - \tilde{F}_n(\mathbf{X}_{n+1}^{(i)}))^2] \quad (3.8)$$

where $\mathbf{X}_n := (X_1, \dots, X_n)$ and $\mathbf{X}_n^{(i)} := (X_1, \dots, X_{i-1}, X_{i+1}, \dots, X_n)$.

We write $\tilde{F}_n(\mathbf{X}_n) - \tilde{F}_n(\mathbf{X}_{n+1}^{(i)}) = \tilde{F}_n(\mathbf{X}_n) - \tilde{F}_{n-1}(\mathbf{X}_n^{(i)}) - (\tilde{F}_n(\mathbf{X}_{n+1}^{(i)}) - \tilde{F}_{n-1}(\mathbf{X}_n^{(i)}))$. By the bound $(a+b)^2 \leq 2a^2 + 2b^2$ (which comes from Jensen's inequality), and (3.8), and the exchangeability of (X_1, \dots, X_{n+1}) ,

$$\mathrm{Var}(F(\mathcal{X}_n)) = \mathrm{Var}[\tilde{F}_n(\mathbf{X}_n)] \leq n\mathbb{E}[(\tilde{F}_n(\mathbf{X}_n) - \tilde{F}_{n-1}(\mathbf{X}_{n-1}))^2],$$

and (3.7) follows. \square

Lemma 3.6 (Quantitative version of Slutsky's theorem). *Suppose X and Y are random variables on the same probability space with $\mathbb{E}[Y] = 0$ and $\mathrm{Var}[Y] < \infty$. Then $d_K(X + Y, N(0, 1)) \leq 3(d_K(X, N(0, 1)) + (\mathrm{Var}[Y])^{1/3})$.*

Proof. Let $t \in \mathbb{R}$ and set $a := (\mathrm{Var}[Y])^{1/3}$. Then by the union bound and Chebyshev's inequality

$$\begin{aligned} |\mathbb{P}[X + Y \leq t] - \mathbb{P}[X \leq t]| &\leq \mathbb{P}[\{X + Y \leq t\} \Delta \{X \leq t\}] \\ &\leq \mathbb{P}[t - a < X \leq t + a] + \mathbb{P}[|Y| \geq a] \\ &\leq \mathbb{P}[t - a < N(0, 1) \leq t + a] + 2d_K(X, N(0, 1)) + a^{-2}\mathrm{Var}[Y] \\ &\leq 3a + 2d_K(X, N(0, 1)), \end{aligned}$$

and the result follows. \square

3.3 Percolation type estimates

For finite $\mathcal{X} \subset \mathbb{R}^d$, and $x \in \mathcal{X}$ and $s > 0$, let $\mathcal{C}_s(x, \mathcal{X})$ denote the vertex set of the component of $G(\mathcal{X}, s)$ containing x , and $|\mathcal{C}_s(x, \mathcal{X})|$ the number of elements of $\mathcal{C}_s(x, \mathcal{X})$ (i.e. the order of this component).

To prove our theorems, we shall need to establish uniqueness of the giant component in $G(\mathcal{X}_n, r)$ or $G(\mathcal{P}_n, r)$ (with $r = r(n)$). The next two lemmas help do this, and are proved using discretization and path-counting (Peierls) arguments of the sort used in the theory of continuum percolation.

The first lemma says that if $nr_n^d \rightarrow \infty$ as $n \rightarrow \infty$, the existence of two components of diameter drastically larger than r_n is extremely unlikely for n large. Throughout, the diameter of a component means the Euclidean (rather than graph-theoretic) diameter of its set of vertices.

Bounds of this sort also arise in the study of connectivity thresholds (which concerns the regime with $I_n \rightarrow c \in (0, \infty)$); see for instance [8, Proposition 3.2]. In the proof, we shall invoke a topological lemma from [8].

Lemma 3.7 (Uniqueness of the large component). *Suppose $(r_n)_{n \geq 1}$ satisfies $nr_n^d \rightarrow \infty$ as $n \rightarrow \infty$. Let ϕ_n be given with $\phi_n \geq \log n$ for all $n \geq 1$ and assume $\phi_n r_n \rightarrow 0$ as $n \rightarrow \infty$. Let \mathcal{U}_n , respectively $\tilde{\mathcal{U}}_n$, denote the event that there exists at most one component of $G(\mathcal{P}_n, r_n)$ (respectively $G(\mathcal{X}_n, r_n)$) with diameter larger than $\phi_n r_n$. Then there exists $c > 0$ depending only on d, A and f such that for all n large enough,*

$$\mathbb{P}[\mathcal{U}_n^c] \leq \exp(-c\phi_n nr_n^d); \quad \mathbb{P}[\tilde{\mathcal{U}}_n^c] \leq \exp(-c\phi_n nr_n^d). \quad (3.9)$$

Proof. First assume that $\partial A \in C^2$. Let $\varepsilon = 1/(99\sqrt{d})$. Given n , partition \mathbb{R}^d into cubes $(Q_{n,i})$ of side length εr_n indexed by \mathbb{Z}^d . To be definite, for $i = (i_1, \dots, i_d) \in \mathbb{Z}^d$, set $Q_{n,i} := \prod_{k=1}^d ((i_k - 1)\varepsilon r_n, i_k \varepsilon r_n]$. We say a set $\sigma \subset \mathbb{Z}^d$ is $*$ -connected if the closure of the union of cubes $Q_{n,i}, i \in \sigma$ is connected. By the deterministic topological lemma [8, Lemma 3.5], there exist $\alpha, \alpha' > 0, n_1 \in \mathbb{N}$ such that for all $n \geq n_1$ and for any finite $\mathcal{X} \subset A$, if U and V are the vertex sets of two components of $G(\mathcal{X}, r_n)$, then there exists a $*$ -connected set $\sigma \subset \mathbb{Z}^d$ enjoying the following properties:

- i) $\mathcal{X} \cap (\cup_{i \in \sigma} Q_{n,i}) = \emptyset$;
- ii) $|\{i \in \sigma : Q_{n,i} \subset A\}| \geq \alpha \#(\sigma)$;
- iii) $\varepsilon r_n \#(\sigma) \geq \min(d^{-1/2} \text{diam}(U), d^{-1/2} \text{diam}(V), \alpha')$.

In (iii), the factor of $d^{-1/2}$ arises because if diam_∞ denotes diameter in the ℓ_∞ sense (as used in [8]) then $\text{diam}_\infty(\cdot) \geq d^{-1/2} \text{diam}(\cdot)$.

We shall apply this lemma to \mathcal{C}_1 and \mathcal{C}_2 which we define to be the vertex sets of the largest and second-largest component (in terms of Euclidean diameter) of $G(\mathcal{P}_n, r_n)$, with diameter ℓ_1, ℓ_2 respectively (so $\ell_1 \geq \ell_2$).

For $n > 0, k \in \mathbb{N}$, define

$$\mathcal{K}_{n,k,\alpha} := \{\sigma \subset \mathbb{Z}^d : \#(\sigma) = k, \sigma \text{ is } * \text{-connected, } \#\{i \in \sigma : Q_{n,i} \subset A\} \geq \alpha k\}, \quad (3.10)$$

and define the events

$$\mathcal{G}_{n,k} := \cup_{\sigma \in \mathcal{K}_{n,k,\alpha}} \{\mathcal{P}_n \cap (\cup_{i \in \sigma} Q_{n,i}) = \emptyset\}; \quad \tilde{\mathcal{G}}_{n,k} := \cup_{\sigma \in \mathcal{K}_{n,k,\alpha}} \{\mathcal{X}_n \cap (\cup_{i \in \sigma} Q_{n,i}) = \emptyset\}. \quad (3.11)$$

If event \mathcal{U}_n^c occurs, then $\ell_2 \geq r_n \phi_n$, so by the lemma, $\mathcal{G}_{n,k}$ occurs for some $k \geq \varepsilon^{-1} d^{-1/2} \phi_n \geq \phi_n$. By a Peierls argument (e.g. [7, Lemma 9.3]) there exists $c = c(d, A) > 0$ such that the family of $*$ -connected sets $\sigma \subset \mathbb{Z}^d$ with $\#(\sigma) = k$ and with $Q_{n,i} \cap A \neq \emptyset$ for some $i \in \sigma$ has cardinality at most $cr_n^{-d} e^{ck}$, which is at most ne^{ck} , provided n is large enough, by the condition $nr_n^d \rightarrow \infty$. Thus by the union bound, for n large enough we have

$$\mathbb{P}[\mathcal{G}_{n,k}^c] \leq n \exp(ck - k\alpha n f_0(\varepsilon r_n)^d) \leq n \exp(-(\alpha f_0 \varepsilon^d / 2) k n r_n^d), \quad (3.12)$$

where we used that $nr_n^d \rightarrow \infty$ again for the last inequality. The same bound holds for $\tilde{\mathcal{G}}_{n,k}$, since the probability of a binomial random quantity taking the value zero is bounded above by the corresponding probability for a Poisson random quantity with the same mean.

By (3.12), for n large enough

$$\begin{aligned} \mathbb{P}[\mathcal{U}_n^c] &\leq \sum_{k \geq \phi_n} \mathbb{P}[\mathcal{G}_{n,k}] \leq 2n \exp(-(\alpha f_0 \varepsilon^d / 2) n r_n^d \phi_n) \\ &\leq \exp(-(\alpha f_0 \varepsilon^d / 4) n r_n^d \phi_n), \end{aligned}$$

where we used the conditions $n r_n^d \rightarrow \infty$ and $\phi_n \geq \log n$, for the last inequality. This gives us the first assertion in (3.9), and the second assertion is obtained similarly using $\tilde{\mathcal{G}}_{n,k}$.

In the case where $A = [0, 1]^2$, we can argue similarly (see [7, Lemma 13.5]). We should now take ε so that the cubes $Q_{n,i}$ fit exactly in the unit cube, which means ε needs to vary with n but we can take $\varepsilon(n)$ to satisfy this condition as well as $\varepsilon \in [1/(99\sqrt{d}), 1/(98\sqrt{d})]$ for all large enough n , and the preceding argument still works. \square

We next provide a bound on the probability of existence of a moderately large components of $G(\mathcal{P}_n, r_n)$ near a given location in A , again measuring ‘size’ of a component \mathcal{C} by the Euclidean diameter of its vertex set $V(\mathcal{C})$. For $x, y, z \in \mathbb{R}^d$ and \mathcal{X} a finite set of points in \mathbb{R}^d , we use the notation

$$\mathcal{X}^x := \mathcal{X} \cup \{x\}; \quad \mathcal{X}^{x,y} := \mathcal{X} \cup \{x, y\}; \quad \mathcal{X}^{x,y,z} := \mathcal{X} \cup \{x, y, z\}. \quad (3.13)$$

Suppose $0 \leq \varepsilon < K \leq \infty$. Given $(r_n)_{n \geq 1}$ we define events

$$\mathcal{M}_{n,\varepsilon,K}(x, \mathcal{X}) := \{\varepsilon r_n < \text{diam}(\mathcal{C}_{r_n}(x, \mathcal{X}^x)) \leq K r_n\}; \quad (3.14)$$

$$\mathcal{M}_{n,\varepsilon,K}^*(x, \mathcal{X}) := \cup_{y \in \mathcal{X} \cap B_{r_n}(x)} \mathcal{M}_{n,\varepsilon,K}(y, \mathcal{X}). \quad (3.15)$$

Lemma 3.8 (Non-existence of moderately large components near a fixed site). *Suppose $n r_n^d \rightarrow \infty$ and $n^{1/(2d)} r_n \rightarrow 0$ as $n \rightarrow \infty$. There exist $\beta, n_1 \in (0, \infty)$ such that for all $n \geq n_1$ and all $x, y \in A$, all $\rho \in [1, n^{1/(2d)}]$, with ξ_n representing any of \mathcal{P}_n , $\mathcal{P}_n \cup \{y\}$, \mathcal{X}_{n-1} , $\mathcal{X}_{n-2} \cup \{y\}$ or $\mathcal{X}_{n-3} \cup \{y\}$, we have*

$$\mathbb{P}[\mathcal{M}_{n,\rho,n^{1/(2d)}}(x, \xi_n)] \leq \exp(-\beta \rho n r_n^d); \quad (3.16)$$

$$\mathbb{P}[\mathcal{M}_{n,\rho,n^{1/(2d)}}^*(x, \xi_n)] \leq \exp(-\beta \rho n r_n^d). \quad (3.17)$$

Proof. Suppose $\xi_n = \mathcal{P}_n$. Assume for now that $\partial A \in C^2$. As in the previous proof, given n we partition \mathbb{R}^d into cubes $Q_{n,i}$ of side εr_n with $\varepsilon = 1/(99\sqrt{d})$. For $n > 0, k \in \mathbb{N}$, with $\mathcal{K}_{n,k,\alpha}$ defined at (3.10) define

$$\mathcal{K}_{n,k,\alpha,x} = \{\sigma \in \mathcal{K}_{n,k,\alpha} : \varepsilon r_n \sigma \cap B_{r_n}(x) \neq \emptyset \text{ or } \cup_{i \in \sigma} \overline{Q_{n,i}} \text{ surrounds } x\},$$

where we say a set $D \subset \mathbb{R}^d$ *surrounds* x if x lies in a bounded component of $\mathbb{R}^d \setminus D$, and $a\sigma := \{az : z \in \sigma\}$ for $a > 0$, and \overline{D} denotes the closure of a set D . Define the event

$$\mathcal{G}_{n,k,x} := \cup_{\sigma \in \mathcal{K}_{n,k,\alpha,x}} \{\mathcal{P}_n \cap (\cup_{i \in \sigma} Q_{n,i}) = \emptyset\}.$$

Let $\rho > 0$. Suppose now that $\mathcal{M}_{n,\rho,n^{1/(2d)}}(x, \mathcal{P}_n)$ occurs, and let $\mathcal{C} := \mathcal{C}_{r_n}(x, \mathcal{P}_n^x)$. In this case we have to look into the proof of [7, Lemma 13.9] (or that of [8, Lemma 3.5], which is the same). There, the set we are calling A is called Ω . In the preamble to that

lemma, a sequence of sets $\Omega_1, \Omega_2, \Omega_3$ is defined; each of these sets is a connected union of closed hypercubes, with $\Omega_i \subset \Omega_{i+1}^o$ for $i = 1, 2$, with $\Omega_3 \subset A^o$, and also with the ‘interior’ set Ω_I (defined in [7]) contained in Ω_1 .

Set $\mathcal{C}' = \mathcal{C} \oplus B_{r_n/2}(o)$; then \mathcal{C}' is a connected compact set. Let \mathcal{D} be the closure of the unbounded component of $\mathbb{R}^d \setminus \mathcal{C}'$, and let $\partial\mathcal{C}' := \mathcal{C}' \cap \mathcal{D}$, which is the external boundary of \mathcal{C}' . Let Σ denote the collection of $i \in \mathbb{Z}^d$ such that $Q_{n,i} \cap \partial\mathcal{C}' \neq \emptyset$.

Then $\partial\mathcal{C}'$ is connected by the unicoherence of \mathbb{R}^d (see e.g. [7]), so Σ is $*$ -connected. Also $\mathcal{P}_n \cap Q_{n,i} = \emptyset$ for all $i \in \Sigma$. Moreover, since $\text{diam}(\mathcal{C}) \leq n^{1/(2d)}r_n$ and $\mathcal{C} \cap B_{r_n}(x) \neq \emptyset$, we have that $\cup_{i \in \Sigma} Q_{n,i} \subset B_{2n^{1/(2d)}r_n}(x)$.

We claim that $\cup_{i \in \Sigma} \overline{Q_{n,i}}$ surrounds x . Indeed, $x \notin \cup_{i \in \Sigma} \overline{Q_{n,i}}$ since $\text{dist}(x, \partial\mathcal{C}') \geq r_n/2$. Since $x \in \mathcal{C} \subset \mathcal{C}'$, any path from x to a point in \mathcal{D} must pass through a point in $\partial\mathcal{C}'$, and the claim follows.

Since $n^{1/(2d)}r_n \rightarrow 0$ as $n \rightarrow \infty$, if $x \in \Omega_3$ then (provided n is large enough) we have $B_{2n^{1/(2d)}r_n}(x) \subset A$ and hence $\cup_{i \in \Sigma} Q_{n,i} \subset A$. If $x \notin \Omega_3$ then (provided n is large enough) we have $\mathcal{C} \cap \Omega_2 = \emptyset$, and by following the second case considered in the proof of [7, Lemma 13.9], we see that $\Sigma \in \mathcal{K}_{n,k,\alpha,x}$ for some k . Also $|\Sigma| \geq \varepsilon^{-1}d^{-1/2}\rho \geq \rho$.

Thus in either case if $\mathcal{M}_{n,\rho,n^{1/(2d)}}(x, \mathcal{P}_n)$ occurs, then event $\mathcal{G}_{n,k}$ (defined at (3.11)) occurs for some $k \geq \rho$. Since there are constants c, c' such that the family of $*$ -connected sets $\sigma \subset \mathbb{Z}^d$ with $\#(\sigma) = k$ and with $\mathbb{R}^d \setminus \overline{\cup_{i \in \sigma} Q_{n,i}}$ surrounding x has cardinality at most $c'e^{ck}$, for n large enough we have

$$\mathbb{P}[\mathcal{G}_{n,k}] \leq c' \exp(ck - k\alpha n f_0(\varepsilon r_n)^d) \leq c' \exp(-(\alpha f_0 \varepsilon^d/2)knr_n^d).$$

By summing over $k \geq \rho$, we deduce that for n large enough,

$$\mathbb{P}[\mathcal{M}_{n,\rho,n^{1/(2d)}}(x, \mathcal{P}_n)] \leq 2c' \exp(-(\alpha f_0 \varepsilon^d/2)\rho nr_n^d).$$

Taking $\beta = \alpha f_0 \varepsilon^d/4$, we obtain (3.16). Then using Markov’s inequality, the Mecke formula (see e.g. [5]) and (3.16) we can deduce that

$$\begin{aligned} \mathbb{P}[\mathcal{M}_{n,\rho,n^{1/(2d)}}^*(x, \mathcal{P}_n)] &\leq n \int_{B_{r_n}(x)} \mathbb{P}[\mathcal{M}_{n,\rho,n^{1/(2d)}}(y, \mathcal{P}_n)] \nu(dy) \\ &= O(nr_n^d \exp(-\beta \rho nr_n^d)), \end{aligned}$$

and on taking a smaller value of β we obtain (3.17).

In the case where $A = [0, 1]^2$, we can argue similarly (see [7, Lemma 13.5]). We should now take ε so that the cubes $Q_{n,i}$ fit exactly in the unit cube, which means ε needs to vary with n but we can take $\varepsilon(n)$ to satisfy this condition as well as $\varepsilon \in [1/(99\sqrt{d}), 1/(98\sqrt{d})]$ for all large enough n , and the preceding argument still works.

We can prove the results for the other choices of ξ_n in the statement of the lemma, by similar arguments. \square

We shall use crossing estimates from the theory of continuum percolation. Consider a homogeneous Poisson process \mathcal{H}_α in \mathbb{R}^d with intensity α . For each $s > 0$, let $\mathcal{H}_{\alpha,s} = \mathcal{H}_\alpha \cap C_s$ where $C_s := [0, s]^d$. We say that the graph $G(\mathcal{H}_{\alpha,s}, 1)$ crosses the cube C_s in the first coordinate (for short: $\text{Cross}_1(s)$ occurs) if there exists a component of $G(\mathcal{H}_{\alpha,s}, 1)$

such that its vertex set \mathcal{C} satisfies $(\mathcal{C} \oplus B_{1/2}(o)) \cap (\{0\} \times [0, s]^{d-1}) \neq \emptyset$ and $(\mathcal{C} \oplus B_{1/2}(o)) \cap (\{s\} \times [0, s]^{d-1}) \neq \emptyset$, namely, we can find a path contained in $\mathcal{C} \oplus B_{1/2}(o)$ which connects two opposite faces of C_s along the first coordinate. Define similarly $\text{Cross}_k(s)$ for each $k \in \{2, \dots, d\}$. We say $\text{Cross}(s)$ occurs if $\text{Cross}_k(s)$ occurs for all $k \in [d]$. Observe that the crossing event defined above is slightly different from the one in Meester and Roy [6] where a crossing in the first coordinate is said to occur if there is a path in $(\mathcal{H}_\alpha \oplus B_{1/2}(o)) \cap C_s$ connecting two opposite faces of C_s along the first coordinate. In other words, in [6], one is allowed to use all the Poisson points to construct a crossing path in C_s , while in our setting, one is restricted to the Poisson points in C_s .

A fundamental fact about continuum percolation is the existence of $\alpha_c \in (0, \infty)$ such that, as $s \rightarrow \infty$, $\mathbb{P}_\alpha[\text{Cross}(s)] \rightarrow 1$ for $\alpha > \alpha_c$ and $\mathbb{P}_\alpha[\text{Cross}(s)] \rightarrow 0$ for $\alpha < \alpha_c$. Here \mathbb{P}_α denotes the law of \mathcal{H}_α . For our purpose, we are concerned with the super-critical phase $\alpha > \alpha_c$. The following estimate taken from [7] quantifies the convergence of the crossing probabilities.

Lemma 3.9 ([7, Lemma 10.5 and Proposition 10.6]). *Let $d \geq 2$ and $\alpha > \alpha_c$. Then there exists a finite constant $c > 0$ such that for all $s \geq 1$,*

$$1 - \mathbb{P}_\alpha[\text{Cross}_1(s)] \leq e^{-cs}.$$

From this we derive a bound for the probability of having a small giant component. Again in the next result, diam refers to the Euclidean metric diameter of the vertex set of a component.

Given finite nonempty $\mathcal{X} \subset \mathbb{R}^d$ and $n \geq 1$, let $\mathcal{L}_n(\mathcal{X})$ and $\mathcal{L}_{n,2}(\mathcal{X})$ denote the vertex set of the component of $G(\mathcal{X}, r_n)$ with largest order and second largest order, respectively (setting $\mathcal{L}_{n,2}(\mathcal{X})$ to be empty if the graph is connected). Choose the left-most one if there is a tie.

Lemma 3.10. *Suppose $nr_n^d \rightarrow \infty$ and $nr_n^d = O(\log n)$ as $n \rightarrow \infty$. Then there exist constants $\delta_3, n_1 \in (0, \infty)$ such that for all $n \geq n_1$, with ξ_n denoting either \mathcal{P}_n or \mathcal{X}_n ,*

$$\mathbb{P}[\text{diam}(\mathcal{L}_n(\xi_n)) < (\log n)^2 r_n] \leq \exp(-\delta_3(n/\log n)^{1/d}). \quad (3.18)$$

Proof. First we show there exist constants $\delta, c' \in (0, \infty)$ such that for all large enough n ,

$$\mathbb{P}[\#(\mathcal{L}_n(\xi_n)) < \delta r_n^{-1}] \leq \exp(-c'(n/\log n)^{1/d}); \quad (3.19)$$

Without loss of generality we can and do choose $\delta > 0$ such that $C_{2\delta} := [0, 2\delta]^d \subset A$. Define the event

$$\mathcal{Y}_n := \{G(\mathcal{P}_{e^{-2n}} \cap C_{2\delta}, r_n) \text{ crosses } C_{2\delta} \text{ in the first coordinate}\}.$$

Since $\mathcal{P}_{e^{-2n}} \subset \mathcal{P}_n$, we have that $\mathcal{Y}_n \subset \{\#(\mathcal{L}_n(\mathcal{P}_n)) \geq \delta/r_n\}$ for n large.

Clearly the graph $G(\mathcal{P}_{e^{-2n}} \cap C_{2\delta}, r_n)$ is isomorphic to $G(r_n^{-1}(\mathcal{P}_{e^{-2n}} \cap C_{2\delta}), 1)$. Also $e^{-2n} f_0 r_n^d > \alpha_c + 1$ for all n large by (1.1). We claim that for such n , we have $\mathbb{P}[\mathcal{Y}_n] \geq \mathbb{P}_{\alpha_c+1}[\text{Cross}_1(2\delta/r_n)]$. Indeed, by the mapping theorem [5, Theorem 5.1], $r_n^{-1}(\mathcal{P}_{e^{-2n}} \cap C_{2\delta})$ is a Poisson process in $C_{2\delta/r_n}$ with intensity measure having a density bounded below by

$e^{-2}nf_0r_n^d$, and hence by $\alpha_c + 1$. By the thinning theorem [5, Corollary 5.9], one can couple \mathcal{H}_{α_c+1} and $\mathcal{P}_{e^{-2}n}$ in such a way that $\mathcal{H}_{\alpha_c+1} \cap C_{2\delta/r_n} \subset r_n^{-1}(\mathcal{P}_{e^{-2}n} \cap C_{2\delta})$. Since the crossing event is increasing in the sense that adding more points to the Poisson process increases the chance of its occurrence, this coupling justifies the claim. Thus by Lemma 3.9,

$$\mathbb{P}[\#(\mathcal{L}_n(\mathcal{P}_n)) \geq \delta/r_n] \geq \mathbb{P}[\mathcal{Y}_n] \geq \mathbb{P}_{\alpha_c+1}[\text{Cross}_1(2\delta/r_n)] \geq 1 - e^{-2c\delta/r_n},$$

For the case of binomial input, note that if $Z_{e^{-2}n} \leq n$ then $\mathcal{P}_{e^{-2}n} \leq \mathcal{X}_n$ and hence if also \mathcal{Y}_n occurs then $\#(\mathcal{L}_n(\mathcal{X}_n)) \geq \delta/r_n$ for n large. Therefore using Lemma 3.4(iii) we have

$$\mathbb{P}[\#(\mathcal{L}_n(\mathcal{X}_n)) < \delta/r_n] \leq \mathbb{P}[\mathcal{Y}_n^c] + \mathbb{P}[Z_{e^{-2}n} > n] \leq e^{-2c\delta/r_n} + e^{-n}.$$

Thus using the assumption $nr_n^d = O(\log n)$, we have (3.19) for both choices of ξ_n .

Now for $n \geq 1$ let $\rho := \rho(n) := \max((\log n)^2, 1)$, and partition \mathbb{R}^d into cubes of side length r_n . Necessarily $\mathcal{L}_n(\mathcal{P}_n)$ intersects one of the cubes with non-empty intersection with A , called Q , and if $\text{diam}(\mathcal{L}_n(\mathcal{P}_n)) < \rho r_n$, then $\mathcal{L}_n(\mathcal{P}_n) \subset Q \oplus B_{\rho r_n}$. If also $\#(\mathcal{L}_n(\mathcal{P}_n)) \geq \delta/r_n$, then $\mathcal{P}_n(Q \oplus B_{\rho r_n}) \geq \delta/r_n$. Since $\rho \geq 1$, we have $\lambda(Q \oplus B_{\rho r_n}) \leq (3\rho r_n)^d$. By the union bound, we have

$$\mathbb{P}[\{\text{diam}(\mathcal{L}_n(\mathcal{P}_n)) < \rho r_n\} \cap \{\#(\mathcal{L}_n(\mathcal{P}_n)) \geq \delta/r_n\}] \leq cr_n^{-d} \mathbb{P}[Z_{3^d \rho^d n f_{\max} r_n^d} \geq \delta/r_n].$$

We can then apply Lemma 3.4(iii) provided $\delta/r_n \geq e^2(3^d \rho^d n f_{\max} r_n^d)$, or in other words $\rho^d \leq (c'nr_n^{d+1})^{-1}$ for some constant c' .

By assumption $nr_n^d = O(\log n)$ so $\rho^d nr_n^{d+1} = O((\log n)^{2d+(d+1)/d} n^{-1/d})$. Hence we can apply Lemma 3.4(iii) to deduce that for n large

$$\mathbb{P}[\{\text{diam}(\mathcal{L}_n(\mathcal{P}_n)) < \rho r_n\} \cap \{\#(\mathcal{L}_n(\mathcal{P}_n)) \geq \delta/r_n\}] \leq cr_n^{-d} \exp(-\delta/r_n) \leq \exp(-\delta/(2r_n)).$$

Since $r_n^d = O((\log n)/n)$, we have $r_n^{-1} = \Omega\left(\left(\frac{n}{\log n}\right)^{1/d}\right)$ so using (3.19) and the union bound we can deduce (3.18) for $\xi_n = \mathcal{P}_n$. We can obtain (3.18) for $\xi_n = \mathcal{X}_n$ by a similar argument, using Lemma 3.4(i) instead of Lemma 3.4(iii). \square

4 The number of isolated vertices

In this section we prove Propositions 2.2 and 2.10. In the uniform case we also demonstrate the asymptotic equivalence of I_n and μ_n , defined at (2.1) and (1.4) respectively.

We continue to make the assumptions on ν and A that we set out at the start of Section 3. Also we assume $r_n \in (0, \infty)$ is given for all $n \geq 1$. Recalling from (2.1) that $I_n := n \int \exp(-n\nu(B_{r_n}(x)))\nu(dx)$, we assume throughout this section that r_n satisfies

$$\lim_{n \rightarrow \infty} nr_n^d = \infty; \tag{4.1}$$

$$\liminf_{n \rightarrow \infty} I_n > 0. \tag{4.2}$$

Also for $s > 0$ we write $A^{(-s)} := \{x \in A : B_s(x) \subset A\}$.

4.1 Mean and variance of the number of isolated vertices

Let S_n (respectively S'_n) denote the number of singletons (i.e. isolated vertices) of $G(\mathcal{X}_n, r_n)$ (resp., of $G(\mathcal{P}_n, r_n)$). That is, set

$$S'_n = \sum_{x \in \mathcal{P}_n} \mathbf{1}\{\mathcal{P}_n \cap B_{r_n}(x) = \{x\}\}; \quad S_n = \sum_{x \in \mathcal{X}_n} \mathbf{1}\{\mathcal{X}_n \cap B_{r_n}(x) = \{x\}\}. \quad (4.3)$$

By the Mecke formula $\mathbb{E}[S'_n] = I_n$. Also define

$$\tilde{I}_n := \mathbb{E}[S_n] = n \int (1 - \nu(B_{r_n}(x)))^{n-1} \nu(dx). \quad (4.4)$$

Lemma 4.1 (Lower bounds on I_n). *Let f_0^+, f_1^+ be constants with $f_0^+ > f_0$ and $f_1^+ > f_1$. Then as $n \rightarrow \infty$,*

$$n \exp(-n\theta f_0^+ r_n^d) = o(I_n); \quad (4.5)$$

$$n^{1-1/d} \exp(-n\theta f_1^+ r_n^d/2) = o(I_n). \quad (4.6)$$

Proof. Assume for now that $\partial A \in C^2$. See [12, Lemma 3.1] for a proof of (4.5). For (4.6), choose $x_0 \in \partial A$ with $f(x_0) < f_1^+$. Using the assumed continuity of f , choose $s_0 > 0$ and $\delta > 0$ such that

$$(1 + \delta)^{d+1} \sup\{f(y) : y \in A \cap B_{2s_0}(x_0)\} \leq f_1^+.$$

By Lemma 3.3, there is a constant $r_1 > 0$ (independent of z) such that $\lambda(B_s(z) \cap A) \leq (1 + \delta)\theta s^d/2$ for all $z \in \partial A, s \in (0, r_1)$.

Let $y \in B_{s_0}(x_0) \cap A \setminus A^{(-\delta r_n)}$ and let z be the closest point of ∂A to y . Then $\|y - z\| \leq \delta r_n$, so provided n is large enough

$$\nu(B_{r_n}(y)) \leq \nu(B_{(1+\delta)r_n}(z)) \leq (1 + \delta)^{d+1} f_1 \theta r_n^d/2 \leq f_1^+ \theta r_n^d/2.$$

Therefore since $\lambda(B_{s_0}(x_0) \cap A \setminus A^{(-s)}) = \Omega(s)$ as $s \downarrow 0$,

$$I_n \geq n \int_{B_{s_0}(x_0) \cap A \setminus A^{(-\delta r_n)}} \exp(-n\nu(B_{r_n}(y))) \nu(dy) = \Omega(nr_n e^{-nf_1^+ \theta r_n^d/2}),$$

and then using (4.1) we obtain (4.6).

In the case where $A = [0, 1]^2$ instead of $\partial A \in C^2$, the preceding proof still works, since we can take x_0 to not be a corner of A . \square

Lemma 4.2 (Upper bound on I_n). *Suppose $b^+ \leq (6/5)b_c$ with b_c given at (2.5). Then given $\varepsilon > 0$,*

$$I_n = O(ne^{-n\theta f_0 r_n^d} + n^{1-1/d} e^{-nr_n^d \theta f_1 (\frac{1}{2} - \varepsilon)}). \quad (4.7)$$

Proof. Note first that for all $x \in A^{(-r_n)}$ we have $\nu(B_{r_n}(x)) \geq \theta r_n^d f_0$, so that writing $I_n(S)$ for $n \int_S \exp(-n\nu(B_{r_n}(x))) \nu(dx)$, we have

$$I_n(A^{(-r_n)}) = n \int_{A^{(-r_n)}} e^{-n\nu(B_{r_n}(x))} \nu(dx) \leq ne^{-n\theta r_n^d f_0}.$$

Let $\varepsilon \in (0, \frac{1}{2})$. Suppose $\partial A \in C^2$. Then by Lemma 3.3 and the continuity of f , for all large enough n and all $x \in A \setminus A^{(-r_n)}$ we have $\nu(B_{r_n}(x)) \geq (1 - \varepsilon)f_1\theta r_n^d/2$; hence

$$I_n(A \setminus A^{(-r_n)}) = O(nr_n e^{-n\theta r_n^d f_1(1-\varepsilon)/2}) = O(n^{1-1/d} e^{-n\theta r_n^d f_1(\frac{1}{2}-\varepsilon)}).$$

Now suppose instead that $d = 2$ and $A = [0, 1]^2$. Let Cor_n denote the set of $x \in A$ lying at an ℓ_∞ distance at most r_n from one of the corners of A . By the same argument as above

$$I_n(A \setminus A^{(-r_n)} \setminus \text{Cor}_n) = O(n^{1/2} e^{-n\theta r_n^2 f_1(\frac{1}{2}-\varepsilon)}).$$

Also $I_n(\text{Cor}_n) \leq 4f_{\max}nr_n^2 \exp(-n\pi r_n^2 f_0/4)$ and using the assumption $b^+ < (6/5)b_c = 6/(5f_0)$, we obtain for large n that $nf_0\pi r_n^2 \leq (5/4)\log n$ so that

$$\frac{I_n(\text{Cor}_n)}{ne^{-n\pi r_n^2 f_0}} \leq 4f_{\max}r_n^2 \exp(3n\pi r_n^2 f_0/4) = O(r_n^2 n^{15/16}) = o(1).$$

Combining all of the preceding estimates we obtain for both cases ($\partial A \in C^2$ or $A = [0, 1]^2$) that (4.7) holds. \square

Proof of Proposition 2.2. If $b^+ < b_c = \max(\frac{1}{f_0}, \frac{2-2/d}{f_1})$ then we claim that $I_n \rightarrow \infty$. Indeed, if $b^+ < 1/f_0$, choose $f_0^+ > f_0$, $\delta > 0$, such that $f_0^+(b^+ + \delta) < 1$. Then for n large $ne^{-n\theta f_0^+ r_n^d} > ne^{-f_0^+(b^+ + \delta)\log n}$ so $I_n \rightarrow \infty$ by Lemma 4.1. If $b^+ < (2 - 2/d)/f_1$ then choose $f_1^+ > f_1$ and $\delta > 0$ with $f_1^+(b^+ + \delta) < 2 - 2/d$. Then for n large, $n^{1-1/d}e^{-n\theta f_1^+ r_n^d/2} > n^{1-1/d}e^{-\frac{1}{2}f_1^+(b^+ + \delta)\log n}$ so again $I_n \rightarrow \infty$ by Lemma 4.1, and the claim follows.

Now suppose $b^- > b_c$. We need to show $I_n \rightarrow 0$ as $n \rightarrow \infty$. Since $n \int_A e^{-n\nu(B_s(x))} \nu(dx)$ is nonincreasing in s , it suffices to prove this under the extra assumption $b^+ \leq 6b_c/5$, which makes Lemma 4.2 applicable. Since $b^- > b_c$ there exists $\varepsilon > 0$ such that for n large enough $\theta f_0 nr_n^d > (1 + \varepsilon)\log n$ and $nr_n^d \theta f_1(\frac{1}{2} - \varepsilon) > (1 - 1/d)(1 + \varepsilon)\log n$, and then we see $I_n \rightarrow 0$ by (4.7).

Finally if $b^+ > b_c$ then by the preceding argument $I_n \rightarrow 0$ as $n \rightarrow \infty$ along some subsequence, so we must have $\liminf_{n \rightarrow \infty} I_n = 0$. \square

Lemma 4.3 (Asymptotic equivalence of I_n and \tilde{I}_n). *There exists $\delta > 0$ such that as $n \rightarrow \infty$ we have $|I_n - \tilde{I}_n| = O(e^{-\delta nr_n^d} I_n)$.*

Proof. For $x \in A$, given n write $p_n(x) := \nu(B_{r_n}(x))$. By the bounds $1 - p_n(x) \leq e^{-p_n(x)}$, and $1 - p_n(x) \geq 1 - f_{\max}\theta r_n^d$, and the condition (4.1),

$$\begin{aligned} \tilde{I}_n &\leq (1 - f_{\max}\theta r_n^d)^{-1} n \int_A e^{-np_n(x)} \nu(dx) \\ &= (1 + O(r_n^d)) I_n. \end{aligned}$$

Also by Taylor's theorem $\log(1 - p) \geq -p - p^2$ for $p > 0$ close to 0, so

$$\begin{aligned} \tilde{I}_n &\geq n \int_A \exp(n \log(1 - p_n(x))) dx \\ &\geq n \int_A \exp(n(-p_n(x) - p_n(x)^2)) dx \\ &\geq e^{-n(f_{\max}\theta r_n^d)^2} I_n. \end{aligned}$$

Combining these two estimates and using (4.1) yields $|\tilde{I}_n - I_n| = O(nr_n^{2d}I_n)$. Therefore it suffices to show $nr_n^{2d}e^{\delta nr_n^d} = O(1)$ for some $\delta > 0$. By (4.2) and Proposition 2.2, $nr_n^d = O(\log n)$ so for δ small enough $e^{\delta nr_n^d} = O(n^{1/2})$, while $nr_n^{2d} = O((\log n)^2 n^{-1})$ so $nr_n^{2d}e^{\delta nr_n^d} = O((\log n)^2 n^{-1/2}) = o(1)$. \square

Proposition 4.4 (Variance asymptotics of the number of singletons: Poisson input). *There exists $c > 0$ such that as $n \rightarrow \infty$,*

$$\text{Var}[S'_n] = I_n(1 + O(e^{-c nr_n^d})).$$

Proof. This is proved in [12, Proposition 4.1, case $k = 1$], for $\partial A \in C^2$, under the extra assumption [12, equation (3.1)] which amounts to $b^+ < 1/\max(f_0, d(f_0 - f_1/2))$ (or $b < 2/(df_0)$ in the uniform case). We need to relax this condition to $\liminf_{n \rightarrow \infty} I_n > 0$ as well as $nr_n^d \rightarrow \infty$. Also we need to consider the case with $d = 2, A = [0, 1]^2$. We find that the proof of [12, Proposition 4.1] still works, except for where it uses [12, Lemma 4.2]. The latter proof does invoke the extra assumption on b^+ , so we provide a different proof of [12, Lemma 4.2, case $k = 1$] not requiring the extra assumption on b^+ , that works for $A = [0, 1]^2$ as well as for $\partial A \in C^2$.

The quantity denoted $J_{2,n}$ in [12] is $O(I_n e^{-c nr_n^d})$ as in [12]. The quantity denoted $J_{1,n}$ in [12] (when $k = 1$) is given by

$$\begin{aligned} J_{1,n} &= n^2 \int_A \int_A e^{-n\nu(B_{r_n}(x) \cup B_{r_n}(y))} \mathbf{1}\{r_n < \|y - x\| \leq 2r_n\} \nu(dy) \nu(dx) \\ &\leq 2n^2 \int_A \int_A e^{-n\nu(B_{r_n}(x) \cup B_{r_n}(y))} \mathbf{1}\{r_n < \|y - x\| \leq 2r_n, x \prec y\} \nu(dy) \nu(dx). \end{aligned}$$

By (3.1) from Lemma 3.1, there exists $\delta_1 > 0$ such that for n large and $x, y \in A$ with $\|x - y\| \geq r_n$ and $x \prec y$ we have $\nu(B_{r_n}(y) \setminus B_{r_n}(x)) \geq 2\delta_1 f_0 r_n^d$. Hence

$$\begin{aligned} J_{1,n} &\leq 2n^2 f_{\max} \theta(2r_n)^d \int_A e^{-n\nu(B_{r_n}(x)) - 2n\delta_1 f_0 r_n^d} \nu(dx) \\ &\leq e^{-n\delta_1 f_0 r_n^d} I_n, \end{aligned}$$

which gives us [12, Lemma 4.2, case $k = 1$] and hence the result. \square

Proposition 4.5 (Variance asymptotics of the number of singletons: binomial input). *There exists $\delta > 0$ such that as $n \rightarrow \infty$,*

$$\text{Var}[S_n] = I_n(1 + O(e^{-\delta nr_n^d})).$$

Proof. See [12, Proposition 4.3] and Lemma 4.3 of the present paper. In the proof of [12, Proposition 4.3] it is assumed that [12, equation (3.1)] holds (i.e. $b^+ < 1/\max(f_0, d(f_0 - f_1/2))$ in our notation here), but the proof carries through to the general case with $nr_n^d \rightarrow \infty$ and $I_n \rightarrow \infty$. Instead of [12, Lemma 4.2] we can use the estimate on $J_{1,n}$ in the proof of Proposition 4.4 above. \square

4.2 Asymptotic distribution of the singleton count

For both the normal and Poisson convergence results, we use the following.

Lemma 4.6 (Poisson approximation for S'_n and S_n). *There exists $\delta > 0$ such that as $n \rightarrow \infty$,*

$$d_{\text{TV}}(S'_n, Z_{I_n}) = O(e^{-\delta nr_n^d}); \quad (4.8)$$

$$\text{if } \partial A \in C^2, \quad d_{\text{TV}}(S_n, Z_{\tilde{I}_n}) = O(e^{-\delta nr_n^d}). \quad (4.9)$$

Proof. The result (4.8) is proved in [12, Proposition 5.4] (taking $k = 1$ there), under the extra assumptions that $\partial A \in C^2$ and that [12, equation (3.1)] holds, i.e. $b^+ \leq 1/\max(f_0, d(f_0 - f_1/2))$ in our notation. We wish to drop this assumption on b^+ , and assume only that $nr_n^d \rightarrow \infty$ and $\liminf_{n \rightarrow \infty}(I_n) > 0$. Also we wish to include the case where $A = [0, 1]^2$.

Going through the proof of [12, Proposition 5.4] we find that it still works under our weaker assumptions, but that it uses [12, Lemma 4.2]. In the proof of Proposition 4.4 we have already provided a the proof of [12, Lemma 4.2] under our weaker assumptions, justifying (4.8).

Similarly (4.9) is proved in [12, Proposition 5.6], taking $k = 1$ there, under the extra assumption that $b^+ \leq 1/\max(f_0, d(f_0 - f_1/2))$, which we wish to drop. Going through the proof of [12, Proposition 5.6], we find that it still works without the extra assumption on b^+ . Since $k = 1$ for us, we can give a simpler proof of [12, Lemma 5.8] than the one given there for general k , as follows.

In our case N_6 is the number of singletons of $G(\mathcal{X}_{n,2}, r_n)$ within distance $2r_n$ of \tilde{X}_1 . Each vertex has probability $O(r_n^d)$ of lying in $B_{2r_n}(\tilde{X}_1) \setminus B_{r_n}(\tilde{X}_1)$, and given its location and that of \tilde{X}_1 , using (3.1) from Lemma 3.1 without assuming $x \prec y$ there (and therefore requiring $\partial A \in C^2$), it has probability at most $e^{-\delta nr_n^d}$ of being isolated, for some $\delta > 0$. Thus we can obtain that $\mathbb{E}[N_6] = O(nr_n^d \exp(-\delta nr_n^d))$ which is what we need. \square

Proof of Proposition 2.10. The assertion $\mathbb{E}[\zeta_n] = I_n(1 + O(e^{-\delta nr_n^d}))$ follows from the Mecke formula (in the case $\zeta_n = S'_n$) and from Lemma 4.3 (in the case $\zeta_n = S_n$). The assertion $\text{Var}[\zeta_n] = I_n(1 + O(e^{-\delta nr_n^d}))$ follows from Proposition 4.4 (if $\zeta_n = S'_n$) and from Proposition 4.5 (if $\zeta_n = S_n$).

By the Berry-Esseen theorem $d_K(t^{-1/2}(Z_t - t), N(0, 1)) = O(t^{-1/2})$ as $t \rightarrow \infty$. Hence, by the triangle inequality for d_K , we have

$$d_K(I_n^{-1/2}(S'_n - I_n), N(0, 1)) \leq d_K(S'_n, Z_{I_n}) + O(I_n^{-1/2}). \quad (4.10)$$

Then using (4.8), and the obvious inequality $d_K \leq d_{\text{TV}}$, we obtain (2.22). Similarly using (4.9) and Lemma 4.3 we obtain (2.23). \square

4.3 Asymptotics for I_n in the uniform case

Throughout this subsection we make the additional assumption that $f \equiv f_0 \mathbf{1}_A$, with $f_0 = 1/\lambda(A)$, and assume as $n \rightarrow \infty$ that r_n satisfy

$$nr_n^d \rightarrow \infty; \quad \limsup(nf_0\theta r_n^d - (2 - 2/d)(\log n - \mathbf{1}\{d \geq 3\} \log \log n)) < \infty. \quad (4.11)$$

We shall demonstrate the asymptotic equivalence of I_n and μ_n , defined at (2.1) and (1.4) respectively. Given n , for $x \in A$ set $p_n(x) := \exp(-n\nu(B_{r_n}(x)))$. Given Borel $S \subset A$, let

$$I_n(S) := nf_0 \int_S p_n(x) dx. \quad (4.12)$$

Proposition 4.7 (The case $d = 2$). *Suppose $d = 2$ and either $\partial A \in C^2$, or $A = [0, 1]^2$. Suppose (4.11) holds. Then we have as $n \rightarrow \infty$ that*

$$I_n = n \exp(-nf_0\pi r_n^2)(1 + O(nr_n^2)^{-1/2}). \quad (4.13)$$

Proof. Case 1: $\partial A \in C^2$. In this case the result follows from Proposition 4.8 below, since the ratio between the two terms in the right hand side of (4.14) is given by

$$\frac{e^{-nf_0\theta r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\partial A|}{ne^{-nf_0\theta r_n^d}} = O\left(\exp\left(\frac{nf_0\pi r_n^2}{2} - \frac{\log n}{2} - \frac{\log(nr_n^2)}{2}\right)\right),$$

which is $O((nr_n^2)^{-1/2})$ by (4.11).

Case 2: $A = [0, 1]^2$. In this case $f_0 = 1$. Define the ‘moat’ $\text{Mo}_n := A \setminus A^{(-r_n)}$.

For $1 \leq i \leq 4$ let $\text{Cor}_{n,i}$ be the region of A within ℓ_∞ -distance r_n of the i th corner of A (a square of side r_n). Then $I_n(\text{Cor}_{n,i}) \leq nr_n^2 e^{-n\pi r_n^2/4}$.

The set $\text{Mo}_n \setminus \cup_{i=1}^4 \text{Cor}_{n,i}$ is a union of rectangular regions $\text{Rec}_{n,i}$, $1 \leq i \leq 4$. For all n large enough, $i \leq 4$ and $x \in \text{Rec}_{n,i}$ we have $\nu(B_{r_n}(x)) \geq r_n^2(\frac{\pi}{2} + a(x)/r_n)$, where $a(x)$ denotes the distance from x to ∂A . Hence

$$I_n(\text{Rec}_{n,i}) \leq nr_n e^{-n\pi r_n^2/2} \int_0^1 e^{-nr_n^2 a} da = O(nr_n e^{-\pi nr_n^2/2} (nr_n^2)^{-1}).$$

Combined with the corner region estimate, and the bound $\pi nr_n^2 \leq \log n + c$ from (4.11), this yields

$$\begin{aligned} I_n(\text{Mo}_n)/I_n(A^{(-r_n)}) &= O(n^{-1}r_n^{-1}e^{n\pi r_n^2/2}) + O(r_n^2 e^{(3/4)n\pi r_n^2}) \\ &= O(n^{-1/2}r_n^{-1}) + O(((\log n)/n)n^{3/4}). \end{aligned}$$

Also $I_n(A^{(-r_n)}) = ne^{-n\pi r_n^2}(1 + O(r_n))$ and $r_n = o((nr_n^2)^{-1/2})$. Putting together these estimates yields (4.13) in Case 2. \square

Proposition 4.8 (The case $\partial A \in C^2$). *Suppose $d \geq 2$ and $\partial A \in C^2$. Suppose $(r_n)_{n \geq 1}$ satisfy (4.11). Then as $n \rightarrow \infty$,*

$$I_n = ne^{-nf_0\theta r_n^d} + e^{-nf_0\theta r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\partial A| \left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right). \quad (4.14)$$

Proof. We shall investigate separately the contributions from different regions of the set A . Since the proof is quite long, we break it into steps.

Step 1: Polytopal approximation to ∂A . We shall approximate A locally by a polytope, i.e. break ∂A into finitely many pieces and approximate each piece by a polytopal surface.

Given $x \in \partial A$ we can express ∂A locally in a neighbourhood of x , after a rotation, as the graph of a C^2 function with zero derivative at x . We shall approximate to that function by the graph of a piecewise affine function.

Here are the details of this approximation, where we are following [10] and [4]. For each $x \in \partial A$, we can find an open neighbourhood \mathcal{N}_x of x , a number $r(x) > 0$ such that $B(x, 3r(x)) \subset \mathcal{N}_x$ and a rotation φ_x about x such that $\varphi_x(\partial A \cap \mathcal{N}_x)$ is the graph of a real-valued C^2 function f defined on an open ball $D \subset \mathbb{R}^{d-1}$, with $\langle f'(u), e \rangle \leq 1/(9d)$ for all $u \in D$ and all unit vectors e in \mathbb{R}^{d-1} , where $\langle \cdot, \cdot \rangle$ denotes the Euclidean inner product in \mathbb{R}^{d-1} and $f'(u) := (\partial_1 f(u), \partial_2 f(u), \dots, \partial_{d-1} f(u))$ is the derivative of f at u . Moreover, by taking a smaller neighbourhood if necessary, we can also assume that there exists $\varepsilon > 0$ and $a \in \mathbb{R}$ such that $f(u) \in [a + \varepsilon, a + 2\varepsilon]$ for all $u \in D$ and also $\varphi_x(A) \cap (D \times [a, a + 3\varepsilon]) = \{(u, z) : u \in D, a \leq z \leq f(u)\}$.

By a compactness argument, we can and do take a finite collection of points $x_1, \dots, x_J \in \partial A$ such that

$$\partial A \subset \cup_{j=1}^J B(x_j, r(x_j)). \quad (4.15)$$

Then there are constants $\varepsilon_j > 0$, and rigid motions φ_j , $1 \leq j \leq J$, such that for each j the set $\varphi_j(\partial A \cap \mathcal{N}_{x_j})$ is the graph of a C^2 function f_j defined on a ball I_j in \mathbb{R}^{d-1} , with $|\langle f'_j(u), e \rangle| \leq 1/9$ for all $u \in I_j$ and all unit vectors $e \in \mathbb{R}^{d-1}$, and also with $\varepsilon_j \leq f_j(u) \leq 2\varepsilon_j$ for all $u \in I_j$ and $\varphi_j(A) \cap (I_j \times [0, 3\varepsilon_j]) = \{(u, z) : u \in I_j, 0 \leq z \leq f(u)\}$.

Let $\Gamma \subset \partial A$ be a closed set such that $\Gamma \subset B(x_j, r(x_j))$ for some $j \in \{1, \dots, J\}$. Assume Γ is ‘nice’ in the sense that as $r \downarrow 0$, the number of balls of radius r required to cover $\partial\Gamma$, the relative boundary of Γ in ∂A , is $O(r^{2-d})$ (if $d = 2$ we can assume $\partial\Gamma$ is finite). To simplify notation we shall assume that $\Gamma \subset B(x_1, r(x_1))$, and moreover that φ_1 is the identity map. Then $\Gamma = \{(u, f_1(u)) : u \in U\}$ for some bounded set $U \subset \mathbb{R}^{d-1}$. Also, writing $\phi(\cdot)$ for $f_1(\cdot)$ from now on, we assume

$$\phi(U) \subset [\varepsilon_1, 2\varepsilon_1] \quad (4.16)$$

and

$$A \cap (U \times [0, 3\varepsilon_1]) = \{(u, z) : u \in U, 0 \leq z \leq \phi(u)\}. \quad (4.17)$$

Divide \mathbb{R}^{d-1} into cubes of dimension $d - 1$ and side $r_n^{3/4}$, and divide each of these cubes into $(d - 1)!$ simplices (we take these simplices to be closed). Let U_n be the union of all those simplices in the resulting tessellation of \mathbb{R}^{d-1} into simplices, that are contained within U .

Let $\psi_n : U_n \rightarrow \mathbb{R}$ be the function that is affine on each of the simplices making up U_n , and agrees with the function ϕ on each of the vertices of these simplices. Choose n_0 so that $U_n \neq \emptyset$ for all $n \geq n_0$. Set

$$K := \sup_{n \geq n_0, u \in U_n} (r_n^{-3/2} |\phi(u) - \psi_n(u)|). \quad (4.18)$$

Then $K < \infty$ by an based on Taylor expansion, similar to [10, Lemma 7.18] and to [4, eq. (5.7)].

Now define $\phi_n(u) := \psi_n(u) - Kr_n^{3/2}$. By (4.18), $\phi_n(u) \leq \phi(u) \leq \phi_n(u) + 2Kr_n^{3/2}$ for all $u \in U_n$. Our approximating surface will be defined by $\Gamma_n := \{(x, \phi_n(x)) : x \in U_n\}$, with the constant K given by (4.18).

Step 2: Decomposition of the range of integration. We define the moat \mathbf{Mo}_n by

$$\mathbf{Mo}_n := \{(u, z) : u \in U_n, z \leq \phi_n(u), \text{dist}((u, z), \Gamma_n) \leq r_n\}.$$

This is the region within distance r_n of Γ_n and below Γ_n . We refer to $\mathbf{Rai}_n := \{(u, z) : u \in U_n, \phi_n(u) \leq z \leq \phi(u)\}$ as the *railing*. We refer to $A^{(-2r_n)}$ as the *bulk*.

We further decompose \mathbf{Mo}_n as follows. By construction, the set Γ_n is composed of $(d-1)$ -dimensional facets, which we denote by $\mathbf{Fac}_{n,i}$, $1 \leq i \leq m(n)$. For each $i \leq m(n)$ let $\mathbf{Pri}_{n,i}$ denote the prism with height r_n and upper $(d-1)$ -dimensional base given by the part of $\mathbf{Fac}_{n,i}$ at distance at least $4r_n$ from the set $\partial\mathbf{Fac}_{n,i}$, which we define to be the union of all the $(d-2)$ -dimensional facets bounding the facet $\mathbf{Fac}_{n,i}$.

We claim that the prisms $\mathbf{Pri}_{n,i}$, $1 \leq i \leq m(n)$, are pairwise disjoint. To see this, note that if $u, v \in U$, then as at equation (7.51) of [10], since $|\langle \phi'(w), e \rangle| \leq 1/9$ for all unit vectors $w \in \mathbb{R}^{d-1}$,

$$\|(u, \phi(u)) - (v, \phi(v))\| \leq \|u - v\| + |\phi(u) - \phi(v)| \leq (9/8)\|u - v\|. \quad (4.19)$$

Hence $\|u - v\| \geq (8/9)\|(u, \phi(u)) - (v, \phi(v))\|$.

Let $\pi : \mathbb{R}^d \rightarrow \mathbb{R}^{d-1}$ denote projection onto the first $d-1$ coordinates. If $i \leq m(n)$ and $x \in \mathbf{Pri}_{n,i}$, and y is the nearest point in $\mathbf{Fac}_{n,i}$ to x , then $\text{dist}(y, \partial\mathbf{Fac}_{n,i}) \geq 4r_n$, and $\text{dist}(\pi(y), \pi(\partial\mathbf{Fac}_{n,i})) \geq (8/9)(4r_n) \geq 3r_n$. Since $\|x - y\| \leq r_n$, also $\|\pi(y) - \pi(x)\| \leq r_n$, and therefore $\text{dist}(\pi(x), \pi(\partial\mathbf{Fac}_{n,i})) \geq 2r_n$. In particular, $\pi(x)$ lies in $\mathbf{Fac}_{n,i}^o$, the relative interior of $\mathbf{Fac}_{n,i}$. Similarly, for any $j \neq i$ with $j \leq m(n)$, we have for any $z \in \mathbf{Pri}_{n,j}$ that $\pi(z) \in \pi(\mathbf{Fac}_{n,j}^o)$, and since $\pi(\mathbf{Fac}_{n,j}), \pi(\mathbf{Fac}_{n,i})$ have disjoint interiors, this justifies the claim.

The preceding argument also shows that if $i, j \leq m(n)$ with $i \neq j$, then

$$\text{dist}(x, \mathbf{Fac}_{n,j}) \geq \text{dist}(\pi(x), \pi(\mathbf{Fac}_{n,j})) \geq 2r_n, \quad \forall x \in \mathbf{Pri}_{n,i}. \quad (4.20)$$

By construction, $\bigcup_{i=1}^{m(n)} \mathbf{Pri}_{n,i} \subset \mathbf{Mo}_n$. The left-over regions in the moat near the $(d-2)$ -dimensional facets of Γ_n (and therefore not in any prism) are called $(\mathbf{Cor}_{n,i})$, the *corner regions*. Notice that when $d \geq 3$ the $\mathbf{Cor}_{n,i}$'s may overlap each other a little near the $(d-3)$ -dimensional facets of Γ_n , and they have empty intersection with any of the prisms. Partition the moat into prisms and corners

$$\mathbf{Mo}_n = \left(\bigcup_{i=1}^{m(n)} \mathbf{Pri}_{n,i} \right) \cup \left(\bigcup_{i=1}^{m(n)} \mathbf{Cor}_{n,i} \right).$$

Step 3: contribution of the bulk, railing and corners. To deal with the bulk, note that for each $x \in A^{(-r_n)}$, we have $p_n(x) = e^{-nf_0\theta r_n^d}$ so that by (4.12),

$$I_n(A^{(-2r_n)}) = nf_0|A^{(-2r_n)}|e^{-nf_0\theta r_n^d} = (1 + O(r_n))ne^{-nf_0\theta r_n^d}. \quad (4.21)$$

We deal next with the railing. Given $\kappa < 1/2$, for n large enough, we have by Lemma 3.3 that $p_n(x) \leq e^{-nf_0\theta r_n^d \kappa}$ for $x \in \text{Rai}_n$. The total volume of the railing is $O(r_n^{3/2})$, so by (4.12),

$$I_n(\text{Rai}_n) = O(nr_n^{3/2} e^{-nf_0\theta r_n^d \kappa}). \quad (4.22)$$

To bound the contribution from the corners, we see from Lemma 3.3 that given $\kappa < 1/2$, for large n we have $p_n(x) \leq e^{-nf_0\theta r_n^d \kappa}$ for $x \in \cup_{i=1}^{m(n)} \text{Cor}_{n,i}$, and there are in total $O(r_n^{(1-d)3/4})$ corner regions, each of which has volume at most $cr_n^{2+(d-2)3/4}$ for some constant c , yielding

$$I_n(\cup_{i=1}^{m(n)} \text{Cor}_{n,i}) = O(nr_n^{5/4} e^{-nf_0\theta r_n^d \kappa}). \quad (4.23)$$

Step 4: contribution of the prisms. For $a > 0$ let $g(a) := \lambda(B_1(o) \cap ([0, a] \times \mathbb{R}^{d-1}))$, and for $x \in \text{Pri}_{n,i}$ let $a(x)$ be the distance from x to the face $\text{Fac}_{n,i}$. For each $i \leq m(n)$ and $x \in \text{Pri}_{n,i}$, using (4.20) we have

$$\begin{aligned} |p_n(x) - e^{-nf_0 r_n^d (\frac{\theta}{2} + g(\frac{a(x)}{r_n}))}| &\leq e^{-nf_0 r_n^d (\frac{\theta}{2} + g(\frac{a(x)}{r_n}))} \\ &\times \left(1 - \exp \left(-nf_0 r_n^d \left(g\left(\frac{a(x) + 2Kr_n^{3/2}}{r_n}\right) - g\left(\frac{a(x)}{r_n}\right) \right) \right) \right). \end{aligned}$$

Using the inequality $1 - e^{-s} \leq s$ for $s \in \mathbb{R}$, and (4.11), and the fact that $g'(a) \leq \theta_{d-1}$ for all a , we obtain that there exists a constant c such that

$$|p_n(x) - e^{-nf_0 r_n^d (\frac{\theta}{2} + g(\frac{a(x)}{r_n}))}| \leq cnr_n^{d+\frac{1}{2}} e^{-nf_0 r_n^d (\frac{\theta}{2} + g(\frac{a(x)}{r_n}))}. \quad (4.24)$$

Define the upper base of $\text{Pri}_{n,i}$ by $\text{BPri}_{n,i} = \text{Pri}_{n,i} \cap \Gamma_n$ (which is a part of $\text{Fac}_{n,i}$), and let $|\text{BPri}_{n,i}|$ be its $(d-1)$ -dimensional Hausdorff measure. Then by (4.24),

$$\begin{aligned} &\left| nf_0 \int_{\text{Pri}_{n,i}} p_n(x) dx - nf_0 |\text{BPri}_{n,i}| r_n \int_0^1 e^{-nf_0 r_n^d (\frac{\theta}{2} + g(a))} da \right| \\ &\leq cn^2 r_n^{d+\frac{1}{2}} |\text{BPri}_{n,i}| r_n \int_0^1 e^{-nf_0 r_n^d (\frac{\theta}{2} + g(a))} da. \end{aligned}$$

Hence, using (4.11) and the fact that $t \int_0^1 e^{-tg(a)} da \rightarrow \theta_{d-1}^{-1}$ as $t \rightarrow \infty$ to handle the last factor, and summing over i , we deduce that

$$\begin{aligned} &\left| nf_0 \int_{\cup_{i=1}^{m(n)} \text{Pri}_{n,i}} p_n(x) (dx) - \sum_{i=1}^{m(n)} |\text{BPri}_{n,i}| \int_0^1 nf_0 r_n e^{-nf_0 r_n^d (\frac{\theta}{2} + g(a))} da \right| \\ &= O(n^2 r_n^{d+3/2} e^{-nf_0\theta r_n^d/2} (nr_n^d)^{-1}) \\ &= O(nr_n^{3/2} e^{-nf_0\theta r_n^d/2}). \end{aligned} \quad (4.25)$$

Next we claim that as $n \rightarrow \infty$,

$$\int_0^1 nf_0 r_n e^{-nf_0 r_n^d (\frac{\theta}{2} + g(a))} da = e^{-nf_0\theta r_n^d/2} \theta_{d-1}^{-1} r_n^{1-d} \left(1 + O\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2 \right), \quad (4.26)$$

and

$$\sum_{i=1}^{m(n)} |\mathbf{BPri}_{n,i}| = |\Gamma| \left(1 + O(r_n^{1/4})\right), \quad (4.27)$$

where $|\Gamma|$ denotes the $(d-1)$ -dimensional Hausdorff measure of Γ .

To prove (4.27), recall that Γ is the graph of a C^2 function ϕ on some $U \subset \mathbb{R}^{d-1}$, and Γ_n is obtained by first triangulating U into $(d-1)$ -dimensional simplices, denoted $(s_{n,i})$, then using the simplex $\mathbf{Fac}_{n,i}$ obtained as the convex hull of the set of points $\hat{\phi}(v)$, where v runs through the vertices of the simplex $s_{n,i}$, and for $u \in U$ we set $\hat{\phi}(u) = (u, \phi(u) - Kr_n^{3/2})$.

For each $i \leq m(n)$, pick $a_{n,i} \in s_{n,i}$; then the affine function $h_{n,i} : s_{n,i} \rightarrow \mathbb{R}$ with graph $\mathbf{Fac}_{n,i}$ satisfies $\nabla h_{n,i}(u) = \nabla h_{n,i}(a_{n,i})$ for all $u \in s_{n,i}^o$. By the Mean Value theorem, for $u \in s_{n,i}$ and $1 \leq j \leq d$, we have $\partial_j \phi(u) - \partial_j \phi(a_{n,i}) = \langle (u - a_{n,i}), \nabla \partial_j \phi(v) \rangle$, for some $v \in [a_{n,i}, u]$, and hence there is a constant c such that $|(\|\nabla \phi(u)\|^2 - \|\nabla \phi(a_{n,i})\|^2)| \leq c\|u - a_{n,i}\|$. Therefore with $|\hat{\phi}(s_{n,i})|$ and $|\mathbf{Fac}_{n,i}|$ denoting the $(d-1)$ -dimensional Hausdorff measure of $\hat{\phi}(s_{n,i})$ and of $\mathbf{Fac}_{n,i}$, respectively, we have for some new constants c, c' that

$$\begin{aligned} |(|\hat{\phi}(s_{n,i})| - |\mathbf{Fac}_{n,i}|)| &= \left| \int_{s_{n,i}} \left(\sqrt{1 + \|\nabla \phi(u)\|^2} - \sqrt{1 + \|\nabla h_{n,i}(u)\|^2} \right) du \right| \\ &\leq c \int_{s_{n,i}} \left| \left(\|\nabla \phi(u)\|^2 - \|\nabla \phi(a_{n,i})\|^2 \right) \right| du \leq c' r_n^{3/4} |s_{n,i}|. \end{aligned}$$

Summing over the simplices leads to an approximation error of $|(|\hat{\phi}(U_n)| - |\Gamma_n|)| \leq cr_n^{3/4}$.

Next, $|\mathbf{BPri}_{n,i}|$ approximates $|\mathbf{Fac}_{n,i}|$ with an error of at most $cr_n^{1+3(d-2)/4}$ and there are $O(r_n^{3(1-d)/4})$ facets, so $|\Gamma_n| - \sum_i |\mathbf{BPri}_i| = O(r_n^{1/4})$.

Also, $\Gamma \setminus \hat{\phi}(U_n) = \hat{\phi}(U) \setminus \hat{\phi}(U_n)$ is a region within $cr_n^{3/4}$ of the boundary of $\partial\Gamma$ for a further constant c , so assuming Γ is ‘nice’ (in the sense described shortly before (4.16)), this has $(d-1)$ -dimensional Hausdorff measure that is $O(r_n^{3/4})$. Combining the last three estimates gives us (4.27).

To prove (4.26), we first notice that $g(a) = \theta_{d-1} \int_0^a (1-t^2)^{(d-1)/2} dt$, $0 \leq a \leq 1$. Therefore, we have (i) $g(0) = 0$, g is increasing and $g(a) \leq \theta_{d-1} a$, (ii) for any $\varepsilon \in (0, 1/2)$ there exists $\delta \in (0, 1)$ such that for all $a \in (0, \delta)$, $g(a) \geq (1-\varepsilon)\theta_{d-1} a$, and (iii) upon choosing a smaller δ in (ii), we also have $g(a) \geq \theta_{d-1}(a - da^3)$ for $a \in (0, \delta)$. By item (i), we have

$$\begin{aligned} n f_0 \theta_{d-1} r_n^d \int_0^1 e^{-n f_0 r_n^d g(a)} da &\geq n f_0 \theta_{d-1} r_n^d \int_0^1 e^{-n r_n^d f_0 \theta_{d-1} a} da \\ &= 1 - e^{-n f_0 \theta_{d-1} r_n^d}. \end{aligned}$$

Let $\delta_n \in (0, \delta)$. By item (iii), we have

$$n f_0 \theta_{d-1} r_n^d \int_0^{\delta_n} e^{-n f_0 r_n^d g(a)} da \leq n f_0 \theta_{d-1} r_n^d \int_0^{\delta_n} e^{-n f_0 \theta_{d-1} r_n^d a (1-d\delta_n^2)} da \leq 1 + c\delta_n^2.$$

By item (ii), we have

$$\begin{aligned} n f_0 \theta_{d-1} r_n^d \int_{\delta_n}^{\delta} e^{-n f_0 r_n^d g(a)} da &\leq n f_0 \theta_{d-1} r_n^d \int_{\delta_n}^{\delta} e^{-n f_0 r_n^d (1-\varepsilon)\theta_{d-1} a} da \\ &\leq 2 \exp(-n f_0 r_n^d \theta_{d-1} (1-\varepsilon)\delta_n). \end{aligned}$$

Moreover, using (4.11) it is easy to see that for n large

$$nf_0\theta_{d-1}r_n^d \int_{\delta}^1 e^{-nf_0r_n^d g(a)} da \leq \exp(-nr_n^d f_0 g(\delta)/2)$$

Combining these four estimates, using (4.11) and choosing $\delta_n = c(\log(nr_n^d))/(nr_n^d)$ with a big constant c , leads to

$$nf_0\theta_{d-1}r_n^d \int_0^1 e^{-nf_0r_n^d g(a)} da = 1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right). \quad (4.28)$$

Since the left side of (4.28), multiplied by $\theta_{d-1}^{-1}r_n^{1-d}e^{-nf_0\theta_{d-1}r_n^d/2}$, comes to the left side of (4.26), (4.28) yields (4.26).

Combining (4.25)–(4.27), using (4.12) and setting $M_n := \cup_{i=1}^{m(n)} \text{Pri}_{n,i}$, leads to

$$\begin{aligned} I_n(M_n) &= e^{-nf_0\theta_{d-1}r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\Gamma|(1 + O(r_n^{1/4}))\left(1 + O\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right) + O\left(nr_n^{3/2}e^{-nf_0\theta_{d-1}r_n^d/2}\right) \\ &= e^{-nf_0\theta_{d-1}r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\Gamma|\left(1 + O\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right), \end{aligned}$$

where for the last line we used that by (4.11), $\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2 = \Omega((\log n)^{-2})$ and $nr_n^{d+1/2} = O(r_n^{1/4}) = o((\log n)^{-2}) = o\left(\left(\frac{\log nr_n^d}{nr_n^d}\right)^2\right)$. Also by (4.23), taking κ sufficiently close to $1/2$ we have that

$$I_n(\cup_{i=1}^{m(n)} \text{Cor}_{n,i})/I_n(M_n) = O\left(nr_n^{d+(1/4)}e^{nf_0\theta_{d-1}r_n^d(\frac{1}{2}-\kappa)}\right) = o\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right).$$

Therefore

$$I_n(\text{Mo}_n) = I_n(M_n) + I_n(\cup_{i=1}^{m(n)} \text{Cor}_{n,i}) = e^{-nf_0\theta_{d-1}r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\Gamma|\left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right). \quad (4.29)$$

Step 5: putting together the contributions. Recall that the set Γ was defined in Step 1, and consider now the region $\Gamma^{(2r_n)} \cap A$, where for $s > 0$ we set $\Gamma^{(s)} := \cup_{x \in \Gamma} B_s(x)$. Most of this is covered by $\text{Mo}_n \cup \text{Rai}_n \cup A^{(-r_n)}$, the exception being a leftover region near $\hat{\phi}(U \setminus U_n)$, where for $u \in U$ we set $\hat{\phi}(u) = (u, \phi(u) - Kr_n^{3/2})$ as before, that is covered by $(\partial\Gamma)^{(9dr_n^{3/4})} \cap A$. Using the fact that Γ is ‘nice’, it can be seen that $|(\partial\Gamma)^{(9dr_n^{3/4})}| = O(r_n^{3/2})$, and therefore given $\kappa < 1/2$, using Lemma 3.3 we have

$$I_n((\partial\Gamma)^{(9dr_n^{3/4})} \cap A) = O(r_n^{3/2}n \exp(-nf_0\kappa\theta_{d-1}r_n^d)).$$

Hence, taking κ close to $1/2$ and using (4.29) we obtain that

$$I_n((\partial\Gamma)^{(9dr_n^{3/4})} \cap A)/I_n(\text{Mo}_n) = O(nr_n^{d+3/2}e^{f_0\theta_{d-1}nr_n^d((1/2)-\kappa)}) = o\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right). \quad (4.30)$$

Also by (4.22),

$$I_n(\text{Rai}_n)/I_n(\text{Mo}_n) = O(nr_n^{d+1/2}e^{nf_0\theta dr_n^d((1/2)-\kappa)}) = o\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right). \quad (4.31)$$

Since $\text{Mo}_n \cup \text{Rai}_n \subset \Gamma^{(2r_n)} \subset \text{Mo}_n \cup \text{Rai}_n \cup (A^{(-r_n)} \setminus A^{(-2r_n)}) \cup ((\partial\Gamma)^{(9dr_n^{3/4})} \cap A)$, and since $I_n(A^{(-r_n)} \setminus A^{(-2r_n)}) = O(r_n I_n(A^{(-r_n)}))$, combining (4.29), (4.30) and (4.31) shows that

$$I_n(\Gamma^{(2r_n)}) = e^{-nf_0\theta r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\Gamma|\left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right) + o(r_n I_n(A^{(-r_n)})). \quad (4.32)$$

Now suppose $y \in \Gamma$ and $z \in \partial A \setminus \Gamma$ with $\|y - z\| \leq 4r_n$. Then $\|\pi(y) - \pi(z)\| \leq 4r_n$. Also $\pi(y) \in U$ and $\pi(z) \in \mathbb{R}^{d-1} \setminus U$, and hence $\text{dist}(\pi(y), \partial U) \leq 4r_n$. Hence using (4.19) we have $\text{dist}(y, \partial\Gamma) \leq 5r_n$.

Using (4.15), we can find a finite collection of sets $\Gamma_1, \dots, \Gamma_J$ each having the properties of the Γ we have just been considering, such that $\partial A \subset \cup_{i=1}^J \Gamma_i$ and $\Gamma_1, \dots, \Gamma_J$ have disjoint interiors (see [10] for further details). Then for $i \neq j$, if $x \in \Gamma_i^{(2r_n)} \cap \Gamma_j^{(2r_n)}$ then there exist $y \in \Gamma_i$ and $z \in \Gamma_j$ with $\|y - x\| \leq 2r_n$ and $\|z - x\| \leq 2r_n$, so $\|y - z\| \leq 4r_n$, so that $\text{dist}(y, \partial\Gamma_i) \leq 5r_n$ and $\text{dist}(x, \partial\Gamma_i) \leq 7r_n$. Hence

$$\Gamma_i^{(2r_n)} \cap \Gamma_j^{(2r_n)} \subset (\partial\Gamma_i)^{(7r_n)}.$$

Hence, using the union bound and the Bonferroni lower bound for the measure $I_n(\cdot)$, and (4.30), we obtain that

$$\sum_{i=1}^J I_n(\Gamma_i^{(2r_n)}) \geq I_n(\cup_{i=1}^J \Gamma_i^{(2r_n)}) \geq \sum_{i=1}^J I_n(\Gamma_i^{(2r_n)}) - o\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2 I_n(\text{Mo}_n)\right).$$

Then using (4.32) we obtain that

$$I_n(A \setminus A^{(-2r_n)}) = I_n(\cup_{i=1}^J \Gamma_i^{(2r_n)}) = e^{-nf_0\theta r_n^d/2}\theta_{d-1}^{-1}r_n^{1-d}|\partial A|\left(1 + o\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right) + O(r_n I_n(A^{(-r_n)})). \quad (4.33)$$

The error term in the last expression satisfies

$$\frac{r_n I_n(A^{(-r_n)})}{e^{-nf_0\theta r_n^d/2}r_n^{1-d}} = O(nr_n^d e^{-nf_0\theta r_n^d/2}) = O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right),$$

Also the error term in (4.21) satisfies a similar bound. Thus combining (4.33) with (4.21) shows that (4.14) holds. \square

5 Proof of first-order asymptotics

Throughout this section we make the same assumptions on ν and A that we set out at the start of Section 3. We also assume that (4.1) and (4.2) hold, i.e. that $nr_n^d \rightarrow \infty$ and $\liminf(I_n) > 0$ as $n \rightarrow \infty$.

We shall prove that if ξ_n denotes any of $K_n - 1, K'_n - 1, R_n$ or R'_n , then both $\mathbb{E}[\xi_n]$ and $\text{Var}[\xi_n]$ are asymptotic to I_n (which was defined at (2.1)) as $n \rightarrow \infty$; we will then be able to prove the first-order convergence results from Section 2, i.e. Theorems 2.1, 2.3, 2.4 and 2.7.

To achieve this goal, we shall consider separately the contributions to ξ_n from non-singleton components of $G(\mathcal{X}_n, r_n)$ or $G(\mathcal{P}_n, r_n)$ that are small, medium or large. Here, given fixed $\rho > \varepsilon > 0$, we say a component is *small* (respectively *medium*, *large*) if its Euclidean diameter is less than εr_n (resp., between εr_n and ρr_n , greater than ρr_n). We shall make appropriate choices of the constants ε, ρ as we go along.

For finite $\mathcal{X} \subset \mathbb{R}^d$, $x \in \mathcal{X}$, and $n \geq 1$ we let $\mathcal{F}_n(x, \mathcal{X})$ denote the event that x is the first element of $\mathcal{C}_{r_n}(x, \mathcal{X})$ (defined in Section 3.3) in the \prec ordering (defined in Section 3.1), i.e.

$$\mathcal{F}_n(x, \mathcal{X}) := \{x \prec y \ \forall \ y \in \mathcal{C}_{r_n}(x, \mathcal{X}) \setminus \{x\}\}. \quad (5.1)$$

Given n and $(r_n)_{n \geq 1}$, for $0 \leq \varepsilon < \rho \leq \infty$ we define $K_{n,\varepsilon,\rho}(\mathcal{X})$ to be the number of components of $G(\mathcal{X}, r_n)$ that have Euclidean diameter in the range $(\varepsilon r_n, \rho r_n]$, and $R_{n,\varepsilon,\rho}$ to be the number of vertices in such components, that is, with event $\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X})$ defined at (3.14),

$$K_{n,\varepsilon,\rho}(\mathcal{X}) := \sum_{x \in \mathcal{X}} \mathbf{1}_{\mathcal{M}_{n,\varepsilon,\rho}(\mathcal{X}) \cap \mathcal{F}_n(x, \mathcal{X})}; \quad R_{n,\varepsilon,\rho}(\mathcal{X}) := \sum_{x \in \mathcal{X}} \mathbf{1}_{\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X})}. \quad (5.2)$$

We then define the random variables $K_{n,\varepsilon,\rho} := K_{n,\varepsilon,\rho}(\mathcal{X}_n)$ and $K'_{n,\varepsilon,\rho} := K_{n,\varepsilon,\rho}(\mathcal{P}_n)$. Also we set $R_{n,\varepsilon,\rho} := R_{n,\varepsilon,\rho}(\mathcal{X}_n)$ and $R'_{n,\varepsilon,\rho} := R_{n,\varepsilon,\rho}(\mathcal{P}_n)$.

5.1 Asymptotics of means

We shall bound the expected number of ‘small’ non-singleton components, $\mathbb{E}[K_{n,0,\varepsilon}]$, using the following lemma.

Lemma 5.1. *There exist $\delta_4 \in (0, 1)$ and $c, n_0 < \infty$ such that for all $n \geq n_0$ and any $x \in A$,*

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n) \cap \{0 < \text{diam}(\mathcal{C}_{r_n}(x, \mathcal{P}_n)) \leq \delta_4 r_n\}] \leq c(nr_n^d)^{1-d} e^{-n\nu(B_{r_n}(x))}; \quad (5.3)$$

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{X}_{n-1}) \cap \{0 < \text{diam}(\mathcal{C}_{r_n}(x, \mathcal{X}_{n-1})) \leq \delta_4 r_n\}] \leq c(nr_n^d)^{1-d} e^{-n\nu(B_{r_n}(x))}. \quad (5.4)$$

Proof. See [11, Lemma 4.2(i)], taking $k = 1$ there. Note that a 0-separating set in \mathcal{X} (as it is called in [11]) is simply a component of $G(\mathcal{X}, r_n)$. Note also that if $A = [0, 1]^2$ the proof of [11, Lemma 4.2(i)] remains applicable, using Lemma 3.1 of the present paper. \square

We shall bound the expected number of ‘medium-sized’ non-singleton components, $\mathbb{E}[K_{n,\varepsilon,\rho}]$, using the following two lemmas (here we use notation such as \mathcal{X}^x from (3.13) and $\mathcal{M}_{n,\varepsilon,K}(x, \mathcal{X})$ from (3.14)).

Lemma 5.2. *Let $\varepsilon, \rho \in (0, \infty)$ with $\varepsilon < \rho$. Then there exists $\delta > 0$ such that for all n large and all distinct $x, y, z \in A$, we have:*

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n^x) \cap \mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n)] \leq e^{-n\nu(B_{r_n}(x)) - \delta nr_n^d}; \quad (5.5)$$

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n^{x,y}) \cap \mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n^y)] \leq e^{-n\nu(B_{r_n}(x)) - \delta nr_n^d}; \quad (5.6)$$

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n^{x,y,z}) \cap \mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n^{y,z})] \leq e^{-n\nu(B_{r_n}(x)) - \delta nr_n^d}; \quad (5.7)$$

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{X}_{n-1}^x) \cap \mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X}_{n-1})] \leq e^{-n\nu(B_{r_n}(x)) - \delta nr_n^d}; \quad (5.8)$$

$$\mathbb{P}[\mathcal{F}_n(x, \mathcal{X}_{n-2}^{x,y}) \cap \mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X}_{n-2}^y)] \leq e^{-n\nu(B_{r_n}(x)) - \delta nr_n^d}. \quad (5.9)$$

Proof. See [11, Lemma 4.3], taking the parameter k there to be 1, for a proof of (5.5) and (5.8). The results (5.6), (5.7) and (5.9) are proved similarly. In the case where $A = [0, 1]^2$, [11, Lemma 4.3] applies only when x is not too close to any of the corners of A , but the remaining cases can be proved by a similar argument using [7, Proposition 5.15]. \square

Lemma 5.3 (Bound on means for moderately large components). *There exists $\rho_1 \in (1, \infty)$ such that $\mathbb{E}[R_{n,\rho_1,(\log n)^2}] = O(e^{-nr_n^d} I_n)$ and $\mathbb{E}[R'_{n,\rho_1,(\log n)^2}] = O(e^{-nr_n^d} I_n)$ as $n \rightarrow \infty$, where I_n is defined at (2.1).*

Proof. Let $\rho > 4$. Given $i \in [n] := \{1, \dots, n\}$, if $\rho r_n < \text{diam}(\mathcal{C}_{r_n}(X_i, \mathcal{X}_n)) \leq (\log n)^2 r_n$, then there is at least one component of $G(\mathcal{X}_{n-1}, r_n)$ with at least one vertex in $B_{r_n}(X_i)$ and with diameter in the range $((\rho - 4)r_n/2, (\log n)^2 r_n]$. Hence by the definition at (3.15),

$$\mathbb{E}[R_{n,\rho,(\log n)^2}] \leq n \int_A \mathbb{P}[\mathcal{M}_{n,(\rho-4)/2,(\log n)^2}^*(x, \mathcal{X}_{n-1})] \nu(dx).$$

Hence by Lemma 3.8 we can choose ρ such that for n large

$$\mathbb{E}[R_{n,\rho,(\log n)^2}] \leq n \exp(-(\theta f_0 + 2)nr_n^d),$$

and hence using Lemma 4.1 we have the result claimed for $\mathbb{E}[R_{n,\rho,(\log n)^2}]$. The result for $\mathbb{E}[R'_{n,\rho,(\log n)^2}]$, is proved similarly, using the Mecke formula. \square

We shall approximate R_n with $S_n + R_{n,0,(\log n)^2}$ and K_n with $S_n + K_{n,0,(\log n)^2} + 1$.

Lemma 5.4. *Let $K > 0$. Then all of $\mathbb{P}[R_n \neq S_n + R_{n,0,(\log n)^2}]$, $\mathbb{P}[R'_n \neq S'_n + R'_{n,0,(\log n)^2}]$, $\mathbb{P}[K_n \neq S_n + K_{n,0,(\log n)^2} + 1]$ and $\mathbb{P}[K'_n \neq S'_n + K'_{n,0,(\log n)^2} + 1]$ are $O(n^{-K} I_n e^{-nr_n^d})$ as $n \rightarrow \infty$.*

Proof. By the assumption $\liminf(I_n) > 0$ and Proposition 2.2 we have $nr_n^d = O(\log n)$ as $n \rightarrow \infty$. Hence for some $\alpha > 0$, we have $I_n e^{-nr_n^d} > n^{-\alpha}$ for all large n . Therefore it suffices to prove that for any $K > 0$, the probabilities under consideration are $O(n^{-K})$ as $n \rightarrow \infty$.

Define event $\tilde{\mathcal{W}}_n$ as in Lemma 3.7, taking $\phi_n = (\log n)^2$. Then recalling the definition of $\mathcal{L}_n(\mathcal{X})$ just before Lemma 3.10, we have the event inclusion

$$\{R_n \neq S_n + R_{n,0,(\log n)^2}\} \cup \{K_n \neq S_n + K_{n,0,(\log n)^2} + 1\} \subset \tilde{\mathcal{W}}_n^c \cup \{\text{diam}(\mathcal{L}_n(\mathcal{X}_n)) \leq (\log n)^2 r_n\}.$$

By Lemma 3.7, there is a constant c such that $\mathbb{P}[\tilde{\mathcal{Z}}_n^c] \leq \exp(-c(\log n)^2 nr_n^d)$ for n large. Combining this with (3.18) from Lemma 3.10 (which is applicable because $nr_n^d = O(\log n)$ as mentioned already) gives us the results for R_n and K_n , and the results for R'_n and K'_n are proved similarly. \square

Proposition 5.5 (Approximation of K_n by $S_n + 1$, K'_n by $S'_n + 1$). *As $n \rightarrow \infty$ we have*

$$\max(\mathbb{E}[|K'_n - S'_n - 1|], \mathbb{E}[|K_n - S_n - 1|]) = O((nr_n^d)^{1-d} I_n). \quad (5.10)$$

Proof. Take δ_4 as in Lemma 5.1 and ρ_1 as in Lemma 5.3. Then $K'_n - S'_n = K'_{n,0,\delta_4} + K'_{n,\delta_4,\rho_1} + K'_{n,\rho_1,(\log n)^2} + K'_{n,(\log n)^2,\infty}$. Taking expectations and using the Mecke formula, we obtain that

$$\begin{aligned} \mathbb{E}[|K'_n - S'_n - 1|] &\leq n \int_A \mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n) \cap \{0 < \text{diam}(\mathcal{C}_{r_n}(x, \mathcal{P}_n)) \leq \delta_4 r_n\}] \nu(dx) \\ &\quad + \int_A \mathbb{P}[\mathcal{F}_n(x, \mathcal{P}_n) \cap \{\delta_4 r_n < \text{diam}(\mathcal{C}_{r_n}(x, \mathcal{P}_n)) \leq \rho_1 r_n\}] n \nu(dx) \\ &\quad + \mathbb{E}[K'_{n,\rho_1,(\log n)^2}] + \mathbb{E}[|K'_{n,(\log n)^2,\infty} - 1|]. \end{aligned} \quad (5.11)$$

By Lemma 5.1, the first term in the right hand side of (5.11) is $O((nr_n^d)^{1-d} I_n)$. By Lemma 5.2 there exists $\delta > 0$ such that the second term in the right hand side of (5.11) is at most $e^{-\delta nr_n^d} I_n$ for all large enough n . By Lemma 5.3, the third term in the right hand side is $O(e^{-nr_n^d} I_n)$. For the fourth term, recalling $\#(\mathcal{P}_n) = Z_n$ is Poisson with mean n , we use the Cauchy-Schwarz inequality, and then Lemma 5.4, to deduce that

$$\mathbb{E}[|K'_{n,(\log n)^2,\infty} - 1|] \leq (\mathbb{E}[Z_n^2 + 1])^{1/2} (\mathbb{P}[K'_{n,(\log n)^2,\infty} \neq 1])^{1/2} = O(e^{-nr_n^d} I_n).$$

Combining these estimates shows that $\mathbb{E}[|K'_n - S'_n - 1|] = O((nr_n^d)^{1-d} I_n)$. The proof that $\mathbb{E}[|K_n - S_n - 1|] = O((nr_n^d)^{1-d} I_n)$ is similar, and thus we have (5.10). \square

Lemma 5.6. *Suppose δ_1, δ_4 are as in Lemma 3.1, Lemma 5.1 respectively, and $0 < \rho < \min(\frac{1}{2}, \delta_4, (\delta_1 f_0 / (f_{\max} \theta))^{1/(d-1)})$. Then as $n \rightarrow \infty$, we have*

$$\max(\mathbb{E}[R_{n,0,\rho}], \mathbb{E}[R'_{n,0,\rho}]) = O((nr_n^d)^{1-d} I_n). \quad (5.12)$$

Proof. If the interpoint distances of \mathcal{X}_n are all distinct and non-zero (an event of probability 1), then $R_{n,0,\rho} = K_{n,0,\rho} + N_{n,1} + N_{n,2}$, where we set

$$\begin{aligned} N_{n,1} := & \sum_{(i,j) \in [n] \times [n]: i \neq j} \mathbf{1}(\mathcal{F}_n(X_i, \mathcal{X}_n) \cap \{0 < \text{diam}(\mathcal{C}_{r_n}(X_i, \mathcal{X}_n)) \leq \rho r_n\} \cap \{X_j \in \mathcal{C}_{r_n}(X_i, \mathcal{X}_n)\} \\ & \cap \{\|X_j - X_i\| = \max_{x \in \mathcal{C}_{r_n}(X_i, \mathcal{X}_n)} \|x - X_i\|\}), \end{aligned}$$

and

$$\begin{aligned} N_{n,2} := & \sum_{(i,j,k) \in [n] \times [n] \times [n]: i \neq j \neq k \neq i} \mathbf{1}(\mathcal{F}_n(X_i, \mathcal{X}_n) \cap \{0 < \text{diam}(\mathcal{C}_{r_n}(X_i, \mathcal{X}_n)) \leq \rho r_n\} \\ & \cap \{\{X_j, X_k\} \subset \mathcal{C}_{r_n}(X_i, \mathcal{X}_n)\} \cap \{\|X_j - X_i\| = \max_{x \in \mathcal{C}_{r_n}(X_i, \mathcal{X}_n)} \|x - X_i\|\}). \end{aligned}$$

Using Lemma 5.1 and the Mecke formula as in (5.11), we have that

$$\mathbb{E}[K_{n,0,\rho}] = O((nr_n^d)^{1-d}I_n). \quad (5.13)$$

Recall the notation $A_x := \{y \in A : x \prec y\}$. By assumption $f_{\max}\theta\rho^{d-1} < \delta_1 f_0$, so by Lemma 3.1, for n large and $x \in A, y \in B(x, \rho r_n) \cap A_x$ we have $\nu(B_{r_n}(y) \setminus B_{r_n}(x)) \geq 2\delta_1 f_0 r_n^{d-1} \|y - x\|$ and $\nu(B_{\|y-x\|}(x)) \leq f_{\max}\theta \|y - x\|^d \leq \delta_1 f_0 r_n^{d-1} \|y - x\|$. Hence

$$\begin{aligned} \mathbb{E}[N_{n,1}] &\leq n^2 \int_A \int_{B(x, \rho r_n) \cap A_x} (1 - \nu[(B_{r_n}(x) \cup B_{r_n}(y)) \setminus B_{\|y-x\|}(x)])^{n-2} \nu(dy) \nu(dx) \\ &\leq 2n^2 \int_A \int_{B(x, \rho r_n) \cap A_x} e^{-n(\nu(B_{r_n}(x)) + \nu(B_{r_n}(y) \setminus B_{r_n}(x)) - \nu(B_{\|y-x\|}(x)))} \nu(dy) \nu(dx) \\ &\leq 2n \int_A \left(\int_{B(x, \rho r_n) \cap A_x} n e^{-n\nu(B_{r_n}(x)) - n\delta_1 f_0 r_n^{d-1} \|y-x\|} \nu(dy) \right) \nu(dx). \end{aligned}$$

In the last expression the inner integral can be bounded by

$$n e^{-n\nu(B_{r_n}(x))} \int_{B(o, \rho r_n)} e^{-\delta_1 f_0 n r_n^{d-1} \|u\|} f_{\max} du = (nr_n^d)^{1-d} e^{-n\nu(B_{r_n}(x))} \int_{B(o, \rho n r_n^d)} e^{-\delta_1 f_0 \|v\|} f_{\max} dv$$

and therefore

$$\mathbb{E}[N_{n,1}] = O((nr_n^d)^{1-d}I_n). \quad (5.14)$$

Next, using Lemma 3.1 and the inequality $f_{\max}\theta\rho^{d-1} < \delta_1 f_0$ again we have that

$$\begin{aligned} \mathbb{E}[N_{n,2}] &\leq n^3 \int_A \int_{B(x, \rho r_n) \cap A_x} \int_{B(x, \|y-x\|)} (1 - \nu[(B_{r_n}(x) \cup B_{r_n}(y)) \setminus B_{\|y-x\|}(x)])^{n-3} \\ &\quad \nu(dz) \nu(dy) \nu(dx) \\ &\leq 2n\theta_d f_{\max} \int_A \left(\int_{B(x, \rho r_n) \cap A_x} n^2 \|y - x\|^d \exp(-n\nu(B_{r_n}(x)) - n\delta_1 f_0 r_n^{d-1} \|y - x\|) \nu(dy) \right) \\ &\quad \nu(dx). \end{aligned}$$

In the last expression the inner integral can be bounded by

$$\begin{aligned} &n^2 e^{-n\nu(B_{r_n}(x))} \int_{B(o, \rho r_n)} \|u\|^d e^{-\delta_1 f_0 n r_n^{d-1} \|u\|} f_{\max} du \\ &= (nr_n^d)^{2-2d} e^{-n\nu(B_{r_n}(x))} \int_{B(o, \rho n r_n^d)} \|v\|^d e^{-\delta_1 f_0 \|v\|} f_{\max} dv, \end{aligned}$$

and therefore

$$\mathbb{E}[N_{n,2}] = O((nr_n^d)^{2-2d}I_n).$$

Combined with (5.13) and (5.14) this shows that $\mathbb{E}[R_{n,0,\rho}] = O((nr_n^d)^{1-d}I_n)$, which is the statement about $\mathbb{E}[R_{n,0,\rho}]$ in (5.12). The corresponding statement for $\mathbb{E}[R'_{n,0,\rho}]$ is proved similarly using the multivariate Mecke formula. \square

For $0 < \varepsilon < \rho < \infty$, recall the definition of event $\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X})$ at (3.14). To deal with the medium-sized components, we shall use the following estimate for the integral of $\mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \xi_n)]$ with $\xi_n = \mathcal{P}_n$ or $\xi_n = \mathcal{X}_{n-1}$. We use notation \mathcal{X}^x from (3.13).

Lemma 5.7 (Estimate on medium clusters). *Let $\rho, \varepsilon \in (0, \infty)$ with $\rho > \varepsilon$. Then there exists $\delta = \delta(\varepsilon)$ in $(0, \infty)$ such that as $n \rightarrow \infty$, we have*

$$n \int_A \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n)] \nu(dx) = O(e^{-\delta nr_n^d} I_n); \quad (5.15)$$

$$n \int_A \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{X}_{n-1})] \nu(dx) = O(e^{-\delta nr_n^d} I_n). \quad (5.16)$$

Proof. If $\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n) \setminus \mathcal{F}_n(x, \mathcal{P}_n^x)$ occurs then for at least one $y \in \mathcal{P}_n \cap B_{\rho r_n}(x)$ we have that $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{P}_n^{x,y})) \in (\varepsilon r_n, \rho r_n]$, and moreover $\mathcal{F}_n(y, \mathcal{P}_n^{x,y})$ occurs and $x \in \mathcal{C}_{r_n}(y, \mathcal{P}_n^{x,y})$. By Markov's inequality and the Mecke formula,

$$n \int_A \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n) \setminus \mathcal{F}_n(x, \mathcal{P}_n^x)] \nu(dx) \leq n^2 \int_A \int_{A \cap B_{\rho r_n}(x)} \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(y, \mathcal{P}_n^x) \cap \mathcal{F}_n(y, \mathcal{P}_n^{x,y}) \cap \{x \in \mathcal{C}_{r_n}(y, \mathcal{P}_n^{x,y})\}] \nu(dy) \nu(dx).$$

By (5.6) from Lemma 5.2, there exists $\delta > 0$ such that for n large the probability inside the integral on the right of the last display is bounded above by $\exp(-n\nu(B_{r_n}(y)) - 2\delta nr_n^d)$. Then using Fubini's theorem we obtain that for n large

$$\begin{aligned} n \int_A \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n) \setminus \mathcal{F}_n(x, \mathcal{P}_n^x)] \nu(dx) &\leq n^2 \int_A \nu(B_{\rho r_n}(y)) \exp(-n\nu(B_{r_n}(y)) - 2\delta nr_n^d) \nu(dy) \\ &= O(nr_n^d I_n e^{-2\delta nr_n^d}) = O(e^{-\delta nr_n^d} I_n). \end{aligned} \quad (5.17)$$

Also using Lemma 5.2 we obtain that

$$\begin{aligned} n \int_A \mathbb{P}[\mathcal{M}_{n,\varepsilon,\rho}(x, \mathcal{P}_n) \cap \mathcal{F}_n(x, \mathcal{P}_n^x)] \nu(dx) &\leq n \int_A \exp(-n\nu(B_{r_n}(x)) - \delta nr_n^d) \nu(dx) \\ &= e^{-\delta nr_n^d} I_n, \end{aligned}$$

and combined with (5.17) this yields (5.15).

The proof of (5.16) is similar. \square

We are now ready to estimate the asymptotic expected values of R_n and R'_n .

Proposition 5.8 (Approximation of R_n, R'_n by S_n, S'_n). *As $n \rightarrow \infty$ we have that*

$$\mathbb{E}[|R_n - S_n|] = O((nr_n^d)^{1-d} I_n); \quad (5.18)$$

$$\mathbb{E}[|R'_n - S'_n|] = O((nr_n^d)^{1-d} I_n). \quad (5.19)$$

Proof. Note that $|R_n - S_n - R_{n,0,(\log n)^2}| \leq n$. Hence by Lemma 5.4,

$$\mathbb{E}[|R_n - S_n - R_{n,0,(\log n)^2}|] \leq n \mathbb{P}[R_n \neq S_n + R_{n,0,(\log n)^2}] = O(e^{-nr_n^d} I_n). \quad (5.20)$$

Using Lemma 5.6, choose $\varepsilon \in (0, 1)$ such that $\mathbb{E}[R_{n,0,\varepsilon}] = O((nr_n^d)^{1-d}I_n)$. Using Lemma 5.3, choose $\rho \in (1, \infty)$ such that $\mathbb{E}[R_{n,\rho,(\log n)^2}] = O(e^{-nr_n^d}I_n)$. By Lemma 5.7, there exists $\delta > 0$ such that

$$\mathbb{E}[R_{n,\varepsilon,\rho}] = n \int_A \mathbb{P}[\varepsilon r < \text{diam } \mathcal{C}_{r_n}(x, \mathcal{X}_{n-1}^x) \leq \rho r_n] \nu(dx) = O(e^{-n\delta r_n^d}I_n).$$

Combining these estimates shows that $\mathbb{E}[R_{n,0,(\log n)^2}] = O((nr_n^d)^{1-d}I_n)$. Then using (5.20) yields (5.18).

The proof of (5.19) is similar; the only difference is that in the step of the argument corresponding to (5.20) we use the inequality $|R'_n - S'_n - R'_{n,0,(\log n)^2}| \leq Z_n \mathbf{1}\{R'_n \neq S'_n + R'_{n,0,(\log n)^2}\}$ and the Cauchy-Schwarz inequality. \square

5.2 Proof of first order limit theorems

Proof of Theorem 2.1. By Proposition 5.5 we have

$$\mathbb{E}[K'_n - 1] = \mathbb{E}[S'_n] + \mathbb{E}[K'_n - 1 - S'_n] = I_n(1 + O(nr_n^d)^{1-d}).$$

By Proposition 5.5 and Lemma 4.3 we also have $\mathbb{E}[K_n - 1] = I_n(1 + O(nr_n^d)^{1-d})$. By Proposition 5.8 we have $\mathbb{E}[R'_n] = I_n(1 + O(nr_n^d)^{1-d})$. By Proposition 5.8 and Lemma 4.3 we have $\mathbb{E}[R_n] = \mathbb{E}[S_n] + O((nr_n^d)^{1-d}I_n) = I_n(1 + O(nr_n^d)^{1-d})$. Thus we have (2.2).

For (2.3), suppose for now that ζ_n is $K_n - 1$ or R_n . Recalling that $\tilde{I}_n := \mathbb{E}[S_n]$, we note that

$$\mathbb{E}[|(\zeta_n/I_n) - 1|] \leq \mathbb{E}[I_n^{-1}|\zeta_n - S_n|] + \mathbb{E}[I_n^{-1}|S_n - \tilde{I}_n|] + I_n^{-1}|\tilde{I}_n - I_n|. \quad (5.21)$$

By Proposition 5.5 (when $\zeta_n = K_n - 1$) or Proposition 5.8 (when $\zeta_n = R_n$) we have $\mathbb{E}[|\zeta_n - S_n|] = O((nr_n^d)^{1-d}I_n)$, so the first term in the right hand side of (5.21) is $O((nr_n^d)^{1-d})$. Moreover by the Cauchy-Schwarz inequality the second term in the right hand side of (5.21) is bounded by $(I_n^{-2}\text{Var}(S_n))^{1/2}$, and by Proposition 4.5 this is $O(I_n^{-1/2})$. The third term in the right hand side of (5.21) is $O((nr_n^d)^{1-d})$ by Lemma 4.3. Thus we have (2.3) when ζ_n is $K_n - 1$ or R_n , and the corresponding result when ζ_n is $K'_n - 1$ or R'_n can be proved similarly. Finally if we add 1 to ζ_n then we should add a term of $1/I_n$ on the right hand side of (5.21), but this term is $o(I_n^{-1/2})$ so we have (2.3) when ζ_n is $K_n, R_n + 1, K'_n$ or $R'_n + 1$ too. \square

Proof of Theorem 2.3. Since we assume $I_n \rightarrow \infty$, by Proposition 2.2 we have $b^+ \leq b_c$. Suppose also $b^+ \leq b'_c$. Let $\delta > 0$. Then by Lemma 4.1,

$$ne^{-n\theta f_0 r_n^d(1+\delta)} = o(I_n). \quad (5.22)$$

For an upper bound on I_n , we shall use Lemma 4.2. Let $\varepsilon > 0$ with $f_1\varepsilon < f_0\delta$. Since $b^+ \leq b'_c$ we have $b^+(f_0 - f_1/2) \leq 1/d$, and hence

$$\frac{n^{1-1/d}e^{-n\theta r_n^d f_1(\frac{1}{2}-\varepsilon)}}{ne^{-n\theta f_0 r_n^d(1-2\delta)}} = O(n^{-1/d}e^{n\theta r_n^d(f_0-(f_1/2)-\delta)}) = O(n^{-1/d}e^{b^+(f_0-f_1/2)-\frac{1}{2}\delta} \log n) = o(1).$$

Therefore both terms in the right hand side of (4.7) are $o(ne^{-n\theta f_0 r_n^d(1-2\delta)})$, so by Lemma 4.2, $I_n = o(ne^{-n\theta r_n^d f_0(1-2\delta)})$. Using this, along with (5.22) and the fact that the L^1 convergence in (2.3) implies convergence in probability also, we obtain that with probability tending to one, $ne^{-n\theta r_n^d f_0(1+2\delta)} < \zeta_n < ne^{-n\theta r_n^d f_0(1-2\delta)}$, which gives us (2.6).

Now suppose $b^- \geq b'_c$. Since also $b^+ \leq b_c$, we have $b'_c \leq b_c < \infty$. Hence $b'_c = (d(f_0 - f_1/2))^{-1}$ so $b^- \geq (d(f_0 - f_1/2))^{-1}$ and $b^-((f_1/2) - f_0) \leq -1/d$. Let $\varepsilon > 0$. Then

$$\frac{ne^{-n\theta f_0 r_n^d}}{n^{1-1/d} e^{-nr_n^d \theta f_1(\frac{1}{2}-\varepsilon)}} = n^{1/d} e^{nr_n^d \theta ((f_1/2) - f_0 - f_1 \varepsilon)} \leq n^{1/d} e^{b^-((f_1/2) - f_0 - f_1 \varepsilon/2) \log n} = o(1),$$

so by Lemma 4.2, $I_n = o(n^{1-1/d} e^{-n\theta r_n^d f_1(\frac{1}{2}-2\varepsilon)})$. Also by Lemma 4.1, $n^{1-1/d} \exp^{-nr_n^d \theta f_1(\frac{1}{2}+\varepsilon)} = o(I_n)$. Hence by the convergence in probability of ζ_n/I_n to 1 which follows from (2.3), with probability tending to 1 we have $n^{1-1/d} \exp^{-nr_n^d \theta f_1(\frac{1}{2}+\varepsilon)} \leq \zeta_n \leq n^{1-1/d} e^{-n\theta r_n^d f_1(\frac{1}{2}-2\varepsilon)}$, and (2.7) follows.

Now suppose $b^+ = b^- = b$ for some $b \geq 0$. Then if $f_0 b \leq (1/d) + f_1 b/2$ we have $(f_0 - f_1/2)b \leq 1/d$ so $b \leq b'_c$ and (2.6) applies. By (2.6) we have $\zeta_n = n^{1-bf_0+o_{\mathbb{P}}(1)}$. Conversely if $f_0 b \geq (1/d) + f_1 b/2$ we have $(f_0 - f_1/2)b \geq 1/d$ and $b \geq b'_c$ so (2.7) applies and tells us that $\zeta_n = n^{1-(1/d)-f_1 b/2+o_{\mathbb{P}}(1)}$. \square

Proof of Theorem 2.4. Here we assume as $n \rightarrow \infty$ that $I_n = \Theta(1)$ (which implies $nr_n^d = \Theta(\log n)$ by Proposition 2.2 and Lemma 4.1). Then by Lemma 4.6, for some $\delta > 0$ we have $d_{\text{TV}}(S'_n, Z_{I_n}) = O(e^{-\delta nr_n^d})$.

By Proposition 5.5 (when $\xi_n = K'_n - 1$) or Proposition 5.8 (when $\xi_n = R'_n$) and Markov's inequality, for both those cases $d_{\text{TV}}(\xi_n, S'_n) \leq \mathbb{P}[\xi_n \neq S'_n] \leq \mathbb{E}[|\xi_n - S'_n|] = O((nr_n^d)^{1-d})$, and therefore by Lemma 4.6 and the triangle inequality, $d_{\text{TV}}(\xi_n, Z_{I_n}) = O((nr_n^d)^{1-d}) = O((\log n)^{1-d})$ in those cases.

Now suppose ξ_n is $K_n - 1$ or R_n . By Proposition 5.5 (when $\xi_n = K_n - 1$) or Proposition 5.8 (when $\xi_n = R_n$) and Markov's inequality, for both those cases $\mathbb{P}[\xi_n \neq S_n] \leq \mathbb{E}[|\xi_n - S_n|] = O((nr_n^d)^{1-d})$, and therefore it suffices to prove that $d_{\text{TV}}(S_n, Z_{I_n}) = O((nr_n^d)^{1-d})$. By Lemma 4.6 we have for some $\delta > 0$ that $d_{\text{TV}}(S'_n, Z_{I_n}) = O(e^{-\delta nr_n^d})$, so it suffices to prove that $\mathbb{E}[|S'_n - S_n|] = O((nr_n^d)^{1-d})$.

Recall that $\mathcal{P}_n = \{X_1, \dots, X_{Z_n}\}$. Let $m = m(n) = \lfloor n^{3/4} \rfloor$. By the Cauchy-Schwarz inequality and the Chernoff bound from Lemma 3.4(ii),

$$\begin{aligned} \mathbb{E}[|S'_n - S_n| \mathbf{1}\{|Z_n - n| > m\}] &\leq (\mathbb{E}[\max(Z_n, n)^2])^{1/2} (\mathbb{P}[|Z_n - n| > m])^{1/2} \\ &\leq (2n^2 + n)^{1/2} \exp(-\Omega(n^{1/2})). \end{aligned} \quad (5.23)$$

For $i = 1, 2, \dots$ write $Y_i := X_{Z_n+i}$ and $Y'_i := X_{n+i}$. Then Y_1, Y_2, \dots are ν -distributed random vectors, independent of each other and of \mathcal{P}_n . Observe that

$$\begin{aligned} |S'_n - S_n| \mathbf{1}\{Z_n \leq n \leq Z_n + m\} &\leq \sum_{i=1}^m (\mathbf{1}\{\mathcal{P}_n \cap B_{r_n}(Y_i) = \emptyset\} \\ &\quad + \sum_{x \in \mathcal{P}_n \cap B_{r_n}(Y_i)} \mathbf{1}\{\mathcal{P}_n \cap B_{r_n}(x) = \{x\}\}). \end{aligned}$$

Therefore using the Mecke formula followed by Fubini's theorem we obtain that

$$\begin{aligned} \mathbb{E}[|S'_n - S_n| \mathbf{1}\{Z_n \leq n \leq Z_n + m\}] &\leq m \int_A e^{-n\nu(B_{r_n}(x))} dx \\ &\quad + mn \int_A \int_{B_{r_n}(x)} e^{-n\nu(B_{r_n}(y))} \nu(dy) \nu(dx) \\ &\leq n^{-1/4} I_n + (nf_{\max} \theta r_n^d) n^{-1/4} I_n = O(n^{-1/4} (\log n)). \end{aligned} \quad (5.24)$$

Also Y'_1, Y'_2, \dots are ν -distributed random vectors, independent of each other and of \mathcal{X}_n . Then since $(1 - \nu(B_{r_n}(x)))^{n-1} \leq 2e^{-n\nu(B_{r_n}(x))}$ for all large enough n and all $x \in A$,

$$\begin{aligned} \mathbb{E}[|S'_n - S_n| \mathbf{1}\{n \leq Z_n \leq n + m\}] &\leq \mathbb{E} \left[\sum_{i=1}^m \left(\mathbf{1}\{\mathcal{X}_n \cap B_{r_n}(Y'_i) = \emptyset\} \right. \right. \\ &\quad \left. \left. + \sum_{x \in \mathcal{X}_n \cap B_{r_n}(Y'_i)} \mathbf{1}\{\mathcal{X}_n \cap B_{r_n}(x) = \{x\}\} \right) \right] \\ &\leq m \int_A (1 - \nu(B_{r_n}(x)))^n \nu(dx) + mn \int_A \int_{B_{r_n}(x)} (1 - \nu(B_{r_n}(y)))^{n-1} \nu(dy) \nu(dx) \\ &= O((nr_n^d) n^{-1/4} I_n) = O(n^{-1/4} \log n). \end{aligned}$$

Combined with (5.23) and (5.24) this shows that $\mathbb{E}[|S'_n - S_n|] = O(n^{-1/4} \log n) = O((nr_n^d)^{1-d})$ as required. \square

Proof of Theorem 2.7. Assume the uniform case applies. We first show that for any $\gamma \in \mathbb{R}$ we have:

$$\text{if } \lim_{n \rightarrow \infty} \gamma_n = \gamma \text{ then } \lim_{n \rightarrow \infty} \mu_n = \begin{cases} e^{-\gamma} & \text{if } d = 2 \\ c_{d,A} e^{-\gamma/2} & \text{if } d \geq 3. \end{cases} \quad (5.25)$$

The case $d = 2$ of (5.25) is obvious because $\mu_n = e^{-\gamma_n}$ in this case. Suppose $d \geq 3$. If $\lim_{n \rightarrow \infty} \gamma_n = \gamma$, then as $n \rightarrow \infty$ the second term in the right hand side of (1.4) satisfies

$$\begin{aligned} \theta_{d-1}^{-1} |\partial A| r_n^{1-d} e^{-n\theta f_0 r_n^d/2} &\sim \theta_{d-1}^{-1} |\partial A| \left(\frac{(2 - 2/d) \log n}{n\theta f_0} \right)^{-1+1/d} e^{-\gamma/2} \left(\frac{n}{\log n} \right)^{-1+1/d} \\ &= \theta_{d-1}^{-1} \sigma_A (\theta / (2 - 2/d))^{1-1/d} e^{-\gamma/2} = c_{d,A} e^{-\gamma/2}, \end{aligned}$$

and moreover the ratio between the two terms in the right hand side of (1.4) satisfies

$$\begin{aligned} \frac{ne^{-n\theta f_0 r_n^d}}{\theta_{d-1}^{-1} |\partial A| r_n^{1-d} e^{-n\theta f_0 r_n^d/2}} &= \theta_{d-1} |\partial A|^{-1} n r_n^{d-1} e^{-n\theta f_0 r_n^d/2} \\ &= O \left((\log n) \left(\frac{n}{\log n} \right)^{-1+2/d} \right) = o(1), \end{aligned}$$

and (5.25) follows.

Now suppose $|\gamma_n| = O(1)$, which implies $nr_n^d = \Theta(\log n)$ as $n \rightarrow \infty$. By (5.25) and a subsequence argument we have that $\mu_n = \Theta(1)$ as $n \rightarrow \infty$. Let ξ_n be any of

$K_n - 1, R_n, K'_n - 1$ or R'_n . By a simple coupling argument for $0 < s < t$ we have $d_{\text{TV}}(Z_s, Z_t) \leq t - s$. Hence by the triangle inequality

$$d_{\text{TV}}(\xi_n, Z_{\mu_n}) \leq d_{\text{TV}}(\xi_n, Z_{I_n}) + |I_n - \mu_n|.$$

If $d = 2$ then by Proposition 4.7 and (1.4) $I_n = \mu_n(1 + O(nr_n^2)^{-1/2})$; hence by Theorem 2.4,

$$d_{\text{TV}}(\xi_n, Z_{\mu_n}) = O((\log n)^{-1}) + O((nr_n^2)^{-1/2}) = O((\log n)^{-1/2}).$$

If $d = 3$ then by Proposition 4.8 and (1.4), $I_n = \mu_n \left(1 + O\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)$; hence by Theorem 2.4,

$$d_{\text{TV}}(\xi_n, Z_{\mu_n}) = O((\log n)^{1-d}) + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right) = O\left(\left(\frac{\log \log n}{\log n}\right)^2\right).$$

Thus we have part (a). In particular, for all $d \geq 2$ we have $d_{\text{TV}}(\xi_n, Z_{\mu_n}) \rightarrow 0$ so that if $\gamma_n \rightarrow \gamma$ then by (5.25) we have $\xi_n \xrightarrow{\mathcal{D}} Z_{e^{-\gamma}}$ if $d = 2$ and $\xi_n \xrightarrow{\mathcal{D}} Z_{c_{d,A}e^{-\gamma/2}}$ if $d \geq 3$, which is part (b).

For part (c), now assume (1.1) and (1.2). First suppose $d = 2$. By Proposition 4.7, as $n \rightarrow \infty$ we have $I_n = \mu_n(1 + O((nr_n^2)^{-1/2}))$ and (2.12) follows from (2.2). Also by (2.3) from Theorem 2.1, and Proposition 4.7,

$$\mathbb{E}\left[\left|\frac{\xi_n}{\mu_n} - 1\right|\right] \leq \mathbb{E}\left[\left|\frac{\xi_n}{I_n}\left(\frac{I_n}{\mu_n} - 1\right)\right|\right] + \mathbb{E}\left[\left|\frac{\xi_n}{I_n} - 1\right|\right] = O((nr_n^2)^{-1/2} + I_n^{-1/2}),$$

and hence (2.13).

Suppose $d \geq 3$. By Proposition 4.8, as $n \rightarrow \infty$ we have $I_n = \mu_n \left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right)$. Hence using (2.2) we have (2.14). Also by (2.3) from Theorem 2.1, and Proposition 4.8, we have

$$\mathbb{E}\left[\left|\frac{\xi_n}{\mu_n} - 1\right|\right] \leq \mathbb{E}\left[\left|\frac{\xi_n}{I_n}\left(\frac{I_n}{\mu_n} - 1\right)\right|\right] + \mathbb{E}\left[\left|\frac{\xi_n}{I_n} - 1\right|\right] = O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2 + I_n^{-1/2}\right),$$

and hence (2.15). □

6 Asymptotics of variances

Throughout this section we make the same assumptions on d, A and f that were set out at the start of Section 3. We also assume that (4.1) and (4.2) hold, i.e. that $nr_n^d \rightarrow \infty$ and $\liminf(I_n) > 0$ as $n \rightarrow \infty$.

We shall prove that if ξ_n denotes any of $K_n - 1, K'_n - 1, R_n$ or R'_n , then $\text{Var}[\xi_n]$ is asymptotic to I_n (which was defined at (2.1)) as $n \rightarrow \infty$; in the case of $\text{Var}[K_n - 1]$ and $\text{Var}[R_n]$ we require the extra condition $d \geq 3$.

Later we shall show that the number of non-singleton components has negligible variance compared to the number of singletons. This goal will be achieved by estimating separately the variance for the number of non-singleton components of small (i.e., smaller than δr_n), medium and large (i.e., larger than ρr_n) diameter, and showing that each of these three variances is $o(I_n)$; the constants δ, ρ will be chosen later.

6.1 Variances for small components: Poisson input

Next we consider for $G(\mathcal{P}_n, r_n)$ the number of small non-singleton components $K'_{n,0,\rho}$ and the number of vertices in such components, $R'_{n,0,\rho}$ (as defined at (5.2)), for suitably small (fixed) ρ .

Proposition 6.1. *There exists $\rho_0 > 0$ such that if $0 < \rho < \rho_0$ then as $n \rightarrow \infty$ we have*

$$\max(\text{Var}[K'_{n,0,\rho}], \text{Var}[R'_{n,0,\rho}]) = O((nr_n^d)^{1-d}I_n). \quad (6.1)$$

We divide the proof of this proposition into a series of lemmas. Given $\rho > 0$ and given n , for $x, y \in A$ define the events $\mathcal{T}_x := \mathcal{M}_{n,0,\rho}(x, \mathcal{P}_n)$ and $\mathcal{T}_{x,y} := \mathcal{M}_{n,0,\rho}(x, \mathcal{P}_n^y)$, where $\mathcal{M}_{n,\varepsilon,K}(\mathcal{X})$ was defined at (3.14). Also, recalling the definition of $\mathcal{F}_n(x, \mathcal{X})$ at (5.1), set $\mathcal{E}_x := \mathcal{T}_x \cap \mathcal{F}_n(x, \mathcal{P}_n^x)$ and $\mathcal{E}_{x,y} := \mathcal{T}_{x,y} \cap \mathcal{F}_n(x, \mathcal{P}_n^{x,y})$. We begin with the following bound based on the Mecke formula.

Lemma 6.2. *Suppose $\rho \in (0, 1)$. Then*

$$\text{Var}[R'_{n,0,\rho}] - \mathbb{E}[R'_{n,0,\rho}] \leq n^2 \int_A \int_{A \cap B_{4r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \nu(dy) \nu(dx); \quad (6.2)$$

$$\text{Var}[K'_{n,0,\rho}] - \mathbb{E}[K'_{n,0,\rho}] \leq n^2 \int_A \int_{A \cap B_{4r_n}(x)} \mathbb{P}[\mathcal{E}_{x,y} \cap \mathcal{E}_{y,x}] \nu(dy) \nu(dx). \quad (6.3)$$

Proof. By the Mecke formula, we have $\mathbb{E}[R'_{n,0,\rho}] = n \int_A \mathbb{P}[\mathcal{T}_x] \nu(dx)$. Using this and the multivariate Mecke formula we obtain that

$$\mathbb{E}[R'_{n,0,\rho}(R'_{n,0,\rho} - 1)] - \mathbb{E}[R'_{n,0,\rho}]^2 = n^2 \int_A \int_A (\mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] - \mathbb{P}[\mathcal{T}_x] \mathbb{P}[\mathcal{T}_y]) \nu(dy) \nu(dx).$$

For $\|y - x\| > 4r_n > 2(1 + \rho)r_n$ we have $\mathbb{P}[\mathcal{T}_{x,y}] = \mathbb{P}[\mathcal{T}_x] \mathbb{P}[\mathcal{T}_y]$, and (6.2) follows.

The proof of (6.3) is identical, with $K'_{n,0,\rho}$ replacing $R'_{n,0,\rho}$ and $\mathcal{E}_{x,y}$ replacing $\mathcal{T}_{x,y}$ throughout. \square

The rest of the proof of Proposition 6.1 is devoted to estimating the double integral at (6.2). We deal separately with the integrals over pairs (x, y) satisfying (i) $\|y - x\| > r_n$; (ii) $\rho r_n < \|y - x\| \leq r_n$, and (iii) $\|y - x\| \leq \rho r_n$. Let δ_1 be as in Lemma 3.1.

Lemma 6.3. *Suppose $0 < \rho < \min((f_0 \delta_1 / (2\theta f_{\max}))^{1/d}, 1)$. Then as $n \rightarrow \infty$ we have*

$$n^2 \int_A \int_{A \cap B_{4r_n}(x) \setminus B_{r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \nu(dy) \nu(dx) = O(I_n \exp(-(\delta_1 f_0 / 3) n r_n^d)). \quad (6.4)$$

Proof. Since $\mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \mathbf{1}\{r_n < \|y - x\| \leq 4r_n\}$ is symmetric in x and y it suffices to prove the estimate for the integral restricted to $(x, y) \in A \times A$ with $x \prec y$, i.e. $y \in A_x$. For such (x, y) , if $\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}$ occurs, then $\mathcal{P}_n \cap (B_{r_n}(x) \cup B_{r_n}(y)) \setminus (B_{\rho r_n}(x) \cup B_{\rho r_n}(y)) = \emptyset$. Hence

$$\begin{aligned} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] &\leq \exp(-n\nu[(B_{r_n}(x) \cup B_{r_n}(y)) \setminus (B_{\rho r_n}(x) \cup B_{\rho r_n}(y))]) \\ &\leq \exp(-n\nu(B_{r_n}(x)) - n\nu(B_{r_n}(y) \setminus B_{r_n}(x)) + 2n\theta f_{\max}(\rho r_n)^d). \end{aligned}$$

By Lemma 3.1, if $\|x - y\| \geq r_n$ then $\nu(B_{r_n}(y) \setminus B_{r_n}(x)) \geq 2f_0\delta_1 r_n^d$. Therefore if we take ρ to be so small that $2\theta f_{\max}\rho^d < f_0\delta_1$, the third (positive) term in the exponent is less than half the second (negative) term. Hence $\mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \leq e^{-n\nu(B_{r_n}(x)) - \delta_1 f_0 n r_n^d}$. It follows that

$$n^2 \int_A \int_{A_x \cap B_{4r_n}(x) \setminus B_{r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \nu(dy) \nu(dx) \leq n I_n \theta f_{\max} (4r_n)^d e^{-\delta_1 f_0 n r_n^d},$$

and (6.4) follows. \square

Lemma 6.4. *Let $x, y \in A$ with $\|x - y\| \in (\rho r_n, r_n]$. Then $\mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] = 0$.*

Proof. The condition on $\|x - y\|$ implies that $y \in \mathcal{C}_{r_n}(x, \mathcal{P}_n^y)$ and $\text{diam}(\mathcal{C}_{r_n}(x, \mathcal{P}_n^y)) > \rho r_n$, which negates the event $\mathcal{T}_{x,y}$. \square

Lemma 6.5. *Suppose $0 < \rho < \min((\delta_1 f_0 / (\theta f_{\max}))^{1/(d-1)}, 1)$. Then as $n \rightarrow \infty$ we have*

$$n^2 \int_A \int_{A \cap B_{\rho r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \nu(dy) \nu(dx) = O((nr_n^d)^{1-d} I_n). \quad (6.5)$$

Proof. Let $x, y \in A$ with $\|x - y\| \in (0, \rho r_n]$. Then $\mathcal{T}_{x,y} = \mathcal{T}_{y,x}$. Define event

$$\mathcal{N}_{x,y} := \{\mathcal{P}_n((B_{r_n}(x) \cup B_{r_n}(y)) \setminus B_{\|y-x\|}(x)) = 0\}.$$

By assumption $f_{\max}\rho^{d-1}\theta < \delta_1 f_0$. If $x \prec y$, using Lemma 3.1 yields

$$\begin{aligned} \mathbb{P}[\mathcal{N}_{x,y}] &\leq \exp(-n\nu(B_{r_n}(x)) - 2n\delta_1 f_0 r_n^{d-1} \|y - x\| + n f_{\max} \theta \|y - x\|^d) \\ &\leq \exp(-n\nu(B_{r_n}(x)) - n\delta_1 f_0 r_n^{d-1} \|y - x\|). \end{aligned} \quad (6.6)$$

Similarly, if $y \prec x$ then

$$\mathbb{P}[\mathcal{N}_{x,y}] \leq \exp(-n\nu(B_{r_n}(y)) - n\delta_1 f_0 r_n^{d-1} \|x - y\|). \quad (6.7)$$

Hence, recalling $A_x := \{y \in A : x \prec y\}$ and using Fubini's theorem we obtain that

$$\begin{aligned} &n^2 \int_A \int_{A \cap B_{\rho r_n}(x)} \mathbb{P}[\mathcal{N}_{x,y}] \nu(dy) \nu(dx) \\ &\leq n^2 \int_A \int_{A_x \cap B_{\rho r_n}(x)} e^{-n\nu(B_{r_n}(x)) - n\delta_1 f_0 r_n^{d-1} \|y-x\|} \nu(dy) \nu(dx) \\ &\quad + n^2 \int_A \int_{A_y \cap B_{\rho r_n}(y)} e^{-n\nu(B_{r_n}(y)) - n\delta_1 f_0 r_n^{d-1} \|x-y\|} \nu(dx) \nu(dy) \\ &\leq 2n I_n f_{\max} \int_{B_{\rho r_n}(o)} e^{-n\delta_1 f_0 r_n^{d-1} \|u\|} du \\ &= 2n f_{\max} I_n (nr_n^d)^{-d} \int_{B_{n\rho r_n^d}(o)} e^{-n\delta_1 f_0 \|v\|} dv = O((nr_n^d)^{1-d} I_n). \end{aligned} \quad (6.8)$$

Next, let z denote the furthest point from x in $\mathcal{C}_{r_n}(x, \mathcal{P}_n^{x,y})$. If $z = y$ then $\mathcal{N}_{x,y}$ occurs. Thus if $\mathcal{T}_{x,y} \setminus \mathcal{N}_{x,y}$ occurs then $z \neq y$ and hence $z \in \mathcal{P}_n$ with $\|y - x\| < \|z - x\| \leq \rho r_n$, and moreover $\mathcal{P}_n \cap (B_{r_n}(x) \cup B_{r_n}(z)) \setminus B_{\|z-x\|}(x) = \emptyset$. That is,

$$\{\mathcal{T}_{x,y} \setminus \mathcal{N}_{x,y}\} \subset \{\exists z \in \mathcal{P}_n \cap B_{\rho r_n}(x) \setminus B_{\|y-x\|}(x) : \mathcal{P}_n((B_{r_n}(x) \cup B_{r_n}(z)) \setminus B_{\|z-x\|}(x)) = 0\}.$$

Hence by Markov's inequality, the Mecke formula and Fubini's theorem,

$$\begin{aligned}
& n^2 \int_A \int_{A \cap B_{\rho r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \setminus \mathcal{N}_{x,y}] \nu(dy) \nu(dx) \\
& \leq n^3 \int_A \int_A \int_{B_{\rho r_n}(x) \setminus B_{\|y-x\|}(x)} e^{-n\nu[(B_{r_n}(x) \cup B_{r_n}(z)) \setminus B_{\|z-x\|}(x)]} \nu(dz) \nu(dy) \nu(dx) \\
& \leq n^3 \int_A \int_{B_{\rho r_n}(x)} e^{-n\nu[(B_{r_n}(x) \cup B_{r_n}(z)) \setminus B_{\|z-x\|}(x)]} (f_{\max} \theta \|z-x\|^d) \nu(dz) \nu(dx).
\end{aligned}$$

By the same estimates as at (6.6) and (6.7) (now with z instead of y), the last expression is bounded by

$$\begin{aligned}
& n^3 f_{\max} \theta \int_A \int_{A_x \cap B_{\rho r_n}(x)} \|z-x\|^d e^{-n\nu(B_{r_n}(x)) - n\delta_1 f_0 r_n^{d-1} \|z-x\|} \nu(dz) \nu(dx) \\
& + n^3 f_{\max} \theta \int_A \int_{A_z \cap B_{\rho r_n}(z)} \|x-z\|^d e^{-n\nu(B_{r_n}(z)) - n\delta_1 f_0 r_n^{d-1} \|x-z\|} \nu(dx) \nu(dz) \\
& \leq 2n^2 f_{\max}^2 \theta I_n \int_{B_{\rho r_n}(o)} e^{-n\delta_1 f_0 r_n^{d-1} \|u\|} \|u\|^d du \\
& = O(n^2 (nr_n^{d-1})^{-2d} I_n) = O((nr_n^d)^{2-2d} I_n).
\end{aligned}$$

Combining this with (6.8) yields (6.5). \square

Proof of Proposition 6.1. Applying Lemmas 6.3, 6.4 and 6.5 we obtain that provided ρ is taken small enough, we have as $n \rightarrow \infty$ that

$$n^2 \int_A \int_{A \cap B_{4r_n}(x)} \mathbb{P}[\mathcal{T}_{x,y} \cap \mathcal{T}_{y,x}] \nu(dy) \nu(dx) = O((nr_n^d)^{1-d} I_n). \quad (6.9)$$

Hence by Lemma 6.2, we obtain that

$$(\text{Var}[R'_{n,0,\rho}] - \mathbb{E}[R'_{n,0,\rho}])_+ = O((nr_n^d)^{1-d} I_n). \quad (6.10)$$

Also, by Lemma 5.6, provided ρ is small enough we have $\mathbb{E}[R'_{n,0,\rho}] = O((nr_n^d)^{1-d} I_n)$. Combining this with (6.10) and using the nonnegativity of variance, we obtain the statement about $R'_{n,0,\rho}$ in (6.1).

Since $\mathcal{E}_{x,y} \subset \mathcal{T}_{x,y}$ we still have (6.9) with $\mathcal{T}_{x,y}$ replaced by $\mathcal{E}_{x,y}$. We can then derive the statement about $K'_{n,0,\rho}$ by a similar argument; instead of Lemma 5.6 we now use part of the proof of Proposition 5.5. \square

6.2 Variances for small components: binomial input

Next we consider for $G(\mathcal{X}_n, r(n))$ the number of small non-singleton components $K_{n,0,\rho}$ and the number of vertices in such components, $R_{n,0,\rho}$ (as defined at (5.2)), for suitably small (fixed) ρ .

While the asymptotic variance for small components in a Poisson sample was obtained above by computing the first two moments and exploiting the spatial independence of the Poisson process, we shall bound the variance for small components in a binomial sample by a very different argument, namely, the Efron-Stein inequality from Lemma 3.5. This does not work so well, in the sense that our bound does the job only in dimension $d \geq 3$.

Proposition 6.6 (Variance estimates for small non-singleton components: binomial input). *If $d \geq 3$ then there exists $\delta_5 > 0$ such that if $0 < \rho \leq \delta_5$ then $\text{Var}(K_{n,0,\rho}) = O((nr_n^d)^{2-d}I_n)$ as $n \rightarrow \infty$, and $\text{Var}(R_{n,0,\rho}) = O((nr_n^d)^{2-d}I_n)$ as $n \rightarrow \infty$.*

Proof. By the Efron-Stein inequality (3.7),

$$\begin{aligned} \text{Var}[R_{n,0,\rho}] &\leq n \int_A \mathbb{E}[(D_x R_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) \\ &= n \int_A \mathbb{E}[(D_x^+ R_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) + n \int_A \mathbb{E}[(D_x^- R_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx). \end{aligned}$$

Similarly

$$\text{Var}[K_{n,0,\rho}] \leq n \int_A \mathbb{E}[(D_x^+ K_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) + n \int_A \mathbb{E}[(D_x^- K_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx).$$

Moreover for all finite $\mathcal{X} \subset \mathbb{R}^d$ and $x \in \mathbb{R}^d \setminus \mathcal{X}$ we have $D_x^+ K_{n,0,\rho}(\mathcal{X}) \leq D_x^+ R_{n,0,\rho}(\mathcal{X})$ and $D_x^- K_{n,0,\rho}(\mathcal{X}) \leq D_x^- R_{n,0,\rho}(\mathcal{X})$. Therefore the result follows from the next two lemmas. \square

Lemma 6.7. *Let ρ be as in Lemma 5.6. Then as $n \rightarrow \infty$ we have*

$$n \int_A \mathbb{E}[(D_x^+ R_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) = O((nr_n^d)^{1-d}I_n). \quad (6.11)$$

Proof. Note $D_x^+ R_{n,0,\rho}(\mathcal{X}_{n-1})$ is non-zero only if $0 < \text{diam } \mathcal{C}_r(x, \mathcal{X}_{n-1}^x) \leq \rho r_n$, in which case $D_x^+ R_{n,0,\rho}(\mathcal{X}_{n-1})$ is either 1 (if $\#(\mathcal{X}_{n-1} \cap B_{\rho r_n}(x)) > 1$) or 2 (if $\#(\mathcal{X}_{n-1} \cap B_{\rho r_n}(x)) = 1$). Hence

$$\begin{aligned} n \int_A \mathbb{E}[(D_x^+ R_{n,0,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) &\leq 4n \int_A \mathbb{P}[0 < \text{diam } \mathcal{C}_{r_n}(x, \mathcal{X}_{n-1}^x) \leq \rho r_n] \nu(dx) \\ &= 4\mathbb{E}[R_{n,0,\rho}(\mathcal{X}_n)] \end{aligned}$$

Then the result follows from Lemma 5.6. \square

Lemma 6.8. *Let δ_1 be as in Lemma 3.1 and suppose $0 < \rho < \min((\delta_1 f_0 / (2f_{\max} \theta))^{1/(d-1)}, 1)$. Then as $n \rightarrow \infty$, we have*

$$n \int_A \mathbb{E}[D_x^- R_{n,0,\rho}(\mathcal{X}_{n-1})] \nu(dx) = O((nr_n^d)^{2-d}I_n). \quad (6.12)$$

Proof. For $x \in A$, observe that $D_x^- R_{n,0,\rho}(\mathcal{X}_{n-1})$ is bounded above by $N_{1,x}$, where $N_{1,x}$ denotes the number of vertices $y \in \mathcal{X}_{n-1}$ such that $\|y-x\| \leq 2r_n$ and $0 < \text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \leq \rho r_n$. Therefore

$$(D_x^- R_{n,0,\rho}(\mathcal{X}_{n-1}))^2 \leq N_{1,x}^2 = N_{1,x} + N_{1,x}(N_{1,x} - 1). \quad (6.13)$$

Let $N_{2,x}$ be the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that $0 < \text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \leq \rho r_n$, and $y \prec u$ for all $u \in \mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}) \setminus \{y\}$, and z is the point in $\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1})$ furthest from y . Let $N_{3,x}$ be the number of ordered triples (z, u, y) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that $0 < \text{diam}(\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1})) \leq \rho r_n$, and $z \prec v$

for all $v \in \mathcal{C}_{r_n}(z, \mathcal{X}_{n-1}) \setminus \{z\}$, and u is the point of $\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1})$ furthest from z , and y is another point of $\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1})$.

Then $N_{1,x} \leq 2N_{2,x} + N_{3,x}$. For n large we have

$$\begin{aligned} \mathbb{E}[N_{2,x}] &\leq n^2 \int_{A \cap B_{2r_n}(x)} \int_{A_y \cap B_{\rho r_n}(y)} (1 - \nu[(B_{r_n}(y) \cup B_{r_n}(z)) \setminus B_{\|z-y\|}(y)])^{n-3} \nu(dz) \nu(dy) \\ &\leq 2n^2 \int_{A \cap B_{2r_n}(x)} \int_{A_y \cap B_{\rho r_n}(y)} e^{-n\nu(B_{r_n}(y)) - \delta_1 f_0 n r_n^{d-1} \|z-y\|} \nu(dz) \nu(dy). \end{aligned}$$

Therefore using Fubini's theorem we obtain that

$$\begin{aligned} n \int_A \mathbb{E}[N_{2,x}] \nu(dx) &\leq 2n^3 \int_A e^{-n\nu(B_{r_n}(y))} \int_{A_y \cap B_{\rho r_n}(y)} e^{-\delta_1 f_0 n r_n^{d-1} \|z-y\|} \int_{B_{2r_n}(y)} \nu(dx) \nu(dz) \nu(dy) \\ &\leq 2^{d+1} \theta f_{\max}^2 n^2 r_n^d I_n \int_{\mathbb{R}^d} e^{-\delta_1 f_0 n r_n^{d-1} \|u\|} du \\ &= O((nr_n^d)^{2-d} I_n). \end{aligned} \tag{6.14}$$

Next, we have that for n large

$$\begin{aligned} \mathbb{E}[N_{3,x}] &\leq n^3 \int_{B_{2r_n}(x)} \int_{B_{\rho r_n}(z) \cap A_z} \int_{B_{\|u-z\|}(z)} (1 - \nu[(B_{r_n}(z) \cup B_{r_n}(u)) \setminus B_{\|u-z\|}(z)])^{n-4} \\ &\quad \nu(dy) \nu(du) \nu(dz) \\ &\leq 2\theta f_{\max} n^3 \int_{B_{2r_n}(x)} \int_{B_{\rho r_n}(z) \cap A_z} \|u-z\|^d e^{-n\nu(B_{r_n}(z)) - \delta_1 f_0 n r_n^{d-1} \|u-z\|} \nu(du) \nu(dz). \end{aligned}$$

Then using Fubini's theorem and a change of variable $v = u - z$ we obtain that

$$\begin{aligned} n \int_A \mathbb{E}[N_{3,x}] \nu(dx) &\leq 2^{d+1} \theta^2 f_{\max}^2 n^4 r_n^d \int_A e^{-n\nu(B_{r_n}(z))} \nu(dz) \int_{\mathbb{R}^d} e^{-\delta_1 f_0 n r_n^{d-1} \|v\|} \|v\|^d dv \\ &= O(n^3 r_n^d I_n (nr_n^{d-1})^{-2d}) = O((nr_n^d)^{3-2d} I_n). \end{aligned} \tag{6.15}$$

Combined with (6.14) this shows that

$$n \int_A \mathbb{E}[N_{1,x}] \nu(dx) = O((nr_n^d)^{2-d} I_n). \tag{6.16}$$

Next consider $N_{1,x}(N_{1,x} - 1)$, which equals the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that both $\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})$ and $\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1})$ have Euclidean diameter in the range $(0, \rho r_n]$. For such (y, z) we cannot have $\rho r_n < \|y - z\| \leq r_n$; we distinguish between the cases where $\|y - z\| \leq \rho r_n$ and where $\|y - z\| > r_n$.

Let $N_{4,x}$ be the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that $\|y - z\| \leq \rho r_n$ and $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \leq \rho r_n$.

Let $N_{5,x}$ be the number of ordered quadruples (u, v, y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that $u \prec w$ for all $w \in \mathcal{C}_{r_n}(u, \mathcal{X}_{n-1})$, and v is the furthest point from u in $\mathcal{C}_{r_n}(u, \mathcal{X}_{n-1})$ and y, z are two further points in $\mathcal{C}_{r_n}(u, \mathcal{X}_{n-1})$ and $\text{diam}(\mathcal{C}_{r_n}(u, \mathcal{X}_{n-1})) \leq \rho r_n$. Then

$$N_{4,x} \leq 2N_{2,x} + 4N_{3,x} + N_{5,x}.$$

For $n \geq 4$ we have that

$$\begin{aligned} \mathbb{E}[N_{5,x}] &\leq n^4 \int_{A \cap B_{2r_n}(x)} \int_{A_u \cap B_{\rho r_n}(u)} (\nu(B_{\|v-u\|}(u)))^2 \\ &\quad \times (1 - \nu[(B_{r_n}(u) \cup B_{r_n}(v)) \setminus B_{\|v-u\|}(u)])^{n-4} \nu(dv) \nu(du), \end{aligned}$$

and hence by Fubini's theorem, for n large

$$\begin{aligned} n \int_A \mathbb{E}[N_{5,x}] \nu(dx) &\leq \theta^3 2^{1+d} f_{\max}^3 n^5 r_n^d \int_A e^{-n\nu(B_{r_n}(u))} \nu(du) \int_{\mathbb{R}^d} e^{-f_0 \delta_1 n r_n^{d-1} \|w\|} \|w\|^{2d} dw \\ &= O(n^4 r_n^d I_n (n r_n^{d-1})^{-3d}) = O((n r_n^d)^{4-3d} I_n). \end{aligned}$$

Combined with (6.14) and (6.15) this shows that

$$n \int_A \mathbb{E}[N_{4,x}] \nu(dx) = O((n r_n^d)^{2-d} I_n). \quad (6.17)$$

Let $N_{6,x}$ be the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{2r_n}(x)$ such that $\|y - z\| > r_n$, $y \prec z$ and both $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}))$ and $\text{diam}(\mathcal{C}_{r_n}(z, \mathcal{X}_{n-1}))$ lie in the range $(0, \rho r_n]$. Then $N_{1,x}(N_{1,x} - 1) = N_{4,x} + N_{6,x}$ and

$$\begin{aligned} \mathbb{E}[N_{6,x}] &\leq n^2 \int_{A \cap B_{2r_n}(x)} \int_{A_y \cap B_{2r_n}(x) \setminus B_{r_n}(y)} (1 - \nu[(B_{r_n}(y) \cup B_{r_n}(z)) \setminus (B_{\rho r_n}(y) \cup B_{\rho r_n}(z))])^{n-3} \\ &\quad \nu(dz) \nu(dy). \end{aligned}$$

By our choice of ρ we have $2f_{\max}\theta\rho^d \leq \delta_1 f_0$. Then by Lemma 3.1, for n large and $y \in A$, $z \in A_y$ with $\|z - y\| > r_n$,

$$\begin{aligned} \nu[(B_{r_n}(y) \cup B_{r_n}(z)) \setminus (B_{\rho r_n}(y) \cup B_{\rho r_n}(z))] &\geq \nu(B_{r_n}(y)) + 2\delta_1 f_0 r_n^d - 2f_{\max}\theta(\rho r_n)^d \\ &\geq \nu(B_{r_n}(y)) + \delta_1 f_0 r_n^d. \end{aligned}$$

Hence for n large,

$$\mathbb{E}[N_{6,x}] \leq 2n^2 \int_{A \cap B_{2r_n}(x)} e^{-n\nu(B_{r_n}(y)) - \delta_1 f_0 n r_n^d} f_{\max} \theta (2r_n)^d \nu(dy),$$

so by Fubini's theorem, for n large

$$\begin{aligned} n \int_A \mathbb{E}[N_{6,x}] \nu(dx) &\leq 2^{1+2d} f_{\max}^2 \theta^2 n^3 r_n^{2d} \int_A e^{-n\nu(B_{r_n}(y)) - \delta_1 f_0 n r_n^d} \nu(dy) \\ &= O((n r_n^d)^2 e^{-\delta_1 f_0 n r_n^d} I_n) = O(e^{-(\delta_1 f_0/2) n r_n^d} I_n). \end{aligned}$$

Combined with (6.17) this shows that

$$n \int_A \mathbb{E}[N_{1,x}(N_{1,x} - 1)] \nu(dx) = O((n r_n^d)^{2-d} I_n)$$

and combined with (6.16) and (6.13) this gives us (6.12). \square

6.3 Variance estimates for medium components

We now consider the ‘medium-size’ component count, denoted $K_{n,\varepsilon,\rho}$ or $K'_{n,\varepsilon,\rho}$ (as defined at (5.2)) with $0 < \varepsilon < \rho < \infty$. We also consider the number of vertices in medium-sized components, denoted $R_{n,\varepsilon,\rho}$ or $R'_{n,\varepsilon,\rho}$. We shall bound the variances of all four of these quantities using Lemma 3.5, i.e. using the Poincaré or Efron-Stein inequality.

Proposition 6.9 (Variance estimates for medium-sized components). *Let $0 < \varepsilon < 1 < \rho < \infty$. Let ξ_n stand for any of $R_{n,\varepsilon,\rho}$, $R'_{n,\varepsilon,\rho}$, $K_{n,\varepsilon,\rho}$ or $K'_{n,\varepsilon,\rho}$. Then there exists $c > 0$ such that $\text{Var}(\xi_n) = O(e^{-cnr_n^d} I_n)$ as $n \rightarrow \infty$.*

Proof. Note that $D^+ K_{n,\varepsilon,\rho}(\mathcal{X}) \leq D^+ R_{n,\varepsilon,\rho}(\mathcal{X})$ and $D^- K_{n,\varepsilon,\rho}(\mathcal{X}) \leq D^- R_{n,\varepsilon,\rho}(\mathcal{X})$. Analogously to the proof of Proposition 6.6, but using the Poincaré inequality instead of the Efron-Stein inequality in the case of the results for $R'_{n,\varepsilon,\rho}$ and $K'_{n,\varepsilon,\rho}$, we can obtain the result from the next two lemmas. \square

Lemma 6.10. *Let $0 < \varepsilon < 1 < \rho < \infty$. Then there exists $c > 0$ such that*

$$n \int_A \mathbb{E}[(D_x^- R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) = O(e^{-cnr_n^d} I_n); \quad (6.18)$$

$$n \int_A \mathbb{E}[(D_x^- R_{n,\varepsilon,\rho}(\mathcal{P}_n))^2] \nu(dx) = O(e^{-cnr_n^d} I_n). \quad (6.19)$$

Proof. Observe that $D_x^- R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1})$ is bounded above by the number of vertices $y \in \mathcal{X}_{n-1} \cap B_{2\rho r_n}(x)$ such that $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \in (\varepsilon r, \rho r_n]$. We denote this quantity by $N_{7,x}$.

Let $N_{8,x}$ be the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap B_{3\rho r_n}(x)$ such that $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \in (\varepsilon r, \rho r_n]$ and $y \prec u$ for all $u \in \mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}) \setminus \{y\}$. Then $N_{7,x} \leq 2N_{8,x}$.

Fix $\delta > 0$ small, as in the proof of [12, Lemma 3.4], and discretize \mathbb{R}^d into cubes of side δr_n as in that proof. Assume $4\delta d^2 < \min(\delta_2/\theta, 1)$, where $\delta_2 = \delta_2(\varepsilon, 2\rho)$ is as in Lemma 3.2 of the present paper. Then

$$\mathbb{E}[N_{8,x}] \leq n^2 \int_{B_{3\rho r_n}(x)} \int_{B_{3\rho r_n}(x)} \sum_{\sigma} (1 - \nu([\sigma \cap A_y] \oplus B_{r(1-\sqrt{d}\delta)}(o)) \setminus (\sigma \cap A_y))^{n-3} \nu(dz) \nu(dy),$$

where the sum is over a finite (and uniformly bounded) number of possible shapes σ that could arise as the union of those cubes in the discretization containing points of $\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})$.

Using Lemma 3.2, the continuity of f , and the bound $(1 - u)^d \geq 1 - du$, we have for n large that

$$\begin{aligned} & \nu([\sigma \cap A_y] \oplus B_{r(1-\sqrt{d}\delta)}(o)) \setminus (\sigma \cap A_y) \\ & \geq (1 - \delta) f(y) [\lambda(B_{r(1-\sqrt{d}\delta)}(y) \cap A) + 2\delta_2 (1 - \sqrt{d}\delta)^d r_n^d] \\ & \geq (1 - \delta) f(y) [\lambda(B_{r_n}(y) \cap A) - (1 - (1 - \sqrt{d}\delta)^d) \theta r_n^d + (3/2) \delta_2 r_n^d] \\ & \geq \nu(B_{r_n}(y)) - 2\delta f(y) \lambda(B_{r_n}(y) \cap A) + (5/4) (1 - \delta) f(y) \delta_2 r_n^d \\ & \geq \nu(B_{r_n}(y)) - 2\theta \delta f(y) r_n^d + (9/8) \delta_2 f(y) r_n^d \\ & \geq \nu(B_{r_n}(y)) + \delta_2 f_0 r_n^d, \end{aligned}$$

and thus there exists a constant $c' > 0$ such that for n large

$$\mathbb{E}[N_{8,x}] \leq c'n^2 \int_{B_{3\rho r_n}(x)} \int_{B_{3\rho r_n}(x)} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dz) \nu(dy). \quad (6.20)$$

Hence by Fubini's theorem there is a constant c'' such that for n large

$$\begin{aligned} n \int_A \mathbb{E}[N_{8,x}] \nu(dx) &\leq c'' n^3 r_n^{2d} \int_A e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy) \\ &= O((n r_n^d)^2 e^{-\delta_2 f_0 n r_n^d} I_n). \end{aligned}$$

Next, let $N_{9,x}$ denote the number of ordered triples (y, z, u) of distinct points of $\mathcal{X}_{n-1} \cap B_{3\rho r_n}(x)$ such that $\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1})) \in (\varepsilon r_n, \rho r_n]$ and $y \prec v$ for all $v \in \mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}) \setminus \{y\}$. Then

$$N_{7,x}(N_{7,x} - 1) \leq 2N_{8,x} + N_{9,x}.$$

Using Lemma 3.2 again, we can find a new constant $c' > 0$ such that for n large

$$\begin{aligned} \mathbb{E}[N_{9,x}] &\leq n^3 \int_{B_{3\rho r_n}(x)} \sum_{\sigma} (1 - \nu([\sigma \cap A_y] \oplus B_{(1-\sqrt{d}\delta)r_n}(o)) \setminus (\sigma \cap A_y))^{n-4} (\nu(B_{3\rho r_n}(x)))^2 \nu(dy) \\ &\leq c' n^3 r_n^{2d} \int_{B_{3\rho r_n}(x)} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy), \end{aligned}$$

and hence by Fubini's theorem there is a further new constant c'' such that

$$\begin{aligned} n \int_A \mathbb{E}[N_{9,x}] \nu(dx) &\leq c'' n^4 r_n^{3d} \int_A e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy) \\ &= O((n r_n^d)^3 e^{-\delta_2 f_0 n r_n^d} I_n). \end{aligned}$$

Combined with (6.20) this shows that

$$\begin{aligned} n \int_A \mathbb{E}[(D_x^- R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) &\leq n \int_A \mathbb{E}[N_{7,x} + N_{7,x}(N_{7,x} - 1)] \nu(dx) \\ &= O((n r_n^d)^3 e^{-\delta_2 f_0 n r_n^d} I_n), \end{aligned}$$

and (6.18) follows. The proof of (6.19) is similar, using the Mecke formula. \square

Lemma 6.11. *Let $0 < \varepsilon < 1 < \rho < \infty$. Then there exists $c > 0$ such that*

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) = O(e^{-c n r_n^d} I_n); \quad (6.21)$$

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\varepsilon,\rho}(\mathcal{P}_n))^2] \nu(dx) = O(e^{-c n r_n^d} I_n); \quad (6.22)$$

Proof. Let δ_2 and δ be as in the previous proof. If $D_x^+ R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}) > 0$ then $\text{diam}(\mathcal{C}_{r_n}(x, \mathcal{X}_{n-1}^x)) \in (\varepsilon r_n, \rho r_n]$. We discretise \mathbb{R}^d into cubes of side δr_n as before. For each possible shape σ (i.e., a union of cubes of side δr_n), let $E_{x,\sigma}$ be the event that σ is the shape induced by $\mathcal{C}_{r_n}(x, \mathcal{X}_{n-1}^x)$, i.e. the union of those cubes in the discretization which contain at least one point of $\mathcal{C}_{r_n}(x, \mathcal{X}_{n-1}^x)$. Given $\mathcal{X}, D \subset \mathbb{R}^d$ with \mathcal{X} finite, let $\mathcal{X}(D) := \#(\mathcal{X} \cap D)$. Then

$$(D_x^+ R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}))^2 \leq \sum_{\sigma} \mathbf{1}_{E_{x,\sigma}} (1 + \mathcal{X}_{n-1}(\sigma))^2,$$

and hence

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\varepsilon,\rho}(\mathcal{X}_{n-1}))^2] \nu(dx) \leq n \int_A \sum_{\sigma: x \in \sigma} (\mathbb{P}[E_{x,\sigma}] + 2\mathbb{E}[\mathcal{X}_{n-1}(\sigma) \mathbf{1}_{E_{x,\sigma}}] + \mathbb{E}[\mathcal{X}_{n-1}(\sigma)^2 \mathbf{1}_{E_{x,\sigma}}]) \nu(dx). \quad (6.23)$$

If $E_{x,\sigma}$ occurs there is a point y of $\mathcal{X}_{n-1} \cap \sigma$ with $y \prec z$ for all $z \in \mathcal{X}_{n-1} \cap \sigma \setminus \{y\}$, so using Lemma 3.2 as in the preceding proof, we obtain for n large that

$$\begin{aligned} \mathbb{P}[E_{x,\sigma}] &\leq (n-1) \int_{\sigma} (1 - \nu([\sigma \cap A_y] \oplus B_{(1-\sqrt{d}\delta)r_n}(o)] \setminus (\sigma \cap A_y))^{n-2} \nu(dy) \\ &\leq 2n \int_{\sigma} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy), \end{aligned}$$

and hence by Fubini's theorem there exist constants c', c'' such that

$$\begin{aligned} n \int_A \sum_{\sigma: x \in \sigma} \mathbb{P}[E_{x,\sigma}] \nu(dx) &\leq 2n^2 \int_A \sum_{\sigma: x \in \sigma} \int_{\sigma} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy) \nu(dx) \\ &= 2n^2 \int_A \sum_{\sigma: y \in \sigma} \int_{\sigma} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dx) \nu(dy) \\ &\leq c' n^2 r_n^d \int_A \sum_{\sigma: y \in \sigma} e^{-n\nu(B_{r_n}(y)) - \delta_2 f_0 n r_n^d} \nu(dy) \\ &\leq c'' n r_n^d I_n e^{-\delta_2 f_0 n r_n^d}, \end{aligned} \quad (6.24)$$

where for the third line we used the fact that $\lambda(\sigma)$ is bounded by a constant times r_n^d , and in the fourth line we used the fact that there are a bounded number of shapes σ that contain y and are consistent with the diameter condition.

Next, let $N_1(\sigma)$ denote the number of ordered pairs (y, z) of distinct points of $\mathcal{X}_{n-1} \cap \sigma$ such that $y \prec u$ for all points of $\mathcal{X}_{n-1} \cap \sigma \setminus \{y\}$. Then $\mathcal{X}_{n-1}(\sigma) \leq 1 + N_1(\sigma)$. Therefore

$$\begin{aligned} \mathbb{E}[(\mathcal{X}_{n-1}(\sigma) - 1) \mathbf{1}_{E_{x,\sigma}}] &\leq \mathbb{E}[N_1(\sigma) \mathbf{1}_{E_{x,\sigma}}] \\ &\leq n(n-1) \int_{\sigma} \int_{\sigma} (1 - \nu([\sigma \cap A_y] \oplus B_{(1-\sqrt{d}\delta)r_n}(o)] \setminus (\sigma \cap A_y))^{n-3} \nu(dz) \nu(dy). \end{aligned}$$

The z -integral is bounded by a constant times r_n^d , and by a similar application of Fubini's theorem to the one at (6.24) we obtain that

$$n \int_A \sum_{\sigma: x \in \sigma} \mathbb{E}[(\mathcal{X}_{n-1}(\sigma) - 1) \mathbf{1}_{E_{x,\sigma}}] \nu(dx) = O((n r_n^d)^2 e^{-\delta_2 f_0 n r_n^d} I_n). \quad (6.25)$$

Next, let $N_2(\sigma)$ denote the number of ordered triples (y, z, u) of distinct points of $\mathcal{X}_{n-1} \cap \sigma$ such that $y \prec v$ for all $v \in \mathcal{X}_{n-1} \cap \sigma \setminus \{y\}$.

Then provided $\mathcal{X}_{n-1}(\sigma) \neq 0$, $(\mathcal{X}_{n-1}(\sigma) - 1)(\mathcal{X}_{n-1}(\sigma) - 2)$ is the number of ordered pairs of vertices of $\mathcal{X}_{n-1} \cap \sigma$, other than the first one in the \prec order, and equals $N_2(\sigma)$. If $E_{x,\sigma}$ occurs the $\mathcal{X}_{n-1}(\sigma) \neq 0$. Hence

$$\begin{aligned} \mathbb{E}[(\mathcal{X}_{n-1}(\sigma) - 1)(\mathcal{X}_{n-1}(\sigma) - 2) \mathbf{1}_{E_{x,\sigma}}] &= \mathbb{E}[N_2(x, \sigma) \mathbf{1}_{E_{x,\sigma}}] \\ &\leq n^3 \int_{\sigma} \int_{\sigma} \int_{\sigma} (1 - \nu([\sigma \cap A_y] \oplus B_{(1-\sqrt{d}\delta)r_n}(o)] \setminus (\sigma \cap A_y))^{n-4} \nu(du) \nu(dz) \nu(dy). \end{aligned}$$

The (z, u) -integral is bounded by a constant times r_n^{2d} , and by a similar application of Fubini's theorem to the one at (6.24) we obtain that

$$n \int_A \sum_{\sigma: x \in \sigma} \mathbb{E}[(\mathcal{X}_{n-1}(\sigma) - 1)(\mathcal{X}_{n-1}(\sigma) - 2)\mathbf{1}_{E_{x,\sigma}}] = O((nr_n^d)^3 I_n e^{-\delta_2 f_0 nr_n^d}).$$

Combining this with (6.23), (6.24) and (6.25) we obtain (6.21).

The proof of (6.22) is similar, using the Mecke formula. \square

6.4 Variance estimates for large components

Proposition 6.12 (Variance estimates for moderately large components). *There exists $\rho \in (4, \infty)$ such that if ξ_n stands for any of $R_{n,\rho,(\log n)^2}$, $R'_{n,\rho,(\log n)^2}$, $K_{n,\rho,(\log n)^2}$, or $K'_{n,\rho,(\log n)^2}$, then $\text{Var}(\xi_n) = O(e^{-nr_n^d} I_n)$ as $n \rightarrow \infty$.*

Proof. Analogously to Proposition 6.9 the result follows from the next two lemmas. \square

Lemma 6.13. *There exists $\rho_0 > 1$ such that for any fixed $\rho \geq \rho_0$ we have as $n \rightarrow \infty$ that*

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\rho,(\log n)^2}(\mathcal{X}_{n-1}))^2] \nu(dx) = O(e^{-nr_n^d} I_n); \quad (6.26)$$

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\rho,(\log n)^2}(\mathcal{P}_n))^2] \nu(dx) = O(e^{-nr_n^d} I_n). \quad (6.27)$$

Proof. Let $\rho > 4$. For $y \in \mathcal{X}_{n-1}$, adding a point at x can only increase the diameter of the component containing y . Therefore if adding a point at x causes y to be in a component of diameter in the range $(\rho r_n, (\log n)^2 r_n]$ when it was not before, then y must previously have been in a component of diameter at most ρr_n , and since also the added point at x affects this component we must have $\|y - x\| \leq (\rho + 1)r_n \leq 2\rho r_n$. Also event $\mathcal{M}_{n,\rho/4,(\log n)^2}^*(x, \mathcal{X}_{n-1})$, defined at (3.15), must occur. Therefore defining $N_x := \#(\mathcal{X}_{n-1} \cap B_{2\rho r_n}(x))$, we have $D_x^+ R_{n,\rho,(\log n)^2}(\mathcal{X}_{n-1}) \leq N_x \mathbf{1}_{\mathcal{M}_{n,\rho/4,(\log n)^2}^*(x, \mathcal{X}_{n-1})}$. Hence by the Cauchy-Schwarz inequality, Lemma 3.8 and a standard moment estimate on the Binomial distribution,

$$\begin{aligned} \mathbb{E}[(D_x^+ R_{n,\rho,(\log n)^2}(\mathcal{X}_{n-1}))^2] &\leq (\mathbb{E}[N_x^4])^{1/2} (\mathbb{P}[\mathcal{M}_{n,\rho/4,(\log n)^2}^*(x, \mathcal{X}_{n-1})])^{1/2} \\ &= O(n^2 r_n^{2d} \exp(-(\beta\rho/8)nr_n^d)), \end{aligned}$$

where β is as in Lemma 3.8. Choosing ρ so that $\beta\rho > 8(\theta f_0 + 3)$, and using Lemma 4.1, we obtain that

$$n \int_A \mathbb{E}[(D_x^+ R_{n,\rho,(\log n)^2}(\mathcal{X}_{n-1}))^2] \nu(dx) = O(ne^{-(\theta f_0 + 2)nr_n^d}) = O(e^{-nr_n^d} I_n),$$

as required for (6.26). The proof of (6.27) is similar. \square

Lemma 6.14. *There exists $\rho_0 > 1$ such that if $\rho \geq \rho_0$ then as $n \rightarrow \infty$,*

$$n \int_A \mathbb{E}[(D_x^- R_{n,\rho,(\log n)^2}(\mathcal{X}_{n-1}))^2] \nu(dx) = O(e^{-nr_n^d} I_n); \quad (6.28)$$

$$n \int_A \mathbb{E}[(D_x^- R_{n,\rho,(\log n)^2}(\mathcal{P}_n))^2] \nu(dx) = O(e^{-nr_n^d} I_n). \quad (6.29)$$

Proof. Let $\rho > 1$. For this proof, given n and given $x \in A$ let N_x denote the number of vertices $y \in \mathcal{X}_{n-1} \cap B_{2(\log n)^2 r_n}(x)$ such that $\mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}) \cap B_{r_n}(x) \neq \emptyset$ and $\text{diam} \mathcal{C}_{r_n}(y, \mathcal{X}_{n-1}) \in (\rho r_n, (\log n)^2 r_n]$. Then $D_x^- R_{n, \rho, (\log n)^2}(\mathcal{X}_{n-1}) \leq N_x$.

We have that $\mathbb{E}[N_x] \leq J_{1,x} + J_{2,x}$, where we set

$$J_{1,x} := \int_{B_{3\rho r_n}(x)} n \mathbb{P}[\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-2}^y)) \in (\rho r_n, (\log n)^2 r_n)] \nu(dy)$$

$$J_{2,x} := \int_{B_{2(\log n)^2 r_n}(x) \setminus B_{3\rho r_n}(x)} n \mathbb{P}[\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-2}^y)) \in (\|y - x\|/2, (\log n)^2 r_n)] \nu(dy).$$

Let β be as in Lemma 3.8. By that result,

$$J_{1,x} \leq n f_{\max} \theta (3\rho r_n)^d \exp(-\beta \rho n r_n^d). \quad (6.30)$$

Also by Lemma 3.8,

$$\begin{aligned} J_{2,x} &\leq n \int_{A \setminus B_{3\rho r_n}(x)} \exp(-\beta(\|y - x\|/2) n r_n^{d-1}) \nu(dy) \\ &\leq n f_{\max} \int_{\mathbb{R}^d \setminus B_{3\rho r_n}(o)} \exp(-\beta(\|u\|/2) n r_n^{d-1}) du \\ &= n f_{\max} \int_{\{v: \|v\| > 3\rho r_n(\beta/2) n r_n^{d-1}\}} e^{-\|v\|} (\beta n r_n^{d-1}/2)^{-d} dv \\ &= (2/\beta)^d f_{\max} (n r_n^d)^{1-d} \int_{\rho \beta n r_n^d}^{\infty} e^{-t} d\theta_d t^{d-1} dt \\ &\leq c \beta^{-1} f_{\max} \rho^{d-1} e^{-\rho \beta n r_n^d}, \end{aligned}$$

where the constant c depends only on d . Combined with (6.30), this shows that if we take $\rho \geq (\theta f_0 + 3)/\beta$ then for n large $\mathbb{E}[N_x] \leq \exp(-(\theta f_0 + 2) n r_n^d)$ for all $x \in A$, and then using Lemma 4.1 we obtain that

$$n \int_A \mathbb{E}[N_x] \nu(dx) = O(e^{-n r_n^d} I_n). \quad (6.31)$$

Next, observe that $\mathbb{E}[N_x(N_x - 1)] \leq J_{3,x} + 2J_{4,x}$ where we set

$$J_{3,x} := \int_{B_{3\rho r_n}(x)} \int_{B_{3\rho r_n}(x)} n^2 \mathbb{P}[\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-3}^{y,z})) \in (\rho r_n, (\log n)^2 r_n)] \nu(dz) \nu(dy);$$

$$J_{4,x} := \int_{B_{2(\log n)^2 r_n}(x) \setminus B_{3\rho r_n}(x)} \int_{B_{\|y-x\|}(x)} n^2 \mathbb{P}[\text{diam}(\mathcal{C}_{r_n}(y, \mathcal{X}_{n-3}^{y,z})) \in (\|y - x\|/2, (\log n)^2 r_n)] \nu(dz) \nu(dy).$$

By Lemma 3.8,

$$J_{3,x} \leq n^2 (f_{\max} \theta (3\rho r_n)^d)^2 e^{-\beta \rho n r_n^d}. \quad (6.32)$$

Also by Lemma 3.8,

$$\begin{aligned}
J_{4,x} &\leq n^2 \int_{A \setminus B_{3\rho r_n}(x)} \exp(-\beta(\|y-x\|/2)nr_n^{d-1})(f_{\max}\theta\|y-x\|^d)\nu(dy) \\
&\leq n^2 f_{\max}^2 \theta \int_{\mathbb{R}^d \setminus B_{3\rho r_n}(o)} \exp(-\beta(\|u\|/2)nr_n^{d-1})\|u\|^d du \\
&= n^2 f_{\max}^2 \theta \int_{\{v:\|v\|>3\rho r_n(\beta/2)nr_n^{d-1}\}} e^{-\|v\|}\|v\|^d (\beta nr_n^{d-1}/2)^{-2d} dv \\
&= (2/\beta)^{2d} f_{\max}^2 (nr_n^d)^{2-2d} \int_{\rho\beta nr_n^d}^{\infty} e^{-t} d\theta_d t^{2d-1} dt \\
&\leq c\beta^{-1} f_{\max}^2 \rho^{2d-1} nr_n^d e^{-\rho\beta nr_n^d},
\end{aligned}$$

where the constant c depends only on d . Combined with (6.32), this shows that if we take $\rho \geq (\theta f_0 + 3)/\beta$ then for n large $\mathbb{E}[N_x(N_x - 1)] \leq \exp(-(\theta f_0 + 2)nr_n^d)$ for all $x \in A$, and then using Lemma 4.1 we obtain that

$$n \int_A \mathbb{E}[N_x(N_x - 1)]\nu(dx) = O(e^{-nr_n^d} I_n).$$

Combined with (6.31) this shows that (6.28) holds. The proof of (6.29) is similar. \square

6.5 Variance estimates: conclusion

Putting together the preceding estimates, we obtain the asymptotic variance for K'_n and (when $d \geq 3$) for K_n :

Proposition 6.15. *Assume that $nr_n^d \rightarrow \infty$ and $\liminf(I_n) > 0$ as $n \rightarrow \infty$. Then*

$$\text{Var}[K'_n] = I_n(1 + O((nr_n^d)^{(1-d)/2})); \quad (6.33)$$

$$\text{if } d \geq 3 \text{ then } \text{Var}[K_n] = I_n(1 + O((nr_n^d)^{1-d/2})). \quad (6.34)$$

Proof. Note $K'_n = S'_n + K'_{n,0,\infty}$, where S'_n and $K'_{n,\varepsilon,\rho}$ were defined at (4.3), (5.2).

Let $\rho \in (4, \infty)$ be as in Proposition 6.12. Let ρ_0 be as in Proposition 6.1. Let $\varepsilon = \rho_0$.

Let $W_n := K'_{n,(\log n)^2,\infty}$. Since $|W_n - 1|$ is bounded by $Z_n + 1$ (where $Z_n = \#(\mathcal{P}_n)$), the Cauchy-Schwarz inequality and Lemma 5.4 yield that

$$\begin{aligned}
\text{Var}[W_n] &= \text{Var}[W_n - 1] \leq \mathbb{E}[(W_n - 1)^2] \leq (\mathbb{E}[(Z_n + 1)^4])^{1/2} (\mathbb{P}[W_n \neq 1])^{1/2} \\
&= O(e^{-\frac{1}{2}nr_n^d} I_n).
\end{aligned} \quad (6.35)$$

Then $K'_{n,0,\infty} = K'_{n,0,\varepsilon} + K'_{n,\varepsilon,\rho} + K'_{n,\rho,(\log n)^2} + W_n$. By the estimate $(u + v + w + x)^2 \leq 4(u^2 + v^2 + w^2 + x^2)$ (a consequence of Jensen's inequality), Propositions 6.1, 6.9 and 6.12, along with (6.35),

$$\begin{aligned}
\text{Var}[K'_{n,0,\infty}] &\leq 4(\text{Var}[K'_{n,0,\varepsilon}] + \text{Var}[K'_{n,\varepsilon,\rho}] + \text{Var}[K'_{n,\rho,(\log n)^2}] + \text{Var}[W_n]) \\
&= O((nr_n^d)^{1-d} I_n).
\end{aligned} \quad (6.36)$$

By Proposition 4.4, $\mathbb{V}\text{ar}[S'_n] = I_n(1 + e^{-\Omega(nr_n^d)})$. Hence by the Cauchy-Schwarz inequality, $\text{Cov}(S'_n, K'_{n,0,\infty}) = O((nr_n^d)^{(1-d)/2}I_n)$, and thus

$$\mathbb{V}\text{ar}(K'_n) = \mathbb{V}\text{ar}(S'_n) + \mathbb{V}\text{ar}(K'_{n,0,\infty}) + 2\text{Cov}(S'_n, K'_{n,0,\infty}) = I_n + O((nr_n^d)^{(1-d)/2}I_n),$$

which is (6.33). The proof of (6.34) is similar, but now using Proposition 6.6 instead of Proposition 6.1, which accounts for the different power of nr_n^d in (6.34). \square

We can now also determine the asymptotic variance for R'_n and (if $d \geq 3$) for R_n .

Proposition 6.16. *Under assumptions (4.1) and (4.2), as $n \rightarrow \infty$ we have*

$$\mathbb{V}\text{ar}[R'_n] = I_n(1 + O((nr_n^d)^{(1-d)/2})); \quad (6.37)$$

$$\text{if } d \geq 3, \quad \mathbb{V}\text{ar}[R_n] = I_n(1 + O((nr_n^d)^{1-d/2})). \quad (6.38)$$

Proof. Let $0 < \varepsilon < \rho$ with $\varepsilon < \rho_0$ and ρ_0 as in Proposition 6.1. By Jensen's inequality and Propositions 6.1, 6.9 and 6.12,

$$\begin{aligned} \mathbb{V}\text{ar}[R'_{n,0,(\log n)^2}] &\leq 3(\mathbb{V}\text{ar}[R'_{n,0,\varepsilon}] + \mathbb{V}\text{ar}[R'_{n,\varepsilon,\rho}] + \mathbb{V}\text{ar}[R'_{n,\rho,(\log n)^2}]) \\ &= O((nr_n^d)^{1-d}I_n). \end{aligned} \quad (6.39)$$

Since $|R'_n - S'_n - R'_{n,0,(\log n)^2}| \leq Z_n$, by the Cauchy-Schwarz inequality and Lemma 5.4,

$$\begin{aligned} \mathbb{E}[|R'_n - S'_n - R'_{n,0,(\log n)^2}|^2] &\leq (\mathbb{E}[Z_n^2])^{1/2}(\mathbb{P}[R'_n \neq S'_n + R'_{n,0,(\log n)^2}])^{1/2} \\ &= O(e^{-nr_n^d/2}I_n). \end{aligned}$$

Then using (6.39) and Jensen's inequality again yields

$$\mathbb{V}\text{ar}[R'_n - S'_n] \leq 2(\mathbb{V}\text{ar}[R'_n - S'_n - R'_{n,0,(\log n)^2}] + \mathbb{V}\text{ar}[R'_{n,0,(\log n)^2}]) = O((nr_n^d)^{1-d}I_n). \quad (6.40)$$

By Proposition 4.4, $\mathbb{V}\text{ar}[S'_n] = I_n(1 + e^{-\Omega(nr_n^d)})$. Using this along with (6.40) and the Cauchy-Schwarz inequality gives us (6.37).

The proof of (6.38) is similar. We use Proposition 6.6 instead of Proposition 6.1, and Proposition 4.5 instead of Proposition 4.4. \square

6.6 Proof of convergence in distribution results

Proof of Theorem 2.5. By Proposition 6.15 we have $\mathbb{V}\text{ar}[K'_n] = I_n(1 + (nr_n^d)^{(1-d)/2})$. By Proposition 6.16 we have $\mathbb{V}\text{ar}[R'_n] = I_n(1 + (nr_n^d)^{(1-d)/2})$. Thus we have (2.8). If $d \geq 3$ then by Proposition 6.15 we have $\mathbb{V}\text{ar}[K_n] = I_n(1 + O((nr_n^d)^{1-d/2}))$, and by Proposition 6.16 we have $\mathbb{V}\text{ar}[R_n] = I_n(1 + O((nr_n^d)^{1-d/2}))$. Thus we have (2.10).

By (6.36) in the proof of Proposition 6.15 if $\xi'_n = K'_n - 1$, or (6.40) in the proof of Proposition 6.16 if $\xi'_n = R'_n$,

$$\mathbb{V}\text{ar}[\xi'_n - S'_n] = O((nr_n^d)^{1-d}I_n).$$

Hence $\text{Var}(I_n^{-1/2}(\xi'_n - S'_n - \mathbb{E}[\xi'_n - S'_n])) = O((nr_n^d)^{1-d})$. Hence by Lemma 3.6,

$$d_K(I_n^{-1/2}(\xi'_n - \mathbb{E}[\xi'_n]), N(0, 1)) = O(d_K(I_n^{-1/2}(S'_n - I_n), N(0, 1)) + (nr_n^d)^{(1-d)/3}),$$

and (2.9) then follows by (2.22).

When $d \geq 3$ we prove (2.11) similarly. In the binomial setting we get $(nr_n^d)^{2-d}$ instead of $(nr_n^d)^{1-d}$ in (6.36) or (6.40), and therefore $\text{Var}(I_n^{-1/2}(\xi_n - S_n - \mathbb{E}[\xi_n - S_n])) = O((nr_n^d)^{2-d})$. Therefore using Lemma 3.6 and (2.23) we have

$$\begin{aligned} d_K(\tilde{I}_n^{-1/2}(\xi_n - \mathbb{E}[\xi_n]), N(0, 1)) &= O(d_K(\tilde{I}_n^{-1/2}(S_n - \tilde{I}_n), N(0, 1)) + (nr_n^d)^{(2-d)/3}) \\ &= O((nr_n^d)^{(2-d)/3} + I_n^{-1/2}). \end{aligned}$$

Using the fact that $\tilde{I}_n = I_n(1 + O(e^{-c'nr_n^d}))$ for some further constant c' by Lemma 4.3, and using Lemma 3.6 again we obtain (2.11). \square

Proof of Theorem 2.8. We assume (1.1), (1.2) and that ν is uniform on A . For $n \geq 1$ define γ_n as at (1.2) and set $a_n := -\gamma_n$, so $a_n := (2 - 2/d)(\log n - \mathbf{1}\{d \geq 3\} \log \log n) - n\theta f_0 r_n^d$. By (1.2), $a_n \rightarrow \infty$ as $n \rightarrow \infty$. We claim $I_n \rightarrow \infty$. Indeed, if $d = 2$ then

$$ne^{-n\pi f_0 r^2} = ne^{a_n - \log n} \rightarrow \infty,$$

so that $I_n \rightarrow \infty$ by Proposition 4.7. If instead $d \geq 3$ then

$$e^{-n\theta f_0 r_n^d/2} r_n^{1-d} = e^{a_n/2} \left(\frac{\log n}{n}\right)^{1-1/d} r_n^{1-d} = e^{a_n/2} \left(\frac{nr_n^d}{\log n}\right)^{(1/d)-1}$$

which tends to infinity because, by (1.2), for n large we have $n\theta f_0 r_n^d \leq 2 \log n$. Therefore by Proposition 4.8, we have $I_n \rightarrow \infty$ in this case too, justifying our claim.

Suppose $d = 2$. By Proposition 4.7 and (1.4), as $n \rightarrow \infty$ we have $I_n = \mu_n(1 + O((nr_n^2)^{-1/2}))$ and then (2.16) follows from (2.8). Also by Lemma 3.6, (2.8) and (2.9),

$$\begin{aligned} d_K\left(\frac{\xi'_n - \mathbb{E}[\xi'_n]}{\mu_n^{1/2}}, N(0, 1)\right) &\leq d_K\left(\frac{\xi'_n - \mathbb{E}[\xi'_n]}{I_n^{1/2}}, N(0, 1)\right) + \left(\text{Var}((\mu_n^{-1/2} - I_n^{-1/2})(\xi'_n - \mathbb{E}[\xi'_n]))\right)^{1/3} \\ &= O((nr_n^2)^{-1/3} + I_n^{-1/2}) + O\left(\left(\left(\frac{I_n}{\mu_n}\right)^{1/2} - 1\right)^{2/3}\right) \\ &= O((nr_n^2)^{-1/3} + \mu_n^{-1/2}), \end{aligned} \tag{6.41}$$

and hence (2.17).

Now suppose $d \geq 3$. By Proposition 4.8, as $n \rightarrow \infty$ we have $I_n = \mu_n \left(1 + O\left(\left(\frac{\log(nr_n^d)}{nr_n^d}\right)^2\right)\right)$. Hence using (2.8) we have (2.18), and using (2.10) we have (2.19).

Also using Lemma 3.6, we can obtain (2.20) from (2.9) and (2.21) from (2.11), in both cases by similar steps to those used at (6.41) to derive (2.17). \square

References

- [1] Bobrowski, O. and Krioukov, D. (2022) Random Simplicial Complexes: Models and Phenomena. *Higher-Order Systems*, Eds F. Battiston and G. Petri, pp. 59–96. *Underst. Complex Syst.* Springer, Cham.
- [2] Boucheron, S., Lugosi, G. and Bousquet, O. (2004) Concentration Inequalities. *Machine Learning 2003* (Eds. O. Bousquet et al.), LNAI 3176, pp. 208–240. Springer, Berlin.
- [3] Ganesan, G. (2013) Size of the giant component in a random geometric graph. *Ann. Inst. Henri Poincaré Probab. Stat.* **49**, 1130–1140.
- [4] Higgs, F., Penrose, M. D. and Yang, X. (2024) Covering one point process with another. arXiv:2401.03832
- [5] Last, G. and Penrose, M. (2018) *Lectures on the Poisson Process*. Cambridge University Press, Cambridge.
- [6] Meester, R. and Roy, R. (1996) *Continuum Percolation*. Cambridge University Press, Cambridge.
- [7] Penrose, M. (2003) *Random Geometric Graphs*. Oxford University Press, Oxford.
- [8] Penrose, M. D. (1999) A strong law for the longest edge of the minimal spanning tree. *Ann. Probab.* **27**, 246–260.
- [9] Penrose, M. D. and Yukich, J. E. (2003) Weak laws of large numbers in geometric probability. *Ann. Appl. Probab.* **13**, 277–303.
- [10] Penrose, M. D. (2023) Random Euclidean coverage from within. *Probab. Theory Related Fields* **185**, 747–814.
- [11] Penrose, M. D. and Yang, X. (2024) Fluctuations of the connectivity threshold. arXiv:2406.00647v3.
- [12] Penrose, M. D. and Yang, X. (2023) On k -clusters of high-intensity random geometric graphs. arXiv:2209.14758v3