

# ARTIN-SCHELTER GORENSTEIN PROPERTY OF HOPF GALOIS EXTENSIONS

RUIPENG ZHU

**ABSTRACT.** This paper investigates the homological properties of the faithfully flat Hopf Galois extension  $A \subseteq B$ . It establishes that when  $B$  is a noetherian affine PI algebra and  $A$  is AS Gorenstein,  $B$  inherits the AS Gorenstein property. Furthermore, we demonstrate that injective dimension serves as a monoidal invariant for AS Gorenstein Hopf algebras. Specifically, if two such Hopf algebras have equivalent monoidal categories of comodules, then their injective dimensions are equal.

## INTRODUCTION

Over two decades ago, Brown and Goodearl embarked on exploring ring-theoretic attributes of infinite-dimensional Hopf algebras, with their findings detailed in the literature [3, 4]. Their investigation revealed that numerous examples of noetherian affine Hopf algebras possess the Gorenstein property. Furthermore, they demonstrated that noetherian affine PI, Gorenstein Hopf algebras exhibit various favorable homological characteristics.

In [3], Brown posed a series of inquiries pertaining to general noetherian (PI) Hopf algebras, among which is the following noteworthy question [3, Question A]:

**Conjecture 0.1** (Brown-Goodearl conjecture). Is every noetherian Hopf algebra AS Gorenstein?

Since then, extensive research has explored the homological characteristics associated with the AS Gorenstein property, including the existence of rigid dualizing complexes for infinite-dimensional noetherian Hopf algebras. More recently, the Brown-Goodearl conjecture has been explored in the context of other classes of noetherian algebras that exhibit similarities to Hopf algebras, such as weak Hopf algebras, as reported in [21]. In this paper, we delve into the Brown-Goodearl question specifically for faithfully flat Hopf Galois extensions, which can be regarded as noncommutative analogues of principal bundles.

Applying a result of Wu and Zhang [30], we have the following result concerning whenever faithfully flat Hopf Galois extension of an AS Gorenstein algebra is also AS Gorenstein, see Theorem 2.11.

**Theorem 0.2.** Let  $H$  be a Hopf algebra, and  $B$  be a noetherian affine PI algebra which is a faithfully flat  $H$ -Galois extension of  $A := B^{coH}$ . If  $A$  is AS Gorenstein, then  $B$  is as well. Additionally, if  $H$  is AS Gorenstein of dimension  $d_H$ , then  $\text{injdim}(B_B) = \text{injdim}(A_A) + d_H$ .

In the subsequent discussion, we delve into a specific finiteness condition that offers valuable insights for group algebras.

Let  $R$  be a ring. An  $R$ -module  $M$  is said to be of type  $\mathbf{FP}_n$  ( $n \geq 0$ ) if there is a projective resolution

$$\cdots \longrightarrow P_{n+1} \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0,$$

---

2020 *Mathematics Subject Classification.* 16E65, 16E10, 16T05.

*Key words and phrases.* Artin-Schelter Gorenstein algebra, Hopf Galois extension, monoidal Morita-Takeuchi equivalence.

where  $P_0, P_1, \dots, P_n$  are finitely generated. Furthermore,  $M$  is said to be of type  $\mathbf{FP}_\infty$  if there is a projective resolution

$$\cdots \longrightarrow P_{n+1} \longrightarrow P_n \longrightarrow \cdots \longrightarrow P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0,$$

where each  $P_n$  is finitely generated for all integers  $n \geq 0$ . An  $R$ -module  $M$  of type  $\mathbf{FP}_\infty$  is said to be of type  $\mathbf{FP}$ , if  $M$  has finite projective dimension.

Let  $H$  be a Hopf algebra over a field  $\mathbb{k}$ . Then  $\mathbb{k}$  is a left  $H$ -module induced by the counit map  $\varepsilon : H \rightarrow \mathbb{k}$ , which is called the trivial left module of  $H$ . We will say that  $H$  is of type  $\mathbf{FP}_n$  ( $n \in \mathbb{N} \cup \{\infty, \emptyset\}$ ) if the trivial left module  $\mathbb{k}$  is of type  $\mathbf{FP}_n$  as an  $H$ -module. Evidently, noetherian Hopf algebras are always of type  $\mathbf{FP}_\infty$ . However, it is worth mentioning that there are numerous non-noetherian group algebras that also belong to the type  $\mathbf{FP}_\infty$ , as explored in Chapter VIII of [6].

In [22], Schauenburg has shown the equivalence of the following assertions:

- (1)  $L$  and  $H$  are *monoidally Morita-Takeuchi equivalent*, that is, their comodule categories are monoidally equivalent;
- (2) there is a  $L$ - $H$ -biGalois extension of  $\mathbb{k}$ .

Recently, several homological properties, including global dimension and skew Calabi-Yau property, have garnered significant attention and have been extensively studied in the context of monoidal Morita-Takeuchi equivalence, see [1, 2, 29]. This research area has produced many fascinating results and questions, some of which are outlined below.

**Theorem 0.3.** Let  $H$  and  $L$  be two monoidally Morita-Takeuchi equivalent Hopf algebras of type  $\mathbf{FP}$ .

- (1) [2, Theorem 8] Then they have the same global dimension, that is,  $\text{gldim}(H) = \text{gldim}(L)$ .
- (2) [29, Theorem 2] If  $H$  is AS regular of dimension  $d$ , so is  $L$ .

**Question 0.4.** Let  $H$  and  $L$  be two monoidally Morita-Takeuchi equivalent Hopf algebras.

- (1) [1, 2] Then  $\text{gldim}(H) = \text{gldim}(L)$ ?
- (2) [29, Question 3] Suppose that  $H$  is of type  $\mathbf{FP}$ . Is  $L$  always of type  $\mathbf{FP}$ ?

The article [2] serves as a fundamental reference for Question 0.4. While the question holds true for a significant number of examples, there exists a counterexample, as demonstrated in Example 4.11. Despite this, we will explore alternative homological conditions that might exhibit monoidal invariance. Notably, Theorem 4.6 will prove that being of type  $\mathbf{FP}_\infty$  is a monoidal invariant for Hopf algebras. This theorem suggests that even in the presence of counterexamples, certain homological properties can still retain their invariance under monoidal Morita-Takeuchi equivalence, providing valuable insights into the structure and behavior of these algebras. Moreover, we have established the following theorem.

**Theorem 0.5.** Let  $H$  and  $L$  be two Hopf algebras that are monoidally Morita-Takeuchi equivalent and of type  $\mathbf{FP}_\infty$ . Suppose that the antipodes of  $H$  and  $L$  are both bijective.

- (1) If  $H$  has finite global dimension  $d$ , then the injective dimension  $\text{injdim}(L) = d$ .
- (2) If  $H$  is AS regular of dimension  $d$ , then  $L$  is AS Gorenstein of dimension  $d$ .

Based on this, we pose the following question.

**Question 0.6.** Let  $H$  and  $L$  be two Hopf algebras that are Morita-Takeuchi equivalent and of type  $\mathbf{FP}_\infty$ . Does it follow that  $\text{injdim}(L) = \text{injdim}(H)$ ?

We are able to answer this question under the AS Gorenstein assumption.

**Theorem 0.7.** Let  $H$  and  $L$  be two Hopf algebras with bijective antipodes that are monoidally Morita-Takeuchi equivalent. If both  $L$  and  $H$  are AS Gorenstein of type  $\mathbf{FP}_\infty$ , then  $\text{injdim}(L) = \text{injdim}(H)$ .

Notice that any noetherian AS Gorenstein Hopf algebra has a bijective antipode, as established in [17]. The following corollary is a direct consequence of the famous result established by Wu and Zhang in [30, Theorem 0.2], which states that any noetherian affine PI Hopf algebra is AS Gorenstein.

**Corollary 0.8.** Let  $L$  and  $H$  be two noetherian affine PI Hopf algebras. If they are monoidally Morita-Takeuchi equivalent, then  $\text{injdim}(L) = \text{injdim}(H)$ .

A confirmation of the Brown-Goodearl conjecture (Conjecture 0.1) would significantly advance the resolution of Question 0.6, especially when both  $L$  and  $H$  are noetherian.

The paper is structured as follows. In Section 1, we introduce the fundamental definitions and characteristics of Hopf Galois extensions. Additionally, we delve into the investigation of homological dimensions by utilizing Stefan's spectral sequence. Section 2 focuses on establishing that a faithfully flat Hopf Galois extension of an AS Gorenstein algebra retains the AS Gorenstein property. Section 3 examines the injective dimension of Ehresmann-Schauburg bialgebroids. This investigation sets the stage for Section 4, where we show that a Hopf algebra  $H$  is AS Gorenstein of dimension  $d$  if it admits a faithfully flat  $H$ -Galois extension  $B$  over the base field  $\mathbb{k}$ , with  $B$  being a skew Calabi-Yau algebra of dimension  $d$ . In Section 4, we also elaborate on the implications of our results, focusing particularly on Theorems 0.5 and 0.7. Furthermore, we offer a counterexample to respond to Question 0.4.

## 1. PRELIMINARIES

Throughout this paper,  $\mathbb{k}$  is a base field, and all vector spaces and algebras are over  $\mathbb{k}$ . Unadorned  $\otimes$  means  $\otimes_{\mathbb{k}}$  and  $\text{Hom}$  means  $\text{Hom}_{\mathbb{k}}$ . Let  $R$  be an algebra over the base field  $\mathbb{k}$ . Let  $R^{op}$  denote the opposite ring of  $R$ , and let  $R^e$  denote the enveloping algebra  $R \otimes R^{op}$ . Observe that the flip map, which sends  $a \otimes b$  to  $b \otimes a$ , constitutes an anti-automorphism of the algebra  $R^e$ . Usually we work with left modules. A right  $R$ -module is viewed as an  $R^{op}$ -module, and an  $R$ - $R$ -bimodule is the same as an  $R^e$ -module.

Let  $M$  be a left module over an algebra  $R$ . Let  $\text{pdim}({}_R M)$  denote the projective dimension of  $M$  and  $\text{injdim}({}_R M)$  the injective dimension of  $M$ . Let  $\text{lgld}(R)$  (resp.,  $\text{rgld}(R)$ ) denote the left (resp., right) global dimension of  $R$ . We say that  $R$  has finite injective dimension if the modules  ${}_R R$  and  $R_R$  have finite injective dimensions which are equal, and denote these numbers by  $\text{injdim}(R)$ . We say that  $R$  has finite global dimension if the left and right global dimensions of  $R$  are finite and equal, and denote these numbers by  $\text{gldim}(R)$ .

Let  $H$  stand for a Hopf algebra  $(H, \Delta, \varepsilon)$  with antipode  $S$ . We use the Sweedler notation  $\Delta(h) = \sum_{(h)} h_1 \otimes h_2$  for all  $h \in H$ . We recommend [19] as a basic reference for the theory of Hopf algebras.

The antipode of a finite-dimensional Hopf algebra is bijective [14]. Skryabin showed that the antipode of a noetherian Hopf algebra is injective and hypothesized it to be bijective [27]. Le Meur confirmed the bijectivity of the antipode for Hopf algebras with Van den Bergh duality [15]. Furthermore, noetherian AS Gorenstein Hopf algebras are guaranteed to have a bijective antipode [17]. Nonetheless, there exist Hopf algebras with non-bijective antipodes, as seen in [28] and [24]. Let  $H$  be a Hopf algebra with a bijective antipode  $S$ . Since the antipode  $S : H \rightarrow H^{op}$  is an algebra isomorphism, left and right injective dimensions of  $H$  coincide. In the rest of this paper, the antipode of the given Hopf algebra is always assumed to be bijective.

**1.1. Faithfully flat Hopf Galois extensions.** Let  $H$  be a Hopf algebra over the field  $\mathbb{k}$ , and  $B$  be a right  $H$ -comodule algebra via an algebra homomorphism  $\rho_B : B \rightarrow B \otimes H$ . The *coinvariant subalgebra* of the  $H$ -coaction on  $B$  is  $\{b \in B \mid \rho_B(b) = b \otimes 1\}$ , denoted by  $B^{coH}$ .

**Definition 1.1.** The extension  $A(:= B^{coH}) \subset B$  is called (right) *H-Galois* if the canonical map

$$\beta : B \otimes_A B \rightarrow B \otimes H, \quad b' \otimes_A b \mapsto b' \rho(b) = \sum_{(b)} b' b_0 \otimes b_1$$

is bijective.

Let  $R$  be an algebra, and  ${}_R\mathcal{M}$  (resp.,  $\mathcal{M}_R$ ) be the category of left (resp., right)  $R$ -modules. If  $R$  is a right  $H$ -comodule algebra via  $\rho_R : R \rightarrow R \otimes H$ , then a relative right-right  $(R, H)$ -Hopf module is a vector space with a right  $R$ -action and a right  $H$ -coaction  $\rho_M$  such that

$$\rho_M(mr) = \rho_M(m)\rho_R(r) = \sum_{(m),(r)} m_0 r_0 \otimes m_1 r_1$$

for all  $m \in M$  and  $r \in R$ . A relative left-right  $(R, H)$ -Hopf module is defined similarly. The category of relative right-right (resp., left-right)  $(R, H)$ -Hopf modules is denoted by  $\mathcal{M}_R^H$  (resp.,  ${}_R\mathcal{M}^H$ ).

We require some known facts about Hopf Galois extensions.

**Lemma 1.2.** [10, (1.6)] [26, Remark 3.3] Let  $B$  be a right  $H$ -comodule algebra. Then the following are equivalent.

- (1)  $B$  is an injective right  $H$ -comodule.
- (2) There is a right  $H$ -colinear and unitary map  $\varphi : H \rightarrow B$ .
- (3) Any Hopf module in  $\mathcal{M}_B^H$  is injective as right  $H$ -comodule.

**Theorem 1.3.** [26, Theorem I] Let  $H$  be a Hopf algebra with a bijective antipode,  $B$  be a right  $H$ -comodule algebra and  $A := B^{coH}$ . Then the following are equivalent.

- (1)  $B$  is injective as right  $H$ -comodule, and  $\beta : B \otimes_A B \rightarrow B \otimes H$  is surjective.
- (2)  $(- \otimes_A B, (-)^{coH})$  is an equivalence between  $\mathcal{M}_A$  and  $\mathcal{M}_B^H$ .
- (3)  $(B \otimes_A -, (-)^{coH})$  is an equivalence between  ${}_A\mathcal{M}$  and  ${}_B\mathcal{M}^H$ .
- (4)  $B$  is a faithfully flat left  $A$ -module, and  $A \subseteq B$  is an  $H$ -Galois extension.
- (5)  $B$  is a faithfully flat right  $A$ -module, and  $A \subseteq B$  is an  $H$ -Galois extension.

Since conditions (4) and (5) in Theorem 1.3 are equivalent, we say a Hopf Galois extension  $A \subseteq B$  is faithfully flat if  $B$  is faithfully flat as a left or right  $A$ -module.

Let  $A(= B^{coH}) \subset B$  be an  $H$ -Galois extension. For any  $A$ - $A$ -bimodule  $M$ , we set

$$M^A := \{m \in M \mid ma = am, \forall m \in M, a \in A\}, \quad M_A := M/[M, A]$$

where  $[M, A]$  is the subspace of  $M$  generated by all commutators  $[m, a] := ma - am$ .

Following [7] we shall call the  $\mathbb{k}$ -linear map

$$\kappa : H \rightarrow (B \otimes_A B)^A, \quad h \mapsto \beta^{-1}(1 \otimes h)$$

the *translation map* associated to the  $H$ -Galois extension  $A \subseteq B$ . For any  $h \in H$  we shall use the notation

$$\kappa(h) = \sum \kappa^1(h) \otimes_A \kappa^2(h).$$

The following are key properties of  $\kappa$  as outlined in [26]:

$$(1.1) \quad \sum \kappa^1(hk) \otimes_A \kappa^2(hk) = \sum \kappa^1(k)\kappa^1(h) \otimes_A \kappa^2(h)\kappa^2(k),$$

$$(1.2) \quad \sum a\kappa^1(h) \otimes_A \kappa^2(h) = \sum \kappa^1(h) \otimes_A \kappa^2(h)a,$$

$$(1.3) \quad \sum_{(h)} \kappa^1(h) \otimes_A \kappa^2(h)_0 \otimes \kappa^2(h)_1 = \sum_{(h)} \kappa^1(h_1) \otimes_A \kappa^2(h_1) \otimes h_2,$$

$$(1.4) \quad \sum_{(h)} \kappa^1(h)_0 \otimes_A \kappa^2(h) \otimes \kappa^1(h)_1 = \sum_{(h)} \kappa^1(h_2) \otimes_A \kappa^2(h_2) \otimes Sh_1,$$

$$(1.5) \quad \sum \kappa^1(h)\kappa^2(h) = \varepsilon(h),$$

Suppose that  $B^{coH} = A$ . Then for any  $B$ - $B$ -bimodule  $M$ ,  $M^A$  is a right  $H$ -module which is defined by

$$(1.6) \quad x \leftarrow h = \sum_{(h)} \kappa^1(h)x\kappa^2(h)$$

for any  $x \in M^A$  and  $h \in H$ ; and  $M_A$  is a left  $H$ -module which is defined by

$$h \rightarrow (x + [A, M]) = \sum_{(h)} \kappa^2(h)x\kappa^1(h) + [A, M]$$

for any  $x \in M$  and  $h \in H$ , see [12, 2.1, 2.2]. In prior literature, specifically [11] and [9], both these structures have been introduced and meticulously examined, where they are recognized as Ulbrich-Miyashita actions. A crucial point to highlight is the functorial nature of these actions; in essence, this means that there exist two well-defined functors given by:

$$(-)^A : {}_B\mathcal{M}_B \longrightarrow \mathcal{M}_H, \quad (-)_A : {}_B\mathcal{M}_B \longrightarrow {}_H\mathcal{M}.$$

**1.2. Stefan's spectral sequence for flat Hopf Galois extensions.** We say a Hopf Galois extension  $A \subseteq B$  is flat if  $B$  is flat both as a left and right  $A$ -module. In this subsection, we give some results about homological properties of flat Hopf Galois extensions by using the Stefan's spectral sequence.

For any given flat Hopf-Galois extension  $A \subseteq B$ , Stefan in his work [9] devised a right  $H$ -module structure on the Hochschild cohomology group  $\mathrm{HH}^i(A, M)$ , which is isomorphic to  $\mathrm{Ext}_{A^e}^i(A, M)$ , for any  $B^e$ -module  $M$ . This construction was achieved by employing universal  $\delta$ -functors and serves as an extension of the Miyashita-Ulbrich action.

The inherent flatness property of  ${}_A B^e$  ensures that every injective  $B^e$ -module also maintains its injectivity when considered as an  $A^e$ -module. Consequently, the derived right  $H$ -module structure on  $\mathrm{Ext}_{A^e}^i(A, M)$  can be effectively realized by computing homological groups from the complex  $\mathrm{Hom}_{A^e}(A, E^\bullet)$ , where  $E^\bullet$  is a  $B^e$ -injective resolution of  $M$ .

Subsequently, Stefan further developed in [9] a spectral sequence that establishes a connection between the Hochschild cohomologies of  $A$  and those of its flat Hopf-Galois extension  $B$ .

**Theorem 1.4.** [9, Theorem 3.3] Let  $A \subseteq B$  be a flat  $H$ -Galois extension. For any  $B^e$ -module  $M$ , there is a convergent spectral sequence

$$\mathrm{Ext}_{H^{op}}^p(\mathbb{k}, \mathrm{Ext}_{A^e}^q(A, M)) \implies \mathrm{Ext}_{B^e}^{p+q}(B, M).$$

Recall that  $\mathrm{pd}({}_H\mathbb{k}) = \mathrm{gldim}(H) = d$  by [4, Corollary 1.4]. Then we have the following immediate consequence.

**Corollary 1.5.** Let  $A \subseteq B$  be a flat  $H$ -Galois extension. Then  $\mathrm{pd}({}_{B^e}B) \leq \mathrm{pd}(\mathbb{k}_H) + \mathrm{pd}({}_{A^e}A) = \mathrm{rgld}(H) + \mathrm{pd}({}_{A^e}A)$ .

In the following, we study the global dimension of faithfully flat Hopf Galois extension.

Let  $M$  and  $N$  be two modules over a  $\mathbb{k}$ -algebra  $R$ . Let  $P_\bullet$  denote a projective resolution of  ${}_R R$ . Given that the sequence  $\cdots \rightarrow P_1 \rightarrow P_0 \rightarrow R \rightarrow 0$  is split exact as an  $R$ -module complex, the complex

$$\cdots \longrightarrow P_1 \otimes_R M \longrightarrow P_0 \otimes_R M \longrightarrow R \otimes_R M (= M) \longrightarrow 0$$

remains exact as well. Owing to the projectivity of each module  $M \otimes_R P_i$  for all  $i$ , the complex  $M \otimes_R P_\bullet$  constitutes a projective resolution of the right  $R$ -module  $M$ . Furthermore, utilizing the isomorphism

$$\mathrm{Hom}_{R^e}(P_\bullet, \mathrm{Hom}(M, N)) \cong \mathrm{Hom}_{R^{op}}(P_\bullet \otimes_R M, N),$$

upon taking homology groups, we obtain the following isomorphism between Ext-groups,

$$(1.7) \quad \text{Ext}_{R^e}^i(R, \text{Hom}(M, N)) \cong \text{Ext}_{R^{op}}^i(M, N).$$

for all  $i \geq 0$ .

**Lemma 1.6.** Let  $A \subseteq B$  be a flat  $H$ -Galois extension. For any right  $B$ -modules  $M$  and  $N$ , there is a convergent spectral sequence

$$\text{Ext}_{H^{op}}^p(\mathbb{k}, \text{Ext}_{A^{op}}^q(M, N)) \implies \text{Ext}_{B^{op}}^{p+q}(M, N).$$

*Proof.* In light of the isomorphism (1.7), for every integer  $i \geq 0$ , we have

$$\text{Ext}_{A^e}^i(A, \text{Hom}(M, N)) \cong \text{Ext}_{A^{op}}^i(M, N).$$

This implies that  $\text{Ext}_{A^{op}}^i(M, N)$  inherits a right  $H$ -module structure through this isomorphism. By Theorem 1.4, there exists a convergent spectral sequence which unfolds as follows:

$$\text{Ext}_{H^{op}}^p(\mathbb{k}, \text{Ext}_{A^{op}}^q(M, N)) \implies \text{Ext}_{B^e}^{p+q}(B, \text{Hom}(M, N)) \cong \text{Ext}_{B^{op}}^{p+q}(M, N).$$

So we get the conclusion.  $\square$

The following result is an immediate consequence of Lemma 1.6.

**Proposition 1.7.** If  $A \subseteq B$  is a flat  $H$ -Galois extension, then  $\text{rgld}(B) \leq \text{gldim}(H) + \text{rgld}(A)$ .

The following example illustrates that within the above proposition, the equality may not be attained.

**Example 1.8.** For a finite-dimensional Hopf algebra  $H$  and its dual  $H^*$ , the Heisenberg double  $B := H \# H^*$  is isomorphic to  $\text{End}(H)$ , a separable algebra that forms an  $H^*$ -Galois extension over its subalgebra  $A = H$ . When  $H$  is not cosemisimple, meaning  $H^*$  is non-semisimple, the following strict inequalities hold:  $\text{pd}_{(B^e)} < \text{gldim}(H^*) + \text{pd}(A_{A^e})$  and  $\text{rgld}(B) < \text{gldim}(H^*) + \text{rgld}(A)$ .

## 2. FAITHFULL FLAT PI HOPF GALOIS EXTENSIONS ARE ALWAYS AS GORENSTEIN

**2.1. Artin-Schelter Gorenstein Hopf algebras.** Recall that an algebra is called Gorenstein when it possesses finite injective dimensions over itself on both the left and right sides. We now turn our attention to a stronger variant of this Gorenstein property for noncommutative rings.

**Definition 2.1.** [21, Definition 1.3] Let  $R$  be a noetherian algebra. We say  $R$  is *left AS Gorenstein* (where AS stands for Artin and Schelter) if

- (1)  ${}_R R$  has finite injective dimension, say  $d$ , and
- (2) for every finite dimensional left  $R$ -module  $V$ ,  $\text{Ext}_R^i(V, R) = 0$  for all  $i \neq d$  and  $\text{Ext}_R^d(V, R)$  is also finite dimensional.

Right AS Gorensteinness is analogously defined. We say  $R$  is *AS Gorenstein* if it is both left and right AS Gorenstein. If further,  $\text{gldim } R = d < +\infty$ , then  $R$  is called *AS regular*.

If  $R$  is an AS Gorenstein algebra of dimension  $d$ , then  $\text{Ext}_R^d(V, R) \neq 0$  for any finite dimensional  $R$ -module  $V$ . Refer to [21, Lemma 1.4] for details.

Let us revisit the definition of skew Calabi-Yau algebras as presented in [20, Definition 0.1], a class intimately related to AS regular algebras. An algebra  $R$  is called homologically smooth when it admits a bounded resolution consisting entirely of finitely generated projective modules over its enveloping algebra  $R^e$ . According to [15, Lemma 3.3], for instance, a Hopf algebra  $H$  is said to be homologically smooth if and only if it is of type **FP**.

An homological smooth algebra  $R$  is called *skew Calabi-Yau* of dimension  $d$  for some integer  $d \geq 0$ , if  $R$  has an automorphism  $\mu$  of  $R$  such that

$$\mathrm{Ext}_{R^e}^i(R, R^e) \cong \begin{cases} 0, & i \neq d \\ R_\mu, & i = d \end{cases}$$

as  $R$ - $R$ -bimodules, where  $R_\mu$  signifies the  $R$ - $R$ -bimodule twisted by  $\mu$ . This bimodule is identical to  $R$  as a  $\mathbb{k}$ -vector space, with the action defined by  $r \cdot x \cdot r' = rx\mu(r')$  for all  $r, r' \in R$  and  $x \in R_\mu$ .

When an algebra  $R$  is AS Gorenstein of dimension  $d$ , for each finite-dimensional simple left module  $V$ , the Ext group  $\mathrm{Ext}_R^d(V, R)$  is also a simple right  $R$ -module, and a similar property holds on the right side, as demonstrated in [21, Lemma 1.4]. This observation substantiates that Definition 2.1 is indeed equivalent to the definition of AS Gorenstein provided in [30, Definition 3.1] for any algebra  $R$  whose simple modules are all finite-dimensional.

In the following, we show that Definition 2.1 is equivalent to the definition of AS Gorenstein in [5, Definition 1.2] for any noetherian Hopf algebra with a bijective antipode.

**Definition 2.2.** Let  $H$  be a Hopf algebra of type  $\mathbf{FP}_\infty$  with a bijective antipode. We say  $H$  is *AS Gorenstein* if

(1)  ${}_H H$  has finite injective dimension, say  $d$ , and

$$(2) \mathrm{Ext}_H^i(\mathbb{k}, H) = \begin{cases} 0, & i \neq d \\ \mathbb{k}, & i = d \end{cases}$$

If further,  $\mathrm{gldim} H = d$ , then  $H$  is called *AS regular*.

Recall the standard actions by which tensor products and Hom-spaces of  $H$ -modules  $U$  and  $V$  become  $H$ -modules

$$h \rightharpoonup (u \otimes v) = \sum_{(h)} (h_1 \rightharpoonup u) \otimes (h_2 \rightharpoonup v), \quad (h \rightharpoonup f)(u) = \sum_{(h)} h_1 \rightharpoonup f(Sh_2 \rightharpoonup u)$$

for all  $h \in H$ ,  $u \in U$ ,  $v \in V$  and  $f \in \mathrm{Hom}(U, V)$ . In case the  $H$ -action on  $V$  is trivial, the second formula simplifies. In particular, for any  $h \in H$ ,  $u \in U$  and  $f \in V^* = \mathrm{Hom}(V, \mathbb{k})$ ,

$$(h \rightharpoonup f)(u) = \sum_{(h)} h_1 \rightharpoonup f(Sh_2 \rightharpoonup u) = \sum_{(h)} \varepsilon(h_1) f(Sh_2 \rightharpoonup u) = f(Sh \rightharpoonup u).$$

Then the natural adjoint isomorphism

$$\mathrm{Hom}(U \otimes V, W) \longrightarrow \mathrm{Hom}(U, \mathrm{Hom}(V, W)),$$

restricts to an isomorphism

$$\mathrm{Hom}_H(U \otimes V, W) \longrightarrow \mathrm{Hom}_H(U, \mathrm{Hom}(V, W)),$$

see [4]. Consequently, if  ${}_H U$  is projective, then  $U \otimes V$  is also a projective  $H$ -module by this adjoint isomorphism.

It is a standard fact that for any left  $H$ -modules  $V$  and  $W$ , the natural map

$$\varphi_{W,V} : W \otimes V^* \longrightarrow \mathrm{Hom}(V, W), \quad w \otimes f \mapsto f(v)w,$$

is an  $H$ -module map. Further,  $\varphi_{W,V}$  is bijective when  $V$  is finite dimensional. There is a morphism of right  $H$ -modules

$$\psi_{U,W} : \mathrm{Hom}_H(U, H) \otimes W \longrightarrow \mathrm{Hom}_H(U, H_H \otimes W), \quad f \otimes w \mapsto (u \mapsto \sum_{(f(u))} f(u)_0 \otimes f(u)_1 \rightharpoonup w)$$

where the right  $H$ -module structure on  $\mathrm{Hom}_H(U, H) \otimes W$  is given by

$$(f \otimes w) \leftarrow h = \sum_{(h)} (f \leftarrow h_1) \otimes (Sh_2 \rightharpoonup w)$$

for all  $f \in \text{Hom}_H(U, H)$ ,  $w \in W$  and  $h \in H$ . Further,  $\psi_{U,W}$  is bijective when  $U$  is finitely presented as an  $H$ -module.

**Lemma 2.3.** Let  $V$  be finite dimensional  $H$ -module, and  $U$  be an  $H$ -module of type  $\mathbf{FP}_\infty$ . Then for all  $i \geq 0$ ,

$$\text{Ext}_H^i(U \otimes V, H) \cong \text{Ext}_H^i(U, H) \otimes V^*$$

as right  $H$ -modules, the right  $H$ -module structure on  $\text{Ext}_H^i(U, H) \otimes V^*$  is given by

$$(x \otimes f) \leftarrow h = \sum_{(h)} (x \leftarrow h_1) \otimes (Sh_2 \rightarrow f)$$

for all  $x \in \text{Ext}_H^i(U, H)$ ,  $f \in V^*$  and  $h \in H$ .

*Proof.* For any finitely generated projective  $H$ -module  $P$ , there exist the following natural isomorphisms of right  $H$ -modules:

$$\text{Hom}_H(P, H) \otimes V^* \xrightarrow{\psi_{P,V^*}} \text{Hom}_H(P, H \otimes V^*) \xrightarrow{\text{Hom}(P, \varphi_{H,V})} \text{Hom}_H(P, \text{Hom}(V, H)) \cong \text{Hom}_H(P \otimes V, H).$$

Consider a projective resolution  $P_\bullet$  of  ${}_H U$  such that each component is a finitely generated  $H$ -module. Then  $P_\bullet \otimes V$  is a projective resolution of  ${}_H(U \otimes V)$ . Given that  $\text{Hom}_H(P_\bullet \otimes V, H)$  is isomorphic to  $\text{Hom}_H(P_\bullet, H) \otimes V^*$ , upon taking homologies, it follows that  $\text{Ext}_H^i(U \otimes V, H) \cong \text{Ext}_H^i(U, H) \otimes V^*$  for all  $i \geq 0$ .  $\square$

By using Ischebeck's spectral sequence, Brown and Zhang proved that the condition (2) of Definition 2.1 can be replaced by a weaker form when  $R = H$  is a Hopf algebra with a bijective antipode.

**Lemma 2.4.** [5, Lemma 3.2] Let  $H$  be a Hopf algebra of type  $\mathbf{FP}_\infty$  with a bijective antipode. If  $H$  has injective dimension  $d$ , then the following are equivalent.

(1) For each finite dimensional  $H$ -module  $V$ ,  $\text{Ext}_H^i(V, H) = 0$  for all  $i \neq d$  and  $\text{Ext}_H^d(V, H)$  is finite dimensional.

(2)  $\text{Ext}_H^i(\mathbb{k}, H) = 0$  for all  $i \neq d$  and  $\text{Ext}_H^d(\mathbb{k}, H)$  is finite dimensional.

(3)  $\text{Ext}_H^i(\mathbb{k}, H) = \begin{cases} 0, & i \neq d \\ \mathbb{k}, & i = d \end{cases}$

*Proof.* (1)  $\Rightarrow$  (2) It is trivial.

(2)  $\Rightarrow$  (3) Given that  $H$  is a Hopf algebra of type  $\mathbf{FP}_\infty$  with finite injective dimension, Ischebeck's spectral sequence yields

$$\text{Ext}_H^p(\text{Ext}_{H^{op}}^{-q}(\mathbb{k}, H), H) \Rightarrow \text{Tor}_{-p-q}^H(\mathbb{k}, H).$$

From Lemma 2.3, we deduce that  $\text{Ext}_H^d(\mathbb{k}, H) \otimes \text{Ext}_{H^{op}}^d(\mathbb{k}, H)^* \cong \text{Ext}_H^d(\text{Ext}_{H^{op}}^d(\mathbb{k}, H), H) \cong \mathbb{k}$ . Thus,  $\dim(\text{Ext}_H^d(\mathbb{k}, H)) = 1$ , implying  $\text{Ext}_H^d(\mathbb{k}, H) = \mathbb{k}$ .

(3)  $\Rightarrow$  (1) By Lemma 2.3, for any finite dimensional  $H$ -module  $V$ ,

$$\text{Ext}_H^i(V, H) \cong \text{Ext}_H^i(\mathbb{k}, H) \otimes V^* \cong \begin{cases} 0, & i \neq d \\ V^*, & i = d. \end{cases}$$

as right  $H$ -modules. This establishes the result.  $\square$



**2.2. Hopf module structure on  $\text{Hom}_{A^{op}}(M, X)$ .** Let  $H^*$  be the linear dual of  $H$ . A left module  $V$  over  $H^*$  is said to be rational if for any  $v \in V$ , the submodule generated by the action of  $H^*$  on  $v$ , denoted as  $H^* \cdot v$ , possesses finite dimension.

For any left  $H^*$ -module  $V$ , it is rational if and only if the  $H^*$ -action can be derived from a compatible right  $H$ -coaction. In other words, that is,  $V$  must be an  $H$ -comodule such that

$$\sum_{(v)} v_0 \langle f, v_1 \rangle = f \rightharpoonup v$$

for any  $v \in V$  and  $f \in H^*$ .

Let  $M$  be a vector space and  $N$  be a right  $H$ -comodule. Then  $\text{Hom}(N, M)$  is a left  $H^*$ -module, under the action

$$(h^* \rightharpoonup f)(m) = h^* \rightharpoonup (f(m))$$

for all  $h^* \in H^*$ ,  $f \in \text{Hom}(M, N)$  and  $m \in M$ .

Then we have the following immediate consequence.

**Lemma 2.5.** Let  $A = B^{coH} \subseteq B$  be an  $H$ -Galois extension, and  $M$  be a right  $A$ -module and  $X \in \mathcal{M}_A^H$ .

- (1) If  $M_A$  is finitely generated, then  $\text{Hom}_{A^{op}}(M, X)$  is an  $H$ -comodule.
- (2) If  $M_A$  is finitely presented, then  $\text{Hom}_{A^{op}}(M, X)^{coH} = \text{Hom}_{A^{op}}(M, X^{coH})$ .

*Proof.* (1) It is straightforward to verify that  $\text{Hom}_{A^{op}}(M, X)$  constitutes an  $H^*$ -submodule of  $\text{Hom}(M, X)$ . Furthermore, when  $M$  is finitely generated as an  $A$ -module,  $\text{Hom}_{A^{op}}(M, X)$  can be viewed as a subset of a direct sum of multiple copies of  $X$ . Given that  $X$  is a rational  $H^*$ -module, it follows that  $\text{Hom}_{A^{op}}(M, X)$  inherits its rationality and thus becomes an  $H$ -comodule itself.

Consequently, the  $H$ -coaction on  $\text{Hom}_{A^{op}}(M, X)$  is characterized by the following equation:

$$(2.1) \quad \sum_{(f)} f_0(m) \otimes f_1 = \sum_{(f(m))} f(m)_0 \otimes f(m)_1$$

for all  $f \in \text{Hom}_{A^{op}}(M, X)$  and  $m \in M$ .

- (2) For any finitely generated  $A$ -module  $N$ , we have a canonical morphism

$$\alpha_N : \text{Hom}_{A^{op}}(N, X^{coH}) \rightarrow \text{Hom}_{A^{op}}(N, X)^{coH}$$

which is functorial in both the variables  $N$  and  $X$ . If further,  $N$  is projective, then  $\alpha_N$  is bijective. Since  $M_A$  is finitely presented, there is a short exact sequence

$$P_1 \longrightarrow P_0 \longrightarrow M \longrightarrow 0,$$

where  $P_0$  and  $P_1$  are finitely generated projective  $A$ -module. Thus, we have a commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Hom}_{A^{op}}(M, X^{coH}) & \longrightarrow & \text{Hom}_{A^{op}}(P_0, X^{coH}) & \longrightarrow & \text{Hom}_{A^{op}}(P_1, X^{coH}) \\ & & \downarrow \alpha_M & & \downarrow \alpha_{P_0} & & \downarrow \alpha_{P_1} \\ 0 & \longrightarrow & \text{Hom}_{A^{op}}(M, X)^{coH} & \longrightarrow & \text{Hom}_{A^{op}}(P_0, X)^{coH} & \longrightarrow & \text{Hom}_{A^{op}}(P_1, X)^{coH} \end{array}$$

in which every row is exact. The map  $\alpha_M$  is also bijective because  $\alpha_{P_0}$  and  $\alpha_{P_1}$  are both bijective.  $\square$

**Lemma 2.6.** Given a  $B$ -module  $M$ , finitely generated over  $A$ , and any Hopf module  $X \in \mathcal{M}_B^H$ , the Hom space  $\text{Hom}_{A^{op}}(M, X)$  naturally admits a Hopf module structure in  $\mathcal{M}_H^H$ .

*Proof.* Notice that the right  $H$ -module structure on  $\text{Hom}_{A^{op}}(M, X)$ , equivalent to  $\text{Hom}(M, X)^A$ , is expressed by (1.6) as:

$$(2.2) \quad (f \leftarrow h)(m) = \sum f(m\kappa^1(h))\kappa^2(h).$$

For all  $f \in \text{Hom}_{A^{op}}(M, X)$ ,  $m \in M$ ,  $h \in H$  and  $h^* \in H^*$ ,

$$\begin{aligned} \sum_{(f \leftarrow h)} (f \leftarrow h)_0(m) \otimes (f \leftarrow h)_1 &\stackrel{(2.1)}{=} \rho_X((f \leftarrow h)(m)) \\ &\stackrel{(2.2)}{=} \rho_X(f(m\kappa^1(h))\kappa^2(h)) \\ &= \sum_{(f(m\kappa^1(h))), (\kappa^2(h))} f(m\kappa^1(h))_0\kappa^2(h)_0 \otimes f(m\kappa^1(h))_1\kappa^2(h)_1 \\ &\stackrel{(1.3)}{=} \sum_{(h), (f(m\kappa^1(h_1)))} f(m\kappa^1(h_1))_0\kappa^2(h_1) \otimes f(m\kappa^1(h_1))_1h_2 \\ &\stackrel{(2.1)}{=} \sum_{(h), (f)} f_0(m\kappa^1(h_1))\kappa^2(h_1) \otimes f_1h_2 \\ &\stackrel{(2.2)}{=} \sum_{(f), (h)} (f_0 \leftarrow h_1)(m) \otimes f_1h_2. \end{aligned}$$

Consequently,  $\text{Hom}_A(M, X)$  is endowed with a Hopf module structure in  $\mathcal{M}_H^H$ .  $\square$

### 2.3. AS Gorenstein property of Hopf Galois extensions.

**Lemma 2.7.** Let  $R$  be a ring and  $V$  be a right  $R$ -module of type  $\mathbf{FP}_\infty$ . If  $I_\lambda$  is an injective  $R$ -module for any  $\lambda \in \Lambda$ , then the direct sum  $\bigoplus_{\lambda \in \Lambda} I_\lambda$  is acyclic with respect to the functor  $\text{Hom}_{R^{op}}(V, -)$ , that is,  $\text{Ext}_{R^{op}}^i(V, \bigoplus_{\lambda \in \Lambda} I_\lambda) = 0$  for all  $i > 0$ .

*Proof.* By assumption,  $V_R$  admits a projective resolution  $P_\bullet$  where each  $R$ -module  $P_i$  is finitely generated. Therefore, we have the following natural isomorphism

$$\text{Hom}_{R^{op}}(P_i, \bigoplus_{\lambda \in \Lambda} I_\lambda) \cong \bigoplus_{\lambda \in \Lambda} \text{Hom}_{R^{op}}(P_i, I_\lambda).$$

Given that for all  $\lambda \in \Lambda$ , the sequence

$$0 \longrightarrow \text{Hom}_{R^{op}}(V, I_\lambda) \longrightarrow \text{Hom}_{R^{op}}(P_0, I_\lambda) \longrightarrow \cdots \longrightarrow \text{Hom}_{R^{op}}(P_i, I_\lambda) \longrightarrow \cdots$$

is exact, it follows that the sequence

$$0 \longrightarrow \text{Hom}_{R^{op}}(V, \bigoplus_{\lambda \in \Lambda} I_\lambda) \longrightarrow \text{Hom}_{R^{op}}(P_0, \bigoplus_{\lambda \in \Lambda} I_\lambda) \longrightarrow \cdots \longrightarrow \text{Hom}_{R^{op}}(P_i, \bigoplus_{\lambda \in \Lambda} I_\lambda) \longrightarrow \cdots$$

also remains exact. Taking homologies then yields  $\text{Ext}_{R^{op}}^i(V, \bigoplus_{\lambda \in \Lambda} I_\lambda) = 0$  for all  $i > 0$ . This confirms that  $\bigoplus_{\lambda \in \Lambda} I_\lambda$  is acyclic with respect to the functor  $\text{Hom}_{R^{op}}(V, -)$ .  $\square$

**Lemma 2.8.** Given a  $B$ -module  $V$ , if  $V_A$  is of type  $\mathbf{FP}_\infty$ , then  $\text{Ext}_{A^{op}}^i(V, B) \cong \text{Ext}_{A^{op}}^i(V, A) \otimes H$  as right  $H$ -modules for  $i \geq 0$ .

*Proof.* For any  $X \in \mathcal{M}_B^H$ , there is an injective right  $B$ -module  $I$  with a  $B$ -module inclusion  $\iota : X \hookrightarrow I$ . Hence there is an injective morphism  $X \hookrightarrow I \otimes H$ ,  $x \mapsto \sum_{(x)} \iota(x_0) \otimes x_1$  in  $\mathcal{M}_B^H$ . By induction, we construct an exact complex  $0 \rightarrow B \rightarrow E^0 \rightarrow E^1 \rightarrow \cdots$  in  $\mathcal{M}_B^H$  where each term  $E^i$  is isomorphic to  $I^i \otimes H$  and  $I^i$  is an injective  $B$ -module for any  $i \in \mathbb{N}$ .

Since  ${}_A B$  is flat, any injective right  $B$ -module  $I$  is also injective as an  $A$ -module. Moreover, it's clear that  $I \otimes H$  can be viewed as a direct sum of copies of  $I$  as  $A$ -modules. By virtue of Lemma 2.7,  $I \otimes H$  is  $\text{Hom}_{A^{op}}(V, -)$ -acyclic. Thus, we infer that for all  $i \geq 0$ ,

$$\text{Ext}_{A^{op}}^i(V, B) \cong \text{H}^i(\text{Hom}_{A^{op}}(V, E^\bullet))$$

as  $H$ -modules.

As  $\text{Hom}_{A^{op}}(V, E^\bullet)$  forms a complex in  $\mathcal{M}_H^H$  by Lemma 2.6, we obtain the following isomorphisms

$$\begin{aligned} \text{H}^i(\text{Hom}_{A^{op}}(V, E^\bullet)) &\cong \text{H}^i(\text{Hom}_{A^{op}}(V, E^\bullet)^{coH}) \otimes H \\ &\cong \text{H}^i(\text{Hom}_{A^{op}}(V, E^\bullet)^{coH}) \otimes H && \text{since } (-)^{coH} \text{ is exact on } \mathcal{M}_H^H \\ &\cong \text{H}^i(\text{Hom}_{A^{op}}(V, (E^\bullet)^{coH})) \otimes H && \text{by Lemma 2.5 (2)}. \end{aligned}$$

Since the functor  $(-)^{coH} : \mathcal{M}_B^H \rightarrow \mathcal{M}_A$  is an equivalence, the sequence

$$0 \longrightarrow A = B^{coH} \longrightarrow (E^0)^{coH} \longrightarrow (E^1)^{coH} \longrightarrow \dots$$

is exact. Furthermore, each term  $(E^i)^{coH} = (I^i \otimes H)^{coH}$  can be identified with  $I^i$  as a right  $A$ -module. Therefore, the complex  $(E^\bullet)^{coH}$  constitutes an injective resolution of  $A_A$ . Then we derive the following isomorphism of right  $H$ -modules,

$$\text{Ext}_{A^{op}}^i(V, B) \cong \text{H}^i(\text{Hom}_{A^{op}}(V, E^\bullet)) \cong \text{H}^i(\text{Hom}_{A^{op}}(V, (E^\bullet)^{coH})) \otimes H \cong \text{Ext}_{A^{op}}^i(V, A) \otimes H.$$

This leads us to our desired conclusion.  $\square$

Recall that an algebra  $R$  is called affine if it is finitely generated as a  $\mathbb{k}$ -algebra. Let  $\text{Kdim}$  denote the Krull dimension. A polynomial identity algebra, or PI algebra for short, is an algebra satisfying a polynomial identity. We refer to [18, MR, Chapter 13] for some basic materials about PI algebras.

Now, let's proceed with the following hypothesis.

**Hypothesis 2.9.** Let  $R$  be a noetherian affine PI algebra. For any  $i \geq 0$ ,

- (1)  $\text{Ext}_{R^{op}}^i(U, R) \neq 0$  if and only if  $\text{Ext}_{R^{op}}^i(V, R) \neq 0$  for all simple  $A$ -modules  $U$  and  $V$ , and
- (2)  $\text{Ext}_{R^{op}}^i(-, R)$  is an exact functor on  $A$ -modules of finite length.

In [30], Wu and Zhang gave a useful criterion for noetherian affine PI algebras to be AS Gorenstein.

**Theorem 2.10.** [30, Corollary 2.10, Propostion 3.2] If the Hypothesis 2.9 holds, then

- (1)  $\text{injdim}({}_R R) = \text{injdim}(R_R) = d = \text{Kdim } R < +\infty$ ;
- (2)  $R$  is AS Gorenstein of dimension  $d$ .

Using the above theorem of Wu and Zhang, we promptly obtain the following theorem.

**Theorem 2.11.** Let  $H$  be a Hopf algebra, and  $B$  is a noetherian affine PI algebra which is a faithfully flat  $H$ -Galois extension of  $A$ . If  $A$  is AS Gorenstein, then  $B$  is also AS Gorenstein. Further, if  $H$  is AS Gorenstein of dimension  $d_H$ , then  $\text{injdim}(B_B) = \text{injdim}(A_A) + d_H$ .

*Proof.* Given that  $B$  is an affine PI algebra, every simple  $B$ -module has finite dimension [18, 13.10.3(i)]. Let  $U$  be a simple right  $B$ -module. Then  $U$  is finite dimensional. Since  $B$  is faithfully flat as a right  $A$ -module,  $A$  is right noetherian. Consequently,  $U$  is an  $A$ -module of type  $\mathbf{FP}_\infty$ . This yields the isomorphism  $\text{Ext}_{A^{op}}^i(U, B) \cong \text{Ext}_{A^{op}}^i(U, A) \otimes_A B$  for all  $i \geq 0$ . If  $A$  is AS Gorenstein of dimension  $d_A$ , then  $\text{Ext}_{A^{op}}^i(U, A) = 0$  for  $i \neq d_A$  and  $\text{Ext}_{A^{op}}^{d_A}(U, A)$  is non-zero

and finite dimensional. Hence  $\text{Ext}_{A^{op}}^i(U, B) \neq 0$  if and only if  $i = d_A$ . By Lemma 1.6, we infer that for any integer  $i \geq 0$ ,

$$\text{Ext}_{B^{op}}^i(U, B) \cong \begin{cases} 0, & i < d_A; \\ \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, \text{Ext}_{A^{op}}^{d_A}(U, B)), & i \geq d_A. \end{cases}$$

According to Lemma 2.8,  $\text{Ext}_{A^{op}}^{d_A}(U, B)$  constitutes a free  $H$ -module of rank  $\dim(\text{Ext}_{A^{op}}^{d_A}(U, A))$ . Therefore,

$$\text{Ext}_{B^{op}}^i(U, B) \neq 0 \iff \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, \text{Ext}_{A^{op}}^{d_A}(U, B)) \neq 0 \iff \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, H) \neq 0.$$

For another simple  $B$ -module  $V$ ,

$$\text{Ext}_{B^{op}}^i(U, B) \neq 0 \iff \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, H) \neq 0 \iff \text{Ext}_{B^{op}}^i(V, B) \neq 0.$$

Thus,  $B$  fulfills Hypothesis 2.9 (1).

Consider an exact sequence  $0 \rightarrow W' \rightarrow W \rightarrow W'' \rightarrow 0$  of finite length left  $B$ -modules. If  $A$  is AS-Gorenstein of dimension  $d_A$ , then we obtain a short exact sequence

$$(2.3) \quad 0 \longrightarrow \text{Ext}_{A^{op}}^{d_A}(W'', B) \longrightarrow \text{Ext}_{A^{op}}^{d_A}(W, B) \longrightarrow \text{Ext}_{A^{op}}^{d_A}(W', B) \longrightarrow 0$$

in  $\mathcal{M}_H^H$ . By Lemma 2.8, each term in (2.3) is a projective  $H$ -module, making (2.3) split as an exact sequence of  $H$ -modules. Hence, the following sequence

$$0 \rightarrow \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, \text{Ext}_{A^{op}}^{d_A}(W'', B)) \rightarrow \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, \text{Ext}_{A^{op}}^{d_A}(W, B)) \rightarrow \text{Ext}_{H^{op}}^{i-d_A}(\mathbb{k}, \text{Ext}_{A^{op}}^{d_A}(W', B)) \rightarrow 0$$

remains exact. Consequently, for all  $i \geq 0$ , we have the exact sequence

$$0 \longrightarrow \text{Ext}_{B^{op}}^i(W'', B) \longrightarrow \text{Ext}_{B^{op}}^i(W, B) \longrightarrow \text{Ext}_{B^{op}}^i(W', B) \longrightarrow 0,$$

showing that  $B$  satisfies Hypothesis 2.9 (2). Therefore, by Theorem 2.10, the desired conclusion is reached.  $\square$

It is a well-established fact that AS Gorenstein noetherian affine PI algebras exhibit numerous favorable homological properties, see [30, Proposition 3.7] for instance.

### 3. HOMOLOGICAL DIMENSION OF THE EHRESMANN-SCHAUBENBURG BIALGEBROIDS

In this section, we shall revisit the definition and essential attributes of Ehresmann-Schaubenburg bialgebroids, further elucidating certain homological properties inherent to these structures.

**3.1. Hopf bimodule categories and Ehresmann-Schaubenburg bialgebroids.** Let  $H$  be a Hopf algebra,  $R$  and  $T$  be two  $H$ -comodule algebras. A relative Hopf bimodule  $X \in {}_R\mathcal{M}_T^H$  is an object of both  ${}_R\mathcal{M}^H$  and  $\mathcal{M}_T^H$  with the same comodule structure and such that the two module structures make it an  $R$ - $T$ -bimodule.

For any  $R \otimes T^{op}$ -module  $M$ ,  $M \otimes H$  can be viewed as a Hopf module in  ${}_R\mathcal{M}_T^H$  through the structure defined by

$$(3.1) \quad r(m \otimes h)t = \sum_{(r),(t)} r_0 m t_0 \otimes r_1 h t_1, \quad \rho(m \otimes h) = \sum_{(h)} m \otimes h_1 \otimes h_2,$$

for any  $m \in M$ ,  $h \in H$ ,  $r \in R$  and  $t \in T$ . The functor  $- \otimes H$  serves as a right adjoint to the forgetful functor from  ${}_R\mathcal{M}_T^H$  to  ${}_R\mathcal{M}_T$ , that is, there exists a natural isomorphism

$$(3.2) \quad \text{Hom}_{{}_R\mathcal{M}_T}(X, M) \longrightarrow \text{Hom}_{{}_R\mathcal{M}_T^H}(X, M \otimes H), \quad f \mapsto \left( x \mapsto \sum_{(x)} f(x_0) \otimes x_1 \right)$$

for any  $X \in {}_R\mathcal{M}_T^H$  and  $M \in {}_R\mathcal{M}_T$ . The inverse map is given by  $g \mapsto (\text{id}_M \otimes \varepsilon) \circ g$ .

It is evident that the left (or right) adjoint functor of an exact functor preserves projective (or injective) objects. Given that  $- \otimes H : {}_R\mathcal{M}_T \rightarrow {}_R\mathcal{M}_T^H$  and the forgetful functor  ${}_R\mathcal{M}_T^H \rightarrow {}_R\mathcal{M}_T$  are both exact, the following lemmas can be derived.

**Lemma 3.1.** (1) If  $X$  is a projective object in  ${}_R\mathcal{M}_T^H$ , then  $X$  is projective  $R \otimes T^{op}$ -module.  
(2) If  $M$  is an injective  $R \otimes T^{op}$ -module, then  $M \otimes H$  is an injective object in  ${}_R\mathcal{M}_T^H$ .

**Lemma 3.2.** Let  $H$  be a Hopf algebra,  $R$  and  $T$  be two  $H$ -comodule algebras.

(1) Then  ${}_R\mathcal{M}_T^H$  has enough injective objects.  
(2) For any  $X \in {}_R\mathcal{M}_T^H$ , there is an exact complex  $0 \rightarrow X \rightarrow I^0 \otimes H \rightarrow I^1 \otimes H \rightarrow \cdots$  in  ${}_R\mathcal{M}_T^H$  where  $I^i$  is an injective  $R \otimes T^{op}$ -module for any  $i \in \mathbb{N}$ .

*Proof.* (1) For any  $X \in {}_R\mathcal{M}_T^H$ , there is an injective  $R \otimes T^{op}$ -module  $I$  with a  $R \otimes T^{op}$ -module inclusion  $\iota : X \hookrightarrow I$ . By Lemma 3.1,  $I \otimes H$  is an injective object. Since the morphism  $X \hookrightarrow I \otimes H$  defined by  $x \mapsto \sum_{(x)} \iota(x_0) \otimes x_1$  is injective,  ${}_R\mathcal{M}_T^H$  has enough injective objects.

(2) can be proved by repeatedly applying (1).  $\square$

For any  $H$ -comodule  $V$ ,  $R \otimes V \otimes T$  can be viewed as a Hopf module in  ${}_R\mathcal{M}_T^H$  via

$$(3.3) \quad r'(r \otimes v \otimes t)t' = r'r \otimes v \otimes tt', \quad \rho(r \otimes v \otimes t) = \sum_{(r),(v),(t)} (r_0 \otimes v_0 \otimes t_0) \otimes r_1 v_1 t_1,$$

for any  $v \in V$ ,  $r, r' \in R$  and  $t, t' \in S$ . One can readily verify that the functor  $R \otimes - \otimes T : \mathcal{M}^H \rightarrow {}_R\mathcal{M}_T^H$  serves as a left adjoint to the forgetful functor from  ${}_R\mathcal{M}_T^H$  to  $\mathcal{M}^H$ . As a result, there exists a natural isomorphism given by

$$(3.4) \quad \text{Hom}_{\mathcal{M}^H}(R \otimes V \otimes T, X) \longrightarrow \text{Hom}_{{}_R\mathcal{M}_T^H}(V, X), \quad f \mapsto (v \mapsto f(1_R \otimes v \otimes 1_T))$$

for any  $X \in {}_R\mathcal{M}_T^H$  and  $V \in \mathcal{M}^H$ . The inverse map is defined by  $g \mapsto (r \otimes v \otimes t \mapsto rg(v)t)$ .

Since  $R \otimes - \otimes T : {}_R\mathcal{M}_T^H \rightarrow {}_R\mathcal{M}_T^H$  and the forgetful functor  ${}_R\mathcal{M}_T^H \rightarrow \mathcal{M}^H$  are both exact, we have the following lemma.

**Lemma 3.3.** (1) If  $X$  is an injective object in  ${}_R\mathcal{M}_T^H$ , then  $X$  is injective  $H$ -comodule.  
(2) If  $V$  is a projective  $H$ -comodule, then  $R \otimes V \otimes T$  is a projective object in  ${}_R\mathcal{M}_T^H$ .

To any Hopf–Galois extension  $A = B^{coH} \subseteq B$ , one associates a  $A$ -coring  $B \otimes H$  and a bialgebroid  $L(B, B, H)$ , see [23] or [8, section 34]. In the remainder of this section,  $H$  is assumed to be a Hopf algebra with a bijective antipode, and  $R$  and  $B$  are considered as right  $H$ -comodule algebras. Put

$$L(R, B, H) := (R \otimes B)^{coH} = R \square_H B^{op}.$$

It is evident that  $L(R, B, H)$  forms a subalgebra of  $R \otimes B^{op}$ . We fix the formal notation

$$l = \sum l^1 \otimes l^2 \in R \otimes B^{op}$$

for  $l \in L(R, B, H)$ , so that  $ll' = \sum l^1 l'^1 \otimes l'^2 l^2$ .

In [23], Schauenburg extended Schneider's structure theorem for relative Hopf modules to a description of the structure of relative Hopf bimodules.

**Theorem 3.4.** [23, Theorem 3.3] Let  $H$  be a Hopf algebra and  $B$  a right faithfully flat  $H$ -Galois extension of  $A := B^{coH}$ . Consider a right  $H$ -comodule algebra  $R$ , and denote  $L := L(R, B, H)$ . There exists a category equivalence

$$F : {}_R\mathcal{M}_B^H \longrightarrow {}_L\mathcal{M}, \quad M \mapsto M^{coH},$$

with quasi-inverse

$$G : {}_L\mathcal{M} \longrightarrow {}_R\mathcal{M}_B^H, \quad N \mapsto N \otimes_A B.$$

The  $L$ -module structure of  $M^{coH}$  is defined by

$$l \triangleright m = \sum l^1 m l^2;$$

For  $N \in {}_L\mathcal{M}$ , the action  $\triangleright$  denotes the module structure, with  $N$  being a right  $A$ -module via  $na = (1 \otimes a) \triangleright n$ . The right  $B$ -module and  $H$ -comodule structures on  $N \otimes_A B$  are induced by those of  $B$ , while the left  $R$ -module structure is given by

$$(3.5) \quad r(n \otimes_A b) = \sum_{(r)} (r_0 \otimes \kappa^1(r_1)) \triangleright n \otimes_A \kappa^2(r_1)b.$$

**3.2. Some homological properties of  $L(R, B, H)$ .** In this subsection, we investigate the homological characteristics of the algebra  $L := L(R, B, H)$  arising from an  $H$ -comodule algebra  $R$  and a faithfully flat  $H$ -Galois extension  $A = B^{coH} \subseteq B$ .

**Lemma 3.5.** The algebra  $L$  is projective as a right  $A$ -module.

*Proof.* Since  $A \subseteq B$  is a faithfully flat  $H$ -Galois extension, there exists a right  $H$ -colinear and right  $A$ -linear splitting of the multiplication  $B \otimes A \rightarrow B$  by [25, Theorem 5.6]. Therefore,  $L = (R \otimes B)^{coH}$  is a direct summand of  $R \otimes A \otimes A_A = (R \otimes B \otimes A)^{coH}$  as a right  $A$ -module, implying that  $L$  is a projective  $A$ -module.  $\square$

For any  $M \in {}_R\mathcal{M}_B$ , the left  $L$ -module structure of  $M$  is obtained by restriction of scalars through the natural embedding map  $L \hookrightarrow R \otimes B^{op}$ . As a result,  $M \otimes_A B$  is identified as a Hopf bimodule within the category  ${}_R\mathcal{M}_B^H$ , as rigorously defined in Theorem 3.4.

**Lemma 3.6.** Let  $M$  be a  $R$ - $B$ -bimodule. Then  $M \otimes_A B \cong M \otimes H$  in  ${}_R\mathcal{M}_B^H$ , where the Hopf module structure on  $M \otimes H$  is given by (3.1).

*Proof.* Utilizing the bijectivity of the Galois map  $\beta : B \otimes_A B \rightarrow B \otimes H$ , the composition map

$$\varphi : M \otimes_A B \xrightarrow{\cong} M \otimes_B B \otimes_A B \xrightarrow{\text{id}_M \otimes \beta} M \otimes H, \quad m \otimes b \mapsto m \otimes 1_B \otimes b \mapsto \sum_{(b)} mb_0 \otimes b_1$$

is also bijective. To establish that  $\varphi$  is a morphism in  ${}_R\mathcal{M}_B^H$ , we recall that  $M \otimes_A B$  is a left  $R$ -module via

$$(3.6) \quad r(m \otimes_A b) \stackrel{(3.5)}{=} \sum_{(r)} (r_0 \otimes \kappa^1(r_1)) \triangleright m \otimes_A \kappa^2(r_1)b = \sum_{(r)} r_0 m \kappa^1(r_1) \otimes_A \kappa^2(r_1)b.$$

For any  $r \in R$ ,  $m \in M$  and  $b \in B$ , we obtain that  $\varphi(r(m \otimes_A b))$  is equal to

$$\begin{aligned} &= \varphi\left(\sum_{(r)} r_0 m \kappa^1(r_1) \otimes_A \kappa^2(r_1)b\right) \\ &= \sum_{(r), (\kappa^2(r_1)), (b)} r_0 m \kappa^1(r_1) \kappa^2(r_1)_0 b_0 \otimes \kappa^2(r_1)_1 b_1 \\ &\stackrel{(1.3)}{=} \sum_{(r), (b)} r_0 m \kappa^1(r_1) \kappa^2(r_1) b_0 \otimes r_2 b_1 \\ &\stackrel{(1.5)}{=} \sum_{(r), (b)} r_0 m b_0 \otimes r_1 b_1 \\ &\stackrel{(3.1)}{=} \sum_{(r)} r \varphi(m \otimes b). \end{aligned}$$

This verifies that  $\varphi$  is a left  $R$ -module morphism. Furthermore, it is straightforward to verify that  $\varphi$  also preserves  $B$ -module and  $H$ -comodule structures. Hence  $M \otimes_A B$  is isomorphic to  $M \otimes H$  as Hopf modules in  ${}_R\mathcal{M}_B^H$ .  $\square$

**Lemma 3.7.** For any left  $L$ -module  $V$ ,  $R \otimes B^{op}$ -module  $M$  and  $i \in \mathbb{N}$ ,

$$\mathrm{Ext}_L^i(V, M) \cong \mathrm{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, M).$$

In particular, if  $M = R \otimes B$ , then  $\mathrm{Ext}_L^i(V, R \otimes B) \cong \mathrm{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, R \otimes B)$  as  $B$ - $R$ -bimodules.

*Proof.* Let  $P_\bullet$  be a projective resolution of the left  $L$ -module  $V$ . Given that  $- \otimes_A B : {}_L\mathcal{M} \rightarrow {}_R\mathcal{M}_B^H$  constitutes an equivalence of categories, it follows that each  $P_i \otimes_A B$  is a projective object in  ${}_R\mathcal{M}_B^H$  and also a projective  $R \otimes B^{op}$ -module, as established by Lemma 3.1. Due to the flatness of  ${}_A B$ , the sequence

$$\cdots \longrightarrow P_1 \otimes_A B \longrightarrow P_0 \otimes_A B \longrightarrow V \otimes_A B \longrightarrow 0$$

is exact, making  $P_\bullet \otimes_A B$  a projective resolution of the left  $R \otimes B^{op}$ -module  $V \otimes_A B$ . By Lemma 3.6 and the adjoint isomorphism (3.2), we have the following isomorphisms,

$$\begin{aligned} \mathrm{Hom}_L(P_\bullet, M) &\cong \mathrm{Hom}_{{}_R\mathcal{M}_B^H}(P_\bullet \otimes_A B, M \otimes_A B) \\ &\cong \mathrm{Hom}_{{}_R\mathcal{M}_B^H}(P_\bullet \otimes_A B, M \otimes H) \\ &\cong \mathrm{Hom}_{R \otimes B^{op}}(P_\bullet \otimes_A B, M). \end{aligned}$$

Upon taking homologies, we derive the isomorphism

$$\mathrm{Ext}_L^i(V, M) \cong \mathrm{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, M)$$

for all  $i$ . The subsequent claim is implied by the naturality of this isomorphism.  $\square$

**Lemma 3.8.** If  $A \subseteq B$  be a faithfully flat  $H$ -Galois extension, then  $R \otimes V \otimes B$  is a projective object in  ${}_R\mathcal{M}_B^H$  for any  $H$ -comodule  $V$ .

*Proof.* Since all objects in  ${}_R\mathcal{M}_B^H$  are injective  $H$ -comodules by Lemma 1.2, any short exact sequence

$$0 \longrightarrow X' \longrightarrow X \longrightarrow X'' \longrightarrow 0$$

in  ${}_R\mathcal{M}_B^H$  splits as a sequence of  $H$ -comodules. Therefore, the induced sequence

$$0 \longrightarrow \mathrm{Hom}_{\mathcal{M}^H}(V, X') \longrightarrow \mathrm{Hom}_{\mathcal{M}^H}(V, X) \longrightarrow \mathrm{Hom}_{\mathcal{M}^H}(V, X'') \longrightarrow 0$$

is also exact. It follows from (3.4) that  $R \otimes V \otimes B$  is a projective object in  ${}_R\mathcal{M}_B^H$ .  $\square$

**Proposition 3.9.** Keep the notation as above. Then the following statements hold.

- (1)  $R \otimes B$  is a projective  $L$ -module.
- (2)  $L$  is a direct summand of  $R \otimes B$  as an  $L$ -module.

*Proof.* Given that the pair  $(- \otimes_A B, (-)^{coH})$  establishes an equivalence between  ${}_L\mathcal{M}$  and  ${}_R\mathcal{M}_B^H$  by Theorem 3.4, it suffices to prove the following:

- (1)  $R \otimes B \otimes_A B$  is a projective object in  ${}_R\mathcal{M}_B^H$ , and
- (2)  $L \otimes_A B$  is a direct summand of  $R \otimes B \otimes_A B$  within the category  ${}_R\mathcal{M}_B^H$ .

- (1) Observe that we have the following isomorphism in  ${}_R\mathcal{M}_B^H$ ,

$$R \otimes H \otimes B \xrightarrow{\cong} R \otimes B \otimes H, \quad r \otimes h \otimes b \mapsto \sum_{(r),(b)} r_0 \otimes b_0 \otimes r_1 h b_1,$$

with the inverse

$$R \otimes B \otimes H \longrightarrow R \otimes H \otimes B, \quad r \otimes b \otimes h \mapsto \sum_{(r),(b)} r_0 \otimes b_0 \otimes (S r_1) h (S^{-1} b_1).$$

The Hopf module structures of  $R \otimes H \otimes B$  and  $R \otimes B \otimes H$  are defined by (3.3) and (3.1), respectively. By Lemma 3.6, we have

$$R \otimes B \otimes H \cong R \otimes B \otimes_A B$$

in  ${}_R\mathcal{M}_B^H$ . Since  $R \otimes H \otimes B$  is a projective object in  ${}_R\mathcal{M}_B^H$  by Lemma 3.8, it follows that  $R \otimes B \otimes_A B$  is also a projective object.

(2) Consider the exact sequence of  $H$ -comodules,

$$0 \longrightarrow \mathbb{k} \longrightarrow H \longrightarrow \overline{H} := H/\mathbb{k}1_H \longrightarrow 0.$$

Applying the functor  $R \otimes - \otimes B$  yields an exact sequence

$$0 \longrightarrow R \otimes \mathbb{k} \otimes B \longrightarrow R \otimes H \otimes B \longrightarrow R \otimes \overline{H} \otimes B \longrightarrow 0.$$

in  ${}_R\mathcal{M}_B^H$ . By Lemma 3.8, this sequence splits, implying that  $R \otimes B (\cong R \otimes \mathbb{k} \otimes B)$  is a direct summand of  $R \otimes B \otimes_A B (\cong R \otimes H \otimes B)$  within  ${}_R\mathcal{M}_B^H$ .

Thus, the proof is concluded.  $\square$

**Lemma 3.10.** Let  $V$  be a left  $L$ -module.

- (1) If  $\text{Ext}_L^i(V, L) \neq 0$ , then  $\text{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, R \otimes B) \neq 0$ .
- (2) Suppose that  $V$  is an  $L$ -module of type  $\mathbf{FP}_\infty$ . If  $\text{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, R \otimes B) \neq 0$ , then  $\text{Ext}_L^i(V, L) \neq 0$ .

*Proof.* (1) If  $\text{Ext}_L^i(V, L) \neq 0$ , then  $\text{Ext}_L^i(V, R \otimes B) \neq 0$  as  $L$  is a direct summand of  $R \otimes B^{op}$ , per Lemma 3.9 (2). By Lemma 3.7, we obtain the isomorphism,

$$\text{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, R \otimes B) \cong \text{Ext}_L^i(V, R \otimes B).$$

Consequently,  $\text{Ext}_{R \otimes B^{op}}^i(V \otimes_A B, R \otimes B) \neq 0$  as well.

(2) As  $R \otimes B$  is a projective  $L$ -module by Lemma 3.9 (1), there exists a free  $L$ -module  $F$  with  $\text{Ext}_L^i(V, F) \neq 0$ . If  $V$  is an  $H$ -module of type  $\mathbf{FP}_\infty$ , then  $\text{Ext}_L^i(V, F)$  is isomorphic to a non-zero direct sum of some copies of  $\text{Ext}_L^i(V, L)$ . Therefore,  $\text{Ext}_L^i(V, L)$  is also non-zero.  $\square$

The following result is a well-established fact, as demonstrated in [6, Proposition VIII. 6.7] for instance.

**Lemma 3.11.** If  $M$  is an  $R$ -module of type  $\mathbf{FP}$ , then  $\text{pd}({}_R M) = \sup\{i \mid \text{Ext}_R^i(M, R) \neq 0\}$ .

*Proof.* Let  $\text{pd}({}_R M) = n < +\infty$ . Consequently, there exists an  $R$ -module  $N$  such that  $\text{Ext}_R^n(M, N) \neq 0$ . Now, let  $F$  be a free  $R$ -module with a surjective  $R$ -module morphism  $F \twoheadrightarrow N$ . Since  $\text{pd}({}_R M) = n$ , the induced morphism  $\text{Ext}_R^n(M, F) \rightarrow \text{Ext}_R^n(M, N)$  is surjective. Hence  $\text{Ext}_R^n(M, F) \neq 0$ . Since  $M$  is a  $R$ -module of type  $\mathbf{FP}$ , we have an isomorphism  $\text{Ext}_R^n(M, R) \otimes_R F \cong \text{Ext}_R^n(M, F)$ . It follows that  $\text{Ext}_R^n(M, R) \otimes_R F$  is also non-zero.  $\square$

**Lemma 3.12.** For any left  $L$ -module  $V$ ,  $\text{pd}_{R \otimes B^{op}}(V \otimes_A B) \leq \text{pd}({}_L V)$ . If further,  $\text{pd}({}_L V) < \infty$ , then  $\text{pd}_{R \otimes B^{op}}(V \otimes_A B) = \text{pd}({}_L V)$ .

*Proof.* Assume that  $\text{pd}({}_L V) = d < \infty$ . Let  $P_\bullet$  be a projective resolution of  $V$  of length  $d$ . Given that the functor  $- \otimes_A B : {}_L\mathcal{M} \rightarrow {}_R\mathcal{M}_B^H$  is an equivalence,  $P_\bullet \otimes_A B$  constitutes a projective resolution of  $V \otimes_A B$  in  ${}_R\mathcal{M}_B^H$ . Furthermore, each  $P_i \otimes_A B$  is a projective  $R \otimes B^{op}$ -module due to Lemma 3.1 (1). Consequently, the projective dimension of  $V \otimes_A B$  over  $R \otimes B^{op}$  is at most  $d$ , i.e.,  $\text{pd}_{R \otimes B^{op}}(V \otimes_A B) \leq d$ .

Since  $\text{Ext}_L^d(V, L) \neq 0$  by Lemma 3.11, the non-vanishing of  $\text{Ext}_{R \otimes B^{op}}^d(V \otimes_A B, R \otimes B^{op})$  follows from Lemma 3.10. Therefore, the projective dimension of  $V \otimes_A B$  as an  $R \otimes B^{op}$ -module is exactly  $d$ .  $\square$

The following proposition is an immediate consequence of Lemma 3.12

**Proposition 3.13.** If  $\text{lgld}(L) < \infty$ , then  $\text{lgld}(L) \leq \text{lgld}(R \otimes B^{op})$ .



**3.3. Injective dimension of the Ehresmann-Schauberg bialgebroids.** Indeed,  ${}_B\mathcal{M}_B^H$  is a monoidal category equipped with the tensor product over  $B$ . While  $L := L(B, B, H)$  is not generally a bialgebra,  ${}_L\mathcal{M}$  is nonetheless a monoidal category, its structure being induced by  ${}_B\mathcal{M}_B^H$ . In fact,  $L$  is a bialgebroid, as defined in [23]. This notion of a bialgebroid extends the concept of a bialgebra to a non-commutative base ring context.  $L(B, B, H)$  is recognized as the Ehresmann-Schauberg bialgebroid, as cited in [8, 34.14].

**Lemma 3.14.** Let  $M$  be a  $B^e$ -bimodule. If  $M_B$  or  ${}_B M$  is projective, then  $\text{pd}({}_{B^e}M) \leq \text{pd}({}_{B^e}B)$ .

*Proof.* Assuming  $\text{pd}({}_{B^e}B) = d < +\infty$ , consider a projective resolution  $P_\bullet$  of  ${}_{B^e}B$  with length  $d$ . If  $M_B$  is projective, then  $B^e \otimes_B M$  or equivalently  $B \otimes M$  becomes a projective  $B^e$ -module. Since  $P_\bullet$  splits as a complex of right  $B$ -modules,  $P_\bullet \otimes_B M$  serves as a projective resolution of  ${}_{B^e}M$ . Consequently,  $\text{pd}({}_{B^e}M) \leq d = \text{pd}_{B^e}(B)$ .  $\square$

**Lemma 3.15.** If  $V$  is a left  $L$ -module, then  $\text{pd}_{B^e}(V \otimes_A B) \leq \text{pd}({}_{B^e}B) + \text{rgld}(A)$ .

*Proof.* Without loss of generality, assume that  $\text{rgld}(A) = d < +\infty$ . Let  $V$  be a left  $L$ -module. Then there is an exact sequence of  $L$ -modules

$$0 \longrightarrow P_d \longrightarrow P_{d-1} \longrightarrow \cdots \longrightarrow P_0 \longrightarrow V \longrightarrow 0,$$

where each  $L$ -module  $P_i$  is projective for all  $i < d$ . Since  $L_A$  is projective by Lemma 3.5 and that  $\text{rgld}(A) = d$ , each  $P_i$  is projective as an  $A$ -module for all  $i \leq d$ . Consequently,  $P_i \otimes_A B$  is a projective  $B$ -module for all  $i \leq d$ . By Lemma 3.14,  $\text{pd}_{B^e}(P_i \otimes_A B) \leq \text{pd}({}_{B^e}B)$  for all  $i \leq d$ . Applying the functor  $- \otimes_A B$ , we obtain the following exact sequence of  $B^e$ -modules

$$0 \longrightarrow P_d \otimes_A B \longrightarrow P_{d-1} \otimes_A B \longrightarrow \cdots \longrightarrow P_0 \otimes_A B \longrightarrow V \otimes_A B \longrightarrow 0.$$

Therefore,  $\text{pd}({}_{B^e}(V \otimes_A B)) \leq \text{pd}({}_{B^e}B) + d = \text{pd}({}_{B^e}B) + \text{rgld}(A)$ .  $\square$

**Proposition 3.16.** Then  $\text{injdim}({}_L L) \leq \text{pd}({}_{B^e}B) + \text{rgld}(A) \leq \text{gldim}(H) + \text{pd}({}_{A^e}A) + \text{rgld}(A)$ .

*Proof.* By Lemma 3.10 (1) and 3.15, we have

$$\begin{aligned} \text{injdim}({}_L L) &= \sup\{i \mid \text{Ext}_L^i(V, L) \neq 0 \text{ for some left } L\text{-module } V\} \\ &\leq \sup\{i \mid \text{Ext}_{B^e}^i(V \otimes_A B, B^e) \neq 0 \text{ for some left } L\text{-module } V\} \\ &\leq \sup\{\text{pd}_{B^e}(V \otimes_A B) \mid \text{for some left } L\text{-module } V\} \\ &\leq \text{pd}({}_{B^e}B) + \text{rgld}(A). \end{aligned}$$

The second inequality follows from Corollary 1.5.  $\square$

#### 4. HOMOLOGICAL PROPERTIES UNDER MONOIDAL MORITA-TAKEUCHI EQUIVALENCES

In this section, we focus specifically on the case where  $A = \mathbb{k}$ . As established in [22], under this condition,  $L := L(B, B, H)$  becomes a Hopf algebra. The comultiplication of  $L$  is given by

$$\Delta(l) = \sum_{(l^1)} l^1_0 \otimes \kappa^1(l^1_1) \otimes \kappa^2(l^1_1) \otimes l^2,$$

where  $l = \sum l^1 \otimes l^2 \in L \subseteq B^e$ . The counit of  $L$  is defined as  $\varepsilon(l) = \sum l^1 l^2$ , and the antipode is given by  $S(l) = \sum l^2_0 \otimes \kappa^1(l^2_1) l^1 \kappa^2(l^2_1)$ .

Furthermore, as detailed in [22],  $B$  is endowed with a left  $L$ -comodule algebra structure via the comodule map  $\delta : B \rightarrow L \otimes B$  defined as

$$(4.1) \quad b \mapsto \sum_{(b)} (b_0 \otimes \kappa^1(b_1)) \otimes \kappa^2(b_1).$$

This structure confers upon  $B$  the property of being an  $L$ - $H$ -biGalois extension of  $\mathbb{k}$ . In other words,  $B$  is both a right  $H$ -Galois extension of  $\mathbb{k}$  and a left  $L$ -Galois of  $\mathbb{k}$  such that the two comodule structures make it an  $L$ - $H$ -bicomodule.

**4.1. Monoidal Morita-Takeuchi equivalences preserve the type  $\mathbf{FP}_\infty$ .** In the following, we demonstrate that the type  $\mathbf{FP}_\infty$  of Hopf algebras is an invariant of monoidal Morita-Takeuchi equivalences.

Recall that for any  $H$ -comodule  $V$ , by (3.3),  $B \otimes V \otimes B$  naturally acquires a structure of a Hopf bimodule in the category  ${}_B\mathcal{M}_B^H$ . The  $B$ - $B$ -bimodule structure is induced by the first and third tensorands, and the  $H$ -comodule structure arises from the diagonal coaction. Consequently, the functor  $B \otimes - \otimes B$  serves as a left adjoint of the forgetful functor from  $\mathcal{M}^H$  to  ${}_B\mathcal{M}_B^H$ , established by the existence of the following natural isomorphism

$$(4.2) \quad \mathrm{Hom}_{\mathcal{M}^H}(V, M) \longrightarrow \mathrm{Hom}_{{}_B\mathcal{M}_B^H}(B \otimes V \otimes B, M), \quad f \mapsto (b \otimes v \otimes b' \mapsto bf(v)b')$$

for any  $V \in \mathcal{M}^H$  and  $M \in {}_B\mathcal{M}_B^H$ .

**Lemma 4.1.** Let  $M$  be a Hopf module in  ${}_B\mathcal{M}_B^H$ .

(1) Then  $M$  is finitely generated as a  $B^e$ -module if and only if there is a finite-dimensional  $H$ -comodule  $V$  with a surjective morphism  $\pi : B \otimes V \otimes B \rightarrow M$  in  ${}_B\mathcal{M}_B^H$ .

(2) If  $M$  is a  $B^e$ -module of type  $\mathbf{FP}_n$ , then there exists an exact sequence

$$\cdots \longrightarrow B \otimes V_i \otimes B \longrightarrow \cdots \longrightarrow B \otimes V_0 \otimes B \longrightarrow M \longrightarrow 0 \quad \text{in } {}_B\mathcal{M}_B^H,$$

where each  $V_i$  is an  $H$ -comodules for  $i \geq 0$  and  $V_i$  is finite-dimensional for  $i \leq n$ .

*Proof.* (1) Let  $\{x_1, \dots, x_m\}$  be a set generators of  ${}_B^e M$ . The Finiteness Theorem [19, 5.1.1] guarantees the existence of a finite-dimensional  $H$ -subcomodule  $V$  of  $M$  that includes  $\{x_1, \dots, x_m\}$ . Consequently, there is a surjective morphism  $\pi : B \otimes V \otimes B \rightarrow M$ ,  $b \otimes v \otimes b' \mapsto bv b'$  in  ${}_B\mathcal{M}_B^H$ .

(2) From (1), there exists a finite-dimensional left  $H$ -comodule  $V_0$  with a surjective morphism  $\pi : B \otimes V_0 \otimes B \rightarrow M$  in  ${}_B\mathcal{M}_B^H$ . Schanuel's lemma implies that the kernel of  $\pi$  is a  $B^e$ -module of type  $\mathbf{FP}_{n-1}$ . The desired conclusion then follows by induction.  $\square$

The subsequent lemma is a well-established fact, and we present a concise proof for it.

**Lemma 4.2.** Let  $M$  be a module over a ring  $R$ . If  $\varinjlim \mathrm{Hom}_R(M, N_i) = 0$  for any direct system  $\{N_i\}$  with  $\varinjlim N_i = 0$ , then  $M$  is finitely generated.

*Proof.* Consider the direct system  $\{M/M'\}$  indexed by the finitely generated submodules  $M'$  of  $M$ . We have  $\varinjlim M/M' = 0$ , which implies  $\varinjlim \mathrm{Hom}_R(M, M/M') = 0$ . Consequently, there exists a finitely generated submodule  $M'$  for which the induced map  $\overline{\mathrm{id}_M}$  is the zero map in  $\mathrm{Hom}_R(M, M/M')$ , where  $\mathrm{id}_M$  denotes the identity map on  $M$ . Therefore, we conclude that  $M = M'$ , establishing that  $M$  is a finitely generated  $R$ -module.  $\square$

**Lemma 4.3.** If  $V$  is a finite dimensional  $H$ -comodule, then  $(B \otimes V \otimes B)^{coH}$  is a finitely generated projective  $L$ -module.

*Proof.* Recall that the functor  $(-)^{coH} : {}_B\mathcal{M}_B^H \rightarrow {}_L\mathcal{M}$  establishes an equivalence. Given that  $B \otimes V \otimes B$  is a projective object in  ${}_B\mathcal{M}_B^H$  by Lemma 3.8,  $(B \otimes V \otimes B)^{coH}$  is a projective  $L$ -module.

To demonstrate that  $(B \otimes V \otimes B)^{coH}$  is finitely generated, we utilize Lemma 4.2, which asserts that it suffices to show that the functor  $\mathrm{Hom}_L((B \otimes V \otimes B)^{coH}, -)$  commutes with direct limits. In other words, for any direct system  $\{N_i\}$  of  $L$ -modules, there must exist a natural isomorphism

$$\mathrm{Hom}_L((B \otimes V \otimes B)^{coH}, \varinjlim N_i) \cong \varinjlim \mathrm{Hom}_L((B \otimes V \otimes B)^{coH}, N_i).$$

By Theorem 3.4, we establish the following isomorphisms

$$\begin{aligned} \mathrm{Hom}_L((B \otimes V \otimes B)^{coH}, \varinjlim N_i) &\cong \mathrm{Hom}_{{}_B\mathcal{M}_B^H}((B \otimes V \otimes B), (\varinjlim N_i) \otimes B) \\ &\cong \mathrm{Hom}_{{}_B\mathcal{M}_B^H}(B \otimes V \otimes B, \varinjlim (N_i \otimes B)) \end{aligned}$$

$$\stackrel{(4.2)}{\cong} \text{Hom}_{\mathcal{M}^H}(V, \varinjlim(N_i \otimes B)).$$

By the definition of direct limit, we have an exact sequence

$$\oplus(N_i \otimes B) \longrightarrow \oplus(N_i \otimes B) \longrightarrow \varinjlim(N_i \otimes B) \longrightarrow 0$$

in  ${}_B\mathcal{M}_B^H$ , which is split exact in  $\mathcal{M}^H$  by Lemma 1.2. Hence we have a commutative diagram with exact rows:

$$\begin{array}{ccccccc} \text{Hom}_{\mathcal{M}^H}(V, \oplus(N_i \otimes B)) & \longrightarrow & \text{Hom}_{\mathcal{M}^H}(V, \oplus(N_i \otimes B)) & \longrightarrow & \text{Hom}_{\mathcal{M}^H}(V, \varinjlim(N_i \otimes B)) & \longrightarrow & 0 \\ \downarrow \pi & & \downarrow \pi & & \downarrow & & \\ \oplus \text{Hom}_{\mathcal{M}^H}(V, N_i \otimes B) & \longrightarrow & \oplus \text{Hom}_{\mathcal{M}^H}(V, N_i \otimes B) & \longrightarrow & \varinjlim \text{Hom}_{\mathcal{M}^H}(V, N_i \otimes B) & \longrightarrow & 0 \end{array}$$

Since  $V$  is finite dimensional, the canonical map  $\pi$  is bijective. This yields an isomorphism between  $\text{Hom}_{\mathcal{M}^H}(V, \varinjlim(N_i \otimes B))$  and  $\varinjlim \text{Hom}_{\mathcal{M}^H}(V, N_i \otimes B)$ . Therefore, we obtain the following isomorphisms,

$$\begin{aligned} \text{Hom}_L((B \otimes V \otimes B)^{coH}, \varinjlim N_i) &\cong \varinjlim \text{Hom}_{\mathcal{M}^H}(V, N_i \otimes B) \\ &\stackrel{(4.2)}{\cong} \varinjlim \text{Hom}_{{}_B\mathcal{M}_B^H}(B \otimes V \otimes B, N_i \otimes B) \\ &\cong \varinjlim \text{Hom}_L((B \otimes V \otimes B)^{coH}, N_i). \end{aligned}$$

In conclusion,  $(B \otimes V \otimes B)^{coH}$  is a finitely generated projective  $L$ -module.  $\square$

Recall that by (3.5) and (4.1), for any left  $L$ -module  $V$ ,  $V \otimes B$  acquires a left  $B^e$ -bimodule structure defined by

$$(b' \otimes b'')(v \otimes b) = \sum_{(b')} b'_1 v \otimes b'_0 b b''.$$

**Lemma 4.4.** Let  $V$  be a left  $L$ -module.

- (1) If  $V$  is projective, then  $V \otimes B$  is a projective  $B^e$ -module.
- (2) If  $V$  is finitely generated, then  $V \otimes B$  is also finitely generated as a  $B^e$ -module.

*Proof.* (1) If  $V$  is projective, then it is a direct summand of some copies of  $L$ . Given that the Galois map  $\beta' : B \otimes B \rightarrow L \otimes B$  is an isomorphism of  $B^e$ -modules,  $V \otimes B$  is also a projective  $B^e$ -module.

(2) If  $V$  is finitely generated, then there exists an integer  $n$  with a surjective morphism  $L^{\oplus n} \twoheadrightarrow V$ . Consequently,  $(B \otimes B)^{\oplus n} \cong L^{\oplus n} \otimes B \twoheadrightarrow V \otimes B$  is a surjective morphism of  $B^e$ -modules, confirming that  $V \otimes B$  is finitely generated.  $\square$

**Proposition 4.5.** Let  $H$  be a Hopf algebra with a bijective antipode, and  $B$  be an  $H$ -Galois extension of  $\mathbb{k}$ . For any  $0 \leq n \leq \infty$ ,  $H$  is of type  $\mathbf{FP}_n$  if and only if  $B$  is a  $B^e$ -module of type  $\mathbf{FP}_n$ .

*Proof.* “ $\Rightarrow$ ” Given that  $H$  is of type  $\mathbf{FP}_n$ , there exists a projective resolution  $P_\bullet$  of  $\mathbb{k}_H$  with each  $P_i$  is finitely generated for  $i \leq n$ . By Lemma 4.4, tensoring  $P_\bullet$  with  $B$  yields a projective resolution of the  $B^e$ -module  $B = \mathbb{k} \otimes B$ , where each  $P_i \otimes B$  is finitely generated for  $i \leq n$ . Therefore,  $B$  is a  $B^e$ -module of type  $\mathbf{FP}_n$ .

“ $\Leftarrow$ ” Since  $B$  is a  $B^e$ -module of type  $\mathbf{FP}_n$ , Lemma 4.1 guarantees the existence of an exact sequence in the category  ${}_B\mathcal{M}_B^H$  of the form

$$\cdots \longrightarrow B \otimes V_i \otimes B \longrightarrow \cdots \longrightarrow B \otimes V_0 \otimes B \longrightarrow B \longrightarrow 0 \quad \text{in } {}_B\mathcal{M}_B^H,$$

where each  $V_i$  is an  $H$ -comodules for  $i \geq 0$  and  $V_i$  is finite-dimensional for  $i \leq n$ . Applying the functor of  $H$ -coinvariants  $(-)^{coH} : {}_B\mathcal{M}_B^H \rightarrow {}_L\mathcal{M}$  yields an exact sequence

$$\cdots \longrightarrow (B \otimes V_i \otimes B)^{coH} \longrightarrow \cdots \longrightarrow (B \otimes V_0 \otimes B)^{coH} \longrightarrow B^{coH} \longrightarrow 0$$

in  ${}_L\mathcal{M}$ . Here,  $B^{coH} = \mathbb{k}$  is the trivial  $L$ -module. By Lemma 4.3,  $(B \otimes V_i \otimes B)^{coH}$  is always finitely generated as an  $L$ -module for all  $i \geq 0$ . Thus,  $H$  is of type  $\mathbf{FP}_n$ .  $\square$

**Theorem 4.6.** Let  $H$  and  $L$  be two monoidally Morita-Takeuchi equivalent Hopf algebras with bijective antipodes. If  $H$  is of type  $\mathbf{FP}_\infty$ , then so is  $L$ .

*Proof.* Let  $B$  be an  $L$ - $H$ -biGalois extension of  $\mathbb{k}$ . Since the antipode  $S$  of  $L$  is bijective as we assumed,  $L^{cop}$  is also a Hopf algebra with opposite comultiplication antipode  $S^{-1}$ , and that  $B$  is a right  $L^{cop}$ -Galois extension of  $\mathbb{k}$ . Then the conclusion follows from Proposition 4.5.  $\square$

**4.2. AS Gorenstein property under monoidal Morita-Takeuchi equivalence.** In the subsequent portion of this section, our objective is to establish a parallel version of Theorem 0.3 for the scenario where Hopf algebras possess finite injective dimension, rather than finite global dimension. The ultimate goal is to substitute the AS regular condition with the AS Gorenstein condition.

According to (1.6),  $B \otimes B$  serves as a right  $B \otimes H$ -module via the action defined by

$$(x \otimes y)(b \otimes h) = \sum \kappa^1(h)xb \otimes y\kappa^2(h).$$

**Lemma 4.7.** Let  $H$  be a Hopf algebra and  $B$  be a right  $H$ -Galois extension of  $\mathbb{k}$ . Then  $B \otimes B \cong B \otimes H$  as right  $B \otimes H$ -modules.

*Proof.*  $B \otimes H$  is an  $H$ -comodule algebra, arising from the coaction  $b \otimes h \mapsto \sum_{(h)} b \otimes h_1 \otimes h_2$ . Verifying that  $B \subseteq B \otimes H$  is a faithfully flat  $H$ -Galois extension is straightforward. Consequently, an equivalence functor  $(-\otimes_B(B \otimes H), (-)^{coH})$  exists between  $\mathcal{M}_B$  and  $\mathcal{M}_{B \otimes H}^H$ . Computing  $(\text{id}_B \otimes \Delta)((x \otimes y)(b \otimes h))$  yields

$$\begin{aligned} &= \sum_{(y), (\kappa^2(h))} \kappa^1(h)xb \otimes y_0\kappa^2(h)_0 \otimes y_1\kappa^2(h)_1 \\ &\stackrel{(1.3)}{=} \sum_{(y), (\kappa^2(h))} \kappa^1(h_1)xb \otimes y_0\kappa^2(h_1) \otimes y_1h_2 \\ &= \sum_{(y), (h)} ((x \otimes y_0)(b \otimes h_1)) \otimes y_1h_2, \end{aligned}$$

This establishes that  $B \otimes B$  is a Hopf bimodule in the category  $\mathcal{M}_{B \otimes H}^H$ . Since  $(B \otimes B)^{coH} = B \otimes B^{coH} \cong B$  as right  $B$ -modules, it follows that

$$B \otimes B \cong (B \otimes B)^{coH} \otimes_B (B \otimes H) = B \otimes H$$

as right  $B \otimes H$ -modules, completing the argument.  $\square$

**Lemma 4.8.** [31, Lemma 2.2] [29, Lemma 2.3.1] Let  $H$  be a Hopf algebra of type  $\mathbf{FP}_\infty$  with a bijective antipode, and  $B$  be a  $H$ -Galois extension of  $\mathbb{k}$ . Then

$$B_B \otimes \text{Ext}_{H^{op}}^i(\mathbb{k}, H) \cong \text{Ext}_{B^e}^i(B, B^e)$$

as right  $B$ -modules for all  $i$ .

*Proof.* Since  $\mathbb{k}_H$  is of type  $\mathbf{FP}_\infty$ , we have

$$\text{Ext}_{H^{op}}^i(\mathbb{k}, B \otimes H) \cong (B \otimes H) \otimes_H \text{Ext}_{H^{op}}^i(\mathbb{k}, H) \cong B \otimes \text{Ext}_{H^{op}}^i(\mathbb{k}, H)$$

for all  $i$ . On the other hand, by Lemma 4.7,  $B^e \cong B \otimes H$  as right  $B \otimes H$ -modules. Consequently, applying Lemma 1.4, we obtain

$$\mathrm{Ext}_{H^{op}}^i(\mathbb{k}, B \otimes H) \cong \mathrm{Ext}_{H^{op}}^i(\mathbb{k}, B^e) \cong \mathrm{Ext}_{B^e}^i(B, B^e)$$

for all  $i$ . By combining the above isomorphisms, we arrive at the desired conclusion.  $\square$

**Theorem 4.9.** Let  $H$  be a Hopf algebra of type  $\mathbf{FP}_\infty$  with a bijective antipode, and  $B$  be a  $H$ -Galois extension of  $\mathbb{k}$ .

- (1) [2, Theorem 7] If  $\mathrm{gldim}(H) = d < \infty$ , then  $\mathrm{pd}_{(B^e B)} = d$ .
- (2) [31, Theorem 2.5] If  $H$  is AS regular of dimension  $d$ , then  $B$  is skew Calabi-Yau of dimension  $d$ .
- (3) If  $\mathrm{pd}_{(B^e B)} = d$ , then the injective dimension  $\mathrm{injdim}(H) = d$ .
- (4) If  $B$  is skew Calabi-Yau of dimension  $d$ , then  $H$  is AS Gorenstein of dimension  $d$ .

*Proof.* (1) Since  $\mathbb{k}_H$  is of type  $\mathbf{FP}_\infty$ , we can deduce from Lemma 3.11 that  $\mathrm{Ext}_{H^{op}}^d(\mathbb{k}, H) \neq 0$ . Hence  $\mathrm{Ext}_{B^e}^d(B, B^e) \neq 0$  by Lemma 4.8. Since  $\mathrm{pd}_{(B^e B)} \leq \mathrm{pd}(\mathbb{k}_H) = d$  by Corollary 1.5, it follows that  $\mathrm{pd}_{(B^e B)} = d$ .

(2) By Lemma 4.8,

$$\mathrm{Ext}_{B^e}^d(B, B^e) \cong \begin{cases} 0, & i \neq d, \\ B \otimes \mathrm{Ext}_{H^{op}}^d(\mathbb{k}, H) \cong B, & i = d. \end{cases}$$

as right  $B$ -modules. Similarly, we can prove that  $\mathrm{Ext}_{B^e}^d(B, B^e) \cong B$  as left  $B$ -modules. Hence there exists an automorphism  $\mu$  of  $B$  such that  $\mathrm{Ext}_{B^e}^d(B, B^e) \cong B_\mu$ . So  $B$  is skew Calabi-Yau of dimension  $d$ .

(3) By Lemma 4.5,  $B^e B$  is of type  $\mathbf{FP}_\infty$ , which ensures that  $\mathrm{Ext}_{B^e}^d(B, B^e) \neq 0$  according to Lemma 3.11. Therefore, Lemma 4.8 implies that  $\mathrm{Ext}_{H^{op}}^d(\mathbb{k}, H) \neq 0$ . On the other hand, since  $\mathrm{injdim}({}_L L) \leq d$  by Proposition 3.16, it follows that  $\mathrm{injdim}({}_L L) = d$ .

(4) If  $B$  is skew Calabi-Yau of dimension  $d$ , then by Lemma 4.8

$$B \otimes \mathrm{Ext}_H^i(\mathbb{k}, H) \cong \mathrm{Ext}_{B^e}^d(B, B^e) \cong \begin{cases} 0, & i \neq d, \\ B, & i = d. \end{cases}$$

as right  $B$ -modules for all  $i$ . Hence  $\mathrm{Ext}_{H^{op}}^d(\mathbb{k}, H)$  is finite dimensional over  $\mathbb{k}$ , see [13, Corollary 1.2] for example. Therefore, Lemma 2.4 implies that  $H$  is AS Gorenstein of dimension  $d$ .  $\square$

**Theorem 4.10.** Let  $H$  and  $L$  be two monoidally Morita-Takeuchi equivalent Hopf algebras of type  $\mathbf{FP}_\infty$ . Suppose that the antipodes of  $H$  and  $L$  are both bijective.

- (1) If  $H$  has finite global dimension  $d$ , then the injective dimension  $\mathrm{injdim}({}_L L) = d$ .
- (2) If  $H$  is AS regular of dimension  $d$ , then  $L$  is AS Gorenstein of dimension  $d$ .
- (3) If  $L$  and  $H$  are both AS Gorenstein, then  $\mathrm{injdim}(L) = \mathrm{injdim}(H)$ .

*Proof.* Observing that  $B$  is a right  $L^{cop}$ -Galois extension of  $\mathbb{k}$ , we can deduce (1) and (2) directly from Theorem 4.9.

(3) According to Lemma 4.8, we have

$$\mathrm{Ext}_{L^{op}}^i(\mathbb{k}, L) \neq 0 \text{ if and only if } \mathrm{Ext}_{B^e}^i(B, B^e) \neq 0 \text{ if and only if } \mathrm{Ext}_{H^{op}}^i(\mathbb{k}, H) \neq 0.$$

Since  $L$  and  $H$  are AS Gorenstein by our assumption, it follows that  $\mathrm{injdim}(L) = \mathrm{injdim}(H)$ .  $\square$

### 4.3. Monoidal Morita-Takeuchi equivalences don't preserve the global dimension.

We conclude our discussion by presenting a counterexample to Question 0.4.

**Example 4.11.** The Liu Hopf algebra  $H := L(n, \xi)$  [16] is generated by  $x, g, g^{-1}$  with the relations

$$gg^{-1} = g^{-1}g = 1, \quad xg = \xi gx, \quad x^n = 1 - g^n,$$

where  $\xi \in \mathbb{k}$  is an  $n$ -th primitive root of unity. The comultiplication, counit and antipode on  $H$  are defined by

$$\Delta(g) = g \otimes g, \quad \Delta(x) = x \otimes 1 + g \otimes x, \quad \varepsilon(g) = 1, \quad \varepsilon(x) = 0, \quad S(g) = g^{-1}, \quad S(x) = -g^{-1}x.$$

Let  $B := \mathbb{k}\langle g, g^{-1}, t \rangle / \langle gg^{-1} - 1, g^{-1}g - 1, tg - \xi gt, t^n + g^n \rangle$ . It serves as a right  $H$ -comodule algebra through the comodule map  $\rho$  defined by

$$\rho(g) = g \otimes g, \quad \rho(t) = t \otimes 1 + g \otimes x.$$

The bijectivity of the  $H$ -comodule map  $\gamma : H \rightarrow B$  given by  $g^i x^j \mapsto g^i t^j$  implies its convolution invertibility with the inverse  $\gamma S$ . Consequently,  $B$  is an  $H$ -cleft extension of  $\mathbb{k}$ , and thus a right  $H$ -Galois extension of  $\mathbb{k}$  according to [19, Theorem 8.2.4].

Let  $L$  be the Hopf algebra which is generated by  $y, g, g^{-1}$  with the relations

$$gg^{-1} = g^{-1}g = 1, \quad yg = \xi gy, \quad y^n = 0.$$

The comultiplication, counit and antipode on  $L$  are defined by

$$\Delta(g) = g \otimes g, \quad \Delta(y) = 1 \otimes y + y \otimes g, \quad \varepsilon(g) = 1, \quad \varepsilon(y) = 0, \quad S(g) = g^{-1}, \quad S(y) = -yg^{-1}.$$

$B$  is also a left  $L$ -comodule algebra, defined through the comodule map  $\delta$  with  $\delta(g) = g \otimes g$  and  $\delta(t) = g \otimes t + y \otimes 1$ . Furthermore,  $B$  qualifies as a left  $L$ -Galois extension of  $\mathbb{k}$ . Verifying that  $B$  is an  $L$ - $H$ -bicomodule is straightforward, leading to the conclusion that  $B$  is an  $L$ - $H$ -biGalois extension of  $\mathbb{k}$ .

The global dimension of  $H$  is 1, as stated in [16, Lemma 2.4], whereas the global dimension of  $L$  is infinite. This observation underscores that monoidal Morita-Takeuchi equivalences do not consistently preserve the global dimension. Nevertheless, since  $H$  and  $L$  are both noetherian affine PI Hopf algebras, they share the same injective dimension, as implied by Corollary 0.8.

#### ACKNOWLEDGMENTS

The research work was supported by the NSFC (project 12301052) and Fundamental Research Funds for the Central Universities (2022110884).

#### REFERENCES

- [1] J. BICHON, *Gerstenhaber-Schack and Hochschild cohomologies of Hopf algebras*, Doc. Math., 21 (2016), pp. 955–986.
- [2] ———, *On the monoidal invariance of the cohomological dimension of Hopf algebras*, C. R. Math. Acad. Sci. Paris, 360 (2022), pp. 561–582.
- [3] K. A. BROWN, *Representation theory of Noetherian Hopf algebras satisfying a polynomial identity*, in Trends in the representation theory of finite-dimensional algebras (Seattle, WA, 1997), vol. 229 of Contemp. Math., Amer. Math. Soc., Providence, RI, 1998, pp. 49–79.
- [4] K. A. BROWN AND K. R. GOODEARL, *Homological aspects of Noetherian PI Hopf algebras of irreducible modules and maximal dimension*, J. Algebra, 198 (1997), pp. 240–265.
- [5] K. A. BROWN AND J. J. ZHANG, *Dualising complexes and twisted Hochschild (co)homology for Noetherian Hopf algebras*, J. Algebra, 320 (2008), pp. 1814–1850.
- [6] K. S. BROWN, *Cohomology of groups*, vol. 87 of Graduate Texts in Mathematics, Springer-Verlag, New York-Berlin, 1982.
- [7] T. BRZEZIŃSKI, *Translation map in quantum principal bundles*, J. Geom. Phys., 20 (1996), pp. 349–370.
- [8] T. BRZEZIŃSKI AND R. WISBAUER, *Corings and comodules*, vol. 309 of London Mathematical Society Lecture Note Series, Cambridge University Press, Cambridge, 2003.
- [9] D. ŞTEFAN, *Hochschild cohomology on Hopf Galois extensions*, J. Pure Appl. Algebra, 103 (1995), pp. 221–233.
- [10] Y. DOI, *Algebras with total integrals*, Comm. Algebra, 13 (1985), pp. 2137–2159.
- [11] Y. DOI AND M. TAKEUCHI, *Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and Azumaya algebras*, J. Algebra, 121 (1989), pp. 488–516.
- [12] P. JARA AND D. ŞTEFAN, *Hopf-cyclic homology and relative cyclic homology of Hopf-Galois extensions*, Proc. London Math. Soc. (3), 93 (2006), pp. 138–174.

- [13] T. Y. LAM, *Lectures on modules and rings*, vol. 189 of Graduate Texts in Mathematics, Springer-Verlag, New York, 1999.
- [14] R. G. LARSON AND M. E. SWEEDLER, *An associative orthogonal bilinear form for Hopf algebras*, Amer. J. Math., 91 (1969), pp. 75–94.
- [15] P. LE MEUR, *Smash products of Calabi-Yau algebras by Hopf algebras*, J. Noncommut. Geom., 13 (2019), pp. 887–961.
- [16] G. LIU, *On Noetherian affine prime regular Hopf algebras of Gelfand-Kirillov dimension 1*, Proc. Amer. Math. Soc., 137 (2009), pp. 777–785.
- [17] J. LÜ, S.-Q. OH, X. WANG, AND X. YU, *A note on the bijectivity of the antipode of a Hopf algebra and its applications*, Proc. Amer. Math. Soc., 146 (2018), pp. 4619–4631.
- [18] J. C. MCCONNELL AND J. C. ROBSON, *Noncommutative Noetherian rings*, vol. 30 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, revised ed., 2001. With the cooperation of L. W. Small.
- [19] S. MONTGOMERY, *Hopf algebras and their actions on rings*, vol. 82 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993.
- [20] M. REYES, D. ROGALSKI, AND J. J. ZHANG, *Skew Calabi-Yau algebras and homological identities*, Adv. Math., 264 (2014), pp. 308–354.
- [21] D. ROGALSKI, R. WON, AND J. J. ZHANG, *A proof of the Brown-Goodearl conjecture for module-finite weak Hopf algebras*, Algebra Number Theory, 15 (2021), pp. 971–997.
- [22] P. SCHAUENBURG, *Hopf bi-Galois extensions*, Comm. Algebra, 24 (1996), pp. 3797–3825.
- [23] ———, *Bialgebras over noncommutative rings and a structure theorem for Hopf bimodules*, Appl. Categ. Structures, 6 (1998), pp. 193–222.
- [24] ———, *Faithful flatness over Hopf subalgebras: counterexamples*, in Interactions between ring theory and representations of algebras (Murcia), vol. 210 of Lecture Notes in Pure and Appl. Math., Dekker, New York, 2000, pp. 331–344.
- [25] P. SCHAUENBURG AND H.-J. SCHNEIDER, *On generalized Hopf Galois extensions*, J. Pure Appl. Algebra, 202 (2005), pp. 168–194.
- [26] H.-J. SCHNEIDER, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math., 72 (1990), pp. 167–195.
- [27] S. SKRYABIN, *New results on the bijectivity of antipode of a Hopf algebra*, J. Algebra, 306 (2006), pp. 622–633.
- [28] M. TAKEUCHI, *Free Hopf algebras generated by coalgebras*, J. Math. Soc. Japan, 23 (1971), pp. 561–582.
- [29] X. WANG, X. YU, AND Y. ZHANG, *Calabi-Yau property under monoidal Morita-Takeuchi equivalence*, Pacific J. Math., 290 (2017), pp. 481–510.
- [30] Q.-S. WU AND J. J. ZHANG, *Noetherian PI Hopf algebras are Gorenstein*, Trans. Amer. Math. Soc., 355 (2003), pp. 1043–1066.
- [31] X. YU, *Hopf-Galois objects of Calabi-Yau Hopf algebras*, J. Algebra Appl., 15 (2016), pp. 1650194, 19.

SCHOOL OF MATHEMATICS, SHANGHAI UNIVERSITY OF FINANCE AND ECONOMICS, SHANGHAI 200433, CHINA

*Email address:* zhuruipeng@sufe.edu.cn