SKEW CALABI-YAU PROPERTY OF FAITHFULLY FLAT HOPF GALOIS EXTENSIONS

RUIPENG ZHU

ABSTRACT. This paper shows that if H is a Hopf algebra and $A \subseteq B$ is a faithfully flat H -Galois extension, then B is skew Calabi-Yau provided A and H are. Specifically, for cleft extensions $A \subseteq B$, the Nakayama automorphism of B can be derived from those of A and H , along with the homological determinant of the H -action on A . This finding is based on the study of the Hopf bimodule structure on $\mathrm{Ext}^i_{A^e}(A,B^e).$

INTRODUCTION

Calabi-Yau algebras, introduced by Ginzburg [\[13\]](#page-40-0), serve as the noncommutative counterparts of coordinate rings for smooth affine Calabi-Yau varieties. Their study has been motivated by applications in algebraic geometry, representation theory, and mathematical physics. Skew Calabi-Yau algebras encompass Artin–Schelter regular algebras [\[2\]](#page-40-1) and noetherian rigid Gorenstein Hopf algebras of finite global dimension [\[5\]](#page-40-2), sharing homological properties akin to Calabi-Yau algebras. Each skew Calabi-Yau algebra is associated with a Nakayama automorphism, which is unique up to composition with an inner automorphism. A skew Calabi–Yau algebra is Calabi-Yau if and only if its Nakayama automorphism is inner.

Numerous studies have explored the Nakayama automorphism and its utilities, see $[8, 23]$ $[8, 23]$ $[8, 23]$ for example. Informally, $[8]$ shows that the Nakayama automorphism governs certain characteristics of the class of Hopf algebras that act on a specified connected graded skew Calabi-Yau algebra. As established in [\[23,](#page-41-0) Theorem 4.2], the Nakayama automorphism plays a pivotal role as a central element in the Picard group. Consequently, gaining a comprehensive understanding and providing an explicit description of the Nakayama automorphism is crucial, as noted in [\[23\]](#page-41-0). Simultaneously, the Nakayama automorphism is a subtle invariant that can be challenging to determine.

Identifying a skew Calabi-Yau algebra can also be difficult. To address this, we propose a somewhat more practical criterion to facilitate its determination.

Proposition 0.1. (Proposition [1.7\)](#page-5-0) Let A be a homologically smooth algebra. If

(1) $\mathrm{Ext}^i_{A^e}(A, A^e) = 0$ for all $i \neq d$, and

(2) $\text{Ext}_{A^e}^d(A, A^e)$ is a finitely generated projective right A-module, and

(3) $\text{Ext}_{A^e}^d(A, A^e)$ is isomorphic to A as a left A-module,

then A is a skew Calabi-Yau algebra of dimension d.

²⁰²⁰ Mathematics Subject Classification. 16E65, 16S35, 16S38.

Key words and phrases. Skew Calabi-Yau algebra, Nakayama automorphism, Hopf Galois extension.

Calabi-Yau algebras also arise from noncommutative crepant resolutions of Goren-stein algebras [\[32\]](#page-41-1). For instance, $\mathbb{C}[x_1,\ldots,x_n]\#G$, where G is a finite subgroup of $SL_n(\mathbb{C})$, serves as a non-commutative crepant resolution of $\mathbb{C}[x_1, \ldots, x_n]^G$. Skew group algebras of polynomial rings by finite groups are not uniformly Calabi-Yau. Nonetheless, they consistently exhibit the skew Calabi-Yau property. The (skew) Calabi-Yau property of the smash product of a skew Calabi-Yau algbera by a Hopf algebra has been examined in $[34]$, $[20]$, $[41]$, $[27]$, $[19]$. Let H be a Hopf algebra and A be an H-module algebra. Le Meur established in $[19]$ that when both A and H are skew Calabi-Yau, the smash product $A\#H$ is also skew Calabi-Yau, and its Nakayama automorphism can be expressed in terms of those of A and H . Additionally, Yu and Zhang investigate the skew Calabi-Yau property of the crossed product $A#_{\sigma}H$, with $\sigma: H \otimes H \to \mathbb{k}$ a 2-cocycle, in [\[40\]](#page-41-7).

Smash products and more general crossed products are inherently Hopf Galois extensions. Let H be a Hopf algebra. A right H -Galois extension is a right H comodule algebra B for which the canonical map $B \otimes_A B \to B \otimes H$ is bijective, where $A = B^{coH}$ is the coinvariant subalgebra. In noncommutative geometry, faithfully flat Hopf Galois extensions can be viewed as noncommutative principal bundles. This paper concentrates on examining the skew Calabi-Yau property of such extensions. Given a faithfully flat Hopf Galois extension $A \subseteq B$ over a Hopf algebra H, Stefan's spectral sequence arises as

$$
\operatorname{Ext}^p_H(\Bbbk,\operatorname{Ext}^q_{A^e}(A,B^e)) \Longrightarrow \operatorname{Ext}^{p+q}_{B^e}(B,B^e)
$$

It is not easy to clarify both the right H -module and left B^e -module structures on $\text{Ext}_{A^e}^n(A, B^e)$ concurrently. However, we discover that $\text{Ext}_{A^e}^i(A, B^e)$ admits an H^e -comodule structure, thereby endowing it with the structure of a Hopf bimodule in the category ${}_{B^e}\mathcal{M}_H^{H^e}$. This enables us to easily verify the conditions stated in Proposition [0.1,](#page-0-0) which in turn allows us to establish our main theorem.

Theorem 0.2. (Theorem [2.15\)](#page-14-0) Let H be a skew Calabi-Yau Hopf algebra of dimension n, and $A \subseteq B$ be a faithfully flat H-Galois extension. If A is skew Calabi-Yau of dimension d, then B is skew Calabi-Yau of dimension $(d + n)$.

The proof of Theorem [0.2](#page-1-0) does not yield a concrete characterization of the Nakayama automorphism. However, by utilizing Schauenburg's structure theorem for relative Hopf bimodules, it is demonstrated that the category ${}_{B^e}\mathcal{M}_H^{H^e}$ is equivalent to a left module category over Λ_1 , where Λ_1 is a subalgebra of B^e . In the case of a smash product $B = A \# H$, Λ_1 is isomorphic to Δ_1 , as discussed in [\[16,](#page-40-4)[19\]](#page-41-6), which has been instrumental in studying the Calabi-Yau property of such products. For a crossed product $\mathcal{A} \#_{\sigma} H$, there exists a natural embedding $H \hookrightarrow \Lambda_1$, allowing the homological determinant to be defined. Consequently, the Nakayama automorphisms of $\mathcal{A} \#_{\sigma} H$ are governed by those of A and H, as well as the homological determinant of the H-action on A.

Theorem 0.3. (Theorem [4.11\)](#page-37-0) Let H be a skew Calabi-Yau algebra of dimension n, $B = A \#_{\sigma} H$ be a crossed product of A with H. If A is skew Calabi-Yau of dimension d with a Nakayama automorphism μ_A , then B is also a skew Calabi-Yau algebra of dimension $n + d$ with a Nakayama automorphism μ_B which is defined by

$$
\mu_B(a \# h) = \sum \mu_A(a) \text{Hdet}(S^{-2}h_1) \# S^{-2}h_2 \chi(Sh_3),
$$

where Hdet is the homological determinant with respect to a μ_A -twisted volume as defined in Definition [4.8](#page-33-0) and $\chi : H \to \mathbb{k}$ is given by $\text{Ext}^n_H(\mathbb{k}, H) \cong {\chi} \mathbb{k}$.

The organization of the paper is as follows.

In Section [1,](#page-2-0) we review essential notation and recap preliminary findings. Theorem [0.1](#page-0-0) presents a criterion for a homologically smooth algebra to be skew Calabi-Yau, and its proof is provided in this section. Section [2](#page-5-1) outlines the basic definitions and properties of faithfully flat Hopf-Galois extensions. We endow $\text{Ext}_{A}^{i}(A, B^{e})$ with a Hopf module structure and, leveraging this framework, establish Theorem [0.2](#page-1-0) using Stefan's spectral sequence. We then consider the Nakayama automorphism for specific Hopf-Galois extension cases. To delve into the Nakayama automorphism for faithfully flat Hopf-Galois extensions $A \subseteq B$, we revisit Schauenburg's structure theorem for relative Hopf bimodules in Section [3,](#page-20-0) followed by an examination of the Hopf module structure on $\mathrm{Ext}_{A^e}^i(A, B^e)$. In Section [4,](#page-27-0) we describe an H-action on $\text{Ext}_{A^e}^i(A, A^e)$ when B is the crossed product $A#_{\sigma}H$. We compute the homological determinant for the crossed product $A#_{\sigma}H$. Subsequently, Theorem [0.3](#page-1-1) is proven, which expresses the Nakayama automorphism of $A#_{\sigma}H$ in terms of the Nakayama automorphisms of A and H, along with the homological determinant of the H-action on A.

1. Preliminaries

Throughout this paper, k is a base field, and all vector spaces and algebras are over k. Unadorned ⊗ means \otimes_k and Hom means Hom_k. Let A be an algebra over the base field k. We denote the opposite algebra of A by A^{op} , and denote the enveloping algebra $A \otimes A^{op}$ of A by A^e . Note that the switching operation $a \otimes b \mapsto b \otimes a$ extends to an anti-automorphism of the algebra A^e . We always assume that k acts centrally on all bimodules, and then we can identify the right (or left) A^e -modules with the A-A-bimodules. Usually we work with right modules, and a left A-module is viewed as a right A^{op} -module.

Given an abelian category A, let $\mathcal{D}(\mathcal{A})$ denote the derived category of A, and for $* = -, +, b$ let $\mathcal{D}^*(\mathcal{A})$ be the appropriate full subcategory (conventions as in [\[33\]](#page-41-8)). Given an algebra A, let $\mathcal{D}(A)$ denote the derived category of (right) A-modules. A complex $X \in \mathcal{D}^{-}(A)$ is called *pseudo-coherent* if L has a bounded-above free resolution $P^{\bullet} = (\cdots \to P^i \to P^{i+1} \to \cdots)$ such that each component P^i is a finitely generated free A-module $[38]$. An A-module M is called *pseudo-coherent* if it is pseudo-coherent as a complex. A complex is called perfect, if it is quasi-isomorphic to a bounded complex of finitely generated projective A-modules. Let $\mathcal{D}_{perf}(A^e)$ denote the full subcategory of $\mathcal{D}^b(A^e)$ consisting of perfect complexes.

For complexes $M, N \in \mathcal{D}(A)$ with either $M \in \mathcal{D}^{-}(A)$ or $N \in \mathcal{D}^{+}(A)$, let $\text{RHom}_{A}(M, N) \in \mathcal{D}(\mathbb{k})$ denote the right derived functor of the Hom functor. For complexes $M \in \mathcal{D}(A)$ and $N \in \mathcal{D}(A^{op})$ with $M \in \mathcal{D}^{-}(A)$ or $N \in \mathcal{D}^{-}(A^{op})$, let $M^{\mathbf{L}}\otimes_A N \in \mathcal{D}(\mathbb{k})$ denote the left derived functor of the tensor functor. For the notations of derived categories and derived functors we refer the reader to [\[33,](#page-41-8) [37\]](#page-41-10).

1.1. Skew Calabi-Yau algebras and Nakayama automorphism. An algebra is called *homologically smooth*, if A is perfect as a complex of A^e -modules. For any

 A^e -module M, the left A^e -module structure on A^e induces a A-A-bimodule structure on $\text{Hom}_{A^e}(M, A^e)$. For any $f \in \text{Hom}_{A^e}(M, A^e)$, $a, a' \in A$ and $m \in M$,

$$
(afa')(x) = (a \otimes a')f(m) \in A^e.
$$

Recall that a k-algebra A has Van den Bergh duality (or VdB duality, for short), if it is homologically smooth, and there exists some invertible A-A-bimodule ω and $d \in \mathbb{N}$, such that

$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \cong \omega[-d]
$$

in $\mathcal{D}^b(A^e)$, and ω is called the *VdB dual module* of A.

Let M be an A-A-bimodule. If μ is an algebra endomorphism of A, then we let M_{μ} denote the bimodule M with normal left action, but right action twisted by μ . That is, $a \cdot m \cdot a' = am\mu(a')$ for all $a, a' \in A$ and $m \in M$.

Definition 1.1. [\[27,](#page-41-5) Definition 0.1] A homologically smooth algebra A is called skew Calabi-Yau of dimension d for some integer $d > 0$, if there exists an algebra automorphism $\mu \in \text{Aut}(A)$ such that

$\operatorname{RHom}_{A^e}(A, A^e) \cong A_\mu[-d]$

in $\mathcal{D}^b(A^e)$. Then μ is called a *Nakayama automorphism* of A. If μ is an inner automorphism, that is, $\text{Ext}_{A^e}^d(A, A^e) \cong A$ as A^e -module, then A is called a Calabi-Yau algebra of dimension d.

The coordinate ring of a smooth affine Calabi-Yau variety is the original and most prototypical example of a Calabi-Yau algebra. Skew Calabi-Yau algebras are commonly exemplified by Artin-Schelter regular and Hopf algebras, as discussed in the following subsection.

1.2. Artin-Schelter regular Hopf algebras. Throughout this paper, H will stand for a Hopf algebra (H, Δ, ε) with a bijective antipode S. We use the Sweedler notation $\Delta(h) = \sum h_1 \otimes h_2$ for all $h \in H$. We recommend [\[24\]](#page-41-11) as a basic reference for the theory of Hopf algebras. We will always assume that the antipode S is bijective. This assumption on S is automatic if H is noetherian AS-Gorenstein $[22,$ Theorem 0.2] or skew Calabi-Yau [\[19,](#page-41-6) Proposition 3.4.2].

Definition 1.2. A Hopf algebra H is called AS-Gorenstein of dimension d, if

- (1) H has left and right injective dimension $d < +\infty$,
- (2) $\mathrm{Ext}^i_H(\mathbb{k}, H) \cong \mathrm{Ext}^i_{H^{op}}(\mathbb{k}, H) \cong$ $\int 0, \quad i \neq d$ $\mathbb{k}, \quad i = d$, where \Bbbk is the trivial $H\text{-module}$ $H/Ker \varepsilon$.

If in addition, H has finite global dimension d , then H is called AS -regular.

Let H be an AS-Gorenstein algebra of dimension d . The right homological integral [\[21\]](#page-41-13) of H is defined to be the 1-dimensional H-H-bimodule $\text{Ext}_{H}^{d}(\mathbb{K}, H)$. Then there exists an algebra homomorphism $\chi : H \to \mathbb{k}$ such that $\text{Ext}^d_H(\mathbb{k}, H)$ is isomorphic to $\chi^{\mathbb{R}}$ as H-H-bimodules, that is, for any $x \in \text{Ext}_{H}^{d}(\mathbb{R}, H)$, $hx = \chi(h)x$ and $xh = \varepsilon(h)x$. One may define the left homological integral $\text{Ext}_{H^{op}}^d(\mathbb{k}, H)$ similarly.

As well known, the skew Calabi-Yau property of Hopf algebra is also equivalent to the AS-regular property.

Theorem 1.3. [\[19,](#page-41-6) Lemma 3.3] The following statements are equivalent for a Hopf algebra H.

 (1) H is homologically smooth.

 (2) k is a perfect H-complex.

Theorem 1.4. [\[19,](#page-41-6) Theorem 1] The following statements are equivalent for a Hopf algebra H with a bijective antipode.

- (1) H has VdB duality.
- (2) H is AS-regular, and the trivial H -module k is pseudo-coherent.
- (3) H is skew Calabi-Yau.

 $\sum S^{-2}h_1\chi(Sh_2)$ for all $h \in H$, where $\text{Ext}^d_H(\mathbb{k}, H) \cong \chi \mathbb{k}$. In each case, μ_H serves as a Nakayama automorphism of H, given by $\mu_H(h)$ =

Here, we present several examples of Skew Calabi-Yau Hopf algebras.

If H is a finite-dimensional Hopf algebra, it is skew Calabi-Yau if and only if it is semisimple.

Any noetherian affine PI Hopf algebra with finite global dimension d is inherently a d-dimensional skew Calabi-Yau algebra [\[5,](#page-40-2) 6.2]. In particular, the coordinate ring $\mathcal{O}(G)$ of an affine algebraic group G over a field k of characteristic zero is always a Calabi-Yau algebra.

According to [\[42,](#page-41-14) Proposition 6.6], any connected Hopf algebra with finite Gelfand-Kirillov dimension d over a field of characteristic 0 is guaranteed to be a skew Calabi-Yau algebra of dimension d.

1.3. A criterion for homologically smooth algebras to be skew Calabi-Yau. In this subsection, we give a useful criterion for homologically smooth algebras to be skew Calabi-Yau.

An object X in $\mathcal{D}^b(A^e)$ is termed a *two-side tilting complex* over A, if there exists an object $X^{\vee} \in \mathcal{D}^b(A^e)$ satisfying $X^{\mathbf{L}} \otimes_A X^{\vee} \cong A \cong X^{\vee} \mathbf{L} \otimes_A X$ in $\mathcal{D}^b(A^e)$. If X is a two-side tilting complex over A, then $X^{\vee} \cong \mathbf{R}\text{Hom}_{A}(X, A) \cong \mathbf{R}\text{Hom}_{A^{\text{op}}}(X, A)$ in $\mathcal{D}(A^e)$, and X is a perfect complex in both $\mathcal{D}(A)$ and $\mathcal{D}(A^{op})$.

Lemma 1.5. Let A be a homologically smooth algebra. If $\Omega := \mathbf{R} \text{Hom}_{A^e}(A, A^e)$ is perfect in both $\mathcal{D}(A)$ and $\mathcal{D}(A^{op})$, then Ω is a two-side tilting complex over A.

Proof. Given that A is homologically smooth, there is a bounded complex P of finitely generated projective A^e -modules such that $A \cong P$ in $\mathcal{D}^b(A^e)$. Thus, in $\mathcal{D}(A^e), \Omega \cong \text{Hom}_{A^e}(P, A^e)$ and $A \cong P \cong \text{Hom}_{A^e}(\text{Hom}_{A^e}(P, A^e), A^e) \cong \text{R}\text{Hom}_{A^e}(\Omega, A^e)$. This implies

$$
A \cong \mathbf{R}\mathrm{Hom}_{A^e}(\Omega, A^e)
$$

\n
$$
\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, \mathbf{R}\mathrm{Hom}_A(\Omega, A^e))
$$

\n
$$
\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e \otimes_A \mathbf{R}\mathrm{Hom}_A(\Omega, A))
$$

\n
$$
\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A \otimes \mathbf{R}\mathrm{Hom}_A(\Omega, A))
$$

\n
$$
\cong (A \otimes \mathbf{R}\mathrm{Hom}_A(\Omega, A))^{\mathbf{L}} \otimes_{A^e} \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e)
$$

\n
$$
\cong \mathbf{R}\mathrm{Hom}_A(\Omega, A)^{\mathbf{L}} \otimes_A \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e)
$$

\n
$$
\cong \mathbf{R}\mathrm{Hom}_A(\Omega, A)^{\mathbf{L}} \otimes_A \Omega.
$$

Similarly, it can be shown that $A \cong \Omega^{\mathbf{L}} \otimes_A \mathbf{R}$ Hom_{$A^{op}(\Omega, A)$} in $\mathcal{D}(A^e)$. Consequently, Ω a two-side tilting complex over A. □

The following result is well known.

Lemma 1.6. Let R be a ring, and ω be an R-R-bimodule. If ω is a free R-module of rank one, then there exists an algebra endmorphism μ of R for which ω is isomorphic to R_{μ} as an R-R-bimodules. Furthermore, if ω is an invertible R-R-bimodule, then μ is an automorphism.

As a consequence of Lemma [1.5](#page-4-0) and [1.6,](#page-5-2) we establish the following criterion for a homologically smooth algebra to be skew Calabi-Yau.

Proposition 1.7. Let A be a homologically smooth algebra. Suppose that there exists an A-A-bimodule ω and an integer $d \in \mathbb{N}$ such that $\mathbf{R}\text{Hom}_{A^e}(A, A^e) \cong \omega[-d]$ in $\mathcal{D}(A^e)$. If ω is finitely generated and projective as both a left and right A-module, then ω is an invertible A-A-bimodule. Moreover, if ω is isomorphic to A as either a left or right A-module, then A is a skew Calabi-Yau algebra of dimension d.

2. The skew Calabi-Yau property of flat Hopf Galois extensions

2.1. Hopf Galois extensions. In this subsection we recall the definition and basic properties of an H-Galois extension. Let B be a right H-comodule algebra via an algebra homomorphism $\rho : B \to B \otimes H$. The *coinvariant subalgebra* of the Hcoaction on B is ${b \in B \mid \rho(b) = b \otimes 1}$, denoted by B^{coH} . If (V, ρ_V) is a right H-comodule, then we use the Sweedler notation $\rho(v) = \sum v_1 \otimes v_2$ for all $h \in H$.

Definition 2.1. Then the extension $A(:= B^{coH}) \subset B$ is called (right) *H*-Galois if the canonical map

$$
\beta: B \otimes_A B \to B \otimes H, \quad b' \otimes_A b \mapsto b' \rho(b) = \sum b' b_0 \otimes b_1
$$

is bijective, and β is also called the Galois map.

Let A be an algebra, and \mathcal{M}_A (resp., $_A\mathcal{M}$) be the category of right (resp., left) Amodules. Let B be a right H-comodule algebra. A relative right-right (B, H) -Hopf module is a vector space with a right B-action and a right H-coaction ρ_M such that

$$
\rho_M(mb) = \rho_M(m)\rho(b) = \sum m_0b_0 \otimes m_1b_1
$$

for all $m \in M$ and $b \in B$. A relative left-right (B, H) -Hopf module is defined similarly. The category of relative right-right (resp., left-right) (B, H) -Hopf modules is denoted by \mathcal{M}_B^H (resp., $_B\mathcal{M}^H$).

We require some known facts about Hopf Galois extensions.

Theorem 2.2. [\[31,](#page-41-15) Theorem I] [\[30,](#page-41-16) Theorem 5.6] Let H be a Hopf algebra with a bijective antipode, B be a right H-comodule algebra and $A := B^{coH}$. Then the following are equivalent.

(1) B is injective as right H-comodule, and $\beta : B \otimes_A B \to B \otimes H$ is surjective.

- (2) $(-\otimes_A B, (-)^{coH})$ is an equivalence between \mathcal{M}_A and \mathcal{M}_B^H .
- (3) $(B \otimes_A -, (-)^{coH})$ is an equivalence between $_A \mathcal{M}$ and $_B \mathcal{M}^H$.
- (4) B is a faithfully flat left A-module, and $A \subseteq B$ is an H-Galois extension.

(5) B is a faithfully flat right A-module, and $A \subseteq B$ is an H-Galois extension.

(6) $A \subseteq B$ is a Hopf Galois extension, and the comodule algebra B is right Hequivariantly projective as a left A -module, i.e., there exists a right H -colinear and left A-linear splitting of the multiplication $A \otimes B \to B$.

Then we say a Hopf Galois extension $A \subseteq B$ is faithfully flat if B is faithfully flat as a left or right A-module. A faithfully flat H-Galois extension is also called a principal homogeneous space in $[31]$ or principal H-extension in $[7]$.

2.2. Ulbrich-Miyashita actions on M^A and M_A . Suppose now that $A \subset B$ is an H-Galois extension. Following $[6]$ we shall call the k-linear map

$$
\kappa: H \to B \otimes_A B, \quad h \mapsto \beta^{-1}(1 \otimes h)
$$

the translation map associated to the H-Galois extension $A \subseteq B$. For any $h \in H$ we shall use the notation

$$
\kappa(h) = \sum \kappa^1(h) \otimes_A \kappa^2(h).
$$

The following are key properties of κ as outlined in [\[31\]](#page-41-15):

(2.1)
$$
\sum \kappa^1(hk) \otimes_A \kappa^2(hk) = \sum \kappa^1(k) \kappa^1(h) \otimes_A \kappa^2(h) \kappa^2(k) \text{ in } B \otimes_A B,
$$

(2.2)
$$
\sum a \kappa^1(h) \otimes_A \kappa^2(h) = \sum \kappa^1(h) \otimes_A \kappa^2(h) a \text{ in } B \otimes_A B,
$$

$$
(2.3) \sum \kappa^1(h) \otimes_A \kappa^2(h)_0 \otimes \kappa^2(h)_1 = \sum \kappa^1(h_1) \otimes_A \kappa^2(h_1) \otimes h_2 \text{ in } B \otimes_A B \otimes H,
$$

$$
(2.4) \sum \kappa^1(h)_0 \otimes_A \kappa^2(h) \otimes \kappa^1(h)_1 = \sum \kappa^1(h_2) \otimes_A \kappa^2(h_2) \otimes Sh_1 \text{ in } B \otimes_A B \otimes H,
$$

(2.5)
$$
\sum b_0 \kappa^1(b_1) \otimes_A \kappa^2(b_1) = 1 \otimes_A b \text{ in } B \otimes_A B,
$$

(2.6)
$$
\sum \kappa^1(h)\kappa^2(h) = \varepsilon(h) \text{ in } B,
$$

Given an H-Galois extension $A \subseteq B$, $B \otimes_A B$ is a left H-module induced by the module structure on $B \otimes H$. More precisely, for all $h \in H$ and $x, y \in B$, the action is defined as

$$
h \rightarrow (x \otimes_A y) := \beta^{-1}(h\beta(x \otimes_A y)) = \beta^{-1}(h \sum xy_0 \otimes y_1) = \beta^{-1}(\sum xy_0 \otimes hy_1).
$$

Using equations (2.3) and (2.6) , we find

$$
\beta\big(\sum x\kappa^1(h)\otimes_A\kappa^2(h)y\big)=\sum x\kappa^1(h)\kappa^2(h)_{0}y_{0}\otimes_A\kappa^2(h)_{1}y_{1}=\sum xy_{0}\otimes hy_{1},
$$

The bijectiveness of β then yields

$$
h \rightarrow (x \otimes_A y) = \beta^{-1}(\sum xy_0 \otimes hy_1) = \sum x \kappa^1(h) \otimes_A \kappa^2(h)y.
$$

Thus, the Galois map $\beta : B \otimes_A B \to B \otimes H$ is a morphism of $H\text{-}B^e$ -bimodules, where the right B^e module structure on $B \otimes H$ is given by

$$
(x\otimes h)(b\otimes b')=\sum b'xb_0\otimes hb_1
$$

for all $h \in H$ and $x, b, b' \in B$.

For any A-A-bimodule M, we set

$$
M^A := \{ m \in M \mid am = ma, \ \forall a \in A \}, \quad \text{and} \quad M_A := M/[A, M],
$$

where $[A, M]$ is the addtive subgroup of M generated by the elements $am - ma$ for all $a \in A, m \in M$. Then one can easy to check that

$$
M^A \cong \text{Hom}_{A^e}(A, M), \quad M_A \cong A \otimes_{A^e} M \cong M \otimes_{A^e} A.
$$

For any left B^e -module M, we have the following isomorphisms:

$$
(2.7) \quad M^A = \text{Hom}_{A^e}(A, M) \cong \text{Hom}_{B^e}(A \otimes_{A^e} B^e, M) \cong \text{Hom}_{B^e}(B \otimes_A B, M).
$$

Thus M^A is a right H-module via the left H-module structure on $B \otimes_A B$. More precisely, for all $h \in H$ and $x \in M^A$,

(2.8)
$$
x - h = \sum \kappa^1(h) x \kappa^2(h).
$$

Similarly, we also have the following isomorphisms:

(2.9)
$$
M_A \cong A \otimes_{A^e} M \cong (A \otimes_{A^e} B^e) \otimes_{B^e} M \cong (B \otimes_A B) \otimes_{B^e} M.
$$

Thus M_A is a left H-module via the left H-module structure on $B \otimes_A B$. More precisely, for all $h \in H$ and $x \in M$,

(2.10)
$$
h \rightarrow (x + [A, M]) = \sum \kappa^{2}(h) x \kappa^{1}(h) + [A, M].
$$

These H-module structures, known as Ulbrich-Miyashita actions, were previously introduced in [\[11\]](#page-40-7) and [\[9\]](#page-40-8). It is worth mentioning that the Ulbrich-Miyashita actions are natural in the sense that, for any B^e -modules M and N, and any map $f \in$ $\text{Hom}_{B^e}(M, N)$, the induced maps

$$
f^0: M^A \longrightarrow N^A
$$
, and $f_0: M_A \longrightarrow N_A$

are both H-linear. Therefore, the assignment $M \mapsto M^A$ constitutes a functor from $_{B}P_{B}M$ (which is equivalent to $_{B}M_{B}$) to \mathcal{M}_{H} .

2.3. Stefan's spectral sequence for flat Hopf Galois extensions. A Hopf Galois extension $A \subseteq B$ is termed flat if B is flat as both a left and right A-module. For any right H -module V , we have

(2.11)
$$
B \otimes V \cong V \otimes_H (B \otimes H) \cong V \otimes_H (B \otimes_A B).
$$

As $B \otimes_A B$ is an H-B^e-bimodule, $B \otimes V$ becomes a B^e-module with the action defined by

$$
(x\otimes v)(b\otimes b')=\sum b'xb_0\otimes vb_1,
$$

for any $b, b', x \in B$ and $v \in V$. The assignment $V \mapsto B \otimes V$ gives a functor from \mathcal{M}_H to $_{B^e}\mathcal{M}$, which is left adjoint to $(-)^A$. Moreover, we have the natural isomorphisms

$$
\text{Hom}_{H}(V, M^{A}) \stackrel{(2.7)}{\cong} \text{Hom}_{H}(V, \text{Hom}_{B^{e}}(B \otimes_{A} B, M))
$$
\n
$$
\cong \text{Hom}_{B^{e}}(V \otimes_{H} (B \otimes_{A} B), M)
$$
\n
$$
\stackrel{(2.11)}{\cong} \text{Hom}_{B^{e}}(B \otimes V, M),
$$

for any left B^e -module M and right H-module V.

For any injective B^e -module I, the adjoint isomorphism (2.12) implies that I^A is an injective H-module as the functor $B \otimes - : \mathcal{M}_H \to {}_{Be}\mathcal{M}$ is exact. Furthermore, since

 $\operatorname{Hom}_{B^e}(B, -) \cong \operatorname{Hom}_H(\mathbb{k}, \operatorname{Hom}_{A^e}(A, -)) = \operatorname{Hom}_H(\mathbb{k}, -) \circ \operatorname{Hom}_{A^e}(A, -),$

we derive the following Grothendieck spectral sequence associated with a flat Hopf-Galois extension, as established by Stefan.

Theorem 2.3. [\[9,](#page-40-8) Theorem 3.3] Let $A \subseteq B$ be a flat H-Galois extension. For any B^e -module M, we have that

$$
\mathbf{R}\mathrm{Hom}_{B^e}(B,M) \cong \mathbf{R}\mathrm{Hom}_H(\mathbb{k},\mathbf{R}\mathrm{Hom}_{A^e}(A,M)).
$$

To see whether a Hopf Galois extension B is skew Calabi-Yau, we need to study the B-B-bimodule structure on the cohomological groups $\text{Ext}_{B^e}^n(B, B^e)$ for all $n \geq 0$. Stefan's spectral sequenceallows us to focus on the B^e -H bimodule structure on $Ext_{A^e}^n(A, B^e)$ for all $n \geq 0$, see the following corollary which is immediate consequence of Theorem [2.3.](#page-8-0) For clarity, we provide a proof here. For the reader's convenience, we include a proof here.

Corollary 2.4. Let $A \subseteq B$ be a flat H-Galois extension. For any B^e -module M, in $\mathcal{D}(B^e)$.

$$
\mathbf{R}\mathrm{Hom}_{B^e}(B,B^e) \cong \mathbf{R}\mathrm{Hom}_H(\mathbb{k},\mathbf{R}\mathrm{Hom}_{A^e}(A,B^e)).
$$

Proof. Take an injective $B^e \otimes (B^e)^{op}$ -module resolution E^{\bullet} of B^e . Since B is flat both as a left and right A-modules, each component $Eⁿ$ is an injective left A^e -module by the natural adjoint isomorphism

 $\text{Hom}_{A^e}(M, E^n) = \text{Hom}_{A^e}(M, \text{Hom}_{B^e}(B^e, E^n)) \cong \text{Hom}_{B^e}(M \otimes_{A^e} B^e, E^n)$

for any left A^e -module M. Then $\mathbf{R}\text{Hom}_{A^e}(A, B^e)$ is isomorphic to $\text{Hom}_{A^e}(A, E^{\bullet})$ in $\mathcal{D}((H \otimes (B^e)^{op})$. Given that $\text{Hom}_{A^e}(A, E^n)$ is an injective H-module, we have

 $\mathbf{R}\mathrm{Hom}_H(\mathbb{k},\mathbf{R}\mathrm{Hom}_{A^e}(A,B^e)) \cong \mathrm{Hom}_H(\mathbb{k},\mathrm{Hom}_{A^e}(A,E^{\bullet})).$

This is isomorphic to $\text{Hom}_{B^e}(B, E^{\bullet})$ due to the adjoint isomorphism [\(2.12\)](#page-7-2), thus completing the proof. \Box

On one hand, taking an injective $B^e \otimes (B^e)^{op}$ -module resolution E^{\bullet} of B^e allows us to understand the H-module structure on $\text{Ext}_{A^e}^n(A, B^e) = \text{H}^n(\text{Hom}_{A^e}(A, E^{\bullet})),$ yet it does not expose the B^e -module structure. On the other hand, a projective A^e -module resolution P_{\bullet} of A provides the opposite insight. Thus, it is challenging to concurrently clarify the right H -module and left B^e -module structures on $\text{Ext}_{A^e}^n(A, B^e)$. Subsequently, we utilize the H-comodule structure of $\text{Ext}_{A^e}^n(A, B^e)$ to investigate its bimodule structures.

2.4. Hopf module structure on M^A . Let $H^* := \text{Hom}(H, \mathbb{k})$ be the dual algebra of H. A left H^{*}-module (V, \rightharpoonup) is called rational if $H^* \rightharpoonup v$ is finite dimensional for any $v \in V$. A left H^* -module V is rational if and only if its H^* -action comes from a right H -coaction, meaning V is an H -comodule satisfying

$$
f \rightarrow v = \sum_{(v)} v_0 f(v_1)
$$

for any $v \in V$ and $f \in H^*$.

Let N be a vector space and M be a right H-comodule. The space $\text{Hom}(N, M)$ forms a left H^* -module via the action

$$
(h^* \to f)(n) = h^* \to (f(n))
$$

for all $h^* \in H^*$, $f \in \text{Hom}(N, M)$ and $n \in N$. If $N \in \mathcal{M}_{A^e}$ and $M \in \mathcal{M}_{A^e}^{H^e}$, then $\text{Hom}_{A^e}(N, M)$ is an H^* -submodule of $\text{Hom}(N, M)$. Moreover, if N is finitely generated as an A^e -module, then $\text{Hom}_{A^e}(N, M)$ is an H^* -submodule of a direct sum of copies of M. Consequently, $\text{Hom}_{A^e}(N, M)$ is a rational H^* -module, that is, implying it is an H-comodule. This leads to the following direct corollary.

Lemma 2.5. If N is a finitely generated A^e -module, then for any $M \in B^e \mathcal{M}_{B^e}^H$, $\text{Hom}_{A^e}(N, M)$ is inherently a Hopf module in $\mathcal{M}_{B^e}^H$.

Let M be a module and comodule over H . We call M an (right-right) Yetter-Drinfeld module if and only if the action and coaction are compatible in the following sence:

$$
(2.13) \ \rho(m-h) = \sum (m-h)_0 \otimes (m-h)_1 = \sum m_0 - h_2 \otimes (Sh_1)m_1h_3,
$$

for all $m \in M$ and $h \in H$. We denote \mathcal{YD}_H^H as the category of right-right Yetter-Drinfeld modules, with morphisms that are both linear and colinear.

For any $M \in {}_B\mathcal{M}_B^H$, it is evident that M^A is an H-subcomodule of M. We recall that M^A carries a right H-module as per (2.8) . This action and coaction fulfill the Yetter-Drinfeld compatibility condition (2.13) , which means $(-)^A$ can be promoted to a functor from ${}_B\mathcal{M}_B^H$ to \mathcal{YD}_H^H , as shown in [\[11,](#page-40-7) Proposition 3.10].

As B^e cannot be inherently viewed as a Hopf bimodule in ${}_B\mathcal{M}_B^H$, we must instead examine the category ${}_{B^e}M^{\check{H}^e}$. Since the antipode S is bijective as we assumed, H^{op} also constitutes a Hopf algebra with antipode S^{-1} . For an H-Galois extension $A \subseteq B$, B^{op} is a right H^{op} -comodule algebra, and B^e is a right H^e -comodule algebra. Consequently, $A^{op} \subseteq B^{op}$ is an H^{op} -Galois extension. Since there is an isomorphism

$$
B\otimes_A B\stackrel{\cong}{\longrightarrow}B^{op}\otimes_{A^{op}}B^{op},\qquad x\otimes_A y\mapsto y\otimes_{A^{op}} x,
$$

the element $\sum \kappa^2(k) \otimes_{A^{op}} \kappa^1(k)$ in $B^{op} \otimes_{A^{op}} B^{op}$ is well defined for any $k \in H$. Furthermore, we have

$$
\beta\left(\sum \kappa^2 (S^{-1}h) \otimes_{A^{op}} \kappa^1 (S^{-1}h)\right) = \sum \kappa^1 (S^{-1}h){}_0 \kappa^2 (S^{-1}h) \otimes \kappa^1 (S^{-1}h)_{1}
$$

$$
\xrightarrow{\underline{\text{(2.4)}}} \sum \kappa^1 (S^{-1}h_1) \kappa^2 (S^{-1}h_1) \otimes h_2
$$

$$
\xrightarrow{\underline{\text{(2.6)}}} 1 \otimes h.
$$

Thus, the translation map $\kappa_{B^{op}}$ for the H^{op} -Galois extension $A^{op} \subseteq B^{op}$ satisfies

$$
\kappa_{B^{op}}(h)=\sum\kappa_{B^{op}}^1(h)\otimes_{A^{op}}\kappa_{B^{op}}^2(h)=\sum\kappa^2(S^{-1}h)\otimes_{A^{op}}\kappa^1(S^{-1}h)\in B^{op}\otimes_{A^{op}}B^{op}.
$$

Then $A^e \subseteq B^e$ is an H^e -Galois extension, with the translation map κ_{B^e} defined as

$$
\kappa_{B^e}(h \otimes k) = \sum \kappa_{B^e}^1(h \otimes k) \otimes \kappa_{B^e}^2(h \otimes k)
$$

\n
$$
= \sum \kappa^1(h) \otimes \kappa_{B^{op}}^1(k) \otimes_{A^e} \kappa^2(h) \otimes \kappa_{B^{op}}^2(k)
$$

\n
$$
= \sum \kappa^1(h) \otimes \kappa^2(S^{-1}k) \otimes_{A^e} \kappa^2(h) \otimes \kappa^1(S^{-1}k) \in B^e \otimes_{A^e} B^e.
$$

In the subsequent discussion, H is considered a right H^e -comodule algebra through the coaction

(2.15)
$$
\rho_H: H \longrightarrow H \otimes H^e, \qquad h \mapsto \sum h_2 \otimes (h_3 \otimes Sh_1).
$$

Lemma 2.6. The mapping $M \mapsto M^A$ constitutes a functor from $\mathcal{M}_{B^e}^{H^e}$ to $\mathcal{M}_H^{H^e}$. Furthermore, $(-)^A : M \mapsto M^A$ extends to a functor form ${}_{B^e} \mathcal{M}_{B^e}^{H^e}$ to ${}_{B^e} \mathcal{M}_H^{H^e}$.

Proof. For any $M \in \mathcal{M}_{Be}^{H^e}$, M is naturally a B-B-bimodule, and M^A is an H^e subcomodule of M. Recall that M^A is endowed with a right H-module structure indicated in [\(2.8\)](#page-7-3). For any $m \in M^A$, observe that $\rho(m) = \sum m_0 \otimes m_1 \in M \otimes H^e$. For any $h \in H$, it follows that

ρ(m ↼ h) [\(2.8\)](#page-7-3) ==== ρ(Xκ 1 (h)mκ² (h)) = Xκ 1 (h)0m0κ 2 (h)⁰ ⊗ κ 1 (h)1m1κ 2 (h)¹ [\(2.3\)](#page-6-0),[\(2.4\)](#page-6-2) ======= Xκ 1 (h2)m0κ 2 (h2) ⊗ (Sh1)m1h³ = X(m⁰ ↼ h2) ⊗ (m1(h³ ⊗ Sh1)) = ρM(m)ρH^e (h),

confirming that M^A belongs to \mathcal{M}_H^{He} . Moreover, for any $M \in {}_{B^e}\mathcal{M}_{Be}^{He}$, M^A is also a left B^e -module and the B^e -action is compatible with the H^e -coaction. Therefore, M^A is a Hopf module in $_{B^e}\mathcal{M}_H^{H^e}$. H^e .

Lemma 2.7. The assignment $M \mapsto M_A = M/[A, M]$ becomes a functor from ${}_{B^e}\mathcal{M}^{H^e}$ to ${}_H\mathcal{M}^{H^e}$. Furthermore, $(-)_A: M \mapsto M_A$ extends to a functor form ${}_{B^e}\mathcal{M}^{H^e}_{B^e}$ to ${}_H{\cal M}^{H^e}_{B^e}.$

Proof. For any $M \in {}_{B^e}{\mathcal{M}^{H^e}}$, M_A is a quotient H^e -comodule of M. Recall that M^A is a right H-module via [\(2.8\)](#page-7-3). For all $m \in M$ and $h \in H$, we have

$$
\rho(h \rightarrow (m + [A, M])) \xrightarrow{\underline{(-1, 0)}} \rho(\sum \kappa^2(h) m \kappa^1(h) + [A, M])
$$

=
$$
\sum (\kappa^1(h)_{0} m_{0} \kappa^2(h)_{0} + [A, M]) \otimes \kappa^1(h)_{1} m_{1} \kappa^2(h)_{1}
$$

$$
\xrightarrow{\underline{(-2, 3), (2, 4)}} \sum (\kappa^1(h_2) m_{0} \kappa^2(h_2) + [A, M]) \otimes (Sh_1) m_{1} h_3
$$

=
$$
\sum (h_2 \rightarrow (m_0 + [A, M])) \otimes ((h_3 \otimes Sh_1) m_1)
$$

=
$$
\rho_{H^e}(h) \rho_M(m + [A, M]).
$$

showing that $M_A \in {}_H{\mathcal{M}}^{H^e}$. Moreover, for any $M \in {}_{B^e}{\mathcal{M}}^{H^e}_{B^e}$, M_A is also a right B^e -module and the B^e -action satisfies the corresponding compatibility condition. Thus M_A is a Hopf module in ${}_H{\mathcal{M}_{Be}^{He}}$. H^e
 B^e . Further, consider the following commutative diagram of forgetful functors:

Notice that $A \otimes B^{op}$ is a trivial $H(= H \otimes \mathbb{k})$ -comodule algebra. Thus, for any $M \in$ $A\otimes B^{op}\mathcal{M}_H^H$, M^{coH} constitutes an $A\otimes B^{op}$ -submodule of M. For any $N\in A\otimes B^{op}\mathcal{M}$, $N \otimes H$ forms a Hopf bimodule in ${}_{A \otimes B^{op}}\mathcal{M}^H_H$, defined by

$$
\rho_H(n \otimes h) = \sum_{(h)} n \otimes h_1 \otimes h_2, \quad (a \otimes b)(n \otimes h)h' = (a \otimes b)n \otimes hh',
$$

where $a \in A$, $b \in B$, $h, h' \in H$ and $n \in N$.

Then we have the following lemma.

Lemma 2.8. Let H be a Hopf algebra and B a right faithfully flat H-Galois extension of $A := B^{coH}$. Then we have the following category equivalences

(1) $_{A\otimes B^{op}}\mathcal{M}_H^H \cong_{A\otimes B^{op}}\mathcal{M}$ via $M \mapsto M^{coH}$ and $N \otimes H \leftrightarrow N$,
(2) $_{\sim} M^H \cong_{\sim} M$ via $X \mapsto X^{coH}$ and $B \otimes_{\sim} N \leftrightarrow N$ (2) $_{B^e}\mathcal{M}^H \cong_{A\otimes B^{op}} \mathcal{M}$ via $X \mapsto X^{coH}$ and $B \otimes_A N \leftrightarrow N$,

Proof. (1) The result is straightforward, given the equivalence $\mathcal{M}_H^H \cong \mathcal{M}_k$ with the mappings $M \mapsto M^{coH}$ and $N \otimes H \leftarrow N$.

(2) As $A \otimes B^{op} \subset B \otimes B^{op}$ is a faithfully flat H-Galois extension (with $H = H \otimes \mathbb{k}$), the assertion is a direct consequence of Theorem [2.2.](#page-5-3)

2.5. Hopf module structure on $\text{Ext}_{A^e}^n(A, B^e)$.

Lemma 2.9. Let H be a Hopf algebra, R and T be two H -comodule algebras. Then ${}_{R}\mathcal{M}^{H}_{T}$ has enough injective objects.

Proof. For any R-T-bimodule M, $M \otimes H$ is a Hopf bimodule in $_R\mathcal{M}_T^H$, where the R-T-bimodule structure defined by $r(m \otimes h)t = \sum r_0 m t_0 \otimes r_1 ht_1$, and the right H-comodule is defined by $\rho(m \otimes h) = \sum m \otimes h_1 \otimes h_2$. For any $M \in {}_R\mathcal{M}_T$ and $X \in {}_R\mathcal{M}_T^H$, there is a natural isomorphism

$$
(2.16) \Phi: \text{Hom}_{R^{\mathcal{M}_T}}(X, M) \xrightarrow{\cong} \text{Hom}_{R^{\mathcal{M}_T^H}}(X, M \otimes H), \quad f \mapsto \big(x \mapsto \sum f(x_0) \otimes x_1\big),
$$

and the inverse map of Φ is given by $g \mapsto (\mathrm{id}_M \otimes \varepsilon) \circ g$. For any $X \in {}_R\mathcal{M}_T^H$, there is an injective $R \otimes T^{op}$ -module I with an $R \otimes T^{op}$ -linear imbedding $\iota : X \hookrightarrow I$. Given that $(id_M \otimes \varepsilon) \circ \Phi(\iota) = \Phi^{-1} \Phi(\iota) = \iota$, the map $\Phi(\iota) : X \hookrightarrow I \otimes H$ is also injective. As $I \otimes H$ is an injective object in ${}_{R}\mathcal{M}^{H}_{T}$ by the natural isomorphism [\(2.16\)](#page-11-0), we conclude that ${}_{R}\mathcal{M}_{T}^{H}$ has enough injective objects. \square

Remark 2.10. Recall that an abelian category A is Grothendieck, if it has exact direct limits and a generator, and that any Grothendieck category has enough injective objects. Since \mathcal{M}^H is a subcategory of $_{H^*}\mathcal{M}$ which is closed under subobjects, quotient objects and coproducts, $\widetilde{\mathcal{M}}^H$ is a Grothendieck category (see [\[12\]](#page-40-9) for example). By analogy, ${}_{R}\mathcal{M}^{H}_{T}$ is also classified as a Grothendieck category.

□

Proposition 2.11. Let H be a Hopf algebra with a bijective antipode, B be a flat H-Galois extension of A.

(1) Then the right derived functor F of $(-)^A = \text{Hom}_{A^e}(A, -) : B^e \mathcal{M}_{B^e}^{H^e} \to$ ${}_{B^e}\mathcal{M}_H^{He}$ exists, such that the following diagram

$$
\begin{CD} \mathcal{D}^+\!\left({_{B^e}}\mathcal{M}_{B^e}^{H^e}\right) \xrightarrow{\qquad F}{} \mathcal{D}\!\left({_{B^e}}\mathcal{M}_{H}^{H^e}\right) \ @V{\qquad \qquad }\\ \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow\\ \mathcal{D}^+\!\left(\mathcal{M}_{A^e}\right) \xrightarrow{\mathbf{R}\mathrm{Hom}_{A^e}(A,-)}{} \mathcal{D}\!\left(\mathcal{M}_\Bbbk\right) \end{CD}
$$

commutes up to natural isomorphism. So the derived functor from ${\cal D}^+(_{B^e}{\cal M}^{H^e}_{B^e})$ to $\mathcal{D}(B_e\mathcal{M}_H^{H^e})$ is also denoted by $\mathbf{R}\text{Hom}_{A^e}(A,-)$.

- (2) For any $X \in {}_{B^e}{\mathcal M}^{H^e}_{A^e}$, **R**Hom_A^e(-, X) is isomorphic to $X^{\mathbf{L}} \otimes_{A^e} \mathbf{R}$ Hom_A^e(-, A^e) as a functor from $\mathcal{D}_{perf}(A^e)^{op}$ to $\mathcal{D}(B^e \mathcal{M}^{H^e}).$
- (3) The derived functor $\mathbf{R} \text{Hom}_{A^e}(-,-)$ is a bifunctor

$$
\text{RHom}_{A^e}(-,-): \mathcal{D}_{perf}(A^e)^{op} \times \mathcal{D}^+(B^e \mathcal{M}_{B^e}^{H^e}) \longrightarrow \mathcal{D}(B^e \mathcal{M}^{H^e}),
$$

and $\mathrm{Ext}^n_{A^e}(Y,X) \cong \mathrm{H}^n(\mathbf{R}\mathrm{Hom}_{A^e}(Y,X)).$

Proof. (1) Since $_{Be}\mathcal{M}_{Be}^{He}$ has enough injective objects by Lemma [2.9,](#page-11-1) the right derived functor

$$
\operatorname{\mathbf{R}Hom}\nolimits_{A^e}(A,-):{\mathcal D}^+({}_B{}_e{\mathcal M}^{H^e}_{B^e})\longrightarrow{\mathcal D}({}_{B^e}{\mathcal M}^{H^e}_H)
$$

exists as per Theorem $[33,$ Theorem 10.5.6. Given that B is flat as both a right and a left A-module, the forgetful functor from ${}_{B^e}\mathcal{M}_{B^e}^{H^e}$ to \mathcal{M}_{A^e} maintains injective objects. Then the diagram's commutativity follows from the uniqueness of the derived functor.

(2) For any $Y \in \mathcal{D}_{perf}(A^e)$, it is quasi-isomorphic to a bounded complex P of finitely generated projective A^e -modules. The conclusion follows from

$$
\begin{aligned} \mathbf{R}\mathrm{Hom}_{A^e}(Y, X) &\cong \mathrm{Hom}_{A^e}(P, X) \\ &\cong X \otimes_{A^e} \mathrm{Hom}_{A^e}(P, A^e) \\ &\cong X^{\mathbf{L}} \otimes_{A^e} \mathrm{Hom}_{A^e}(P, A^e) \\ &\cong X^{\mathbf{L}} \otimes_{A^e} \mathbf{R}\mathrm{Hom}_{A^e}(Y, A^e). \end{aligned}
$$

(3) This follows from a similar proof of $[33,$ Theorem 10.7.4]. \Box

2.6. Skew Calabi-Yau property of flat Hopf-Galois extensions. In this subsection, let H be a Hopf algebra with a bijective antipode, B be a faithfully flat H-Galois extension of A.

Proposition 2.12. If A is homologically smooth, then

$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, B^e) \cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B \otimes H \text{ in } \mathcal{D}(\mathcal{M}_{B \otimes H}).
$$

Further, if H is also homologically smooth, then

 $\mathbf{R}\text{Hom}_{B^e}(B, B^e) \cong \mathbf{R}\text{Hom}_{A^e}(A, A^e) \otimes_A B \otimes \mathbf{R}\text{Hom}_H(\mathbb{k}, H)$ in $\mathcal{D}(\mathcal{M}_B)$.

Proof. Since B is faithfully flat both as a left and right A-modules, it follows that B^e is faithfully flat as an A^e-module. **R**Hom_A^e(A, $\overline{B^e}$) is a complex in $\mathcal{D}(B^e \mathcal{M}_H^{He})$ by Proposition [2.11](#page-12-0) (1). Given that A is homologically smooth, Proposition 2.11 (2), yields, in $\mathcal{D}(B_e \mathcal{M}^{H^e}),$

(2.17)
\n
$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, B^e) \cong B^e \mathbf{L}_{\otimes_{A^e}} \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e)
$$
\n
$$
\cong B^e \otimes_{A^e} \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e)
$$
\n
$$
\cong B \otimes_A (\mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B).
$$

Viewing $\mathbf{R} \text{Hom}_{A^e}(A, B^e)$ as a complex in $\mathcal{D}(B^e \mathcal{M}^H)$, Lemma [2.8](#page-11-2) (2) gives

(2.18)
$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, B^e)^{coH} \stackrel{(2.17)}{\cong} (B \otimes_A \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B)^{coH}
$$

$$
\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B
$$

in $\mathcal{D}_{A\otimes B^{op}}\mathcal{M}$). Further, by Lemma [2.8](#page-11-2) (1), in $\mathcal{D}_{A\otimes B^{op}}\mathcal{M}_{H}^{H}$), we have

(2.19)
\n
$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, B^e) \cong \mathbf{R}\mathrm{Hom}_{A^e}(A, B^e)^{coH} \otimes H
$$
\n
$$
\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B \otimes H.
$$

Considering $\mathbf{R}\text{Hom}_{A^e}(A, B^e)$ in $\mathcal{D}(B^{op}\mathcal{M}_H)(=\mathcal{D}(\mathcal{M}_{B\otimes H}))$, Corollary [2.4](#page-8-1) yields

 $\mathbf{R}\text{Hom}_{B^e}(B, B^e) \cong \mathbf{R}\text{Hom}_H(\mathbb{k}, \mathbf{R}\text{Hom}_{A^e}(A, B^e))$ $\overset{(2.19)}{\cong} \mathbf{R}\mathrm{Hom}_{H}(\Bbbk,\mathbf{R}\mathrm{Hom}_{A^{e}}(A,A^{e})\otimes_{A}B\otimes H)$ $\overset{(2.19)}{\cong} \mathbf{R}\mathrm{Hom}_{H}(\Bbbk,\mathbf{R}\mathrm{Hom}_{A^{e}}(A,A^{e})\otimes_{A}B\otimes H)$ $\overset{(2.19)}{\cong} \mathbf{R}\mathrm{Hom}_{H}(\Bbbk,\mathbf{R}\mathrm{Hom}_{A^{e}}(A,A^{e})\otimes_{A}B\otimes H)$ (since H is homologically smooth.) \cong (**R**Hom_A^e(*A*, *A*^e) ⊗_A *B* ⊗ *H*)^{**L**}⊗_H **R**Hom_H(**k**, *H*) \cong **R**Hom_A^e(*A*, *A*^e) ⊗_A *B* ⊗ **R**Hom_{*H*}(**k**, *H*)

in $\mathcal{D}(B)$. This completes the proof. \Box

Similarly, we obtain the following result.

Proposition 2.13. If A is homologically smooth, then

$$
\mathbf{R}\mathrm{Hom}_{A^e}(A, B^e) \cong B \otimes_A \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes H \text{ in } \mathcal{D}({}_B\mathcal{M}_H).
$$

Further, if H is also homologically smooth, then

 $\mathbf{R}\text{Hom}_{B^e}(B, B^e) \cong B \otimes_A \mathbf{R}\text{Hom}_{A^e}(A, A^e) \otimes \mathbf{R}\text{Hom}_H(\mathbb{k}, H)$ in $\mathcal{D}_B(\mathcal{M})$.

Liu, Wu, and Zhu demonstrated in [\[20\]](#page-41-3) that for any H-module algebra R, the smash product $R\#H$ is homologically smooth if both R and H are homologically smooth. Their proof is directly applicable to Hopf Galois extensions with no modifications. Consequently, we can assert the following analogous result.

Proposition 2.14. [\[20,](#page-41-3) Proposition 2.11] Let $A \subseteq B$ be a flat H-Galois extension. If A and H are homologically smooth, then so is B .

Now we can prove our main result in this paper.

Theorem 2.15. Let H be a n-dim skew Calabi-Yau Hopf algebra, and $A \subseteq B$ be a faithfully flat H-Galois extension.

- (1) If A has d-dim VdB duality, then B has $(d + n)$ -dim VdB duality.
- (2) If A is d-dim skew Calabi-Yau, then B is $(d + n)$ -dim skew Calabi-Yau.

Proof. Let ω be a VdB dual module of A. Hence \mathbf{R} Hom_{A^e} $(A, A^e) \cong \omega[-d]$. By Proposition [2.12,](#page-12-1) in $\mathcal{D}(\mathcal{M}_B)$,

$$
\begin{aligned} \mathbf{R}\mathrm{Hom}_{B^e}(B, B^e) &\cong \mathbf{R}\mathrm{Hom}_{A^e}(A, A^e) \otimes_A B \otimes \mathbf{R}\mathrm{Hom}_H(\mathbb{k}, H) \\ &\cong \omega[-d] \otimes_A B \otimes \mathbf{R}\mathrm{Hom}_H(\mathbb{k}, H) \\ &\cong \omega \otimes_A B[-d - n]. \end{aligned}
$$

Similarly, we can prove that in $\mathcal{D}(B\mathcal{M})$,

$$
\mathbf{R}\mathrm{Hom}_{B^e}(B,B^e) \cong B \otimes_A \omega[-d-n].
$$

Then the conclusion follows from Proposition [1.7.](#page-5-0) \Box

In the aforementioned theorem, we did not answer the following question.

Question 2.16. What is the precise form of the Nakayama automorphism of B in Theorem [2.15](#page-14-0) (2) ?

Note that the Nakayama automorphism of any commutative skew Calabi-Yau algebra is the identity map, meaning such algebras are always Calabi-Yau. Hence we have the following result.

Corollary 2.17. Let H be a *n*-dim skew Calabi-Yau Hopf algebra with a bijective antipode, and $A \subseteq B$ be a faithfully flat H-Galois extension of commutative algebras. If A is a Calabi-Yau algebra, then so is B.

The geometric version of this corollary states that any principal fiber bundle over a Calabi-Yau variety with a smooth affine algebraic group is also Calabi-Yau. However, the noncommutative Hopf Galois extension does not preserve the Calabi-Yau property, even when the coinvariant subalgebra is commutative. See the following example.

Example 2.18. Let $\mathfrak{g} = \mathbb{k} x \oplus \mathbb{k} y$ be a Lie algebra with $[y, x] = x$, B be the enveloping algebra $\mathcal{U}(\mathfrak{g})$ of \mathfrak{g} . The quotient Hopf algebra $B/(x)$ is denoted by H. Then B is a right H-Galois extension of coinvariant subalgebra $A = \mathbb{k}[x]$. The Nakayama automorphism μ of B is given by $\mu(x) = x, \mu(y) = y + 1$ (see [\[36\]](#page-41-17) or $[5]$, which is not a inner automorphism. Although both H and A are Calabi-Yau algebras, B is not a Calabi-Yau algebra.

A flat (but not necessarily faithfully flat) Hopf Galois extension may maintain the skew Calabi-Yau property, as demonstrated by the following example.

Example 2.19. [\[3,](#page-40-10) Example 2.4] Consider the Hopf algebra $H = \mathbb{k}[x]$, where x a primitive element. Define

$$
B = \mathcal{O}(SL(2)) = \mathbb{k}[a, b, c, d]/(ad - bc - 1),
$$

which is a 3-dim Calabi-Yau algebra. B is an H -comodule algebra structure via the coaction $\rho: B \to B \otimes H$ given on generators by

 $\rho(a) = a \otimes 1 + c \otimes x$, $\rho(b) = b \otimes 1 + d \otimes x$, $\rho(c) = c \otimes 1$, $\rho(d) = d \otimes 1$.

Note that $B^{coH} = A = \mathbb{k}[c, d]$. Then $A \subseteq B$ is a flat not faithfully flat H-Galois extension, as detailed in [\[3,](#page-40-10) Example 2.4].

We do not know whether Theorem [2.15](#page-14-0) still holds in the flat case.

Question 2.20. Let H be a skew Calabi-Yau Hopf algebra with a bijective antipode, and $A \subseteq B$ be a flat H-Galois extension. If A is skew Calabi-Yau, must B also be skew Calabi-Yau ?

2.7. Nakayama automorphisms for certain Hopf Galois extensions. We now present the Nakayama automorphism for specific instances of Hopf-Galois extensions.

2.7.1. Nakayama automorphisms of α -Frobenius extensions. Kreimer and Takeuchi demonstrated that every Hopf-Galois extension over a finite-dimensional Hopf al-gebra is always a Frobenius extension [\[18,](#page-41-18) Theorem 1.7]. Recall the notion of α -Frobenius extensions, introduced by Nakayama and Tsuzuku [\[25,](#page-41-19)[26\]](#page-41-20), which are also referred to as Frobenius extensions of the second kind. These extensions represent a generalization of the classical Frobenius extensions.

Definition 2.21. Consider an algebra extension $A \subseteq B$ and an algebra automorphism $\alpha : A \to A$. The extension $A \subseteq B$ is termed an α -Frobenius extension if it satisfies

- (1) B is a finitely generated projective right A-module, and
- (2) B is isomorphic to $_{\alpha}$ Hom $_{A}(B, A)$ as A-B-bimodules.

This is equivalent to saying that B is a projective left A -module, and B is isomorphic to $\text{Hom}_{A^{op}}(B, A)_{\alpha^{-1}}$ as a B-A-bimodule. If α is the identity map on A, then $A \subseteq B$ is referred to as a Frobenius extension.

Lemma 2.22. Let A be a subalgebra of an algebra B. Suppose that both A and B are homologically smooth.

(1) If B is a pseudo-coherent right A-module, then in $\mathcal{D}(B \otimes A^{op}),$

 $\mathbf{R}\mathrm{Hom}_A(B,A) \mathbf{L}_{\bigotimes_B} \mathbf{R}\mathrm{Hom}_{B^e}(B,B^e) \cong \mathbf{R}\mathrm{Hom}_{A^e}(A,A^e) \mathbf{L}_{\bigotimes_A B}.$

(2) If B is a pseudo-coherent left A-module, then in $\mathcal{D}(A \otimes B^{op}),$

 $\mathbf{R}\mathrm{Hom}_{B^e}(B,B^e) \perp \otimes_B \mathbf{R}\mathrm{Hom}_{A^{op}}(B,A) \cong B^{\mathbf{L}} \otimes_A \mathbf{R}\mathrm{Hom}_{A^e}(A,A^e).$

Proof. (1) In $\mathcal{D}(B \otimes A^{op})$, we have the following isomorphisms

 $\operatorname{\mathbf{R}Hom}\nolimits_{A}(B, A) \operatorname{\mathbf{L}}\nolimits_{\otimes_{B} \operatorname{\mathbf{R}Hom}\nolimits_{B^{e}}\nolimits(B, B^{e})$

 \cong RHom_{Be} $(B, \text{RHom}_{A}(B, A) \otimes B)$ since B is homologically smooth

 \cong **RH**om_{B^{e}}(B, **RH**om_A(B, A \otimes B)) since B is a pseudo-coherent A-module</sub>

 \cong **RHom**_{A⊗B}_{op} (B, A ⊗ B)

 \cong **RH**om_{Ae} $(A, \mathbf{R}$ Hom_{Bop} $(B, A \otimes B))$

- \cong **RHom**_{Ae} $(A, A \otimes B)$
- \cong **R**Hom_Ae (A, A^e) since A is homologically smooth.

(2) The proof for the second part is analogous to the first. \Box

We will show that any homologically smooth α -Frobenius extension B of a skew Calabi-Yau algebra A inherently possesses the skew Calabi-Yau property. Furthermore, we will establish that the restriction of a Nakayama automorphism of B to A constitutes a Nakayama automorphism of A.

Proposition 2.23. Let $A \subseteq B$ be an α -Frobenius extension. Suppose that B is homologically smooth. If A is a skew Calabi-Yau algebra of dimension d , then B is also a skew Calabi-Yau algebra of dimension d. Moreover, there exists a Nakayama automorphism μ of B such that $(\mu \circ \alpha)|_A$ is a Nakayama automorphism of A.

Proof. Since $A \subseteq B$ is an α -Frobenius extension for some $\alpha \in Aut(A)$, we have ${}_{\alpha}$ Hom_A $(B, A) \cong B$ as A-B-bimodules and Hom_{A^{op}} $(B, A)_{\alpha^{-1}} \cong B$ as B-A-bimodules.</sub> By Lemma 2.22 , we obtain the following isomorphisms of $A-B$ -bimodules

$$
\text{Ext}_{B^e}^i(B, B^e) \cong {}_{\alpha}\text{Hom}_A(B, A) \otimes_B \text{Ext}_{B^e}^i(B, B^e) \cong {}_{\alpha}\text{Ext}_{A^e}^i(A, A^e) \otimes_A B.
$$

Given that A is a skew Calabi-Yau algebra of dimension d with a Nakayama automor-

phism μ_A , we find $\text{Ext}_{B^e}^i(B, B^e) \cong$ $\int 0, \quad i \neq d$ $_{\alpha\mu_A^{-1}}B, \quad i=d$ as A-B-bimodules. Similarly,

 $\operatorname{Ext}_{B^e}^i(B, B^e) = \begin{cases} 0, & i \neq d \\ B & i \neq d \end{cases}$ $B_{\mu_A\alpha^{-1}}, \quad i = d$ as $B-A$ -bimodules. This implies that B is a skew

Calabi-Yau algebra by Proposition [1.7.](#page-5-0) Let μ_B be a Nakayama automorphism of B. The B-A-bimodule $B_{\mu_B\alpha}$ is isomorphic to B_{μ_A} . Thus, there exists an invertible element $u \in B$ for which $u^{-1}\mu_B(\alpha(a))u = \mu_A(a)$ holds for all $a \in A$. Defining μ by $b \mapsto u^{-1}\mu_B(b)u$, we confirm that μ is also a Nakayama automorphism of B, and $(\mu \circ \alpha)|_A$ is a Nakayama automorphism of A.

Applying the previous proposition to a Hopf Galois extension over a finite dimensional Hopf algebra, we obtain the following results.

Corollary 2.24. Let H be a finite dimensional semisimple Hopf algebra, and $A \subseteq B$ be an H-Galois extension. If A is a skew Calabi-Yau algebra of dimension d, then B is also a skew Calabi-Yau algebra of dimension d , equipped with a Nakayama automorphism μ such that $\mu|_A$ is a Nakayama automorphism of A.

Proof. Given that the H-Galois extension $A \subseteq B$ is a Frobenius extension and B is homologically smooth by Proposition [2.14,](#page-13-3) the result is a direct consequence of Proposition [2.23.](#page-16-0) \Box

Additionally, we establish the following connection between an α -Frobenius extension and the possession of a VdB dual module.

Proposition 2.25. Let $A \subseteq B$ be an α -Frobenius extension. Suppose that A is homologically smooth, and B is a faithfully flat A-module on both sides. If B has a VdB dual module of dimension d, then A also has one.

Proof. Given that B is faithfully flat as a right A-module, Lemma [2.22](#page-15-0) yields

$$
\text{Ext}_{B^e}^i(B, B^e) \cong {}_{\alpha}\text{Hom}_A(B, A) \otimes_B \text{Ext}_{B^e}^i(B, B^e) \cong {}_{\alpha}\text{Ext}_{A^e}^i(A, A^e) \otimes_A B
$$

as A-B-bimodules. This implies $\mathrm{Ext}_{A^e}^i(A, A^e) = 0$ for all $i \neq d$ and $\mathrm{Ext}_{A^e}^d(A, A^e) \otimes_A$ $B \cong \alpha^{-1} \text{Ext}_{B^e}^d(B, B^e)$. Write $\omega = \text{Ext}_{A^e}^d(A, A^e)$. Since A is a direct summand of B as a left A-module by [\[28,](#page-41-21) Proposition 2.11.29], ω is a direct summand of the projective left A-module $_{\alpha^{-1}}\text{Ext}_{B^e}^d(B, B^e)$. Thus, ω is a finitely generated projective left A-module. Analogously, ω is a finitely generated projective right A-module. According to Proposition [1.7,](#page-5-0) A has a VdB dual module ω of dimension d. \Box

We now proceed to demonstrate that an algebra extension between skew Calabi-Yau algebras is nearly an α -Frobenius extension.

Proposition 2.26. Let $A \subseteq B$ be an algebra extension. Suppose that A and B are both skew Calabi-Yau algebras of dimension d . If B is a pseudo-coherent right A module, then B is a projective right A -module, and there exists an algebra morphism $\alpha: A \to B$ such that $_{\alpha}$ Hom $_{A}(B, A) \cong B$ as A-B-bimodules.

Proof. Let μ_A and μ_B be Nakayama automorphisms of A and B, respectively. By Lemma [2.22,](#page-15-0) in $\mathcal{D}(B \otimes A^{op}),$

$$
\operatorname{RHom}_A(B, A) \mathbf{L}_{\otimes_B} B_{\mu_B}[-d] \cong A_{\mu_A}[-d] \mathbf{L}_{\otimes_A} B.
$$

This yields $\mu_B^{-1} \mu_A \text{Hom}_A(B, A) \cong B$ as A-B-bimodules and $\text{Ext}^i_A(B, A) = 0$ for any $i > 0$. Let n be the projective dimension of the right A-module B. Since A has finite global dimension by our assumption, there exists a right A -module N , such that $\text{Ext}_{A}^{n}(B, N) \neq 0$. Let F be a free A-module with a surjective A-module morphism $F \to N$. This induces a short exact sequence $\text{Ext}_{A}^{n}(B, F) \to \text{Ext}_{A}^{n}(B, N) \to 0$, leading to $\text{Ext}_{A}^{n}(B, F) \neq 0$. As B is a pseudo-coherent right A-module, we have $\text{Ext}_{A}^{n}(B,A)\otimes_{A}^{A}F\cong \text{Ext}_{A}^{n}(B,F)\neq 0.$ Consequently, $n=0$, indicating that B is a projective right A-module. \Box

2.7.2. Nakayama automotphisms of Hopf Galois extensions of strongly separable algebras. In the rest of this section, we maintain the assumption that the coinvariant ring $A := B^{coH}$ is a strongly separable algebra over k, that is, there exists an element $e := \sum_{n=1}^{\infty}$ $i=1$ $e'_i \otimes e''_i \in A^e$ satisfying

$$
\sum_{i=1}^{n} ae'_i \otimes e''_i = \sum_{i=1}^{n} e'_i \otimes e''_i a, \qquad \sum_{i=1}^{n} e'_i e''_i = 1, \qquad \sum_{i=1}^{n} e''_i \otimes e'_i = e.
$$

Such an e is referred to as a symmetric separability idempotent of A^e .

It is evident that $\mathbb k$ is strongly separable over itself. Any Hopf Galois extension of the base field k is commonly termed a Hopf Galois object of H . In [\[39\]](#page-41-22), Yu demonstrated that Hopf-Galois objects of a skew Calabi-Yau Hopf algebra retain their skew Calabi-Yau nature and explicitly provided their Nakayama automorphism.

We will use the shorthand notation $x \otimes y$ and $x \otimes_A y$ to represent arbitrary elements in $B \otimes B$ and $B \otimes_A B$, respectively. Before introducing the Nakayama automorphisms of Hopf Galois extensions of strongly separable algebras, we require the following result.

Lemma 2.27. [\[1,](#page-40-11) Theorem 4.1] The map $\tau : (B \otimes B)^A \to B \otimes_A B$, $x \otimes y \mapsto y \otimes_A x$ is bijective.

Observe that $B \otimes_A B$ is also equipped with a right H-module defined by

(2.20)
\n
$$
(x \otimes_A y) \leftarrow h := \tau(\tau^{-1}(x \otimes_A y) \leftarrow h) = \tau((y \otimes x) \leftarrow h)
$$
\n
$$
\xrightarrow{\text{(2.8)}} \tau(\sum \kappa^1(h)y \otimes x\kappa^2(h))
$$
\n
$$
= \sum x\kappa^2(h) \otimes_A \kappa^1(h)y.
$$

Thus, τ preserves the H-module structure.

In the subsequent discussion, we posit that

$$
\sum \kappa^2(h) \otimes \kappa^1(h) = \tau^{-1} \beta^{-1} (1 \otimes h) \in (B \otimes B)^A.
$$

The following are useful properties of κ :

(2.21)
$$
\sum \kappa^2(hk) \otimes \kappa^1(hk) = \sum \kappa^2(h)\kappa^2(k) \otimes \kappa^1(k)\kappa^1(h),
$$

(2.22)
$$
\sum \kappa^2(h)_0 \otimes \kappa^1(h) \otimes \kappa^2(h)_1 = \sum \kappa^2(h_1) \otimes \kappa^1(h_1) \otimes h_2,
$$

(2.23)
$$
\sum \kappa^2(h) \otimes \kappa^1(h)_0 \otimes \kappa^1(h)_1 = \sum \kappa^2(h_2) \otimes \kappa^1(h_2) \otimes Sh_1.
$$

Lemma 2.28. If V is a left H-module, then $B \otimes V$ is a B-B-bimodule which is defined by

$$
b'(x \otimes v)b = \sum b' x \kappa^{2} (S^{-2}b_1) \kappa^{1} (S^{-2}b_1) b_0 \otimes S^{-1}b_2 \to v.
$$

There is a natural isomorphism $\Phi : (B \otimes_A B) \otimes_H V \to B \otimes V$ in ${}_B\mathcal{M}_B$, given by

$$
\Phi(x \otimes_A y \otimes_H v) = \sum x \kappa^2 (S^{-2} y_1) \kappa^1 (S^{-2} y_1) y_0 \otimes S^{-1} y_2 \to v,
$$

Proof. First, we demonstrate that Φ is well defined. Given $\kappa^2(h) \otimes \kappa^1(h) \in (B \otimes B)^A$, it is inferred that $\kappa^2(h)\kappa^1(h)$ is an element of the center of B. Then we have

$$
\Phi(x \otimes_A ay \otimes_H v) = \sum x \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) ay_0 \otimes S^{-1}y_2 \to v
$$

=
$$
\sum x a \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) y_0 \otimes S^{-1}y_2 \to v
$$

=
$$
\Phi(x \otimes_A ay \otimes_H v)
$$

for any $a \in A$. For any $h \in H$, it holds that

$$
\Phi((x \otimes_A y) \leftarrow h \otimes_H v)
$$
\n
$$
\stackrel{(2.20)}{=} \Phi\left(\sum x\kappa^2(h) \otimes_A \kappa^1(h)y \otimes_H v\right)
$$
\n
$$
\stackrel{(2.23)}{=} \sum x\kappa^2(h_3)\kappa^2((S^{-1}h_2)(S^{-2}y_1))\kappa^1((S^{-1}h_2)(S^{-2}y_1))\kappa^1(h_3)y_0 \otimes (h_1S^{-1}y_2) \to v
$$
\n
$$
\stackrel{(2.21)}{=} \sum x\kappa^2(S^{-2}y_1)\kappa^1(S^{-2}y_1)y_0 \otimes h \to (S^{-1}y_2 \to v)
$$
\n
$$
= \Phi(x \otimes_A y \otimes_H (h \to v)).
$$

Next, we show that $\Psi : B \otimes V \to (B \otimes_A B) \otimes_H V$, $x \otimes v \mapsto x \otimes_A 1 \otimes_H v$ is the inverse of Φ as k-linear maps. One can easily to check that $\Phi \Psi = id_{B \otimes V}$. Note that

$$
(\beta \otimes id_H) \Big(\sum x \kappa^2 (S^{-2}y_1) \otimes_A \kappa^1 (S^{-2}y_1) y_0 \otimes y_2 \Big)
$$

= $\sum x \kappa^2 (S^{-2}y_2) \kappa^1 (S^{-2}y_2)_{0} y_0 \otimes \kappa^1 (S^{-2}y_2)_{1} y_1 \otimes y_3$

$$
\xrightarrow{(2.23)} \sum x \kappa^2 (S^{-2}y_3) \kappa^1 (S^{-2}y_3) y_0 \otimes (S^{-1}y_2) y_1 \otimes y_4
$$

= $\sum x \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) y_0 \otimes 1 \otimes y_2$
= $(\beta \otimes id_H) \Big(\sum x \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) y_0 \otimes 1 \otimes y_2 \Big).$

Since the canonical map β is bijective, it follows that

(2.24)
$$
\sum x \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) y_0 \otimes_A 1 \otimes y_2
$$

$$
= \sum x \kappa^2 (S^{-2}y_1) \otimes_A \kappa^1 (S^{-2}y_1) y_0 \otimes y_2.
$$

On the other hand, we can verify that

$$
\Psi\Phi(x \otimes y \otimes_H v) = \sum x \kappa^2 (S^{-2}y_1) \kappa^1 (S^{-2}y_1) y_0 \otimes_A 1 \otimes_H S^{-1}y_2 \to v
$$

$$
\xrightarrow{\underline{(^{2.24)}}} \sum x \kappa^2 (S^{-2}y_1) \otimes_A \kappa^1 (S^{-2}y_1) y_0 \otimes_H S^{-1}y_2 \to v
$$

$$
\xrightarrow{\underline{(^{2.20)}}} \sum x \otimes_A y_0 \otimes_H S^{-2}y_1 \to (S^{-1}y_2 \to v)
$$

$$
= x \otimes_A y \otimes_H v.
$$

Hence $\Psi \Phi = \mathrm{id}_{(B \otimes_A B) \otimes_H V}$. So $B \otimes V$ is a B-B-bimodule via

$$
b'(x \otimes v)b := \Phi(b'\Psi(x \otimes v)b) = \Phi(b'(x \otimes A \otimes v)b) = \Phi(b'x \otimes b \otimes_H v)
$$

=
$$
\sum b'x\kappa^2(S^{-2}b_1)\kappa^1(S^{-2}b_1)b_0 \otimes S^{-1}b_2 \to v,
$$

and Φ is an isomorphism of B-B-bimodules. \Box

The following lemma follows immediately from Lemma [2.28.](#page-18-3)

Lemma 2.29. The right H-module $B \otimes_A B$ is faithfully flat.

Theorem 2.30. Let H be a skew Calabi-Yau algebra of dimension d, and B be an H -Galois extension of a strongly separable algebra A .. Then B is skew Calabi-Yau of dimension d with a Nakayama automorphism μ , which is defined by

$$
\mu(b) = \sum \kappa^2 (S^{-2}b_1) \kappa^1 (S^{-2}b_1) b_0 \chi(Sb_2),
$$

where $\chi: H \to \mathbb{k}$ is given by $\mathrm{Ext}^d_H(\mathbb{k}, H) \cong {}_\chi \mathbb{k}$.

Proof. Since A is separable, A is a direct summand of B as an A-module. Hence, B is a faithfully flat A-module. By Theorem [2.15,](#page-14-0) B is a skew Calabi-Yau algebra of dimension d. Consider the following isomorphisms:

$$
\begin{aligned}\n&\mathbf{R}\text{Hom}_{B^e}(B, B^e) \\
\cong \mathbf{R}\text{Hom}_H(\mathbb{k}, \mathbf{R}\text{Hom}_{A^e}(A, B^e)) \\
&\cong \mathbf{R}\text{Hom}_{A^e}(A, B^e) \cdot \mathbf{L}_{\otimes_H} \mathbf{R}\text{Hom}_H(\mathbb{k}, H)\n\end{aligned}\n\quad \text{by Theorem 2.3}
$$

$$
\cong \text{Hom}_{A^e}(A, B^e) \cdot^L \otimes_H \mathbf{R} \text{Hom}_H(\mathbb{k}, H) \qquad \text{since } A \text{ is (strongly) separable} \n\cong (B \otimes_A B) \cdot^L \otimes_H \mathbf{R} \text{Hom}_H(\mathbb{k}, H) \qquad \text{by Lemma 2.27} \n\cong (B \otimes_A B) \otimes_H \text{Ext}^d_H(\mathbb{k}, H)[-d] \qquad \text{since } H \text{ is skew Calabi-Yau} \n\cong (B \otimes_A B) \otimes_H \text{Ext}^d_H(\mathbb{k}, H)[-d] \qquad \text{by Lemma 2.28} \n\cong B \otimes \text{Ext}^d_H(\mathbb{k}, H)[-d]
$$

Thus, there there is an algebra endomorphism μ of B such that $B \otimes \text{Ext}_{H}^{d}(\mathbb{k}, H) \cong B_{\mu}$ as B - B -bimodules, where μ is given by

$$
\mu(b) = \sum \kappa^2 (S^{-2}b_1) \kappa^1 (S^{-2}b_1) b_0 \chi(S^{-1}b_2) = \sum \kappa^2 (S^{-2}b_1) \kappa^1 (S^{-2}b_1) b_0 \chi(Sb_2)
$$

for all $b \in B$. Since B is a skew Calabi-Yau algebra, it follows that B_μ is an invertible B-B-bimodule. This implies that μ is an automorphism. Then μ is a Nakayama automorphism of B. \Box

3. Some structure theorems of Hopf bimodule categories

Throughout this section, H is a Hopf algebra with a bijective antipode S , and $A \subseteq B$ is a faithfully flat H-Galois extension.

3.1. A structure theorem of Hopf bimodule category. Let R and T be two right H-comodule algebra. Put $L(R, T, H) := (R \otimes T)^{coH}$. We will abuse notation by using simple tensors $s \otimes t$ instead of arbitrary elements of $R \otimes T$, and note for later use that

(3.1)
$$
L(R,T,H) = \{r \otimes t \mid \sum r_0 \otimes t_0 \otimes r_1 t_1 = r \otimes t \otimes 1\}
$$

$$
= \{r \otimes t \mid \sum r_0 \otimes t \otimes Sr_1 = \sum r \otimes t_0 \otimes t_1\}.
$$

as in [\[31,](#page-41-15) Lemma 3.1].

It is straightforward to confirm that $L(R, T, H)$ is a subalgebra of $R \otimes T^{op}$, with the multiplication defined as $(r \otimes t)(r' \otimes t') = rr' \otimes t't$. In general, $L(R, T, H)$ does not form a subalgebra of $R \otimes T$. Obviously, $R^{coH} \otimes (T^{coH})^{op}$ is contained within $L(R, T, H)$ as a subalgebra.

Let us revisit Schauenburg's structure theorem for relative Hopf bimodules which is a generalization of Theorem [2.2.](#page-5-3)

Theorem 3.1. [\[29,](#page-41-23) Theorem 3.3] Let H be a Hopf algebra and B a right faithfully flat H-Galois extension of $A := B^{coH}$. Let R be a right H-comodule algebra, and put $L := L(R, B, H)$. Then we have quasi-inverse category equivalences

$$
{}_{R}\mathcal{M}_{B}^{H} \longrightarrow {}_{L}\mathcal{M}, \ N \mapsto N^{coH}, \qquad {}_{L}\mathcal{M} \longrightarrow {}_{R}\mathcal{M}_{B}^{H}, \ M \mapsto M \otimes_{A} B,
$$

where

(1) the L-module structure of N^{coH} is defined by

$$
l \rhd n = rnb
$$
, $\forall l = r \otimes b \in L \subseteq R \otimes B^{op};$

(2) for $M \in L^1$ we use \triangleright to denote the module structure, N is a right A-module via $ma = (1 \otimes a) \triangleright m$, and the right B-module and H-comodule structures

on $M \otimes_A B$ are those induced by the right tensorand, and the left R-module stucture is given by

$$
r(m \otimes_A b) = \sum (r_0 \otimes \kappa^1(r_1)) \rhd m \otimes_A \kappa^2(r_1) b.
$$

Theorem 3.2. [\[29,](#page-41-23) Theorem 3.3] Let R be a right H-comodule algebra, and put $L := L(B, R, H)$. Then we have quasi-inverse category equivalences

$$
{}_B\mathcal{M}_R^H \longrightarrow {}_L\mathcal{M}, \ N \mapsto N^{coH}, \qquad {}_L\mathcal{M} \longrightarrow {}_B\mathcal{M}_R^H, \ N \mapsto B \otimes_A M,
$$

where

(1) the L-module structure of N^{coH} is defined by

$$
l \rhd n = bnr
$$
, $\forall l = b \otimes r \in L \subseteq B \otimes R^{op}$;

(2) for $N \in L^{\mathcal{M}}$ we use \triangleright to denote the module structure, M is a left A-module via $am = (a \otimes 1) \triangleright m$, and the left B-module and H-comodule structures on $B \otimes_A M$ are those induced by the left tensorand, and the right R-module stucture is given by

(3.2)
$$
(b \otimes_A m)r = \sum b\kappa^1(S^{-1}r_1) \otimes_A (\kappa^2(S^{-1}r_1) \otimes r_0) \rhd m.
$$

The following lemmas will be required subsequently.

Lemma 3.3. Let H be a Hopf algebra, B a right faithfully flat H -Galois extension of $A := B^{coH}$, and R a right H-comodule algebra. Then

- (1) $L(B, R, H)$ is projective as a left A-module, and
- (2) $L(R, B, H)$ is projective as a right A-module.

Proof. (1) By Theorem [2.2,](#page-5-3) a right H-colinear and left A-linear map $g : B \to A \otimes B$ exists, satisfying $g(ab) = a \otimes b$ for any $a \in A, b \in B$. Consequently, a right Hcolinear and left A-linear map $g \otimes id_R : B \otimes R \to A \otimes B \otimes R$ is induced, with the property that $(g \otimes id_R)(ab \otimes r) = a \otimes b \otimes r$. This yields $L(B, R, H) = (B \otimes R)^{coH}$ as a direct summand of the free A-module $(A \otimes B \otimes R)^{coH} = A \otimes L(B, R, H)$. Therefore, $L(B, R, H)$ is a projective left A-module.

(2) Applying (1) to the faithfully flat H^{op} -Galois extension $A^{op} \subseteq B^{op}$, we deduce that $L(R, B, H)$ which equals $L(B^{op}, R^{op}, H^{op})$, is projective as a left A^{op} -module, and consequently, as a right A-module. \Box

Lemma 3.4. Let H be a Hopf algebra, B a right faithfully flat H -Galois extension of $A := B^{coH}$, and R a right H-comodule algebra.

(1) If M is a left $L(R, B, H)$ -module, then $\text{Hom}_A(M, A)$ is a left $L(B, R, H)$ module, defined by

(3.3)
$$
((b \otimes r) \triangleright f)(m) = \sum bf((r_0 \otimes \kappa^1(r_1)) \triangleright m)\kappa^2(r_1),
$$

for any $b \otimes r \in L(B, R, H)$, $f \in \text{Hom}_A(M, A)$ and $m \in M$.

(2) For any left $L(R, B, H)$ -module M and $N \in {}_B\mathcal{M}_B^H$, Hom $_A(M, N)$ is a Hopf bimodule in ${}_B\mathcal{M}_R^H$, where the right R-module is given by

(3.4)
$$
(fr)(m) = \sum f((r_0 \otimes \kappa^1(r_1)) \rhd m)\kappa^2(r_1),
$$

for any $b \otimes r \in L(B, R, H)$, $f \in \text{Hom}_A(M, N)$ and $m \in M$.

(3) The canonical map

$$
\Phi: N \otimes_A \text{Hom}_A(M, A) \longrightarrow \text{Hom}_A(M, N), \qquad n \otimes_A f \mapsto (m \mapsto nf(m))
$$

is a *B-R*-bimodule morphism.

Proof. (1) For any $r \in R$, we have

$$
\sum r_0 \otimes \kappa^1(r_2)_0 \otimes_A \kappa^2(r_2) \otimes r_1 \kappa^1(r_2)_1
$$

$$
\xrightarrow{(2.4)} \sum r_0 \otimes \kappa^1(r_3) \otimes_A \kappa^2(r_3) \otimes r_1 Sr_2
$$

$$
= \sum r_0 \otimes \kappa^1(r_1) \otimes_A \kappa^2(r_1) \otimes 1,
$$

which implies $\sum r_0 \otimes \kappa^1(r_1) \otimes_A \kappa^2(r_1) \in L(R, B, H) \otimes_A B$.

For any $b \otimes r \in L(B, R, H), f \in Hom_A(M, A)$ and $m \in M$, we assert that

$$
((b \otimes r) \rhd f)(m) = \sum bf((r_0 \otimes \kappa^1(r_1)) \rhd m)\kappa^2(r_1) \in A \subseteq B.
$$

This is shown by the equality

$$
((b \otimes r) \rhd f)(m)_0 \otimes ((b \otimes r) \rhd f)(m)_1
$$

= $\sum b_0 f((r_0 \otimes \kappa^1(r_1)) \rhd m) \kappa^2(r_1)_0 \otimes b_1 \kappa^2(r_1)_0$

$$
\xrightarrow{(2.3)} \sum b_0 f((r_0 \otimes \kappa^1(r_1)) \rhd m) \kappa^2(r_1) \otimes b_1 r_2
$$

= $\sum bf((r_0 \otimes \kappa^1(r_1)) \rhd m) \kappa^2(r_1) \otimes 1$ since $b \otimes r \in (B \otimes R)^{coH}$,

implying $\sum bf\Big(\big(r_0\otimes \kappa^1(r_1)\big)\rhd m\Big)\kappa^2(r_1)\in A.$ Given $a \in A$, we have

$$
((b \otimes r) \rhd f)(ma) = bf((r_0 \otimes \kappa^1(r_1)) \rhd (ma))\kappa^2(r_1)
$$

= $bf((r_0 \otimes a\kappa^1(r_1)) \rhd m)\kappa^2(r_1)$

$$
\xrightarrow{(2.2)} bf((r_0 \otimes \kappa^1(r_1)) \rhd m)\kappa^2(r_1)a
$$

= $((b \otimes r) \rhd f)(m)a,$

demonstrating that $(b \otimes r) \triangleright f$ is A-linear.

For any $b \otimes r, b' \otimes r' \in L(B, R, H)$, we calculate

$$
\begin{aligned}\n&\left((b \otimes r) \triangleright ((b' \otimes r') \triangleright f)\right)(m) \\
&= b\big((b' \otimes r')f\big)\Big((r_0 \otimes \kappa^1(r_1)) \triangleright m\Big) \kappa^2(r_1) \\
&= bb'f\Big((r'_0 \otimes \kappa^1(r'_1)) \triangleright (r_0 \otimes \kappa^1(r_1) \triangleright m)\Big) \kappa^2(r'_1) \kappa^2(r_1) \\
&= bb'f\Big((r'_0r_0 \otimes \kappa^1(r_1) \kappa^1(r'_1)) \triangleright m\Big) \kappa^2(r'_1) \kappa^2(r_1) \\
&\xrightarrow{(2.1)} bb'f\Big((r'_0r_0 \otimes \kappa^1(r'_1r_1)) \triangleright m\Big) \kappa^2(r'_1r_1) \\
&= \big((bb' \otimes r'r) \triangleright f\big)(m).\n\end{aligned}
$$

verifying that the module structure [\(3.3\)](#page-21-0) is well defined.

(2) Given any $f \in \text{Hom}_A(M, N)$, $r \in R$, $m \in M$ and $a \in A$.

$$
(fr)(ma) = f((r_0 \otimes \kappa^1(r_1)) \rhd (ma))\kappa^2(r_1)
$$

= $f((r_0 \otimes a\kappa^1(r_1)) \rhd m)\kappa^2(r_1)$

$$
\xrightarrow{(2.2)} f((r_0 \otimes \kappa^1(r_1)) \rhd m)\kappa^2(r_1)a
$$

= $(fr)(m)a$,

showing that fr is A-linear.

For any $r, r' \in R$, we find

$$
((fr)r')(m)
$$

= $(fr)\Big((r'_0 \otimes \kappa^1(r'_1)) \rhd m\Big) \kappa^2(r'_1)$
= $f\Big((r_0 \otimes \kappa^1(r_1)) \rhd (r'_0 \otimes \kappa^1(r'_1) \rhd m)\Big) \kappa^2(r_1) \kappa^2(r'_1)$
= $f\Big((r_0r'_0 \otimes \kappa^1(r'_1) \kappa^1(r_1)) \rhd m\Big) \kappa^2(r_1) \kappa^2(r'_1)$

$$
\xrightarrow{(2.1)} f\Big((r_0r'_0 \otimes \kappa^1(r_1r'_1)) \rhd m\Big) \kappa^2(r_1r'_1)
$$

= $(f(rr'))(m).$

confirming that the module structure [\(3.4\)](#page-21-1) is well defined.

(3) Since $\text{Hom}_A(M, A)$ is a left $L(B, R, H)$ -module, $B \otimes_A \text{Hom}_A(M, A)$ is a Hopf bimodule in ${}_B\mathcal{M}_R^H$ by Theorem [3.2.](#page-21-2) Consequently, $N \otimes_A \text{Hom}_A(M, A) \cong N \otimes_B B \otimes_A$ $\text{Hom}_A(M, A)$ is indeed a Hopf bimodule in ${}_B\mathcal{M}^H_R$.

It is evident that Φ is a left B-module morphism. To demonstrate that Φ is also a right R-module morphism, we calculate

$$
\Phi((n \otimes_A f)r)(m)
$$
\n
$$
\stackrel{(3.2)}{=} \Phi\left(\sum n\kappa^1 (S^{-1}r_1) \otimes_A (\kappa^2 (S^{-1}r_1) \otimes r_0) \triangleright f\right)(m)
$$
\n
$$
= \sum n\kappa^1 (S^{-1}r_1) \Big((\kappa^2 (S^{-1}r_1) \otimes r_0) \triangleright f\Big)(m)
$$
\n
$$
\stackrel{(3.3)}{=} \sum n\kappa^1 (S^{-1}r_2) \kappa^2 (S^{-1}r_2) f\Big((r_0 \otimes \kappa^1(r_1)) \triangleright m\Big) \kappa^2(r_1)
$$
\n
$$
\stackrel{(2.6)}{=} \sum n f\Big((r_0 \otimes \kappa^1(r_1)) \triangleright m\Big) \kappa^2(r_1)
$$
\n
$$
\stackrel{(3.4)}{=} (\Phi(n \otimes_{A^e} f)r)(m).
$$

This shows that Φ is indeed right R-linear. \Box

3.2. Some subalgebras Λ_i of B^e . To further study categories ${}_{B^e}\mathcal{M}_H^{H^e}$ and ${}_H\mathcal{M}_{B^e}^{H^e}$, we now introduce a class of algebras Λ_i .

Let (R, ρ) be a right H-comodule algebra and any Hopf algebra map $\sigma : H \to K$, then R also has a K-comodule strutrue via $(id_R \otimes \sigma) \circ \rho$ such that R is also a

right K-comodule algebra, denoted by R^{σ} . It is obvious that S^2 is a Hopf algebra automorphism of H.

Lemma 3.5. Let R and T be two right H -comodule algebra.

- (1) For any Hopf automorphism σ of H, $L(R, T, H) = L(R^{\sigma}, T^{\sigma}, H)$.
- $(2) L(R, T, H) \cong L(T, R^{S^2}, H)^{op}.$

Proof. (1) From (3.1) , we deduce that

$$
L(R, T, H) = \{r \otimes t \mid \sum r_0 \otimes t \otimes Sr_1 = \sum r \otimes t_0 \otimes t_1\}
$$

= $\{r \otimes t \mid \sum r_0 \otimes t \otimes \sigma Sr_1 = \sum r \otimes t_0 \otimes \sigma t_1\}$
= $\{r \otimes t \mid \sum r_0 \otimes t \otimes S\sigma r_1 = \sum r \otimes t_0 \otimes \sigma t_1\}$
= $L(R^{\sigma}, T^{\sigma}, H)$

 (2) According to (3.1) ,

$$
L(R, T, H) = \{r \otimes t \mid \sum r_0 \otimes t \otimes Sr_1 = \sum r \otimes t_0 \otimes t_1\}
$$

and

$$
L(T, R^{S^2}, H) = \{t \otimes r \mid \sum_{\alpha} t_0 \otimes r \otimes St_1 = \sum_{\alpha} t \otimes r_0 \otimes S^2 r_1\}.
$$

Since S is bijective, we have $L(T, R^{S^2}, H) = \{t \otimes r \mid \sum t_0 \otimes r \otimes t_1 = \sum t \otimes r_0 \otimes S r_1\}.$ Thus, the map

$$
L(R, T, H) \longrightarrow L(T, R^{S^2}, H)^{op}, \qquad r \otimes t \mapsto t \otimes r
$$

constitutes an algebra isomorphism.

Recall that H is a right H^e-comodule algebra via (2.15) . For any $t \otimes r \otimes h \in$ $L(T \otimes R^{op}, H, H^e) \subseteq T \otimes R^{op} \otimes H$, then

(3.5)
$$
\sum t_0 \otimes r_0 \otimes h_2 \otimes t_1 h_3 \otimes (Sh_1) r_1 = t \otimes r \otimes h \otimes 1 \otimes 1.
$$

Hence

$$
t \otimes r \otimes h = \sum t_0 \otimes r_0 \otimes \varepsilon(h_2) \otimes (St_1)t_2h_3 \otimes \varepsilon((Sh_1)r_1)
$$

$$
\xrightarrow{\underline{\underline{(3.5)}}} \sum t_0 \otimes r \otimes \varepsilon(h) \otimes St_1 \otimes \varepsilon(1)
$$

$$
= \varepsilon(h) \sum t_0 \otimes r \otimes St_1.
$$

and

$$
t \otimes r \otimes h = \sum t_0 \otimes r_0 \otimes \varepsilon(h_2) \otimes \varepsilon(t_2 h_3) \otimes (S^{-2} r_1) S^{-1}((Sh_1) r_2)
$$

$$
\xrightarrow{\underline{(3.5)}} \sum t \otimes r_0 \otimes \varepsilon(h) \otimes \varepsilon(1) \otimes S^{-2} r_1
$$

$$
= \varepsilon(h) \sum t \otimes r_0 \otimes S^{-2} r_1.
$$

It follows from [\(3.1\)](#page-20-1) that $\varepsilon(h)t \otimes r \in L(T, R^{S^{-2}}, H) = L(T^{S^2}, R, H)$. This gives an algebra isomorphism $L(T \otimes R^{op}, H, H^e) \to L(T^{S^2}, R, H)$ defined by $t \otimes r \otimes h \mapsto$ $\varepsilon(h)t \otimes r$, and the inverse map is given by $t \otimes r \mapsto \sum t_0 \otimes r \otimes St_1$.

$$
\qquad \qquad \Box
$$

Similarly, there is an algebra isomorphism $L(H, R \otimes T^{op}, H^e) \to L(T, R, H)$ defined by $h \otimes r \otimes t \mapsto \varepsilon(h)t \otimes r$, and the inverse map is given by $t \otimes r \mapsto \sum t_1 \otimes r \otimes t_0$. For any $h \otimes r \otimes t \in L(H, R \otimes T^{op}, H^e) \subseteq H \otimes R \otimes T^{op}$, then

$$
\sum h_2 \otimes r_0 \otimes t_0 \otimes h_3r_1 \otimes t_1Sh_1 = h \otimes r \otimes t \otimes 1 \otimes 1.
$$

Hence

$$
r \otimes t \otimes h = \sum \varepsilon(h_2) \otimes r_0 \otimes t_0 \otimes h_3 r_2 S^{-1} r_1 \otimes \varepsilon(t_1 Sh_1)
$$

=
$$
\sum \varepsilon(h) \otimes r_0 \otimes t \otimes S^{-1} r_1 \otimes \varepsilon(1)
$$

=
$$
\varepsilon(h) \sum r_0 \otimes t \otimes S^{-1} r_1.
$$

and

$$
r \otimes t \otimes h = \sum \varepsilon(h_2) \otimes r_0 \otimes t_0 \otimes \varepsilon(h_3 r_1) \otimes S^{-1}(t_2 Sh_1)t_1
$$

=
$$
\sum \varepsilon(h) \otimes r \otimes t_0 \otimes \varepsilon(1) \otimes t_1
$$

=
$$
\varepsilon(h) \sum r \otimes t_0 \otimes t_1.
$$

It follows from (3.1) that $\varepsilon(h)t \otimes r \in L(T, R, H)$.

Definition 3.6. Define $\Lambda_i := L(B^{S^{2i}}, B, H)$ for $i \in \mathbb{Z}$ and let $\Lambda := \Lambda_0$.

For any $i \in \mathbb{Z}$, by Lemma [3.5,](#page-24-1)

$$
\Lambda_i = L(B^{S^{2i}}, B, H) \cong L(B, B^{S^{2(i+1)}}, H)^{op} \cong L(B^{S^{-2(i+1)}}, B, H)^{op} = (\Lambda_{-i-1})^{op}.
$$

It is evident that $A^e \subseteq \Lambda_i$ implying A^e is a subalgebra of Λ_i . Note that A can be seen as a left Λ_i -module via $(x \otimes y) \triangleright a = xay$ for any $a \in A$ and $x \otimes y \in \Lambda_i$.

According to the explanation preceding Definition [3.6,](#page-25-0) we have the following result.

Lemma 3.7. (1) There is an algebra isomorphism $\Lambda_i \stackrel{\cong}{\to} L(B^{S^{2(i-1)}} \otimes B^{op}, H, H^e)$ defined by $x \otimes y \mapsto \sum x_0 \otimes y \otimes Sx_1$.

(2) There is an algebra isomorphism $\Lambda_i \stackrel{\cong}{\to} L(H, B \otimes (B^{S^{2i}})^{op}, H^e)$ defined by $x \otimes y \mapsto \sum x_1 \otimes y \otimes x_0.$

The subsequent theorems are consequences of Theorem [3.1](#page-20-2) and [3.2.](#page-21-2)

Theorem 3.8. Let H be a Hopf algebra and B a right faithfully flat H -Galois extension of $A := B^{coH}$. Then we have quasi-inverse category equivalences

$$
{}_{B^e}{\mathcal M}^{H^e}_H\longrightarrow {}_{\Lambda_1}{\mathcal M},\;\;N\mapsto N^{coH^e},\qquad {}_{\Lambda_1}{\mathcal M}\longrightarrow {}_{B^e}{\mathcal M}^{H^e}_H,\;\;M\mapsto B^e\otimes_{A^e}M,
$$

where

(1) the Λ_1 -module structure of N^{coH^e} is defined by

$$
\lambda \triangleright n = \sum (x_0 \otimes y)n(Sx_1), \qquad \forall \lambda = x \otimes y \in \Lambda_1 \subseteq B^e;
$$

(2) for $M \in \Lambda_1\mathcal{M}$ we use \triangleright to denote the module structure, M is a left A^e -module via $(a \otimes a')m = (a \otimes a' \otimes 1) \triangleright m$, and the left B^e -module and H^e -comodule

structures on $B^e \otimes_{A^e} M$ are those induced by the left tensorand, and the right H-module stucture is given by

$$
(3.6) \ \ (x \otimes y \otimes_{A^e} m)h = \sum x \kappa^1(S^{-1}h_2) \otimes \kappa^2(Sh_1)y \otimes_{A^e} (\kappa^2(S^{-1}h_2) \otimes \kappa^1(Sh_1)) \rhd m.
$$

Proof. Recall that $A^e \subseteq B^e$ is an H^e -Galois extension. The conclusion is then drawn from Theorem [3.2](#page-21-2) and Lemma [3.7.](#page-25-1) For the reader's benefit, we will examine the Hmodule structure on $B^e \otimes_{A^e} M$. The isomorphism $L(B^e, H, H^e) \to \Lambda_1$, $x \otimes y \otimes h \mapsto$ $\varepsilon(h)x \otimes y$ as stated in Lemma [3.7,](#page-25-1) indicates that M is also endowed with a left $L(B^e, H, H^e)$ -module structure. We denote this module structure by \triangleright as well. Thus, our notation leads to the expression $(x \otimes y \otimes h) \triangleright m = \varepsilon(h)(x \otimes y) \triangleright m$. Given $x, y \in B$, $h \in H$ and $m \in M$, we compute

$$
(x \otimes y \otimes_{A^e} m)h
$$

\n
$$
\stackrel{(3.2)}{\longrightarrow} \sum (x \otimes y) \kappa_{Be}^1 (S_{He}^{-1}(h_3 \otimes Sh_1)) \otimes_{A^e} (\kappa_{Be}^2 (S_{He}^{-1}(h_3 \otimes Sh_1)) \otimes h_2) \rhd m
$$

\n
$$
= \sum (x \otimes y) \kappa_{Be}^1 (S^{-1}h_3 \otimes S^2h_1) \otimes_{A^e} (\kappa_{Be}^2 (S^{-1}h_3 \otimes S^2h_1) \otimes h_2) \rhd m
$$

\n
$$
\stackrel{(2.14)}{\longrightarrow} \sum (x \otimes y) (\kappa^1 (S^{-1}h_3) \otimes \kappa^2 (Sh_1)) \otimes_{A^e} (\kappa^2 (S^{-1}h_3) \otimes \kappa^1 (Sh_1) \otimes h_2) \rhd m
$$

\n
$$
= \sum x \kappa^1 (S^{-1}h_2) \otimes \kappa^2 (Sh_1) y \otimes_{A^e} (\kappa^2 (S^{-1}h_2) \otimes \kappa^1 (Sh_1)) \rhd m.
$$

Theorem 3.9. Let H be a Hopf algebra and B a right faithfully flat H -Galois extension of $A := B^{coH}$. Then we have quasi-inverse category equivalences

$$
{}_H{\mathcal M}^{H^e}_{B^e}\longrightarrow {}_\Lambda{\mathcal M},\ N\mapsto N^{coH^e},\qquad {}_\Lambda{\mathcal M}\longrightarrow {}_H{\mathcal M}^{H^e}_{B^e},\ M\mapsto M\otimes_{A^e}B^e,
$$

where

(1) the Λ -module structure of N^{coH^e} is defined by

$$
\lambda \rhd n = \sum x_1 n(y \otimes x_0), \qquad \forall \ \lambda = x \otimes y \in \Lambda \subseteq B^e;
$$

(2) for $M \in \Lambda \mathcal{M}$ we use \triangleright to denote the module structure, M is a right A^e module via $m(a \otimes a') = (a \otimes a) \triangleright m$, and the right B^e -module and H^e comodule structures on $M \otimes_{A^e} B^e$ are those induced by the right tensorand, and the left H-module stucture is given by

$$
h(m\otimes_{A^e}x\otimes y)=\sum(\kappa^2(h_1)\otimes\kappa^1(h_2))\rhd m\otimes_{A^e}\kappa^2(h_2)x\otimes y\kappa^1(h_1).
$$

Proof. The proof parallels that of Theorem [3.8.](#page-25-2) \Box

The subsequent lemma is a direct consequence of Lemma [3.4,](#page-21-4) Lemma [3.7](#page-25-1) Theorem [3.8](#page-25-2) and theorem [3.9.](#page-26-0)

Lemma 3.10. Let H be a Hopf algebra and B a right faithfully flat H-Galois extension of $A := B^{coH}$.

(1) For any left Λ -module M, $\text{Hom}_{A^e}(M, A^e)$ is a left Λ_1 -module, defined by

(3.7)
$$
\begin{aligned} ((x \otimes y) \triangleright f)(m) \\ &= \sum (x_0 \otimes y) f(\kappa^2(Sx_2) \otimes (\kappa^1(Sx_1)) \triangleright m)(\kappa^2(Sx_1) \otimes \kappa^1(Sx_2)), \end{aligned}
$$

for any $x \otimes y \in \Lambda$, $f \in \text{Hom}_{A^e}(M, A^e)$ and $m \in M$.

(2) For any $M \in M$ and $N \in {}_{B^e}{\mathcal M}^{H^e}_{B^e}$, Hom_{$A^e(M, N)$} is a B^e -H-bimodule, where the right H -module is given by

(3.8)
$$
(fh)(m) = \sum f\left((\kappa^2(h_1) \otimes \kappa^1(h_2)) \rhd m\right) (\kappa^2(h_2) \otimes \kappa^1(h_1))
$$

for any $f \in \text{Hom}_{A^e}(M, N)$, $h \in H$ and $m \in M$.

- (3) The canonical map
- $\Phi: N \otimes_{A^e} \text{Hom}_{A^e}(M, A^e) \longrightarrow \text{Hom}_{A^e}(M, N), \qquad n \otimes_{A^e} f \mapsto (m \mapsto nf(m))$ is a B^e -H-bimodule morphism.

For any $N \in B^e \mathcal M_{B^e}^{H^e}$ and $f \in \text{Hom}_{A^e}(A,N)$, we have

$$
(fh)(1_A) = \sum f((\kappa^2(h_1) \otimes \kappa^1(h_2)) \triangleright 1_A)(\kappa^2(h_2) \otimes \kappa^1(h_1))
$$

= $\sum \kappa^1(h_1) f(\kappa^2(h_1) 1_A \kappa^1(h_2)) \kappa^2(h_2)$
(since $\sum \kappa^1(h_1) \otimes_A \kappa^2(h_1) \kappa^1(h_2) \otimes_A \kappa^2(h_2) \in B \otimes_A A \otimes_A B)$
= $\sum \kappa^1(h_1) f(1_A) \kappa^2(h_1) \kappa^1(h_2) \kappa^2(h_2)$
 $\xrightarrow{(2.6)} \sum \kappa^1(h) f(1_A) \kappa^2(h)$
 $\xrightarrow{(2.8)} f(1_A) \leftarrow h$

Consequently, the right H-module structure on $\text{Hom}_{A^e}(A, N) \cong N^A$, as defined in (3.8) , aligns with that in (2.8) .

Proposition 3.11. Let H be a Hopf algebra and B a right faithfully flat H-Galois extension of $A := B^{coH}$.

- (1) For any $M \in \Lambda \mathcal{M}$ and $i > 0$, $\text{Ext}_{A^e}^i(M, A^e)$ is a left Λ_1 -module.
- (2) If M is perfect as an A^e-complex, then $\operatorname{Ext}_{A^e}^i(M, B^e) \cong B^e \otimes_{A^e} \operatorname{Ext}_{A^e}^i(M, A^e)$ as B^e -H-bimodules, where the right H-module on $B^e \otimes_{A^e} \text{Ext}_{A^e}^i(M, A^e)$ is given by (3.6) .

Proof. (1) Taking a projective resolution P_{\bullet} of the Λ -module M, by Lamma [3.10](#page-26-2) (1), Hom_{Ae}(P_{\bullet} , A^e) is a complex of Λ_1 -modules. Since Λ is a projective A^e -module, $\text{Ext}_{A^e}^i(M, A^e) \cong H^i(\text{Hom}_{A^e}(P_{\bullet}, A^e)).$ Thus, $\text{Ext}_{A^e}^i(M, A^e)$ is a left Λ_1 -module.

(2) Note that the canonical chain map $\Phi_{\bullet}: B^e \otimes_{A^e} \text{Hom}_{A^e}(P_{\bullet}, A^e) \to \text{Hom}_{A^e}(P_{\bullet}, B^e)$ is both B^e -linear and H-linear, as per [3.10](#page-26-2) (3). Given that M is perfect, P_{\bullet} can be selected as a bounded complex of finitely generated projective A^e -modules. Hence, Φ_{\bullet} is a quasi-isomoprhism. Taking homologies yields the desired result. \Box

4. Homological determinats and Nakayama automotphisms for crossed products

Let H be a Hopf algebra and A be an algebra. Suppose that H measures A, that is, there is a k-linear map $H \otimes A \to A$, given by $h \otimes a \mapsto h \cdot a$, such that

$$
h \cdot 1 = \varepsilon(h)1
$$
, and $h \cdot (ab) = \sum (h_1 \cdot a)(h_2 \cdot b)$,

for all $h \in H$, $a, b \in A$. Let σ be a convolution invertible map in Hom $(H \otimes H, A)$, meaning there exists $\sigma^{-1} \in \text{Hom}(H \otimes H, A)$ satisfying $\sum \sigma(h_1, k_1) \sigma^{-1}(h_2, k_2)$ = $\varepsilon(hk)$ for all $h, k \in H$. Let $A \#_{\sigma} H$ be the (in general nonassociative) algebra whose underlying vector space is $A \otimes H$ with multiplication

$$
(a \# h)(b \# k) = \sum a(h_1 \cdot b)\sigma(h_2, k_1) \# h_3 k_2
$$

for all $a, b \in A$, $h, k \in H$. The element $a \otimes h$ will usually be written $a \# h$. The algebra $A\#_{\sigma}H$ is called *crossed product* of A with H if it is associative with $1\#1$ as identity element. Recall from [\[4\]](#page-40-12) and [\[10\]](#page-40-13) that $A\#_{\sigma}H$ is a crossed product if and only if the following two conditions are satisfied:

(1) A is a twisted H-module. That is, $1 \cdot a = a$, for all $a \in A$, and

(4.1)
$$
h \cdot (k \cdot a) = \sum \sigma(h_1, k_1)(h_2 k_2 \cdot a) \sigma^{-1}(h_3, k_3)
$$

for all $h, k \in H$, $a \in A$.

(2) σ is normal cocycle. That is, $\sigma(h, 1) = \sigma(1, h) = \varepsilon(h)1$, for all $h \in H$, and

(4.2)
$$
\sum (h_1 \cdot \sigma(k_1, l_1)) \sigma(h_2, k_2 l_2) = \sum \sigma(h_1, k_1) \sigma(h_2 k_2, l)
$$
for all $h \ge l \in H$

for all $h, k, l \in H$.

Deriving from (4.2) , we have the following identities for all $h, k, l \in H$:

(4.3)
$$
\sum \sigma^{-1}(h_1, k_1) (h_2 \cdot \sigma(k_2, l)) = \sum \sigma(h_1 k_1, l_1) \sigma^{-1}(h_2, k_2 l_2),
$$

(4.4)
$$
\sum \sigma^{-1}(h_1k_1, l)\sigma^{-1}(h_2, k_2) = \sum \sigma^{-1}(h_1, k_1l_1)(h_2 \cdot \sigma^{-1}(k_2, l_2)),
$$

Clearly, each crossed product $A#_{\sigma}H$ is a right H-comodule algebra such that $A \subseteq A \#_{\sigma} H$ is a faithfully flat H-Galois extension.

Let B be a right H-comodule algebra. The extension $A(:=B^{coH})\subset B$ is termed H-cleft if there exists a convolution invertible right H-comodule map $\gamma : H \to B$.

On one hand, consider the crossed product $A#_{\sigma}H$. Define the map $\gamma : H \rightarrow$ $A#_{\sigma}H$ by $\gamma(h) = 1#h$, which is convolution invertible with the inverse given by

(4.5)
$$
\gamma^{-1}(h) = \sum \sigma^{-1}(Sh_2, h_3) \# Sh_1,
$$

Thus, $A \subseteq A \#_{\sigma} H$ forms an H-cleft extension, as shown in Proposition 7.2.7 of [\[24\]](#page-41-11).

On the other hand, every H-cleft extension $A \subseteq B$ is a crossed product of A with H, where the action is defined by

(4.6)
$$
h \cdot a = \sum \gamma(h_1) a \gamma^{-1}(h_2), \ \forall a \in A, h \in H,
$$

and the convolution invertible map $\sigma : H \otimes H \to A$ is given by

(4.7)
$$
\sigma(h,k) = \sum \gamma(h_1) \gamma(k_1) \gamma^{-1}(h_2 k_2), \ \forall \, h, k \in H,
$$

This result is established in Proposition 7.2.3 [\[24\]](#page-41-11).

Given a cleft H-extension $A \subseteq B$ and a convolution invertible right H-comodule map $\gamma : H \to B$, for any $h \in H$, the following equality holds:

$$
\beta(\sum \gamma^{-1}(h_1) \otimes \gamma(h_2)) = \beta(\sum \sigma^{-1}(Sh_2, h_3) \# Sh_1 \otimes 1 \# h_4)
$$

= $\sum \sigma^{-1}(Sh_3, h_4) \sigma(Sh_2, h_5) \#(Sh_1) h_6 \otimes h_7$

$$
=1\#h.
$$

Consequently, the translation map for the extension is given by

(4.8)
$$
\kappa(h) = \sum \kappa^1(h) \otimes \kappa^2(h) = \sum \gamma^{-1}(h_1) \otimes \gamma(h_2).
$$

Lemma 4.1. Let B be a crossed product $A\#_{\sigma}H$. Then the algebra Λ_i , as defined in Definition [3.6,](#page-25-0) is identified with $\{\sum x \# h_1 \otimes y \# S^{2i+1} h_2 \mid x, y \in A, h \in H\} \subseteq B^e$.

Proof. The inclusion $\{\sum x \# h_1 \otimes y \# S^{2i+1} h_2 \mid x, y \in A, h \in H\} \subseteq \Lambda_i$ is evident. For any $x\#h \otimes y\#k \in \Lambda_i$, the definition of Λ_i yields

$$
\sum x \# h_1 \otimes y \# k_1 \otimes (S^{2i}h_2)k_2 = x \# h \otimes y \# k \otimes 1.
$$

Thus, we have

$$
x \# h \otimes y \# k \otimes S^{2i+1} h_2 = \sum x \# h_1 \otimes y \# k_1 \otimes (S^{2i+1} h_2)(S^{2i} h_3) k_2
$$

= $\sum x \# h \otimes y \# k_1 \otimes k_2.$

Thus, $x\#h \otimes y\#k = x\#h \otimes y\# \varepsilon(k_1)k_2 = x\#h \otimes y\# \varepsilon(k)S^{2i+1}h_2$. This implies $\Lambda_i \subseteq \{ \sum x \# h_1 \otimes y \# S^{2i+1} h_2 \mid x, y \in A, h \in H \}.$ Therefore, the result follows.

Smash products are a specialized form of cleft extensions. Consider a right Hmodule algebra A with action denoted by "·" and a cocycle $\sigma(h, k) = \varepsilon(h) \varepsilon(k)$ for all $h, k \in H$. In this setting, the crossed product $A \#_{\sigma} H$ constitutes a smash product, denoted $A#H$. Let Δ_i be the algebra whose underlying vector space is $A^e \otimes H$, and whose multiplication is defined by

$$
(x\otimes y\otimes h)(x'\otimes y'\otimes h')=\sum x(h_1\cdot x')\otimes (S^{2i}h_3\cdot y')y\otimes h_2h'.
$$

The algebra Δ_i , introduced in [\[16\]](#page-40-4) and [\[19\]](#page-41-6), has been utilized to examine the Calabi-Yau property of smash products, as detailed in [\[20\]](#page-41-3), [\[27\]](#page-41-5), [\[19\]](#page-41-6).

Subsequently, we will show that Λ_i is essentially Δ_i in the context of smash products.

Lemma 4.2. The map $F: \Delta_i \to \Lambda_i$ defined by

$$
x \otimes y \otimes h \mapsto \sum x \# h_1 \otimes S^{2i+1} h_3 \cdot y \# S^{2i+1} h_2
$$

is an algebra isomorphism.

Proof. The map F is evidently bijective, with the inverse map given by

$$
\sum x \# h_1 \otimes y \# S^{2i+1} h_2 \mapsto \sum x \otimes (S^{2i} h_2 \cdot y) \otimes h_1.
$$

To show that F is an algebra isomorphism, we verify that it preserves the algebra structure. We have

$$
F((x \otimes y \otimes h)(x' \otimes y' \otimes h'))
$$

= $\varphi(\sum x(h_1 \cdot x') \otimes (S^{2i}h_3 \cdot y')y \otimes h_2h')$
= $\sum x(h_1 \cdot x') \# h_2h'_1 \otimes S^{2i+1}(h_4h'_3) \cdot ((S^{2i}h_5 \cdot y')y) \# S^{2i+1}(h_3h'_2)$
= $\sum x(h_1 \cdot x') \# h_2h'_1 \otimes (S^{2i+1}h'_4 \cdot y') (S^{2i+1}(h_4h'_3) \cdot y) \# S^{2i+1}(h_3h'_2)$

$$
= \sum x(h_1 \cdot x') \# h_2 h'_1 \otimes (S^{2i+1}h'_4 \cdot y')((S^{2i+1}h'_3)(S^{2i+1}h_4) \cdot y) \# (S^{2i+1}h'_2)(S^{2i+1}h_3)
$$

=
$$
(\sum x \# h_1 \otimes S^{2i+1}h_3 \cdot y \# S^{2i+1}h_2)(\sum x' \# h'_1 \otimes S^{2i+1}h'_3 \cdot y' \# S^{2i+1}h'_2)
$$

=
$$
F(x \otimes y \otimes h)F(x' \otimes y' \otimes h'),
$$

which demonstrates that F is an algebra isomorphism. \Box

Subsequently, we consider B as a crossed product $A\#_{\sigma}H$ with the comodule map $\gamma: H \to A$ defined by $\gamma(h) = 1 \# h$.

Lemma 4.3. For any $h, k \in H$,

(4.9)
$$
\gamma^{-1}(k)\gamma^{-1}(h) = \gamma^{-1}(h_1k_1)(\sigma^{-1}(h_2,k_2)\#1).
$$

Proof. We proceed with the calculation as follows:

$$
\gamma^{-1}(k)\gamma^{-1}(h)
$$
\n
$$
\frac{(4.5)}{2}\sum \sigma^{-1}(Sk_2, k_3) \# Sk_1) (\sigma^{-1}(Sh_2, h_3) \# Sh_1)
$$
\n
$$
= \sum \sigma^{-1}(Sk_4, k_5) (Sk_3 \cdot \sigma^{-1}(Sh_3, h_4)) \sigma(Sk_2, Sh_2) \# (Sk_1)(Sh_1)
$$
\n
$$
= \sum \sigma^{-1}(Sk_5, k_6) \sigma^{-1}(Sk_4, (Sh_4)h_5) (Sk_3 \cdot \sigma^{-1}(Sh_3, h_6)) \sigma(Sk_2, Sh_2) \# S(h_1k_1)
$$
\n
$$
\frac{(4.4)}{2}\sum \sigma^{-1}(Sk_5, k_6) \sigma^{-1}((Sk_4)(Sh_4), h_5) \sigma^{-1}(Sk_3, Sh_3) \sigma(Sk_2, Sh_2) \# S(h_1k_1)
$$
\n
$$
= \sum \sigma^{-1}(S(h_3k_3)h_4, k_4) \sigma^{-1}(S(h_2k_2), h_5) \# S(h_1k_1)
$$
\n
$$
\frac{(4.4)}{2}\sum \sigma^{-1}(S(h_3k_3), h_4k_4) (S(h_2k_2) \cdot \sigma^{-1}(h_5, k_5)) \# S(h_1k_1)
$$
\n
$$
= \sum \sigma^{-1}(S(h_2k_2), h_3k_3) \# S(h_1k_1) (\sigma^{-1}(h_4, k_4) \# 1)
$$
\n
$$
= \gamma^{-1}(h_1k_1) (\sigma^{-1}(h_2, k_2) \# 1)
$$

This completes the proof. \Box

In general, H cannot inherently be a subalgebra Λ_i unless σ is trivial. However, there exists an embedding map $\pi_i: H \to \Lambda_i$ defined by $h \mapsto \sum \gamma(h_1) \otimes \gamma^{-1}(S^{2i}h_2)$.

The following properties hold for this map.

Lemma 4.4. For any $h, k \in H$, in Λ_i ,

(1) $\pi_i(1) = 1 \otimes 1$, (2) $\pi_i(h)(a\#1 \otimes a'\#1) = \sum (h_1 \cdot a\#1 \otimes S^{2i}h_3 \cdot a'\#1)\pi_i(h_2),$ (3) $\pi_i(h)\pi_i(k) = \sum (\sigma(h_1, k_1) \# 1 \otimes \sigma^{-1}(S^{2i}h_3, S^{2i}k_3) \# 1)\pi_i(h_2k_2).$

Proof. (1) This follows from $\gamma(1_H) = 1_B$. (2) We compute

$$
\pi_i(h)(a \otimes a') = (\sum \gamma(h_1) \otimes \gamma^{-1}(S^{2i}h_2))(a \otimes a')
$$

= $\sum \gamma(h_1)a \otimes a'\gamma^{-1}(S^{2i}h_2)$
= $\sum \gamma(h_1)a\gamma^{-1}(h_2)\gamma(h_3) \otimes \gamma^{-1}(S^{2i}h_4)\gamma(S^{2i}h_5)a'\gamma^{-1}(S^{2i}h_6)$
= $\sum (\gamma(h_1)a\gamma^{-1}(h_2) \otimes \gamma(S^{2i}h_5)a'\gamma^{-1}(S^{2i}h_6))(\gamma(h_3) \otimes \gamma^{-1}(S^{2i}h_4))$

$$
= \sum (h_1 \cdot a \otimes S^{2i} h_3 \cdot a') \pi_i(h_2).
$$

(3) Using $\gamma(h)\gamma(k) = (1 \# h)(1 \# k) = \sum \sigma(h_1, k_1)\gamma(h_2k_2)$, and (4.9), we get

$$
\pi_i(h)\pi_i(k) = \left(\sum \gamma(h_1) \otimes \gamma^{-1}(S^{2i} h_2)\right) \left(\sum \gamma(k_1) \otimes \gamma^{-1}(S^{2i} k_2)\right)
$$

$$
= \sum \gamma(h_1)\gamma(k_1) \otimes \gamma^{-1}(S^{2i} k_2)\gamma^{-1}(S^{2i} h_2)
$$

$$
\xrightarrow{\underline{(4.9)}} \sum (\sigma(h_1, k_1) \# 1) \gamma(h_2k_2) \otimes \gamma^{-1}(S^{2i}(h_3k_3)) \left(\sigma^{-1}(S^{2i} h_4, S^{2i} k_4) \# 1\right)
$$

$$
= \sum (\sigma(h_1, k_1) \# 1 \otimes \sigma^{-1}(S^{2i} h_3, S^{2i} k_3) \# 1) \pi_i(h_2k_2).
$$

The proof is completed. \Box

For any $M \in \Lambda_1 \mathcal{M}, h \in H$ and $m \in M$, we define

(4.10)
$$
h \cdot m := \pi_1(h)m = \left(\sum \gamma(h_1) \otimes \gamma^{-1}(S^2h_2)\right) \rhd m.
$$

Observe that M is generally not an H-module under the action defined above. Building upon Lemma [4.4,](#page-30-1) we establish the following:

Lemma 4.5. For all $h, k \in H$, $a, a' \in A$, and $m \in M$,

$$
(4.11) \t\t\t 1_H \cdot m = m,
$$

(4.12)
$$
h \cdot (ama') = \sum (h_1 \cdot a)(h_2 \cdot m)(S^2 h_2 \cdot a'),
$$

(4.13)
$$
h \cdot (k \cdot m) = \sum \sigma(h_1, k_1) (h_2 k_2 \cdot m) \sigma^{-1} (S^2 h_3, S^2 k_3).
$$

4.1. Homological determinant. Inspired by the Nakayama automorphism's expression for smash products of skew Calabi-Yau algebras via homological determinants, we introduce the concept of homological determinants for crossed products.

Let A be a skew Calabi-Yau algebra of dimension d with a Nakayama automorphism μ_A . Hence there exists a generator e of the A^e-module $\text{Ext}_{A^e}^{\phi}(A, A^e) (\cong A_{\mu_A})$ such that $ea = \mu_A(a)e$ for any $a \in A$. The generator e will be referred to a μ_A -twisted *volume* on A. As established in Proposition [3.11,](#page-27-2) $\text{Ext}_{A}^{d}(A, A^{e})$ is a Λ_{1} -module. Since e also generates $\text{Ext}_{A^e}^d(A, A^e)$ as a free left A-module of rank one, there exists a klinear map $\varphi : H \to A$ defined by

(4.14)
$$
h \cdot e = \pi_1(h)e = \varphi(h)e.
$$

Then we can deduce the subsequent outcome from Lemma [4.5.](#page-31-0)

Lemma 4.6. For any $h, k \in H$, $a \in A$, then

$$
\varphi(1_H) = 1_A,
$$

(4.16)
$$
\sum (h_1 \cdot \mu_A(a)) \varphi(h_2) = \sum \varphi(h_1) \mu_A(S^2 h_2 \cdot a),
$$

(4.17)
$$
\sum (h_1 \cdot \varphi(k)) \varphi(h_2) = \sum \sigma(h_1, k_1) \varphi(h_2 k_2) \mu_A(\sigma^{-1}(S^2 h_3, S^2 k_3)).
$$

Proof. (1) Note that e is a generator of a free left A-module $\text{Ext}_{A}^{d}(A, A^{e})$. We have $\varphi(1_H) = 1_A$ because $e = 1_H \cdot e = \varphi(1_H)e$ by [\(4.11\)](#page-31-1). (2) We compute

$$
\sum (h_1 \cdot \mu_A(a)) \varphi(h_2) e \xrightarrow{\frac{(4.14)}{2}} \sum (h_1 \cdot \mu_A(a))(h_2 \cdot e)
$$

$$
\xrightarrow{\frac{(4.12)}{2}} h \cdot (\mu_A(a)e) = h \cdot (ea)
$$

$$
\xrightarrow{\frac{(4.14)}{2}} \sum (\mu_1 \cdot e)(S^2 h_2 \cdot a)
$$

$$
= \sum \varphi(h_1) \mu_A(S^2 h_2 \cdot a)e
$$

implies that [\(4.16\)](#page-31-4) holds.

(3) We calculate

$$
\sum (h_1 \cdot \varphi(k)) \varphi(h_2) e \stackrel{(4.14)}{=} \sum (h_1 \cdot \varphi(k)) (h_2 \cdot e)
$$

$$
\stackrel{(4.12)}{=} h \cdot (\varphi(k)e) \stackrel{(4.14)}{=} h \cdot (k \cdot e)
$$

$$
\stackrel{(4.13)}{=} \sum \sigma(h_1, k_1) ((h_2 k_2) \cdot e) \sigma^{-1} (S^2 h_3, S^2 k_3)
$$

$$
\stackrel{(4.14)}{=} \sum \sigma(h_1, k_1) \varphi(h_2 k_2) \mu_A (\sigma^{-1} (S^2 h_3, S^2 k_3)) e
$$

showing that (4.17) is valid. This completes the proof. \Box

We now demonstrate that φ is convolution invertible, with its inverse, referred to as the homological determinant of the H-action on A.

Lemma 4.7. The map φ defined by [\(4.14\)](#page-31-2) is convolution invertible with the convolution inverse ψ which is defined by

(4.18)
$$
h \mapsto \sum \sigma^{-1}(h_5, S^{-1}h_4)h_6 \cdot \left(\varphi(S^{-1}h_3)\mu_A\big(\sigma(Sh_2, S^2h_1)\big)\right).
$$

Proof. For the convolution product $(\psi * \varphi)(h)$, we have

$$
= \sum \psi(h_1)\varphi(h_2)
$$

\n
$$
= \sum \sigma^{-1}(h_5, S^{-1}h_4)h_6 \cdot \left(\varphi(S^{-1}h_3)\mu_A(\sigma(Sh_2, S^2h_1))\right)\varphi(h_7)
$$

\n
$$
= \sum \sigma^{-1}(h_5, S^{-1}h_4)\left(h_6 \cdot \varphi(S^{-1}h_3)\right)\left(h_7 \cdot \mu_A(\sigma(Sh_2, S^2h_1))\right)\varphi(h_8)
$$

\n
$$
\xrightarrow{(4.16)} \sum \sigma^{-1}(h_5, S^{-1}h_4)\left(h_6 \cdot \varphi(S^{-1}h_3)\right)\varphi(h_7)\mu_A(S^2h_8 \cdot \sigma(Sh_2, S^2h_1))
$$

\n
$$
\xrightarrow{(4.17)} \sum \sigma^{-1}(h_7, S^{-1}h_6)\sigma(h_8, S^{-1}h_5)\varphi\left(h_9(S^{-1}h_4)\right)\mu_A(\sigma^{-1}(S^2h_{10}, Sh_3))
$$

\n
$$
\mu_A(S^2h_{11} \cdot \sigma(Sh_2, S^2h_1))
$$

\n
$$
= \sum \varphi(1_H)\mu_A(\sigma^{-1}(S^2h_4, Sh_3))\mu_A(S^2h_5 \cdot \sigma(Sh_2, S^2h_1))
$$

\n
$$
\xrightarrow{(4.15)} \sum \mu_A(\sigma^{-1}(S^2h_4, Sh_3)(S^2h_5 \cdot \sigma(Sh_2, S^2h_1))
$$

$$
\frac{(4.3)}{} \sum \mu_A \Big(\sigma \big((S^2 h_5)(Sh_4), S^2 h_1 \big) \sigma^{-1} \big(S^2 h_6, (Sh_3)(S^2 h_2) \big) \Big)
$$
\n
$$
= \sum \mu_A \big(\sigma (1, S^2 h_1) \sigma^{-1} (S^2 h_2, 1) \big)
$$
\n
$$
= \varepsilon (h).
$$

Analogously, for $(\varphi * \psi)(h)$

$$
= \sum \varphi(h_1)\psi(h_2)
$$

\n
$$
= \sum \varphi(h_1)\sigma^{-1}(h_6, S^{-1}h_5)h_7 \cdot (\varphi(S^{-1}h_4)\mu_A(\sigma(Sh_3, S^2h_2)))
$$

\n
$$
= \sum \sigma^{-1}(h_9, S^{-1}h_8)\sigma(h_{10}, S^{-1}h_7)(h_{11}S^{-1}h_6 \cdot \varphi(h_1))\sigma^{-1}(h_{12}, S^{-1}h_5)
$$

\n
$$
h_{13} \cdot (\varphi(S^{-1}h_4)\mu_A(\sigma(Sh_3, S^2h_2)))
$$

\n
$$
\xrightarrow{(4.1)}
$$

$$
\sum \sigma^{-1}(h_7, S^{-1}h_6)h_8 \cdot (S^{-1}h_5 \cdot \varphi(h_1))h_9 \cdot (\varphi(S^{-1}h_4)\mu_A(\sigma(Sh_3, S^2h_2)))
$$

\n
$$
= \sum \sigma^{-1}(h_7, S^{-1}h_6)h_8 \cdot ((S^{-1}h_5 \cdot \varphi(h_1))\varphi(S^{-1}h_4)\mu_A(\sigma(Sh_3, S^2h_2)))
$$

\n
$$
\xrightarrow{(4.17)}
$$

$$
\sum \sigma^{-1}(h_{10}, S^{-1}h_9)
$$

\n
$$
h_{11} \cdot ((\sigma(S^{-1}h_8, h_1)\varphi((S^{-1}h_7)h_2)\mu_A(\sigma^{-1}(Sh_6, S^2h_3))\mu_A(\sigma(Sh_5, S^2h_4)))
$$

\n
$$
= \sum \sigma^{-1}(h_4, S^{-1}h_3)h_5 \cdot (\sigma(S^{-1}h_2, h_1)\varphi(1_H))
$$

\n
$$
\xrightarrow{(4.15)}
$$

$$
\sum \sigma^{-1}(h_4, S^{-1}h_3)h_5 \cdot \sigma(S^{-1}h_2, h_1)
$$

\n
$$
= \sum \sigma(1, h_1)\sigma^{-1}(h_2, 1)
$$

\n
$$
= \varepsilon(h).
$$

Thus, ψ is the convolution inverse of φ .

Definition 4.8. The k-linear map $\psi : H \to A$ from Lemma [4.7](#page-32-0) is termed the homological determinant of the H-action on $A = B^{coH}$ with respect to the μ_A twisted volume e, and is denoted by Hdet.

Given a crossed product $B := A#_{\sigma}H$ of A with H, if A is a right H-module algebra under the action denoted by "",", then the homological determinant defined as in Definition [4.8,](#page-33-0) coincides with that described in $[17, 19]$ $[17, 19]$ $[17, 19]$. For instance, A is always an H-module algebra when H is cocommutative and $\sigma(H \otimes H) \subseteq Z(A)$, where $Z(A)$ is the center of A.

4.2. Nakayama automorphisms of crossed products. In this subsection, we will always assume that B is a crossed product $A#_{\sigma}H$ with the comodule map $\gamma: H \to A$ defined by $\gamma(h) = 1 \# h$.

Lemma 4.9. For any $M \in \Lambda_1 \mathcal{M}$, $B \otimes_A M \otimes H$ is a Hopf bimodule in ${}_{B^e} \mathcal{M}_H^{H^e}$, where the right H-module structure on $B \otimes_A M \otimes H$ is given by

$$
(4.19) \t\t\t(b \otimes_A m \otimes h) \leftarrow k = b \otimes_A m \otimes (Sk)h,
$$

the left B^e -module structure on $B \otimes_A M \otimes H$ is given by (4.20) $(x \otimes y)(b \otimes_A m \otimes h) = \sum x b \gamma^{-1} (S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m) \gamma (S^{-1}y_1) y_0 \otimes hy_4,$ the H^e -comodule structure on $B \otimes_A M \otimes H$ is given by

(4.21)
$$
\rho(b \otimes_A m \otimes h) = \sum b_0 \otimes_A m \otimes h_2 \otimes b_1(S^{-1}h_1) \otimes h_3
$$

for any $b, x, y \in B$, $h, k \in H$ and $m \in M$.

Proof. The proof is straightforward and thus omitted. \Box

Recall that for any $M \in \Lambda_1\mathcal{M}$, by Theorem [3.8,](#page-25-2) $B^e \otimes_{A^e} M$ is a Hopf bimodule in ${}_{B^e}\mathcal{M}_H^{H^e}$, and the right H-module structure is given by

$$
(x \otimes y \otimes_{A^e} m)h
$$

\n
$$
\xrightarrow{\text{(3.6)}} \sum x \kappa^1 (S^{-1}h_2) \otimes \kappa^2 (Sh_1) y \otimes_{A^e} (\kappa^2 (S^{-1}h_2) \otimes \kappa^1 (Sh_1)) \rhd m
$$

\n
$$
\xrightarrow{\text{(4.8)}} \sum x \gamma^{-1} (S^{-1}h_4) \otimes \gamma (Sh_1) y \otimes_{A^e} (\gamma (S^{-1}h_3) \otimes \gamma^{-1} (Sh_2)) \rhd m
$$

\n
$$
\xrightarrow{\text{(4.10)}} \sum x \gamma^{-1} (S^{-1}h_3) \otimes \gamma (Sh_1) y \otimes_{A^e} S^{-1}h_2 \cdot m
$$

for any $x, y \in B$, $h \in H$ and $m \in M$.

Lemma 4.10. For any $M \in \Lambda_1\mathcal{M}$, there is an isomorphism

$$
F: B^e \otimes_{A^e} M \stackrel{\cong}{\longrightarrow} B \otimes_A M \otimes H
$$

in ${}_{B^e}\mathcal{M}_H^{H^e}$, which is defined by

(4.23)
$$
F(x \otimes y \otimes_{A^e} m) = \sum x \gamma^{-1} (S^{-3} y_3) \otimes_A (S^{-3} y_2 \cdot m) \gamma (S^{-1} y_1) y_0 \otimes y_4.
$$

where $x, y \in B$, and $m \in M$.

Proof. Note that M is an A-A-bimodule since A^e is a subalgebra of Λ_1 . For any $b := a \# h \in B$,

$$
\sum \gamma(S^{-1}b_1)b_0 = \sum (1 \# S^{-1}h_2)(a \# h_1) = \sum (S^{-1}h_5 \cdot a)\sigma(S^{-1}h_4, h_1) \# (S^{-1}h_3)h_2
$$

=
$$
\sum (S^{-1}h_3 \cdot a)\sigma(S^{-1}h_2, h_1) \# 1 \in A.
$$

Thus, for any $x, y \in B$, and $m \in M$, the expression

$$
\sum x\gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0 \otimes y_4
$$

is unambiguous. For any $a, a \in A$, we have

$$
F(xa \otimes a'y \otimes_{A^e} m)
$$

= $\sum x a \gamma^{-1} (S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m) \gamma (S^{-1}y_1) a'y_0 \otimes y_4$
= $\sum x \gamma^{-1} (S^{-3}y_7) \gamma^{-1} (S^{-3}y_6) a \gamma^{-1} (S^{-3}y_5) \otimes_A$
 $(S^{-3}y_4 \cdot m) \gamma (S^{-1}y_3) a' \gamma^{-1} (S^{-1}y_2) \gamma (S^{-1}y_1) y_0 \otimes y_8$
 $\xrightarrow{(4.6)} \sum x \gamma^{-1} (S^{-3}y_5) (S^{-3}y_4 \cdot a) \otimes_A (S^{-3}y_3 \cdot m) (S^{-1}y_2 \cdot a') \gamma (S^{-1}y_1) y_0 \otimes y_6$
= $\sum x \gamma^{-1} (S^{-3}y_5) \otimes_A (S^{-3}y_4 \cdot a) (S^{-3}y_3 \cdot m) (S^{-1}y_2 \cdot a') \gamma (S^{-1}y_1) y_0 \otimes y_6$

$$
\frac{(4.12)}{4.12} \sum_{y} x \gamma^{-1} (S^{-3} y_3) \otimes_A (S^{-3} y_2 \cdot (ama')) \gamma (S^{-1} y_1) y_0 \otimes y_4
$$

= $F(x \otimes y \otimes_{A^e} ama').$

Therefore, F is well defined.

Next, we verify that F is a morphism in ${}_{B^e}\mathcal{M}_H^{H^e}$. To show that F is a right H-module morphism, we calculate

$$
F((x \otimes y \otimes_{A^e} m)h)
$$

\n
$$
\frac{(4.22)}{2} F(\sum x \gamma^{-1} (S^{-1}h_3) \otimes \gamma (Sh_1) y \otimes_{A^e} S^{-1}h_2 \cdot m)
$$

\n
$$
= \sum x \gamma^{-1} (S^{-1}h_7) \gamma^{-1} (S^{-3}((Sh_2) y_3)) \otimes_A S^{-3}((Sh_3) y_2) \cdot (S^{-1}h_6 \cdot m)
$$

\n
$$
\gamma (S^{-1}((Sh_4) y_1)) \gamma (Sh_5) y_0 \otimes (Sh_1) y_4
$$

\n
$$
= \sum x \gamma^{-1} (S^{-1}h_7) \gamma^{-1} ((S^{-3} y_3) (S^{-2}h_2)) \otimes_A ((S^{-3} y_2) (S^{-2}h_3)) \cdot (S^{-1}h_6 \cdot m)
$$

\n
$$
\gamma ((S^{-1} y_1)h_4) \gamma (Sh_5) y_0 \otimes (Sh_1) y_4
$$

\n
$$
\xrightarrow{(4.13)} \sum x \gamma^{-1} (S^{-1}h_{11}) \gamma^{-1} ((S^{-3} y_5) (S^{-2}h_2)) \otimes_A \sigma ((S^{-3} y_4) (S^{-2}h_3), S^{-1}h_{10})
$$

\n
$$
((S^{-3} y_3) (S^{-2}h_4) (S^{-1}h_9) \cdot m) \sigma^{-1} ((S^{-1} y_2)h_5, Sh_8) \gamma ((S^{-1} y_1)h_6) \gamma (Sh_7) y_0
$$

\n
$$
\otimes (Sh_1) y_6
$$

\n
$$
\xrightarrow{(4.9)} \sum x \gamma^{-1} ((S^{-3} y_3) (S^{-2}h_2) (S^{-1}h_7)) \otimes_A ((S^{-3} y_2) (S^{-2}h_3) (S^{-1}h_6) \cdot m)
$$

\n
$$
\gamma ((S^{-1} y_1)h_4 (Sh_5)) y_0 \otimes (Sh_1) y_4
$$

\n
$$
= \sum x \gamma^{-1} (S^{-3} y_3) \otimes_A (S^{-3} y_2 \cdot m) \gamma (S^{-1} y_1) y_0 \otimes (Sh_1) y_4
$$

To show that F is a left B^e -module morphism, we compute

$$
F((b' \otimes b)(x \otimes y \otimes_{A^e} m))
$$

= $F((b'x \otimes yb \otimes_{A^e} m))$
= $\sum b'x\gamma^{-1}(S^{-3}(y_3b_3)) \otimes_A (S^{-3}(y_2b_2) \cdot m)\gamma(S^{-1}(y_1b_1))y_0b_0 \otimes y_4b_4$
 $\xrightarrow{(4.9)}$ $\sum b'x\gamma^{-1}(S^{-3}y_5)\gamma^{-1}(S^{-3}b_5)\sigma(S^{-3}b_4, S^{-3}y_4) \otimes_A (S^{-3}(y_3b_3) \cdot m)$
 $\sigma^{-1}(S^{-1}b_2, S^{-1}y_2)\gamma(S^{-1}b_1)\gamma(S^{-1}y_1)y_0b_0 \otimes y_6b_6$
 $\xrightarrow{(4.13)}$ $\sum b'x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-3}b_5) \otimes_A (S^{-3}b_4 \cdot (S^{-3}y_2 \cdot m))$
 $\gamma(S^{-1}b_3)\gamma(S^{-1}y_1)y_0\gamma^{-1}(S^{-1}b_2)\gamma(S^{-1}b_1)b_0 \otimes y_4b_6$
 $\xrightarrow{(4.12)}$ $\sum b'x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-3}b_3) \otimes_A S^{-3}b_2 \cdot ((S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0)$
 $\gamma(S^{-1}b_1)b_0 \otimes y_4b_4$
 $\xrightarrow{(4.20)}$ $(b' \otimes b)F(x \otimes y \otimes_{A^e} m).$

To show that F is an H^e -comodule morphism, we calculate

$$
(F \otimes id_{H^e})\rho(x \otimes y \otimes_{A^e} m)
$$

= $F(\sum x_0 \otimes y_0 \otimes_{A^e} m) \otimes x_1 \otimes y_1$
= $\sum x_0 \gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0 \otimes y_4 \otimes x_1 \otimes y_5$
= $\sum x_0 \gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0 \otimes y_6 \otimes x_1(S^{-2}y_4)(S^{-1}y_5) \otimes y_7$
 $\xrightarrow{(4.21)} \rho(\sum x_0 \gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0 \otimes y_4)$
= $\rho F(x \otimes y \otimes_{A^e} m).$

This implies that F is a morphism in $_{B^e}\mathcal{M}_H^{H^e}$.

A well-defined map $G : B \otimes_A M \otimes H \to B^e \otimes_{A^e} M$ is given by

$$
b\otimes_A m\otimes h\mapsto \sum b\gamma^{-1}(S^{-2}h_1)\otimes \gamma(h_3)\otimes_{A^e} S^{-2}h_2\cdot m.
$$

We demonstrate that G is the inverse of F by computing

$$
FG(b \otimes_A m \otimes h)
$$

= $F(\sum b\gamma^{-1}(S^{-2}h_1) \otimes \gamma(h_3) \otimes_{A^e} S^{-2}h_2 \cdot m)$
= $\sum b\gamma^{-1}(S^{-2}h_1)\gamma^{-1}(S^{-3}h_6) \otimes_A (S^{-3}h_5 \cdot (S^{-2}h_2 \cdot m))\gamma(S^{-1}h_4)\gamma(h_3) \otimes h_7$
 $\xrightarrow{(4.9)}$ $\sum b\gamma^{-1}((S^{-3}h_{10})(S^{-2}h_1))\sigma(S^{-3}h_9, S^{-2}h_2) \otimes_A (S^{-3}h_8 \cdot (S^{-2}h_3 \cdot m))$
 $\sigma(S^{-1}h_7, h_4)\gamma((S^{-1}h_6)h_5) \otimes h_{11}$
= $\sum b\gamma^{-1}((S^{-3}h_8)(S^{-2}h_1)) \otimes_A \sigma(S^{-3}h_7, S^{-2}h_2)(S^{-3}h_6 \cdot (S^{-2}h_3 \cdot m))$
 $\sigma(S^{-1}h_5, h_4) \otimes h_9$
 $\xrightarrow{(4.13)}$ $\sum b\gamma^{-1}((S^{-3}h_4)(S^{-2}h_1)) \otimes_A ((S^{-3}h_3)(S^{-2}h_2)) \cdot m \otimes h_5$
= $b \otimes_A (1_H \cdot m) \otimes h$
 $\xrightarrow{(4.11)}$ $b \otimes_A m \otimes h$,

and

$$
GF(x \otimes y \otimes_{A^e} m)
$$

= $G(\sum x\gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0 \otimes y_4)$
= $\sum x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-2}y_4) \otimes \gamma(y_6) \otimes_{A^e} S^{-2}y_5 \cdot ((S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0)$

$$
\xrightarrow{(4.12)} \sum x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-2}y_4) \otimes \gamma(y_7) \otimes_{A^e} (S^{-2}y_5 \cdot (S^{-3}y_2 \cdot m))
$$

 $y_6 \cdot (\gamma(S^{-1}y_1)y_0)$
= $\sum x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-2}y_4) \otimes (y_6 \cdot (\gamma(S^{-1}y_1)y_0))\gamma(y_7) \otimes_{A^e}$

$$
(S^{-2}y_5 \cdot (S^{-3}y_2 \cdot m))
$$

= $\sum x\gamma^{-1}(S^{-3}y_3)\gamma^{-1}(S^{-2}y_4) \otimes \gamma(y_6)\gamma(S^{-1}y_1)y_0\gamma^{-1}(y_7)\gamma(y_8) \otimes_{A^e}$

$$
(S^{-2}y_5 \cdot (S^{-3}y_2 \cdot m))
$$

\n
$$
\xrightarrow{(4.13)} \sum x \gamma^{-1} (S^{-3}y_5) \gamma^{-1} (S^{-2}y_6) \otimes \gamma(y_{10}) \gamma (S^{-1}y_1) y_0 \otimes_{A^e}
$$

\n
$$
\sigma (S^{-2}y_7, S^{-3}y_4) ((S^{-2}y_8)(S^{-3}y_3) \cdot m) \sigma^{-1}(y_9, S^{-1}y_2)
$$

\n
$$
= \sum x \gamma^{-1} (S^{-3}y_5) \gamma^{-1} (S^{-2}y_6) \sigma (S^{-2}y_7, S^{-3}y_4) \otimes \sigma^{-1}(y_9, S^{-1}y_2) \gamma(y_{10}) \gamma (S^{-1}y_1) y_0
$$

\n
$$
\otimes_{A^e} ((S^{-2}y_8)(S^{-3}y_3) \cdot m)
$$

\n
$$
\xrightarrow{(4.9)} \sum x \gamma^{-1} ((S^{-2}y_4)(S^{-3}y_3)) \otimes \gamma(y_6(S^{-1}y_1)) y_0 \otimes_{A^e} ((S^{-2}y_5)(S^{-3}y_2) \cdot m)
$$

\n
$$
= \sum x \otimes y \otimes_{A^e} (1_H \cdot m)
$$

\n
$$
\xrightarrow{(4.11)} \sum x \otimes y \otimes_{A^e} m.
$$

This completes the proof. \Box

The Nakayama automorphisms of the crossed product $\mathcal{A} \#_{\sigma} H$ are determined by those of A and H , along with the homological determinant of the H -action on A .

Theorem 4.11. Let H be a skew Calabi-Yau algebra of dimension $n, B = A \#_{\sigma} H$ be a crossed product of A with H. If A is skew Calabi-Yau of dimension d with a Nakayama automorphism μ_A , then B is also a skew Calabi-Yau algebra of dimension $n + d$ with a Nakayama automorphism μ_B which is defined by

$$
\mu_B(a \# h) = \sum \mu_A(a) \text{Hdet}(S^{-2}h_1) \# S^{-2}h_2 \chi(Sh_3),
$$

where Hdet is the homological determinant of the H -action on A with respect to a μ_A -twisted volume as defined in Definition [4.8](#page-33-0) and $\chi : H \to \mathbb{k}$ is given by $\text{Ext}^n_H(\mathbb{k}, H) \cong {\mathbb{k}}.$

Proof. Observe that B is a faithfully flat left A -module. According to Theorem [2.15,](#page-14-0) B is a skew Calabi-Yau algebra of dimension $n + d$. Let $\gamma : H \to B$ be the map defined by $h \mapsto 1 \# h$. We have the following B^e -module isomorphisms

$$
\operatorname{Ext}_{B^e}^{d+n}(B, B^e)
$$
\n
$$
\cong \operatorname{Ext}_{H}^{n}(\mathbb{k}, \operatorname{Ext}_{A^e}^d(A, B^e))
$$
\n
$$
\cong \operatorname{Ext}_{A^e}^d(A, B^e) \otimes_H \operatorname{Ext}_{H}^n(\mathbb{k}, H)
$$
\n
$$
\cong (B^e \otimes_{A^e} \operatorname{Ext}_{A^e}^d(A, A^e)) \otimes_H \operatorname{Ext}_{H}^n(\mathbb{k}, H)
$$
\n
$$
\cong B \otimes_A \operatorname{Ext}_{A^e}^d(A, A^e) \otimes H \otimes_H \operatorname{Ext}_{H}^n(\mathbb{k}, H)
$$
\n
$$
\cong B \otimes_A \operatorname{Ext}_{A^e}^d(A, A^e) \otimes H \otimes_H \operatorname{Ext}_{H}^n(\mathbb{k}, H)
$$
\n
$$
\cong B \otimes_A \operatorname{Ext}_{A^e}^d(A, A^e) \otimes H \otimes_H \chi \mathbb{k}
$$
\n
$$
\xrightarrow{(4.19)} B \otimes_A \operatorname{Ext}_{A^e}^d(A, A^e)_{\chi S^{-1}},
$$

where the B^e -module structure on $B \otimes_A \mathrm{Ext}_{A^e}^d(A, A^e)_{\chi S^{-1}}$ is given by

$$
(x \otimes y)(b \otimes_A m)
$$

= $\sum xb\gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0\chi(S^{-1}(hy_4))$
= $\sum xb\gamma^{-1}(S^{-3}y_3) \otimes_A (S^{-3}y_2 \cdot m)\gamma(S^{-1}y_1)y_0\chi(S(hy_4))$

for any $x, y, b \in B$ and $m \in \text{Ext}_{A^e}^d(A, A^e)$. Let e be a μ_A -twisted volume on A. Then $1 \otimes e$ is a generator of $B \otimes_A \mathrm{Ext}^d_{A^e}(A, A^e)_{\chi S^{-1}}$ as a left B-module. We obtain

$$
(1 \otimes e)b = \sum \gamma^{-1}(S^{-3}b_3) \otimes_A (S^{-3}b_2 \cdot e)\gamma(S^{-1}b_1)b_0\chi(5b_4)
$$

=
$$
\sum \chi(Sb_4)\gamma^{-1}(S^{-3}b_3) \otimes_A \varphi(S^{-3}b_2)\mu_A(\gamma(S^{-1}b_1)b_0)e
$$

=
$$
\sum \chi(Sb_4)\gamma^{-1}(S^{-3}b_3)\varphi(S^{-3}b_2)\mu_A(\gamma(S^{-1}b_1)b_0) \otimes_A e.
$$

Hence, $\text{Ext}_{B^e}^{d+n}(B, B^e)$ is isomorphic to B_{μ_B} as B-B-bimodules, where μ_B is an algebra morphism of B defined by

$$
\mu_B(b) = \sum \chi(Sb_4)\gamma^{-1}(S^{-3}b_3)\varphi(S^{-3}b_2)\mu_A(\gamma(S^{-1}b_1)b_0).
$$

Then the Nakayama automorphism μ_B be represented by the homological determinant Hdet and Nakayama automorphism μ_A and μ_H as follow:

$$
\mu_{B}(a \# h)
$$
\n
$$
= \sum_{i} \chi(Sh_{5})\gamma^{-1}(S^{-3}h_{4})\varphi(S^{-3}h_{3})\mu_{A}(\gamma(S^{-1}h_{2})(a \# h_{1}))
$$
\n
$$
= \sum_{i} \chi(Sh_{7})\left(\sigma^{-1}(S^{-2}h_{5}, S^{-3}h_{4}) \# S^{-2}h_{6}\right)\left(\varphi(S^{-3}h_{3}) \# 1\right)\mu_{A}\left((1 \# S^{-1}h_{2})(a \# h_{1})\right)
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{5})S^{-2}h_{7} \cdot \left(\varphi(S^{-3}h_{4})\mu_{A}\left((S^{-1}h_{3} \cdot a)\sigma(S^{-1}h_{2}, h_{1})\right)\right)
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{5})S^{-2}h_{7} \cdot \left(\varphi(S^{-3}h_{4})\mu_{A}(S^{-1}h_{3} \cdot a)\mu_{A}(\sigma(S^{-1}h_{2}, h_{1}))\right)
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{5})S^{-2}h_{7} \cdot \left(\varphi(S^{-3}h_{4})\mu_{A}(S^{-1}h_{3} \cdot a)\mu_{A}(\sigma(S^{-1}h_{2}, h_{1}))\right)
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{5})S^{-2}h_{7} \cdot \left((S^{-3}h_{4} \cdot \mu_{A}(a))\varphi(S^{-3}h_{3})\mu_{A}(\sigma(S^{-1}h_{2}, h_{1}))\right)
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{5})S^{-2}h_{7} \cdot (S^{-3}h_{4} \cdot \mu_{A}(a))
$$
\n
$$
= \sum_{i} \sigma^{-1}(S^{-2}h_{6}, S^{-3}h_{7})\sigma(S^{-2}h_{9}, S^{-3}h_{6})\left((S^{-2}h_{10})(S^{-3}h_{5})\right) \cdot \mu_{A}(a)
$$
\n
$$
\sigma^{-1}(S^{-2}h_{11}, S^{-3}h_{4})S^{-2}h_{12
$$

Since $\text{Ext}_{B^e}^{d+n}(B, B^e)$ is an invertible B-B-bimodule, it follows that μ_B is an isomorphism. Thus, we arrive at the desired conclusion. \Box

In the following, let's consider the Galois extension over a connected Hopf algebra, where the coradical has dimension one.

Lemma 4.12. [\[3,](#page-40-10) Proposition 1.3] Let B be a right comodule algebra over a connected Hopf algebra H. Then the following statements are equivalent.

- (1) $B^{coH} \subseteq B$ is a cleft extension.
- (2) $B^{coH} \subset B$ is a faithfully flat H-Galois extension.
- (3) There is a total integral $\gamma : H \to B$, that is, γ is H-colinear and $\gamma(1) = 1$.

Corollary 4.13. Assume that \mathbb{k} is of characteristic zero. Let H be a connected Hopf algebra of finite GK dimension d, and B be a cleft H-Galois extension of k. Then B is a skew Calabi-Yau algebra of dimension d. Moreover, B is Calabi-Yau if and only if H is Calabi-Yau.

Proof. By [\[42,](#page-41-14) Proposition 6.6], H is skew Calabi-Yau of dimension d. Let μ_H denote the Nakayama automorphism of H as defined in Theorem [1.4.](#page-4-1) Given that $B^{coH} = \mathbb{k}$, the homological determinant is trivial, that is, Hdet = ε . Consequently, Theorem [4.11](#page-37-0) implies that B is a skew Calabi-Yau algebra of dimension d with a Nakayama automorphism μ_B defined as

$$
\mu_B(h) = \sum 1 \# S^{-2} h_1 \chi(Sh_2) = 1 \# \mu_H(h) \in \mathbb{k} \#_{\sigma} H.
$$

Consider the coradical filtration $\{H_i\}_{i\in\mathbb{N}}$ of the connected Hopf algebra H, and let gr H denote the associated graded algebra $\bigoplus H_i/H_{i-1}$ with $H_{-1} = 0$. Since gr H is a i∈**N**

connected graded Hopf algebra, it is an iterated Ore extension of \Bbbk by [\[43,](#page-42-0) Theorem B], hence a connected graded domain. This indicates that every invertible element of H lies in $H_0 = \mathbb{k}$. Therefore, H is Calabi-Yau if and only if μ_H is the identity map on H. Define $\gamma: H \to B$ by $h \mapsto 1 \# h$. It is clear that $\{\gamma(H_i)\}_{i \in \mathbb{N}}$ forms an algebra filtration of B. As $\Delta(H_n) \subseteq \sum^{n}$ $i=0$ $H_i \otimes H_{n-i}$, it follows that $\Delta(h) - 1 \otimes h \in H_n \otimes H_{n-1}$ for any $h \in H_n$. Consequently,

$$
\gamma(h)\gamma(k) - \gamma(hk) = \sum \sigma(h_1, k_1) \# h_2 k_2 - 1 \# hk \in \gamma(H_{i+j-1})
$$

for any $h \in H_i \setminus H_{i-1}$ and $k \in H_j \setminus H_{j-1}$. Consequently, the associated graded algebra gr B is also a connected graded domain. So every invertible element of B also lies in k. This implies that B is Calabi-Yau if and only if μ_B is the identity map on B, establishing the second assertion. \Box

Let $(\mathfrak{g}, [-,-])$ be a finite dimensional Lie algebra, and $f \in Z^2(\mathfrak{g}, \mathbb{k})$ be a 2-cocycle, satisfying

$$
f(x, x) = 0
$$
 and $f(x, [y, z]) + f(y, [z, x]) + f(z, [x, y]) = 0$

for all $x, y, z \in \mathfrak{g}$. The Sridharan enveloping algebra of \mathfrak{g} with respect to f is defined to be the associative algebra

$$
\mathcal{U}_f(\mathfrak{g}) := T\mathfrak{g}/\langle x \otimes y - y \otimes x - [x, y] - f(x, y) \mid x, y \in \mathfrak{g} \rangle.
$$

There exists a canonical k-linear map $\overline{(-)} : \mathfrak{g} \to \mathcal{U}(\mathfrak{g}), x \mapsto \overline{x}$ such that $\mathcal{U}_f(\mathfrak{g})$ is a $U(\mathfrak{g})$ -comodule algebra via

$$
\rho: \mathcal{U}_f(\mathfrak{g}) \longrightarrow \mathcal{U}_f(\mathfrak{g}) \otimes \mathcal{U}(\mathfrak{g}), \qquad \overline{x} \mapsto \overline{x} \otimes 1 + 1 \otimes x, \text{ for all } x \in \mathfrak{g}.
$$

Given a basis $\{x_1, x_2, \ldots, x_d\}$ of \mathfrak{g} , a basis for $\mathcal{U}_f(\mathfrak{g})$ is given by $\{1\}$ and all monomials $\overline{x}_{i_1}\overline{x}_{i_2}\cdots\overline{x}_{i_n}$ with $i_1 \leq i_2 \leq \cdots \leq i_n$. Since the map

$$
\gamma: \mathcal{U}(\mathfrak{g}) \longrightarrow \mathcal{U}_f(\mathfrak{g}), \qquad x_{i_1} x_{i_2} \cdots x_{i_n} \mapsto \overline{x}_{i_1} \overline{x}_{i_2} \cdots \overline{x}_{i_n}
$$

is an H-comodule map, Lemma 4.12 implies that γ is convolution invertible, hence, $\Bbbk \subset \mathcal{U}_f(\mathfrak{g})$ is a cleft extension [\[15\]](#page-40-14). Observe that $\mathcal{U}(\mathfrak{g})$ is a skew Calabi-Yau algebra of dimension dim \mathfrak{g} , as stated in [\[36,](#page-41-17) Theorem A]. Moreover, $\mathcal{U}(\mathfrak{g})$ is Calabi-Yau if and only if g is unimodular, which means $tr(\text{ad}_{\mathfrak{a}}(x)) = 0$ for all $x \in \mathfrak{g}$. Consequently, Corollary [4.13](#page-39-1) yields the following conclusion.

Corollary 4.14. [\[14,](#page-40-15) Theorem 5.3] [\[35,](#page-41-25) Corollary 1.1] The Sridharan enveloping algebra $\mathcal{U}_f(\mathfrak{g})$ is Calabi-Yau if and only if the universal enveloping algebra $\mathcal{U}(\mathfrak{g})$ is Calabi-Yau if and only if g is unimodular.

ACKNOWLEDGMENTS

The research work was supported by the the NSFC (project 12301052) and Fundamental Research Funds for the Central Universities (project 2022110884).

REFERENCES

- [1] M. Aguiar, A note on strongly separable algebras, vol. 65, 2000, pp. 51–60. Colloquium on Homology and Representation Theory (Spanish) (Vaquerías, 1998).
- [2] M. Artin and W. F. Schelter, Graded algebras of global dimension 3, Adv. in Math., 66 (1987), pp. 171–216.
- [3] A. D. BELL, Comodule algebras and Galois extensions relative to polynomial algebras, free algebras, and enveloping algebras, Comm. Algebra, 28 (2000), pp. 337–362.
- [4] R. J. BLATTNER, M. COHEN, AND S. MONTGOMERY, Crossed products and inner actions of Hopf algebras, Trans. Amer. Math. Soc., 298 (1986), pp. 671–711.
- [5] K. A. BROWN AND J. J. ZHANG, Dualising complexes and twisted Hochschild (co)homology for Noetherian Hopf algebras, J. Algebra, 320 (2008), pp. 1814–1850.
- [6] T. BRZEZIŃSKI, *Translation map in quantum principal bundles*, J. Geom. Phys., 20 (1996), pp. 349–370.
- [7] T. BRZEZIŃSKI AND P. M. HAJAC, The Chern-Galois character, C. R. Math. Acad. Sci. Paris, 338 (2004), pp. 113–116.
- [8] K. Chan, C. Walton, and J. Zhang, Hopf actions and Nakayama automorphisms, J. Algebra, 409 (2014), pp. 26–53.
- [9] D. STEFAN, *Hochschild cohomology on Hopf Galois extensions*, J. Pure Appl. Algebra, 103 (1995), pp. 221–233.
- [10] Y. DOI AND M. TAKEUCHI, *Cleft comodule algebras for a bialgebra*, Comm. Algebra, 14 (1986), pp. 801–817.
- [11] $____\$ Hopf-Galois extensions of algebras, the Miyashita-Ulbrich action, and Azumaya algebras, J. Algebra, 121 (1989), pp. 488–516.
- [12] S. DASCALESCU, C. NASTASESCU, AND C. RAIANU, *Hopf algebras*, vol. 235 of Monographs and Textbooks in Pure and Applied Mathematics, Marcel Dekker, Inc., New York, 2001. An introduction.
- [13] V. GINZBURG, *Calabi-yau algebras*, (2007).
- [14] J.-W. He, F. Van Oystaeyen, and Y. Zhang, Cocommutative Calabi-Yau Hopf algebras and deformations, J. Algebra, 324 (2010), pp. 1921–1939.
- [15] P. JARA AND D. STEFAN, *Hopf-cyclic homology and relative cyclic homology of Hopf-Galois* extensions, Proc. London Math. Soc. (3), 93 (2006), pp. 138–174.
- [16] A. Kaygun, Hopf-Hochschild (co)homology of module algebras, Homology Homotopy Appl., 9 (2007), pp. 451–472.

- [17] E. Kirkman, J. Kuzmanovich, and J. J. Zhang, Gorenstein subrings of invariants under Hopf algebra actions, J. Algebra, 322 (2009), pp. 3640–3669.
- [18] H. F. KREIMER AND M. TAKEUCHI, *Hopf algebras and Galois extensions of an algebra*, Indiana Univ. Math. J., 30 (1981), pp. 675–692.
- [19] P. Le Meur, Smash products of Calabi-Yau algebras by Hopf algebras, J. Noncommut. Geom., 13 (2019), pp. 887–961.
- [20] L.-Y. Liu, Q.-S. Wu, and C. Zhu, Hopf action on Calabi-Yau algebras, in New trends in noncommutative algebra, vol. 562 of Contemp. Math., Amer. Math. Soc., Providence, RI, 2012, pp. 189–209.
- [21] D.-M. Lu, Q.-S. Wu, AND J. J. ZHANG, Homological integral of Hopf algebras, Trans. Amer. Math. Soc., 359 (2007), pp. 4945–4975.
- $[22]$ J. Lü, S.-Q. Oh, X. WANG, AND X. Yu, A note on the bijectivity of the antipode of a Hopf algebra and its applications, Proc. Amer. Math. Soc., 146 (2018), pp. 4619–4631.
- [23] J.-F. LÜ, X.-F. MAO, AND J. J. ZHANG, Nakayama automorphism and applications, Trans. Amer. Math. Soc., 369 (2017), pp. 2425–2460.
- [24] S. MONTGOMERY, *Hopf algebras and their actions on rings*, vol. 82 of CBMS Regional Conference Series in Mathematics, Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993.
- [25] T. Nakayama and T. Tsuzuku, On Frobenius extensions. I, Nagoya Math. J., 17 (1960), pp. 89–110.
- [26] \ldots , On Frobenius extensions. II, Nagoya Math. J., 19 (1961), pp. 127–148.
- [27] M. Reyes, D. Rogalski, and J. J. Zhang, Skew Calabi-Yau algebras and homological identities, Adv. Math., 264 (2014), pp. 308–354.
- [28] L. H. Rowen, Ring theory. Vol. I, vol. 127 of Pure and Applied Mathematics, Academic Press, Inc., Boston, MA, 1988.
- [29] P. SCHAUENBURG, Bialgebras over noncommutative rings and a structure theorem for Hopf bimodules, Appl. Categ. Structures, 6 (1998), pp. 193–222.
- [30] P. SCHAUENBURG AND H.-J. SCHNEIDER, On generalized Hopf Galois extensions, J. Pure Appl. Algebra, 202 (2005), pp. 168–194.
- [31] H.-J. SCHNEIDER, *Principal homogeneous spaces for arbitrary Hopf algebras*, Israel J. Math., 72 (1990), pp. 167–195.
- [32] M. VAN DEN BERGH, Non-commutative crepant resolutions, in The legacy of Niels Henrik Abel, Springer, Berlin, 2004, pp. 749–770.
- [33] C. A. WEIBEL, An introduction to homological algebra, vol. 38, Cambridge: Cambridge University Press, 1994.
- [34] Q.-S. WU AND C. ZHU, Skew group algebras of Calabi-Yau algebras, J. Algebra, 340 (2011), pp. 53–76.
- [35] Q. S. WU AND C. ZHU, *Poincaré-Birkhoff-Witt deformation of Koszul Calabi-Yau algebras*, Algebr. Represent. Theory, 16 (2013), pp. 405–420.
- [36] A. YEKUTIELI, The rigid dualizing complex of a universal enveloping algebra, J. Pure Appl. Algebra, 150 (2000), pp. 85–93.
- [37] , Derived categories, vol. 183 of Cambridge Studies in Advanced Mathematics, Cambridge University Press, Cambridge, 2020.
- [38] A. YEKUTIELI AND J. J. ZHANG, *Homological transcendence degree*, Proc. London Math. Soc. (3), 93 (2006), pp. 105–137.
- [39] X. Yu, *Hopf-Galois objects of Calabi-Yau Hopf algebras*, J. Algebra Appl., 15 (2016), pp. 1650194, 19.
- [40] X. Yu, F. Van Oystaeyen, and Y. Zhang, Cleft extensions of Koszul twisted Calabi-Yau algebras, Israel J. Math., 214 (2016), pp. 785–829.
- [41] X. YU AND Y. ZHANG, *Calabi-Yau pointed Hopf algebras of finite Cartan type*, J. Noncommut. Geom., 7 (2013), pp. 1105–1144.
- [42] G.-S. Zhou, Coideal subalgebras of pointed and connected Hopf algebras, Trans. Amer. Math. Soc., 377 (2024), pp. 2663–2709.

[43] G.-S. ZHOU, Y. SHEN, AND D.-M. LU, The structure of connected (graded) Hopf algebras, Adv. Math., 372 (2020), pp. 107292, 31.

SCHOOL OF MATHEMATICS, SHANGHAI UNIVERSITY OF FINANCE AND ECONOMICS, SHANGHAI 200433, China

Email address: zhuruipeng@sufe.edu.cn