

Chemical Distance for the Level Sets of the Gaussian Free Field

TAL PERETZ*

January 7, 2025

Abstract

We consider the Gaussian free field φ on \mathbb{Z}^d for $d \geq 3$ and study the level sets $\{\varphi \geq h\}$ in the percolating regime. We prove upper and lower bounds for the probability that the chemical distance is much larger than Euclidean distance. Our proof uses a renormalization scheme combined with a bootstrap argument.

Keywords and phrases. Gaussian free field; percolation; chemical distance; large deviations.
MSC 2020 subject classifications. 60K35, 82B43.

1. Introduction

Percolation is one of the central topics in probability theory and over the past two decades there has been active research in such models which have long-range correlation. In this article we study a canonical example, the level-sets of the Gaussian free field (GFF) on \mathbb{Z}^d for $d \geq 3$. This subject was first studied in [2], and later reintroduced in [9]. Since then, there has been much progress in understanding its percolation properties.

Denote $\{\varphi_x : x \in \mathbb{Z}^d\}$ to be the GFF on \mathbb{Z}^d , whose distribution we denote by \mathbb{P} . More concretely, this is the centered Gaussian field such that $\mathbb{E}[\varphi_x \varphi_y] = g(x, y)$, where g is the Green's function of the simple random walk on \mathbb{Z}^d , see (2.1) for the definition. For a fixed height $h \in \mathbb{R}$, we are interested in the set

$$E^{\geq h} = \{x \in \mathbb{Z}^d : \varphi_x \geq h\},$$

which we consider as a subgraph of \mathbb{Z}^d . Let $\{0 \overset{\geq h}{\longleftrightarrow} \infty\}$ denote the event there exists an infinite connected subset of $E^{\geq h}$ which contains the origin, and define the critical height

$$h_* = h_*(d) = \inf\{h \in \mathbb{R} : \mathbb{P}[0 \overset{\varphi \geq h}{\longleftrightarrow} \infty] = 0\}.$$

Rodriguez and Sznitman in [9] showed that this parameter is critical in the following sense:

- for $h < h_*$, \mathbb{P} -a.s. $E^{\geq h}$ contains a unique infinite connected component,
- for $h > h_*$, \mathbb{P} -a.s. $E^{\geq h}$ consists only of finite connected components.

Today much is known about h_* , including that $h_* \in (0, \infty)$, see [3, 9], and $h_*(d) \sim \sqrt{2 \log d}$ as $d \rightarrow \infty$, see [5]. Furthermore, from [6, 7] we have that the level sets are in a strongly supercritical regime when $h < h_*$, and in a strongly subcritical regime when $h > h_*$: for all $h \neq h_*$

$$\mathbb{P}[0 \overset{\geq h}{\longleftrightarrow} \partial B_N, 0 \overset{\geq h}{\not\longleftrightarrow} \infty] \leq \begin{cases} e^{-cN/\log N} & \text{for } d = 3 \\ e^{-cN} & \text{for } d \geq 4 \end{cases}, \quad (1.1)$$

*Technion - Israel Institute of Technology. E-mail: tal.peretz@campus.technion.ac.il

where the event refers to the origin lying in a finite connected component of $E^{\geq h}$ which intersects the boundary of $B_N = \{x \in \mathbb{Z}^d : |x|_\infty \leq N\}$.

This article is interested in the graph distance on the level sets in the supercritical regime $h < h_*$. We define the chemical distance of $x, y \in E^{\geq h}$ as

$$\rho_h(x, y) = \inf\{n \in \mathbb{N} : \exists z_1, \dots, z_n \in E^{\geq h} \text{ s.t. } |z_{i+1} - z_i|_1 = 1, z_1 = x, z_n = y\},$$

where we use the convention $\inf \emptyset = \infty$. For $h < h_*$, let $\mathcal{S}_N(h)$ be the vertices in $E^{\geq h}$ which are in connected components with $|\cdot|_\infty$ -diameter greater than N . The chemical distance of $E^{\geq h}$, as well as for many other percolation models, was first studied in [4]. The authors showed that with very high probability the chemical distance of the GFF is comparable to the Euclidean norm: for $h < h_*$ there exists constants $\Delta = \Delta(d, h) > 0$, $C = C(d, h) > 0$ and $c = c(d, h) > 0$ such that

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > CN] \leq \exp(-c(\log N)^{1+\Delta}). \quad (1.2)$$

Their proof is a multiscale argument which is robust enough to apply to many percolation models with long-range correlation. However, the stretched exponential bound is a byproduct of their methods and is not expected to be sharp for our case.

Theorem 1.1. *For $d \geq 3$ and $h < h_*$, there exists $c = c(d, h) > 0$ and $C = C(d, h) > 0$ such that*

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > CN] \leq \exp(-cN^{1-2/d}).$$

We also provide complementary lower bounds.

Theorem 1.2. *For $h < h_*$ and $\alpha > 1$, there exists $c = c(d, h, \alpha) > 0$ such that*

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > \alpha N] \geq \begin{cases} e^{-cN/\log N} & \text{for } d = 3 \\ e^{-cN} & \text{for } d \geq 4 \end{cases}.$$

The upper bound improves on (1.2), however there remains a gap between the upper and lower bounds. We give intuition on the exponents in both theorems and for the discrepancy between them. Following the proof of Theorem 1.1, one sees that the upper bound is dominated by the event the chemical distance is larger than N in a box of radius $N^{1/d}$, which has exponential cost $N^{1-2/d}$. On the other hand, the lower bound is derived by forcing the path between two vertices to make a large detour of size N , which has exponential cost $N/\log N$ and N for $d = 3$ and $d \geq 4$, respectively. As a point of comparison, for Bernoulli percolation the probability of the chemical distance being larger than Euclidean distance decays exponentially, see [1].

The technical contribution of this work is to use a renormalization scheme from [10] in order to bootstrap the estimate (1.2). We will partition B_N into boxes U of side-length L , which we will eventually take to infinity, and check the local connectivity of the level sets inside each box. To decouple the field, we will use the Markov property of the GFF: $\varphi = \psi^U + \xi^U$, where ψ^U is a local independent GFF, and ξ^U is a harmonic average field. We define a box U being good if the local field ψ^U has typical chemical distance, and that the harmonic average ξ^U is not too small inside the box. By construction, the chemical distance ρ_h behaves typically inside a box which is good with respect to ψ^U and ξ^U . Using (1.2) and the independence of the fields ψ^U , we can show that most boxes are good with respect to the local field. In order to estimate the number of good boxes with respect to the harmonic average, we will use a Gaussian estimate from [10], see Lemma 2.3. On the event the number of bad boxes is not too large, we will be able to bound the chemical distance between any two connected points in B_N .

The article is organized in the following way. In Section 2, we introduce notation and preliminary results for the level sets of GFF. In Section 3, we set up a renormalization scheme. We define notions of good and bad boxes, and prove bounds on the probability of having many bad boxes. In Section 4, we show that on the event of having few bad boxes, the chemical distance in $E^{\geq h}$ behaves well. In Section 5 we gather all of the results to prove Theorem 1.1, and in Section 6 we prove Theorem 1.2.

Throughout the rest of this text, we denote c, c', C, C', \dots to be generic numbers in $(0, \infty)$ which change from line to line, while numbered constants c_1, c_2, \dots will be fixed throughout the text. We will note if they depend on parameters, with the exception of the dimension d . Lastly, some of our inequalities will only hold for large N and L .

2. Preliminaries

2.1. Notation. For a field $\chi : \mathbb{Z}^d \rightarrow \mathbb{R}$, we write $\{\chi \geq h\} = \{z \in \mathbb{Z}^d : \chi_z \geq h\}$. For two sets $U, V \subset \mathbb{Z}^d$, we define $\{U \xleftrightarrow{\chi \geq h} V\}$ to be the event there exists a nearest-neighbor path in $\{\chi \geq h\}$ starting in U and ending in V . For shorthand, we sometimes write $\{U \xleftrightarrow{\geq h} V\} = \{U \xleftrightarrow{\varphi \geq h} V\}$. For $x \in \mathbb{Z}^d$ we denote $\{x \xleftrightarrow{\geq h} \infty\}$ to be the event x lies in the unique infinite connected component of $E^{\geq h}$. We also define the complement of these events

$$\{U \not\xleftrightarrow{\geq h} V\} = \{U \xleftrightarrow{\geq h} V\}^c, \quad \{x \not\xleftrightarrow{\geq h} \infty\} = \{x \xleftrightarrow{\geq h} \infty\}^c.$$

Denote $|\cdot|_p$ for $p \in \{1, 2, \infty\}$ to be the usual ℓ_p norm on \mathbb{Z}^d . Define $B_N(x) = \{y \in \mathbb{Z}^d : |x - y|_\infty \leq N\}$ and $B_N = B_N(0)$. For finite $U \subset \mathbb{Z}^d$, we define its boundary $\partial U = \{x \in U^c : \exists y \in U, |x - y|_1 = 1\}$.

For positive sequences a_n and b_n , we write $a_n \ll b_n$ if $\lim_{n \rightarrow \infty} a_n/b_n = 0$, $a_n = O(b_n)$ if $\limsup_{n \rightarrow \infty} a_n/b_n < \infty$ and $a_n \asymp b_n$ if $0 < \liminf_{n \rightarrow \infty} a_n/b_n \leq \limsup_{n \rightarrow \infty} a_n/b_n < \infty$.

Let $\{X_n : n \in \mathbb{N}\}$ denote the simple random walk on \mathbb{Z}^d , and let P^x denote its law conditioned on starting at $x \in \mathbb{Z}^d$. For $d \geq 3$, let

$$g(x, y) = \sum_{n=0}^{\infty} P^x[X_n = y] \quad \text{for } x, y \in \mathbb{Z}^d \tag{2.1}$$

denote its Green's function. We recall the well known fact

$$g(x, y) \asymp |x - y|_\infty^{2-d} \quad \text{as } |x - y|_\infty \rightarrow \infty. \tag{2.2}$$

For $U \subset \mathbb{Z}^d$, let $T_U = \inf\{n \geq 0, X_n \notin U\}$ denote the exit time of U , and let $H_U = \inf\{n \geq 1 : X_n \in U\}$ be the hitting time of U . Define the Green's function killed outside of U

$$g_U(x, y) = \sum_{n=0}^{\infty} P^x(X_n = y, n < T_U) \quad \text{for } x, y \in \mathbb{Z}^d.$$

For $U \subset \subset \mathbb{Z}^d$, define the equilibrium measure of U

$$e_U(x) = P^x[H_U = \infty] \quad \text{for } x \in \mathbb{Z}^d,$$

and the capacity of U

$$\text{Cap}(U) = \sum_{x \in U} e_U(x).$$

For $K \subset\subset U$, define the equilibrium measure of K relative to U

$$e_{K,U}(x) = P^x[H_K > T_U] \quad \text{for } x \in \mathbb{Z}^d,$$

and the capacity of K relative to U

$$\text{Cap}_U(K) = \sum_{x \in K} e_{K,U}(x).$$

We observe that $\text{Cap}_{\mathbb{Z}^d}(K) = \text{Cap}(K)$.

2.2. Covariance Structure. A central element of our proof is the Gibbs-Markov decomposition which is expressed in the following lemma, see for instance [10].

Proposition 2.1. *For finite $U \subset \mathbb{Z}^d$, we have*

$$\varphi_x = \psi_x^U + \xi_x^U$$

where

- ψ^U is a centered Gaussian field with covariance $g_U(\cdot, \cdot)$, independent of $\sigma(\varphi_z : z \in U^c)$.
- ξ^U is the unique harmonic function on U with boundary condition $\xi|_{U^c} = \varphi|_{U^c}$:

$$\xi_x^U = E^x[\varphi_{X_{T_U}}] = \sum_{y \in \mathbb{Z}^d} P^x[X_{T_U} = y] \varphi_y.$$

In particular, ψ^U and ξ^U are independent of each other.

We introduce integers $L \geq 1$ and $K \geq 100$. We will eventually let L go to infinity and let K be some large fixed constant. Define the lattice

$$\mathbb{L} = L\mathbb{Z}^d.$$

For $z \in \mathbb{L}$, define the boxes

$$C_z = z + [0, L]^d \subset D_z = z + [-3L, 4L]^d \subset U_z = z + [-KL + 1, L + KL - 1]^d.$$

Given a subset $\mathcal{C} \subset \mathbb{L}$, the next two lemmas will be used to decouple the fields φ^{U_z}, ξ^{U_z} for $z \in \mathcal{C}$.

Lemma 2.2 ([10, Lemma 4.1]). *Let $\mathcal{C} \subset \mathbb{L}$ be a collection of sites with mutual $|\cdot|_\infty$ -distance at least $(2K + 1)L$. Then the fields $\psi^{U_z}, z \in \mathcal{C}$, are independent.*

Lemma 2.3 ([10, Corollary 4.4]). *For all $K \geq 100$ and $a > 0$, there exist constants $c = c(K)$ and $c' = c'(K)$ such that*

$$\limsup_L \sup_{\mathcal{C}} \left\{ \log \mathbb{P} \left[\bigcap_{z \in \mathcal{C}} \left\{ \inf_{y \in D_z} \xi_y^{U_z} \leq -a \right\} \right] + c' \left(a - c \cdot \sqrt{\frac{|\mathcal{C}|}{\text{Cap}(\Sigma)}} \right)_+^2 \text{Cap}(\Sigma) \right\} \leq 0$$

where $\Sigma = \bigcup_{z \in \mathcal{C}} C_z$, and the supremum runs over all $\mathcal{C} \subset \mathbb{L}$ with mutual $|\cdot|_\infty$ -distance at least $(2K + 1)L$. We also have for every $z \in \mathbb{Z}^d$

$$\mathbb{P} \left[\sup_{y \in D_z} |\xi_y^{U_z}| \geq a \right] \leq 2 \exp \left(-c' L^{d-2} \left(a - \frac{c}{L^{(d-2)/2}} \right)_+^2 \right).$$

2.3. Connectivity Estimates. We recall connective properties of the percolation of the level sets of φ .

Definition 2.4. Denote χ to be either ψ^{U_z} or φ . For $L \geq 1$ and $h_1, h_2 \in \mathbb{R}$ such that $h_1 \leq h_2$, define $\text{LocUniq}(\chi, z, h_1, h_2)$ as the intersection of the events

$$\text{Exist}(\chi, z, h_2) = \left\{ \begin{array}{l} \text{there exists a connected component in} \\ \{\chi \geq h_2\} \cap C_z \text{ with diameter at least } L/10 \end{array} \right\}$$

and

$$\text{Unique}(\chi, z, h_1, h_2) = \left\{ \begin{array}{l} \text{for any } x \in \mathbb{L} \text{ with } |z - x|_1 = L, \text{ any connected clusters in} \\ \{\chi \geq h_2\} \cap C_x \text{ and } \{\chi \geq h_2\} \cap C_z \text{ having diameter at least } L/10 \\ \text{are connected to each other in } \{\chi \geq h_1\} \cap D_z \end{array} \right\}.$$

Lemma 2.5 ([4],[6]). *For $h_1 \leq h_2 < h_*$, there exists constants $c_1 = c_1(h_1, h_2)$ and $c = c(h_1, h_2)$ such that for any $z \in \mathbb{Z}^d$*

$$\mathbb{P}[\text{LocUniq}(\varphi, z, h_1, h_2)] > 1 - e^{-cL^{c_1}}.$$

The next result, which was already mentioned in the introduction, will be our initial estimate for controlling the chemical distance.

Lemma 2.6 ([4],[6]). *For $h_1 \leq h_2 < h_*$, there exists $C_1 = C_1(h_1, h_2)$, $c = c(h_1, h_2)$, and $c_2 = c_2(h_1, h_2)$ such that for any $z \in \mathbb{Z}^d$*

$$\mathbb{P}[\forall x, y \in \mathcal{S}_{L/10}(h_2) \cap D_z, \rho_{h_1}(x, y) \leq C_1 L] > 1 - e^{-c(\log L)^{1+c_2}}.$$

3. Renormalization

In this section, we setup a renormalization argument, which we will use to prove the upper bound in Theorem 1.1.

3.1. Good and Bad Boxes. We define notion of good and bad vertices in \mathbb{L} .

Definition 3.1. For $\varepsilon > 0$, we say $z \in \mathbb{L}$ is ξ -good at level ε if

$$\inf_{x \in D_z} \xi_x^{U_z} > -\varepsilon.$$

Else, we say z is ξ -bad at level ε .

Denote $\mathcal{S}_z^\psi(h)$ to be connected components in $\{x \in \mathbb{Z}^d : \psi_x^{U_z} \geq h\}$ with diameter at least $L/10$. We define the chemical distance with respect to the field ψ^{U_z} : for $x, y \in \{w \in \mathbb{Z}^d : \psi_w^{U_z} \geq h\}$,

$$\eta_{z,h}(x, y) = \inf \{n \in \mathbb{N} : \exists z_1, \dots, z_n \in \mathcal{S}_z^\psi(h) \text{ s.t. } |z_{i+1} - z_i|_1 = 1, z_1 = x, z_n = y\}.$$

Definition 3.2. For $h_1 \leq h_2$, we say $z \in \mathbb{L}$ is ψ -good at level (h_1, h_2) if the event

$$\text{LocUniq}(\psi^{U_z}, z, h_1, h_2) \cap \{\forall x, y \in \mathcal{S}_z^\psi(h_2) \cap D_z, \eta_{z,h_1}(x, y) \leq C_1 L\}$$

occurs. Else, we say z is ψ -bad at level (h_1, h_2) .

Finally, we will define a notion of a vertex being bad with respect to both the independent field and the harmonic part.

Definition 3.3. We say $z \in \mathbb{L}$ is good at level (ε, h_1, h_2) if it is both ξ -good at level ε and ψ -good at level (h_1, h_2) . Else, we say it is bad at level (ε, h_1, h_2) .

3.2. Bounding the number of bad boxes. The main result of this section is the following proposition, which estimates the number of bad vertices in a large box. We let $N \in \mathbb{N}$, which we will eventually take to infinity, and fix $h_1 < h_2 < h_*$ and $\varepsilon > 0$. Denote $\mathcal{B}_N = \mathcal{B}_N(\varepsilon, h_1, h_2) = \{x \in \mathbb{L} \cap B_N : x \text{ is } (\varepsilon, h_1, h_2)\text{-bad}\}$.

Proposition 3.4. *There exist constants $c = c(K)$, $c' = c'(K)$ and $C = C(K)$ such that for $L < N$ and $m \in \mathbb{N}$ satisfying $m^{2/d}/L^{d-2} \leq c'\varepsilon^2$, we have*

$$\mathbb{P}[|\mathcal{B}_N(\varepsilon, h_1, h_2)| \geq m] \leq \exp(Cm \log N - cm(\log L)^{1+c_2}) + \exp(Cm \log N - c\varepsilon^2 m^{1-2/d} L^{d-2}).$$

To prove this proposition, we will first estimate the probability of any fixed configuration of vertices in \mathbb{L} are either ψ -bad or ξ -bad.

Lemma 3.5. *Suppose $\mathcal{C} \subset \mathbb{L}$ is a set of points with mutual distance at least $(2K+1)L$. There exists $c = c(\varepsilon, h)$ independent of \mathcal{C} such that*

$$\mathbb{P}[\forall z \in \mathcal{C}, z \text{ is } \psi\text{-bad at level } (h_1, h_2)] \leq \exp(-c \cdot |\mathcal{C}|(\log L)^{1+c_2}).$$

Proof. By Lemma 2.2 and the definition of being ψ -bad, the events $\{z \text{ is } \psi\text{-bad at level } (h_1, h_2)\}$ for $z \in \mathcal{C}$ are independent. By the union bound,

$$\begin{aligned} \mathbb{P}[z \text{ is } \psi\text{-bad at level } (h_1, h_2)] &\leq \mathbb{P}[\text{LocUniq}(\psi^{U_z}, z, h_1, h_2)^c] \\ &\quad + \mathbb{P}[\exists x, y \in \mathcal{S}_z^\psi(h_2) \cap D_z, \eta_{z, h_1}(x, y) > C_1 L], \end{aligned} \tag{3.1}$$

and so we need to bound both terms on the right hand side. By translation invariance of \mathbb{P} , we only need to consider the case where z is the origin. Let

$$\delta = \frac{(h_2 - h_1) \wedge (h_* - h_2)}{4},$$

which satisfies

$$h_1 + \delta < h_2 - \delta, \quad h_2 + \delta < h_*. \tag{3.2}$$

We make the following observation: on the event

$$A_\delta = \left\{ \sup_{x \in D_0} |\varphi_x - \psi_x^{U_0}| \leq \delta \right\},$$

we have for any $h' \in \mathbb{R}$

$$\{x \in D_0 : \varphi_x \geq h' + \delta\} \subset \{x \in D_0 : \psi_x^{U_0} \geq h'\} \subset \{x \in D_0 : \varphi_x \geq h' - \delta\} \tag{3.3}$$

We bound the first term on the right hand side of (3.1). From the union bound and Lemma 2.3, we have

$$\begin{aligned} \mathbb{P}[\text{LocUniq}(\psi^{U_0}, 0, h_1, h_2)^c] &\leq \mathbb{P}[\text{Exist}(\psi^{U_0}, 0, h_1, h_2)^c, A_\delta] \\ &\quad + \mathbb{P}[\text{Unique}(\psi^{U_0}, 0, h_1, h_2)^c, A_\delta] \\ &\quad + \exp(-c\delta^2 L^{d-2}). \end{aligned} \tag{3.4}$$

Since $\text{Exist}(\chi, 0, h_2)$ is decreasing in h_2 , by (3.3) we have

$$\mathbb{P}[\text{Exist}(\psi^{U_0}, 0, h_2)^c, A_\delta] \leq \mathbb{P}[\text{Exist}(\varphi, 0, h_2 + \delta)^c] \leq e^{-cL^{c_1}},$$

where the last inequality follows from Lemma 2.5 and (3.2).

To bound the second term on the right hand side of (3.4), observe that on A_δ , the first inclusion in (3.3) implies that for $x, y \in \mathbb{Z}^d$

$$\{x \xleftrightarrow{\varphi \geq h_1 + \delta} y\} \subset \{x \xleftrightarrow{\psi^{U_0} \geq h_1} y\}, \quad (3.5)$$

while the second inclusion implies $\mathcal{S}_0^\psi(h_2) \subset \mathcal{S}_{L/10}(h_2 - \delta)$. Hence on the event A_δ

$$\text{Unique}(\varphi, 0, h_1 + \delta, h_2 - \delta) \subset \text{Unique}(\psi^{U_0}, 0, h_1, h_2),$$

and so by Lemma 2.5 and (3.2) we have

$$\mathbb{P}[\text{Unique}(\psi^{U_0}, 0, h_1, h_2)^c] \leq e^{-cL^{c_1}}.$$

Thus $\mathbb{P}[\text{LocUniq}(\psi^{U_0}, 0, h_1, h_2)^c] \leq e^{-cL^{c_1}}$.

We now bound the second term on the right hand side of (3.1). On the event A_δ we have $\eta_{0, h_1}(x, y) \leq \rho_{h_1 + \delta}(x, y)$ by (3.5), and also $\mathcal{S}_0^\psi(h_2) \subset \mathcal{S}_{L/10}(h_2 - \delta)$. Hence on A_δ we have

$$\{\exists x, y \in \mathcal{S}_0^\psi(h_2) \cap D_0, \eta_{0, h_1}(x, y) > C_1 L\} \subset \{\exists x, y \in \mathcal{S}_{L/10}(h_2 - \delta) \cap D_x, \rho_{h_1 + \delta}(x, y) > C_1 L\}$$

and so

$$\begin{aligned} & \mathbb{P}[\exists x, y \in \mathcal{S}_0^\psi(h_2) \cap D_x, \eta_{x, h_1}(x, y) > C_1 L] \\ & \leq \mathbb{P}[\sup_{y \in D_0} |\varphi_y - \psi_y^{U_0}| > \delta] + \mathbb{P}[\exists x, y \in \mathcal{S}_{L/10}(h_2 - \delta) \cap C_x, \rho_{h_1 + \delta}(x, y) > C_1 L]. \end{aligned}$$

By Lemmas 2.3 and 2.6, and (3.2) we have

$$\mathbb{P}[\exists x, y \in \mathcal{S}_{L/10}(h_2 - \delta) \cap C_x, \rho_{h_1 + \delta}(x, y) > C_1 L] \leq \exp(-cL^{d-2}) + \exp(-c(\log L)^{1+c_2}),$$

which finishes the proof. \square

Before we prove the equivalent result for the harmonic components, we will need the following lower bound on the capacity of separated boxes.

Lemma 3.6. *Let $m \in \mathbb{N}$, and suppose $\{z_1, \dots, z_m\} \subset \mathbb{L}$ is a subset of points at mutual distance at least $(2K + 1)L$. Then there exists $c_3 = c_3(K)$ independent of L and m such that*

$$\text{Cap}(\cup_{i=1}^m C_{z_i}) \geq c_3 \cdot m^{1-2/d} L^{d-2}.$$

Proof. Let $U = \cup_{i=1}^m C_{z_i}$. From [7, (2.6)], we have

$$\text{Cap}(U) \geq \frac{|U|}{\max_{x \in U} \sum_{y \in U} g(x, y)},$$

and so we need to bound $\max_{x \in U} \sum_{y \in U} g(x, y)$. By (2.2), without loss of generality we can assume $U \subset B_{\lfloor C m^{1/d} L \rfloor}(z)$ for some $z \in \mathbb{Z}^d$ and large C .

For any $x \in U$, we have the bound

$$\sum_{y \in U} g(x, y) = \sum_{r=0}^{\infty} \sum_{y \in \mathcal{D}_r(x)} g(x, y)$$

where $\mathcal{D}_r(x) = U \cap \{y \in \mathbb{Z}^d : \bar{L}r \leq |x - y|_\infty < \bar{L}(r+1)\}$ for $\bar{L} = (1+2K)L$. Since $|z_i - z_j|_\infty \geq \bar{L}$ for $i \neq j$, we have

$$\{z_1, \dots, z_m\} \cap \{x \in \mathbb{Z}^d : \bar{L}r \leq |z - x|_\infty \leq \bar{L}(r+1)\} \leq Cr^{d-1}.$$

In particular, we have $|\mathcal{D}_r \cap U| \leq Cr^{d-1}L^d$. Since $|x - y|_\infty \geq \bar{L}r$ for $y \in \mathcal{D}_r$, by (2.2) we have $g(x, y) \leq C(rL)^{2-d}$. Hence

$$\sum_{y \in U} g(x, y) \leq \sum_{r=0}^{\lfloor C'm^{1/d} \rfloor} C(rL)^{2-d} r^{d-1} L^d \leq Cm^{2/d} L^2,$$

and we conclude that

$$\text{Cap}(U) \geq \frac{cmL^d}{m^{2/d}L^2} = c \cdot m^{1-2/d} L^{d-2}.$$

□

Lemma 3.7. *Suppose \mathcal{C} is a subset of \mathbb{L} of points at mutual distance at least $(2K+1)L$. There exist $c = c(K)$ and $c' = c'(K)$ such that for every $\varepsilon > 0$ and $m, L \in \mathbb{N}$ satisfying $m^{2/d}/L^{d-2} \leq c'\varepsilon^2$, and any $\{z_1, \dots, z_m\} \subset \mathcal{C}$,*

$$\mathbb{P}[z_i \text{ is } \xi\text{-bad at level } \varepsilon \text{ for } i = 1, \dots, m] \leq \exp(-c\varepsilon^2 \text{Cap}(\cup_{i=1}^m C_{z_i})).$$

Proof. We apply Lemma 2.3

$$\begin{aligned} & \mathbb{P}[z_i \text{ is } \xi\text{-bad at level } \varepsilon \text{ for } i = 1, \dots, m] \\ &= \mathbb{P}\left[\inf_{x \in D_{z_i}} \xi_x^{U_{z_i}} \leq -\varepsilon, \text{ for } i = 1, \dots, m\right] \\ &\leq \exp\left(-c' \left(\varepsilon - c\sqrt{\frac{m}{\text{Cap}(\cup_{i=1}^m C_{z_i})}}\right)_+^2 \text{Cap}(\cup_{i=1}^m C_{z_i})\right). \end{aligned}$$

Combining Lemma 3.6 with the assumption $m^{2/d} \leq c'\varepsilon^2 L^{d-2}$, we have

$$\frac{m}{\text{Cap}(\cup_{i=1}^m C_{z_i})} \leq \frac{m}{c_3 m^{1-2/d} L^{d-2}} \leq \frac{c'\varepsilon^2}{c_3}.$$

Hence for small enough c' , we have

$$\mathbb{P}[z_i \text{ is } \xi\text{-bad at level } \varepsilon \text{ for } i = 1, \dots, m] \leq \exp(-c\varepsilon^2 \text{Cap}(\cup_{i=1}^m C_{z_i})).$$

This finishes the proof. □

With Lemmas 3.5 and 3.7, we will prove in a straightforward manner Proposition 3.4.

Proof of Proposition 3.4. Define the events

$$\mathcal{E} = \{\exists z_1, \dots, z_{\lfloor m/2 \rfloor} \in \mathbb{L} \cap B_N : z_1, \dots, z_{\lfloor m/2 \rfloor} \text{ are } \psi\text{-bad at level } (h_1, h_2)\}$$

and

$$\mathcal{F} = \{\exists z_1, \dots, z_{\lfloor m/2 \rfloor} \in \mathbb{L} \cap B_N : z_1, \dots, z_{\lfloor m/2 \rfloor} \text{ are } \xi\text{-bad at level } \varepsilon\},$$

so that by the union bound

$$\mathbb{P}[|\mathcal{B}_N(\varepsilon, h_1, h_2)| \geq m] \leq \mathbb{P}[\mathcal{E}] + \mathbb{P}[\mathcal{F}].$$

Define $\mathcal{A}_{N,L} = |\{z \in \mathbb{L} : C_z \cap B_N \neq \emptyset\}|$, and note that $\mathcal{A}_{N,L} \leq CN^d/L^d$. We bound the first term on the right hand side. We observe that if there are $\lfloor m/2 \rfloor$ boxes, we can choose in some fixed deterministic way a subset of $m' = Cm/(2K+1)^d$ boxes which are $(2K+1)L$ separated. We thus have

$$\mathbb{P}[\mathcal{E}] \leq \binom{\mathcal{A}_{N,L}}{m'} \sup_{\{z_1, \dots, z_{m'}\}} \mathbb{P}[z_1, \dots, z_{m'} \text{ are } \psi\text{-bad at level } (h_1, h_2)],$$

where the supremum is over all $\{z_1, \dots, z_{m'}\} \subset \mathbb{L} \cap B_N$ with mutual distance at least $(2K+1)L$. Applying Lemma 3.5 and the bounds $\binom{n}{k} \leq n^k$ and $m' \leq Cm/(2K+1)^d$ yields us

$$\mathbb{P}[\mathcal{E}] \leq \exp(C'm(\log m + \log N) - cm(\log L)^{1+c_2})$$

for $C' = C/(2K+1)^d$. We now bound the second term. The previous separation argument, combined with Lemmas 3.6 and 3.7, implies

$$\mathbb{P}[\mathcal{E}] \leq \binom{\mathcal{A}_{N,L}}{m'} \exp(-c\varepsilon^2 m^{1-2/d} L^{d-2}) \leq \exp(C'm \log N - c\varepsilon^2 m^{1-2/d} L^{d-2}).$$

This finishes the proof. □

4. Connectivity

In this section, we will construct a deterministic path between any two points in the same connected component based on arguments from [1, Section 3]. First, we show that we can construct a path in $E^{\geq h}$ along a sequence of good boxes.

Proposition 4.1. *If $\{z_1, \dots, z_n\} \subset \mathbb{L}$ is a sequence of nearest-neighbor points in \mathbb{L} which are all good at level (ε, h_1, h_2) , then there exists a path in*

$$E^{\geq h_1 - \varepsilon} \cap (\cup_{i=1}^n D_{z_i})$$

starting at C_{z_1} and ending in C_{z_n} whose length is bounded by $C_1 L \cdot n$.

Proof. Suppose z_j and z_{j+1} are nearest-neighbor points in \mathbb{L} . Since they are both ψ -good at level (h_1, h_2) , this implies both C_{z_j} and $C_{z_{j+1}}$ contain connected components of $\{\psi^{U_{z_j}} \geq h_2\}$ and $\{\psi^{U_{z_{j+1}}} \geq h_2\}$, respectively, with diameter greater than $L/10$. Furthermore, they are connected in $\{\psi^{U_z} \geq h_1\} \cap D_{z_j}$ and the chemical distance η_{z_j, h_1} between any two points in any of these connected components in D_{z_j} is bounded by $C_1 L$. Since z_j and z_{j+1} are both ξ -good at level ε , these connectivity properties extend to $(D_{z_j} \cup D_{z_{j+1}}) \cap E^{\geq h_1 - \varepsilon}$. Using induction finishes the proof. □

We introduce more notation. We say $\gamma \subset \mathbb{L}$ is a $*$ -connected path if $\gamma = (z_1, \dots, z_n)$ and $|z_{j+1} - z_j|_\infty = L$ for $j = 1, \dots, n-1$. We say a set $U \subset \mathbb{L}$ is $*$ -connected if for any $x, y \in U$, there exists a $*$ -connected path between x and y contained in U . Denote \mathcal{C} to be the collection of $*$ -connected components of $\{z \in \mathbb{L} : z \text{ is } (\varepsilon, h_1, h_2) \text{ bad}\}$. For $z \in \mathbb{L}$, denote \mathbf{C}_z to be the element of \mathcal{C} containing z . If z is good, denote $\mathbf{C}_z = \emptyset$ and $\partial \mathbf{C}_z = \{z\}$. For $x \in \mathbb{Z}^d$, let $\ell(x) \in \mathbb{L}$ be the unique point such that $x \in C_{\ell(x)}$. The following result follows from [1, Proposition 3.1].

Proposition 4.2 ([1, Proposition 3.1]). *Fix $x, y \in \mathbb{Z}^d$ and a $*$ -connected path $\sigma = (\sigma_1, \dots, \sigma_n) \subset \mathbb{L}$ with $\sigma_1 = \ell(x)$ and $\sigma_n = \ell(y)$. On the event $\{x \xleftrightarrow{\geq h_1 - \varepsilon} y\}$, there exists a self-avoiding path $\gamma \subset E^{\geq h_1 - \varepsilon}$ connecting x and y such that*

$$\gamma \subset W = \bigcup_{j=1}^n \bigcup_{z \in \overline{\mathbf{C}}_{\sigma_j}} D_z$$

where $\overline{\mathbf{C}}_{\sigma_j} = \mathbf{C}_{\sigma_j} \cup \partial \mathbf{C}_{\sigma_j}$

Remark 4.3. This proposition follows from the proof of [1, Proposition 3.1]. While their setting is Bernoulli bond percolation, their proof is a deterministic construction of amending a nearest-neighbor path inside a bond percolation cluster such that it lies in the boundary and interior of a cluster of bad boxes. Their proof does not rely on the distribution of the cluster, rather on the macroscopic properties of good boxes, which they denote as a ‘white boxes’. Their definition of a good box, see [1, (2.9)], is not the same as ours. However, the only property the authors use of good boxes is [1, (2.13)], which we can replace with Proposition 4.1.

5. Proof of Theorem 1.1

Fix $h < h_*$, $\varepsilon \in (0, (h_* - h)/4)$ and $N \in \mathbb{N}$. Let $h_1 = h + \varepsilon < h_*$ and $h_2 = h + 2\varepsilon < h_*$. Let $M > C/\varepsilon^2$ for some large C , and let

$$m_N = \left\lfloor \frac{N^{1-2/d}}{\log N} \right\rfloor, \quad L_N = \lfloor M(N/m_N)^{1/d} \rfloor = \lfloor M(N^{2/d} \log N)^{1/d} \rfloor.$$

Note that by our choice $m_N L_N^d \asymp N$.

Lemma 5.1. *Fix $x, y \in B_N$. On the event*

$$\{x \xleftrightarrow{\geq h} y\} \cap \{|\mathcal{B}_{2N}(\varepsilon, h_1, h_2)| \leq m_N\}$$

there exists $C_2 = C_2(h, \varepsilon)$ such that

$$\rho_h(x, y) \leq C_2 N.$$

Proof. Let $x, y \in B_N$ in the same cluster, and let $\sigma = (\sigma_1, \dots, \sigma_n)$ be a $*$ -connected path of vertices in \mathbb{L} such that $\sigma_1 = \ell(x)$ and $\sigma_n = \ell(y)$. Since $x, y \in B_N$, we can assume $n \leq CN/L_N$. By Proposition 4.2, there exists a path $\gamma \subset E^{\geq h_1 - \varepsilon} = E^{\geq h}$ connecting x and y such that $\gamma \subset W$, where

$$\gamma \subset W = \bigcup_{j=1}^n \bigcup_{z \in \overline{\mathbf{C}}_{\sigma_j}} D_z.$$

Since $x, y \in B_N$ and B_{2N} has at most m_N (ε, h_1, h_2) -bad boxes, we infer that the diameter of $\overline{\mathbf{C}}_{\sigma_j}$ is at most $7u_N L_N$ for $j = 1, \dots, n$. Since $7u_N L_N \ll N$, we conclude that $W \subset B_{2N}$. Since B_{2N} contains W and has at most m_N bad boxes, this implies that W intersects at most $(2^d + 1)m_N$ boxes which are either (ε, h_1, h_2) -bad or $*$ -neighbors of one. Hence the path crosses at most $CN/L_N + (2^d + 1)m_N$ good boxes, and at most m_N bad boxes. By Proposition 4.1, the




Figure 1: The red boxes represent bad boxes, while the blue line represents a connected subset of the level sets that connects the origin to Ne_1 . Crossing a bad box, which is unavoidable in this example, incurs a cost of $O(L^d)$, whereas avoiding bad boxes costs $O(L)$.

chemical distance inside a good box is bounded by $C_1 L_N$, while for a bad box a trivial upper bound for the chemical distance is $C' L_N^d$ for some C' . We thus get

$$|\gamma| \leq \sum_{j=1}^n \sum_{z \in \overline{C}_{\sigma_j}} \max_{x,y \in D_z} \rho_h(x,y) \leq (CN/L_N + (2^d + 1)m_N) \times C_1 L + m_N \times C' L^d \leq C_2 N,$$

for some $C_2 = C_2(h, \varepsilon)$ since $m_N L_N^d \asymp N$. □

Proof of upper bound in Theorem 1.1. We first decompose our probability

$$\begin{aligned} \mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > CN] &\leq \mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\not\leftrightarrow} y] \\ &\quad + \mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\leftrightarrow} y, \rho_h(x, y) > CN]. \end{aligned}$$

To bound the first term, we make a few observations. First, if $\{x \overset{\geq h}{\not\leftrightarrow} y\}$ occurs, then necessarily either x or y does not lie in the infinite connected component. Second, if x is in a connected component of diameter at least $N/10$, then it is connected to the boundary of $B(x, N/10)$. Hence we have

$$\begin{aligned} \{\exists x, y \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\not\leftrightarrow} y\} &\subset \{\exists x \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\not\leftrightarrow} \infty\} \\ &\subset \{\exists x \in B_N \text{ s.t. } x \overset{\geq h}{\leftrightarrow} \partial B(x, N/10), x \overset{\geq h}{\not\leftrightarrow} \infty\}. \end{aligned}$$

From (1.1), a union bound, and translation invariance, we have for $d \geq 3$

$$\mathbb{P}[\exists x \in B_N \text{ s.t. } x \overset{\geq h}{\leftrightarrow} \partial B(x, N/10), x \overset{\geq h}{\not\leftrightarrow} \infty] \leq \exp(-cN/\log N)$$

which implies

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\not\leftrightarrow} y] \leq e^{-cN/\log N}.$$

We bound the second term. By Lemma 5.1, we get

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, x \overset{\geq h}{\longleftrightarrow} y, \rho_h(x, y) > C_2 N] \leq \mathbb{P}[|\mathcal{B}_{2N}(\varepsilon, h_1, h_2)| \geq m_N].$$

By our choice of m_N and L_N we have

$$\frac{m_N^{2/d}}{L_N^{d-2}} \leq \frac{m_N^{2/d}}{M \log N \cdot m_N^{2/d}} = o(1),$$

and so we can apply Proposition 3.4. We then get

$$\begin{aligned} \mathbb{P}[|\mathcal{B}_{2N}(\varepsilon, h)| \geq m_N] &\leq \exp(Cm_N \log N - cm_N(\log L_N)^{1+c_2}) + \exp(Cm_N \log N - c\varepsilon^2 m_N^{1-2/d} L_N^{d-2}) \\ &\leq \exp(-cN^{1-2/d}(\log N)^{c_2}) + \exp(-c\varepsilon^2 N^{1-2/d}) \\ &\leq 2 \exp(-c\varepsilon^2 N^{1-2/d}) \end{aligned}$$

for large enough N , where the second inequality follows from $\varepsilon^2 m_N^{1-2/d} L_N^{d-2} \geq Cm_N \log N$ by our choice of m_N and L_N . We thus have

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > C_2 N] \leq \exp(-cN/\log N) + 2 \exp(-c\varepsilon^2 N^{1-2/d})$$

which finishes the proof. \square

6. Proof of Theorem 1.2

In this section we will prove the lower bounds using techniques from [7]. We will need the following general result for lower bounds for the GFF. For $U \subset \mathbb{Z}^d$, denote \mathbb{P}_U to be the law of ψ^U . Given $A \in \mathcal{B}(\mathbb{R}^K)$, $K \subset \mathbb{Z}^d$, and $h \in \mathbb{R}$, we define

$$A^h = \{\varphi|_K - h \in A\} \tag{6.1}$$

where $\varphi|_K - h$ refers to the field restricted to K shifted by $-h$ coordinate-wise.

Lemma 6.1 ([7, Lemma 3.2]). *Let $U_N \subset\subset V_N \subset \mathbb{Z}^d$ be subsets with $\text{Cap}_{V_N}(U_N) \rightarrow \infty$. Let $A_N \in \mathcal{B}(\mathbb{R}^{U_N})$ and $I \subset \mathbb{R}$ be an interval such that, for every $h' \in I$,*

$$\mathbb{P}_{V_N}[A_N^{h'}] \rightarrow 1.$$

Then for every $h \notin I$,

$$\liminf_{N \rightarrow \infty} \frac{1}{\text{Cap}_{V_N}(U_N)} \log \mathbb{P}_{V_N}[A_N^h] \geq -\frac{1}{2}d(h, I)^2.$$

Our strategy to prove the lower bound will be to create a long path between the two points inside $E^{\geq h}$ which is insulated by $\{\varphi < h\}$. To decouple the increasing event the points are connected, and the decreasing event the path is insulated, we will use the Gibbs-Markov decomposition.

Proof of Theorem 1.2. We first consider the more involved case $d = 3$. For $r \geq 0$ and $n \in \mathbb{N}$, define the r -neighborhood of the line segment connecting 0 to ne_1

$$P_{n,r}^{(1)} = [-r, n+r] \times [-r, r]^{d-1} \cap \mathbb{Z}^d$$




Figure 2: The event $D_N(h)$ creates a long path (in blue) inside $U_N \cap E^{\geq h}$, while the event $F_N(h)$ insulates (in red) the path, forcing the chemical distance to be at least αN .

and the r -neighborhood of the line segment connecting 0 to ne_2

$$P_{n,r}^{(2)} = [-r, r] \times [-r, n+r] \times [-r, r]^{d-2} \cap \mathbb{Z}^d.$$

We fix $\alpha, N \in \mathbb{N}$ and define the set

$$P_r = P_r(N, \alpha) = P_{\alpha N, r}^{(2)} \cup (\alpha N e_2 + P_{N, r}^{(1)}) \cup (N e_1 + P_{\alpha N, r}^{(2)}).$$

We fix $\varepsilon > 0$ and define the sets $U_N \subset V_N \subset W_N$ by $U_N = P_{\lfloor N^\varepsilon \rfloor}$, $V_N = P_{\lfloor N^{2\varepsilon} \rfloor}$ and $W_N = P_{\lfloor N^{3\varepsilon} \rfloor}$.

For a field $\chi : \mathbb{Z}^d \rightarrow \mathbb{R}$, define the events

$$D_N(\chi, h) = \{B_{\lfloor N^{\varepsilon/2} \rfloor}(0) \text{ and } B_{\lfloor N^{\varepsilon/2} \rfloor}(N e_1) \text{ are connected by a path in } \{\chi \geq h\} \cap U_N\}$$

and

$$F_N(h) = \{\partial V_N \xrightarrow{\geq h} W_N\}.$$

Since

$$D_N(h) \cap F_N(h) \subset \bigcup_{x \in \partial B_{\lfloor N^{\varepsilon/2} \rfloor}(0)} \bigcup_{y \in \partial B_{\lfloor N^{\varepsilon/2} \rfloor}(N e_1)} \{x \xrightarrow{\geq h} y, \rho_h(x, y) \geq (2\alpha + 1)N - 2\lfloor N^{\varepsilon/2} \rfloor\},$$

we have

$$D_N(h) \cap F_N(h) \subset \{\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > \alpha N\}$$

for large enough N .

To derive a lower bound for $\mathbb{P}[D_N(h) \cap F_N(h)]$, fix $\delta > 0$ and define the event

$$E_N(h, \delta) = \left\{ \inf_{y \in U_N} \xi_y^{V_N} \geq -(h_* - h + \delta) \right\}.$$

Since D_N is an increasing event, and by Proposition 2.1, we have

$$\begin{aligned}\mathbb{P}[D_N(\varphi, h) \cap E_N(h, \delta) \cap F_N(h)] &\geq \mathbb{P}[D_N(\psi^{V_N}, h_* + \delta) \cap E_N(h) \cap F_N(h)] \\ &= \mathbb{P}[D_N(\psi^{V_N}, h_* + \delta)]\mathbb{P}[E_N(h, \delta) \cap F_N(h)].\end{aligned}$$

We will now derive lower bounds for both terms. We first claim that for $h \in (h_*, h_* + \delta)$, $\mathbb{P}[E_N(h, \delta) \cap F_N(h)] \rightarrow 1$ as $N \rightarrow \infty$. From (1.1), we have for $R \in \mathbb{N}$ and $h > h_*$

$$\mathbb{P}[0 \xleftrightarrow{\geq h} \partial B_R] \leq C e^{-cR/\log R}.$$

Applying a union bound and translation invariance with this estimate for $R = \lfloor N^{3\varepsilon} \rfloor - \lfloor N^{2\varepsilon} \rfloor$, we have $\mathbb{P}[F_N(h)] \rightarrow 1$ for $h > h_*$. To bound $\mathbb{P}[E_N(h, \delta)]$, we first bound the variance of $\xi_x^{V_N}$ for $x \in U_N$. Following the computation from [7], see the equation below (3.15), we have

$$\mathbb{E}[(\xi_x^{V_N})^2] \leq C \cdot \text{dist}(x, V_N^c)^{-(d-2)} \leq C N^{-2\varepsilon(d-2)}.$$

Using Gaussian tail estimates and a union bound implies

$$\mathbb{P}\left[\inf_{y \in U_N} \xi_y^{V_N} \geq -(h_* - h + \delta)\right] \rightarrow 1$$

for $h \in (h_*, h_* + \delta)$. We conclude that $\mathbb{P}[E_N(h, \delta) \cap F_N(h)] \rightarrow 1$ for $h \in (h_*, h_* + \delta)$. Note that the events $E_N(h, \delta)$ and $F_N(h)$ take the form in the assumption for Lemma 6.1. For $E_N(h, \delta)$, this is because we can write

$$\left\{ \inf_{y \in U_N} \xi_y^{V_N} \geq -(h_* - h + \delta) \right\} = \left\{ \inf_{y \in U_N} \sum_{x \in V_N} P^y[X_{T_{V_N}} = x](\varphi_x - h) \geq -(h_* + \delta) \right\}.$$

Hence we can apply Lemma 6.1 with $V_N = \mathbb{Z}^d$ and $U_N = W_N$ and conclude that for $h < h_*$ and $\delta > 0$

$$\mathbb{P}[E_N(h, \delta) \cap F_N(h)] \geq \exp(-c \text{Cap}(W_N)).$$

To derive an upper bound for $\text{Cap}(W_N)$, recall that for $A, B \subset \mathbb{Z}^d$

$$\text{Cap}(A \cup B) \leq \text{Cap}(A) + \text{Cap}(B),$$

see [8, Proposition 2.2.1]. From [7, Lemmas 2.2, 2.5], we have that for any $\varepsilon > 0$ and $i \in \{1, 2\}$,

$$\text{Cap}(P_{N, \lfloor N^\varepsilon \rfloor}^{(i)}) \leq CN/\log N.$$

We conclude that

$$\text{Cap}(W_N) \leq 2\text{Cap}(P_{\alpha N, \lfloor N^{3\varepsilon} \rfloor}^{(1)}) + \text{Cap}(P_{\alpha N, \lfloor N^{3\varepsilon} \rfloor}^{(2)}) \leq CN \log N,$$

which implies

$$\mathbb{P}[E_N(h) \cap F_N(h)] \geq e^{-cN/\log N}.$$

Next we bound $\mathbb{P}[D_N(\psi^{V_N}, h_* + \delta)]$. We first claim that for $h < h_*$, $\mathbb{P}[D_N(\varphi, h)] \rightarrow 1$ as $N \rightarrow \infty$. From [10], we have for $h < h_*$ and $R \in \mathbb{N}$

$$\mathbb{P}[B_R \xleftrightarrow{\geq h} \partial B_{2R}] \leq C e^{-cR^{d-2}}.$$

Applying a union bound and translation invariance with this estimate for $R = \lfloor N^{\varepsilon/2} \rfloor$ proves the claim. We then have for $h < h_*$ and $\bar{\delta} < (h_* - h)/2$

$$\mathbb{P}[D_N(\psi^{V_N}, h)] \geq \mathbb{P}[D_N(\varphi, h + \bar{\delta}), \sup_{x \in U_N} \xi^{V_N} \leq \bar{\delta}] \geq \mathbb{P}[D_N(\varphi, h + \bar{\delta})] - \mathbb{P}[\sup_{x \in U_N} \xi^{V_N} > \bar{\delta}].$$

Since $h + \bar{\delta} < h_*$, we have $\mathbb{P}[D_N(\varphi, h + \bar{\delta})] \rightarrow 1$ by the earlier claim. By an earlier calculation, we also have $\mathbb{P}[\sup_{x \in U_N} \xi^{V_N} > \bar{\delta}] \rightarrow 0$. We conclude that $\mathbb{P}[D_N(\psi^{V_N}, h)] \rightarrow 1$ for $h < h_*$. We can now apply Lemma 6.1, and conclude that

$$\mathbb{P}[D_N(\psi^{V_N}, h_* + \delta)] \geq \exp(-c \text{Cap}_{V_N}(U_N)).$$

We are left to derive an upper bound for $\text{Cap}_{V_N}(U_N)$. Since $H_{A \cup B} \leq H_A$, we have $P^x[H_{A \cup B} > T_U] \leq P^x[H_A > T_U]$. This implies that for $A, B \subset U$

$$\begin{aligned} \text{Cap}_U(A \cup B) &= \sum_{x \in A \cup B} P^x[H_{A \cup B} > T_U] \leq \sum_{x \in A} P^x[H_{A \cup B} > T_U] + \sum_{x \in B} P^x[H_{A \cup B} > T_U] \\ &\leq \sum_{x \in A} P^x[H_A > T_U] + \sum_{x \in B} P^x[H_B > T_U] \\ &= \text{Cap}_U(A) + \text{Cap}_U(B). \end{aligned}$$

Hence we have

$$\text{Cap}_{V_N}(U_N) \leq \text{Cap}_{V_N}(P_{\alpha N, \lfloor N^\varepsilon \rfloor}^{(2)}) + \text{Cap}_{V_N}(\alpha N e_2 + P_{N, \lfloor N^\varepsilon \rfloor}^{(1)}) + \text{Cap}_{V_N}(N e_1 + P_{\alpha N, \lfloor N^\varepsilon \rfloor}^{(2)}).$$

Note that for $K \subset U_1 \subset U_2$, $\text{Cap}_{U_2}(K) \subset \text{Cap}_{U_1}(K)$, which implies

$$\text{Cap}_{V_N}(P_{\alpha N, \lfloor N^\varepsilon \rfloor}^{(2)}) \leq \text{Cap}_{P_{\alpha N, \lfloor N^{2\varepsilon} \rfloor}^{(2)}}(P_{\alpha N, \lfloor N^\varepsilon \rfloor}^{(2)}).$$

Applying a similar bound to the remaining terms, and using [7, Lemmas 2.2 and 2.5] we conclude that

$$\text{Cap}_{V_N}(U_N) \leq CN / \log N.$$

To summarize, for $h < h_*$

$$\mathbb{P}[\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > \alpha N] \geq \mathbb{P}[D_N(h) \cap E_N(h) \cap F_N(h)] \geq e^{-cN / \log N},$$

which finishes the proof for $d = 3$.

For $d \geq 4$, define the set $Z_n = P_0$, where we use the notation from the beginning of the proof. We define the event

$$G_N(h) = \{\forall x \in P_0, \varphi_x \geq h, \forall y \in \partial P_0, \varphi_y < h\}$$

and observe that

$$G_N(h) \subset \{0 \overset{\geq h}{\longleftrightarrow} N e_1, \rho_h(x, y) > \alpha N\} \subset \{\exists x, y \in \mathcal{S}_N(h) \cap B_N, \rho_h(x, y) > \alpha N\}.$$

Rerunning the FKG-inequality argument in [7, §3.3] yields

$$\mathbb{P}[G_N(h)] \geq e^{-cN}$$

for $d \geq 4$. This finishes the proof of the theorem. \square

Acknowledgements. I would like to thank Ron Rosenthal for advising me throughout this project and Pierre-François Rodriguez for an insightful discussion.

References

- [1] ANTAL, P., AND PISZTORA, A. On the chemical distance for supercritical Bernoulli percolation. *Ann. Probab.* 24, 2 (1996), 1036–1048.
- [2] BRICMONT, J., LEBOWITZ, J. L., AND MAES, C. Percolation in strongly correlated systems: the massless Gaussian field. *J. Statist. Phys.* 48, 5-6 (1987), 1249–1268.
- [3] DREWITZ, A., PRÉVOST, A., AND RODRIGUEZ, P.-F. The sign clusters of the massless Gaussian free field percolate on \mathbb{Z}^d , $d \geq 3$ (and more). *Comm. Math. Phys.* 362, 2 (2018), 513–546.
- [4] DREWITZ, A., RÁTH, B., AND SAPOZHNIKOV, A. On chemical distances and shape theorems in percolation models with long-range correlations. *J. Math. Phys.* 55, 8 (2014), 083307, 30.
- [5] DREWITZ, A., AND RODRIGUEZ, P.-F. High-dimensional asymptotics for percolation of Gaussian free field level sets. *Electron. J. Probab.* 20 (2015), no. 47, 39.
- [6] DUMINIL-COPIN, H., GOSWAMI, S., RODRIGUEZ, P.-F., AND SEVERO, F. Equality of critical parameters for percolation of Gaussian free field level sets. *Duke Math. J.* 172, 5 (2023), 839–913.
- [7] GOSWAMI, S., RODRIGUEZ, P.-F., AND SEVERO, F. On the radius of Gaussian free field excursion clusters. *Ann. Probab.* 50, 5 (2022), 1675–1724.
- [8] LAWLER, G. F. *Intersections of random walks.* Modern Birkhäuser Classics. Birkhäuser/Springer, New York, 2013. Reprint of the 1996 edition.
- [9] RODRIGUEZ, P.-F., AND SZNITMAN, A.-S. Phase transition and level-set percolation for the Gaussian free field. *Comm. Math. Phys.* 320, 2 (2013), 571–601.
- [10] SZNITMAN, A.-S. Disconnection and level-set percolation for the Gaussian free field. *J. Math. Soc. Japan* 67, 4 (2015), 1801–1843.