

BIFURCATIONS AND STABILITY OF SYNCHRONIZED SOLUTIONS IN THE KURAMOTO MODEL WITH UNIFORMLY SPACED NATURAL FREQUENCIES

KAZUYUKI YAGASAKI

ABSTRACT. We consider the classical Kuramoto model (KM) with natural frequencies and its continuum limit (CL), and discuss the existence of synchronized solutions and their bifurcations and stability. We specifically assume that the frequency function is symmetric and linear in the CL, so that the natural frequencies are evenly spaced in the KM. We show that in the KM, $O(2^n)$ one-parameter families of synchronized solutions are born and $O(2^n)$ saddle-node and pitchfork bifurcations occur at least, when the node number n is odd and tends to infinity. Moreover, we prove that the family of synchronized solutions obtained in the previous work suffers a saddle-node bifurcation at which its stability changes from asymptotically stable to unstable and the other families of synchronized solutions are unstable in the KM. For the CL, we show that the one-parameter family of synchronized solutions obtained in the previous work is the only continuous one and there exist uncountably many one-parameter families of noncontinuous synchronized solutions and that the former is asymptotically stable and the latter are unstable.

1. INTRODUCTION

We consider the classical Kuramoto model (KM) [28,29] with natural frequencies,

$$\frac{d}{dt}u_i^n(t) = \omega_i^n + \frac{K}{n} \sum_{j=1}^n \sin(u_j^n(t) - u_i^n(t)), \quad i \in [n] := \{1, 2, \dots, n\}, \quad (1.1)$$

where $u_i^n(t) \in \mathbb{S}^1 = \mathbb{R}/2\pi\mathbb{Z}$ and $\omega_i^n \in \mathbb{R}$ stand for the phase and natural frequency of the oscillator at the node $i \in [n]$ and K is a coupling constant. The model (1.1) and its generalizations have provided mathematical models of various problems arising in many fields including physics, biology, chemistry, engineering, and economics, and have extensively been studied. See [2, 4, 19, 40–43] for the reviews of the enormous research works.

In the previous work [26], coupled oscillator networks including (1.1) were studied and shown to be well approximated by the corresponding continuum limits (CLs), for instance, which is given by

$$\frac{\partial}{\partial t}u(t, x) = \omega(x) + K \int_I \sin(u(t, y) - u(t, x))dy \quad (1.2)$$

Date: January 8, 2025.

2020 Mathematics Subject Classification. 34C15; 45J05; 34D06; 34D20; 45M10; 37G10; 34C23.

Key words and phrases. Kuramoto model; continuum limit; synchronization; equilibrium; stationary solution; bifurcation; stability.

for (1.1) with

$$\omega_i^n = n \int_{I_i^n} \omega(x) dx, \quad i \in [n], \quad (1.3)$$

where $I = [0, 1]$ and

$$I_i^n := \begin{cases} [(i-1)/n, i/n] & \text{for } i < n; \\ [(n-1)/n, 1] & \text{for } i = n. \end{cases}$$

We call $\omega(x) \in L^2(I)$ a *frequency function*. More general cases in which networks of coupled oscillators are defined on multiple graphs which may be deterministic or random, and dense or sparse were discussed in [26]. Similar results for such networks which are defined on single graphs and do not have natural frequencies depending on each node were obtained earlier in [27, 34–36] although they are not applicable to (1.1) and (1.2). The same CL as (1.2) was also adopted for the KM (1.1) without a rigorous mathematical guarantee much earlier in [20], while similar CLs were utilized for the KM with nonlocal coupling and a single or zero natural frequency in [1, 22, 47].

A different approach for approximation of coupled oscillators like (1.1) by integro-partial differential equations called the *Vlasov equations* was more frequently used although its mathematical foundations were provided recently in [8, 11] (see also [9, 10] for further extensions). See [2, 15, 19, 42, 43] and references therein for its applications. Compared with those results, where probability density functions are treated, two advantages of the approach of [26] are to deal with a complete deterministic case in which the natural frequencies are fixed like (1.1), and to give more direct description on the dynamics of the coupled oscillators without using probability density functions.

Moreover, it was shown in [26] that the CL (1.2) has the synchronized solutions

$$u(t, x) = U(x) + \Omega t + \theta, \quad U(x) = \arcsin \left(\frac{\omega(x) - \Omega}{KC} \right), \quad (1.4)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant, the range of the function \arcsin is $[-\frac{1}{2}\pi, \frac{1}{2}\pi]$ and

$$\Omega = \int_I \omega(x) dx,$$

if there exists a constant $C > 0$ such that


$$C = \int_I \sqrt{1 - \left(\frac{\omega(x) - \Omega}{KC} \right)^2} dx. \quad (1.5)$$

This is easily confirmed by substituting (1.4) into (1.2). A solution of the form (1.4) to (1.2) was also obtained in [20] although the constant term was not contained. Similarly, the KM (1.1) is shown to have the synchronized solutions

$$u_i^n(t) = \Omega_D t + U_i + \theta, \quad U_i = \arcsin \left(\frac{\omega_i^n - \Omega_D}{KC_D} \right), \quad (1.6)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant and

$$\Omega_D = \frac{1}{n} \sum_{i=1}^n \omega_i^n,$$


 FIGURE 1. Dependence of C on K/a in (1.11).

if there exists a constant $C_D > 0$ such that

$$C_D = \frac{1}{n} \sum_{i=1}^n \sqrt{1 - \left(\frac{\omega_i^n - \Omega_D}{KC_D} \right)^2}. \quad (1.7)$$

This is easily checked by substituting (1.6) into (1.1). We also see that

$$U_i \rightarrow U(x), \quad C_D \rightarrow C \quad \text{as } n \rightarrow \infty, \quad (1.8)$$

where $i/n \rightarrow x$ as $n \rightarrow \infty$. See Section 3.1 of [26] for more details.

We specifically consider the case in which the frequency function $\omega(x)$ is symmetric and linear, i.e.,

$$\omega(x) = a(x - \frac{1}{2}), \quad (1.9)$$

where $a > 0$ is a constant.

Then the natural frequencies are placed equally, and

$$\omega_i^n = \frac{a}{2n}(2i - n - 1).$$

We have $\Omega = 0$ and compute (1.5) as

$$C = \frac{KC}{a} \left(\arcsin \left(\frac{a}{2KC} \right) + \frac{a}{2KC} \sqrt{1 - \left(\frac{a}{2KC} \right)^2} \right), \quad (1.10)$$

which yields

$$\frac{a}{K} = \varphi \left(\frac{a}{2KC} \right), \quad (1.11)$$

where $\varphi(\eta) = \arcsin \eta + \eta \sqrt{1 - \eta^2}$. Since $\varphi(\eta)$ is monotonically increasing on $(0, 1)$ and

$$\varphi(0) = 0, \quad \varphi(1) = \frac{1}{2}\pi,$$

the synchronized solutions (1.4) with $\Omega = 0$ exist if $K/a \geq 2/\pi$, and it is continuous in x . The dependence of C on a/K which is calculated from (1.11) is displayed in Fig. 1. Hence, the synchronized solutions (1.4) suddenly appear, i.e., a ‘‘bifurcation’’ occurs, at $K = K_c := 2a/\pi$ when the coupling constant K is taken as the control parameter and it is increased, say, from zero. Moreover, for n sufficiently large, the synchronized solutions (1.6) with $\Omega_D = 0$ suddenly appear near $K = K_c$ in the

KM (1.1), which is well approximated by the CL (1.2) as stated above, when the coupling constant K is increased, See Section 3.2 of [26] for the details.

The above observation seems strange from the viewpoint of dynamical systems theory [21, 46]. For finite-dimensional systems (and most infinite-dimensional systems), if an equilibrium state suddenly appears when a control parameter is changed, then a saddle-node bifurcation yielding a pair of equilibria of different stability types generally occurs. according to the theory. So we conjecture that such a bifurcation occurs in the KM (1.1) when the synchronized solutions (1.6) appear. On the other hand, in the CL (1.2), no other continuous synchronized solutions are found except for (1.4) (see Theorem 7.1). One of our main objects is to explain the reason why such a phenomenon occurs in the CL (1.2).

In this paper, we discuss synchronized solutions and their bifurcations and stability in the KM (1.1) whose natural frequencies correspond to the symmetric linear frequency function (1.9), so that they are evenly spaced. Our discussions can apply to asymmetric linear frequency functions after the system (1.1) is transformed in the rotational frame with the rotational speed Ω_D . The approaches used here can be extended to more general cases of natural frequencies although we restrict ourselves to such a simple case.

The KM (1.1) with a finite node number n have been studied in many but fewer references than the infinite node number case. The critical values of K for the existence and stability of synchronized solutions were investigated in [18, 37–39, 44, 45]. In [38, 39], it was assumed that the natural frequencies are evenly spaced as in our setting (and additionally n is odd in [38]). Relationships between the natural frequencies and synchronization were studied in [7, 18], and complete synchronization states were discussed in [13, 17, 23] for a large coupling strength. The occurrence of partial synchronization was also argued in [3, 6, 24]. For the cases of small node numbers of $n \leq 5$, bifurcations and chaotic attractors were detected and found mainly numerically in [30–32].

Here we assume that the node number n is odd as in [38], and show that $O(2^n)$ families of equilibria are born and $O(2^n)$ saddle-node and pitchfork bifurcations occur at least as $n \rightarrow \infty$ (see Remark 5.2(iii) and Proposition 5.3 for more precise statements). Moreover, we prove that the family of synchronized solutions (1.6) suffers a saddle-node bifurcation at which its stability changes from asymptotically stable to unstable and the other families of synchronized solutions are unstable. For the CL (1.2), we show that the family of synchronized solutions (1.4) is the only continuous one and there exist uncountably many one-parameter families of noncontinuous synchronized solutions, and the former is asymptotically stable but the latter are unstable.

The outline of this paper is as follow: In Section 2, the previous results of [26] are reviewed in the context of the KM (1.1) and its CL (1.2). We also give a new result on a relationship of instability between the KM (1.1) and CL (1.2). After analyzing the case of $n = 3$ in Section 3, we provide our results for the existence of synchronized solutions in Section 4, and their bifurcations and stability in Sections 5 and 6, respectively. Finally, we discuss the implications of the results in Sections 4–6 for the CL (1.2) in Section 7. We extensively use dynamical systems theory throughout the paper, and recommend the readers to consult the textbooks [21, 46] if they are not familiar with the theory.

2. PREVIOUS AND NEW FUNDAMENTAL RESULTS

We first review the results of [26] in the context of (1.1) and (1.2) and give new fundamental results on relationships between the KM (1.1) and CL (1.2), which can be extended to more general coupled oscillator networks as in [26]. See Section 2 and Appendices A and B of [26] for more details including the proofs of the theorems stated below except for Theorems 2.4 and 2.7.

Let $g(x) \in L^2(I)$. We have the following on the existence and uniqueness of solutions to the initial value problem (IVP) of the CL (1.2) (see Theorem 2.1 of [26]).

Theorem 2.1. *For any $T \in (0, \infty)$, there exists a unique solution $\mathbf{u}(t) \in C^1([0, T], L^2(I))$ to the IVP of (1.2) with*

$$u(0, x) = g(x).$$

Moreover, the solution depends continuously on g .

We next consider the IVP of the KM (1.1) and turn to the issue on convergence of solutions in (1.1) to those in the CL (1.2). Since the right-hand side of (1.1) is Lipschitz continuous in u_i^n , $i \in [n]$, we see by a fundamental result of ordinary differential equations (e.g., Theorem 2.1 of Chapter 1 of [14]) that the IVP of (1.1) has a unique solution. Let $\mathbf{u} : \mathbb{R} \rightarrow L^2(I)$ stand for an $L^2(I)$ -valued function. Given a solution $u_n(t) = (u_1^n(t), u_2^n(t), \dots, u_n^n(t))$ to the IVP of (1.1), we define an $L^2(I)$ -valued function $\mathbf{u}_n : \mathbb{R} \rightarrow L^2(I)$ as

$$\mathbf{u}_n(t) = \sum_{i=1}^n u_i^n(t) \mathbf{1}_{I_i^n},$$

where $\mathbf{1}_{I_i^n}$ represents the characteristic function of I_i^n , $i \in [n]$. Let $\|\cdot\|$ denote the norm in $L^2(I)$. We have the following from Theorem 2.3 of [26].

Theorem 2.2. *If $\mathbf{u}_n(t)$ is the solution to the IVP of (1.1) such that*

$$\lim_{n \rightarrow \infty} \|\mathbf{u}_n(0) - \mathbf{u}(0)\| = 0, \quad (2.1)$$

then for any $T > 0$ we have

$$\lim_{n \rightarrow \infty} \max_{t \in [0, T]} \|\mathbf{u}_n(t) - \mathbf{u}(t)\| = 0,$$

where $\mathbf{u}(t)$ represents the solution to the IVP of the CL (1.2).

Remark 2.3. *The condition (2.1) holds if one takes*

$$u_i^n(0) = u_{i0}^n := n \int_{I_i^n} u(0, x) dx \quad (2.2)$$

for $n > 0$ sufficiently large. Equation (2.2) was assumed instead of (2.1) in the original statement of Theorem 2.3 in [26], which can be modified as in Theorem 2.2. See the proof of Theorem 2.3 in Appendix B of [26].

For $\theta \in \mathbb{S}^1$, let $\boldsymbol{\theta}$ represent the constant function $u = \theta$. If $\bar{\mathbf{u}}_n(t)$ is a solution to the KM (1.1), then so is $\bar{\mathbf{u}}_n(t) + \boldsymbol{\theta}$ for any $\theta \in \mathbb{S}^1$. Similarly, if $\bar{\mathbf{u}}(t)$ is a solution to the CL (1.2), then so is $\bar{\mathbf{u}}(t) + \boldsymbol{\theta}$ for any $\theta \in \mathbb{S}^1$. Let $\mathcal{U}_n = \{\bar{\mathbf{u}}_n(t) + \boldsymbol{\theta} \mid \theta \in \mathbb{S}^1\}$ and $\mathcal{U} = \{\bar{\mathbf{u}}(t) + \boldsymbol{\theta} \mid \theta \in \mathbb{S}^1\}$ denote the families of solutions to (1.1) and (1.2) like (1.6) and (1.4), respectively. We say that \mathcal{U}_n (resp. \mathcal{U}) is *stable* if solutions starting in its (smaller) neighborhood remain in its (larger) neighborhood for $t \geq 0$, and *asymptotically stable* if \mathcal{U}_n (resp. \mathcal{U}) is stable and the distance between such

solutions and \mathcal{U}_n (resp. \mathcal{U}) converges to zero as $t \rightarrow \infty$. We obtain the following result.

Theorem 2.4. *Suppose that the KM(1.1) and CL (1.2) have solutions $\bar{\mathbf{u}}_n(t)$ and $\bar{\mathbf{u}}(t)$, respectively, such that*

$$\lim_{n \rightarrow \infty} \|\bar{\mathbf{u}}_n(t) - \bar{\mathbf{u}}(t)\| = 0 \quad (2.3)$$

for any $t \in [0, \infty)$. Then the following hold:

- (i) *If \mathcal{U}_n is stable (resp. asymptotically stable) for $n > 0$ sufficiently large, then \mathcal{U} is also stable (resp. asymptotically stable).*
- (ii) *If \mathcal{U} is stable, then for any $\varepsilon, T > 0$ there exists $\delta > 0$ such that for $n > 0$ sufficiently large, if*

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(0) - \bar{\mathbf{u}}_n(0) - \boldsymbol{\theta}\| < \delta, \quad (2.4)$$

then

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}_n(t) - \boldsymbol{\theta}\| < \varepsilon \quad (2.5)$$

for any $t \in [0, T]$, where $\mathbf{u}_n(t)$ is a solution to the KM (1.1). Moreover, if \mathcal{U} is asymptotically stable, then

$$\lim_{t \rightarrow \infty} \lim_{n \rightarrow \infty} \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}_n(t) - \boldsymbol{\theta}\| = 0, \quad (2.6)$$

where $\mathbf{u}_n(t)$ is any solution to (1.1) such that $\mathbf{u}_n(0)$ is contained in the basin of attraction for \mathcal{U} .

Proof. We only give a proof of part (ii) since part (i) is obtained from Theorem 2.7(i) in [26]. Suppose that \mathcal{U} is stable and let $\varepsilon > 0$ and $T > 0$ be sufficiently small and large, respectively. Then there exists $\delta > 0$ such that if $\mathbf{u}(t)$ is a solution to the CL (1.2) satisfying

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}(0) - \bar{\mathbf{u}}(0) - \boldsymbol{\theta}\| < 3\delta < \frac{1}{2}\varepsilon,$$

then

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}(t) - \bar{\mathbf{u}}(t) - \boldsymbol{\theta}\| < \varepsilon - 2\delta$$

for any $t \in [0, T]$. By Theorem 2.2 and our assumption, we can take $N > 0$ sufficiently large such that if $n > N$, then

$$\|\mathbf{u}_n(t) - \mathbf{u}(t)\|, \|\bar{\mathbf{u}}_n(t) - \bar{\mathbf{u}}(t)\| < \delta.$$

Hence, if Eq. (2.4) holds and $n > N$, then

$$\begin{aligned} \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}(0) - \bar{\mathbf{u}}(0) - \boldsymbol{\theta}\| &< \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(0) - \bar{\mathbf{u}}_n(0) - \boldsymbol{\theta}\| \\ &+ \|\mathbf{u}_n(0) - \mathbf{u}(0)\| + \|\bar{\mathbf{u}}_n(0) - \bar{\mathbf{u}}(0)\| < 3\delta, \end{aligned}$$

so that

$$\begin{aligned} \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}_n(t) - \boldsymbol{\theta}\| &< \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}(t) - \bar{\mathbf{u}}(t) - \boldsymbol{\theta}\| \\ &+ \|\mathbf{u}_n(t) - \mathbf{u}(t)\| + \|\bar{\mathbf{u}}_n(t) - \bar{\mathbf{u}}(t)\| < \varepsilon, \end{aligned}$$

which yields (2.5). Moreover, if \mathcal{U} is asymptotically stable, then we have

$$\lim_{n \rightarrow \infty} \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}_n(t) - \boldsymbol{\theta}\| = \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}(t) - \bar{\mathbf{u}}(t) - \boldsymbol{\theta}\|,$$

in which we take the limit as $t \rightarrow \infty$ to obtain (2.6). \square

Remark 2.5. *In the definition of stability and asymptotic stability of solutions to the CL (1.2), we cannot distinguish two solutions that are different only in a set with the Lebesgue measure zero.*

From the proof of Theorem 2.4 we immediately get the following without assuming the existence of the solution $\bar{\mathbf{u}}_n(t)$ to the KM (1.1) satisfying (2.3).

Corollary 2.6. *For any $\varepsilon > 0$ there exists $\delta > 0$ such that for $n > 0$ sufficiently large, if*

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(0) - \bar{\mathbf{u}}(0) - \theta\| < \delta,$$

then

$$\inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}(t) - \theta\| < \varepsilon,$$

and that if \mathcal{U} is asymptotically stable, then

$$\lim_{t \rightarrow \infty} \liminf_{n \rightarrow \infty} \inf_{\theta \in \mathbb{S}^1} \|\mathbf{u}_n(t) - \bar{\mathbf{u}}(t) - \theta\| = 0$$

in Theorem 2.4(ii).

Finally, we prove the following theorem.

Theorem 2.7. *Suppose that the hypothesis of Theorem 2.4 holds. Then the following hold:*

- (i) *If \mathcal{U}_n is unstable for $n > 0$ sufficiently large and does not converge to the same family of solutions to the CL (1.2) as a stable family to the KM (1.1) when $n \rightarrow \infty$, then \mathcal{U} is unstable;*
- (ii) *If \mathcal{U} is unstable, then so is \mathcal{U}_n for $n > 0$ sufficiently large.*

Proof. Part (ii) follows from the contraposition of Theorem 2.4(i). So we only give a proof part (i).

Suppose that \mathcal{U}_n is unstable and does not converge to the same family of solutions to the CL as a stable family of solutions to the KM in the sense of $L^2(I)$ when $n \rightarrow \infty$. Then for any $\varepsilon, \delta, T > 0$ there exists a solution $\mathbf{u}_n(t)$ to the KM (1.1) such that

$$\|\mathbf{u}_n(0) - \bar{\mathbf{u}}_n(0)\| < \frac{1}{3}\delta$$

and

$$\inf_{\theta \in \mathbb{R}} \|\mathbf{u}_n(T) - \bar{\mathbf{u}}_n(T) - \theta\| > 3\varepsilon. \quad (2.7)$$

Let $\mathbf{u}(t)$ be a solution to the CL (1.2) with

$$\|\mathbf{u}(0) - \mathbf{u}_n(0)\| < \frac{1}{3}\delta.$$

By Theorem 2.2, for any $\varepsilon > 0$ there exists an integer $N > 0$ such that

$$\|\mathbf{u}(T) - \mathbf{u}_n(T)\| < \varepsilon$$

for $n > N$. Moreover, by our assumption, we have

$$\|\bar{\mathbf{u}}(0) - \bar{\mathbf{u}}_n(0)\|, \|\bar{\mathbf{u}}(T) - \bar{\mathbf{u}}_n(T)\| < \min(\varepsilon, \frac{1}{3}\delta)$$

for $n > 0$ sufficiently large. Hence,

$$\|\mathbf{u}(0) - \bar{\mathbf{u}}(0)\| < \|\mathbf{u}(0) - \mathbf{u}_n(0)\| + \|\mathbf{u}_n(0) - \bar{\mathbf{u}}_n(0)\| + \|\bar{\mathbf{u}}_n(0) - \bar{\mathbf{u}}(0)\| < \delta$$

and

$$\begin{aligned} & \| \mathbf{u}(T) - \bar{\mathbf{u}}(T) - \boldsymbol{\theta} \| \\ & > \| \mathbf{u}_n(T) - \bar{\mathbf{u}}_n(T) - \boldsymbol{\theta} \| - \| \mathbf{u}(T) - \mathbf{u}_n(T) \| - \| \bar{\mathbf{u}}_n(T) - \bar{\mathbf{u}}(T) \| \\ & > \| \mathbf{u}_n(T) - \bar{\mathbf{u}}_n(T) - \boldsymbol{\theta} \| - 2\varepsilon, \end{aligned}$$

which yields

$$\inf_{\boldsymbol{\theta} \in \mathbb{R}} \| \mathbf{u}(T) - \bar{\mathbf{u}}(T) - \boldsymbol{\theta} \| > \varepsilon.$$

Thus, we conclude that the family \mathcal{U} is unstable. \square

Remark 2.8. *In the proof of Theorem 2.7, if \mathcal{U}_n converged to a stable family of solutions in the sense of $L^2(I)$ as $n \rightarrow \infty$, Eq. (2.7) would not hold.*

3. CASE $n = 3$

We now concentrate on the KM (1.1) and begin with the case of $n = 3$:

$$\begin{aligned} \dot{u}_1 &= -\nu + \frac{K}{3}(\sin(u_2 - u_1) + \sin(u_3 - u_1)), \\ \dot{u}_2 &= \frac{K}{3}(\sin(u_1 - u_2) + \sin(u_3 - u_2)), \\ \dot{u}_3 &= \nu + \frac{K}{3}(\sin(u_1 - u_3) + \sin(u_2 - u_3)), \end{aligned} \quad (3.1)$$

where $\nu = \frac{1}{3}a$. For the KM (1.1) with $n = 3$, a more general case in which the natural frequencies are not evenly spaced was discussed in [30–32]. Condition (1.7) becomes

$$C_D = \frac{1}{3} + \frac{2}{3} \sqrt{1 - \left(\frac{a}{3KC_D} \right)^2}. \quad (3.2)$$

As stated in Section 1, if there exists a constant C_D satisfying (3.2), then the system (3.1) has the synchronized solutions

$$u_1 = -\arcsin\left(\frac{a}{3KC_D}\right) + \theta, \quad u_2 = \theta, \quad u_3 = \arcsin\left(\frac{a}{3KC_D}\right) + \theta, \quad (3.3)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant. Moreover, $C_D \geq \frac{1}{3}$ and

$$\frac{K}{a} \geq \kappa_0 := \frac{16\sqrt{2/(15 + \sqrt{33})}}{4 + \sqrt{34 - 2\sqrt{33}}} = 0.56812\dots$$

since by (3.2)

$$\frac{a}{K} = \psi_3\left(\frac{a}{3KC_D}\right), \quad \psi_3(\xi) = \xi + 2\xi\sqrt{1 - \xi^2},$$

and $\psi_3(\xi)$ has a unique extreme (maximum) κ_0^{-1} at

$$\xi = \sqrt{\frac{15 + \sqrt{33}}{32}}$$




FIGURE 2. Dependence of C_D and \hat{C}_D on K/a in (3.2) and (3.11). The solid and dashed lines represent it for C_D and \hat{C}_D , respectively.

on $(0, 1)$. In particular, $K/a = 1$ and $\arcsin(a/3KC_D) = \frac{1}{2}\pi$ when $C_D = \frac{1}{3}$, and

$$C_D = C_{D0} := \frac{1}{3} + \frac{1}{12}(\sqrt{33} - 1) = 0.72871\dots,$$

$$\arcsin\left(\frac{a}{3KC_D}\right) = \arcsin\sqrt{\frac{15 + \sqrt{33}}{32}} = 0.93592\dots$$

when $K/a = \kappa_0$. See Fig. 2 for the dependence of C_D on K/a in (3.2).

Let $v_1 = u_1 - u_2$ and $v_2 = u_3 - u_2$. We rewrite (3.1) as

$$\begin{aligned} \dot{v}_1 &= -\nu - \frac{K}{3}(2\sin v_1 + \sin v_2 - \sin(v_2 - v_1)), \\ \dot{v}_2 &= \nu - \frac{K}{3}(\sin v_1 + 2\sin v_2 + \sin(v_2 - v_1)). \end{aligned} \quad (3.4)$$

The equilibria in (3.4) satisfy

$$\sin v_1 + \sin v_2 = 0, \quad -\sin v_1 + \sin v_2 + 2\sin(v_2 - v_1) = \frac{6\nu}{K}$$

i.e.,

$$v_1 = -v_2, \quad \sin v_2(1 + 2\cos v_2) = \frac{a}{K} \quad (3.5)$$

or

$$v_1 = v_2 - \pi, \quad v_2 = \arcsin\left(\frac{a}{K}\right) \text{ or } \pi - \arcsin\left(\frac{a}{K}\right). \quad (3.6)$$

The Jacobian matrix for the vector field of (3.4) at the equilibria is computed as

$$A = -\frac{K}{3} \begin{pmatrix} 2\cos v_1 + \cos(v_2 - v_1) & \cos v_2 - \cos(v_2 - v_1) \\ \cos v_1 - \cos(v_2 - v_1) & 2\cos v_2 + \cos(v_2 - v_1) \end{pmatrix}.$$

Hence, A only has eigenvalues with negative real parts and the equilibria are asymptotically stable if

$$\begin{aligned} -\cos v_1 - \cos v_2 - \cos(v_2 - v_1) &< 0 \\ \text{and } \cos v_1 \cos v_2 + (\cos v_1 + \cos v_2) \cos(v_2 - v_1) &> 0, \end{aligned} \quad (3.7)$$

and it has an eigenvalue with positive real parts and they are unstable if

$$\begin{aligned} & -\cos v_1 - \cos v_2 - \cos(v_2 - v_1) > 0 \\ \text{or } & \cos v_1 \cos v_2 + (\cos v_1 + \cos v_2) \cos(v_2 - v_1) < 0 \end{aligned} \quad (3.8)$$

Solutions to (3.1) corresponding to these equilibria in (3.4) are given by

$$u_1 = v_1 + \theta, \quad u_2 = \theta, \quad u_3 = v_2 + \theta,$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant, since

$$\dot{u}_2 = \frac{K}{3}(\sin v_1 + \sin v_2) = 0.$$

One of the equilibria satisfying (3.5) is given by

$$v_1 = -\arcsin\left(\frac{a}{3KC_D}\right), \quad v_2 = \arcsin\left(\frac{a}{3KC_D}\right), \quad (3.9)$$

which corresponds to the synchronized solution (3.3) in (3.1), and the other is given by

$$v_1 = \arcsin\left(\frac{a}{3K\hat{C}_D}\right) - \pi, \quad v_2 = -\arcsin\left(\frac{a}{3K\hat{C}_D}\right) + \pi, \quad (3.10)$$

where

$$\hat{C}_D = \frac{1}{3} - \frac{2}{3}\sqrt{1 - \left(\frac{a}{3K\hat{C}_D}\right)^2}. \quad (3.11)$$

We easily see that $\hat{C}_D \leq \frac{1}{3}$ and that $K/a = 1$ and $v_2 = -v_1 = \pi/2$ when $\hat{C}_D = \frac{1}{3}$. Moreover, $\hat{C}_D \rightarrow 0$ as $K/a \rightarrow \infty$, and

$$\frac{a}{K} \geq \hat{\kappa}_0 := -\frac{16\sqrt{2/(15 - \sqrt{33})}}{4 - \sqrt{34 + 2\sqrt{33}}} = 2.70996\dots$$

since by (3.11)

$$\frac{a}{K} = \hat{\psi}_3\left(\left|\frac{a}{3K\hat{C}_D}\right|\right), \quad \hat{\psi}_3(\xi) = \xi - 2\xi\sqrt{1 - \xi^2}$$

and $\hat{\psi}_3(\xi)$ has a unique extreme (maximum) $\hat{\kappa}_0^{-1}$ at

$$\xi = \sqrt{\frac{15 - \sqrt{33}}{32}}$$

on $(0, 1)$. In particular,

$$\hat{C}_D = \hat{C}_{D0} := \frac{1}{3} - \frac{1}{12}\sqrt{34 + 2\sqrt{33}} = -0.22871\dots,$$

$$\arcsin\left(\frac{a}{3K\hat{C}_D}\right) = -\arcsin\sqrt{\frac{15 - \sqrt{33}}{32}} = -0.56782\dots$$

when $K/a = \hat{\kappa}_0$. See Fig. 2 for the dependence of \hat{C}_D on K/a in (3.11).

On the other hand, the equilibria satisfying (3.6) exist when $K/a \geq 1$ and coalesce at $K/a = 1$. Thus, a saddle-node bifurcation where the equilibria given by (3.9) are born occurs at $K/a = \hat{\kappa}_0$ and a pitchfork bifurcation where the equilibria




FIGURE 3. Numerically computed bifurcation diagram of equilibria in (3.4).




FIGURE 4. Dependence of C_D and \hat{C}_D^1 on K/a in (4.2): (a) $n = 5$; (b) $n = 11$. The solid and dashed lines represent it for C_D and \hat{C}_D^1 , respectively, where $\sigma = \{-1, 1, 1, -1\}$ and $\{-1, 1, \dots, 1, -1\}$.

given by (3.6) and (3.10) are born occurs at $K/a = 1$. Moreover, the equilibria given by (3.9) are asymptotically stable if $v_2 = -v_1$ is less than

$$v_0 = \arcsin \left(\sqrt{\frac{15 + \sqrt{33}}{32}} \right),$$

and they are unstable if it is greater than v_0 , while the other equilibria are always unstable, as shown in Appendix A.

A numerically computed bifurcation diagram of equilibria in (3.4) is displayed in Fig. 3. Here the computer tool called AUTO [16] was used. The solid and dashed lines represent stable and unstable equilibria, respectively. The red and black lines, respectively, corresponds to the equilibrium given by (3.9) and the others, which are given by (3.6) and (3.10). Only the equilibrium given by (3.9) can be stable.

4. SYNCHRONIZED SOLUTIONS

In the next three sections we consider the case in which the node number n is odd and greater than three for (1.1), and discuss the existence of synchronized solutions and their bifurcations and stability.

Let $n = 2n_0 + 1$ with $n_0 > 1$. Equation (1.1) becomes

$$\dot{u}_i = (i - n_0 - 1)\nu + \frac{K}{n} \sum_{j=1}^n \sin(u_j - u_i), \quad i \in [n], \quad (4.1)$$

where $\nu = a/n$. Condition (1.7) becomes

$$C_D = \frac{1}{n} + \frac{2}{n} \sum_{j=1}^{n_0} \sqrt{1 - \left(\frac{ja}{nKC_D}\right)^2}. \quad (4.2)$$

As stated in Section 1, if there exists a constant C_D satisfying (4.2), then the system (4.1) has the synchronized solutions

$$u_i = \arcsin\left(\frac{(i - n_0 - 1)a}{nKC_D}\right) + \theta, \quad i \in [n], \quad (4.3)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant. Moreover,

$$C_D \geq C_{D1} := \frac{1}{n} + \frac{2}{n} \sum_{j=1}^{n_0} \sqrt{1 - \left(\frac{j}{n_0}\right)^2},$$

which is $\frac{1}{5}(1 + \sqrt{3}) = 0.5464\dots$ for $n = 5$ and $\frac{1}{55}(19 + 4\sqrt{6} + 2\sqrt{21}) = 0.69023\dots$ for $n = 11$, and $K/a \geq \kappa_0 \approx 0.60670\dots$ for $n = 5$ and $0.62791\dots$ for $n = 11$, since by (4.2)

$$\frac{a}{K} = \psi_n\left(\frac{n_0 a}{nKC_D}\right), \quad \psi_n(\xi) = \frac{1}{n_0} \left(\xi + 2\xi \sum_{j=1}^{n_0} \sqrt{1 - \left(\frac{j}{n_0}\xi\right)^2} \right)$$

and $\psi_n(\xi)$ has a unique extreme (maximum) κ_0^{-1} at $\xi = 0.88209\dots$ for $n = 5$ and $0.94573\dots$ for $n = 11$. We also have

$$\frac{K}{a} = \kappa_1 := \frac{n_0}{nC_{D1}},$$

which is $\sqrt{3} - 1 = 0.73205\dots$ for $n = 5$ and $25/(19 + 4\sqrt{6} + 2\sqrt{21}) = 0.65853\dots$ for $n = 11$ when $C_D = C_{D1}$, and $C_{D0} = 0.74741\dots$ for $n = 5$ and $0.76543\dots$ for $n = 11$ when $K/a = \kappa_0$. See Fig. 4 for the dependence of C_D on K/a in (4.2) for $n = 5$ and 11 .

Let

$$v_i = \begin{cases} u_i - u_{n_0+1} & \text{for } i \leq n_0; \\ u_{i+1} - u_{n_0+1} & \text{for } n_0 < i \leq 2n_0. \end{cases}$$

We rewrite (4.1) as

$$\begin{aligned} \dot{v}_i &= (i - n_0 - 1)\nu - \frac{K}{n} \left(2 \sin v_i + \sum_{j=1, j \neq i}^{2n_0} (\sin v_j - \sin(v_j - v_i)) \right), \quad i \leq n_0, \\ \dot{v}_i &= (i - n_0)\nu - \frac{K}{n} \left(2 \sin v_i + \sum_{j=1, j \neq i}^{2n_0} (\sin v_j - \sin(v_j - v_i)) \right), \quad n_0 < i \leq 2n_0, \end{aligned} \quad (4.4)$$

where $\nu = a/n$. We easily see that the system (4.4) has an equilibrium

$$v_i = \begin{cases} \arcsin \left(\frac{(i - n_0 - 1)a}{nK C_D} \right) & \text{for } i \leq n_0; \\ \arcsin \left(\frac{(i - n_0)a}{nK C_D} \right) & \text{for } n_0 < i \leq 2n_0, \end{cases} \quad (4.5)$$

which corresponds to the synchronized solution (4.3) in (4.1) and satisfies the relations

$$v_i = -v_{2n_0-i+1} \quad (4.6)$$

and

$$v_i \in [-\frac{1}{2}\pi, 0), \quad v_{n_0+i} \in (0, \frac{1}{2}\pi] \quad (4.7)$$

for $i \in [n_0]$.

Let $\sigma = \{\sigma_i\}_{i=1}^{2n_0}$ be a sequence of length $2n_0$ with $\sigma_i \in \{-1, 1\}$, $i \in [2n_0]$, and let

$$\Sigma_{n_0} = \{ \sigma = \{\sigma_i\}_{i=1}^{2n_0} \mid \sigma_i \in \{-1, 1\}, i \in [2n_0] \}.$$

For each $\sigma \in \Sigma_{n_0}$, we define \hat{C}_D^σ such that it satisfies

$$\begin{aligned} \hat{C}_D^\sigma &= \frac{1}{n} + \frac{1}{n} \sum_{i=1}^{n_0} \sigma_i \sqrt{1 - \left(\frac{(i - n_0 - 1)a}{nK \hat{C}_D^\sigma} \right)^2} \\ &\quad + \frac{1}{n} \sum_{i=n_0+1}^{2n_0} \sigma_i \sqrt{1 - \left(\frac{(i - n_0)a}{nK \hat{C}_D^\sigma} \right)^2}, \end{aligned} \quad (4.8)$$

and write

$$v_i^\sigma = \begin{cases} \phi_i & \text{if } \sigma_i = 1; \\ \pi - \phi_i & \text{if } \sigma_i = -1 \text{ and } \phi_i > 0; \\ -\phi_i - \pi & \text{if } \sigma_i = -1 \text{ and } \phi_i < 0, \end{cases} \quad (4.9)$$

where

$$\phi_i = \begin{cases} \arcsin \left(\frac{(i - n_0 - 1)a}{nK \hat{C}_D^\sigma} \right) & \text{for } i \leq n_0; \\ \arcsin \left(\frac{(i - n_0)a}{nK \hat{C}_D^\sigma} \right) & \text{for } n_0 < i \leq 2n_0. \end{cases} \quad (4.10)$$

Note that $\phi_i \neq 0$, $i \in [2n_0]$. We have the following.

Theorem 4.1.

- (i) For each $\sigma \in \Sigma_{n_0}$ $v = v^\sigma \in \mathbb{T}^{2n_0} := \prod_{i=1}^{2n_0} \mathbb{S}^1$ gives an equilibrium in (4.4), when \hat{C}_D^σ satisfies (4.8). Moreover, no other equilibrium exists in (4.4).

- (ii) Fix $K/a > 0$ and $i \in [n_0]$. If the equilibrium $v = v^\sigma$ with $\sigma_i = -1$ and $\sigma_{2n_0-i+1} = 1$ exists, then so does $v = v^{\bar{\sigma}}$ with $\bar{\sigma}_i = 1$, $\bar{\sigma}_{2n_0-i+1} = -1$ and $\bar{\sigma}_j = \sigma_j$, $j \in [2n_0] \setminus \{i, 2n_0 - i + 1\}$, and vice versa, where $\chi^\sigma(\xi) = \chi^{\bar{\sigma}}(\xi)$ and their absolute values equals K/a .

Proof. Let $w = (w_1, \dots, w_{2n_0})$ denote an equilibrium of (4.4). We begin with the following two lemmas.

Lemma 4.2. *We have*

$$\sum_{i=1}^{2n_0} \sin w_i = 0.$$

Proof. Since the right-hand sides of (4.4) are zero when $v_i = w_i$, $i \in [2n_0]$, we sum up them to obtain the desired result. \square

Let

$$C_w = \sum_{i=1}^{2n_0} \cos w_i,$$

and let $i' = 2n_0 - i + 1$ for $i \in [n_0]$.

Lemma 4.3. *If $C_w \neq -1$, then*

$$\sin w_i + \sin w_{i'} = 0.$$

Proof. Adding two of the right-hand sides of (4.4) for i and i' and using Lemma 4.2, we have

$$(C_w + 1)(\sin w_i + \sin w_{i'}) = 0.$$

This yields the desired result. \square

Using Lemma 4.3 and (4.4), we have

$$\begin{aligned} (i - n_0 - 1)\nu - \frac{K}{n}(C_w + 1) \sin w_i &= 0 & \text{for } i \leq n_0; \\ (i - n_0)\nu - \frac{K}{n}(C_w + 1) \sin w_i &= 0 & \text{for } n_0 < i \leq 2n_0. \end{aligned}$$

Obviously, $C_w \neq -1$ since the above relations do not hold otherwise. Hence, $w_i = \phi_i$ or $w_i = \pi - \phi_i$, $i \in [2n_0]$, where ϕ_i is given by (4.10) with $\hat{C}_D^\sigma = (C_w + 1)/n$, since $\nu = a/n$. This yields part (i). By (4.8) we have $\hat{C}_D^\sigma = \hat{C}_D^{\bar{\sigma}}$, from which part (ii) follows. Thus, we complete the proof. \square

In particular, $v = v^\sigma$ is the same as the equilibrium given by (4.5) and $\hat{C}_D^\sigma = C_D$ when $\sigma = \{1, \dots, 1\}$. Moreover, $\hat{C}_D^\sigma = \hat{C}_D$ when $\sigma = \{-1, -1\}$ in the case of $n = 3$. See Fig. 4 for the dependence of \hat{C}^σ on K/a when $\sigma = \{-1, 1, 1, -1\}$ and $\{-1, 1, \dots, 1, -1\}$ for $n = 5$ and $n = 11$, respectively. Solutions to (4.1) corresponding to these equilibria in (4.4) are given by

$$u_{n_0+1} = \theta, \quad u_i = \begin{cases} v_i + \theta & \text{for } i \leq n_0; \\ v_{i-1} + \theta & \text{for } i \geq n_0 + 2, \end{cases} \quad (4.11)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant, since

$$\dot{u}_{n_0+1} = \frac{K}{n} \sum_{i=1}^{2n_0} \sin v_i = 0$$

by Lemma 4.2.

For $\sigma \in \Sigma_{n_0}$, we define

$$\chi^\sigma(\xi) = \frac{\xi}{n_0} \left(1 + \sum_{i=1}^{n_0} \sigma_i \sqrt{1 - \left(\frac{i - n_0 - 1}{n_0} \xi \right)^2} + \sum_{i=n_0+1}^{2n_0} \sigma_i \sqrt{1 - \left(\frac{i - n_0}{n_0} \xi \right)^2} \right),$$

where $\xi \in [0, 1]$. Obviously, $\chi^\sigma(0) = 0$ for any $\sigma \in \Sigma_{n_0}$. In general, we can choose an integer $\ell \in [n_0]$ and a monotonically increasing sequence of integers $\{i_k\}_{k=1}^\ell$ with $i_k \in [n_0]$, $k \in [\ell]$, such that $\sigma_j = \sigma_{2n_0-j+1}$ if $j = i_k$ and $\sigma_j \neq \sigma_{2n_0-j+1}$ if $j \neq i_k$ for any $k \in [\ell]$. So we write

$$\begin{aligned} \chi^\sigma(\xi) &= \frac{\xi}{n_0} \left(1 + \sum_{k=1}^\ell (\sigma_{i_k} + \sigma_{2n_0-i_k+1}) \sqrt{1 - \left(\frac{i_k - n_0 - 1}{n_0} \xi \right)^2} \right) \\ &= \frac{\xi}{n_0} \left(1 + 2 \sum_{k=1}^\ell \sigma_{i_k} \sqrt{1 - \left(\frac{i_k - n_0 - 1}{n_0} \xi \right)^2} \right). \end{aligned} \quad (4.12)$$

Letting $\xi = n_0 a / nK |\hat{C}_D^\sigma|$, we rewrite the relation (4.8) as

$$\frac{a}{K} = |\chi^\sigma(\xi)|. \quad (4.13)$$

Hence, we have $\xi \neq 0$ for any $K > 0$ since $\chi^\sigma(0) = 0$. In particular, $\psi_n(\xi) = \chi^\sigma(\xi)$ when $\sigma_j = 1$, $j \in [2n_0]$. From Theorem 4.1(i) we immediately obtain the following corollary.

Corollary 4.4. *If $\xi \in (0, 1]$ satisfies (4.13) for $\sigma \in \Sigma_{n_0}$ and $K > 0$, then v^σ given by (4.9) with*

$$\phi_i = \begin{cases} \pm \arcsin \left(\frac{(i - n_0 - 1)\xi}{n_0} \right) & \text{for } i \leq n_0; \\ \pm \arcsin \left(\frac{(i - n_0)\xi}{n_0} \right) & \text{for } n_0 < i \leq 2n_0, \end{cases} \quad (4.14)$$

instead of (4.10) is an equilibrium in (4.4), where the upper or lower sign is taken, depending on whether $\chi^\sigma(\xi)$ is positive or not.

Remark 4.5.

- (i) It is clear that $v^\sigma \neq v^{\bar{\sigma}}$ if $\sigma \neq \bar{\sigma}$ and $\xi \neq 1$.
- (ii) Mirollo and Strogatz [37], and Verwoerd and Mason [45] used an expression similar to the function $\chi^\sigma(\xi)$ (and equivalently to Eq. (4.8)) to obtain equilibria in the KM (1.1) when ω_i , $i \in [n]$, are not necessarily evenly spaced. In particular, the former half of Theorem 4.1(i) is equivalent to a special case of Theorem 2 of [45] although Theorem 4.1 contains stronger statements than it.

5. BIFURCATIONS

We next discuss bifurcations of the equilibria in (4.4) detected by Theorem 4.1 and equivalently by Corollary 4.4. The relation (4.13) gives a branch of equilibria when K is taken as a control parameter. So we have the following result.

Theorem 5.1.

- (i) The equilibrium v^σ suffers a supercritical (resp. subcritical) saddle-node bifurcation at

$$K = \frac{a}{|\chi^\sigma(\xi_0)|} \quad (5.1)$$

in (4.4) with $\xi = \xi_0$ on $(0, 1)$ if and only if $|\chi^\sigma(\xi)|$ has a local maximum (resp. minimum). In particular, if

$$\sigma_1, \sigma_{2n_0} = 1, \quad \chi^\sigma(1) \geq 0 \quad (5.2)$$

or

$$\sigma_1, \sigma_{2n_0} = -1, \quad \chi^\sigma(1) \leq 0, \quad (5.3)$$

then a supercritical saddle-node bifurcation occurs. Moreover, if $\sigma_i = 1$, $i \in [2n_0]$, then $\chi^\sigma(\xi)$ has a unique maximum and no local minimum, and v^σ suffers only one saddle-node bifurcation.

- (ii) Let $v^{\sigma^{\pm\pm}}$ be four equilibria in (4.4) such that

$$\sigma_i^{++} = \sigma_i^{--} = \sigma_i^{+-} = \sigma_i^{-+}, \quad i \neq 1, 2n_0,$$

and

$$\sigma_1^{++}, \sigma_1^{+-} = 1, \quad \sigma_1^{-+}, \sigma_1^{--} = -1, \quad \sigma_{2n_0}^{++}, \sigma_{2n_0}^{-+} = 1, \quad \sigma_{2n_0}^{+-}, \sigma_{2n_0}^{--} = -1.$$

If $\chi^\sigma(1) \neq 0$, then a pitchfork bifurcation where $v^{\sigma^{++}}$ changes to $v^{\sigma^{--}}$ and where $v^{\sigma^{+-}}$ and $v^{\sigma^{-+}}$ are born occurs at

$$K = \frac{a}{|\chi^\sigma(1)|}, \quad (5.4)$$

where any of $\sigma^{\pm\pm}$ may be chosen as σ . Moreover, the bifurcation is super- or subcritical, depending on whether

$$\chi^\sigma(1) \frac{d\chi^\sigma}{d\xi}(1) \quad \text{with } \sigma = \sigma^{+-} \text{ and } \sigma^{-+} \quad (5.5)$$

is positive or negative, where any of σ^{+-} and σ^{-+} may be chosen as σ .

Proof. We easily see that if $\xi \in (0, 1)$ satisfies (4.13), then there exists the equilibrium v^σ with $\hat{C}_D^\sigma = n_0 a / n K \xi$. Hence, If $|\chi^\sigma(\xi)|$ has a local maximum (resp. minimum) at $\xi = \xi_0$ on $(0, 1)$, then near (5.1), no value of ξ satisfies (4.13) for smaller (resp. larger) values of K than it but two values of ξ satisfies it for larger (resp. smaller) values of K than it, so that the equilibrium v^σ suffers a supercritical (resp. subcritical) saddle-node bifurcation there.

If condition (5.2) or (5.3) holds, then $|\chi^\sigma(\xi)|$ has a local maximum, since $\chi^\sigma(0) = 0$ and

$$\frac{d\chi^\sigma}{d\xi}(\xi) \rightarrow -\infty \quad \text{or} \quad \frac{d\chi^\sigma}{d\xi}(\xi) \rightarrow +\infty \quad (5.6)$$

as $\xi \rightarrow 1$, depending on whether $\sigma_1, \sigma_{2n_0} = 1$ or -1 . Moreover, when $\sigma_i = 1$, $i \in [2n_0]$,

$$\chi^\sigma(\xi) = \frac{\xi}{n_0} \left(1 + 2 \sum_{i=1}^{n_0} \sqrt{1 - \left(\frac{i}{n_0} \xi \right)^2} \right) > 0$$

and

$$\begin{aligned} \frac{d\chi^\sigma}{d\xi}(\xi) &= \frac{1}{n_0} \left(1 + 2 \sum_{i=1}^{n_0} \left(1 - 2 \left(\frac{i}{n_0} \xi \right)^2 \right) / \sqrt{1 - \left(\frac{i}{n_0} \xi \right)^2} \right) \\ &= \frac{1}{n_0} \left(1 + 2 \sum_{i=1}^{n_0} \left(2 \sqrt{1 - \left(\frac{i}{n_0} \xi \right)^2} - 1 / \sqrt{1 - \left(\frac{i}{n_0} \xi \right)^2} \right) \right) \end{aligned}$$

is monotonically decreasing on $(0, 1)$ and goes to $-\infty$ as $\xi \rightarrow 1$, so that $\chi^\sigma(\xi)$ has a unique maximum and no minimum. Thus, we prove part (i).

We turn to the proof of part (ii). When $\xi = 1$, the four equilibria coincide and the corresponding functions $\chi^{\sigma^{\pm\pm}}(\xi)$ have the same value. To obtain the desired result, we only have to notice that σ^{++} and σ^{--} , respectively, satisfy the first and second equations of (5.6) as $\xi \rightarrow 1$, and that $v^{\sigma^{+-}}$ and $v^{\sigma^{-+}}$ exist in a pair by Theorem 4.1(ii) for larger or smaller values of K than and near (5.4), depending on whether the quantity (5.5), which takes the same value for both of $\sigma = \sigma^{+-}$ and σ^{-+} , is positive or negative. \square

Remark 5.2.

- (i) *It follows from Theorem 5.1(i) that one of the equilibria v^σ with $\sigma = \sigma^{++}$ or σ^{--} in Theorem 5.1(ii) suffers a saddle-node bifurcation. Moreover, the equilibrium v^σ with $\sigma = \sigma^{+-}$ and σ^{-+} can suffer it. Indeed, when $\sigma_i \neq \sigma_{2n_0-i+1}$, $i \in [n_0] \setminus \{2\}$ and $\sigma_2 = \sigma_{2n_0-1} = 1$, we have*

$$\chi^\sigma(\xi) = \frac{\xi}{n_0} \left(1 + 2 \sqrt{1 - \left(\frac{n_0-1}{n_0} \xi \right)^2} \right) > 0$$

on $(0, 1]$ (see (4.12)) and

$$\frac{d\chi^\sigma}{d\xi}(\xi) = \frac{1}{n_0} \left(1 + 2 \left(1 - 2 \left(\frac{n_0-1}{n_0} \xi \right)^2 \right) / \sqrt{1 - \left(\frac{n_0-1}{n_0} \xi \right)^2} \right),$$

which is monotonically decreasing. If $n_0 \geq 6$, then $\chi^\sigma(\xi)$ has a maximum since $(d\chi^\sigma/d\xi)(1) < 0$. This means the claim by Theorem 5.1(i).

- (ii) *In Theorem 5.1(ii), if $(d^j \chi^\sigma / d^j \xi)(1) = 0$, $j \in [\ell]$, for some $\ell \in \mathbb{N}$, then Eq. (5.5) is replaced by*

$$\chi^\sigma(1) \frac{d^{\ell+1} \chi^\sigma}{d\xi^{\ell+1}}(1) \quad \text{with } \sigma = \sigma^{+-} \text{ and } \sigma^{-+}.$$

- (iii) *From Theorems 4.1 and 5.1 we see that in the reduced system (4.4) (resp. in the KM (4.1)), 2^{2n_0} or $3 \cdot 2^{2(n_0-1)}$ equilibria (resp. families of equilibria) are born, depending on whether ones with $\sigma = \sigma^{++}$ and σ^{-+} in Theorem 5.1(ii) are distinguished or not, and that $2^{2(n_0-1)}$ saddle-node bifurcations occur at least since a supercritical one occurs if Eq. (5.2) or (5.3) holds.*

Moreover, we have the following on the number of pitchfork bifurcations to occur in these systems.

Proposition 5.3. *In the reduced system (4.4) and equivalently in the KM (4.1), the following hold on the number of pitchfork bifurcations:*

- (i) $2^{2n_0-3} + 2^{n_0-2}n_0$ pitchfork bifurcations occur at least;
- (ii) $2^{2(n_0-1)}$ pitchfork bifurcations occur at least if n_0 is prime.

Proof. We begin with the proof of part (i). By Theorem 5.1(ii) and Remark 5.2(ii), we only have to estimate the number of $\sigma \in \Sigma_{n_0}$ such that $\chi^\sigma(1) \neq 0$. Let $4\bar{n}$ denote the number. If for some $\ell \in [n_0 - 1]$ there exists a sequence $\{i_k\}_{k=1}^\ell$ such that $i_k \in [n_0] \setminus \{1\}$, $\sigma_j = \sigma_{2n_0-j+1}$ for $j = i_k$ and $\sigma_j \neq \sigma_{2n_0-j+1}$ if $j \neq i_k$, then

$$\chi^\sigma(1) = \frac{1}{n_0} \left(1 + 2 \sum_{k=1}^{\ell} \sigma_{i_k} \sqrt{1 - \left(\frac{i_k - n_0 - 1}{n_0} \right)^2} \right). \quad (5.7)$$

Obviously, if $\ell = 1$, then $\chi^\sigma(1) > 0$. Assume that $\ell \geq 2$ and let $\sigma^\pm \in \Sigma_{n_0}$ be such sequences satisfying $\sigma_{i_\ell}^\pm = \pm 1$ and $\sigma_{i_j}^+ = \sigma_{i_j}^-$ for $j \in [\ell - 1]$. If $\chi^{\sigma^+}(1) = 0$, then $\chi^{\sigma^-}(1) \neq 0$, and vice versa. The number of how to choose the pair σ^\pm is $2^{\ell-1}$ for each sequence $\{i_j\}_{j=1}^\ell$. Hence,

$$\begin{aligned} \bar{n} &\geq 2^{n_0-1} + 2^{n_0-2} \cdot 2(n_0 - 1) + \sum_{\ell=2}^{n_0-1} \frac{(n_0 - 1)!}{(n_0 - \ell - 1)! \ell!} 2^{\ell-1} 2^{n_0-\ell-1} \\ &= 2^{2n_0-3} + 2^{n_0-2}n_0, \end{aligned}$$

which yields part (i).

We turn to the proof of part (ii). Again, by Theorem 5.1(ii) and Remark 5.2(ii), we only have to show that $\chi^\sigma(1) \neq 0$ for any $\sigma \in \Sigma_{n_0}$ when n_0 is prime. We prove it by contradiction and assume that $\chi^\sigma(1)$ is given by (5.7) and $\chi^\sigma(1) = 0$ for some $\sigma \in \Sigma_{n_0}$ as well as n_0 is prime. For some $i_k \neq 1$ and relatively prime positive integers q, p with $p \geq 2$, we must have

$$\sqrt{1 - \left(\frac{i_k - n_0 - 1}{n_0} \right)^2} = \frac{q}{p} \in (0, 1),$$

i.e.,

$$n_0^2 - (i_k - n_0 - 1)^2 = \frac{q^2 n_0^2}{p^2}, \quad p > q > 0, \quad (5.8)$$

since square roots of non-square integers are linearly independent over \mathbb{Z} (see, e.g., [5]). Moreover, if $(p, q) = (p_{j_k}, q_{j_k})$, $k \in [m]$, satisfy (5.8) for some $m \in [\ell]$, then we find a sequence $\{s_j\}_{j=1}^m$ such that $s_j \in \{0, \pm 1\}$, $k \in [m]$, and

$$\sum_{k=1}^m s_k \frac{p_{j_k}}{q_{j_k}} = -\frac{1}{2}.$$

Since n_0 is prime, it follows from (5.8) that $n_0 = p$. So we have

$$q^2 + (p - i_k + 1)^2 = p^2. \quad (5.9)$$

Hence, $(q, p - i_k + 1, p)$ is a *Pythagorean triple* [33]. We need the following elementary result on Pythagorean triples.

Lemma 5.4. *Let $\ell_j \in \mathbb{N}$, $j = 1, 2, 3$, and ℓ_1, ℓ_3 are relatively prime. The triple (ℓ_1, ℓ_2, ℓ_3) is Pythagorean, i.e., $\ell_1^2 + \ell_2^2 = \ell_3^2$, if and only if there exist relatively prime integers m_1, m_2 with $m_1 > m_2 > 0$ such that m_1, m_2 are of opposite parity, i.e., one is even and the other is odd,*

$$(\ell_1, \ell_2) = (m_1^2 - m_2^2, 2m_1m_2) \text{ or } (2m_1m_2, m_1^2 - m_2^2)$$

and $\ell_3 = m_1^2 + m_2^2$.

See, e.g., Section 13.2 of [25] or Chapter 1 and Appendix B of [33] for a proof of Lemma 5.4. Note that if the triple (ℓ_1, ℓ_2, ℓ_3) is Pythagorean and ℓ_1, ℓ_3 are relatively prime, then so are ℓ_1, ℓ_2, ℓ_3 .

We also need the following elementary result on a sum of two squares (see, e.g., Section 15.1 of [25] for a proof).

Lemma 5.5. *Let $\ell_3 > 2$ be a prime integer. There exist $m_1, m_2 \in \mathbb{N}$ such that $\ell_3 = m_1^2 + m_2^2$ and $m_1 > m_2$ if and only if $\ell_3 \equiv 1 \pmod{4}$. Moreover, the pair (m_1, m_2) is unique.*

We return to the proof of part (ii) in Proposition 5.3. It follows from Lemmas 5.4 and 5.5 that if Eq. (5.9) holds, then there uniquely exists a pair (m_1, m_2) of integers such that $m_1 > m_2 > 0$ and

$$p = m_1^2 + m_2^2, \quad q = 2m_1m_2 \text{ or } m_1^2 - m_2^2.$$

Hence,

$$s_1 \frac{2m_1m_2}{m_1^2 + m_2^2} + s_2 \frac{m_1^2 - m_2^2}{m_1^2 + m_2^2} = -\frac{1}{2}$$

for some $s_j \in \{0, \pm 1\}$, but this never occurs. So we have a contradiction. This completes the proof of Proposition 5.3. \square

We leave our discussion on saddle-node and pitchfork bifurcations in (4.4).

Proposition 5.6. *For any $\sigma \in \Sigma_{n_0}$, the equilibrium v^σ suffers no Hopf bifurcation in (4.4).*

Proof. We first notice that the one-parameter family of equilibria given by (4.11) in (4.1) has the same stability type as the corresponding equilibrium in (4.4). Moreover, the former exhibits Hopf bifurcations if and only if so does the latter. So we analyze the KM (4.1) instead of (4.4) in the following.

We compute each element of the Jacobian matrix A for the vector field of (4.1) as

$$A_{ij} = \begin{cases} -\frac{K}{n} \left(\cos v_i + \sum_{j=1, j \neq i}^{2n_0} \cos(v_j - v_i) \right) & \text{if } i = j; \\ \frac{K}{n} \cos(v_j - v_i) & \text{if } i \neq j \neq n_0 + 1; \\ \frac{K}{n} \cos v_i & \text{if } i \neq j = n_0 + 1 \end{cases} \quad (5.10)$$

for $i \neq n_0 + 1$ and

$$A_{n_0+1, j} = \begin{cases} -\frac{K}{n} \sum_{j=1}^{2n_0} \cos v_j & \text{if } j = n_0 + 1; \\ \frac{K}{n} \cos v_j & \text{if } j \neq n_0 + 1. \end{cases} \quad (5.11)$$

Thus, the matrix A is symmetric and consequently only has real eigenvalues. This implies the desired result. \square

Remark 5.7. *From Remark 4.5 and Proposition 5.6 we see that no other bifurcation than ones detected in Theorem 5.1 occurs for equilibrium in the system (4.1).*




FIGURE 5. Function $\chi^\sigma(\xi)$: (a) $n = 5$; (b) and (c) $n = 11$.

The graph of $\chi^\sigma(\xi)$ is displayed for $n = 5$ and 11 in Fig. 5. It is plotted as a solid line if $\chi^\sigma(\xi)$ has a maximum or minimum, and as a dotted or dashed line otherwise, depending on whether $\sigma_1 \neq \sigma_{2n_0}$ or not. In Fig. 5(a), the black and blue lines represent the graph for $(\sigma_2, \sigma_3) = (1, 1)$ and $(-1, -1)$, respectively, while the red line for $(\sigma_2, \sigma_3) = (1, -1)$ or $(-1, 1)$. In Figs. 5(b) and (c), respectively, the graphs are plotted for $n = 11$ and $\sigma \in \Sigma_{n_0}$ with $\sigma_i = \sigma_{2n_0-i+1}$ for any $i \neq 1$ and with $\sigma_i \neq \sigma_{2n_0-i+1}$ for some $i \neq 1, 2n_0$. See Appendix B for the meaning of the line colors in Figs. 5(b) and (c).

6. STABILITY

For the KM (4.1) and reduced system (4.4) we finally discuss the stability of the equilibria detected by Theorem 4.1 and equivalently by Corollary 4.4. Unlike (3.4), it is not easy in general to determine the stability of the equilibria in (4.4). However, we can prove the following theorem.

Theorem 6.1.

- (i) *The equilibrium v^σ with $\sigma_i = 1$ for all $i \in [2n_0]$ is asymptotically stable if $\xi = n_0 a / n K \hat{C}_B^g < \xi_0$ and unstable if $\xi > \xi_0$, where ξ_0 is the unique maximum of $\chi^\sigma(\xi)$ detected in Theorem 5.1(i).*
- (ii) *The equilibrium v^σ is unstable if $\sigma_i = -1$ for some $i \in [2n_0]$.*

Proof. Fix $\sigma \in \Sigma_{n_0}$ and increase $\xi \in (0, 1)$ along a branch of equilibria, v^σ with $\chi^\sigma(\xi)$. We have the following lemma.

Lemma 6.2. *Suppose that the number of zero eigenvalues of A for v^σ with $\chi^\sigma(\xi)$, $\sigma \in \Sigma_{n_0}$, increases at $\xi = \xi_0$ when the value of $\xi \in (0, 1)$ is increased. Then $\chi^\sigma(\xi)$ has an extremum at $\xi = \xi_0$ and the increasing number is one at most.*

Proof. Let $\sigma \in \Sigma_{n_0}$ and $a > 0$ be fixed. We write $f(v; K)$ for the the vector field of (4.4) and $v^\sigma(\xi) = v^\sigma \in \mathbb{T}^{n_0}$. The Jacobian matrix $Df(v^\sigma(\xi); K)$ is symmetric like that of the KM (1.1), so that its eigenvalues are real and geometrically simple. In particular, if the eigenvalues change their sign, then they must become zero (see the proof of Proposition 5.6).

Assume that $D_v f(v^\sigma(\xi); K)$ has a simple zero eigenvalue when Eq. (4.13) holds and $\xi = \xi_0 \in (0, 1)$. Let $\bar{e} \in \mathbb{R}^{2n_0}$ denote the associated eigenvector. Then the equilibrium $v^\sigma(\xi_0)$ has a one-dimensional center manifold [21, 46] on which the system (4.4) reduces to

$$\dot{v}_c = c_1 v_c^j + \cdots + c_2(K - K_0) + \cdots, \quad v_c \in \mathbb{R}, \quad (6.1)$$

where $K_0 = a/|\chi^\sigma(\xi_0)|$, $c_1, c_2 \in \mathbb{R}$ are constants and $j > 1$ is an even integer, since the eigenvalue does not change its sign if j is odd. If \bar{e} is linearly independent of $(dv/d\xi)(\xi_0)$, then by (6.1) there exists a different family of equilibria from $v^\sigma(\xi)$ near $\xi = \xi_0$, but coincide with it at $\xi = \xi_0$. This contradicts that the equilibrium $v^\sigma(\chi^\sigma(\xi_0))$ is isolated for $\xi \in (0, 1)$. Hence, we can take $\bar{e} = (dv/d\xi)(\xi_0)$, so that by (4.13)

$$\frac{d\chi^\sigma}{d\xi}(\xi_0) = 0$$

since

$$\begin{aligned} & \left. \frac{d}{d\xi} f\left(v^\sigma(\xi); \frac{a}{|\chi^\sigma(\xi)|}\right) \right|_{\xi=\xi_0} \\ &= D_v f(v^\sigma(\xi_0); K_0) \frac{dv}{d\xi}(\xi_0) \mp \frac{\partial f}{\partial K}(v^\sigma(\xi_0); K_0) \frac{a}{|\chi^\sigma(\xi_0)|^2} \frac{d\chi^\sigma}{d\xi}(\xi_0) = 0, \end{aligned}$$

where the upper or lower sign is taken, depending on whether $\chi^\sigma(\xi_0)$ is positive or negative. Similarly, we can show that $D_v f(v^\sigma(\xi_0); \xi_0)$ never has a non-simple zero eigenvalue even when $d\chi^\sigma(\xi_0)/d\xi = 0$, since the equilibrium $v^\sigma(\chi^\sigma(\xi_0))$ is isolated for $\xi \in (0, 1)$. Thus, we obtain the desired result from Remark 5.7. \square

We turn to the proof of Theorem 6.1. As in the proof of Proposition 5.6, we first analyze the KM (4.1) since the one-parameter family of equilibria given by (4.11) in (4.1) has the same stability type as the corresponding equilibrium in (4.4).

Fix $\sigma \in \Sigma_{n_0}$, and let n_+ and n_- , respectively, denote the numbers of $\sigma_i = 1$ and -1 , $i \in [2n_0]$. Take the limit $\xi = n_0 a/nK|\hat{C}_D| \rightarrow 0$. Then $K \rightarrow \infty$ and $\phi_i \rightarrow 0$, so that by (4.9)

$$v_i^\sigma \rightarrow \begin{cases} 0 & \text{if } \sigma_i = 1; \\ \pi & \text{if } \sigma_i = -1 \text{ and } i > n_0 + 1. \end{cases}$$

Hence,

$$\cos v_i \rightarrow \begin{cases} 1 & \text{if } \sigma_i = 1; \\ -1 & \text{if } \sigma_i = -1, \end{cases} \quad \cos(v_j - v_i) \rightarrow \begin{cases} 1 & \text{if } \sigma_i = \sigma_j; \\ -1 & \text{if } \sigma_i \neq \sigma_j \end{cases}$$

eigenvalue as $\xi \rightarrow 0$. Since $d\chi^\sigma(\xi)/d\xi = 0$ only at $\xi = \xi_0$, as shown in the proof of Theorem 5.1(i), we obtain part (i) by Lemma 6.2. Note that at $\xi = \xi_0$ the matrix A keeps the zero eigenvalue and one of the negative eigenvalues becomes positive.

We turn to the proof of part (ii). Let $\sigma \in \Sigma_{n_0}$ and $a > 0$ be fixed. As in the proof of Lemma 6.2, we write $f(v; K)$ for the vector field of (4.4) and $v^\sigma(\xi) = v^\sigma \in \mathbb{T}^{n_0}$. We compute

$$\frac{dv_i^\sigma}{d\xi}(\xi) = \begin{cases} \pm\sigma_i \left(\frac{i-n_0-1}{n_0} \right) / \sqrt{1 - \left(\frac{i-n_0-1}{n_0} \xi \right)^2} & \text{for } i \leq n_0; \\ \pm\sigma_i \left(\frac{i-n_0}{n_0} \right) / \sqrt{1 - \left(\frac{i-n_0}{n_0} \xi \right)^2} & \text{for } n_0 < i \leq 2n_0 \end{cases}$$

and

$$\frac{\partial f_i}{\partial K}(v^\sigma(\xi); K) = \begin{cases} -(i-n_0-1)\nu/K & \text{for } i \leq n_0; \\ -(i-n_0)\nu/K & \text{for } n_0 < i \leq 2n_0 \end{cases}$$

for $i \in [2n_0]$ when $K = a/|\chi^\sigma(\xi)|$, where $v_i^\sigma(\xi)$ and $f_i(v; K)$ are the i th elements of $v^\sigma(\xi)$ and $f(v; K)$, respectively, and the upper or lower sign is taken, depending on whether $\chi^\sigma(\xi)$ is positive or not. Here we have used the relations $f_i(v^\sigma(\xi); K) = 0$, $i \in [n]$.

Suppose that $(d\chi^\sigma/d\xi)(\xi)$ has a zero at $\xi = \xi_0$ and let $K_0 = a/|\chi^\sigma(\xi_0)|$ be bounded. Then the Jacobian matrix $D_v f(v^\sigma(\xi_0); K_0)$ has a simple zero eigenvalue and $(dv/d\xi)(\xi_0)$ is the associated eigenvector. The equilibrium $v^\sigma(\xi_0)$ has a one-dimensional center manifold on which the system (4.4) reduces to (6.1). See the proof of Lemma 6.2. In addition, c_1/c_2 is negative or positive, whether $|\xi^\sigma(\xi)|$ has a local maximum or minimum at $\xi = \xi_0$, since by Theorem 5.1(i) a supercritical or subcritical saddle-node bifurcation occurs. Moreover, substituting $v = (dv^\sigma/d\xi)(\xi_0)v_c + v^\sigma(\xi_0)$ into (4.4) and taking the inner product of the resulting equation with $(dv/d\xi)(\xi_0)$, we see that

$$c_2 = \frac{dv_i^\sigma}{d\xi}(\xi_0) \cdot \frac{\partial f_i}{\partial K}(v^\sigma(\xi_0); K_0) / \left| \frac{dv_i^\sigma}{d\xi}(\xi_0) \right|^2$$

is positive or negative, depending on whether $h^\sigma(\xi_0)\chi^\sigma(\xi_0)$ is negative or positive, where the dot “ \cdot ” represents the standard inner product and

$$h^\sigma(\xi) = \frac{1}{n_0} \left(\sum_{i=1}^{n_0} \sigma_i \left(\frac{i-n_0-1}{n_0} \right)^2 / \sqrt{1 - \left(\frac{i-n_0-1}{n_0} \xi \right)^2} + \sum_{i=n_0+1}^{2n_0} \sigma_i \left(\frac{i-n_0}{n_0} \right)^2 / \sqrt{1 - \left(\frac{i-n_0}{n_0} \xi \right)^2} \right).$$

We can easily determine the sign of $h^\sigma(\xi)$ as follows.

Lemma 6.4. $h^\sigma(\xi_0)\chi^\sigma(\xi_0) > 0$.

Proof. We easily see that

$$\begin{aligned} \frac{d\chi^\sigma}{d\xi}(\xi) &= \frac{1}{n_0} \left(1 + \sum_{i=1}^{n_0} \sigma_i \left(1 - 2 \left(\frac{i-n_0-1}{n_0} \xi \right)^2 \right) \right) / \sqrt{1 - \left(\frac{i-n_0-1}{n_0} \xi \right)^2} \\ &\quad + \sum_{i=n_0+1}^{2n_0} \sigma_i \left(1 - 2 \left(\frac{i-n_0}{n_0} \xi \right)^2 \right) / \sqrt{1 - \left(\frac{i-n_0}{n_0} \xi \right)^2} \\ &= \frac{\chi^\sigma(\xi)}{\xi} - h^\sigma(\xi) \xi^2 \end{aligned}$$

for $\xi \in (0, 1)$, so that

$$h^\sigma(\xi_0) = \frac{\chi^\sigma(\xi_0)}{\xi_0^3}.$$

Hence, If $\chi^\sigma(\xi_0)$ is positive (resp. negative), then so is $h^\sigma(\xi_0)$. This yields the desired result. \square

We also need the following lemma.

Lemma 6.5. *Suppose that $\chi^\sigma(\xi) = 0$ for some $\sigma \in \Sigma_{n_0}$ and $\xi \in (0, 1)$. Then $v_i^\sigma(\xi) = v_i^\sigma(0)$, $i \in [2n_0]$, for some $\bar{\sigma} \in \Sigma_{n_0}$.*

Proof. Let u_i , $i \in [n]$, represent an equilibrium given by (4.11) with $v = v^\sigma(\xi)$ in the KM (4.1). Since $a/K = |\chi^\sigma(\xi)| = 0$, we have

$$\sum_{j=1}^n \sin(u_j - u_i) = \sum_{j=1}^{2n_0} \sin(v_j^\sigma(\xi) - v_i^\sigma(\xi)) = -\sin v_i^\sigma(\xi) \sum_{j=1}^{2n_0} \cos v_j^\sigma(\xi) = 0$$

by Lemma 4.2. Hence,

$$\sin v_i^\sigma(\xi) = 0, \quad i \in [2n_0],$$

which implies the desired result. \square

From the expression (4.12) we see that $\chi^\sigma(\xi) > 0$ on $(0, 1]$ if $\sigma_i = 1$ except for only one $i \in [2n_0]$, i.e., $\ell = n - 1$ and $\sigma_{i_k} = 1$ for any $k \in [\ell]$ in (4.12), as in part (i). Hence, it follows from Lemma 6.5 that $D_v f(v^\sigma(\xi); K)$ with $K = a/\chi^\sigma(\xi)$ has two positive eigenvalues at least as $\chi^\sigma(\xi) \rightarrow 0$ but $\xi \not\rightarrow 0$ if $\sigma \in \Sigma_{n_0}$ satisfies the hypothesis of Lemma 6.5.

Now we are in position to complete the proof of Theorem 6.1(ii). From (6.1) and Lemma 6.4 we see that a negative (resp. positive) eigenvalue of the Jacobian matrix $D_v f(v^\sigma(\xi); K)$ with $K = a/|\chi^\sigma(\xi)|$ becomes positive (resp. negative) when $|\chi^\sigma(\xi)|$ has a local maximum (resp. minimum) with $\chi^\sigma(\xi) > 0$ at $\xi = \xi_0$. Using the above observation on the number of its positive eigenvalues as $\chi^\sigma(\xi) \rightarrow 0$, we see via Lemma 6.3 that $D_v f(v^\sigma(\xi); K)$ with $K = a/\chi^\sigma(\xi)$ has one positive eigenvalue at least for any $\xi \in (0, 1]$ if $\sigma_i = -1$ for some $i \in [2n_0]$. This yields the desired result. \square

Remark 6.6. *In Section 3 of [37], the statement of Theorem 6.1(ii) was given in more general setting although their proof was not complete.*

Numerically computed bifurcation diagrams of equilibria in (4.4) for $n = 5$ and 11 are displayed in Figs. ?? and 6, respectively. Here the computer tool AUTO [16] was used, again. The solid and dashed lines represent stable and unstable equilibria, respectively. The red and black lines, respectively, correspond to the equilibrium




FIGURE 6. Bifurcation diagram of equilibria in (4.4) with $n = 5$: (a) v_3 -component; (b) v_4 -component.




FIGURE 7. Numerically computed bifurcation diagram of equilibria in (4.4) with $n = 11$: (a) v_9 -component; (b) v_{10} -component.

given by (4.5) and the others. They are plotted for all equilibria in Fig. 6, but only for $(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, 1, 1, 1)$,

$$(\sigma_6, \sigma_7, \sigma_8, \sigma_9) = (1, 1, 1, 1), (1, 1, 1, -1) \text{ or } (1, 1, -1, 1)$$

and $\sigma_1, \sigma_{10} = 1$ or -1 in Fig. 7. We observe that only the equilibrium with $\sigma_i = 1$, $i \in [2n_0]$, i.e., given by (4.5), can be stable, as in the case of $n = 3$.

7. CONTINUUM LIMITS

Finally we consider the CL (1.2) with the frequency function (1.9) and discuss the implications of the above results to it.

The synchronized solution (1.4) becomes

$$u(t, x) = U(x) + \theta, \quad U(x) = \arcsin\left(\frac{a(x - \frac{1}{2})}{KC}\right), \quad (7.1)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant and the constant C satisfies (1.10). From Section 3 of [26], we immediately obtain the following.

Theorem 7.1. *Equation (7.1) gives a unique family of continuous stationary solutions to the CL (1.2) with the natural frequency (1.9).*

Theorem 7.1 does not deny the existence of other noncontinuous stationary solutions to (1.2). Indeed, Eq. (1.2) has infinitely many one-parameter families of noncontinuous stationary solutions of the form

$$u(t, x) = \begin{cases} U(x) + \theta & \text{for } x \in [0, 1] \setminus \hat{I}; \\ \pi - U(x) + \theta & \text{for } x \in \hat{I}_j^-, j = [m_-]; \\ -U(x) - \pi + \theta & \text{for } x \in \hat{I}_j^+, j = [m_+], \end{cases} \quad (7.2)$$

where $\theta \in \mathbb{S}^1$ is an arbitrary constant, m_{\pm} are nonnegative integers and may be infinite, $\hat{I}_j^{\pm} \subset [0, 1]$, $j \in [m_{\pm}]$, are intervals such that $\hat{I}_j^- \in [0, \frac{1}{2}]$, $\hat{I}_j^+ \in [\frac{1}{2}, 1]$ and $\hat{I}_j^- \cap \hat{I}_k^-, \hat{I}_j^+ \cap \hat{I}_k^+ = \emptyset$,

$$\hat{I} = \bigcup_{j=1}^{m_-} \hat{I}_j^- \cup \bigcup_{j=1}^{m_+} \hat{I}_j^+,$$

and the constant C in $U(x)$ satisfies

$$C = \int_{[0,1] \setminus \hat{I}} \sqrt{1 - \left(\frac{\omega(x) - \Omega}{KC} \right)^2} dx - \int_{\hat{I}} \sqrt{1 - \left(\frac{\omega(x) - \Omega}{KC} \right)^2} dx.$$

Here the interiors of \hat{I}_j^{\pm} , $j \in [m_{\pm}]$, may be empty. We see that the one-parameter family of synchronized solutions given by (4.11) converge to (7.2) when

$$\sigma_i = \begin{cases} 1 & \text{if } i/n \in [0, 1] \setminus \hat{I}; \\ -1 & \text{if } i/n \in \hat{I}. \end{cases} \quad (7.3)$$

Using Theorems 2.4 and 2.7, we prove the following

Theorem 7.2.

- (i) *The family of stationary solutions given by (7.1) is asymptotically stable.*
- (ii) *The family of stationary solutions given by (7.2) is unstable if it is different from (7.1) in the sense of $L^2(I)$.*

Proof. We easily see that the family of synchronized solutions (4.11) for $v = v^{\sigma}$ with $\sigma_i = 1$, $i \in [2n_0]$, which is asymptotically stable for $\xi < \xi_0$ by Theorem 6.1(i), in the KM (1.1) converges to the family of stationary solutions (7.1) as $n \rightarrow \infty$. Recall that at $\xi = \xi_0$ $\chi^{\sigma}(\xi)$ has a unique extremum (maximum). Moreover, since

$$\chi^{\sigma}(\xi) \rightarrow \frac{n}{n_0} \xi \int_0^1 \sqrt{1 - \xi^2 x^2} dx = \frac{n}{2n_0} (\xi \sqrt{1 - \xi^2} + \arcsin \xi) = \frac{n}{2n_0} \varphi(\xi),$$

we have $\xi_0 \rightarrow 1$ as $n \rightarrow \infty$. Using Theorem 2.4(i), we obtain part (i).

We turn to the proof of part (ii). The family of synchronized solutions (4.11) for $v = v^{\sigma}$ with (7.3), which is unstable by Theorem 6.1(ii), converges to the family of stationary solutions (7.2) in $L^2(I)$ as $n \rightarrow \infty$. Hence, by Theorem 2.7, if it is different from (7.1) in the sense of $L^2(I)$, then the family (7.2) is unstable. \square

From Theorems 5.1 and 6.1 we obtain the following result for the KM (1.1). The family of synchronized solutions (4.11) for v^{σ} with $\sigma_i = 1$, $i \in [2n_0]$, suffers a saddle-node bifurcation at $\xi = \xi_0$, i.e., $K = n_0 a / n \chi^{\sigma}(\xi_0)$ (see Eq. (4.13)), and it turns unstable from stable. In addition, it suffers a pitchfork bifurcation at

$\xi = 1$, i.e., $K = n_0 a / n \chi^\sigma(1)$, and it changes to the family with $\sigma_1 = \sigma_{2n_0} = -1$ and $\sigma_j = 1$, $i \in [2n_0] \setminus \{1, n_0\}$, while two families with $\sigma_1 = -1$, $\sigma_{2n_0} = 1$ or $\sigma_1 = 1$, $\sigma_{2n_0} = -1$ and $\sigma_j = 1$, $i \in [2n_0] \setminus \{1, n_0\}$, are born. When $n \rightarrow \infty$, the saddle-node and pitchfork bifurcations collide, since $\xi_0 \rightarrow 1$. Moreover, these four families converge to four families of stationary solutions that are the same in the sense of $L^2(I)$ and asymptotically stable as a family of $L^2(I)$ solutions by Theorem 7.2. Thus, bifurcation behavior in the CL (1.2) is very subtle, compared with finite-dimensional dynamical systems such as the KM (1.1).

ACKNOWLEDGEMENTS

This work was partially supported by the JSPS KAKENHI Grant Number 23K22409. The author thanks for the anonymous referees for their helpful comments and constructive suggestions, which have improved this work.

APPENDIX A. STABILITY OF THE EQUILIBRIA IN (3.4)

Let

$$z = \sqrt{1 - \left(\frac{a}{K}\right)^2} \in (0, 1).$$

For (3.6), we have

$$-\cos v_1 - \cos v_2 - \cos(v_2 - v_1) = 1 > 0$$

and

$$\cos v_1 \cos v_2 + (\cos v_1 + \cos v_2) \cos(v_2 - v_1) = -z^2 < 0.$$

By (3.8) the equilibria given by (3.6) are unstable.

Let

$$z = \sqrt{1 - \left(\frac{a}{3KC_D}\right)^2} \in (0, 1).$$

For (3.9), we have

$$-\cos v_1 - \cos v_2 - \cos(v_2 - v_1) = -2z^2 - 2z + 1 = -2\left(z + \frac{1}{2}\right)^2 + \frac{3}{2}$$

which is negative when $z > (\sqrt{3} - 1)/2$, and

$$\cos v_1 \cos v_2 + (\cos v_1 + \cos v_2) \cos(v_2 - v_1) = 4z^3 + z^2 - 2z,$$

which is positive or negative, depending on whether z is greater or less than $(\sqrt{33} - 1)/8 > (\sqrt{3} - 1)/2$. By (3.7) and (3.8), the equilibria given by (3.9) are asymptotically stable (resp. unstable) if $v_2 = -v_1$ is less (resp. greater) than v_0 .

Let

$$z = \sqrt{1 - \left(\frac{a}{3K\hat{C}_D}\right)^2} \in (0, 1).$$

For (3.10), we have

$$-\cos v_1 - \cos v_2 - \cos(v_2 - v_1) = -2z^2 + 2z + 1 = -2\left(z - \frac{1}{2}\right)^2 + \frac{3}{2} > 0.$$

By (3.8), the equilibria given by (3.10) are unstable.

APPENDIX B. LINE COLORS IN FIGS. 5(B) AND (C)

In this appendix, we provide the meaning of line colors in Figs. 5(b) and (c), where $n = 11$ and $n_0 = 5$. For each line color, σ is specified for the graphs of the equilibria below in the same order as the corresponding graphs in Fig. 5(b) or (c).

B.1. Fig. 5(b). For $\sigma \in \Sigma_5$ in Fig. 5(b), we have $\sigma_{11-i} = \sigma_i$ for any $i \neq 1$ and $(\sigma_1, \sigma_{10}) = (1, 1), (-1, -1), (-1, 1)$ or $(1, -1)$, depending on whether the line is solid, dashed or dotted. So we only give the values of σ_i for $i \in [5] \setminus \{1\}$ below.

- Black lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, 1, 1, 1), (-1, -1, 1, 1), (-1, -1, -1, -1).$$

- Red lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (-1, 1, 1, 1), (-1, 1, -1, 1), (1, -1, -1, -1).$$

- Blue lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, -1, 1, 1), (-1, 1, 1, -1), (-1, 1, -1, -1).$$

- Purple lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, 1, -1, 1), (1, -1, -1, 1), (-1, -1, 1, -1).$$

- Green lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, 1, 1, -1), (1, -1, 1, -1), (-1, -1, -1, 1).$$

- Orange lines:

$$(\sigma_2, \sigma_3, \sigma_4, \sigma_5) = (1, 1, -1, -1).$$

B.2. Fig. 5(c). For $\sigma \in \Sigma_5$ in Fig. 5(c), we have $\sigma_i \neq \sigma_{11-i}$ for some $i \neq 1$ and $(\sigma_1, \sigma_{10}) = (1, 1), (-1, -1), (-1, 1)$ or $(1, -1)$, depending on whether the line is solid, dashed or dotted. We only give the values of σ_i for $i \in [5] \setminus \{1\}$ below, where $\sigma_i, \sigma_{11-i} = 1$ or -1 and $\sigma_i \neq \sigma_{11-i}$ if $\sigma_i = \pm 1$ and $\sigma_i = \sigma_{11-i}$ otherwise.

- Black lines:

$$\begin{aligned} (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (\pm 1, 1, 1, 1), (\pm 1, \pm 1, 1, 1), (-1, \pm 1, 1, 1), \\ & (\pm 1, 1, -1, 1), (1, \pm 1, \pm 1, \pm 1), (-1, 1, \pm 1, \pm 1), \\ & (-1, \pm 1, \pm 1, \pm 1), (1, -1, \pm 1, \pm 1), (\pm 1, -1, 1, -1), \\ & (1, \pm 1, -1, -1), (\pm 1, \pm 1, -1, -1), (\pm 1, -1, -1, -1). \end{aligned}$$

- Red lines:

$$\begin{aligned} (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (1, \pm 1, 1, 1), (\pm 1, 1, \pm 1, 1), (-1, 1, \pm 1, 1), \\ & (\pm 1, 1, \pm 1, \pm 1), (1, \pm 1, 1, -1), (\pm 1, -1, \pm 1, 1), \\ & (-1, \pm 1, -1, 1), (\pm 1, 1, \pm 1, -1), (\pm 1, -1, \pm 1, \pm 1), \\ & (1, -1, \pm 1, -1), (\pm 1, -1, \pm 1, -1), (-1, \pm 1, -1, -1). \end{aligned}$$

- Blue lines:

$$\begin{aligned} (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (1, 1, \pm 1, 1), (\pm 1, 1, 1, \pm 1), (-1, 1, 1, \pm 1), \\ & (1, -1, \pm 1, 1), (1, 1, -1, \pm 1), (\pm 1, -1, 1, \pm 1), \\ & (\pm 1, 1, -1, \pm 1), (-1, -1, 1, \pm 1), (-1, 1, \pm 1, -1), \\ & (1, -1, -1, \pm 1), (\pm 1, -1, -1, \pm 1), (-1, -1, \pm 1, -1). \end{aligned}$$

- Purple lines:

$$\begin{aligned}
 (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (1, 1, 1, \pm 1), (1, \pm 1, \pm 1, 1), (\pm 1, -1, 1, 1), \\
 & (\pm 1, 1, 1, -1), (1, 1, \pm 1, -1)(\pm 1, \pm 1, -1, 1), \\
 & (\pm 1, \pm 1, 1, -1), (-1, -1, \pm 1, 1), (\pm 1, -1, -1, 1), \\
 & (\pm 1, 1, -1, -1), (-1, \pm 1, \pm 1, -1), (-1, -1, -1, \pm 1).
 \end{aligned}$$

- Green lines:

$$\begin{aligned}
 (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (1, \pm 1, 1, \pm 1), (\pm 1, \pm 1, \pm 1, 1), (1, -1, 1, \pm 1), \\
 & (-1, \pm 1, \pm 1, 1), (1, \pm 1, \pm 1, -1), (-1, 1, -1, \pm 1), \\
 & (\pm 1, \pm 1, \pm 1, -1), (-1, \pm 1, -1, \pm 1).
 \end{aligned}$$

- Orange lines:

$$\begin{aligned}
 (\sigma_2, \sigma_3, \sigma_4, \sigma_5) = & (1, 1, \pm 1, \pm 1), (\pm 1, \pm 1, 1, \pm 1), (1, \pm 1, -1, \pm 1), \\
 & (-1, \pm 1, 1, \pm 1), (1, \pm 1, -1, \pm 1), (-1, \pm 1, \pm 1), \\
 & (\pm 1, \pm 1, -1), (-1, -1, \pm 1, \pm 1).
 \end{aligned}$$

REFERENCES

- [1] D.M. Abrams and S.H. Strogatz, Chimera states in a ring of nonlocally coupled oscillators, *Internat. J. Bifur. Chaos*, **16** (2006), 21–37.
- [2] J.A. Acebrón, L.L. Bonilla, C.J.P. Vicente, F. Ritort and R. Spigler, The Kuramoto model: A simple paradigm for synchronization phenomena, *Rev. Mod. Phys.*, **77** (2005), 137–185.
- [3] D. Aeyels and J.A. Rogge, Existence of partial entrainment and stability of phase locking behavior of coupled oscillators *Progr. Theoret. Phys.*, **112** (2004), 921–942.
- [4] A. Arenas, A. Diaz-Guilera, J. Kurths, Y. Moreno and C. Zhou, Synchronization in complex networks, *Phys. Rep.*, **469**(2008), 93–153.
- [5] A.S. Besicovitch, On the linear independence of fractional powers of integers, *J. London Math. Soc.*, **15** (1940), 3–6.
- [6] J.C. Bronski and L. Wang, Partially phase-locked solutions to the Kuramoto model, *J. Stat. Phys.*, **183** (2021), 46.
- [7] J. Bronski, T. Carty and L. Deville, Synchronisation conditions in the Kuramoto model and their relationship to seminorms, *Nonlinearity*, **34** (2021), 5399–5433.
- [8] H. Chiba, A proof of the Kuramoto conjecture for a bifurcation structure of the infinite-dimensional Kuramoto model, *Ergodic Theory Dynam. Systems*, **35** (2015), 762–834.
- [9] H. Chiba and G.S. Medvedev, The mean field analysis of the Kuramoto model on graphs I: The mean field equation and transition point formulas, *Discrete Contin. Dyn. Syst.*, **39** (2019a), 131–155.
- [10] H. Chiba and G.S. Medvedev, The mean field analysis of the Kuramoto model on graphs II: Asymptotic stability of the incoherent state, center manifold reduction, and bifurcations, *Discrete Contin. Dyn. Syst.*, **39** (2019), 3897–3921.
- [11] H. Chiba and I. Nishikawa, Center manifold reduction for large populations of globally coupled phase oscillators, *Chaos*, **21** (2011), 043103.
- [12] H. Chiba and D. Pazó, Stability of an $[N/2]$ -dimensional invariant torus in the Kuramoto model at small coupling, *Phys. D*, **238** (2009), 1068–1081.
- [13] Y.-P. Choi, S.-Y. Ha, S. Jung and Y. Kim, Asymptotic formation and orbital stability of phase-locked states for the Kuramoto model, *Phys. D*, **241** (2012), 735–754.
- [14] E.A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
- [15] H. Dietert and B. Fernandez, The mathematics of asymptotic stability in the Kuramoto model, *Proc. A.*, **474** (2018), 20180467.
- [16] E. Doedel and B.E. Oldeman, *AUTO-07P: Continuation and Bifurcation Software for Ordinary Differential Equations*, 2012, available online from <http://cmvl.cs.concordia.ca/auto>.

- [17] J.-G. Dong and X. Xue, Synchronization analysis of Kuramoto oscillators, *Commun. Math. Sci.*, **11** (2013), 465–480.
- [18] F. Dörfler and F. Bullo, On the critical coupling for Kuramoto oscillators, *SIAM J. Appl. Dyn. Syst.*, **10** (2011), 1070–1099.
- [19] F. Dörfler and F. Bullo, Synchronization in complex networks of phase oscillators: A survey, *Automatica*, **50** (2014), 1539–1564.
- [20] G.B. Ermentrout, Synchronization in a pool of mutually coupled oscillators with random frequencies *J. Math. Biol.*, **23** (1985), 55–74.
- [21] J. Guckenheimer and P. Holmes, *Nonlinear Oscillations, Dynamical Systems, and Bifurcations of Vector Fields*, Springer, New York, 1983.
- [22] T. Girnyk, M. Hasler and Y. Maistrenko, Multistability of twisted states in non-locally coupled Kuramoto-type models, *Chaos*, **22** (2012), 013114.
- [23] S.-Y. Ha, H.K. Kim and J. Park, Remarks on the complete synchronization of Kuramoto oscillators, *Nonlinearity*, **28** (2015), 1441–1462.
- [24] S.-Y. Ha and S.-Y. Ryoo, Asymptotic phase-locking dynamics and critical coupling strength for the Kuramoto model *Comm. Math. Phys.*, **377** (2020), 811–857.
- [25] G.H. Hardy and E.M. Wright, *An Introduction to the Theory of Numbers*, 5th ed., Oxford University Press, Oxford, 1979.
- [26] R. Ihara and K. Yagasaki, Continuum limits of coupled oscillator networks depending on multiple sparse graphs, *J. Nonlinear Sci.*, **33** (2023), 62.
- [27] D. Kaliuzhnyi-Verbovetskyi and G. S. Medvedev, The semilinear heat equation on sparse random graphs, *SIAM J. Math. Anal.*, **49** (2017), no. 2, 1333–1355.
- [28] Y. Kuramoto, Self-entrainment of a population of coupled non-linear oscillators, in *International Symposium on Mathematical Problems in Theoretical Physics*, H. Araki (ed.), Springer, Berlin, 1975, pp. 420–422.
- [29] Y. Kuramoto, *Chemical Oscillations, Waves, and Turbulence*, Springer, Berlin, 1984.
- [30] Y. Maistrenko, O.V. Popovych, O. Burylko and P.A. Tass, Mechanism of desynchronization in the finite-dimensional Kuramoto model, *Phys. Rev. Lett.*, **93** (2004), 084102.
- [31] Y. Maistrenko, O.V. Popovych and P.A. Tass, Desynchronization and chaos in the Kuramoto model, in *Dynamics of Coupled Map Lattices and of Related Spatially Extended Systems*, J.-R. Chazottes and B. Fernandez (eds.), Lecture Notes in Physics Vol. 671, 2005, pp. 285–306.
- [32] Y. Maistrenko, O.V. Popovych and P.A. Tass, Chaotic attractor in the Kuramoto model, *Internat. J. Bifur. Chaos*, **15** (2005), 3457–3466.
- [33] E. Maor, *The Pythagorean Theorem: A 4,000-Year History*, Princeton University Press, Princeton, 2007.
- [34] G.S. Medvedev, The nonlinear heat equation on dense graphs and graph limits, *SIAM J. Math. Anal.*, **46** (2014), 2743–2766.
- [35] G.S. Medvedev, The nonlinear heat equation on W -random graphs, *Arch. Ration. Mech. Anal.*, **212** (2014), 781–803.
- [36] G.S. Medvedev, The continuum limit of the Kuramoto model on sparse random graphs, *Comm. Math. Sci.*, **17** (2019), no. 4, 883–898.
- [37] R.E. Mirollo and S.H. Strogatz, The spectrum of the locked state for the Kuramoto model of coupled oscillators, *Phys. D*, **205** (2005), 249–266.
- [38] B. Ottino-Löffler and S.H. Strogatz, Kuramoto model with uniformly spaced frequencies: Finite- N asymptotics of the locking threshold, *Phys. Rev. E*, **93** (2016), 062220.
- [39] D. Pazó, Thermodynamic limit of the first-order phase transition in the Kuramoto model, *Phys. Rev. E*, **72** (2005), 046211.
- [40] A. Pikovsky and M. Rosenblum, Dynamics of globally coupled oscillators: Progress and perspectives, *Chaos*, **25** (2015), 097616.
- [41] A. Pikovsky, M. Rosenblum, and J. Kurths, *Synchronization: A Universal Concept in Non-linear Sciences*, Cambridge University Press, Cambridge, 2001.
- [42] F.A. Rodrigues, T.K.D.M. Peron, P. Ji and J. Kurths, The Kuramoto model in complex networks, *Phys. Rep.*, **610** (2016), 1–98.
- [43] S.H. Strogatz, From Kuramoto to Crawford: Exploring the onset of synchronization in populations of coupled oscillators, *Phys. D*, **143** (2000), 1–20.
- [44] J.L. van Hemmen and W.F. Wreszinski, Lyapunov function for the Kuramoto model of nonlinearly coupled oscillators, *J. Stat. Phys.*, **72** (1993), 145–166.

- [45] M. Verwoerd and O. Mason, Global phase-locking in finite populations of phase-coupled oscillators, *SIAM J. Appl. Dyn. Syst.*, **7** (2008), 134–160.
- [46] S. Wiggins, *Introduction to Applied Nonlinear Dynamical Systems and Chaos*, Springer, New York, 2003.
- [47] D.A. Wiley, S.H. Strogatz and M. Girvan, The size of the sync basin, *Chaos*, **16** (2006), 015103.

DEPARTMENT OF APPLIED MATHEMATICS AND PHYSICS, GRADUATE SCHOOL OF INFORMATICS,
KYOTO UNIVERSITY, YOSHIDA-HONMACHI, SAKYO-KU, KYOTO 606-8501, JAPAN

Email address: `yagasaki@amp.i.kyoto-u.ac.jp`