

On Counting H -Intersecting Families and Graph Homomorphisms

Igal Sason

Abstract. This work leverages Shearer’s inequalities to derive a new upper bound on the maximum cardinality of a family of graphs on a fixed number of vertices, in which every pair of graphs shares a fixed common subgraph. The derived bound is expressed in terms of the chromatic number of the shared subgraph. Additionally, Shearer’s inequalities, in conjunction with properties of the Shannon entropy, are employed to establish bounds related to the enumeration of graph homomorphisms, providing further insights into the interplay between combinatorial structures and information-theoretic principles.

Keywords. Entropy, counting problems, intersecting families of graphs, graph homomorphisms, Shearer’s lemma.

2020 Mathematics Subject Classification (MSC). 05C30, 05C60, 94A15.

I. INTRODUCTION

Families of intersecting graphs are collections of simple graphs (i.e., graphs with no self loops or multiple edges between vertices), where every two graphs of that family intersect in a specified way, such as sharing at least one edge or another common subgraph. These families play a central role in extremal combinatorics and graph theory, and determining their maximum possible size has been a longstanding challenge. A pivotal conjecture, posed by Simonovits and Sós, concerned the maximum size of triangle-intersecting graph families - those in which every pair of graphs shares a triangle. Their foundational work, initially presented in [34], along with other results on intersection theorems for families of graphs where the shared subgraphs are cycles or paths, was surveyed by these authors in [35]. The first significant progress on this conjecture was made by Chung *et al.* [6], who utilized Shearer’s inequality to establish a non-trivial bound on the largest possible cardinality of a family of triangle-intersecting graphs with a fixed number of vertices. That was ”midway” between the trivial bound and the conjectured bound (or more formally, their geometric mean). The conjecture was ultimately resolved by Ellis,

Igal Sason is with the Viterbi Faculty of Electrical and Computer Engineering, and also with the Department of Mathematics (secondary appointment), Technion - Israel Institute of Technology, Haifa 3200003, Israel.

Correspondence: Igal Sason. Email: eeigal@technion.ac.il; Tel: +97248294699.

Filmus, and Friedgut [8], who proved that the largest triangle-intersecting family comprises all graphs containing a fixed triangle. Building on the spectral approach in [8] (see also Section 4 in [9]), a recent work by Berger and Zhao [3] extended the investigation to K_4 -intersecting graph families, addressing analogous questions for graph families where every pair of graphs intersects in a complete subgraph of size four. Additionally, Keller and Lifshitz [19] provided high-probability results for constructing, for every graph H , families of large random graphs with a common vertex set such that every pair of graphs contains a subgraph isomorphic to H . These are referred to as families of H -intersecting graphs. These contributions highlight the rich interplay of combinatorial, probabilistic, and algebraic methods in advancing the understanding of intersecting graph families.

The interplay between Shannon entropy and extremal combinatorics has significantly enhanced our understanding of the structural and quantitative properties of combinatorial objects through information-theoretic methods. Entropy serves as a versatile and powerful tool to derive concise, often elegant proofs of classical results in extremal combinatorics (see, e.g., Chapter 37 of [1], [12], Chapter 22 of [16], [25] [26], and [28]). Notable examples include Radhakrishnan’s entropy-based proof of Bregman’s theorem on permanents of matrices [27] and the application of Shearer’s lemma to upper-bound the size of the largest triangle-intersecting families of graphs with a fixed number of vertices [6]. Beyond this specific context, Shearer’s inequalities have found extensive applications across diverse areas, including finite geometry, graph theory, the analysis of Boolean functions, and large deviations ([6], [10], [13], [17], [18], [23], [28], [29], [30]). The reader is referred to a recent talk by the author on Shearer’s inequalities and their various combinatorial applications [33]. The first part of this paper builds on the entropy-based proof technique introduced in [6], extending the combinatorial version of Shearer’s inequalities to derive a new upper bound on the cardinality of families of H -intersecting graphs with a fixed number of vertices.

Graph homomorphisms serve as a versatile framework to understand graph mappings, which facilitate studies of structural properties, colorings, and symmetries. Their applications span

various fields, including statistical physics, where they model spin systems [5], and computational complexity, where they underpin constraint satisfaction problems [15]. Recent research has yielded profound insights into counting graph homomorphisms, a problem with deep theoretical and practical relevance ([4], [11], [12], [22], [37], [38]). The second part of this paper relies on Shearer's inequalities and properties of the Shannon entropy to obtain two bounds on the number of graph homomorphisms.

The paper is structured as follows: Section II presents essential preliminary material for this paper, including three variations of Shearer's inequalities. In Section III, the combinatorial version of Shearer's lemma is employed for upper bounding the size of H -intersecting families of graphs. Section IV focuses on entropy-based proofs, also incorporating a probabilistic version of Shearer's lemma, to obtain bounds on the number of graph homomorphisms.

II. PRELIMINARIES

A. Shearer's Inequalities

The following subsection introduces three versions of Shearer's inequalities that are useful in the analysis presented in this paper. The first version serves as a foundation for proving the other two, which are directly applied in this work. Familiarity with the Shannon entropy and its basic properties is assumed, following standard notation (see, e.g., Chapter 3 of [7]).

Proposition 1 (Shearer's Lemma). Let

- $n, m, k \in \mathbb{N}$,
- X_1, \dots, X_n be discrete random variables,
- $[n] \triangleq \{1, \dots, n\}$,
- $\mathcal{S}_1, \dots, \mathcal{S}_m \subseteq [n]$ be subsets such that each element $i \in [n]$ belongs to at least $k \geq 1$ of these subsets.
- $X^n \triangleq (X_1, \dots, X_n)$, and $X_{\mathcal{S}_j} \triangleq (X_i)_{i \in \mathcal{S}_j}$ for all $j \in [m]$.

Then,

$$k \mathbb{H}(X^n) \leq \sum_{j=1}^m \mathbb{H}(X_{\mathcal{S}_j}). \quad (1)$$

Proof.

- By assumption, $d(i) \geq k$ for all $i \in [n]$, where

$$d(i) \triangleq |\{j \in [m] : i \in \mathcal{S}_j\}|. \quad (2)$$

- Let $\mathcal{S} = \{i_1, \dots, i_\ell\}$, $1 \leq i_1 < \dots < i_\ell \leq n \implies |\mathcal{S}| = \ell$, $\mathcal{S} \subseteq [n]$.
- Let $X_{\mathcal{S}} \triangleq (X_{i_1}, \dots, X_{i_\ell})$.
- By the chain rule and the fact that conditioning reduces entropy,

$$\begin{aligned} \mathbb{H}(X_{\mathcal{S}}) &= \mathbb{H}(X_{i_1}) + \mathbb{H}(X_{i_2}|X_{i_1}) + \dots + \mathbb{H}(X_{i_\ell}|X_{i_1}, \dots, X_{i_{\ell-1}}) \\ &\geq \sum_{i \in \mathcal{S}} \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \\ &= \sum_{i=1}^n \left\{ \mathbb{1}\{i \in \mathcal{S}\} \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \right\}. \end{aligned} \quad (3)$$

Consequently, we get

$$\begin{aligned} \sum_{j=1}^m \mathbb{H}(X_{\mathcal{S}_j}) &\geq \sum_{j=1}^m \sum_{i=1}^n \left\{ \mathbb{1}\{i \in \mathcal{S}_j\} \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \right\} \\ &= \sum_{i=1}^n \left\{ \sum_{j=1}^m \mathbb{1}\{i \in \mathcal{S}_j\} \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \right\} \\ &= \sum_{i=1}^n \left\{ d(i) \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \right\} \\ &\geq k \sum_{i=1}^n \mathbb{H}(X_i|X_1, \dots, X_{i-1}) \\ &= k \mathbb{H}(X^n), \end{aligned} \quad (4)$$

where inequality (4) holds due to the nonnegativity of the conditional entropies of discrete random variables, and under the assumption that $d(i) \geq k$ for all $i \in [n]$. \square

Remark 1. If every element $i \in [n]$ belongs to *exactly* k of the subsets \mathcal{S}_j ($j \in [m]$), then Shearer's lemma also applies to continuous random variables X_1, \dots, X_n , with entropy replaced by the differential entropy. Hence, Shearer's lemma yields the subadditivity property of the Shannon entropy for discrete and continuous random variables.

Example 1. [Special case 1: Subadditivity of the Shannon entropy] Let $n = m$ with $n \in \mathbb{N}$, and $\mathcal{S}_i = \{i\}$ (singletons) for all $i \in [n]$, so every element $i \in [n]$ belongs to a single set among $\mathcal{S}_1, \dots, \mathcal{S}_n$ (i.e., $k = 1$). By Shearer's Lemma, it follows that

$$H(X^n) \leq \sum_{j=1}^n H(X_j),$$

which is the subadditivity property of the Shannon entropy for discrete random variables. This also holds for continuous random variables, where the entropy is replaced by differential entropy since every element $i \in [n]$ is contained in exactly one subset (see Remark 1).

Example 2. Special case: Han's Inequality [14] For all $\ell \in [n]$, let $\mathcal{S}_\ell = [n] \setminus \{\ell\}$. By Shearer's Lemma (Proposition 1) applied to these n subsets of $[n]$, since every element $i \in [n]$ is contained in exactly $k = n - 1$ of these subsets,

$$(n - 1) H(X^n) \leq \sum_{\ell=1}^n H(X_1, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_n) \leq n H(X^n). \quad (5)$$

An equivalent form of (5) is given by

$$0 \leq \sum_{\ell=1}^n \left\{ H(X^n) - H(X_1, \dots, X_{\ell-1}, X_{\ell+1}, \dots, X_n) \right\} \leq H(X^n). \quad (6)$$

The equivalent forms in (5) and (6) are known as Han's inequality. Note that, by Remark 1, the right-hand side inequality of (6) remains valid for continuous random variables as well.

In the combinatorial version of Shearer's lemma [6], the concept of entropy is hidden.

Proposition 2 (Combinatorial Shearer's Lemma). Consider the following setting:

- Let \mathcal{F} be a finite multiset of subsets of $[n]$ (possibly with repetitions), where each element $i \in [n]$ is included in at least $k \geq 1$ sets of \mathcal{F} .
- Let \mathcal{M} be a set of subsets of $[n]$.
- For every set $\mathcal{S} \in \mathcal{F}$, let the trace of \mathcal{M} on \mathcal{S} , denoted $\text{trace}_{\mathcal{S}}(\mathcal{M})$, be the set of all possible intersections of elements of \mathcal{M} with \mathcal{S} , i.e.,

$$\text{trace}_{\mathcal{S}}(\mathcal{M}) \triangleq \{\mathcal{A} \cap \mathcal{S} : \mathcal{A} \in \mathcal{M}\}, \quad \forall \mathcal{S} \in \mathcal{F}. \quad (7)$$

Then,

$$|\mathcal{M}| \leq \prod_{S \in \mathcal{F}} |\text{trace}_S(\mathcal{M})|^{\frac{1}{k}}. \quad (8)$$

Proof.

- Let $\mathcal{X} \subseteq [n]$ be a set that is selected uniformly at random from \mathcal{M} .
- Represent \mathcal{X} by the random vector $X^n = (X_1, \dots, X_n)$, where X_i (for all $i \in [n]$) denotes the indicator function of the event $\{i \in \mathcal{X}\}$.
- For every $S \in \mathcal{F}$, let $X_S = (X_i)_{i \in S}$. Then,

$$\mathbb{H}(X_S) \leq \log |\text{trace}_S(\mathcal{M})|. \quad (9)$$

- Applying Shearer's lemma (Proposition 1) gives

$$k \mathbb{H}(X^n) \leq \sum_{S \in \mathcal{F}} \log |\text{trace}_S(\mathcal{M})|. \quad (10)$$

- $\mathbb{H}(X^n) = \log |\mathcal{M}|$ since X^n is in one-to-one correspondence with \mathcal{X} , which is a set selected uniformly at random from \mathcal{M} . Hence,

$$\log |\mathcal{M}| \leq \frac{1}{k} \sum_{S \in \mathcal{F}} \log |\text{trace}_S(\mathcal{M})|, \quad (11)$$

and exponentiation of both sides of (11) gives (8).

□

Proposition 3 (Shearer's Lemma: Probabilistic Version). Let X^n be a discrete n -dimensional random vector, and let $\mathcal{S} \subseteq [n]$ be a random subset of $[n]$, independent of X^n , with an arbitrary probability mass function $P_{\mathcal{S}}$. If there exists $\theta > 0$ such that

$$\Pr[i \in \mathcal{S}] \geq \theta, \quad \forall i \in [n], \quad (12)$$

then,

$$\mathbb{E}_{\mathcal{S}}[\mathbb{H}(X_{\mathcal{S}})] \geq \theta \mathbb{H}(X^n). \quad (13)$$

Proof.

$$\begin{aligned} \mathbb{E}_{\mathcal{S}}[\mathbb{H}(X_{\mathcal{S}})] &= \sum_{\mathcal{S} \subseteq [n]} P_{\mathcal{S}}(\mathcal{S}) \mathbb{H}(X_{\mathcal{S}}) \\ &\geq \sum_{\mathcal{S} \subseteq [n]} \left\{ P_{\mathcal{S}}(\mathcal{S}) \sum_{i=1}^n \mathbb{1}\{i \in \mathcal{S}\} \mathbb{H}(X_i | X_1, \dots, X_{i-1}) \right\} \end{aligned} \quad (14)$$

$$= \sum_{i=1}^n \left\{ \sum_{\mathcal{S} \subseteq [n]} \left\{ P_{\mathcal{S}}(\mathcal{S}) \mathbb{1}\{i \in \mathcal{S}\} \right\} \mathbb{H}(X_i | X_1, \dots, X_{i-1}) \right\} \quad (15)$$

$$\begin{aligned} &= \sum_{i=1}^n \Pr[i \in \mathcal{S}] \mathbb{H}(X_i | X_1, \dots, X_{i-1}) \\ &\geq \theta \sum_{i=1}^n \mathbb{H}(X_i | X_1, \dots, X_{i-1}) \end{aligned} \quad (16)$$

$$= \theta \mathbb{H}(X^n).$$

where (14) holds by (3) that is valid for any set $\mathcal{S} \subseteq [n]$; (15) holds by swapping the order of summation, and (16) holds by the assumption that the random variables $\{X_i\}$ are discrete (so, the conditional entropies are nonnegative) and by the condition in (12), and the last equality holds by the chain rule for the Shannon entropy. \square

Remark 2. Similarly to Remark 1, if $\Pr[i \in \mathcal{S}] = \theta$ for all $i \in [n]$, then inequality (16) holds with equality. Hence, if the condition in (12) is satisfied with equality for all $i \in [n]$, then (13) extends to continuous random variables, with entropies replaced by differential entropies.

B. Intersecting Families of Graphs

We start by considering triangle-intersecting families of graphs, which was the problem in extremal combinatorics address in [6].

Definition 1 (Triangle-Intersecting Families of Graphs). Let \mathcal{G} be a family of graphs on the vertex set $[n]$, with the property that for every $G_1, G_2 \in \mathcal{G}$, the intersection $G_1 \cap G_2$ contains a triangle (i.e, there are three vertices $i, j, k \in [n]$ such that each of $\{i, j\}$, $\{i, k\}$, $\{j, k\}$ is in the edge sets of both G_1 and G_2). The family \mathcal{G} is referred to as a *triangle-intersecting* family of graphs on n vertices.

Question 1. (*Simonovits and Sós, 1976 [34]*) How large can \mathcal{G} (a family of triangle-intersecting graphs) be?

The family \mathcal{G} can be as large as $2^{\binom{n}{2}-3}$. To that end, consider the family \mathcal{G} of all graphs on n vertices that include a particular triangle. On the other hand, $|\mathcal{G}|$ cannot exceed $2^{\binom{n}{2}-1}$. The latter upper bound holds since, in general, a family of distinct subsets of a set of size m , where any two of these subsets have a non-empty intersection, can have a cardinality of at most 2^{m-1} (\mathcal{A} and \mathcal{A}^c cannot be members of this family). The edge sets of the graphs in \mathcal{G} satisfy this property, with $m = \binom{n}{2}$.

Proposition 4 (Ellis, Filmus and Friedgut (2012), [8]). The size of a family \mathcal{G} of triangle-intersecting graphs on n vertices satisfies $|\mathcal{G}| \leq 2^{\binom{n}{2}-3}$, and this upper bound is attained by the family of all graphs with a common vertex set of n vertices, and with a fixed common triangle.

This result was proved by using discrete Fourier analysis to obtain the sharp bound in Proposition 4, as conjectured by Simonovits and Sós [34].

The first significant progress towards proving the Simonovits–Sós conjecture came from an information-theoretic approach [6]. Using the combinatorial Shearer lemma (Proposition 4), a simple and elegant upper bound on the size of \mathcal{G} was derived in [6]. That bound is equal to $2^{\binom{n}{2}-2}$, falling short of the Simonovits–Sós conjecture by a factor of 2.

Proposition 5 (Chung, Graham, Frankl, and Shearer (1986)). Let \mathcal{G} be a family of K_3 -intersecting graphs on a common vertex set $[n]$. Then, $|\mathcal{G}| \leq 2^{\binom{n}{2}-2}$.

We next consider more general intersecting families of graphs.

Definition 2 (H-intersecting Families of Graphs). Let \mathcal{G} be a family of graphs on a common vertex set. Then, it is said that \mathcal{G} is H-intersecting if for every two graphs $G_1, G_2 \in \mathcal{G}$, the graph $G_1 \cap G_2$ contains H as a subgraph.

Example 3. Let $H = K_t$ with $t \geq 2$. Then,

- $t = 2$ means that \mathcal{G} is intersecting,

- $t = 3$ means that \mathcal{G} is triangle-intersecting.

Question 2. [*Problem in Extremal Combinatorics*] Given H and n , what is the maximum size of an H -intersecting family of graphs on n labeled vertices?

Conjecture 1. (*Ellis, Filmus, and Friedgut, 2012*) Every K_t -intersecting family of graphs on a common vertex set $[n]$ has size at most $2^{\binom{n}{2} - \binom{t}{2}}$, with equality for the family of all graphs containing a fixed clique on t vertices.

- For $t = 2$, it is trivial (since K_2 is an edge).
- For $t = 3$, it was proved by Ellis, Filmus & Friedgut [8].
- For $t = 4$, it was recently proved by Berger and Zhao [3].
- For $t \geq 5$, this problem is left open.

C. Counting Graph Homomorphisms

In the sequel, let $V(H)$ and $E(H)$ denote the vertex and edge sets of a graph H , respectively. Further, let T and G be finite, simple, and undirected graphs, and denote the edge connecting a pair of adjacent vertices $u, v \in V(H)$ by an edge $e = \{u, v\} \in E(H)$.

Definition 3. A *homomorphism* from T to G , denoted by $T \rightarrow G$, is a mapping of the vertices of T to those of G , $\sigma: V(T) \rightarrow V(G)$, such that every edge in T is mapped to an edge in G :

$$\{u, v\} \in E(T) \implies \{\sigma(u), \sigma(v)\} \in E(G). \quad (17)$$

On the other hand, non-edges in T may be mapped to the same vertex, a non-edge, or an edge in G .

Example 4. There is a homomorphism from every bipartite graph G to K_2 . To that end, let $V(G) = \mathcal{X} \cup \mathcal{Y}$, where \mathcal{X} and \mathcal{Y} are the two partite sets. Then, a mapping that maps every vertex in \mathcal{X} to '0', and every vertex in \mathcal{Y} to '1' is a homomorphism $G \rightarrow K_2$. This holds because every edge in G is mapped to the edge $\{0, 1\}$ in K_2 .

- Every non-edge in \mathcal{X} or in \mathcal{Y} is mapped to the same vertex in K_2 .

- Every non-edge between two vertices in \mathcal{X} and \mathcal{Y} is mapped to $\{0, 1\}$.

The following connects graph homomorphisms to graph invariants. Let $\omega(G)$ and $\chi(G)$ denote the clique number and chromatic number of a finite, simple, and undirected graph G . Then,

- $\omega(G)$ is the largest integer k for which a homomorphism $K_k \rightarrow G$ exists. This holds because the image of a complete graph under a homomorphism is a complete graph of the same size. This is valid because a homomorphism preserves adjacency, and for a complete graph K_k , all pairs of vertices are adjacent. To preserve this property, the image of K_k under the homomorphism must also be a complete graph of size k .
- A graph G is k -colorable if and only if it has a homomorphism to the complete graph K_k ; this is because k -coloring assigns one of k colors to each vertex such that adjacent vertices receive different colors, which is equivalent to mapping the vertices of G to the k vertices of K_k in a way that adjacency is preserved. Consequently, it follows by definition that $\chi(G)$ is the smallest integer k for which there exists a homomorphism $G \rightarrow K_k$.

Let $\text{Hom}(T, G)$ denote the set of all the homomorphisms $T \rightarrow G$, and let

$$\text{hom}(T, G) \triangleq |\text{Hom}(T, G)| \tag{18}$$

denote the number of these homomorphisms.

Definition 4 (Fractional independence number). The fractional independence number of a graph G , denoted as $\alpha_f(G)$, is a fractional relaxation of the independence number $\alpha(G)$. It is defined as the optimal value of the following linear program:

- Optimization variables: x_v for every vertex $v \in V(G)$.
- Objective: Maximize $\sum_{v \in V(G)} x_v$.
- Constraints: $x_v \geq 0$ for all $v \in V(G)$, and $\sum_{v \in \mathcal{C}} x_v \leq 1$ for every clique $\mathcal{C} \subseteq V(G)$.

This relaxation allows fractional values for x_v , in contrast to the integer programming formulation for $\alpha(G)$, where x_v must be binary (either 0 or 1) for all $v \in V(G)$. Consequently, $\alpha_f(G) \geq \alpha(G)$.

The following result was obtained by Alon [2] and by Friedgut and Khan [11], where the latter provides an entropy-based proof that relies on Shearer's lemma with an extension of the

result on the number of homomorphisms for hypergraphs.

Proposition 6. Let T and G be finite, simple, and undirected graphs, having no isolated vertices.

Then,

$$\text{hom}(T, G) \leq (2|E(G)|)^{\alpha_f(T)}. \quad (19)$$

Furthermore, the upper bound in (19) is essentially tight for a fixed graph T in the sense that there exists a graph G such that

$$\text{hom}(T, G) \geq \left(\frac{|E(G)|}{|E(T)|} \right)^{\alpha_f(T)}. \quad (20)$$

III. INTERSECTING FAMILIES OF GRAPHS

The following result generalizes Proposition 5 and it extends the concept of proof in [6] to hold for every family of H -intersecting graphs on a common vertex set. It provides a new upper bound on the maximum cardinality of a family of graphs on a fixed number of vertices, in which every pair of graphs shares a fixed common subgraph. The derived bound is expressed in terms of the chromatic number of the shared subgraph.

Proposition 7. Let H be a non-empty graph, and let \mathcal{G} be a family of H -intersecting graphs on a common vertex set $[n]$. Then, $|\mathcal{G}| \leq 2^{\binom{n}{2} - (\chi(H)-1)}$.

Proof.

- Identify $G \in \mathcal{G}$ with its edge set $E(G)$, and let $\mathcal{M} = \{E(G) : G \in \mathcal{G}\}$ (all these graphs have the common vertex set $[n]$).
- Let $\mathcal{U} = E(K_n)$. For every $G \in \mathcal{G}$, we have $E(G) \subseteq \mathcal{U}$, and $|\mathcal{U}| = \binom{n}{2}$.
- Let $t \triangleq \chi(H)$. For every unordered equipartition of $[n]$ into $t-1$ subsets, i.e., $\bigcup_{j=1}^{t-1} \mathcal{A}_j = [n]$, which satisfies $||\mathcal{A}_i| - |\mathcal{A}_j|| \leq 1$ for all $1 \leq i < j \leq t-1$, let $\mathcal{U}(\{\mathcal{A}_j\}_{j=1}^{t-1})$ be the subset of \mathcal{U} consisting of all those edges that lie entirely inside one of the subsets $\{\mathcal{A}_j\}_{j=1}^{t-1}$.
- We apply Proposition 2 with $\mathcal{F} = \{\mathcal{U}(\{\mathcal{A}_j\}_{j=1}^{t-1})\}$ taken over all unordered equipartition of $[n]$, $\{\mathcal{A}_j\}_{j=1}^{t-1}$, as above.

- Let $m = |\mathcal{U}(\{\mathcal{A}_j\}_{j=1}^{t-1})|$, which is independent of the equipartition since

$$m = \begin{cases} (t-1)\binom{n/(t-1)}{2} & \text{if } (t-1)|n, \\ (t-2)\binom{\lfloor n/(t-1) \rfloor}{2} + \binom{\lceil n/(t-1) \rceil}{2} & \text{if } (t-1)|(n-1), \\ \vdots \\ \binom{\lfloor n/(t-1) \rfloor}{2} + (t-2)\binom{\lceil n/(t-1) \rceil}{2} & \text{if } (t-1)|(n-(t-2)). \end{cases} \quad (21)$$

- By (21) with $t \triangleq \chi(\mathbf{H})$, it follows that

$$m \leq \frac{1}{\chi(\mathbf{H}) - 1} \binom{n}{2}. \quad (22)$$

Proof. The graph \mathbf{H} is non-empty, so $t = \chi(\mathbf{H}) \geq 2$. If $(t-1)|n$, then,

$$\begin{aligned} (t-1)\binom{n/(t-1)}{2} &= \frac{n(n-(t-1))}{2(t-1)} \\ &\leq \frac{1}{2(t-1)} \binom{n}{2}. \end{aligned}$$

Otherwise, $(t-1)|(n-j)$ for some integer $j \in [t-2]$, so $n = j + r(t-1)$ with $r \in \mathbb{N}$.

Consequently, (21) gives

$$\begin{aligned} m &= (t-1-j)\binom{r}{2} + j\binom{r+1}{2} \\ &= \frac{r}{2} [(t-1-j)(r-1) + r+1] \\ &= \frac{n-j}{2(t-1)} \left[(t-1-j)\left(\frac{n-j}{t-1} - 1\right) + \frac{n-j}{t-1} + 1 \right] \\ &= \frac{n-j}{2(t-1)^2} [(t-1-j)(n-j-t+1) + n-j+t-1] \\ &= \frac{(n-j)[n(t-1) - (t-1)(t-2) - (j-1)(n-j)]}{2(t-1)^2} \\ &\leq \frac{(n-j)n(t-1)}{2(t-1)^2} \\ &\leq \frac{n(n-1)}{2(t-1)} \\ &= \frac{1}{t-1} \binom{n}{2}. \end{aligned}$$

□

- By a simple double-counting argument in regard to the edges of the complete graph K_n (the set \mathcal{U}), if k is the number of elements of \mathcal{F} in which each element of \mathcal{U} occurs, then

$$m|\mathcal{F}| = \binom{n}{2}k. \quad (23)$$

- Let $\mathcal{S} \in \mathcal{F}$. Observe that $\text{trace}_{\mathcal{S}}(\mathcal{M})$, as defined in (7), forms an intersecting family of subsets of \mathcal{S} . Indeed,

- 1) Let us assign to each vertex in $[n]$ the index of the subset \mathcal{A}_j ($1 \leq j \leq \chi(H) - 1$) in the partitioning of $[n]$ that corresponds to \mathcal{S} . Let these assignments be associated with $\chi(H) - 1$ color classes of the vertices.
- 2) For any $G, G' \in \mathcal{G}$, the graph $G \cap G'$ has a subgraph H (by assumption).
- 3) By the definition of the chromatic number of H as the smallest number of colors that are required such that any two adjacent vertices in H are assigned different colors, it follows that there exists an edge in H whose two vertices are assigned the same index (color). Hence, that edge belongs to the set \mathcal{A}_j , for some $j \in [\chi(H) - 1]$, so it belongs to \mathcal{S} .
- 4) The complement of \mathcal{S} (in \mathcal{U}) is therefore H -free (viewed as a graph with the vertex set $[n]$).

Consequently, since $|\mathcal{S}| = m$, we get

$$|\text{trace}_{\mathcal{S}}(\mathcal{M})| \leq 2^{m-1}. \quad (24)$$

- By Proposition 2 (and the one-to-one correspondence between \mathcal{G} and \mathcal{M}),

$$|\mathcal{G}| = |\mathcal{M}| \leq (2^{m-1})^{\frac{|\mathcal{F}|}{k}} \quad (25)$$

$$= 2^{\binom{n}{2} \left(1 - \frac{1}{m}\right)} \quad (26)$$

$$\leq 2^{\binom{n}{2} - (\chi(H) - 1)}, \quad (27)$$

where (25) relies on (8) and (24), then (26) relies on (23), and (27) holds due to (22).

□

The family \mathcal{G} of H -intersecting graphs on n vertices can be as large as $2^{\binom{n}{2}-|E(H)|}$. To that end, consider the family \mathcal{G} of all graphs on n vertices that include a particular H subgraph. Combining this with Proposition 7 gives that the largest family \mathcal{G} of H -intersecting graphs on n vertices satisfies

$$2^{\binom{n}{2}-|E(H)|} \leq |\mathcal{G}| \leq 2^{\binom{n}{2}-\chi(H)-1}. \quad (28)$$

Specialization of Proposition 7 to complete subgraphs gives the following.

Corollary 1. Let \mathcal{G} be a family of K_t -intersecting graphs, with $t \geq 2$, on a common vertex set $[n]$. Then, $|\mathcal{G}| \leq 2^{\binom{n}{2}-(t-1)}$.

Proof. For a complete subgraph $H = K_t$, we have $\chi(H) = t$. □

Remark 3. The bound in Corollary 1 falls short of the conjectured result by Ellis, Filmus, and Friedgut (2012), which states that for every K_t -intersecting family of graphs on a common vertex set $[n]$ has size at most $2^{\binom{n}{2}-\frac{t(t-1)}{2}}$, with equality for the family of all graphs containing a fixed clique on t vertices. It, however, generalizes Proposition 5, which is a specialized result with $H = K_3$ (triangle-intersecting graphs), and it uniformly improves the trivial bound of $2^{\binom{n}{2}-1}$ for $H = K_t$ for all $t \geq 3$.

Remark 4. If H is a bipartite graph, then $\chi(H) = 2$. In that case, our result for an H -intersecting family of graphs on n vertices is specialized to the trivial bound $|\mathcal{G}| \leq 2^{\binom{n}{2}-1}$.

The computational complexity of the chromatic number of a graph is in general NP-hard. This poses a problem in calculating the upper bound in Proposition 7 on the cardinality of H -intersecting families of graphs on a fixed number of vertices. This bound can be however loosened, expressing it in terms of the Lovász ϑ -function of the complement graph \overline{H} (see Corollary 3 in [20]).

Corollary 2. Let H be a graph, and let \mathcal{G} be a family of H -intersecting graphs on a common vertex set $[n]$. Then, $|\mathcal{G}| \leq 2^{\binom{n}{2}-\lceil\vartheta(\overline{H})\rceil-1}$.

Proof. The Lovász ϑ -function of the complement graph \overline{H} satisfies the inequality

$$\omega(H) \leq \vartheta(\overline{H}) \leq \chi(H), \quad (29)$$

so it is bounded between the clique and chromatic numbers of H , which are both NP-hard to compute. Since the chromatic number $\chi(H)$ is an integer, we also have $\chi(H) \geq \lceil \vartheta(\overline{H}) \rceil$. \square

The Lovász ϑ -function of the complement graph \overline{H} , which appears in Corollary 2, can be efficiently computed in polynomial time by solving the following semidefinite (SDP) problem:

$$\begin{array}{l} \text{maximize } \text{Tr}(\mathbf{B} \mathbf{J}_p) \\ \text{subject to} \\ \left\{ \begin{array}{l} \mathbf{B} \in \mathcal{S}_+^p, \quad \text{Tr}(\mathbf{B}) = 1, \\ A_{i,j} = 0 \Rightarrow B_{i,j} = 0, \quad i, j \in [p], i \neq j. \end{array} \right. \end{array} \quad (30)$$

where

- $\mathbf{A} = \mathbf{A}(H)$ is the $p \times p$ adjacency matrix of H , with $p \triangleq |\mathbf{V}(H)|$;
- \mathbf{J}_p is the all-ones $p \times p$ matrix;
- \mathcal{S}_+^p is the set of all $p \times p$ positive semidefinite matrices.

The reader is referred to an account of properties of the Lovász ϑ -function in Chapter 11 of [21], and a recent presentation in Section 2.5 of [32].

The following result generalizes Corollary 1 by relying on properties of the Lovász ϑ -function.

Proposition 8. Let \mathcal{G} be an H -intersecting family of graphs on n vertices, where H is a non-empty graph on t vertices.

(1)

$$\log_2 |\mathcal{G}| \leq \binom{n}{2} - \left\lceil \max_{\mathbf{T}} \frac{\lambda_{\max}(\mathbf{T})}{|\lambda_{\min}(\mathbf{T})|} \right\rceil, \quad (31)$$

where the maximization on the right-hand side of (31) is taken over all symmetric nonzero $t \times t$ matrices $\mathbf{T} = (T_{i,j})$ with $T_{i,j} = 0$ for all $i, j \in [t]$ such that $\{i, j\} \notin E(H)$ or $i = j$ (e.g., the adjacency matrix of H), and $\lambda_{\max}(\mathbf{T})$ and $\lambda_{\min}(\mathbf{T})$ denote the largest and smallest (real) eigenvalues of \mathbf{T} , respectively.

(2) Specifically, if H is a d -regular graph and $d \in [t - 1]$, then

$$\log_2 |\mathcal{G}| \leq \binom{n}{2} - \left\lceil \frac{d}{|\lambda_{\min}(H)|} \right\rceil, \quad (32)$$

where $\lambda_{\min}(H)$ is the smallest eigenvalue of the adjacency matrix of the graph H .

(3) If H is a connected strongly d -regular graph with exactly λ common neighbors for every two adjacent vertices and exactly μ common neighbors for every two non-adjacent vertices, then

$$\log_2 |\mathcal{G}| \leq \binom{n}{2} - \left\lceil \frac{2d}{\sqrt{(\lambda - \mu)^2 + 4(d - \mu)} - \lambda + \mu} \right\rceil. \quad (33)$$

Proof. By Corollary 2,

$$\log_2 |\mathcal{G}| \leq \binom{n}{2} - (\lceil \vartheta(\overline{H}) \rceil - 1). \quad (34)$$

Item 1 then holds by the property that for every finite, simple, and undirected graph G on n vertices

$$\vartheta(G) = 1 + \max_{\mathbf{T}} \frac{\lambda_{\max}(\mathbf{T})}{|\lambda_{\min}(\mathbf{T})|}, \quad (35)$$

where the maximization on the right-hand side of (35) is taken over all symmetric nonzero $n \times n$ matrices $\mathbf{T} = (T_{i,j})$ with $T_{i,j} = 0$ for all $i, j \in [n]$ such that $\{i, j\} \in E(G)$ or $i = j$. Equality (35) is then applied to $G = \overline{H}$, so the maximization on the right-hand side of (31) is needed to be taken over all symmetric nonzero $t \times t$ matrices $\mathbf{T} = (T_{i,j})$ with $T_{i,j} = 0$ for all $i, j \in [t]$ such that $\{i, j\} \notin E(H)$ or $i = j$. This includes in particular the adjacency matrix of the graph H , i.e., $\mathbf{T} = \mathbf{A}(H)$.

We next prove Item 2, which refers to non-empty d -regular graphs on t vertices. Relaxing the bound in (31) by selecting $\mathbf{T} = \mathbf{A}(H)$ gives $\lambda_{\max}(\mathbf{T}) = d$, and $\lambda_{\min}(\mathbf{T}) = \lambda_{\min}(H)$, which gives the relaxed bound in (32).

Item 3 is then obtained from Item 2 by relying on the closed-form expression of the smallest eigenvalue of the adjacency matrix of a strongly d -regular graph H on t vertices with exactly λ common neighbors for every two adjacent vertices and exactly μ common neighbors for every two non-adjacent vertices. In that case, the largest eigenvalue of the adjacency matrix is

$\lambda_1(H) = d$ with multiplicity 1, and the other two distinct eigenvalues of its adjacency matrix are given by

$$p_{1,2} = \frac{1}{2} \left(\lambda - \mu \pm \sqrt{(\lambda - \mu)^2 + 4(d - \mu)} \right), \quad (36)$$

with the respective multiplicities

$$m_{1,2} = \frac{1}{2} \left(t - 1 \mp \frac{2d + (t - 1)(\lambda - \mu)}{\sqrt{(\lambda - \mu)^2 + 4(d - \mu)}} \right). \quad (37)$$

Specifically, by (36), the absolute value of the smallest eigenvalue of H is given by

$$|\lambda_{\min}(H)| = \frac{1}{2} \left(\sqrt{(\lambda - \mu)^2 + 4(d - \mu)} + \mu - \lambda \right). \quad (38)$$

Finally, substituting (38) into (32) gives (33). \square

It is natural to ask the following question:

Question 3. Is there a graph H (apart of an edge) for which the bound provided in Proposition 7 is tight for a largest H-intersecting family of graphs?

We provide a partial reply to Question 3 by comparing the leftmost and rightmost terms in (28), which is equivalent to comparing $|E(H)|$ and $\chi(H) - 1$. According to the inequality

$$\chi(H) \leq \Delta(H) + 1, \quad (39)$$

where $\Delta(H)$ is the maximum degree of the vertices in the graph H, it follows that unless H is an edge, there is a gap between the size of the graph ($|E(H)|$) and the chromatic number minus one ($\chi(H) - 1$). Furthermore, according to Brooks' Theorem, for connected, undirected graphs H that are neither complete nor odd cycles, the chromatic number satisfies

$$\chi(H) \leq \Delta(H), \quad (40)$$

which provides a tighter bound in comparison to (39), further increasing the gap between $|E(H)|$ and $\chi(H) - 1$ unless H is an edge. It is also noted that the chromatic number satisfies

$$\chi(H) \leq \frac{1}{2} (1 + \sqrt{1 + 8|E(H)|}), \quad (41)$$

with an equality if and only if H is a complete graph. This bound arises by observing that the chromatic number cannot exceed the largest integer k such that $\binom{k}{2} \leq |E(H)|$. Hence, as $m = |E(H)|$ gets large, the smallest possible gap between $\chi(H) - 1$ and $|E(H)|$ is increased.

IV. NUMBER OF GRAPH HOMOMORPHISMS

Proposition 9. Let G be a finite, simple, and undirected graph on n vertices, and let m_ℓ be the number of cliques of order $\ell \in \mathbb{N}$ in G . Then, for all $s, t \in \mathbb{N}$ with $2 \leq s < t \leq n$,

$$(t! m_t)^s \leq (s! m_s)^t. \quad (42)$$

We next provide a generalization of Proposition 9.

Proposition 10. Let G be a finite, simple, and undirected graph on n vertices, let $s, t \in \mathbb{N}$ with $s < t < n$, let T be an induced subgraph of G on t vertices, and let $m(H, G)$ denote the number of copies of a subgraph H in the graph G . Then,

$$(t! m(T, G))^s \leq \max_S (s! m(S, G))^t, \quad (43)$$

where the maximization on the right-hand side of (43) is with respect to all induced subgraphs S of T on s vertices. Equivalently, in terms of homomorphism counts,

$$\left(\frac{t!}{\text{aut}(T)} \cdot \text{hom}(T, G) \right)^s \leq \max_S \left(\frac{s!}{\text{aut}(S)} \cdot \text{hom}(S, G) \right)^t, \quad (44)$$

where the maximization on the right-hand side of (44) is as in (43), and $\text{aut}(H)$ denotes the size of the automorphism group of H .

Proof.

- Label the vertices of G by the elements of $[n]$, and let T be an induced subgraph of G with $|V(T)| = t < n$.
- Select a copy of T in G uniformly at random, and then choose a uniform random ordering of the vertices in that copy. This process produces a random vector (X_1, \dots, X_t) , representing the selected order of the vertices.

- Let $m(\mathsf{T}, \mathsf{G})$ denote the number of copies of T in G . Then,

$$\mathsf{H}(X_1, \dots, X_t) = \log(t! m(\mathsf{T}, \mathsf{G})), \quad (45)$$

as the vertices of each copy of T in G can be ordered in $t!$ equally probable ways.

- Let $\mathcal{S} \subseteq [t]$ be chosen uniformly at random from all subsets of $[t]$ of fixed size s , where $1 \leq s < t$. Then,

$$\Pr[i \in \mathcal{S}] = \frac{s}{t}, \quad \forall i \in [t]. \quad (46)$$

- By Proposition 3 and equalities (45) and (46), it follows that

$$\mathbb{E}_{\mathcal{S}}[\mathsf{H}(X_{\mathcal{S}})] \geq \frac{s}{t} \cdot \log(t! m(\mathsf{T}, \mathsf{G})). \quad (47)$$

- The random subvector $X_{\mathcal{S}}$ corresponds to a copy, in G , of an induced subgraph $\mathsf{S} \subseteq \mathsf{T}$ with s vertices. All $s!$ permutations of the subvector $X_{\mathcal{S}}$ correspond to the same copy of S in G , and there are $m(\mathsf{S}, \mathsf{G})$ such copies of S in G .
- The entropy of the random subvector $X_{\mathcal{S}}$ therefore satisfies

$$\mathsf{H}(X_{\mathcal{S}}) \leq \log(s! m(\mathsf{S}, \mathsf{G})), \quad (48)$$

where $m(\mathsf{S}, \mathsf{G})$ denotes the number of copies of a graph S in G , and $s!$ accounts for the $s!$ permutations of the vector $X_{\mathcal{S}}$ that corresponds to the same copy of S in G .

- By (48), it follows that

$$\mathbb{E}_{\mathcal{S}}[\mathsf{H}(X_{\mathcal{S}})] \leq \max_{\mathsf{S}} \log(s! m(\mathsf{S}, \mathsf{G})), \quad (49)$$

where the maximization on the right-hand side of (49) is taken over all induced subgraphs S of T on s vertices.

- Combining (47) and (49) yields

$$\frac{s}{t} \cdot \log(t! m(\mathsf{T}, \mathsf{G})) \leq \max_{\mathsf{S}} \log(s! m(\mathsf{S}, \mathsf{G})), \quad (50)$$

and exponentiating both sides (50) gives (43).

- Let $\text{aut}(H)$ denote the size of automorphism group of a graph H , defined as the number of vertex permutations that preserve the graph's structure, i.e., its adjacency and non-adjacency relations. Then,

$$\text{hom}(H, G) = \text{aut}(H) m(H, G), \quad (51)$$

which gives inequality (44) by combining (43) and (51). Furthermore, by (51), inequalities (43) and (44) are equivalent.

□

Remark 5 (Specialization of Proposition 10 to Proposition 9). Proposition 9 follows as a corollary of Proposition 10 by setting $T = K_t$ (a clique of order t), for which every induced subgraph of T on s vertices is a clique S of order s ($S = K_s$). In that case, $\text{aut}(T) = t!$ and $\text{aut}(S) = s!$. Consequently, the maximization on the right-hand side of (44) is performed over the single graph K_s , which gives

$$\text{hom}(K_t, G)^s \leq \text{hom}(K_s, G)^t, \quad 1 \leq s < t < n. \quad (52)$$

By (51), we have

$$\text{hom}(K_t, G) = t! m_t, \quad (53)$$

$$\text{hom}(K_s, G) = s! m_s, \quad (54)$$

where m_t and m_s denote, respectively, the number of cliques of orders t and s in G . Combining (52), (53), and (54) then gives

$$(t! m_t)^s \leq (s! m_s)^t, \quad 1 \leq s < t < n. \quad (55)$$

This reproduces Proposition 9, which establishes a relationship between the numbers of cliques of two different orders in a finite, simple, undirected graph G .

We next rely on the Shannon entropy to derive a lower bound on the number of homomorphisms from any complete bipartite graph to any bipartite graph, and examine its tightness by a derivation of an upper bound that relies on Proposition 6.

Proposition 11. Let G be a simple bipartite graph having no isolated vertices. Let its partite vertex sets be of sizes n_1 and n_2 , and let its number of edges be equal to $\alpha n_1 n_2$ with $\alpha \in (0, 1]$. Then, the number of homomorphisms from the complete bipartite graph $K_{s,t}$ to G , where $s, t \in \mathbb{N}$, satisfies

$$\alpha^{st} n_1^s n_2^t \leq \text{hom}(K_{s,t}, G) \leq (2\alpha n_1 n_2)^{\max\{s,t\}}. \quad (56)$$

Proof. We first provide an entropy-based proof of the lower bound in the leftmost term of (56). Let \mathcal{U} and \mathcal{V} denote the partite vertex sets of the simple bipartite graph G , where $|\mathcal{U}| = n_1$ and $|\mathcal{V}| = n_2$. By assumption, $|E(G)| = \alpha n_1 n_2$ where $\alpha \in (0, 1]$.

Let (U, V) be a random vector taking values in $\mathcal{U} \times \mathcal{V}$, and distributed uniformly at random on the edges of G . Then, its entropy is $H(U, V) = \log(\alpha n_1 n_2)$. The random vector (U, V) can be sampled by first sampling $U = u$ from the marginal probability mass function (PMF) of U (P_U), and then sampling V from the conditional PMF $P_{V|U}(\cdot|u)$.

Next, consider the random vector (U_1, V_1, \dots, V_t) by first sampling $U_1 = u_1$ from the marginal PMF P_U , and then sampling $V_1 = v_1, \dots, V_t = v_t$ independently at random from the conditional PMF $P_{V|U}(\cdot|u_1)$. Then,

$$\begin{aligned} H(U_1, V_1, \dots, V_t) &= H(U_1) + \sum_{i=1}^t H(V_i|U_1) \\ &= H(U) + t H(V|U) \\ &= t H(U, V) - (t-1) H(U) \\ &= t \log(\alpha n_1 n_2) - (t-1) H(U) \\ &\geq t \log(\alpha n_1 n_2) - (t-1) \log n_1 \\ &= \log(\alpha^t n_1^s n_2^t). \end{aligned} \quad (57)$$

Further, consider the random vector $(U_1, \dots, U_s, V_1, \dots, V_t)$ by first sampling $(V_1, \dots, V_t) = (v_1, \dots, v_t)$ from the joint PMF P_{V_1, \dots, V_t} , and then sampling U_1, \dots, U_s independently from the

conditional PMF $P_{U|V_1, \dots, V_t}(\cdot | v_1, \dots, v_t)$, where

$$P_{U|V_1, \dots, V_t}(u | v_1, \dots, v_t) = \frac{P_U(u) \prod_{i=1}^t P_{V|U}(v_i | u)}{\sum_{u \in \mathcal{U}} \left\{ P_U(u) \prod_{i=1}^t P_{V|U}(v_i | u) \right\}}, \quad u \in \mathcal{U}. \quad (58)$$

Then,

$$\begin{aligned} & H(U_1, \dots, U_s, V_1, \dots, V_t) \\ &= H(V_1, \dots, V_t) + \sum_{i=1}^s H(U_i | V_1, \dots, V_t) \\ &= H(V_1, \dots, V_t) + s H(U_1 | V_1, \dots, V_t) \\ &= s [H(V_1, \dots, V_t) + H(U_1 | V_1, \dots, V_t)] - (s-1) H(V_1, \dots, V_t) \\ &= s H(U_1, V_1, \dots, V_t) - (s-1) H(V_1, \dots, V_t) \\ &\geq s \log(\alpha^t n_1 n_2^t) - (s-1) H(V_1, \dots, V_t) \quad (\text{by (57)}) \\ &\geq s \log(\alpha^t n_1 n_2^t) - (s-1) \log(n_2^t) \\ &= \log(\alpha^{st} n_1^s n_2^t). \end{aligned} \quad (59)$$

The vector $(U_1, \dots, U_s, V_1, \dots, V_t)$ corresponds to a homomorphism from $K_{s,t}$ to G (this holds by construction, see (58) where P_{UV} is uniformly distributed over the edges of the graph G , P_U is the marginal PMF of U , and $P_{V|U}$ is the conditional PMF of V given U). Hence,

$$H(U_1, \dots, U_s, V_1, \dots, V_t) \leq \log \text{hom}(K_{s,t}, G). \quad (60)$$

Combining (59) and (60) yields

$$\text{hom}(K_{s,t}, G) \geq \alpha^{st} n_1^s n_2^t. \quad (61)$$

We next prove the upper bound on the number of homomorphisms in the rightmost term of (56). By Proposition 6, with $|E(G)| = \alpha n_1 n_2$, we get

$$\text{hom}(K_{s,t}, G) \leq (2\alpha n_1 n_2)^{\alpha_t(K_{s,t})}. \quad (62)$$

Every bipartite graph is a perfect graph, so $\alpha(K_{s,t}) = \alpha_t(K_{s,t}) = \chi(\overline{K_{s,t}})$. The independence number of $K_{s,t}$ is the size of the largest among the left and right vertex sets, so it satisfies

$\alpha(K_{s,t}) = \max\{s, t\}$. Similarly, $\overline{K_{s,t}} = K_s \cup K_t$ is the disjoint union of the complete graphs K_s and K_t , so $\chi(\overline{K_{s,t}}) = \max\{s, t\}$. Since $\alpha(H) \leq \alpha_f(H) \leq \chi(\overline{H})$ for every graph H , it follows that $\alpha_f(K_{s,t}) = \max\{s, t\}$, whose substitution into (62) gives the rightmost inequality in (56). \square

Setting $s = t$ in Proposition 11 gives the following.

Corollary 3. Let G be a bipartite graph having no isolated vertices. Then, in the setting of Proposition 11,

$$\alpha^{s^2}(n_1 n_2)^s \leq \text{hom}(K_{s,s}, G) \leq (2\alpha)^s (n_1 n_2)^s, \quad (63)$$

so, for a fixed $\alpha \in (0, 1)$, it follows from (63) that the number of homomorphisms from the complete bipartite graph $K_{s,t}$ to G scales like $(n_1 n_2)^s$.

Acknowledgment

A stimulating discussion with Yuval Peled, which took place while the author gave a seminar talk on the subject at the Einstein Institute of Mathematics in the Hebrew University of Jerusalem, is gratefully acknowledged. The author also appreciates the hospitality during the seminar, organized by Yuval.

Conflict of interest

The author declares no conflicts of interest.

REFERENCES

- [1] M. Aigner and G. M. Ziegler, *Proofs from THE BOOK*, Sixth Edition, Springer, Berlin, 2018. <https://doi.org/10.1007/978-3-662-57265-8>
- [2] N. Alon, “On the number of subgraphs of prescribed type of graphs with a given number of edges,” *Israel Journal of Mathematics*, vol. 38, pp. 116–130, 1981. <https://doi.org/10.1007/BF02761855>
- [3] A. Berger and Y. Zhao, “ K_4 -intersecting families of graphs,” *Journal of Combinatorial Theory, Series B*, vol. 163, pp. 112–132, November 2023. <https://doi.org/10.1016/j.jctb.2023.07.005>
- [4] C. Borgs, J. Chayes, L. Lovász, V. T. Sós, and K. Vesztegombi, “Counting graph homomorphisms,” *Topics in Discrete Mathematics*, pp. 315–371, Springer, 2006. https://doi.org/10.1007/3-540-33700-8_18
- [5] G. Brightwell and P. Winkler, “Graph homomorphisms and phase transitions,” *Journal of Combinatorial Theory, Series B*, vol. 77, no. 2, pp. 221–262, 1999. <https://doi.org/10.1006/jctb.1999.1899>

- [6] F. R. K. Chung, L. R. Graham, P. Frankl and J. B. Shearer, “Some intersection theorems for ordered sets and graphs,” *Journal of Combinatorial Theory, Series A*, vol. 43, no. 1, pp. 23–37, 1986. [https://doi.org/10.1016/0097-3165\(86\)90019-1](https://doi.org/10.1016/0097-3165(86)90019-1)
- [7] T. M. Cover and J. A. Thomas, *Elements of Information Theory*, second edition, John Wiley & Sons, 2006. <https://doi.org/10.1002/047174882X>
- [8] D. Ellis, Y. Filmus, and E. Friedgut, “Triangle-intersecting families of graphs,” *Journal of the European Mathematical Society*, vol. 14, no. 3, pp. 841–855, 2012. <https://doi.org/10.4171/JEMS/320>
- [9] D. Ellis, “Intersection problems in extremal combinatorics: theorems, techniques and questions - old and new,” *Surveys in Combinatorics 2022*, pp. 115–173. London Mathematical Society Lecture Note Series, Cambridge University Press, 2022. <https://doi.org/10.1017/9781009093927.005>
- [10] E. Friedgut, “Hypergraphs, entropy, and inequalities,” *The American Mathematical Monthly*, vol. 111, no. 9, pp. 749–760, November 2004. <https://doi.org/10.2307/4145187>
- [11] E. Friedgut and J. Kahn, “On the number of copies of one hypergraph in another,” *Israel Journal of Mathematics*, vol. 105, pp. 251–256, 1998. <https://doi.org/10.1007/BF02780332>
- [12] D. Galvin, “Three tutorial lectures on entropy and counting,” *Proceedings of the 1st Lake Michigan Workshop on Combinatorics and Graph Theory*, Kalamazoo, MI, USA, March 2014. <https://doi.org/10.48550/arXiv.1406.7872>
- [13] D. Gavinsky, S. Lovett, M. Saks and S. Srinivasan, “A tail bound for read- k families of functions,” *Random Structures and Algorithms*, vol. 47, no. 1, pp. 99–108, August 2015. <http://doi.org/10.1002/rsa.20532>
- [14] T. S. Han, “Nonnegative entropy measures of multivariate symmetric correlations,” *Information and Control*, vol. 36, no. 2, pp. 133–156, February 1978. [https://doi.org/10.1016/S0019-9958\(78\)90275-9](https://doi.org/10.1016/S0019-9958(78)90275-9)
- [15] P. Hell and J. Neštril, *Graphs and Homomorphisms*, Oxford Lecture Series in Mathematics and Its Applications, Oxford University Press, 2004. <https://doi.org/10.1093/acprof:oso/9780198528173.001.0001>
- [16] S. Jukna, *Extremal Combinatorics with Applications in Computer Science*, second edition, Springer, 2011. <https://doi.org/10.1007/978-3-642-17364-6>
- [17] J. Kahn, “An entropy approach to the hard-core model on bipartite graphs,” *Combinatorics, Probability and Computing*, vol. 10, no. 3, pp. 219–237, May 2001. <https://doi.org/10.1017/S0963548301004631>
- [18] J. Kahn, “Entropy, independent sets and antichains: a new approach to Dedekind’s problem,” *Proceedings of the American Mathematical Society*, vol. 130, no. 2, pp. 371–378, June 2001. <https://doi.org/10.1090/S0002-9939-01-06058-0>
- [19] N. Keller and N. Lifshitz, “A note on large H -intersecting families,” *SIAM Journal on Discrete Mathematics*, vol. 33, no. 1, pp. 398–401, 2019. <https://doi.org/10.1137/18M1220765>
- [20] L. Lovász, “On the Shannon capacity of a graph,” *IEEE Transactions on Information Theory*, vol. 25, no. 1, pp. 1–7, January 1979. <https://doi.org/10.1109/TIT.1979.1055985>
- [21] L. Lovász, *Graphs and Geometry*, American Mathematical Society, volume 65, 2019. <https://doi.org/10.1090/coll/065>
- [22] L. Lovász, *Large Networks and Graph Limits*, American Mathematical Society, 2012. <https://doi.org/10.1090/coll/060>
- [23] M. Madiman, A. W. Marcus and P. Tetali, “Entropy and set cardinality inequalities for partition-determined functions,” *Random Structures and Algorithms*, vol. 40, no. 4, pp. 399–424, July 2012. <https://doi.org/10.1002/rsa.20385>

- [24] M. Madiman and P. Tetali, “Information inequalities for joint distributions, interpretations and applications,” *IEEE Transactions on Information Theory*, vol. 56, no. 6, pp. 2699–2713, June 2010. <https://doi.org/10.1109/TIT.2010.2046253>
- [25] N. Pippenger, “An information-theoretic method in combinatorial theory,” *Journal of Combinatorial Theory, Series A*, vol. 23, no. 1, pp. 99–104, July 1977. [https://doi.org/10.1016/0097-3165\(77\)90083-8](https://doi.org/10.1016/0097-3165(77)90083-8)
- [26] N. Pippenger, “Entropy and enumeration of Boolean functions,” *IEEE Transactions on Information Theory*, vol. 45, no. 6, pp. 2096–2100, September 1999. <https://doi.org/10.1109/18.782146>
- [27] J. Radhakrishnan, “An entropy proof of Bregman’s theorem,” *Journal of Combinatorial Theory, Series A*, Elsevier Science, vol. 77, no. 1, pp. 161–164, January 1997. <https://doi.org/10.1006/jcta.1996.2727>
- [28] J. Radhakrishnan, “Entropy and counting,” *Proceedings of the IIT Kharagpur, Golden Jubilee Volume on Computational Mathematics, Modelling and Algorithms*, Narosa Publishers, New Delhi, India, pp. 1–25, 2001. <https://www.tcs.tifr.res.in/~jaikumar/Papers/EntropyAndCounting.pdf>
- [29] I. Sason, “A generalized information-theoretic approach for bounding the number of independent sets in bipartite graphs,” *Entropy*, vol. 23, no. 3, paper 270, pp. 1–14, 2021. <https://doi.org/10.3390/e23030270>
- [30] I. Sason, “Information inequalities via submodularity, and a problem in extremal graph theory,” *Entropy*, vol. 24, no. 5, paper 597, pp. 1–31, 2022. <https://doi.org/10.3390/e24050597>
- [31] I. Sason, “Observations on the Lovász ϑ -function, graph capacity, eigenvalues, and strong products,” *Entropy*, vol. 25, no. 1, paper 104, pp. 1–40, January 2023. <https://doi.org/10.3390/e25010104>
- [32] I. Sason, “Observations on graph invariants with the Lovász ϑ -function,” *AIMS Mathematics*, vol. 9, no. 6, pp. 15385–15468, April 2024. <https://doi.org/10.3934/math.2024747>
- [33] I. Sason, “Combinatorial applications of the Shearer and Han inequalities in graph theory and Boolean functions,” *Workshop on Information Theory, Boolean Functions, and Lattice Problems*, Hausdorff Research Institute for Mathematics (HIM), Bonn, Germany, November 18–22, 2024. The recorded talk is available at <https://www.youtube.com/watch?v=D-poDm7AnLU>
- [34] M. Simonovits and V. T. Sós, “Intersection theorems for graphs,” *Problèmes combinatoires et théorie des graphes (Colloques internationaux du Centre National de la Recherche Scientifique (CNRS))*, University of Orsay, Orsay, France, July 1976), pp. 389–391, *Colloques internationaux du CNRS*, no. 260, Paris, France, 1978. <https://users.renyi.hu/~miki/OrsayB.pdf>
- [35] M. Simonovits and V. T. Sós, “Intersection theorems on structures,” *Annals of Discrete Mathematics*, vol. 6, pp. 301–313, 1980. [https://doi.org/10.1016/S0167-5060\(08\)70715-5](https://doi.org/10.1016/S0167-5060(08)70715-5)
- [36] C. E. Shannon, “The zero error capacity of a noisy channel,” *IEEE Transactions on Information Theory*, vol. 2, no. 3, pp. 8–19, September 1956. <https://doi.org/10.1109/TIT.1956.1056798>
- [37] Z. Wang, J. Tu, and R. Lang, “Entropy, graph homomorphisms, and dissociation sets,” *Entropy*, vol. 25, paper 163, pp. 1–11, January 2023. <https://www.mdpi.com/1099-4300/25/1/163>
- [38] Y. Zhao, *Graph Theory and Additive Combinatorics: Exploring Structures and Randomness*, Cambridge University Press, 2023. <https://doi.org/10.1017/9781009310956>