

YAMADA-WATANABE UNIQUENESS RESULTS FOR SPDES DRIVEN BY WIENER AND PURE JUMP PROCESSES

KISTOSIL FAHIM, ERIKA HAUSENBLAS, AND KENNETH H. KARLSEN

ABSTRACT. The Yamada-Watanabe theory provides a robust framework for understanding stochastic equations driven by Wiener processes. Despite its comprehensive treatment in the literature, the applicability of the theory to SPDEs driven by Poisson random measures or, more generally, Lévy processes remains significantly less explored, with only a handful of results addressing this context. In this work, we leverage a result by Kurtz to demonstrate that the existence of a martingale solution combined with pathwise uniqueness implies the existence of a unique strong solution for SPDEs driven by both a Wiener process and a Poisson random measure. Our discussion is set within the variational framework, where the SPDE under consideration may be nonlinear. This work is influenced by earlier research conducted by the second author alongside de Bouard and Ondreját.

1. INTRODUCTION

The development of the Yamada-Watanabe uniqueness theory for stochastic equations primarily focuses on those driven by Wiener processes. The literature becomes more sparse when considering equations influenced by both Wiener noise and Poisson random measures, particularly in the infinite-dimensional noise case. Notably, in [4], the theory is extended to stochastic differential equations driven by this dual noise setup on a locally compact space, using the original Yamada and Watanabe method. Similarly, [31] applies the theory to variational solutions of PDEs driven solely by a Poisson random measure, again on a locally compact space, employing the same basic methods. Reference [11] presents the theory within a semigroup framework, focusing exclusively on a Poisson random measure.

Our research aims to broaden this scope by investigating the Yamada-Watanabe theory on Banach spaces, accommodating both Poisson random measures and Wiener processes in infinite dimensions within a variational setting.

In their foundational work [30, 29], Yamada and Watanabe presented a proof leveraging the concept of a regular version of conditional probabilities. Their methodology has since proven to be exceptionally robust and versatile, finding applications well beyond stochastic differential equations. The development of an abstract Yamada-Watanabe theory by Kurtz in 2007 [20] marked an important

Date: January 7, 2025.

2020 Mathematics Subject Classification. Primary: 60H15; Secondary: 60G57.

Key words and phrases. Yamada-Watanabe theory, Wiener processes, Lévy processes, pathwise uniqueness, pure jump process, SPDEs in variational form.

This work was supported by the project Pure Mathematics in Norway, funded by Trond Mohn Foundation and Tromsø Research Foundation, and by the Research Council of Norway under project 351123 (NASTRAN). Erika Hausenblas and Kistosil Fahim was supported by the ASEA network of the Austrian Exchange Service.

expansion of the theory to a broad array of stochastic problems. Kurtz's original argument, rooted in the Skorokhod representation theorem, abstracted the Yamada-Watanabe principle to a new level of generality. It was further explored by Kurtz in [19].

Applying Kurtz's abstract framework, Bouard, Hausenblas, and Ondrejat in [11] established that for stochastic evolution equations driven by a Poisson random measure, the pathwise uniqueness and the existence of a martingale solution implies the existence of a unique, strong solution. Our work seeks to advance this line of inquiry by adapting these concepts to the variational framework and by incorporating also a Wiener process into the analysis. Our initial motivation arose from the requirement in [14] to establish specific uniqueness results for a Lévy-driven stochastic bidomain model in electrophysiology. However, given its broad applicability, we believe that a readily citable result of this kind would be valuable in different contexts, which led us to write this paper.

The remainder of this paper is structured as follows: In Section 2, we review key concepts from stochastic analysis, including cylindrical Wiener processes, Poisson random measures, and Lévy processes. Sections 3 and 4 are dedicated to the development of the Yamada-Watanabe theory. In Section 3, we explain Kurtz's abstract framework within the context of SPDEs, while Section 4 demonstrates how weak existence and pathwise uniqueness together imply uniqueness in law.

2. PRELIMINARY MATERIAL

Before going into the stochastic preliminaries, let us first establish some notations that will be utilized throughout this paper. We denote the set of real numbers as \mathbb{R} , with $\mathbb{R}^+ := \{x \in \mathbb{R} : x > 0\}$ and $\mathbb{R}_0^+ := \mathbb{R}^+ \cup \{0\}$. The set of natural numbers, including 0, is denoted by \mathbb{N} . If $(\mathcal{F}_t)_{t \in [0, T]}$ represents a filtration and θ is a measure, we use \mathcal{F}_t^θ to denote the augmentation of \mathcal{F}_t with the θ -null sets contained in \mathcal{F}_T^θ .

A measurable space (S, \mathcal{S}) is called *Polish* if there exists a metric ϱ on S such that (S, ϱ) is a complete separable metric space and $\mathcal{S} = \mathcal{B}(S)$ (i.e., \mathcal{S} = the Borel sets of S with respect to ϱ). For a Polish space (S, \mathcal{S}) , we denote by $\mathcal{B}(S)$ the set of all Borel measurable mappings $F : S \rightarrow \mathbb{R}$ for which F is $\mathcal{B}(S)/\mathcal{B}(\mathbb{R})$ measurable. The collection of all finite non-negative measures on a Polish space (S, \mathcal{S}) is denoted by $M^+(S)$. Additionally, $\mathcal{P}(S)$ represents the set of probability measures on S . As S is a separable metric space, $\mathcal{P}(S)$ can be metrized as a separable metric space [24, Theorem 6.2, page 43]. Moreover, $\mathcal{P}(S)$ is a compact metric space if and only if S is compact [24, Theorem 6.4, page 45].

If a family of sets $\{S_n \in \mathcal{S} : n \in \mathbb{N}\}$ satisfies $S_n \uparrow S$, then $M_{\mathbb{N}}(\{S_n\})$ denotes the family of all $\mathbb{N} \cup \{\infty\}$ -valued measures μ on \mathcal{S} such that $\mu(S_n) < \infty$ for every $n \in \mathbb{N}$. By $\mathcal{M}_{\mathbb{N}}(\{S_n\})$, we denote the σ -algebra on $M_{\mathbb{N}}(\{S_n\})$ generated by the functions $i_B : M_{\mathbb{N}}(\{S_n\}) \ni \mu \mapsto \mu(B) \in \mathbb{N}$, $B \in \mathcal{S}$. The following simple result can be proven directly by constructing the metric.

Lemma 2.1. *Let (S, \mathcal{S}) be a Polish space and consider a family $\{S_n \in \mathcal{S}\}$ of sets satisfying $S_n \uparrow S$. Then $(M_{\mathbb{N}}(\{S_n\}), \mathcal{M}_{\mathbb{N}}(\{S_n\}))$ is a Polish space.*

Proof. Fix any $n \in \mathbb{N}$. Let $\mu \in M_{\mathbb{N}}^+(\{S_n\})$ be a mapping $\mu : \mathcal{B}(S_n) \rightarrow \mathbb{N}$ that is measurable and satisfies $\mu(S_n) < \infty$, where the $+$ superscript indicates the positive elements in $M_{\mathbb{N}}(\{S_n\})$. Define $F := \{f_n \in C(S) : n \in \mathbb{N}\}$ as a family of functions such that $|f_n|_{C(S)} \leq 1$ and F separates points in S . Here, $C(S)$ denotes the set of

continuous real-valued functions defined on S , equipped with the supremum norm $\|f\|_{C(S)}$. By the Hahn-Banach Theorem, and given that $\mathcal{P}(S)$ is separable, there exists such a countable set that separates points. Additionally, let $\{\lambda_n : n \in \mathbb{N}\}$ be such that $\sum_{n \in \mathbb{N}} \lambda_n < \infty$. Now, define the following metric for $\mu_1, \mu_2 \in M_{\mathbb{N}}^+(S_n)$:

$$d_{S_n}(\mu_1, \mu_2) := \sum_{n \in \mathbb{N}} \lambda_n \frac{|\langle \mu_1, f_n \rangle - \langle \mu_2, f_n \rangle|}{1 + |\langle \mu_1, f_n \rangle - \langle \mu_2, f_n \rangle|}.$$

This defines a metric on $M_{\mathbb{N}}^+(S_n)$. For a measure $\mu : \mathcal{M}(\{S_n\}) \rightarrow \mathbb{R}_0^+$, define $\mu|_{S_n}(A) := \mu(A \cap S_n)$ for $A \in \mathcal{B}(S)$. Next, we define a metric on $M_{\mathbb{N}}^+(\{S_n\})$:

$$d_S(\mu_1, \mu_2) := \sum_{n \in \mathbb{N}} \lambda_n \frac{d_{S_n}(\mu_1|_{S_n}, \mu_2|_{S_n})}{1 + d_{S_n}(\mu_1|_{S_n}, \mu_2|_{S_n})}, \quad \mu_1, \mu_2 \in M_{\mathbb{N}}^+(\{S_n\}).$$

It is straightforward to verify that d_S defines a metric on $M_{\mathbb{N}}^+(\{S_n\})$. It remains to show that this metric is complete and that $M_{\mathbb{N}}(\{S_n\})$ is separable.

Note that if $\mu^k, \mu \in M_{\mathbb{N}}^+(\{S_n\})$ and $\mu^k \rightarrow \mu$ in the weak topology, then we have $\mu^k|_{S_n}(f) \rightarrow \mu|_{S_n}(f)$ for each f . Hence, $d_{S_n}(\mu^k|_{S_n}, \mu|_{S_n}) \rightarrow 0$, and therefore $d_S(\mu^k, \mu) \rightarrow 0$. Finally, separability follows again from the Hahn-Banach Theorem. \square

Let us transition to the stochastic framework. All stochastic processes are defined on a filtered probability space. Throughout the paper, we denote this filtered probability space by

$$\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P}), \quad (2.1)$$

where \mathbb{P} is complete on (Ω, \mathcal{F}) , and $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ is a filtration satisfying the *usual conditions*:

- (i) for each $t \in [0, T]$, \mathcal{F}_t contains all $(\mathcal{F}, \mathbb{P})$ -null sets;
- (ii) the filtration \mathbb{F} is right-continuous.

A Lévy process encompasses both a cylindrical Wiener process and a pure jump process, for both of which we will provide detailed definitions. To understand the characteristics of a pure jump process, we utilize Poisson random measures.

2.1. The cylindrical Wiener process. Let \mathcal{H} denote a separable Hilbert space. Consider \mathbb{W} as a cylindrical Wiener process evolving over \mathcal{H} , defined on \mathfrak{A} . By using the spectral decomposition theorem, the Wiener process can be expressed as

$$\mathbb{W}(t) := \int_0^t \sum_{k=1}^{\infty} h_k d\beta_k(t), \quad t \geq 0,$$

where $\{\beta_k : k \in \mathbb{N}\}$ is a family of mutually independent Brownian motions, and $\{h_k : k \in \mathbb{N}\}$ forms an orthonormal basis in \mathcal{H} . Referencing Proposition 4.7 in [10, p. 85], we note that this representation can be considered without loss of generality.

To facilitate our later discussions, let us introduce the concept of Hilbert-Schmidt operators between two Hilbert spaces \mathcal{H} and H . We define the space of all Hilbert-Schmidt operators $\mathcal{L}_{\text{HS}}(\mathcal{H}, H)$ as follows:

$$\mathcal{L}_{\text{HS}}(\mathcal{H}, H) := \left\{ L : \mathcal{H} \rightarrow H : \sum_{i=1}^{\infty} |Le_i|_H^2 = \sum_{i,k=1}^{\infty} |\langle Le_i, f_k \rangle|^2 < \infty \right\},$$

where (e_i) and $(f_i)_i$ denote orthonormal bases in \mathcal{H} and H , respectively.

2.2. Poisson random measures and Lévy processes. To begin, we revisit [17, Definition I.8.1]. Given the diverse and not always consistent approaches to defining a time-homogeneous Poisson random measure in the literature, we present our own definition for clarity and precision in this context.

Definition 2.2 ([17, Definition I.8.1]). *Let (S, \mathcal{S}) be a Polish space and ν be a σ -finite measure on (S, \mathcal{S}) , where σ -finite means that there exists a nested sequence $\{S_n \in \mathcal{S}\}_{n \in \mathbb{N}}$ such that $S_n \uparrow S$ and $\nu(S_n) < \infty$ for every $n \in \mathbb{N}$.*

A time homogenous Poisson random measure η over a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$, is a measurable function

$$\eta : (\Omega, \mathcal{F}) \rightarrow (M_{\mathbb{N}}(\{S_n \times [0, T]\}), \mathcal{M}_{\mathbb{N}}(\{S_n \times [0, T]\})),$$

such that

- (i) for each $B \in \mathcal{S} \otimes \mathcal{B}([0, T])$ with $\mathbb{E} \eta(B) < \infty$, $\eta(B) := i_B \circ \eta : \Omega \rightarrow \mathbb{N}^1$ is a Poisson random variable with parameter $\mathbb{E} \eta(B)$, otherwise $\eta(B) = \infty$ a.s.
- (ii) η is independently scattered, i.e., if the sets $B_j \in \mathcal{S} \otimes \mathcal{B}([0, T])$, $j = 1, \dots, n$, are disjoint, then the random variables $\eta(B_j)$, $j = 1, \dots, n$, are mutually independent;
- (iii) for each $U \in \mathcal{S}$, the \mathbb{N} -valued process $(N(t, U))_{t \in [0, T]}$ defined by

$$N(t, U) := \eta(U \times (0, t]), \quad t \in [0, T]$$

is \mathbb{F} -adapted, and its increments are stationary and independent of the past, i.e., if $t > s \geq 0$, $N(t, U) - N(s, U) = \eta(U \times (s, t])$ is independent of \mathcal{F}_s .

Remark 2.3. *If ν is a finite measure on a Polish space (S, \mathcal{S}) , then for any $U \in \mathcal{S}$, $N(t, U)$ is a Poisson-distributed random variable with parameter $t\nu(U)$. In particular, the number of jumps is finite. However, if ν is only σ -finite and $\nu(S_n) \rightarrow \infty$ as $n \rightarrow \infty$, then $N(t, S) = \infty$, meaning there are infinitely many jumps within any time interval $[t_1, t_2]$, where $t_1 < t_2$.*

In Definition 2.2, the assignment

$$\nu : \mathcal{S} \ni A \mapsto \mathbb{E}[\eta(A \times (0, 1))] \tag{2.2}$$

defines a uniquely determined measure, called the *intensity measure* of the Poisson random measure η . Moreover, it turns out that the *compensator* γ of η is uniquely determined by

$$\gamma : \mathcal{S} \times \mathcal{B}([0, T]) \ni (A, I) \mapsto \nu(A) \times \text{Leb}_{[0, T]}(I),$$

where $\text{Leb}_{[0, T]}$ denotes the Lebesgue measure on $[0, T] \subset \mathbb{R}$. The difference between a time-homogeneous Poisson random measure η and its compensator γ , i.e.,

$$\tilde{\eta} := \eta - \gamma,$$

is called the *time-homogeneous compensated Poisson random measure*.

Remark 2.4. *Fix $n \in \mathbb{N}$. If S_n is a metric space, then the process*

$$\Omega \times [0, T] \ni (\omega, t) \mapsto \eta(\cdot \times [0, t])(\omega) \in M_{\mathbb{N}}(S_n)$$

is càdlàg and belongs to $\mathbb{D}([0, T]; M_{\mathbb{N}}(S_n))$.

If $S_n \subset E$, E is a Banach space of type p , $1 \leq p \leq 2$, and

$$\lim_{n \rightarrow \infty} \int_{S_n} |z|^p \nu(dz) < \infty,$$

¹Here, the mapping i_B is defined by $i_B : M_{\mathbb{N}}(\{S_n\}) \ni \mu \mapsto \mu(B) \in \mathbb{N}$, $B \in \mathcal{S}$.

then

$$L : \Omega \times [0, T] \ni (\omega, t) \mapsto \int_S z \tilde{\eta}(\cdot \times [0, t])(\omega) \in E$$

is càdlàg and belongs to $\mathbb{D}([0, T]; E)$.

Poisson random measures arise in a natural way by means of a Lévy process.

Definition 2.5. *Let E be a Banach space. A stochastic process $\{L(t) : t \geq 0\}$ is a Lévy process if the following conditions are met:*

- for any $l \in \mathbb{N}$ and $0 \leq t_0 < t_1 < \dots < t_l$, the random variables $L(t_0)$, $L(t_1) - L(t_0)$, \dots , $L(t_l) - L(t_{l-1})$ are independent;
- $L_0 = 0$ a.s.;
- For all $0 \leq s < t$, the distribution of $L(t+s) - L(s)$ does not depend on s ;
- L is stochastically continuous;
- the trajectories of L are a.s. càdlàg on E .

The characteristic function of a Lévy process is uniquely determined by the Lévy-Khinchin formula. Before describing this formula, let us first introduce the concept of the Lévy measure ν .

Definition 2.6 ([21, Chapter 5.4]). *Let E be a separable Banach space with dual E' . A symmetric σ -finite Borel measure λ on E is called a Lévy measure if and only if (i) $\lambda(\{0\}) = 0$ and (ii) the function²*

$$E' \ni a \mapsto \exp \left(\int_E (\cos \langle x, a \rangle - 1) \lambda(dx) \right)$$

is a characteristic function of a Radon measure on E [21, p. 17].

In the measure space $(E, \mathcal{B}(E))$, a σ -finite Borel measure λ is termed a Lévy measure if its symmetric part $\frac{1}{2}(\lambda + \lambda^-)$, where $\lambda^-(A) = \lambda(-A)$ for $A \in \mathcal{B}(E)$, qualifies as a Lévy measure. The collection of all Lévy measures on $(E, \mathcal{B}(E))$ is denoted by $\mathcal{L}(E)$.

For the definition of Banach spaces of (Rademacher) type p , where $p \in [1, 2]$, see [16, p. 54] and [21, p. 40]. Common examples of such Banach spaces include L^q spaces with $p \leq q < \infty$, defined over bounded domains, as well as the corresponding Besov spaces (see [6] for further details). Moreover, if a Banach space E is of type p , then it is also of type q for all $q \geq p$.

The Lévy-Khintchine formula establishes that for any E -valued Lévy process $\{L(t) : t \geq 0\}$, there exist a positive operator $Q : E' \rightarrow E$, a non-negative measure ν concentrated on $E \setminus \{0\}$ with the property that $\int_E 1 \wedge |z|_E^p \nu(dz) < \infty$, and an element $m \in E$ such that (see [1, 2] or [21, Theorem 5.7.3])

$$\begin{aligned} \mathbb{E} \left[e^{i \langle L(1), x \rangle} \right] &= \exp \left(i \langle m, x \rangle - \frac{1}{2} \langle Qx, x \rangle \right. \\ &\quad \left. - \int_E \left(1 - e^{i \langle y, x \rangle} + \mathbf{1}_{(-1,1)}(|y|_E) i \langle y, x \rangle \right) \nu(dy) \right), \end{aligned} \tag{2.3}$$

²As remarked in [21, Chapter 5.4], we do not need to suppose that $\int_E (\cos \langle x, a \rangle - 1) \lambda(dx)$ is finite. However, if λ is a symmetric Lévy measure, then, for each $a \in E'$, the integral is finite, see Corollary 5.4.2 in [21].

for each $x \in E'$. We refer to the measure ν as the *Lévy measure* of the Lévy process L . Furthermore, the triplet (Q, m, ν) uniquely characterizes the law of L .

We now proceed to construct a Poisson random measure whose intensity measure is determined by a Lévy measure ν . Let \mathfrak{A} be a filtered probability space, see (2.1), and let E be a p -stable Banach space for some $p \in [1, 2]$. Consider an E -valued Lévy process $\{L(t) : t \geq 0\}$, defined on \mathfrak{A} , which is of pure jump type³ with Lévy measure ν . To this process, we associate a counting measure η_L , also defined on \mathfrak{A} , as follows:

$$\mathcal{B}(E) \times \mathcal{B}((0, T]) \ni (B, I) \mapsto \eta_L(B \times I) := \#\{s \in I \mid \Delta_s L \in B\} \in \mathbb{N}.$$

Here, the jump process $\Delta L = \{\Delta_t L : 0 \leq t < \infty\}$ linked to L is defined by

$$\Delta_t L := L(t) - L(t-) = L(t) - \lim_{\varepsilon \rightarrow 0} L(t - \varepsilon), \quad t > 0, \quad \Delta_0 L = 0.$$

If ν is symmetric and supported within the unit ball, then η_L function as a time-homogeneous Poisson random measure with intensity measure ν . Furthermore, we can express $L(t)$ through the integral

$$L(t) = \int_0^t \int_Z z \tilde{\eta}_L(dz, ds), \quad t \geq 0,$$

where $\tilde{\eta}_L$ denotes the compensated version of η_L .

Conversely, when given a time-homogeneous Poisson random measure η on a p -stable Banach space E , with $p \in [1, 2]$, it is possible to construct a Lévy process L . The integral

$$\int_I \int_Z z \tilde{\eta}(dz, ds)$$

is well-defined for any $I \in \mathcal{B}([0, T])$ if and only if the intensity measure ν of η is a Lévy measure, see [12, p. 123, Theorem (2.1)].

For further information about the connection between Poisson random measures and Lévy processes, we direct the reader to Applebaum [1], Ikeda and Watanabe [17], and Peszat and Zabczyk [25].

The formulation of stochastic integrals within arbitrary Banach spaces present notable challenges depending significantly on the geometric structure of the space [28]. Our discussion in what follows is therefore confined to UMD Banach spaces of type $p \in [1, 2]$ (and thus of martingale type p), where the p depends on the integrability properties of the specific Lévy measure being used. Standard examples of Banach spaces with martingale type $p \in [1, 2]$ include Hilbert spaces (martingale type 2), L^p spaces, and uniformly convex spaces (which have martingale type 2). We refer to [15, Chapters 3.5 & 4] for details.

We define the space of possible integrands as follows:

$$\mathcal{M}^p([0, T]; L^p(Z, \nu; E)) := \left\{ \xi : [0, T] \times \Omega \rightarrow L^p(Z, \nu; E) \mid \right. \\ \left. \xi \text{ is progressively measurable and } \mathbb{E} \int_0^T |\xi(t)|_{L^p(Z, \nu; E)}^p dt < \infty \right\}.$$

In [7], the second author and Brzeźniak demonstrated the existence and uniqueness of a continuous linear operator that assigns to each progressively measurable process

³A Lévy process is of pure jump type if $Q = 0$.

$\xi \in \mathcal{M}^p([0, T]; L^p(Z, \nu; E))$ an adapted cádlág E -valued process. This stochastic integral process is denoted by

$$\int_0^\cdot \int_Z \xi(r, x) \tilde{\eta}(dx, dr),$$

and it satisfies the following: for a random step process $\xi \in \mathcal{M}([0, T], L^p(Z, \nu; E))$ with the form

$$\xi(r, z) = \sum_{j=1}^l \mathbf{1}_{(t_{j-1}, t_j]}(r) \xi_j(z), \quad r \geq 0,$$

where $\{0 = t_0 < t_1 < \dots < t_l\}$ is a finite partition of $[0, T]$ and $\xi_j(z)$ is an E -valued, $\mathcal{F}_{t_{j-1}}$ -measurable, p -summable random variable for each j , then

$$\int_0^t \int_Z \xi(r, z) \tilde{\eta}(dz, dr) = \sum_{j=1}^l \int_Z \xi_j(z) \tilde{\eta}(dz, (t_{j-1} \wedge t, t_j \wedge t]).$$

Furthermore, the operator

$$I : \mathcal{M}^p([0, T]; L^p(Z, \nu; E)) \ni \xi \mapsto \int_0^\cdot \int_Z \xi(r, x) \tilde{\eta}(dx, dr) \in \mathcal{M}^p([0, T]; E)$$

is continuous. That is, there exists a constant $C = C(E)$, independent of ξ , η , and ν , such that

$$\mathbb{E} \left| \int_0^t \int_Z \xi(r, z) \tilde{\eta}(dz, dr) \right|^p \leq C \mathbb{E} \int_0^t \int_Z |\xi(r, z)|^p \nu(dz) dr, \quad t \geq 0.$$

When working with a Poisson random measure, the associated solution process can be cádlág and predictable in one space while being only progressively measurable in another. This is why we make a distinction between cádlág behavior and predictability versus progressive measurability. For completeness, we provide a simple example to illustrate this point.

Example 2.7. *First, let us construct a space-time Poisson random measure with a prescribed intensity measure ν on \mathbb{R}^d (compare also with [25, Proposition 7.21]). To this end, let us specify ν_0 as a Lévy measure on $\mathbb{R} \setminus \{0\}$, assuming that for any $n \in \mathbb{N}$, the measure ν_0 is finite on $\mathbb{R} \setminus [-\frac{1}{n}, \frac{1}{n}]$ and $\int_{[-1, 1] \setminus \{0\}} |z|^2 \nu_0(dz) < \infty$. Set*

$$S_n := (\mathbb{R} \setminus [-\frac{1}{n}, \frac{1}{n}]) \times \mathbb{R}^d$$

and let us define a measure ν_n on S_n by

$$\nu_n(A \times B) := \nu_0(A) \text{Leb}(B), \quad A \in \mathcal{B}(\mathbb{R} \setminus [-\frac{1}{n}, \frac{1}{n}]), \quad B \in \mathcal{B}(\mathbb{R}^d),$$

where Leb denotes the Lebesgue measure on \mathbb{R}^d . To ensure that ν_n is well-defined as a measure on S_n , we note that the collection of rectangular sets $A \times B$, where $A \in \mathcal{B}(\mathbb{R} \setminus [-\frac{1}{n}, \frac{1}{n}])$ and $B \in \mathcal{B}(\mathbb{R}^d)$, forms a π -system. Since ν_n is finitely additive and satisfies the properties of a Dynkin system, it extends uniquely to the σ -algebra $\mathcal{B}(S_n)$. Introduce the σ -algebra $\mathcal{S} := \bigvee_{n \in \mathbb{N}} \mathcal{B}(S_n)$.⁴ We define the measure ν on \mathcal{S} as the unique extension of the sequence of measures ν_n defined on $\mathcal{B}(S_n)$ for $n \in \mathbb{N}$. This extension is well-defined since any set in \mathcal{S} can be approximated from below by its restrictions to the truncated domains S_n .

⁴Let Z be a set, and let \mathcal{A} and \mathcal{B} be collections of subsets of Z . We define $\mathcal{A} \vee \mathcal{B}$ as the smallest σ -algebra on X that contains every set in both \mathcal{A} and \mathcal{B} , that is, $\mathcal{A} \vee \mathcal{B}$ is generated by the union $\mathcal{A} \cup \mathcal{B}$: $\mathcal{A} \vee \mathcal{B} = \sigma(\mathcal{A} \cup \mathcal{B})$.

The space-time Poisson noise on $(\mathbb{R}^d, \mathcal{B}(\mathbb{R}^d))$ with jump intensity measure ν , defined over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, is a $\mathcal{F}/\mathcal{M}(M_{\mathbb{N}}(\{S_n \times [0, T]\}))$ -measurable mapping⁵

$$\eta : \Omega \rightarrow \mathcal{M}(M_{\mathbb{N}}(\{S_n \times [0, T]\})), \quad (2.4)$$

such that:

- For any $U \in \mathcal{S} \otimes \mathcal{B}([0, T])$ with $(\nu \times \text{Leb}_{[0, T]})(U) < \infty$, the random variable $\eta(U) := i_U \circ \eta$ is Poisson-distributed with parameter $(\nu \otimes \text{Leb}_{[0, T]})(U)$.
- If $U_1 \in \mathcal{S} \otimes \mathcal{B}([0, T])$ and $U_2 \in \mathcal{S} \otimes \mathcal{B}([0, T])$ are disjoint, then the random variables $\eta(U_1)$ and $\eta(U_2)$ are independent, and $\eta(U_1 \cup U_2) = \eta(U_1) + \eta(U_2)$ almost surely.

The space-time Poisson random measure can be represented as a function-valued Poisson process with a Poisson random measure on the space $E_0 := B_{2, \infty}^{-\frac{d}{2}}(\mathbb{R}^d)$.⁶ To verify this claim, let us first define the mapping

$$f_n : S_n \longrightarrow E_0 : (z, x) \mapsto f_n(z, x) = z \delta_x,$$

which maps points $(z, x) \in S_n$ to elements of the function space E_0 . The set

$$E_n := \{f_n(z, x) \mid (z, x) \in S_n\} \subset E_0,$$

represents the image of S_n under f_n , for each $n \in \mathbb{N}$. Since E_0 is not of type 2, we embed E_0 into the (type 2) Besov space $E := B_{2, 2}^{-\gamma}(\mathbb{R}^d)$, where $\gamma > \frac{d}{2}$ is arbitrary. For each $n \in \mathbb{N}$, the measure μ_n on E is then defined as the pushforward of the measure ν_n under f_n , i.e.,

$$\mu_n(B) := \nu_n(f_n^{-1}(B \cap E_n)), \quad B \in \mathcal{B}(E).$$

It is straightforward to verify that μ_n converges to a Lévy measure on E as $n \rightarrow \infty$. With a slight abuse of notation, we will denote the limit measure by ν .

Now, let us consider the solutions ξ and ξ_n , $n \in \mathbb{N}$, to the following SPDEs:

$$(*) \quad d\xi(t) - \Delta \xi(t) dt = \int_E z \tilde{\eta}(dz, dt), \quad (*)_n \quad d\xi_n(t) - \Delta \xi_n(t) dt = \int_{E_n} z \tilde{\eta}(dz, dt),$$

where $\tilde{\eta}$ is the compensator of the random measure η defined in (2.4), and we use additive noise for simplicity of presentation. It follows that

$$[0, T] \ni t \mapsto \xi(t) = \int_0^t \int_E e^{-(t-s)\Delta} z \tilde{\eta}(dz, ds)$$

and

$$[0, T] \ni t \mapsto \xi_n(t) = \int_0^t \int_{E_n} e^{-(t-s)\Delta} z \tilde{\eta}(dz, ds),$$

are the unique solutions to $(*)$ and $(*)_n$, respectively, where $(e^{\Delta t})_{t \geq 0}$ denotes the heat semigroup. Consequently, the processes ξ and $\{\xi_n : n \in \mathbb{N}\}$ are càdlàg in the “large space” $E = B_{2, 2}^{-\gamma}(\mathbb{R}^d)$. More precisely, the process ξ_n is càdlàg in E , and

⁵Let (X_i, \mathcal{X}_i) , $i = 1, 2$, be two measurable spaces. A function $f : X_1 \rightarrow X_2$ is $\mathcal{X}_1/\mathcal{X}_2$ -measurable if, for all $A \in \mathcal{X}_2$, the pre-image $f^{-1}(A) := \{x \in X_1 : f(x) \in A\}$ belongs to \mathcal{X}_1 .

⁶The choice of E_0 is motivated by the fact that the Dirac measure δ belongs to the Besov space $B_{2, \infty}^{-\frac{d}{2}}(\mathbb{R}^d)$ ($= E_0$) (see [27, p. 34]). Since the Besov embedding $B_{2, 2}^{-\frac{d}{2}}(\mathbb{R}^d) \hookrightarrow B_{2, \infty}^{-\frac{d}{2}}(\mathbb{R}^d)$ holds (see [27, p. 30]), the Dirac measure δ does not belong to the Sobolev space $H_2^{-\frac{d}{2}}(\mathbb{R}^d)$, which coincides with the Lizorkin-Triebel space $F_{2, 2}^{-\frac{d}{2}}(\mathbb{R}^d)$ and the Besov space $B_{2, 2}^{-\frac{d}{2}}(\mathbb{R}^d)$. However, δ belongs to the Sobolev space $H_2^\gamma(\mathbb{R}^d)$ for $\gamma < -\frac{d}{2}$.

for all $t \in [0, T]$, $\lim_{s \uparrow t} \xi_n(s)$ is predictable. Moreover, the process $[0, T] \ni t \mapsto \lim_{s \uparrow t} \xi(s)$ is also càdlàg and predictable in this space.

In the “smaller space” setting of $E_0 = B_{2, \infty}^{-\frac{d}{2}}(\mathbb{R}^d)$, the processes ξ_n remain càdlàg, due to the finiteness of the Lévy measure. In particular, \mathbb{P} -a.s., $\xi_n \in \mathbb{D}([0, T]; E_0)$. In addition, for $\tilde{E} := B_{2, 2}^{-\frac{d}{2}}(\mathbb{R}^d) \subset E$,

$$\sup_{0 \leq t \leq T} \mathbb{E} |\xi_n(t)|_{\tilde{E}}^2 < \infty, \quad n \in \mathbb{N}.$$

However, ξ is not càdlàg in E_0 , as the space is not of type p . Nevertheless, we have

$$\mathbb{E} \int_0^T |\xi(s)|_{\tilde{E}}^2 ds < \infty,$$

and ξ is progressively measurable in \tilde{E} . In particular, for all $s \in [-\frac{d}{2}, \frac{d}{2} + 1)$, there exists a constant $C > 0$, independent of $n \in \mathbb{N}$, such that

$$\mathbb{E} \int_0^T |\xi_n(s)|_{B_{2, 2}^{\frac{d}{2}}}^2 ds \leq C, \quad \mathbb{E} \int_0^T |\xi(s)|_{B_{2, 2}^{\frac{d}{2}}}^2 ds \leq C.$$

In closing, let us mention that to establish the progressive measurability of a specific version of ξ_n , where $n \in \mathbb{N}$, one may employ the k -th order shifted Haar projection, denoted as $\xi_n^{(k)}$. This approach use piecewise constant approximations of ξ_n to construct a sequence $(\xi_n^{(k)})_{k \in \mathbb{N}}$ that is predictable. Moreover, one can show that as $k \rightarrow \infty$,

$$\xi_n^{(k)} \rightarrow \xi_n \quad \text{in } L^2(\Omega; L^2([0, T], \tilde{E})).$$

For a proof, see [9, Appendix C].

3. KURTZ’S FRAMEWORK

Our goal is to extend the Yamada-Watanabe theorem, which links the existence and uniqueness of weak and strong solutions to stochastic equations, to a broader class of Lévy-driven SPDEs using Kurtz’s abstract principles [19, 20]. Following [22] and [8], we revisit the variational SPDE framework and demonstrate how filtration and regularity conditions can be incorporated into Kurtz’s approach.

Let us turn our attention to the abstract framework introduced in [19] and [20]. Consider B_1 and B_2 , both metric spaces, and a Borel measurable function $\Gamma : B_1 \times B_2 \rightarrow \mathbb{R}$. Let there be a random variable Y (the input of the model), taking values in B_2 , with law ρ , and being defined on a given probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Our focus is on identifying a solution to the equation $\Gamma(X, Y) = 0$. Specifically, we seek a random variable X , taking values in B_1 and being defined over $(\Omega, \mathcal{F}, \mathbb{P})$, that satisfies

$$\Gamma(X, Y) = 0 \quad \text{in the sense that} \quad \mathbb{P}(\{\Gamma(X, Y) = 0\}) = 1. \quad (3.1)$$

We introduce the concept of a strong solution by defining it as follows:

Definition 3.1. *A pair (X, Y) constitutes a strong solution to the problem (3.1) defined by (Γ, ρ) , where $\text{Law}(Y) = \rho$, if there exists a Borel measurable function $F : B_1 \rightarrow B_2$ such that (3.1) holds with $X = F(Y)$, \mathbb{P} -a.s.*

If a strong solution exists over some probability space $(\Omega, \mathcal{F}, \mathbb{P})$, then (X, Y) has a joint distribution. In particular, there exists a probability measure

$$\mu : \mathcal{B}(B_1 \times B_2) \rightarrow [0, 1],$$

such that $\mu(B_1 \times A) = \rho(A)$ for all $A \in \mathcal{B}(B_2)$ and $\mathbb{P}(X \in A) = \int_{B_2} \mu(A, y) \rho(dy)$ for all $A \in \mathcal{B}(B_1)$. Let us denote this joint distribution of the two random variables X, Y by $\mu_{X,Y}$. If (X, Y) is a solution to (3.1), then $\mu_{X,Y}$ is determined by the distribution ρ of the input Y and the mapping F , see [19, Lemma 1.3].

To further elaborate on the problem from the perspective of probability laws, we are interested in finding a joint distribution—also called a joint solution measure— $\mu \in \mathcal{P}(B_1 \times B_2)$. This probability measure μ should satisfy $\mu(B_1 \times A) = \rho(A)$ for all measurable subsets $A \in \mathcal{B}(B_2)$, along with the requirement that:

$$\int_{B_1 \times B_2} |\Gamma(x, y)| \mu(dx, dy) = 0. \quad (3.2)$$

Given Γ and ρ , we define $\mathcal{S}_{\Gamma, \rho}$ as follows:

$$\begin{aligned} \mathcal{S}_{\Gamma, \rho} \text{ is the set of all joint solution measures } \mu \in \mathcal{P}(B_1 \times B_2) \\ \text{for which the constraint (3.2) is satisfied and } \mu(B_1 \times \cdot) = \rho. \end{aligned} \quad (3.3)$$

In this way, we can speak also of a weak solution to the problem (ρ, Γ) :

Definition 3.2. *A weak solution of the problem (Γ, ρ) is a pair of random variables (X, Y) defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ such that Y has distribution ρ and (X, Y) meets the constraints in Γ , that is, $\mu_{X,Y} \in \mathcal{S}_{\Gamma, \rho}$, see (3.3).*

Let us now consider the setting in which we encounter SPDEs in the variational framework. Let H be a separable Hilbert space with inner product $\langle \cdot, \cdot \rangle$ and let H' denote its dual. Consider V as a reflexive Banach space embedded continuously and densely into H . Through the Riesz isomorphism, which identifies H with H' , we establish the Gelfand triple

$$V \subset H \equiv H' \subset V' \text{ as continuous and dense.} \quad (3.4)$$

The duality pairing between V' and V , defined as

$${}_{V'} \langle z, v \rangle_V := z(v), \quad z \in V', \quad v \in V,$$

satisfies

$${}_{V'} \langle z, v \rangle_V = \langle z, v \rangle_H, \quad z \in H, \quad v \in V.$$

In the following, we consider a filtered probability space \mathfrak{A} as given by (2.1). Let \mathbb{W} be a cylindrical Wiener process on a given Hilbert space \mathcal{H} , expressed as

$$\mathbb{W}(t) = \sum_{k=1}^{\infty} h_k \beta_k(t), \quad t \in [0, T], \quad (3.5)$$

where $\{h_k \mid k \in \mathbb{N}\}$ is an orthonormal basis in \mathcal{H} and $\{\beta_k \mid k \in \mathbb{N}\}$ is a sequence of mutually independent Brownian motions defined over \mathfrak{A} .

Let η be a time-homogeneous Poisson random measure, with a given intensity measure ν defined on a given Polish space (S, \mathcal{S}) . It is assumed that η is independent of the Wiener process \mathbb{W} and defined over the same filtered probability space \mathfrak{A} . See Definition 2.2 for further details.

Moving forward, we are presented with the following measurable mappings:

- $b : [0, T] \times V \rightarrow V'$, mapping into the dual space of V ;
- $\sigma : [0, T] \times H \rightarrow \mathcal{L}_{\text{HS}}(\mathcal{H}, V')$, which maps into the space of Hilbert-Schmidt operators from \mathcal{H} to V' ;
- $c : [0, T] \times \{S_n\} \times V_d \rightarrow V'$, where the sets $\{S_n\}$ are defined in Lemma 2.1, and $V_d \subset V'$ is dense.

Let E_1 and E_2 be Banach spaces, with $E_2 \hookrightarrow E_1$ and $V \hookrightarrow E_1$. Suppose E_1 is a UMD space of type 2, which implies that E_1 is also of martingale type 2 [15].⁷ We consider general SPDEs of the form

$$\begin{aligned} dU(t) &= b(t, U) dt + \sigma(t, U(t)) dW(t) + \int_S c(t, z, U(t)) \tilde{\eta}(dz, dt), \\ U(0) &= U_0 \in E_2, \end{aligned} \quad (3.6)$$

where b , σ , and c are the ‘‘coefficients’’ given above. We refer to a stochastic process $U : \Omega \times [0, T] \rightarrow E_1$ as a solution of (3.6) if the equation

$$\begin{aligned} \langle U(t), \varphi \rangle &= \langle U_0, \varphi \rangle + \int_0^t \langle b(s, U(s)), \varphi \rangle ds \\ &+ \int_0^t \sum_{k=1}^{\infty} \langle \sigma(s, U(s)) [h_k], \varphi \rangle d\beta^k(s) + \int_0^t \int_S \langle c(s, z, U(s)), \varphi \rangle \tilde{\eta}(dz, ds), \end{aligned} \quad (3.7)$$

is satisfied \mathbb{P} -a.s., for each $t \in [0, T]$, and for each test function $\varphi \in V$.

We now formulate the above SPDE problem (3.6), (3.7) in an abstract setting. Before proceeding, let us recall that all random variables are defined over a filtered probability space \mathfrak{A} (see (2.1)). Let Y represent the Wiener process \mathbb{W} , the Poisson random measure ν , and the given initial condition, such that $Y = (\mathbb{W}, \nu, U_0)$. Furthermore, we introduce the following space linked to Y :

$$B_2 = C_b([0, T]; \mathcal{H}) \times M_{\mathbb{N}}(\{S_n \times [0, T]\}) \times E_2. \quad (3.8)$$

At the same time, in the abstract formulation under consideration, let X denote the solution variable U belonging to the Skorohod space of càdlàg functions with values in E_1 , which is denoted by

$$B_1 = \mathbb{D}([0, T]; E_1). \quad (3.9)$$

Example 3.3. *To demonstrate the applicability of our results, we give an example of an SPDE that fits within our framework (3.6), (3.7), along with a specification of the relevant spaces. Let⁸*

$$V := L^p(\mathcal{O}) \subset H := H_0^{1,2}(\mathcal{O}) \subset (L^p(\mathcal{O}))' =: V'$$

be a Gelfand triple equipped with the following scalar product

$${}_V \langle u, v \rangle_V := \int \langle \nabla u(x), \nabla v(x) \rangle dx \quad u, v \in H_0^{1,2}(\mathcal{O}).$$

Here, \mathcal{O} denotes a bounded open subset of \mathbb{R}^d . Let $b : V \rightarrow V'$ be the porous medium operator defined by

$$b(u) := \Delta (|u|^{p-2}u), \quad u \in L^p(\mathcal{O}), \quad p \geq 2.$$

Note that b is hemicontinuous, locally monotone, coercive, and bounded; see, for example, [3] or [22, p. 87].

Let $\mathcal{H} = \mathbb{R}^m$, and assume that $\{W_t\}_{t \geq 0}$ is an m -dimensional Wiener process on the filtered probability space (2.1), represented as in (3.5). Additionally, let η

⁷If one deals only with a Poisson random measure, it is also possible to consider a UMD Banach space of type p , with $p \in [1, 2]$, provided the small jumps are p -integrable. However, the stochastic Itô integral with respect to a Wiener process is defined only on a UMD Banach space of type 2.

⁸The dual is defined with respect to the Hilbert space $H_0^{1,2}(\mathcal{O})$.

denote the space-time Poisson random measure constructed in Example 2.7, with intensity measure $\nu_0((a, b)) = \int_a^b |z|^{-\alpha} e^{-|z|} dz$, for $a, b \in \mathbb{R}$ with $a < b$ and $a, b \neq 0$.

The example is now provided by the following SPDE:

$$dU(t) = b(U(t)) dt + \sigma(U(t)) dW(t) + \int_S c(z, U(t)) \tilde{\eta}(dz, dt),$$

where U_0 is an \mathcal{F}_0 -measurable random variable and $\sigma(u)[h] := uh$ for all $u \in V$ and $h \in \mathcal{H}$. Regarding the jump-noise amplitude c , we assume

$$c : B_{2,\infty}^{-\frac{d}{2}}(\mathcal{O}) \times H_0^{1,2}(\mathcal{O}) \ni (z, x) \mapsto x(I - \Delta)^{-\frac{d}{2}-2} z \in H_0^{1,2}(\mathcal{O}),$$

which implies that, in the abstract setup (3.6), the sets $\{S_n\}$ have been replaced by $B_{2,\infty}^{-\frac{d}{2}}(\mathcal{O})$ (see Example 2.7), and V_d has been replaced by $H_0^{1,2}(\mathcal{O})$. As $H_0^{1,2}(\mathcal{O}) \subset V'$, the noise operator c maps into V' , satisfying the required condition. Moreover, note that the coefficients b , σ , and c are assumed here to be independent of t .

Finally, to complete the identification with the abstract framework (3.6), let E_1 be a Banach space such that $V' \hookrightarrow E_1$ continuously; for example, $E_1 = H^{-\frac{d}{2}-1,2}(\mathcal{O})$, and define

$$B_1 = \mathbb{D}([0, T]; E_1),$$

Setting $E_2 = H_0^{1,2}(\mathcal{O})$ (for example), we may define the space B_2 as

$$B_2 = C_b([0, T]; \mathbb{R}^m) \times M_{\mathbb{N}}(\{\mathbb{R} \setminus [-\frac{1}{n}, \frac{1}{n}] \times \mathcal{O} \times [0, T]\}) \times H_0^{1,2}(\mathcal{O}).$$

Now, one must construct a mapping $\Gamma : B_1 \times B_2 \rightarrow \mathbb{R}$ such that the solution X of the abstract equation (3.1) coincides with the solution of the SPDE (3.6). In (3.1), the solution is defined as a random variable that satisfies a constraint given by the mapping $\Gamma : B_1 \times B_2 \rightarrow \mathbb{R}$. In our SPDE example, however, the solution is defined as a process that satisfies a collection of equations or constraints.

To be more precise, let $V_d := \{\varphi_k : k \in \mathbb{N}\}$ be a dense countable subset of V (see (3.4)), and let $\mathbb{Q}_T := \mathbb{Q} \cap [0, T]$. Then, for each $k \in \mathbb{N}$ and $t \in \mathbb{Q}_T$, we define a constraint $\Gamma_{\varphi_k, t} : B_1 \times B_2 \rightarrow \mathbb{R}$ based on (3.7) with $\varphi = \varphi_k$. Within a given probability space, see (2.1), we define

$$\Gamma = \{\Gamma_{\varphi, t} : \varphi \in V_d, t \in \mathbb{Q}_T\} \quad (3.10)$$

by

$$\begin{aligned} \Gamma_{\varphi, t}(U, (\mathbb{W}, \eta, U_0)) &= \langle U_0, \varphi \rangle + \int_0^t \langle b(s, U(s)), \varphi \rangle ds \\ &\quad + \int_0^t \sum_{k=1}^{\infty} \langle \sigma(s, U(s))[h_k], \varphi \rangle d\beta_k(s) \\ &\quad + \int_0^t \int_S \langle c(s, x, U(s)), \varphi \rangle \tilde{\eta}(dx, ds) \\ &\quad - \langle U(t), \varphi \rangle. \end{aligned} \quad (3.11)$$

Omitting the probability variable, we write $\Gamma_{\varphi, t}(x, y)$, where $\Gamma_{\varphi} : B_1 \times B_2 \rightarrow \mathbb{R}$ is defined in (3.11), and B_1 and B_2 are defined in (3.9) and (3.8), respectively. Here, x resides in B_1 , symbolizing the variable U , and y , located in B_2 , represents the input triplet (\mathbb{W}, η, U_0) . Whenever we want to emphasize the dependence on t , we write $\Gamma_{\varphi}(x, y)(t)$ instead of $\Gamma_{\varphi, t}(x, y)$.

Remark 3.4. We will consider probabilistic weak solutions, also known as martingale solutions. Typically, the probability space associated with these solutions does not coincide with the probability space fixed in (2.1); instead, the focus is primarily on the solution measure. To formulate the problem, it is sufficient to specify:

- the Hilbert space \mathcal{H} , where the cylindrical Wiener process is defined,
- the Polish space (S, \mathcal{S}) (along with a sequence $\{S_n\}$ of sets) and the intensity measure ν defined on (S, \mathcal{S}) , which characterizes the Lévy process,
- the distribution ρ_0 defined on E_2 , characterizing the initial condition.

Additionally, the coefficients b , σ , and c of the SPDE (3.6) are required. From \mathcal{H} , ν , and ρ_0 , one can construct a filtered probability space $(\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ with a cylindrical Wiener process \mathbb{W} on \mathcal{H} , a Poisson random measure η on S , and an initial condition U_0 (with law ρ_0). Here, U_0 is \mathcal{F}_0 -measurable, and η and \mathbb{W} are independent, both adapted to the filtration $(\mathcal{F}_t)_{t \in [0, T]}$.

After introducing the problem, we proceed to define the concept of a solution as it will be applied in the subsequent sections. Occasionally, we need to assume additional regularity properties that, while not part of the formal definition of the solution, are crucial for ensuring pathwise uniqueness. These regularity properties can be introduced through additional mappings:

$$\theta_i^{\alpha_i} : \mathbb{D}([0, T]; E_1) \rightarrow \mathbb{R}, \quad \alpha_i \in A_i, \quad i = 0, 1,$$

where A_i , $i = 0, 1$, are two index sets. We define

$$\mathcal{X} := \left\{ U \in B_1 \mid \mathbb{E} \theta_0^{\alpha_0}(U) \leq R, \mathbb{P}(\theta_1^{\alpha_1}(U) < \infty) = 1, \right. \\ \left. \text{for all } \alpha_i \in A_i, i = 0, 1 \right\}, \quad (3.12)$$

for some given $R > 0$. The set \mathcal{X} allows us to incorporate this additional information.

Remark 3.5. To demonstrate how additional regularity assumptions can be applied to the solution U through the mappings $\theta_0^{\alpha_0}$ and $\theta_1^{\alpha_1}$, let us consider an example. Specifically, define the first functional $\theta_0^{\alpha_0}(U)$ as

$$\theta_0^{\alpha_0}(U) := \|U\|_{L^2(0, T; V)}^2, \quad \forall \alpha_0.$$

Imposing the condition $\mathbb{E} \theta_0^{\alpha_0}(U) \leq R$, for some constant R , encodes a boundedness constraint on U . This approach facilitates the enforcement of regularity and additional bounds on the solutions, extending beyond the requirements of the solution concept but necessary for a well-posedness analysis.

Furthermore, non-negativity constraints can be incorporated using the second functional by defining, e.g.,

$$\theta_1^{\alpha_1}(U(t, x)) := \begin{cases} \infty & \text{if } U(t, x) < 0, \\ 0 & \text{elsewhere.} \end{cases}$$

We then require that the probability of the event $\theta_1^{\alpha_1}(U) < \infty$ is equal to one. In this manner, if, for instance, the solution space $E_1 = L^2(\mathcal{O})$ in (3.6) is used, where $\mathcal{O} \subset \mathbb{R}^d$ is bounded open, the set in (3.12) transforms into the measurable set:

$$\mathcal{X} := \left\{ U : [0, T] \rightarrow L^2(\mathcal{O}) \mid \mathbb{E} \|U\|_{L^2(0, T; L^2(\mathcal{O}))}^2 \leq R, \right. \\ \left. \mathbb{P}(\{\text{Leb}(\{(t, x) : U(t, x) < 0\}) = 0\}) = 1 \right\}.$$

We now present the precise definition of a solution within Kurtz's framework. We consider a Gelfand triple (V, H, V') , see (3.4), along with Banach spaces E_1 and E_2 . Here, E_2 is continuously embedded into E_1 , and V is continuously embedded into E_1 . Additionally, E_1 is a UMD space of type 2.

Definition 3.6. *Given a Hilbert space \mathcal{H} , an intensity measure ν over a Polish space (S, \mathcal{S}) , see (2.2) and (2.3), and a distribution ρ_0 on E_2 , we consider a tuple $(\mathfrak{A}, U, \mathbb{W}, \eta, U_0)$ to be a probabilistic weak solution of the SPDE (3.6), under the following conditions: The tuple consists of:*

- (i) *A filtered probability space $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ denotes the filtration.*
- (ii) *A cylindrical Wiener process \mathbb{W} on \mathcal{H} , defined over \mathfrak{A} and adhering to the representation (3.5).*
- (iii) *A time-homogeneous Poisson random measure η on (S, \mathcal{S}) , with intensity measure ν , defined over \mathfrak{A} .*
- (iv) *An initial condition U_0 , which is an E_2 -valued random variable over \mathfrak{A} (with law ρ_0) and is \mathcal{F}_0 -measurable.*
- (v) *A process U on $[0, T]$, which is \mathbb{F} -progressively measurable in H and exhibits càdlàg behavior in E_1 .*

This setup satisfies the following conditions:

- (vi) *For all $t \in [0, T]$, $\mathbb{P}(U(t) \in E_1) = 1$, and $U \in \mathcal{X}$ —see (3.12).*
- (vii) *The integrals*

$$\begin{aligned} & \int_0^t |\langle b(s, U(s)), \varphi \rangle| ds + \int_0^t \sum_{k=1}^{\infty} |\langle \sigma(s, U(s)) [h_k], \varphi \rangle|^2 ds \\ & + \int_0^t \int_{\{x \in S \mid |\langle c(s, x, U(s)), \varphi \rangle|_{V'} < 1\}} |\langle c(s, x, U(s)), \varphi \rangle|^p \nu(dx) ds \\ & + \int_0^t \int_{\{x \in S \mid |\langle c(s, x, U(s)), \varphi \rangle|_{V'} \geq 1\}} |\langle c(s, x, U(s)), \varphi \rangle| \nu(dx) ds \end{aligned} \quad (3.13)$$

are finite, \mathbb{P} -a.s., for every $t \in [0, T]$ and every $\varphi \in V$.

- (viii) *The process U satisfies (3.7), \mathbb{P} -a.s., $\forall t \in [0, T]$ and $\forall \varphi \in V$.*

The typical well-posedness approach begins with establishing the existence of a martingale solution. Once existence is ensured, the focus shifts to proving the uniqueness of the solution. A key challenge in this step is that the concept of solution depends on the chosen definition of the stochastic integral. The choice of integral—whether Itô, Stratonovich, or Marcus—can result in different solution processes, depending on the noise coefficients σ and c . Therefore, before addressing uniqueness, one must specify the type of stochastic integral being used. In the context of Itô calculus, especially dealing with Lévy processes, concepts such as adaptivity, progressively measurability, and predictability must be considered to properly define the solution process. For details, see, e.g., [26, Chapter].

In the framework of the Itô integral, an essential condition is that the increments $\mathbb{W}(s) - \mathbb{W}(t)$, where $s > t \geq 0$, are independent of the solution process $U(t)$. The same requirement applies to the increments of the Poisson process. To address this, the following definition is crucial:

Definition 3.7. Let $\mathbb{W} \in C_b([0, \infty); \mathcal{H})$ and $\eta \in M_{\mathbb{N}}(\{S_n \times [0, T]\})$ be the Wiener process and the Poisson random measure introduced before. Then we define

$$\begin{aligned}\mathscr{W}_t(h) &= \sigma(\{\langle \mathbb{W}(s), h \rangle : 0 \leq s \leq t\}), \quad h \in \mathcal{H}, \\ \mathscr{W}^t(h) &= \sigma(\{\langle \mathbb{W}(s) - \mathbb{W}(t), h \rangle : t \leq s \leq T\}), \quad h \in \mathcal{H},\end{aligned}\tag{3.14}$$

and

$$\begin{aligned}\eta_t(V) &= \eta(V \cap (S \times [0, t])), \quad V \in \mathcal{S} \otimes \mathcal{B}([0, T]), \\ \eta^t(V) &= \eta(V \cap (S \times (t, T])), \quad V \in \mathcal{S} \otimes \mathcal{B}([0, T]).\end{aligned}\tag{3.15}$$

The proof of the next lemma is straightforward and will, therefore, be omitted.

Lemma 3.8. If \mathbb{W} is a Wiener process and η is a time-homogeneous Poisson random measure over a filtered probability space \mathfrak{A} (2.1). Then, for every $t \in [0, T]$, $\mathscr{W}_t(h)$ and $\eta_t(V)$ are \mathcal{F}_t -measurable random variables. In addition, $\mathscr{W}^t(h)$ and $\eta^t(V)$ are independent of \mathcal{F}_t .

In the context of SPDEs, incorporating time into the Kurtz framework requires extending the probability space framework to include a filtration. We work on a filtered probability space $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$, where $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ represents the filtration (see (2.1)). Specifically, the solution process is typically progressively measurable with respect to the filtration generated by both the Wiener process and the Poisson random measure. To handle this additional complexity, Kurtz introduced the concept of *compatibility*. Before defining this concept, however, it is necessary to present some additional definitions.

In our setting, the initial condition U_0 and the processes \mathbb{W} and η are given, where \mathbb{W} is $C_b([0, T]; \mathcal{H})$ -valued and η is $M(\{S_n \times [0, T]\})$ -valued. The initial condition U_0 is assumed to be \mathcal{F}_0 -measurable. The processes \mathbb{W} and η naturally generate a filtration $(\mathcal{F}_t^Y)_{t \in [0, T]}$, where $Y = (\mathbb{W}, \eta, U_0)$, on the underlying probability space Ω . In the following definition, we introduce what is known as the induced filtration on $\mathbb{D}([0, T]; E)$, where E is a Banach space.

Definition 3.9. Let Z be a $\mathbb{D}([0, T]; E)$ -valued random variable on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$. Denote by $Z_t : \mathbb{D}([0, T]; E) \rightarrow E$ the evaluation map $Z \mapsto Z_t$. For $t \in [0, T]$, define $\mathcal{B}_t^Z = \sigma(\{Z_s : s \leq t\})$ as the σ -algebra on $\mathbb{D}([0, T]; E)$ generated by the values of Z up to time t . Now, let us introduce the σ -algebra of the preimages of \mathcal{B}_t^Z on Ω . For any $t \in [0, T]$, let \mathcal{F}_t^Z be the coarsest σ -algebra with respect to which the mapping

$$Z : (\Omega, \mathcal{F}_t^Z) \rightarrow (Z_t, \mathcal{B}_t^Z)$$

is measurable. We refer to the filtration $(\mathcal{F}_t^Z)_{t \in [0, T]}$ as the filtration induced by the random variable Z on the probability space $(\Omega, \mathcal{F}, \mathbb{P})$.

Remark 3.10. A σ -algebra is said to be generated by a family of sets if the family of sets and the σ -algebra are part of the same Borel σ -algebra. In the definition above, $\{Z_s : s \leq t\}$ and \mathcal{B}_t^Z belong to $\mathcal{B}(\mathbb{D}([0, T]; E))$.

Let X_1 and X_2 be Banach spaces, and let $f : X_1 \rightarrow X_2$ be a map. A σ -algebra is said to be induced by f if the family of preimages $\{f^{-1}(A) : A \in \mathcal{B}(X_2)\}$ generates a σ -algebra on X_1 . Here, $f^{-1}(A)$ denotes the set $\{x \in X_1 : f(x) \in A\}$. In the context of Definition 3.9, the random variable Z serves as the mapping f .

Define $Y = (\mathbb{W}, \eta, U_0)$, and let $(\mathcal{F}_t^Y)_{t \in [0, T]}$ be the filtration induced by Y on the underlying probability space Ω . Let X represent a solution to the SPDE (3.6),

taking values in $\mathbb{D}([0, T]; E_1)$. Due to the definition of the Itô integral, the solution process X must be progressively measurable with respect to the induced filtration $(\mathcal{F}_t^Y)_{t \in [0, T]}$. Furthermore, since the system is autonomous, meaning that the only external influences come from the processes \mathbb{W} , η , and the initial data U_0 , the information contained in $(\mathcal{F}_t^Y)_{t \in [0, T]}$ is sufficient to determine the process X almost surely with respect to \mathbb{P} . In particular, for any $t \in [0, T]$ and any bounded Borel measurable function $h : \mathbb{D}([0, T]; E_1) \rightarrow \mathbb{R}$, the following identity holds:⁹

$$\mathbb{E} \left[h(Y) \mid \mathcal{F}_t^{(X, Y)} \right] = \mathbb{E} \left[h(Y) \mid \mathcal{F}_t^Y \right],$$

recalling that a strong solution X can be represented as $F(Y)$ for some function F (see Definition 3.1). This motivates the definition of *temporal compatibility*, as given in Definition 2.1 of [19], which is presented here for generic processes and is not specific to the X and Y associated with the above SPDE.

Definition 3.11. *Let (E_1, \mathcal{E}_1) and (E_2, \mathcal{E}_2) be two Polish spaces, and let X and Y be defined on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, taking values in $B_1 := \mathbb{D}([0, T]; E_1)$ and $B_2 := \mathbb{D}([0, T]; E_2)$, respectively. We say that the process X is temporally compatible with Y , if for every bounded $h \in \mathcal{B}(B_2)$ and for all times $t \in [0, T]$, the following equality holds almost surely:*

$$\mathbb{E} \left[h(Y) \mid \mathcal{F}_t^{(X, Y)} \right] = \mathbb{E} \left[h(Y) \mid \mathcal{F}_t^Y \right],$$

where, as before, $(\mathcal{F}_t^{(X, Y)})_{t \in [0, T]}$ and $(\mathcal{F}_t^Y)_{t \in [0, T]}$ denote the filtrations induced by (X, Y) and Y , respectively (see also Definition 3.9).

We say that a probability measure μ is temporally compatible if, for any pair of random variables X and Y with joint law μ , the variable X is temporally compatible with Y .

Remark 3.12. *If X is temporally compatible with Y , then all relevant information is provided by the σ -algebra $(\mathcal{F}_t^Y)_{t \in [0, T]}$. In other words, once at time $t \in [0, T]$ \mathcal{F}_t^Y is known, knowing \mathcal{F}_t^X does not provide any additional information for calculating the expectation of $h(Y)$. Alternatively, at every time $t \in [0, T]$, knowing the history of both X and Y together (via $\mathcal{F}_t^{(X, Y)}$) does not improve our ability to predict $h(Y)$ beyond just knowing the history of Y alone (via \mathcal{F}_t^Y). If Y has independent increments, then X is temporally compatible with Y if $Y(t+\cdot) - Y(t)$ is independent of $\mathcal{F}_t^{(X, Y)}$ for all $t \in [0, T]$, see [20, Lemma 3.2].*

Remark 3.13. *If the Stratonovich integral is considered, the solution U of an SPDE at time $t \in [0, T]$ inherently depends on the future behavior of the process $t \mapsto \mathbb{W}_t$. Consequently, the solution to an SPDE interpreted in the Stratonovich sense does not satisfy the compatibility condition. Only the Itô integral satisfies the compatibility condition described above. However, in practical applications, it is often possible to convert a Stratonovich-driven SPDE into an Itô-driven SPDE, allowing the above concept to be applied.*

Remark 3.14. *Recall that $\mathcal{S}_{\Gamma, \rho}$, which is defined in (3.3), represents the set of all joint solution measures $\mu_{X, Y} \in \mathcal{P}(B_1 \times B_2)$ that satisfy the constraint (3.2) and*

⁹For two σ -algebras \mathcal{G}_1 and \mathcal{G}_2 , the σ -algebra $\mathcal{G}_1 \vee \mathcal{G}_2$ is defined as $\sigma(\mathcal{G}_1 \cup \mathcal{G}_2)$. We have $\mathcal{F}_t^{(X, Y)} = \mathcal{F}_t^X \vee \mathcal{F}_t^Y$, where \mathcal{F}_t^Z , with $Z = (X, Y)$, $Z = X$, or $Z = Y$, are the induced filtrations (see Definition 3.9).

for which $\mu(B_1 \times \cdot) = \rho$ (ρ is the distribution of the input data Y). Since temporal compatibility is an additional property of the solution that must be satisfied, we introduce the so-called Kurtz set $\mathcal{S}_{\Gamma, \rho, \mathcal{T}}$:

$$\begin{aligned} \mathcal{S}_{\Gamma, \rho, \mathcal{T}} \text{ is the set of joint solution measures } \mu_{X, Y} \in \mathcal{S}_{\Gamma, \rho} \\ \text{that are temporally compatible (see Definition 3.11).} \end{aligned} \quad (3.16)$$

We have defined the set of solutions in (3.10) using a countable dense set \mathbb{Q} of times $t \in [0, T]$ and a countable dense set of test functions φ . Similarly, the concept of compatibility can be extended to a countable set or a sequence of random variables, as detailed in the following lemma. The proof of this lemma is similar to [11, Lemma 5.7] and is therefore not provided here.

Lemma 3.15. *Consider a cylindrical Wiener process \mathbb{W} and a Poisson random measure η with intensity measure ν , both defined over a probability space $(\Omega, \mathcal{F}, \mathbb{P})$ as before. Suppose \mathbb{W} and η are adapted to a filtration $(\mathcal{F}_t)_{t \in [0, T]}$ and let U_0 be an \mathcal{F}_0 -measurable E_2 -valued random variable. Then a sequence of $\mathbb{D}([0, T]; E_1)$ -valued random variables $\{X_j : j \in \mathbb{N}\}$ is temporally compatible with $Y = (\mathbb{W}, \eta, U_0)$ if and only if*

$$\mathcal{F}_t^{X_{I_1}} \vee \dots \vee \mathcal{F}_t^{X_{I_l}} \vee \mathcal{F}_t^Y \text{ is } \mathbb{P}\text{-independent of } \sigma(\mathcal{W}^t) \text{ and } \sigma(\eta^t),$$

where \mathcal{W}^t and η^t are defined in (3.14) and (3.15), respectively. Here, the notation I_1, I_2, \dots, I_l refers to elements of an arbitrary index set I from the power set of \mathbb{N} .

4. YAMADA-WATANABE UNIQUENESS THEORY

In Section 3, we translated the abstract framework of Kurtz [19, 20] into the setting of SPDEs in variational form. Now, we turn our attention to uniqueness of solutions. For SPDEs, one distinguishes between three types of uniqueness: (i) *Uniqueness in law* means that if two solutions start from initial conditions with the same distribution, then the resulting solution processes will have the same distribution. (ii) *Pathwise uniqueness* asserts that if two solutions are given on the same (but arbitrary) filtered probability space and start from the same initial condition, then the two solutions are indistinguishable. (iii) *Strong uniqueness* states that, given a filtered probability space (2.1) where the Wiener process, the Poisson random measure, and the initial condition are all defined, any solution is almost surely unique.

We will establish a key technical result (Lemma 4.5), which asserts that if a process shares the same law as a variational solution—whether strong or weak, as per Definitions 3.1 and 3.2—then this process also qualifies as a variational solution. With the help of Lemma 4.5 and Kurtz’s generalization of the Yamada-Watanabe theorem (see the upcoming Theorem 4.6), we will obtain our primary uniqueness result (Theorem 4.8).

In the following two definitions, pathwise (pointwise) uniqueness and uniqueness in law (distribution) are formulated in the abstract setting (cf. [19, Definition 1.4]).

Definition 4.1. *Pathwise uniqueness is said to hold for the abstract equation (3.1) if, for any processes X_1, X_2 , and Y defined on the same probability space $(\Omega, \mathcal{F}, \mathbb{P})$ and associated with the joint measures $\mu_{X_1, Y}$ and $\mu_{X_2, Y} \in \mathcal{S}_{\Gamma, \rho, \mathcal{T}}$ (cf. Remark 3.14), respectively, it holds that*

$$\mathbb{P}(\{X_1 = X_2\}) = 1.$$

Definition 4.2. *Joint uniqueness in law (or weak joint uniqueness) is said to hold for the abstract equation (3.1) if $\mathcal{S}_{\Gamma, \rho, \mathcal{T}}$ contains at most one measure. Uniqueness in law (or weak uniqueness) holds if all solution measures $\mu \in \mathcal{S}_{\Gamma, \rho, \mathcal{T}}$ have the same marginal distribution on B_1 (cf. Remark 3.14).*

In our context, uniqueness as described in Definition 4.1 aligns with the standard notion of pathwise uniqueness defined below.

Definition 4.3. *Whenever $(\mathfrak{A}, U^{(i)}, \mathbb{W}, \eta, U_0^{(i)})$, $i = 1, 2$, are two solutions to the SPDE (3.6) that adheres to Definition 3.6 and Assumption 4.4, such that \mathbb{W} is a cylindrical Wiener process evolving over \mathcal{H} , ν is the intensity measure of η , ρ_0 is the law of $U_0^{(i)}$, $i = 1, 2$, and*

$$\mathbb{P} \left(\left\{ U^{(1)}(0) = U^{(2)}(0) \right\} \right) = 1,$$

then it holds that

$$\mathbb{P} \left(\left\{ U^{(1)}(t) = U^{(2)}(t) \right\} \right) = 1, \quad \forall t \in [0, T].$$

Often, one is only able in a first step to prove the existence of a probabilistic weak solution in the sense of Definition 3.6, and, in a second step is to verify pathwise uniqueness. However, pathwise uniqueness is often only achievable under additional regularity conditions on the solution process. These additional regularity properties are not inherent to the definition of a solution but are needed for proving uniqueness. They are introduced through two abstract mappings, $\theta_0^{\alpha_0}$ and $\theta_1^{\alpha_1}$, where α_0 and α_1 belong to some index sets. See the discussion leading to (3.12).

Assumption 4.4. *Let $\{\theta_i^{\alpha_i} : \mathbb{D}([0, T], E_1) \rightarrow [0, \infty] \mid \alpha_i \in A_i\}$, $i = 0, 1$, be two families of mappings, where A_0 and A_1 are index sets. The solution U satisfies the condition $U \in \mathcal{X}$, where the set \mathcal{X} , defined in (3.12), depends on these families of mappings.*

Given a filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$ on a probability space $(\Omega, \mathcal{F}, \mathbb{P})$, the augmented filtration $\mathbb{F}^{\mathbb{P}} = (\mathcal{F}_t^{\mathbb{P}})_{t \in [0, T]}$ is defined for each $t \geq 0$ as $\mathcal{F}_t^{\mathbb{P}} = \sigma(\mathcal{F}_t \cup \mathcal{N})$, where \mathcal{N} is the collection of all \mathbb{P} -null sets in \mathcal{F} . The augmented filtration is complete (i.e., it contains all null sets and is right-continuous).

The next lemma demonstrates that although the tuple $(U, \mathbb{W}, \eta, U_0)$, defined on a given filtered probability space \mathfrak{A} as in (2.1), is not initially assumed to solve the SPDE, the equality of the laws, together with the existence of a solution on a different filtered probability space $\bar{\mathfrak{A}}$, implies that $(U, \mathbb{W}, \eta, U_0)$ must satisfy the SPDE. Recall that pathwise uniqueness ensures that two solutions with the same driving noise (Wiener process and Poisson random measure) and initial conditions are indistinguishable. Therefore, when pathwise uniqueness is known, even if the solutions are defined on different filtered probability spaces, they can be related through their distributions. This connection allows us to conclude the existence of a unique solution on the original space \mathfrak{A} .

Lemma 4.5. *Consider a filtered probability space $\mathfrak{A} = (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ with filtration $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$, along with the following additional elements:*

- U , a $\mathbb{D}([0, T]; E_1)$ -valued random variable,
- \mathbb{W} , a cylindrical Wiener process on \mathcal{H} , represented as (3.5),
- η , a random variable with values in $M_{\mathbb{N}}(\{S_n \times [0, T]\})$, and

- U_0 , a \mathcal{F}_0 -measurable E_2 -valued random variable,

all defined on \mathfrak{A} . In addition, suppose that for any $t \in [0, T]$, $U(t)$, \mathscr{W}_t , and η_t are \mathcal{F}_t -measurable, and that \mathscr{W}^t and η^t are independent of \mathcal{F}_t (see Definition 3.7 and Lemma 3.8).

Suppose there exists a solution

$$(\bar{\mathfrak{A}}, \bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0),$$

where $\bar{\mathfrak{A}} := (\bar{\Omega}, \bar{\mathcal{F}}, \bar{\mathbb{P}}, \bar{\mathbb{P}})$, $\bar{\mathbb{F}} = (\bar{\mathcal{F}}_t)_{t \in [0, T]}$, is a potentially different filtered probability space, and this solution to the SPDE (3.6) adheres to the conditions outlined in Definition 3.6 and Assumption 4.4, such that the law of $(U, \mathbb{W}, \eta, U_0)$ coincides with the law of $(\bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ on $\mathbb{D}([0, T]; E_1) \times C_b(0, T; \mathcal{H}) \times M_{\mathbb{N}}(\{S_n \times [0, T]\}) \times E_2$.

Then the tuple $(\mathfrak{A}^{\mathbb{P}}, U, \mathbb{W}, \eta, U_0)$, where $\mathfrak{A}^{\mathbb{P}} := (\Omega, \mathcal{F}, \mathbb{F}^{\mathbb{P}}, \mathbb{P})$, is a solution to the SPDE (3.6), according to Definition 3.6 and Assumption 4.4.

Proof. The proof is similar to the proof of Lemma 4.4 in [11] with the modification of incorporating the Wiener process and the variational setting. Given the equality of laws and the premise that $(\bar{\mathfrak{A}}, \bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ constitutes a solution, the procedure of verifying that $(\mathfrak{A}^{\mathbb{P}}, U, \mathbb{W}, \eta)$ meets the criteria (i) through (vii) outlined in Definition 3.6 and conforms to Assumption 4.4 follows a routine argument. Consequently, our focus below is to verify condition (viii). This involves demonstrating that the triple $(U, \mathbb{W}, \eta, U_0)$ satisfies the specified SPDE (3.6) in the sense of (3.7).

Consider a process U that is progressively measurable over \mathfrak{A} and càdlàg in the space E_1 , meaning that for each time $t \geq 0$, $U(t)$ takes values in E_1 , with well-defined left limits, and satisfies $\lim_{s \downarrow t} U(s) = U(t)$ in E_1 almost surely with respect to \mathbb{P} . Suppose this process satisfies the condition given in (3.13). For every test function $\varphi \in V$ (see (3.4)) and time $t \in [0, T]$, we introduce a nonlinear mapping $\mathcal{K}_{\mathfrak{A}}^{\varphi}$ as follows:

$$\begin{aligned} \mathcal{K}_{\mathfrak{A}}^{\varphi}(U, \mathbb{W}, \eta, U_0)(t) &:= \langle U_0, \varphi \rangle + \int_0^t \langle b(s, U(s)), \varphi \rangle \\ &\quad + \int_0^t \sum_{k=1}^{\infty} \langle \sigma(s, U(s))[h_k], \varphi \rangle d\beta^k(s) \\ &\quad + \int_0^t \int_S \langle c(s, x, U(s)), \varphi \rangle \tilde{\eta}(dx, ds). \end{aligned}$$

Note that $\mathcal{K}_{\mathfrak{A}}^{\varphi}(V, \mathbb{W}, \eta, U_0)$ is contingent upon \mathbb{W} through the Hilbert space \mathcal{H} and on η via its compensator. Consequently, it also depends on the probability measure \mathbb{P} . The objective is to demonstrate that if, for all $\varphi \in V$ and $t \in [0, T]$,

$$\bar{\mathbb{P}}\left(\mathcal{K}_{\bar{\mathfrak{A}}}^{\varphi}(\bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)(t) - \bar{U}(t) = 0\right) = 1,$$

then it necessarily follows that, for all $\varphi \in V$ and $t \in [0, T]$,

$$\mathbb{P}\left(\mathcal{K}_{\mathfrak{A}}^{\varphi}(U, \mathbb{W}, \eta, U_0)(t) - U(t) = 0\right) = 1. \quad (4.1)$$

To establish this, we examine each component of the operators $\mathcal{K}_{\mathfrak{A}}^{\varphi}(U, \mathbb{W}, \eta, U_0)$ and $\mathcal{K}_{\bar{\mathfrak{A}}}^{\varphi}(\bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$, demonstrating their equivalence in distribution. This equivalence directly implies that (4.1) holds, or equivalently, the tuple $(\mathfrak{A}, U, \mathbb{W}, \eta, U_0)$ serves as a solution to the SPDE (3.6).

Initially, it is evident that the components of $\mathcal{K}_{\mathfrak{A}}^\varphi(\bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ adhere to (3.13). Given that the laws of $(U, \mathbb{W}, \eta, U_0)$ and $(\bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ are equal, and considering the functions b , σ , and c are measurable, it naturally follows that the components of $\mathcal{K}_{\mathfrak{A}}^\varphi(U, \mathbb{W}, \eta, U_0)$ equally fulfill (3.13). Recall also that the initial data U_0, \bar{U}_0 are given and share the same law.

Define the following real-valued processes:

$$\bar{\mathfrak{b}}_\varphi(s, \bar{\omega}) := \langle b(s, \bar{U}(s, \bar{\omega})), \varphi \rangle, \quad s \in [0, T], \bar{\omega} \in \bar{\Omega},$$

and

$$\mathfrak{b}_\varphi(s, \omega) := \langle b(s, U(s, \omega)), \varphi \rangle, \quad s \in [0, T], \omega \in \Omega.$$

Given the underlying assumptions, we have $\text{Law}(U) = \text{Law}(\bar{U})$ in the intersection space $\mathbb{D}([0, T]; E) \cap L^2(0, T; V)$. Consequently, \mathfrak{b}_φ and $\bar{\mathfrak{b}}_\varphi$ share identical laws on $\mathbb{D}([0, T]; \mathbb{R})$ (for every $\varphi \in V$). Leveraging [23, Theorem 8.3], it follows that for any time $t \in [0, T]$, the laws

$$\text{Law}\left(\bar{U}, \int_0^t \bar{\mathfrak{b}}_\varphi(s) ds\right) \quad \text{and} \quad \text{Law}\left(U, \int_0^t \mathfrak{b}_\varphi(s) ds\right)$$

are identical on the space $\mathbb{D}([0, T]; E) \times \mathbb{R}$ (across all $\varphi \in V$).

Define the following real-valued processes:

$$\bar{\mathfrak{s}}_\varphi^k(s, \bar{\omega}) := \langle \sigma(s, \bar{U}(s, \bar{\omega}))[h_k], \varphi \rangle, \quad s \in [0, T], \bar{\omega} \in \bar{\Omega}$$

and

$$\mathfrak{s}_\varphi^k(s, \omega) := \langle \sigma(s, U(s, \omega))[h_k], \varphi \rangle, \quad s \in [0, T], \omega \in \Omega.$$

The two processes $\{\bar{\mathfrak{s}}_\varphi^k(s) \mid s \in [0, T]\}$ and $\{\mathfrak{s}_\varphi^k(s) \mid s \in [0, T]\}$ are adapted to the filtrations $(\bar{\mathcal{F}}_s)_{s \in [0, T]}$ and $(\mathcal{F}_s)_{s \in [0, T]}$, respectively. Besides, the laws of (U, \mathbb{W}) and $(\bar{U}, \bar{\mathbb{W}})$ are identical. Drawing upon [23, Theorem 8.6], we infer that for any $t \in [0, T]$, the laws

$$\text{Law}\left(\bar{U}, \int_0^t \sum_{k=1}^{\infty} \bar{\mathfrak{s}}_\varphi^k(s) d\beta^k(s)\right) \quad \text{and} \quad \text{Law}\left(U, \int_0^t \sum_{k=1}^{\infty} \mathfrak{s}_\varphi^k(s) d\beta^k(s)\right)$$

are equal on the space $(\mathbb{D}([0, T]; E_1) \cap L^2(0, T; V)) \times \mathbb{R}$ (for all $\varphi \in V$).

In the final step, define the processes through

$$\bar{\mathfrak{c}}_\varphi(s, \bar{\omega}) := \langle c(s, \bar{U}(s, \bar{\omega})), \varphi \rangle, \quad s \in [0, T], \bar{\omega} \in \bar{\Omega}$$

and

$$\mathfrak{c}_\varphi(s, \omega) := \langle c(s, U(s, \omega)), \varphi \rangle, \quad s \in [0, T], \omega \in \Omega,$$

which are adapted to the filtrations $(\bar{\mathcal{F}}_s)_{s \in [0, T]}$ and $(\mathcal{F}_s)_{s \in [0, T]}$, respectively. By the assumption (3.13) (and the equality of laws), $\bar{\mathfrak{c}}_\varphi$ and \mathfrak{c}_φ belong a.s. to $L^p([0, T]; \mathbb{R})$. Proposition B.1 in [11] guarantees the progressive measurability of $\bar{\mathfrak{c}}_\varphi$ and \mathfrak{c}_φ . This is explained by employing a sequence of shifted Haar projections, which approximates (in L^p) the original processes with simpler processes that possess the necessary measurability. Define the stochastic integrals

$$\bar{\mathcal{I}}_\varphi(t) = \int_0^t \int_S \langle c(s, \bar{U}(s, z)), \varphi \rangle \bar{\eta}(dz, ds), \quad t \in [0, T],$$

and

$$\mathcal{I}_\varphi(t) = \int_0^t \int_S \langle c(s, U(s, z)), \varphi \rangle \eta(dz, ds), \quad t \in [0, T].$$

Applying [11, Theorem A.1], we conclude that $(\bar{\mathcal{I}}_\varphi(t), \bar{U}, \bar{\eta})$ and $(\mathcal{I}_\varphi(t), U, \eta)$ possess the same law on $\mathbb{R} \times \mathbb{D}([0, T]; E) \times \mathcal{M}_{\mathbb{N}}(\{S_n \times [0, T]\})$.

Hence, the lemma is proved. \square

In the abstract setting, the following generalisation of the Yamada-Watanabe theorem holds [19, Theorem 1.5].

Theorem 4.6. *The following are equivalent:*

- $\mathcal{S}_{\Gamma, \rho, \mathcal{T}} \neq \emptyset$, see (3.16), and pathwise uniqueness holds.
- There exists a strong solution (Definition 3.1) and joint uniqueness in law (Definition 4.2) holds.

The abstract Theorem 4.6 can be tailored to our specific SPDE context. We now outline the detailed setup and reformulate Theorem 4.6 within this framework. Consider a filtered probability space \mathfrak{A} (see (2.1)) with a cylindrical Wiener process \mathbb{W} on \mathcal{H} (see Section 2.1), a Poisson random measure η on S (see Section 2.2), and an initial condition $U_0 \in E_2$, where U_0 is \mathcal{F}_0 -measurable. Here, η and \mathbb{W} are independent and adapted to the filtration $(\mathcal{F}_t)_{t \in [0, T]}$. Define $Y = (\mathbb{W}, \eta, U_0)$ on B_2 (see (3.8)), and let $X = U$ denote a solution to the SPDE (3.6) within B_1 (see (3.9)). The mappings $\Omega \ni \omega \mapsto X(\omega)$ and $\Omega \ni \omega \mapsto Y(\omega)$ induce two filtrations $(\mathcal{F}_t^X)_{t \in [0, T]}$ and $(\mathcal{F}_t^Y)_{t \in [0, T]}$ on \mathfrak{A} (see Definition 3.11). Here, the notion of temporal compatibility becomes relevant.

To clarify further, let us specify $\mathcal{B}_t^{B_i}$ for $i = 1, 2$, where $i = 1$ corresponds to X , and $i = 2$ corresponds to Y . For the solution process X , as defined in Definition 3.9, the σ -algebra $\mathcal{B}_t^{B_1}$ is generated by the coordinate map $\pi_s : z \in B_1 \mapsto z(s) \in E_1$ for $s \leq t$, $t \in [0, T]$. The input data Y takes values in the path space $B_2 = C([0, T]; \mathcal{H}) \times M_{\mathbb{N}}(\{S_n \times [0, T]\}) \times E_2$. On this path space B_2 , we define the σ -algebra generated by the input data Y as $\mathcal{B}_t^{B_2} = \sigma(\mathbb{W}_t) \otimes \sigma(R_t) \otimes \sigma_B(U_0)$, where \mathbb{W}_t and R_t , for $t \in [0, T]$, denote the canonical restriction mappings defined as follows:

- \mathbb{W}_t restricts a Wiener process \mathbb{W} to the interval $[0, t]$, i.e., \mathbb{W}_t , mapping from $C([0, T]; \mathcal{H})$ to itself, is defined by $\mathbb{W} \mapsto \mathbb{W}\mathbf{1}_{[0, t]}$.
- R_t restricts the Poisson random measure η to the set $\{S_n\} \times (0, t]$, i.e., R_t , which maps from $M_{\mathbb{N}}(\{S_n \times [0, T]\})$ to itself, is defined as

$$R_t : \eta(A \times I) \mapsto \eta(A \times (I \cap (0, t])),$$

where $A \in \mathcal{B}(S_n)$ for some $n \in \mathbb{N}$ and $I \in \mathcal{B}([0, T])$.

Since the initial condition U_0 is known at the outset and remains unchanged thereafter, we have $\mathcal{B}_0^{B_2} = \sigma(U_0)$. Moreover, as U_0 is \mathcal{F}_0 -measurable by definition, we set $\mathcal{F}_0 := \sigma(\{U_0^{-1}(B) : B \in \mathcal{B}(E_2)\})$, ensuring that U_0 is measurable with respect to both \mathcal{F}_0 and $\mathcal{B}(E_2)$.

Let $\rho_{\mathbb{W}}$ denote the law of the cylindrical Wiener process \mathbb{W} over $C([0, T]; \mathcal{H})$. Similarly, let ρ_ν represent the law of the Poisson random measure η equipped with the Lévy measure ν on $M_{\mathbb{N}}(\{S_n \times [0, T]\})$. Furthermore, let ρ_0 symbolize the law of the \mathcal{F}_0 -measurable random variable U_0 over E_2 . Given the independence of \mathbb{W} , η , U_0 , we proceed to define their joint law ρ as the product measure

$$\rho = \rho_{\mathbb{W}} \times \rho_\nu \times \rho_0. \quad (4.2)$$

Let $V_d = \{\varphi_k : k \in \mathbb{N}\}$ be a dense countable subset of V (see (3.4)), and recall the set Γ of mappings given by 3.10 and (3.11). The SPDE (3.6) (via (3.7)) defines the mappings in Γ in (3.10), (3.11). Specifically, given the filtered probability space

\mathfrak{A} described in (2.1), we define Γ by associating a real number $\Gamma_{\varphi,t}(U, (\mathbb{W}, \eta, U_0))$ with an arbitrary process U and any triplet (\mathbb{W}, η, U_0) on \mathfrak{A} , in accordance with (3.11).

Using the notation introduced above, we can now reformulate the SPDE (3.6) as a property of a solution measure on the path spaces B_1 and B_2 , expressed in the format (3.1). To be more precise, let X represent the solution U , and Y denote the input data (\mathbb{W}, η, U_0) , then X is a solution if $\mathbb{P}(\Gamma(X, Y) = 0) = 1$. Our objective is to identify a solution measure $\mu \in \mathcal{S}_{\Gamma, \rho, \mathcal{T}}$ on $B_1 \times B_2$ (see (3.16)) that adheres to the condition

$$\int_{B_1 \times B_2} |\Gamma_{\varphi}(x, y)(t)| \mu(dx, dy) = 0, \quad (4.3)$$

for all $\varphi \in V_d$ and $t \in \mathbb{Q}_T$ (compare with (3.2)). Pathwise uniqueness, however, necessitates further constraints on the solution beyond those defined in (4.3). The additional properties and regularity specified in Assumption 4.4 must be conveyed as a condition on the solution measure. Specifically, the solution measure μ is required to satisfy:

$$\mu(\{(x, y) \in B_1 \times B_2 : \theta_0^{\alpha_0}(x) < \infty\}) = 1, \quad \forall \alpha_0 \in A_0, \quad (4.4)$$

$$\int_{B_1 \times B_2} \theta_1^{\alpha_1}(x) \mu(dx \times dy) < \infty, \quad \forall \alpha_1 \in A_1. \quad (4.5)$$

Remark 4.7. *Although we will not make use of this, the set of solution measures μ on $B_1 \times B_2$ satisfying (4.4) and (4.5) is convex [20, Lemma 3.8].*

We use the notation Γ^{θ} to represent the combination of constraints specified in (4.3), (4.4), and (4.5). The superscript θ indicates the additional constraints (4.4) and (4.5). When μ denotes a Borel probability measure that is the joint law of the random vector $(U, (\mathbb{W}, \eta, U_0))$, we say that μ adheres to the Kurtz convexity constraint Γ^{θ} .

Following [20, p. 958], we now explicitly define the Kurtz set $\mathcal{S}_{\Gamma^{\theta}, \rho, \mathcal{T}}$ (see (3.16)) within the present SPDE context. This set consists of all solution measures that meet the specified criteria. Indeed, a probability measure $\mu \in \mathcal{P}(B_1 \times B_2)$ belongs to $\mathcal{S}_{\Gamma^{\theta}, \rho, \mathcal{T}}$ if and only if it satisfies the following conditions:

- μ is in compliance with the convexity constraint Γ^{θ} , see (4.3), (4.4), and (4.5), and μ is temporally compatible (see Definition 3.11);
- For every A in the Borel σ -algebra $\mathcal{B}(B_2)$, $\mu(B_1 \times A) = \rho(A)$, where ρ is the joint law of (\mathbb{W}, η, U_0) , see (4.2).

An alternative description of the Kurtz set $\mathcal{S}_{\Gamma^{\theta}, \rho, \mathcal{T}}$ is as follows: $\mu \in \mathcal{P}(B_1 \times B_2)$ belongs to $\mathcal{S}_{\Gamma^{\theta}, \rho, \mathcal{T}}$ if and only if the following criteria are met:

- there exists a solution tuple $(\bar{\mathfrak{A}}, \bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ to the SPDE (3.6), where $\bar{\mathfrak{A}}$ is a filtered probability space, satisfying Definition 3.6 and Assumption 4.4, and
 - $\bar{\mathbb{W}}$ is a cylindrical Wiener process evolving over the Hilbert space \mathcal{H} (see Section 2.1);
 - $\bar{\eta}$ is a Poisson random measure with the intensity measure ν (see (4.2) and Section 2.2);
 - \bar{U}_0 has distribution ρ_0 (see (4.2));
- μ is the joint law of $(\bar{U}, (\bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0))$ on $B_1 \times B_2$ (see (3.9) and (3.8)).

Given the new constraints, the definitions of temporal compatibility and pointwise uniqueness must be adjusted accordingly. In particular, the temporal compatibility condition requires that, for $t \in [0, T]$, the solution $\bar{U}(t)$ is independent of $\bar{\mathcal{W}}^t$ and $\bar{\eta}^t$ (see Definition 3.7) and is measurable with respect to $\bar{\mathcal{W}}_t$, $\bar{\eta}_t$, and $\sigma(\bar{U}_0)$.

Our main result is presented in the next theorem, which establishes a connection between the existence of a weak solution (Definition 3.2) with pathwise uniqueness (Definition 4.1) and the existence of a unique strong solution (see Definition 3.1). Using the results developed earlier in this section, the proof follows an adaptation of [11], with the only modification being the inclusion of the Wiener process. Due to the similarity in reasoning, a detailed proof is omitted.

Theorem 4.8. *Consider a Gelfand triple (V, H, V') , see (3.4), along with Banach spaces E_1 and E_2 . Here, E_2 is continuously embedded into E_1 , and V is continuously embedded into E_1 . Additionally, E_1 is a UMD space of type 2. Let ρ_0 be a Borel probability measure on E_2 , see (4.2). Suppose \mathcal{H} is a Hilbert space, ν is an intensity measure over a Polish space (S, \mathcal{S}) , see (2.2) and (2.3), and assume that*

- *there exists a solution $(\bar{\mathfrak{A}}, \bar{U}, \bar{\mathbb{W}}, \bar{\eta}, \bar{U}_0)$ to the SPDE (3.6) that adheres to Definition 3.6 and Assumption 4.4, such that $\bar{\mathbb{W}}$ is a cylindrical Wiener process evolving over \mathcal{H} , see (3.5), ν is the intensity measure of $\bar{\eta}$, and ρ_0 is the probability law of \bar{U}_0 ;*
- *pathwise uniqueness, as defined in Definition 4.3, is satisfied.*

Then there exists a Borel measurable mapping

$$F : C([0, T]; \mathcal{H}) \times M_{\mathbb{N}}(\{S_n \times [0, T]\}) \times E_2 \rightarrow \mathbb{D}([0, T]; E_1),$$

depending on \mathcal{H} , ν and ρ_0 , such that

- *if $(\mathfrak{A}, U, \mathbb{W}, \eta, U_0)$ is a solution to the SPDE (3.6), in the sense of Definition 3.6 and Assumption 4.4, such that \mathbb{W} is a cylindrical Wiener process on \mathcal{H} with the representation (3.5), ν is the intensity measure of η , and ρ_0 is the law of $U(0)$, then*

$$U = F(\mathbb{W}, \eta, U_0) \quad \mathbb{P}\text{-almost surely,}$$

and U is progressively measurable with respect to the \mathbb{P} -augmentation of the filtration (cf. Definition 3.7)

$$\left(\sigma(\{\mathcal{W}_t(h) : h \in \mathcal{H}\}), \sigma(\{\eta_t(V) : V \in S \otimes \mathcal{B}([0, T])\}), \sigma(U_0) \right)_{t \in [0, T]};$$

- *if $\mathfrak{A} := (\Omega, \mathcal{F}, (\mathcal{F}_t)_{t \in [0, T]}, \mathbb{P})$ is a filtered probability space, U_0 is an E_2 -valued \mathcal{F}_0 -measurable random variable with law ρ_0 , \mathbb{W} is a cylindrical Wiener process on \mathcal{H} with the representation (3.5), and η is a time-homogeneous Poisson random measure with intensity ν , then*

$$U = F(\mathbb{W}, \eta, U_0)$$

is adapted to the augmented filtration $(\mathcal{F}_t^{\mathbb{P}})_{t \in [0, T]}$, $U(0) = U_0$, \mathbb{P} -a.s., and

$$(\mathfrak{A}^{\mathbb{P}}, U, \mathbb{W}, \eta, U_0), \quad \text{where } \mathfrak{A}^{\mathbb{P}} := (\Omega, \mathcal{F}, (\mathcal{F}_t^{\mathbb{P}})_{t \in [0, T]}, \mathbb{P}),$$

constitutes a solution to the SPDE (3.6) in the sense of Definition 3.6, fulfilling Assumption 4.4.

We briefly outline the proof of Theorem 4.8. From the assumptions of Theorem 4.8, it follows that the Kurtz set $\mathcal{S}_{\Gamma^\theta, \rho, \mathcal{T}}$ is non-empty. Furthermore, pathwise uniqueness is assumed. Applying Theorem 4.6, we conclude the existence of a strong solution and joint uniqueness in law. In particular, there exists a Borel measurable function

$$F : C_b([0, T]; \mathcal{H}) \times M_{\mathbb{N}}(\{S_n \times [0, T]\}) \times E_2 \rightarrow \mathbb{D}([0, T]; E_1),$$

such that, for a given filtered probability space $\mathfrak{A} := (\Omega, \mathcal{F}, \mathbb{F}, \mathbb{P})$ with $\mathbb{F} = (\mathcal{F}_t)_{t \in [0, T]}$, where U_0 is an E_2 -valued, \mathcal{F}_0 -measurable random variable with law ρ_0 , \mathbb{W} is a cylindrical Wiener process on \mathcal{H} with the representation (3.5), and η is a time-homogeneous Poisson random measure with intensity ν , the solution U is given \mathbb{P} -almost surely by $U = F(\mathbb{W}, \eta, U_0)$. Moreover, due to pathwise uniqueness and Lemma 4.5, if the triplet (\mathbb{W}, η, U_0) is defined on any other filtered probability space satisfying the constraints in Lemma 4.5, then the solution U (given by $F(\mathbb{W}, \eta, U_0)$) on the specified probability space is unique. Hence, we conclude that there exists a unique strong solution.

Remark 4.9. *A consequence of Theorem 4.8 is that if $(\mathfrak{A}^{(i)}, U^{(i)}, \mathbb{W}^{(i)}, \eta^{(i)}, U_0^{(i)})$, where $\mathfrak{A}^{(i)} = (\Omega^{(i)}, \mathcal{F}^{(i)}, \mathbb{F}^{(i)}, \mathbb{P}^{(i)})$, $\mathbb{F}^{(i)} = (\mathcal{F}_t^{(i)})_{t \in [0, T]}$, $i = 1, 2$, are two solutions to the SPDE (3.6) in the sense of Definition 3.6 and Assumption 4.4, such that $\mathbb{W}^{(i)}$ are cylindrical Wiener processes evolving over \mathcal{H} , ν is the intensity measure of $\eta^{(i)}$, and ρ_0 is the law of $U_0^{(i)}$, $i = 1, 2$, then we have uniqueness in law:*

$$\text{Law} \left(U^{(1)}, \mathbb{W}^{(1)}, \eta^{(1)} \right) \text{ coincides with } \text{Law} \left(U^{(2)}, \mathbb{W}^{(2)}, \eta^{(2)} \right).$$

APPENDIX A. THE SKOROHOD SPACE

For an introduction to the Skorokhod space, we refer the reader to [5, 13, 18]. In this section, we recall only a few definitions that are essential for our work. Let $(Y, |\cdot|_Y)$ be a separable Banach space. The space $\mathbb{D}(0, T; Y)$ denotes the set of all right-continuous functions $x : [0, T] \rightarrow Y$ with left-hand limits. Let Λ denote the class of all strictly increasing continuous functions $\lambda : [0, T] \rightarrow [0, T]$ such that $\lambda(0) = 0$ and $\lambda(T) = T$. Clearly, any $\lambda \in \Lambda$ is a homeomorphism of $[0, T]$ onto itself. We now define the Skorokhod topology. First, let

$$\|\lambda\|_{\log} := \sup_{\substack{t, s \in [0, T] \\ t \neq s}} \left| \log \left(\frac{\lambda(t) - \lambda(s)}{t - s} \right) \right| \sim \text{ess sup}_{t \in [0, T]} |\log(\lambda'(t))|.$$

Introducing the function class

$$\Lambda_{\log} := \left\{ \lambda \in \Lambda : \|\lambda\|_{\log} < \infty \right\},$$

the metric d_0 between x and y in $\mathbb{D}(0, T; Y)$ is given by

$$d_0(x, y) := \inf_{\lambda \in \Lambda_{\log}} \left\{ \|\lambda\|_{\log} \wedge \sup_{t \in [0, T]} |x(t) - y(\lambda(t))|_Y \right\}.$$

The metric d_0 generates the Skorokhod topology on $\mathbb{D}([0, T]; Y)$. Moreover, the space $\mathbb{D}([0, T]; Y)$, equipped with the metric d_0 , is a complete and separable metric space.

The metric d_0 is particularly useful for comparing discontinuous functions because it enables small adjustments in the time axis to better align their discontinuities. This flexibility ensures that even if two functions exhibit similar behavior but

with slight variations in the timing of their jumps, the d_0 metric accurately reflects their closeness, while the L^∞ metric does not.

REFERENCES

- [1] D. Applebaum. *Lévy processes and stochastic calculus*, volume 116 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 2009. [5](#), [6](#)
- [2] A. Araujo and E. Giné M. Type, cotype and Lévy measures in Banach spaces. *Ann. Probab.*, 6(4):637–643, 1978. [5](#)
- [3] V. Barbu, G. Da Prato, and M. Röckner. *Stochastic porous media equations*, volume 2163 of *Lecture Notes in Mathematics*. Springer, 2016. [11](#)
- [4] M. Barczy, Z. Li, and G. Pap. Yamada-Watanabe results for stochastic differential equations with jumps. *Int. J. Stoch. Anal.*, 2015:23, 2015. Id/No 460472. [1](#)
- [5] P. Billingsley. *Convergence of probability measures*. Wiley Series in Probability and Statistics: Probability and Statistics. John Wiley & Sons, Inc., New York, second edition, 1999. A Wiley-Interscience Publication. [24](#)
- [6] Z. Brzeźniak and D. Gątarek. Martingale solutions and invariant measures for stochastic evolution equations in banach spaces. *Stochastic Processes and their Applications*, 84(2):187–225, 1999. [5](#)
- [7] Z. Brzeźniak and E. Hausenblas. Maximal regularity for stochastic convolutions driven by Lévy processes. *Probab. Theory Related Fields*, 145(3-4):615–637, 2009. [6](#)
- [8] Z. Brzeźniak, W. Liu, and J. Zhu. Strong solutions for SPDE with locally monotone coefficients driven by Lévy noise. *Nonlinear Anal. Real World Appl.*, 17:283–310, 2014. [9](#)
- [9] Z. Brzeźniak and M. Ondreját. Weak solutions to stochastic wave equations with values in Riemannian manifolds. *Comm. Partial Differential Equations*, 36(9):1624–1653, 2011. [9](#)
- [10] G. Da Prato and J. Zabczyk. *Stochastic equations in infinite dimensions*, volume 152 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, second edition, 2014. [3](#)
- [11] A. de Bouard, E. Hausenblas, and M. Ondreját. Uniqueness of the nonlinear Schrödinger equation driven by jump processes. *NoDEA Nonlinear Differential Equations Appl.*, 26(3):Paper No. 22, 31, 2019. [1](#), [2](#), [17](#), [19](#), [20](#), [21](#), [23](#)
- [12] E. Dettweiler. A characterization of the Banach spaces of type p by Lévy measures. *Math. Z.*, 157(2):121–130, 1977. [6](#)
- [13] S. N. Ethier and T. G. Kurtz. *Markov processes: Characterization and convergence*. John Wiley & Sons, Inc., New York, 1986. [24](#)
- [14] E. Hausenblas, K. Fahim, and K. Karlsen. On the cardiac bidomain model and the effect of Lévy noise: An analysis of strong solutions. in preparation, 2024. [2](#)
- [15] T. Hytönen, J. van Neerven, M. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. I. Martingales and Littlewood-Paley theory*. Springer, Cham, 2016. [6](#), [11](#)
- [16] T. Hytönen, J. van Neerven, M. Veraar, and L. Weis. *Analysis in Banach spaces. Vol. II*. Springer, Cham, 2017. [5](#)
- [17] N. Ikeda and S. Watanabe. *Stochastic differential equations and diffusion processes*, volume 24 of *North-Holland Mathematical Library*. North-Holland Publishing Co., Amsterdam, 1981. [4](#), [6](#)
- [18] J. Jacod and A. N. Shiryaev. *Limit theorems for stochastic processes*, volume 288 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, second edition, 2003. [24](#)
- [19] T. Kurtz. Weak and strong solutions of general stochastic models. *Electron. Commun. Probab.*, 19(58):1–16, 2014. [2](#), [9](#), [10](#), [16](#), [17](#), [21](#)
- [20] T. G. Kurtz. The Yamada-Watanabe-Engelbert theorem for general stochastic equations and inequalities. *Electron. J. Probab.*, 12:951–965, 2007. [1](#), [9](#), [16](#), [17](#), [22](#)
- [21] W. Linde. *Probability in Banach spaces—stable and infinitely divisible distributions*. A Wiley-Interscience Publication. John Wiley & Sons Ltd., Chichester, second edition, 1986. [5](#)
- [22] W. Liu and M. Röckner. *Stochastic partial differential equations: an introduction*. Springer, Cham, 2015. [9](#), [11](#)
- [23] M. Ondreját. Uniqueness for stochastic evolution equations in banach spaces. *Dissertationes Mathematicae*, 426:1–63, 01 2004. [20](#)

- [24] K. R. Parthasarathy. *Probability measures on metric spaces*. Probability and Mathematical Statistics, No. 3. Academic Press Inc., New York, 1967. [2](#)
- [25] S. Peszat and J. Zabczyk. *Stochastic partial differential equations with Lévy noise. An evolution equation approach*. Cambridge University Press, Cambridge, 2007. [6](#), [7](#)
- [26] D. Revuz and M. Yor. *Continuous martingales and Brownian motion*, volume 293 of *Grundlehren der Mathematischen Wissenschaften*. Springer-Verlag, Berlin, third edition, 1999. [14](#)
- [27] T. Runst and W. Sickel. *Sobolev spaces of fractional order, Nemytskij operators, and nonlinear partial differential equations*, volume 3 of *De Gruyter Series in Nonlinear Analysis and Applications*. Walter de Gruyter & Co., Berlin, 1996. [8](#)
- [28] J. van Neerven, M. Veraar, and L. Weis. Stochastic integration in Banach spaces—a survey. In *Stochastic analysis: a series of lectures*, volume 68 of *Progr. Probab.*, pages 297–332. Birkhäuser/Springer, Basel, 2015. [6](#)
- [29] S. Watanabe and T. Yamada. On the uniqueness of solutions of stochastic differential equations. II. *J. Math. Kyoto Univ.*, 11:553–563, 1971. [1](#)
- [30] T. Yamada and S. Watanabe. On the uniqueness of solutions of stochastic differential equations. *J. Math. Kyoto Univ.*, 11:155–167, 1971. [1](#)
- [31] H. Zhao. Yamada-Watanabe theorem for stochastic evolution equation driven by Poisson random measure. *ISRN Probab. Stat.*, 2014:7, 2014. Id/No 982190. [1](#)

(Kistosil Fahim) DEPARTMENT OF MATHEMATICS, INSTITUT TEKNOLOGI SEPULUH NOPEMBER, KAMPUS ITS SUKOLILO, SURABAYA 60111, INDONESIA
Email address: kfahim@matematika.its.ac.id

(Erika Hausenblas) DEPARTMENT OF MATHEMATICS, MONTANUNIVERSITY LEOBEN, 8700 LEOBEN, AUSTRIA
Email address: erika.hausenblas@unileoben.ac.at

(Kenneth Hvistendahl Karlsen) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OSLO, NO-0316 OSLO, NORWAY
Email address: kennethk@math.uio.no