

# Surrogate modeling of gravitational waves microlensed by spherically symmetric potentials

Uddeepa Deka<sup>1,\*</sup>, Gopalkrishna Prabhu<sup>2,†</sup>, Md Arif Shaikh<sup>3,4,‡</sup>,  
Shasvath J. Kapadia<sup>2,§</sup>, Vijay Varma<sup>5,¶</sup> and Scott E. Field<sup>5,\*\*</sup>

<sup>1</sup>*International Centre for Theoretical Sciences, Tata Institute of Fundamental Research, Bangalore 560089, India*

<sup>2</sup>*Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411007, India*

<sup>3</sup>*Department of Physics, Vivekananda Satavarshiki Mahavidyalaya (affiliated to Vidyasagar University), Manikpara 721513, West Bengal, India*

<sup>4</sup>*Department of Physics and Astronomy, Seoul National University, Seoul 08826, Korea*

<sup>5</sup>*Department of Mathematics, Center for Scientific Computing and Visualization Research, University of Massachusetts Dartmouth, Massachusetts 02747, USA*

(Dated: January 7, 2025)

The anticipated observation of the gravitational microlensing of gravitational waves (GWs) promises to shed light on a host of astrophysical and cosmological questions. However, extracting the parameters of the lens from the modulated GWs requires accurate modeling of the lensing amplification factor, accounting for wave-optics effects. Analytic solutions to the lens equation have not been found to date, except for a handful of simplistic lens models. While numerical solutions to this equation have been developed, the time and computational resources required to evaluate the amplification factor numerically make large-scale parameter estimation of the lens (and source) parameters prohibitive. On the other hand, surrogate modeling of GWs has proven to be a powerful tool to accurately, and rapidly, produce GW templates at arbitrary points in parameter space, interpolating from a finite set of available waveforms at discrete parameter values. In this work, we demonstrate that surrogate modeling can also effectively be applied to the evaluation of the time-domain microlensing amplification factor  $\tilde{F}(t)$ , for arbitrary lens configurations. We show this by constructing  $\tilde{F}(t)$  for two lens models, viz. point-mass lens, and singular isothermal sphere. We benchmark the evaluation times, and accuracy, of the surrogate microlensed templates, and show that indeed surrogate modeling is a powerful tool to make large-scale parameter estimation in the context of microlensed GWs practicable.

## I. INTRODUCTION

The LIGO-Virgo-Kagra (LVK) gravitational-wave (GW) detector network [1–3] has detected  $\sim 90$  compact binary coalescence (CBC) events across three observing runs (O1, O2, O3) [4–7]. The overwhelming majority of these are stellar-mass binary black holes (BBHs), although binary neutron star (BNS) [8, 9] and neutron star black hole (NSBH) mergers have also been observed [10].

These observations have enabled tests of general relativity (GR) in the strong-field regime [11], provided distance-ladder independent measurements of the Hubble constant [12], given hints of the population properties of BBHs [13], and allowed probes of matter at densities not accessible in current human-made laboratories via constraints on the NS equation of state [14]. The ongoing<sup>1</sup> observing run (O4) will likely triple the number of observed events, thus further strengthening the astrophysical and cosmological inferences made from O1–O3. Moreover, the increased detections will also improve our chances of seeing a gravitationally lensed GW event, which, to date, has not been detected [15].

Gravitational lensing of GWs, akin to that of light, occurs when they encounter agglomerations of matter in their paths. However, unlike the lensing of electromagnetic (EM) waves,

that of GWs can be categorized into two regimes. If the wavelength of the GWs ( $\lambda_{\text{GW}}$ ) is significantly smaller than the gravitational radius of the lens ( $\lambda_{\text{GW}} \ll GM_{\text{L}}/c^2$ ), then two or more temporally resolved GW images will be formed, with identical frequency evolution as the source, but differing amplitudes [16–18]. The time delay of the images could range from minutes to months for galaxy-scale lenses [19–21], and weeks to years for cluster-scale lenses [22–26]. Conversely, if  $\lambda_{\text{GW}} \sim GM_{\text{L}}/c^2$ , a single modulated image with wave optics effects (such as beating and diffraction patterns) will be produced [27–34]. Lensing promises to significantly enhance the science that can be done with GWs, such as enhanced GW early-warning of electromagnetically bright CBCs [35], additional unique tests of GR [36–40], precisely measuring the expansion rate of the universe [41–46], enable constraints on the structure of the lens (including probing properties such as electric charge) [47], as well as on the fraction of dark matter in the form of massive compact objects [48, 49].

Gravitational microlensing of the LVK’s GWs falls in the wave-optics regime. Examples of corresponding microlenses include isolated intermediate-mass black holes (IMBHs), whose masses lie in the range  $M \in [10^2, 10^4]M_{\odot}$ . Currently, searches for microlensing/wave-optics patterns (see, e.g., [15]) in detected CBC events assume a point-mass lens. This assumption is made for two reasons. The first is that an exact, analytical form of the frequency domain microlensing amplification factor,  $F(f)$ , is known (see, e.g., [29]). The evaluation of this amplification factor, for arbitrary GW frequencies and

\* uddeepa.deka@icts.res.in

† gopal.prabhu@iucaa.in

‡ arifshaikh.astro@gmail.com

§ shasvath.kapadia@iucaa.in

¶ vijay.varma392@gmail.com

\*\* sfield@umassd.edu

<sup>1</sup> at the time of writing

<sup>2</sup>  $M_{\text{L}}$  is the mass of the lens,  $G$  is the Newton’s gravitational constant and  $c$  is the speed of light in vacuum.

lens parameters (viz., lens mass and impact parameter), is sufficiently rapid to attempt large scale parameter estimation runs without prohibitively taxing computational resources [50]. The second is that IMBHs as microlenses have been argued to be well-modelled by the point-mass-lens [31].

However, other models of microlenses exist, many of which do not currently have any known analytical form for  $F(f)$ . An example of particular astrophysical relevance is microlens(es) embedded in a macro potential. It has been argued that a microlens, such as a massive compact object, lying in the halo of an intervening galaxy that provides the macro-potential, could result in the production of resolvable images with wave-optics effects imprinted on each of them [51]. Moreover, multiple microlenses in the macropotential would additionally produce interference patterns between the images [52–55]. The resulting amplification factor  $F(f)$ , containing beating, diffraction and interference patterns, has no known analytical form to date.

Constructing such realistic  $F(f)$ 's requires solving the lens equation, with the appropriate superposition of lensing potentials, numerically. This has been achieved (see, e.g., [55]) using the method described in [56]. However, producing these  $F(f)$ 's, for a single set of lens parameters, typically takes several seconds or longer per waveform, making large-scale GW parameter estimation (PE) of such lens configurations unfeasible.

In this work, we adopt a widely used interpolation technique, called ‘‘surrogate modeling’’. This technique has found its application in the field of GWs for modeling waveforms in the context of CBCs [57–68]. In addition, a recent study has demonstrated the application of surrogate modeling to describe waveforms for hyperbolic encounters between black holes [69]. The main appeal of surrogate modeling lies in the fact that one can obtain a fast and accurate prediction of a high-dimensional, complicated function, at arbitrary points in parameter space, interpolating from a finite (and usually small) set of points in that space where the function is known. To achieve this, surrogate models take advantage of the underlying similarity among the numerical solutions across the parameter space. Surrogate models thus provide a drastic increase in the evaluation speed of the function, compared to slow numerical solutions, while only negligibly deviating from them. We apply surrogate modelling, *for the first time*, to the rapid and accurate construction of time domain lensing amplification factors  $\tilde{F}(t)$ . We benchmark the accuracy and production speeds of the surrogate microlensed GWs in the frequency domain.

As a proof of principle, we consider the following lens models: point-mass lens and singular isothermal sphere (SIS). We numerically evaluate the time domain amplification factor  $\tilde{F}(t)$  at a few discrete points in the lensing parameter space. From these, we construct surrogate amplification factors  $\tilde{F}^s(t)$  for each of the lensing configurations. After Fourier transforming and producing microlensed GW waveforms in the frequency domain, we evaluate the production-time and accuracy of the surrogate microlensed waveforms. We find that we’re able to achieve mismatches of the order of  $\mathcal{O}(10^{-7} - 10^{-3})$  with respect to the waveforms evaluated numerically, with evaluation times of  $\mathcal{O}(10^{-1} - 10^{-2})$ s. These benchmarking tests showcase the efficacy of surrogate modeling applied to microlensed GWs.

They also suggest that surrogate modeling can be feasibly used for GW parameter estimation in the context of GW microlensing.

The rest of the paper is organised as follows. Section II delineates the lensing wave-optics-driven amplification factor for the three lensing configurations considered in this work. It also introduces the basics of surrogate modelling and its application to the construction of time-domain lensing amplification factors. Section III presents the results which demonstrate the accuracy and evaluation speed of the surrogate microlensed waveforms. Section IV summarises the results and suggests future work.

## II. METHOD

### A. Lensing Amplification Factor

In this section, we describe the formalism for computing the lensing amplification factor  $F(f)$ , which is defined as the ratio between the lensed and the unlensed GW waveform,

$$F(f) \equiv h^{\text{SL}}(f)/h^{\text{UL}}(f). \quad (1)$$

Under the thin lens approximation, the frequency-domain amplification factor is given by the Kirchhoff diffraction integral [70]:

$$F(f) = \frac{D_S \xi_0^2 (1 + z_L) f}{c D_L D_{LS}} \frac{1}{i} \int d^2 \vec{x} \exp [2\pi i f t_d(\vec{x}, \vec{y})], \quad (2)$$

where  $\xi_0$  is a typical length scale in the lens plane,  $z_L$  is the lens redshift,  $D_L$  is the angular diameter distance to the lens,  $D_S$  is the angular diameter distance to the source and  $D_{LS}$  is the angular diameter distance between the source and the lens (see Fig. 1 for an illustration). The integral is over the dimensionless length vector on the lens plane  $\vec{x} \equiv \vec{\xi}/\xi_0$ . Similarly,  $\vec{y} \equiv (D_L/\xi_0 D_S) \vec{\eta}$  is the dimensionless impact parameter vector on the source plane indicating the position of the source. The time delay function,  $t_d$ , as measured by an observer is a combination of  $t_{\text{geom}}$  and  $t_{\text{grav}}$ , where the former is the time delay due to the extra path length of the deflected GW signal relative to an unperturbed null geodesic and the latter is the time delay due to gravitational time dilation:

$$t_d(\vec{x}, \vec{y}) \equiv t_{\text{geom}} + t_{\text{grav}} = \frac{D_S \xi_0^2 (1 + z_L)}{c D_L D_{LS}} \left[ \frac{|\vec{x} - \vec{y}|^2}{2} - \psi(\vec{x}) \right]. \quad (3)$$

Here, the lensing potential  $\psi(\vec{x})$  is the appropriately scaled, projected Newtonian potential on the two-dimensional lens plane that depends on the geometry of the lens. Further, it is convenient to introduce the dimensionless angular frequency  $w$  defined as:

$$w \equiv \frac{D_S}{c D_L D_{LS}} \xi_0^2 (1 + z_L) 2\pi f. \quad (4)$$

The amplification factor in eq.(2) then becomes:

$$F(w) = \frac{w}{2\pi i} \int d^2 \vec{x} \exp [i w \tau_d(\vec{x}, \vec{y})], \quad (5)$$




Figure 1. Lensing geometry for the source (a compact binary merger) at redshift  $z_s$ , the lens (compact object within the thin lens approximation) at redshift  $z_L$ , and the ground-based detector.  $\xi$  denotes the lens plane coordinates. The source is located at  $\vec{\eta}$  measured from the optical axis (dotted line).  $D_L$ ,  $D_S$  and  $D_{LS}$  are the angular diameter distances between the detector and the lens, between the detector and the source, and between the lens and the source, respectively. The dashed line denotes the deflected path of the incoming signal.

where  $\tau_d$  is the dimensionless time delay, given by:

$$\tau_d(\vec{x}, \vec{y}) = \frac{|\vec{x} - \vec{y}|^2}{2} - \psi(\vec{x}). \quad (6)$$

The integrand in Eq. (5) is highly oscillatory for large values of  $w\tau_d$ . With the exception of the point mass lens (PML) model, for which there exists a closed-form analytic  $F(w)$ , it is computationally expensive to solve the integral using conventional numerical integration techniques for other astrophysically relevant lens models. Therefore, we resort to computing the amplification factor in time domain, similar to the idea proposed in [56]. For this purpose we define  $\tilde{F}(t)$  as the inverse Fourier transform of  $[2\pi i F(w)/w]$ , which gives:

$$\tilde{F}(t) = \int d^2\vec{x} \delta[\tau_d(\vec{x}, \vec{y}) - t] = \frac{dS}{dt}, \quad (7)$$

where we identify that the contribution to  $\tilde{F}(t)$  comes from the area  $dS$  on the lens plane, between the curves of constant time delays  $t$  and  $t + dt$ . Therefore, the computation of  $\tilde{F}(t)$  boils down to finding areas between nearby time delay curves, which is done using the method described in [47].

We can then obtain  $F(w)$  by Fourier transforming  $\tilde{F}(t)$  as shown below:

$$F(w) = \frac{w}{2\pi i} \times \text{FFT}[\tilde{F}(t)], \quad (8)$$

where  $\text{FFT}[\cdot]$  refers to the fast Fourier transform routine. Finally, we can go from the dimensionless angular frequency  $w$  to the dimension-full frequency  $f$  by setting the length scale  $\xi_0$  in Eq. (4) appropriately.

According to Fermat's principle, in the geometric optics limit ( $\lambda_{\text{GW}} \ll GM_L/c^2$ , or,  $w \gg 1$ ), the primary contribution to Eq. (5) comes from the stationary points of  $\tau_d$ . These stationary points are the location of images, and can be determined using:

$$\vec{\nabla}_x \tau_d(\vec{x}, \vec{y}) = 0. \quad (9)$$

For axially symmetric lenses, that is,  $\psi(\vec{x}) = \psi(x)$ , it can be seen that the images are collinear with  $\vec{y}$ . Therefore, without loss of generality, we choose  $\vec{y} = (y, 0)$ . Throughout this work, we restrict ourselves to the lens parameter space which results in two images.

Below, we describe the various lens models for which we build the surrogate lensing waveforms.

### 1. Point Mass Lens

For a point mass lens (PML) with mass  $M_L$ , the projected surface mass density on the lens plane is given by [70]:

$$\Sigma(\vec{x}) = \frac{M_L}{\xi_0^2} \delta^2(\vec{x}) = \frac{M_L}{\xi_0^2} \frac{1}{2\pi x} \delta(x). \quad (10)$$

To determine the lensing potential, we solve the Poisson equation in two-dimensions:

$$\nabla_x^2 \psi(\vec{x}) = \frac{2\Sigma(\vec{x})}{\Sigma_{\text{cr}}}, \quad (11)$$

where:

$$\Sigma_{\text{cr}} \equiv \frac{c^2}{4\pi G} \frac{D_S}{D_L D_{LS}}. \quad (12)$$

If we set  $\xi_0$  to be the *Einstein radius* of the lens,

$$\xi_0^2 = 4 \frac{GM_L}{c^2} \frac{D_L D_{LS}}{D_S}, \quad (13)$$

the lensing potential for PML is found to be:

$$\psi_{\text{PML}}(x) = \ln(x). \quad (14)$$

As mentioned earlier, the amplification factor for the PML can be analytically computed [71]:

$$F(w) = \exp\left[\frac{\pi w}{4} + \frac{iw}{2} \left(\ln\left(\frac{w}{2}\right) - 2\phi_m(y)\right)\right] \times \Gamma\left(1 - \frac{i}{2}w\right) {}_1F_1\left(\frac{i}{2}w, 1; \frac{i}{2}wy^2\right), \quad (15)$$

where:

$$\phi_m(y) = \frac{(x_m - y)^2}{2} - \ln x_m, \\ x_m = \frac{y + \sqrt{y^2 + 4}}{2},$$




Figure 2. *Top panel:* Time-domain amplification factor,  $\tilde{F}(t)$ , and its transformed version  $\tilde{F}(f)$  due to a point mass lens for various impact parameters  $y$ . *Bottom panel:* Amplitude of the lensed waveforms (left) considering a GW150914-like source and the amplitude and phase of the frequency-domain amplification factor  $F(f)$  (right) for corresponding values of impact parameters computed numerically (solid lines) and with the surrogate models (dashed lines).

$\Gamma$  is the standard gamma function and  ${}_1F_1$  is the Kummer's confluent hypergeometric function of the first kind. In the geometric optics limit, the images are formed at:

$$x_{\pm} = \frac{1}{2} \left( y \pm \sqrt{y^2 + 4} \right), \quad (16)$$

where  $x_+$  is the location of the minima image and  $x_-$  is the location of the saddle image. The magnifications of these images with respect to the unlensed signal are,

$$\mu_{\pm} = \frac{1}{2} \pm \frac{y^2 + 2}{2y \sqrt{y^2 + 4}}. \quad (17)$$

The lensing time delay of the saddle image with respect to the minima image can be found to be,

$$t_{\text{peak}} = \frac{1}{2} y \sqrt{y^2 + 4} + \ln \left( \frac{\sqrt{y^2 + 4} + y}{\sqrt{y^2 + 4} - y} \right). \quad (18)$$

Fig. 2 shows the amplification factor in time domain (top right panel) and in frequency domain (bottom right panel) for various impact parameters  $y$ , computed numerically and using the surrogate model for PML (see section III for details). The resulting lensed GW waveforms are shown in the bottom left panel. The logarithmic peaks in  $\tilde{F}(t)$  on the left panel occur

at  $t = t_{\text{peak}}$  above. This feature is characteristic of the saddle image.

## 2. Singular Isothermal Sphere

The singular isothermal sphere (SIS) models a spherically symmetric mass agglomeration with its constituents behaving like an ideal gas in hydrostatic equilibrium. The surface mass density of the SIS is determined by its velocity dispersion  $\sigma_v$  and is expressed as [70]:

$$\Sigma(\vec{x}) = \frac{\sigma_v^2}{2G\xi_0 x}. \quad (19)$$

By choosing:

$$\xi_0 = \frac{4\pi\sigma_v^2 D_L D_{LS}}{c^2 D_S}, \quad (20)$$

and using equations (11), (12) and (19), the lensing potential for SIS can be derived as:

$$\psi_{\text{SIS}}(x) = x \quad (21)$$

Unlike the PML, SIS does not have a closed-form analytic solution for  $F(f)$ . However, there exists a summation form for




Figure 3.  $\tilde{F}(t)$  for various impact parameters  $y$  for the SIS lens model computed using the numerical method (solid lines) and the surrogate model (dashed lines).

the amplification factor for SIS [72],

$$F(w) = \exp\left(\frac{i}{2}wy^2\right) \sum_{n=0}^{\infty} \frac{\Gamma(1+n/2)}{n!} (2w)^{n/2} \exp\left(\frac{i3\pi n}{4}\right) {}_1F_1\left(1+\frac{n}{2}, 1; -\frac{i}{2}wy^2\right). \quad (22)$$

Fig. 3 illustrates the time-domain amplification factor,  $\tilde{F}(t)$ , computed for various impact parameters using both the numerical method and the surrogate model for SIS. For  $y < 1$ , two images form in the geometric optics limit at the locations:

$$x_{\pm} = y \pm 1. \quad (23)$$

Similar to PML, one of the images (at  $x_+$ ) is a minima and the other (at  $x_-$ ) is a saddle image. The magnifications of the minima and the saddle images with respect to the unlensed waveform are,

$$\mu_{\pm} = 1 \pm \frac{1}{y} \quad (24)$$

respectively, for  $y < 1$ . The lensing time delay between the images is,

$$t_{\text{peak}} = 2y, \quad (25)$$

which is where the logarithmic peak due to the saddle image can be seen. However, for  $y > 1$ , only a single image is present, which is the *weak lensing* regime. In this work, we restrict to the two image parameter space for the SIS lenses.

We define the characteristic mass scale of the SIS lens as the mass contained within  $\xi_0$ , and is given by:

$$M_L = \frac{4\pi^2 \sigma_v^4 D_L D_{LS}}{c^2 G D_S}. \quad (26)$$

In the following sections, the lens mass of the SIS model will refer to the mass defined in Eq. (26) above.





Figure 4. Illustration of improvement with peak reconstruction. *Top*: Time domain amplification factor with and without peak reconstruction for PML lens. The dashed vertical lines denote the window in which peak correction is applied. *Bottom*: Amplitude of the frequency domain amplification factor with and without peak reconstruction. Additionally, the analytically computed  $|F(f)|$  is shown for comparison.

## B. Peak reconstruction

Accurate computation of the time-domain amplification factor via Eq. (7) demands sufficiently high resolution in the numerical integration. Nonetheless, even at high resolution, the numerical solution may struggle to capture the peak of  $\tilde{F}(t)$  due to the logarithmic divergence at the peak. This limitation can impact the accuracy of the amplitude of the frequency-domain amplification factor  $F(w)$ , as defined in Eq. (8) (see Fig. 4 for an illustration).

To overcome the limitation of numerical methods in resolving the peak feature, we utilize the analytical expressions for the peak location and the behavior of  $\tilde{F}(t)$  in the vicinity of the peak given by [56],

$$F_{\text{approx}}(t) = \sqrt{\mu_+} - \frac{1}{\pi} \sqrt{|\mu_-|} \log(|t - t_{\text{peak}}|), \quad (27)$$

where the values of  $\mu_+$ ,  $\mu_-$  and  $t_{\text{peak}}$  are given by Eq. (17), (18) for PML and (24), (25) for SIS lenses. We use the following steps to reconstruct the peak features by smoothly hybridizing the numerical and analytical approximation on either side of the peak:

- (a) Identify the peak location,  $t_{\text{peak}}$ , using the analytical expression given in Eq. (18) for PML and Eq. (25) for SIS.
- (b) Split the numerical  $\widetilde{F}(t)$  at  $t_{\text{peak}}$  into a left segment,  $F_{\text{num}}^{\text{l}}(t)$ , and a right segment,  $F_{\text{num}}^{\text{r}}(t)$ .
- (c) Using the analytical approximation of  $\widetilde{F}(t)$  near the peak (Eq. (27)) evaluate the left and right approximations,  $F_{\text{approx}}^{\text{l}}(t)$  for  $t < t_{\text{peak}}$  and  $F_{\text{approx}}^{\text{r}}(t)$  for  $t > t_{\text{peak}}$ .
- (d) Define a reconstruction window,  $[t_{\text{match}}^{\text{l}}, t_{\text{match}}^{\text{r}}]$ , around  $t_{\text{peak}}$  to match and reconstruct the peak. A window spanning a time interval equal to 5% on each side of  $t_{\text{peak}}$  is found to be effective.
- (e) Rescale the approximated segments  $F_{\text{approx}}^{\text{l}}(t)$  and  $F_{\text{approx}}^{\text{r}}(t)$  to match the numerical  $\widetilde{F}(t)$  values at  $t_{\text{match}}^{\text{l}}$  and  $t_{\text{match}}^{\text{r}}$ , respectively.
- (f) Replace the portion of the numerical  $\widetilde{F}(t)$  within the reconstruction window with the rescaled approximated segments. This results in the reconstructed  $\widetilde{F}(t)$ :

$$\widetilde{F}(t) = \begin{cases} F_{\text{num}}^{\text{l}}(t), & t \leq t_{\text{match}}^{\text{l}} \\ F_{\text{approx}}^{\text{l}}(t), & t_{\text{match}}^{\text{l}} < t < t_{\text{peak}} \\ F_{\text{approx}}^{\text{r}}(t), & t_{\text{peak}} < t < t_{\text{match}}^{\text{r}} \\ F_{\text{num}}^{\text{r}}(t), & t \geq t_{\text{match}}^{\text{r}} \end{cases}.$$

Fig. 4 shows that reconstructing the peak in the  $\widetilde{F}(t)$  (top panel) consistently provides a better match (bottom panel) with the analytic expression (Eq. (15)) as compared to the vanilla numerical  $\widetilde{F}(t)$ .

### C. Time-domain amplitude transformation

The logarithmic singularity at  $t_{\text{peak}}$  presents a challenge for accurately modeling the region near the peak of  $\widetilde{F}(t)$  using a surrogate. To address this, we apply a transformation to the reconstructed  $\widetilde{F}(t)$  (see Sec. II B for details on the reconstruction) and construct a surrogate model based on this peak-reconstructed, transformed time-domain amplification factor,  $\widetilde{\mathbb{F}}(t)$ . The following transformation suppresses the peak contribution:

$$\widetilde{\mathbb{F}}(t) = 1 - \exp \left[ -\pi \frac{(\widetilde{F}(t) - 1)}{\sqrt{\mu_-}} \right], \quad (28)$$

where  $\mu_-$  is the magnification due to the saddle image, given by Eq. (17) for PML and (24) for SIS lens models. The transformation effectively gets rid of the singularity, and makes the derivative of the amplification factor continuous. The top left panel in Fig. 2 compares the transformed  $\widetilde{\mathbb{F}}(t)$  for PML computed for various impact parameters using the numerical method and the surrogate model for PML. After evaluating the surrogate, an inverse transformation is applied to obtain the time domain amplification factor  $\widetilde{F}(t)$  from  $\widetilde{\mathbb{F}}(t)$ .

### D. Surrogate Modeling

In this section, we introduce the basics of surrogate modeling, its application in computing the lensing amplification factor for lensed GW waveforms and the setup used in this work. Surrogate modeling has been used extensively in building fast and accurate GW waveforms, especially in building surrogate models for Numerical Relativity (NR) waveforms [57, 73]. Surrogate modeling is most beneficial in cases where the number of waveforms are limited because of prohibitively large computation time resulting in a small training data set.

In the following, we will briefly discuss how surrogate modeling reduces the dimensionality of the problem by using reduced basis and empirical interpolation method. These two methods ensure that with only a limited number of waveforms (chosen appropriately), one can build an accurate model for predicting solutions outside the training set of parameters. Because these two steps can be precomputed, it also ensures that the model, evaluated at arbitrary points in parameter space, is orders of magnitude faster than numerical solutions.

#### 1. Surrogate modeling basics

We follow the methods described in Sec. II and III of Ref. [57] and refer the reader to it for a detailed overview of building a surrogate model. In the following, we briefly outline building a surrogate model.

1. The first step is to find a minimal set of solutions called the *reduced basis* (RB) [74] in terms of which solutions at other parameters can be expressed. Given a known set of solutions (also called the training set), the RB is found using a greedy search algorithm. For example, with a RB with  $m$  basis elements,  $\{e_i\}_{i=1}^m$ , a solution  $\widetilde{\mathbb{F}}(t; \lambda)$  in the training set at the lens parameter values  $\lambda$  can be well approximated as:

$$\widetilde{\mathbb{F}}(t; \lambda) \approx \sum_{i=1}^m c_i(\lambda) e_i(t). \quad (29)$$

This approximation is also good at other parameters outside the training set as long as the training set is dense enough. However, getting a dense training set can itself be very expensive and sometimes prohibitive. In such cases, one can build the training set using a greedy search method which will provide an optimal training set for a given desired accuracy of the surrogate prediction. We come back to this in sec. II D 3 where we discuss the surrogate modeling setup specific to our problem.

Since the number of RB,  $m$ , is usually very small compared to the number of solutions in the training set, say  $M$ , the dimensionality of the problem reduces by a factor of  $m/M$  which is usually  $\ll 1$ .

2. The next step is to find the most representative times, the empirical times or nodes  $\{T_i\}_{i=1}^m$  to construct an interpolant in time using *empirical interpolant method*

(EIM) [57, 75–77] for a given parameter  $\lambda$ :

$$\mathcal{I}_m[\widetilde{\mathbb{F}}](t; \lambda) = \sum_{i=1}^m C_i(\lambda) e_i(t). \quad (30)$$

The coefficients  $C_i(\lambda)$  are defined by requiring that the interpolant becomes equal to the value of the solutions at the empirical nodes:

$$\sum_{i=1}^m C_i(\lambda) e_i(T_j) = \widetilde{\mathbb{F}}(T_j; \lambda), \quad j = 1, \dots, m \quad (31)$$

or:

$$\sum_{i=1}^m V_{ji} C_i(\lambda) = \widetilde{\mathbb{F}}(T_j; \lambda), \quad j = 1, \dots, m \quad (32)$$

The coefficients  $C_i(\lambda)$  can be obtained by solving the above  $m$ -by- $m$  system with interpolation matrix:

$$V \equiv \begin{pmatrix} e_1(T_1) & e_2(T_1) & \dots & e_m(T_1) \\ e_1(T_2) & e_2(T_2) & \dots & e_m(T_2) \\ \vdots & \vdots & \ddots & \vdots \\ e_1(T_m) & e_2(T_m) & \dots & e_m(T_m) \end{pmatrix}. \quad (33)$$

Thus:

$$C_i = \sum_{j=1}^m (V^{-1})_{ij} \widetilde{\mathbb{F}}(T_j; \lambda). \quad (34)$$

Substituting Eq. (34) in Eq. (30), the empirical interpolant can be rewritten as:

$$\mathcal{I}_m[\widetilde{\mathbb{F}}](t; \lambda) = \sum_{j=1}^m B_j(t) \widetilde{\mathbb{F}}(T_j; \lambda), \quad (35)$$

with:

$$B_j(t) \equiv \sum_{i=1}^m e_i(t) (V^{-1})_{ij}. \quad (36)$$

Note that  $B_j(t)$  is independent of  $\lambda$  and can be constructed offline using only the information contained in the RB. Similar to the RB, the empirical nodes are also obtained using a greedy search algorithm. The number of empirical nodes is the same as the number of RB. Since the number of RB is usually minimal, it ensures that the EIM is using only a very small number of points in time instead of using a uniform grid of points which could significantly increase computational cost. Thus the dimensionality of the problem scales down by a factor of  $m/L$ , where  $m$  is the number of RB (= number of empirical nodes) and  $L$  is the number of points in the case of an interpolant on a uniform grid.

The use of RB and EIM reduces the dimensionality of the problem by a factor of  $(m \times m)/(M \times L) \ll 1$ .

3. With the interpolant  $\mathcal{I}_m[\widetilde{\mathbb{F}}](t; \lambda)$  in Eq. (35) at hand, we need to know the values of the solutions at the empirical nodes  $\{T_i\}_{i=1}^m$ . Therefore, the third step is to construct parametric fits at each empirical node across the parameter space.
4. Finally, the surrogate model prediction at a new parameter  $\lambda_*$  outside the training data set is computed by first using the parametric fits (constructed in the third step) to evaluate the values of  $\widetilde{\mathbb{F}}(t; \lambda_*)$  at the empirical nodes  $\{T_i\}_{i=1}^m$  and then building the interpolant  $\mathcal{I}_m[\widetilde{\mathbb{F}}](t; \lambda)$  using Eq. (35).

The first three steps are performed offline (i.e. precomputed), which drastically expedites the evaluation of the surrogate model at arbitrary points in parameter space.

## 2. Surrogate for microlensed waveforms

We apply the technique of surrogate modeling to obtain a model of the GWs waveform undergoing microlensing. However, since the effect of microlensing is a frequency-dependent modulation of the GW amplitude as explained in Sec. II A, one needs only to model this amplification factor provided a fast unlensed model exists already. Therefore, instead of modeling the microlensed waveform directly, we build a surrogate model  $\widetilde{\mathbb{F}}^S(t)$  for the transformed lens amplification factor  $\widetilde{\mathbb{F}}(t)$  in time domain. We then obtain the frequency-dependent amplification factor  $F^S(f)$  by taking a Fourier transform of the back-transformed  $\widetilde{\mathbb{F}}^S(t)$  (see Sec. II C). The surrogate microlensed waveform  $\hat{h}^{\text{SL}}(f)$  is then obtained by multiplying the unlensed waveform  $\hat{h}^{\text{UL}}(f)$  by the frequency-dependent amplification factor  $F^S(f)$ :

$$\hat{h}^{\text{SL}}(f) = F^S(f) \times \hat{h}^{\text{UL}}(f). \quad (37)$$

## 3. Surrogate modeling setup

The RB is found by using a greedy algorithm that guarantees that the error  $\sigma_m$

$$\sigma_m \equiv \max_{\lambda} \min_{c_i \in C} \left\| \widetilde{\mathbb{F}}(\cdot; \lambda) - \sum_{i=1}^m c_i(\lambda) e_i(\cdot) \right\| \quad (38)$$

$$\text{with } \left\| \widetilde{\mathbb{F}}(\cdot; \lambda) \right\| \equiv \int_{t_{\min}}^{t_{\max}} \left| \widetilde{\mathbb{F}}(t; \lambda) \right|^2 dt \quad (39)$$

associated with the approximation in Eq. (29) (minimized over the coefficients  $c_i$  and maximized over the parameter  $\lambda$ ) is less than a desired value, usually called the basis tolerance. Using an optimal basis tolerance when building the RB is crucial in making a surrogate model as fast as possible. Using a very small basis tolerance may result in a larger number of bases which in turn results in a larger number of empirical time nodes. While evaluating the surrogate model, a significant portion of the computation time involves evaluating the fits at the empirical nodes. Thus, a smaller basis tolerance is likely to make

the surrogate model slower. Therefore, we use an optimal basis tolerance of  $\sim 5 \times 10^{-4}$  for building the RB so that the surrogate models achieve the required accuracy compared to the microlensed waveform computed using the numerical solutions without causing any overfitting. To obtain the training data set, we employ a greedy search algorithm as described in [58]. In the greedy search, we start with an initial training set consisting of only the corner points of the parameter range of interest. At each step, we build the surrogate model with a specific basis tolerance ( $\sim 5 \times 10^{-4}$  in our case) and then validate it against a large number of numerical solutions generated randomly within the parameter range. This validation set excludes the parameters already in the training set. The validation involves measuring the normalised  $L_2$  error between the surrogate prediction  $\widetilde{\mathbb{F}}^S(t)$  and the numerical solution  $\widetilde{\mathbb{F}}(t)$  defined as:

$$L_2 = \frac{\sqrt{\int_{t_1}^{t_2} |\widetilde{\mathbb{F}}^S(t) - \widetilde{\mathbb{F}}(t)|^2 dt}}{\sqrt{\int_{t_1}^{t_2} |\widetilde{\mathbb{F}}(t)|^2 dt}}, \quad (40)$$

where  $t_1$  and  $t_2$  denote the range of the time segment within which the error is computed.

We pick the parameter where the  $L_2$  error is the maximum and add it to the training set to be used in the next step and repeat the procedure until we achieve the maximum  $L_2$  error to be smaller than a desired value which in our case is  $10^{-2}$ . With such greedy search, we find our training data sets that consist of  $\sim 35 - 100$  data points depending on the particular microlens model.

For constructing the fits at the empirical nodes, we use the Gaussian Process Regression (GPR) method as described in the supplemental material of [65].

In a nutshell, the training data  $\widetilde{\mathbb{F}}(t)$  is obtained by applying a peak reconstruction followed by a transformation operation on  $\widetilde{F}(t)$ . This training data is used to build the surrogate  $\widetilde{\mathbb{F}}^S(t)$ . Then an inverse transformation is applied followed by a peak reconstruction to obtain  $\widetilde{F}^S(t)$ .

### III. RESULTS

In order to validate our model, we compute the *match*  $\mathcal{M}$  between the surrogate microlensed waveform model and the corresponding numerical model used for training. Match between two given waveforms  $h_1$  and  $h_2$  is defined as [78]:

$$\mathcal{M} \equiv \arg \max_{t_c, \phi_c} \frac{\langle h_1, h_2 \rangle}{\sqrt{\langle h_1, h_1 \rangle \langle h_2, h_2 \rangle}}, \quad (41)$$

where  $t_c$  and  $\phi_c$  are the coalescence time and phase, respectively, and the overlap  $\langle h_1, h_2 \rangle$  is:

$$\langle h_1, h_2 \rangle = 4\mathcal{R} \int_{f_{\text{low}}}^{f_{\text{high}}} \frac{\widetilde{h}_1^*(f) \widetilde{h}_2(f)}{S_n(f)} df. \quad (42)$$

Here,  $f$  is the frequency,  $\widetilde{h}_{1,2}$  are the Fourier transforms of the corresponding time domain waveforms,  $\star$  represents the com-

plex conjugate and  $\mathcal{R}$  represents the real part.  $f_{\text{low}}$  and  $f_{\text{high}}$  denote the lower and higher frequency cutoff of the detector band and  $S_n(f)$  denote the one-sided power spectral density (PSD) of the detector noise. We use `pycbc.filter.matchedfilter` module from the PyCBC package [79] to compute the match. We make use of the next-generation ground-based detector Einstein Telescope (ET) PSD, pertaining to the ET design sensitivity [80]. The reason for choosing this particular PSD is two fold: the frequency bin is larger than the current ground based detectors, giving rise to conservative mismatches; and statistically, we expect to see significant number of lensed GW events during the observing runs of next-generation detectors such as ET.

Note that, the numerical amplification factors used to train the model for PML and SIS are calibrated to the analytic form given in Eq. (15) for PML and the summation form given in Eq. (22) for SIS, such that the maximum mismatch ( $1 - \mathcal{M}$ ) between the numerically computed lensed waveform and the analytically obtained lensed waveform is  $< 10^{-4}$  for all possible points in the parameter space.

We present mismatches between the surrogate microlensed GWs and the numerically evaluated microlensed GWs used to construct the surrogate model. We ensure that the lensing parameters used for training the model, and those used for testing (i.e, evaluating the mismatches), are different. The lensing parameters are chosen over a grid, and the mismatches evaluated for each point on that grid, as well as different unlensed CBC waveforms. Evaluation times per microlensed frequency-domain waveform are also measured. These are then compared with corresponding times pertaining to other methods, viz., a frequency domain interpolation method employed by the LVK collaboration to search for GWs microlensed by PMLs, as well as numerical methods to acquire amplification factor for the SIS lens. The waveforms are sampled at frequencies with bin width,  $\delta f = 1/32$ , resulting in the total number of waveform evaluation points in the frequency domain to range from  $\sim 1.3 \times 10^4$  (for total binary mass  $M_{\text{tot}} = 100M_{\odot}$ ) to  $\sim 6.5 \times 10^4$  (for total binary mass  $M_{\text{tot}} = 20M_{\odot}$ ).

The surrogate model for PML is created using 27 basis vectors with basis tolerance of  $5 \times 10^{-4}$ , while for SIS it is created using 15 basis vectors with the same basis tolerance. We use the `numpy.fft` module from NumPy [81] package to compute FFT. All computations related to creating and evaluating the surrogate model were performed on a 32-core workstation with Intel Xeon E5-2650 v2 chip, OS: Linux. Note that no parallelization was used during evaluation; thus, the computations effectively used a single core.

*a. Point Mass Lens (PML):* We create a surrogate model for the transformed time-domain amplification factor  $\widetilde{\mathbb{F}}(t)$ . It is a single parameter model parameterized by the impact parameter  $y$ . Note that the lens mass  $M_L$  sets the time scale while converting the dimensionless angular frequency  $w$  to the dimension-full frequency  $f$ . Therefore, it can be factored out while creating the model. We use the numerically computed  $\widetilde{\mathbb{F}}(t)$  (as shown in Fig. 2) to train our model. Fig. 5 shows the improvement in the maximum  $L_2$  between the numerically obtained  $\widetilde{\mathbb{F}}(t)$  and the surrogate  $\widetilde{\mathbb{F}}^S(t)$  with an increase in the number of greedy parameters. The error versus greedy para-






Figure 5. Improvement in the  $L_2$  with increasing number of greedy parameters for the PML model (top) and the SIS model (bottom). The  $L_2$  is computed between the validation set and the surrogate model.

rameters curve plateaus for greedy parameter counts greater than approximately 30 for PML. Therefore, we use the first 35 greedy parameters to construct the surrogate model. The greedy parameter space of  $y$  spans from  $y = 0.1$  to  $y = 2$ .

The current LVK microlensing analysis makes use of an interpolation table (“lookup table”) in  $w - y$  plane to compute  $F(w)$  given in Eq. (15) [15]. Fig. 6 shows the evaluation time for the surrogate model compared to the lookup table for the same number of evaluation points. We also indicate the time taken to generate the numerically obtained  $F(f)$ . As seen from the figure, the surrogate performs faster than the lookup table throughout the parameter space. Note that a part of the lookup table evaluation uses information from the geometric optics approximation, whereas the numerical method and therefore, the surrogate model is based completely on the wave optics computation.

Fig. 7 shows the mismatch between the surrogate microlensed waveform and the microlensed waveform obtained using the numerical method for a GW150914-like signal for various values of  $M_L$  and  $y$ . The unlensed waveform is generated using the IMRPhenomXP approximant [82]. As evident from the figure, the mismatches are better than 0.03%.

Further, we show the dependence of the accuracy of our model on the source mass of the binary by simulating equal component mass, non-spinning, binary black hole (BBH) unlensed waveforms for different total binary masses ( $M_{\text{tot}} =$




Figure 6. Evaluation time for waveforms lensed by point mass lens for various impact parameters  $y$  for a CBC source with  $M_{\text{tot}} = 20M_{\odot}$ . We compare the evaluation times for the numerical amplification factor ( $F(f)$ ), surrogate model in the time domain ( $\mathbb{F}^S(t)$ ) and after FFT ( $F^S(f)$ ) with the time taken using the lookup table.




Figure 7. Mismatch in the  $M_L - y$  plane between the numerical method and the surrogate model for a lensed GW150914-like signal. We assume the lens to be a point mass lens.

{20, 40, 60, 80, 100} $M_{\odot}$ ) by computing mismatch in the  $M_L - y$  grid spanning  $100 \leq M_L/M_{\odot} \leq 1000$  and  $0.1 \leq y \leq 2$ . We consider equal mass binaries. The top left panel in Fig. 8 shows the histogram of mismatch values for the different source masses between the lensed waveforms generated using the numerical method and the surrogate model. As can be seen, the mismatches are lower than 0.05%. Moreover, the top right panel compares the evaluation times for the surrogate with the lookup table for the same grid and source masses.

*b. Singular Isothermal Sphere (SIS):* Similar to the PML model, this model is also a single-parameter model parameterized by  $y$ . As shown in Fig. 5, the error versus greedy parameters curve plateaus for greedy parameter counts greater than approximately 85 for SIS. Therefore, to construct the surrogate model for SIS, we use the first 100 greedy parameters. The greedy parameter space of  $y$  spans from  $y = 0.01$  to  $y = 0.95$ . We compute mismatch and compare evaluation times




Figure 8. *Top panel: (Left)* Histograms of mismatch between the lensed waveforms due to PML computed using the surrogate model and the numerical method. Each histogram corresponds to a different source with the total source mass,  $M_{\text{tot}} = \{20, 40, 60, 80, 100\}M_{\odot}$ . The mismatches are computed in a grid of  $M_L - y$ , where  $100 \leq M_L/M_{\odot} \leq 1000$  and  $0.1 \leq y \leq 2$ . *(Right)* Histograms of the evaluation time for the surrogate model (solid) and for the lookup table (dashed) in the same  $M_L - y$  grid for PML. *Bottom panel: (Left)* Similar to the top left plot but for SIS lens with the range of  $y$  being  $0.01 \leq y \leq 0.95$ . *(Right)* Histograms of the evaluation time for the surrogate model (solid) and for the numerical method (dashed) in the same  $M_L - y$  grid for SIS.

between the numerically obtained microlensed waveforms and the surrogate microlensed waveforms. The bottom left panel in Fig. 8 shows the histogram of mismatch values for various source masses, similar to the case for PML. Here, we restrict the  $y$  values between  $0.01 \leq y \leq 0.95$  because for  $y > 1$  we are no longer in the two-image regime. Similar to PML, the mismatch values are within 0.05%, showing very promising results. The bottom right panel compares the evaluation times for the surrogate with the numerically obtained microlensed waveform. The surrogate model is evidently faster than the numerically obtained microlensed waveforms by orders of magnitude.

#### IV. SUMMARY AND DISCUSSION

Gravitational microlensing of GWs is likely to be detected in the near future, if not in O4, then in forthcoming observing runs of the LVK detector network and the next generation of ground based detectors. Such detections promise to shed light on a host of questions pertaining to astrophysics, cosmology and fundamental physics. Probing for signatures of microlensing-related wave-optics effects in detected GW events requires

large scale Bayesian PE runs, whose feasibility crucially relies on accurate and rapidly producible microlensed waveforms. Apart from GWs microlensed by a PML and a few other simplistic lensing models, no closed form analytical solutions to the lensing equation exist. Numerical methods are therefore the only recourse to generate microlensed waveforms for most realistic lensing configurations.

Evaluating microlensed GWs numerically is time consuming, with waveform generation taking up to several seconds (per waveform). Indeed, even efficient numerical methods (see, e.g: [83]) are likely to be too slow to enable large-scale PE campaigns in a cost-effective manner. One way to mitigate this issue is to use interpolation, where the microlensing amplification factor is evaluated at a few discrete points in the parameter space of the lens, and then interpolated to any arbitrary point. This has been attempted for the PML in the frequency domain, and found to be faster than directly evaluating the complicated amplification factor analytically [15]. It has also been attempted for other spherically symmetric lensing configurations in the time domain, where the amplification factor is then Fourier transformed to the frequency domain [84]. Both these methods require reading-off precomputed interpolation data from a table, and are referred to in this work

as the “lookup table” methods. For other efficient methods for rapid evaluation of lensing amplification factors, we refer the reader to [85].

Given the impressive success of surrogate modelling to accurately and rapidly produce (unlensed) CBC templates [57, 59–66], in this work, we assess its applicability in the context of generating microlensed GWs. We construct surrogate models of GWs microlensed by a PML and SIS lens. We consider a range of realistic lens masses and impact parameters. We achieve this by building time-domain surrogate amplification factors, Fourier transforming them, and then multiplying them with the unlensed frequency-domain GW waveform.

To assess the performance of the surrogate models, we compute mismatches between the surrogate microlensed waveforms and the corresponding, numerically evaluated, waveforms that were used to build the surrogate models, ensuring that the mismatches are not evaluated at lens parameter values used for surrogate construction. We do so over a grid of lens parameter values, and find that we’re able to achieve mismatches in the range  $\sim 10^{-8} - 10^{-4}$ . Moreover, we measure the evaluation times of the microlensed waveforms in the frequency domain and find that they’re typically  $\mathcal{O}(10^{-2} - 10^{-1})$ s for number of sample points ranging from  $\sim (1.3 - 6.5) \times 10^4$ . These benchmarking tests demonstrate the power of surrogate modelling applied to microlensed GWs, suggesting that large-scale PE runs can be feasibly conducted using surrogate amplification factors.

It is worth pointing out that the rate-determining step in the computation of surrogate microlensed GWs in the frequency domain is, in fact, the Fourier transform, for which we employ NumPy’s FFT package [81]. The evaluation times of the time-domain surrogate waveforms are smaller than the FFT computation by a couple of orders of magnitude. More sophisticated FFT algorithms may further reduce the evaluation times. Nevertheless, we still find that, for the PML, the microlensed waveforms computed using the lookup table method in the frequency domain is still slower than the corresponding surrogate waveforms by about an order of magnitude for large portions of the lensing parameter space considered.

The main advantage of surrogate modelling of microlensed GWs is that it does not intrinsically rely on any symmetries of the lensing potential. Indeed, if numerical solutions are able to provide time-domain amplification factors, it should be

possible to construct corresponding surrogate models. Thus, in principle, our work should be readily extendable to asymmetric lensing configurations that better model those found in nature, such as a microlens embedded in a macropotential (e.g: an intermediate mass black hole inside a galaxy-scale lens). However, constructing surrogate models for realistic lenses needs to mitigate an important problem.

Caustics – curves in the source plane where the image magnification formally diverges – demarcate distinct regions in that plane. These regions differ from each other in the number of images<sup>3</sup> that will be produced if the source is positioned in those regions. Abrupt changes in the number of images when crossing over caustics correspond to changes in the shape of the time-domain amplification factor as a function of the impact parameter that are not smooth. Indeed, even in the case of SIS lens, we restricted ourselves to lens parameters that ensure that exactly two images are produced. A possible workaround is to build separate surrogate models for each region in the lens plane containing a fixed number of images. However, an additional challenge pertaining to this workaround is to appropriately model the shape of the caustics. This should be straightforward for lensing potentials with symmetries, but might not be so for asymmetric lens systems, especially those that are an aggregation of multiple microlensing potentials embedded in a macropotential. We leave the surrogate modelling of such lensing configurations for future work.

## ACKNOWLEDGMENTS

We are grateful to Tousif Islam for the careful review of the manuscript and useful comments. We thank the members of the astrophysical relativity group at ICTS for their valuable input. We acknowledge the support of the Department of Atomic Energy, Government of India, under project no. RTI4001. M.A.S.’s research was supported by the National Research Foundation of Korea under grant No. NRF-2021M3F7A1082056. S.J.K gratefully acknowledges support from SERB grant SRG/2023/000419. All computations were performed using the computational facilities at the International Centre for Theoretical Sciences and the Inter-University Centre for Astronomy and Astrophysics.

- 
- [1] J. Aasi *et al.* (LIGO Scientific), “Advanced LIGO,” *Class. Quant. Grav.* **32**, 074001 (2015), arXiv:1411.4547 [gr-qc].
- [2] F. Acernese *et al.* (Virgo), “Advanced Virgo: a second-generation interferometric gravitational wave detector,” *Class. Quant. Grav.* **32**, 024001 (2015), arXiv:1408.3978 [gr-qc].
- [3] T. Akutsu *et al.* (KAGRA), “Overview of KAGRA: Detector design and construction history,” *PTEP* **2021**, 05A101 (2021), arXiv:2005.05574 [physics.ins-det].

- [4] B. P. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-1: A Gravitational-Wave Transient Catalog of Compact Binary Mergers Observed by LIGO and Virgo during the First and Second Observing Runs,” *Phys. Rev. X* **9**, 031040 (2019), arXiv:1811.12907 [astro-ph.HE].
- [5] R. Abbott *et al.* (LIGO Scientific, Virgo), “GWTC-2: Compact Binary Coalescences Observed by LIGO and Virgo During the First Half of the Third Observing Run,” *Phys. Rev. X* **11**, 021053 (2021), arXiv:2010.14527 [gr-qc].
- [6] R. Abbott *et al.* (LIGO Scientific, VIRGO), “GWTC-2.1: Deep extended catalog of compact binary coalescences observed by LIGO and Virgo during the first half of the third observing run,”

---

<sup>3</sup> in the geometric optics limit

- Phys. Rev. D **109**, 022001 (2024), arXiv:2108.01045 [gr-qc].
- [7] R. Abbott *et al.* (KAGRA, VIRGO, LIGO Scientific), “GWTC-3: Compact Binary Coalescences Observed by LIGO and Virgo during the Second Part of the Third Observing Run,” *Phys. Rev. X* **13**, 041039 (2023), arXiv:2111.03606 [gr-qc].
- [8] Benjamin P. Abbott *et al.* (LIGO Scientific, Virgo), “GW170817: Observation of Gravitational Waves from a Binary Neutron Star Inspiral,” *Phys. Rev. Lett.* **119**, 161101 (2017), arXiv:1710.05832 [gr-qc].
- [9] B.P. Abbott *et al.* (LIGO Scientific, Virgo), “GW190425: Observation of a Compact Binary Coalescence with Total Mass  $\sim 3.4M_{\odot}$ ,” *Astrophys. J. Lett.* **892**, L3 (2020), arXiv:2001.01761 [astro-ph.HE].
- [10] R. Abbott *et al.* (LIGO Scientific, KAGRA, VIRGO), “Observation of Gravitational Waves from Two Neutron Star–Black Hole Coalescences,” *Astrophys. J. Lett.* **915**, L5 (2021), arXiv:2106.15163 [astro-ph.HE].
- [11] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), “Tests of General Relativity with GWTC-3,” (2021), arXiv:2112.06861 [gr-qc].
- [12] R. Abbott *et al.* (LIGO Scientific, Virgo, KAGRA), “Constraints on the Cosmic Expansion History from GWTC-3,” *Astrophys. J.* **949**, 76 (2023), arXiv:2111.03604 [astro-ph.CO].
- [13] R. Abbott *et al.* (KAGRA, VIRGO, LIGO Scientific), “Population of Merging Compact Binaries Inferred Using Gravitational Waves through GWTC-3,” *Phys. Rev. X* **13**, 011048 (2023), arXiv:2111.03634 [astro-ph.HE].
- [14] B. P. Abbott *et al.* (LIGO Scientific, Virgo), “GW170817: Measurements of neutron star radii and equation of state,” *Phys. Rev. Lett.* **121**, 161101 (2018), arXiv:1805.11581 [gr-qc].
- [15] R. Abbott *et al.* (LIGO Scientific, VIRGO, KAGRA), “Search for gravitational-lensing signatures in the full third observing run of the LIGO-Virgo network,” (2023), arXiv:2304.08393 [gr-qc].
- [16] Yun Wang, Albert Stebbins, and Edwin L. Turner, “Gravitational lensing of gravitational waves from merging neutron star binaries,” *Phys. Rev. Lett.* **77**, 2875–2878 (1996).
- [17] Liang Dai and Tejaswi Venumadhav, “On the waveforms of gravitationally lensed gravitational waves,” (2017), arXiv:1702.04724 [gr-qc].
- [18] Jose María Ezquiaga, Daniel E. Holz, Wayne Hu, Macarena Lagos, and Robert M. Wald, “Phase effects from strong gravitational lensing of gravitational waves,” *Phys. Rev. D* **103**, 064047 (2021).
- [19] Ken K. Y. Ng, Kaze W. K. Wong, Tom Broadhurst, and Tjonnie G. F. Li, “Precise ligo lensing rate predictions for binary black holes,” *Phys. Rev. D* **97**, 023012 (2018).
- [20] Shun-Sheng Li, Shude Mao, Yuetong Zhao, and Youjun Lu, “Gravitational lensing of gravitational waves: A statistical perspective,” *Monthly Notices of the Royal Astronomical Society* **476**, 2220–2229 (2018).
- [21] Masamune Oguri, “Effect of gravitational lensing on the distribution of gravitational waves from distant binary black hole mergers,” *Monthly Notices of the Royal Astronomical Society* **480**, 3842–3855 (2018).
- [22] Graham P Smith, Mathilde Jauzac, John Veitch, Will M Farr, Richard Massey, and Johan Richard, “What if ligo’s gravitational wave detections are strongly lensed by massive galaxy clusters?” *Monthly Notices of the Royal Astronomical Society* **475**, 3823–3828 (2018).
- [23] Rory JE Smith, Gregory Ashton, Avi Vajpeyi, and Colm Talbot, “Massively parallel bayesian inference for transient gravitational-wave astronomy,” *Monthly Notices of the Royal Astronomical Society* **498**, 4492–4502 (2020).
- [24] Graham P. Smith, Andrew Robertson, Matteo Bianconi, and Mathilde Jauzac (LSST), “Discovery of Strongly-lensed Gravitational Waves - Implications for the LSST Observing Strategy,” (2019), arXiv:1902.05140 [astro-ph.HE].
- [25] Andrew Robertson, Graham P Smith, Richard Massey, Vincent Eke, Mathilde Jauzac, Matteo Bianconi, and Dan Ryczanowski, “What does strong gravitational lensing? the mass and redshift distribution of high-magnification lenses,” *Monthly Notices of the Royal Astronomical Society* **495**, 3727–3739 (2020).
- [26] Dan Ryczanowski, Graham P Smith, Matteo Bianconi, Richard Massey, Andrew Robertson, and Mathilde Jauzac, “On building a cluster watchlist for identifying strongly lensed supernovae, gravitational waves and kilonovae,” *Monthly Notices of the Royal Astronomical Society* **495**, 1666–1671 (2020).
- [27] Shuji Deguchi and William D. Watson, “Wave effects in gravitational lensing of electromagnetic radiation,” *Phys. Rev. D* **34**, 1708–1718 (1986).
- [28] Takahiro T. Nakamura, “Gravitational lensing of gravitational waves from inspiraling binaries by a point mass lens,” *Phys. Rev. Lett.* **80**, 1138–1141 (1998).
- [29] Ryuichi Takahashi and Takashi Nakamura, “Wave effects in the gravitational lensing of gravitational waves from chirping binaries,” *The Astrophysical Journal* **595**, 1039 (2003).
- [30] Zhoujian Cao, Li-Fang Li, and Yan Wang, “Gravitational lensing effects on parameter estimation in gravitational wave detection with advanced detectors,” *Phys. Rev. D* **90**, 062003 (2014).
- [31] Kwun-Hang Lai, Otto A. Hannuksela, Antonio Herrera-Martín, Jose M. Diego, Tom Broadhurst, and Tjonnie G. F. Li, “Discovering intermediate-mass black hole lenses through gravitational wave lensing,” *Phys. Rev. D* **98**, 083005 (2018).
- [32] Pierre Christian, Salvatore Vitale, and Abraham Loeb, “Detecting stellar lensing of gravitational waves with ground-based observatories,” *Phys. Rev. D* **98**, 103022 (2018).
- [33] Liang Dai, Shun-Sheng Li, Barak Zackay, Shude Mao, and Youjun Lu, “Detecting lensing-induced diffraction in astrophysical gravitational waves,” *Phys. Rev. D* **98**, 104029 (2018).
- [34] Jose M. Diego, “Constraining the abundance of primordial black holes with gravitational lensing of gravitational waves at ligo frequencies,” *Phys. Rev. D* **101**, 123512 (2020).
- [35] Sourabh Magare, Shasvath J Kapadia, Anupreeta More, Mukesh Kumar Singh, Parameswaran Ajith, and AN Ramprakash, “Gear up for the action replay: Leveraging lensing for enhanced gravitational-wave early warning,” *The Astrophysical Journal Letters* **955**, L31 (2023).
- [36] Tessa Baker and Mark Trodden, “Multimessenger time delays from lensed gravitational waves,” *Phys. Rev. D* **95**, 063512 (2017).
- [37] Thomas E. Collett and David Bacon, “Testing the speed of gravitational waves over cosmological distances with strong gravitational lensing,” *Phys. Rev. Lett.* **118**, 091101 (2017).
- [38] Xi-Long Fan, Kai Liao, Marek Biesiada, Aleksandra Piórkowska-Kurpas, and Zong-Hong Zhu, “Speed of gravitational waves from strongly lensed gravitational waves and electromagnetic signals,” *Phys. Rev. Lett.* **118**, 091102 (2017).
- [39] Srashti Goyal, K. Haris, Ajit Kumar Mehta, and Parameswaran Ajith, “Testing the nature of gravitational-wave polarizations using strongly lensed signals,” *Phys. Rev. D* **103**, 024038 (2021).
- [40] Jose María Ezquiaga and Miguel Zumalacárregui, “Gravitational wave lensing beyond general relativity: Birefringence, echoes, and shadows,” *Phys. Rev. D* **102**, 124048 (2020).
- [41] M Sereno, Ph Jetzer, Alberto Sesana, and Marta Volonteri, “Cosmography with strong lensing of lisa gravitational wave sources,” *Monthly Notices of the Royal Astronomical Society* **415**, 2773–2781 (2011).

- [42] Kai Liao, Xi-Long Fan, Xuheng Ding, Marek Biesiada, and Zong-Hong Zhu, “Precision cosmology from future lensed gravitational wave and electromagnetic signals,” *Nature Communications* **8**, 1148 (2017).
- [43] Shuo Cao, Jingzhao Qi, Zhoujian Cao, Marek Biesiada, Jin Li, Yu Pan, and Zong-Hong Zhu, “Direct test of the flrw metric from strongly lensed gravitational wave observations,” *Scientific Reports* **9**, 11608 (2019).
- [44] Yufeng Li, Xilong Fan, and Lijun Gou, “Constraining cosmological parameters in the flrw metric with lensed gw+ em signals,” *The Astrophysical Journal* **873**, 37 (2019).
- [45] Otto A Hannuksela, Thomas E Collett, Mesut Çalıřkan, and Tjonnie GF Li, “Localizing merging black holes with sub-arcsecond precision using gravitational-wave lensing,” *Monthly Notices of the Royal Astronomical Society* **498**, 3395–3402 (2020).
- [46] Souvik Jana, Shasvath J. Kapadia, Tejaswi Venumadhav, and Parameswaran Ajith, “Cosmography using strongly lensed gravitational waves from binary black holes,” *Phys. Rev. Lett.* **130**, 261401 (2023).
- [47] Uddepta Deka, Sumanta Chakraborty, Shasvath J. Kapadia, Md Arif Shaikh, and Parameswaran Ajith, “Probing black hole charge with gravitational microlensing of gravitational waves,” (2024), [arXiv:2401.06553 \[gr-qc\]](https://arxiv.org/abs/2401.06553).
- [48] S. Basak, A. Ganguly, K. Haris, S. Kapadia, A. K. Mehta, and P. Ajith, “Constraints on Compact Dark Matter from Gravitational Wave Microlensing,” *Astrophys. J.* **926**, L28 (2022), [arXiv:2109.06456 \[gr-qc\]](https://arxiv.org/abs/2109.06456).
- [49] A. Barsode, S. J. Kapadia, and P. Ajith, “Constraints on Compact Dark Matter from the Nonobservation of Gravitational-wave Strong Lensing,” *Astrophys. J.* **975**, 48 (2024), [arXiv:2405.15878 \[gr-qc\]](https://arxiv.org/abs/2405.15878).
- [50] OA Hannuksela, K Haris, KKY Ng, S Kumar, AK Mehta, D Keitel, TGF Li, and P Ajith, “Search for gravitational lensing signatures in ligo-virgo binary black hole events,” *The Astrophysical Journal Letters* **874**, L2 (2019).
- [51] Ashish Kumar Meena and J. S. Bagla, “Gravitational lensing of gravitational waves: wave nature and prospects for detection,” *Mon. Not. Roy. Astron. Soc.* **492**, 1127–1134 (2020), [arXiv:1903.11809 \[astro-ph.CO\]](https://arxiv.org/abs/1903.11809).
- [52] José María Diego, Otto A Hannuksela, Patrick L Kelly, G Pagano, Tom Broadhurst, K Kim, Tjonnie Guang Feng Li, and George Fitzgerald Smoot, “Observational signatures of microlensing in gravitational waves at ligo/virgo frequencies,” *Astronomy & Astrophysics* **627**, A130 (2019).
- [53] Giulia Pagano, Otto A Hannuksela, and Tjonnie GF Li, “lensinggw: a python package for lensing of gravitational waves,” *Astronomy & Astrophysics* **643**, A167 (2020).
- [54] Mark HY Cheung, Joseph Gais, Otto A Hannuksela, and Tjonnie GF Li, “Stellar-mass microlensing of gravitational waves,” *Monthly Notices of the Royal Astronomical Society* **503**, 3326–3336 (2021).
- [55] Anuj Mishra, Ashish Kumar Meena, Anupreeta More, Sukanta Bose, and Jasjeet Singh Bagla, “Gravitational lensing of gravitational waves: effect of microlens population in lensing galaxies,” *Mon. Not. Roy. Astron. Soc.* **508**, 4869–4886 (2021), [arXiv:2102.03946 \[astro-ph.CO\]](https://arxiv.org/abs/2102.03946).
- [56] Andrew Ulmer and Jeremy Goodman, “Femtolensing: Beyond the Semiclassical Approximation,” *Astrophys. J.* **442**, 67 (1995), [arXiv:astro-ph/9406042 \[astro-ph\]](https://arxiv.org/abs/astro-ph/9406042).
- [57] Scott E. Field, Chad R. Galley, Jan S. Hesthaven, Jason Kaye, and Manuel Tiglio, “Fast prediction and evaluation of gravitational waveforms using surrogate models,” *Phys. Rev. X* **4**, 031006 (2014), [arXiv:1308.3565 \[gr-qc\]](https://arxiv.org/abs/1308.3565).
- [58] Jonathan Blackman, Scott E. Field, Mark A. Scheel, Chad R. Galley, Daniel A. Hemberger, Patricia Schmidt, and Rory Smith, “A surrogate model of gravitational waveforms from numerical relativity simulations of precessing binary black hole mergers,” *Physical Review D* **95** (2017), [10.1103/physrevd.95.104023](https://doi.org/10.1103/physrevd.95.104023).
- [59] Jonathan Blackman, Scott E. Field, Mark A. Scheel, Chad R. Galley, Christian D. Ott, Michael Boyle, Lawrence E. Kidder, Harald P. Pfeiffer, and Béla Szilágyi, “Numerical relativity waveform surrogate model for generically precessing binary black hole mergers,” *Phys. Rev. D* **96**, 024058 (2017), [arXiv:1705.07089 \[gr-qc\]](https://arxiv.org/abs/1705.07089).
- [60] Vijay Varma, Scott E. Field, Mark A. Scheel, Jonathan Blackman, Davide Gerosa, Leo C. Stein, Lawrence E. Kidder, and Harald P. Pfeiffer, “Surrogate models for precessing binary black hole simulations with unequal masses,” *Phys. Rev. Research* **1**, 033015 (2019), [arXiv:1905.09300 \[gr-qc\]](https://arxiv.org/abs/1905.09300).
- [61] Tousif Islam, Vijay Varma, Jackie Lodman, Scott E. Field, Gaurav Khanna, Mark A. Scheel, Harald P. Pfeiffer, Davide Gerosa, and Lawrence E. Kidder, “Eccentric binary black hole surrogate models for the gravitational waveform and remnant properties: comparable mass, nonspinning case,” *Phys. Rev. D* **103**, 064022 (2021), [arXiv:2101.11798 \[gr-qc\]](https://arxiv.org/abs/2101.11798).
- [62] Jooheon Yoo, Vijay Varma, Matthew Giesler, Mark A. Scheel, Carl-Johan Haster, Harald P. Pfeiffer, Lawrence E. Kidder, and Michael Boyle, “Targeted large mass ratio numerical relativity surrogate waveform model for GW190814,” *Phys. Rev. D* **106**, 044001 (2022), [arXiv:2203.10109 \[gr-qc\]](https://arxiv.org/abs/2203.10109).
- [63] Jooheon Yoo *et al.*, “Numerical relativity surrogate model with memory effects and post-Newtonian hybridization,” *Phys. Rev. D* **108**, 064027 (2023), [arXiv:2306.03148 \[gr-qc\]](https://arxiv.org/abs/2306.03148).
- [64] Davide Gerosa, François Hébert, and Leo C. Stein, “Black-hole kicks from numerical-relativity surrogate models,” *Phys. Rev. D* **97**, 104049 (2018), [arXiv:1802.04276 \[gr-qc\]](https://arxiv.org/abs/1802.04276).
- [65] Vijay Varma, Davide Gerosa, Leo C. Stein, François Hébert, and Hao Zhang, “High-accuracy mass, spin, and recoil predictions of generic black-hole merger remnants,” *Phys. Rev. Lett.* **122**, 011101 (2019), [arXiv:1809.09125 \[gr-qc\]](https://arxiv.org/abs/1809.09125).
- [66] Tousif Islam, Scott E. Field, and Gaurav Khanna, “Remnant black hole properties from numerical-relativity-informed perturbation theory and implications for waveform modeling,” *Phys. Rev. D* **108**, 064048 (2023), [arXiv:2301.07215 \[gr-qc\]](https://arxiv.org/abs/2301.07215).
- [67] Katie Rink, Ritesh Bachhar, Tousif Islam, Nur E. M. Rifat, Kevin Gonzalez-Quesada, Scott E. Field, Gaurav Khanna, Scott A. Hughes, and Vijay Varma, “Gravitational wave surrogate model for spinning, intermediate mass ratio binaries based on perturbation theory and numerical relativity,” (2024), [arXiv:2407.18319 \[gr-qc\]](https://arxiv.org/abs/2407.18319).
- [68] Tousif Islam, Scott E. Field, Scott A. Hughes, Gaurav Khanna, Vijay Varma, Matthew Giesler, Mark A. Scheel, Lawrence E. Kidder, and Harald P. Pfeiffer, “Surrogate model for gravitational wave signals from nonspinning, comparable-to large-mass-ratio black hole binaries built on black hole perturbation theory waveforms calibrated to numerical relativity,” *Physical Review D* **106** (2022), [10.1103/physrevd.106.104025](https://doi.org/10.1103/physrevd.106.104025).
- [69] Joan Fontbuté, Tomas Andrade, Raimon Luna, Juan Calderón Bustillo, Gonzalo Morrás, Santiago Jaraba, Juan García-Bellido, and Germán López Izquierdo, “A numerical-relativity surrogate model for hyperbolic encounters of black holes: challenges in parameter estimation,” (2024), [arXiv:2409.16742 \[gr-qc\]](https://arxiv.org/abs/2409.16742).
- [70] P. Schneider, J. Ehlers, and E.E. Falco, *Gravitational Lenses*, Astronomy and Astrophysics Library (Springer Berlin Heidelberg, 2013).
- [71] P. C. Peters, “Index of refraction for scalar, electromagnetic, and gravitational waves in weak gravitational fields,” *Phys. Rev. D* **9**,

- 2207–2218 (1974).
- [72] Norihito Matsunaga and Kazuhiro Yamamoto, “The finite source size effect and the wave optics in gravitational lensing,” *JCAP* **01**, 023 (2006), [arXiv:astro-ph/0601701](#).
- [73] Manuel Tiglio and Aarón Villanueva, “Reduced order and surrogate models for gravitational waves,” *Living Rev. Rel.* **25**, 2 (2022), [arXiv:2101.11608 \[gr-qc\]](#).
- [74] Scott E. Field, Chad R. Galley, Frank Herrmann, Jan S. Hesthaven, Evan Ochsner, and Manuel Tiglio, “Reduced basis catalogs for gravitational wave templates,” *Phys. Rev. Lett.* **106**, 221102 (2011), [arXiv:1101.3765 \[gr-qc\]](#).
- [75] Maxime Barrault, Yvon Maday, Ngoc Cuong Nguyen, and Anthony T Patera, “An ‘empirical interpolation’ method: application to efficient reduced-basis discretization of partial differential equations,” *Comptes Rendus Mathématique* **339**, 667–672 (2004).
- [76] Yvon Maday, Ngoc Cuong Nguyen, Anthony T. Patera, and S. H. Pau, “A general multipurpose interpolation procedure: the magic points,” (2009).
- [77] Jan S. Hesthaven, Benjamin Stamm, and Shun Zhang, “Efficient greedy algorithms for high-dimensional parameter spaces with applications to empirical interpolation and reduced basis methods,” *ESAIM: Mathematical Modelling and Numerical Analysis* **48**, 259–283 (2014).
- [78] Benjamin P Abbott *et al.* (LIGO Scientific, Virgo), “A guide to LIGO–Virgo detector noise and extraction of transient gravitational-wave signals,” *Class. Quant. Grav.* **37**, 055002 (2020), [arXiv:1908.11170 \[gr-qc\]](#).
- [79] Alex Nitz, Ian Harry, Duncan Brown, Christopher M. Biwer, Josh Willis, Tito Dal Canton, Collin Capano, Thomas Dent, Lorne Pekowsky, Andrew R. Williamson, Soumi De, Miriam Cabero, Bernd Machenschalk, Duncan Macleod, Prayush Kumar, Steven Reyes, Francesco Pannarale, Gareth S Cabourn Davies, dfinstad, Sumit Kumar, Márton Tápai, Leo Singer, Sebastian Khan, Stephen Fairhurst, Alex Nielsen, Shashwat Singh, Thomas Massinger, Koustav Chandra, Shasvath, and Veronicavilla, “[gwastro/pycbc: v2.0.4 release of pycbc](#),” (2022).
- [80] S. Hild *et al.*, “Sensitivity Studies for Third-Generation Gravitational Wave Observatories,” *Class. Quant. Grav.* **28**, 094013 (2011), [arXiv:1012.0908 \[gr-qc\]](#).
- [81] Charles R. Harris *et al.*, “Array programming with NumPy,” *Nature* **585**, 357–362 (2020), [arXiv:2006.10256 \[cs.MS\]](#).
- [82] Geraint Pratten *et al.*, “Computationally efficient models for the dominant and subdominant harmonic modes of precessing binary black holes,” *Phys. Rev. D* **103**, 104056 (2021), [arXiv:2004.06503 \[gr-qc\]](#).
- [83] Giovanni Tambalo, Miguel Zumalacárregui, Liang Dai, and Mark Ho-Yeuk Cheung, “Lensing of gravitational waves: Efficient wave-optics methods and validation with symmetric lenses,” *Phys. Rev. D* **108**, 043527 (2023).
- [84] Mark Ho-Yeuk Cheung, Ken K. Y. Ng, Miguel Zumalacárregui, and Emanuele Berti, “Probing minihalo lenses with diffracted gravitational waves,” *Phys. Rev. D* **109**, 124020 (2024), [arXiv:2403.13876 \[gr-qc\]](#).
- [85] Hector Villarrubia-Rojo, Stefano Savastano, Miguel Zumalacárregui, Lyla Choi, Srashti Goyal, Liang Dai, and Giovanni Tambalo, “GLoW: novel methods for wave-optics phenomena in gravitational lensing,” (2024), [arXiv:2409.04606 \[gr-qc\]](#).