Classifier Weighted Mixture models

Elouan Argouarc'h, François Desbouvries Senior Member, IEEE, Éric Barat, Eiji Kawasaki, Thomas Dautremer

Abstract—This paper proposes an extension of standard mixture stochastic models, by replacing the constant mixture weights with functional weights defined using a classifier. *Classifier Weighted Mixtures* enable straightforward density evaluation, explicit sampling, and enhanced expressivity in variational estimation problems, without increasing the number of components nor the complexity of the mixture components.

Index Terms—mixture models, classifier, variational estimation, reparameterization gradients.

I. INTRODUCTION

Mixture distributions $\sum_{k=1}^{K} \pi_k p_k(x)$ with $\pi_k \in (0, 1)$ and $\sum_{k=1}^{K} \pi_k = 1$ are powerful tools in probabilistic modeling. For instance, mixture of Gaussians distributions are widely used for density estimation and implicit clustering [1], because their structure makes them well suited for representing distributions with several modes, and because of the availability of a potent closed-form Expectation-Maximization (EM) algorithm [2][3]. Moreover, mixtures are a particular type of latent variable model (LVM) where the latent variable is discrete and finite. As a result, not only can mixture distributions be sampled from easily, but they also benefit from a tractable probability density function (PDF), making them versatile tools for variational estimation problems.

Modern machine learning problems involve increasingly complex target distributions, which, from a variational estimation perspective, highlights the challenge of ensuring sufficient expressivity in the class of approximating surrogate distributions. This leads us to the limitations of classical mixtures. On the one hand, a mixture of K simple distributions (e.g. Gaussians) enables to get arbitrarily close in theory [4] to a given distribution, provided K is large enough; but in practice, a high number of components leads to computational burdens and numerical instability. On the other hand, one can increase the expressiveness of a mixture by considering more sophisticated component distributions (denoted by \mathcal{P}_k). For instance, [5] investigated mixtures of Normalizing-Flows (NF), where each mixture component is the result of a complex invertible mapping of some base distribution (denoted \mathcal{P}_0). Even though this approach has been successfully applied in practice on density estimation tasks [6], building a mixture of K differents NFs is theoretically equivalent to applying a unique mapping to a base distribution which itself is a K-mixture, so the advantage of this approach remains to be investigated.

Our aim in this paper is to extend the expressivity of classical mixtures, while maintaining the critical computational features of a tractable PDF and a straightforward sampling scheme. Rather than increasing K or considering complex components \mathcal{P}_k , our approach consists in replacing the constant mixture weights π_k by functions $\pi_k(x)$. In practice, we will achieve this goal by using a classifying function, yielding a new approach for building parameterized distributions which we refer to as *Classifier Weighted Mixtures* (CWM). The paper is organized as follows. First, in section II, we detail the construction of the CWM PDF and unravel the underlying latent variable structure. Next, in section III, we discuss the parameterization and applications to variational estimation problems, and compare our approach to alternative models through simulations ¹.

II. FROM STANDARD MIXTURES TO CWM

Our goal is to extend a mixture by replacing the constant weights $\pi_k \in (0, 1)$ by functions $\pi_k : x \to \pi_k(x)$:

$$\sum_{k=1}^{K} \pi_k p_k(x) \longrightarrow \sum_{k=1}^{K} \pi_k(x) p_k(x).$$
(1)



Fig. 1: Classical mixtures (left) can be extended to mixtures with functional and non-necessarily constant weights (right).

Of course, these functions cannot be arbitrary. In section II-A, we thus propose sufficient conditions on $\pi_k(x)$ such that $\sum_{k=1}^{K} \pi_k(x) p_k(x)$ is a valid PDF, which leads us to the CWM solution. In section II-B, we unravel its underlying latent variable structure, which shows that CWM can easily be sampled from. Finally in section II-C, we decompose our stochastic model into its three fundamental components.

A. A classifier based construction

Let us focus on the right hand side of (1). For it to be a valid PDF, this function needs to be non-negative and sum to 1. We therefore aim to constrain functions $\pi_k(x)$ such that:

$$\sum_{k=1}^{K} \pi_k(x) p_k(x) \ge 0;$$
 (2)

$$\int \sum_{k=1}^{K} \pi_k(x) p_k(x) dx = 1.$$
 (3)

¹Code available at https://github.com/ElouanARGOUARCH/Classifier-Weighted-Mixtures

Elouan Argouarc'h and François Desbouvries are with SAMOVAR, Télécom SudParis, Institut Polytechnique de Paris, 91120 Palaiseau, France (e-mail: elouan.argouarch,francois.desbouvries@telecom-sudparis.eu). Elouan Argouarc'h, Éric Barat, Eiji Kawasaki and Thomas Dautremer are with Université Paris Saclay, CEA, List F-91120 Palaiseau, France (e-mail: elouan.argouarch, eric.barat, eiji.kawasaki, thomas.dautremer@cea.fr)

Firstly, (2) is satisfied if all addends $\pi_k(x)p_k(x) \ge 0$, and thus if $\pi_k(x) \ge 0$. To fulfill the second condition, we use P_k the cumulative distribution function (CDF) of \mathcal{P}_k . With the change of variable $u = P_k(x)$, we rewrite the left-hand side of (3) as:

$$\int \sum_{k=1}^{K} \pi_k(x) p_k(x) \mathrm{d}x = \int_{[0,1]} \sum_{k=1}^{K} \pi_k(P_k^{-1}(u)) \mathrm{d}u.$$
(4)

If $\sum_{k=1}^{K} \pi_k(P_k^{-1}(u)) = 1$, the previous integral reduces to 1 and (3) is satisfied. We have hence identified sufficient conditions on $\pi_k(x)$ such that $\sum_{k=1}^{K} \pi_k(x)p_k(x)$ is indeed a PDF. Finally, to see how a classifier is involved, we set $\alpha_k(u) \stackrel{\Delta}{=} \pi_k(P_k^{-1}(u))$, which enables us to rewrite the two previous conditions about $\pi_k(x)$ as:

$$\sum_{k=1}^{K} \alpha_k(u) = 1 \text{ and } \alpha_k(u) \ge 0 \text{ for all } u \in [0,1].$$
 (5)

It is now clear that functions $\alpha_k(u)$ define a vector of probabilities, so these functions are nothing but a classifier. Therefore, with such a K-label classifying function, the construction:

$$\sum_{k=1}^{K} \underbrace{\alpha_k(P_k(x))}_{\pi_k(x)} p_k(x) \stackrel{\Delta}{=} p_{\text{CWM}}(x); \tag{6}$$

is indeed a valid PDF. Note that $\sum_{k=1}^{K} \alpha_k(u) = 1$, but $\sum_{k=1}^{K} \pi_k(x) \neq 1$: weights $\pi_k(x)$ are defined via a classifier (whence the name CWM), but, unless all functions are constant (in which case $p_{\text{CWM}}(x)$ reduces to a classical mixture), are not themselves a classifier. In figure 2, we display the classifying functions $\alpha_k(u)$ and the corresponding weights $\pi_k(x)$ which produced the CWM distribution in figure 1.



Fig. 2: Classifier $\alpha_k(u)$ and corresponding weights $\pi_k(x)$.

B. The underlying latent variable construction

In the previous section we built a mixture weighted by functions $\pi_k(x)$ built from classifying functions $\alpha_k(z)$. We now explain that, even though our construction is more intricate than a classical mixture model, it can nonetheless be described as a latent variable model. Therefore, in addition to a straightforward PDF evaluation mechanism (see equation (6)), the resulting distribution also benefits from an explicit sampling scheme.

PDF (6) is expressed as a sum over values k = 1, ..., K, and is thus the marginal over a couple (X, R), where R is categorical, and their joint PDF are the added terms in (6):

$$p_{\text{CWM}}(x, R = k) = \alpha_k(P_k(x))p_k(x).$$
(7)

As it stands, this joint PDF is not factorized as a marginal over (R) multiplied by a conditional (X|R) (unlike in usual

mixture models). However, augmenting this joint PDF with the additional variable $u \in [0, 1]$ and applying the change of variable $X = P_k^{-1}(U)$ yields a PDF over (X, R, U) which indeed is factorized as (U) times (R|U) times (X|R, U):

$$p_{\text{CWM}}(x, R = k, u) = \underbrace{\mathbb{1}_{[0,1]}(u)}_{(U)} \underbrace{\alpha_k(u)}_{(R|U)} \underbrace{\delta_{P_k^{-1}(u)}(x)}_{(X|R,U)}.$$
 (8)

This joint PDF has an appropriate factorization and we can hence deduce the corresponding directed graph (see figure 3).



Fig. 3: Directed graph associated with the CWM construction

From this directed graph, it is easy to deduce a sampling scheme. Indeed, LVMs benefit from built-in procedures: by following the directed graph we can sequentially sample the latent variables, and then sample the observed variable. In our case, we can obtain samples from this distribution with:

$$U \sim \text{Uniform}([0,1]),$$
 (9)

$$R \sim \text{Categorical}(\alpha(U)),$$
 (10)

$$X = P_B^{-1}(U),$$
 (11)

where $\alpha(U) = [\alpha_1(U), ..., \alpha_K(U)]^T$.

C. Change of base distribution and practical construction

In section II-A, we proposed a construction based on a uniform base distribution, and on the CDFs P_k of the mixture components \mathcal{P}_k . But indeed we can replace the uniform distribution by another one without additional conditions about $\pi_k(x)$. To see this, let us replace U by $Z \sim \mathcal{P}_0$. This corresponds to augmenting graph 3 into:



If we denote $w_k(z) \triangleq \alpha_k(P_0(z))$, then $w_k(z)$ and $\alpha_k(u)$ are both equivalently classifying functions. On the other hand, the inverse CDFs P_k^{-1} (resp. $P_0 \circ P_k^{-1}$) are simply used as a change of variable in order to transform the base distribution Uniform([0, 1]) (resp. \mathcal{P}_0) into \mathcal{P}_k . So in practice (see section III-A3 below) in order to adjust these mixture components, we shall consider invertible parameterized mappings T_k . With J_{T_k} denote the Jacobian matrix of T_k , the CWM PDF finally reads:

$$p_{\text{CWM}}(x) = \sum_{k=1}^{K} w_k(T_k(x)) p_0(T_k(x)) |\det(\mathbf{J}_{T_k}(x))|.$$
(12)

III. APPLICATION TO VARIATIONAL ESTIMATION PROBLEMS

We now apply our CWM model to variational problems. In Section III-A we build our model step-by-step, considering issues such as sampling, PDF computation, and reparameterization. In the mean time, we detail the construction that we used in our experiments, in which, because we want to highlight the potential expressivity induced by functional weights $\pi_k(x)$ as compared to constant ones π_k , we purposefully use simple Gaussian mixture components (see III-A4). Next in section III-B we apply our model in the context of variational problems. Section III-C displays simulation results.

A. Parameterization

1) Base distribution: Remember from section II-C that we can theoretically start from any parameterized base distribution. However, if we need to sample from the corresponding CWM model (12) (resp. evaluate its PDF), the base distribution must be sampled from easily (resp. benefit from a closedform PDF). Moreover, application to variational estimation problems might call for a reparameterization of the gradient procedure (see section III-B2). In turn this means that the base distribution should benefit from a sampling procedure which can be reparameterized, enabling to propagate gradients with respect to the parameters of the base distribution through its samples.

Now, it is common practice in LVMs [7][8][9] to consider parameter-free base distributions; moreover, in our experiments, we will focus on comparing our model to a Gaussian mixture model (GMM) and to NFs (see section III-C). For all these reasons, we choose a standard, parameter-free, normal distribution:

$$\mathcal{P}_0 = \mathcal{N}(0, \mathbf{I}). \tag{13}$$

2) Classifying function: Recall that it is sufficient that functions w_k compute positive values which add to 1; therefore, any classifying function works [10][11]. In this paper we experiment with a fully-connected feed-forward Neural Network (NN), say f, which outputs K logit values:

$$w(z) = \text{Softmax}(f(z)). \tag{14}$$

The weights and biases parameters will be adjusted during the variational estimation procedure, see section III-B.

3) Invertible mapping: We use invertible mappings T_k to transform the base distribution \mathcal{P}_0 into the mixture components with PDF given by $p_0(T_k(x))|\det J_{T_k}(x)|$, and which can be sampled from with $T_k^{-1}(z), z \sim \mathcal{P}_0$. We can parameterize these invertible mappings using NF architectures (see [12][13][14][15][16] or [17][18] for reviews of NF methods). In our experiments, we aim to highlight the advantage of using functional weights $\pi_k(x)$ rather than constant weights π_k , so we build a mixture of simple Gaussians with:

$$T_k(x) = \Sigma_k^{-1/2} (x - \mu_k);$$
(15)

where means μ_k and covariance matrices Σ_k are parameters to be adjusted during the optimization. Moreover in practice, to ensure that Σ_k remain positive-definite during gradient updates, we can consider diagonal matrices and parameterize the coefficients using unconstrained log-variance parameters.

4) A resulting model: Specific choices of base distribution, classifying function and invertible mappings, as discussed above, lead to different architectures. For instance, in our simulations (see section III-C), we will use the construction based on (13), (14) and (15); in that case equation (12) reduces to:

$$p_{\text{CWM}}(x) = \sum_{k=1}^{K} w_k (\Sigma_k^{-1/2} (x - \mu_k)) \mathcal{N}(x; \mu_k, \Sigma_k).$$
(16)

It is indeed a mixture of Gaussian distributions, but weighted by a NN classifier function.

$$X \qquad Z \sim \mathcal{N}(0, \mathbf{I})$$

$$R \sim \text{Categorical}(w(Z)) \quad (17)$$

$$X = \Sigma_R^{1/2} Z + \mu_R$$

B. Application to Variational estimation problems

We now see how CWM can be used for solving variational estimation problems. We focus on two types of problems: density estimation in section III-B1, and reparameterization gradients in section III-B2. Let θ denote the set of all parameters (of the base distribution \mathcal{P}_0 , of the weight function $w_k(.)$ and of the invertible mappings T_k), and let $p_{\theta}(x)$ the resulting CWM PDF (see equation (12)).

1) Density Estimation: In density estimation, we dispose of recorded samples $\mathcal{D} = \{x_1, ..., x_{|\mathcal{D}|}\}$ from a distribution of interest, say \mathcal{P} , and the goal is to build a surrogate distribution \mathcal{P}_{θ^*} that closely resembles \mathcal{P} by finding appropriate parameters θ^* . In practice, this is usually achieved with maximum likelihood estimation (MLE) (which is equivalent to minimizing a Monte Carlo (MC) estimate of the Kullback-Leibler divergence between \mathcal{P} and \mathcal{P}_{θ}) approximated using gradient ascent. In this case, the log-likelihood of $\mathcal{D}, \sum_{i=1}^{|\mathcal{D}|} \log(p_{\theta}(x_i))$, is differentiable with respect to θ and we can thus solve the MLE problem by gradient ascent (at least approximately towards a local maxima).

Now, when considering the specific parameterization discussed in the previous section (see (16) (17)), forcing the classifier $w_k(z)$ to constant values $\pi_k \in (0, 1)$ (which can be achieved by setting all the weights of the last layer of the classifier f to 0 and its biases to $\log(\pi_k)$) results in a standard mixture of Gaussians. Therefore, we can easily use the EM algorithm for Gaussian mixtures as a pre-training step, and next use the (constant weights) mixture produced by that EM algorithm as the initial point of the gradient-based optimization of the CWM model, in which the classifier $w_k(z)$ will no longer predict constant values.

2) Reparameterization Gradients: Many variational tasks [19][9][20] can be expressed as minimizing over θ , by gradient descent, expectation $\mathbb{E}_{X \sim \mathcal{P}_{\theta}}(h(X))$, where h is a measurable function. It is therefore of interest to compute, or at least

estimate with MC, the gradient of $\mathbb{E}_{X \sim \mathcal{P}_{\theta}}(h(X))$ with respect to θ . However, computing the gradient of the crude estimate:

$$\mathbb{E}_{X \sim \mathcal{P}_{\theta}}(h(X)) \approx \frac{1}{M} \sum_{i=1}^{M} h(x_i), x_i \stackrel{iid}{\sim} \mathcal{P}_{\theta}, \qquad (18)$$

is difficult since the samples depend on parameters θ .

Practitioners often rely on reparameterization tricks [9][21] that consist of rewriting samples using a bivariate function, differentiable with respect to θ , and that detangles the randomness of the parameters. We now discuss the *reparameterization gradients* of a CWM model.

First, as we have mentioned in section II-C, it is necessary to parameterize the base distribution appropriately in this context, and we suppose that one can reparameterize its samples in order to propagate the gradients with respect to parameters of the base distribution. But, since sampling from a CWM involves the discrete variable R, samples $X = T_R^{-1}(Z)$ cannot be reparameterized using a differentiable function because $R = \sum_{k=1}^{K} \mathbb{1}(U \leq \sum_{j=1}^{k} w_j(Z)), U \sim \mathcal{U}_{[0,1]}$ is not continuous and we cannot propagate the gradients through this variable. Besides the reinforce approach [22] which is suited for both discrete and continuous variables (but leads to high variance estimates [23]), notorious workarounds include the Gumbell-Softmax [24][25] for categorical variables. It consists in approximating $R = \arg \max_k(g_k + \log(w_k(Z))), g_k \overset{iid}{\sim}$ Gumbell(0, 1) by replacing the argmax operator with a softmax, which yields a differentiable, but biased estimate.

In our case, since the categorical variable is a latent variable, we will show that a differentiable (approximate) reparameterization of R is not required. Instead, we can build an estimate where this variable is marginalized out, following the Rao-Blackwellization (RB) [26] principle. RB is based on $\mathbb{E}(h(X)) = \mathbb{E}(\mathbb{E}(h(X)|Z))$ [27]. Since we can compute $\mathbb{E}(h(X)|Z) = \sum_{k=1}^{K} w_k(Z)h(T_k^{-1}(Z))$, only the outer expectation calls for an MC approximation. This leads to the estimate:

$$\mathbb{E}_{X \sim \mathcal{P}_{\theta}}(h(X)) \approx \frac{1}{M} \sum_{\substack{i=1\\z_i \sim \mathcal{P}_0}}^{M} \underbrace{\sum_{k=1}^{K} w_k(z_i) h(T_k^{-1}(z_i))}_{\mathbb{E}(h(X)|Z=z_i)}.$$
 (19)

The interest of (19) is twofold. First, $\mathbb{V}ar(\mathbb{E}(h(X)|Z)) = \mathbb{V}ar(h(X)) - \mathbb{E}(\mathbb{V}ar(h(X)|Z))$ [28], so (19) has lower variance than the crude estimate. Next, computing $\mathbb{E}(h(X)|Z)$ marginalizes out R, so (19) does not involve sampling from this variable; the estimate is thus differentiable and no longer relies on a reparameterization of the categorical samples.

C. Experiments

In this illustrative section, we compare our CWM model to NF architectures (RNVP [14], NSF [16]) and to GMMs on a density estimation task. We draw samples (b) (an histogram is displayed in the figure) from the 2-dimensional distribution associated with a grey-scale image (a); for NFs and CWM, the density is estimated via gradient ascent of the log-likelihood (after pre-training, as mentioned in section III-B1); for GMM we used the EM algorithm. We summarize our results by displaying the PDF of each model (compared to the original image) in figure 4, and the likelihood scores in table I.



Fig. 4: Image - samples - CWM - RNVP - NSF - GMM

	CWM $(K = 50)$	RNVP	NSF	$GMM \ (K = 50)$
# Parameters	39925	85780	79560	343
Likelihood	$0.221\pm1.6\mathrm{e}-3$	$0.133 \pm 1.1e - 2$	$0.182 \pm 3.9e - 3$	$0.150 \pm 2.2e - 3$

Table I: Architectures, parameters and log-likelihood

Our approach outperforms the other architectures in terms of likelihood score, and with fewer number of parameters (at least compared to NF models). This is confirmed visually: the resulting PDF closely resembles the original image, which means that our stochastic model can easily represent distributions with disjoint elements of mass, fine details and sharp edges.

IV. CONCLUSION

In this paper we propose to extend classical mixtures of distributions, where the weights are constant, by mixtures where the weights become functional. More precisely, the CWM solution we propose is a construction where the functional weights are built using a classifying function. This approach allows the use of flexible (possibly NN-based) classifiers, increasing the expressiveness of our stochastic model in variational estimation problems, as compared to standard mixtures models, while still ensuring a tractable PDF and an explicit sampling scheme. As such, CWM models are suited for density estimation tasks but also allow for reparameterization gradients using RB of its discrete latent variable. Finally, our experiments showcase the interest of our approach compared to other architectures.

REFERENCES

- G. Celeux and G. Govaert, "A classification EM algorithm for clustering and two stochastic versions," *Computational statistics & Data analysis*, vol. 14, no. 3, pp. 315–332, 1992.
- [2] A. P. Dempster, N. M. Laird, and D. B. Rubin, "Maximum likelihood from incomplete data via the EM algorithm," *Journal of the royal statistical society: series B* (*methodological*), vol. 39, no. 1, pp. 1–22, 1977.
- [3] C. J. Wu, "On the convergence properties of the EM algorithm," *The Annals of statistics*, pp. 95–103, 1983.
- [4] K. N. Plataniotis and D. Hatzinakos, "Gaussian mixtures and their applications to signal processing," in *Advanced Signal Processing Handbook* (S. Stergiopoulos, ed.), pp. 89–124, Boca Raton, London, New York, Washington D.C.: CRC Press, 1st edition, 2000.
- [5] J. Postels, M. Liu, R. Spezialetti, L. Van Gool, and F. Tombari, "Go with the flows: Mixtures of normalizing flows for point cloud generation and reconstruction," in 2021 International Conference on 3D Vision (3DV), (London, United Kingdom), pp. 1249–1258, IEEE, 2021.
- [6] Y. Yang and T. M. Hospedales, "Mixture of normalizing flows for European option pricing," in *Proceedings of the Thirty-Ninth Conference on Uncertainty in Artificial Intelligence* (R. J. Evans and I. Shpitser, eds.), vol. 216 of *Proceedings of Machine Learning Research*, pp. 2390– 2399, PMLR, 31 Jul–04 Aug 2023.
- [7] D. M. Blei, A. Y. Ng, and M. I. Jordan, "Latent dirichlet allocation," J. Mach. Learn. Res., vol. 3, p. 993–1022, Mar. 2003.
- [8] I. Goodfellow, J. Pouget-Abadie, M. Mirza, B. Xu, D. Warde-Farley, S. Ozair, A. Courville, and Y. Bengio, "Generative adversarial nets," *Advances in neural information processing systems*, vol. 27, 2014.
- [9] D. P. Kingma and M. Welling, "Auto-Encoding Variational Bayes," in 2nd International Conference on Learning Representations, ICLR 2014, Banff, AB, Canada, April 14-16, 2014, Conference Track Proceedings, 2014.
- [10] J. A. Nelder and R. W. Wedderburn, "Generalized linear models," *Journal of the Royal Statistical Society Series A: Statistics in Society*, vol. 135, no. 3, pp. 370–384, 1972.
- [11] Y. LeCun, L. Bottou, Y. Bengio, and P. Haffner, "Gradient-based learning applied to document recognition," *Proceedings of the IEEE*, vol. 86, no. 11, pp. 2278– 2324, 1998.
- [12] D. Rezende and S. Mohamed, "Variational inference with normalizing flows," in *International conference on machine learning*, pp. 1530–1538, PMLR, 2015.
- [13] D. P. Kingma, T. Salimans, R. Jozefowicz, X. Chen, I. Sutskever, and M. Welling, "Improved variational inference with inverse autoregressive flow," *Advances in neural information processing systems*, vol. 29, 2016.
- [14] L. Dinh, J. Sohl-Dickstein, and S. Bengio, "Density estimation using real NVP," in *International Conference* on Learning Representations, 2017.
- [15] G. Papamakarios, T. Pavlakou, and I. Murray, "Masked

autoregressive flow for density estimation," Advances in neural information processing systems, vol. 30, 2017.

- [16] C. Durkan, A. Bekasov, I. Murray, and G. Papamakarios, "Neural spline flows," *Advances in neural information processing systems*, vol. 32, 2019.
- [17] I. Kobyzev, S. J. Prince, and M. A. Brubaker, "Normalizing flows: An introduction and review of current methods," *IEEE transactions on pattern analysis and machine intelligence*, vol. 43, no. 11, pp. 3964–3979, 2020.
- [18] G. Papamakarios, E. Nalisnick, D. J. Rezende, S. Mohamed, and B. Lakshminarayanan, "Normalizing flows for probabilistic modeling and inference," *Journal of Machine Learning Research*, vol. 22, no. 57, pp. 1–64, 2021.
- [19] R. S. Sutton, D. McAllester, S. Singh, and Y. Mansour, "Policy gradient methods for reinforcement learning with function approximation," *Advances in neural information processing systems*, vol. 12, 1999.
- [20] Y. Ganin, E. Ustinova, H. Ajakan, P. Germain, H. Larochelle, F. Laviolette, M. March, and V. Lempitsky, "Domain-adversarial training of neural networks," *Journal of machine learning research*, vol. 17, no. 59, pp. 1–35, 2016.
- [21] M. Figurnov, S. Mohamed, and A. Mnih, "Implicit reparameterization gradients," *Advances in neural information processing systems*, vol. 31, 2018.
- [22] R. J. Williams, "Simple statistical gradient-following algorithms for connectionist reinforcement learning," *Machine learning*, vol. 8, pp. 229–256, 1992.
- [23] M. Xu, M. Quiroz, R. Kohn, and S. A. Sisson, "Variance reduction properties of the reparameterization trick," in *The 22nd international conference on artificial intelligence and statistics*, pp. 2711–2720, PMLR, 2019.
- [24] E. Jang, S. Gu, and B. Poole, "Categorical reparameterization with Gumbel-Softmax," in *International Conference on Learning Representations*, 2017.
- [25] M. Balog, N. Tripuraneni, Z. Ghahramani, and A. Weller, "Lost relatives of the Gumbel trick," in *Proceedings of the 34th International Conference on Machine Learning* (D. Precup and Y. W. Teh, eds.), vol. 70 of *Proceedings of Machine Learning Research*, pp. 371–379, PMLR, 06– 11 Aug 2017.
- [26] G. Casella and C. P. Robert, "Rao-blackwellisation of sampling schemes," *Biometrika*, vol. 83, no. 1, pp. 81– 94, 1996.
- [27] D. Blackwell, "Conditional expectation and unbiased sequential estimation," *The Annals of Mathematical Statistics*, pp. 105–110, 1947.
- [28] C. R. Rao, "Information and the accuracy attainable in the estimation of statistical parameters," in *Breakthroughs* in *Statistics: Foundations and basic theory*, pp. 235–247, Springer, 1992.