## Why a Bose-Einstein condensate cannot exist in a system of interacting bosons at ultrahigh temperatures

Maksim D. Tomchenko\*

Bogolyubov Institute for Theoretical Physics, 14b, Metrolohichna Str., Kyiv 03143, Ukraine

(Dated: January 7, 2025)

It is well known that a Bose-Einstein (BE) condensate of atoms exists in a system of interacting Bose atoms at  $T \leq T_c^{(i)}$ , where  $T_c^{(i)}$  is the BE condensation temperature of an ideal gas. It is also generally accepted that BE condensation is impossible at "ultrahigh" temperatures  $T \gg T_c^{(i)}$ . While the latter property has been theoretically proven for an ideal gas, no such proof exists for an interacting system, to our knowledge. In this paper, we propose an approximate mathematical proof for a finite, nonrelativistic, periodic system of N spinless interacting bosons. The key point is that, at  $T \gg T_c^{(i)}$ , the main contribution to the occupation number  $N_0 = \frac{1}{Z} \sum_{\wp} e^{-E_{\wp}/k_B T} \langle \Psi_{\wp} | \hat{a}_0^+ \hat{a}_0 | \Psi_{\wp} \rangle$ , corresponding to atoms with zero momentum, originates from the states containing N elementary quasiparticles. These states do not contain the BE condensate of zero-momentum atoms, implying that an ultrahigh temperature should "blur" such a condensate.

Keywords: Bose-Einstein condensate, ultrahigh temperatures.

## I. INTRODUCTION

It is known that in a system of interacting bosons, Bose-Einstein (BE) condensation occurs at temperatures below or of the order of the BE condensation temperature  $T_c^{(i)}$  of an ideal gas [1, 2]. The properties of a BE condensate of atoms have been well studied, first for liquid  ${}^{4}He$  and later for trapped Bose gases. Moreover, a condensate of interacting mesons was theoretically predicted long ago (see reviews [3, 4]), but, as far as we know, there is no reliable experimental confirmation of its existence. In addition, BE condensation has been predicted and experimentally detected for photons [10] and quasiparticles such as magnons [5, 6], excitons [7, 8], and polaritons [9] (see also review [11] and books [1, 2]). Furthermore, while BE condensation in liquid  ${}^{4}He$  and trapped gases occurs at low (~ 1 K) and ultralow (  $\leq 10^{-5} K$ ) temperatures, respectively, BE condensation of quasiparticles and mesons occurs at high temperatures,  $\sim 100 K$  and  $\sim 10^{12}\,K$  respectively.

Despite the significant variation in temperature, in all these four cases, the condensate corresponds to temperatures below or of the order of the BE condensation temperature of an infinite ideal gas,  $T_c^{(i)} \approx 3.3\hbar^2 n^{2/3}/m$  [12, 13] (this is the formula for the nonrelativistic case; m and n = N/V denote the boson mass and density). Since the physical measure of temperature is precisely the value of  $T_c^{(i)}$ , all the BE condensates mentioned above are low-temperature ones. However, there are also temperatures  $T \gg T_c^{(i)}$ , which we will call ultrahigh temperatures, to distinguish them from "high" temperatures,

which are typically below  $T_c^{(i)}$ ). For an infinite threedimensional ideal gas, it has been theoretically found that  $N_0(T)/N = 1 - (T/T_c^{(i)})^{3/2}$  at  $T \leq T_c^{(i)}$  and  $N_0 = 0$  at  $T > T_c^{(i)}$ , where  $N_0$  is the number of zero-momentum atoms [12, 13]. It is widely accepted, as supported by experimental evidence, that at  $T \gg T_c^{(i)}$  there is no condensate also in an interacting Bose gas. However, to our knowledge, this rather obvious property has not been rigorously proven theoretically. In what follows, we propose a sketchy mathematical proof, partly grounded in physical considerations, that BE condensation of zeromomentum atoms is impossible for a system of interacting spinless bosons at ultrahigh temperatures.

## II. MAIN PART

For an ideal Bose gas, the condensate is typically evaluated by the method proposed as early as by A. Einstein [14]: the system is treated as a collection of particles, and the occupation of the single-particle states is analysed. Such a method, however, is not applicable to a system of *N* interacting bosons, since each state of such a system is described by an *N*-particle wave function (WF)  $\Psi(\mathbf{r}_1, \ldots, \mathbf{r}_N)$ , and single-particle states are not well-defined. In this case, the condensate can be calculated using a more general and universal approach based on the diagonal expansion of the single-particle density matrix  $F_1(\mathbf{r}, \mathbf{r'})$  in the complete orthonormalised basis  $\{\phi_j(\mathbf{r})\}$  [12]:

$$F_1(\mathbf{r}, \mathbf{r}') = \sum_{j=1}^{\infty} \lambda_j \phi_j^*(\mathbf{r}') \phi_j(\mathbf{r}).$$
(1)

Here,  $\lambda_j$  are the occupation numbers of the single-particle states  $\phi_j(\mathbf{r})$ , and  $\lambda_j/N$  is the probability that a particle occupies the state  $\phi_j(x)$  (we use the normalisation of  $F_1(\mathbf{r}, \mathbf{r}')$ , for which  $\lambda_1 + \ldots + \lambda_\infty = N$ ). The state  $\phi_1(\mathbf{r})$ corresponds to the condensate if  $\lambda_1 \sim N$ . This is the definition of the condensate of Bose particles for a system

<sup>\*</sup>Electronic address: mtomchenko@bitp.kyiv.ua

of many interacting bosons. This definition is also valid for free bosons, then each  $\lambda_j$  can take natural values or be zero.

Consider a nonrelativistic, equilibrium, uniform system of interacting spinless bosons with a fixed number N of particles at temperature T. For simplicity, we use periodic boundary conditions. In the formalism of second quantisation, we have

$$F_1(\mathbf{r}, \mathbf{r}') = \langle \hat{\psi}^+(\mathbf{r}') \hat{\psi}(\mathbf{r}) \rangle, \qquad (2)$$

$$\hat{\psi}(\mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{a}_{\mathbf{k}} e^{i\mathbf{k}\mathbf{r}}, \ \hat{\psi}^+(\mathbf{r}) = \frac{1}{\sqrt{V}} \sum_{\mathbf{k}} \hat{a}_{\mathbf{k}}^+ e^{-i\mathbf{k}\mathbf{r}}.$$
 (3)

Here and below,  $\langle \rangle$  stands for the statistical average over the canonical ensemble [13]:

$$\begin{aligned} \langle \hat{A} \rangle_{T>0} &= \frac{1}{Z} \int d\mathbf{r}_1 \dots d\mathbf{r}_N \sum_{\wp} e^{-E_{\wp}/k_B T} \Psi_{\wp}^* \hat{A} \Psi_{\wp} \\ &\equiv \frac{1}{Z} \sum_{\wp} e^{-E_{\wp}/k_B T} \langle \Psi_{\wp} | \hat{A} | \Psi_{\wp} \rangle, \end{aligned} \tag{4}$$

where  $Z = \sum_{\wp} e^{-E_{\wp}/k_B T}$ ,  $\{\Psi_{\wp}\}$  represents a complete orthonormalised set of WFs of the system with a fixed number N of particles, and  $\wp$  numerates the physically different states. The WFs  $\Psi_{\wp}$  and  $\Psi_{\wp}$  describe the same stationary state in the coordinate representation and the second quantised representation, respectively. From Eqs. (2)–(4) we obtain

$$F_{1}(\mathbf{r},\mathbf{r}') = \frac{1}{V} \sum_{\mathbf{k},\mathbf{q}} \langle \hat{a}_{\mathbf{q}}^{+} \hat{a}_{\mathbf{k}} \rangle e^{i\mathbf{k}\mathbf{r} - i\mathbf{q}\mathbf{r}'}$$
$$= \frac{1}{V} \sum_{\mathbf{k}} \langle \hat{a}_{\mathbf{k}}^{+} \hat{a}_{\mathbf{k}} \rangle e^{i\mathbf{k}(\mathbf{r}-\mathbf{r}')} \equiv \sum_{\mathbf{k}} N_{\mathbf{k}} \phi_{\mathbf{k}}^{*}(\mathbf{r}') \phi_{\mathbf{k}}(\mathbf{r}), \quad (5)$$

because  $\langle \Psi_{\wp} | \hat{a}_{\mathbf{q}}^{+} \hat{a}_{\mathbf{k}} | \Psi_{\wp} \rangle = 0$  if  $\mathbf{q} \neq \mathbf{k}$ ; here  $N_{\mathbf{k}} = \langle \hat{a}_{\mathbf{k}}^{+} \hat{a}_{\mathbf{k}} \rangle$ and  $\phi_{\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}}/\sqrt{V}$ .

Thus, for the periodic system, the expansion of  $F_1(\mathbf{r}, \mathbf{r}')$  directly yields a diagonal form, and the number of zero-momentum atoms is given by the formula

$$N_0 = \langle \hat{a}_{\mathbf{0}}^+ \hat{a}_{\mathbf{0}} \rangle = \frac{1}{Z} \sum_{\wp} e^{-E_{\wp}/k_B T} \langle \Psi_{\wp} | \hat{a}_{\mathbf{0}}^+ \hat{a}_{\mathbf{0}} | \Psi_{\wp} \rangle.$$
(6)

Here, each number  $\wp$  corresponds to a set of elementary quasiparticles with momenta  $\mathbf{p}_1, \mathbf{p}_2, \ldots$ ;  $\wp \equiv \{n_{\mathbf{p}_1}, n_{\mathbf{p}_2}, \ldots\}$ , where  $n_{\mathbf{p}_j}$  is the number of quasiparticles with momentum  $\mathbf{p}_j$ . Let us denote  $\langle \Psi_{\wp} | \hat{a}_{\mathbf{0}}^{\dagger} \hat{a}_{\mathbf{0}} | \Psi_{\wp} \rangle = N_0^{(\wp)}$ , then

$$N_0 = \frac{\sum\limits_{\wp} N_0^{(\wp)} e^{-E_{\wp}/k_B T}}{\sum\limits_{\wp} e^{-E_{\wp}/k_B T}},$$
(7)

where each term in the numerator and denominator is non-negative:  $e^{-E_{\wp}/k_BT} > 0$ ,  $N_0^{(\wp)} \ge 0$ . For a small number of quasiparticles, one has

$$E_{\wp} = E_0 + \sum_{\mathbf{p}_j} n_{\mathbf{p}_j} \epsilon(\mathbf{p}_j), \qquad (8)$$

where  $E_0$  is the ground-state energy and  $\epsilon(\mathbf{p}_i)$  is the energy of the quasiparticle with momentum  $\mathbf{p}_i$ . When the number of quasiparticles is large ( $\sim N$ ), interactions between them must be accounted for, leading to a modification of the dispersion relation in (8):  $\epsilon(\mathbf{p}_i) \rightarrow \epsilon(\mathbf{p}_i)$  $\epsilon(\mathbf{p}_j; n_{\mathbf{p}_1}, n_{\mathbf{p}_2}, \ldots)$ . Each state of the system of N interacting bosons is an N-particle state. Accordingly, (i) one should describe such a system in the language of collective excitations (elementary quasiparticles), even at ultrahigh temperatures. Experiment shows that at ultrahigh temperatures the matter is in a gaseous, nonsuperfluid state. Therefore, we assume that our system is such a gas, and that the Bogoliubov formula [15]  $\epsilon_B(\mathbf{k}) = \sqrt{K^2(\mathbf{k}) + 2n_0\nu(k)K(\mathbf{k})}$ , where  $K(\mathbf{k}) =$  $\hbar^2 \mathbf{k}^2/2m$  and  $n_0 = N_0/V$ , approximately describes the quasiparticle dispersion law. It is well known that (ii) such a gas can be treated as a collection of N interacting atoms with the Maxwell-Boltzmann momentum distribution (note that known to us methods of deriving this distribution imply continuity and differentiability of the distribution function, and thus the absence of condensate). We are also aware that (iii) the free energy of the system,  $F = -k_B T \ln Z$ , and other thermodynamic quantities can be derived from the statistical sum [13]. Properties (i), (ii) and (iii) jointly imply that at ultrahigh temperatures, the main contribution to the statistical sum  $Z = \sum_{\wp} e^{-E_{\wp}/k_B T}$  should come from the states, containing N quasiparticles with the dispersion law  $\epsilon(\mathbf{p}) \approx \hbar^2 \mathbf{p}^2 / 2m$ , which follows from the Bogoliubov formula  $\epsilon_B(\mathbf{p})$  at  $|\mathbf{p}| \to \infty$ . This establishes a correspondence between the quasiparticle and particle descriptions of the system. We henceforth use the language of quasiparticles as more rigorous. It is worth noting that although only single-quasiparticle states [16, 17] are usually described quantum mechanically, it is also possible to accurately describe two- [18] and multi-quasiparticle [18, 19] states.

The formula (7), being *exact*, directly implies that a BE condensate should be absent at ultrahigh temperatures. Indeed, at T = 0, only the ground state (i.e., the state without quasiparticles) contributes to the formula (7), yielding  $N_0 = N_0^{(0)}$ . The structure of the ground-state WF of a Bose gas and a Bose liquid ensures the existence of a BE condensate of atoms with zero momentum:  $N_0 = N_0^{(0)} \sim N$  [13, 20–24]. At high temperatures,  $T \gtrsim T_c^{(i)}$ , a huge number of non-negative terms contribute to the sums in the numerator and denominator of the right-hand side of Eq. (7). Therefore, at high T: (i) the change in  $N_0$  with increasing T must be very smooth, and (ii) for  $N_0 \sim N$  to hold, it is necessary that  $N_0^{(\wp)} \sim N$  be valid for the majority of the significant terms in (7).

At  $T \gg T_c^{(i)}$ , the main contribution to the statistical sum comes from states, containing N quasiparticles with large momenta  $|\mathbf{p}|$  (see above; this is apparently due to the fact that such states constitute the majority among all possible states; interestingly, the maximum possible number of elementary quasiparticles is also N [18, 19]). Consequently, in order to get  $N_0 \sim N$  at  $T \gg T_c^{(i)}$ , it is necessary and sufficient that  $N_0^{(\wp)} \sim N$  holds for states with N quasiparticles with large momenta  $|\mathbf{p}|$ . (The sufficiency of this condition is evident; and its necessity follows from the fact that if  $N_0^{(\wp)} \sim N$  holds only for states containing < N quasiparticles, the condensate would be suppressed by the statistical sum. This is because the main contribution to the sum is provided by states with N quasiparticles, which greatly outnumber states with fewer quasiparticles.) The following qualitative reasoning shows that this is impossible.

The WF of the ground state of a Bose gas or a Bose liquid has the form [18, 25]

$$\Psi_0(\mathbf{r}_1,\dots,\mathbf{r}_N) = A_0 e^{S(\mathbf{r}_1,\dots,\mathbf{r}_N)},\tag{9}$$

$$S = \sum_{\mathbf{q}_{1}\neq 0} \frac{c_{2}(\mathbf{q}_{1})}{2!} \rho_{\mathbf{q}_{1}} \rho_{-\mathbf{q}_{1}} + \sum_{\mathbf{q}_{1},\mathbf{q}_{2}\neq 0}^{\mathbf{q}_{1}+\mathbf{q}_{2}\neq 0} \frac{c_{3}(\mathbf{q}_{1},\mathbf{q}_{2})}{3!N^{1/2}} \rho_{\mathbf{q}_{1}} \rho_{\mathbf{q}_{2}} \rho_{-\mathbf{q}_{1}-\mathbf{q}_{2}} + \dots + \sum_{\mathbf{q}_{1},\dots,\mathbf{q}_{N-1}\neq 0}^{\mathbf{q}_{1}+\dots+\mathbf{q}_{N-1}\neq 0} \frac{c_{N}(\mathbf{q}_{1},\dots,\mathbf{q}_{N-1})}{N!N^{(N-2)/2}} \times \times \rho_{\mathbf{q}_{1}}\dots\rho_{\mathbf{q}_{N-1}} \rho_{-\mathbf{q}_{1}-\dots-\mathbf{q}_{N-1}},$$
(10)

where  $\rho_{\mathbf{k}} = \frac{1}{\sqrt{N}} \sum_{j=1}^{N} e^{-i\mathbf{k}\mathbf{r}_{j}}$ . Any excited state (with total momentum  $\hbar \mathbf{p}$  and any number  $\leq N$  of quasiparticles) is described by the WF [16, 18]

$$\Psi_{\mathbf{p}}(\mathbf{r}_1,\ldots,\mathbf{r}_N) = A_{\mathbf{p}}\psi_{\mathbf{p}}\Psi_0,\tag{11}$$

$$\psi_{\mathbf{p}} = b_{1}(\mathbf{p})\rho_{-\mathbf{p}} + \sum_{\mathbf{q}_{1}\neq0}^{\mathbf{q}_{1}+\mathbf{p}\neq0} \frac{b_{2}(\mathbf{q}_{1};\mathbf{p})}{2!N^{1/2}}\rho_{\mathbf{q}_{1}}\rho_{-\mathbf{q}_{1}-\mathbf{p}} \\ + \sum_{\mathbf{q}_{1},\mathbf{q}_{2}\neq0}^{\mathbf{q}_{1}+\mathbf{q}_{2}+\mathbf{p}\neq0} \frac{b_{3}(\mathbf{q}_{1},\mathbf{q}_{2};\mathbf{p})}{3!N}\rho_{\mathbf{q}_{1}}\rho_{\mathbf{q}_{2}}\rho_{-\mathbf{q}_{1}-\mathbf{q}_{2}-\mathbf{p}} + \dots \\ + \sum_{\mathbf{q}_{1}+\dots+\mathbf{q}_{N-1}+\mathbf{p}\neq0}^{\mathbf{q}_{1}+\dots+\mathbf{q}_{N-1}+\mathbf{p}\neq0} \frac{b_{N}(\mathbf{q}_{1},\dots,\mathbf{q}_{N-1};\mathbf{p})}{N!N^{(N-1)/2}} \times \\ \times \rho_{\mathbf{q}_{1}}\dots\rho_{\mathbf{q}_{N-1}}\rho_{-\mathbf{q}_{1}-\dots-\mathbf{q}_{N-1}-\mathbf{p}}.$$
(12)

The formulae (9)–(12) are exact: (10) and (12) are simply expansions of the functions  $\ln \Psi_0$  and  $\Psi_p$  in the complete set of Bose-symmetric functions  $1, \rho_{-\mathbf{q}_1}, \rho_{-\mathbf{q}_1}\rho_{-\mathbf{q}_2}, \ldots, \rho_{-\mathbf{q}_1}\rho_{-\mathbf{q}_2}\cdots\rho_{-\mathbf{q}_N}$ , each associated with a specific total momentum. Consider a dilute Bose gas. In this case [26],

$$\Psi_0 = A_0 e^S, \quad S \approx \sum_{\mathbf{k} \neq 0} \frac{c_2(\mathbf{k})}{2!} \rho_{\mathbf{k}} \rho_{-\mathbf{k}}, \tag{13}$$

$$c_2(\mathbf{k}) \approx 1/2 - m\epsilon_B(\mathbf{k})/(\hbar^2 k^2).$$
(14)

It follows from the treatment in works [23, 27] that for liquid He II for configurations  $(\mathbf{r}_1, \ldots, \mathbf{r}_N)$  corresponding to a uniform atomic distribution  $(|\rho_{\mathbf{k}}| \sim 1)$ , one has  $S \equiv S_l \sim -N$ . Consequently, for a gaseous <sup>4</sup>He it should be  $S \equiv S_g \sim -|a|N$ , where |a| is a small number that depends on the densities of the gas and liquid. Since  $|S_g| \ll |S_l|$ , almost all atoms of the gas are in the condensate:  $N_0 \approx N$ ,  $N - N_0 \ll N$  [15]. Therefore, when calculating the condensate in the gas at T = 0, it suffices to assume  $S_g = 0$  in a zero-order approximation, yielding

$$\Psi_0(\mathbf{r}_1,\ldots,\mathbf{r}_N)\approx 1. \tag{15}$$

(In (15) and below, we omit normalisation constants; note that the approximation (15) is unsuitable for the derivation of the dispersion law, as it results in  $\epsilon(\mathbf{k}) \sim k^2$ for all k.) A state with a single Bogoliubov's quasiparticle with momentum **p** is approximately described by the WF [16–18, 28]

$$\Psi_{\mathbf{p}}(\mathbf{r}_1,\ldots,\mathbf{r}_N)\approx\rho_{-\mathbf{p}}\Psi_0.$$
(16)

Given (15), we may write

$$\Psi_{\mathbf{p}}(\mathbf{r}_1, \dots, \mathbf{r}_N) \approx \rho_{-\mathbf{p}}.$$
(17)

Similarly, the state containing a quasiparticle with momentum  $\mathbf{p}_1$  and another quasiparticle with momentum  $\mathbf{p}_2$ , in the zero-order approximation can be described by the WF [18, 28]

$$\Psi_{\mathbf{p}_1\mathbf{p}_2}(\mathbf{r}_1,\ldots,\mathbf{r}_N) \approx \rho_{-\mathbf{p}_1}\rho_{-\mathbf{p}_2},\tag{18}$$

and a state with N quasiparticles with momenta  $\mathbf{p}_1, \mathbf{p}_2, \ldots, \mathbf{p}_N$  is described in the same approximation by the WF

$$\Psi_{\mathbf{p}_1\mathbf{p}_2\ldots\mathbf{p}_N}(\mathbf{r}_1,\ldots,\mathbf{r}_N)\approx\rho_{-\mathbf{p}_1}\rho_{-\mathbf{p}_2}\cdots\rho_{-\mathbf{p}_N}.$$
 (19)

There are two ways of showing that the maximum possible number of elementary quasiparticles is N (see appendix 1 in [18] and section 7 in [29]). For a Bose gas, this can be seen without using formulae: quasiparticles with large momentum  $|\mathbf{p}|$  have the energy  $\epsilon(\mathbf{p}) \approx \hbar^2 \mathbf{p}^2/2m$ [15] and are similar to free atoms; the number of the latter is equal to N, hence the maximum possible number of elementary quasiparticles with large  $|\mathbf{p}|$  is also equal to N. The treatment in [18, 29] shows that this conclusion holds for elementary quasiparticles with arbitrary momenta. Interestingly, the number of rotons in <sup>4</sup>He at  $T = T_{\lambda}$  is of the order of N: The density of free rotons,  $n_r$ , can be determined in two ways: (1) from the relations  $\rho_n = Q_r^2 n_r/(3k_BT)$  [30] and  $\rho_n = \rho$ , and (2) from the formula  $n_r = 2Q_r^2(\mu_r k_B T)^{1/2}e^{-\Delta/k_b T}((2\pi)^{3/2}\hbar^3)^{-1}$ [30]. This gives  $n_r \approx n/7$  and  $n_r \approx n/6$ , respectively (here n = N/V; the parameters  $\rho(T_{\lambda}) = 0.1462 \, g/cm^3$ ,  $\Delta(T_{\lambda}) \approx 0.56 \, meV$ ,  $\mu_r(T_{\lambda}) \approx 0.135 \, m_4$ , and  $Q_r(T_{\lambda}) \approx \hbar \cdot 1.93 \, \text{\AA}^{-1}$  are taken from [31, 32] and correspond to the saturated vapour pressure). Accounting for roton interactions, we obtain  $n_r \simeq (0.1-1)n$ .

So, for a dilute Bose gas, we have the following picture in the zero-order approximation. The ground state (15)does not contain quasiparticles; in this case, all atoms remain in the BE condensate of zero-momentum atoms. The state (16) corresponds to a single Bogoliubov's quasiparticle; in this case, N-1 atoms have zero momentum and one atom has momentum  $\mathbf{p}$ . The state (17) corresponds to two quasiparticles with momenta  $\mathbf{p}_1$  and  $\mathbf{p}_2$ ; in that case, N-2 atoms have zero momentum, one atom has momentum  $\mathbf{p}_1$ , and another has momentum  $\mathbf{p}_2$ . And so on. Finally, the state (19) represents N quasiparticles with momenta  $\mathbf{p}_1, \ldots, \mathbf{p}_N$ ; in this case, N atoms have the same momenta  $\mathbf{p}_1, \ldots, \mathbf{p}_N$ . Thus, with an increase in the number of elementary quasiparticles, the BE condensate of zero-momentum atoms gradually fades away. And when the number of quasiparticles becomes close to N, this condensate disappears completely.

Note that the average number of quasiparticles is given by the formula

$$\bar{N}_Q(T) = \frac{1}{Z} \int d\mathbf{r}_1 \dots d\mathbf{r}_N \sum_{\wp} e^{-E_{\wp}/k_B T} \Psi_{\wp}^* \hat{N}_{Qp} \Psi_{\wp}$$
$$\equiv \frac{1}{Z} \sum_{\wp} e^{-E_{\wp}/k_B T} N_{Qp}^{(\wp)}, \qquad (20)$$

where  $N_{Qp}^{(\wp)}$  is the number of quasiparticles for a state  $\Psi_{\wp}$ . Since states containing N quasiparticles correspond to the relation  $N_{Qp}^{(\wp)} = N$  and form the majority of all possible states, at  $T \gg T_c^{(i)}$  we should have  $\bar{N}_Q(T) \approx N$ .

The formulae (15)–(19) do not take into account the interatomic interaction. The latter leads to a complication in the form of the WFs in accordance with relations (9)–(12) and to an additional "blurring" of the condensate (since terms with a larger number of  $\rho_{-\mathbf{p}_i}$  have to be considered in the expansion of  $\Psi_{\mathbf{p}_1...\mathbf{p}_j}(\mathbf{r}_1,\ldots,\mathbf{r}_N)$  in  $\rho_{-\mathbf{p}_i}$ ). It implies that our conclusions above are valid not only for dilute gas but also for dense gas and liquid. This reasoning shows that gas states containing Nelementary quasiparticles are characterized by the property  $N_0^{(\wp)} \sim 1 \ll N$ . This means that, according to the formula (7) and the ensuing treatment, at  $T \gg T_c^{(i)}$ there is no BE condensate of zero-momentum atoms:  $N_0 \sim 1 \ll N$ .

We also remark that WFs (9)–(12) can be written in a different form. Since for a system of N Bose particles the functions  $1, \rho_{-\mathbf{q}_1}, \rho_{-\mathbf{q}_1}\rho_{-\mathbf{q}_2}, \ldots, \rho_{-\mathbf{q}_1}\rho_{-\mathbf{q}_2}\cdots\rho_{-\mathbf{q}_N}$  form a complete (though non-orthogonal) set of basis functions [25], any system state (ground or excited) with total mo-

mentum  $\hbar \mathbf{p}$  can be written as

$$\Psi_{\mathbf{p}}(\mathbf{r}_{1},\ldots,\mathbf{r}_{N}) = d_{1}(\mathbf{p})\rho_{-\mathbf{p}}$$

$$+ \sum_{\mathbf{q}_{1}\neq0}^{\mathbf{q}_{1}+\mathbf{p}\neq0} \frac{d_{2}(\mathbf{q}_{1};\mathbf{p})}{2!N^{1/2}}\rho_{\mathbf{q}_{1}}\rho_{-\mathbf{q}_{1}-\mathbf{p}} + \ldots$$

$$+ \sum_{\mathbf{q}_{1},\ldots,\mathbf{q}_{N-1}\neq0}^{\mathbf{q}_{1}+\ldots+\mathbf{q}_{N-1}+\mathbf{p}\neq0} \frac{d_{N}(\mathbf{q}_{1},\ldots,\mathbf{q}_{N-1};\mathbf{p})}{N!N^{(N-1)/2}} \times$$

$$\times \rho_{\mathbf{q}_{1}}\ldots \rho_{\mathbf{q}_{N-1}}\rho_{-\mathbf{q}_{1}-\ldots-\mathbf{q}_{N-1}-\mathbf{p}}.$$
(21)

For the ground state,  $\mathbf{p} = 0$ . If the ground state of a Bose system is a liquid or a gas, then both the formulae (9)–(12) and (21) are applicable, but the formulae (9)– (12) are preferable for several reasons. However, which WFs can adequately describe the gaseous or liquid phase when the ground state of the system is a crystal? We have not found an answer in the literature. The WF (21) is suitable for this case, while the solutions (11), (12) yield a non-isotropic dispersion law, because of the anisotropy of the crystalline ground-state WF  $\Psi_0$ .

The WFs (9)–(12) and (21) can be represented in the second quantised form  $\Psi_{\wp}$ , by replacing  $\rho_{\mathbf{k}\neq 0}$  with  $\frac{1}{\sqrt{N}}\sum_{\mathbf{q}} \hat{a}^+_{\mathbf{q}-\mathbf{k}} \hat{a}_{\mathbf{q}}$  [33] in (9)–(12), (21) and adding the factor  $[\hat{a}^+_0]^N |0_{bare}\rangle$  at the end of the right-hand side of formulae (9) and (21) [19].

Note that in addition to the condensate  $N_0$  of zeromomentum atoms, the condensate  $N_{\mathbf{p}\neq 0}$  of atoms with non-zero momentum  $\mathbf{p}$  can also exist. According to the treatment in [29], such a condensate is only possible if there is a corresponding condensate of elementary quasiparticles. It is clear that ultrahigh temperature should also suppress such a condensate of atoms. However, it appears feasible to create it by means of an external field [29], possibly even at ultrahigh temperatures. But we cannot imagine how the condensate  $N_0$  could be produced at ultrahigh temperatures.

It is also interesting to note that although a BE condensate is typically associated with a large number of particles  $(N_0, N \gg 1)$ , it is possible — and reasonable — to define a condensate even for systems with a *small* number of particles  $(N \ge 2)$  [34]. This is because the Gross-Pitaevskii equation describes few-particle systems more precisely than many-particle systems [34–36]. A few-particle condensate can be introduced based on the density matrix (1) using the criterion  $\lambda_1 \gg \lambda_2 + \ldots + \lambda_N$ [34]. However, our analysis above, as well as the very concept of temperature, assumes that  $N \gg 1$ .

## III. CONCLUSION

Recapitulating, we have proposed a mathematical reasoning showing that a BE condensate of zero-momentum atoms should be absent in a nonrelativistic system of many spinless interacting bosons at ultrahigh temperatures,  $T \gg T_c^{(i)}$ . This reasoning is not rigorous. Although we have considered the canonical ensemble, the

formula (7) can be generalised to the case of a grand canonical ensemble, and the conclusions should remain the same.

This research was supported in part by the Na-

- Novel Superfluids, K. H. Bennemann and J. B. Ketterson (eds.), Oxford University Press, Oxford (2013), Vol. 1. https://doi.org/10.1093/acprof:oso/9780199585915.001.0001
- [2] Novel Superfluids, K. H. Bennemann and J. B. Ketterson (eds.), Oxford University Press, Oxford (2014), Vol. 2. https://doi.org/10.1093/acprof:oso/9780198719267.001.0001
- [3] A. B. Migdal, E. E. Saperstein, M. A. Troitsky, and D. N. Voskresensky, *Phys. Rep.* **192**, 179 (1990). https://doi.org/10.1016/0370-1573(90)90132-L
- [4] M. Mannarelli, Particles 2, 411 (2019). https://doi.org/10.3390/particles2030025
- [5] S. O. Demokritov, V. E. Demidov, O. Dzyapko, G. A. Melkov, A. A. Serga, B. Hillebrands, and A. N. Slavin, *Nature* 443, 430 (2006). https://doi.org/10.1038/nature05117
- [6] L. H. Bennett and E. Della Torre, J. Mod. Phys. 5, 693 (2014). https://doi.org/10.4236/jmp.2014.58081
- [7] A. Kogar, M. S. Rak, S. Vig, et al., Science 358, 1314 (2017). https://doi.org/10.1126/science.aam6432
- [8] D. Wang, N. Luo, W. Duan, and X. Zou, *J. Phys. Chem. Lett.* **12**, 5479 (2021). https://doi.org/10.1021/acs.jpclett.1c01370
- [9] T. Byrnes, N. Y. Kim, and Y. Yamamoto, *Nature Physics* 10, 803 (2014). https://doi.org/10.1038/nphys3143
- [10] J. Klaers, J. Schmitt, F. Vewinger, and M. Weitz, *Nature* 468, 545 (2010). https://doi.org/10.1038/nature09567
- [11] J. B. Ketterson and K. H. Bennemann, Survey of Some Novel Superfluids, in [1], pp. 74–155. https://doi.org/10.1093/acprof:oso/9780199585915.003.0002
- [12] C. J. Pethick and H. Smith, Bose-Einstein Condensation in Dilute Gases, Cambridge University Press, New York (2008), Chaps. 2, 13, 15.
- [13] K. Huang, *Statistical Mechanics*, Wiley, New York (1963).
- [14] A. Einstein, Sitzungsber. Preuss. Akad. Wiss., Phys.-Math. Kl., p. 3 (1925).
- [15] N. N. Bogoliubov, J. Phys. USSR 11, 23 (1947).
- [16] I. A. Vakarchuk and I. R. Yukhnovskii, *Theor. Math. Phys.* 42, 73 (1980). https://doi.org/10.1007/BF01019263
- [17] R. P. Feynman, Statistical Mechanics: A Set of Lectures, Benjamin, Massachusetts (1972).
- [18] M. Tomchenko, J. Low Temp. Phys. 201, 463 (2020). https://doi.org/10.1007/s10909-020-02498-z

tional Academy of Sciences of Ukraine (Project No. 0121U109612). The author thanks the Simons Foundation for additional financial support.

- [19] M. Tomchenko, arXiv:2501.00893 [cond-mat.quant-gas].
- [20] L. Reatto and G. V. Chester, Phys. Rev. 155, 88 (1967). https://doi.org/10.1103/PhysRev.155.88
- W. E. Parry and C. R. Rathbone, Proc. Phys. Soc. 91, 273 (1967). https://doi.org/10.1088/0370-1328/91/2/302
- [22] L. Reatto and G. L. Masserini, *Phys. Rev. B* 38, 4516 (1988). https://doi.org/10.1103/PhysRevB.38.4516
- M. Tomchenko, Fiz. Nizk. Temp. 32, 53
   (2006) [Low Temp. Phys. 32, 38 (2006)]. https://doi.org/10.1063/1.2160507
- [24] A. A. Rovenchak and I. O. Vakarchuk, J. Phys. Stud. 11, 404 (2007). https://doi.org/10.30970/jps.11.404
- [25] I. A. Vakarchuk and I. R. Yukhnovskii, *Theor. Math. Phys.* **40**, 626 (1979). https://doi.org/10.1007/BF01019246
- [26] N. N. Bogoliubov and D. N. Zubarev, Sov. Phys. JETP 1, 83 (1956).
- [27] M. D. Tomchenko, JETP 102, 137 (2006). https://doi.org/10.1134/S106377610601016X
- [28] R. P. Feynman, Phys. Rev. 94, 262 (1954). https://doi.org/10.1103/PhysRev.94.262
- M. Tomchenko, J. Phys. A: Math. Theor. 57, 495205
   (2024) https://doi.org/10.1088/1751-8121/ad9188
- [30] I. M. Khalatnikov, An Introduction to the Theory of Superfluidity, Perseus, New York (2000).
- [31] B. N. Esel'son, V. N. Grigor'ev, V. G. Ivantsov, E. Ya. Rudavskii, D. N. Sanikadze, and I. A. Serbin, *Properties of Liquid and Solid Helium*, Izd. Standartov, Moscow (1978) [in Russian].
- [32] M. R. Gibbs, K. H. Andersen, W. G. Stirling, and H. Schober, J. Phys.: Condens. Matter 11, 603 (1999). https://doi.org/10.1088/0953-8984/11/3/003
- [33] D. Pines and P. Noziéres, The Theory of Quantum Liquids: Normal Fermi Liquids, CRC Press, Boca Raton (1989). https://doi.org/10.4324/9780429492662
- [34] M. Tomchenko, J. Phys. A: Math. Theor. 57, 495202 (2024). https://doi.org/10.1088/1751-8121/ad9187
- [35] D. Blume and C. H. Greene, *Phys. Rev. A* 63, 063601 (2001). https://doi.org/10.1103/PhysRevA.63.063601
- [36] D. Blume, Rep. Prog. Phys. 75, 046401 (2012). https://doi.org/10.1088/0034-4885/75/4/046401