FROBENIUS-SCHUR INDICATORS OF DUAL FUSION CATEGORIES AND SEMISIMPLE PARTIALLY DUALIZED QUASI-HOPF ALGEBRAS

KANGQIAO LI

Abstract. Frobenius-Schur indicators of objects in pivotal monoidal categories were defined and formulated by Ng and Schauenburg in 2007. In this paper, we introduce and study an analogous formula for indicators in the dual category $\mathcal{C}_{\mathcal{M}}^*$ to a spherical fusion category $\mathcal C$ (with respect to an indecomposable semisimple module category $\mathcal M$) over $\mathbb C$. Our main theorem is a relation between indicators of specific objects in $\mathcal{C}_{\mathcal{M}}^*$ and \mathcal{C} . As consequences: 1) We obtain equalities on the indicators between certain representations and the exponents of a semisimple complex Hopf algebra as well as its partially dualized quasi-Hopf algebra; 2) We show that the property in Kashina's conjecture on the exponent holds for a class of semisimple Hopf algebras fitting into abelian extensions with assumptions weaker than split.

CONTENTS

[^{5.} Applications: Indicators and exponents of semisimple Hopf algebras and its left partial dualization](#page-27-0) 28

Key words and phrases. Frobenius-Schur indicator; Fusion category; Dual tensor category; Hopf algebra; Partial dualization; Exponent.

²⁰²⁰ Mathematics Subject Classification. 18M15, 16T05, 18M20.

[†] This work was supported by National Natural Science Foundation of China [grant number 12301049].

- [5.1. Canonical pivotal structures of integral fusion categories,](#page-27-1) and indicators for semisimple Hopf algebras
- [5.2. Semisimple left partially dualized quasi-Hopf algebras and the reconstruction](#page-28-0) 29
- [5.3. Relations between the indicators of a semisimple Hopf algebra and](#page-32-0) its left partial duals 33
- [5.4. Invariance for the exponent of semisimple \(quasi-\)Hopf algebras under partial dualities](#page-37-0) 38

[Acknowledgement](#page-40-0) 41

[References](#page-40-1) 41

1. INTRODUCTION

The notion of (higher) Frobenius-Schur indicators ν_n of modules over a finite-dimensional Hopf algebras or objects in pivotal monoidal categories is a classical invariant under equivalences. It was defined firstly by Linchenko and Montgomery [\[LM00\]](#page-41-0) for a semisimple Hopf algebra H , and they provided a criteria for a simple H -module V being self-dual with the value of $\nu_2(V)$. Afterwards in [\[KSZ06\]](#page-41-1), Kashina, Sommerhäuser and Zhu described various invariants corresponding to simple modules with properties of the higher Frobenius-Schur indicators for semisimple Hopf algebras, including some essential formulas. One of the formulas was later expressed to define the regular Frobenius-Schur indicators of a finite-dimensional Hopf algebra H (not necessarily semisimple) by Kashina, Montgomery and Ng in [\[KMN12\]](#page-41-2), which was generalized and studied further by Shimizu [\[Shi15\]](#page-42-0).

Moreover for a k-linear pivotal monoidal category \mathcal{C} , Ng and Schauenburg defined in [\[NS07a\]](#page-41-3) the Frobenius-Schur indicators of objects in C by a categorical version of another formula in [\[KSZ06\]](#page-41-1). The definition indeed generalizes the indicators for a semisimple Hopf algebra H, as they coincide when $\mathcal{C} = \text{Rep}(H)$ is the category of finite-dimensional modules over H. Furthermore, if C is a spherical fusion category over the complex field $\mathbb C$, they obtained in [\[NS07b\]](#page-42-1) a formula on the Frobenius-Schur indicators, which is described by the pivotal trace "ptr(−)" of the power of the ribbon structure θ on the center $\mathcal{Z}(\mathcal{C})$: For any object $X \in \mathcal{C}$, its n-th Frobenius-Schur indicator equals to

$$
\nu_n(X) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta_{I(X)}^n\right),\tag{1.1}
$$

where dim(C) denotes the categorical dimension of C, and $I: \mathcal{C} \to \mathcal{Z}(\mathcal{C})$ is a two-sided adjoint to the forgetful functor. This formula was used to estimate the values of indicators and Frobenius-Schur exponents. As for non-pivotal categories, a theory of Frobenius-Schur indicators was developed by Shimizu [\[Shi17a\]](#page-42-2) with the construction of the pivotal cover.

Clearly, the results on Frobenius-Schur indicators for categories would help to find properties and applications of indicators for semisimple (quasi-)Hopf algebras. One might see [\[MN05,](#page-41-4) [Sch04,](#page-42-3) [NS08\]](#page-42-4) and so on for the case of quasi-Hopf algebras.

To the knowledge of the author, most common invariance properties of Frobenius-Schur indicators for categories were studied under tensor equivalences or center constructions. Our goal of the paper to find some invariance properties under the dual of (semisimple) tensor categories, or to find a categorical version of the results such as

$$
\nu_n(H^*) = \nu_n(H) \tag{1.2}
$$

in the case of representations of a Hopf algebra H , which were shown in $[\text{Shi12}]$ and $[\text{Shi15}]$. This motivates us to consider indicators of objects in $\mathcal{C}_{\mathcal{M}}^* := \text{Rex}_{\mathcal{C}}(\mathcal{M})^{\text{rev}}$, the dual category to C with respect to some C-module category M, since we know that $\mathsf{Rep}(H)_{\mathsf{Vec}}^* \approx \mathsf{Rep}(H^*)$ as tensor categories in the cases over Hopf algebras.

However, since we do not know when $\mathcal{C}_{\mathcal{M}}^*$ becomes pivotal, the Frobenius-Schur indicators are not well-defined in the original sense by Ng and Schauenburg. Thus we assume that $\mathcal C$ is a spherical fusion category, and attempt to redefine $\nu_n(\mathbf{F})$ for each object $\mathbf{F} \in C^*_{\mathcal{M}}$ with a formula which is completely similar to Equation (1.1) :

$$
\nu_n(\mathbf{F}) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta_{K(\mathbf{F})}^n\right),\,
$$

where K is a two-sided adjoint to the composition functor

$$
\mathcal{Z}(\mathcal{C}) \xrightarrow{\text{Schauenburg's equivalence}} \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \xrightarrow{\text{forgetful functor}} \mathcal{C}_{\mathcal{M}}^*.
$$

Fortunately, there is a description of the right adjoint functor K in $[\text{Shi20}]$ via the construction of ends, namely,

$$
K(\mathbf{F}) = \left(\int_{M \in \mathcal{M}} \underline{\text{Hom}}(M, F(M)), \sigma^{\mathbf{F}} \right) \in \mathcal{Z}(\mathcal{C}).
$$

With the help of this functor constructed, we are able to compute indicators $\nu_n(\mathbf{F})$ in some situations and obtain our main result($=$ Theorem [4.9\)](#page-25-0) in Section [4.3:](#page-21-0)

Theorem 1.1. *Let* C *be a spherical fusion category over* C*. Suppose that* A *is an algebra in* C such that $M = C_A$ is an indecomposable C-module category which is semisimple. Then for *any* $M \in \mathcal{C}_A$, the *n*-th indicator of $-\otimes M \in \mathcal{C}_{\mathcal{C}_A}^*$ is

$$
\nu_n(-\otimes M) = \nu_n(M) \qquad (\forall n \ge 1),
$$

where:

- The C-module functor $-\otimes M$ is canonically isomorphic to $(-) \otimes_A (A \otimes M)$ with *structure as the associativity constraint of* C*;*
- *The object* M *in the right-hand side is regarded as the image in* C *(under the forgetful functor from* C_A *to* C *)*.

We remark that any finite (left) C-module category M must be equivalent to C_A , the category of right A-modules in \mathcal{C} , for some algebra $A \in \mathcal{C}$. As a consequence, Theorem [1.1](#page-2-0) provides a relation between indicators in $\mathcal{C}_{\mathcal{M}}^*$ and \mathcal{C} .

Moreover, this result is applied to the following algebraic case: $C = \mathsf{Rep}(H)$ is the integral fusion category of representations of a semisimple complex Hopf algebra, and the indecomposable exact Rep(H)-module category $\mathcal{M} = \text{Rep}(B)$ holds for left coideal subalgebra B of H. As it is reconstructed in [\[Li23\]](#page-41-5) the so-called *(left) partially dualized quasi-Hopf algebra* $(H/B^+H)^* \#B$ from the dual tensor category $\mathsf{Rep}(H)^*_{\mathsf{Rep}(B)}$, we show by Theorem [1.1](#page-2-0) that $(=\mathrm{Theorem~5.12})$:

Theorem 1.2. *Let* H *be a semisimple Hopf algebra over* C*. Suppose* B *is a left coideal* $subalgebra, and $(H/B^+H)^* \#B$ is a left partially dualized quasi-Hopf algebra of H. Then for$ $any L \in \mathsf{Rep}(B)$,

$$
\nu_n((H/B^+H)^*\#L) = \nu_n(L\square_{B^*}H^*) \qquad (\forall n \ge 1),
$$

 $where \ (H/B^+H)^*#L \in \mathsf{Rep}((H/B^+H)^*#B) \ and \ L\square_{B^*}H^* \in \mathsf{Rep}(H) \ with \ structures \ defined$ *in Subsection [5.3.](#page-32-0)*

In particular, the indicators of the regular representations of $(H/B^+H)^* \#B$ and H coincide. Since the left partial dual $(H/B^+H)^* \# B$ is regarded as a generalization of the dual Hopf algebra H^* in a sense, we could obtain Equation (1.2) directly from Theorem [1.2.](#page-2-1)

Another gauge invariant for semisimple (quasi-)Hopf algebras is the *exponent* ([\[Kas99,](#page-41-6) [EG99,](#page-41-7) Eti02 etc.), which is in fact the period of the sequence of indicators (of the regular representation). It is conjectured by Kashina [\[Kas99,](#page-41-6) [Kas00\]](#page-41-9) that

$$
\exp(H) | \dim(H) \tag{1.3}
$$

holds for each semisimple and cosemisimple Hopf algebras. As an application of our results, we show a method to prove with the help of Theorem [1.2](#page-2-1) that a certain class of semisimple Hopf algebras satisfy the property (1.3) :

Theorem 1.3. *Let* H *be a semisimple Hopf algebra over* C *with a commutative subbialgebra* B*. Suppose that* B⁺H *is a biideal such that* H/B⁺H *is cocommutative, Then*

 $\exp(H)$ | dim (H)

if either of the following conditions hold:

- (1) *There is a left* B-module coalgebra map $\zeta : H \to B$ which is a retraction of the *inclusion* $B \rightarrowtail H$ *, or:*
- (2) *There is a right* H/B^+H -comodule algebra map $\gamma : H/B^+H \to H$ which is a section *of the quotient map* $H \rightarrow H/B^{+}H$.

This result(=Theorem [5.23\)](#page-39-0) generalizes a result stated in $[KSZ06]$ that every semisimple and cosemisimple Hopf algebra H fitting into a split abelian extension has property (1.3) . As an application, we show that a class of semisimple Hopf algebras H with a certain kind of commutative Hopf subalgebra with prime index satisfies [\(1.3\)](#page-3-1).

The paper is organized as follows: In Section [2,](#page-3-0) we collect some necessary notions of tensor categories and module categories, especially pivotal structures preserved by Schauenburg's equivalence $\Omega : \mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$, as well as constructions of adjoint functors to the forgetful functors from centers. The definition of Frobenius-Schur indicators in $\mathcal{C}_{\mathcal{M}}^{*}$ is defined in Section [3,](#page-11-0) and some basic invariance properties of indicators in various senses are discussed then. Section [4](#page-18-0) is devoted to our main theorem. Finally we introduce applications to indicators and exponent for semisimple Hopf algebras and its left partially dualized quasi-Hopf algebras in Section [5.](#page-27-0)

2. Preliminaries on monoidal categories, module categories and centers

We refer to [\[EGNO15,](#page-41-10) Chapters 1 to 4] for the elementary definitions and properties about monoidal categories and tensor categories.

Let C be a monoidal category with tensor product functor \otimes and unit object 1. In this paper, associativity and unit constraints in $\mathcal C$ are always abbreviated without the loss of generality, according to the coherence theorem [\[MacL63\]](#page-41-11).

For any object $X \in \mathcal{C}$, we would denote its left and right dual objects in C by X^* and *X respectively when exist. Also, the evaluation and coevaluation for the left dual X^* of X are denoted by

 $\operatorname{\mathsf{ev}}_X}: X^* \otimes X \to \mathbf{1} \quad \text{and} \quad \operatorname{\mathsf{coev}}_X: \mathbf{1} \to X \otimes X^*.$

When C is rigid, the *left categorical trace* could be defined for a morphism $f: X \to X^{**}$ in $\mathcal C$ as

$$
\mathrm{tr}(f):=\mathrm{ev}_{X^*}\circ (f\otimes \mathrm{id}_{X^*})\circ \mathrm{coev}_X\in \mathrm{End}_{\mathcal{C}}(\mathbf{1}).
$$

If C has furthermore a pivotal structure j, denote the *right pivotal trace* ([\[NS07b,](#page-42-1) Section 2]) of an endomorphism $g: X \to X$ in C by

$$
\operatorname{ptr}(g) := \operatorname{tr}(j_X \circ g).
$$

We remark that pivotal traces of (the powers of) a natural endomorphism are independent on the choices of objects in \mathcal{C} :

Lemma 2.1. Let C be a rigid monoidal category with pivotal structure j. Suppose θ is a *natural endomorphism on the identity functor* $\text{Id}_{\mathcal{C}}$. If two objects $X, Y \in \mathcal{C}$ are isomorphic, *then*

$$
\operatorname{ptr}\left(\theta_{X}^{n}\right)=\operatorname{ptr}\left(\theta_{Y}^{n}\right)
$$

holds for each positive integer n*.*

Proof. Suppose $\phi : X \cong Y$ is an isomorphism in C. Then the naturalities of θ and the pivotal structure j imply respectively that

$$
\theta_Y \circ \phi = \phi \circ \theta_X
$$
 and $j_Y \circ \phi = \phi^{**} \circ j_X$.

Thus for any positive integer n , we obtain that

$$
\begin{array}{rcl}\n\text{ptr} \,(\theta_Y^n) & = & \text{ptr} \left(\phi \circ \theta_X^n \circ \phi^{-1} \right) \\
& = & \text{tr} \left(\phi^{**} \circ j_X \circ \theta_X^n \circ \phi^{-1} \right) \\
& = & \text{tr} \left(\phi^{**} \circ j_X \circ \theta_X^n \circ \phi^{-1} \right) \\
& = & \text{tr} \left(j_X \circ \theta_X^n \right) \\
& = & \text{tr} \left(j_X \circ \theta_X^n \right) \\
& = & \text{ptr} \left(\theta_X^n \right),\n\end{array}
$$

where the forth equality is due to [\[EGNO15,](#page-41-10) Proposition 4.7.3(4)].

Moreover, recall in [\[EGNO15,](#page-41-10) Definition 4.7.14] that a pivotal structure j of a rigid monoidal category C is said to be *spherical*, if $tr(j_X) = tr(j_{X^*})$ holds for all $X \in \mathcal{C}$. In fact, it is equivalent to the condition that (right) pivotal traces of any endomorphism and its dual morphisms are equal:

Lemma 2.2. *(*[\[BW99,](#page-40-2) Definition 2.5] *or* [\[Shi12,](#page-42-5) Section 2.1]*)Let* C *be a rigid monoidal category with pivotal structure* j*. Then* j *is spherical if and only if*

$$
ptr(f) =ptr(f^*)
$$
\n(2.1)

holds for each endomorphism $f: X \to X$ *in* C.

2.1. Monoidal equivalences preserving pivotal structure. Since pivotal structures are crucial in defining and studying Frobenius-Schur indicators for monoidal categories, we should pay special attention to monoidal functors said to preserve the pivotal structure, which are introduced in [\[NS07a\]](#page-41-3).

Recall that if $\Phi : \mathcal{C} \to \mathcal{D}$ is a (strong) monoidal functor with structures

$$
J: \Phi(-) \otimes \Phi(-) \cong \Phi(- \otimes -) \quad \text{and} \quad \varphi: \mathbf{1} \cong \Phi(\mathbf{1})
$$

between rigid monoidal categories, then there must be a canonical isomorphism

$$
\chi_X: \Phi(X^*) \xrightarrow{\cong} \Phi(X)^*,\tag{2.2}
$$

which is called the *duality transformation* of Φ, such that the following diagrams both commute for each $X \in \mathcal{C}$:

$$
\Phi(X^*) \otimes \Phi(X) \xrightarrow{\chi_X \otimes id} \Phi(X)^* \otimes \Phi(X)
$$
\n
$$
\Phi(X^* \otimes X)
$$
\n
$$
\Phi(X^* \otimes X)
$$
\n
$$
\Phi(1) \xrightarrow{\varphi^{-1}} \Phi(1)
$$
\n
$$
\Phi(2) \xrightarrow{\varphi^{-1}} \Phi(X) \qquad \text{coev}_{\Phi(X)}
$$
\n
$$
\Phi(X) \otimes \Phi(X)^* \xrightarrow{\chi_{X,X^*}^{-1}} \Phi(X) \otimes \Phi(X^*).
$$
\n(2.3)

By [\[NS07a,](#page-41-3) Lemma 1.1], χ is in fact a monoidal transformation natural in $X \in \mathcal{C}$, and it is unique as

$$
\chi_X = (\Phi(\text{ev}_X) \otimes \text{id}_{\Phi(X)^*}) \circ (\text{id}_{\Phi(X^*)} \otimes \text{coev}_{\Phi(X)}),
$$

where monoidal structures J and φ are omitted.

Notation 2.3. For convenience in this paper, we denote by $\xi : \Phi((-)^{**}) \cong \Phi(-)^{**}$ the *monoidal natural isomorphism from defined as follows:*

$$
\xi_X: \Phi(X^{**}) \xrightarrow{Xx^*} \Phi(X^*)^* \xrightarrow{(\chi_X^*)^{-1}} \Phi(X)^{**},
$$
\n(2.4)

where χ *is the duality transformation* [\(2.2\)](#page-4-1) *of* Φ *.*

It is shown in [\[NS07a\]](#page-41-3) that monoidal functors Φ preserve categorical traces in the following sense, which is described with terms of χ (or ξ). Here we recall this result without identifying $\Phi(1)$ with the unit object:

Lemma 2.4. *([\[NS07a,](#page-41-3) Lemma 6.1]<i>) Let* $\Phi : C \rightarrow D$ *be a monoidal functor between rigid monoidal categories with structures* J and φ : $\mathbf{1} \cong \Phi(\mathbf{1})$ *. Suppose* $f : X \to X^{**}$ *is a morphism in* C*. Then:*

- (1) $\Phi(\text{tr}(f)) = \varphi \circ \text{tr}(\xi_X \circ \Phi(f)) \circ \varphi^{-1} \in \text{End}_{\mathcal{D}}(\Phi(1));$
- (2) *If* C *and* D *are tensor categories over a field* k *and* Φ *is a tensor functor, then*

$$
\operatorname{tr}(f) = \operatorname{tr}(\xi_X \circ \Phi(f)) \in \mathbb{k}
$$

under the usual identifications $\text{End}_{\mathcal{C}}(\mathbf{1}) = \text{End}_{\mathcal{D}}(\mathbf{1}) = \mathbb{k}$.

The notion of monoidal functors preserving the pivotal structure is introduced as follows:

Definition 2.5. *(*[\[NS07a,](#page-41-3) Section 1]*) Let* C *and* D *be rigid monoidal categories with pivotal structures j* and *j'* respectively. A monoidal functor $\Phi : \mathcal{C} \to \mathcal{D}$ is said to preserve the pivotal *structure, if the diagram*

commutes for each $X \in \mathcal{C}$ *.*

Conversely, if the rigid monoidal category $\mathcal C$ is assumed to be pivotal, then a tensor equivalence $\Phi : \mathcal{C} \to \mathcal{D}$ would provide \mathcal{D} a pivotal structure preserved by Φ .

Furthermore, we have the following proposition involving spherical structures:

Proposition 2.6. Let C and D be rigid monoidal categories with pivotal structures j and j' *respectively. Suppose a faithful monoidal functor* $\Phi : \mathcal{C} \to \mathcal{D}$ *preserves the pivotal structure. If* j ′ *is spherical, then* j *is also spherical.*

Proof. Suppose $X \in \mathcal{C}$. In order to prove

$$
tr(j_X) = tr(j_{X^*}) \in End_{\mathcal{C}}(1), \qquad (2.6)
$$

let us compare their images in $\text{End}_{\mathcal{D}}(\Phi(\mathbf{1}))$ under the functor Φ which is faithful.

In fact according to Lemma $2.4(1)$ $2.4(1)$, one could find that

$$
\Phi(\text{tr}(j_X)) = \varphi \circ \text{tr}(\xi_X \circ \Phi(j_X)) \circ \varphi^{-1} = \varphi \circ \text{tr}\left(j'_{\Phi(X)}\right) \circ \varphi^{-1} \in \text{End}_{\mathcal{D}}(1),\tag{2.7}
$$

where ξ_X is defined in [\(2.4\)](#page-5-1) for Φ , and the latter equality is followed by

$$
\xi_X \circ \Phi(j_X) = j'_{\Phi(X)} \in \text{Hom}_{\mathcal{D}}(\Phi(X), \Phi(X)^{**})
$$

as Φ preserves the pivotal structure with the commutativity of (2.5) .

Similarly, another equation is obtained as

$$
\Phi(\text{tr}(j_{X^*})) = \varphi \circ \text{tr}\left(\xi_{X^*} \circ \Phi(j_{X^*})\right) \circ \varphi^{-1} = \varphi \circ \text{tr}\left(j'_{\Phi(X^*)}\right) \circ \varphi^{-1} \in \text{End}_{\mathcal{D}}(1). \tag{2.8}
$$

Furthermore, the fact that j' : Id $_D \cong (-)^{**}$ is a natural isomorphism provides the following commutative diagram in D:

$$
\Phi(X^*) \xrightarrow{j'_{\Phi(X^*)}} \Phi(X^*)^{**}
$$
\n
$$
\chi_X \downarrow \qquad \qquad \chi_X^{**}
$$
\n
$$
\Phi(X)^* \xrightarrow[j'_{\Phi(X)^*}]{} \Phi(X)^{***},
$$

where χ is the duality transformation of Φ . This is equivalent to $j'_{\Phi(X)^*} = (\chi_X^{-1})^{**} \circ j'_{\Phi(X)^*} \circ \chi_X$, and consequently

$$
\Phi(\mathrm{tr}(j_{X^*})) \stackrel{(2.8)}{=} \varphi \circ \mathrm{tr}\left(j'_{\Phi(X^*)}\right) \circ \varphi^{-1} = \varphi \circ \mathrm{tr}\left((\chi_X^{-1})^{**} \circ j'_{\Phi(X)^*} \circ \chi_X\right) \circ \varphi^{-1}
$$

$$
= \varphi \circ \mathrm{tr}\left(j'_{\Phi(X)^*} \circ \chi_X \circ \chi_X^{-1}\right) \circ \varphi^{-1} = \varphi \circ \mathrm{tr}\left(j'_{\Phi(X)^*}\right) \circ \varphi^{-1} \qquad (2.9)
$$

with the usage of [\[EGNO15,](#page-41-10) Proposition 4.7.3(4)].

However, it is clear that $tr(j'_{\Phi(X)}) = tr(j'_{\Phi(X)^*})$ since j' is a spherical structure of \mathcal{D} . Thus we conclude by Equations [\(2.7\)](#page-6-1) and [\(2.9\)](#page-6-2) that $\Phi(\text{tr}(j_X)) = \Phi(\text{tr}(j_{X^*}))$, which implies that $tr(j_X) = tr(j_{X^*})$ because Φ is faithful.

We end this subsection by noting a direct consequence of Lemma [2.4](#page-5-0) for monoidal functors preserving the pivotal structure:

Corollary 2.7. Let C and D be pivotal rigid monoidal categories, and let $\Phi : C \to D$ be a *monoidal functor with structures* J and $\varphi : \mathbf{1} \cong \Phi(\mathbf{1})$ *, which preserves the pivotal structure. Suppose* $f: X \to X^{**}$ *is a morphism in* C. Then:

- (1) $\Phi(\text{ptr}(f)) = \varphi \circ \text{ptr}(\Phi(f)) \circ \varphi^{-1} \in \text{End}_{\mathcal{D}}(\Phi(\mathbf{1}));$
- (2) *If* C *and* D *are tensor categories over a field* k *and* Φ *is a tensor functor, then*

$$
ptr(f) =ptr(\Phi(f)) \in \mathbb{k}
$$
\n(2.10)

under the usual identifications $\text{End}_{\mathcal{C}}(\mathbf{1}) = \text{End}_{\mathcal{D}}(\mathbf{1}) = \Bbbk$.

2.2. Braided monoidal categories with pivotal (or spherical) structures. In this subsection, suppose C is a rigid monoidal category with braiding σ , which is a family of isomorphisms

$$
\sigma^X_Y:X\otimes Y\cong Y\otimes X
$$

natural in $X, Y \in \mathcal{C}$, satisfying the equations

$$
(\mathrm{id}_Y \otimes \sigma_Z^X) \circ (\sigma_Y^X \otimes \mathrm{id}_Z) = \sigma_{Y \otimes Z}^X, \quad (\sigma_Z^X \otimes \mathrm{id}_Y) \circ (\mathrm{id}_X \otimes \sigma_Z^Y) = \sigma_Z^{X \otimes Y} \qquad (\forall X, Y, X \in \mathcal{C}).
$$

The detailed definition and basic properties of *braided monoidal categories* is given in [\[JS93,](#page-41-12) Section 2].

Moreover, when C is rigid, recall in [\[BK01,](#page-40-3) Section 2.2] that the *Drinfeld morphism* u of C and its inverse are defined as follows:

$$
u_X = (\text{ev}_X \otimes \text{id}_{X^*}) \circ (\text{id}_{X^*} \otimes (\sigma_{X^{**}}^X)^{-1}) \circ (\text{coev}_{X^*} \otimes \text{id}_X) \quad : \quad X \to X^{**}, \tag{2.11}
$$

$$
u_X^{-1} = (\mathrm{id}_X \otimes \mathrm{ev}_{X^*}) \circ (\sigma_X^{X^{**}} \otimes \mathrm{id}_{X^*}) \circ (\mathrm{id}_{X^{**}} \otimes \mathrm{coev}_X) \qquad : \quad X^{**} \to X, \tag{2.12}
$$

which are isomorphisms natural in $X \in \mathcal{C}$. Now let us study how a braided monoidal functor transfers the Drinfeld morphisms:

Lemma 2.8. Let C and D be braided rigid monoidal categories. Suppose $\Phi : \mathcal{C} \to \mathcal{D}$ is a *braided monoidal functor. Denote by* u *and* u ′ *respectively the Drinfeld morphisms of* C *and* D*. Then*

$$
u'_{\Phi(X)} = \xi_X \circ \Phi(u_X)
$$

holds for each object $X \in \mathcal{C}$ *, where* ξ *is the monoidal natural isomorphism defined in* (2.4) *.*

Proof. We denote the structures of Φ by $J : \Phi(-) \otimes \Phi(-) \cong \Phi(- \otimes -)$ and $\varphi : \mathbf{1} \cong \Phi(\mathbf{1})$ as usual.

Suppose $X \in \mathcal{C}$. Let us explain at first the following diagram commutes in \mathcal{D} :

The reasons for the commutativity of the cells in this diagram are stated as follows:

- The top and top-right quadrangle: \otimes is a bifunctor on \mathcal{D} ;
- The middle pentagon: The definition (2.3) of the duality transformation χ ;
- The bottom-left trapezoid: The naturality of $J_{X^{**},-}$;
- The bottom-right quadrangle: The monoidal structure axiom for J.

On the other hand, we could obtain the commutativity of another diagram:

where three cells commute due to similar reasons above, except:

• The top quadrangle: The definition of the left dual morphism χ_X^* and the equation

$$
(\mathrm{id}_{\Phi(X^*)}\otimes \mathsf{ev}_{\Phi(X^*)})\circ (\mathsf{coev}_{\Phi(X^*)}\otimes \mathrm{id}_{\Phi(X^*)})=\mathrm{id}_{\Phi(X^*)};
$$

• The left triangle: The definition (2.4) of ξ .

Moreover, since Φ is a braided monoidal functor, the diagram

$$
\Phi(X)^{**} \otimes \Phi(X) \otimes \Phi(X)^{*} \overset{\sigma^{\Phi}(X)^{**}}{\longrightarrow} \Phi(X) \otimes \Phi(X)^{**} \otimes \Phi(X)^{*}
$$
\n
$$
\stackrel{id \otimes id \otimes \chi_X^{-1}}{\longrightarrow} \Phi(X) \otimes \Phi(X)^{**} \otimes id \qquad \qquad \downarrow \text{id} \otimes id \otimes \chi_X^{-1}
$$
\n
$$
\Phi(X)^{**} \otimes \Phi(X) \otimes \Phi(X^*) \overset{\sigma^{\Phi}(X)^{**}}{\longrightarrow} \Phi(X) \otimes \Phi(X)^{**} \otimes \Phi(X^*)
$$
\n
$$
\stackrel{\xi_X \otimes id}{\longrightarrow} \Phi(X) \otimes \Phi(X^*) \overset{\sigma^{\Phi}(X^{**})}{\longrightarrow} \Phi(X) \otimes \Phi(X^*) \otimes id \qquad \qquad \downarrow \text{id} \otimes \xi_X id \otimes id
$$
\n
$$
\Phi(X^{**}) \otimes \Phi(X) \otimes \Phi(X^*) \overset{\sigma^{\Phi}(X^{**})}{\longrightarrow} \Phi(X) \otimes \Phi(X^{**}) \otimes \Phi(X^*)
$$
\n
$$
\stackrel{J_X \times \ast, \chi \otimes id}{\longrightarrow} \Phi(X^* \otimes X) \otimes \Phi(X^*) \overset{\sigma^{\Phi}(X^{**})}{\longrightarrow} \Phi(X \otimes X^{**}) \otimes \Phi(X^*)
$$
\n
$$
\stackrel{J_X \otimes X^{**} \otimes id}{\longrightarrow} \Phi(X^{**} \otimes X) \otimes \Phi(X^*) \overset{J_X \otimes X^{**} \otimes id}{\longrightarrow} \Phi(X^{**} \otimes X \otimes X^*)
$$
\n
$$
\stackrel{\sigma^{\Phi}(X^{**} \otimes id)}{\longrightarrow} \Phi(X^{**} \otimes X \otimes X^*)
$$
\n
$$
(2.15)
$$

commutes as well. Here the braiding on $\mathcal D$ is also denoted by σ without confusions.

As a conclusion of the commuting diagrams (2.13) , (2.14) and (2.15) , we find the equation

$$
\begin{array}{ll} & \left(\operatorname{id}_{\Phi(X)} {\otimes} \mathrm{ev}_{\Phi(X)^*}\right) \circ \left(\sigma^{\Phi(X)^{**}}_{\Phi(X)} \otimes \operatorname{id}_{\Phi(X)^*}\right) \circ \left(\operatorname{id}_{\Phi(X)^{**}} {\otimes} \mathrm{coev}_{\Phi(X)}\right) \\ \\ = & \Phi(\operatorname{id}_{X^{**}} {\otimes} \mathrm{ev}_{X^*}) \circ \Phi(\sigma^{X^{**}}_X \otimes \operatorname{id}_{X^*}) \circ \Phi(\operatorname{id}_{X^{**}} {\otimes} \mathrm{coev}_X) \circ \xi_X^{-1} \\ \\ = & \Phi\Big((\operatorname{id}_{X^{**}} {\otimes} \mathrm{ev}_{X^*}) \circ (\sigma^{X^{**}}_X \otimes \operatorname{id}_{X^*}) \circ (\operatorname{id}_{X^{**}} {\otimes} \mathrm{coev}_X)\Big) \circ \xi_X^{-1}, \end{array}
$$

which is equivalent to $(u'_{\Phi(X)})^{-1} = \Phi(u_X^{-1}) \circ \xi_X^{-1}$ according to the definition of the Drinfeld morphism. Thus our desired equation is followed. \square 10 K. LI

Now suppose further that the braided rigid monoidal category $\mathcal C$ has a spherical structure j. Then the *ribbon structure* θ on $\mathcal C$ is the natural automorphism defined as follows:

$$
\theta_X = u_X^{-1} \circ j_X : X \cong X \quad (\forall X \in \mathcal{C}).
$$

It is stated in [\[NS07b,](#page-42-1) Section 2] that a braided monoidal functor preserving the pivotal structure would also preserve the ribbon structure. We recall this claim as a consequence of Lemma [2.8:](#page-7-2)

Corollary 2.9. *(cf.* [\[NS07b,](#page-42-1) Section 2]*) Let* C *and* D *be braided monoidal categories, with spherical structures j* and *j'* respectively. Suppose $\Phi : \mathcal{C} \to \mathcal{D}$ is a braided monoidal functor *preserving the pivotal structure. Denote by* θ *and* θ ′ *respectively the ribbon structures of* C *and* D*. Then*

$$
\theta'_{\Phi(X)} = \Phi(\theta_X)
$$

holds for each object $X \in \mathcal{C}$ *.*

Proof. One might find by the definition of the ribbon structures as well as Lemma [2.8](#page-7-2) that

$$
\begin{array}{rcl}\n\theta'_{\Phi(X)} & = & u'_{\Phi(X)}^{-1} \circ j'_{\Phi(X)} = \Phi(u_X)^{-1} \circ \xi_X^{-1} \circ j'_{\Phi(X)} \stackrel{(2.5)}{=} \Phi(u_X)^{-1} \circ \Phi(j_X) \\
& = & \Phi(u_X^{-1} \circ j_X) = \Phi(\theta_X)\n\end{array}
$$

holds for any $X \in \mathcal{C}$.

2.3. Module categories and dual tensor categories. The definition of (left) module categories over monoidal categories could be found in [\[EGNO15,](#page-41-10) Sections 7.1 to 7.3]. In this subsection, we recall some elementary properties of module categories over a finite tensor category $\mathcal C$ over an algebraically closed field $\mathbb k$, as well as the corresponding dual categories in the literature.

Lemma 2.10. *Let* M *be a module category over a finite tensor category* C*.*

- (1) *(*[\[EO04,](#page-41-13) Lemma 3.4]*) If* M *is indecomposable and exact as a* C*-module category, then* M *is finite.*
- (2) *(*[\[EGNO15,](#page-41-10) Section 7.5]*) If* M *is semisimple, then* M *is exact;*

Remark 2.11. *Note in* [\[EO04,](#page-41-13) Section 3.1] *that an exact module category* M *over a finite tensor category* C *is defined to have only finitely many isoclasses of simple objects.*

Suppose the left C-module category M is exact. The *dual tensor category* of C with respect to M is defined as

$$
\mathcal{C}_\mathcal{M}^*:=\mathrm{Rex}_\mathcal{C}(\mathcal{M})^\mathrm{rev},
$$

the category of k-linear right exact C-module endofunctors on \mathcal{M} . We should remark that the tensor product on $\mathcal{C}_{\mathcal{M}}^*$ in this paper is chosen to be opposite to the composition of $\mathcal{C}\text{-module}$ endofunctors, which would differ with [\[EO04\]](#page-41-13) and [\[EGNO15\]](#page-41-10), etc. Denote each object in $\mathcal{C}_{\mathcal{M}}^*$ by a pair $\mathbf{F} = (F, s)$, where s is a C-module structure of k-linear endofunctor $F : \mathcal{M} \to \mathcal{M}$ with notation

$$
s_{X,M}: X \otimes F(M) \cong F(X \otimes M) \qquad (\forall X \in \mathcal{C}, M \in \mathcal{M}).
$$

where $\mathcal{D}: \mathcal{C} \times \mathcal{M} \to \mathcal{M}$ denotes the module product functor.

Lemma 2.12. *(*[\[EO04,](#page-41-13) Section 3.3]*) Suppose that* M *is an exact module category over a* $finite$ tensor category C . Then $C^*_{\mathcal{M}}$ is a finite multitensor category, which would be a tensor *category if* M *is indecomposable as a* C*-module category.*

In particular, we would pay more attention to the case when $\mathcal C$ and $\mathcal M$ are both semisimple. Note at first that when C is a fusion category, it is remarked in $[EO04, Example 3.3(iii)]$ that M is exact if and only if it is semisimple.

Lemma 2.13. *Let* C *be a fusion category over an algebraically closed field* k *of characteristic* 0*. Suppose* M *is an indecomposable* C*-module category which is semisimple (or exact). Then:*

- (1) *(*[\[EO04,](#page-41-13) Section 3.3]*)* M *is finite as a* k*-linear abelian category;*
- (2) $([ENO05, Theorem 2.15])$ $([ENO05, Theorem 2.15])$ $([ENO05, Theorem 2.15])$ $\mathcal{C}_{\mathcal{M}}^{*}$ *is a fusion category.*

2.4. The centers as rigid monoidal categories and Schauenburg's equivalence. For any monoidal category C , it is well-known that its (left) *center* $\mathcal{Z}(C)$ (see [\[JS91\]](#page-41-15) e.g.) consists of objects $V = (V, \sigma^V)$, where $V \in \mathcal{C}$ and the *half-braiding* σ^V is a family of isomorphisms

$$
\sigma^V_X: V \otimes X \cong X \otimes V
$$

natural in $X \in \mathcal{C}$, such that $(\sigma_X^V \otimes id_Y) \circ (id_X \otimes \sigma_Y^V) = \sigma_{X \otimes Y}^V$ holds for all $X, Y \in \mathcal{C}$.

Also, $\mathcal{Z}(\mathcal{C})$ is a braided monoidal category with the braiding defined by

$$
\sigma_W^V: V \otimes W \cong W \otimes V
$$

for any $\mathbf{V} = (V, \sigma^V)$ and $\mathbf{W} = (W, \sigma^W)$ in $\mathcal{Z}(\mathcal{C})$. Moreover, when C is rigid, recall that $\mathcal{Z}(\mathcal{C})$ becomes rigid as well, and each object $\mathbf{V} = (V, \sigma^V)$. has left and right duals:

$$
\mathbf{V}^* = (V^*, (\sigma^V_{*(-)})^*) \text{ and } ^* \mathbf{V} = (*V, ^* (\sigma^V_{(-)^*})).
$$

Moreover, it is clear that the Drinfeld morphism u of $\mathcal{Z}(\mathcal{C})$ and its inverse are determined as follows:

$$
u_{\mathbf{V}} = (\text{ev}_V \otimes \text{id}_{V^*}) \circ (\text{id}_{V^*} \otimes (\sigma_{V^{**}}^V)^{-1}) \circ (\text{coev}_{V^*} \otimes \text{id}_V) : V \to V^{**},
$$

$$
(u_{\mathbf{V}})^{-1} = (\text{id}_V \otimes \text{ev}_{V^*}) \circ (\sigma_V^{V^{**}} \otimes \text{id}_{V^*}) \circ (\text{id}_{V^{**}} \otimes \text{coev}_V) : V^{**} \to V,
$$

for any object $\mathbf{V} = (V, \sigma^V) \in \mathcal{Z}(\mathcal{C})$.

Now suppose that C is a finite tensor category over a field k, and M is a finite (left) \mathcal{C} -module category. Then by [\[Sch01,](#page-42-7) Theorem 3.3], there is an equivalence $\Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ of k-linear braided monoidal categories, which is referred as *Schauenburg's equivalence*.

We recall the details of the functor Ω according to [\[Shi20,](#page-42-6) Section 3]: For each object $\mathbf{V} = (V, \sigma^V) \in \mathcal{Z}(\mathcal{C}), \text{ set } \Omega(\mathbf{V}) = (V \otimes -, \varsigma^{V \otimes -}) \in \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*), \text{ where the left } \mathcal{C}\text{-module structure}$ of $V \otimes -$ is defined as the collection

$$
(\sigma_X^V)^{-1} \otimes \mathrm{id}_M : X \otimes V \otimes M \cong V \otimes X \otimes M \qquad (X \in \mathcal{C}, M \in \mathcal{M}).
$$

The half-braiding $\zeta^{V\otimes -}$ of $\Omega(V)$ is given as follows: For each $F \in C^*_{\mathcal{M}}$, denote its C-module structure by $s : - \otimes F(-) \cong F(- \otimes -)$, then let

$$
(\varsigma_F^{V\otimes -})_M = s_{V,M}^{-1} : (\mathbf{F} \circ \Omega(\mathbf{V}))(M) = F(V \otimes M) \xrightarrow{\cong} V \otimes F(M) = (\Omega(\mathbf{V}) \circ \mathbf{F})(M) \tag{2.16}
$$

for $M \in \mathcal{M}$, and hence we write $\varsigma_F^{V \otimes -} = s_{V, -}^{-1}$. Also, one could verify that the monoidal structure of Ω is given by:

$$
\sigma_V^W \otimes \mathrm{id}_{(-)}: \ \Omega({\mathbf{W}}) \circ \Omega({\mathbf{V}}) = W \otimes V \otimes (-) \ \xrightarrow{\cong} \ V \otimes W \otimes (-) = \Omega({\mathbf{V}} \otimes {\mathbf{W}})
$$

for any objects $\mathbf{V} = (V, \sigma^V)$ and $\mathbf{W} = (W, \sigma^W)$ in $\mathcal{Z}(\mathcal{C})$.

However for the remaining of this paper, we only focus on the case when the C -module category M is indecomposable, and consequently $\mathcal{C}_{\mathcal{M}}^*$ is a tensor category by Lemma [2.12.](#page-9-1) In this case, Schauenburg's equivalence Ω might be noted again as:

Lemma 2.14. *(cf.* [\[Shi20,](#page-42-6) Theorem 3.13]*) Let* C *be a finite tensor category, and let* M *be an exact indecomposable* C-module category. Then Ω : $\mathcal{Z}(\mathcal{C}) \to \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ *is an equivalence of braided finite tensor categories.*

For the remaining of this subsection, we describe the spherical structure and ribbon structure on $\mathcal{Z}(\mathcal{C}^*_\mathcal{M})$ when $\mathcal C$ is moreover spherical, such that Schauenburg's equivalence Ω preserves the pivotal structure.

Let us denote the spherical structure on a finite tensor category $\mathcal C$ by j, and let $\mathcal M$ be an exact indecomposable C -module category as usual. Clearly, it follows at first that the center $\mathcal{Z}(\mathcal{C})$ has a spherical structure ([Müg03, Proposition 3.9] e.g.):

$$
j_{\mathbf{V}} : \mathbf{V} = (V, \sigma) \stackrel{j_V}{\cong} (V^{**}, (\sigma_{**}{}_{(-)})^{**}) = \mathbf{V}^{**} \in \mathcal{Z}(\mathcal{C}), \tag{2.17}
$$

for each $\mathbf{V} = (V, \sigma) \in \mathcal{Z}(\mathcal{C})$. Consequently, the ribbon structure θ of the braided spherical tensor category $\mathcal{Z}(\mathcal{C})$ is the family of natural automorphism $\theta_{\mathbf{V}} = u_{\mathbf{V}}^{-1} \circ j_{\mathbf{V}}$ $(\mathbf{V} \in \mathcal{Z}(\mathcal{C}))$, and it also satisfies

$$
(\theta_{\mathbf{V}})^* = \theta_{\mathbf{V}^*} \quad (\forall \mathbf{V} \in \mathcal{Z}(\mathcal{C})) \tag{2.18}
$$

according to [\[EGNO15,](#page-41-10) Proposition 8.10.12].

We end this section by applying Corollary [2.9](#page-9-2) to Schauenburg's equivalence $\Omega : \mathcal{Z}(\mathcal{C}) \approx$ $\mathcal{Z}(\mathcal{C}^*_\mathcal{M})$:

Corollary 2.15. *Let* C *be a finite tensor category with spherical structure* j*, and let* M *be a finite* C-module category. Denote the ribbon structure of $\mathcal{Z}(\mathcal{C})$ by $\theta = u^{-1}j$. If $\mathcal{C}_{\mathcal{M}}^*$ is pivotal *and* Ω *preserves the pivotal structure, then*

$$
\Theta_{\Omega(\mathbf{V})} = \Omega(\theta_{\mathbf{V}}) = \theta_{\mathbf{V}} \otimes \mathrm{id}_{(-)}
$$
\n(2.19)

for each $V \in \mathcal{Z}(\mathcal{C})$, where Θ *is the ribbon structure of* $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$.

Proof. This is a direct consequence of Corollary [2.9.](#page-9-2)

3. Frobenius-Schur indicators of the dual category to spherical fusion **CATEGORIES**

3.1. Definition and compatibility. Let $\mathcal C$ be a linear pivotal monoidal category with pivotal structure j over a field. The n-th Frobenius-Schur indicators ν_n for C is defined in [\[NS07b,](#page-42-1) Definition 3.1] as follows: For any object $X \in \mathcal{C}$, define the linear transformation

$$
\begin{array}{rcl}\nE_X^{(n)} & : & \operatorname{Hom}\nolimits_{\operatorname{\mathcal C}\nolimits}(\mathbf{1},X^{\otimes n}) & \to & \operatorname{Hom}\nolimits_{\operatorname{\mathcal C}\nolimits}(\mathbf{1},X^{\otimes n}), \\
f & \mapsto & (\operatorname{ev}\nolimits_X \otimes \operatorname{id}\nolimits_X^{\otimes (n-1)} \otimes j_X^{-1}) \circ (\operatorname{id}\nolimits_{X^*} \otimes f \otimes \operatorname{id}\nolimits_{X^{**}}) \circ \operatorname{coev}\nolimits_{X^*},\n\end{array}
$$

where n is a positive integer. Then the n-th *Frobenius-Schur indicator* of V is the scalar

$$
\nu_n(X) := \text{Tr}\left(E_X^{(n)}\right) \quad (n \ge 1),\tag{3.1}
$$

which denotes the trace of the linear transformation $E_X^{(n)}$ on the finite-dimensional vector space $\text{Hom}_{\mathcal{C}}(\mathbf{1}, X^{\otimes n})$.

For the remaining of this paper, we always assume that $\mathcal C$ is furthermore spherical and fusion over the complex field C. Note that the left and right pivotal traces of each endomorphism in C are identical, which are regarded as complex numbers since End_C(1) $\cong \mathbb{C}$ in this case.

$$
\Box
$$

Besides, the *categorical dimension* of C is defined as

$$
\dim(\mathcal{C}) = \sum_{X \in \mathcal{O}(\mathcal{C})} \text{ptr}(\text{id}_X) \text{ptr}(\text{id}_{X^*}),\tag{3.2}
$$

where $\mathcal{O}(\mathcal{C})$ denotes the set of isoclasses of simple objects in C. It was proved in [\[NS07b,](#page-42-1) Theorem 4.1 that indicators could be computed through the ribbon structure θ of the center $\mathcal{Z}(\mathcal{C})$, which is also a fusion category:

Lemma 3.1. *(NS07b, Theorem 4.1)* Let C be any spherical fusion category over \mathbb{C} with *spherical structure j. Suppose* u *is the Drinfeld morphism of* $\mathcal{Z}(\mathcal{C})$ *. For any* $V \in \mathcal{C}$ *,*

$$
\nu_n(V) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta_{I(V)}^n\right) \qquad (\forall n \ge 1),\tag{3.3}
$$

where $\theta = u^{-1}j$ *is the ribbon structure of* $\mathcal{Z}(\mathcal{C})$ *, and I is a two-sided adjoint to the forgetful functor* $\mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ *.*

Remark 3.2. As noted in [\[NS07b,](#page-42-1) Section 4], the adjoint $I: \mathcal{C} \rightarrow \mathcal{Z}(\mathcal{C})$ to the forgetful *functor is supposed to be chosen by a result of* [Müg03, Proposition 8.1] *such that*

$$
I(X) \cong \bigoplus_{(V,\sigma^V)\in \mathcal{O}(\mathcal{Z}(\mathcal{C}))} \dim(\text{Hom}_{\mathcal{C}}(V,X))\cdot (V,\sigma^V),
$$

where $\mathcal{O}(\mathcal{Z}(\mathcal{C}))$ *denotes the set of isoclasses of simple objects in* $\mathcal{Z}(\mathcal{C})$ *. We would explain in the next paragraph that* I *could be replaced by an arbitrary (left or right) adjoint to the forgetful functor.*

In fact as a fusion category, C must be unimodular ([\[ENO04,](#page-41-17) Corollary 6.4]), which means that the distinguished invertible object is the unit object 1. Thus according to [\[Shi17a,](#page-42-2) Theorem 4.10], the forgetful functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ is Frobenius because the base field $\mathbb C$ is perfect, which follows that its left and right adjoints are naturally isomorphic. We conclude by Lemma [2.1](#page-4-2) that the functor I in (3.3) could be chosen as an arbitrary adjoint of the forgetful functor.

In order to study the indicators of objects in the dual category $\mathcal{C}_{\mathcal{M}}^*$, we attempt to introduce an analogous formula, which is supposed to be regarded as a definition. This is because that we do not know whether $\mathcal{C}_{\mathcal{M}}^*$ is naturally pivotal, even though $\mathcal C$ is a spherical fusion category.

Definition 3.3. Let C be a spherical fusion category over C with spherical structure j, and let M be an indecomposable C-module category which is semisimple. Denote by $\mathbf{U}_\mathcal{M} : \mathcal{Z}(\mathcal{C}_\mathcal{M}^*) \to$ $\mathcal{C}_{\mathcal{M}}^*$ the forgetful functor, and suppose u is the Drinfeld morphism of $\mathcal{Z}(\mathcal{C})$. For any $\mathbf{F} \in \mathcal{C}_{\mathcal{M}}^*$, *the* n*-th Frobenius-Schur indicator of* F *is the defined as the scalar*

$$
\nu_n(\mathbf{F}) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta_{K(\mathbf{F})}^n\right) \in \mathbb{C} \qquad (\forall n \ge 1), \tag{3.4}
$$

where $\theta = u^{-1}j$ is the ribbon structure of $\mathcal{Z}(\mathcal{C})$, and K is an arbitrary (left or right) adjoint *to* $\mathbf{U}_{\mathcal{M}} \circ \Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}_{\mathcal{M}}^*$.

Similarly, we should explain the scalar ptr $(\theta_{K(\mathbf{F})}^n)$ is independent on the choice of the functor $K: \mathcal{C}^*_{\mathcal{M}} \to \mathcal{Z}(\mathcal{C})$ as a left or right adjoint to $\mathbf{U}_{\mathcal{M}} \circ \Omega$:

Lemma 3.4. *Let* C*,* M *and* θ *be as in Definition [3.3.](#page-12-1) Suppose* K *and* K′ *are left or right adjoints to the functor* $\mathbf{U}_{\mathcal{M}} \circ \Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}_{\mathcal{M}}^*$. Then

$$
\text{ptr}\big(\theta^n_{K(\mathbf{F})}\big) = \text{ptr}\big(\theta^n_{K'(\mathbf{F})}\big) \qquad (\forall n \ge 1)
$$

holds for each $\mathbf{F} \in C_{\mathcal{M}}^*$.

Proof. At first $C^*_{\mathcal{M}}$ is a fusion category by [\[ENO05,](#page-41-14) Theorem 2.15], as the C-module category ${\mathcal M}$ is assumed to be indecomposable and semisimple.

Since $\Omega: \mathcal{Z}(\mathcal{C}) \to \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ is a tensor equivalence, $\Omega \circ K$ and $\Omega \circ K'$ are both (left or right) adjoint functors to the forgetful functor $U_{\mathcal{M}}: \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \to \mathcal{C}_{\mathcal{M}}^*$, and hence they are naturally isomorphic according to [\[Shi17a,](#page-42-2) Theorem 4.10] as well. Consequently, we could know that $K(\mathbf{F}) \cong K'(\mathbf{F})$ holds for each $\mathbf{F} \in C^*_{\mathcal{M}}$. Our desired equality could be obtained with the help of Lemma [2.1.](#page-4-2)

For later use, we usually choose the adjoint functor K in the view of Lemma $4.5(1)$ $4.5(1)$ as

$$
\mathcal{Z}(\rho^{\text{ra}}): \mathcal{C}_{\mathcal{M}}^* \to \mathcal{Z}(\mathcal{C}), \ \ \mathbf{F} = (F, s) \mapsto \left(\int_{M \in \mathcal{M}} \underline{\text{Hom}}(M, F(M)), \sigma^{\mathbf{F}} \right), \tag{3.5}
$$

where the detailed constructions would be recalled in Subsections [4.1](#page-18-1) and [4.2.](#page-19-0) We also note the following fact, which is a direct consequence of Lemma [2.1](#page-4-2) as well:

Remark 3.5. *Isomorphic objects in* $\mathcal{C}_{\mathcal{M}}^*$ *always have the same n-th indicators in the sense of Definition [3.3.](#page-12-1)*

Now recall again from [\[ENO05,](#page-41-14) Theorem 2.15] that if M is a semisimple and indecomposable module category over a complex fusion category \mathcal{C} , then $\mathcal{C}_{\mathcal{M}}^{*}$ is also a fusion category with

$$
\dim(\mathcal{C}_{\mathcal{M}}^*) = \dim(\mathcal{C}).\tag{3.6}
$$

Thus we might formulate indicators of objects in $\mathcal{C}_{\mathcal{M}}^*$ with the ribbon structure Θ of its center $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ similarly to [\(3.3\)](#page-12-0), as long as $\mathcal{C}_{\mathcal{M}}^*$ already has a pivotal structure such that Schauenburg's equivalence Ω preserves the pivotal structure:

Proposition 3.6. Let C be any spherical fusion category over $\mathbb C$ with spherical structure j, and let M be an indecomposable C -module category which is semisimple. Suppose $C^*_{\mathcal{M}}$ is also *spherical and* Ω : $\mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ preserves the pivotal structure. Then for any $\mathbf{F} = (F, s) \in$ $\mathcal{C}_{\mathcal{M}}^{*}$, *its n*-th indicator in the sense of Definition [3.3](#page-12-1) is

$$
\nu_n(\mathbf{F}) = \frac{1}{\dim(\mathcal{C}_{\mathcal{M}}^*)} \text{ptr}\left(\Theta_{I_{\mathcal{M}}(\mathbf{F})}^n\right),\tag{3.7}
$$

where Θ *is the ribbon structure of* $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^{*})$ *, and* $I_{\mathcal{M}}$ *is an adjoint of the forgetful functor* $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^{*}) \to \mathcal{C}_{\mathcal{M}}^{*}$. In other words, the indicators of any $\mathbf{F} \in \mathcal{C}_{\mathcal{M}}^{*}$ in the senses of [\(3.3\)](#page-12-0) and [\(3.4\)](#page-12-2) *coincide.*

Proof. Since Ω : $\mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ preserves the pivotal structure, we obtain the equation

$$
\xi \mathbf{v} \circ \Omega(j\mathbf{v}) = j'_{\Omega(\mathbf{V})} \qquad (\forall \mathbf{V} \in \mathcal{Z}(\mathcal{C})) \tag{3.8}
$$

according to Definition [2.5,](#page-5-4) where ξ [\(2.4\)](#page-5-1) is determined by the duality transformation of Ω as usual.

Now with usage of Lemma [2.4\(](#page-5-0)2), we compute for any $n \geq 1$ that

$$
\text{ptr}\left(\theta_{K(\mathbf{F})}^n\right) \stackrel{(2.10)}{=} \text{ptr}\left(\Omega(\theta_{K(\mathbf{F})}^n)\right) = \text{ptr}\left(\Omega(\theta_{K(\mathbf{F})})^n\right) \stackrel{(2.19)}{=} \text{ptr}\left(\Theta_{\Omega(K(\mathbf{F}))}^n\right).
$$

However, it is clear that $\Omega \circ K$ is also an adjoint to to the forgetful functor $U_{\mathcal{M}} : \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \to \mathcal{C}_{\mathcal{M}}^*$. The same argument with Lemma [3.4](#page-12-3) would imply that

$$
\operatorname{ptr}\left(\Theta^n_{\Omega(K(\mathbf{F}))}\right)=\operatorname{ptr}\left(\Theta^n_{I_{\mathcal{M}}(\mathbf{F})}\right).
$$

As a conclusion, we find by Equation (3.6) that

$$
\frac{1}{\dim(\mathcal{C})}\text{ptr}\left(\theta_{K(\mathbf{F})}^{n}\right) = \frac{1}{\dim(\mathcal{C}_{\mathcal{M}}^{*})}\text{ptr}\left(\Theta_{I_{\mathcal{M}}(\mathbf{F})}^{n}\right)
$$
as complex numbers for each $n \geq 1$.

3.2. Invariance of adjoints in the dual category. In this subsection, let us prove that indicators in the sense of Definition [3.3](#page-12-1) are invariant under the dualities on objects in $\mathcal{C}_{\mathcal{M}}^*$.

Before that, we recall some descriptions in [\[Shi20\]](#page-42-6) on adjoint pairs between dual categories and centers: Assume that M is a finite left module category over a finite tensor category C . Define the *action functor*

$$
\rho: \mathcal{C} \to \text{Rex}(\mathcal{M}), \ \ X \mapsto X \otimes -,
$$

and note that ρ is an exact monoidal functor. Thus it has both left adjoint ρ^{la} and right adjoint ρ^{ra} . Moreover, they could be "lifted" to adjoint pairs between the centers:

Lemma 3.7. *(*[\[Shi20,](#page-42-6) Section 3.6]*) Suppose that* M *is exact as a* C*-module category. Then there are are adjoint pairs*

$$
\left(\mathcal{Z}(\rho^{\text{la}}): \text{Rex}_{\mathcal{C}}(\mathcal{M}) \to \mathcal{Z}(\mathcal{C}), \quad \mathcal{Z}(\rho): \mathcal{Z}(\mathcal{C}) \to \text{Rex}_{\mathcal{C}}(\mathcal{M})\right)
$$
(3.9)

and

$$
\left(\mathcal{Z}(\rho):\mathcal{Z}(\mathcal{C})\to\text{Rex}_{\mathcal{C}}(\mathcal{M}),\quad\mathcal{Z}(\rho^{\text{ra}}):\text{Rex}_{\mathcal{C}}(\mathcal{M})\to\mathcal{Z}(\mathcal{C})\right).
$$
(3.10)

Proof. We omit here the detailed constructions of functors $\mathcal{Z}(\rho)$, $\mathcal{Z}(\rho^{\text{la}})$ and $\mathcal{Z}(\rho^{\text{ra}})$, but note that the latter adjunction desired is [\[Shi20,](#page-42-6) Theorem 3.11], while the former one could also be obtained by applying the 2-functor $\mathcal Z$ to ρ and its left adjoint ρ^{la} . Details are founded in $[\text{Shi20, Section 3.6}].$

Note again that in this paper, the tensor product bifunctor on the dual tensor category $\mathcal{C}^*_{\mathcal{M}} := \text{Rex}_{\mathcal{C}}(\mathcal{M})^{\text{rev}}$ is chosen as

$$
(\mathbf{F},\mathbf{G})\mapsto \mathbf{G}\circ \mathbf{F},
$$

which is opposite to the composition of C -module endofunctors. Thus the left and right duals of object $\mathbf{F} \in \mathcal{C}_{\mathcal{M}}^*$ would be its left and right adjoint functors, denoted by \mathbf{F}^{la} and \mathbf{F}^{ra} respectively. Now we show that their indicators are equal as long as well-defined:

Proposition 3.8. *Let* C *be a spherical fusion category over* C*, and let* M *be an indecomposable C*-module category which is semisimple. Then for any $\mathbf{F} \in C_{\mathcal{M}}^*$,

$$
\nu_n(\mathbf{F}^{\text{la}}) = \nu_n(\mathbf{F}) = \nu_n(\mathbf{F}^{\text{ra}}) \qquad (\forall n \ge 1).
$$

Proof. At first by a fact stated in [\[Ost03,](#page-42-8) Section 2.2], our assumptions imply that the left dual \mathbf{F}^{la} and right dual \mathbf{F}^{ra} are isomorphic as objects in the fusion category $\mathcal{C}_{\mathcal{M}}^*$. Thus $\nu_n(\mathbf{F}^{\text{la}})$ = $\nu_n(\mathbf{F}^{\text{ra}})$ holds for all $n \geq 1$ due to Remark [3.5](#page-13-1) and it suffices to show $\nu_n(\mathbf{F}^{\text{la}}) = \nu_n(\mathbf{F})$.

Now with the same argument as the proof of $[\text{Shi20}, \text{Equation (3.2)}]$, we try to show that

$$
^{\ast}\mathcal{Z}(\rho^{\mathrm{la}})(\mathbf{F}^{\mathrm{la}}) \cong \mathcal{Z}(\rho^{\mathrm{ra}})(\mathbf{F})
$$
\n(3.11)

as objects in $\mathcal{Z}(\mathcal{C})$, where the adjoint functors $\mathcal{Z}(\rho^{\text{la}})$ and $\mathcal{Z}(\rho^{\text{rad}})$ are denoted in Lemma [3.7.](#page-14-1)

Indeed, the duality adjunctions follow the isomorphism

$$
\operatorname{Hom}_{\mathcal{Z}(\mathcal{C})}\left(\mathbf{V},{}^*\mathcal{Z}(\rho^{\mathrm{la}})(\mathbf{F}^{\mathrm{la}})\right) \quad \cong \quad \operatorname{Hom}_{\mathcal{Z}(\mathcal{C})}\left(\mathcal{Z}(\rho^{\mathrm{la}})(\mathbf{F}^{\mathrm{la}}),\mathbf{V}^*\right)
$$

 16 K. LI

 $\stackrel{(3.9)}{\cong}$ $\stackrel{(3.9)}{\cong}$ $\stackrel{(3.9)}{\cong}$ \cong Nat_{Rexc}(M) $(\mathbf{F}^{\text{la}}, \mathcal{Z}(\rho)(\mathbf{V}^*))$,

natural in $\mathbf{V} \in \mathcal{Z}(\mathcal{C})$. Moreover, one could verify that the functor $\mathcal{Z}(\rho)$ is (strong) monoidal, since ρ is so. Therefore we obtain isomorphisms $\mathcal{Z}(\rho)(\mathbf{V}^*) \cong \mathcal{Z}(\rho)(\mathbf{V})^{\text{la}}$ and

$$
\begin{array}{rcl}\n\operatorname{Nat}_{\operatorname{Rexc}(\mathcal{M})}(\mathbf{F}^{\mathrm{la}}, \mathcal{Z}(\rho)(\mathbf{V}^*)) & \cong & \operatorname{Nat}_{\operatorname{Rexc}(\mathcal{M})}(\mathbf{F}^{\mathrm{la}}, \mathcal{Z}(\rho)(\mathbf{V})^{\mathrm{la}}) \\
\\ \cong & \operatorname{Nat}_{\operatorname{Rexc}(\mathcal{M})}(\mathcal{Z}(\rho)(\mathbf{V}), \mathbf{F}) \stackrel{(3.10)}{\cong} \operatorname{Hom}_{\mathcal{Z}(\mathcal{C})}(\mathbf{V}, \mathcal{Z}(\rho^{\mathrm{ra}})(\mathbf{F})),\n\end{array}
$$

which are natural in $V \in \mathcal{Z}(\mathcal{C})$. Thus according to Yoneda Lemma, the natural isomorphism $\text{Hom}_{\mathcal{Z}(\mathcal{C})}(\mathbf{V},^*\mathcal{Z}(\rho^{\text{la}})(\mathbf{F}^{\text{la}})) \cong \text{Hom}_{\mathcal{Z}(\mathcal{C})}(\mathbf{V}, \mathcal{Z}(\rho^{\text{rad}})(\mathbf{F}))$ concluded above implies [\(3.11\)](#page-14-4), which follows also that

$$
\mathcal{Z}(\rho^{\text{la}})(\mathbf{F}^{\text{la}}) \cong \mathcal{Z}(\rho^{\text{ra}})(\mathbf{F})^*
$$
\n(3.12)

On the other hand, more adjunctions could be obtained by the adjoint pairs

$$
\left(\mathcal{Z}(\rho^{\text{la}}),\mathcal{Z}(\rho)\right),\quad \left(\mathcal{Z}(\rho),\mathcal{Z}(\rho^{\text{ra}})\right)\quad\text{and}\quad \left(\mathbf{U}_{\mathcal{M}}\circ\Omega,\mathcal{Z}(\rho^{\text{ra}})\right),
$$

according to [\[MacL98,](#page-41-18) Theorem 1 in Section IV.8]: Both $\mathcal{Z}(\rho) \circ \Omega^{-1}$ and the forgetful functor $U_{\mathcal{M}}$ are both left adjoints of $\Omega \circ \mathcal{Z}(\rho^{ra})$, and hence they would be naturally isomorphic. Therefore, as the left and right adjoints of the forgetful functor $U_{\mathcal{M}}$ are naturally isomorphic in this case (by [\[Shi17a,](#page-42-2) Theorem 4.10]), we find that $\Omega \circ \mathcal{Z}(\rho^{\text{la}})$ and $\Omega \circ \mathcal{Z}(\rho^{\text{ra}})$ are both adjoints to $U_{\mathcal{M}}$. It follows that

$$
\mathcal{Z}(\rho^{\text{la}})
$$
 and $\mathcal{Z}(\rho^{\text{ra}})$

are both (left or right) adjoints to $\mathbf{U}_{\mathcal{M}} \circ \Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}_{\mathcal{M}}^*$.

Finally, since the definition of ribbon structure provides Equation [\(2.18\)](#page-11-3) that $\theta_{V^*} = (\theta_V)^*$ for any $\mathbf{V} \in \mathcal{Z}(\mathcal{C})$, we find that

$$
\text{ptr}\left(\theta_{\mathbf{V}^*}^n\right) = \text{ptr}\left(\left(\theta_{\mathbf{V}}^n\right)^*\right) \stackrel{(2.1)}{=} \text{ptr}\left(\theta_{\mathbf{V}}^n\right) \quad (\forall n \ge 1)
$$
\n(3.13)

always holds in the spherical braided category $\mathcal{Z}(\mathcal{C})$. Thus for any $n \geq 1$, we could compute by Lemma [2.1](#page-4-2) that

$$
\operatorname{ptr}\left(\theta^{n}_{\mathcal{Z}(\rho^{\text{la}})(\mathbf{F}^{\text{la}})}\right)\stackrel{(3.12)}{=}\operatorname{ptr}\left(\theta^{n}_{\mathcal{Z}(\rho^{\text{ra}})(\mathbf{F})^{*}}\right)\stackrel{(3.13)}{=}\operatorname{ptr}\left(\theta^{n}_{\mathcal{Z}(\rho^{\text{ra}})(\mathbf{F})}\right).
$$

Consequently, the desired invariance property $\nu_n(\mathbf{F}^{la}) = \nu_n(\mathbf{F})$ is obtained due to Definition [3.3,](#page-12-1) which is independent of the choices of adjoint functors $\mathcal{Z}(\rho^{\text{la}})$ or $\mathcal{Z}(\rho^{\text{ra}})$.

3.3. Invariance under tensor equivalences. In this subsection, let us check in brief the invariance of indicators in $\mathcal{C}_{\mathcal{M}}^*$ under certain tensor equivalences between $\mathcal C$ and other spherical fusion categories. Suppose that $\Phi : \mathcal{C} \to \mathcal{D}$ is a tensor functor and M is an exact D-module category via bifunctor \odot . Recall in [\[EO04,](#page-41-13) Section 3.5] that M becomes naturally a C-module category via Φ as follows:

- The module product functor is $C \times \mathcal{M} \to \mathcal{M}$, $(X, M) \mapsto \Phi(X) \otimes M$;
- The module associativity constraint is $J^{-1} \otimes id$, where J is the monoidal structure of Φ. Specifically,

$$
J_{X,Y}^{-1} \otimes id_M : \Phi(X \otimes Y) \otimes M \cong \Phi(X) \otimes \Phi(Y) \otimes M. \tag{3.14}
$$

(Here we abbreviate the module associativity of the $\mathcal{D}\text{-module category } \mathcal{M}.$)

If the C-module category M determined as above is also exact (in this case (Φ, \mathcal{M}) is called an *exact pair* by [\[EO04,](#page-41-13) Definition 3.44]), then we have the *dual tensor functor*

$$
\Phi^* : \mathcal{D}^*_{\mathcal{M}} \to \mathcal{C}^*_{\mathcal{M}},\tag{3.15}
$$

which sends $\mathbf{F} = (F, s) \in \mathcal{D}_{\mathcal{M}}^*$ to the C-module functor $(F, s_{\Phi(-), -}) \in \mathcal{C}_{\mathcal{M}}^*$.

Proposition 3.9. *Let* C *and* D *be spherical fusion categories over* C*, and let* M *be an indecomposable* D -module category which is semisimple. Suppose that $\Phi : C \to D$ *is a tensor equivalence preserving the pivotal structure. Then for any* $\mathbf{F} = (F, s) \in \mathcal{D}_{\mathcal{M}}^*$,

$$
\nu_n(\Phi^*(\mathbf{F})) = \nu_n(\mathbf{F}) \qquad (\forall n \ge 1).
$$

Proof. Evidently, it follows by Lemma [2.10\(](#page-9-3)2) that the semisimple category M is also exact as a C-module category. Thus the indicators of objects $\Phi^*(\mathbf{F}) \in C^*_{\mathcal{M}}$ are well-defined for all $\mathbf{F} \in \mathcal{D}_{\mathcal{M}}^*$.

Let us explain at first how Φ induces an equivalence between centers $\mathcal{Z}(\mathcal{C})$ and $\mathcal{Z}(\mathcal{D})$, which is mentioned in [\[NS07b,](#page-42-1) Section 2]. For the purpose, suppose $\psi : \text{Id}_{\mathcal{D}} \cong \Phi \Phi^{-1}$ is a monoidal natural isomorphism, where Φ[−]¹ denotes a *quasi-inverse* of Φ.

For each $\mathbf{V} = (V, \sigma^V) \in \mathcal{Z}(\mathcal{C})$, we define $\Phi(\mathbf{V}) = (\Phi(V), \sigma^{\Phi(V)}) \in \mathcal{Z}(\mathcal{D})$, where the halfbraiding $\sigma^{\Phi(V)}$ is defined through the commuting diagram for all $X' \in \mathcal{D}$:

$$
\Phi(V) \otimes X' \xrightarrow{\operatorname{id}_{\Phi(V)} \otimes \psi_{X'}} \Phi(V) \otimes \Phi \Phi^{-1}(X') \xrightarrow{J_{V, \Phi^{-1}(X')}} \Phi(V \otimes \Phi^{-1}(X')) \qquad (3.16)
$$
\n
$$
\sigma_{X'}^{\Phi(V)} \downarrow^{\Phi(\sigma_{\Phi^{-1}(X')}^V)} \Phi(V) \xrightarrow{\Psi_{\Phi_{X'} \otimes \operatorname{id}_{\Phi(V)}}} \Phi \Phi^{-1}(X') \otimes \Phi(V) \xrightarrow{J_{\Phi^{-1}(X'), V}} \Phi(\Phi^{-1}(X') \otimes V) .
$$

Morphisms in $\mathcal{Z}(\mathcal{C})$ are mapped to $\mathcal{Z}(\mathcal{D})$ the same with $\Phi : \mathcal{C} \to \mathcal{D}$. Indeed, one could find that the diagram

$$
\Phi(V) \otimes \Phi(X) \xrightarrow{J_{V,X}} \Phi(V \otimes X) \tag{3.17}
$$
\n
$$
\sigma_{\Phi(X)}^{\Phi(V)} \downarrow \qquad \qquad \downarrow \Phi(\sigma_X^V)
$$
\n
$$
\Phi(X) \otimes \Phi(V) \xrightarrow{J_{X,V}} \Phi(X \otimes V)
$$

commutes for all $X \in \mathcal{C}$ as a consequence of the commuting diagram [\(3.16\)](#page-16-0), and hence Φ is an equivalence between braided tensor categories.

Now we conclude relevant fusion categories and tensor functors with the following diagram of categories and functors:

$$
\mathcal{Z}(\mathcal{C}) \xrightarrow{\Omega} \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \xrightarrow{\mathbf{U}_{\mathcal{M}}} \mathcal{C}_{\mathcal{M}}^*,
$$
\n
$$
\downarrow^{\Phi^*}
$$
\n
$$
\mathcal{Z}(\mathcal{D}) \xrightarrow{\Omega'} \mathcal{Z}(\mathcal{D}_{\mathcal{M}}^*) \xrightarrow{\mathbf{U}_{\mathcal{M}}'} \mathcal{D}_{\mathcal{M}}^*
$$
\n(3.18)

where Ω' and $\mathbf{U}'_{\mathcal{M}}$ denote respectively Schauenburg's equivalence and the forgetful functor in the same ways. Here we try to verify that the diagram [\(3.18\)](#page-16-1) is commutative.

In fact according to Subsection [2.4,](#page-10-0) the composition $U_M \circ \Omega$ sends each object $V =$ $(V, \sigma^V) \in \mathcal{Z}(\mathcal{C})$ to the endofunctor $\Phi(V) \otimes -$ on M with C-module structure determined as the left vertical arrow of the following commuting diagram:

$$
\Phi(X) \otimes \Phi(V) \otimes M \xrightarrow{J_{X,V} \otimes id_M} \Phi(X \otimes V) \otimes M
$$
\n
$$
\cong \qquad \qquad \downarrow \Phi((\sigma_X^V)^{-1}) \otimes id_M
$$
\n
$$
\Phi(V) \otimes \Phi(X) \otimes M \xrightarrow{J_{V,X} \otimes id_M} \Phi(V \otimes X) \otimes M
$$
\n
$$
(3.19)
$$

for all $X \in \mathcal{C}$ and $\mathcal{M} \in M$. This is because horizontal arrows are module associativity constraints (3.14) of the C-module category M . However, according to the commuting diagram (3.17) , the left vertical arrow of (3.19) is $(\sigma_{\Phi(X)}^{\Phi(V)})$ $(\frac{\Phi(V)}{\Phi(X)})^{-1} \otimes \text{id}_M$. As a conclusion:

$$
\mathbf{U}_{\mathcal{M}} \circ \Omega \left(V, \sigma^{V} \right) = \left(\Phi(V) \otimes -, (\sigma_{\Phi(-)}^{\Phi(V)})^{-1} \otimes \mathrm{id}_{-} \right) \in \mathcal{C}_{\mathcal{M}}^{*}.
$$
 (3.20)

On the other hand, we obtain similarly that $\mathbf{U}'_{\mathcal{M}} \circ \Omega' \circ \Phi$ sends $\mathbf{V} = (V, \sigma^V) \in \mathcal{Z}(\mathcal{C})$ to the endofunctor $\Phi(V) \otimes -$ on M with D-module structure $(\sigma_-^{\Phi(V)})^{-1} \otimes id$. Afterwards, it is followed by the definition of the dual tensor equivalence Φ^* [\(3.15\)](#page-16-3) that

$$
\Phi^*\left(\Phi(V)\otimes-,({\sigma^{\Phi(V)}_+})^{-1}\otimes\operatorname{id}_-\right)=\left(\Phi(V)\otimes-,({\sigma^{\Phi(V)}_{\Phi(-)}})^{-1}\otimes\operatorname{id}_-\right)\in\mathcal{C}_\mathcal{M}^*,
$$

which coincides with (3.20) . Therefore, we have shown that (3.18) is commutative.

In order to compute the indicators desired, suppose that K' is a (right) adjoint to $\mathbf{U}'_{\mathcal{M}} \circ \Omega'$. Then we apply [\[MacL98,](#page-41-18) Theorem 1 in Section IV.8] to find that

 $(\Phi^* \circ \mathbf{U}'_{\mathcal{M}} \circ \Omega' \circ \Phi, \ \ \Phi^{-1} \circ K' \circ (\Phi^*)^{-1}),$

or identically (by the commutativity of [\(3.18\)](#page-16-1)),

$$
\left(\mathbf{U}_{\mathcal{M}}\circ\Omega,\ \ \Phi^{-1}\circ K'\circ(\Phi^*)^{-1}\right)
$$

would be an adjoint pair. Denote $K := \Phi^{-1} \circ K' \circ (\Phi^*)^{-1}$ for simplicity, and hence there is a natural isomorphism

$$
K' \cong \Phi \circ K \circ \Phi^*.
$$
\n^(3.21)

On the other hand, it is known in [\[NS07b,](#page-42-1) Section 2] that the equivalence $\Phi : \mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{D})$ between braided tensor categories also preserves the ribbon structure, namely,

$$
\theta'_{\Phi(\mathbf{V})} = \Phi(\theta_{\mathbf{V}}) \qquad (\forall \mathbf{V} \in \mathcal{Z}(\mathcal{C})), \tag{3.22}
$$

where θ and θ' are the ribbon structures of $\mathcal{Z}(\mathcal{C})$ and $\mathcal{Z}(\mathcal{D})$ respectively. Therefore, we find by Lemma [2.1](#page-4-2) that

$$
\begin{aligned} \operatorname{ptr}\left(\theta'^{n}_{K'(\mathbf{F})}\right) \stackrel{(3.21)}{=} \operatorname{ptr}\left(\theta'^{n}_{\Phi\circ K\circ\Phi^{*}(\mathbf{F})}\right) \stackrel{(3.22)}{=} \operatorname{ptr}\left(\Phi\left(\theta_{K\circ\Phi^{*}(\mathbf{F})}\right)^{n}\right) \\ = \operatorname{ptr}\left(\Phi\left(\theta^{n}_{K\circ\Phi^{*}(\mathbf{F})}\right)\right) \stackrel{(2.10)}{=} \operatorname{ptr}\left(\theta^{n}_{K\circ\Phi^{*}(\mathbf{F})}\right) \end{aligned}
$$

holds for each $\mathbf{F} \in \mathcal{D}_{\mathcal{M}}^*$.

Finally, since Φ is a tensor equivalence between fusion categories preserving the pivotal structure,

$$
\dim(\mathcal{C}) \quad \stackrel{(3.2)}{=} \sum_{X \in \mathcal{O}(\mathcal{C})} \text{ptr}(\text{id}_X) \text{ptr}(\text{id}_{X^*}) \stackrel{(2.10)}{=} \sum_{X \in \mathcal{O}(\mathcal{C})} \text{ptr}(\Phi(\text{id}_X)) \text{ptr}(\Phi(\text{id}_{X^*}))
$$
\n
$$
= \sum_{X \in \mathcal{O}(\mathcal{C})} \text{ptr}(\text{id}_{\Phi(X)}) \text{ptr}(\text{id}_{\Phi(X^*)}) = \sum_{X' \in \mathcal{O}(\mathcal{D})} \text{ptr}(\text{id}_{X'}) \text{ptr}(\text{id}_{X'^*}) \stackrel{(3.2)}{=} \dim(\mathcal{D}).
$$

As a conclusion, for each $n \geq 1$,

$$
\nu_n(\Phi^*(\mathbf{F})) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta''_{K'(\mathbf{F})}\right) = \frac{1}{\dim(\mathcal{D})} \text{ptr}\left(\theta''_{K \circ \Phi^*(\mathbf{F})}\right) = \nu_n(\mathbf{F}).
$$

4. Invariance for certain objects under the duality of categories

4.1. Ends and the central Hopf comonad. The notions of ends play an essential role in the (right) adjoint of the forgetful functor from the center of a finite tensor category. We start by recalling relevant definitions, which could be found in [\[MacL98,](#page-41-18) Chapter IX].

Suppose that C and D are categories, and $S, T : C^{\vee} \times C \to D$ are functors, where C^{\vee} denotes the category having same objects but reversed directions of morphisms with C. A *dinatural transformation* from S to T is a family

$$
\varphi = {\varphi_X : S(X, X) \to T(X, X)}_{X \in \mathcal{C}}
$$

of morphisms in $\mathcal D$ satisfying

$$
T(\mathrm{id}_X, f) \circ \varphi_X \circ S(f, \mathrm{id}_X) = T(f, \mathrm{id}_Y) \circ \varphi_Y \circ S(\mathrm{id}_Y, f)
$$

for all morphisms $f: X \to Y$ in C.

An *end* of S is an object $E \in \mathcal{D}$ equipped with a dinatural transformation π from E (as a constant functor) to S such that: For every dinatural transformation π' from a constant functor E' to S, there is a unique morphism $g: E' \to E$ satisfying $\pi'_X = \pi_X \circ g$ for all $X \in \mathcal{C}$. An end of S is denoted by $\int_{X \in \mathcal{C}} S(X, X)$, which is unique up to a unique isomorphism.

The definition of a *coend* $\int^{X \in \mathcal{C}} S(X, X)$ of S is completely dual to an end. We omit the details, since the coend construction seldom in this paper. See [\[MacL98,](#page-41-18) Chapter IX] for the details.

In particular, for any $V \in \mathcal{C}$, consider the functor

$$
\mathcal{C}^{\vee} \times \mathcal{C} \to \mathcal{C}, \ \ (X, Y) \mapsto Y \otimes V \otimes X^*.
$$
\n
$$
(4.1)
$$

We would denote the end of [\(4.1\)](#page-18-2) by $Z(V) = \int_{X \in \mathcal{C}} X \otimes V \otimes X^*$, which makes Z an endofunctor called the *central Hopf comonad* on C (see [\[Shi17b,](#page-42-9) Definition 3.2]) if such ends exist for all $V \in \mathcal{C}$. In this case, the universal dinatural transformations would be denoted by

$$
\pi_X^{\mathsf{Z}(V)}:\mathsf{Z}(V)\to X\otimes V\otimes X^* \quad (X\in\mathcal{C}).
$$

Furthermore, the comonad structure of Z means that there are morphisms

$$
\delta_V : \mathsf{Z}(V) \to \mathsf{Z}^2(V) \quad \text{and} \quad \epsilon_V : \mathsf{Z}(V) \to V \quad (V \in \mathcal{C})
$$

uniquely determined by the property that $\epsilon_V = \pi_V^{\mathsf{Z}(1)}$ $V^{(1)}$ and

$$
\left(\operatorname{id}_X\otimes\pi_Y^{{\sf Z}(V)}\otimes\operatorname{id}_{X^*}\right)\circ\pi_X^{{\sf Z}^2(V)}\circ\delta_V=\pi_{X\otimes Y}^{{\sf Z}(V)}
$$

hold for all objects $V, X, Y \in \mathcal{C}$ (see [\[Shi17b,](#page-42-9) Section 3.2] for details).

Lemma 4.1. *(*[\[Shi17b,](#page-42-9) Section 3] *or* [\[Shi19,](#page-42-10) Section 3.3]*) Let* C *be a finite tensor category. Then the forgetful functor* $U : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ *has a right adjoint:*

$$
I: \mathcal{C} \to \mathcal{Z}(\mathcal{C}), \ \ V \mapsto \left(\mathsf{Z}(V), \sigma^{\mathsf{Z}(V)}\right),
$$

 20 K. LI

where the half-braiding $\sigma^{Z(V)}$ *is*

$$
\sigma_X^{\mathsf{Z}(V)}:\mathsf{Z}(V)\otimes X\xrightarrow{\delta_V\otimes \mathrm{id}_X}\mathsf{Z}^2(V)\otimes X\xrightarrow{\pi_X^{\mathsf{Z}^2(V)}\otimes \mathrm{id}_X}X\otimes \mathsf{Z}(V)\otimes X^*\otimes X\xrightarrow{\mathrm{id}_X\otimes \mathrm{id}_{\mathsf{Z}(V)}\otimes \text{ev}_X}X\otimes \mathsf{Z}(V).
$$

The half-braiding $\sigma^{Z(V)}$ determined in Lemma [4.1](#page-18-3) is referred to as the *canonical halfbraiding* for $Z(V)$. Here we provide another description for later use:

Lemma 4.2. Let C be a finite tensor category. Suppose $V \in \mathcal{C}$, and $\sigma^{Z(V)}$ is the canonical *half-braiding for* Z(V)*. Then*

$$
\left(\mathrm{id}_X \otimes \pi_Y^{\mathsf{Z}(V)}\right) \circ \sigma_X^{\mathsf{Z}(V)} = \left(\mathrm{id}_X \otimes \mathrm{id}_Y \otimes \mathrm{id}_V \otimes \mathrm{id}_Y \ast \otimes \mathsf{ev}_X\right) \circ \left(\pi_{X \otimes Y}^{\mathsf{Z}(V)} \otimes \mathrm{id}_X\right) \tag{4.2}
$$

holds for all $X, Y \in \mathcal{C}$.

Proof. Note that the canonical half-braiding for $Z(V)$ is defined in Lemma [4.1](#page-18-3) as

$$
\sigma_X^{\mathsf{Z}(V)} = (\mathrm{id}_X \otimes \mathrm{id}_{\mathsf{Z}(V)} \otimes \mathsf{ev}_X) \circ (\pi_X^{\mathsf{Z}^2(V)} \otimes \mathrm{id}_X) \circ (\delta_V \otimes \mathrm{id}_X).
$$

Thus our goal is to explain the commutativity of the diagram

for all $X, Y \in \mathcal{C}$: The top-right quadrangle commutes due to [\[Shi17b,](#page-42-9) Equation (3.8)], while the bottom quadrangle is commutative since \otimes is a bifunctor on C.

Consequently, Equation (4.2) is a result of the commutativity of the outside hexagon of this diagram. \square

Remark 4.3. *(*[\[ENO05,](#page-41-14) Proposition 5.4]*) If* C *is furthermore a fusion category, then*

$$
\mathbf{U}\circ I(X)\cong \bigoplus_{Y\in\mathcal{O}(\mathcal{C})} Y\otimes X\otimes Y^*
$$

holds for each $X \in \mathcal{C}$ *, where* I *is a (right) adjoint functor to the forgetful functor* $\mathbf{U} : \mathcal{Z}(\mathcal{C}) \rightarrow$ C*.*

4.2. The internal Hom functor and adjoints to the forgetful functor. Suppose $\mathcal M$ is an exact and indecomposable left module category over a finite tensor category \mathcal{C} . Then \mathcal{M} is known to be *closed*, which mean that the functor

$$
\mathcal{C} \to \mathcal{M}, \ \ X \mapsto -\otimes M
$$

has a right adjoint for each $M \in \mathcal{M}$. Consequently the *internal Hom functor*

$$
\underline{\mathrm{Hom}}(-,-): \mathcal{M}^{\vee} \times \mathcal{M} \to \mathcal{C}
$$

could be defined ($[EO04, Section 3.2]$ $[EO04, Section 3.2]$). It is furthermore described in $[Shi20]$ etc. that there exist C -module structures

$$
\begin{aligned}\n\mathfrak{a}_{X,M,N} &\; : \quad X \otimes \underline{\text{Hom}}(M,N) \xrightarrow{\cong} \underline{\text{Hom}}(M,X \otimes N) &\quad (X \in \mathcal{C}), \\
\mathfrak{b}_{M,N,Y} &\; : \quad \underline{\text{Hom}}(M,N) \otimes Y^* \xrightarrow{\cong} \underline{\text{Hom}}(Y \otimes M,N) &\quad (Y \in \mathcal{C}), \\
\mathfrak{c}_{X,M,N,Y} &\; : \quad X \otimes \underline{\text{Hom}}(M,N) \otimes Y^* \xrightarrow{\cong} \underline{\text{Hom}}(Y \otimes M,X \otimes N) &\quad (X,Y \in \mathcal{C})\n\end{aligned}
$$

with relations listed in the following lemma, and they are all dinatural in $(M, N) \in \mathcal{M}^{\vee} \times \mathcal{M}$. Note that $\mathfrak{c}_{X,M,N,Y}$ is also dinatural in $(Y, X) \in \mathcal{C}^{\vee} \times \mathcal{C}$. Here our notations follow [\[Shi24,](#page-42-11) Section 2.3] as a reference for a number of formulas among them.

Lemma 4.4. *With assumptions and notations above. We have:*

(1) *For all* $X, Y \in \mathcal{C}$ *and* $M, N \in \mathcal{M}$,

$$
\mathfrak{c}_{X,M,N,Y} = \mathfrak{a}_{X,Y \otimes M,N} \circ (\mathrm{id}_X \otimes \mathfrak{b}_{M,N,Y}) = \mathfrak{b}_{M,X \otimes N,Y} \circ (\mathfrak{a}_{X,M,N} \otimes \mathrm{id}_{Y^*});
$$

(2) *For all* $X_1, X_2, Y_1, Y_2 \in \mathcal{C}$ *and* $M, N \in \mathcal{M}$ *,*

$$
\mathfrak{a}_{X_1 \otimes X_2,M,N} = \mathfrak{a}_{X_1,M,X_2 \otimes N} \circ (\mathrm{id}_{X_1} \otimes \mathfrak{a}_{X_2,M,N}), \tag{4.3}
$$

$$
\mathfrak{b}_{M,N,Y_1 \otimes Y_2} = \mathfrak{b}_{Y_2 \otimes M,N,Y_1} \circ (\mathfrak{b}_{M,N,Y_2} \otimes \mathrm{id}_{Y_1^*})
$$
(4.4)

and

$$
\mathfrak{c}_{X_1 \otimes X_2, M, N, Y_2^* \otimes Y_1^*} = \mathfrak{c}_{X_1, Y_2 \otimes M, X_2 \otimes N, Y_1} \circ (\mathrm{id}_{X_1} \otimes \mathfrak{c}_{X_2, M, N, Y_2} \otimes \mathrm{id}_{Y_1^*}). \tag{4.5}
$$

- *Proof.* (1) The latter equality could be found as [\[Shi24,](#page-42-11) Equation (2.10)], while the former one is in fact the definition of \mathfrak{c} (See [\[Shi20,](#page-42-6) Equation (2.18)] for example).
	- (2) Equation [\(4.3\)](#page-20-0) is introduced in $[\text{Shi20}, \text{Equation (2.10)}]$ as the left C-module structure of the functor $\underline{\text{Hom}}(M, -)$. As for Equation [\(4.4\)](#page-20-0), it is essentially proved in [\[Shi20,](#page-42-6) Lemmas 2.3 and A.1], where the notion of b appearing is mutually inverse with [\[Shi24\]](#page-42-11) and ours in this paper.

In order to show Equation [\(4.5\)](#page-20-1), let us verify the commutativity of the following diagram:

$$
\begin{array}{c}\nX_1 \otimes X_2 \otimes \underline{\mathrm{Hom}}(M,N) \otimes Y_2^* \otimes Y_1^* \xrightarrow{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{M,N,Y_2} \otimes \mathrm{id}} X_1 \otimes X_2 \otimes \underline{\mathrm{Hom}}(Y_2 \otimes M,N) \otimes Y_1^* \\
\downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{M,N,Y_1 \otimes Y_2}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N,Y_1}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N \otimes \mathrm{id}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N \otimes \mathrm{id}}}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N \otimes \mathrm{id}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N \otimes \mathrm{id}}}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N \otimes \mathrm{id}} \downarrow^{\mathrm{id} \otimes \mathrm{id} \otimes \mathrm{b}_{Y_2 \otimes M,N}} \downarrow^{\mathrm{id} \otimes \mathrm{id
$$

In fact, the commutativity of the two triangles on the left are due to Equations [\(4.3\)](#page-20-0) and [\(4.4\)](#page-20-0), and the right-sided quadrangle commutes since both compositions equal to $id_{X_1} \otimes \mathfrak{c}_{X_2,Y_2\otimes M,N,Y_1}$ according to (1).

Finally by (1) again, the compositions of the outside hexagon are respectively $\mathfrak{c}_{X_1 \otimes X_2,M,N,Y_2^* \otimes Y_1^*}$ and $\mathfrak{c}_{X_1,Y_2 \otimes M,X_2 \otimes N,Y_1} \circ (\mathrm{id}_{X_1} \otimes \mathfrak{c}_{X_2,M,N,Y_2} \otimes \mathrm{id}_{Y_1^*})$. Thus Equation [\(4.5\)](#page-20-1) is implied by the commutativity of the diagram above.

 \Box

With the languages of ends, the internal Hom functor and the module structures above, we could choose some $\mathcal{Z}(\rho^{ra})$ appearing in Lemma [3.7](#page-14-1) as the right adjoint to the forgetful functor

 $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \to \mathcal{C}_{\mathcal{M}}^*$ in [\[Shi20,](#page-42-6) Section 3.7]. Recall that for an object $\mathbf{F} = (F, s) \in \text{Rex}_{\mathcal{C}}(\mathcal{M})$, we have $\mathcal{Z}(\rho^{\text{ra}})(\mathbf{F}) = (\rho^{\text{ra}}(F), \sigma^{\mathbf{F}}) \in \mathcal{Z}(\mathcal{C})$, where

$$
\rho^{\text{ra}}(F) = \int_{M \in \mathcal{M}} \underline{\text{Hom}}(M, F(M)) \in \mathcal{C}
$$
\n(4.6)

is an end of $\underline{\text{Hom}}(-, F(-))$, and the half-braiding $\sigma^{\mathbf{F}}$ is a unique natural transformation such that the following diagram commutes in C for all objects $X \in \mathcal{C}$ and $M \in \mathcal{M}$:

$$
\rho^{\text{ra}}(F) \otimes X \xrightarrow{\pi_{X \otimes M} \otimes \text{id}} \longrightarrow \text{Hom}(X \otimes M, F(X \otimes M)) \otimes X
$$
\n
$$
\downarrow \qquad \qquad \downarrow \qquad
$$

according to [\[Shi20,](#page-42-6) Lemma 3.12].

At the final of this subsection, we recall the following lemma in order to provide our usual choice of an adjoint functor K in Definition [3.3:](#page-12-1)

Lemma 4.5. *(*[\[Shi20,](#page-42-6) Theorem 3.14] *and* [Shi20, Corollary 3.15]*) Let* $\mathbf{U}_{\mathcal{M}} : \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \to \mathcal{C}_{\mathcal{M}}^*$ *be the forgetful functor. Then:*

(1)
$$
(\mathbf{U}_{\mathcal{M}} \circ \Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}_{\mathcal{M}}^*, \quad \mathcal{Z}(\rho^{\text{ra}}) : \mathcal{C}_{\mathcal{M}}^* \to \mathcal{Z}(\mathcal{C})
$$
 is an adjoint pair;
\n(2) $(\mathbf{U}_{\mathcal{M}} : \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*) \to \mathcal{C}_{\mathcal{M}}^*, \quad \Omega \circ \mathcal{Z}(\rho^{\text{ra}}) : \mathcal{C}_{\mathcal{M}}^* \to \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ is an adjoint pair.

4.3. An equality between indicators in $\mathcal{C}_{\mathcal{M}}^*$ and \mathcal{C} . As we know that if $\mathcal M$ is a finite C-module category, we could assume that $\mathcal{M} = \mathcal{C}_A$, the category of right A-modules in C, for some algebra $A \in \mathcal{C}$. In this case, there is a tensor equivalence

$$
{}_{A}\mathcal{C}_{A} \to \mathcal{C}_{\mathcal{M}}^{*}, \quad P \mapsto (-) \otimes_{A} P, \tag{4.8}
$$

according to the proof of [\[EO04,](#page-41-13) Proposition 3.23].

Let us consider indicators of certain objects in $\mathcal{C}_{\mathcal{M}}^*$ under this equivalence. Specifically, for each object $M \in \mathcal{C}_A$, there is an A-bimodule structure on $A \otimes M \in \mathcal{C}$ via the regular left A-module and the right A-module structure on M. This would determine a functor $(-) \otimes_A (A \otimes M) \in C^*_{\mathcal{M}}$ under [\(4.8\)](#page-21-2), which is canonically isomorphic to $-\otimes M$ with C-module structure as the associativity constraint of $\mathcal C$ (and hence abbreviated).

Lemma 4.6. *Let* C *be a finite tensor category. Suppose that* A *is an algebra in* C *such that* $\mathcal{M} = \mathcal{C}_A$ *is an indecomposable* \mathcal{C} -module category which is exact. For any $\mathbf{M} = (M, \mu) \in \mathcal{C}_A$, *denote*

$$
\mathsf{E} := \rho^{\text{ra}}(-\otimes M) = \int_{N \in \mathcal{C}_A} \underline{\text{Hom}}(N, N \otimes M) \in \mathcal{C}
$$

as in [\(4.6\)](#page-21-3) with half-braiding σ^E defined such that Diagram [\(4.7\)](#page-21-4) commutes. Then the follow*ing diagram commutes for each* $X \in \mathcal{C}$ *:*

$$
\begin{array}{c}\n\mathsf{E} \otimes X \xrightarrow{\pi_{X \otimes N}^{\mathsf{E}} \otimes \mathrm{id}_{X}} \longrightarrow \mathrm{Hom}(X \otimes N, X \otimes N \otimes M) \otimes X \\
\downarrow^{\mathsf{c}_{X, N, N \otimes M, X}^{\mathsf{I}} \otimes \mathrm{id}_{X}} \\
\downarrow^{\mathsf{c}_{X, N, N \otimes M, X}^{\mathsf{I}} \otimes \mathrm{id}_{X}} \\
X \otimes \mathrm{Hom}(N, N \otimes M) \otimes X^{*} \otimes X \\
\downarrow^{\mathrm{id}_{X} \otimes \mathrm{id}_{\mathrm{Hom}(N, N \otimes M)} \otimes \mathrm{ev}_{X}} \\
X \otimes \mathsf{E} \xrightarrow{\mathrm{id}_{X} \otimes \pi_{X}^{\mathsf{E}}} X \otimes \mathrm{Hom}(N, N \otimes M) .\n\end{array} \tag{4.9}
$$

Proof. By comparison to Diagram (4.7) , it suffices to show that the following diagram commutes for all $X \in \mathcal{C}$ and $N \in \mathcal{C}_A$:

$$
\begin{array}{c}\n\underbrace{\operatorname{Hom}}(X \otimes N, X \otimes N \otimes M) \otimes X \\
\stackrel{\mathfrak{b}_{X \otimes N, X \otimes N \otimes M, X} \otimes \mathrm{id}_{X}}\n\downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \qquad \qquad \downarrow \
$$

where the C-module structure of $-\otimes M$ is abbreviated. Specifically, the top triangle is commutative due to Lemma [4.4\(](#page-20-2)1), and the bottom quadrangle commutes as \otimes is a bifunctor on $\mathcal{C}.$

Proposition 4.7. *Let* C *be a finite tensor category. Suppose that* A *is an algebra in* C *such that* $M = C_A$ *is an indecomposable* C-module category which is exact. Then for any $M \in C_A$,

$$
\left(\int_{N \in \mathcal{C}_A} \underline{\text{Hom}}(N, N \otimes M), \sigma^{-\otimes M}\right) \cong \left(\mathsf{Z}(\underline{\text{Hom}}(A, M)), \sigma^{\mathsf{Z}(\underline{\text{Hom}}(A, M))}\right)
$$

as objects in $\mathcal{Z}(\mathcal{C})$ *.*

Proof. Recall from $[Ost03, Lemma 3.2]$ that there is an adjoint pair (L, R) defined as

$$
\begin{cases}\nL: \mathcal{C} \to \mathcal{C}_A, & X \mapsto X \otimes A; \\
R: \mathcal{C}_A \to \mathcal{C}, & N \mapsto N\n\end{cases}
$$
 (the forgetful functor), (4.11)

(4.10)

with the unit and counit

$$
\eta: \mathrm{Id}_{\mathcal{C}} \to -\otimes A = RL, \quad \mu: -\otimes A = LR \to \mathrm{Id}_{\mathcal{C}_A}.
$$
\n
$$
(4.12)
$$

We remark that μ could be chosen such that μ_N is the right A-module structure on every $N \in \mathcal{C}_A$ according to the proof of [\[EGNO15,](#page-41-10) Lemma 7.8.12], and thus

$$
id_X \otimes \mu_N = \mu_{X \otimes N} \qquad (\forall X \in \mathcal{C}, \ \forall N \in \mathcal{C}_A)
$$
\n(4.13)

as the right A-module structure of $X \otimes N$.

 24 K. LI

Suppose $M \in C_A$. We apply [\[Shi20,](#page-42-6) Lemma 2.2] (which is the dual form of [\[BV12,](#page-40-4) Lemma 3.9]) to this adjunction and the functor

$$
\mathcal{C} \times (\mathcal{C}_A)^{\vee} \to \mathcal{C}, \quad (X, N) \mapsto \underline{\text{Hom}}(N, X \otimes M),
$$

and then obtain an isomorphism f :

$$
\mathsf{E} := \int_{N \in \mathcal{C}_A} \underline{\operatorname{Hom}}(N, N \otimes M) \stackrel{f}{\cong} \int_{X \in \mathcal{C}} \underline{\operatorname{Hom}}(X \otimes A, X \otimes M) = \mathsf{E}'. \tag{4.14}
$$

Note that these ends do exist according to [\[Shi20,](#page-42-6) Theorem 3.4].

However, recall that there is an isomorphism

 $\mathfrak{c}_{Y,A,M,X}: Y \otimes \underline{\mathrm{Hom}}(A,M) \otimes X^* \cong \underline{\mathrm{Hom}}(X \otimes A, Y \otimes M),$

introduced before Lemma [4.4,](#page-20-2) which is dinatural in $(X, Y) \in C^{\vee} \times C$. Consequently, another isomorphism g:

$$
\mathsf{E}' = \int_{X \in \mathcal{C}} \underline{\operatorname{Hom}}(X \otimes A, X \otimes M) \stackrel{g}{\cong} \int_{X \in \mathcal{C}} X \otimes \underline{\operatorname{Hom}}(A, M) \otimes X^* = \mathsf{Z}(\underline{\operatorname{Hom}}(A, M)) = \mathsf{C} \tag{4.15}
$$

between ends could be obtained.

Furthermore, let us explain that the isomorphisms $f(4.14)$ $f(4.14)$ and $g(4.15)$ $g(4.15)$ between ends could be chosen such that the following diagram commutes for all objects $N \in \mathcal{C}_A$ and $X \in \mathcal{C}$:

$$
\underline{\text{Hom}}(N, N \otimes M) \xleftarrow{\pi_N^{\mathbb{F}}} \int_{N \in \mathcal{C}_A} \underline{\text{Hom}}(N, N \otimes M) \xrightarrow{\pi_{X \otimes A}^{\mathbb{F}}} \underline{\text{Hom}}(X \otimes A, X \otimes A \otimes M)
$$
\n
$$
\underline{\text{Hom}}(\mu_N, \text{id}) \downarrow \qquad \qquad \downarrow \
$$

where

 $\pi^{\mathsf{E}} : \mathsf{E} \to \underline{\mathrm{Hom}}(-, -\otimes M), \ \ \pi^{\mathsf{E}'} : \mathsf{E}' \to \underline{\mathrm{Hom}}(-\otimes A, -\otimes M) \ \text{ and } \ \pi^{\mathsf{C}} : \mathsf{C} \to (-)\otimes \underline{\mathrm{Hom}}(A, M) \otimes (-)^*$ denote the corresponding universal dinatural transformations for the ends respectively: More explicitly, the commutativity of the top rectangles of [\(4.16\)](#page-23-2) could be ensured due to the paragraph below [\[Shi20,](#page-42-6) Lemma 2.2], while the bottom rectangles of [\(4.16\)](#page-23-2) could be obtained by [\[MacL98,](#page-41-18) Proposition 1 in Section IX.7].

We point out the left rectangles of the commuting diagram [\(4.16\)](#page-23-2) as follows, where the isomorphism $g \circ f$ is denoted by h for simplicity:

$$
\mathsf{E} \longrightarrow \mathsf{C} \tag{4.17}
$$

\n
$$
\pi_N^{\mathsf{E}} \downarrow \qquad \qquad \mathsf{Hom}(N, N \otimes M) \longrightarrow \mathsf{Hom}(N \otimes A, N \otimes M) \longrightarrow \mathsf{C}_{\mathsf{N}, A, M, N} \otimes \mathsf{Hom}(A, M) \otimes N^*.
$$

On the other hand, one might apply [\[MacL98,](#page-41-18) Theorem 1(ii) in Section IV.1] to the adjunction [\(4.12\)](#page-22-0) to obtain that

$$
\mu_N \circ \eta_N = \text{id}_N
$$
, and hence $\underline{\text{Hom}}(\eta_N, \text{id}) \circ \underline{\text{Hom}}(\mu_N, \text{id}) = \text{id} \ (\forall N \in C_A).$ (4.18)

Now we focus on the following diagram [\(4.19\)](#page-24-0) and try to prove its commutatively for all X ∈ C and N ∈ C_A. Please note that due to Equations [\(4.18\)](#page-23-3), if the arrow id ⊗Hom(μ_N , id) in Diagram [\(4.19\)](#page-24-0) is replaced by the dotted arrow id ⊗Hom(η_N , id) with inverse direction, then the new diagram will be also commutative:

Our reasons for the commutativity of the cells in this diagram (with original arrow id $\otimes \underline{\text{Hom}}(\mu_N, id)$) are stated respectively as follows:

- The two (top and bottom) pentagons: Diagram (4.17) commutes;
- The upper rectangle: This is obtained by the naturality of $c_{X,-,N,N\otimes M,X}$ for the morphism μ_N in C, where Equation [\(4.13\)](#page-22-1) will be used;
- The triangle: Equation (4.5) ;
- The remaining two quadrangles (the right trapezoid and the lower rectangle): ⊗ is a bifunctor on C .

If we put the following two commuting diagrams - [\(4.9\)](#page-22-2) and [\(4.2\)](#page-19-1) where $\mathsf{Z}(\text{Hom}(A, M)) = \mathsf{C}$ - at the left and right sides of [\(4.16\)](#page-23-2) respectively:

then we could conclude by the commutativity (with the dotted arrow id $\otimes \underline{\text{Hom}}(\eta_N, id)$ instead) that

$$
(\mathrm{id}_X\otimes\pi_N^{\mathsf{E}})\circ\sigma_X^{\mathsf{E}}=(\mathrm{id}_X\otimes\pi_N^{\mathsf{E}})\circ[(\mathrm{id}_X\otimes h)\circ\sigma_X^{\mathsf{C}}\circ(h^{-1}\otimes\mathrm{id}_X)]\quad \ (\forall X\in\mathcal{C},\ \forall N\in\mathcal{C}_A).
$$

 26 K. LI

It follows by Fubini Theorem and the universal property for ends that

$$
\sigma_X^{\mathsf{E}} \circ (h \otimes \mathrm{id}_X) = (\mathrm{id}_X \otimes h) \circ \sigma_X^{\mathsf{C}} \qquad (\forall X \in \mathcal{C}).
$$

Finally as a result, (E, σ^E) and (C, σ^C) are isomorphic objects in $\mathcal{Z}(\mathcal{C})$ as desired.

The following corollary could be a generalization of the last isomorphism in [\[FS23,](#page-41-19) Remark 6]:

Corollary 4.8. *Let* C *be a finite tensor category. Suppose that* A *is an algebra in* C *such that* $\mathcal{M} = \mathcal{C}_A$ *is an indecomposable* $\mathcal{C}\text{-module category which is exact. Then for any $M \in \mathcal{C}_A$,$

$$
\mathcal{Z}(\rho^{\text{ra}})(-\otimes M) = \left(\int_{N \in \mathcal{C}_A} \underline{\text{Hom}}(N, N \otimes M), \sigma^{-\otimes M} \right) \cong (\mathcal{Z}(M), \sigma^{\mathcal{Z}(M)}),
$$

where $\mathcal{Z}(\rho^{ra})$ *is defined in* [\(4.6\)](#page-21-3) *and* [\(4.7\)](#page-21-4) *as a right adjoint of* $\mathbf{U}_{\mathcal{C}_A} \circ \Omega : \mathcal{Z}(\mathcal{C}) \to \mathcal{C}_{\mathcal{C}_A}^*$.

Proof. As the central Hopf comonad **Z** is an endofunctor on C, the isomorphism $\sigma_X^{Z(V)}$ defined in Lemma [4.1](#page-18-3) will be natural in $V, X \in \mathcal{C}$. Thus by Proposition [4.7,](#page-22-3) we only need to show that $M \cong Hom(A, M)$ as objects in C. In fact due to Yoneda Lemma, it is obtained by the following isomorphism

$$
\operatorname{Hom}_{\mathcal{C}}(X, M) \stackrel{(4.11)}{\cong} \operatorname{Hom}_{\mathcal{C}_A}(X \otimes A, M) \cong \operatorname{Hom}_{\mathcal{C}}(X, \underline{\operatorname{Hom}}(A, M))
$$

which is natural in $X \in \mathcal{C}$.

Theorem 4.9. *Let* C *be a spherical fusion category over* C*. Suppose that* A *is an algebra in* C such that $M = C_A$ is an indecomposable C-module category which is semisimple. Then for *any* $M \in \mathcal{C}_A$, the *n*-th indicator of $-\otimes M \in \mathcal{C}_{\mathcal{C}_A}^*$ is

$$
\nu_n(-\otimes M) = \nu_n(M) \qquad (\forall n \ge 1),
$$

where:

- *The* C-module functor $-\otimes M$ *is canonically isomorphic to* $(-) \otimes_A (A \otimes M)$ *with structure as the associativity constraint of* C*;*
- *The object* M *in the right-hand side is regarded as the image in* C *(under the forgetful functor from* C_A *to* C *)*.

Proof. We compute the indicators in the left-hand side of this equation by Definition [3.3.](#page-12-1) Choose $K_{\mathcal{C}_A} = \mathcal{Z}(\rho_{\mathcal{C}_A}^{ra})$ as in Corollary [4.8,](#page-25-1) which is a two-sided adjoint of $U_M \circ \Omega$. Thus for any $n \geq 1$,

$$
\nu_n(-\otimes M) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta_{\mathcal{Z}(\rho^{ra})(-\otimes M)}^n\right). \tag{4.20}
$$

On the other hand, the *n*-th indicator of $M \in \mathcal{C}$ is computed according to Lemma [3.1](#page-12-5) as

$$
\nu_n(M) = \frac{1}{\dim(\mathcal{C})} \text{ptr}\left(\theta^n_{(\mathsf{Z}(M), \sigma^{\mathsf{Z}(M)})}\right),\tag{4.21}
$$

because we know by Lemma [4.1](#page-18-3) that $I: \mathcal{C} \to \mathcal{Z}(\mathcal{C}), V \mapsto (\mathcal{Z}(V), \sigma^{\mathcal{Z}(V)})$ is a right adjoint to the forgetful functor $\mathbf{U}: \mathcal{Z}(\mathcal{C}) \to \mathcal{C}$.

Finally, it follows by Lemma [2.1](#page-4-2) that the right-hand sides of Equations [\(4.20\)](#page-25-2) and [\(4.21\)](#page-25-3) coincide, since $\mathcal{Z}(\rho^{ra})(-\otimes M) \cong (\mathcal{Z}(M), \sigma^{\mathcal{Z}(M)})$ in $\mathcal{Z}(\mathcal{C})$ due to Corollary [4.8.](#page-25-1)

We end this subsection by considering the particular case when $\mathcal M$ is chosen as $\mathcal C$ with the regular left C-module category. There is a tensor equivalence

$$
\mathcal{C} \to \mathcal{C}_{\mathcal{C}}^*, \ \ X \mapsto - \otimes X,
$$

according to [\[EGNO15,](#page-41-10) Example 7.12.3] for example. Under this equivalence, we remark that indicators of objects in $\mathcal{C}_{\mathcal{C}}^*$ coinsides with those in a spherical fusion category \mathcal{C} which are originally defined by Ng and Schauenburg.

Corollary 4.10. *Let* \mathcal{C} *be a spherical fusion category over* \mathbb{C} *. Then for each* $X \in \mathcal{C}$ *,*

$$
\nu_n(-\otimes X) = \nu_n(X) \qquad (\forall n \ge 1), \tag{4.22}
$$

where $-\otimes X$ is an object in $\mathcal{C}_{\mathcal{C}}^*$ with $\mathcal{C}\text{-module}$ structure as the inverse of the associativity *constraint in* C*.*

Proof. We show this with the help of Theorem [4.9.](#page-25-0) Since left regular C-module category could be identified with the category C_1 of right 1-modules in C, where the unit object $1 \in C$ is trivially an algebra. Meanwhile, the functor $-\otimes X$ is clearly isomorphic to $-\otimes_1 (1 \otimes X)$ for each $X \in \mathcal{C}$.

 \Box

4.4. Frobenius-Schur exponents and dimensions of objects in $\mathcal{C}_{\mathcal{M}}^*$. Frobenius-Schur exponents of a pivotal monoidal category and its objects are defined in [\[NS07b,](#page-42-1) Definition 5.1]. In order to recall these definitions, let us use convention $\min \emptyset = \infty$ in this paper.

For any object X in a pivotal monoidal category C, the *Frobenius-Schur exponent of* X is defined as

$$
\text{FSexp}(X) := \min\{n \ge 1 \mid \nu_n(X) = \text{ptr}(\text{id}_X)\},\tag{4.23}
$$

where ptr_l denotes the left pivotal trace. Moreover, the *Frobenius-Schur exponent* of C is defined to be

$$
\text{FSexp}(\mathcal{C}) := \min\{n \ge 1 \mid \forall X \in \mathcal{C}, \ \nu_n(X) = \text{ptr}(\text{id}_X)\}. \tag{4.24}
$$

However, due to the reason that $\mathcal{C}_{\mathcal{M}}^{*}$ is not known to be consequently pivotal, the Frobenius-Schur exponents of its objects might not be defined frequently. However, when the dual fusion category $\mathcal{C}_{\mathcal{M}}^*$ is assumed to be pivotal such that Schauenburg's equivalence Ω preserves the pivotal structure, we could compare the Frobenius-Schur exponents of \mathcal{C} and $\mathcal{C}_{\mathcal{M}}^*$ by a combination of [\[NS07a,](#page-41-3) Corollary 4.4] and [\[NS07b,](#page-42-1) Corollary 7.8]:

Corollary 4.11. *Let* C *be any spherical fusion category over* C *with spherical structure* j*,* and let M be an indecomposable C -module category which is semisimple. Suppose $C^*_{\mathcal{M}}$ is also *spherical and* Ω : $\mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^{*})$ *preserves the pivotal structure. Then*

$$
\text{FSexp}(\mathcal{C}_{\mathcal{M}}^*) = \text{FSexp}(\mathcal{C}),\tag{4.25}
$$

where $\text{FSexp}(\mathcal{C}_{\mathcal{M}}^*)$ *is the Frobenius-Schur exponent of the spherical fusion category* $\mathcal{C}_{\mathcal{M}}^*$ *in the sense of [\(4.24\)](#page-26-1).*

Proof. We know according to [\[NS07a,](#page-41-3) Corollary 4.4] that Ω : $\mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ would preserves the indicators of all the corresponding objects, as it is assumed to preserve the pivotal structure. Consequently, the definition of the Frobenius-Schur exponent [\(4.24\)](#page-26-1) implies that

$$
\operatorname{FSexp}(\mathcal{Z}(\mathcal{C}_\mathcal{M}^*))=\operatorname{FSexp}(\mathcal{Z}(\mathcal{C})).
$$

On the other hand, it is clear by [\[NS07b,](#page-42-1) Corollary 7.8] that

$$
\operatorname{FSexp}(\mathcal{C}_\mathcal{M}^*) = \operatorname{FSexp}(\mathcal{Z}(\mathcal{C}_\mathcal{M}^*)) \qquad \text{ and } \qquad \operatorname{FSexp}(\mathcal{C}) = \operatorname{FSexp}(\mathcal{Z}(\mathcal{C}))
$$

both hold. Therefore one might conclude to obtain Equation (4.25) .

5. Applications: Indicators and exponents of semisimple Hopf algebras and its left partial dualization

In this section, our results are applied to the case the category $C = \text{Rep}(H)$ of representations over a semisimple Hopf algebra H over C, whose counit is denoted by ε , and the Sweedler notation is frequently used for the comultiplication.

5.1. Canonical pivotal structures of integral fusion categories, and indicators for semisimple Hopf algebras. Recall in [\[EGNO15,](#page-41-10) Definition 9.6.1] that a fusion category is said to be *integral* if the Frobenius-Perron dimensions of all the objects in C are integers. An integral fusion category is evidently *weakly integral*. Moreover, one could combine [\[ENO05,](#page-41-14) Propositions 8.25 and 8.23] to know that a fusion category over C is *pseudo-unitary* ([\[ENO05,](#page-41-14) Section 8.4]) and hence admits a unique spherical structure satisfying

$$
ptr(id_X) = FPdim(X) \qquad (\forall X \in \mathcal{C}), \tag{5.1}
$$

which is referred by $[NS08, Section 2]$ as the *canonical pivotal structure* of C (it is also called the canonical spherical structure in [\[EGNO15,](#page-41-10) Section 9.5]).

Lemma 5.1. Let C be a integral fusion category over $\mathbb C$ with canonical pivotal structure j. *Then:*

- (1) *Its center* Z(C) *is integral as well;*
- (2) *The spherical structure on* Z(C) *induced by* j *via [\(2.17\)](#page-11-4) is also the canonical pivotal structure.*
- *Proof.* (1) This is mentioned in the proof of [\[EGNO15,](#page-41-10) Proposition 9.6.11], as the forgetful functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ preserves Frobenius-Perron dimensions.
	- (2) Recall in [\(2.17\)](#page-11-4) that the spherical structure on $\mathcal{Z}(\mathcal{C})$ induced is the same as j. Thus for each object $\mathbf{V} = (V, \sigma^V) \in \mathcal{Z}(\mathcal{C})$, we find

$$
ptr(id_{\mathbf{V}}) = tr(j_{\mathbf{V}}) = tr(j_V) = FPdim(V) = FPdim(\mathbf{V}),
$$

where the last equality holds since the forgetful functor $\mathcal{Z}(\mathcal{C}) \to \mathcal{C}$ preserves Frobenius-Perron dimensions as well.

$$
\Box
$$

Proposition 5.2. *Let* C *be an integral fusion category over* C *with canonical pivotal structure* j*. Suppose* M *is an indecomposable* C*-module category which is semisimple. Then the indicators of any* $\mathbf{F} \in \mathcal{C}_{\mathcal{M}}^*$ *in the senses of Lemma [3.1](#page-12-5) and Definition [3.3](#page-12-1) coincide.*

Proof. Since C is an integral fusion category, we conclude by [\[ENO05,](#page-41-14) Theorem 8.35 and Proposition 8.24] that its dual fusion category $\mathcal{C}_{\mathcal{M}}^*$ will be also integral and hence pseudounitary with canonical structure j' (which is in fact spherical).

According to Lemma [5.1,](#page-27-2) the centers $\mathcal{Z}(\mathcal{C})$ and $\mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ are both pseudo-unitary with canonical pivotal structures j and j' respectively. It follows by [\[NS07a,](#page-41-3) Corollary 6.2] that Schauenburg's equivalence $\Omega : \mathcal{Z}(\mathcal{C}) \approx \mathcal{Z}(\mathcal{C}_{\mathcal{M}}^*)$ will preserve their canonical pivotal structures. Thus the claim desired is obtained with the help of Proposition [3.6.](#page-13-2) \Box

From now on, we would focus on the case when $\mathcal{C} = \mathsf{Rep}(K)$ is the category of a finite-dimensional quasi-Hopf algebra K over C. It is known in [\[ENO05,](#page-41-14) Theorem 8.33] that $\mathsf{Rep}(K)$ is an integral fusion category (with Frobenius-Perron dimensions of objects being their dimensions as vector spaces), and hence $\mathsf{Rep}(K)$ admits the canonical pivotal structure. Details could be found in [\[NS08\]](#page-42-4), where the notion of higher Frobenius-Schur indicators for a semisimple quasi-Hopf algebra over $\mathbb C$ is then defined as a consequence:

Definition 5.3. *Let* K *be a finite-dimensional semisimple quasi-Hopf algebra over* C*. Regard* Rep(K) *as the integral fusion category with the canonical pivotal structure.*

- (1) *(*[\[NS08,](#page-42-4) Definition 3.1]*) For any left* K*-module* V *, its* n*-th Frobenius-Schur indicator is defined to be* $\nu_n(V)$ *in the sense of Equation [\(3.1\)](#page-11-5), where* V *is regarded as an object in* Rep(K)*;*
- (2) *(*[\[NS07b,](#page-42-1) Proposition 5.3]*) The Frobenius-Schur exponent* FSexp(K) *of* K *is defined in the sense of* (4.23) *for the regular module* $K \in \text{Rep}(K)$ *, which coincides with* F Sexp(Rep(K)) *in the sense of* (4.24) *as well.*

Remark 5.4. *Suppose the semisimple complex quasi-Hopf algebra* K *is in fact a Hopf algebra. Then it is shown in* [\[NS08,](#page-42-4) Remark 3.4] *that the n-th indicator of* $V \in \text{Rep}(K)$ *defined above coincides with* [\[KSZ06,](#page-41-1) Definition 2.3] *for semisimple Hopf algebras:*

$$
\nu_n(V) = \chi_V\left(\sum \Lambda_{(1)}\Lambda_{(2)}\cdots \Lambda_{(n)}\right),\tag{5.2}
$$

where χ_V *is the character of the* K-module V, and Λ *is the integral of* K *with counit* 1.

5.2. Semisimple left partially dualized quasi-Hopf algebras and the reconstruction. Let H be a finite-dimensional Hopf algebra over a field k, and let $\mathsf{Rep}(H)$ be the category of finite-dimensional representations of H. A well-known result [\[AM07,](#page-40-5) Proposition 1.19] states that each indecomposable exact $\mathsf{Rep}(H)$ -module category has form $\mathsf{Rep}(B)$ for some indecomposable exact left H -comodule algebra B . When B is in particular a left coideal subalgebra of H , it is reconstructed in [\[Li23\]](#page-41-5) a quasi-Hopf algebra from the dual tensor category $\mathsf{Rep}(H)_{\mathsf{Rep}(B)}^*$.

In this subsection, we recall some of the relevant notions and results. The first one is [\[Li23,](#page-41-5) Definition 2.6] stated as follows, where only the case $C = H/B^{+}H$ is necessary in this paper (see [\[Li23,](#page-41-5) Remark 2.7] for details):

Definition 5.5. *Let* H *be a finite-dimensional Hopf algebra. Suppose* B *is a left coideal subalgebra with inclusion* $\iota : B \to H$, and denote by $\pi : H \to H/B^+H$ the projection (which *is a map right* H*-module coalgebras). Then the pair of* k*-linear diagrams*

$$
B \xleftarrow[\frac{\iota}{\zeta} \frac{\ast}{\tau}] \xleftarrow[\pi]{} H \xleftarrow[\pi]{} H \qquad and \qquad (H/B^+H)^* \xleftarrow[\pi]{} H^* \xleftarrow[\pi]{} H^* \xleftarrow[\pi]{} H^* \xleftarrow[\pi]{} H \xleftarrow
$$

is said to be a partially admissible mapping system for ι *, denoted by* (ζ, γ^*) *for simplicity, if all the conditions*

- (1) ζ *and* γ *have convolution inverses* $\overline{\zeta}$ *and* $\overline{\gamma}$ *respectively;*
- (2) ζ preserves left B-actions, and γ preserves right H/B^+H -coactions;
- (3) ζ *and* γ *preserve both the units and counits, meaning that*

$$
\zeta(1_H) = 1_B, \quad \varepsilon \mid_B \circ \zeta = \varepsilon, \quad \gamma[\pi(1_H)] = 1_H \quad \text{and} \quad \varepsilon \circ \gamma = \varepsilon_{H/B+H};
$$
\n
$$
(4) \ (\iota \circ \zeta) * (\gamma \circ \pi) = id_H,
$$

and the dual forms of (1) to (4) hold equivalently.

Remark 5.6. With notations in Definition [5.5,](#page-28-1) one could find that $\pi[\iota(b)] = \varepsilon(b)\pi(1)$ for all $b \in B$.

The partially admissible mapping system is not unique for a left coideal subalgebra ι : $B \hookrightarrow H$, but each one would determine a quasi-Hopf algebra denoted by $(H/B^+H)^* \# B$. We collect necessary descriptions of it as the following definition, which is a combination of some contents in [\[Li23,](#page-41-5) Theorem 3.1, Definition 3.3 and Remark 3.4].

For the purpose, note that $(H/B^+H)^*$ can be regarded as a right coideal subalgebra of H^* via π^* , and we would use notations for $b \in B$ and $f \in (H/B^+H)^*$ that

$$
\sum b_{(1)} \otimes b_{(2)} \in H \otimes B \quad \text{and} \quad \sum f_{(1)} \otimes f_{(2)} \in (H/B^{+}H)^{*} \otimes H^{*} \tag{5.4}
$$

to represent the structures of the left and right coideals (or comodules) respectively. Then we could also write equations

$$
\sum \iota(b)_{(1)} \otimes \iota(b)_{(2)} = \sum b_{(1)} \otimes \iota(b_{(2)}) \text{ and } \sum \pi^*(f)_{(1)} \otimes \pi^*(f)_{(2)} = \sum \pi^*(f_{(1)}) \otimes f_{(2)}.
$$
 (5.5)

Definition 5.7. *Let* H *be a finite-dimensional Hopf algebra. Suppose that* B *is a left coideal subalgebra of* H *with a partially admissible mapping system* $(ζ, γ^*)$ *. Then the left partially dualized quasi-Hopf algebra (or left partial dual)* (H/B⁺H) [∗]#B *determined by the partially admissible mapping system* $(ζ, γ^*)$ *is defined with the following structures:*

(1) As an algebra, $(H/B^+H)^* \# B$ is the smash product algebra with underlying vector space $(H/B^+H)^* \otimes B$: The multiplication is given by

$$
(f \# b)(g \# c) := \sum f(b_{(1)} \to g) \# b_{(2)} c \qquad (\forall f, g \in (H/B^+H)^*, \ \forall b, c \in B), \tag{5.6}
$$

and the unit element is $\varepsilon \#1$;

- (2) *The "comultiplication"* ∆ *and "counit" are described in* [\[Li23,](#page-41-5) Remark 3.4(2) and Definition 3.1(2)]*;*
- (3) The associator ϕ is the inverse of the element

$$
\phi^{-1} = \sum_{i,j} \left(\varepsilon \# \zeta [\gamma(x_i) \gamma(x_j)_{(1)}] \right) \otimes \left(x_i^* \# \zeta [\gamma(x_j)_{(2)}] \right) \otimes (x_j^* \# 1), \tag{5.7}
$$

where $\{x_i\}$ *is a linear basis of* H/B^+H *with dual basis* $\{x_i^*\}$ *of* $(H/B^+H)^*$;

(4) *The antipodes are described in* [\[Li23,](#page-41-5) Definition 3.1(4)]*.*

If the associator ϕ *is trivial, then the left partial dual* $(H/B^+H)^* \#B$ *is called a left partial dualized Hopf algebra.*

The *reconstruction theorem for left partial duals* should be remarked as [\[Li23,](#page-41-5) Theorem 4.22]. Specifically, a left partial dualized quasi-Hopf algebra $(H/B^+H)^* \# B$ reconstructs the dual tensor category $\mathsf{Rep}(H)_{\mathsf{Rep}(B)}^*$. On the other hand, since $\mathsf{Rep}(B)$ would be an indecompos-able and exact Rep(H)-module category, one could know by Lemma [2.13](#page-10-1) that $\mathsf{Rep}(H)_{\mathsf{Rep}(B)}^*$ would be fusion as long as $\mathsf{Rep}(H)$ is fusion in characteristic 0. In other words:

Corollary 5.8. *Let* H *be a semisimple Hopf algebra over an algebraically closed field* k *of characteristic* 0*. Then every left partial dualized quasi-Hopf algebra of* H *is semisimple.*

Finally, let us study some properties of left partial duals determined by more particular partially admissible mapping systems for subsequent uses. The following lemma is an analogue to [\[Li23,](#page-41-5) Lemma 6.3]:

Lemma 5.9. *Let* H *be a finite-dimensional Hopf algebra with left coideal subalgebra* B*. Suppose*

$$
B \xrightarrow{\iota} H \xrightarrow{\pi} H/B^+H \qquad and \qquad (H/B^+H)^* \xrightarrow{\pi^*} H^* \xrightarrow{\iota^*} B^*
$$

is a partially admissible mapping system for the inclusion $\iota : B \hookrightarrow H$ *. Then the left partial dual* $(H/B^+H)^* \# B$ *determined by* (ζ, γ^*) *is a Hopf algebra, if either of the following conditions hold:*

- (1) B *is a subbialgebra of* H*, and* ι*,* ζ *are both coalgebra maps;*
- (2) B^+H *is a bideal of H, and* π *,* γ *are both algebra maps;.*

Proof. We aim to prove that $(H/B+H)^* \#B$ is a Hopf algebra under the assumption (2) at first:

(2) Assume that π are γ are both algebra maps, which follow that

$$
\pi(xy) = \pi(x)\pi(y) \quad \text{and} \quad \gamma(xy) = \gamma(x)\gamma(y) \qquad (\forall x, y \in H/B^+H). \tag{5.8}
$$

It suffices to show that its associator ϕ is trivial. Our reasons are the equation

$$
\iota \circ \zeta = \mathrm{id}_{H} *(\overline{\gamma} \circ \pi) \in \mathrm{End}_{\mathbb{k}}(H) \tag{5.9}
$$

(which is equivalent to Definition [5.5\(](#page-28-1)4)) as well as the fact that γ is a right $H/B^{+}H^{-}$ comodule map:

$$
\sum \gamma(x)_{(1)} \otimes \pi[\gamma(x)_{(2)}] = \sum \gamma(x_{(1)}) \otimes x_{(2)} \quad (\forall x \in H/B^{+}H). \tag{5.10}
$$

In fact, recall in Equation [\(5.11\)](#page-30-0) that the inverse ϕ^{-1} of the associator could be written as

$$
\phi^{-1} = \sum_{i,j} \left(\varepsilon \# \zeta [\gamma(x_i) \gamma(x_j)_{(1)}] \right) \otimes \left(x_i^* \# \zeta [\gamma(x_j)_{(2)}] \right) \otimes \left(x_j^* \# 1 \right) \in \left((H/B^+ H)^* \# B \right)^{\otimes 3}, \tag{5.11}
$$

where $\{x_i\}$ is a linear basis of H/B^+H with dual basis $\{x_i^*\}$ of $(H/B^+H)^*$. Now we compute the image of ϕ^{-1} [\(5.11\)](#page-30-0) under the injection

 $(\mathrm{id} \otimes \iota) \otimes (\mathrm{id} \otimes \iota) \otimes (\mathrm{id} \otimes \mathrm{id}) : ((H/B^+H)^* \# B)^{\otimes 3} \to ((H/B^+H)^* \# H)^{\otimes 2} \otimes ((H/B^+H)^* \# B)$ as follows:

$$
\sum_{i,j} \left(\varepsilon \# \iota \left(\zeta [\gamma(x_i) \gamma(x_j)_{(1)}] \right) \right) \otimes \left(x_i^* \# \iota \left(\zeta [\gamma(x_j)_{(2)}] \right) \right) \otimes (x_j^* \# 1)
$$
\n
$$
\stackrel{(5.9)}{=} \sum_{i,j} \left(\varepsilon \# \gamma(x_i)_{(1)} \gamma(x_j)_{(1)} \overline{\gamma} \left(\pi [\gamma(x_i)_{(2)} \gamma(x_j)_{(2)}] \right) \right)
$$
\n
$$
\otimes \left(x_i^* \# \gamma(x_j)_{(2)} \overline{\gamma} \left(\pi [\gamma(x_i)_{(1)}] \right) \right) \otimes (x_j^* \# 1)
$$
\n
$$
\stackrel{(5.8)}{=} \sum_{i,j} \left(\varepsilon \# \gamma(x_i)_{(1)} \gamma(x_j)_{(1)} \overline{\gamma} \left(\pi [\gamma(x_i)_{(2)}] \pi [\gamma(x_j)_{(2)}] \right) \right)
$$
\n
$$
\otimes \left(x_i^* \# \gamma(x_j)_{(3)} \overline{\gamma} \left(\pi [\gamma(x_j)_{(4)}] \right) \right) \otimes (x_j^* \# 1)
$$
\n
$$
\stackrel{(5.10)}{=} \sum_{i,j} \left(\varepsilon \# \gamma(x_{i(1)}) \gamma(x_{j(1)}) \overline{\gamma}(x_{i(2)} x_{j(2)}) \right) \otimes \left(x_i^* \# \gamma(x_{j(3)}) \overline{\gamma}(x_{j(4)}) \right) \otimes (x_j^* \# 1)
$$
\n
$$
\stackrel{(5.8)}{=} \sum_{i,j} \left(\varepsilon \# \gamma(x_{i(1)} x_{j(1)}) \overline{\gamma}(x_{i(2)} x_{j(2)}) \right) \otimes \left(x_i^* \# \gamma(x_{j(3)}) \overline{\gamma}(x_{j(4)}) \right) \otimes (x_j^* \# 1)
$$

$$
32 \t\t K. LI
$$

$$
= \sum_{i,j} \left(\varepsilon \# \langle \varepsilon, x_{i(1)} x_{j(1)} \rangle 1_H \right) \otimes \left(x_i^* \# \langle \varepsilon, x_{j(2)} \rangle 1_H \right) \otimes (x_j^* \# 1)
$$

$$
= \left(\varepsilon \# 1_H \right) \otimes (\varepsilon \# 1_H) \otimes (\varepsilon \# 1),
$$

where $(H/B^+H)^* \# H$ denotes the vector space $(H/B^+H)^* \otimes H$. Therefore, ϕ is trivial since ι is an injection of algebras. It follows that the left partial dual $(H/B^+H)^* \#B$ is indeed a Hopf algebra.

(1) We know by [\[Li23,](#page-41-5) Proposition 2.10] that (γ^*, ζ) is a partially admissible mapping system for $\pi^*: (H/B^+H)^*$ ^{op cop} $\rightarrow H^{* \text{op cop}}$, and it determines the left partially dualized Hopf algebra B^* ^{op cop}# $(H/B^+H)^*$ ^{op cop} of H^* ^{op cop}. Moreover, according to [\[Li23,](#page-41-5) Proposition 5.1], there is an isomorphism

$$
(H/B^{+}H)^{*} \# B)^{\text{op cop}} \cong B^{* \text{op cop}} \# (H/B^{+}H)^{* \text{op cop}}, \quad f \# b \mapsto b \# f \tag{5.12}
$$

of quasi-Hopf algebras.

Now assume that B is a subbialgebra of H, and ι , ζ are both coalgebra maps. Then ι^* and ζ^* are algebra maps (between bialgebras $H^{* \text{op cop}}$ and $B^{* \text{op cop}}$). It would follow by (2) that the left partial dual $B^{* \text{op cop}} \# (H/B^+H)^{* \text{op cop}}$ has trivial associator, and we could find via the isomorphism (5.12) that $(H/B⁺H)[*] \#B$ is also a Hopf algebra with trivial associator.

 \Box

Lemma 5.10. *Let* H *be a finite-dimensional Hopf algebra. Suppose* B *is a subbialgebra of* H *such that* B⁺H *is a biideal of* H*. Then every left partial dualized quasi-Hopf algebra*

$$
(H/B^+H)^*\#B=(H/B^+H)^*\otimes B
$$

as an algebra. In particular, if B *is commutative and* H/B⁺H *is cocommutative, then* $(H/B+H)^*$ #B *is commutative.*

Proof. Note here that $\iota : B \rightarrow H$ and $\pi : H \rightarrow H/B^+H$ are bialgebra maps, and hence the right H^* -comodule structure of $(H/B^+H)^*$ and the left H-comodule structure of B are respectively:

$$
(H/B^+H)^* \rightarrow (H/B^+H)^* \otimes H^* \qquad \text{and} \qquad B \rightarrow H \otimes B f \rightarrow \sum f_{(1)} \otimes \pi^*(f_{(2)}) \qquad \text{and} \qquad b \rightarrow \sum \iota(b_{(1)}) \otimes b_{(2)},
$$

where the Sweedler notations stand for the comultiplications of respective Hopf algebras. Then we compute in the left partially dualized quasi-Hopf algebra $(H/B^+H)^* \# B$ with the formula [\(5.6\)](#page-29-0) that: For all $f, g \in (H/B^+H)^*$ and $b, c \in B$,

$$
(f \# b)(g \# c) = \sum f(\iota(b_{(1)}) \to g) \# b_{(2)} c = \sum f g_{(1)} \langle \pi^*(g_{(2)}), \iota(b_{(1)}) \rangle \# b_{(2)} c
$$

=
$$
\sum f g_{(1)} \langle g_{(2)}, \pi[\iota(b_{(1)})] \rangle \# b_{(2)} c \stackrel{\text{Remark 5.6}}{=} \sum f g_{(1)} \langle g_{(2)}, \varepsilon(b_{(1)}) \rangle \# b_{(2)} c
$$

=
$$
fg \# bc.
$$

Furthermore, if B is commutative and H/B^+H is cocommutative, then the tensor product $(H/B^+H)^* \otimes B$ of algebras becomes commutative, since $(H/B^+H)^*$ is commutative as well. \Box

In particular, this lemma holds for partially admissible mapping systems which are Hopf algebra extensions [\[Mas94,](#page-41-20) Definitions 1.3 and 2.4].

5.3. Relations between the indicators of a semisimple Hopf algebra and its left partial duals. This subsection is devoted to applying Theorem [4.9](#page-25-0) to the reconstruction $\mathsf{Rep}(H)_{\mathsf{Rep}(B)}^* \approx \mathsf{Rep}((H/B^+H)^* \# B)$ mentioned in the previous subsection, where H is a semisimple Hopf algebra over $\mathbb C$ with left coideal subalgebra B . Our result would be an equation on indicators of two related modules over $(H/B^+H)^* \#B$ and H respectively. At first we introduce the structures desired in the following, some of which are known in [\[Mas94\]](#page-41-20) for example:

Suppose L is a finite-dimensional left B-module, or $L \in \mathsf{Rep}(B)$ for simplicity. We denote by $(H/B^+H)^* \# L$ the left $(H/B^+H)^* \# B$ -module whose underlying vector space is $(H/B^+H)^* \otimes$ L, and its action is defined as

$$
[(H/B^{+}H)^{*} \# B] \otimes [(H/B^{+}H)^{*} \# L] \rightarrow (H/B^{+}H)^{*} \# L
$$
\n
$$
(f \# b) \otimes (g \# l) \rightarrow \sum f(b_{(1)} \rightarrow g) \# b_{(2)} l
$$
\n
$$
(5.13)
$$

with notations in Subsection [5.2.](#page-28-0) Furthermore, this follows a restricted action of the subalgebra $B \cong \varepsilon \# B$ on $(H/B^+H)^* \# L$, which induces then a right B^* -comodule structure

$$
[(H/B^{+}H)^{*}\#L] \to [(H/B^{+}H)^{*}\#L] \otimes B^{*}, \ f\#L \mapsto \sum (f\#L)_{(0)} \otimes (f\#L)_{(1)}
$$

satisfies that the equation $\sum (f \# l)_{(0)} \langle (f \# l)_{(1)}, b \rangle = (\varepsilon \# b)(f \# l) = \sum (b_{(1)} \to f) \# b_{(2)}l$ holds for each $b \in B$.

On the other hand, $L \in \mathsf{Rep}(B)$ becomes similarly a right B^* -comodule, such that the coaction

$$
L \to L \otimes B^*, \quad l \mapsto \sum l_{\langle 0 \rangle} \otimes l_{\langle 1 \rangle} \tag{5.14}
$$

satisfies that the equation $\sum l_{\langle 0 \rangle} \langle l_{\langle 1 \rangle}, b \rangle = bl$ holds for each $b \in B$. Meanwhile, if we regard H^* as a left B^* -comodule induced by the coalgebra map $\iota^*: H^* \to B^*$, then the *cotensor product* ([\[Tak77,](#page-42-12) Section 0] e.g.) of L and H^* could be defined as

$$
L\square_{B^*}H^* := \Big\{ \sum_i l_i \otimes h_i^* \in L \otimes H^* \mid \sum_i l_{i(0)} \otimes l_{i(1)} \otimes h_i^* = \sum_i l_i \otimes \iota^* (h_{i(1)}^*) \otimes h_{i(2)}^* \Big\}. \tag{5.15}
$$

Then $L\square_{B^*}H^*$ is a right H^* -comodule via the comultiplication on the latter tensorand H^* (similarly as $[Li23, Equation (4.14)]$), which induces naturally a left $H\text{-action}$:

$$
H \otimes (L\square_{B^*} H^*) \to L\square_{B^*} H^*, \ \ h \otimes \left(\sum_i l_i \otimes h_i^*\right) \mapsto \sum_i l_i \otimes (h \to h_i^*). \tag{5.16}
$$

Besides, $L\square_{B^*}H^*$ is a right $(H/B^+H)^*$ -module (also as in [\[Li23,](#page-41-5) Equation (4.14)]) with structure

$$
(L\square_{B^*}H^*)\otimes (H/B^+H)^* \to L\square_{B^*}H^*, \ \left(\sum_i l_i\otimes h_i^*\right)\otimes f \mapsto \sum_i l_i\otimes h_i^*\pi^*(h^*). \tag{5.17}
$$

Lemma 5.11. *With notations above, there is an isomorphism*

$$
\psi : (H/B^{+}H)^{*} \otimes (L\Box_{B^{*}}H^{*}) \cong [(H/B^{+}H)^{*} \# L] \Box_{B^{*}}H^{*},
$$
\n
$$
f \otimes \left(\sum_{i} l_{i} \otimes h_{i}^{*}\right) \mapsto \sum_{i} (f_{(1)} \# l_{i}) \otimes f_{(2)}h_{i}^{*}
$$
\n
$$
(5.18)
$$

 in $_{(H/B+H)*}$ Rep (H) _{$(H/B+H)*$}, where:

• $(H/B^+H)^* \otimes (L\square_{B^*}H^*) \in \mathsf{Rep}(H)$ *via the diagonal H-actions, and it is similar* to [\(5.16\)](#page-32-1)^{*f*} that $[(H/B+H)^*#L] \square_{B^*} H^* \in \text{Rep}(H)$ *via the hit* H-action on the last *tensorand* H[∗] *;*

• *The left and right* $(H/B^+H)^*$ -module structures on $(H/B^+H)^* \otimes (L\Box_{B^*}H^*)$ are de*termined on respective tensorands. However as for* $[(H/B+H)*H] \square_{B*} H^*$, *its right* $(H/B^+H)^*$ -action is the same as (5.17) , while the left $(H/B^+H)^*$ -action is diagonal *at the cotensor product (*[\[Li23,](#page-41-5) Equation (4.13)]*).*

Proof. Firstly, one might prove directly by the definition of the cotensor product (5.15) that

$$
\sum_{i} l_i \otimes h_i^* \in L \square_{B^*} H^* \quad \iff \quad \forall b \in B, \ \sum_{i} bl_i \otimes h_i^* = \sum_{i} l_i \otimes (h_i^* \leftarrow \iota(b)) \tag{5.19}
$$

for $l_i \in L$ and $h_i^* \in H^*$. Thus in order to show that ψ [\(5.18\)](#page-32-4) is well-defined, suppose $f \in (H/B^+H)^*$ and $\sum_i l_i \otimes h_i^* \in L\square_{B^*}H^*$. We compute for any $b \in B$ that

$$
\sum_{i} (\varepsilon \# b)(f_{(1)} \# l_{i}) \otimes f_{(2)} h_{i}^{*} \stackrel{(5.13)}{=} \sum_{i} [(b_{(1)} \to f_{(1)}) \# b_{(2)} l_{i}] \otimes f_{(2)} h_{i}^{*}
$$

\n
$$
\sum_{i} [(b_{(1)} \to f_{(1)}) \# l_{i}] \otimes f_{(2)} (h_{i}^{*} \gets \iota(b_{(2)}))
$$

\n
$$
= \sum_{i} (f_{(1)} \langle f_{(2)}, b_{(1)} \rangle \# l_{i}) \otimes f_{(3)} (h_{i}^{*} \gets \iota(b_{(2)}))
$$

\n
$$
= \sum_{i} (f_{(1)} \# l_{i}) \otimes (f_{(2)} \gets b_{(1)}) (h_{i}^{*} \gets \iota(b_{(2)}))
$$

\n
$$
\sum_{i} (f_{(1)} \# l_{i}) \otimes (f_{(2)} \gets \iota(b)_{(1)}) (h_{i}^{*} \gets \iota(b)_{(2)})
$$

\n
$$
= \sum_{i} (f_{(1)} \# l_{i}) \otimes (f_{(2)} h_{i}^{*} \gets \iota(b)),
$$

which means that $\sum_i (f_{(1)} \# l_i) \otimes f_{(2)} h_i^* \in L \square_{B^*} H^*$ according to [\(5.19\)](#page-33-0).

Now let us verify that ψ is a morphism in $(H/B+H)^*$ Rep $(H)(H/B+H)^*$: Suppose $f \in$ $(H/B^+H)^*$ and $\sum_i l_i \otimes h_i^* \in L\square_{B^*}H^*.$

• (ψ preserves left H-actions) For any $h \in H$,

$$
\psi\left(h \cdot \left[f \otimes \left(\sum_{i} l_{i} \otimes h_{i}^{*}\right)\right]\right) = \psi\left(\sum(h_{(1)} \rightarrow f) \otimes \left[\sum_{i} l_{i} \otimes (h_{(2)} \rightarrow h_{i}^{*})\right]\right)
$$

$$
= \sum_{i} \psi\left(f_{(1)} \otimes \left(\sum_{i} l_{i} \otimes h_{i}^{*}(1)\right) \langle f_{(2)} h_{i}^{*}(2), h \rangle\right)
$$

$$
= \sum_{i} (f_{(1)} \# l_{i}) \otimes f_{(2)} h_{i}^{*}(1) \langle f_{(3)} h_{i}^{*}(2), h \rangle
$$

$$
= \sum_{i} (f_{(1)} \# l_{i}) \otimes (h \rightarrow f_{(2)} h_{i}^{*})
$$

$$
= h \cdot \psi\left(f \otimes \left(\sum_{i} l_{i} \otimes h_{i}^{*}\right)\right);
$$

- (ψ preserves left $(H/B^+H)^*$ -actions) This is easy to see by the definition of ψ ;
- (ψ preserves right $(H/B^+H)^*$ -actions) This is direct to verify, as the right $(H/B^+H)^*$ module structures of both sides of [\(5.18\)](#page-32-4) are determined by the rightmost tensorand H^* via the injection π^* of algebras.

Finally, it is straightforward to verify that the linear map

$$
[(H/B^+H)^*\#L] \square_{B^*} H^* \rightarrow (H/B^+H)^* \otimes (L\square_{B^*} H^*),
$$

$$
\sum_i (f_i\#l_i) \otimes h_i^* \rightarrow \sum_i f_{i(1)} \otimes [l_i \otimes S(f_{i(2)})h_i^*]
$$

would be a left inverse of ψ , which implies that ψ is a monomorphism in $(H/B+H)^*$ **Rep** $(H)(H/B+H)^*$. However, one could find the following linear isomorphisms

$$
L\square_{B^*}H^*\cong L\square_{B^*}[B^*\otimes (H/B^+H)^*]\cong (L\square_{B^*}B^*)\otimes (H/B^+H)^*\cong L\otimes (H/B^+H)^* \quad (5.20)
$$

for every right B^* -comodule L, where the the first one is mentioned in [\[Li23,](#page-41-5) Equation (4.39)], and the second and third ones are mentioned in [\[Tak77,](#page-42-12) Section 0] as properties of cotensor products. It would imply that

$$
\dim \left([(H/B^{+}H)^{*} \# L] \Box_{B^{*}} H^{*} \right) \stackrel{(5.20)}{=} \dim \left([(H/B^{+}H)^{*} \# L] \otimes (H/B^{+}H)^{*} \right)
$$

\n
$$
= \dim \left((H/B^{+}H)^{*} \right) \dim \left(L \otimes (H/B^{+}H)^{*} \right)
$$

\n
$$
\stackrel{(5.20)}{=} \dim \left((H/B^{+}H)^{*} \right) \dim \left(L \Box_{B^{*}} H^{*} \right)
$$

\n
$$
= \dim \left((L \Box_{B^{*}} B^{*}) \otimes (H/B^{+}H)^{*} \right).
$$

Thus ψ is indeed an isomorphism as claimed.

Theorem 5.12. *Let* H *be a semisimple Hopf algebra over* C*. Suppose* B *is a left coideal* $subalgebra, and $(H/B^+H)^* \#B$ is a left partially dualized quasi-Hopf algebra of H. Then for$ $any L \in \mathsf{Rep}(B)$,

$$
\nu_n((H/B^+H)^*\#L) = \nu_n(L\square_{B^*}H^*) \qquad (\forall n \ge 1),
$$

where $(H/B^+H)^*\#L \in \text{Rep}((H/B^+H)^*\#B)$ and $L\square_{B^*}H^* \in \text{Rep}(H)$. (5.21)

Proof. Denote $\mathcal{M} = \mathsf{Rep}(H)_{(H/B^+H)^*}$, the category of right $(H/B^+H)^*$ -modules in $\mathsf{Rep}(H)$. This is an indecomposable $\mathsf{Rep}(H)$ -module category, which is exact and then semisimple. It is followed by [\[Mas94,](#page-41-20) Lemma 1.8] that $L\Box_{B^*}H^*$ becomes an object in $\mathsf{Rep}(H)_{(H/B^+H)^*}$ with the left H-module structure [\(5.16\)](#page-32-1) and right $(H/B+H)^*$ -module structure [\(5.17\)](#page-32-2).

It is mentioned in [\[Li23,](#page-41-5) Corollary 4.23] that the dual category $\mathsf{Rep}(H)^*_{\mathcal{M}}$ is tensor equivalent to $\mathsf{Rep}((H/B^+H)^*\#B)$. Let us describe a tensor equivalence

$$
\Psi : \mathsf{Rep}((H/B^+H)^* \# B) \approx \mathsf{Rep}(H)^*_{\mathcal{M}} \tag{5.22}
$$

in details, which is essentially the composition of $[Li23, (4.20)$ and $(4.5)]$: Specifically, for each $V \in \mathsf{Rep}((H/B^+H)^* \# B)$, we have

$$
\Psi(V)=-\otimes_{(H/B^+H)^*}(V\Box_{B^*}H^*)\in\operatorname{Rep}(H)^*_{\mathcal{M}}.
$$

Thus the tensor equivalence Ψ will send the object $(H/B^+H)^*\#L$ considered to

$$
\Psi((H/B^{+}H)^{*} \# L) = - \otimes_{(H/B^{+}H)^{*}} \left(\left[(H/B^{+}H)^{*} \# L \right] \square_{B^{*}} H^{*} \right)
$$
\n
$$
\stackrel{(5.18)}{\cong} - \otimes_{(H/B^{+}H)^{*}} (H/B^{+}H)^{*} \otimes (L\square_{B^{*}}H^{*})
$$
\n
$$
\cong - \otimes (L\square_{B^{*}}H^{*}) \in \text{Rep}(H)^{*}_{\mathcal{M}}.
$$
\n
$$
(5.23)
$$

We remark that $L\square_{B^*}H^*$ in [\(5.23\)](#page-34-2) is indeed an object in $\mathcal{M} = \mathsf{Rep}(H)_{(H/B^+H)^*}$, and the module structure of $-\otimes (L\square_{B^*}H^*)$ is the associativity constraint in $\mathsf{Rep}(H)$.

On the other hand, note according to [\[ENO05,](#page-41-14) Theorem 8.33] that $\mathsf{Rep}(H)$ (as well as $\mathsf{Rep}((H/B^+H)^*\#B))$ is an integral fusion category. By [\[ENO05,](#page-41-14) Theorem 8.35 and Proposition 8.24], its dual fusion category $\mathsf{Rep}(H)^*_{\mathcal{M}}$ will be also pseudo-unitary. It follows by [\[NS07a,](#page-41-3)

 36 K. LI

Corollary 6.2 that Ψ [\(5.22\)](#page-34-3) will preserves their canonical pivotal structures. Consequently, one could apply [\[NS07a,](#page-41-3) Corollary 4.4] to [\(5.23\)](#page-34-2) and obtain

$$
\nu_n((H/B^+H)^* \# L) = \nu_n(- \otimes (L\square_{B^*} H^*)) \qquad (\forall n \ge 1), \tag{5.24}
$$

where both indicators are in the sense of Lemma [3.1](#page-12-5) with respect to canonical pivotal structures.

However, we know by Proposition [5.2](#page-27-3) that the indicator $\nu_n(- \otimes (L\Box_{B^*} H^*))$ in Equation [\(5.24\)](#page-35-0) could be identified in the sense of Definition [3.3.](#page-12-1) Therefore due to Theorem [4.9,](#page-25-0) we have

$$
\nu_n(-\otimes (L\square_{B^*}H^*)) = \nu_n(L\square_{B^*}H^*) \qquad (\forall n \ge 1). \tag{5.25}
$$

At final, Equation [\(5.21\)](#page-34-4) is followed as a combination of Equations [\(5.24\)](#page-35-0) and [\(5.25\)](#page-35-1). \Box

We could point out two extreme situations for $L \in \mathsf{Rep}(B)$ being the regular module B and the trivial module C1:

Corollary 5.13. *Let* H *be a semisimple Hopf algebra over* C*. Suppose that* B *is a left coideal subalgebra of* H*. Then:*

(1) *The indicators of regular representations in* $\mathsf{Rep}((H/B^+H)^* \# B)$ *and* $\mathsf{Rep}(H)$ *coincide:*

$$
\nu_n((H/B^+H)^*\#B) = \nu_n(H) \qquad (\forall n \ge 1);
$$

(2) *The indicators of* $(H/B^+H)^*$ *regarded as objects in* $\mathsf{Rep}(H/B^+H)^* \#B$ *and* $\mathsf{Rep}(H)$ *coincide. Specifically, with the notations in* (5.4) *, the module structure on* $(H/B^+H)^* \in$ $\mathsf{Rep}((H/B^+H)^*\#B)$ *is*

$$
[(H/B^{+}H)^{*} \# B] \otimes (H/B^{+}H)^{*} \rightarrow (H/B^{+}H)^{*},
$$

\n
$$
(f \# b) \otimes g \rightarrow \sum f g_{(1)} \langle g_{(2)}, \iota(b) \rangle,
$$
\n
$$
(5.26)
$$

while the module structure on $(H/B^+H)^* \in \text{Rep}(H)$ *is*

$$
H \otimes (H/B^{+}H)^{*} \to (H/B^{+}H)^{*}, \quad h \otimes g \mapsto \sum g_{(1)}\langle g_{(2)}, h \rangle. \tag{5.27}
$$

Proof. (1) If we choose L to be the regular left B-module in Theorem [5.12,](#page-34-0) then the equation

> $\nu_n((H/B^+H)^*\#B) = \nu_n(B\Box_{B^*}H^*$ $(\forall n \geq 1)$

is obtained, where the module $(H/B^+H)^* \# B$ with structure as in (5.13) is exactly the regular representation in $\mathsf{Rep}((H/B^+H)^*\#B)$. Now let us focus on the left H-module $B\square_{B^*}H^*$:

Since the right B^* -coation on B is defined as in (5.14) , it is indeed isomorphic to the regular right B^{*}-comodule B^{*} via $b \mapsto \sum \varepsilon(b_{(0)})b_{(1)}$. Thus according to one of the canonical isomorphisms in $[Tak77, Section 0]$ on cotensor products, we might conclude a liner isomorphism

$$
B\square_{B^*}H^*\cong B^*\square_{B^*}H^*\cong H^*,\ \ \sum_i b_i\otimes h_i^*\mapsto \sum_i \varepsilon(b_i)h_i^*,
$$

by which the left H-action defined in [\(5.16\)](#page-32-1) is mapped to the hit action \rightarrow on H. However, it is known in [\[Mon93,](#page-41-21) Theorem 2.1.3(3)] that (H^*, \rightarrow) is also isomorphic to the regular left H-module. Therefore we find the equation

$$
\nu_n(B\Box_{B^*}H^*) = \nu_n(H) \qquad (\forall n \ge 1)
$$

on the indicators of objects in $\mathsf{Rep}(H)$, and hence the desired claim holds as a consequence.

(2) If we choose L to be the trivial left B-module C1 (induced by the counit ε restricted on B) in Theorem [5.12,](#page-34-0) then the equation

$$
\nu_n((H/B^+H)^*\#1) = \nu_n(\mathbb{C}1\square_{B^*}H^*) \qquad (\forall n \ge 1)
$$

is obtained. Here the left $(H/B^+H)^* \#B$ -module $(H/B^+H)^* \#1$ could be directly verified isomorphic to $(H/B^+H)^*$ with structure [\(5.26\)](#page-35-2). As for the left H-module $\mathbb{C}1\square_{B^*}H^*$, we could also use the isomorphism (5.20) to know that

$$
\mathbb{C}1\square_{B^*}H^* \cong \mathbb{C}1\square_{B^*}[B^*\otimes (H/B^+H)^*] \cong (\mathbb{C}1\square_{B^*}B^*)\otimes (H/B^+H)^* \cong \mathbb{C}1\otimes (H/B^+H)^* \cong (H/B^+H)^*, 1\otimes \pi^*(f) \leftarrow f.
$$

According to the latter equation in (5.5) , this isomorphism preserves left H -actions if $(H/B+H)^*$ is regarded to have the structure [\(5.27\)](#page-35-3).

 \Box

If the left coideal subalgebra B is furthermore chosen to be the trivial Hopf subalgebra $\mathbb{C}1$, then a semisimple case of [\[Shi12,](#page-42-5) Theorem 3.13(f)] or [\[Shi15,](#page-42-0) Proposition 3.4(1)] could be followed:

Corollary 5.14. *Let* H *be a semisimple Hopf algebra over* C*. Then*

$$
\nu_n(H^*) = \nu_n(H) \qquad (\forall n \ge 1).
$$

Proof. It is known in [\[Li23,](#page-41-5) Corollary 6.1(2)] that the dual Hopf algebra H^* is also a left partial dual of H . Thus the desired claim is implied by Corollary $5.13(1)$ $5.13(1)$.

 \Box

Corollary [5.13](#page-35-4) could be applied to show that the indicators of regular representations for the group $F \bowtie G$ defined by a matched pair $(F, G, \triangleright, \triangleleft)$ of finite groups as well as the bismash product $\mathbb{C}^G \# \mathbb{C} F$:

Corollary 5.15. *(*[\[KMN12,](#page-41-2) Corollary 5.7]*) Let* (F, G) *be a matched pair of finite groups. Then*

$$
\nu_n(\mathbb{C}^G \# \mathbb{C} F) = \nu_n(F \bowtie G) \qquad (\forall n \ge 1)?
$$

Proof. This could be shown according to [\[Li23,](#page-41-5) Proposition 6.8] that $\mathbb{C}^G \# \mathbb{C} F$ is a left partial dualized Hopf algebra of the group algebra $\mathbb{C}(F \bowtie G)$.

At the end of this subsection, let us mention a proposition on the Frobenius-Schur exponents as a consequence of Corollary [5.13\(](#page-35-4)1) as follows:

Proposition 5.16. *Let* H *be a semisimple Hopf algebra over* C*. Suppose that* B *is a left coideal subalgebra of* H*. Then*

$$
FSexp((H/B^+H)^*\#B) = FSexp(H).
$$

Proof. As Rep($(H/B^+H)^* \#B$) and Rep(H) are regarded as integral fusion categories with canonical pivotal structures, the pivotal traces [\(5.1\)](#page-27-4) of the identity morphisms on their regular modules would be

$$
\operatorname{ptr}(\operatorname{id}_{(H/B^+H)^* \# B}) = \operatorname{FPdim}((H/B^+H)^* \# B) = \dim((H/B^+H)^* \# B)
$$

and

$$
ptr(id_H) = FPdim(H) = dim(H),
$$

respectively. Moreover, it is implied by [\[Mas92,](#page-41-22) Theorem 2.1(6)] that $dim((H/B^+H)^*\#B)$ = $dim(H)$, and hence

$$
ptr(id(H/B+H)*#B) =ptr(idH)
$$
\n(5.28)

holds.

On the other hand, we know in Corollary [5.13\(](#page-35-4)1) that the regular modules $(H/B^+H)^* \#B$ and H have the same n-th indicators for each $n \geq 1$. We could combine the definition of Frobenius-Schur exponent [\(4.23\)](#page-26-3) and Equation [\(5.28\)](#page-37-1) to find that $\text{FSexp}((H/B^+H)^* \# B) =$ F Sexp (H) hold.

We remark according to [\[NS07b,](#page-42-1) Proposition 5.3] that: If $\mathcal C$ is a pseudo-unitary fusion category, its Frobenius-Schur exponent $\text{FSexp}(\mathcal{C})$ equals to the Frobenius-Schur exponent of an object containing every simple object of \mathcal{C} . The indicators of the regular object in \mathcal{C} were studied in [\[Shi12\]](#page-42-5). It should be worth comparing indicators between the regular objects of C and $\mathcal{C}_{\mathcal{M}}^*$. However, the indicators of the regular object in $\mathcal{C}_{\mathcal{M}}^*$ might not be defined analogously, since we do not know whether $\mathcal{C}_{\mathcal{M}}^*$ becomes pivotal.

5.4. Invariance for the exponent of semisimple (quasi-)Hopf algebras under partial **dualities.** Let C be a fusion category over \mathbb{C} . The *exponent* of C is defined in [\[Eti02,](#page-41-8) Remark] 6] as the order of the square of the braiding σ for its left center $\mathcal{Z}(\mathcal{C})$. Namely,

$$
\exp(\mathcal{C}) := \min\{n \ge 1 \mid \forall V, W \in \mathcal{Z}(\mathcal{C}), \ (\sigma_V^W \circ \sigma_W^V)^n = \mathrm{id}_V \otimes \mathrm{id}_W\}.
$$
 (5.29)

It is a finite positive integer, which is also related with the Frobenius-Schur exponent according to [\[NS07b,](#page-42-1) Section 6].

However in this subsection, we concentrate on the case for $exp(Rep(H))$ where H is a semisimple Hopf algebra over C, and finally show that a certain class of semisimple Hopf algebras satisfy the property in a conjecture by Kashina in [\[Kas99,](#page-41-6) [Kas00\]](#page-41-9).

Note at first that $\exp(\mathsf{Rep}(H))$ equals to

$$
\exp(H) := \min\big\{n \geq 1 \mid \forall h \in H, \sum h_{(1)}h_{(2)}\cdots h_{(n)} = \varepsilon(h)1_H\big\},\
$$

which is referred in [\[EG99\]](#page-41-7) to be the *exponent* of the semisimple Hopf algebra H (with involutory antipode). It is also a finite positive integer, and it divides $\dim(H)^3$ according to [\[EG99,](#page-41-7) Theorem 4.3]. Moreover in character 0, the number $\exp(H)$ is known to be the period of the sequence

$$
\nu_n(H), \quad n = 1, 2, \cdots \tag{5.30}
$$

of indicators (5.2) for the regular representation H. This is a consequence of the results in [\[KSZ06\]](#page-41-1), and some specific descriptions could be found in [\[NS07b,](#page-42-1) Section 1].

Besides, we have mentioned in Remark [5.4](#page-28-3) the fact stated in [\[NS08,](#page-42-4) Remark 3.4]: If the semisimple Hopf algebra H is regarded as a quasi-Hopf algebra, then the regular indicators $\nu_n(H)$ in the senses of Definition [5.3\(](#page-28-4)1) and Equation [\(5.2\)](#page-28-2) coincide. Furthermore, it follows by [\[NS07b,](#page-42-1) Proposition 5.3] that $FSexp(H)$ is indeed the period of the sequence [\(5.30\)](#page-37-2), since

the regular module H is clearly a projective generator of $\mathsf{Rep}(H)$. Thus we could conclude that:

Lemma 5.17. *(cf.* [\[NS07b\]](#page-42-1)*) Let* H *be a finite-dimensional semisimple Hopf algebra over* C*. Then its exponent and Frobenius-Schur exponent are equal. Namely,* $\exp(H) = \text{F}\text{S}\exp(H)$.

Then we could find the following consequence of Proposition [5.16,](#page-36-0) which might be an algebraic version of [\[Eti02,](#page-41-8) Proposition 6.3(2)] for the left partial dualization:

Corollary 5.18. *(cf.* [\[Eti02,](#page-41-8) Proposition 6.3(2)]*) Let* H *be a semisimple Hopf algebra over* C*. Suppose that* B *is a left coideal subalgebra of* H*, and* (H/B⁺H) [∗]#B *is a left partial dualized Hopf algebra of* H*. Then*

$$
\exp((H/B^+H)^*\#B) = \exp(H).
$$

Note in [\[Li23,](#page-41-5) Proposition 6.8] that for a matched pair (F, G) of finite groups, the bismash product $\mathbb{C}^{\tilde{G}}$ # $\mathbb{C}F$ is a left partial dualized Hopf algebra of the group algebra $\mathbb{C}(F \bowtie G)$. One might apply Corollary [5.18](#page-38-0) to obtain the result [\[LMS06,](#page-41-23) Proposition 3.1] over C:

Corollary 5.19. *([\[LMS06,](#page-41-23) Proposition 3.1]) For a matched pair* $(F, G, \triangleright, \triangleleft)$ *of finite groups, we have*

$$
\exp(\mathbb{C}^G \# \mathbb{C} F) = \exp(\mathbb{C}(F \bowtie G)) = \exp(F \bowtie G),\tag{5.31}
$$

where $\exp(F \bowtie G)$ *denotes the exponent of the group* $F \bowtie G$ *.*

In fact, the bismash product Hopf algebra $\mathbb{C}^G \# \mathbb{C} F$ arises from the *split abelian extension* (whose definition is provided in the next paragraph)

$$
0 \to \mathbb{C}^G \to \mathbb{C}^G \# \mathbb{C} F \to \mathbb{C} F \to 0.
$$

Conversely, every semisimple split abelian extension

$$
0 \to B \to H \to C \to 0
$$

of Hopf algebras over $\mathbb C$ determines a matched pair $(F, G, \triangleright, \triangleleft)$ of finite groups, such that $B \cong \mathbb{C}^G$, $C \cong \mathbb{C}F$ and $H \cong \mathbb{C}^G \# \mathbb{C}F$. The relevant results are concluded in [\[Mas02\]](#page-41-24).

Let us recall the definition of abelian extensions with our notations for convenience:

Definition 5.20. *(cf.* [\[Sch02,](#page-42-13) Definition 6.5.2]*) Let* H*,* B *and* C *be Hopf algebras. Then an extension*

$$
0 \to B \xrightarrow{\iota} H \xrightarrow{\pi} C \to 0 \tag{5.32}
$$

of Hopf algebras is said to be split, if there exist a left B-module coalgebra map $\zeta : H \to B$ *and a right* C-comodule algebra map $\gamma : C \to H$, such that $(\iota \circ \zeta) * (\gamma \circ \pi) = id_H$.

Furthermore, we say that the extension [\(5.32\)](#page-38-1) is abelian, if B *is commutative and* C *is cocommutative.*

Remark 5.21. *With the notations in Definition* [5.20,](#page-38-2) *it is clear that* ζ *preserves the counits, and* γ *preserves the units. Moreover according to the arguments in the proof of* [\[DT86,](#page-40-6) Theorem 9] *and* [\[Mas92,](#page-41-22) Lemma 2.15]*, we could assume that* ζ *and* γ *both preserve the units and the counits since* ζ(1) *must be a group-like element. Details are also found in the paragraphs before* [\[Li23,](#page-41-5) Lemma 2.5]*.*

Consequently, a split extension [\(5.32\)](#page-38-1) of finite-dimensional Hopf algebras follows a partially admissible mapping system (ζ, γ^*) *of ι*, *such that* ζ *is a left* B-module coalgebra map and γ *is a right* C*-comodule algebra map.*

On the other hand, Kashina's conjecture on the exponent states that:

Conjecture 5.22. *(*[\[Kas99,](#page-41-6) [Kas00\]](#page-41-9)*) Let* H *be a semisimple and cosemisimple Hopf algebra over a field. Then* $\exp(H) \mid \dim(H)$.

Thus, one could conclude by Equation [\(5.31\)](#page-38-3) to know that: $\exp(H) | \dim(H)$ holds when H is a semisimple split abelian extension Hopf algebra over \mathbb{C} , as the exponent of a finite group must divide its order. Now we try to generalize this result a little bit, where C in (5.32) is replaced by the quotient bialgebra H/B^+H without the loss of generality:

Theorem 5.23. *Let* H *be a semisimple Hopf algebra over* C *with a commutative subbialgebra* B*. Suppose that* B⁺H *is a biideal such that* H/B⁺H *is cocommutative, and that there is a partially admissible mapping system*

$$
B \xrightarrow{\iota} H \xrightarrow{\pi} H/B^+H \qquad and \qquad (H/B^+H)^* \xrightarrow{\pi^*} H^* \xrightarrow{\iota^*} B^*
$$

where ι *and* π *are bialgebra maps. Then*

$$
\exp(H) | \dim(H)
$$

if either of the following conditions hold:

- (1) ζ *is a coalgebra map, or*
- (2) γ *is an algebra map.*

Proof. We could conclude by Lemma [5.9](#page-30-4) that the left partial dual $(H/B^+H)^* \#B$ is in fact a Hopf algebra, if either conditions (1) or (2) holds. Then it follows by Corollary [5.18](#page-38-0) that $\exp((H/B^+H)^*\#B) = \exp(H)$ as a consequence.

On the other hand, as B is commutative and H/B^+H is cocommutative, we know by Lemma [5.10](#page-31-1) that $(H/B^+H)^* \# B$ is a commutative semisimple Hopf algebra, which is isomorphic to the dual Hopf algebra of a finite group algebra. Therefore,

 $\exp((H/B^+H)^*\#B)$ | dim $((H/B^+H)^*\#B)$.

However, it is clear by [\[Li23,](#page-41-5) Corollary 2.8] that $\dim((H/B^+H)^*\#B) = \dim(H)$. Thus $\exp(H)$ | dim(H) holds as a conclusion.

Remark 5.24. *For semisimple Hopf algebras admitting non-split abelian extensions, the indicators are computed in* [\[Shi12,](#page-42-5) Section 4.2]*, and there are results in* [\[Nat07,](#page-41-25) Section 5.2] *on the upper bounds of the exponent.*

This theorem could be applied to give a slightly analogous result to [\[Kas03,](#page-41-26) Theorem 6.3] for the case over C:

Proposition 5.25. *Let* H *be a semisimple Hopf algebra over* C*. Suppose that* H *has a commutative Hopf subalgebra B of index p, which is the smallest prime divisor of* $dim(H)$. *If* $B \cong \mathbb{C}^G$ *as Hopf algebras for some group G whose Sylow subgroups are all cyclic, then* $\exp(H)$ | dim (H) .

Proof. We know by [\[KM97,](#page-41-27) Proposition 2] that B is normal in H , and hence the quotient $H/B+H$ is a Hopf algebra of dimension p. Then it follows by [\[Zhu94,](#page-42-14) Theorem 2] that H/B^+H is the group algebra $\mathbb{C}C_p$ of a cyclic group C_p with order p.

Now we consider the abelian extension

$$
0 \to \mathbb{C}^G \to H \to \mathbb{C}C_p \to 0,\tag{5.33}
$$

and apply some results in [\[Mas08,](#page-41-28) Page 444] (for example). At first, the extension [\(5.33\)](#page-39-1) gives rise to a matched pair $(C_p, G, \triangleright, \triangleleft)$ of finite groups, where the action \triangleright must be trivial due to [\[Mas00,](#page-41-29) Lemma 1.1]. Furthermore, such a matched pair would determine a weak action

$$
\succ: C_p \times \mathbb{C}^G \to C_p, \quad (x, e_g) \mapsto e_{g \triangleleft x^{-1}} \qquad (\forall g \in G)
$$

and a trivial weak coaction. On the other hand, it is known by [\[Nat99,](#page-41-30) Proposition 2.6] and our assumption that H is in fact isomorphic to a crossed product Hopf algebra $\mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p$ (with trivial dual cocycle), where $\sigma : \mathbb{C}C_p \otimes \mathbb{C}C_p \to \mathbb{C}^G$ is a cocycle corresponds to a normalized 2-cocycle.

In conclusion, the extension [\(5.33\)](#page-39-1) is equivalent to

$$
0 \to \mathbb{C}^G \xrightarrow{\iota} \mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p \xrightarrow{\pi} \mathbb{C}C_p \to 0.
$$

Here,

$$
\iota = \mathrm{id}_{\mathbb{C}^G} \otimes 1_{\mathbb{C}C_p}, \qquad \pi = \varepsilon_{\mathbb{C}^G} \otimes \mathrm{id}_{\mathbb{C}C_p},
$$

and the operations on the Hopf algebra $\mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p$ are given as follows: For any $b, c \in \mathbb{C}^G$ and $x, y \in \mathbb{C}C_p$,

- Multiplication: $(b \rtimes x)(c \rtimes y) = \sum b(x_{(1)} \succ c)\sigma(x_{(2)}, y_{(1)}) \rtimes x_{(3)}y_{(2)};$
- Unit element: $1_{\mathbb{C}^G} \rtimes 1_{\mathbb{C}C_p}$;
- Comultiplication: $b \rtimes x \mapsto \sum (b_{(1)} \rtimes x_{(1)}) \otimes (b_{(2)} \otimes x_{(2)})$;
- Counit element: $\varepsilon_{\mathbb{C}^G} \otimes \varepsilon_{\mathbb{C}C_p}$.

These structures are introduced in [\[AD95,](#page-40-7) Equations (2.5) and (2.15)] for the situation when the weak coaction and the dual cocycle are both trivial.

Consequently, it is straightforward to show that

$$
\zeta : \mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p \to \mathbb{C}^G, \quad b \rtimes x \mapsto b\varepsilon(x)
$$

is a biunitary left \mathbb{C}^G -module coalgebra map with convolution inverse $\overline{\zeta} = S_{\mathbb{C}^G} \otimes \varepsilon_{\mathbb{C}C_p}$. Thus we could formulate a partially admissible mapping system

$$
\mathbb{C}^G \xrightarrow[\zeta]{\iota} \mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p \xrightarrow[\gamma]{\pi} \mathbb{C}C_p \quad \text{and} \quad \mathbb{C}^{C_p} \xrightarrow[\gamma]{\pi^*} (\mathbb{C}^G \rtimes_{\sigma} \mathbb{C}C_p)^* \xrightarrow[\zeta^*]{\iota^*} \mathbb{C}G
$$

satisfying the conditions in Theorem [5.23\(](#page-39-0)1), and hence $\exp(H)$ | dim(H) holds.

Acknowledgement

The author is extremely grateful to Professor Kenichi Shimizu, who provided the motivation and question, as well as taught author the methods for this paper with so much patience. The author would also like to thank Professor Gongxiang Liu for useful discussions and suggestions.

REFERENCES

- [AD95] N. Andruskiewitsch, J. Devoto, Extensions of Hopf algebras. Algebra i Analiz 7 (1995), no. 1, 22-61. [AM07] N. Andruskiewitsch, J. M. Mombelli, On module categories over finite-dimensional Hopf algebras. J. Algebra 314 (2007), no. 1, 383-418.
- [BK01] B. Bakalov, A. Kirillov Jr. Lectures on tensor categories and modular functors. University Lecture Series, 21. American Mathematical Society, Providence, RI, 2001. x+221 pp.

[BW99] J. W. Barrett, B. W. Westbury, Spherical categories. Adv. Math. 143 (1999), no. 2, 357-375.

[BV12] A. Bruguières, A. Virelizier, Quantum double of Hopf monads and categorical centers. Trans. Amer. Math. Soc. 364 (2012), no. 3, 1225-1279.

[DT86] Y. Doi, M. Takeuchi, Cleft comodule algebras for a bialgebra. Comm. Algebra 14 (1986), no. 5, 801-817.

- [Eti02] P. Etingof, On Vafa's theorem for tensor categories. Math. Res. Lett. 9 (2002), no. 5-6, 651-657.
- [EG99] P. Etingof and S. Gelaki, On the exponent of finite-dimensional Hopf algebras. Math. Res. Lett. 6 (1999), no. 2, 131-140.
- [EGNO15] P. Etingof, S. Gelaki, D. Nikshych, V. Ostrik, Tensor Categories. Mathematical Surveys and Monographs, 205. American Mathematical Society, Providence, RI, 2015. xvi+343 pp.
- [ENO04] P. Etingof, D. Nikshych, V. Ostrik, An analogue of Radford's $S⁴$ formula for finite tensor categories. Int. Math. Res. Not. 2004, no. 54, 2915-2933.
- [ENO05] P. Etingof, D. Nikshych, V. Ostrik, On fusion categories. Ann. of Math. (2) 162 (2005), no. 2, 581-642.
- [EO04] P. Etingof, V. Ostrik, Finite tensor categories. Mosc. Math. J. 4 (2004), no. 3, 627-654, 782-783.
- [FS23] J. Fuchs, C. Schweigert, Internal natural transformations and Frobenius algebras in the Drinfeld center. Transform. Groups 28 (2023), no. 2, 733-768.
- [JS91] A. Joyal and R. Street. Tortile Yang-Baxter operators in tensor categories. J. Pure Appl. Algebra 71 (1991), no. 1, 43-51.

[JS93] A. Joyal and R. Street. *Braided tensor categories*. Adv. Math. 102 (1993), no. 1, 20-78.

- [Kas99] Y. Kashina, On the order of the antipode of Hopf algebras in ${}_H^H YD$. Comm. Algebra 27 (1999), no. 3, 1261-1273.
- [Kas00] Y. Kashina, A generalized power map for Hopf algebras. Hopf algebras and quantum groups (Brussels, 1998), 159-175, Lecture Notes in Pure and Appl. Math., 209, Dekker, New York, 2000.
- [Kas03] Y. Kashina, On semisimple Hopf algebras of dimension 2m. Algebr. Represent. Theory 6 (2003), no. 4, 393-425.
- [KMN12] Y. Kashina, S. Montgomery, S.-H. Ng, On the trace of the antipode and higher indicators. Israel J. Math. 188 (2012), 57-89.
- [KSZ06] Y. Kashina, Y. Sommerhäuser, Y. Zhu, On higher Frobenius-Schur indicators. Mem. Amer. Math. Soc. 181 (2006), no. 855, viii+65 pp.
- [KM97] T. Kobayashi, A. Masuoka, A result extended from groups to Hopf algebras. Tsukuba J. Math. 21 (1997), no. 1, 55-58.
- [LMS06] R. Landers, S. Montgomery, P. Schauenburg, Hopf powers and orders for some bismash products. J. Pure Appl. Algebra 205 (2006), no. 1, 156-188.
- [Li23] K. Li, Partially dualized quasi-Hopf algebras reconstructed from dual tensor categories to finitedimensional Hopf algebras. preprint, arXiv:2309.04886.
- [LM00] V. Linchenko, S. Montgomery, A Frobenius-Schur theorem for Hopf algebras. Algebr. Represent. Theory 3 (2000), no. 4, 347-355.
- [MacL63] S. Mac Lane. Natural associativity and commutativity. Natural associativity and commutativity. Rice Univ. Stud. 49 (1963), no. 4, 28-46.
- [MacL98] S. Mac Lane, Categories for the Working Mathematician. Second edition. Graduate Texts in Mathematics, 5. Springer-Verlag, New York, 1998. xii+314 pp.
- [MN05] G. Mason, S.-H. Ng. Central invariants and Frobenius-Schur indicators for semisimple quasi-Hopf algebras. Adv. Math. 190 (2005), no. 1, 161-195.
- [Mas92] A. Masuoka, Freeness of hopf algebras over coideal subalgebras. Comm. Algebra 20 (1992), no. 5, 1353-1373.
- [Mas94] A. Masuoka, *Coideal subalgebras in finite Hopf algebras*. J. Algebra 163 (1994), no. 3, 819-831.
- [Mas00] A. Masuoka, Cocycle deformations and Galois objects for some cosemisimple Hopf algebras of finite dimension. New trends in Hopf algebra theory (La Falda, 1999), 195-214, Contemp. Math., 267, Amer. Math. Soc., Providence, RI, 2000.
- [Mas02] A. Masuoka, Hopf algebra extensions and cohomology. New directions in Hopf algebras, 167-209, Math. Sci. Res. Inst. Publ., 43, Cambridge Univ. Press, Cambridge, 2002.
- [Mas08] A. Masuoka, *Classification of semisimple Hopf algebras*. Handbook of algebra. Vol. 5, 429-455, Handb. Algebr., 5, Elsevier/North-Holland, Amsterdam, 2008.
- [Mon93] S. Montgomery, *Hopf algebras and their actions on rings*. CBMS Regional Conference Series in Mathematics, 82. Published for the Conference Board of the Mathematical Sciences, Washington, DC; by the American Mathematical Society, Providence, RI, 1993. xiv+238 pp.
- [Müg03] M. Müger, From subfactors to categories and topology. II. The quantum double of tensor categories and subfactors. J. Pure Appl. Algebra 180 (2003), no. 1-2, 159-219.
- [Nat99] S. Natale, On semisimple Hopf algebras of dimension pq^2 . J. Algebra 221 (1999), no. 1, 242-278.
- [Nat07] S. Natale, On the exponent of tensor categories coming from finite groups. Israel J. Math. 162 (2007), 253-273.
- [NS07a] S.-H. Ng, P. Schauenburg, Higher Frobenius-Schur indicators for pivotal categories. Hopf algebras and generalizations, 63-90, Contemp. Math., 441, Amer. Math. Soc., Providence, RI, 2007.
- [NS07b] S.-H. Ng, P. Schauenburg, Frobenius-Schur indicators and exponents of spherical categories. Adv. Math. 211 (2007), no. 1, 34-71.
- [NS08] S.-H. Ng, P. Schauenburg, Central invariants and higher indicators for semisimple quasi-Hopf algebras. Trans. Amer. Math. Soc. 360 (2008), no. 4, 1839-1860.
- [Ost03] V. Ostrik, Module categories, weak Hopf algebras and modular invariants. Transform. Groups 8 (2003), no. 2, 177-206.
- [Sch01] P. Schauenburg, *The monoidal center construction and bimodules.* J. Pure Appl. Algebra 158 (2001), no. 2-3, 325-346.
- [Sch02] P. Schauenburg, Hopf bimodules, coquasibialgebras, and an exact sequence of Kac. Adv. Math. 165 (2002), no. 2, 194-263.
- [Sch04] P. Schauenburg, On the Frobenius-Schur indicators for quasi-Hopf algebras. J. Algebra 282 (2004), no. 1, 129-139.
- [Shi12] K. Shimizu, Some computations of Frobenius-Schur indicators of the regular representations of Hopf algebras. Algebr. Represent. Theory 15 (2012), no. 2, 325-357.

[Shi15] K. Shimizu, On indicators of Hopf algebras. Israel J. Math. 207 (2015), no. 1, 155-201.

- [Shi17a] K. Shimizu, On unimodular finite tensor categories. Int. Math. Res. Not. IMRN 2017, no. 1, 277-322. [Shi17b] K. Shimizu, The monoidal center and the character algebra. J. Pure Appl. Algebra 221 (2017), no. 9, 2338-2371.
- [Shi19] K. Shimizu, Integrals for finite tensor categories. Algebr. Represent. Theory 22 (2019), no. 2, 459-493.
- [Shi20] K. Shimizu, Further results on the structure of (co)ends in finite tensor categories. Appl. Categ. Structures 28 (2020), no. 2, 237-286.
- [Shi24] K. Shimizu, Quasi-Frobenius algebras in finite tensor categories. preprint, arXiv:2402.02929.
- [Tak77] M. Takeuchi, Morita theorems for categories of comodules. J. Fac. Sci. Univ. Tokyo Sect. IA Math. 24 (1977), no. 3, 629-644.
- [Zhu94] Y. Zhu, Hopf algebras of prime dimension. Internat. Math. Res. Notices 1994, no. 1, 53-59.

School of Mathematics, Hangzhou Normal University, Hangzhou 311121, China

Email address: kqli@hznu.edu.cn