QUASI-SHADOWING PROPERTY FOR NONUNIFORMLY PARTIALLY HYPERBOLIC SYSTEMS

GANG LIAO AND XUETONG ZU

Abstract

In this paper, we establish a new quasi-shadowing property for any nonuiformly partially hyperbolic set of a $C^{1+\alpha}$ diffeomorphism, which is adaptive to the movement of the pseudo-orbit. Moreover, the quasi-specification property and quasi-closing property are also investigated. As an application of quasi-closing property, we extend Katok's reslut on the growth of periodoc orbits for hyperbolic ergodic measure to any ergodic measure: the number of quasi-periodic points grows exponentially at least the metric entropy.

1. INTRODUCTION

The mathematical models used in practical applications are often simplified compared to reality, yet still need to be effective, thus such properties that remain unchanged or can be approxianted to some extent under small perturbations attract people usually. In this paper, we are concerned with the approximation mechanism of pseudo-orbits by real orbits, i.e., the shadowing property of dynamical system. The shadowing property is a poweful tool in the structrual anlysis of differentiable dynamical system, see Liao [16], Mañé [17], Wen [22] and Gan [5] for the investigation of stability, and Bowen [3], Katok [10] and Sarig [19] for the investigation of complexity.

(1) Shadowing for uniformly hyperbolic systems. Throughout, let M be a compact boundaryless Riemannian manifold and, $\|\cdot\|$ and d be the metric and norm induced by the Riemannian metric. Let f be a diffeomorphism on M and Λ be a uniformly hyperbolic set: there exist an Df-invariant splitting $T_{\Lambda}M = E^s \oplus E^u$, constants $C \ge 1$ and $0 < \lambda < 1$ such that

(i) $||D_x f^n(v)|| \le C\lambda^n ||v||, \quad \forall x \in \Lambda, v \in E^s(x);$

(ii) $||D_x f^{-n}(v)|| \le C\lambda^n ||v||, \quad \forall x \in \Lambda, v \in E^u(x).$

For $\delta > 0$, we refer to $\{x_n\}_{n \in \mathbb{Z}}$ as a δ pseudo-orbit if $d(f(x_n), x_{n+1}) < \delta$, $n \in \mathbb{Z}$. It holds the following shadowing property (Anosov [1], Bowen [4] and Sinai [21]): For any $\varepsilon > 0$, there exists $\delta > 0$ such that for any δ pseudo-orbit $\{x_n\}_{n \in \mathbb{Z}} \subset \Lambda$, there exists a point x such that

$$d(x_n, f^n(x)) < \varepsilon, \ n \in \mathbb{Z}.$$

Besides, there exists $\varepsilon_0 > 0$ such that if $\varepsilon < \varepsilon_0$, the shadowing point x is unique. By considering periodic pseudo-orbit, there is closing property: for any $\varepsilon > 0$, there exists $\beta > 0$ such that for any x with $d(x, f^p(x)) < \beta$ for some $p \in \mathbb{Z}^+$, there exists a point z with $f^p(z) = z$ such that

$$d(x,z) < \varepsilon.$$

Gan [6] proved a generalized shadowing lemma for a closed invariant set with a continuous invariant splitting. Wen et al. [23] proved that the chain component that contains a hyperbolic periodic point is hyperbolic if it is C^1 -stably shadowable.

Key words and phrases. Nonuiformly partially hyperbolic set, quasi-shadowing property, the growth of quasi-periodic points.

²⁰¹⁰ Mathematics Subject Classification. 37C50, 37D25, 37D30.

School of Mathematical Sciences, Center for Dynamical Systems and Differential Equations, Soochow University, Suzhou 215006, China; This research was partially supported by National Key R&D Program of China (2022YFA1005802), NSFC (12071328, 12122113) and BK20211595.

(2) Shadowing for nonuniformly hyperbolic systems. For nonuniformly hyperbolic systems, the hyperbolicity is not uniform, thus the scale of shadowing property needs to be adjusted to the level of hyperbolicity.

Let f be a $C^{1+\alpha}$ diffeomorphism on M. Given positive numbers λ , μ and ε with $\lambda, \mu \gg \varepsilon$ and a positive integer k, define $\Lambda_k = \Lambda_k(\lambda, \mu, \varepsilon)$ to be all points $x \in M$ for which there is an Df-invariant decomposition $T_{\operatorname{orb}(x)}M = E^s \oplus E^u$ satisfying

- (i) $\|Df^n(v)\| \le e^{\varepsilon k} e^{-(\lambda-\varepsilon)n} e^{\varepsilon|m|} \|v\|, \ \forall n \ge 0, m \in \mathbb{Z}, v \in E^s(f^m(x));$
- (ii) $\|Df^{-n}(v)\| \le e^{\varepsilon k} e^{-(\mu-\varepsilon)n} e^{\varepsilon |m|} \|v\|, \ \forall n \ge 0, m \in \mathbb{Z}, v \in E^u(f^m(x)).$

We refer to Λ_k as a hyperbolic block or Pesin block [18]. It is veried that $\Lambda_k \subseteq \Lambda_{k+1}$ from the definition. The hyperbolicity of Λ_k is weaker as k becomes larger. Denote $\Lambda = \bigcup_{k>1} \Lambda_k(\lambda, \mu, \varepsilon)$

and we call Λ a nonuniformly hyperbolic set. One may see that if there is some ergodic hyperbolic measure, then we can get a nonuniformly hyperbolic set of full measure by taking λ as the absolute value of the maximal negative Lyapunov exponent and μ as the minimum positive Lyapunov exponent.

For a sequence of numbers $\{\delta_k\}_{k\in\mathbb{Z}^+}$, if there exist a sequence of positive integers $\{s_n\}_{n\in\mathbb{Z}}$ and a sequence of points $\{x_n\}_{n\in\mathbb{Z}} \subset \Lambda$ such that

- (a) $x_n \in \Lambda_{s_n}, \forall n \in \mathbb{Z};$
- (b) $|s_n s_{n+1}| \leq 1, \forall n \in \mathbb{Z};$
- (c) $d(f(x_n), x_{n+1}) < \delta_{s_n}, \forall n \in \mathbb{Z},$

we refer to $\{x_n\}_{n\in\mathbb{Z}}\subset\Lambda$ as a $\{\delta_k\}_{k\in\mathbb{Z}^+}$ pseudo-orbit of f.

Katok [10] proved the following shadowing property: for any $\eta > 0$, there exists a sequence of positive numbers $\{\delta_k\}_{k \in \mathbb{Z}^+}$ such that for any $\{\delta_k\}_{k \in \mathbb{Z}^+}$ pseudo-orbit $\{x_n\}_{n \in \mathbb{Z}} \subset \Lambda$, there exists a point x satisfying:

$$d(x_n, f^n(x)) < \eta \epsilon_{s_n},$$

where $\epsilon_{s_n} = \epsilon_0 e^{-s_n \cdot \frac{\varepsilon}{\alpha}}$ and ϵ_0 is a number independent of s_n . Besides, if $\eta < 1$, the shadowing point x is unique. There is also closing property according to the scale of hyperbolic block Λ_k . For each $k \in \mathbb{Z}^+$ and any $\eta > 0$, there exists a positive number $\beta = \beta(k, \eta)$ such that for any point $x \in \Lambda_k$ with $d(x, f^p(x)) < \beta$, there exists a point z with $f^p(z) = z$ such that:

 $d(x,z) < \eta.$

(3) Quasi-shadowing for partially hyperbolic systems.

Comparing to hyperbolic systems, the partially hyperbolic systems has additional central direction. The shadowing property does not exist for all partially hyperbolic systems any more [11]. Consequently, the research turns to the concept of quasi-shadowing property and sequences of shadowing points (not orbits in general) are allowed to make minor adjustments in the central direction.

Let f be a diffeomorphism on M and Λ be a (uniformly) partially hyperbolic set: there exist an Df-invariant splitting $T_{\Lambda}M = E^s \oplus E^c \oplus E^u$, constants $C \ge 1$, $0 < \lambda < 1 < \mu$ and $0 < \lambda < \lambda' \le \mu' < \mu < 1$ such that

- (i) $||D_x f^n(v)| \le C\lambda^n ||v||, \quad \forall x \in \Lambda, v \in E^s(x);$
- (ii) $\frac{1}{C}(\lambda')^n \|v\| \le \|D_x f^n(v)\| \le C(\mu')^n \|v\|, \quad \forall x \in \Lambda, v \in E^c(x);$
- (iii) $||D_x f^{-n}(v)| \le C\mu^{-n} ||v||, \quad \forall x \in \Lambda, v \in E^u(x).$

Take a constant $\rho_0 > 0$ such that for any $x \in M$ the standard exponential map $\exp_x : \{v \in T_x M : ||v|| < \rho_0\} \to M$ is a C^{∞} diffeomorphism to its image. Hu, Zhou and Zhu [9] proved that on partially hyperbolic set, f has the following quasi-shadowing property: for any $\varepsilon \in (0, \rho_0)$, there exists $\delta > 0$ such that for any δ -pseudo orbit $\{x_n\}_{n \in \mathbb{Z}}$, there exist a sequence of points

 $\{y_n\}_{n\in\mathbb{Z}}$ and a sequence of vectors $\{u_k\in E^c(x_k)\}_{n\in\mathbb{Z}}$ such that

$$d(x_n, y_n) < \varepsilon, \, \forall \, n \in \mathbb{Z}$$

where

$$y_n = \exp_{x_n}(u_n + \exp_{x_n}^{-1}(f(y_{n-1}))).$$

Besides, $\{y_n\}_{n\in\mathbb{Z}}$ and $\{u_n\}_{n\in\mathbb{Z}}$ can be chosen uniquely so as to satisfy

$$y_n \in \exp_{x_n}(E^s(x_n) + E^u(x_n)).$$

Here $u_k \in E^c(x_k)$ is taken into account as a motion along the center direction.

if for any pseudo orbit, there is a sequence of points $\{y_k\}_{k\in\mathbb{Z}}$ tracing it, in which y_{k+1} is obtained from $f(x_k)$ by a motion along the center direction. In [13], Li et al. studied the quasi-shadowing property for partially hyperbolic flows. In [12], Li et al. proved the quasi-shadowing property for quasi-partially hyperbolic pseudo-orbit made by quasi-partially hyperbolic strings.

(4)Quasi-shadowing for nonuniformly partially hyperbolic systems.

Nonuniformly partially hyperbolic system is a generalization of nonuniformly hyperbolic system by taking into account the center boundle and also a generalization of uniformly partially hyperbolic system by modifying the uniformity. Nonuniformly partially hyperbolic sets always exist, as we will see.

Let f be a $C^{1+\alpha}$ diffeomorphism on M.

Definition 1. Given numbers $\lambda, \mu > 0, -\lambda < -\lambda' < \mu' < \mu, \varepsilon > 0$ with $\varepsilon \ll \min\{-\lambda, \mu, |\lambda - \lambda'|, |\mu - \mu'|\}$, and a positive integer k, we define $\Lambda_k = \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$ to be all points $x \in M$ for which there is a Df-invariant decomposition $T_{\operatorname{orb}(x)}M = E^s \oplus E^c \oplus E^u$ along the orbit of x satisfying

(i)
$$\|Df^n(v)\| \le e^{\varepsilon k} e^{-(\lambda-\varepsilon)n} e^{\varepsilon |m|} \|v\|, \ \forall n \ge 0, m \in \mathbb{Z}, \ v \in E^s(f^m(x));$$

(ii) $\|Df^n(v)\| \le e^{\varepsilon k} e^{(\mu'+\varepsilon)n} e^{\varepsilon |m|} \|v\|,$ $\|Df^{-n}(v)\| \le e^{\varepsilon k} e^{(\lambda'+\varepsilon)n} e^{\varepsilon |m|} \|v\|, \ \forall n \ge 0, m \in \mathbb{Z}, \ v \in E^c(f^m(x));$

(iii)
$$\|Df^{-n}(v)\| \le e^{\varepsilon k} e^{-(\mu-\varepsilon)n} e^{\varepsilon |m|} \|v\|, \ \forall n \ge 0, m \in \mathbb{Z}, \ v \in E^u(f^m(x))$$

We refer to Λ_k as a partially hyperbolic block or Pesin block in nonuniformly partially hyperbolic sense. Denote by $\kappa = \kappa(x)$ the minimum value of the k that satisfy the aforementioned conditions for x and we refer to it as the index of partial hyperbolicity. Denote $\Lambda = \Lambda(\lambda, \mu, \lambda', \mu', \varepsilon) = \bigcup_{k \ge 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$ and we call Λ a nonuniformly partially hyperbolic set.

Remark. Nonuniformly partially hyperbolic set always exist. Given an ergodic measure which always exists, we can get a nonuniformly partially hyperbolic set of full measure by taking λ as the absolute value of the maximal negative Lyapunov exponent, μ as the minimum positive Lyapunov exponent and $\lambda' < \lambda$, $\mu' < \mu$. Here E^s or E^u may be trivial if μ has no negative Lyapunov exponent or positive Lyapunov exponent.

In this paper, we shall study the quasi-shadowing property for nonuniformly partially hyperbolic systems. For any $x \in M$ and a positive integer n, denote the orbit segment

$$[x, f^{n}(x)] := \{x, f(x), \cdots, f^{n}(x)\}\$$

On the orbit segment, there is no movement and it is real orbit. We want to show that if it is an orbit segment in the pseudo-orbit, it is shadowed by an orbit segment in the shadowing orbit.

There exists $\{\delta_k\}_{k\in\mathbb{Z}^+}$ such that for a $\{\delta_k\}_{k\in\mathbb{Z}^+}$ pseudo-orbit $\{z_i\}$, denoting $\tau(z_i) = s_i$, there exist fake stable foliation \mathcal{W}^s , fake center foliation \mathcal{W}^c and fake unstable foliation \mathcal{W}^u inside $B(z_i, \epsilon_0 e^{-\tau(z_i)\frac{\varepsilon}{\alpha}})$, where ϵ_0 is independent of $\tau(z_i)$. We refer to Section 2 for more details on the fake foliations. We have the following quasi-shadowing property adaptive to the movement of the pseudoorbit, i.e., the shadowing orbit has movement only if the pseudo-orbit has movement.

Theorem A. Let f be a $C^{1+\alpha}$ diffeomorphism on M and $\Lambda = \bigcup_{k\geq 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$ be a nonuniformly partially hyperbolic set. Then f has the quasi-shadowing property for Λ in the following sense: for any $\eta > 0$ there exists a sequence of positive numbers $\{\delta_k\}_{k\in\mathbb{Z}^+}$ such that for any $\{\delta_k\}_{k\in\mathbb{Z}^+}$ pseudo-orbit $\{z_i\} = \{\cdots, [x_{-n}, f^{a_{-n}-1}(x_{-n})], \cdots, [x_0, f^{a_0-1}(x_0)], \cdots, [x_n, f^{a_n-1}(x_n)], \cdots \} \subset$ Λ , there exists a sequence of points $\{y_n\}_{n\in\mathbb{Z}}$ satisfying:

(i) $f^{a_n}(y_n) \in \mathcal{W}^c(y_{n+1}), n \in \mathbb{Z}$, where \mathcal{W}^c is the fake center foliation of f;

(ii) $d(f^i(x_n), f^i(y_n)) < \eta \epsilon_{\tau(x_n)}, \ 0 \le i \le a_n, \ n \in \mathbb{Z}.$




FIGURE 1. Quasi-shadowing

In the category of invariant sets, periodic orbits are important invariant sets, and in the context of invariant measures, periodic measures are significant measures. Regarding periodic points, many scholars have already derived some results. Sigmund [20] proved that in nuniformly hyperbolic systems, periodic measures are dense in the set of invariant measure. Hirayama [7] proved that in nonuniformly hyperbolic systems, periodic measures are dense in the set of invariant measures supported on a total measure set with respect to a hyperbolic mixing measure. Liang et al [14] replaced the assumption of hyperbolic mixing measure by a more natural and weaker assumption of hyperbolic ergodic measures can be well approximated by periodic measures.

For n, p > 0, denote by $P'_{n,p}(f) = \{y : \text{ there exist a sequence of number } \{n_i\}_{i=0}^{p-1} \text{ satisfying } \sum_{i=0}^{p-1} n_i = n \text{ and a sequence of point } \{z_i\}_{i=1}^{p-1} \text{ such that } z_1 \in \mathcal{W}^c(f^{n_0}(x)), z_{i+1} \in \mathcal{W}^c(f^{n_i}(z_i)), i = 1, \cdots, p-2, x \in \mathcal{W}^c(f^{n_{p-1}}(z_{p-1}))\}, \text{ where } \mathcal{W}^c \text{ is the fake center foliation of } f. We refer to <math>P'_{n,p}(f)$ as the p quasi-periodic points of f with period n.

We have the following quasi-specification property concerning periodic shadowing.

Theorem B. Let $f: M \to M$ be a $C^{1+\alpha}$ diffeomorphism on a boundless compact smooth Riemannian manifold and there are a nonuniformly partially hyperbolic set $\Lambda = \bigcup_{k \ge 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$.

Let m be a Borel probability f-invariant ergodic measure and denote $\tilde{\Lambda}_k$ as the support of μ restricted on Λ_k and $\tilde{\Lambda} = \bigcup_{k\geq 1} \tilde{\Lambda}_k$. Then f has the quasi-specification property for $\tilde{\Lambda}$ in the fol-

lowing sense: for any $\eta > 0$, there exists a set of positive integers $\{M_{k,t} = M_{k,t}(\eta)\}_{k,t\in\mathbb{Z}^+}$ such that for any sequence of orbit segment $[x_1, f^{a_1}(x_1)], \cdots, [x_l, f^{a_l}(x_l)]$ with $x_i, f^{a_i}(x_i) \in \tilde{\Lambda}_{k_i}$ for $1 \leq i \leq l$, there exist two sequences of points $\{y_i\}_{i=i}^l$ and $\{\hat{y}_i\}_{i=1}^l$, and a sequence of positive integers $\{X_{1,2}, \cdots, X_{l-1,l}, X_{l,1}\}$ with $X_{i,i+1} \leq M_{k_i,k_{i+1}}, 1 \leq i \leq l-1$, and $X_{l,1} \leq M_{k_l,k_1}$ satisfying:

- (i) $f^{a_i}(y_i) \in \mathcal{W}^c(\hat{y}_i), f^{X_{i,i+1}}(\hat{y}_i) \in \mathcal{W}^c(y_{i+1}), i = 1, \cdots, l-1, f^{a_l}(y_l) \in \mathcal{W}^c(\hat{y}_l), and f^{X_{l,1}}(\hat{y}_l) \in \mathcal{W}^c(y_1), where \mathcal{W}^c$ is the fake center foliation of f, i.e., y_1 is a 2l quasi-periodic point of f;
- (ii) $d(f^{j}(y_{i}), f^{j}(x_{i})) \leq \eta \epsilon_{k_{i}}, \ 0 \leq j \leq a_{i}, \ i = 1, \cdots, l.$




FIGURE 2. Quasi-specification

The above quasi-specification property can be derived from the quasi-shadowing property as follows.

For any $\eta > 0$ and $k \ge 1$, take δ_k as in Theorem A. Let $\{M_{k,l} = M_{k,l}(\eta)\}_{k,l \in \mathbb{Z}^+}$ be as the constrction in [15] (Page 567). For each $1 \le i \le l$, there exist $z_i \in B(f^{a_i}(x_i), \delta_{k_i}) \cap \Lambda_{k_i}$ and a sequence of positive integers $\{X_{1,2}, \cdots, X_{l-1,l}, X_{l,1}\}$ with $X_{i,i+1} \le M_{k_i,k_{i+1}}, 1 \le i \le l-1, X_{l,1} \le M_{k_l,k_1}$ satisfying

$$f^{X_{i,i+1}}(z_i) \in B(x_{i+1}, \delta_{k_{i+1}}) \cap \Lambda_{k_{i+1}}, \quad 1 \le i \le l-1,$$

$$f^{X_{l,1}}(z_l) \in B(x_1, \delta_{k_1}) \cap \Lambda_{k_1},$$

and

$$\left\{ \begin{array}{c} x_1, \cdots, f^{a_1-1}(x_1), z_1, \cdots, f^{X_{1,2}-1}(z_1), x_2, \cdots, f^{a_2-1}(x_2), z_2, \cdots, f^{X_{2,3}-1}(z_2), \\ \cdots, x_l, \cdots, f^{a_l-1}(x_l), z_l, \cdots, f^{X_{l,1}-1}(z_l) \right\}^{\infty} \right\}$$

is a periodic $\{\delta_k\}_{k\in\mathbb{Z}^+}$ -pseudo-orbit. Theorem B is obtained by applying Theorem A.

The statement of the quasi-closing property is as follows:

Theorem C. Let $f: M \to M$ be a $C^{1+\alpha}$ diffeomorphism on a boundless compact smooth Riemannian manifold M and there is a nonuniformly partially hyperbolic set $\Lambda = \bigcup_{k\geq 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$.

Then f has the quasi-closing property for Λ in the following sense: for any $\eta > 0$ and $k \ge 1$ there exists $\beta = \beta(k, \eta) > 0$ such that for any point $x \in \Lambda_k$ if for some $p \in \mathbb{N}$ one has

$$f^p(x) \in \Lambda_k$$

and

$$d(x, f^p(x)) < \beta,$$

then there exists $y \in M$ such that:

- (i) $f^p(y) \in \mathcal{W}^c(y)$, where \mathcal{W}^c is the fake center foliation of f;
- (ii) $d(f^i(x), f^i(y)) < \eta \epsilon_k, \ i = 0, 1, \cdots, p-1.$

For any $\eta > 0$ and $k \ge 1$, take δ_k as in Theorem A and let $\beta(k, \eta) = \delta_k$. Then Theorem C is obtained by taking periodic pseudo-orbit

$$\left\{x, f(x), \cdots, f^{p-1}(x)\right\}^{\infty}.$$




FIGURE 3. Quasi-closing

Entropy is a concept widely applied across various fields including physics, information theory, statistics, and mathematics. Initially introduced by physicist Clausius in the mid-19th century, it describes the degree of disorder in thermodynamic systems. In different disciplines, entropy has various meanings and applications. In the theory of dynamical systems, entropy serves as a measure of the complexity of the system's dynamics. The larger the entropy of a dynamical system is, the less predictable the system is. Topological entropy and metric entropy are two distinct types of entropy. The concept of metric entropy was introduced by Kolmogorov in 1958 and is closely related to entropy in information theory, while topological entropy and topological entropy can be converted or complement each other, providing us with different perspectives to understand and analyze the complexity of dynamical systems. Interestingly, topological entropy and metric entropy are related to the number of periodic points.

Denote by $\#P_n(f)$ the number of periodic points with period n. Katok [10] uesd the closing property proved an estimation of asymptotic growth of the numbers of $\#P_n(f)$ by the metric entropy:

$$\overline{\lim_{n \to \infty}} \, \frac{\ln \# P_n(f)}{n} \ge h_m(f),$$

where m is a Borel probability f-invariant measure with non-zero Lyapunov exponents and #A is the cardinality of a set A. There have been someone who have demonstrated the relationship between the topological entropy and the exponential growth rate of periodic points. For any uniformly hyperbolic system, Bowen [3] proved that the asymptotical exponential growth of the number of periodic points was determined by the topological entropy h_{top} :

$$\lim_{n \to \infty} \frac{\ln \# P_n(f)}{n} = h_{\text{top}}(f)$$

As the application of Theorem C, we extend Katok's result of hyperbolic ergodic measure to any ergodic measure.

Recall that

$$P'_{n,1}(f) = \{ x : f^n(x) \in \mathcal{W}^c(x) \}.$$

and let

$$P_{n,1}'(f,\varepsilon) = \max \#E$$

with $E \subseteq P'_{n,1}(f)$ satisfying:

$$\forall x \neq y \in E : \exists i \in [0, n), d(f^{i}(x), f^{i}(y)) > \varepsilon.$$

Theorem D. Let $f: M \to M$ be a nonuniformly partially hyperbolic $C^{1+\alpha}$ diffeomorphism on a boundaryless compact smooth Riemannian manifold and there is a nonuniformly partially hyperbolic set $\Lambda = \bigcup_{k\geq 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$. Let m be an ergodic f-invariant measure with full measure for Λ . Then

$$\lim_{\varepsilon \to 0} \overline{\lim_{n \to \infty}} \frac{\ln P'_{n,1}(f,\varepsilon)}{n} \ge h_m(f).$$

2. UNIFORM PARTIAL HYPERBOLICITY

Let M be a boundaryless compact smooth Riemannian manifold, and $f: M \to M$ be a $C^{1+\alpha}$ diffeomorphism which has a nonuniformly partially hyperbolic set $\Lambda = \bigcup_{k \ge 1} \Lambda_k(\lambda, \mu, \lambda', \mu', \varepsilon)$.

By the definition of nonuniformly partially hyperbolic diffeomorphism, the time to admit contracting or expanding in E_x^s or E_x^u is related to the position of x. Replacing $\|\cdot\|$ with another metric $\|\cdot\|''$, which is equivalent to $\|\cdot\|$ on any given Λ_k , we can get uniformly contracting or expanding on Λ . Further details are provided below.

Let

$$\begin{cases} \lambda_1 = \lambda - 2\varepsilon, \\ \mu_1 = \mu - 2\varepsilon, \\ \lambda'_1 = \lambda' + 2\varepsilon, \\ \mu'_1 = \mu' + 2\varepsilon. \end{cases}$$

For $x \in \Lambda$, define

$$\| v_s \|'_s = \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_x f^n(v_s) \|, v_s \in E_x^s,$$

$$\| v_c \|'_c = \sum_{n=0}^{\infty} e^{-\mu'_1 n} \| D_x f^n(v_c) \| + \sum_{n=1}^{\infty} e^{-\lambda'_1 n} \| D_x f^{-n}(v_c) \|, v_c \in E_x^c,$$

$$\| v_u \|'_u = \sum_{n=0}^{\infty} e^{\mu_1 n} \| D_x f^{-n}(v_u) \|, v_u \in E_x^u.$$

Then let

$$\|\cdot\|'\colon T_{\Lambda}M\to\mathbb{R}$$

$$|| v ||'_{x} = \max\{|| v_{s} ||'_{s}, || v_{c} ||'_{c}, || v_{u} ||'_{u}\},\$$

where $v = v_s + v_c + v_u \in T_x M$ with $v_i \in E_x^i$, i = s, c, u. With respect to $x \in \Lambda_k$, it holds that

$$\sum_{n=0}^{\infty} e^{\lambda_1 n} \parallel D_x f^n(v_s) \parallel \leq \sum_{n=0}^{\infty} e^{\lambda_1 n} e^{-(\lambda-\varepsilon)} e^{\varepsilon k} \parallel v_s \parallel \leq \sum_{n=0}^{\infty} e^{-n\varepsilon} e^{k\varepsilon} \parallel v_s \parallel < \infty.$$

Similarly, we have $\sum_{n=0}^{\infty} e^{-\mu_1' n} \| D_x f^n(v_c) \| + \sum_{n=1}^{\infty} e^{-\lambda_1' n} \| D_x f^{-n}(v_c) \| < \infty$ and $\sum_{n=0}^{\infty} e^{\mu_1 n} \| D_x f^{-n}(v_u) \| < \infty$.

With the metric $\|\cdot\|', f: \Lambda \to \Lambda$ behaves uniformly partially hyperbolic: **Proposition 2.**

$$\begin{aligned} \frac{1}{\|Df^{-1}\|} \|v_s\|' &\leq \| Df(v_s) \|' &\leq e^{-\lambda_2} \| v_s \|', v_s \in E_x^s; \\ e^{-\lambda_2'} \| v_c \|' &\leq \| Df(v_c) \|' &\leq e^{\mu_2''} \| v_c \|', v_c \in E_x^c; \\ \frac{1}{\|Df\|} \|v_u\|' &\leq \| Df^{-1}(v_u) \|' &\leq e^{-\mu_2} \| v_u \|', v_u \in E_x^u. \end{aligned}$$

Proof. (1) E^s :

$$\| D_x f(v_s) \|' = \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_x f^{n+1}(v_s) \|$$

= $e^{-\lambda_1} \sum_{n=0}^{\infty} e^{\lambda_1 (n+1)} \| D_x f^{n+1}(v_s) \|$
 $\leq e^{-\lambda_1} \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_x f^n(v_s) \|$
 $\leq e^{-\lambda_1} \| v_s \|'.$

Besides,

$$\| D_x f(v_s) \|' = \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_{f^n(x)} f \circ D_x f^n(v_s) \|$$

$$\geq \frac{1}{\|Df^{-1}\|} \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_x f^n(v_s) \|$$

$$= \frac{1}{\|Df^{-1}\|} \| v_s \|'.$$

(2) E^c :

$$\| D_x f(v_c) \|' = \sum_{n=0}^{\infty} e^{-\mu'_1 n} \| D_x f^{n+1}(v_c) \| + \sum_{n=1}^{\infty} e^{-\lambda'_1 n} \| D_x f^{-(n-1)}(v_c) \|$$

$$= e^{\mu'_1} \sum_{n=0}^{\infty} e^{-\mu'_1(n+1)} \| D_x f^{n+1}(v_c) \| + e^{-\lambda'_1} \sum_{n=1}^{\infty} e^{-\lambda'_1(n-1)} \| D_x f^{-(n-1)}(v_c) \|,$$

which implies

$$e^{-\lambda'_1} \parallel v_c \parallel' \leq \parallel D_x f(v_c) \parallel' \leq e^{\mu'_1} \parallel v_c \parallel'.$$

(3) E^u : Similarly, we have

$$\frac{1}{\|Df\|} \|v_u\|' \le \|D_x f^{-1}(v_u)\|' \le e^{-\mu_1} \|v_u\|'.$$

If $||v||' = ||v_s||'_s$, then

.

$$\| v \|' = \| v_s \|'_s = \sum_{n=0}^{\infty} e^{\lambda_1 n} \| D_x f^n(v_s) \|$$

$$\leq \sum_{n=0}^{\infty} e^{\lambda_1 n} e^{\varepsilon k} e^{-(\lambda - \varepsilon)n} \| v_s \|$$

$$= e^{\varepsilon k} \sum_{n=0}^{\infty} e^{-\varepsilon n} \| v_s \| .$$

8

If $||v||' = ||v_c||'_c$, then

$$\| v \|' = \| v_c \|'_c = \sum_{n=0}^{\infty} e^{-\mu'_1 n} \| D_x f^n(v_c) \| + \sum_{n=1}^{\infty} e^{-\lambda'_1 n} \| D_x f^{-n}(v_c) \|$$

$$\leq \sum_{n=0}^{\infty} e^{-\mu'_1 n} e^{\varepsilon k} e^{(\mu'+\varepsilon)n} \| v_c \| + \sum_{n=1}^{\infty} e^{-\lambda'_1 n} e^{\varepsilon k} e^{(\lambda'+\varepsilon)n} \| v_c \|$$

$$\leq e^{\varepsilon k} \sum_{n=-\infty}^{\infty} e^{-\varepsilon |n|} \| v_c \| .$$

Similarly, if $||v||' = ||v_u||'_u$, then $||v||' = ||v_u||'_u \le e^{\varepsilon k} \sum_{n=0}^{\infty} e^{-\varepsilon n} ||v_u||.$

Denote
$$C = \sum_{n=-\infty}^{\infty} e^{-\varepsilon |n|}$$
, then $||v||' \le Ce^{\varepsilon k} ||v||$. Moreover,
 $||v||^2 = \langle v_s + v_c + v_u, v_s + v_c + v_u \rangle$
 $= ||v_s||^2 + ||v_c||^2 + ||v_u||^2 + 2\langle v_s, v_c \rangle + 2\langle v_s, v_u \rangle + 2\langle v_s, v_u \rangle$
 $\le 3(||v_s||^2 + ||v_c||^2 + ||v_u||^2) \le 3[(||v_s||'_s)^2 + (||v_c||'_c)^2 + (||v_u||'_u)^2]$
 $\le 9(||v||')^2.$

Thus we have $||v|| \leq 3 ||v||'$. Consequently, we obtain

$$\frac{1}{3} \parallel v \parallel \leq \parallel v \parallel' \leq C e^{\varepsilon k} \parallel v \parallel, \tag{1}$$

which means metric $\|\cdot\|'$ is equivalent to $\|\cdot\|$ on Λ_k . But metric $\|\cdot\|'$ is not equivalent to $\|\cdot\|$ on the whole $\Lambda = \bigcup_{k\geq 1} \Lambda_k$, as $\lim_{k\to\infty} Ce^{\varepsilon k} = +\infty$.

By taking small neighborhood U of any $x \in M$ with uniform size, we can trivialize the tangent bundle over U by identifying $T_U M = U \times \mathbb{R}^{\dim M}$. For any point $y \in U$ and tangent vector $v \in T_y M$, we can use the identification $T_U M = U \times \mathbb{R}^{\dim M}$ to translate the vector v to a corresponding vector $\bar{v} \in T_x M$. For $x \in \Lambda_k$, define

$$\|\cdot\|'': T_U M \to \mathbb{R}$$
$$\|v\|_{u}''=\|\bar{v}\|_{x}'$$

and a new splitting $T_y M = E_y^{s''} \oplus E_y^{c''} \oplus E_y^{u''}$ by translating the splitting $T_x M = E_x^s \oplus E_x^c \oplus E_x^u$. Denoting $E_y^{cs''} = E_y^{s''} \oplus E_y^{c''}$ and $E_y^{cu''} = E_y^{c''} \oplus E_y^{u''}$, we have the splittings $T_y M = E_y^{cs''} \oplus E_y^{u''}$ and $E_y^{cs} = E_y^{c''} \oplus E_y^{u''}$. Similarly, for a small neighborhood V of f(x) and any $z \in V$, we can define a metric $\|\cdot\|_z'$ by $\|\cdot\|_{f(x)}'$ and a new splitting $T_z M = E_z^{s''} \oplus E_z^{c''} \oplus E_z^{u''}$ by translating the splitting $T_{f(x)} M = E_{f(x)}^s \oplus E_{f(x)}^c \oplus E_{f(x)}^u$.

For a splitting $F = F_1 \oplus F_2$ of an Euclidean space F with norm || ||, and $\xi > 0$, we denote by $Q_{|| ||}(F_1,\xi)$ the cone of width ξ of F_1 in || ||, i.e. the set $\{v = v_1 + v_2 \in F : v_1 \in F_1, v_2 \in F_2, ||v_2|| \le \xi ||v_1||\}$.

Proposition 3. Let $\epsilon = \varepsilon/\alpha$. Take $\lambda_2 = \lambda_1 - \varepsilon$, $\mu_2 = \mu_1 - \varepsilon$, $\lambda'_2 = \lambda'_1 + \varepsilon$ and $\mu'_2 = \mu'_1 + \varepsilon$; $\lambda_3 = \lambda_2 - \varepsilon$, $\mu_3 = \mu_2 - \varepsilon$, $\lambda'_3 = \lambda'_2 + \varepsilon$ and $\mu'_3 = \mu'_2 + \varepsilon$.

(i) There exists $\epsilon_0 > 0$ such that for any $x \in \Lambda_k$, $y \in B(x, \epsilon_0 e^{-k\epsilon})$, one has

$$\left(\frac{1}{\|Df^{-1}\|} - (e^{\varepsilon} - 1)\right) \|v_s\|'' \le \|Df(v_s)\|'' \le e^{-\lambda_2} \|v_s\|'', v_s \in E_y^{s''}; e^{-\lambda_2'} \|v_c\|' \le \|Df(v_c)\|'' \le e^{\mu_2'} \|v_c\|'', v_c \in E_y^{c''}; \left(\frac{1}{\|Df\|} - (e^{\varepsilon} - 1)\right) \|v_u\|'' \le \|Df^{-1}(v_u)\|'' \le e^{-\mu_2} \|v_u\|'', v_u \in E_y^{u''}.$$

(ii) For any small $\xi > 0$,

$$\begin{split} \| Df(v) \|'' &\leq e^{-\lambda_3} \| v \|'', v \in Q_{\|\|_y''}(E_y^{s''},\xi); \\ e^{-\lambda_3'} \| v \|' \leq & \| Df(v) \|'' &\leq e^{\mu_3'} \| v \|'', v \in Q_{\|\|_y''}(E_y^{c''},\xi); \\ & \| Df^{-1}(v) \|'' &\leq e^{-\mu_3} \| v \|'', v_u \in Q_{\|\|_y''}(E_y^{u''},\xi). \end{split}$$

Moreover, there exist $\varsigma \in (0,1)$ and $a_{\xi} > 0$ such that for any $x \in \Lambda_k$, $y \in B(x, a_{\xi} \epsilon_0 e^{-k\epsilon})$,

$$D_y f(Q_{\parallel\parallel_y''}(E_y^*,\xi)) \subset Q_{\parallel\parallel_{f(y)}''}(E_{f(y)}^*,\varsigma\xi), \quad * \in \{u'', cu''\},$$

$$D_y f^{-1}(Q_{\parallel\parallel_y''}(E_y^*,\xi)) \subset Q_{\parallel\parallel_{f^{-1}(y)}''}(E_{f^{-1}(y)}^*,\varsigma\xi), \quad * \in \{s'', cs''\}.$$

Proof. Note that f is a $C^{1+\alpha}$ diffeomorphism and so f^{-1} is also $C^{1+\alpha}$. There exists a constant K > 0 such that

 $\| D_x f(v) - D_y f(v) \| \le K |x - y|^{\alpha} \| v \| \quad \text{and} \quad \| D_x f^{-1}(v) - D_y f^{-1}(v) \| \le K |x - y|^{\alpha} \| v \|.$ (i) Denote by B(x, r) the ball centered at x with radius r. Let

$$\epsilon_k = \min\left\{1, \left(\frac{e^{-\lambda_2} - e^{-\lambda_1}}{3Ce^{\varepsilon(k+1)}K}\right)^{\frac{1}{\alpha}}, \left(\frac{e^{\lambda_2'} - e^{\lambda_1'}}{3Ce^{\varepsilon(k+1)}K}\right)^{\frac{1}{\alpha}}, \left(\frac{(e^{-\mu_2} - e^{-\mu_1})}{3Ce^{\varepsilon(k+1)}K}\right)^{\frac{1}{\alpha}}, \left(\frac{(e^{\mu_2'} - e^{\mu_1'})}{3Ce^{\varepsilon(k+1)}K}\right)^{\frac{1}{\alpha}}\right\},$$

then for any $y \in B(x, \epsilon_\kappa)$, we have

(1) E^s :

$$\| D_y f(v_s) \|_{f(y)}^{\prime\prime} = \| D_y f(v_s) \|_{f(x)}^{\prime}$$

$$\leq \| D_x f(v_s) \|_{f(x)}^{\prime} + \| D_y f(v_s) - D_x f(v_s) \|_{f(x)}^{\prime}$$

$$\leq e^{-\lambda_1} \| v_s \|_x^{\prime} + C e^{\varepsilon (k+1)} K |y - x|^{\alpha} \| v_s \|_x$$

$$\leq (e^{-\lambda_1} + 3C e^{\varepsilon (k+1)} K |y - x|^{\alpha}) \| v_s \|_x^{\prime}$$

$$\leq e^{-\lambda_2} \| v_s \|_y^{\prime\prime}, v_s \in E_y^{s^{\prime\prime}}.$$

Besides,

$$\| D_y f(v_s) \|_{f(y)}^{\prime\prime} = \| D_y f(v_s) \|_{f(x)}^{\prime}$$

$$\geq \| D_x f(v_s) \|_{f(x)}^{\prime} - \| D_y f(v_s) - D_x f(v_s) \|_{f(x)}^{\prime}$$

$$\geq \frac{1}{\|Df^{-1}\|} \| v_s \|_x^{\prime} - Ce^{\varepsilon(k+1)} K |y - x|^{\alpha} \| v_s \|_x$$

$$\geq \frac{1}{\|Df^{-1}\|} \| v_s \|_x^{\prime} - 3Ce^{\varepsilon(k+1)} K |y - x|^{\alpha} \| v_s \|_x^{\prime}$$

$$\geq \left(\frac{1}{\|Df^{-1}\|} - (e^{\varepsilon} - 1) \right) \| v_s \|_x^{\prime}, \ v_s \in E_y^{s^{\prime\prime}}.$$

Similarly, we can further have that

(2)
$$E^{c}$$
:
 $e^{-\lambda'_{2}} \parallel v_{c} \parallel''_{y} \leq \parallel D_{y}f(v_{c}) \parallel''_{f(y)} \leq e^{\mu'_{2}} \parallel v_{c} \parallel''_{y}, v_{c} \in E_{y}^{c'};$
(3) E^{u} :
 $\left(\frac{1}{\parallel Df \parallel} - (e^{\varepsilon} - 1)\right) \parallel v_{u} \parallel''_{y} \leq \parallel D_{y}f^{-1}(v_{u}) \parallel''_{f^{-1}(y)} \leq e^{-\mu_{2}} \parallel v_{u} \parallel''_{y}, v_{u} \in E_{y}^{u'}$

Let $\epsilon = \frac{\epsilon}{\alpha}$. At most let ϵ_k small with finite k, we can take $\epsilon_0 > 0$, which is independent of k, such that $\epsilon_k = \epsilon_0 e^{-\epsilon k}$.

(ii) The first part on the inequalities can be obtained by (i) by taking small ξ . We are going to show the cone properties. Using the invariance of E^u and the domination property at

x there exists $\varsigma_1 \in (0,1)$ independent of x satisfying $D_x f(Q_{\parallel\parallel\parallel_x'}(E_x^u,\xi)) \subset Q_{\parallel\parallel\parallel_{f(x)}'}(E_{f(x)}^u,\varsigma_1\xi)$. Then for any $y \in B(x, a_{\xi}\epsilon_k)$, we get by (1) and the choice of ϵ_k ,

$$\begin{aligned} \|D_x f - D_y f\|_x'' &= \max_{\|v\|_x''=1} \|D_x f(v) - D_y f(v)\|_{f(x)}'' \\ &\leq C e^{(k+1)\varepsilon} \|D_x f - D_y f\| \\ &\leq C e^{(k+1)\varepsilon} K(a_{\xi} \epsilon_k)^{\alpha} \\ &\leq a_{\xi}^{\alpha}. \end{aligned}$$

For small ε , by (i) one has also

$$\frac{1}{2\|Df^{-1}\|} \le \min_{\|v\|_{x}^{''}=1} \|D_{y}f(v)\|_{f(x)}^{''} \le \max_{\|v\|_{x}^{''}=1} \|D_{y}f(v)\|_{f(x)}^{''} \le 2\|Df\|.$$

Take $\varsigma \in (\varsigma_1, 1)$. It follows that for $||v||_x'' = 1$, the angle $\angle''(D_y f(v), D_x f(v))$ with respect to $||\cdot||_{f(x)}''$ is less than $\arctan(\varsigma\xi) - \arctan(\varsigma_1\xi)$ for a_{ξ} small enough. We conclude that $D_y f\left(Q_{\|\|_x''}(E_y^u, \xi) \subset Q_{\|\|_{f(y)}'}\left(E_{f(y)}^u, \varsigma\xi\right)$ for any $y \in B(x, a_{\xi}\epsilon_k)$. The cone invariance property for $E^{cu''}$, $E^{s''}$ and $E^{cs''}$ can be obtained similarily.

2.1. Hölder continuity of Oseledets splitting. A k-dimensional distribution E on a smooth manifold M is a family of k-dimensional subspaces $E(x) \subset T_x M$. A Riemannian metric on M naturally induces distances in TM and in the space of k-dimensional distributions on TM. The Hölder continuity of a distribution E can be defined using these distances. By the Whitney Embedding Theorem, M can be embedded in an Euclidead space \mathbb{R}^N . Since M is compact, the Riemannian metric on M is equivalent to the distance ||x - y|| by the embedding for $x, y \in M$. Let $A \subset \mathbb{R}^N$ be a subspace and $v \in \mathbb{R}^N$ be a vector, set

$$d(v, A) = \min_{w \in A} \|v - w\|.$$

For subspaces A and B in \mathbb{R}^N , define

$$d(A,B) = \max\left\{\max_{v \in A, \|v\|=1} d(v,B), \max_{v \in B, \|v\|=1} d(w,A)\right\}$$

Let $D \subset \mathbb{R}^N$ be a subset and let E be a k-dimensional distribution. The distribution E is said to be Hölder continuous with Hölder exponent $\beta \in (0, 1]$ and Hölder constant $L \ge 0$ if there exists $\varepsilon_0 > 0$ such that

$$d(E(x), E(y)) \le L ||x - y||^{\beta}$$

for every $x, y \in D$ with $||x - y|| \le \varepsilon_0$.

Considering $E^s \oplus E^{cu}$, for $x \in \Lambda_k$, let $\theta_n = \angle (E^s(f^n(x)), E^{cu}(f^n(x)))$ for $n \in \mathbb{N}$. For any unit vector $v \in (E^s(x))^{\perp}$, denote $v = v_s + v_{cu}$ with $v_s \in E^s(x)$, $v_{cu} \in E^{cu}(x)$. It holds that $\|v_{cu}\| \ge \frac{1}{|\sin \theta_0|} \ge 1$. Then

$$\begin{aligned} \|Df^{n}v\| &\geq \|Df^{n}v_{cu}\| |\sin \theta_{n}| \\ &\geq e^{-k\varepsilon}e^{-n(\lambda'+\varepsilon)} |\sin \theta_{n}| \\ &\geq e^{-k\varepsilon}e^{-n(\lambda'+\varepsilon)}e^{-(k+n)\varepsilon} \\ &= e^{-2k\varepsilon}e^{-n(\lambda'+2\varepsilon)}. \end{aligned}$$

Besides, for unit vector $v \in E^s(x)$,

$$\|Df^n v\| \leq e^{k\varepsilon} e^{-n(\lambda-\varepsilon)},$$

and by Lemma 4.16 of [2] for every $a > \max\{\|D_z f\|^{1+\alpha}, \|D_z f^{-1}\|^{1+\alpha} : z \in M\}$, there exists D > 1 such that for every $n \in \mathbb{N}$ and every $x, y \in M$ we have

$$\|D_x f^{\pm n} - D_y f^{\pm n}\| \leqslant Da^n \|x - y\|^{\alpha}$$

By Lemma 4.14 of [2], we have for $x, y \in \Lambda_k$,

$$d(E^{s}(x), E^{s}(y)) \leqslant 3(e^{2k\varepsilon})^{2} \frac{e^{-(\lambda'+2\varepsilon)}}{e^{-(\lambda-\varepsilon)}} (D\|x-y\|^{\alpha})^{\frac{-(\lambda'+2\varepsilon)+\lambda-\varepsilon}{\ln a+\lambda-\varepsilon}}$$
$$= 3e^{4k\varepsilon} e^{\lambda-\lambda'-3\varepsilon} (D\|x-y\|^{\alpha})^{\frac{\lambda-\lambda'-3\varepsilon}{\ln a+\lambda-\varepsilon}}$$

Similarly, consider $E^{cs} \oplus E^u$. For any unit vector $v \in (E^{cs}(x))^{\perp}$, it holds that

$$\|Df^n v\| \ge e^{-2k\varepsilon} e^{n(\mu - 2\varepsilon)}.$$

Besides, for unit vector $v \in E^{cs}(x)$,

$$\|Df^n v\| \leq e^{k\varepsilon} e^{n(\mu' + \varepsilon)}.$$

We have for $x, y \in \Lambda_k$,

$$d(E^{cs}(x), E^{cs}(y)) \leqslant 3(e^{2k\varepsilon})^2 \frac{e^{\mu-2\varepsilon}}{e^{\mu'+\varepsilon}} (D\|x-y\|^{\alpha})^{\frac{\mu-2\varepsilon-(\mu'+\varepsilon)}{\ln a-(\mu'+\varepsilon)}}$$
$$= 3e^{4k\varepsilon} e^{\mu-\mu'-3\varepsilon} (D\|x-y\|^{\alpha})^{\frac{\mu-\mu'-3\varepsilon}{\ln a-\mu'-\varepsilon}}.$$

Moreover, considering f^{-1} , we can get that for $x, y \in \Lambda_k$,

$$d(E^{u}(x), E^{u}(y)) \leqslant 3(e^{2k\varepsilon})^{2} \frac{e^{-(\mu'+2\varepsilon)}}{e^{-(\mu-\varepsilon)}} (D||x-y||^{\alpha})^{\frac{-(\mu'+2\varepsilon)+(\mu-\varepsilon)}{\ln a+(\mu-\varepsilon)}}$$
$$= 3e^{4k\varepsilon} e^{\mu-\mu'-3\varepsilon} (D||x-y||^{\alpha})^{\frac{\mu-\mu'-3\varepsilon}{\ln a+\mu-\varepsilon}},$$

and

$$d(E^{cu}(x), E^{cu}(y)) \leqslant 3(e^{2k\varepsilon})^2 \frac{e^{\lambda - 2\varepsilon}}{e^{\lambda' + \varepsilon}} (D \|x - y\|^{\alpha})^{\frac{\lambda - 2\varepsilon - (\lambda' + \varepsilon)}{\ln a - (\lambda' + \varepsilon)}}$$
$$= 3e^{4k\varepsilon} e^{\lambda - \lambda' - 3\varepsilon} (D \|x - y\|^{\alpha})^{\frac{\lambda - \lambda' - 3\varepsilon}{\ln a - \lambda' - \varepsilon}}.$$

Denote

$$a_1 = \max\left\{\frac{\lambda - \lambda' - 3\varepsilon}{\ln a + \lambda - \varepsilon}, \ \frac{\mu - \mu' - 3\varepsilon}{\ln a - \mu' - \varepsilon}, \ \frac{\mu - \mu' - 3\varepsilon}{\ln a + \mu - \varepsilon}, \ \frac{\lambda - \lambda' - 3\varepsilon}{\ln a - \lambda' - \varepsilon}\right\}$$

Then E^s , E^{cs} , E^u , E^{cu} on Λ_k are $a_1\alpha$ -Hölder continuous with Hölder constant $b_1e^{4k\varepsilon}$, where b_1 is independent of k and n. Since $E^c = E^{cs} \cap E^{cu}$, we also get that E^c on Λ_k is $a_1\alpha$ -Hölder continuous with Hölder constant $b_1e^{4k\varepsilon}$.

Now we consider the Hölder continuity with respect to <,>'.

Proposition 4. E^s, E^{cu}, E^u and E^{cs} on Λ_k are $a_1\alpha$ -Hölder continuous with Hölder constant $b_2e^{6k\varepsilon}$ for some constant b_2 , with respect to <,>'.

Proof. For $x, y \in \Lambda_k$, let $e_s \in E^s(x)$, $e_{cu} \in E^{cu}(x)$ and $v \in E^s(y)$ be the unit vectors with respect to $\langle \rangle'$. Denote by γ' the angle between e_s and v with respect to $\langle \rangle'_x$ then

$$v = \cos \gamma' \cdot e_s + \sin \gamma' \cdot e_{cu}$$

Denote by γ the angle between e_s and v with respect to the Riemannian metric of M, then

$$\begin{aligned} |\tan \gamma| &= \frac{\|\sin \gamma' \sin \theta_0 \cdot e_{cu}\|}{\|\cos \gamma' \cdot e_s\| + |\cos \theta_0| \|\sin \gamma' \cdot e_{cu}|} \\ &\geqslant \frac{|\sin \gamma'| e^{-k\varepsilon} C^{-1} e^{-k\varepsilon}}{3+3}, \end{aligned}$$

which implies

$$\gamma' \leqslant C_1 e^{2k\varepsilon} \gamma$$

for some constant C_1 .




FIGURE 4.

Note that

$$\gamma \leqslant b_1 e^{4k\varepsilon} \|x - y\|^{a_1\alpha} \leqslant b_1 e^{4k\varepsilon} (3\|x - y\|')^{a_1\alpha}.$$

For two subspaces with small angle, any one of the distance, the angle, tangent and sine are uniformly bounded by any other. Therefore, we get that

$$d'(E^s(x), E^s(y)) \leqslant b_2 e^{6k\varepsilon} (\|x-y\|')^{a_1\alpha}$$

for some constant b_2 , where d' is the distance of subspaces with respect to \langle , \rangle' . The Hölder continuity of E^{cu}, E^u and E^{cs} can be proved analogously.

2.2. Invariant foliations. Note that for any $\eta > 0$, $B(x, \eta)$ contains a ball centered at x with radius $\frac{1}{3}\eta$ in the distance d'', which we denote by $B''(x, \frac{1}{3}\eta)$. By Proposition 3, it holds uniform partial hyperbolicity for $T_yM = E_y^{s''} \oplus E_y^{c''} \oplus E_y^{u''}$ in $\|\cdot\|''$ for $y \in B''(x, \frac{1}{3}a_{\xi}\epsilon_0e^{-k\epsilon}) \subseteq B(x, a_{\xi}\epsilon_0e^{-k\epsilon}), x \in \Lambda_k$. By taking the exponential map at x we can assume without loss of generality that we are working in \mathbb{R}^d . Let ξ and a_{ξ} be as in Proposition 3. For any $x \in \Lambda^*$, we can extend $f \mid_{B(x, a_{\xi}\epsilon_0e^{-k\epsilon})}$ to a diffeomorphism $\tilde{f}_x : \mathbb{R}^d \to \mathbb{R}^d$ such that

- $\tilde{f}_x(y) = f(y)$ for $y \in B(x, a_{\xi} \epsilon_0 e^{-k\epsilon})$;
- $||D_y \tilde{f}_x D_x f||_x'' \le 2a_{\varepsilon}^{\alpha}$ for $y \in \mathbb{R}^d$.

By taking a_{ξ} smaller, the properties of Proposition 3 hold with respect to \tilde{f}_x for all $y \in \mathbb{R}^d$. Moreover, there exists L > 1 such that

$$||D_y \tilde{f}_x^{\pm}||'' \le L, \quad \forall y \in \mathbb{R}^{\mathsf{d}}, \ x \in \Lambda^*.$$

Let

$$\delta_k = ((1-\varsigma)\xi(b_2e^{6k\varepsilon}))^{-a_1\alpha}$$
 for any $k \in \mathbb{Z}_+$.

For a sequence $\{a_n\}_{n\in\mathbb{Z}}$ of positive integers, let $\{\cdots, [x_{-n}, f^{a_{-n}-1}(x_{-n})], \cdots, [x_0, f^{a_0-1}(x_0)], \cdots, [x_n, f^{a_n-1}(x_n)], \cdots \}$ Λ be a sequence of orbit segments which is a $\{\delta_k\}_{k\in\mathbb{Z}^+}$ pseudo-orbit, i.e., there exist two sequences of positive integers $\{s_n^-\}_{n\in\mathbb{Z}}$ and $\{s_n^+\}_{n\in\mathbb{Z}}$ such that

(a)
$$x_n \in \Lambda_{s_n}^-, f^{a_n}(x_n) \in \Lambda_{s_n}^+, \forall n \in \mathbb{Z};$$

(b) $|s_n^+ - \bar{s_{n+1}}| \le 1, \forall n \in \mathbb{Z};$

(c)
$$d(f^{a_n}(x_n), x_{n+1}) < \delta_{s_n}, \forall n \in \mathbb{Z}.$$

Then by the Hölder property of Proposition 4,

$$d(E^*(f^{a_n}(x_n), E^*(x_{n+1}))) < (1-\varsigma)\xi, \ * \in \{s, cs, cu, u\}$$

Therefore,

$$Q_{\|\|_{f^{a_n}(x_n)}^{\prime\prime}}(E^*_{f^{a_n}(x_n)},\varsigma\xi) \subset Q_{\|\|_{x_{n+1}}^{\prime\prime}}(E^*_{x_{n+1}},\xi), \ * \in \{s,cs,cu,u\}$$

Denote

$$\{z_i\}_{i=-\infty}^{+\infty} = \{\cdots, x_{-n}, \cdots, f^{a_{-n}-1}(x_{-n}), \cdots, x_0, \cdots, f^{a_0-1}(x_0), \cdots, x_n, f^{a_n-1}(x_n), \cdots\}$$

with $z_0 = x_0$. Let Ξ be the disjoint union given by

$$\Xi = \prod_{i=-\infty}^{+\infty} \{z_i\} \times \mathbb{R}^{\mathsf{d}}.$$

Then $\tilde{f} = (\tilde{f}_{z_i})_{i=-\infty}^{+\infty}$ can be viewed as a map from Ξ to itself by letting $\tilde{f}(x,v) = \left(f(x), \tilde{f}_x(v)\right)$.

Note that the global splitting $\coprod_{i=-\infty}^{+\infty} \{z_i\} \times \mathbb{R}^d = \coprod_{i=-\infty}^{+\infty} \{z_i\} \times (E_{z_i}^s \oplus \oplus E_{z_i}^c \oplus E_{z_i}^u)$ is dominated with respect to \tilde{f} . By [8] 5 and , we can obtain a family $\{\mathcal{Y}_{:}^{cs} x \in \Lambda^*\}$ of global invariant foliations C^1 submanifolds in \mathbb{R}^d which are C^1 graphs defined on E_x^{cs} such that we have for all $x \in \Lambda^*$:

$$\{x\} \times \mathcal{Y}_x^{cs} = \bigcap_{n=0}^{+\infty} \tilde{f}^{-n} \left(\{f^n(x)\} \times Q_{\parallel\parallel_x^{\prime\prime}}(E_{f^n(x)}^{cs},\xi)\right),$$
$$\forall y \in \mathbb{R}^{\mathsf{d}}, \ T_y \mathcal{Y}_x^{cs} \subset Q_{\parallel\parallel_x^{\prime\prime}}(E_x^{cs},\xi).$$

In particular we get $\tilde{f}^{\pm}(\{x\} \times \mathcal{Y}_x^{cs}) \subset \{f^{\pm}(x)\} \times \mathcal{Y}_{f^{\pm}(x)}^{cs}$. Since we have $\tilde{f}|_{\{x\} \times B(x, a_{\xi} \gamma_{\kappa(x)})} = f|_{B(x, b_{\xi} \gamma_{\kappa(x)})}$, one concludes the proof by considering $\mathcal{W}_x^{cs} = \mathcal{Y}_x \cap B(x, a_{\xi} \gamma_{\kappa(x)})$ and taking much smaller b_{ξ} than a_{ξ} .

Proposition 5. Any $\{z_i\} \times \mathbb{R}^d$ with $-\infty < i < +\infty$ is foliated by foliations $\mathcal{W}_{z_i}^s$, $\mathcal{W}_{z_i}^{cs}$, $\mathcal{W}_{z_i}^{cu}$, $\mathcal{W}_{z_i}^c$, and $\mathcal{W}_{z_i}^u$ with the following properties: for $* \in \{s, cs, cu, c, u\}$ and for each $\tilde{y} = \{z_i\} \times \{y\} \in \{z_i\} \times \mathbb{R}^d$,

- (i) almost tangency: the leaf $\mathcal{W}_{z_i}^*(y)$ is C^1 , and the tangent space $T_y \mathcal{W}_{z_i}^*(z)$ lies in a cone of radius ϵ about $E_{z_i}^*$.
- (ii) invariance: $f_{z_i}(\mathcal{W}^*_{z_i}(y)) = \mathcal{W}^*_{z_{i+1}}(f_{z_i}(y)).$
- (iii) exponential growth bounds:
 - (a) for any $y_1, y_2 \in \mathcal{W}^s_{z_i}(y), d''(f_{z_i}(y_1), f_{z_i}(y_2)) \leq e^{-\lambda_3} d''(y_1, y_2);$
 - (b) for any $y_1, y_2 \in \mathcal{W}_{z_i}^c(y), \ e^{-\lambda'_3} d''(y_1, y_2) \le d''(f_{z_i}(y_1), f_{z_i}(y_2)) \le e^{\mu'_3} d''(y_1, y_2);$
 - (c) for any $y_1, y_2 \in \mathcal{W}^u_{z_i}(y), d''(f_{z_i}(y_1), f_{z_i}(y_2)) \ge e^{\mu_3} d''(y_1, y_2).$
- (iv) coherence: $\mathcal{W}_{z_i}^s$ and $\mathcal{W}_{z_i}^c$ subfoliate $\mathcal{W}_{z_i}^{cs}$, and $\mathcal{W}_{z_i}^u$ and $\mathcal{W}_{z_i}^c$ subfoliate $\mathcal{W}_{z_i}^{cu}$.
- (v) regularity: the foliations W* and their tangent distributions are uniformly Hölder continuous.

For $y, z \in \mathbb{R}^d$, write

- $\{[x,y]_{cs,u}\} = \mathcal{W}^{cs}_{z_i}(x) \cap \mathcal{W}^u_{z_i}(y),$
- $\{[x,y]_{s,cu}\} = \mathcal{W}^s_{z_i}(x) \cap \mathcal{W}^{cu}_{z_i}(y).$

There exists $b \ge 1$ such that

$$\begin{aligned} d_{\mathcal{W}_{z_{i}}^{cs}(x)}(x, [x, y]_{cs, u}) &\leq bd(x, y), \qquad d_{\mathcal{W}_{z_{i}}^{u}(x)}(y, [x, y]_{cs, u}) \leq bd(x, y), \\ d_{\mathcal{W}_{z_{i}}^{s}(x)}(x, [x, y]_{s, cu}) &\leq bd(x, y), \qquad d_{\mathcal{W}_{z_{i}}^{cu}(x)}(y, [x, y]_{s, cu}) \leq bd(x, y). \end{aligned}$$

By letting ϵ small, we can let b close to 1 such that

$$\frac{b}{2}(1+e^{-\min\{\lambda_3,\,\mu_3\}+\epsilon})<1.$$

By letting ϵ small, we can let b close to 1 such that

$$\frac{b}{2}(1+e^{-\min\{\lambda_3,\,\mu_3\}+\theta^{-1}\epsilon})<1.$$

By the regularity property (v), $\mathcal{W}_{z_i}^*$ is θ -Hölder continuous for some $\theta \in (0,1)$ and the constant \tilde{C} can be chosen uniformly for uniformly bounded regions.

3. Quasi-shadowing property

Proof of Theorem A. For any z_i , $-\infty < i < +\infty$, we take cylinders

$$C^{cu}(z_i, \gamma e^{-\theta^{-1}\tau(z_i)\epsilon}) = \bigcup_{x \in \mathcal{W}^s_{z_i}(z_i, \gamma e^{-\theta^{-1}\tau(z_i)\epsilon})} \mathcal{W}^{cu}_{z_i}(x),$$
$$C^{cs}(z_i, \gamma e^{-\theta^{-1}\tau(z_i)\epsilon}) = \bigcup_{x \in \mathcal{W}^u_{z_i}(z_i, \gamma e^{-\theta^{-1}\tau(z_i)\epsilon})} \mathcal{W}^{cs}_{z_i}(x),$$

where γ will be assigned later. Note that

$$\begin{split} \tilde{f}(C^{cu}(z_{i},\gamma e^{-\theta^{-1}\tau(z_{i})\epsilon})) &\subset C^{cu}(\tilde{f}(z_{i}),e^{-\lambda_{3}}\gamma e^{-\theta^{-1}\tau(z_{i})\epsilon}).\\ \text{Take } \delta_{i} &= \frac{\gamma}{2}(1-e^{-\lambda_{3}+\theta^{-1}\epsilon})C^{-1}e^{-(1+\theta^{-1})\tau(z_{i})\epsilon}. \text{ Since } d(\tilde{f}(z_{i}),z_{i+1}) < \delta_{i+1} &= \frac{\gamma}{2}(1-e^{-\lambda_{3}+\theta^{-1}\epsilon})C^{-1}e^{-(1+\theta^{-1})\tau(z_{i+1})\epsilon},\\ d''(\tilde{f}(z_{i}),z_{i+1}) &< \frac{1}{2}(1-e^{-\lambda_{3}+\theta^{-1}\epsilon})\gamma e^{-\theta^{-1}\tau(z_{i+1})\epsilon}. \end{split}$$

For any $y \in \mathcal{W}^s_{f(z_i)}(e^{-\lambda_3}\gamma e^{-\theta^{-1}\tau(z_i)\epsilon})$, we have

$$d''(y, z_{i+1}) < e^{-\lambda_3} \gamma e^{-\theta^{-1} \tau(z_i)\epsilon} + \frac{1}{2} (1 - e^{-\lambda_3 + \theta^{-1}\epsilon}) \gamma e^{-\theta^{-1} \tau(z_{i+1})\epsilon},$$

$$\leq e^{-\lambda_3 + \theta^{-1}\epsilon} \gamma e^{-\theta^{-1} \tau(z_{i+1})\epsilon} + \frac{1}{2} (1 - e^{-\lambda_3 + \theta^{-1}\epsilon}) \gamma e^{-\theta^{-1} \tau(z_{i+1})\epsilon},$$

$$= \frac{1}{2} (1 + e^{-\lambda_3 + \theta^{-1}\epsilon}) \gamma e^{-\theta^{-1} \tau(z_{i+1})\epsilon}.$$

Therefore,

$$d_{\mathcal{W}^{s}(z_{i+1})}''(z_{i+1}, [y, z_{i+1}]_{cu,s}) < \frac{b}{2}(1 + e^{-\lambda_{3} + \theta^{-1}\epsilon})\gamma e^{-\theta^{-1}\tau(z_{i+1})\epsilon} \le \gamma e^{-\theta^{-1}\tau(z_{i+1})\epsilon},$$

which implies

$$\tilde{f}(C^{cu}(z_i, \gamma e^{-\theta^{-1}\tau(z_i)\epsilon})) \subset C^{cu}(z_{i+1}, \gamma e^{-\theta^{-1}\tau(z_{i+1})\epsilon}).$$

In the same manner, one can get that

$$\tilde{f}^{-1}(C^{cs}(z_i,\gamma e^{-\theta^{-1}\tau(z_i)\epsilon})) \subset C^{cs}(z_{i-1},\gamma e^{-\theta^{-1}\tau(z_{i-1})\epsilon}).$$

Denote

$$C^{cu}_{\infty}(z_i) = \bigcap_{j=0}^{+\infty} \tilde{f}^j (C^{cu}(z_{i-j}, \gamma e^{-\theta^{-1}\tau(z_{i-j})\epsilon})),$$

$$C^{cs}_{\infty}(z_i) = \bigcap_{-\infty}^{+\infty} \tilde{f}^{-j} (C^{cs}(z_{i+j}, \gamma e^{-\theta^{-1}\tau(z_{i+j})\epsilon})).$$

$$C_{\infty}(z_i) = \prod_{j=0}^{\infty} f^{-j} (C^{-j}(z_{i+j}, \gamma e^{-j(z_{i+j})}))$$

which consist of cu-leaves and cs-leaves, respectively. It holds that

$$\tilde{f}(C^{cu}_{\infty}(z_i)) = C^{cu}_{\infty}(z_{i+1}),$$

$$\hat{f}^{-1}(C^{cs}_{\infty}(z_i)) = C^{cs}_{\infty}(z_{i-1}).$$

Take one cu-leaf $F_{z_0} \subset C^{cu}_{\infty}(z_0)$ and one cs-leaf $G_{z_0} \subset C^{cs}_{\infty}(z_0)$. Denote

$$F_{z_i} = \tilde{f}^i(F_{z_0}) \subset C^{cu}_{\infty}(z_i) \quad \text{and} \quad G_{z_i} = \tilde{f}^i(G_{z_0}) \subset C^{cs}_{\infty}(z_i), \quad \forall i \in \mathbb{Z}.$$

Then F_{z_i} and G_{z_i} are *cu*-leaf and *cs*-leaf, respectively.

Denote

$$\{p(z_i)\} = F_{z_i} \cap \mathcal{W}_{z_i}^s(z_i), \quad \{q(z_i)\} = G_{z_i} \cap \mathcal{W}_{z_i}^u(z_i), \quad \{t(z_i)\} = G_{z_i} \cap \mathcal{W}_{z_i}^u(p(z_i)).$$

It holds that

$$t(z_i) \in F_{z_i} \cap G_{z_i} = \mathcal{W}_{z_i}^c(t(z_i)).$$

Moreover, by the Hölder continuity of $\mathcal{W}^{cs}_{z_i},$

$$d_{\mathcal{W}_{z_i}^u(p(z_i))}(p(z_i), t(z_i)) \leq \tilde{C}(d_{\mathcal{W}_{z_i}^u(z_i)}(z_i, q(z_i)))^{\theta} \leq \tilde{C}(\gamma e^{-\theta^{-1}\tau(z_i)\epsilon})^{\theta} = \tilde{C}\gamma^{\theta} e^{-\tau(z_i)\epsilon}.$$

And

$$d_{\mathcal{W}_{z_i}^s(z_i)}(z_i, p(z_i)) \le \gamma e^{-\theta^{-1}\tau(z_i)\epsilon}.$$

Thus

$$d''(z_i, t(z_i)) \le \tilde{C}\gamma^{\theta} e^{-\tau(z_i)\epsilon} + \gamma e^{-\theta^{-1}\tau(z_i)\epsilon} \le C_1 \gamma^{\theta} e^{-\tau(z_i)\epsilon},$$
(2)

for some constant $C_1 > 1$.

Now correspond to the pseudo-orbit

$$\{\cdots, [x_{-n}, f^{a_{-n}-1}(x_{-n})], \cdots, [x_0, f^{a_0-1}(x_0)], \cdots, [x_n, f^{a_n-1}(x_n)], \cdots\} \subset \Lambda.$$

(I) the distance along orbit segment $[x_n, f^{a_n-1}(x_n)]$ for each n.

By the invariance of fake foliations,

$$\tilde{f}^{a_n-1}(t(x_n)) = t(\tilde{f}^{a_n-1}(x_n)) \in F_{\tilde{f}^{a_n-1}(x_n)} \cap G_{\tilde{f}^{a_n-1}(x_n)} = \mathcal{W}^c_{t(\tilde{f}^{a_n-1}(x_n))}(t(\tilde{f}^{a_n-1}(x_n))).$$
By the Hölder continuity of WC8

By the Hölder continuity of $\mathcal{W}^{cs}_{\tilde{f}^{a_n-1}(x_n)}$,

$$d_{\mathcal{W}^{u}_{\tilde{f}^{a_{n}-1}(x_{n})}(p(\tilde{f}^{a_{n}-1}(x_{n}))}(p(\tilde{f}^{a_{n}-1}(x_{n})), t(f\tilde{f}^{a_{n}-1}(x_{n})))$$

$$\leq \tilde{C}(d_{\mathcal{W}^{u}_{\tilde{f}^{a_{n}-1}(x_{n})}(\tilde{f}^{a_{n}-1}(x_{n}))}(f^{a_{n}-1}(x_{n}), q(\tilde{f}^{a_{n}-1}(x_{n})))^{\theta}$$

$$\leq \tilde{C}(\gamma e^{-\theta^{-1}\tau(\tilde{f}^{a_{n}-1}(x_{n}))\epsilon})^{\theta} \leq C_{1}\gamma^{\theta}e^{-\tau(\tilde{f}^{a_{n}-1}(x_{n}))\epsilon}.$$

Moreover,

$$d_{\mathcal{W}^{u}_{\tilde{f}^{a_{n}}(x_{n})}(p(\tilde{f}^{a_{n}}(x_{n}))}(p(\tilde{f}^{a_{n}}(x_{n})), t(\tilde{f}^{a_{n}}(x_{n})))$$

$$\leq d_{\mathcal{W}^{u}_{\tilde{f}^{a_{n}-1}(x_{n})}(p(\tilde{f}^{a_{n}-1}(x_{n}))}(p(\tilde{f}^{a_{n}-1}(x_{n})), t(\tilde{f}^{a_{n}-1}(x_{n})))L$$

$$< C_{1}\gamma^{\theta}e^{\epsilon}e^{-\tau(\tilde{f}^{a_{n}}(x_{n}))\epsilon}L.$$

Note that $p(\tilde{f}^{a_n-1}(x_n)) = \tilde{f}^{a_n-1}(p(x_n))$ and $t(\tilde{f}^{a_n-1}(x_n)) = \tilde{f}^{a_n-1}(t(x_n))$. It follows that for $0 \le j \le a_n - 1$,

$$\begin{aligned} &d_{\mathcal{W}^{u}_{\tilde{f}^{j}(x_{n})}(\tilde{f}^{j}(p(x_{n})))}(\tilde{f}^{j}(p(x_{n})),\tilde{f}^{j}(t(x_{n})))\\ &\leq \quad d_{\mathcal{W}^{u}_{\tilde{f}^{a_{n}-1}(x_{n})}(p(\tilde{f}^{a_{n}-1}(x_{n}))}(p(\tilde{f}^{a_{n}-1}(x_{n})),t(\tilde{f}^{a_{n}-1}(x_{n})))\cdot e^{-\mu_{3}(a_{n}-1-j)}\\ &\leq \quad C_{1}\gamma^{\theta}e^{-\tau(\tilde{f}^{a_{n}-1}(x_{n}))\epsilon}e^{-\mu_{3}(a_{n}-1-j)}\leq C_{1}\gamma^{\theta}e^{-\tau(\tilde{f}^{j}(x_{n}))\epsilon}.\end{aligned}$$

Besides, we have

$$d_{\mathcal{W}^{s}_{\tilde{f}^{j}(x_{n})}(\tilde{f}^{j}(x_{n}))}(\tilde{f}^{j}(x_{n}), \tilde{f}^{j}(p(x_{n})))$$

$$\leq d_{\mathcal{W}^{s}_{x_{n}}(x_{n})}(x_{n}, p(x_{n})) \cdot e^{-\lambda_{3}j}$$

$$\leq \gamma e^{-\theta^{-1}\tau(x_{n})\epsilon} e^{-\lambda_{3}j} \leq C_{1}\gamma^{\theta} e^{-\tau(\tilde{f}^{j}(x_{n}))\epsilon}.$$

Thus,

$$d''(f^{j}(x_{n}), f^{j}(t(x_{n})))$$

$$\leq d_{\mathcal{W}^{s}_{\tilde{f}^{j}(x_{n})}(\tilde{f}^{j}(x_{n}))}(\tilde{f}^{j}(x_{n}), \tilde{f}^{j}(p(x_{n}))) + d_{\mathcal{W}^{u}_{\tilde{f}^{j}(x_{n})}(\tilde{f}^{j}(p(x_{n})))}(\tilde{f}^{j}(p(x_{n})), \tilde{f}^{j}(t(x_{n})))$$

$$\leq 2C_{1}\gamma^{\theta}e^{-\tau(\tilde{f}^{j}(x_{n}))\epsilon}.$$

(II) the distance along center leaf.

It holds that

$$d''(t(x_{n+1}), \tilde{f}^{a_n}(t(x_n)))$$

$$\leq d''(t(x_{n+1}), x_{n+1}) + d''(\tilde{f}^{a_n}(x_n), x_{n+1}) + d''(\tilde{f}^{a_n}(x_n), \tilde{f}^{a_n}(t(x_n))).$$

Note that

$$d''(t(x_{n+1}), x_{n+1}) \leq C_1 \gamma^{\theta} e^{-\tau(x_{n+1})\epsilon},$$

$$d''(f^{a_n}(x_n), x_{n+1}) \leq \gamma C_0 e^{-(1+\theta^{-1})\tau(x_{n+1})\epsilon},$$

and

$$d''(\tilde{f}^{a_n}(x_n), \tilde{f}^{a_n}(t(x_n))) \leq d''(\tilde{f}^{a_n-1}(x_n), \tilde{f}^{a_n-1}(t(x_n)))L$$

$$\leq 2C_1 \gamma^{\theta} e^{-\tau(\tilde{f}^{a_n-1}(x_n))\epsilon}L$$

$$\leq 2C_1 \gamma^{\theta} e^{\varepsilon} e^{-\tau(\tilde{f}^{a_n}(x_n))\epsilon}L$$

$$\leq 2C_1 \gamma^{\theta} e^{2\varepsilon} e^{-\tau(x_{n+1})\epsilon}L.$$

Therefore,

$$\mathcal{L}''(t(x_{n+1}), \tilde{f}^{a_n}(t(x_n))) \le C_2 \gamma^{\theta} e^{-\tau(x_{n+1})\epsilon},$$

for some constant $C_2 > 2LC_1 > 1$. For any $\eta > 0$, let $\gamma = \left(\frac{\epsilon_0 \eta}{3C_2}\right)^{\frac{1}{\theta}}$, we can get the estimate in Theorem A. Moreover, there exists small $\eta_0 > 0$ such that for any $\gamma \in \left(0, \left(\frac{\epsilon_0 \eta_0}{C_2}\right)^{\frac{1}{\theta}}\right)$, the above estimates are done in the corresponding $B''(z, \frac{1}{3}a_{\xi}\epsilon_0e^{-\tau(z)\epsilon})$ thus in $B(z, a_{\xi}\epsilon_0e^{-\tau(z)\epsilon})$ for any $z = z_i$, where $\tilde{f} = f$.

4. THE GROWTH OF QUASI-PERIODIC ORBITS

Firstly, we introduce the following Katok.s definition of metric entropy ([10]). For every $0 < \delta < 1$ and an ergodic *f*-invariant measure *m*, denote $d_n^f(x, y) = \max_{0 \le i \le n-1} d(f^i(x), f^i(y))$ and denote by $N_f(n, \gamma, \delta)$ the minimal number of γ -balls in the d_n^f -metric which cover the set of measure more than or equal to $1 - \delta$. Then

$$h_m(f) = \lim_{\gamma \to 0} \lim_{n \to \infty} \frac{\ln N_f(n, \gamma, \delta)}{n} = \lim_{\gamma \to 0} \lim_{n \to \infty} \frac{\ln N_f(n, \gamma, \delta)}{n}$$

Recall the set of quasi-periodic points: $P'_n(f) = \{x : f^n(x) \in \mathcal{W}^c(x)\}.$

Proof of Theorem D. Fix a positive integer k. Then for any small positive number γ , any positive number l and any positive integer n, we shall construct a finite set $K_n = K_n(\gamma, l)$ satisfying the following properties (for some n the set $K_n(\gamma, l)$ may be empty):

- (i) $K_n \subset \Lambda_k$;
- (ii) if $x, y \in K_n$ and $x \neq y$, then $d_n^f(x, y) > l^{-1}$;
- (iii) for every $x \in K_n$ there exists a number $m(x) : n \leq m(x) \leq (1+\gamma)n$ such that $f^{m(x)}(x) \in \Lambda_k$ and

$$d(x, f^{m(x)}(x)) \le \beta(k, (3l)^{-1});$$

(iv) for every $\gamma > 0$, denote by $\#K_n(\gamma, l)$ the cardinality of $K_n(\gamma, l)$,

$$\lim_{l \to \infty} \lim_{n \to \infty} \frac{\ln \# K_n(\gamma, l)}{n} \ge h_m(f)$$

To construct the set K_n , we choose a finite partition ξ satisfying

diam
$$\xi < \beta(k, 1/(3l))$$

and

$$\xi > \{\Lambda_k, M \setminus \Lambda_k\},\$$

which means that every element of ξ either belongs to the set Λ_k or is disjoint from this set.

Let $\Lambda_{k,n} = \{x \in \Lambda_k : \exists m \in [n, (1 + \gamma)n] \text{ such that the points } x \text{ and } f^m(x) \text{ belong to the same element of } \xi\}$. Define K_n as the subset of the set $\Lambda_{k,n}$ with the maximal cardinality satisfying the property (ii). The properties (i) and (iii) are true by definition.

By the property (ii), we know that the union of l^{-1} -balls in the d_n^f -metric around points of K_n covers the set $\Lambda_{k,n}$. Then we have

$$#K_n(\gamma, l) \ge N_f(n, l^{-1}, 1 - m(\Lambda_{k,n})).$$

Lemma 6.

$$\lim_{n \to \infty} m(\Lambda_{k,n}) = m(\Lambda_k)$$

Proof. We fix an element $A \in \xi$ which is a subset of the set Λ_k and set

$$A_{n,\gamma} = \{ x \in A : \sum_{k=0}^{n-1} \chi_A(f^k(x)) < nm(A)(1+\frac{\gamma}{3}), \sum_{k=0}^{[n(1+\gamma)]} \chi_A(f^k(x)) > nm(A)(1+\frac{2\gamma}{3}) \},$$

where χ_D is a characteristic function of a set D.

Obviously $A_{n,\gamma} \subset \Lambda_{k,n} \cap A$. By the ergodic theorem we have $m(A \setminus A_{n,\gamma}) \to 0$, as $n \to \infty$. Applying these arguments to every element $A \in \xi$ belonging to Λ_k we obtain that

$$\lim_{n \to \infty} m(\Lambda_{k,n}) = m(\Lambda_k).$$

By Lemma 6, we can get that for every $\delta > 1 - m(\Lambda_k)$, we have

$$\lim_{n \to \infty} \frac{\ln \# K_n}{n} \ge \lim_{n \to \infty} \frac{\ln N_f(n, l^{-1}, \delta)}{n}.$$
(3)

By Katok.s definition of metric entropy, we know that the property (iv) is true.

From Theorem C, for every point $x \in K_n$ we can find a quasi-periodic point z(x) with period m(x) satisfying $d(f^i(x), f^i(z(x))) < (3l)^{-1}, i = 0, 1, \dots, m(x) - 1$. If $x, y \in K_n$ and $x \neq y$ then

$$\begin{split} & \max_{0 \le i \le n-1} d(f^i(z(x)), f^i(z(y))) \\ & \ge \max_{0 \le i \le n-1} d(f^i(x), f^i(y)) - \max_{0 \le i \le n-1} d(f^i(x), f^i(z(x))) - \max_{0 \le i \le n-1} d(f^i(y), f^i(z(y))) \\ & \ge \max_{0 \le i \le n-1} d(f^i(x), f^i(y)) - \max_{0 \le i \le m(x)-1} d(f^i(x), f^i(z(x))) - \max_{0 \le i \le m(x)-1} d(f^i(y), f^i(z(y))) \\ & \ge l^{-1} - (3l)^{-1} - (3l)^{-1} \\ & \ge (3l)^{-1}. \end{split}$$

So that z(x) and z(y) are different. Consequently, we get

$$\sum_{m=n}^{[(1+\gamma)n]} \# P'_{m,1}(f,(3l)^{-1}) \ge \# K_n(\gamma,n)$$

and

$$\max_{\leq m \leq (1+\gamma)n} \# P'_{m,1}(f, (3l)^{-1}) \geq \frac{\# K_n(\gamma, l)}{\gamma n + 1}.$$

Thus, we can find a sequence of integers $\{m_n\}$ where $n \leq m_n \leq (1+\gamma)n$ such that

$$\underbrace{\lim_{n \to \infty} \frac{\ln \# P'_{m_n,1}(f,(3l)^{-1})}{m_n}}_{n \to \infty} \ge \underbrace{\lim_{n \to \infty} \frac{n}{m_n} \cdot \frac{\ln \# K_n(\gamma,l) - \ln(\gamma n + 1)}{n}}_{1 \to \gamma} \\
\ge \frac{1}{1 + \gamma} \underbrace{\lim_{n \to \infty} \frac{\ln \# K_n(\gamma,l)}{n}}_{n}.$$

By (3), we have

$$\underbrace{\lim_{l \to \infty} \lim_{n \to \infty} \frac{\ln \# P'_{m_n,1}(f, (3l)^{-1})}{m_n}}_{l \to \infty} = \underbrace{\lim_{l \to \infty} \lim_{n \to \infty} \frac{\ln \# P'_{m_n,1}(f, (3l)^{-1})}{m_n}}_{l \to \infty}$$

$$\geq \frac{1}{1+\gamma} \cdot \underbrace{\lim_{l \to \infty} \lim_{n \to \infty} \frac{\ln \# K_n(\gamma, l)}{n}}_{l \to \infty}}_{n \to \infty}$$

$$= \frac{1}{1+\gamma} h_m(f).$$

Let $\gamma \to 0$, we can get the result.

 $n \leq 1$

References

- D. V. Anosov. Geodesic flows on closed riemannian manifolds of negative curvature. Pro. Steklov Inst. Math., 90, 1967.
- [2] L. Barreira and Y. Pesin. Introduction to smooth ergodic theory, volume 148 of Grad. Stud. Math. Amer. Math. Soc., Providence, RI, 2013.
- [3] R. Bowen. Topological entropy and axiom A. In Global Analysis Proc. Sympos. Pure Math. 1970.
- [4] R. Bowen. Periodic points and measures for Axiom A diffeomorphisms. Trans. Amer. Math. Soc., 154:377– 397, 1971.
- [5] S. Gan. The star systems \mathcal{X}^* and a proof of the C^1 ω -stability conjecture for flows. J. Differential Equations, 163:1–17, 2000.
- [6] S. Gan. A generalized shadowing lemma. Discrete Contin. Dyn. Syst., 8:627–632, 2002.
- [7] M. Hirayama. Periodic probability measures are dense in the set of invariant measures. Discrete Contin. Dyn. Syst., 9:1185–1192, 2003.
- [8] M. Hirsch, C. Pugh, and M. Shub. Invariant manifolds, volume 583 of Lect. Notes in Math. 1977.
- H. Hu, Y. Zhou, and Y. Zhu. Quasi-shadowing for partially hyperbolic diffeomorphisms. Ergod. Th. & Dynam. Sys., 35:412–430, 2015.
- [10] A. Katok. Lyapunov exponents, entropy and periodic orbits for diffeomorphisms. Inst. Hautes Études Sci. Publ. Math., 51:137–173, 1980.
- S. Kryzhevich and S. Tikhomirov. Partial hyperbolicity and central shadowing. Discrete Contin. Dyn. Syst., 33:2901–2909, 2013.
- [12] Z. Li and Y. Zhou. Quasi-shadowing and limit quasi-shadowing for quasi-partially hyperbolic strings of flows. J. Differential Equations, 269:11062–11085, 2020.
- [13] Z. Li and Y. Zhou. Quasi-shadowing for partially hyperbolic flows. Discrete Contin. Dyn. Syst., 40:2089– 2103, 2020.
- [14] C. Liang, G. Liu, and W. Sun. Approximation properties on invariant measure and oseledec splitting in non-uniformly hyperbolic systems. *Trans. Amer. Math. Soc.*, 361:1543–1579, 2009.
- [15] C. Liang, W. Sun, and X. Tian. Ergodic properties of invariant measures for C^{1+α} non-uniformly hyperbolic system. Ergod. Th. & Dynam. Sys., 33:560–584, 2013.
- [16] S. Liao. An existence theorem for periodic orbits. Acta Sci. Natur. Univ. Pekinensis, 1:1–20, 1979.
- [17] R. Mañé. A proof of the C¹ stability conjecture. Publ. Math. IHES, 66:161–210, 1988.
- [18] Y. Pesin. Families of invariant manifolds corresponding to nonzero characteristic exponents. Math. USSR-Izv., 40:1261–1305, 1976.
- [19] O. Sarig. Symbolic dynamics for surface diffeomorphisms with positive entropy. J. Amer. Math. Soc., 26:341–426, 2013.
- [20] K. Sigmund. Generic properties of invariant measures for axiom a diffeomorphisms. Invent. Math., 11:99– 109, 1970.
- [21] Y. G. Sinai. Gibbs measures in ergodic theory. Uspekhi Mat. Nauk, 27:21–64, 1972.

GANG LIAO AND XUETONG ZU

- $\left[22\right]$ L. Wen. On the preperiodic set. Discrete Contin. Dynam. Systems, $6:\!237\!-\!241,\,2000.$
- [23] X. Wen, S. Gan, and L. Wen. C¹-stably shadowable chain components are hyperbolic. J. Differential Equations, 246:340–357, 2009.

Email address: lg@suda.edu.cn Email address: zuxuetong@163.com