

Predictions for Bottomonium from a Relativistic Screened Potential Model

Chaitanya Anil Bokade^{1,*} and Bhaghyesh^{1,†}

¹*Department of Physics, Manipal Institute of Technology
Manipal Academy of Higher Education, Manipal, 576104, Karnataka, India*

In this work, a comprehensive analysis of the mass spectra and decay properties of bottomonium states using a relativistic screened potential model is carried out. The mass spectrum, decay constants, $E1$ transitions, $M1$ transitions, and annihilation decay widths are evaluated. The interpretation of $\Upsilon(10355)$, $\Upsilon(10580)$, $\Upsilon(10860)$, and $\Upsilon(11020)$ as $S - D$ mixed bottomonium states are analysed. The $\Upsilon(10355)$ state is considered to be $3S - 2D$, $\Upsilon(10580)$ state is considered to be $4S - 3D$ mixed state, the $\Upsilon(10753)$ is obtained as purely $\Upsilon_1(3D)$ bottomonium state, and the $\Upsilon(10860)$ and $\Upsilon(11020)$ are deemed to be $5S - 4D$ mixed states.

Keywords: Bottomonium, Relativistic Potential Model, Screened Potential, Radiative Decays, $S - D$ Mixing.

I. INTRODUCTION

The study of heavy quarkonium, specifically bottomonium, has emerged as a captivating and influential field in contemporary particle physics. The allure of this research lies not only in the experimental endeavors aimed at unraveling the intricate properties of these heavy quark systems [1, 2], but also its rich theoretical framework which allows to understand intricate interplay of perturbative and non-perturbative quantum chromodynamics (QCD) phenomena across a broad energy spectrum [3]. The history of bottomonium traces back to $\Upsilon(1S)$, the first discovered bottomonium particle, which had shed light on the existence of a new heavy quark. $\Upsilon(1S)$ was initially observed by the E288 Collaboration at Fermilab along with $\Upsilon(2S)$ and $\Upsilon(3S)$ in 1977 [4, 5]. This marked the beginning of a journey that unfolded with subsequent breakthroughs, including the identification of excited bottomonium states. In 1982, the $\chi_{bJ}(2P)$ states were observed from the data obtained by CUSB detector at CESR in the $E1$ transition $\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)$ for $(J = 0, 1, 2)$ [6, 7]. The $\chi_{bJ}(1P)$ states were discovered in 1983 for the first time in $\eta_b(2S) \rightarrow \gamma\chi_{bJ}(1P)$ and $\chi_{bJ}(1P) \rightarrow \gamma\eta_b(1S)$ decay processes with $(J = 0, 1, 2)$ [8]. The centre of gravity, fine structure splitting, and the branching ratios were also measured [8]. $\chi_{bJ}(1P)$ were later confirmed in the same year by CESR from the data in the $\Upsilon(2S) \rightarrow \gamma\chi_{bJ}(1P) \rightarrow \gamma\Upsilon(1S)$ reaction [9]. Later in 2005, most precise measurement of branching fractions and photon energies of $\chi_{bJ}(1P)$ and $\chi_{bJ}(2P)$ for $(J = 0, 1, 2)$ were conducted by the CLEO Collaboration in the same decay process of $\Upsilon(3S) \rightarrow \gamma\chi_{bJ}(2P)$ [10]. For the first time in 1980, CESR observed a peak above the $B\bar{B}$ threshold from the data obtained by e^+e^- collisions and suggested it to be $\Upsilon(4S)$ [11]. Later in 1985, the data from the CLEO detector at CESR above the $B\bar{B}$ threshold for e^+e^- collisions were analyzed and in addition to $\Upsilon(4S)$, the observations of $\Upsilon(10860)$ and $\Upsilon(11020)$ resonances were also reported [12]. $\Upsilon(10860)$ and $\Upsilon(11020)$ are generally considered as the $\Upsilon(5S)$ and $\Upsilon(6S)$ bottomonium states. Most recently, Belle Collaboration in 2019 conducted new measurements of $e^+e^- \rightarrow \Upsilon(1, 2, 3S)\pi^+\pi^-$ cross sections [13]. They measured the mass and decay width of $10885.3 \pm 1.5_{-0.9}^{+2.2}$ MeV and $36.6_{-3.9-1.1}^{+4.5+0.5}$ MeV for $\Upsilon(10860)$, and mass and decay width of $11000.0_{-4.5-1.3}^{+4.0+1.0}$ MeV and $23.8_{-6.8-1.8}^{+8.0+0.7}$ MeV for $\Upsilon(11020)$ [13]. In 2004, CLEO Collaboration in the photon cascade reaction of $\Upsilon(3S) \rightarrow \gamma\chi_b(2P)$, $\chi_b(2P) \rightarrow \gamma\Upsilon(1D)$, $\Upsilon(1D) \rightarrow \gamma\chi_b(1P)$ and $\chi_b(1P) \rightarrow \gamma\Upsilon(1S)$, with $\Upsilon(1S)$ annihilating into l^+l^- (where $l = e$ or μ), observed the $\Upsilon(1D)$ state [14]. The peak value is observed at $10161.1 \pm 0.6 \pm 1.6$ MeV, which is predicted to be $\Upsilon_2(1D)$ state [14]. BABAR Collaboration later confirmed observation of $\Upsilon_J(1D)$ triplet in the $\Upsilon(3S) \rightarrow \gamma\Upsilon(1D) \rightarrow \gamma\gamma\pi^+\pi^-\Upsilon(1S)$ decay mode with significance of 5.8σ for $\Upsilon_2(1D)$ state [15]. There is also possibility of observing $\Upsilon_1(1D)$ and $\Upsilon_3(1D)$ state with the mass value around 10.13 GeV and 10.18 GeV, respectively, but their significance is very low [15]. After decades, a breakthrough came in the bottomonium sector with the discovery of $\eta_b(1S)$ in 2008 from the photon energy spectrum with the BABAR detector at the PEP-II B factory at SLAC [16]. The peak was observed with significance of 10σ and is interpreted as a monochromatic photon from the radiative transition $\Upsilon(3S) \rightarrow \gamma\eta_b(1S)$ [16], which was later confirmed by CLEO Collaboration [17, 18]. CLEO Collaboration also identified $\eta_b(2S)$ with 5σ significance from the $\Upsilon(2S) \rightarrow \gamma\eta_b(2S)$ decay process [18]. Belle Collaboration observed $\eta_b(2S)$ for the first time in $h_b(2P) \rightarrow \gamma\eta_b(2S)$ decay process with most precise measurement of mass value of $9999.0 \pm 3.5_{-1.9}^{+2.8}$ MeV and hyperfine splitting of $m[\Upsilon(2S)] - m[\eta_b(2S)] = 24.3_{-4.5}^{+4.0}$ MeV [19]. In 2011, the first observation of $h_b(1P)$ was recorded by BABAR Collaboration in cascade decay process of $\Upsilon(3S) \rightarrow \pi^0 h_b(1P) \rightarrow \pi^0 \gamma \eta_b(1S)$ with significance of 3.1σ [20].

* anshul497@gmail.com

† bhaghyesh.mit@manipal.edu; Corresponding author; ORCID iD: 0000-0003-3994-9945

Subsequently the Belle Collaboration, in the $\Upsilon(5S) \rightarrow \pi^+\pi^-h_b(1P)$ decay process gave the confirmation of $h_b(1P)$ [21]. They also observed the $h_b(2P)$ state for the first time with significance of 11.2σ having mass value of $10259.8 \pm 0.6_{-1.0}^{+1.4}$ MeV [21]. In 2012, ATLAS Collaboration at LHC observed decay modes of $\chi_b(nP) \rightarrow \gamma\Upsilon(1S, 2S)$ in the production of $\chi_b(nP)$ states with proton-proton collision [22]. Among detected modes, they also observed decays consistent with $\chi_b(3P) \rightarrow \gamma\Upsilon(1S, 2S)$, having mass barycentre at $10530 \pm 5 \pm 9$ MeV [22]. In the same year D0 Collaboration observed the same structure in $\chi_b(3P) \rightarrow \gamma\Upsilon(1S)$ decay with significance of 5.6σ having mass barycentre at $10551 \pm 14 \pm 17$ MeV [23]. Later in 2018, using the data from proton-proton collisions CMS Collaboration observed $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ in the $\gamma\Upsilon(3S)$ decay mode [24]. The mass values were determined to be $\chi_{b1}(3P) = 10513.42 \pm 0.41 \pm 0.18$ MeV and $\chi_{b2}(3P) = 10524.02 \pm 0.57 \pm 0.17$ MeV and having mass difference of $10.60 \pm 0.64 \pm 0.17$ MeV [24]. In 2012, from the data on $\Upsilon(5S)$ decays to $\Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) and $h_b(mP)\pi^+\pi^-$ ($m = 1, 2$) Belle Collaboration observed two charged structures, $Z_b(10610)$ and $Z_b(10650)$ [25]. Due to their charged property, they cannot be described in conventional quarkonium picture and require a four-quark configuration description such as hadronic molecules [26, 27], tetraquarks [28, 29], etc. In 2019, BELLE Collaboration using the data of $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) decay process from the Belle detector at KEKB detected a new structure $\Upsilon(10753)$ with mass value of $10752.7 \pm 5.9_{-1.1}^{+0.7}$ MeV and decay width of $35.5_{-11.3-3.3}^{+17.6+3.9}$ MeV with significance of 5.2σ [30]. $\Upsilon(10753)$ is also observed from the calculation of Born cross section, dressed cross section and the total hadronic cross section for the e^+e^- annihilation in the bottomonium region from the data collected by BABAR and BELLE experiments [31]. Even with significant progress in experimental domain there is still lack of key details, such as the total width and mass values of higher resonance, branching ratios for significant decay modes, etc. Unlike the rich data on charmonium-like XYZ states, bottomonium sector has only discovered limited number of states ($\Upsilon(10580), \Upsilon(10860), \Upsilon(11020), Z_b(10610)$ and $Z_b(10650)$), thus a parallel abundance of novel states is yet to be explored. There is also no experimental information on X_b , the counterpart of $X(3872)$ in bottomonium sector [32]. The pursuit of similar exotic states in the bottomonium system, such as the X_b whose existence is predicted in multiple models [33, 34], holds promise for understanding the nature of the internal structure of $X(3872)$. Exotic hadron investigations have predominantly relied on e^+e^- annihilation experiments, exemplified by BESIII, Belle, BaBar, and CLEO, owing to their pristine experimental conditions and diverse production mechanisms [3]. The Belle II experiment, situated at the SuperKEKB e^+e^- collider, occupies a distinctive role in bottomonium physics, boasting comprehensive detector capabilities and with the ambitious goal of reaching a peak luminosity of $8 \times 10^{35} \text{cm}^2 \text{s}^{-1}$ by 2025. Belle II plans to extend operations until 2027, accumulating over 50ab^{-1} of invaluable data [3]. Following the LHCb Upgrade I, the data from Run-3 is poised to be of significant importance. Additionally, the PANDA experiment, focusing on antiproton-nucleon interaction, offers an alternative avenue for the exploration of these states. Furthermore, various projects centered on super τ - charm factories are in the planning stages, promising intriguing prospects in the near future [2, 35]. In view of the potential for discovering new states in the bottomonium sector, we use a relativistic screened potential model that has proven effective for charmonium [36]. The proposed model provides a robust theoretical framework by considering relativistic effects and the screening of the potential. Our model can facilitate the identification and characterization of new and exotic states within the bottomonium spectrum.

In this paper we have conducted a comprehensive study of bottomonium in a relativistic screened potential model. In section II, we discuss the theoretical model used to describe the bottomonium bound system and the numerical approach used to solve the relativistic Schrodinger equation. Decay constants are discussed in section III. In sections IV and V we present the various decays that are investigated. In section VI, $S - D$ mixing of bottomonium states is discussed. In section VII a thorough investigation of our evaluation and interpretation of bottomonium states are conducted, along with comparison between the experimental results and other theoretical models. In section VIII we present our conclusion.

II. METHODOLOGY

A relativistic potential model is developed to investigate various bottomonium properties. We utilize the spinless Salpeter equation, which is a relativistic extension of the Schrodinger equation [37]

$$H = \sqrt{-\nabla_q^2 + m_q^2} + \sqrt{-\nabla_{\bar{q}}^2 + m_{\bar{q}}^2} + V(r), \quad (1)$$

where $\vec{r} = \vec{x}_{\bar{q}} - \vec{x}_q$, $\vec{x}_{\bar{q}}$ and \vec{x}_q are the coordinates of the quarks and operators ∇_q^2 and $\nabla_{\bar{q}}^2$ are the partial derivatives of those coordinates respectively. m_q and $m_{\bar{q}}$ are the masses of quark and an anti-quark respectively. The interaction potential $V(r)$ between the quark and antiquark is composed of two components: $V_V(r)$, representing the one-gluon-exchange coulomb potential term that is dominant at short distance, and $V_S(r)$, which represents the linear confining

term adjusted to account for colour screening effects at longer distances [38]:

$$V_V(r) = -\frac{4}{3} \frac{\alpha_s(r)}{r}, \quad (2)$$

$$V_S(r) = \lambda \left(\frac{1 - e^{-\mu r}}{\mu} \right) + V_0, \quad (3)$$

$$V(r) = V_V(r) + V_S(r). \quad (4)$$

Here λ is the linear potential slope and μ is the screening factor which regulates the behaviour of the long-range component of $V(r)$, causing it to flatten out as r becomes much larger than $1/\mu$ and exhibit a linear increase as r becomes much smaller than $1/\mu$. The $V(r)$ converges to Cornell potential as $\mu \rightarrow 0$ [38]. $\alpha_s(r)$ is the running coupling constant in coordinate space obtained by Fourier transformation of coupling constant in momentum space $\alpha_s(Q^2)$ [37] and is given by

$$\alpha_s(r) = \sum_i \alpha_i \frac{2}{\sqrt{\pi}} \int_0^{\gamma_i r} e^{-x^2} dx, \quad (5)$$

where α_i 's are the free parameters to imitate the short-distance behaviour of $\alpha_s(Q^2)$ as predicted by QCD. The parameters values are taken as $\alpha_1 = 0.15$, $\alpha_2 = 0.15$, $\alpha_3 = 0.20$, and $\gamma_1 = 1/2$, $\gamma_2 = \sqrt{10}/2$, $\gamma_3 = \sqrt{1000}/2$ [37, 39]. The Hamiltonian H is solved as an eigenvalue equation using the method developed in [39, 40]. The Hamiltonian Eq. (1) as a eigen equation is given by

$$E\Psi(\vec{r}) = \left[\sqrt{-\nabla_q^2 + m_q} + \sqrt{-\nabla_{\bar{q}}^2 + m_{\bar{q}}} + V(r) \right] \Psi(\vec{r}). \quad (6)$$

The wave function can be expanded using spectral integration, which allows us to express the wave function as an integral over the eigenstates of the Hamiltonian H :

$$\Psi(\vec{r}) = \int d^3r' \int \frac{d^3k}{(2\pi)^3} e^{i\vec{k}(\vec{r}-\vec{r}')} \Psi(\vec{r}'). \quad (7)$$

Eq. (1) can be rewritten as

$$E\Psi(\vec{r}) = \int d^3r' \frac{d^3k}{(2\pi)^3} \left(\sqrt{k^2 + m_q} + \sqrt{k^2 + m_{\bar{q}}} \right) e^{i\vec{k}(\vec{r}-\vec{r}')} \Psi(\vec{r}') + V(r)\Psi(\vec{r}). \quad (8)$$

The exponential term can be expanded in terms of spherical harmonics as

$$e^{i\vec{k}\cdot\vec{r}} = 4\pi \sum_{nl} Y_{nl}^*(\hat{k}) Y_{nl}(\hat{r}) j_l(kr) i^l, \quad (9)$$

where j_l is the spherical Bessel function, $Y_{nl}^*(\hat{k})$ and $Y_{nl}(\hat{r})$ are the spherical harmonics with normalization condition $\int d\Omega Y_{n_1 l_1}(\hat{k}) Y_{n_2 l_2}(\hat{r}) = \delta_{n_1 n_2} \delta_{l_1 l_2}$, \hat{k} and \hat{r} are unit vectors along the \vec{k} and \vec{r} direction, respectively. The wave function can be factorized into radial $R_l(r)$ and angular $Y_{nl}(r)$ parts, and substituting Eq. (9) in (8) and simplifying, we get [39, 40]

$$E u_l(r) = \frac{2}{\pi} \int dk k^2 \int dr' r r' \left(\sqrt{k^2 + m_q^2} + \sqrt{k^2 + m_{\bar{q}}^2} \right) j_l(kr) j_l(kr') u_l(r') + V(r) u_l(r), \quad (10)$$

where $u_l(r)$ is the reduced radial wave function ($R_l(r) = u_l(r)/r$). When the separation distance grows for a quark-antiquark bound state, the wavefunction gradually decreases and eventually approaches zero at sufficiently large distance. In order to represent this behaviour, a characteristic distance scale L is introduced, confining the bound state's wavefunction within the spatial interval of $0 < r < L$. Next, one can expand the reduced wavefunction $u_l(r)$ in terms of spherical Bessel function for angular momentum l as

$$u_l(r) = \sum_{n=1}^{\infty} c_n \frac{a_n r}{L} j_l \left(\frac{a_n r}{L} \right), \quad (11)$$

where c_n 's are the expansion coefficients, a_n are the n -th root of the spherical Bessel function, $j_l(a_n) = 0$. For large value of N Eq. (11) can be truncated. The momentum is discretized as a result of confinement of space, which

TABLE I. Parameters used in our model

m_q (GeV)	σ (GeV ²)	λ (GeV)	μ (GeV)	Λ (GeV)
4.744	4.967	0.240	0.039	0.17

allows us to replace $a_n/L \leftrightarrow k$ and the integration in Eq. (10) can be replaced by $\int dk \rightarrow \sum_n \Delta a_n/L$, where, $\Delta a_n = a_n - a_{n-1}$. For finite space interval, $0 < r, r' < L$, incorporating all the changes in the Eq. (10), we get the final equation in terms of the coefficients c_n 's as [39, 40]

$$Ec_m = \sum_{n=1}^N \frac{a_n}{N_m^2 a_m} \int_0^L dr V(r) r^2 j_l \left(\frac{a_m r}{L} \right) \left(\frac{a_n r}{L} \right) c_n + \frac{2}{\pi L^3} \left[\sqrt{\left(\frac{a_m}{L} \right)^2 + m_q^2} + \sqrt{\left(\frac{a_m}{L} \right)^2 + m_{\bar{q}}^2} \right] \Delta a_m a_m^2 N_m^2 c_m, \quad (12)$$

where N_m is module of spherical Bessel function

$$N_m^2 = \int_0^L dr' r'^2 j_l \left(\frac{a_m r'}{L} \right)^2. \quad (13)$$

When L and N attain sufficiently large values, the solution tends to become nearly stationary [39, 40]. Eq. (12) is the eigenvalue equation in the matrix form, which is solved numerically. The eigenvalues correspond to masses of spin averaged states and the eigenvectors correspond to the wavefunction of these states. The spin-dependent interactions are introduced perturbatively. The spin dependent interaction potential is given by [41, 42]

$$V_{SD}(r) = V_{SS}(r) \vec{S}_q \cdot \vec{S}_{\bar{q}} + V_{LS}(r) \vec{L} \cdot \vec{S} + V_T(r) S_{12}. \quad (14)$$

V_{SS} is the spin singlet-triplet hyperfine splitting term given by

$$V_{SS}(r) = \frac{32\pi\alpha_s(r)}{9m_q^2} \tilde{\delta}_\sigma(r). \quad (15)$$

Here $\tilde{\delta}_\sigma(r) = \left(\frac{\sigma}{\sqrt{\pi}} \right)^3 e^{-\sigma^2 r^2}$ is the smeared delta function [43, 44]. To regularize the non-zero hyperfine splitting, smearing of delta function as Gaussian of width $1/\sigma$ is necessary [43, 44]. The spin orbit term V_{LS} and the tensor term V_T which describe the fine structure splitting of the states are given by

$$\begin{aligned} V_{LS}(r) &= \frac{1}{2m_q^2 r} \left(3V_V'(r) - V_S'(r) \right), \\ V_T(r) &= \frac{1}{2m_q^2} \left(\frac{V_V'(r)}{r} - V_V''(r) \right). \end{aligned} \quad (16)$$

The tensor operator $S_{12} = 3 \left(\vec{S}_q \cdot \hat{r} \right) \left(\vec{S}_{\bar{q}} \cdot \hat{r} \right) - \vec{S}_q \cdot \vec{S}_{\bar{q}}$, has non-vanishing diagonal matrix elements only between $L > 0$ spin-triplet states. The spin-dependent interactions are diagonal in a $|J, L, S\rangle$ basis with matrix elements given by [43–45]

$$\begin{aligned} \langle \vec{S}_q \cdot \vec{S}_{\bar{q}} \rangle &= \frac{1}{2} S^2 - \frac{3}{4}, \\ \langle \vec{L} \cdot \vec{S} \rangle &= \frac{1}{2} [J(J+1) - L(L+1) - S(S+1)], \\ \langle S_{12} \rangle &= \begin{cases} -\frac{L}{6(2L+3)} & J = L+1 \\ \frac{1}{6} & J = L \\ -\frac{L+1}{6(2L-1)} & J = L-1. \end{cases} \end{aligned} \quad (17)$$

Using the obtained normalized wave functions for the spin-averaged states, the spin-dependent corrections are evaluated perturbatively. The parameters used in our model are listed in Table I. The masses of S, P, D, F and G states of $b\bar{b}$ bound system are presented in Tables II-V, respectively.

III. DECAY CONSTANTS

Decay constants are fundamental parameters that characterize the strength of the weak interaction responsible for the decay processes, and it measures the probability amplitude to decay into lighter hadrons. The decay constant of pseudoscalar (f_P) and vector (f_V) states can be calculated using the Van Royen Weisskopf formula [46]

$$f_{P/V} = \sqrt{\frac{3|R_{P/V}(0)|^2}{\pi M_{P/V}}} \bar{C}(\alpha_s), \quad (18)$$

where $R_{P/V}(0)$ is the radial wavefunction at the origin for pseudoscalar (vector) meson state, $M_{P/V}$ is the mass of the pseudoscalar (vector) meson state and $\bar{C}(\alpha_s)$ is the QCD correction given by [47]

$$\bar{C}^2(\alpha_s) = 1 - \frac{\alpha_s(\mu)}{\pi} \left(\delta^{P,V} - \frac{m_q - m_{\bar{q}}}{m_q + m_{\bar{q}}} \ln \frac{m_q}{m_{\bar{q}}} \right), \quad (19)$$

where $\delta^P = 2$ and $\delta^V = 8/3$. The decay constant of P -wave states can be evaluated using [48, 49]

$$\begin{aligned} f_{\chi_0} &= 12 \sqrt{\frac{3}{8\pi m_q}} \left(\frac{|R'_{\chi_0}(0)|}{M_{\chi_0}} \right), \\ f_{\chi_1} &= 8 \sqrt{\frac{9}{8\pi m_q}} \left(\frac{|R'_{\chi_1}(0)|}{M_{\chi_1}} \right). \end{aligned} \quad (20)$$

Here M_{χ_0} and M_{χ_1} are the masses of χ_0 and χ_1 states, respectively. The decay constants for pseudoscalar f_P and vector f_V are presented in Table VI and decay constants for f_{χ_0} and f_{χ_1} are presented in Table VII.

IV. ANNIHILATION DECAYS

Bottomonium annihilation decays leave distinct signals in experimental data, allowing bottomonium states to be identified and characterized in high-energy collider experiments and precision spectroscopic investigations. The following section outlines and discusses various annihilation decays explored in this study:

A. Di-Leptonic Decays

The leptonic decay formula for S -wave (n^3S_1) and D -wave (n^3D_1) states are calculated using Van Royen-Weisskopf formula along with the QCD correction factor [46, 50–52]

$$\begin{aligned} \Gamma(n^3S_1 \rightarrow l^+l^-) &= \frac{4\alpha^2 e_q^2}{M(n^3S_1)^2} |R_{nS}(0)|^2 \left[1 - \frac{16\alpha_s(\mu)}{3\pi} \right], \\ \Gamma(n^3D_1 \rightarrow l^+l^-) &= \frac{25\alpha^2 e_q^2}{2m_q^4 M(n^3D_1)^2} |R''_{nD}(0)|^2, \end{aligned} \quad (21)$$

where $R'_{nL}(0)$ is the value of radial wavefunction at origin for nL state and $(')$ represents the order of derivative and $M(n^{2S+1}L_J)$ is the mass of $n^{2S+1}L_J$ state.

B. Di-Photonic Decays

The annihilation decays for S -wave (n^1S_0) and P -wave (n^3P_0 and n^3P_2) into two photons ($\gamma\gamma$) with first order QCD correction factors are given by [50, 51]

$$\begin{aligned}\Gamma(n^1S_0 \rightarrow \gamma\gamma) &= \frac{2^2 3\alpha^2 e_q^4}{M(n^1S_0)^2} |R_{nS}(0)|^2 \left[1 - \frac{3.4\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^3P_0 \rightarrow \gamma\gamma) &= \frac{2^4 27\alpha^2 e_q^4}{M(n^3P_0)^4} |R'_{nP}(0)|^2 \left[1 + \frac{0.2\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^3P_2 \rightarrow \gamma\gamma) &= \frac{2^4 36\alpha^2 e_q^4}{5M(n^3P_2)^4} |R'_{nP}(0)|^2 \times \left[1 - \frac{16\alpha_s(\mu)}{3\pi} \right].\end{aligned}\quad (22)$$

C. Tri-Photonic Decays

The annihilation decays for S -wave (n^3S_1) states into three photons ($\gamma\gamma\gamma$) with first order QCD correction factors are given by [50, 51]

$$\Gamma(n^3S_1 \rightarrow \gamma\gamma\gamma) = \frac{2^{24}(\pi^2 - 9)\alpha^3 e_q^6}{3\pi M(n^3S_1)^2} |R_{nS}(0)|^2 \left[1 - \frac{12.6\alpha_s(\mu)}{\pi} \right].\quad (23)$$

D. Di-Gluonic Decays

The annihilation decays for S -wave (n^1S_0), P -wave (n^3P_0 and n^3P_2), D -wave (n^1D_2), F -wave (n^3F_2 , n^3F_3 and n^3F_4) and G -wave (n^1G_4) states into two gluons (gg) with first order QCD correction factors are given by [50, 51, 53]

$$\begin{aligned}\Gamma(n^1S_0 \rightarrow gg) &= \frac{2^2 2\alpha_s^2(\mu)}{3M(n^1S_0)^2} |R_{nS}(0)|^2 \left[1 + \frac{4.8\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^3P_0 \rightarrow gg) &= \frac{2^4 6\alpha_s^2(\mu)}{M(n^3P_0)^4} |R'_{nP}(0)|^2 \left[1 + \frac{10\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^3P_2 \rightarrow gg) &= \frac{2^4 8\alpha_s^2(\mu)}{5M(n^3P_2)^4} |R'_{nP}(0)|^2 \left[1 - \frac{0.1\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^1D_2 \rightarrow gg) &= \frac{2^6 2\alpha_s^2(\mu)}{3\pi M(n^1D_2)^6} |R''_{nD}(0)|^2 \left[1 - \frac{2.2\alpha_s(\mu)}{\pi} \right], \\ \Gamma(n^3F_2 \rightarrow gg) &= \frac{2^8 919\alpha_s^2(\mu)}{135M(n^3F_2)^8} |R'''_{nF}(0)|^2, \\ \Gamma(n^3F_3 \rightarrow gg) &= \frac{2^8 20\alpha_s^2(\mu)}{27M(n^3F_3)^8} |R'''_{nF}(0)|^2, \\ \Gamma(n^3F_4 \rightarrow gg) &= \frac{2^8 20\alpha_s^2(\mu)}{27M(n^3F_4)^8} |R'''_{nF}(0)|^2, \\ \Gamma(n^1G_4 \rightarrow gg) &= \frac{2^{10} 2\alpha_s^2(\mu)}{3\pi M(n^1G_4)^{10}} |R''''_{nG}(0)|^2.\end{aligned}\quad (24)$$

E. Tri-Gluonic Decays

The annihilation decays for S -wave (n^3S_1), P -wave (n^1P_1) and D -wave (n^3D_1, n^3D_2 and n^3D_3) states into three gluons (ggg) with first order QCD correction factors are given by [50, 51, 54]

$$\begin{aligned}
\Gamma(n^3S_1 \rightarrow ggg) &= \frac{2^2 10 (\pi^2 - 9) \alpha_s^3(\mu)}{81 \pi M (n^3S_1)^2} |R_{nS}(0)|^2 \left[1 - \frac{4.9 \alpha_s(\mu)}{\pi} \right], \\
\Gamma(n^1P_1 \rightarrow ggg) &= \frac{2^4 20 \alpha_s^3(\mu)}{9 \pi M (n^1P_1)^4} |R'_{nP}(0)|^2 \ln(m_q \langle r \rangle), \\
\Gamma(n^3D_1 \rightarrow ggg) &= \frac{2^6 760 \alpha_s^3(\mu)}{81 \pi M (n^3D_1)^6} |R''_{nD}(0)|^2 \ln(4m_q \langle r \rangle), \\
\Gamma(n^3D_2 \rightarrow ggg) &= \frac{2^6 10 \alpha_s^3(\mu)}{9 \pi M (n^3D_2)^6} |R''_{nD}(0)|^2 \ln(4m_q \langle r \rangle), \\
\Gamma(n^3D_3 \rightarrow ggg) &= \frac{2^6 40 \alpha_s^3(\mu)}{9 \pi M (n^3D_3)^6} |R''_{nD}(0)|^2 \ln(4m_q \langle r \rangle).
\end{aligned} \tag{25}$$

F. Mixed Annihilation Decays

The annihilation decays for S -wave (n^3S_1) states via strong and electromagnetic interaction, into a photon and two gluons (γgg) are given by [44, 50, 55]

$$\Gamma(n^3S_1 \rightarrow \gamma gg) = \frac{2^2 8 (\pi^2 - 9) e_q^2 \alpha_s^3(\mu)}{9 \pi M (n^3S_1)^2} |R_{nS}(0)|^2 \left[1 - \frac{7.4 \alpha_s(\mu)}{\pi} \right]. \tag{26}$$

The annihilation decays for P -wave (n^3P_1) into a light flavour meson and a gluon ($q\bar{q}g$) are given by [50, 51]

$$\Gamma(n^3P_1 \rightarrow q\bar{q}g) = \frac{2^4 8 n_f \alpha_s^3(\mu)}{9 \pi M (n^3P_1)^4} |R'_{nP}(0)|^2 \ln(m_q \langle r \rangle), \tag{27}$$

where n_f is the number of flavors.

For all decays the strong coupling constant $\alpha_s(\mu)$ is calculated using the expression

$$\alpha_s(\mu) = \frac{4\pi}{\beta_0 \ln \frac{\mu^2}{\Lambda^2}} \left[1 - \frac{\beta_1 \ln \left(\ln \frac{\mu^2}{\Lambda^2} \right)}{\beta_0^2 \ln \frac{\mu^2}{\Lambda^2}} \right], \tag{28}$$

where $\beta_0 = 11 - (2/3)n_f$, $\beta_1 = 102 - (18/3)n_f$, Λ is the QCD constant, and μ is the reduced mass. All annihilation decay widths for $b\bar{b}$ bound system are presented in Tables VIII-XIV, respectively.

V. RADIATIVE TRANSITION

The bottomonium states have substantially heavier masses and their intrinsic compactness is more pronounced [56]. As a result, radiative transitions in bottomonium are expected to be dominant because of the favourable conditions for photon emission or absorption [56]. Radiative transitions serves as an effective means to detect, especially for states with higher quantum numbers that are hard to observe with traditional techniques. The leading-order electromagnetic transitions, the electric dipole ($E1$) and magnetic dipole ($M1$) transitions are discussed below.

A. E1 Transitions

The $E1$ radiative partial widths between the states ($n_i^{2S+1}L_{J_i}^i \rightarrow \gamma + n_f^{2S+1}L_{J_f}^f$) are given by [57, 58]

$$\Gamma_{E1}(i \rightarrow \gamma + f) = \frac{4\alpha_e^2}{3} E_\gamma^3 \frac{E_f}{M_i} C_{fi} \delta_{S_f S_i} |\epsilon_{fi}|^2. \tag{29}$$

Here $\alpha = 1/137$ is the fine structure constant, e_q is the quark charge, E_f is the energy of the final state, M_i is the mass of the initial state, $E_\gamma = (M_i^2 - M_f^2)/2M_i$ is the emitted photon energy. M_f is the mass of the final state. E_f/M_i is the relativistic phase factor and C_{fi} is the statistical factor given by

$$C_{fi} = \max(L_i, L_f) (2J_f + 1) \left\{ \begin{matrix} J_i & 1 & J_f \\ L_f & S & L_i \end{matrix} \right\}^2, \quad (30)$$

where $\{:::\}$ are the $6j$ symbol. In Eq. (29), ϵ_{fi} is the overlapping integral determined using the initial $R_{n_i l_i}(r)$ and final state $R_{n_f l_f}(r)$ wavefunctions:

$$\epsilon_{fi} = \frac{3}{E_\gamma} \int_0^\infty dr R_{n_i l_i}(r) R_{n_f l_f}(r) \left[\frac{E_\gamma r}{2} j_0 \left(\frac{E_\gamma r}{2} \right) - j_1 \left(\frac{E_\gamma r}{2} \right) \right] \quad (31)$$

The $E1$ transitions widths for S, P, D, F and G wave states for $b\bar{b}$ bound system are presented in Tables XV-XIX, respectively.

B. M1 Transitions

The $M1$ radiative partial widths between the states $(n_i^{2S_i+1} L_{J_i} \rightarrow \gamma + n_f^{2S_f+1} L_{J_f})$ are given by [43, 57, 59]

$$\Gamma_{M1}(i \rightarrow \gamma + f) = \frac{4\alpha\mu_q^2}{3} \frac{2J_f + 1}{2L + 1} E_\gamma^3 \frac{E_f}{M_i} |m_{fi}|^2 \delta_{L_f L_i} \delta_{S_f S_i \pm 1}, \quad (32)$$

where m_{fi} is given by

$$m_{fi} = \int_0^\infty dr R_{n_i l_i}(r) R_{n_f l_f}(r) \left[j_0 \left(\frac{E_\gamma r}{2} \right) \right], \quad (33)$$

and μ_q is the magnetic dipole moment given by [55]

$$\mu_q = \frac{m_{\bar{q}} e_q - m_q e_{\bar{q}}}{2m_q m_{\bar{q}}}. \quad (34)$$

The $M1$ transitions widths for S and P wave states for $b\bar{b}$ bound system are presented in Table XX.

VI. S-D MIXING

In bottomonium, the proximity of energy levels of higher excited states with same J^{PC} can result in mixing of states. The mixing is caused by the tensor potential term, but it is not strong enough to induce substantial mixing [60, 61]. However, for the states above open flavor threshold the mixing can be caused by coupled-channel dynamics, threshold effects, meson exchange, and multi-gluon exchange interactions [62–65]. These effects can modify the wavefunctions, causing mass shift and mixing between states, as well as affecting decay properties such as open channel strong decay, leptonic decays etc. Consequently, the conventional representation of bottomonium states as pure S and D wavefunctions breaks down, and the states are instead identified as admixtures of both components. The mixed states can be represented in terms of pure $|nS\rangle$ and $|n'D\rangle$ states as [61]

$$\begin{aligned} |\phi\rangle &= \cos\theta |nS\rangle + \sin\theta |n'D\rangle, \\ |\phi'\rangle &= -\sin\theta |nS\rangle + \cos\theta |n'D\rangle, \end{aligned} \quad (35)$$

where $|\phi\rangle$ and $|\phi'\rangle$ are the mixed states, and θ is the mixing angle. The masses of the mixed states can be calculated using [61]

$$\begin{aligned} M_\phi &= \left[\left(\frac{M_{nS} + M_{n'D}}{2} \right) + \left(\frac{M_{nS} - M_{n'D}}{2 \cos 2\theta} \right) \right], \\ M_{\phi'} &= \left[\left(\frac{M_{nS} + M_{n'D}}{2} \right) + \left(\frac{M_{n'D} - M_{nS}}{2 \cos 2\theta} \right) \right]. \end{aligned} \quad (36)$$

Here M_ϕ and $M_{\phi'}$ are the masses of the mixed states, and M_{nS} and $M_{n'D}$ are the masses of the corresponding pure S and D states.

The leptonic decay widths of the mixed states are given by [61, 66]

$$\begin{aligned}\Gamma_\phi &= \left[\frac{2\alpha e_q}{M_{nS}} |R_{nS}(0)| \cos\theta + \frac{5\alpha e_q}{\sqrt{2}m_q^2 M_{n'D}} |R''_{n'D}(0)| \sin\theta \right]^2, \\ \Gamma_{\phi'} &= \left[\frac{5\alpha e_q}{\sqrt{2}m_q^2 M_{n'D}} |R''_{n'D}(0)| \cos\theta - \frac{2\alpha e_q}{M_{nS}} |R_{nS}(0)| \sin\theta \right]^2.\end{aligned}\quad (37)$$

The leptonic decay of the mixed states is fitted to the experimental data to obtain the mixing angle, which is then used to calculate the masses of the mixed states. Our results of $S - D$ mixing are presented in Table XXI.

VII. RESULTS AND DISCUSSION

In this study, a screened potential model within a relativistic framework is employed to compute the mass spectrum and decay widths of $b\bar{b}$ bound system. The fitted parameters presented in Table I are determined using the chi-square fitness test.

The masses of S -wave states are presented in Table II and are compared with the experimental data and other theoretical models. The $1S$ and $2S$ triplet and singlet states of bottomonium have been firmly established through experiments and serve as valuable tools for validating theoretical models. The masses of $1S$ bottomonium states in our model are evaluated to be $\eta_b(1S) = 9406.4$ MeV and $\Upsilon(1S) = 9451.1$ MeV, and for $2S$ bottomonium states are, $\eta_b(2S) = 9998.9$ MeV and $\Upsilon(2S) = 10023.8$ MeV. The hyperfine mass splitting of singlet and triplet states is given by $\Delta m(nS) = m[\Upsilon(nS)] - m[\eta(nS)]$. Experimentally the hyperfine mass splitting are determined to be $\Delta m(1S) = 62.3 \pm 3.2$ MeV and $\Delta m(2S) = 24.3 \pm 3.5^{+2.8}_{-1.9}$ MeV [67]. In our model these value are evaluated to be $\Delta m(1S) = 44.7$ MeV and $\Delta m(2S) = 24.9$ MeV, consistent with the experimentally observed values. The identification of $\Upsilon(10355)$ as the $\Upsilon(3S)$ bottomonium state has been well established in the literature. In our model the mass values evaluated for $\Upsilon(3S)$ bottomonium state is 10394.2 MeV. The mass difference, $m[\Upsilon(3S)] - m[\Upsilon(nS)] = 331.50 \pm 0.02 \pm 0.13$ MeV is determined experimentally [67], which comes out to be 370.4 MeV in our model. Traditionally, the $\Upsilon(10580)$ is recognized as the $\Upsilon(4S)$ bottomonium state in the literature [56, 58, 59]. In our model the mass of $\Upsilon(4S)$ is calculated to be 10688.1 MeV which is overestimated by 108.7 MeV compared to experimental value. This discrepancy of overestimating the mass value is consistently observed across all the potential models [58, 59]. In Ref. [68], a 3P_0 model was employed to analyze $\Upsilon(10580)$. They concluded that $\Upsilon(10580)$ exhibits a significant meson-meson component due its proximity with the $B^*\bar{B}^*$ states. As outlined in Ref. [69], the state discovered by CLEO Collaboration with mass of $10684 \pm 10 \pm 8$ MeV and labelled as $b\bar{b}g$ hybrid [70], is deemed a more fitting assignment for $\Upsilon(10580)$ bottomonium state. This finding is also corroborated by our model. The presence of intermediate $B^*\bar{B}^*$ states could potentially lead to observable $S - D$ mixing within the $\Upsilon(10580)$ state [68]. This possibility is studied in Ref. [61] and they predicted $\Upsilon(10580)$ to be $\Upsilon(4S) - \Upsilon(3D)$ mixture state with a substantial mixing angle. The $\Upsilon(10860)$ and $\Upsilon(11020)$ states are typically associated with the $\Upsilon(5S)$ and $\Upsilon(6S)$ bottomonium states, respectively [58]. In our potential model, the mass of $\Upsilon(5S)$ and $\Upsilon(6S)$ bottomonium states are computed to be 10938.9 MeV and 11160.9 MeV, respectively, which is overestimated by 53.7 MeV and 160.9 MeV, respectively. In most theoretical model it is observed that the mass values of $\Upsilon(5S)$ and $\Upsilon(6S)$ often exhibit discrepancies, either through overestimation or underestimation. Various interpretations have been explored in the literature to address this issue. The $\Upsilon(10860)$ is considered as mixture of $\Upsilon(5S)$ and lowest P wave hybrid [71]. In a lattice QCD static-light potential they could not confirm if $\Upsilon(11020)$ corresponds to $\Upsilon(S)$ or $\Upsilon(D)$ state [72]. In a 3P_0 model study, they concluded that $\Upsilon(10860)$ and $\Upsilon(11020)$ are structures are mainly $b\bar{b}$ states with small $S - D$ mixing component [68]. This aspect was analyzed in [73], proposing that $\Upsilon(10860)$ can be a $\Upsilon(5S) - \Upsilon(4D)$ mixture state, and in [61] it is suggested that both $\Upsilon(10860)$ and $\Upsilon(11020)$ are $\Upsilon(5S) - \Upsilon(4D)$ mixture state. Nevertheless, more experimental data is required to understand the nature of $\Upsilon(10860)$ and $\Upsilon(11020)$ states. We discuss the possibility of $S - D$ mixing in $\Upsilon(10580)$, $\Upsilon(10860)$ and $\Upsilon(11020)$ later in this section.

Masses of P -wave states are presented in Table III and compared with the experimental data and other theoretical models. The $1P$ and $2P$ states of bottomonium have been observed experimentally and our evaluated masses of these states are in accordance with the experimentally determined values. The experimentally determined mass difference are $m[\chi_{b2}(1P)] - m[\chi_{b1}(1P)] = 19.10 \pm 0.25$ MeV, $m[\chi_{b1}(1P)] - m[\chi_{b0}(1P)] = 32.49 \pm 0.93$ MeV, $m[\chi_{b2}(2P)] - m[\chi_{b1}(2P)] = 13.10 \pm 0.24$ MeV and $m[\chi_{b1}(2P)] - m[\chi_{b0}(2P)] = 23.8 \pm 1.7$ MeV [67]. In our model these values are evaluated to be 19.9 MeV, 27 MeV, 16.1 MeV and 21.3 MeV, respectively, exhibiting good agreement with the experimental data. Among $3P$ states of bottomonium only $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ have been identified. The masses of

TABLE II. S wave mass spectra of $b\bar{b}$ states (in MeV)

States	Ours	Exp[67]	[58]	[53]	[59]	[56]	[74]
1^1S_0	9406.4	9398.7	9398	9402	9423	9412.22	9390
2^1S_0	9998.9	9999.0	9989	9976	9983	9995.48	9990
3^1S_0	10374.9		10336	10336	10342	10339.00	10326
4^1S_0	10671.8		10597	10635	10638	10572.49	10584
5^1S_0	10924.5		10810	10869	10901	10746.76	10800
6^1S_0	11147.9		10991	11097	11140	11064.47	10988
1^3S_1	9451.1	9460.3	9463	9465	9463	9460.75	9460
2^3S_1	10023.8	10023.3	10017	10003	10001	10026.22	10015
3^3S_1	10394.2	10355.1	10356	10354	10354	10364.65	10343
4^3S_1	10688.1	10579.4	10612	10635	10650	10594.47	10597
5^3S_1	10938.9	10885.2	10822	10878	10912	10766.14	10811
6^3S_1	11160.9	11000.0	11001	11102	11151	11081.70	10997

TABLE III. P wave mass spectra of $b\bar{b}$ states (in MeV)

States	Ours	Exp[67]	[58]	[53]	[59]	[56]	[74]
1^1P_1	9872.9	9899.3	9894	9882	9899	9874.56	9909
2^1P_1	10271.7	10259.8	10259	10250	10268	10270.00	10254
3^1P_1	10582.7		10530	10541	10570	10526.50	10519
4^1P_1	10845.6		10751	10790		10714.80	
5^1P_1	11077.1		10938	11016		10863.00	
1^3P_0	9838.7	9859.4	9858	9847	9874	9849.61	9864
2^3P_0	10244.9	10232.5	10235	10226	10248	10252.54	10220
3^3P_0	10559.4		10513	10522	10551	10512.88	10490
4^3P_0	10824.5		10736	10775		10703.56	
5^3P_0	11057.4		10926	11004		10853.38	
1^3P_1	9865.7	9892.8	9889	9876	9894	9871.47	9903
2^3P_1	10266.2	10255.5	10255	10246	10265	10267.86	10249
3^3P_1	10578.1	10513.4	10527	10538	10567	10524.84	10515
4^3P_1	10841.5		10749	10788		10713.44	
5^3P_1	11073.3		10936	11014		10861.83	
1^3P_2	9885.6	9912.2	9910	9897	9907	9881.40	9921
2^3P_2	10282.3	10268.6	10269	10261	10274	10274.77	10264
3^3P_2	10592.3	10524.0	10539	10550	10576	10530.21	10528
4^3P_2	10854.6		10758	10798		10717.86	
5^3P_2	11085.6		10944	11022		10865.62	

$\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ in our model are evaluated to be 10578.1 MeV and 10592.3 MeV, respectively. Our mass values for $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ states are higher by 64.7 MeV and 68.9 MeV, respectively. This discrepancy can be due to proximity of these states with open flavor threshold $B\bar{B}^*$, which could cause mixing in these states [75, 76]. The experimentally measured mass difference $m[\chi_{b2}(3P)] - m[\chi_{b1}(3P)] = 10.60 \pm 0.64 \pm 0.17$ MeV [67], which is calculated to be 14.2 MeV in our model.

Masses of D -wave states are presented in Table IV and compared with the experimental data and other theoretical models. The mass of $\Upsilon_1(1D)$ bottomonium state in our model is evaluated to be 10147.9 MeV, showing a slight deviation of 15.8 MeV from the experimentally measured value [67]. The $\Upsilon_2(1D)$ and $\Upsilon_3(1D)$ states are also anticipated to be observed experimentally, with an estimated mass values of 10.13 GeV and 10.18 GeV, respectively [15]. The values obtained from our calculations for these states are 10139.4 MeV and 10154.2 MeV, respectively. The recently observed $\Upsilon(10753)$ is commonly associated with $\Upsilon_1(3D)$ according to existing literature [58, 76], along with alternative interpretations such as a tetraquark [77, 78], hybrid meson [3], etc. The mass of $\Upsilon_1(3D)$ bottomonium state in our model is evaluated to be 10741.3 MeV, which is in alignment with experimental value [67]. Through a reanalysis of BABAR data, the mass of $\Upsilon_1(2D)$ was estimated to be 10495 ± 5 MeV with significance of 10.7σ [79]. Our model's estimation of 10467.3 MeV is consistent with their prediction.

Masses of F and G - wave states are presented in Table V and compared with other theoretical models. Presently there is no experimental data for F and G states. The masses for lower F and G states in our models are consistent

TABLE IV. D wave mass spectra of $b\bar{b}$ states (in MeV)

States	Ours	Exp[67]	[58]	[53]	[59]	[56]	[74]
1^1D_2	10149.1		10163	10148	10149	10153.80	10153
2^1D_2	10476.3		10450	10450	10465	10456.60	10432
3^1D_2	10749.9		10681	10706	10740	10664.70	
4^1D_2	10989.2		10876	10935	10988	10823.00	
5^1D_2	11204.1		11046			10952.60	
1^3D_1	10139.4		10153	10138	10145	10144.99	10146
2^3D_1	10467.3		10442	10441	10462	10450.23	10425
3^3D_1	10741.3	10752.7	10675	10698	10736	10659.68	
4^3D_1	10981.0		10871	10928	10985	10818.83	
5^3D_1	11196.1		11041			10949.01	
1^3D_2	10147.9	10163.7	10162	10147	10149	10152.77	10153
2^3D_2	10475.0		10450	10449	10465	10455.86	10432
3^3D_2	10748.7		10681	10705	10740	10664.12	
4^3D_2	10987.9		10876	10934	10988	10822.52	
5^3D_2	11202.8		11045			10951.59	
1^3D_3	10154.2		10170	10155	10150	10158.31	10157
2^3D_3	10481.1		10456	10455	10466	10459.85	10436
3^3D_3	10754.6		10686	10711	10741	10667.25	
4^3D_3	10993.7		10880	10939	10990	10825.12	
5^3D_3	11208.5		11049			10954.42	

TABLE V. F and G wave mass spectra of $b\bar{b}$ states (in MeV)

States	Ours	[58]	[53]	States	Ours	[58]	[53]
1^1F_3	10366.8	10366	10355	1^1G_4	10552.9	10534	10530
2^1F_3	10652.2	10609	10619	2^1G_4	10809.8	10747	10770
3^1F_3	10900.0	10812	10853	3^1G_4	11038.2	10929	
4^1F_3	11121.5	10988		4^1G_4	11245.3		
5^1F_3	11323.0			5^1G_4	11435.6		
1^3F_2	10363.6	10362	10350	1^3G_3	10552.8	10533	10529
2^3F_2	10648.8	10605	10615	2^3G_3	10809.2	10745	10769
3^3F_2	10896.5	10809	10850	3^3G_3	11037.3	10928	
4^3F_2	11117.8	10985		4^3G_3	11244.1		
5^3F_2	11319.2			5^3G_3	11434.3		
1^3F_3	10366.8	10366	10355	1^3G_4	10553.4	10535	10531
2^3F_3	10652.2	10609	10619	2^3G_4	10810.1	10747	10770
3^3F_3	10899.9	10812	10853	3^3G_4	11038.4	10929	
4^3F_3	11121.3	10988		4^3G_4	11245.5		
5^3F_3	11322.7			5^3G_4	11435.8		
1^3F_4	10368.5	10369	10358	1^3G_5	10552.6	10536	10532
2^3F_4	10654.1	10612	10622	2^3G_5	10809.8	10748	10772
3^3F_4	10902.1	10815	10856	3^3G_5	11038.5	10931	
4^3F_4	11123.7	10990		4^3G_5	11245.9		
5^3F_4	11325.3			5^3G_5	11436.4		

with other theoretical models, but for higher excited states there is deviation. The masses for different J states are very close to each other which could make it harder to differentiate between these states experimentally.

Decay constants of pseudoscalar (f_P), vector (f_V) and tensor (f_{χ_0}, f_{χ_1}) states are presented in Tables VI and VII, respectively, and are compared with the experimental data and other theoretical models. Our calculated values for the vector decay constants (f_V) are in accord with experimental values and shows more consistency over other theoretical models.

The di-leptonic decay widths $\Gamma(l^+l^-)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor for $\Upsilon(nS)$ and $\Upsilon(nD)$ states are presented in Table VIII and are compared with experimental data and other theoretical models. The di-leptonic decay widths of $\Upsilon(nS)$ states evaluated without the correction term are more in agreement with the experimental value. On the contrary, the incorporation of the correction factor results in significant suppres-

TABLE VI. Pseudoscalar and vector decay constants (in MeV)

States	$f_{P/V}$	Exp[67]	[59]	[56]	[80]	[81]
1^1S_0	655.9		529	578.21	646.025	744
2^1S_0	489.2		317	499.48	518.803	577
3^1S_0	431.8		280	450.35	474.954	511
4^1S_0	398.6		264	413.93	449.654	471
5^1S_0	375.5		255	385.68	432.072	443
6^1S_0	357.6		249	360.93	418.645	422
1^3S_1	640.2	715 ± 5	530	551.53	647.250	706
2^3S_1	478.0	498 ± 8	317	477.05	519.436	547
3^3S_1	422.0	430 ± 4	280	430.42	475.440	484
4^3S_1	389.7	336 ± 18	265	395.80	450.066	446
5^3S_1	349.7		255	368.91	432.437	419
6^3S_1	335.3		249	345.40	418.977	399

TABLE VII. Decay constants of P -wave states (in MeV)

States	f_{X_0}	States	f_{X_1}
1^3P_0	227.8	1^3P_1	262.4
2^3P_0	248.7	2^3P_1	286.6
3^3P_0	257.3	3^3P_1	296.6
4^3P_0	262.0	4^3P_1	302.1
5^3P_0	264.8	5^3P_1	305.4

sion in the decay widths. The di-leptonic decay widths of $\Upsilon(nD)$ are comparatively smaller than those of $\Upsilon(nS)$ by factor of 1000. This discrepancy acts as a distinguishing factor between $\Upsilon(nS)$ and $\Upsilon(nD)$ in most models [56, 58]. This difference in di-leptonic decay widths between $\Upsilon(nS)$ and $\Upsilon(nD)$ states is used as a justification for assigning $\Upsilon(10580)$, $\Upsilon(10860)$ and $\Upsilon(11020)$ states to the $\Upsilon(4S)$, $\Upsilon(5S)$, and $\Upsilon(6S)$, respectively, in potential models. This assignment is substantiated by their large di-leptonic decay widths of 0.272 ± 0.029 keV, 0.31 ± 0.07 keV and 0.13 ± 0.03 keV, respectively. Given that $\Upsilon(10580)$, $\Upsilon(10860)$ and $\Upsilon(11020)$ cannot be purely D states but exhibit characteristics of S states, their widths are often overestimated in models, suggesting a potential for $S - D$ mixing component [61]. For $n \geq 3$ the probability of $S - D$ mixing in bottomonium states increases significantly and even a small mixing angle can increase the di-leptonic decay widths of $\Upsilon(nD)$ by order of 2 [82]. To study $S - D$ mixing in our model, the di-leptonic decay widths without the correction factor are utilized to obtain the mixing angle.

The di-photonic decay widths $\Gamma(\gamma\gamma)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor for $\eta_b(nS)$, $\chi_{b0}(nP)$ and $\chi_{b2}(nP)$ states are listed in Table IX and are compared with other theoretical models. Our di-photonic decay widths are comparable to the results obtained in Ref. [59, 80], but compared to results in Ref. [53, 58], our decay widths are lower in magnitude. The di-photonic decay width of $\eta_b(1S)$ is not seen experimentally and we predict it to be 0.344 keV as our estimate.

TABLE VIII. Di-leptonic decay widths (in keV for S states and in eV for D states)

States	Γ	Γ_{cf}	Exp[67]	[61]	[58]	[53]	[56]	[59]
1^3S_1	1.268	0.883	1.34 ± 0.018	1.370	1.65	1.44	0.7700	0.582
2^3S_1	0.666	0.464	0.612 ± 0.011	0.626	0.821	0.73	0.5442	0.197
3^3S_1	0.501	0.349	0.443 ± 0.008	0.468	0.569	0.53	0.4288	0.149
4^3S_1	0.415	0.289	0.272 ± 0.029	0.393	0.431	0.39	0.3549	0.129
5^3S_1	0.360	0.251	0.31 ± 0.07	0.346	0.348	0.33	0.3035	0.117
6^3S_1	0.320	0.223	0.13 ± 0.03	0.313	0.286	0.27	0.2586	0.109
1^3D_1	1.149			2.0	1.88	1.38	5.0	1.65
2^3D_1	2.166			3.0	2.81	1.99	5.8	2.42
3^3D_1	3.059			5.0	3.00	2.38	5.9	3.19
4^3D_1	4.573			6.0	3.00	2.18	5.8	3.97
5^3D_1	5.219			8.0	3.02		5.7	

TABLE IX. Di-photonic decay widths (in keV)

States	Γ	Γ_{cf}	[58]	[53]	[56]	[80]	[59]
1^1S_0	0.426	0.344	1.05	0.94	0.3035	0.387	0.2361
2^1S_0	0.223	0.180	0.489	0.41	0.2122	0.263	0.0896
3^1S_0	0.168	0.135	0.323	0.29	0.1668	0.229	0.0726
4^1S_0	0.139	0.112	0.237	0.20	0.1378	0.212	0.0666
5^1S_0	0.120	0.097	0.192	0.17	0.1176	0.201	0.0636
6^1S_0	0.107	0.086	0.152	0.14	0.1000	0.193	0.0619
1^3P_0	0.042	0.042	0.199	0.15	0.1150	0.0196	0.0168
2^3P_0	0.046	0.047	0.205	0.15	0.1014	0.0195	0.0172
3^3P_0	0.046	0.047	0.180	0.13	0.0875	0.0194	0.0192
4^3P_0	0.045	0.046	0.157	0.13	0.0768	0.0192	
5^3P_0	0.044	0.046	0.146		0.0686	0.0191	
1^3P_2	0.011	0.007	0.0106	0.0093	0.0147	0.0052	0.0024
2^3P_2	0.012	0.008	0.0133	0.012	0.0131	0.0052	0.0025
3^3P_2	0.012	0.009	0.0141	0.013	0.0114	0.0051	0.0027
4^3P_2	0.012	0.008	0.0142	0.015	0.0100	0.0051	
5^3P_2	0.011	0.008	0.0143		0.0090	0.0050	

TABLE X. Tri-photonic decay widths (in 10^{-3} eV)

States	Γ	Γ_{cf}	[58]	[53]	[83]	[59]
1^3S_1	42.16	11.92	19.4	17.0	3.44	30.67
2^3S_1	22.17	6.27	10.9	9.8	2.00	11.58
3^3S_1	16.66	4.71	8.04	7.6	1.55	9.376
4^3S_1	13.81	3.91	6.36	6.0	1.29	8.590
5^3S_1	11.98	3.39	5.43		1.10	8.206
6^3S_1	10.65	3.01	4.57		0.96	7.982

The tri-photonic decay widths $\Gamma(\gamma\gamma\gamma)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor are calculated for $\Upsilon(nS)$ states in Table X and are compared with other theoretical models. The values of tri-photonic decay widths vary significantly among models. Experimental data is needed for validation.

The di-gluonic decay widths $\Gamma(gg)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor are calculated for $\eta_b(nS)$, $\chi_{b0}(nP)$, $\chi_{b2}(nP)$, and $\eta_b(nD)$ in Table XI and $\chi_{b2}(nF)$, $\chi_{b3}(nF)$, $\chi_{b4}(nF)$ and $\eta_b(nG)$ in Table XII and are compared with other theoretical models. Our di-gluonic decay widths of S , P and D states are comparable to [56]. Although they are 2-4 times smaller in comparison to results from references [53, 58, 59, 83]. In the case of the lower-lying $\eta_b(nS)$ states, the di-gluonic decay widths constitute approximately $\sim 100\%$ of their total width, due to suppression of OZI-allowed two-body strong decay channels [58]. Our di-gluonic decay width of $\eta_b(1S)$ is evaluated to be 5.763 MeV, which close to the error margin of total decay width 10.0^{+5}_{-4} MeV [67].

The tri-gluonic decay widths $\Gamma(ggg)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor are estimated for $\Upsilon(nS)$, $h_b(nP)$, $\Upsilon_1(nD)$, $\Upsilon_2(nD)$ and $\Upsilon_3(nD)$ in Table XIII and are compared with experimental data and other theoretical models. The tri-gluonic decay width for $\Upsilon(1S)$ is lower by 13.95 MeV, but the widths for $\Upsilon(2S)$ and $\Upsilon(3S)$ are in good agreement with the experimental results. The tri-gluonic decay widths values for P and D states are lower compared to predictions of other models.

The photo-gluonic decay widths $\Gamma(\gamma gg)$ for $\Upsilon(nS)$ and quark-gluonic decay width $\Gamma(q\bar{q}g)$ for $\chi_{b1}(nP)$ of bottomonium states with (Γ_{cf}) and without (Γ) the correction factor are evaluated in Table XIV and are compared with experimental data and other theoretical models. The photo-gluonic decay widths of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ are in accordance with the experimental data. The multi-gluon or hybrid $q\bar{q}g$ decays are dominant channel for $\chi_{b1}(1P)$ state [58]. Our predictions for quark-gluonic decay width for $\chi_{b1}(1P)$ are observed to be lower compared to other models.

The S wave $E1$ transitions widths in Table XV are calculated for $\Gamma(2S \rightarrow \gamma P)$, $\Gamma(3S \rightarrow \gamma P)$ and $\Gamma(4S \rightarrow \gamma P)$. The transition widths of $\Gamma(2S \rightarrow \gamma\chi_b(1P))$ in our model exhibit consistency with experimental results. The transition widths for $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(P))$ presents a complex scenario, due to discrepancies in the predictions of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(2P))$ and $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(1P))$ across various models [58, 83]. In our model, the transition width $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(2P))$ are estimated slightly higher than the experimentally measured values. The $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(1P))$ transition widths are highly suppressed. This is typical characteristic of $E1$ transitions among states separated by two radial nodes, which makes them susceptible to relativistic corrections [84, 85]. This suppression is also evident in our

TABLE XI. Di-gluonic decay widths of S , P (in MeV) and D (in keV) states

States	Γ	Γ_{cf}	[58]	[53]	[83]	[59]	[56]
1^1S_0	4.608	5.763	17.9	16.6	20.18	11.326	6.8520
2^1S_0	2.412	3.016	8.33	7.2	10.64	4.301	5.2374
3^1S_0	1.811	2.264	5.51	4.9	7.94	3.485	4.3182
4^1S_0	1.500	1.876	4.03	3.4		3.193	3.6829
5^1S_0	1.301	1.627	3.26			3.051	3.2196
6^1S_0	1.156	1.446	2.59			2.968	2.8519
1^3P_0	0.454	0.713	3.37	2.6	2.00	1.34	1.4297
2^3P_0	0.499	0.783	3.52	2.6	2.37	1.39	1.2358
3^3P_0	0.503	0.789	3.10	2.2	2.46	1.54	1.0539
4^3P_0	0.496	0.779	2.73	2.1			0.9175
5^3P_0	0.486	0.763	2.54				0.8127
1^3P_2	0.119	0.118	0.165	0.147	0.837	0.209	0.2370
2^3P_2	0.131	0.130	0.220	0.207	0.104	0.215	0.2064
3^3P_2	0.132	0.132	0.243	0.227	0.111	0.240	0.1767
4^3P_2	0.131	0.130	0.251	0.248			0.1543
5^3P_2	0.128	0.127	0.258				0.1370
1^1D_2	0.321	0.281	0.657	1.8	0.37	0.489	
2^1D_2	0.534	0.468	1.22	3.3	0.67	0.764	
3^1D_2	0.679	0.595	1.59	4.7		1.06	
4^1D_2	0.785	0.686	1.86			1.38	
5^1D_2	0.861	0.754	2.13				

TABLE XII. Di-gluonic decay widths of F (in keV) and G (in eV) states

States	Γ	[58]	[53]	States	Γ	[58]	[53]
1^3F_2	0.282	0.834	0.70	1^3F_4	0.031	0.05	0.048
2^3F_2	0.618	2.04	1.77	2^3F_4	0.067	0.126	0.13
3^3F_2	0.946	3.17		3^3F_4	0.102	0.210	
4^3F_2	1.248			4^3F_4	0.135		
5^3F_2	1.517			5^3F_4	0.164		
1^3F_3	0.031	0.0672	0.060	1^1G_4	0.289	0.661	2.3
2^3F_3	0.067	0.167	0.16	2^1G_4	0.778		
3^3F_3	0.103	0.270		3^1G_4	1.383		
4^3F_3	0.135			4^1G_4	2.044		
5^3F_3	0.165			5^1G_4	2.723		

evaluation, where $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(1P)) = 0.057$ keV and $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P)) = 0.139$ keV show good alignment with the experimental results, whereas the estimate of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P)) = 0.115$ keV exceeds the experimental value. This atypical hierarchy of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b2}(1P)) > \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b0}(1P)) > \Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))$ mentioned in ref. [86] is also observed in our model. This is said to arise from the mixing of $\chi_{b1}(1P)$ with the $\chi_b(2P)$ and $\chi_b(3P)$ states, resulting in further suppression of the transition width of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_{b1}(1P))$. In ref. [61], the $S - D$ mixing in the $\Upsilon(3S)$ state is considered to explain the $E1$ transitions widths of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(2P))$, which enabled them to reproduce the experimental values in their estimations. This explanation may also be extended for analysis of $\Gamma(\Upsilon(3S) \rightarrow \gamma\chi_b(1P))$ transition widths. We also evaluate the $\Gamma(4S \rightarrow \gamma P)$ for future reference.

The P wave $E1$ transitions widths are presented for $\Gamma(1P \rightarrow \gamma S)$, $\Gamma(2P \rightarrow \gamma S)$, $\Gamma(2P \rightarrow \gamma D)$, $\Gamma(3P \rightarrow \gamma S)$ and $\Gamma(3P \rightarrow \gamma D)$ in Table XVI and XVII. The transition width of $\Gamma(1P \rightarrow \gamma S)$ calculated in our model are consistent with the experimental results and other theoretical models. Using the measured branching ratios of $B[\chi_{b0}(1P) \rightarrow \gamma\Upsilon(1S)] = 1.94 \pm 0.27\%$, $B[\chi_{b1}(1P) \rightarrow \gamma\Upsilon(1S)] = 35.2 \pm 2.0\%$, and $B[\chi_{b2}(1P) \rightarrow \gamma\Upsilon(1S)] = 18.0 \pm 1.0\%$ [67], we calculate the total decay width to be 1.19 MeV for $\chi_{b0}(1P)$, 79.0 keV for $\chi_{b1}(1P)$, and 177.0 keV for $\chi_{b2}(1P)$, respectively. Our evaluation of total decay width for $\chi_{b0}(1P)$ is consistent with value of 1.3 ± 0.9 MeV and the condition $\Gamma_{total} < 2.4$ MeV predicted by Belle Collaboration [87]. The $h_b(1P) \rightarrow \gamma\eta_b(1S)$ is the primary transition of $h_b(1P)$ and its branching ratio is measured to be $B[h_b(1P) \rightarrow \gamma\eta_b(1S)] = 52^{+6}_{-5}\%$ [67]. From the branching ratio we determine the total decay width of $h_b(1P)$ to be 74.0 keV, which is consistent with the Ref. [74]. The evaluated transition width of $h_b(2P) \rightarrow \gamma\eta_b(2S)$ is lower than the experimental value. This trend is observed in most of the potential models.

TABLE XIII. Tri-gluonic decay widths (in keV)

States	Γ	Γ_{cf}	Exp[67]	[58]	[53]	[56]	[83]
1^3S_1	41.85	30.18	44.13±1.09	50.8	47.6	28.5	41.63
2^3S_1	22.00	15.86	18.8±1.59	28.4	26.3	19.3	24.25
3^3S_1	16.54	11.92	7.25±0.85	21.0	19.8	14.8	18.76
4^3S_1	13.71	9.89		16.7	15.1	12.1	15.58
5^3S_1	11.89	8.58		14.2	13.1	10.2	13.33
6^3S_1	10.57	7.62		12.0	11.0	8.5	11.57
1^1P_1	20.10			44.7	37.0	35.7	35.26
2^1P_1	26.99			64.6	54.0	34.6	52.70
3^1P_1	30.23			71.1	59.0	33.1	62.16
4^1P_1	31.99			73.2	64.0	32.7	
5^1P_1	32.97			76.2		30.9	
1^3D_1	3.11			10.4	8.11	10.6	9.97
2^3D_1	5.61			20.1	14.8	11.9	9.69
3^3D_1	7.54			26.0	21.2	11.8	
4^3D_1	9.05			30.4		11.3	
5^3D_1	1.02			34.7		10.8	
1^3D_2	0.37			0.821	0.69		0.62
2^3D_2	0.66			1.65	1.4		0.61
3^3D_2	0.89			2.27	2.0		
4^3D_2	0.11			2.75			
5^3D_2	0.12			3.23			
1^3D_3	1.46			2.19	2.07	6.0	0.22
2^3D_3	2.64			4.56	4.3	5.6	1.25
3^3D_3	3.54			6.65	6.6	5.5	
4^3D_3	4.26			8.38		5.3	
5^3D_3	4.82			10.1		5.1	

TABLE XIV. Photo-gluon decay widths of S states and quark-gluon decay widths of P states (in keV)

States	Γ	Γ_{cf}	Exp[67]	[58]	[83]	[59]	[56]
1^3S_1	1.37	0.79	1.19±0.33	1.32	0.79	0.903	0.7220
2^3S_1	0.72	0.42	0.60±0.10	0.739	0.46	0.341	0.4982
3^3S_1	0.54	0.31	0.20±0.04	0.547	0.36	0.276	0.3874
4^3S_1	0.45	0.26		0.433	0.30	0.253	0.3176
5^3S_1	0.39	0.22		0.370	0.25	0.242	0.2698
6^3S_1	0.34	0.19		0.311	0.22	0.235	0.2272
1^3P_1	32.25			81.7	71.53	45.55	57.9585
2^3P_1	43.28			117.0	106.14	56.16	55.3966
3^3P_1	48.46			126.0	124.53	68.97	52.9585
4^3P_1	51.27			128.0			52.4466
5^3P_1	52.82			132.0			49.5181

From the measured branching ratios of $B[h_b(2P) \rightarrow \gamma\eta_b(2S)] = 48 \pm 13\%$ and $B[h_b(2P) \rightarrow \gamma\eta_b(1S)] = 22 \pm 5\%$ [67], the total decay width of $h_b(2P)$ is evaluated to be 46.0 keV and 50.0 keV, respectively. The average comes out to be 48.0 keV, which is smaller compared to estimate of Ref. [74]. The transition width for $\Gamma(\chi_{b0}(1P) \rightarrow \gamma\Upsilon(2S))$ is overestimated across most of the models. From the measured branching ratios of $B[\chi_{b0}(2P) \rightarrow \gamma\Upsilon(2S)] = 1.38 \pm 0.30\%$ and $B[\chi_{b0}(2P) \rightarrow \gamma\Upsilon(1S)] = (3.8 \pm 1.7) \times 10^{-3}$ [67] the branching ratios and the total decay width for $\chi_{b0}(2P)$ is determined to be 0.88 MeV and 2.20 MeV, respectively. These values exhibit significant variance, but the latter value aligns with the prediction (~ 2.5 MeV) of Ref. [53]. The transition width for $\Gamma(\chi_{b1}(2P) \rightarrow \gamma\Upsilon(S))$ and $\Gamma(\chi_{b2}(2P) \rightarrow \gamma\Upsilon(S))$ predicted by our model, demonstrate exceptional agreement with experimental results. From the measured branching ratios of $B[\chi_{b1}(2P) \rightarrow \gamma\Upsilon(2S)] = 18.1 \pm 1.9\%$ and $B[\chi_{b1}(2P) \rightarrow \gamma\Upsilon(1S)] = 9.9 \pm 1.0\%$ [67], we determine the total decay width of $\chi_{b1}(2P)$ to be 87.0 keV and 91.0 keV, respectively, with average value of 89.0 keV. Our evaluated total decay width of $\chi_{b1}(2P)$ is consistent with the previously reported value of 96 ± 16 keV, determined by CLEO Collaboration [88]. Using the branching ratio of $B[\chi_{b2}(2P) \rightarrow \gamma\Upsilon(1S)] = 6.6 \pm 0.8\%$ [67], the total decay width of $\chi_{b2}(2P)$ in our model is calculated to be 146.0 ± 18.0 keV, in agreement with the value of 138 ± 19 keV

TABLE XV. $E1$ transition widths (in keV) and photon energies (in MeV) of S wave states

Initial State	Final State	Ours E_γ	Ours Γ_{E1}	Exp[67]	[58]	[53]	[83]	[74]
2^1S_0	1^1P_1	125.3	4.769		2.467	2.48	2.85	3.41
2^3S_1	1^3P_0	183.4	1.632	1.22 ± 0.11	0.907	0.91	1.09	1.09
	1^3P_1	156.9	3.092	2.21 ± 0.22	1.60	1.63	1.84	2.17
	1^3P_2	137.7	3.472	2.29 ± 0.22	1.86	1.88	2.08	2.62
3^1S_0	2^1P_1	102.7	6.596		2.88	2.96	2.60	4.25
	1^1P_1	489.9	0.226		1.12	1.3	0.0084	0.67
3^3S_1	2^3P_0	148.2	2.157	1.20 ± 0.12	1.06	1.03	1.21	1.21
	2^3P_1	127.2	4.132	2.56 ± 0.26	1.96	1.91	2.13	2.61
	2^3P_2	111.3	4.651	2.66 ± 0.27	2.37	2.30	2.56	3.16
	1^3P_0	540.6	0.057	0.055 ± 0.01	0.0099	0.01	0.15	0.097
	1^3P_1	515.1	0.115	0.018 ± 0.01	0.0363	0.05	0.16	0.0005
	1^3P_2	496.1	0.139	0.2 ± 0.03	0.359	0.45	0.0827	0.14
4^1S_0	3^1P_1	88.6	7.329		1.50	1.24		
	2^1P_1	392.5	0.718			0.732		
	1^1P_1	768.9	0.022		0.688			
4^3S_1	3^3P_0	127.9	2.384		0.587	0.48	0.61	
	3^3P_1	109.5	4.544		1.14	0.84	1.17	
	3^3P_2	95.4	5.057		1.16	0.82	1.45	
	2^3P_0	434.0	0.160		0.0137		0.17	
	2^3P_1	413.6	0.344		0.0138		0.18	
	2^3P_2	398.1	0.440		0.226		0.11	
	1^3P_0	815.7	0.007		5.12×10^{-4}		0.0588	
	1^3P_1	790.8	0.012		0.0507		0.0474	
	1^3P_2	772.4	0.013		0.219		0.012	

obtained by CLEO Collaboration [88]. There is currently no experimental data for the $\Gamma(3P \rightarrow \gamma S)$ and $\Gamma(3P \rightarrow \gamma D)$ transitions, although the detections of $\Gamma(\chi_{b1}(3P) \rightarrow \gamma \Upsilon(1S))$, $\Gamma(\chi_{b1}(3P) \rightarrow \gamma \Upsilon(2S))$, $\Gamma(\chi_{b1}(3P) \rightarrow \gamma \Upsilon(3S))$, and $\Gamma(\chi_{b2}(3P) \rightarrow \gamma \Upsilon(3S))$ have been reported. We estimate the transition width for these decays to be 4.754 keV, 5.876 keV, 12.895 keV, and 15.899 keV, respectively.

$E1$ transitions widths for D wave states are calculated for $\Gamma(1D \rightarrow \gamma P)$, $\Gamma(2D \rightarrow \gamma P)$ and $\Gamma(2D \rightarrow \gamma F)$. in Table XVIII. In $1D$ states of bottomonium only the transition $\Gamma(\Upsilon_2(1D) \rightarrow \gamma \chi_b(1P))$ has been reportedly observed. The transition widths for $\Gamma(\Upsilon_2(1D) \rightarrow \gamma \chi_{b2}(1P))$ and $\Gamma(\Upsilon_2(1D) \rightarrow \gamma \chi_{b1}(1P))$ in our model is calculated to be 6.194 keV and 22.890 keV, respectively, consistent with other theoretical models. As outlined in Ref. [53, 58, 83], the $\eta_b(1D)$ and $\Upsilon_3(1D)$ states are predicted to have their total decay width equivalent to transition widths $\Gamma(\eta_b(1D) \rightarrow \gamma h_b(1P))$ and $\Gamma(\Upsilon_3(1D) \rightarrow \gamma h_b(1P))$, respectively. We estimate these transition widths to be 28.719 keV and 26.518 keV, respectively, in accordance with other models. The $\Upsilon_1(1D)$ state is predicted to be detected in $\Gamma(\Upsilon_1(1D) \rightarrow \gamma \chi_{b0}(1P))$ and $\Gamma(\Upsilon_1(1D) \rightarrow \gamma \chi_{b1}(1P))$ transitions because of their large branching ratio [53, 58, 83]. We compute the transition widths of these decays to be 20.292 keV and 11.661 keV, respectively. In $2D$ states of bottomonium, $\Gamma(\eta_b(2D) \rightarrow \gamma h_b(2P))$ is predicted to have a large branching ratio, making it feasible to detect the $\eta_b(2D)$ unobserved state [53, 58]. In our model the transition width of $\Gamma(\eta_b(2D) \rightarrow \gamma h_b(2P))$ is estimated to be 21.128 keV. The transition widths pertaining to the $2D$ states of bottomonium in our model are found to align with the predictions of other theoretical models.

$E1$ transitions widths for F and G wave states are estimated for $\Gamma(1F \rightarrow \gamma D)$, $\Gamma(2F \rightarrow \gamma D)$, $\Gamma(2F \rightarrow \gamma G)$ and $\Gamma(1G \rightarrow \gamma F)$ in Table XIX. The transition widths for F and G wave states in our model are a bit higher compared to other potential models.

The $M1$ transition widths for S and P wave states are presented in Table XX and are compared with the available experimental data and other theoretical models. There are noticeable differences for $M1$ transition widths values in our model in contrast to the predictions in Ref. [53, 56, 58]. Our estimates of the $\Gamma(\Upsilon(nS) \rightarrow \gamma \eta_b(nS))$ transition widths are significantly lower compared to other models, yet they show a closer alignment with the experimental results. The decay widths values are highly dependent on the wavefunction, thereby resulting in considerable discrepancies among models.

The masses and leptonic decay widths of $S - D$ mixed states are presented in Table XXI for $3S - 2D$, $4S - 3D$, and $5S - 4D$ bottomonium states. The mixed states are also assigned to experimentally observed states. The $\Upsilon(10355)$ state is considered to be $3S - 2D$ mixed state with a small mixing component, having mass and leptonic decay width

TABLE XVI. $E1$ transition widths (in keV) and photon energies (in MeV) of $1P$ and $2P$ wave states

Initial State	Final State	Ours E_γ	Ours Γ_{E1}	Exp[67]	[58]	[53]	[83]	[74]
1^1P_1	1^1S_0	455.4	38.692	35.77	34.4	35.7	43.66	35.8
1^3P_0	1^3S_1	379.9	23.099		22.8	23.8	28.07	27.5
1^3P_1	1^3S_1	405.8	27.901	32.544	28.3	29.5	35.66	31.9
1^3P_2	1^3S_1	424.9	31.805	34.38	31.4	32.8	39.15	31.8
2^1P_1	1^1D_2	121.8	3.604		1.81	1.7	5.36	2.24
	2^1S_0	269.1	21.962	40.32	15.0	14.1	17.60	16.2
	1^1S_0	828.8	11.071		10.8	13.0	14.90	16.1
2^3P_0	1^3D_1	104.9	2.316		1.05	1.0	0.74	1.77
	2^3S_1	218.7	12.165	1.2×10^{-4}	11.1	10.9	12.80	14.4
	1^3S_1	763.0	8.198		2.31	2.5	5.44	5.54
2^3P_1	1^3D_1	126.0	0.995		0.511	0.5	0.41	0.56
	1^3D_2	117.6	2.436		1.25	1.2	1.26	0.50
	2^3S_1	239.6	15.790	19.4 ± 5	13.7	13.3	15.89	15.3
	1^3S_1	782.7	8.991	8.9 ± 2.2	5.09	5.5	9.13	10.8
2^3P_2	1^3D_1	141.9	0.056		0.0267	0.03	0.0209	0.026
	1^3D_2	133.5	0.708		0.339	0.3	0.35	0.42
	1^3D_3	127.3	3.442		1.61	1.5	2.06	2.51
	2^3S_1	255.2	18.908	15.1 ± 5.6	14.6	14.3	17.50	15.3
	1^3S_1	797.6	9.626	9.8 ± 2.3	7.86	8.4	11.38	12.5

TABLE XVII. $E1$ transition widths (in keV) and photon energies (in MeV) of $3P$ states

Initial State	Final State	Ours E_γ	Ours Γ_{E1}	[58]	[53]	[83]	[74]
3^1P_1	2^1D_2	105.9	5.482	1.44	1.6	4.72	4.21
	1^1D_2	424.7	0.208	0.0585	0.081	0.35	0.17
	3^1S_0	205.8	18.156	9.94	8.9	12.27	14.1
	2^1S_0	567.7	7.175	4.60	8.2	6.86	7.63
	1^1S_0	1110.9	5.592	3.91	3.6	7.96	10.7
3^3P_0	2^3D_1	91.8	3.593	0.966	1.0	3.50	2.20
	1^3D_1	411.7	0.163	0.189	0.20	3.59×10^{-2}	0.15
	3^3S_1	163.9	9.527	7.15	6.9	8.50	7.95
	2^3S_1	522.0	5.156	1.26	1.7	2.99	2.55
	1^3S_1	1050.1	4.462	0.427	0.3	1.99	1.87
3^3P_1	2^3D_1	110.2	1.541	0.425	0.47	1.26	1.07
	2^3D_2	102.5	3.738	0.950	1.1	3.34	0.94
	1^3D_1	429.6	0.056	0.00418	7.0×10^{-3}	4.80×10^{-2}	0.010
	1^3D_2	421.4	0.147	0.0615	0.080	0.11	0.015
	3^3S_1	182.3	12.897	8.36	8.4	9.62	10.3
	2^3S_1	539.8	5.876	2.49	3.1	4.58	5.63
	1^3S_1	1066.9	4.754	1.62	1.3	4.17	6.41
3^3P_2	2^3D_1	124.3	0.088	0.0248	0.027	0.18	0.049
	2^3D_2	116.6	1.090	0.295	0.32	0.79	0.78
	2^3D_3	110.6	5.229	1.37	1.5	4.16	4.60
	1^3D_1	443.2	0.003	1.15×10^{-4}		3.38×10^{-3}	0.047
	1^3D_2	435.1	0.037	3.11×10^{-4}		4.41×10^{-2}	0.068
	1^3D_3	429.0	0.188	0.0288	0.046	0.21	0.12
	3^3S_1	196.2	15.889	9.30	9.3	10.38	10.8
	2^3S_1	553.2	6.478	3.66	4.5	5.62	6.72
	1^3S_1	1079.7	4.986	3.17	2.8	5.65	8.17

TABLE XVIII. $E1$ transition widths (in keV) and photon energies (in MeV) of D states

Initial State	Final State	Ours E_γ	Ours Γ_{E1}	[58]	[53]	[83]	[74]
1^1D_2	1^1P_1	272.5	28.719	24.3	24.9	17.23	30.3
1^3D_1	1^3P_0	296.3	20.292	16.3	16.5	20.98	19.8
	1^3P_1	270.0	11.661	9.51	9.7	12.29	13.3
	1^3P_2	250.6	0.626	0.550	0.56	0.65	1.02
1^3D_2	1^3P_1	278.3	22.890	18.8	19.2	21.95	21.8
	1^3P_2	258.9	6.194	5.49	5.6	6.23	7.23
1^3D_3	1^3P_2	265.1	26.518	23.9	24.3	24.74	32.1
2^1D_2	1^1F_3	108.9	2.640	1.35	1.8	2.20	
	2^1P_1	202.6	21.128	16.8	16.5	11.66	15.6
	1^1P_1	586.0	6.313	3.36	3.0	4.15	5.66
2^3D_1	1^3F_2	103.1	2.246	1.18	1.6	2.05	
	2^3P_0	220.0	14.846	11.0	10.6	8.35	9.58
	2^3P_1	199.1	8.380	6.71	6.5	4.84	6.74
	2^3P_2	183.3	0.441	0.40	0.4	0.24	0.47
	1^3P_0	609.7	4.118	2.99	2.9	3.52	5.56
	1^3P_1	584.3	2.599	1.03	0.9	1.58	2.17
	1^3P_2	565.5	0.152	0.030	0.02	0.061	0.44
2^3D_2	1^3F_2	110.8	0.308	0.164	0.21	0.24	
	1^3F_3	107.6	2.265	1.21	1.5	1.93	
	2^3P_1	206.7	16.793	13.1	12.7	9.10	11.4
	2^3P_2	190.9	4.462	3.96	3.8	2.55	3.75
	1^3P_1	591.6	4.921	2.81	2.6	3.43	4.00
	1^3P_2	572.9	1.441	0.489	0.4	0.80	1.11
2^3D_3	1^3F_2	116.8	0.007	0.004	0.005	0.005	
	1^3F_3	113.6	0.237	0.125	0.16	0.19	
	1^3F_4	112.0	2.632	1.37	1.7		
	2^3P_2	196.9	19.488	16.8	16.4	10.70	17.0
	1^3P_2	578.6	5.997	2.99	2.6	3.80	5.22

values of 10374.9 MeV and 0.440 keV, respectively. The $\Upsilon(10580)$ state is considered to be $4S - 3D$ mixed state with substantial mixing component. Due to its narrow leptonic decay width, the partner of $\Upsilon(10580)$ with mass value of 10856.3 MeV is not regarded as $\Upsilon(10860)$. Instead, the $\Upsilon(10860)$ and $\Upsilon(11020)$ are assigned to be $5S - 4D$ mixed states, which is also supported by Ref. [61]. The mass and leptonic decay width values of the $\Upsilon(10860)$ and $\Upsilon(11020)$ are consistent with the experimental results. After taking into account the $S - D$ mixing, our final assignments are presented in Table XXII.

TABLE XIX. $E1$ transition widths (in keV) and photon energies (in MeV) of F and G states.

Initial State	Final State	Ours E_γ	Ours Γ_{E1}	[58]	[53]
1^1F_3	1^1D_2	215.3	27.505	22.0	18.8
1^3F_2	1^3D_1	221.8	25.149	19.4	16.4
	1^3D_2	213.4	4.171	3.26	2.7
	1^3D_3	207.3	0.109	0.0852	0.070
1^3F_3	1^3D_2	216.6	24.873	19.7	16.7
	1^3D_3	210.4	28.592	2.26	1.9
1^3F_4	1^3D_3	212.0	26.299	21.2	18.0
2^1F_3	1^1G_4	98.8	2.069	1.06	1.5
	2^1D_2	174.5	22.171	17.4	19.9
	1^1D_2	491.2	4.759	1.99	1.6
2^3F_2	1^3G_3	95.6	1.877	0.946	1.4
	2^3D_1	180.0	20.371	15.1	17.5
	2^3D_2	172.4	3.332	2.55	3.0
	2^3D_3	166.4	0.086	0.0681	0.080
	1^3D_1	497.2	4.214	1.95	1.6
	1^3D_2	489.1	0.727	0.224	0.16
	1^3D_3	483.1	0.019	0.00367	0.002
2^3F_3	1^3G_3	98.9	0.129	0.0664	0.10
	1^3G_4	98.2	1.910	0.957	1.4
	2^3D_2	175.6	20.091	15.4	17.9
	2^3D_3	169.7	2.275	1.80	2.1
	1^3D_2	492.3	4.273	1.83	1.4
	1^3D_3	486.3	0.506	0.145	0.1
2^3F_4	1^3G_3	100.8	0.002	8.80×10^{-4}	0.001
	1^3G_4	100.2	0.104	0.0535	0.080
	1^3G_5	100.9	2.098	1.05	1.5
	2^3D_3	171.6	21.148	16.9	19.6
	1^3D_3	488.2	4.635	0.126	1.4
1^1G_4	1^1F_3	184.5	27.129	21.1	23.1
1^3G_3	1^3F_2	187.5	26.098	20.1	22.3
	1^3F_3	184.3	2.174	1.67	1.8
	1^3F_4	182.7	0.034	0.0256	0.028
1^3G_4	1^3F_3	184.9	25.586	20.1	22.0
	1^3F_4	183.3	0.034	0.0256	0.028
1^3G_5	1^3F_4	182.6	26.304	21.1	23.1

TABLE XX. $M1$ transition widths (in keV) and photon energies (in MeV) of S and P states

Initial State	Final State	Ours E_γ	Ours Γ_{M1}	Exp[67]	[58]	[53]	[83]	
1^3S_1	1^1S_0	44.5	4.228		9.52	10.0	9.34	
2^1S_0	1^3S_1	532.8	4.848		70.6	68.0	45.0	
2^3S_1	2^1S_0	24.8	0.732		0.582	0.590	0.580	
	1^1S_0	598.3	3.569	12.5 ± 4.9	68.8	81.0	56.50	
3^1S_0	2^3S_1	345.1	2.015		11.1	9.10	9.20	
	1^3S_1	882.6	5.342		73.2	74.0	5.10	
3^3S_1	3^1S_0	19.3	0.343		0.337	0.250	0.658	
	2^1S_0	387.7	1.488	<13	11.8	19.0	11.0	
	1^1S_0	940.8	2.836	10 ± 2	60.4	60.0	57.0	
2^1P_1	1^3P_0	423.9	0.439		5.56	0.320	36.40	
	1^3P_1	398.0	0.857		1.30	1.10	1.280	
	1^3P_2	378.9	1.018		0.992	2.20	0.007	
2^3P_0	1^1P_1	365.2	0.475		5.21	9.70	2.390	
2^3P_1	1^1P_1	385.8	0.692		3.90×10^{-6}	2.20	0.167	
2^3P_2	1^1P_1	401.2	0.905		3.86	0.240	1.780	
3^1P_1	2^3P_0	332.5	0.366		2.16		1.710	
	2^3P_1	311.8	0.709		0.559		0.597	
	2^3P_2	269.2	0.831		0.407		0.007	
	1^3P_0	717.9	0.279		5.10	0.980	3.770	
	1^3P_1	692.8	0.639		1.01	0.930	1.230	
3^3P_0	1^3P_2	674.2	0.867		1.48	0.140	0.051	
	2^1P_1	283.4	0.372		2.05			
	1^1P_1	664.2	0.465		6.23			
	3^3P_1	2^1P_1	301.9	0.569		9.80×10^{-4}		
		1^1P_1	681.7	0.566		0.032		
3^3P_2	2^1P_1	315.7	0.772		1.74			
	1^1P_1	694.9	0.654		3.53			

TABLE XXI. $S - D$ mixed states with the masses of mixed states M_ϕ and $M_{\phi'}$ (in MeV) and their di-leptonic decay widths Γ_ϕ and $\Gamma_{\phi'}$ (in keV)

$S - D$ States	M_S M_D	θ	θ [61]	M_ϕ $M_{\phi'}$	M_{exp} [67]	Γ_ϕ $\Gamma_{\phi'}$	Γ_{exp} [67]
$3S$	10394.2	19.28	-9.0	10374.9	10355.1	0.440	0.443 ± 0.008
$2D$	10467.3			10486.5		0.036	
$4S$	10688.1	-30.72	-12.5	10573.2	10579.4	0.282	0.272 ± 0.029
$3D$	10741.3			10856.3		0.142	
$5S$	10938.9	44.55	-38.0	10909.8	10885.2	0.291	0.31 ± 0.07
$4D$	10981.0			11010.1	11000.0	0.142	0.13 ± 0.03

TABLE XXII. Our assignments for $b\bar{b}$ states with masses (in MeV) and di-leptonic decay widths (in keV)

States	Assignment	M_{exp} [67]	M_{cal}	Γ_{exp}^{ee} [67]	Γ_{cal}^{ee}
$\eta_b(1S)$	$\eta_b(1S)$	9398.7 ± 2	9406.4		
$\Upsilon(1S)$	$\Upsilon(1S)$	$9460.4 \pm 0.09 \pm 0.04$	9451.1	1.34 ± 0.018	1.268
$\chi_{b0}(1P)$	$\chi_{b0}(1P)$	$9859.44 \pm 0.42 \pm 0.31$	9838.7		
$\chi_{b1}(1P)$	$\chi_{b1}(1P)$	$9892.78 \pm 0.26 \pm 0.31$	9865.7		
$h_b(1P)$	$h_b(1P)$	9899.3 ± 0.8	9872.9		
$\chi_{b2}(1P)$	$\chi_{b2}(1P)$	$9912.21 \pm 0.26 \pm 0.31$	9885.6		
$\eta_b(2S)$	$\eta_b(2S)$	$9999.0 \pm 3.5^{+2.8}_{-1.9}$	9998.9		
$\Upsilon(2S)$	$\Upsilon(2S)$	10023.4 ± 0.5	10023.3	0.612 ± 0.011	0.666
$\Upsilon_2(1D)$	$\Upsilon_2(1D)$	10163.7 ± 1.4	10147.9		
$\chi_{b0}(2P)$	$\chi_{b0}(2P)$	$10232.5 \pm 0.4 \pm 0.5$	10244.9		
$\chi_{b1}(2P)$	$\chi_{b1}(2P)$	$10255.46 \pm 0.22 \pm 0.5$	10266.2		
$h_b(2P)$	$h_b(2P)$	$10259.8 \pm 0.5 \pm 1.1$	10271.7		
$\chi_{b2}(2P)$	$\chi_{b2}(2P)$	$10268.65 \pm 0.22 \pm 0.5$	10282.3		
$\Upsilon(10355)$	$\Upsilon(3S) - \Upsilon(2D)$	10355.1 ± 0.5	10374.9	0.443 ± 0.008	0.440
$\chi_{b1}(3P)$	$\chi_{b1}(3P)$	$10513.42 \pm 0.41 \pm 0.53$	10578.1		
$\chi_{b2}(3P)$	$\chi_{b2}(3P)$	$10524.02 \pm 0.57 \pm 0.53$	10592.3		
$\Upsilon(10580)$	$\Upsilon(4S) - \Upsilon(3D)$	10579.4 ± 1.2	10573.2	0.272 ± 0.029	0.282
$\Upsilon(10753)$	$\Upsilon_1(3D)$	$10752.7 \pm 5.9^{+0.7}_{-1.1}$	10741.3		0.003
$\Upsilon(10860)$	$\Upsilon(5S) - \Upsilon(4D)$	$10885.2 \pm 1.6^{+2.6}_{-1.6}$	10909.8	0.31 ± 0.07	0.291
$\Upsilon(11020)$	$\Upsilon(5S) - \Upsilon(4D)$	11000 ± 4	11010.1	0.13 ± 0.03	0.142

VIII. CONCLUSION

In this study, we explored the bottomonium system using a screened potential model within a relativistic framework to compute the mass spectrum of S, P, D, F and G wave, decay widths, and $E1$ and $M1$ transition widths, along with mass and leptonic decay widths of $S - D$ mixed states. Our calculated mass values demonstrate good agreement with experimental data and our $E1$ transition widths are utilized to estimate the total decay widths for higher bottomonium states. Our model provides better predictions for vector decay constants, tri-gluonic decays, radiative decays, and the total decay width of bottomonium states, demonstrating significant improvements in alignment with experimental data over previous potential models. A recurring challenge in potential models has been achieving theoretical predictions that align with experimental data for both the mass and leptonic decays of higher bottomonium states, such as $\Upsilon(10355)$, $\Upsilon(10580)$, $\Upsilon(10860)$ and $\Upsilon(11020)$. Our model addresses this issue by considering these states as $S - D$ mixed states resulting in predictions consistent with experimental observations. This research paves the way for future investigations, particularly in the exploration of higher excited states and the effects of $S - D$ mixing.

-
- [1] F.-K. Guo, H. Peng, J.-J. Xie, and X. Zhou, Hadron spectroscopy at STCF, arXiv:2203.07141 [hep-ph].
 - [2] D. M. Asner, H. Atmacan, S. Banerjee, J. V. Bennett, M. Bertemes, *et al.*, Belle II Executive Summary, arXiv:2203.10203v2 [hep-ex].
 - [3] N. Brambilla, S. Eidelman, C. Hanhart, A. Nefediev, C.-P. Shen, *et al.*, The XYZ states: Experimental and theoretical status and perspectives, Phys. Rep. **873**, 1 (2020).
 - [4] S. W. Herb, D. C. Hom, L. M. Lederman, J. C. Sens, H. D. Snyder, *et al.*, Observation of a Dimuon Resonance at 9.5 GeV in 400-GeV Proton-Nucleus Collisions, Phys. Rev. Lett. **39**, 252 (1977).
 - [5] W. R. Innes, J. A. Appel, B. C. Brown, C. N. Brown, K. Ueno, *et al.*, Observation of Structure in the Υ Region, Phys. Rev. Lett. **39**, 1240 (1977).
 - [6] K. Han, T. Böhringer, P. Franzini, G. Mageras, D. Peterson, *et al.*, Observation of P -Wave $b\bar{b}$ Bound States, Phys. Rev. Lett. **49**, 1612 (1982).
 - [7] G. Eigen, G. Blunar, H. Dietl, E. Lorenz, F. Pauss, *et al.*, Evidence for χ'_b Production in the Exclusive Reaction $\Upsilon'' \rightarrow \gamma\chi'_b \rightarrow (\gamma\gamma\Upsilon' \text{ or } \gamma\gamma\Upsilon)$, Phys. Rev. Lett. **49**, 1616 (1982).
 - [8] C. Klopfenstein, J. E. Horstkotte, J. Lee-Franzini, R. D. Schamberger, M. Sivertz, *et al.*, Observation of the Lowest P -Wave $b\bar{b}$ Bound States, Phys. Rev. Lett. **51**, 160 (1983).
 - [9] F. Pauss, H. Dietl, G. Eigen, E. Lorenz, G. Mageras, *et al.*, Observation of χ_b production in the exclusive reaction $\Upsilon' \rightarrow \gamma\gamma\chi_b \rightarrow \gamma\gamma\Upsilon \rightarrow \gamma\gamma(e^+e^- \text{ or } \mu^+\mu^-)$, Phys. Lett. B **130**, 439 (1983).
 - [10] CLEO Collaboration, M. Artuso, C. Boulahouache, S. Blusk, J. Butt, *et al.*, Photon Transitions in $\Upsilon(2S)$ and $\Upsilon(3S)$ Decays, Phys. Rev. Lett. **94**, 032001 (2005).
 - [11] D. Andrews, K. Berkelman, R. Cabenda, D. G. Cassel, J. W. DeWire, *et al.*, Observation of a Fourth Upsilon State in e^+e^- Annihilations, Phys. Rev. Lett. **45**, 219 (1980).
 - [12] D. Besson, J. Green, R. Namjoshi, F. Sannes, P. Skubic, *et al.*, Observation of New Structure in the e^+e^- Cross Section above the $\Upsilon(4S)$, Phys. Rev. Lett. **54**, 381 (1985).
 - [13] R. Mizuk, A. Bondar, I. Adachi, H. Aihara, D. M. Asner, *et al.*, Observation of a new structure near 10.75 GeV in the energy dependence of the $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) cross sections, J. High Energy Phys. **2019** (10), 1.
 - [14] G. Bonvicini, D. Cinabro, M. Dubrovin, A. Bornheim, E. Lipeles, *et al.*, First observation of a $\Upsilon(1D)$ state, Phys. Rev. D **70**, 032001 (2004).
 - [15] BABAR Collaboration, P. del Amo Sanchez, J. P. Lees, V. Poireau, E. Prencipe, *et al.*, Observation of the $\Upsilon(1^3D_J)$ bottomonium state through decays to $\pi^+\pi^-\Upsilon(1S)$, Phys. Rev. D **82**, 111102 (2010).
 - [16] BABAR Collaboration, B. Aubert, M. Bona, Y. Karyotakis, J. P. Lees, *et al.*, Observation of the Bottomonium Ground State in the Decay $\Upsilon(3S) \rightarrow \gamma\eta_b$, Phys. Rev. Lett. **101**, 071801 (2008).
 - [17] S. Dobbs, Z. Metreveli, A. Tomaradze, T. Xiao, and K. K. Seth, Observation of the $\eta_b(2S)$ Meson in $\Upsilon(2S) \rightarrow \gamma\eta_b(2S)$, $\eta_b(2S) \rightarrow$ Hadrons and Confirmation of the $\eta_b(1S)$ Meson, Phys. Rev. Lett. **109**, 082001 (2012).
 - [18] S. Dobbs and The CLEO Collaboration, A New Measurement of $\eta_b(1S)$ From $\Upsilon(3S)$ Radiative Decay at CLEO, AIP Conf. Proc. **1257**, 408 (2010).
 - [19] Belle Collaboration, R. Mizuk, D. M. Asner, A. Bondar, T. K. Pedlar, *et al.*, Evidence for the $\eta_b(2S)$ and Observation of $h_b(1P) \rightarrow \eta_b(1S)\gamma$ and $h_b(2P) \rightarrow \eta_b(1S)\gamma$, Phys. Rev. Lett. **109**, 232002 (2012).
 - [20] The BABAR Collaboration, J. P. Lees, V. Poireau, E. Prencipe, V. Tisserand, *et al.*, Evidence for the $h_b(1P)$ meson in the decay $\Upsilon(3S) \rightarrow \pi^0 h_b(1P)$, Phys. Rev. D **84**, 091101 (2011).
 - [21] Belle Collaboration, I. Adachi, H. Aihara, K. Arinstein, D. M. Asner, *et al.*, First Observation of the P -Wave Spin-Singlet Bottomonium States $h_b(1P)$ and $h_b(2P)$, Phys. Rev. Lett. **108**, 032001 (2012).
 - [22] ATLAS Collaboration, G. Aad, B. Abbott, J. Abdallah, A. A. Abdelalim, *et al.*, Observation of a New χ_b State in Radiative Transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS, Phys. Rev. Lett. **108**, 152001 (2012).

- [23] D0 Collaboration, V. M. Abazov, B. Abbott, B. S. Acharya, M. Adams, *et al.*, Observation of a narrow mass state decaying into $\Upsilon(1S) + \gamma$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV, *Phys. Rev. D* **86**, 031103 (2012).
- [24] CMS Collaboration, A. M. Sirunyan, A. Tumasyan, W. Adam, F. Ambrogio, *et al.*, Observation of the $\chi_{b1}(3P)$ and $\chi_{b2}(3P)$ and Measurement of their Masses, *Phys. Rev. Lett.* **121**, 092002 (2018).
- [25] Belle Collaboration, A. Bondar, A. Garmash, R. Mizuk, D. Santel, *et al.*, Observation of Two Charged Bottomoniumlike Resonances in $\Upsilon(5S)$ Decays, *Phys. Rev. Lett.* **108**, 122001 (2012).
- [26] F.-K. Guo, C. Hanhart, U.-G. Meißner, Q. Wang, Q. Zhao, *et al.*, Hadronic molecules, *Rev. Mod. Phys.* **90**, 015004 (2018).
- [27] Q. Wang, V. Baru, A. A. Filin, C. Hanhart, A. V. Nefediev, *et al.*, Line shapes of the $Z_b(10610)$ and $Z_b(10650)$ in the elastic and inelastic channels revisited, *Phys. Rev. D* **98**, 074023 (2018).
- [28] E. S. Swanson, Z_b and Z_c exotic states as coupled channel cusps, *Phys. Rev. D* **91**, 034009 (2015).
- [29] L. Maiani, A. D. Polosa, and V. Riquer, A theory of X and Z multiquark resonances, *Phys. Lett. B* **778**, 247 (2018).
- [30] R. Mizuk, A. Bondar, I. Adachi, H. Aihara, D. M. Asner, *et al.*, Observation of a new structure near 10.75 GeV in the energy dependence of the $e^+e^- \rightarrow \Upsilon(nS)\pi^+\pi^-$ ($n = 1, 2, 3$) cross sections, *J. High Energy Phys.* **2019** (220), 1.
- [31] X.-K. Dong, X.-H. Mo, P. Wang, and C.-Z. Yuan, Hadronic cross section of e^+e^- annihilation at bottomonium energy region, *Chin. Phys. C* **44**, 083001 (2020).
- [32] W.-S. Hou, Searching for the bottom counterparts of $X(3872)$ and $Y(4260)$ via $\pi^+\pi^-\Upsilon$, *Phys. Rev. D* **74**, 017504 (2006).
- [33] A. Ali, C. Hambrook, I. Ahmed, and M. J. Aslam, A case for hidden $b\bar{b}$ tetraquarks based on $e^+e^- \rightarrow b\bar{b}$ cross section between $\sqrt{s} = 10.54$ and 11.20 GeV, *Phys. Lett. B* **684**, 28 (2010).
- [34] F.-K. Guo, C. Hidalgo-Duque, J. Nieves, and M. Pavón Valderrama, Consequences of heavy-quark symmetries for hadronic molecules, *Phys. Rev. D* **88**, 054007 (2013).
- [35] G. Mezzadri and S. Spataro, XYZ states: An experimental point-of-view, *Rev. Phys.* **8**, 100070 (2022).
- [36] C. A. Bokade and Bhaghyesh, Charmonium: Conventional and XYZ States in a Relativistic Screened Potential Model, arXiv:2408.06759 [hep-ph].
- [37] S. Godfrey and N. Isgur, Mesons in a relativized quark model with chromodynamics, *Phys. Rev. D* **32**, 189 (1985).
- [38] D. Yibing, C. Kuangta, and Q. Danhua, Screened $Q - \bar{Q}$ Potential and Spectrum of Heavy Quarkonium, *Chin. Phys. Lett.* **10**, 460 (1993).
- [39] J.-B. Liu and M.-Z. Yang, Spectrum of the charmed and b-flavored mesons in the relativistic potential model, *J. High Energy Phys.* **2014** (7), 1.
- [40] M.-Z. Yang, Wave functions and decay constants of B and D mesons in the relativistic potential model, *Eur. Phys. J. C* **72**, 1 (2012).
- [41] E. Eichten and F. Feinberg, Spin-dependent forces in quantum chromodynamics, *Phys. Rev. D* **23**, 2724 (1981).
- [42] M. B. Voloshin, Charmonium, *Prog. Part. Nucl. Phys.* **61**, 455 (2008).
- [43] T. Barnes, S. Godfrey, and E. S. Swanson, Higher charmonia, *Phys. Rev. D* **72**, 054026 (2005).
- [44] R. F. Lebed and E. S. Swanson, Quarkonium h states as arbiters of exoticity, *Phys. Rev. D* **96**, 056015 (2017).
- [45] B.-Q. Li and K.-T. Chao, Higher charmonia and X, Y, Z states with screened potential, *Phys. Rev. D* **79**, 094004 (2009).
- [46] R. Van Royen and V. F. Weisskopf, Hadron decay processes and the quark model, *Nuovo Cim. A* **50**, 617 (1967).
- [47] E. Braaten and S. Fleming, QCD radiative corrections to the leptonic decay rate of the B_c meson, *Phys. Rev. D* **52**, 181 (1995).
- [48] J. P. Lansberg and T. N. Pham, Effective Lagrangian for two-photon and two-gluon decays of P -wave heavy quarkonium $\chi_{c0,2}$ and $\chi_{b0,2}$ states, *Phys. Rev. D* **79**, 094016 (2009).
- [49] N. Akbar, Decay Properties of Conventional and Hybrid Charmonium Mesons, *J. Korean Phys. Soc.* **77**, 17 (2020).
- [50] W. Kwong and J. L. Rosner, D-wave quarkonium levels of the Υ family, *Phys. Rev. D* **38**, 279 (1988).
- [51] W. Kwong, P. B. Mackenzie, R. Rosenfeld, and J. L. Rosner, Quarkonium annihilation rates, *Phys. Rev. D* **37**, 3210 (1988).
- [52] V. A. Novikov, L. B. Okun, M. A. Shifman, A. I. Vainshtein, M. B. Voloshin, *et al.*, Charmonium and gluons, *Phys. Rep.* **41**, 1 (1978).
- [53] S. Godfrey and K. Moats, Bottomonium mesons and strategies for their observation, *Phys. Rev. D* **92**, 054034 (2015).
- [54] G. Bélanger and P. Moxhay, Three-gluon annihilation of D-wave quarkonium, *Phys. Lett. B* **199**, 575 (1987).
- [55] W. A. Bardeen, E. J. Eichten, and C. T. Hill, Chiral multiplets of heavy-light mesons, *Phys. Rev. D* **68**, 054024 (2003).
- [56] B. Pandya, M. Shah, and P. C. Vinodkumar, Importance of confinement in instanton induced potential for bottomonium spectroscopy, *Eur. Phys. J. C* **81**, 1 (2021).
- [57] E. Eichten, K. Gottfried, T. Kinoshita, K. D. Lane, and T.-M. Yan, Charmonium: The model, *Phys. Rev. D* **17**, 3090 (1978).
- [58] J.-Z. Wang, Z.-F. Sun, X. Liu, and T. Matsuki, Higher bottomonium zoo, *Eur. Phys. J. C* **78**, 1 (2018).
- [59] V. Kher, R. Chaturvedi, N. Devlani, and A. K. Rai, Bottomonium spectroscopy using Coulomb plus linear (Cornell) potential, *Eur. Phys. J. Plus* **137**, 1 (2022).
- [60] S. F. Radford and W. W. Repko, Potential model calculations and predictions for heavy quarkonium, *Phys. Rev. D* **75**, 074031 (2007).
- [61] Z. Zhao, K. Xu, A. Limphirat, W. Sreethawong, N. Tagsinsit, *et al.*, Mass spectrum of 1^{--} heavy quarkonium, *Phys. Rev. D* **109**, 016012 (2024).
- [62] K. Heikkilä, N. A. Törnqvist, and S. Ono, Heavy $c\bar{c}$ and $b\bar{b}$ quarkonium states and unitarity effects, *Phys. Rev. D* **29**, 110 (1984).
- [63] Y. Lu, M. N. Anwar, and B.-S. Zou, Coupled-channel effects for the bottomonium with realistic wave functions, *Phys. Rev. D* **94**, 034021 (2016).

- [64] R. Machleidt, K. Holinde, and Ch. Elster, The bonn meson-exchange model for the nucleon-nucleon interaction, *Phys. Rep.* **149**, 1 (1987).
- [65] L. Ya. Glozman and D. O. Riska, The spectrum of the nucleons and the strange hyperons and chiral dynamics, *Phys. Rep.* **268**, 263 (1996).
- [66] J.-Z. Wang and X. Liu, Identifying a characterized energy level structure of higher charmonium well matched to the peak structures in $e^+e^- \rightarrow \pi^+D^0D^{*-}$, *Phys. Lett. B* **849**, 138456 (2024).
- [67] Particle Data Group, R. L. Workman, V. D. Burkert, V. Crede, E. Klempt, *et al.*, Review of Particle Physics, *Prog. Theor. Exp. Phys.* **2022**, 083C01 (2022).
- [68] W.-H. Liang, N. Ikeno, and E. Oset, $\Upsilon(nI)$ decay into $B^{(*)}\bar{B}^{(*)}$, *Phys. Lett. B* **803**, 135340 (2020).
- [69] E. van Beveren and G. Rupp, Production of hadron pairs in e^+e^- annihilation near the K^+K^- , $D\bar{D}$, $B\bar{B}$, and $\Lambda_c^+\Lambda_c^-$ thresholds, *Phys. Rev. D* **80**, 074001 (2009).
- [70] D. Besson, J. Green, R. Namjoshi, F. Sannes, P. Skubic, *et al.*, Observation of New Structure in the e^+e^- Cross Section above the $\Upsilon(4S)$, *Phys. Rev. Lett.* **54**, 381 (1985).
- [71] R. Bruschini and P. González, A plausible explanation of $\Upsilon(10860)$, *Phys. Lett. B* **791**, 409 (2019).
- [72] P. Bicudo, N. Cardoso, L. Mueller, and M. Wagner, Study of $I = 0$ bottomonium bound states and resonances in S , P , D , and F waves with lattice QCD static-static-light-light potentials, *Phys. Rev. D* **107**, 094515 (2023).
- [73] Q. Li, M.-S. Liu, Q.-F. Lü, L.-C. Gui, and X.-H. Zhong, Canonical interpretation of $Y(10750)$ and $\Upsilon(10860)$ in the Υ family, *Eur. Phys. J. C* **80**, 1 (2020).
- [74] W.-J. Deng, H. Liu, L.-C. Gui, and X.-H. Zhong, Spectrum and electromagnetic transitions of bottomonium, *Phys. Rev. D* **95**, 074002 (2017).
- [75] M. Karliner and J. L. Rosner, $X(3872)$, X_b , and the $\chi_{b1}(3P)$ state, *Phys. Rev. D* **91**, 014014 (2015).
- [76] B. Chen, A. Zhang, and J. He, Bottomonium spectrum in the relativistic flux tube model, *Phys. Rev. D* **101**, 014020 (2020).
- [77] A. Ali, L. Maiani, A. Y. Parkhomenko, and W. Wang, Interpretation of $Y_b(10750)$ as a tetraquark and its production mechanism, *Phys. Lett. B* **802**, 135217 (2020).
- [78] Z.-G. Wang, Vector hidden-bottom tetraquark candidate: $Y(10750)$, *Chin. Phys. C* **43**, 123102 (2019).
- [79] E. van Beveren and G. Rupp, Observation of the Upsilon(2 3D1) and indication of the Upsilon(1 3D1), arXiv 10.48550/arXiv.1009.4097 (2010), 1009.4097.
- [80] N. R. Soni, B. R. Joshi, R. P. Shah, H. R. Chauhan, and J. N. Pandya, $Q\bar{Q}$ ($Q \in \{b, c\}$) spectroscopy using Cornell potential, *Eur. Phys. J. C* **78**, 1 (2018).
- [81] B. Patel and P. C. Vinodkumar, Properties of $Q\bar{Q}$ ($Q \in \{b, c\}$) mesons in Coulomb plus Power potential, *J. Phys. G: Nucl. Part. Phys.* **36**, 035003 (2009).
- [82] A. M. Badalian, B. L. G. Bakker, and I. V. Danilkin, Possibility to observe higher n^3D_1 bottomonium states in the e^+e^- processes, *Phys. Rev. D* **79**, 037505 (2009).
- [83] J. Segovia, P. G. Ortega, D. R. Entem, and F. Fernández, Bottomonium spectrum revisited, *Phys. Rev. D* **93**, 074027 (2016).
- [84] A. K. Grant and J. L. Rosner, Dipole transition matrix elements for systems with power-law potentials, *Phys. Rev. D* **46**, 3862 (1992).
- [85] E. Eichten, S. Godfrey, H. Mahlke, and J. L. Rosner, Quarkonia and their transitions, *Rev. Mod. Phys.* **80**, 1161 (2008).
- [86] A. M. Badalian and B. L. G. Bakker, Dominant spin-orbit effects in radiative decays $\Upsilon(3S) \rightarrow \gamma + \chi_{bJ}(1P)$, *Phys. Rev. D* **86**, 074001 (2012).
- [87] A. Abdesselam, I. Adachi, K. Adamczyk, H. Aihara, S. A. Said, *et al.*, Study of $\chi_{\{bJ\}}(1P)$ Properties in the Radiative $\Upsilon(2S)$ Decays, arXiv:1606.01276 [hep-ex].
- [88] C. Cawfield, B. I. Eisenstein, I. Karliner, D. Kim, N. Lowrey, *et al.*, Experimental study of $\chi_b(2P) \rightarrow \pi\pi\chi_b(1P)$, *Phys. Rev. D* **73**, 012003 (2006).