Velocity Addition/Subtraction in Special Relativity

Domenico Giulini Institute for Theoretical Physics Leibniz University of Hannover, Germany giulini@itp.uni-hannover.de

January 7, 2025

Abstract

We reconsider velocity addition/subtraction in Special Relativity (SR) and rederive its well-known non-commutative and non-associative algebraic properties in a self contained way, including various explicit expressions for the Thomas angle, the derivation of which will be seen to be not as challenging as often suggested. All this is based on the polar-decomposition theorem in the traditional component language, in which Lorentz transformations are ordinary matrices. In the second part of this paper we offer a less familiar alternative geometric view, that leads to an invariant definition of the concept of relative velocity between two states of motion, which is based on the boost-link-theorem, of which we also offer an elementary proof that does not seem to be widely known in the relativity literature. Finally we compare this to the corresponding geometric definitions in Galilei-Newton spacetime, emphasising similarities and differences. Regarding the presentation of the material we will pursue an uncompromising pedagogical strategy, willingly accepting repetitions and occasional redundancies if deemed beneficial for clarity and the avoidance of anticipated misunderstandings. An appendix with four sections includes some mathematical details on results needed in the main text, as well some recollections on notions like semi-direct products of groups and affine spaces.

Keywords: Special Relativity, velocity addition, relative velocity, Thomas rotation, boost-link problem

Mathematics Subject Classification (MSC): 83A05, 20N05, 57K32 Physics and Astronomy Classification Scheme(PACS): 03.30.+p

Contents

1	Introduction		3
2	Velocity addition/subtraction: the traditional story		4
	2.1	A reminder on the elementary text-book derivation	4
	2.2	Lorentz transformations in the traditional matrix representation	6
	2.3	The generalised orthogonal group and its Lie algebra	7
	2.4	Restriction to four dimensions and Lorentzian signature	10
	2.5	Polar decomposition of boost products	13
	2.6	On the magnitude of Thomas rotation	17
	2.7	On the algebraic structure of Einstein addition	18
3	3 Velocity subtraction: the modern (geometric) story		24
	3.1	States of motion and the non-naturalness of polar decoposition \ldots .	24
	3.2	The kinematical setting	26
	3.3	Defining link velocity	31
	3.4	Solving the boost-link problem	33
	3.5	Base-point dependence of the link-velocity	36
4	Comparison with Galilei-Newton spacetimes		39
	4.1	Boost-Rotation decomposition in SR rephrased	39
	4.2	Galilei-Newton spacetime	40
	4.3	Automorphims of Galilei-Newton spacetime	42
	4.4	Boost-Rotation decomposition for the Galilei group	45
A	Polar decomposition		46
B	B Parallel transport along geodesics on state space		49
С	C Semi-direct products of groups		
D	Affine structures		54
	D.1	Affine spaces	55
	D.2	Affine maps and groups	57
	D.3	Affine bases and charts	59

1 Introduction

This paper attempts to give a comprehensive account of the algebraic and geometric aspects connected with the composition and linking properties of boost transformations in Special Relativity (henceforth abbreviated by SR). It contains many known results¹ but also offers new aspects. Some of the proofs of the known results appear to be new and - hopefully - easier than those in the published literature. The style of our presentation is largely shaped by an educational intent and clearly reveals that the following remarks originate from a lecture manuscript.²

In classical mechanics we are used to the fact that the relative velocity between two states of motion is independent of the third state of motion (of the "observer") from which it is reckoned. In Special Relativity this ceases to be true in a twofold way: regarding its measure and regarding the space of which it is an element of. Nevertheless, all assignments are naturally fully covariant and hence consistent with the principle of relativity.

All of these aspects will be clarified in this paper, whose main result is stated and fully proved in Section 3. It may be summarised as follows: Let S denote the set of *states of motion*, which is a maximally symmetric 3-dimensional Riemannian manifold of constant negative curvature (given by the set of unit timelike future pointing vectors). For any ordered pair (s_1, s_2) of elements in S the *relative velocity* between them is a section $s \mapsto \beta(s, s_1, s_2) \in TS$ in the tangent bundle over S. The expression $\beta(s, s_1, s_2)$ is a rational function of its arguments (formula (130a)) which is equivariant under the action of the Lorentz group, in the sense that for any Lorentz transformation L it obeys $\beta(Ls, Ls_1, Ls_2) = L\beta(s, s_1, s_2)$. Its geometric interpretation is that of the velocity β (in units of c) of the unique boost $B(\beta)$ relative to s that links s_1 with s_2 . The uniqueness statement here is known as *boost-link-theorem*, of which we give an elementary and constructive proof. The analytic expression of the corresponding boost map (formula (134)) is likewise a rational function of (s, s_1, s_2) .

The organisation of the paper is as follows. Section 2 presents in a self-contained fashion the known story on Einstein's law of velocity addition/subtraction in component form. Polar decomposition is used to decompose elements in the Lorentz group into a boost and a rotation. It is emphasised that this decomposition is not natural but rather dependent of a distinguished state of motion. Explicit formulae for the polar decomposition of the composition of two boosts are given to derive once more Einstein addition and also Thomas Rotation, the properties of which are derived. In particular, uniqueness of polar decomposition is used to prove that Einstein addition endows the open unit ball in \mathbb{R}^3 with the algebraic structure of a non-associative quasigroup with identity, called a *loop*.

¹ See Gourgoulhon (2013) for a comprehensive modern account.

² Lectures on Special Relativity, delivered at Leibniz University Hannover at irregular intervals since 2009.

Section 3 replaces the common matrix notation by a proper geometric setting in which the analytic expressions merely contain states of motion and their scalar products. The boost-link-theorem is proved and taken as basis for the definition of "relative velocity", namely as the velocity of the boost that links the two given states (s_1, s_2) . We will call it the "link velocity". Next to the two states it also depends on the state s relative to which we identify the linking Lorentz transformation as a pure boost. Since the group generated by pure boosts is the entire Lorentz group, that reference to s is necessary. We emphasise that this latter reference renders the notion of relative velocity a ternary (rather than binary) relation, and that this fact is *not* in conflict with the relativity principle as sometimes claimed.

Section 4 compares the findings of Section 3 to the case of Galilei-Newton spacetime, which we again characterise in proper geometric terms. Here boosts form indeed a 3-dimensional subgroup, which is even abelian and normal, such that the relative velocity between two states (s_1, s_2) of motion—now defined in terms of the geometric structures given here—does *not* require the specification of a third reference state relative to which the linking Galilei transformation is identified as pure boost.

Finally, an Appendix of four parts contains the statement and proof of the polardecomposition theorem, a proof of the statement (used in the main text) that paralleltransport along geodesics on the space S of states is equivalent to boost transformations, and two summaries of elementary mathematical concepts used in the text, namely semi-direct products (for groups) and affine structures (on sets).

Notation and conventions: Throughout spacetime is a 4-dimensional real affine space with associated vector space V and dual space V^{*}. By *the Minkowski metric*, denoted by η , we understand a non-degenerate symmetric bilinear form on V of signature (-, +, +, +).

Acknowledgements: I thank Philip Schwartz and Lea Zybur for many conversations that helped me develop and refine some of the ideas presented here.

2 Velocity addition/subtraction: the traditional story

In this section we recall the standard, matrix-representation-based derivation of velocity addition/subtraction, including an analysis of the less widely known algebraic structure it defines. We shall give complete proofs of all relevant statements. Some of these proofs appear shorter and yet easier to follow than those in the published literature.

2.1 A reminder on the elementary text-book derivation

A frequently-seen derivation of the velocity addition in SR ist this: A particle P moves along a spatial trajectory $\mathbf{x}'(t')$ relative to an inertial system I' with whose

coordinates are labelled by (t', \mathbf{x}') . Relative to another inertial system, I, whose coordinates are labelled by (t, \mathbf{x}) , the system I' moves with constant velocity **v**. According to SR, the boost transformation between I and I' is

$$\mathbf{t} = \gamma(\mathbf{t}' + \mathbf{v} \cdot \mathbf{x}'/\mathbf{c}^2), \qquad (1a)$$

$$\mathbf{x} = \mathbf{x}' + (\gamma - 1)\mathbf{n}(\mathbf{n} \cdot \mathbf{x}') + \gamma \mathbf{v} \mathbf{t}', \tag{1b}$$

where

$$\gamma := (1 - \|\mathbf{v}\|^2 / c^2)^{-1/2}, \quad \mathbf{n} := \mathbf{v} / \|\mathbf{v}\|.$$
 (2)

The particle's P trajectory relative to I is then obtained by inserting $\mathbf{x}'(t')$ for \mathbf{x}' into these equations. Both expressions then become a functions of t'. Taking the differential, setting $d\mathbf{x}'/dt' := \mathbf{u}'$ and $d\mathbf{x}/dt := \mathbf{u}$, we get:

$$dt = \gamma (1 + \mathbf{v} \cdot \mathbf{u}') dt', \qquad (3a)$$

$$\mathbf{dx} = \left[\mathbf{u}' + (\gamma - 1)\mathbf{n}(\mathbf{n} \cdot \mathbf{u}') + \gamma \mathbf{v}\right] \mathbf{dt}'. \tag{3b}$$

Hence, the particle's velocity relative to I is

$$\mathbf{u} = \frac{\mathbf{v} + \mathbf{u}_{\parallel}' + \gamma^{-1} \mathbf{u}_{\perp}'}{1 + \mathbf{v} \cdot \mathbf{u}'/c^2}, \qquad (4)$$

where

$$\mathbf{u}_{\parallel}' := \mathbf{n}(\mathbf{n} \cdot \mathbf{u}') \quad \text{and} \quad \mathbf{u}_{\perp}' := \mathbf{u}' - \mathbf{u}_{\parallel}'$$
 (5)

are the orthogonal projections of \mathbf{u}' parallel and perpendicular to \mathbf{v} , respectively. Alternative ways to write (4) are

$$\mathbf{u} = \frac{1}{1 + \frac{\mathbf{v} \cdot \mathbf{u}'}{c^2}} \left\{ \mathbf{v} + \mathbf{u}' + \frac{\gamma}{c^2(\gamma + 1)} \, \mathbf{v} \times (\mathbf{v} \times \mathbf{u}') \right\}$$
(6a)

$$= \frac{1}{1 + \frac{\mathbf{v} \cdot \mathbf{u}'}{c^2}} \left\{ \mathbf{v} + \boldsymbol{\gamma}^{-1} \boldsymbol{u}' + \frac{\boldsymbol{\gamma}}{c^2(\boldsymbol{\gamma} + 1)} \, \mathbf{v} \left(\mathbf{v} \cdot \mathbf{u}' \right) \right\} \,. \tag{6b}$$

The first term takes a simple form if \mathbf{v} is parallel to \mathbf{u}' (in which case the term involving the ×-products vanishes) and the second an even simpler form if \mathbf{v} is perpendicular to \mathbf{u}' (in which case the term in the numerator as well as that in the denominator involving the inner product $\mathbf{v} \cdot \mathbf{u}'$ vanishes). These formulae show that \mathbf{u} is a rational function of \mathbf{v} and \mathbf{u}' which is C^{∞} as long as $\mathbf{v} \cdot \mathbf{u}' > -c^2$, which is the case as long as $\|\mathbf{v}\| \le c$ and $\|\mathbf{u}'\| \le c$ without simultaneous equality. In that case one has $\|\mathbf{u}'\| \le c$ with equality if and only if either $\|\mathbf{v}\| = c$ or $\|\mathbf{u}'\| = c$. This immediately follows from the equation

$$\gamma(\mathbf{u}) = \gamma(\mathbf{v})\gamma(\mathbf{u}')\left(1 + \mathbf{v} \cdot \mathbf{u}'/c^2\right)$$
(7)

where the γ -factors are now defined for any of the three velocities involved; i.e., $\gamma(\mathbf{u}) := (1 - \|\mathbf{u}\|^2 / c^2)^{-1/2}$, etc. Equation (7) follows easiest from taking the square of (4). We shall encounter another proof below.

Formulae (4), (6), and (7) summarise Einstein's law of "velocity addition" as usually discussed. The interesting algebraic structure behind it will be the subject of the next section. It has been analysed in detail before by others (Ungar, 1988, 1989, 1997, 2005; Urbantke, 2003), but our treatment here will be different, self contained, and direct. The third and main part of this paper addresses and resolves the following concern: How can the addition of different relative velocities possibly make sense in view of the fact that the vector space of relative velocities depends on the inertial frame it refers to? In our example, \mathbf{u}' seems to be defined relative to I', whereas \mathbf{v} and \mathbf{u} are relative to I. However, there is no obvious way to add vectors in the space of velocities relative to I' to vectors in the (different) space of velocities relative to I; and yet, this is precisely what seems to have been done above. How can that be a meaningful operation? The answer we shall give is that, albeit \mathbf{u}' does represent the velocity of I'' against I', it does so with reference to I rather than I'. How this additional reference to I, which renders the notion of "relative velocity" a ternary rather than binary relation, is to be understood properly will be explained in detail. Related observations had been made before in (Matolcsi and Goher, 2001; Matolcsi and Matolcsi, 2005; Urbantke, 2003) but not in the manifestly covariant form that we shall present here.

2.2 Lorentz transformations in the traditional matrix representation

We begin by slightly reformulating the previous formulae in a way that is easier to memorise. First, we take all relative velocities in units of c and call them (as is usual in the SR-literature) by β . Setting $\beta_1 := \mathbf{v}/c$, $\beta_2 := \mathbf{u}'/c$ and $\beta := \mathbf{u}/c$, and also $\beta = \|\beta\|$ etc. for the modulus, as well as $\gamma := \gamma(\beta) = 1/\sqrt{1-\beta^2}$ and correspondingly for $\gamma_i := \gamma(\beta_i)$. Then (4) and (6) read

$$\boldsymbol{\beta} \coloneqq \boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2} = \frac{1}{1 + \boldsymbol{\beta}_{1} \cdot \boldsymbol{\beta}_{2}} \left\{ \boldsymbol{\beta}_{1} + \boldsymbol{\beta}_{2}^{\parallel} + \boldsymbol{\gamma}_{1}^{-1} \boldsymbol{\beta}_{2}^{\perp} \right\}$$
(8a)

$$= \frac{1}{1+\beta_1\cdot\beta_2} \left\{ \beta_1 + \beta_2 + \frac{\gamma_1}{1+\gamma_1}\beta_1 \times (\beta_1 \times \beta_2) \right\}$$
(8b)

$$= \frac{1}{1+\beta_1\cdot\beta_2} \left\{ \beta_1 + \gamma_1^{-1}\beta_2 + \frac{\gamma_1}{1+\gamma_1}\beta_1 \left(\beta_1\cdot\beta_2\right) \right\}, \quad (8c)$$

where we defines the binary \oplus -operation – the "Einstein addition law" – by the expressions on the right hand side, which, apart from the square-root in γ_1 , involves the velocities only in a rational fashion. The superscripts || and \perp now refer to the Euclidean orthogonal projections parallel and perpendicular to β_1 , which, using $\mathbf{n}_1 := \beta_1/\beta_1$, are given in analogy to (5) by

$$\boldsymbol{\beta}_2^{\parallel} := \mathbf{n}_1(\mathbf{n}_1 \cdot \boldsymbol{\beta}_2) \quad \text{and} \quad \boldsymbol{\beta}_2^{\perp} := \boldsymbol{\beta}_2 - \boldsymbol{\beta}_2^{\parallel}.$$
 (9)

Equation (7) reads

$$\gamma = \gamma_1 \gamma_2 (1 + \beta_1 \cdot \beta_2), \qquad (10)$$

showing that $\gamma < \infty$ if $\gamma_i < \infty$ (i = 1, 2), and that $\gamma \to \infty$ if and only if at least one $\gamma_i \to \infty$. This implies that

$$\oplus: \mathring{B}_1(\mathbb{R}^3) \times \mathring{B}_1(\mathbb{R}^3) \to \mathring{B}_1(\mathbb{R}^3), \qquad (11)$$

where $\mathring{B}_1(\mathbb{R}^3)$ denotes the open ball of unit radius in \mathbb{R}^3 . In this section we will clarify the properties of the binary operation that \oplus puts on $\mathring{B}_1(\mathbb{R}^3)$. This would be a tedious problem to do from the algebraic form alone. In fact, we will see that this algebraic form derives from an underlying group law, even though \oplus is not itself a group multiplication. This results in \oplus being – in some sense – a minimal non-associative generalisation of group multiplication.

In order to appreciate this later insight we encourage the reader to work on the following two exercises (doing the first, trying the second):

Exercise 1. Use the expression (8a) for $\beta = \beta_1 \oplus \beta_2$ and solve it for β_2 , showing that

$$\boldsymbol{\beta}_2 = (-\boldsymbol{\beta}_1) \oplus \boldsymbol{\beta} = \frac{1}{1 - \boldsymbol{\beta}_1 \cdot \boldsymbol{\beta}} \left\{ -\boldsymbol{\beta}_1 + \boldsymbol{\beta}^{\parallel} + \boldsymbol{\gamma}_1^{-1} \boldsymbol{\beta}^{\perp} \right\}.$$
(12)

Hint: Equation (8a) can be used to express β^{\parallel} and β^{\perp} as functions of β_1 and β_2 , and also to express $\beta \cdot \beta_1$ in terms of $\beta_1 \cdot \beta_2$ and γ_1 . The rest is straightforward computation.

Definition 1. *The expression* (12) *is called the* velocity difference *between* β *and* β_{1} .

Exercise 2. *Try to repeat Exercise1, now solving* $\beta = \beta_1 \oplus \beta_2$ *for* β_1 *(as function of* β *and* β_2 *) rather than* β_2 *(as function of* β *and* β_1 *) as above. There is a unique solution!*

It is likely that you will not find the solution to Exercise 2 at this point (it is given in (83) below). It is *not* given by $\beta \oplus (-\beta_2)$ (as one might have expected at first) unless β and β_2 are parallel. Hence there is no immediate analogue of Definition 1 for the "other" velocity difference, i.e. that between β and β_2 .

2.3 The generalised orthogonal group and its Lie algebra

Let us first consider the general case where V be a real n-dimensional vector space, V^{*} its dual space, and $\eta \in V^* \otimes V^*$ a symmetric, non-generate, bilinear form of any signature. Recall that the *signature* of such a form is given by a pair (n_-, n_+) of non-negative integers which denote the dimensions of maximal linear subspaces V_- and V_+ of V restricted to which η is negative- and positive-definite, respectively.³ Whereas n_{\pm} are uniquely defined by η , V_{\pm} are not. Note that $V = V_- \oplus V_+$,

³ Our convention is mostly shared in the physics literature, whereas in the mathematics literature The word "signature" is often used differently. For example, Greub (1975, p. 269) calls n_+ the *index* and $n_+ - n_-$ the *signature*.

hence $n = n_- + n_+$, with V_+ and $V_- \eta$ -orthogonal to each other. If $n_- = 1$ and $n_+ = n - 1$ (or vice versa; though we will stick to the "mostly plus" convention in this paper) η is called a *Minkowski metric* in n dimensions. All statements in this subsection apply to general n > 1 and (n_-, n_+) , unless stated otherwise.

Definition 2. *The* (generalised) orthogonal group *of* (V,η) *, denoted by* $O(V,\eta)$ *, is defined by the following subgroup of the group* Gl(V) *of linear isomorphsims of* V*:*

$$O(V,\eta) = \left\{ L \in GL(V) : \eta(L\nu, Lw) = \eta(\nu, w) ; \forall \nu, w \in V \right\}.$$
(13)

It follows from this definition that $det(L) = \pm 1$ for all $L \in O(V, \eta)$. The special (generalised) orthogonal group is the subgroup of orientation preserving elements in $O(V, \eta)$:

$$SO(V,\eta) = \left\{ L \in O(V,\eta) : det(L) = 1 \right\}.$$
(14)

If η is either positive or negative definite, i.e. if either $n_- = 0$ or $n_+ = 0$, the group $O(V,\eta)$ contains two connected components with $SO(V,\eta)$ being the component containing the group identity (usually called the "identity component"). In all other cases with, i.e. for $n_{\pm} \ge 1$, the group $O(V,\eta)$ decomposes (as set) into the disjoint union (denoted by \sqcup) of four connected components (see, e.g., the book by O'Neill (1983, pp. 237)),

$$O(V,\eta) = \underbrace{O_{(+,+)}(V,\eta) \sqcup O_{(-,-)}(V,\eta)}_{SO(V,\eta)} \sqcup O_{(-,+)}(V,\eta) \sqcup O_{(+,-)}(V,\eta) , \quad (15)$$

where the \pm in the (first, second) slot indicates whether the orientation amongst the (negative, positive)—definite subspaces is preserved or reversed.

If $\{e_1, \dots, e_n\}$ is a basis for V with dual basis $\{\theta^1, \dots, \theta^n\}$ for V^{*}, so that $\theta^a(e_b) = \delta^a_b$, we can write

$$\mathsf{L} = \mathsf{L}^{\mathfrak{a}}_{\ \mathfrak{b}} \ \mathfrak{e}_{\mathfrak{a}} \otimes \mathfrak{\theta}^{\mathfrak{b}} \tag{16}$$

and

$$\eta = \eta_{ab} \,\theta^a \otimes \theta^b \tag{17}$$

We recall that the metric η defines an isomorphism

$$\eta_{\downarrow}: V \to V^*, \quad \nu \mapsto \eta_{\downarrow}(\nu) := \eta(\nu, \cdot), \tag{18}$$

that has an inverse

$$\eta_{\uparrow}: V^* \to V, \quad \lambda \mapsto \eta_{\uparrow}(\lambda) := (\eta_{\downarrow})^{-1}(\lambda).$$
 (19)

Hence η defines a likewise non-degenerate symmetric bilinear form in the dual space V^{*}, which we call η^* . Its basis-idependent definition is

$$\eta^*(\lambda,\sigma) := \eta\big(\eta_{\uparrow}(\lambda), \eta_{\uparrow}(\sigma)\big) = \lambda\big(\eta_{\uparrow}(\sigma)\big) = \sigma\big(\eta_{\uparrow}(\lambda)\big) \,. \tag{20}$$

With respect to the pair auf dual bases this leads to

$$\eta^* = \eta^{ab} e_a \otimes e_b \tag{21}$$

where η^{ab} are the components of the matrix that is inverse to η_{ab} . Hence we can write $\eta_{ab} = \eta(e_a, e_b)$ and $\eta^{ab} = \eta^*(\theta^a, \theta^b)$. Equivalent to (13) is then

$$O(V,\eta) = \left\{ L \in GL(V) : \eta^* \left(L^\top \lambda, L^\top \sigma \right) = \eta^*(\lambda, \sigma) ; \, \forall \lambda, \sigma \in V^* \right\}.$$
(22)

In terms of components (13) and (22) read, respectively,

$$\mathcal{L}^{a}_{\ c}\mathcal{L}^{b}_{\ d}\eta_{ab} = \eta_{cd}\,,\tag{23a}$$

$$L^{a}_{\ c}L^{b}_{\ d}\eta^{cd} = \eta^{ab}.$$
^(23b)

Taking the s-derivative at s = 0 of (13), where L = L(s) is a differentiable curve in $O(V, \eta)$ with $L(s = 0) = id_V$, we obtain for $\ell = dL/ds|_{s=0}$ the defining relation for the Lie algebra:

$$\mathfrak{o}(V,\eta) = \left\{ \ell \in \operatorname{End}(V) : \eta(\ell \nu, w) = -\eta(\nu, \ell w) ; \, \forall \nu, w \in V \right\}.$$
(24)

In other words, $o(V, \eta)$ is given by the η -antisymmetric endomorphisms of V.

The given isomorphism (18) provides a specific identification of V^{*} with V via replacing any $\alpha \in V^*$ with $\nu = \eta_{\uparrow}(\alpha)$. Hence we may also identify End(V), which is *naturally* isomorphic to $V \otimes V^*$, with $V \otimes V$. This will notationally simplify later expressions. In particular, $\mathfrak{o}(V, \eta)$ is identified with $V \wedge V$. The natural Lie product on End(V), which is just given by the commutator, induces a Lie product on $V \otimes V$, which on pure tensor products is given by

$$\left[\nu \otimes w, \nu' \otimes w'\right] = \eta(w, \nu') \nu \otimes w' - \eta(w', \nu) \nu' \otimes w$$
(25)

with unique bilinear extension to all of $V \otimes V$. On $\mathfrak{o}(V, \eta)$ this becomes (recall $v \wedge w = v \otimes w - w \otimes v$):

$$\begin{bmatrix} \nu \wedge w, \nu' \wedge w' \end{bmatrix} = \eta(\nu, w') w \wedge \nu' + \eta(w, \nu') \nu \wedge w' - \eta(\nu, \nu') w \wedge w' - \eta(w, w') \nu \wedge \nu'.$$
(26)

If $\{e_1, \dots, e_n\}$ is a basis for V with $\eta_{ab} := \eta(e_a, e_b)$ and $M_{ab} := e_a \wedge e_b$, (26) takes the form found in many physics-textbooks:

$$[M_{ab}, M_{cd}] = \eta_{ad} M_{bc} + \eta_{bc} M_{ad} - \eta_{ac} M_{bd} - \eta_{bd} M_{ad}.$$
(27)

The set $\{M_{\alpha}b : 1 \le \alpha < b \le n\}$ forms a basis for the $\frac{1}{2}n(n-1)$ - dimensional Lie algebra $\mathfrak{o}(V,\eta)$.

2.4 Restriction to four dimensions and Lorentzian signature

From now on we restrict to the special-relativistic case in which $(n_-, n_+) = (1,3)$.⁴ In this case the (full) *orthogonal group*, $O(V,\eta)$, is called the *Lorentz group*, abbreviated by Lor. As we have seen above, it has four components. The component containing the identity is $O_{(+,+)}(V,\eta)$ and usually called the *proper or-thochronous Lorentz group*, where the "proper" stands for "overall-orientation preserving", meaning that one restricts to $L \in SO(V,\eta) = O_{(+,+)}(V,\eta) \sqcup O_{(-,-)}(V,\eta)$, i.e. to those L with det(L) = 1. "Orthochronous" stands for "time-orientation preserving", which means that one restricts to $L \in O_{(+,+)}(V,\eta) \sqcup O_{(+,-)}(V,\eta)$, i.e. to those L with $\eta(Lv,v) < 0$ for all timelike v. Note that $O_{(+,+)}(V,\eta) \sqcup O_{(-,+)}(V,\eta)$, too, is a two-component subgroup of $O(V,\eta)$ which is sometimes called the *orthochronous Lorentz group*; compare, e.g., (Streater and Wightman, 1964, p. 11).

Remark 3. In what follows we will restrict attention to the component of the identity only and simply write

Lor :=
$$O_{(+,+)}(V,\eta)$$
, for $(n_-, n_+) = (1,3)$. (28)

Note that often Lor (or just L) stands for the whole Lorentz group whereas the proper orthochronous subgroup is denoted by Lor^{\uparrow}_+ (or L^{\uparrow}_+), where the subscript + represents the "proper" and the superscript \uparrow the "orthochronous". But since here we do consider time- or space-reversing maps we simplify the notation as indicated. We remark that everything we are going to say simply extends to all time-orientation preserving Lorentz transformations, i.e. to $O_{(+,-)}(V,\eta) \sqcup O_{(+,-)}(V,\eta)$ (or to $\text{Lor}^{\uparrow}_+ \sqcup \text{Lor}^{\uparrow}_-$). In order to also include time-orientation reversing transformations we would have to generalise our later Definition 7 of the set of states of motion to also include the other connected component (called V_1^-) of the hyperboloid of unit timelike vectors. We avoid these complications since they are irrelevant for what is to follow.

According to (23), the matrices representing Lorentz transformations must satisfy (23) with the basis $\{e_0, e_1, e_2, e_3\}$ so chosen that

 $1 = -\eta_{00} = \eta_{11} = \eta_{22} = \eta_{33} \quad \text{and} \quad \eta_{ab} = 0 \quad \text{for} \quad a \neq b \,. \eqno(29)$

If we write a general matrix with components L^{a}_{b} in (1+3)-decomposed form as⁵

$$\left\{ L^{a}_{b} \right\} = \begin{pmatrix} \mathbf{c} & \mathbf{a}^{\top} \\ \mathbf{b} & \underline{\mathbf{M}} \end{pmatrix}, \qquad (30)$$

where $c \in \mathbb{R}$, $a, b \in \mathbb{R}^3$, and $\underline{M} \in \text{End}(\mathbb{R}^3)$ is a (3×3) -matrix⁶, equations (23a)

 ⁴ Or, alternatively, (n₋, n₊) = (3, 1). But in this paper we shall adopt the "mostly plus" convention.
 ⁵ The symbol c for the upper-left matrix entry in (30) should not be confused with the velocity of light, which does not explicitly appear in any of the following formulae.

⁶ Generally, in this section elements in R³ are written by bold-faced letters and elements in End(R³) (i.e. 3 × 3 matrices) by underlined bold-faced letters. The transposed object of either of these is indicated by a superscript ⊤.

and (23b) are equivalent to, respectively,

$$\|\mathbf{b}\|^2 = c^2 - 1, \qquad (31a)$$

$$\mathbf{\underline{M}}^{\mathsf{T}}\mathbf{b} = \mathbf{c}\mathbf{a}, \qquad (31b)$$

$$\underline{\mathbf{M}}^{\top}\underline{\mathbf{M}} = \underline{\mathbf{E}}_3 + \mathbf{a} \otimes \mathbf{a}^{\top}, \qquad (31c)$$

and

$$\|\mathbf{a}\|^2 = \mathbf{c}^2 - 1, \qquad (32a)$$

$$\underline{\mathbf{M}}\mathbf{a} = \mathbf{c}\mathbf{b}\,,\tag{32b}$$

$$\underline{\mathbf{M}}\,\underline{\mathbf{M}}^{\top} = \underline{\mathbf{E}}_3 + \mathbf{b} \otimes \mathbf{b}^{\top}, \qquad (32c)$$

where $\underline{\mathbf{E}}_3$ denotes the (3×3) unit matix. Note that, according to (23), (23b) follow from equations (23a) by replacing L with its transposed (due to the numerical equality of η_{ab} with η^{ab}). Accordingly, the set (32) follows from the set (31) by exchanging **a** with **b** and <u>M</u> with $\underline{\mathbf{M}}^{\mathsf{T}}$.

Clearly, spatial rotations and pure boosts (1) satisfy these equations. In fact, "spatial rotation" here means to embed $R : SO(3) \hookrightarrow Lor, \underline{D} \mapsto R(\underline{D})$, according to

$$\mathsf{R}(\underline{\mathbf{D}}) := \begin{pmatrix} 1 & \mathbf{0}^\top \\ \mathbf{0} & \underline{\mathbf{D}} \end{pmatrix} \,. \tag{33}$$

With respect to the same basis (for which $x^0 = ct$), a boost (1) corresponds to

$$B(\boldsymbol{\beta}) := \begin{pmatrix} \boldsymbol{\gamma} & \boldsymbol{\gamma}\boldsymbol{\beta}^{\mathsf{T}} \\ \boldsymbol{\gamma}\boldsymbol{\beta} & \underline{\mathbf{E}}_{3} + (\boldsymbol{\gamma} - 1)\mathbf{n} \otimes \mathbf{n}^{\mathsf{T}} \end{pmatrix}, \qquad (34)$$

where we abbreviate $\mathbf{n} := \beta/\beta$ and each entry in the matrix on the right-hand side is to be regarded as function of β . In particular, since $\beta^2 = 1 - \gamma^{-2}$, we have $(\gamma - 1)\mathbf{n} \otimes \mathbf{n}^{\top} = \frac{\gamma^2}{1+\gamma}\beta \otimes \beta^{\top}$, which is a continuously differentiable function of β .

For (33) satisfaction of (31) is immediate (with (32) following by replacing $\underline{\mathbf{D}}$ with its transposed (= inverse)). Likewise, noting that $\underline{\mathbf{E}}_3 + (\gamma - 1)\mathbf{n} \otimes \mathbf{n}^\top = \mathbf{P}_\perp + \gamma \mathbf{P}_\parallel$, with \mathbf{P}_\parallel and \mathbf{P}_\perp the projectors in \mathbb{R}^3 perpendicular and parallel to \mathbf{n} , respectively, we immediately see that (31b) and (31c) are satisfied. Equation (31a) is equivalent to the identity $\beta^2 \gamma^2 = \gamma^2 - 1$. The converse to this result is the following:

Theorem 4. Any Lorentz transformation (30) that is 'proper', i.e. satisfies det(L) = +1 and 'orthochronous', i.e. $c \ge 1$, can be composed into the product of a rotation and a boost. This decomposition is just the "polar decomposition" with respect to the standard Euclidean metric in \mathbb{R}^4 (compare Appendix A) and hence unique if the order of rotation-boost-multiplication is fixed. The reversed order has the same rotation but a boost velocity that differs from the first by that rotation. That is,

$$\left\{L^{a}_{b}\right\} = \begin{pmatrix} c & a^{\dagger} \\ b & \underline{\mathbf{M}} \end{pmatrix} = B(\boldsymbol{\beta})R(\underline{\mathbf{D}}) = R(\underline{\mathbf{D}})B(\boldsymbol{\beta}'), \qquad (35a)$$

where $\underline{\mathbf{D}} \in SO(3)$, $\boldsymbol{\beta} \in \mathring{B}_1(\mathbb{R}^3)$, and are $\boldsymbol{\beta}' \in \mathring{B}_1(\mathbb{R}^3)$ are determined as rational functions of $\gamma, \mathbf{a}, \mathbf{b}$, and $\underline{\mathbf{M}}$ by

$$\underline{\mathbf{D}} := \underline{\mathbf{M}} - \frac{\mathbf{b} \otimes \mathbf{a}^{\top}}{\gamma + 1}, \qquad (35b)$$

$$\boldsymbol{\beta} := \mathbf{b}/\mathbf{c}, \quad \boldsymbol{\beta}' = \mathbf{a}/\mathbf{c}.$$
 (35c)

We further have $\beta = \underline{D}\beta'$ *; compare* (37) *and* (42) *below.*

Proof. We only need to show the second equality in (35b), as the third then follows from the proof of the first. We proceed by proving three things: 1) That the product $B(\beta)R(\underline{D})$ with \underline{D} and β as in (35b) and (35c) indeed equals $\{L^{\alpha}_{b}\}$; 2) that $B(\beta) \in GL(\mathbb{R}^{4})$ is symmetric and positive definite; and 3) that $\underline{D} \in SO(3)$. Now, the product of $B(\beta)$ given by (34) and $R(\underline{D})$ given by (33) is

$$\begin{pmatrix} \gamma & \gamma(\underline{\mathbf{D}}^{\top}\boldsymbol{\beta})^{\top} \\ \gamma\boldsymbol{\beta} & \underline{\mathbf{D}} + (\gamma - 1)\mathbf{n} \otimes (\underline{\mathbf{D}}^{\top}\mathbf{n})^{\top} \end{pmatrix}.$$
 (36)

This must equal (30), which for the upper-left entry implies $\gamma = c$ and the lowerleft entry $\gamma \beta = \mathbf{b}$. Since $\|\gamma \beta\|^2 = \gamma^2 \beta^2 = \gamma^2 - 1$ this is only consistent with the first equation if $\|\mathbf{b}\|^2 = c^2 - 1$, which is just guaranteed by (31a). Given $\gamma \beta = \mathbf{b}$, equality of the upper-right entry of (36) and (30) is equivalent to $\mathbf{\underline{D}}^{\top}\mathbf{b} = \mathbf{a}$. This is indeed satisfied by (35b) since then

$$\underline{\mathbf{D}}^{\top}\mathbf{b} = \underline{\mathbf{M}}^{\top}\mathbf{b} - \mathbf{a}\frac{\|\mathbf{b}\|^2}{\gamma+1} = \mathbf{a}\left(\mathbf{c} - \frac{\mathbf{c}^2 - 1}{\gamma+1}\right) = \mathbf{a}$$
(37)

where we used (31a) and (31b) for the second and $c = \gamma$ for the third equality. This also shows equality of the lower-right entries, which requires

$$\underline{\mathbf{D}} = \underline{\mathbf{M}} - (\gamma - 1)\mathbf{n} \otimes (\underline{\mathbf{D}}^{\top}\mathbf{n})^{\top} = \underline{\mathbf{M}} - (\gamma - 1)\frac{\mathbf{b} \otimes (\underline{\mathbf{D}}^{\top}\mathbf{b})^{\top}}{\beta^2 \gamma^2}, \qquad (38)$$

which is equivalent to (35b) in view of (37) and $\beta^2 \gamma^2 = \gamma^2 - 1$. Next we check orthogonality of **D**:

$$\underline{\mathbf{D}}^{\top} \underline{\mathbf{D}} = \left(\underline{\mathbf{M}}^{\top} - \frac{\mathbf{a} \otimes \mathbf{b}^{\top}}{\gamma + 1} \right) \left(\underline{\mathbf{M}} - \frac{\mathbf{b} \otimes \mathbf{a}^{\top}}{\gamma + 1} \right)
= \underline{\mathbf{M}}^{\top} \underline{\mathbf{M}} - \frac{\underline{\mathbf{M}}^{\top} \mathbf{b} \otimes \mathbf{a} + \mathbf{a} \otimes (\underline{\mathbf{M}}^{\top} \mathbf{b})^{\top}}{\gamma + 1} + \|\mathbf{b}\|^{2} \frac{\mathbf{a} \otimes \mathbf{a}^{\top}}{(\gamma + 1)^{2}}
= \underline{\mathbf{E}}_{3} + \mathbf{a} \otimes \mathbf{a}^{\top} \left(1 - \frac{2\mathbf{c}}{\gamma + 1} + \frac{\mathbf{c}^{2} - 1}{(\gamma + 1)^{2}} \right)
= \underline{\mathbf{E}}_{3},$$
(39)

where we used (31b) and (31c) in the third and $c = \gamma$ in the last step. Similarly we could have proven $\underline{\mathbf{D}} \underline{\mathbf{D}}^{\top} = \underline{\mathbf{E}}_3$ using the relations (32). Finally, symmetry of B($\boldsymbol{\beta}$)

is obvious and positive definiteness follows from ist eigenvalues, which depend on the modulus β of β but not on its direction. Hence we may choose, e.g., $\beta = \beta e_1$ to calculate its characteristic polynomial, which reads

$$\mathsf{P}(\lambda) = \left[(\gamma - \lambda)^2 - \beta^2 \gamma^2 \right] (1 - \lambda)^2 \,. \tag{40}$$

Hence the eigenvalues are

$$\lambda_1 = \sqrt{\frac{1+\beta}{1-\beta}}, \qquad \lambda_2 = \sqrt{\frac{1-\beta}{1+\beta}}, \qquad \lambda_{3,4} = 1.$$
(41)

Finally, given orthogonality of $R(\underline{\mathbf{D}})$ and symmetry as well as positive-definiteness of $B(\beta)$, we see that the right-hand side of (35a) ist just the polar decomposition of $\{L^{\alpha}_{b}\} \in GL(\mathbb{R}^{4})$, which is unique – if we agree on the order of the two terms on the right hand side, i.e. the orthogonal matrix $R(\underline{\mathbf{D}})$ to the right of the positive-definite symmetric one. Had we chosen the opposite order, as in the third equality in (35a), the two matrices would again be uniquely determined with the same rotation but a different (rotated) boost $\beta' = \underline{\mathbf{D}}^{-1}\beta$. This is a simple consequence of the general relation

$$R(\underline{\mathbf{D}})B(\boldsymbol{\beta})R(\underline{\mathbf{D}}^{-1}) = B(\underline{\mathbf{D}}\boldsymbol{\beta}), \qquad (42)$$

that is an immediate consequence of (33) and (34). Indeed, since $\beta = \mathbf{b}/\mathbf{c}$ and $\underline{\mathbf{D}}^{-1} = \underline{\mathbf{D}}^{\top}$, equation (37) precisely shows just that.

2.5 Polar decomposition of boost products

As in section (2.4) we pick a state of motion $s \in S$ which we take as our zeroth (time-like) basis vector for V; i.e. we choose $e_0 = s$ and complement this to an orthonormal basis $\{e_0, e_1, e_2, e_3\}$ of V. With respect to that choice we shall now speak of pure boosts, whose matrix representatives look like (34) and, and also of "polar decomposition". The task is to polar decompose the product of boosts $B(\beta_1)$ and $B(\beta_2)$:

$$B(\boldsymbol{\beta}_1)B(\boldsymbol{\beta}_2) = B(\boldsymbol{\beta})R(\underline{\mathbf{D}}).$$
(43)

The boost parameter β and the rotation matrix \underline{D} on the right-hand side are uniquely determined by β_1 and β_2 . In other words, there are functions

$$\boldsymbol{\beta}: \mathring{B}_1(\mathbb{R}^3) \times \mathring{B}_1(\mathbb{R}^3) \to \mathring{B}_1(\mathbb{R}^3), \qquad (44a)$$

$$\underline{\mathbf{D}}: \mathring{B}_1(\mathbb{R}^3) \times \mathring{B}_1(\mathbb{R}^3) \to \operatorname{SO}(3), \qquad (44b)$$

which we now determine. For that we just need to follow the procedure outlined in section 2.4. Using the matrix representation (34) for $B(\beta_1)$ and $B(\beta_2)$ we obtain a

matrix of the form (30) with

$$\mathbf{c} = \gamma_1 \gamma_2 (1 + \beta_1 \beta_2), \qquad (45a)$$

$$\mathbf{a} = \gamma_1 \gamma_2 \left(\beta_2 + \beta_1^{\parallel} + \gamma_2^{-1} \beta_1^{\perp} \right), \tag{45b}$$

$$\mathbf{b} = \gamma_1 \gamma_2 \left(\beta_1 + \beta_2^{\parallel} + \gamma_1^{-1} \beta_2^{\perp} \right), \tag{45c}$$

$$\underline{\mathbf{M}} = \underline{\mathbf{E}}_3 + (\gamma_1 - 1) \, \mathbf{n}_1 \otimes \mathbf{n}_1^\top + (\gamma_2 - 1) \, \mathbf{n}_2 \otimes \mathbf{n}_2^\top + \left[\beta_1 \gamma_1 \beta_2 \gamma_2 + (\gamma_1 - 1)(\gamma_2 - 1)(\mathbf{n}_1 \cdot \mathbf{n}_2) \right] \mathbf{n}_1 \otimes \mathbf{n}_2^\top, \qquad (45d)$$

where the superscripts \parallel and \perp on β_1 and β_2 refer to the projections parallel and perpendicular to the "other" β , i.e. β_2 and β_1 respectively. From $c = \gamma$ and (35c) we get

$$\boldsymbol{\beta} = \mathbf{b}/\mathbf{c} = \frac{\boldsymbol{\beta}_1 + \boldsymbol{\beta}_2^{\parallel} + \boldsymbol{\gamma}_1^{-1} \boldsymbol{\beta}_2^{\perp}}{1 + \boldsymbol{\beta}_1 \cdot \boldsymbol{\beta}_2} = \boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2, \qquad (46a)$$

$$\boldsymbol{\beta}' = \mathbf{a}/\mathbf{c} = \frac{\boldsymbol{\beta}_2 + \boldsymbol{\beta}_1^{\parallel} + \boldsymbol{\gamma}_2^{-1} \boldsymbol{\beta}_1^{\perp}}{1 + \boldsymbol{\beta}_1 \cdot \boldsymbol{\beta}_2} = \boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_1, \qquad (46b)$$

and, since $c = \gamma$,

$$\gamma = \gamma(\beta) = \gamma(\beta') = \gamma_1 \gamma_2 (1 + \beta_1 \cdot \beta_2) = \gamma_1 \gamma_2 (1 + \beta_1 \beta_2 \cos(\varphi)), \quad (46c)$$

if φ denotes the angle between β_1 and β_2 . Hence the boost contained in the polar decomposition of the product of boosts with parameters β_1 and β_2 is just the boost with Einstein added parameters $\beta = \beta_1 \oplus \beta_2$.

Next we turn to the rotation $\underline{\mathbf{D}}$. Like β it is a function of β_1 and β_2 and is given a special name:

Definition 5. The rotation matrix $\underline{\mathbf{D}}$ resulting from polar decomposition (43) of the product of two boosts $B(\beta_1) B(\beta_2)$ is called the Thomas rotation and denoted by $\underline{\mathbf{T}}(\beta_1, \beta_2)$.

The Thomas rotation can be written down by forming the expression on the right-hand side of (35b) using (46), but the algebraic expression is complex and not immediately telling. It can be understood in simple terms as follows. First we observe that it is a linear combination of $\underline{\mathbf{E}}_3$, $\mathbf{n}_1 \otimes \mathbf{n}_1$, $\mathbf{n}_2 \otimes \mathbf{n}_2$, $\mathbf{n}_1 \otimes \mathbf{n}_2$, and $\mathbf{n}_2 \otimes \mathbf{n}_1$. Hence it maps the plane $\text{Span}\{\mathbf{n}_1, \mathbf{n}_2\} = \text{Span}\{\beta_1, \beta_2\}$ into itself and leaves the orthogonal complement pointwise fixed; in other words: it is a rotation in the plane spanned by the two boost velocities. That plane contains \mathbf{a} and \mathbf{b} and we know from (37) that $\mathbf{b} = \underline{\mathbf{D}}\mathbf{a}^{.7}$ Hence the rotation angle θ is the angle between \mathbf{a} and \mathbf{b} ,

$$\cos(\theta) = \frac{\mathbf{a} \cdot \mathbf{b}}{\|\mathbf{a}\| \|\mathbf{b}\|} = \frac{\mathbf{a} \cdot \mathbf{b}}{\gamma^2 - 1},$$
(47)

⁷ Note that whereas $\mathbf{b} = \underline{\mathbf{D}}\mathbf{a}$ is always true, $\underline{\mathbf{D}}$ need generally not be a rotation in the plane spanned by \mathbf{a} and \mathbf{b} . For the Lorentz transformation resulting from the composition of two boosts, however, this is the case.

counted positively with respect to the orientation given by the ordered pair $\{\mathbf{a}, \mathbf{b}\}$. Here we used $\|\mathbf{a}\| = \|\mathbf{b}\| = \sqrt{\gamma^2 - 1}$ (compare (31a) and (31b), using $\mathbf{c} = \gamma$) in the last step.

On the other hand, according to the general formula for the rotation angle, we have

$$1 + 2\cos(\theta) = \operatorname{trace}(\underline{\mathbf{D}}) = \operatorname{trace}(\underline{\mathbf{M}}) - \frac{\mathbf{a} \cdot \mathbf{b}}{\gamma + 1}, \qquad (48)$$

using (35b) in the last step. Replacing $\mathbf{a} \cdot \mathbf{b}$ according to (47) leads to an equation for $\cos(\theta)$ that we can solve:

$$\cos(\theta) = \frac{\operatorname{trace}(\underline{\mathbf{M}}) - 1}{\gamma + 1} \,. \tag{49}$$

Now, (45d) gives

$$\operatorname{trace}(\underline{\mathbf{M}}) - 1 = 3 + (\gamma_1 - 1) + (\gamma_2 - 1) + \beta_1 \gamma_1 \beta_2 \gamma_2 \cos(\varphi) + (\gamma_1 - 1)(\gamma_2 - 1) \cos^2(\varphi) = 2 + \gamma_1 \gamma_2 (1 + \beta_1 \beta_2 \cos(\varphi)) - (\gamma_1 - 1)(\gamma_2 - 1) \sin^2(\varphi) = 1 + (\gamma + 1) - (\gamma_1 - 1)(\gamma_2 - 1) \sin^2(\varphi) .$$
 (50)

where, as in (46c), ϕ denotes the angle between the velocities, i.e.

$$\cos(\varphi) := \mathbf{n}_1 \cdot \mathbf{n}_2 \,. \tag{51}$$

Hence (49) becomes

$$\cos(\theta) = 1 - \frac{(\gamma_1 - 1)(\gamma_2 - 1)\sin^2(\varphi)}{1 + \gamma_1\gamma_2 + \sqrt{(\gamma_1^2 - 1)(\gamma_2^2 - 1)}\cos(\varphi)}.$$
 (52)

We have deliberately written the denominator in the second term, which is just $1 + \gamma$, in terms of γ_1 , γ_2 , and φ , so as to explicitly display θ as function of the moduli of the two boost velocities (i.e. their γ -factors) and the angle φ between them.

Alternatively, instead of $(\gamma_1, \gamma_2, \varphi)$ we may express $\cos(\theta)$ as function of $(\gamma_1, \gamma_2, \gamma)$. This is achieved by writing $\sin^2(\varphi) = 1 - \cos^2(\varphi)$ and replacing all occurrences of $\cos(\varphi)$ in (52) with the expression that follows from (46c):

$$\cos(\varphi) = \frac{\gamma - \gamma_1 \gamma_2}{\sqrt{(\gamma_1^2 - 1)(\gamma_2^2 - 1)}} \,.$$
(53)

This gives after a few elementary steps

$$\cos(\theta) = \frac{(1+\gamma+\gamma_1+\gamma_2)^2}{(1+\gamma)(1+\gamma_1)(1+\gamma_2)} - 1.$$
 (54)



Figure 1: Non commutativity of velocity addition due to Thomas rotation. Shown ist the addition of two perpendicular velocities of equal magnitude $\beta = 0.8$ so that $\gamma_*^{-1} = 0.6$. The Thomas rotation $T(\beta_1, \beta_2)$ rotates by an positive angle θ in the oriented plane spanned by the ordered pair {**a**, **b**}. Since **a** is proportional to $\beta_2 \oplus \beta_1$ and **b** to $\beta_1 \oplus \beta_2$, this rotation is in the clockwise – i.e. negative – orientation with respect to the ordered pair { β_1, β_2 }.

This expression is remarkable for its simple structure and permutation symmetry in $\{\gamma, \gamma_1, \gamma_2\}$. An alternative way to write it is in terms of $\cos(\theta/2) = \sqrt{(\cos(\theta) + 1)/2}$ and $\cosh(\rho) := \gamma$, so that $(1 + \gamma) = 2 \cosh^2(\rho/2)$, as well as the corresponding equations for γ_i in terms of ρ_i $(i = 1, 2)^8$

$$\cos(\theta/2) = \frac{1 + \cosh(\rho) + \cosh(\rho_1) + \cosh(\rho_2)}{4 \cosh(\rho/2) \cosh(\rho_1/2) \cosh(\rho_2/2)}.$$
 (55)

These formulae, as as well as (52), are well known in the literature; an early appearance of (55) being (Macfarlane, 1962, formula (124)). Elementary derivations are often claimed to be unduly tedious or even a "herculean" task" (Ungar, 2002, p. 28), and more elegant ways using Clifford algebras have been given (Urbantke, 1990). However, we believe that the derivation given here is sufficiently easy for presentation in, say, a basic lecture on SR.

It is important to note that the right hand side of (54) depends on the reference state which respect to which the β 's and hence the γ 's are defined. In fact, the $(-\gamma)$'s are just the Minkowskian scalar products between the corresponding states of motion with the reference state. In this way (54) can be written as rational function of scalar products of states of motion with the reference state, thereby displaying the dependence on the latter in a basis independent way. We will later adopt precisely that strategy: to reduce all statements to functions of scalar products of states or motion.

 $^{^{8}}$ The greek letter ρ is chosen because this quantity is usually referred to as "rapidity", introduced by Robb (1911).

2.6 On the magnitude of Thomas rotation

In Fig. 1 we show the Thomas rotation for the special case of mutually perpendicular velocities of equal magnitude; i.e. $\cos(\varphi) = 0$ and $\gamma_1 = \gamma_2 = \gamma_*$. In this case (52) reduces to

$$\cos(\theta) = \frac{2\gamma_*}{1 + \gamma_*^2},\tag{56}$$

which is a monotonically decreasing function of $\gamma_* \in (1, \infty)$ ranging from 1 to zero and corresponding to a monotonically increasing angle from 0 to $\pi/2$.

More generally, we may ask for the angle φ at which θ is largest for given $\gamma_{1,2}$. This can be answered using either expression (52) or (54), if in the latter case we regard γ again as function of $\gamma_{1,2}$ and φ according to (46c). This latter possibility turns out to be slightly more convenient, so let's follow this strategy and consider (54). Since cos is monotonically decreasing in $[0, \pi]$, a maximum of θ corresponds to a minimum of $\cos(\theta)$, which we now seek. The only dependence of $\cos(\theta)$ on φ is through γ . The φ derivative of the latter is $\gamma' := d\gamma/d\varphi = -\beta_1\gamma_1\beta_2\gamma_2\sin(\varphi)$, and $d\cos(\theta)/d\varphi = (d\cos(\theta)/d\gamma)\gamma'$. Hence stationary points exist either for $\gamma' = 0$, which is for the boundary values $\varphi = 0$ or $\varphi = \pi$ that correspond to aligned and anti-alingned velocities and which correspond to $\theta = 0$ (hence a minimum), or for $d\cos(\varphi)/d\gamma = 0$, containing the maxima. In view of (54), the latter equation is equivalent to

$$\frac{d}{d\gamma} \frac{(1+\gamma+\gamma_1+\gamma_2)^2}{(1+\gamma)} = \frac{1+\gamma+\gamma_1+\gamma_2}{(1+\gamma)^2} \left(1+\gamma-\gamma_1-\gamma_2\right) = 0.$$
(57)

Since all γ 's are positive, this is in turn equivalent to

$$\gamma = \gamma_{\max} := \gamma_1 + \gamma_2 - 1 \tag{58}$$

which, according to (53), corresponds to an angle ϕ_{max} between the velocities that satisfies

$$\cos(\varphi_{\max}) = \frac{\gamma_{\max} - \gamma_1 \gamma_2}{\sqrt{(\gamma_1^2 - 1)(\gamma_2^2 - 1)}} = -\sqrt{\frac{(\gamma_1 - 1)(\gamma_2 - 2)}{(\gamma_1 + 1)(\gamma_2 + 2)}}.$$
 (59)

The fact that this is negative means that $\varphi_{max} > \pi/2$. This means that the maximal Thomas angle is only obtained for obtuse angles between the velocities. The value

of the maximal Thomas angle is given by inserting (58) into (54):

$$\begin{aligned}
\cos(\theta_{\max}) &= \frac{(1 + \gamma_{\max} + \gamma_1 + \gamma_2)^2}{(1 + \gamma_{\max})(1 + \gamma_1)(1 + \gamma_2)} - 1 \\
&= \frac{3(\gamma_1 + \gamma_2) - \gamma_1\gamma_2 - 1}{(1 + \gamma_1)(1 + \gamma_2)} \\
&= 1 - 2\frac{(\gamma_1 - 1)(\gamma_2 - 1)}{(\gamma_1 + 1)(\gamma_2 + 1)} \\
&= 1 - 2\cos^2(\phi_{\max}) \\
&= -\cos(2\phi_{\max}).
\end{aligned}$$
(60)

From this we infer

$$\cos(\theta_{max}) + \cos(2\phi_{max}) = 2 \cos\left(\frac{\theta_{max} + 2\phi_{max}}{2}\right) \cos\left(\frac{\theta_{max} - 2\phi_{max}}{2}\right) = 0,$$
(61)

which in turn implies that either $\theta_{max} + 2\varphi_{max}$ or $\theta_{max} - 2\varphi_{max}$ is an integer multiple of π . But φ_{max} is obtuse, i.e. between $\pi/2$ and π , as follows from (59). If either γ_1 and/or γ_2 approach the value 1 from above $\cos(\varphi_{max})$ approaches the value zero from below and hence φ_{max} approaches $\pi/2$ from above. In this case θ_{max} approaches zero from below if we refer both angles to the orientation given to the plane of rotation by the ordered pair { β_1, β_2 } (compare Fig. 1). This shows that with that convention our unique solution to (61) is given by

$$\theta_{\max} = \pi - 2\varphi_{\max} \,. \tag{62}$$

So θ_{max} varies between 0 and $-\pi$ and its modulus exceeds the right angle $-\pi/2$ if $\cos(\theta_{max})$ turns negative. This, according to the third equation in (60), is the case if

$$\frac{(\gamma_1 - 1)(\gamma_2 - 1)}{(\gamma_1 + 1)(\gamma_2 + 1)} > \frac{1}{2}.$$
(63)

For equal velocities, i.e. $\gamma_1 = \gamma_2 = \gamma_* = 1/\sqrt{1-\beta_*^2}$ this happens for

$$\gamma_* > \frac{\sqrt{2} + 1}{\sqrt{2} - 1} \approx 5.828$$
 (64)

corresponding to

$$\beta_* > \frac{2^{5/4}}{2^{1/2} + 1} \approx 0.985, \tag{65}$$

that is, 98.5% of the velocity of light.

2.7 On the algebraic structure of Einstein addition

Using Definition 5 we shall now write

$$B(\boldsymbol{\beta}_1)B(\boldsymbol{\beta}_2) = B(\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2)).$$
(66)

Taking the transposed of that equation and using the symmetry of B, the orthogonality of R, and equivariance property (42), we get⁹

$$\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2 = \underline{\mathbf{T}}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2) \big(\boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_1 \big), \tag{67}$$

showing non-commutativity for non-collinear velocities. That we already know from $\mathbf{b} = \underline{\mathbf{D}}\mathbf{a}$ – see line above (47) – and (46). An immediate consequence of (67) is

$$\underline{\mathbf{T}}(\boldsymbol{\beta}_2, \boldsymbol{\beta}_1) = \underline{\mathbf{T}}^{-1}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2)$$
(68)

We also note the following: The vector $\beta_1 \oplus \beta_2$ is a linear combination of \mathbf{n}_1 and \mathbf{n}_2 with coefficients that only involve scalar products of these vectors; that is, the coefficients are invariant under $(\mathbf{n}_1, \mathbf{n}_2) \rightarrow (-\mathbf{n}_1, -\mathbf{n}_2)$ (even functions). The rotation matrix $T(\beta_1, \beta_2)$ is a linear combination of the identity and terms proportional to $\mathbf{n}_a \otimes \mathbf{n}_b^{\top}$ ($a, b \in \{1, 2\}$) with coefficients also depending only on the scalar products of these vectors. This implies

$$(-\boldsymbol{\beta}_1) \oplus (-\boldsymbol{\beta}_2) = -(\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2), \qquad (69a)$$

$$\underline{\mathbf{T}}(-\boldsymbol{\beta}_1, -\boldsymbol{\beta}_2) = \underline{\mathbf{T}}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2).$$
(69b)

Other immediate consequences of the same remark are:

$$(\underline{\mathbf{D}}\boldsymbol{\beta}_1) \oplus (\underline{\mathbf{D}}\boldsymbol{\beta}_2) = \underline{\mathbf{D}}(\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2), \qquad (70a)$$

$$\underline{\mathbf{T}}[\underline{\mathbf{D}}\boldsymbol{\beta}_1,\underline{\mathbf{D}}\boldsymbol{\beta}_2] = \underline{\mathbf{D}}\,\underline{\mathbf{T}}[\boldsymbol{\beta}_1,\boldsymbol{\beta}_2]\underline{\mathbf{D}}^{-1},\tag{70b}$$

for any $D \in SO(3)$.

Now, a general Lorentz transformation is written as

$$L(\boldsymbol{\beta}, \underline{\mathbf{D}}) := B(\boldsymbol{\beta}) R(\underline{\mathbf{D}}).$$
(71)

The composition of two such transformations is then given by

$$L(\boldsymbol{\beta}, \underline{\mathbf{D}}) = L(\boldsymbol{\beta}_1, \underline{\mathbf{D}}_1) L(\boldsymbol{\beta}_2, \underline{\mathbf{D}}_2)$$

= B(\beta_1) R(\beta_1) B(\beta_2) R(\beta_2)
= B(\beta_1) R(\beta_1) B(\beta_2) R(\beta_1^{-1}) R(\beta_1) R(\beta_2) (72)
= B(\beta_1) B(\beta_1 \beta_2) R(\beta_1 \beta_2)
= B(\beta_1 \oplus \beta_1 \beta_2) R(\beta_1 [\beta_1, \beta_1 \beta_2] \beta_1 \beta_2),

⁹ Historically it is interesting to note that this formula has already been written down (in different but easy-to-translate notation) by Silberstein (1914, p. 169, formula (7)). Without explicitly relating it to any orthogonal transformation, mere non-commutativity has been noted from the very beginning; e.g., by Einstein (1905, pp. 905-6) and Laue (1911, p. 44). Einstein remarked (p. 905) that the parallelogram-law for velocity addition is only approximately valid and at the same time finds it "remarkable" (p. 906) that the expression for the modulus of the composed velocities is a symmetric function in these velocities. Laue, after writing down the law of velocity addition, states: "The strangest (merkwürdigste) thing is that the two velocitied to be added do not enter on equal footing (gleichberechtigt)". Laue's text also contains an explicit discussion of the special case of orthogonal velocities and a figure equivalent to our Fig. 1 (Laue, 1911, p. 44, Fig. 4).

so that

$$\boldsymbol{\beta} = \boldsymbol{\beta}_1 \oplus \underline{\mathbf{D}}_1 \boldsymbol{\beta}_2, \qquad (73a)$$

$$\underline{\mathbf{D}} = \underline{\mathbf{T}}[\boldsymbol{\beta}_1, \boldsymbol{\beta}_2] \, \underline{\mathbf{D}}_1 \, \underline{\mathbf{D}}_2 \,. \tag{73b}$$

This should be compared with the group composition law of the semi-direct product $\mathbb{R}^3 \rtimes SO(3)$, which would be given by $\beta = \beta_1 + \underline{\mathbf{D}}_1 \beta_2$ and $\underline{\mathbf{D}} = \underline{\mathbf{D}}_1 \underline{\mathbf{D}}_2$, i.e. \oplus replaced by + and the Thomas rotation always the identity, as it it is the case for the Galilei group (with β replaced by **v**).

In order to deduce the parameters for the inverse transformation $L^{-1}(\boldsymbol{\beta}, \underline{\mathbf{D}})$ we first note that $B^{-1}(\boldsymbol{\beta}) = B(-\boldsymbol{\beta})$, as is already obvious from (34). Taking the inverse of both sides of (71) and using (70b) then gives

$$L^{-1}(\boldsymbol{\beta}, \underline{\mathbf{D}}) := R(\underline{\mathbf{D}}^{-1})B(-\boldsymbol{\beta}) = B(-\underline{\mathbf{D}}^{-1}\boldsymbol{\beta}) R(\underline{\mathbf{D}}^{-1}) = L(-\underline{\mathbf{D}}^{-1}\boldsymbol{\beta}, \underline{\mathbf{D}}^{-1})$$
(74)

Again this can be compared with the inverse of a semi-direct product $\mathbb{R}^3 \rtimes SO(3)$, which would be given by just the same formula.

The Thomas rotation, which we already identified in (67) as directly responsible for the non-commutativity of velocity composition, is also responsible for other remarkable properties. One of them is the fact that the inverse of a composed velocity differs from the transposed composition of the inverse velocities. In fact, the inverse of the combination $B(\beta_1)B(\beta_2)$ is obviously $B(-\beta_2)B(-\beta_1)$, the polar decomposition of which will then, contain a boost with velocity $(-\beta_2) \oplus (-\beta_1)$. The latter differs from the inverse velocity $-(\beta_1 \oplus \beta_2)$ of the original combination $B(\beta_1)B(\beta_2)$ by a Thomas rotation:

$$-(\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}) = (-\boldsymbol{\beta}_{1}) \oplus (-\boldsymbol{\beta}_{2})$$

$$= \underline{\mathbf{T}}(-\boldsymbol{\beta}_{1}, -\boldsymbol{\beta}_{2}) ((-\boldsymbol{\beta}_{2}) \oplus (-\boldsymbol{\beta}_{1}))$$

$$= \underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2}) ((-\boldsymbol{\beta}_{2}) \oplus (-\boldsymbol{\beta}_{1}))$$

$$\neq (-\boldsymbol{\beta}_{2}) \oplus (-\boldsymbol{\beta}_{1}),$$
(75)

Here we used (69a) in the first, (67) in the second, and (69b) in the third equality. This has often been considered paradoxical, following Mocanu (1986). It is known in the literature as "Mocanu Paradox" (Ungar, 1989).

Another consequence of the Thomas rotation is the failure of associativity of Einstein addition.¹⁰ To see this let us start from associativity of composition of Lorentz transformations with three pure boosts:

$$B(\boldsymbol{\beta}_1)(B(\boldsymbol{\beta}_2)B(\boldsymbol{\beta}_3)) = (B(\boldsymbol{\beta}_1)B(\boldsymbol{\beta}_2))B(\boldsymbol{\beta}_3)$$
(76)

Applying polar decomposition (43) with $\beta = \beta_1 \oplus \beta_2$ and $\underline{D} = \underline{T}(\beta_1, \beta_2)$ on on each side gives for the left-hand side

$$B(\boldsymbol{\beta}_1) B(\boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_3) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_2, \boldsymbol{\beta}_3)) = B(\boldsymbol{\beta}_1 \oplus (\boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_3)) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_3)) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_2, \boldsymbol{\beta}_3)).$$
(77a)

¹⁰ The first to explicitly note non-associativity and the rôle of Thomas rotation in it seems to have been Ungar (1989).

and, slightly more complicated, for the right-hand side

$$B(\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2})) B(\boldsymbol{\beta}_{3})$$

$$= B(\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}) B(\underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2})\boldsymbol{\beta}_{3}) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2}))$$

$$= B((\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}) \oplus \underline{\mathbf{T}}[\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2}]\boldsymbol{\beta}_{3}) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}, \underline{\mathbf{T}}[\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2})\boldsymbol{\beta}_{3})) R(\underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2})),$$
(77b)

where we also used (70b) from the first to the second line. As both sides are in polar decomposed form, boost and rotation parts must separately be equal, leading for the boosts to

$$\boldsymbol{\beta}_1 \oplus (\boldsymbol{\beta}_2 \oplus \boldsymbol{\beta}_3) = (\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2) \oplus \underline{\mathbf{T}}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2) \boldsymbol{\beta}_3, \quad (78a)$$

$$(\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2) \oplus \boldsymbol{\beta}_3 = \boldsymbol{\beta}_1 \oplus \left(\boldsymbol{\beta}_2 \oplus \underline{\mathbf{T}}(\boldsymbol{\beta}_2, \boldsymbol{\beta}_1) \boldsymbol{\beta}_3\right), \tag{78b}$$

where (78b) follows immediately from (78a) by setting $\beta'_3 := \underline{\mathbf{T}}(\beta_1, \beta_2)\beta_3$, so that, according to (68), $\beta_3 := \underline{\mathbf{T}}(\beta_2, \beta_1)\beta'_3$. Dropping the prime on β'_3 then gives (78b).

Equations (78) show explicitly how the existence of the Thomas precession obstructs associativity. Formula (78a) and (78b) are is identical with the "right weak associative law of velocity composition" and the "left weak associative law of velocity composition", respectively, stated by Ungar (1988, p. 71, expression iia,b). There it is also stated that the proof of such identities "is lengthy and, hence, requires the use of computer algebra" (Ungar, 1988, p. 72). As we have just seen, this is an exaggeration.

Another interesting and immediate consequence from (76) is obtained by specialising to $\beta_1 = \beta_3$. In this case the triple product is $B(\beta_1)B(\beta_2)B(\beta_1)$, which is symmetric and positive definite, hence already polar decomposed. Therefore the rotational part on the right-hand side of (77a) must be the identity, which leads to

$$\underline{\mathbf{T}}(\beta_1,\beta_2) = \underline{\mathbf{T}}(\beta_1,\beta_2\oplus\beta_1) = \underline{\mathbf{T}}(\beta_1\oplus\beta_2,\beta_2),$$
(79)

where we also used (68) and the second equation follows from the first by inversion and exchange the indices 1 and 2.

Even though Einstein addition fails associativity, it is remarkable that it does maintains a property that is usually implied by it. To explain this, let us first make the obvious observation that, like in ordinary vector addition, the neutral element for Einstein addition is still the zero velocity and the unique left- and right inverse of β is $(-\beta)$. Hence for all β we have

$$\boldsymbol{\beta} \oplus \boldsymbol{0} = \boldsymbol{0} \oplus \boldsymbol{\beta} = \boldsymbol{\beta} \tag{80}$$

and

$$\boldsymbol{\beta} \oplus (-\boldsymbol{\beta}) = (-\boldsymbol{\beta}) \oplus \boldsymbol{\beta} = \boldsymbol{0}.$$
(81)

Assume for a moment that associativity did hold. We could then uniquely solve an equation like

$$\boldsymbol{\beta}_1 \oplus \boldsymbol{\beta}_2 = \boldsymbol{\beta}_3 \tag{82}$$

for β_1 given β_2 and β_3 , or for β_2 given β_1 and β_3 . The way to achieve this would in the first case be to \oplus -multiply (82) from the right with $(-\beta_2)$ and then use associativity to show that the left-hand side is just β_1 whereas the right-hand side is $(\beta_3) \oplus (-\beta_2)$. Alternatively, left \oplus -multiplication with $-\beta_1$ would determine β_2 as $(-\beta_1) \oplus \beta_3$. Now, in reality, we do not have associativity and we cannot proceed in this way. But – and that is a remarkable fact – we can still write down explicit expressions solving (82) for either β_1 or β_2 . Moreover, at least for β_2 , the expression is just that we would have derived on account of associativity, as just discussed. We have

Theorem 6. The unique solutions to (82) are

$$\boldsymbol{\beta}_1 = \boldsymbol{\beta}_3 \oplus \left(-\underline{\mathbf{T}}(\boldsymbol{\beta}_3, \boldsymbol{\beta}_2)\boldsymbol{\beta}_2\right), \tag{83a}$$

$$\boldsymbol{\beta}_2 = (-\boldsymbol{\beta}_1) \oplus \boldsymbol{\beta}_3 \,. \tag{83b}$$

Proof. The proof of (83b) is just given by left \oplus -multiplication with $(-\beta_1)$. We now use formula (78) in which we replace β_1 with $(-\beta_1)$, β_2 with β_1 , and β_3 with β_2 . The Thomas term $\underline{T}(\beta_1, \beta_2)$ then turns into $\underline{T}(-\beta_1, \beta_1)$ which is the identity. Hence, in this special case, we may proceed as if associativity holds and get (83b). For the proof of (83a) we have to go a little further and start from the general relation

$$L(\boldsymbol{\beta}_1, \underline{\mathbf{D}}_1) L(\boldsymbol{\beta}_2, \underline{\mathbf{D}}_2) = L(\boldsymbol{\beta}_3, \underline{\mathbf{D}}_3), \qquad (84a)$$

which reads in terms of parameters, according to (73),

$$\boldsymbol{\beta}_3 = \boldsymbol{\beta}_1 \oplus \underline{\mathbf{D}}_1 \boldsymbol{\beta}_2, \qquad (84b)$$

$$\underline{\mathbf{D}}_3 = \mathsf{T}(\boldsymbol{\beta}_1, \underline{\mathbf{D}}_1 \boldsymbol{\beta}_2) \underline{\mathbf{D}}_1 \underline{\mathbf{D}}_2.$$
(84c)

On the group level we know how to solve (84a) for $L(\beta_1, \underline{D}_1)$ through rightmultiplication with $L^{-1}(\beta_2, \underline{D}_2)$. From (74) and (73) we also know the respective parameter expressions for inversion and multiplication. Hence we get

$$L(\boldsymbol{\beta}_{1}, \underline{\mathbf{D}}_{1}) = L(\boldsymbol{\beta}_{3}, \underline{\mathbf{D}}_{3})L(-\underline{\mathbf{D}}_{2}^{-1}\boldsymbol{\beta}_{2}, \underline{\mathbf{D}}_{2}^{-1}) = L(\boldsymbol{\beta}_{3} \oplus (-\underline{\mathbf{D}}_{3}\underline{\mathbf{D}}_{2}^{-1}\boldsymbol{\beta}_{2}), \underline{\mathbf{T}}(\boldsymbol{\beta}_{3}, -\underline{\mathbf{D}}_{2}^{-1}\boldsymbol{\beta}_{2})\underline{\mathbf{D}}_{3}\underline{\mathbf{D}}_{2}^{-1}),$$
(85a)

so that for the parameters we get

$$\boldsymbol{\beta}_1 = \boldsymbol{\beta}_3 \oplus \left(-\underline{\mathbf{D}}_3 \underline{\mathbf{D}}_2^{-1} \boldsymbol{\beta}_2\right), \tag{85b}$$

$$\underline{\mathbf{D}}_{1} = \mathsf{T}(\boldsymbol{\beta}_{3}, -\underline{\mathbf{D}}_{3}\underline{\mathbf{D}}_{2}^{-1}\boldsymbol{\beta}_{2})\underline{\mathbf{D}}_{3}\underline{\mathbf{D}}_{2}^{-1}.$$
(85c)

We note that the equations (84) and (85) are equivalent sets and valid for all $(\beta_i, \underline{\mathbf{D}}_i)$, i = 1, 2, 3. Next we observe that (82) is just (84b) for the special case in

which $\underline{\mathbf{D}}_1$ is the identity. Hence we consider (84) and (85) for $\underline{\mathbf{D}}_1$ being the identity. In that case (84c) becomes

$$\underline{\mathbf{D}}_{3}\underline{\mathbf{D}}_{2}^{-1} = \underline{\mathbf{T}}(\boldsymbol{\beta}_{1}, \boldsymbol{\beta}_{2})
= \mathbf{T}(\boldsymbol{\beta}_{1} \oplus \boldsymbol{\beta}_{2}, \boldsymbol{\beta}_{2}) \quad \text{(because of (79))}
= \mathbf{T}(\boldsymbol{\beta}_{3}, \boldsymbol{\beta}_{2}) \quad \text{(because of (84b))}.$$
(86)

Inserting this into (85b) proves (83a).

We can summarise the algebraic structure realised by \oplus on the open ball $\mathring{B}_1(\mathbb{R}^3)$ in modern terminology as follows; see also Fig. 2:

- Let M be a set and $\phi : M \times M$ a map, denoted by $(a, b) \mapsto \phi(a, b) =: a \cdot b$. Then the pair (M, ϕ) is called a *magma* (sometimes also *groupoid*). Hence $(\mathring{B}_1(\mathbb{R}^3), \oplus)$ is a magma.
- A magma (M, ϕ) is called a *semigroup* if ϕ is associative, i.e. $a \cdot (b \cdot c) = (a \cdot b) \cdot c$ for all $a, b, c \in M$. Hence $(\mathring{B}_1(\mathbb{R}^3), \oplus)$ is *not* a semigroup.
- A magma (M, φ) is called a *quasigroup* if for each pair (a, b) ∈ M × M there is a unique pair (x, y) ∈ M × M such that a ⋅ x = b and y ⋅ a = b. This property is also called *divisibility*. Theorem 6 then shows that (B₁(ℝ³), ⊕) is a quasigroup.
- A magma (M, ϕ) is said to have an *identity* if there exists an $e \in M$ such that $e \cdot a = a \cdot e = a$ for all $a \in M$. Such an e is necessarily unique (even if (M, ϕ) is not a quasigroup). As for Einstein addition **0** is an identity (compare (80)), we infer that $(\mathring{B}_1(\mathbb{R}^3), \oplus)$ is a *quasigroup with identity*.
- In a quasigroup with identity each element a has a unique left and a unique right inverse, namely that x and that y satsifying $x \cdot a = e$ and $a \cdot y = e$. x and y need not be identical. If they are, we write $x = y =: a^{-1}$. From (81) we see that $(\mathring{B}_1(\mathbb{R}^3), \oplus)$ is a *quasigroup with identity and coinciding left-right inverses*.
- A quasigroup with identity is called a *loop*. An associative loop is a group. Hence (B₁(ℝ³), ⊕) is a loop that is not a group, but is special insofar as left-and right-inverse elements coincide.



Figure 2: Hierarchy of algebraic structures. Einstein addition \oplus endows the open ball $\mathring{B}_1(\mathbb{R}^3)$ with the structure of a loop, which is just short of being a group by its failure to satisfy associativity. (Picture source: https://commons.wikimedia.org/wiki/File:Magma_to_group3.svg. Picture attribution: Tomruen, CC0, via Wikimedia Commons)

3 Velocity subtraction: the modern (geometric) story

In this section, which is the heart of this paper, we will present a geometric view of Lorentz transformations and how to decompose them as boosts and rotations. This will clarify the invariant meaning behind the Einstein law of velocity addition. All this will result in a geometrically satisfying definition of the notion of "relative velocity" between two states of motion which we will call their "link velocity", indicating that this definition rests on the so-called-called "boost link theorem" that we state and prove. The conceptually important point to keep in mind is that the link-velocity between two states of motion needs to be referred to a third state of motion s. All constructions are "geometric" insofar as all expressions involve only vectors and their scalar products.

3.1 States of motion and the non-naturalness of polar decoposition

Above we used the polar decomposition to decompose a Lorentz transformation into a spatial rotation and a boost. It is important to realise that this operation is *not* natural. It depends on a preferred state of motion which in the matrix formulation above was given by the choice of the timelike vector e_0 of the chosen basis. Any rotation $R(\underline{D})$ takes place in the corresponding "rest space" that is the orthogonal complement of e_0 . Hence, any of our rotations $R(\underline{D})$ act in a spacelike 2-plane within that "space" and pointwise fixes the timleike 2-plane orthogonal to it. The latter always contains e_0 . Similarly, any Boosts $B(\beta)$ take place in a timelike 2-plane containing e_0 and pointwise fixes the spacelike orthogonal complement. Decompositions with respect to different choices of states of motion are *a priori* incomparable, as they refer to different "spaces" between which no natural identification (isomorphism) exists.

Before we continue let us give the definition of the term "state of motion" that we just used:

Definition 7. Let

$$V_1 := \{ \nu \in V : \eta(\nu, \nu) = -1 \}$$
(87)

be the set of unit timelike vectors in V. It consists of two connected components, *i.e.*

$$V_1 := V_1^+ \cup V_1^-, \quad V_1^+ \cap V_1^- = \emptyset.$$
 (88)

If $v \in V_1^+$ then $-v \in V_1^-$. On V_1 we consider the equivalence relation $v \sim w \Leftrightarrow v = \pm w$. Elements in the quotient set

$$\mathcal{S} := V_1 / \sim \tag{89}$$

are called states of motion. They may be faithfully represented by, say, elements of V_1^+ , e.g. as follows: pick any element $s \in = V_1^+$; then

$$\mathcal{S} = \{ \nu \in V_1 : \eta(\nu, s) \le -1 \}.$$

$$\tag{90}$$

As remarked, polar decomposition ist not a natural operation on Lor. The additional structure that is needed is a state of motion or, equivalently, a *Euclidean* inner product g on V. The equivalence is seen as follows: We identify S with V_1^+ ; then any $s \in S$ defines a Euclidean inner product on (V, η) by

$$g := \eta + 2\sigma \otimes \sigma, \tag{91}$$

where $\sigma := \eta_{\downarrow}(s)$. Conversely, given V and a Lorentzian metric η as well as a Euclidean metric g, we consider the Euclidean sphere $S^3 := \{v \in V : g(v, v) = 1\}$ (which is compact) and on it the function $Q : S^3 \to \mathbb{R}, v \mapsto \eta(v, v)$. It has precisely two negative minima on S^3 corresponding to a pair of antipodal timelike vectors $\pm v \in V$ and hence, upon normalisation, determine a unique state of motion.

We now give the general definition of "polar decomposition"¹¹ for Lorentz transformations:

Definition 8. Given $L \in Lor$, its polar decomposition relative to a state of motion $s \in S$ is as follows: Let $\sigma := \eta_{\downarrow}(s)$ and g defined by means of s and η as in (91). Write L = BR where R is orthogonal with respect to g and where B is symmetric and positive definite with respect to g; that is, g(Bv, w) = g(v, Bw) for all $v, w \in V$ and g(Bv, Bv) > 0 for all $v \in V \setminus \{0\}$. For given L the factors R and B always exist, are unique, and are both again elements of Lor.

¹¹ We review the general theory of polar decomposition for finite-dimensional real vector spaces in Appendix A.

Note that the map R fixes s; in fact, it pointwise fixes a timelike subspace of V containing s and acts non-trivially (if $R \neq id_V$) on the two-dimensional spacelike plane orthogonal to it (the plane of rotation). Likewise, B pointwise fixes a two-dimensional spacelike subspace of V within the orthogonal complement of s and acts non-trivially (if $B \neq id_V$) on the two-dimensional timelike subspace of V containing s. This leads us to

Definition 9. We call $L \in \text{Lor } a$ (spatial) rotation relative to $s \in S$ *iff it pointwise fixes a two-dimensional timelike subspace of* V *containing* s. We call L a boost relative to $s \in S$ *iff it pointwise fixes a two-dimensional spacelike subspace of* V *within the orthogonal complement of* s.

Suppose $L \in Lor$ is a pure rotation relative to $s \in S$. Then it is easy to see that L is also a pure rotation relative $s' \neq s$ iff s' lies in the timelike orthogonal complement of the spacelike plane of rotation, i.e. if the timelike plane Span $\{s, s'\}$ is pointwise fixed. Likewise, if $L \in Lor$ is a pure boost relative to $s \in S$, it is also a pure boost relative $s' \neq s$ iff s' lies in the timelike plane of boost, i.e. if the spacelike orthogonal complement of Span $\{s, s'\}$ is pointwise fixed by L.

To say that the polar decomposition of Lor is "non-natural" means that it only exists relative to an additional structural input, here the choice of some $s \in S$. We will see below that given that choice the decomposition into boost and rotation can be easily formulated without ever talking about polar decomposition. This alternative formulation will be better suited for comparison of the Galilei-Newton case that we perform in the final section.

3.2 The kinematical setting

We recall that the space of states S is identified with a connected component of the spacelike hyperboloid of unit timelike vectors in (V, η) . Elements of S shall be denoted by the letter s, possibly with lower-case indices for distinction. We shall simplify notation by denoting the Minkowski inner product by a dot, i.e. $\eta(u, v) =: u \cdot v$. Further we write $u^2 := u \cdot u$ and $||u|| := \sqrt{|u^2|}$. The latter does not define a norm in V due to the indefiniteness of η , but its does on any spacelike subspace $T_s S$.

We shall identify the tangent space to S at s with the η -orthogonal complement of s in V:

$$\mathsf{T}_{\mathsf{s}}\mathcal{S} = \mathsf{R}_{\mathsf{s}} := \{ \mathsf{u} \in \mathsf{V} : \mathsf{u} \cdot \mathsf{s} = \mathsf{0} \}.$$
(92)

In order to notationally distinguish tangent vectors to S from other vectors in V we shall set them in bold.

Associated to any $s \in S$ are projection endomorphisms in V parallel (||) and

perpendicular (\perp) to s:

$$\mathbf{P}_{\mathbf{s}}^{\parallel} = -\mathbf{s} \otimes \mathbf{s}, \qquad (93a)$$

$$\mathbf{P}_{s}^{\perp} = \mathrm{id}_{\nu} + s \otimes s \,. \tag{93b}$$

As already explained in section 2.3 below equation (24), we shall identify $End(V) = V \otimes V^*$ with $V \otimes V$, so that, e.g., an element $u \otimes v \in V \otimes V$ corresponds to the endomorphism $V \ni w \mapsto (v \cdot w) u \in V$. Hence

$$\mathbf{P}_{\mathbf{s}}^{\parallel}(\mathbf{v}) = -(\mathbf{v} \cdot \mathbf{s}) \, \mathbf{s} \quad \text{and} \quad \mathbf{P}_{\mathbf{s}}^{\perp}(\mathbf{v}) = \mathbf{v} + (\mathbf{v} \cdot \mathbf{s}) \, \mathbf{s} \,. \tag{94}$$

Definition 10. *Given two states of motion* s *and* s_1 *. The* relative velocity between s_1 and s, judged from s, *is defined by*

$$\boldsymbol{\beta}(\boldsymbol{s},\boldsymbol{s}_1) := \frac{\mathbf{P}_{\boldsymbol{s}}^{\perp}(\boldsymbol{s}_1)}{\|\mathbf{P}_{\boldsymbol{s}}^{\parallel}(\boldsymbol{s}_1)\|} \in \mathsf{T}_{\boldsymbol{s}}\mathcal{S}.$$
(95)

This, in geometric terms, is just the ordinary definition of velocity (in units of c) in SR.

The apparently redundant second reference to s expressed in the phrase "judged from s" will be justified below. It turns out to be necessary because relative velocities between two states need reference to a third one, as already emphasised above, and that third one needs not be any of the given two ones.

Noting that $\|P_s^{\parallel}(s_1)\| = -(s \cdot s_1) > 0$, the expression in (95) is just

$$\beta(s, s_1) = -s - \frac{s_1}{s \cdot s_1}$$
 (96)

The squared modulus is

$$\beta^{2}(s, s_{1}) := \|\beta(s, s_{1})\|^{2} = 1 - (s \cdot s_{1})^{-2}, \qquad (97)$$

or (again taking into account that $(s \cdot s_1) < 0$)

$$\gamma(s, s_1) := -(s \cdot s_1) = \frac{1}{\sqrt{1 - \beta^2(s, s_1)}} \,. \tag{98}$$

This is just the usual "gamma-factor" associated to any relative velocity. Note that the modulus (97) is symmetric in its arguments, i.e. $\beta(s, s_1) = \beta(s_1, s)$, whereas the vectors $\beta(s, s_1) \in T_s S$ and $\beta(s_1, s) \in T_{s_1} S$ lie in different vector spaces and cannot be compared directly. In particular, a reciprocity statement, like " $\beta(s, s_1) = -\beta(s_1, s)$ ", would make no sense. We will see below how the reference to a common reference state will eventually render such a statement meaning-ful.

It follows from our earlier discussion of boost transformation that there is a unique boost relative to s transforming s to s_1 . We will denote it either by $B(s, s_1)$ or $B(s, \beta)$, with $\beta \in T_s S$ given by (96). In fact, its form can just be read off (by abstraction) from (34), now setting $\mathbf{n} := \beta/\beta$ and $\gamma = (1 - \beta^2)^{-1/2}$:

$$B(s,\beta) = id_{V} + (\gamma - 1)(-s \otimes s + \mathbf{n} \otimes \mathbf{n}) + \beta\gamma(s \otimes \mathbf{n} - \mathbf{n} \otimes s)$$
(99a)

$$= \mathbf{P}_{(\mathbf{s},\mathbf{n})}^{\perp} + \gamma \mathbf{P}_{(\mathbf{s},\mathbf{n})}^{\parallel} + \beta \gamma \, \mathbf{s} \wedge \mathbf{n} \,. \tag{99b}$$

The second line results from the first by observing that $(-s \otimes s + \mathbf{n} \otimes \mathbf{n})$ is just the η -orthogonal projector onto span $\{s, \mathbf{n}\}$ which we denoted by $P_{(s,\mathbf{n})}^{\parallel}$. Accordingly, $P_{(s,\mathbf{n})}^{\perp} := id_V - P_{(s,\mathbf{n})}^{\parallel}$. Now, any of the expressions (99) immediately implies

$$\mathsf{B}(\mathbf{s},\boldsymbol{\beta})\mathbf{s} = \boldsymbol{\gamma}(\mathbf{s}+\boldsymbol{\beta}\mathbf{n})\,,\tag{100a}$$

$$B(s,\beta)\mathbf{n} = \gamma(\mathbf{n} + \beta s), \qquad (100b)$$

$$B(s, \beta)\mathbf{v} = \mathbf{v} (\forall \mathbf{v} : \mathbf{v} \cdot \mathbf{s} = \mathbf{v} \cdot \mathbf{n} = \mathbf{0}), \qquad (100c)$$

which clearly qualifies $B(s, \beta)$ uniquely as the boost in the s-n plane with velocity $\beta = \beta n$. Replacing n according to (96) by $n = (1/\beta\gamma)(s_1 - \gamma s)$ in (99a) leads after a short calculation to

$$B(s,s_1) = id_V + \frac{s \otimes s + s_1 \otimes s_1 + s \wedge s_1 - 2\gamma s_1 \otimes s}{\gamma + 1}.$$
 (101)

Since $\gamma = -(s \cdot s_1)$, this shows that $B(s_1, s)$ is a rational function of s and s_1 . This is in contrast to (99a), which, if expressed in terms of $\beta = \beta \mathbf{n}$, still has terms proportional to γ (rather than γ^2) which is not rational in β .

Even though it is obvious from its derivation, we can verify directly that (101) is a boost in the plane spanned by s and s_1 mapping s to s_1 . In fact, expression (101) immediately leads to

$$B(s, s_1)s = s_1,$$
 (102a)

$$B(s, s_1)s_1 = -s + 2\gamma s_1, \qquad (102b)$$

$$B(s, s_1)u = u (\forall u : u \cdot s = u \cdot s_1 = 0).$$
 (102c)

From (102a,102b) we easily infer by short calculations that all three scalar products $s \cdot s = -1$, $s_1 \cdot s_1 = -1$, and $s \cdot s_1 = -\gamma$ are left invariant under B(s, s₁), which together with (102c) shows that indeed all scalar products are left invariant and that hence B(s₁, s) is the said boost.

Equations (102a,102b) also allow us to easily determine the action of $B(s, s_1)$ onto $\beta(s, s_1) \in T_s S$, which necessarily results in an element of $T_{s_1}S$. From (96)

we infer:

$$B(s, s_1)\beta(s, s_1) = B(s, s_1)\left(-s - \frac{s_1}{(s \cdot s_1)}\right)$$
$$= \left(-s_1 - \frac{-s - 2(s \cdot s_1)s_1}{s \cdot s_1}\right)$$
$$= -\left(-s_1 - \frac{s}{s_1 \cdot s}\right)$$
$$= -\beta(s_1, s),$$
(103)

where we used once more (96) in the last step. This is the right form to state a reciprocity of relative velocities at this point¹² if both refer to different reference states. Namely, whereas $\beta(s, s_1) \in T_s S$ and $\beta(s_1, s) \in T_{s_1} S$ are incomparable because they lie in different tangent spaces, $B(s, s_1)\beta(s, s_1)$ can be compared to $\beta(s_1, s)$ (both lying in $T_{s_1}S$) and likewise $B(s_1, s)\beta(s_1, s)$ can be compared to $\beta(s, s_1)$ (both lying in $T_s S$) with the result that these are, respectively, the negative of each other. Moreover, as we will show in Appendix B, the linear isometry $B(s, s_1) : T_s S \to T_{s_1} S$ is just that resulting from parallel transport along the unique geodesic within S (with respect to the metric η restricted to TS) connecting s with s_1 . Hence reciprocity can also be stated with $B(s, s_1)$ being interpreted as parallel transport along connecting geodesics.

At this point we recall that any boost transformation (99) is in the image of the exponential map. In fact

Proposition 11. Let $\beta = \beta \mathbf{n}$ and $\rho = \rho \mathbf{n}$; then

$$\exp(s \wedge \rho) = B(s, \beta), \qquad (104a)$$

where
$$\rho = \tanh^{-1}(\beta)$$
. (104b)

Proof. We start by noting that

$$(\mathbf{s} \wedge \mathbf{n})^2 = (\mathbf{s} \otimes \mathbf{n} - \mathbf{n} \otimes \mathbf{s}) \circ (\mathbf{s} \otimes \mathbf{n} - \mathbf{n} \otimes \mathbf{s}) = -\mathbf{s} \otimes \mathbf{s} + \mathbf{n} \otimes \mathbf{n}$$

= $\mathbf{P}_{(\mathbf{s},\mathbf{n})}^{\parallel}$. (105)

Hence, decomposing the exponential series into even and odd powers gives

$$\exp(\rho \, \mathbf{s} \wedge \mathbf{n}) = \sum_{k=0}^{\infty} \frac{\rho^k}{k!} (\mathbf{s} \wedge \mathbf{n})^k$$

=
$$\sum_{k=0}^{\infty} \frac{\rho^{2k}}{(2k)!} (\mathbf{s} \wedge \mathbf{n})^{2k} + \sum_{k=0}^{\infty} \frac{\rho^{2k+1}}{(2k+1)!} (\mathbf{s} \wedge \mathbf{n})^{2k+1}$$
(106)

¹² We will later state another one in which both "velocities" (more precisely: "link-velocities", see below) refer to the same reference state; see Corollary17.

In view of (105), each term in the first (even) sum is proportional to $P_{(s,n)}^{\parallel}$ except the first (k = 0), which equals $id_V = P_{(e,n)}^{\parallel} + P_{(e,n)}^{\perp}$. In the second (odd) sum, again because of (105) and also because of $P_{(s,n)}^{\parallel} \circ (s \wedge n) = (s \wedge n) \circ P_{(s,n)}^{\parallel} = s \wedge n$ we have $(s \wedge n)^{2k+1} = (s \wedge n)$ for all $k \ge 0$. Hence

$$\exp(\rho \, \mathbf{s} \wedge \mathbf{n}) = \mathbf{P}_{(\mathbf{s},\mathbf{n})}^{\perp} + \cosh(\rho) \, \mathbf{P}_{(\mathbf{s},\mathbf{n})}^{\parallel} + + \sinh(\rho) \, \mathbf{s} \wedge \mathbf{n} \,, \tag{107}$$

which is just (99b) taking into account (104b), i.e. $\gamma = 1/\sqrt{1-\beta^2} = \cosh(\rho)$ and $\beta\gamma = \sinh(\rho)$.

Finally we show how to repeat the initial matrix-calculation that led to the addition formula, now in a geometric fashion making manifest that all the velocities appearing in it refer to the same reference state s and are hence elements of the same vector space. Using (99a) and $s \cdot \beta_i = 0$ (i = 1, 2) we get

$$(B(s, \beta_1) \circ B(s, \beta_2))s = B(s, \beta_1)[\gamma_2(s + \beta_2)]$$

$$= \gamma_2 B(s, \beta_1)s + \gamma_2 B(s, \beta_1)\beta_2$$

$$= \gamma_1 \gamma_2(s + \beta_1)$$

$$+ \gamma_2 [\beta_2 + (\gamma_1 - 1)(\mathbf{n}_1 \cdot \beta_2) \mathbf{n}_1 + \gamma_1(\beta_1 \cdot \beta_2) s]$$

$$= \gamma_1 \gamma_2 (1 + \beta_1 \cdot \beta_2) s$$

$$+ \gamma_1 \gamma_2 [\beta_1 + (\mathbf{n}_1 \cdot \beta_2) \mathbf{n}_1] + \gamma_2 [\beta_2 - (\mathbf{n}_1 \cdot \beta_2) \mathbf{n}_1]$$

$$= \gamma_1 \gamma_2 (1 + \beta_1 \cdot \beta_2) \left(s + \frac{\beta_1 + \beta_2^{\parallel} + \gamma_1^{-1} \beta_2^{\perp}}{1 + \beta_1 \cdot \beta_2} \right),$$

$$(108)$$

where, as before, the superscripts \parallel and \perp refer to the projections parallel and perpendicular to \mathbf{n}_1 . This is of the form of a pure boost relative to s acting on s:

$$B(s,\beta)s = \gamma(s+\beta)$$
(109)

with

$$\gamma = \gamma_1 \gamma_2 (1 + \beta_1 \cdot \beta_2), \qquad (110a)$$

$$\beta = \frac{\beta_1 + \beta_2^{\parallel} + \gamma_1^{-1} \beta_2^{\perp}}{1 + \beta_1 \cdot \beta_2} = \beta_1 \oplus \beta_2.$$
(110b)

Note that we only calculated the action of $B(\beta_1, s)B(\beta_2, s)$ on s, not the map as such, which also contains the Thomas rotation in the spacelike plane perpendicular to s and which hence acts as the identity on s. This is why the Thomas rotation does not appear here.

As a final remark in this subsection we wish to point out an interesting relation with hyperbolic geometry. We fix a reference state s and set $s_i := B(s, \beta_i)s$ for i = 1, 2. We consider the open unit ball in the tangent space of S at s,

$$\mathcal{U} := \{ \mathbf{v} \in \mathsf{T}_{s} \mathcal{S} : \mathbf{v} \cdot \mathbf{v} < 1 \}, \tag{111}$$

in which the velocities relative to s take their values. On \mathcal{U} we define the hyperbolic distance function $d : \mathcal{U} \times \mathcal{U} \to \mathbb{R}_{\geq 0}$ which assigns to each pair $\beta_{1,2} \in \mathcal{U}$ the Riemannian geodesic distance¹³ between $s_{1,2} = \gamma_{1,2}(s + \beta_{1,2})$. In this way (\mathcal{U}, d) becomes isometric to (\mathcal{S}, d_h) , where d_h is the distance function induced by the Riemannian metric h on \mathcal{S} . Explicitly, the distance function d is given by

$$d(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2) := \arccos\left(-s_1 \cdot s_2\right) = \arccos\left(\gamma_1 \gamma_2 (1 - \boldsymbol{\beta}_1 \cdot \boldsymbol{\beta}_2)\right). \tag{112}$$

Now, (108-110) show that for any boost $B(s, \beta)$ relative to s we have

$$s'_{i} := B(s, \beta)s_{i} = B(s, \beta \oplus \beta_{i})s$$
(113)

But as boosts preserve scalar products we have $s'_1 \cdot s'_2 = s_1 \cdot s_2$, which implies

$$\mathbf{d}(\boldsymbol{\beta} \oplus \boldsymbol{\beta}_1, \, \boldsymbol{\beta} \oplus \boldsymbol{\beta}_2) = \mathbf{d}(\boldsymbol{\beta}_1, \boldsymbol{\beta}_2), \quad (114)$$

for any triple $(\beta, \beta_1, \beta_2)$ of points in \mathcal{U} . This equations says that, with respect to d, velocity addition \oplus defines an isometric "action" of \mathcal{U} on itself. The origin of this lies of course in the proper isometric action of the Lorentz group on (\mathcal{S}, h) . One may ask to what extent relativistic addition is characterised by (114). In other words: what is the most general map $f: \mathcal{U} \times \mathcal{U} \to \mathcal{U}$ that satisfies

$$d(f(\boldsymbol{\beta},\boldsymbol{\beta}_1),f(\boldsymbol{\beta},\boldsymbol{\beta}_2)) = d(\boldsymbol{\beta}_1,\boldsymbol{\beta}_2), \qquad (115)$$

for any triple $(\beta, \beta_1, \beta_2)$ of points in \mathcal{U} ? This has been solved by Benz (2000, Theorem 2) who proved that f equals \oplus iff f satisfies the following two conditions: 1) $\|\mathbf{f}(\beta, \mathbf{0})\| = \|\boldsymbol{\beta}\|$ and 2) $\gamma(\mathbf{f}(\beta_1, \beta_2)) \mathbf{f}(\beta_1, \beta_2) - \gamma(\beta_2)\beta_2 = c\beta_1$, where c > 0. The first seems obvious and the second acquires a straightforward physical meaning if we recall that the momentum relative to s of a particle at velocity $\boldsymbol{\beta} \in$ $T_s S$ is proportional to $\gamma(\beta)\beta$. The second condition then just says that boosting a particle with boost-velocity $\boldsymbol{\beta}$ adds to it a momentum positively proportional to $\boldsymbol{\beta}$ (all with reference to s). Using this momentum-interpretation, the two conditions may be replaced by others that also include the non-relativistic case (Benz, 2002).

3.3 Defining link velocity

Given three states of motion, s, s_1 , and s_2 . It turns out that there exists a unique boost $B(\beta, s)$ relative to s that maps s_1 to s_2 :

$$B(s,\beta)s_1 = s_2. \tag{116}$$

¹³ Equivalently: hyperbolic angle or rapidity.

This is the affirmative answer to the so-called "boost-link-problem" that we shall prove in an elementary fashion in the next subsection. More precisely, the results of the next subsection can be summarised as follows:

Theorem 12. For any three given states of motion (s, s_1, s_2) there exists a unique $\beta \in T_s S$, given by a rational function $\beta = \beta(s, s_1, s_2)$ of (s, s_1, s_2) , satisfying (116) (see (130a) below). The boost $B(s, s_1, s_2) := B(s, \beta(s, s_1, s_2))$ is then also a rational function of (s, s_1, s_2) (see (134) below). The function β is Lorentz equivariant; i.e. for any S-preserving (no time reversal) Lorentz transformation L we have

$$\beta(Ls, Ls_1, Ls_2) = L\beta(s, s_1, s_2).$$
(117)

As an immediate consequence we note that for special s we have already encountered the expression for the boost linking s_1 and s_2 :

Corollary 13. Let (s, s_1, s_2) and (s', s_1, s_2) be two triple of states where s as well as s' lie in the plane Span $\{s_1, s_2\}$. Then the boosts $B(s, \beta)$ and $B(s', \beta')$ which according to Theorem 12 satisfy (116) are identical and given by

$$B(s,\beta) = B(s',\beta') = id_V + \frac{s_1 \otimes s_1 + s_2 \otimes s_2 + s_1 \wedge s_2 - 2\gamma_{12}s_2 \otimes s_1}{1 + \gamma_{12}}, (118)$$

where $\gamma_{12} := -(\mathbf{s}_1 \cdot \mathbf{s}_2)$. Note that the right-hand side of (118) is independent of s and given by the expression (101) with arguments (s, s_1) changed to (s_1, s_2) . Hence all boosts linking s_1 and s_2 coincide as long as their reference state s lies in the plane spanned by s_1 and s_2 .

Proof. We already know that expression (101) with arguments (s, s_1) changed to (s_1, s_2) gives the unique boost in the plane Span $\{s_1, s_2\}$ mapping s_1 to s_2 . But that plane contains s and hence is also a boost relative to s. Uniqueness then implies the statement.

Based on Theorem 12 we now make the following

Definition 14. Given three states of motion s, s_1 , and s_2 . The link-velocity between s_1 and s_2 relative to s is defined as that (unique!) $\beta \in T_s S$ solving (116). We will speak of it as "the velocity of s_2 against s_1 relative to s" or "the velocity of s_2 relative to s_1 judged from s", or similar, so as to in any case avoid a double appearance of the word "relative".

In view of Theorem 12 we can characterise the link-velocity in the following way:

Theorem 15. The link-velocity is a function

$$\begin{aligned} \boldsymbol{\beta} : \boldsymbol{\mathcal{S}} \times \boldsymbol{\mathcal{S}} \to \boldsymbol{\Gamma} \boldsymbol{\Gamma} \boldsymbol{\mathcal{S}} \\ (s_1, s_2) &\mapsto \boldsymbol{\beta} [s_1, s_2] \end{aligned}$$
(119)

that maps any ordered pair (s_1, s_2) of states to a section $\beta[s_1, s_2]$ in the tangentbundle of S. That section is such that the value of $\beta[s_1, s_2]$ at $s \in S$ equals $\beta(s, s_1, s_2)$ as defined in Theorem 12, the explicit expression of which is given below in (130a).

Remark 16. The condition of equivariance (117) is equivalent to the statement that the map (119) is invariant under Lor. Indeed, Lor acts on the domain $S \times S$ by taking the cartesian-product of its natural action on S:

$$\begin{aligned} \Phi : \mathrm{Lor} \times (\mathcal{S} \times \mathcal{S}) &\to (\mathcal{S} \times \mathcal{S}) \\ \left(\mathsf{L}, (\mathsf{s}_1, \mathsf{s}_2) \right) &\mapsto \Phi_{\mathsf{L}}(\mathsf{s}_1, \mathsf{s}_2) := (\mathsf{Ls}_1, \mathsf{Ls}_2) \,. \end{aligned}$$
 (120a)

Moreover, Lor also acts on the co-domain (the target) ΓTS as follows:

$$\begin{split} \Psi : \operatorname{Lor} \times \Gamma T \mathcal{S} &\to \Gamma T \mathcal{S} \\ (L, \sigma) &\mapsto \Psi_{L}(\sigma) := L \circ \sigma \circ L^{-1} \,. \end{split} \tag{120b}$$

Note that L acts directly on the image of σ via its defining representation on V, since according to (92) we identified $T_s S$ with $R_s := \{u \in V : u \cdot s = 0\}$. Restricting the defining action of L on V to $R_s \subset V$ results in an isometry between $R_s S$ and $R_{Ls}S$, i.e. between $T_s S$ and $T_{Ls}S$. Now, by standard construction (already applied in (120b)), actions on domains and codomains always combine to an action on the set of maps. Applied to the map β in (119) with actions (120a) on the domain and (120b) on the co-domain, we get an action of Lor on the set of maps $S \times S \to \Gamma T S$, given by

$$\mathsf{T}_{\mathsf{L}}(\boldsymbol{\beta}) := \Psi_{\mathsf{L}} \circ \boldsymbol{\beta} \circ \Phi_{\mathsf{L}}^{-1} \,. \tag{121}$$

Equation (117) is then equivalent to the statement that the map (119) is Lorinvariant:

$$\mathsf{T}_{\mathsf{L}}(\boldsymbol{\beta}) = \boldsymbol{\beta} \quad (\forall \, \mathsf{L} \in \mathrm{Lor}) \,. \tag{122}$$

The fact that the link-velocity $\beta(s, s_1, s_2)$ is a ternary has been discussed before by a particular school following Oziewicz (2006, 2007, 2011); Oziewicz and Page (2011), who interpret this fact as an "astonishing conflict of the Lorentz group with relativity" (Oziewicz, 2011); see also (Celakoska, 2008; Celakoska, Chakmakov and Petrushevski, 2015; Koczan, 2023). But (122) shows that this is an unwarranted complaint.

3.4 Solving the boost-link problem

In this section we shall prove and elaborate on Theorem 12, using the following notation: The element $\beta \in T_s S$ that we wish (116) to solve for is again written as $\beta = \beta \mathbf{n}$ with $\mathbf{n} \in T_s S$ a unit vector. The norm β of β can equivalently be expresses

as usual by $\gamma := 1/\sqrt{1-\beta^2}$, i.e. $\beta = \sqrt{1-\gamma^{-2}}$. Next to that we define the other "gammas" via the three possible scalar products between the {s, s₁, s₂}:

$$\gamma_1 := -s \cdot s_1 \quad \Rightarrow \quad \beta_1 := \sqrt{1 - \gamma_1^{-2}}, \quad (123a)$$

$$\gamma_2 := -\mathbf{s} \cdot \mathbf{s}_2 \quad \Rightarrow \quad \beta_2 := \sqrt{1 - \gamma_2^{-2}}, \tag{123b}$$

$$\gamma_{12} := -\mathbf{s}_1 \cdot \mathbf{s}_2 \quad \Rightarrow \quad \beta_{12} := \sqrt{1 - \gamma_{12}^{-2}} \,. \tag{123c}$$

In what follows (s, s_1, s_2) and hence $(\gamma_1, \gamma_2, \gamma_{12})$ and $(\beta_1, \beta_2, \beta_{12})$ are considered given, whereas (β, \mathbf{n}) , or equivalently (γ, \mathbf{n}) , are to be determined as functions of the former. This is the task to which we now turn.

Inserting the expression (99a) for $B(\beta, s)$ into (116) leds to

$$s_{2} - s_{1} = s \left[(\gamma - 1)\gamma_{1} + \beta\gamma(\mathbf{n} \cdot s_{1}) \right] + \mathbf{n} \left[(\gamma - 1)(\mathbf{n} \cdot s_{1}) + \beta\gamma\gamma_{1} \right].$$
(124)

The right-hand side gives the components of $(s_2 - s_1)$ parallel and perpendicular to s. The parallel component of the left-hand side is $P_s^{\parallel}(s_2 - s_1) = s(\gamma_2 - \gamma_1)$. Equating this to the s-term of the right-hand side allows to express $(\mathbf{n} \cdot s_1)$ as follows:

$$\mathbf{n} \cdot \mathbf{s}_1 = \frac{\gamma_2 - \gamma \gamma_1}{\beta \gamma} \,. \tag{125}$$

The perpendicular component of the left-hand side is $P_s^{\perp}(s_2 - s_1)$ which we equate to the **n**-term on the right-hand side. In the latter we replace $\mathbf{n} \cdot s_1$ with the expression just found in (125). This leads, after a short calculation, to

$$\mathbf{n} = \sqrt{\frac{\gamma+1}{\gamma-1}} \frac{\mathbf{P}_{s}^{\perp}(s_{2}-s_{1})}{\gamma_{1}+\gamma_{2}}.$$
(126)

This is not yet the solution since an unknown, γ , still appears on the right-hand side. But we can determine γ by using the fact that **n** has unit norm. Recalling that

$$\mathbf{P}_{\mathbf{s}}^{\perp}(\mathbf{s}_{2}-\mathbf{s}_{1}) \,=\, \mathbf{s}_{2}-\mathbf{s}_{1}-(\gamma_{2}-\gamma_{1})\,\mathbf{s}\,, \tag{127a}$$

$$\|P_s^{\perp}(s_2 - s_1)\|^2 = \gamma_1^2 + \gamma_2^2 + 2(\gamma_{12} - \gamma_1\gamma_2 - 1), \qquad (127b)$$

we can take the square of (126) and solve the ensuing equation for γ , which gives

$$\gamma = \gamma(s, s_1, s_2) := \frac{\gamma_1^2 + \gamma_2^2 + \gamma_{12} - 1}{1 + 2\gamma_1\gamma_2 - \gamma_{12}},$$
(128)

where we think of the right-hand side as rational function in the scalar products (123).

Note that now all terns of the right-hands side of (128), and hence also on the right-hand side of (126), are expressed in terms of given quantities. Hence we succeeded in proving existence and uniqueness of the solution for the boost-link-problem and also justified Definition 14.

In the sequel we shall use the abbreviation

$$(\mathbf{s_2} - \mathbf{s_1})_{\perp} := \mathbf{P}_{\mathbf{s}}^{\perp}(\mathbf{s_2} - \mathbf{s_1})$$
 (129)

which we write in bold so stress that this is an element in T_sS . It is just a simple linear combination of s, s_1 , and s_2 as shown in (127a), but it will be notationally more compact and also geometrically more transparent to write $(s_2 - s_1)_{\perp}$. Now, multiplying (126) with $\beta = \gamma^{-1}\sqrt{\gamma^2 - 1}$ we get, using (128),

$$\boldsymbol{\beta} = \boldsymbol{\beta}(s, s_1, s_2) := \frac{\gamma_1 + \gamma_2}{\gamma_1^2 + \gamma_2^2 + \gamma_{12} - 1} (s_2 - s_1)_{\perp}, \qquad (130a)$$

$$\gamma \beta = \gamma(s, s_1, s_2) \beta(s, s_1, s_2) = \frac{\gamma_1 + \gamma_2}{1 + 2\gamma_1 \gamma_2 - \gamma_{12}} (s_2 - s_1)_{\perp}.$$
 (130b)

Note that in view of (127a) the right-hand sides are linear combinations of s, s_1 , and s_2 with coefficients that are rational functions in the scalar products between these states. This is what we mean by saying that (130a) is itself a rational function of (s, s_1, s_2) . It is the first such function mentioned in Theorem 12. The equivariance condition (117) is obvious from these remarks.

We also note that the fraction of the right-hand side of (130a) is a symmetric function in (s_1, s_2) , whereas $(s_2 - s_1)_{\perp}$ is clearly antisymmetric. Denoting the right-hand side of (130a) with $\beta(s_2, s_1; s)$, we have

Corollary 17. Link-velocities obey the reciprocity relation

$$\beta(s, s_2, s_1) = -\beta(s, s_1, s_2).$$
(131)

Note that (131) makes sense since both sides refer to the same reference-state s, i.e. both sides are elements of the same vector space $T_s S$.

We now compute the boost as function of (s, s_1, s_2) , i.e. we insert the expressions (126) for **n**, (128) for γ , and (130b) for $\gamma\beta$ into (99a) and obtain

$$B(s, s_1, s_2) := B(s, \beta(s, s_1, s_2))$$

= id_V + (\gamma - 1)(-s \otimes s + **n** \otimes **n**) + s \lapha \gamma \beta . (132)

The various terms simplify as follows: From (128) we get

$$-(\gamma - 1)s \otimes s = -\frac{(\gamma_1 - \gamma_2)^2 + 2(\gamma_{12} - 1)}{1 + 2\gamma_1\gamma_2 - \gamma_{12}} s \otimes s.$$
(133a)

From (126) and again (128)

$$(\gamma - 1)\mathbf{n} \otimes \mathbf{n} = \frac{\gamma + 1}{(\gamma_1 + \gamma_2)^2} (\mathbf{s}_2 - \mathbf{s}_1)_\perp \otimes (\mathbf{s}_2 - \mathbf{s}_1)_\perp$$

=
$$\frac{(\mathbf{s}_2 - \mathbf{s}_1)_\perp \otimes (\mathbf{s}_2 - \mathbf{s}_1)_\perp}{1 + 2\gamma_1\gamma_2 - \gamma_{12}}.$$
 (133b)

And from (130b)

$$\mathbf{s} \wedge \boldsymbol{\gamma} \boldsymbol{\beta} = \frac{\gamma_1 + \gamma_2}{1 + 2\gamma_1 \gamma_2 - \gamma_{12}} \, \mathbf{s} \wedge (\mathbf{s_2} - \mathbf{s_1})_{\perp} \,. \tag{133c}$$

Using (127a), the sum of (133b) and (133c) is, abbreviating $s_{21} := (s_2 - s_1)$,

$$(\gamma - 1)\mathbf{n} \otimes \mathbf{n} + \mathbf{s} \wedge \gamma \boldsymbol{\beta} = \frac{(\gamma_1 - \gamma_2)^2 \mathbf{s} \otimes \mathbf{s} + \mathbf{s}_{21} \otimes \mathbf{s}_{21} + 2\gamma_1 \mathbf{s} \otimes \mathbf{s}_{21} - 2\gamma_2 \mathbf{s}_{21} \otimes \mathbf{s}}{1 + 2\gamma_1 \gamma_2 - \gamma_{12}}$$
(133d)

Finally, adding this to (133a) we obtain for (132)

$$B(s, s_1, s_2) = id_V + \frac{2(1 - \gamma_{12}) s \otimes s + s_{21} \otimes s_{21} + 2\gamma_1 s \otimes s_{21} - 2\gamma_2 s_{21} \otimes s}{1 + 2\gamma_1 \gamma_2 - \gamma_{12}}.$$
 (134)

This is the second rational function mentioned in Theorem 12.

A few easy checks reassure us that (134) is indeed the right expression. First of all, the tensor structure of the right-hand side makes it immediately evident that $B(s_2, s_1; s)s_1 = s_2$ maps the 2-dimensional timelike plane Span{ s, s_{12} } to itself and fixes points in the 2-dimensional spacelike orthogonal complement. Second, a simple computation gives $B(s_2, s_1; s)s_1 = s_2$ and, third, a another simple computation for the special case where $s = s_1$ (hence $\gamma_1 = 1$ and $\gamma_2 = \gamma_{12}$) turns expression (134) into (118).

Finally we mention an alternative way to derive (134), based on the Cartan-Dieudonné theorem¹⁴, according to which we can write the linking boost by the composition of two reflections; see (Urbantke, 2003).

3.5 Base-point dependence of the link-velocity

We have just seen that the boost relative to s that links s_1 with s_2 takes place in the plane spanned by s and $(s_2 - s_1)$, the velocity being given by (130a). For fixed s_1 and s_2 the velocities $\beta(s, s_1, s_2)$ vary with s in a twofold way. First, their "directions" differ in the sense that they are elements on the tangent spaces $T_s S$ depending on s. Second, the magnitude also depends on s in an interesting way

¹⁴ Let (V, η) be an n-dimensional real vector space with non-degenerate symmetric bilinear form $\eta \in V^* \otimes V^*$. Let $O(V, \eta)$ be the corresponding orthogonal group, defined as in (13). The *Cartan-Dieudonné theorem* states that any $L \in O(V, \eta)$ is the composition of at most n reflections at non-null (non-degenerate) hyperplanes. We recall that if $u \in V$ is non-null, i.e. $\eta(u, u) \neq 0$, so that we may assume $\eta(u, u) = \varepsilon \in \{1, -1\}$, the reflection $\rho_u \in O(V, \eta)$ at the hyperplane $u^{\perp} \subset V$ perpendicular to u is defined by $\rho_u(v) := v - 2\varepsilon\eta(v, u) u$. We refer to (Jacobson, 1985, Chap. 6.6) for a proof of the general Cartan-Dieudonné theorem. If one relaxes the upper bound on the number of reflections from n to 2n - 1, the proof becomes much easier; see, e.g., (Giulini, 2006, p. 101, Theorem 6).

that we now wish to elaborate on. We do this by studying $\gamma(s_2, , s_1; s)$ as a function of s.

From Corollary 13 we know that $B(s, s_1, s_2)$, and hence $\gamma(s, s_1, s_2)$ does not depend on s as long as $s \in \text{Span}\{s_1, s_2\}$. In that case we obviously have $\gamma = \gamma_{12}$. This suggests to compute the deviation of γ from γ_{12} in dependence of the amount by which s "sticks out" of $\text{Span}\{s_1, s_2\}$; that is, by the norm of the η -orthogonal projection into the orthogonal complement of $\text{Span}\{s_1, s_2\}$. In order to determine the latter, we note

Lemma 18. Let $s_1, s_2 \in S$; the η -orthogonal projector into Span $\{s_1, s_2\}$ is

$$P_{(s_1,s_2)}^{\parallel} = \frac{s_1 \otimes s_1 + s_2 \otimes s_2 - \gamma_{12}(s_1 \otimes s_2 + s_2 \otimes s_1)}{\gamma_{12}^2 - 1} \,. \tag{135}$$

Proof. We note that $P_{(s_1,s_2)}^{\parallel}$ is a linear combination of tensor products of elements taken from the set $\{s_1, s_2\}$ which is symmetric under exchange of s_1 and s_2 . Hence it is a η -symmetric linear map whose kernel contains the η -orthogonal complement of Span $\{s_1, s_2\}$. An easy calculation shows $P_{(s_1,s_2)}^{\parallel}s_1 = s_1$, hence also $P_{(s_1,s_2)}^{\parallel}s_2 = s_2$ (symmetry) and therefore $P_{(s_1,s_2)}^{\parallel} \circ P_{(s_1,s_2)}^{\parallel} = P_{(s_1,s_2)}^{\parallel}$.

In passing we note that expression (135) immediately leads to

$$s_{\parallel} := \mathbf{P}_{(s_1, s_2)}^{\parallel} \mathbf{s} = \frac{(\gamma_{12}\gamma_2 - \gamma_1)\mathbf{s}_1 + (\gamma_{12}\gamma_1 - \gamma_2)\mathbf{s}_2}{\gamma_{12}^2 - 1} \,. \tag{136}$$

Moreover, the η -orthogonal projector into the η -orthogonal complement of Span $\{s_1, s_2\}$ is

$$\mathbf{P}_{(s_1,s_2)}^{\perp} = \mathrm{id}_{\mathbf{V}} - \mathbf{P}_{(s_1,s_2)}^{\parallel}, \qquad (137)$$

so that the squared norm of the projection

$$\mathbf{s}_{\perp} := \mathbf{P}_{(\mathbf{s}_1, \mathbf{s}_2)}^{\perp} \mathbf{s} \tag{138}$$

becomes

$$|s_{\perp}||^{2} = s_{\perp} \cdot s_{\perp} = s \cdot s_{\perp} = s \cdot s - s \cdot s_{\parallel}$$

= $\frac{1 - \gamma_{1}^{2} - \gamma_{2}^{2} - \gamma_{12}^{2} + 2\gamma_{1}\gamma_{2}\gamma_{12}}{\gamma_{12}^{2} - 1}$, (139)

where we used (136) in the last step. We rewrite this as

$$\gamma_1^2 + \gamma_2^2 - 1 = -\|s_{\perp}\|^2 (\gamma_{12}^2 - 1) - \gamma_{12}^2 + 2\gamma_{12}\gamma_1\gamma_2, \qquad (140)$$

which we use to eliminate $\gamma_1^2 + \gamma_2^2 - 1$ in the denominator on the right-hand side of (128). This leads to the desired final formula

$$\gamma = \gamma_{12} - \|\mathbf{s}_{\perp}\|^2 \frac{\gamma_{12}^2 - 1}{1 + 2\gamma_1\gamma_2 - \gamma_{12}}.$$
(141)

We note that the fraction on the right-hand side is non-negative and zero iff $\gamma_{12} = 1$, i.e. iff $s_1 = s_2$. Indeed, writing

$$s_1 = \gamma_1(s + \beta_1)$$
 and $s_2 = \gamma_2(s + \beta_2)$, (142)

with $\beta_{1,2} \in T_s S$, we have $\gamma_{12} = -s_1 \cdot s_2 = \gamma_1 \gamma_2 (1 - \beta_1 \cdot \beta_2)$ and hence

$$1 + 2\gamma_1\gamma_2 - \gamma_{12} = 1 + \gamma_1\gamma_2(1 + \beta_1 \cdot \beta_2) > 1.$$
 (143)

Equation (141) therefore shows that $\gamma \leq \gamma_{12}$ with equality iff $s_{\perp} = 0$, i.e. $s \in \text{span}\{s_1, s_2\}$.

A specific example may be helpful to develop some intuition of how γ may deviate from γ_{12} . For that we assume that s is tilted symmetrically against s₁ and s₂, i.e that

$$\gamma_1 = \gamma_2 \eqqcolon \gamma_* \,. \tag{144}$$

In that case (128) immediately leads to

$$\gamma = \frac{2\gamma_*^2 + \gamma_{12} - 1}{2\gamma_*^2 - \gamma_{12} + 1} \,. \tag{145}$$

Note that from (143) we already know that the denominator is > 1. For fixed γ_{12} this gives γ as monotonically decreasing function of γ_* that asymptotically approaches $\gamma = 1$ (and hence the corresponding $\beta = \sqrt{1 - \gamma^{-2}}$ approaching $\beta = 0$) as γ_* tends to infinity; the graph is shown to the left in Fig. 3. γ assumes its maximum for the minimal γ_* , which is that where $s \in \text{Span}\{s_1, s_2\}$ and due to $\gamma_1 = \gamma_2$ just bisects s_1 and s_2 . Hence $s = (s_1 + s_2)/\sqrt{2(1 + \gamma_{12})}$ and the minimal γ_* is $\gamma_* = -s \cdot s_1 = \sqrt{(1 + \gamma_{12})/2}$ at which γ takes its maximal value $\gamma = \gamma_{12}$.



Figure 3: Graphs of $\gamma(\gamma_*)$ and $\gamma(\phi)$ showing how $\gamma(s, s_1, s_2)$ decreases for increasing "tilt" of s against Span{ s_1, s_2 }.

Another way to parametrise γ is to use (142) where $\beta_1 = \beta_2 =: \beta_*$ and φ the angle between β_1 and β_2 . We have

$$\gamma_{12} = -s_1 \cdot s_2 = \gamma_*^2 (1 - \beta_*^2 \cos \varphi) = \gamma_*^2 (1 - \cos \varphi) + \cos \varphi \,. \tag{146}$$

using $\beta_*^2 = (1-\gamma_*^{-2})$ in the last step. Solving this for γ_*^2

$$\gamma_*^2 = \frac{\gamma_{12} - \cos \varphi}{1 - \cos \varphi} \tag{147}$$

we can use it to eliminate γ_*^2 in (145) in favour of φ :

$$\gamma = \frac{\gamma_{12} (3 - \cos \varphi) - \cos \varphi - 1}{\gamma_{12} (1 + \cos \varphi) - 3 \cos \varphi + 1}.$$
 (148)

Now γ is a strictly monotonically increasing function in $\varphi \in [0, \pi]$ connecting the minimum at $(\varphi = 0, \gamma = 1)$ with the maximum at $(\varphi = \pi, \gamma = \gamma_{12})$. The graph is shown to the right in Fig. 3.

4 Comparison with Galilei-Newton spacetimes

It is interesting to compare the geometric developments above to the corresponding one in for Galilei-Newton spacetime. For that we start with a summary of the decomposition of a Lorentz transformation into a boost and a rotation, without using the notion of a polar decomposition, which is really not essential here.

4.1 Boost-Rotation decomposition in SR rephrased

Given $L \in Lor$ we do the following:

- 1. Choose a state $s \in S$ and let $s_1 := Ls$.
- Let B := B(s, s₁) as in (101), i.e. the unique Lorentz transformation that maps s to s₁ and pointwise fixes the η-orthogonal complement to Span{s, s₁}. The corresponding β(s, s₁) is then given by (96).
- 3. Define

$$\mathbf{R} := \mathbf{B}^{-1} \circ \mathbf{L} \,. \tag{149}$$

Clearly, R is again a Lorentz transformation that fixes s: Rs = s. Hence it is an element of the stabiliser subgroup

$$\operatorname{Stab}_{s}(\operatorname{Lor}) := \left\{ L \in \operatorname{Lor} : Ls = s \right\}, \tag{150}$$

which consists of Lorentz transformations that map the orthogonal complement to s (which is a 3-dimensional vector space with Euclidean inner product induced from η) isometrically into itself. Hence each Stab_s(Lor) is isomorphic to SO(3), but for $s \neq s'$ the corresponding stabiliser subgroups differ. In fact, they are conjugate subgroups in Lor,

$$\operatorname{Stab}_{s'}(\operatorname{Lor}) = \operatorname{B}(s, s') \circ \operatorname{Stab}_{s}(\operatorname{Lor}) \circ [\operatorname{B}(s, s')]^{-1}, \quad (151)$$

where B(s, s') is as in (101) for $s_1 = s'$.

4. Rewrite (149) as

$$\mathbf{L} = \mathbf{B} \circ \mathbf{R} \tag{152}$$

with both factors, B and R, depending on s. The s-dependent decomposition (152) may now be read as a polar decomposition with respect to the s-dependent Euclidean metric $g = \eta + 2\sigma \otimes \sigma$, where $\sigma := \eta_{\downarrow}(s)$, but that interpretation may be regarded as redundant.

In Section 4.4 below will describe the analogous procedure in the Galilei-Newton setting, where we will also find a decomposition like (152). There, in contrast, the factor B will *not* depend on the choice of s whereas the situation for the "rotational" factor R is just as in the SR case. But now this factorisation has no obvious interpretation as polar decomposition.

4.2 Galilei-Newton spacetime

Like Minkowski spacetime, Galilei-Newton spacetime is an affine space (M, V, +).¹⁵ However, their geometric structures differ. In case of Minkowski spacetime we had $(\eta, \uparrow) =$ (spacetime metric, time-orientation), which for Galilei-Newton spacetime is replaced by a pair (τ, h) . Here $\tau \in V^*$ is an oriented time-difference function and $h \in [\text{Ker}(\tau)]^* \otimes [\text{Ker}(\tau)]^*$ is a symmetric, positive definite inner product on the Ker (τ) . We will explain these structures in turn.

τ is an oriented time metric, i.e. it allows to assign an oriented time difference to any ordered pair (p, q) of spacetime points, given by τ(p − q). It can assume positive as well as negative values – hence "oriented". A vector v ∈ V is called future-pointing if τ(v) > 0 and past-pointing iff τ(v) < 0. Correspondingly, p ∈ M is called to the future or past of q ∈ M if (p − q) is future-pointing or past-pointing, respectively. Two events p, q are called simultaneous iff their time difference vanishes, τ(p − q) = 0. Simultaneity defines an absolute¹⁶ equivalence relation given by: p ~ q ⇔ (p − q) ∈ Ker(τ). The equivalence class [p] of a point p is then simply given by the 3-dimensional affine hyperplane [p] = p+Ker(τ). For the sake of notational ease we shall write

$$V_0 := \operatorname{Ker}(\tau) = \left\{ \nu \in V : \tau(\nu) = 0 \right\}.$$
(153)

Its associated dual space will be called V_0^* . An overall orientation of V will induce a orientation of V_0 in view of τ .¹⁷

¹⁵ For readers unfamiliar with this notation, we explain the meaning of an affine space characterised by the triple (M, V, +) in Appendix D. Suffice it to say here that M is a set (of events) and V a real vector space that acts simply transitively by an action called "+" on M; i.e. $M \times V \ni (m, v) \mapsto$ $m + v \in M$.

¹⁶ i.e. invariant under the automorphism group of spacetime; compare (Giulini, 2001).

¹⁷ A basis $\{e_1, e_2, e_3\}$ of V₀ is positively oriented iff its completion $\{e_0, e_1, e_2, e_3\}$ to a basis of V with $\tau(e_0) > 0$ is positively oriented in V.

h ∈ V₀^{*} ⊗ V₀^{*} is a positive definite symmetric bilinear form, i.e. a Euclidean metric, on V₀. It defines a proper distance function on any equivalence class of mutually simultaneous events through ||p−q|| := √h(p−q, p−q). No spatial distance is associated to non-simultaneous events.

In analogy to Definition 7 we now define the "unit timelike vectors" V_1 , the "future-oriented unit timelike vectors" V_1^+ , and the "states of motion" as follows:

$$V_1 := \{ \nu \in V : |\tau(\nu)| = 1 \},$$
(154a)

$$V_1^+ := \{ \nu \in V : \tau(\nu) = 1 \}.$$
(154b)

Definition 19. *The set* S *of* states of motion *is identified with* V_1^+ :

$$\mathcal{S} := V_1^+ \,. \tag{155}$$

It is a 3-dimensional real affine space over the Euclidean vector space (V_0, h) . At the same time it can also be regarded as a 3-dimensional Riemannian manifold with a flat Riemannian metric h and hence a notion of global parallelism.

Whereas there clearly is a natural embedding

$$i: V_0 \hookrightarrow V, \tag{156}$$

there is no *naturally* given projection map $V \to V_0$. The selection of such a map is equivalent to picking an element $s \in S$. The corresponding projections from V onto Span{s} and onto V₀ are then, respectively, given by¹⁸

$$\mathbf{P}_{\mathbf{s}}^{\parallel} = \mathbf{s} \otimes \boldsymbol{\tau}, \qquad (157a)$$

$$\mathbf{P}_{\mathbf{s}}^{\top} = \mathrm{id}_{\mathbf{V}} - \mathbf{s} \otimes \boldsymbol{\tau} \,. \tag{157b}$$

These should be compared to equations (93). An important difference is that all P_s^{\top} in (157b) project onto the same vector space V_0 , independent of the state s. In contrast, P_s^{\perp} in (93b) projects onto the η -orthogonal complement of s, which clearly does depend on s. This is the reason why we made the notational change from P_s^{\perp} in (93b) to P_s^{\top} in (157b): V_0 here is "transversal" (denoted by \top) to Span{s} but not in any defined sense "orthogonal" (as \perp would suggest).

In analogy to Definition 10, the relative velocity between $s_1 \in S$ and $s \in S$ is now defined as in (95), except that the length of vectors in Span{s} is measured by τ , which is always 1:

¹⁸ In contrast to section 3.2, where due to the existence of a non-degenerate bilinear form η we could identify End(V) with V \otimes V in order to simplify notation, we here use the natural identification End(V) = V \otimes V^{*}.

Definition 20. *Given two states of motion* s *and* s_1 *. The* relative velocity between s_1 and s, judged from s, *is defined by*

$$\mathbf{v}(s,s_1) := \frac{\mathbf{P}_s^{\top}(s_1)}{\tau(\mathbf{P}_s^{\parallel}(s_1))} = s_1 - s \in V_0.$$
(158)

This, in geometric terms, is just the ordinary definition of velocity in Newtonian physics.

The second reference to s expressed in the phrase "judged from s" is now indeed redundant due to the fact that all such relative velocities are members of the same space V₀. Clearly, we could have considered the right-hand side of (158) as element of T_sS, as in (95). But due to the flatness of (S, h) there are now natural isomorphisms between all tangent spaces T_sS given by (path-independent) parallel transport.

Remark 21. The geometric formulation of a Galilei-Newton structure goes back to Weyl (1918, § 18). Later generalisations are due to Friedrichs (1928), Dombrowski and Horneffer (1964), and also Künzle (1972). Sometimes instead of τ only $\tau \otimes \tau$ is prescribed, i.e. a degenerate "time metric" of rank one, which is equivalent to τ without the time orientation. Also, often $h \in V_0^* \otimes V_0^*$ is replaced with a $\bar{h} \in V \otimes V$ whose kernel is just Span{ τ }. In fact, most modern texts follow this formulation, presumably in order to not deal with tensor fields over proper subspaces. Algebraically both formulations are equivalent. In fact, there is a bijection between the sets of non-degenerate bilinear forms on V₀ and simply degenerate bilinear forms on V^{*} whose kernel is Span{ τ }. The bijective correspondence is this: Consider the natural isomorphism $V_0^* \otimes V_0^* \cong Lin(V_0, V_0^*)$ und accordingly regard h als Element von of $Lin(V_0, V_0^*)$. Since h is non-degenerate there exists the inverse map $h^{-1} \in Lin(V_0^*, V_0)$. Let further $i : V_0 \to V$ be the natural embedding and $i^*: V^* \to V_0^*$ its dual. The latter is given by $\lambda \mapsto i^*(\lambda) := \lambda \circ i$. Hence the kernel of i^* is just given by Span{ τ }. Now we define $\bar{h} := i \circ h^{-1} \circ i^* \in Lin(V^*, V) \cong V \otimes V$. Since i is injective and h^{-1} an isomorphism, the kernel of \bar{h} equals the kernel of i^* , i.e. Span{ τ }.

4.3 Automorphims of Galilei-Newton spacetime

Like ILor $\cong V \rtimes Lor$, the identity component of the inhomogeneous Lorentz group (also called the proper orthochronous Poincaré group), is the automorphism group of $(M, V, +, \eta, o_V, o_T)$, where o_V is the overall orientation of V and o_T is the time orientation, we define the inhomogeneous Galilei group to be the automorphism group of the geometric Galilei-Newton structure.

Definition 22. *The* inhomogeneous Galilei group *is the automorphism group of* $(M, V, +, \tau, h, o_V)$; we write IGal := Aut $(M, V, +, \tau, h, o_V)$. Here o_V stands for an orientation of V. Unlike in the special-relativistic case we do not need to also

specify a time orientation o_T , since that is already provided by τ . This group is a subgroup of the affine group Aut(M, V, +). Again it is isomorphic to a semi-direct product $IGal \cong V \rtimes Gal$, where Gal may be considered as subgroup of GL(V):

$$\begin{aligned} \text{Gal} &:= \text{Aut}(V, \tau, h, o_V) \\ &:= \left\{ G \in \text{GL}(V) : \tau \circ G = \tau, \ h \circ G|_{V_0} \times G|_{V_0} = h, \ \text{det}(G) > 0 \right\}. \end{aligned} \tag{159}$$

Note that any $G \in GL(V)$ that preserves τ also preserves $V_0 = Ker(\tau)$ i.e. it defines an element $G|_{V_0} \in GL(V_0)$ by restricting G to V_0 . That element is then required to be in SO(V_0 , h) by the second and third condition in (159) (preserving overall- and time-orientation implies that it also preserves space orientation). It is easy to see that this restriction-map defines a surjective homomorphism of groups, i.e. a "group projection":

$$\pi: \operatorname{Gal} \to \operatorname{SO}(V_0, h), \quad \operatorname{G} \to \pi(\operatorname{G}) := \operatorname{G}|_{V_0}.$$
(160)

Definition 23. *Elements in* $\text{Ker}(\pi)$ *are called* boosts.

Proposition 24. The set of boosts is given by

$$\operatorname{Gal}_{\mathsf{B}} := \operatorname{Ker}(\pi) = \left\{ \operatorname{id}_{\mathsf{V}} + \mathbf{v} \otimes \tau : \mathbf{v} \in V_0 \right\}.$$
(161)

 Gal_B is an abelian normal subgroup of Gal which is isomorphic to V_0 (considered as abelian group with group multiplication given by vector addition).

Proof. Being the kernel of a group homomorphism Gal_B is clearly a normal subgroup. Let { e_0, e_1, e_2, e_3 } be an "adapted" basis of V, which means that $\tau(e_0) = 1$ and $\tau(e_a) = 0$ for a = 1, 2, 3. Let further { $\theta^0, \theta^1, \theta^2, \theta^3$ } be the dual basis, i.e. $\theta^{\alpha}(e_{\beta}) = \delta^{\alpha}_{\beta}$; then $\theta^0 = \tau$. Writing

$$\mathbf{G} = \mathbf{G}^{\alpha}_{\ \beta} \, \mathbf{e}_{\alpha} \otimes \mathbf{\theta}^{\beta} \,, \tag{162}$$

the condition $\tau \circ G = \tau$ is equivalent to $G^0_{\ \beta}\theta^{\beta} = \theta^0$, i.e. $G^0_{\ 0} = 1$ and $G^0_{\ \alpha} = 0$ $(\alpha = 1, 2, 3)$. Moreover,

$$\pi(\mathbf{G}) = \mathbf{G}|_{V_0} = \mathbf{G}^a{}_b e_a \otimes \theta^b \quad (a, b = 1, 2, 3),$$
(163)

so that $G \in \text{Ker}(\pi)$ iff $G^a{}_b = \delta^a_b$. Taken together, the most general element in $\text{Ker}(\pi)$ is of the form

$$\mathbf{G} = e_0 \otimes \theta^0 + \delta^a_b e_a \otimes \theta^b + \mathbf{G}^a_{\ 0} e_a \otimes \theta^0 = \mathrm{id}_V + \mathbf{v} \otimes \tau, \qquad (164)$$

where $\mathbf{v} := G^{a}_{\ 0} e_{a} \in V_{0}$. Such elements form an abelian subgroup isomorphic to V_{0} inside Gal. The corresponding embedding (injective group homomorphism) is

$$B: V_0 \to Gal, \quad B(\mathbf{v}) := id_V + \mathbf{v} \otimes \tau, \tag{165}$$

which maps isomorphically onto its image $Gal_B \subset Gal$. The relations $B(\mathbf{0}) = id_V$ and $B(\mathbf{v}_1) \circ B(\mathbf{v}_2) = B(\mathbf{v}_1 + \mathbf{v}_2)$ are quite obviously satisfied. **Proposition 25.** For any $s \in S$ there is an embedding (injective group homomorphisms)

$$\sigma_{s}: SO(V_{0}, h) \to Gal, \quad D \mapsto \sigma_{s}(D) := s \otimes \tau + D \circ P_{s}^{\top}, \quad (166a)$$

such that

$$\pi \circ \sigma_{\rm s} = {\rm id}_{{\rm SO}(V_0, h)} \,. \tag{166b}$$

Proof. We first check hat σ_s is a group homomorphisms. From (157b) one immediately sees that $\sigma_s(id_{V_0}) = id_V$, and from (166a) that

$$\sigma_{s}(D_{1}) \circ \sigma_{s}(D_{2}) = (s \otimes \tau + D_{1} \circ P_{s}^{\top}) \circ (s \otimes \tau + D_{2} \circ P_{s}^{\top})$$

$$= s \otimes \tau + (D_{1} \circ D_{2}) \circ P_{s}^{\top}$$

$$= \sigma_{s}(D_{1} \circ D_{2}), \qquad (167)$$

where we used $\tau \circ D_2 \circ P_s^\top = 0$, $P_s^\top(s) = 0$, and $P_s^\top \circ D_2 \circ P_s^\top = D_2 \circ P_s^\top$. Finally,

$$\pi(\sigma_{s}(D)) = (s \otimes \tau + D \circ P_{s}^{\top})|_{V_{0}} = D, \qquad (168)$$

since $\tau|_{V_0} = 0$ and $P_s^\top|_{V_0} = id_{V_0}$.

Corollary 26. There is a splitting exact sequence

$$\{1\} \longrightarrow V_0 \xrightarrow{B} Gal \xrightarrow{\pi} SO(V_0, h) \longrightarrow \{1\}.$$
(169)

Here {1} denotes the trivial group with only one element. "Exact" means that at each node of the sequence of maps the image of the arriving equals the kernel of the departing map. For (169) this is equivalent to the three statements that 1) B is an injective group homomorphism (also called an "embedding", symbolised by a tailed arrow), that 2) π is a surjective group homomorphism (symbolised by a double-headed arrow), and that 3) Im(B) = Gal_B = Ker(π). That the exact sequence is "splitting" means that there is an injective group homomorphism – here denoted by σ_s – from SO(V₀, h) to Gal, such that $\pi \circ \sigma_s = id_{SO(V_0,h)}$ All these properties taken together are equivalent to the statement that Gal is a semi-direct product of V₀ and SO(V₀, h), usually denoted by

$$Gal = V_0 \rtimes SO(V_0, h).$$
(170)

Note that the abelian normal subgroup of boosts, $B(V_0) = Gal_B$, is defined independent of a choice of $s \in S$. In contrast, any selection of a subgroup of orthogonal spatial transformations within Gal, i.e. the image of $SO(V_0, h)$ under σ_s in Gal, does depend on a choice of $s \in S$.

The group of boosts, Gal_B , has a simple transitive action on $S = V_1$; that is, for any ordered pair (s_1, s_2) of states there is precisely one boost that maps s_1 to s_2 :

$$B(\mathbf{v})s_1 = s_2$$
 for $\mathbf{v} = \mathbf{v}(s_1, s_2) := s_2 - s_1$. (171)

Hence we call $s_2 - s_1$ the link-velocity between s_1 and s_2 . There is now no need to add the phrase "judged from s_1 " since there is exactly one boost linking s_1 with s_2 , whereas in the SR case there were many, one for each additional reference state s, the choice of which determined the set of boosts to choose the linking transformation from. Note that since the composition of pure boosts result in the simple addition of vectors in V₀, we have, e.g.,

$$\mathbf{v}(s, s_3) = \mathbf{v}(s, s_1) + \mathbf{v}(s, s_2), = (s_1 - s) + (s_2 - s).$$
(172)

This can be easily solved for $v(s, s_2)$ or $v(s, s_1)$, unlike the special-relativistic case, which could only be solved for β_2 in an elementary fashion, as shown in (12). The corresponding solution in the Galilei-Newton case is

$$\mathbf{v}(s,s_2) = \mathbf{v}(s,s_3) - \mathbf{v}(s,s_1) = (s_3 - s) - (s_1 - s) = s_3 - s_1, \quad (173)$$

where we used the laws of differences in affine space (compare the discussion below equations (212) of Appendix D.1). This shows explicitly how the dependence on s drops out – in contrast to the special-relativistic case!

4.4 Boost-Rotation decomposition for the Galilei group

The steps outlined in Section 4.1 for the SR case can now be translated almost literally to the Galilei-Newton case. Indeed, given $G \in Gal$, we do the following:

- 1. Choose a state $s \in S$ and let $s_1 := Gs$.
- 2. Let B := B(v) as in (165) with $v = s_1 s$, i.e.

$$B := B(s, s_1) = (s_1 - s) \otimes \tau + id_V.$$
(174)

3. Define

$$\mathbf{R} := \mathbf{B}^{-1} \circ \mathbf{G} \,. \tag{175}$$

Note that $(B(\mathbf{v}))^{-1} = B(-\mathbf{v})$. Clearly, R is again a Galilei transformation that fixes s: Rs = s. Hence

$$\operatorname{Stab}_{s}(\operatorname{Gal}) := \left\{ G \in \operatorname{Gal} : \operatorname{Gs} = s \right\}, \tag{176}$$

consist of Galilei transformations that map (V_0, h) isometrically into itself preserving orientation. Hence $Stab_s(Gal) \cong SO(V_0, h)$, but for $s \neq s'$ the corresponding stabiliser subgroups are conjugate subgroups in Gal:

$$\operatorname{Stab}_{s'}(\operatorname{Gal}) = \operatorname{B}(s, s') \circ \operatorname{Stab}_{s}(\operatorname{Gal}) \circ [\operatorname{B}(s, s')]^{-1}, \quad (177)$$

where B(s, s') is as in (174) for $s_1 = s'$.

4. Rewrite (175) as

$$G = B \circ R \tag{178}$$

with R – but, unlike the SR case, not B – depending on s.

The decomposition (178) is just the one that follows generally from the semidirect product structure of Gal, i.e. the splitting of (169), as explained in Appendix C. Again, as the splitting σ_s in (169) depends on $s \in S$, so does the decomposition (as far is R is concerned).

Now, according to a general argument also recalled in Appendix C, we consider for any $G \in Gal$ the element $\sigma_s(\pi(G)) \in Gal$. According to (160) and (166a) we have

$$\sigma_{s} \circ \pi(G) = \sigma_{s}(G|_{V_{0}})$$

$$= s \otimes \tau + G|_{V_{0}} \circ P_{s}^{\top}$$

$$= s \otimes \tau + G \circ P_{s}^{\top}$$

$$= G - (Gs - s) \otimes \tau.$$
(179)

Here we used (160) in the first line, (166a) in the second, the fact that we can simply drop the restriction to V_0 if we right-compose G with the projection P_s^{\top} onto V_0 in the third, and, finally, (157b) in the fourth. Hence,

$$\sigma_{s}(\pi(G)) \circ G^{-1} = \mathrm{id}_{V} - (Gs - s) \otimes \tau = B(-\mathbf{v}), \qquad (180)$$

where for the fist equality we used $\tau \circ G^{-1} = \tau$ (obvious since $G^{-1} \in Gal$) and (165) for the second, denoting $\mathbf{v} := Gs - s$. Hence, since $(B(\mathbf{v}))^{-1} = B(-\mathbf{v})$, we get (178) with

$$R := \sigma_s(\pi(G))$$
 and $B = B(\mathbf{v}) = G \circ (\sigma_s(\pi(G)))^{-1}$. (181)

We leave it to the reader to find out whether there is a (s-dependent) euclidean metric g_s on V (built from τ and h in an s-dependent fashion) with respect to which $G = B \circ R$ becomes a polar decomposition.

A Polar decomposition

Let V be an n-dimensional real vector space and $g: V \times V \to \mathbb{R}$ a positive-definite symmetric bilinear form (also known as Euclidean inner product). Let End(V) denote the linear space of all endomorphisms $V \to V$ and GL(V) the subset of all invertible elements in End(V).

$$GL(V) := \{A \in End(V) : det(A) \neq 0\}.$$
(182)

The set GL(V) is a group under composition of maps and is called the *General* Linear Group.

The Euclidean inner product g defines a map

$$\begin{aligned} \dagger : \mathrm{GL}(\mathrm{V}) &\to \mathrm{GL}(\mathrm{V}) \\ & & & & \\ & & & \\ & & & & \\ & & & & \\ & & & \\ & & & \\ & & & \\ & & & & \\ & & & \\ & & & & \\ & & & &$$

through

$$g(Av, w) = g(v, A^{\dagger}W) \quad \forall v, w \in V.$$
(184)

Symmetry of g implies that \dagger is an involution, i.e. $\dagger \circ \dagger = id_{GL(\nu)}$, or $(A^{\dagger})^{\dagger} = A$ for any $A \in GL(V)$. Also, \dagger is a group anti-isomorphism of GL(V), i.e. for any $A, B \in GL(V)$ we have¹⁹ $(id_V)^{\dagger} = id_V$ and $(AB)^{\dagger} = B^{\dagger}A^{\dagger}$ (the "anti-" denoting the reversal of orders).

The subset of fixed points in GL(V) under \dagger is

$$\operatorname{Sym}(V,g) := \{ A \in \operatorname{GL}(V) : A = A^{\dagger} \}.$$
(185)

It is called the set of *symmetric elements* in GL(V). Note that Sym(V, g) is not a subgroup, i.e. if $A = A^{\dagger}$ and $B = B^{\dagger}$ then $(AB)^{\dagger} = B^{\dagger}A^{\dagger} = BA$ which does not equal AB unless A and B commute.

Another subset is

$$\operatorname{Pos}(V,g) := \left\{ A \in \operatorname{GL}(V) : g(\nu, A\nu) > 0 \,\,\forall \nu \in V \setminus \{0\} \right\}. \tag{186}$$

Note that $A \in Pos(V, g)$ implies $A^{\dagger} \in Pos(V, g)$, i.e. \dagger maps Pos(V, g) to itself.

Of interest to us is the intersection

$$PS(V,g) := Pos(V,g) \cap Sym(V,g)$$
(187)

of elements in GL(V) which are at the same time *positive* and *symmetric*. A standard result is

Lemma 27. For any $A \in PS(V,g)$ there exists a unique $B \in PS(V,g)$ such that $A = B^2$, called its "square root". B is also denoted by \sqrt{A} or $A^{1/2}$.

Proof. Since A is symmetric there exists an orthonormal basis $\{e_1, \ldots, e_n\}$ of V that diagonalises A; i.e. $g(e_a, e_b) = \delta_{ab}$ and $A(e_a) = \lambda_a e_a$ (no summation over a). Since A is positive all eigenvalues λ_a are positive. We define B through $B(e_a) = \sqrt{\lambda_a} e_a$ (no summation over a), which clearly satisfies $A = B^2$, showing existence. To prove uniqueness, note that the B just defined commutes with A, i.e. AB = BA. Hence there is a polynomial function p of degree at most n - 1 $(n = \dim(V))$ such that p(A) = B. Indeed, if $(\lambda_1, \cdots, \lambda_k)$ with $k \le n$ is the

¹⁹ In this subsection we abbreviate the composition of maps A, B in GL(V) by juxtaposition, i.e. we write AB instead of A \circ B. Accordingly, B² means BB = B \circ B, etc.

maximal number of pairwise distinct eigenvalues of A, we can choose the interpolating Lagrange polynomial for the k pairs $\{(\lambda_1, \sqrt{\lambda_1}), \dots, (\lambda_k, \sqrt{\lambda_k})\}$ ²⁰ Now, if C is any other element in PS(V, g) satisfying $A = C^2$, we have $B = p(C^2)$, implying that B and C commute and hence that there exists an orthonormal basis diagonalising both of them simultaneously. $A = B^2 = C^2$ then implies that the squares of the eigenvalues and hence the eigenvalues themselves coincide (since they are positive). This shows B = C and hence uniqueness.

Finally we mention the *orthogonal group* of O(V, g), which is a subgroup of GL(V) and defined by

$$O(V,g) := \{A \in GL(V) : g(Av, Aw) = g(v, w) \,\forall v, w \in V\}$$

= $\{A \in GL(V) : A^{-1} = A^{\dagger}\}.$ (188)

We can now state the main theorem underlying polar decomposition:

Theorem 28 (Existence and uniqueness of polar decomposition). For any $A \in GL(V)$ and given Euclidean inner product g there exists a unique $B \in PS(V,g)$ and a unique $R \in O(V,g)$, such that

$$A = BR. (189)$$

Proof. For A ∈ GL(V) have AA[†] ∈ PS(V, g). Set B := $\sqrt{AA^{\dagger}}$ and R := B⁻¹A. Then R[†]R = A[†]B⁻¹B⁻¹A = A[†](AA[†])⁻¹A = id_V. Hence R⁻¹ = R[†] and R ∈ O(V, g), showing existence. To show uniqueness assume (B₁, R₁) and (B₂, R₂) both satisfy A = B₁R₁ = B₂R₂. Then B₂⁻¹B₁ = R₂R₁⁻¹ ∈ O(V, g) (since O(V, g) is a group). Hence, since B₁[†] = B₁, (B₂⁻¹B₁)[†] = (B₂⁻¹B₁)⁻¹ is equivalent to B₁B₂⁻¹ = B₁⁻¹B₂ or to B₁² = B₂². Hence B₁ and B₂ are both square roots of the same element in PS(V, g). Lemma 27 now implies B₁ = B₂ and hence also R₁ = R₂.

Remark 29. On the right-hand side of (189) we have put B to the left of R. We could have chosen the reversed order and proven a corresponding existence and uniqueness result. Then A = BR = R'B', with uniquely determined B, B' \in PS(V, g) and R, R' \in O(V, g). But since BR = R(R⁻¹BR) with R⁻¹BR = R[†]BR \in PS(V, g), uniqueness shows R' = R and B' = R⁻¹BR; that is, the orthogonal factor in the polar decomposition does indeed not depend on the convention concerning the order of the factors, whereas the positive symmetric part does depend on it and varies by conjugation with an orthogonal transformation.

²⁰ The general construction of such a polynomial is as follows: let $\{(x_1, y_1), \dots, (x_k, y_k)\} \subset \mathbb{R}^2$ any k points with pairwise different x-values; i.e. $i \neq j \Rightarrow x_i \neq x_j$. For each $1 \leq j \leq k$ we define $\ell_j(x) := \prod_{i=1, i\neq j}^k (x - x_i)/(x_j - x_i)$, which is a polynomial of degree (k - 1) satisfying $\ell_j(x_i) = \delta_{ij}$. Hence $\ell := \sum_{j=1}^k y_j \ell_j$ is a polynomial of degree at most (k-1) satisfying $\ell(x_i) = y_i$ for each $i \in \{1, \dots, k\}$. It is called the *Lagrange interpolating polynomial* for the given set of k points in \mathbb{R}^2 .

Remark 30. The way in which we calculated B and R from A shows that both are continuous functions of A on the domain GL(V). The inverse map $(B, R) \rightarrow A =:$ BR is trivially also continuous. Hence we have a topological equivalence²¹

$$GL(V) \cong PS(V,g) \times O(V,g)$$
. (190)

As PS(V, g) is contractible (being an open convex cone in a vector space) and O(V, g) is compact, all global topological features of GL(V) reside in the latter.

Remark 31. If $G \subset GL(V)$ is a subgroup then polar decomposition of $A \in G$ will result in some B and R in GL(V) which need not necessarily again be elements of the subset G. Whether or not that will be the case may depend on the chosen g. However, as we have seen in Section 3.1 for the Lorentz group, polar decomposition with respect to the Euclidean metric $g = \eta + 2\sigma \otimes \sigma$ (compare (91)) will again result in factors lying within the Lorentz group.

B Parallel transport along geodesics on state space

We consider state space, i.e. the 3-dimensional Riemannian manifold (S, h) as defined in Section 3.1. We recall that its metric h is just that induced from the flat Minkowski metric η of the ambient V into which σ is embedded as a spacelike hypersurface, i.e. $h = \eta|_{TS}$. In the paragraph below equation (103) we made a statement equivalent to the following:

Proposition 32. Let s_1 and s_2 be two (non-coinciding) points in S and $B(s_1, s_2)$ the unique boost in the plane $\text{Span}\{s_1, s_2\}$ mapping s_1 to s_2 . Let further $\gamma : \mathbb{R} \ni [\sigma_1, \sigma_2] \to S$, $\gamma(\sigma_i) = s_i$, be the unique geodesic on S with respect to the Levi-Civita connection for h. Then parallel transport of any $Y_1 \in T_{s_1}S$ along γ results in $Y_2 = B(s_1, s_2)Y_1 \in T_{s_2}S$.

Proof. Rather than engaging in a direct calculation, we will here follow a more geometric reasoning which we partition into the following four steps.

- The timelike 2-plane Span{s, s1} intersects S in a geodesic. This is because each point of this intersection is a fixed point of the isometry resulting from the reflection in V at Span{s, s1}. But fixed-point sets of isometries are totally geodesic (meaning that any geodesic starting in and tangential to that set remains within it). In particular, if the fixed-point set is a curve, it must itself be a geodesic.
- 2. By its very definition, the Levi-Civita covariant derivative intrinsic to an isometrically embedded hypersurface in Euclidean or semi-Euclidean²² space

²¹ In fact, this topological equivalence is a C^{∞} diffeomorphism in the natural differentiable structures that these manifolds carry.

²² In the semi-Euclidean case, the hypersurface is assumed to be nowhere lightlike.

is given by the extrinsic covariant derivative (in flat embedding space) followed by orthogonal projection tangent to the hypersurface; see, e.g. (Weyl, 1991, § 12). In our case this means that if Y is tangent to S, its covariant derivative along a curve γ in S is given by $\nabla_{\dot{\gamma}} Y = P_s^{\perp}(\partial_{\dot{\gamma}} Y)$, where $\partial_{\dot{\gamma}}$ is the Levi-Civita covariant derivative in (V, η) (i.e. "ordinary" flat derivative) and P_s^{\perp} the projector in V perpendicular to s (compare (93b)). Note that since $\dot{\gamma}$ is tangent to S, we do not need to know Y outside S in order to compute $\partial_{\dot{\gamma}} Y$. It follows that Y is parallely transported along γ within S iff at each parameter value σ the ambient (flat) covariant derivative $\partial_{\dot{\gamma}} Y$ in V is proportional to the normal to S at that point, which is just $\gamma(\sigma) \in V$. This we write as

$$\nabla_{\dot{\gamma}} Y = 0 \, \Leftrightarrow \, \partial_{\dot{\gamma}} Y \propto \gamma \,. \tag{191}$$

- 3. This implies that if $[\sigma_1, \sigma_2] \ni \sigma \mapsto Y(\sigma) \in V$ is a one-parameter family of vectors in V that obey the law (191) of parallel transport, any two $Y(\sigma)$ and $Y(\sigma')$ for $\sigma, \sigma' \in [\sigma_1, \sigma_2]$ differ only by vectors in the plane $\text{Span}\{\gamma(\sigma), \gamma(\sigma')\} = \text{Span}\{s_1, s_2\}$. In particular, the component $Y_{\perp}(\sigma) \in V$ in the orthogonal complement of $\text{Span}\{s_1, s_2\}$ is constant (independent of σ).
- 4. By definition, the boost B(s, s₁) acts in the plane Span{s, s₁} and pointwise fixes its orthogonal complement in V. Hence, applied to any Y(σ₁) ∈ T_{s1}S, it keeps its component Y_⊥ orthogonal to Span{s, s₁} fixed and maps the component Y_{||}(s) within Span{s, s₁} in such a way so as to preserve its length and keeping it tangent to S and hence tangent to the intersection S ∩ Span{s, s₁}, which is just the image of γ. But this is precisely what parallel propagation does, which completes the proof.

C Semi-direct products of groups

Semi-direct products are special examples of *group extensions*. Let us therefore first explain this more general concept.

Given two groups, called N and Q, we will combine the two into a new group, called G, such that G contains a unique normal subgroup N' isomorphic to N with quotient G/N' isomorphic to Q. Formally this is expressed by arranging the triple N-G-Q into a so-called *short exact sequence*:²³

$$\{e\} \longrightarrow \mathsf{N} \xrightarrow{i} \mathsf{G} \xrightarrow{\pi} \mathsf{Q} \longrightarrow \{e\}.$$
(192)

²³ The "short" refers to the fact that three is the smallest number of groups for which an exact sequence makes a non-trivial statement. Two groups related by an exact sequence are merely isomorphic.

Here $\{e\}$ stands for the trivial group consisting only of the identity element *e*, and all arrows denote group homomorphisms. That this sequence be *exact* means that at each node (N, G, Q) the image of the arriving map equals the kernel of the departing one. Clearly, the image of the first map from $\{e\}$ to N must be the neutral element in N so that exactness implies that the following map, i, is injective, i.e. an embedding of N into G. This is indicated by the tailed arrow from N to G. Also, as the last map from Q to $\{e\}$ has all of Q in its kernel, exactness implies that the map π is surjective. This we indicated by a double-headed arrow from G to Q.

In group-theorists' terminology, there are two ways to express the relation (192) for (N, G, Q): one either says that G is an *upward extension* of Q by N or, alternatively, that G is a *downward extension* of N by Q; see, e.g., Conway et al. (1985, p. XX). Note that simply speaking of "an extension" of one group by another is ambiguous, as it does not tell which of the two is going to have a normal embedding in the group to be constructed.

Now, a *semi-direct product* is a special case of (192), which is characterised by the existence of an injective homomorphism (an embedding)

$$\sigma: \mathbf{Q} \to \mathbf{G}$$
 such that $\pi \circ \sigma = \mathrm{id}_{\mathbf{Q}}$. (193)

In that case we write instead of (192)

$$1 \longrightarrow \mathsf{N} \xrightarrow{i} \mathsf{G} \xrightarrow[\sigma]{\pi} \mathsf{Q} \longrightarrow 1 \tag{194}$$

and say that the sequence *splits*. The map σ (193) is then called a *splitting homo-morphism*.

Definition 33 (First definition of semi-direct product). A group G is called the semi-direct product of groups N and Q if they can be arranged in a splitting short exact sequence (194). We write

$$\mathbf{G} = \mathbf{N} \rtimes_{\sigma} \mathbf{Q} \,. \tag{195}$$

The images of i and σ define subgroups of G which we denote by

$$N' := i(N) \subset G$$
 and $Q' := \sigma(Q) \subset G$. (196)

Since N' is the kernel of π it is clearly normal in G. If $q \in Q$ is different from the neutral element then $q' := \sigma(q) \notin N'$ since $\pi(q')$ must be q and hence q' cannot be in the kernel of π . Hence

$$N' \cap Q' = \{e\},$$
 (197)

where $e \in G$ denotes the neutral element in G. Moreover, any $g \in G$ is the unique product of an element $n' \in N'$ and an element $q' \in Q'$. To see this, consider the homomorphism

$$p := \sigma \circ \pi : G \to Q', \quad g \mapsto q' := \sigma(\pi(g)).$$
 (198)

It satisfies $p|_{Q'} = id_{Q'}$ and $p \circ p = p$ due to (193); i.e. it is a projection homomorphism from G onto the subgroup Q'. Now, for any $g \in G$, define $q' \in Q'$ as above and $n' := gq'^{-1} \in N'$ (which is indeed in $N' = \text{Ker}(\pi)$ since $\pi(q') = \pi(g)$). This decomposition is unique, for if (n'_1, q'_1) and (n'_2, q'_2) both satisfy $g = n'_1q'_1 = n'_2q'_2$ it follows that $n'_2^{-1}n'_1 = q'_2q'_1^{-1}$. But the lefthand side is in N' and the right-hand side in Q', so that (197) implies that both sides must equal e, hence $n'_1 = n'_2$ and $q'_1 = q'_2$.

The discussion in the previous paragraph gives rise to two alternative definitions of semi-direct products:

Definition 34 (Second definition of semi-direct product). A group G is called the semi-direct product of its subgroups N' and Q' if the set N'Q' := $\{n'q' : n' \in N', q' \in Q'\}$ equals G, N' $\cap Q' = \{e\}$, and N' is normal in G.

Definition 35 (Third definition of semi-direct product). A group G is called the semi-direct product of its subgroups N' and Q' if there is a projection homomorphism $p : G \rightarrow Q'$ with kernel N'. (Projection meaning: $p|_{Q'} = id_{Q'}$ and $p \circ p = p$.)

Composing σ with the map $Ad_{n'} : G \to G$, $g \mapsto Ad_{n'}(g) := n'gn'^{-1}$, where $n' \in N$ is some element from N', clearly gives another splitting homomorphism $\sigma' := Ad_{n'} \circ \sigma$ satisfying $\pi \circ \sigma' = id_Q$ since n' is in the kernel of π . Hence, in general, neither σ nor Q' are unique. For example, the group E(3) of euclidean motions is a semi-direct product of translations $N = \mathbb{R}^3$ with rotations Q = SO(3). Any splitting embedding $s : SO(3) \to E(3)$ is characterised by the point $\sigma \in \mathbb{R}^3$ (the "origin") about which the elements of SO(3) rotate, i.e. which is fixed under the action of all elements of SO(3). And any two SO(3) subgroups in E(3) differ by a conjugation with the translation mapping the origin of the first into the origin of the second rotation group.

Let us consider Definition 34. The multiplication of $g_1 = n'_1 q'_1$ with $g_2 = n'_2 q'_2$ is

$$g_1g_2 = n'_1q'_1n'_2q'_2 = n'_1q'_1n'_2q'_1^{-1}q'_1q'_2 =: n'_3q'_3,$$
(199a)

where

$$n'_{3} := n'_{1}(q'_{1}n'_{2}q'_{1}^{-1}) = n'_{1}Ad_{q'_{1}}(n'_{2}) \text{ and } q'_{3} = q'_{1}q'_{2}$$
 (199b)

Note that $n'_3 \in N'$ since N' is normal and that, for any $q' \in Q'$, $Ad_{q'|N'} \in Aut(N')$. In fact, it is obvious that the map $Q' \to Aut(N')$, $q' \mapsto Ad_{q'|N'}$ is a homomorphism. Identifying N' with N via the homomorphism $i : N \to G$ which is an isomorphism onto its image N', and likewise and identifying Q with Q' via the homomorphism $\sigma : Q \to G$ which is an isomorphism onto its image Q', we get yet another definition of a semi-direct product that, like Definition 33 is in terms of N and Q:

Definition 36 (Fourth definition of semi-direct product). Let N and Q be groups. Let further

$$\alpha: Q \to Aut(N), \quad q \mapsto \alpha_q \tag{200}$$

be a homomorphism. Hence $\alpha_q = id_N$ if $q = e_Q$ (neutral element in Q) and $\alpha_{q_1} \circ \alpha_{q_2} = \alpha_{q_1q_2}$. Then the set $G := N \times Q$ is made into a group by defining the multiplication through

$$(n_1, q_1)(n_2, q_2) = (n_1 \alpha_{q_1}(n_2), q_1 q_2), \qquad (201)$$

which is called the semi-direct product of N with Q relative to α and denoted by

$$G = N \rtimes_{\alpha} Q. \tag{202}$$

We note the following more or less immediate consequences of this definition:

1. If e_N and e_Q denote the neutral elements of N and Q, respectively, the neutral element of G is

$$\boldsymbol{e}_{\mathrm{G}} = (\boldsymbol{e}_{\mathrm{N}} \,, \, \boldsymbol{e}_{\mathrm{Q}}) \tag{203}$$

and the inverse element of $(n, q) \in G$ is

$$(n,q)^{-1} = (\alpha_{q^{-1}}(n), q^{-1}).$$
 (204)

2. It is immediate from (201) that the map

$$\pi: \mathbf{G} \to \mathbf{Q}, \quad (\mathbf{n}, \mathbf{q}) \mapsto \pi(\mathbf{m}, \mathbf{q}) \coloneqq \mathbf{q}$$
(205)

is a surjective homomorphism whereas the projection onto the first factor $G \to N$, $(n, q) \mapsto n$ fails to be a homomorphism unless α is trivial, that is, $\alpha_q = id_N$ for all $q \in Q$ and $G = N \times Q$ is a proper direct product of groups.

3. N and Q can be embedded into G via the injective homomorphisms

$$\mathbf{i}: \mathbf{N} \to \mathbf{G}, \mathbf{n} \mapsto \mathbf{i}(\mathbf{n}) := (\mathbf{n}, \mathbf{e}_{\mathbf{Q}}), \qquad (206a)$$

$$\sigma: Q \to G, q \mapsto \sigma(n) := (e_N, q), \qquad (206b)$$

the images of which are

$$i(N) =: N' = \{(n, e_Q) : n \in N\} \subset G,$$
 (207a)

$$\mathfrak{r}(\mathbf{N}) \coloneqq \mathbf{N}' = \{(\mathbf{n}, e_{\mathbf{Q}}) : \mathbf{n} \in \mathbf{N}\} \subset \mathbf{G}, \qquad (20/a)$$
$$\sigma(\mathbf{Q}) \coloneqq \mathbf{Q}' = \{(e_{\mathbf{N}}, q) : q \in \mathbf{Q}\} \subset \mathbf{G}. \qquad (207b)$$

Obviously $N' = \text{Ker}(\pi)$ and $\pi \circ \sigma = \text{id}_Q$. Therefore, G defined in (202), together with its subgroups defined in (207) and maps defined in (205) and (206), are related by a short exact sequence (194), leading us back to Definition Definition 33

4. From (201) have $(h, e_Q)(e_H, q) = (h, q)$ and hence

$$H' \cap Q' = e_G$$
 and $H'Q' = G$, (208)

leading us back to Definition 34 Similarly for Definition 35, since the maps π from (205) and σ from (206b) combine to $p := \sigma \circ \pi$, which is the required projection homomorphism $G \to Q'$.

5. Definition 36 shows that any possible homomorphism (200) from Q into Aut(N) can be realised in a semi-direct product, in which the automorphisms of N' (which are usually outer) then appear as restrictions of inner automorphism of G to its normal subgroup N', as in (199b):

$$q_{1}'n_{2}'q_{1}'^{-1} = (e_{N}, q_{1})(n_{2}, e_{Q})(e_{N}, q_{1})^{-1}$$

= $(\alpha_{q_{1}}(n_{2}), q_{1})(\alpha_{q_{1}^{-1}}(e_{N}), q_{1}^{-1})$ (209)
= $(\alpha_{q_{1}}(n_{2}), e_{Q}),$

using (201), (204), and that $\alpha_q(e_N) = e_N$ for all $q \in Q$.

Remark 37. Note that Definitions 33 and 36 use groups N and Q to construct a new group G, whereas Definitions 34 and 35 consider G as given an characterise it in terms of subgroups N' \subset G and Q' \subset G. Therefore, Definitions 33 and 36 are often said to define an *outer* and Definitions 34 and 35 an *inner* semi-direct product.

D Affine structures

In this appendix we review the notion of affine spaces which underlies Minkowski spacetime in Special Relativity and also Galilei-Newton spacetime. Most likely, most of what is being said here will be known to the reader in one form or another, though perhaps it is useful to recall the essential structural properties in a way adapted to the language used in this paper.

In the main text we characterised Minkowski spacetime by the following 6tuple $(M, V, +, \eta, o_V, o_T)$, where (M, V, +) is a 4-dimensional real affine space, $\eta \in V^* \otimes V^*$ is a symmetric, non-degenerate, bilinear form on V of signature (-, +, +, +), o_V denotes an overall orientation of V, and, finally, o_T denotes a time-orientation of (V, η) . Similarly we characterised Galilei-Newton spacetime as a 5-tuple $(M, V, +, \tau, h, o_V)$, where $\tau \in V^*$ is an oriented time-distance function and $h \in [\text{Ker}(\tau)]^* \otimes [\text{Ker}(\tau)]^*$ is a Euclidean metric (symmetric positive definite bilinear form) on $\text{Ker}(\tau)$. Note that $\text{Ker}(\tau)$ receives an orientation from o_V and τ (the latter defines a time-orientation). In this appendix we recall the meaning of (M, V, +) and also explain the notion of affine automorphisms and affine bases. There will be no need to restrict the dimension n which we keep general. Also, we could have easily generalised the underlying number field to \mathbb{C} , but for definiteness we stick with \mathbb{R} . Hence we start with a general definition of the notion of a real affine space.

D.1 Affine spaces

Definition 38. A real affine space of dimension n is a triple (M, V, ϕ) , where M is a set, V is an n-dimensional real vector space, and ϕ is a simply transitive action of V – considered as abelian group – on M. Here "action" means that there is a group-homomorphism from V into the group Bij(M) of bijections of M (with group multiplication being given by composition of maps, denoted by \circ , and the neutral element being the identity map). Hence there is a map

$$\phi: V \to \operatorname{Bij}(M), \quad \nu \mapsto \phi_{\nu}, \tag{210}$$

which for all $v', v \in V$ obeys,

$$\phi_{\nu=0} = \mathrm{id}_V, \qquad (211a)$$

$$\phi_{\nu'+\nu} = \phi_{\nu'} \circ \phi_{\nu} \,. \tag{211b}$$

That this action is "simply transitive" means that for any ordered pair (m, m') of points in M there exists a unique $v \in V$ such that $\phi_v(m) = m'$. Note that ordinary transitivity would be that without the uniqueness statement.

In passing we remark that for any group action on a set we generally have that simple transitivity implies ordinary transitivity, as well as that the action is free. The latter means that the homeomorphism (210) is injective, i.e. $\phi_{\nu} = id_V \Rightarrow \nu = 0$. Now, for abelian groups the converse holds: A free and transitive action is necessarily simply transitive. Hence we could have required ϕ to be free and transitive. We leave the simple proof of the equivalence to the reader.

Recall that group multiplication in V is vector addition "+". Now, it is general practice, although this may be confusing at first, to denote the action of V on M by the very same symbol, i.e. to write m + v instead of $\phi_v(m)$, and hence to eliminate all explicit reference to ϕ . For a given group action ϕ , this does not lead to ambiguities since whether a "+" means group action on M or vector addition in V is uniquely determined by whether the "+" stands between an element of M and an element of V or between two elements of V, respectively. Equations (211) then assumes the simple form

$$\mathfrak{m} + \mathfrak{0} = \mathfrak{m}, \qquad (212a)$$

$$(m + v) + v' = m + (v + v').$$
 (212b)

Note that in (212b) both "+" on the left-hand side are group actions whereas the first "+" on the right-hand side is a group acton and the second is vector addition.

Since for any given two m, $m' \in M$ there exists a unique $v \in V$ so that m' = m + v, we may write m' - m = v. Then, trivially, m' = m + (m' - m). There are other "obvious" relations, like (m' - p) + (p - m) = m' - m or m + (m' - p) = m' + (m - p), valid for all m', m, p in M. Likewise, one has (m' - m) = -(m - m'), where here the two "-" on the right-hand side have two different meanings: In the bracket it denotes the difference operation in M, in front of the bracket scalar multiplication with $(-1) \in \mathbb{R}$ in V. Other obvious notational simplifications apply, like m - v := m + (-v).

As an alternative to Definition 38 above, affine spaces can be defined via the difference map just introduced:

$$\Delta: \mathsf{M} \times \mathsf{M} \to \mathsf{V}, \quad (\mathfrak{m}', \mathfrak{m}) \mapsto \Delta(\mathfrak{m}', \mathfrak{m}) = \mathfrak{m}' - \mathfrak{m}.$$
(213)

Definition 39 (alternative to Definition 38). A real affine space of dimension n is a triple (M, V, Δ) , where M is a set, V is an n-dimensional real vector space, and $\Delta : M \times M \rightarrow V$ is a map that satisfies the following two conditions for any o, m, m', m'' in V:

$$\Delta_{\mathbf{o}}: \mathbf{M} \to \mathbf{V}, \quad \mathbf{m} \mapsto \Delta_{\mathbf{o}}(\mathbf{m}) \coloneqq \Delta(\mathbf{m}, \mathbf{o}) \quad is \ a \ bijection \ , \tag{214a}$$

$$\Delta(\mathfrak{m}'',\mathfrak{m}') + \Delta(\mathfrak{m}',\mathfrak{m}) = \Delta(\mathfrak{m}'',\mathfrak{m}).$$
(214b)

That the existence of (M, V, ϕ) implies (M, V, Δ) has been shown above. Conversely, given (M, V, Δ) with Δ satisfying the axioms above, we deduce a simply-transitive action ϕ of V on M by setting $\phi_{\nu}(m) := \Delta_m^{-1}(\nu)$. Indeed, from (214b) we get for m'' = m' = m that $\Delta(m, m) = 0$, which is equivalent to $\Delta_m^{-1}(0) = m$ for all m, which in turn implies (211a). The second condition (211b) can be deduced as follows: (214b) is equivalent to $\Delta_{m'}(m'') + \Delta_m(m') = \Delta_m(m'')$, which in turn is equivalent to $\Delta_m^{-1}(\Delta_{m'}(m'') + \Delta_m(m')) = m''$ for all m'', m', m. Setting $\Delta_{m'}(m'') =: \nu'$ and $\Delta_m(m') =: \nu$ in order to replace in that equation m'' and m' by ν' and ν , this is equivalent to $\Delta_m^{-1}(\nu' + \nu) = \Delta_{m'}^{-1}(\nu')$. Finally, setting $m' = \Delta_m^{-1}(\nu)$ on the right-hand side, this is seen to be equivalent to $\phi_{\nu'+\nu}(m) = \phi_{\nu'}(\phi_{\nu}(m))$ for all m, ν, ν' , and hence to (211b).

Summing up we can say that in an affine space we can add vectors to points and take differenced of points according to the rules given above. However, points cannot be added. To be sure, any point $o \in M$ defines a bijection $\phi_o : M \to V$ via $m \mapsto \phi_o(m) := (m - o)$. But the linear structure thereby pulled back to M, which is given by $m + m' := \phi_o^{-1} (\phi_o(m) + \phi_o(m')) = o + (m - o) + (m' - o)$, depends on the choice of o. In fact, through an appropriate choice of o *any* point p of M can be obtained as result of such an "addition" of m and m': just choose o = m + (m' - p) = m' + (m - p).

D.2 Affine maps and groups

Definition 40. Let $A = (M, V, \phi)$ and $A' = (M', V', \phi')$ be two affine spaces. An *affine map* from A to A' consists of a pair (F, f) of maps,

$$F: M \to M', \quad f: V \to V' \quad linear,$$
 (215a)

such that

$$\mathsf{F} \circ \mathbf{\Phi} = \mathbf{\Phi}' \circ (\mathsf{f} \times \mathsf{F}) \,. \tag{215b}$$

Here we explicitly displayed the action ϕ of V on M and likewise ϕ' of V' on M', which we consider as maps $\phi : V \times M \to M$, $(v, \mathfrak{m}) \mapsto \phi_v(\mathfrak{m})$, and $\phi' : V \times M' \to M'$, $(v, \mathfrak{m}') \mapsto \phi'_v(\mathfrak{m}')$, respectively. In our simplified notation, in which both actions are written by a common (+)-sign, this reads

$$F(p + v) = F(p) + f(v)$$
, (215c)

for all $p \in M$ and all $v \in V$.

Before we proceed we explicitly check that (215c) makes sense, i.e. leads to the same result independent of how we represent a point $p \in M$ as "sum" of a point with a vector. So let $p = p_1 + v_1 = p_2 + v_2$; then $F(p_1 + v_1) = F(p_1) + f(v_1)$ and $F(p_2+v_2) = F(p_2)+f(v_2)$. But $F(p_2) = F(p_1+(p_2-p_1)) = F(p_1)+f(p_2-p_1) = F(p_1) + f(v_1 - v_2)$ and linearity of f shows that indeed both sides are equal. Note that condition (215c) says that once we know the map f and the value F(q) of the map F for a single point q, we know the map F, i.e. F(p) for any p, namely F(p = q + (p - q)) = F(q) + f(p - q).

In view of the alternative definition of affine spaces in terms of the difference map Δ , we could also have given a corresponding alternative definition of an affine map:

Definition 41 (alternative to Definition 40). Let $A = (M, V, \Delta)$ and $A' = (M', V', \Delta')$ be two affine spaces. An affine map from A to A' consists of a pair (F, f) of maps,

$$F: M \to M', \quad f: V \to V' \quad linear,$$
 (216a)

such that

$$\Delta' \circ (\mathsf{F} \times \mathsf{F}) = \mathsf{f} \circ \Delta \,. \tag{216b}$$

In our simplified notation, in which the difference-map is written by a (-)-sign, this reads

$$\Delta'(F(p) - F(q)) = f(p - q), \qquad (216c)$$

for all $q, p \in M$.

Definition 42. An affine map between affine spaces is called an **affine isomorphism** iff the map $F : M \to M'$ is a bijection. This is equivalent so the requirement

for $f: V \to V'$ to be a bijection (and hence a vector-space isomorphism). An affine isomorhism of an affine space to itself is called an **affine automorphism**. The set of affine automorphisms of an affine space A forms a group under composition which is called the **general affine group**, denoted by GA(A).

In order to understand the group-theoretic structure of GA(A) we first need a label-set that faithfully labels each of its elements. This can be obtained in the following way: Choose a reference point $o \in M$ and use it to label F by the pair (ν, f) , where $\nu \in V$ is defined by $F(o) = o + \nu$, or o := F(o) - o. The action of F on a general point p is then

$$F(p) = F(o + (p - o)) = o + v + f(p - o).$$
(217)

Note that the argument of f is always the difference between the argument p and the chosen base-point o, so that no f moves o. In this way we identify GL(V) with that subgroup of GA(A) which stabilises o. We will see below how this identification behaves under changes of o.

Suppose now that we have two affine automorphisms F_1 and F_2 , which we label by (v_1, f_1) and (v_2, f_2) as just explained, with reference to the same point o. The action of the composition $F_1 \circ F_2$ on a general point can then be calculated:

$$F_1 \circ F_2(p) = F_1(o + \nu_2 + f_2(p - o)) = o + \nu_1 + f_1(\nu_2) + f_1 \circ f_2(p - o).$$
(218)

Form that we infer the multiplication law for GA(A) in the chosen parametrisation

$$(v_1, f_1)(v_2, f_2) = (v_1 + f_1(v_2), f_1f_2)$$
(219)

where composition of maps are now written by simple juxtaposition in order to stress that it is group multiplication (in GA(A) and GL(V)). We infer that GA(A) is isomorphic to the semi-direct product of the abelian group V with GL(V):

$$GA(A) \simeq V \rtimes GL(V)$$
. (220)

The homomorphism α : $GL(V) \rightarrow Aut(V)$, whic we need in order to define a semidirect product, is just the identity if we use the isomorphism $Aut(V) \simeq GL(V)$.

However, it is important to keep in mind that this isomorphism depends on the chosen basepoint o. Had we chosen another one, say o' = o + w, then

$$F(p) = o + v + f(p - o)$$

= o' + v + (o - o') + f(p - o' + (o' - o))
= o' + v + [-w + f(p - o' + w)]
= o' + v + [T_{-w} \circ f \circ T_{w}](p - o'), (221)

where $T_w : V \to V, v \mapsto v + w$ denotes the translation-action of V onto itself. This means that the same affine map F that with respect to the basepoint o is represented

by the pair (v, f) will be represented with respect to o' = o + w by the pair (v', f') where v' = v and $f' = T_{-w} \circ f \circ T_w$. This is intuitively obvious, since the representation (220) of GA(A) by the semi-direct product selects amongst all subgroups GL(V) that one which fixes the selected base-point o. In changing o to o' we also change the subgroup in GA(A) from the stabiliser subgroup of o to that of o' = o + w. These two stabiliser subgroups are clearly related by conjugation with a translation.

An invariant characterisation of of the affine group GA(A) in terms of V and GL(V) would be so say that GA(A) is a "splitting downward extension of GL(V) by V" or, equivalently, a "splitting upward extension of V by GL(V))" (compare AppendixC), which in any case means that we have a splitting short exact sequence with normal subgroup V (translations) and quotient group Q = GL(V):

$$\{1\} \longrightarrow V \xrightarrow{i} GA(A) \xrightarrow[\sigma_o]{\pi} GL(V) \longrightarrow \{1\}.$$
 (222)

The splitting homomorphism (embedding) $\sigma_o : GL(V) \to GA(A)$ depends on the chosen base-point $o \in M$ and is not natural (i.e. no one is distinguished). And two different ones, say σ_o and $\sigma_{o'}$ with o'-o = w, result in different image-subgroups in GA(A) which are conjugate by an element in the image of i (the embedding of V into GA(A)):

$$\sigma_{o+w} \Big(\mathrm{GL}(\mathbf{V}) \Big) = \mathfrak{i}_{w} \circ \sigma_{o} \Big(\mathrm{GL}(\mathbf{V}) \Big) \circ \mathfrak{i}_{(-w)} \,. \tag{223}$$

This means that whereas it makes sense to speak of *the* translations, since they form a normal subgroup, it does *not* make sense to speak of *the* subgroup of homogeneous transformations: GL(V) is a *quotient*- not a sub-group. It may be considered as a subgroup, though not uniquely. There are as many *different* subgroups isomorphic to GL(V) in GA(A) as there are points in V. None of them is intrinsically distinguished.

D.3 Affine bases and charts

Definition 43. Let $A = (M, V, \phi)$ be n affine space. An affine basis B of A is a tuple B = (o, b), where $o \in M$ and $b := \{e_1, \dots, e_n\}$ is a basis for V. We note that the basis b of V uniquely determines a dual basis $b^* = \{\theta^1, \dots, \theta^n\}$ of V^* , the dual vector space to V. It satisfies $\theta^a(e_b) = \delta_b^a$.

Remark 44. Note that an affine basis determines (n + 1) points $\{0, 0 + e_1, \dots, 0 + e_n\}$ which are not contained in any m-dimensional affine subspace with m < n. Hence we may equivalently characterise an affine basis of an n-dimensional affine space by (n + 1) points $\{p_0, p_1, \dots, p_n\} \subset M$ which are independent in the sense of not being contained in any m-dimensional affine subspace for m < n. It is then obviously true that the differences $e_a := p_a - p_0$ form a basis for V, independently of which of the (n + 1) points is selected as p_0 .

Definition 45. An *affine chart* is a bijection $\phi_B : M \to \mathbb{R}^n$ induced by an affine basis B in the following way: the value of the a-th component in \mathbb{R}^n of $\phi_B(p)$ is

$$\left[\phi_{B}(p)\right]^{\mathfrak{a}} \coloneqq \theta^{\mathfrak{a}}(p-o) \,. \tag{224a}$$

Its inverse is (summation convention)

$$\phi_{\rm B}^{-1}(x^1, \cdots, x^n) = o + x^a e_a.$$
(224b)

Suppose now that we have two affine bases, B = (o, b) and B' = (o', b'), with $b := \{e_1, \dots e_n\}$ and $b' := \{e'_1, \dots e'_n\}$, and corresponding dual bases $\{\theta^1, \dots \theta^n\}$ and $\{\theta'^1, \dots \theta'^n\}$, respectively. Then

$$o' = o + a, \quad e'_b = L^a_{\ b} e_a, \quad \theta'^a = [L^{-1}]^a_{\ b} \theta^b.$$
 (225)

Definition 46. The transition function between the affine charts ϕ_B and $\phi_{B'}$ is the bijection

$$\phi_{BB'} \coloneqq \phi_B \circ \phi_{B'}^{-1} : \mathbb{R}^n \to \mathbb{R}^n \,. \tag{226a}$$

Using (224) and (225), the a-th component in \mathbb{R}^n of the transition function is given by

$$\Phi^{\mathfrak{a}}_{\mathsf{B}\mathsf{B}'}(x^1,\cdots,x^n) = \mathfrak{a}^{\mathfrak{a}} + L^{\mathfrak{a}}_{\phantom{\mathfrak{b}}b}x^{\mathfrak{b}}, \qquad (226b)$$

where $a^{\alpha} := \theta^{\alpha}(\alpha)$ is the α -th component of α in the basis $\{e_1, \cdots, e_n\}$ of V.

If we write the coordinates of the point p in the chart B' by $x'^{\alpha}(p)$ and that in the chart B by $x^{\alpha}(p)$, (226b) reads

$$x^{\mathfrak{a}}(\mathfrak{p}) = \mathfrak{a}^{\mathfrak{a}} + L^{\mathfrak{a}}_{\ \mathfrak{b}} x'^{\mathfrak{b}}(\mathfrak{p}), \qquad (227)$$

To be distinguished from that relation between the coordinates of one and the same point p in two different charts is the coordinate representation of an affine map in a single chart B. Given an affine automorphism (F, f) of $A = (M, V, \phi)$ and an affine basis $B = (o, \{e_1, \dots, e_n\})$, such that F(o) = o + a and $f(e_b) = L^a_{\ b}e_a$, then the coordinate representation of the affine map is

$$F_{B} := \phi_{B} \circ F \circ \phi_{B}^{-1} : \mathbb{R}^{n} \to \mathbb{R}^{n} .$$
(228a)

Using (224b), (215c), and (224a) this leads to

$$F_{B}^{\mathfrak{a}}(x^{1},\cdots,x^{n}) = \mathfrak{a}^{\mathfrak{a}} + L^{\mathfrak{a}}_{\ b}x^{\mathfrak{b}}.$$
(228b)

This has the same analytic form as (226b), but the meaning is clearly different. To state it once more:

Remark 47. Whereas (228b) relates the different coordinates of two *different points* in M with respect to the *same chart*, (226b) relates the different coordinates of the *same point* in two *different charts*. The former is sometimes called a *passive* and the latter an *active* coordinate transformation.

Remark 48. Finally we point out that the existence of preferred charts endows affine spaces with a differentiable and even analytic structure. A function $f: M \to \mathbb{R}$ is called differentiable/analytic, if $f_B := f \circ \varphi_B^{-1} : \mathbb{R}^n \to \mathbb{R}$ is. This is independent of what chart B we use, since obviously $f_{B'} = f_B \circ \varphi_{BB'}$. As $\varphi_{BB'}$ is, according to (226b), an affine-linear map, hence in particular analytic, we infer that $f_{B'}$ is smooth/analytic iff f_B is.

References

- Benz, Walter. 2000. "A Characterization of Relativistic Addition." Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg 70:251–258. url: https://doi.org/10.1007/BF02940916
- Benz, Walter. 2002. "A Common Characterization of Classsical and Relativistic Addition." *Journal of Geometry* 74:38–43. url: https://doi.org/10.1007/PL00012536
- Celakoska, Emilija, Dushan Chakmakov and Mirko Petrushevski. 2015. "On Parameterization of Lorentz Boost Links." *International Journal of Contemporary Mathematical Sciences* 10(2):85–90.

```
url: http://www.m-hikari.com/ijcms/ijcms-2015/1-4-2015/
celakoskaIJCMS1-4-2015.pdf
```

- Celakoska, Emilija G. 2008. "On Isometry Links between 4-Vectors of Velocity." Novi Sad Journal of Mathematics 38(3):165-172. url: https://www.emis.de/journals/NSJOM/Papers/38_3/ NSJOM_38_3_165_172.pdf
- Conway, John H. et al. 1985. *ATLAS of Finite Groups*. Oxford: Oxford University Press.
- Dombrowski, Hainz Dieter and Klaus Horneffer. 1964. "Die Differentialgeometrie des Galileischen Relativitätsprinzips." *Mathematische Zeitschrift* 86(4):291– 311.

url: https://doi.org/10.1007/BF01110404

- Einstein, Albert. 1905. "Zur Elektrodynamik bewegter Körper." Annalen der Physik 323(13):891–921. Reprinted with annotations in (Einstein, 1989, Doc. 23, p. 275-310). url: https://doi.org/10.1002/andp.19053221004
- Einstein, Albert. 1989. The Collected Papers of Albert Einstein. Volume 2, The Swiss Years: Writings 1900-1909. Edited by John Stachel. Princeton: Princeton University Press.
- Friedrichs, Kurt. 1928. "Eine invariante Formulierung des Newtonschen Gravitationsgesetzes und des Grenzüberganges vom Einsteinschen zum Newtonschen Gesetz." *Mathematische Annalen* 98(1):566–575. url: https://doi.org/10.1007/BF01451608
- Giulini, Domenico. 2001. "Uniqueness of Simultaneity." *Britisch Journal for the Philosophy of Science* 52(4):651–670. url: https://doi.org/10.1093/bjps/52.4.651

Giulini, Domenico. 2006. Algebraic and Geometric Structures in Special Relativity. In Special Relativity: Will it Survive the Next 101 Years?, ed. Claus Lämmerzahl and Jürgen Ehlers. Vol. 702 of Lecture Notes in Physics Berlin: Springer Verlag pp. 45–111.

url: https://doi.org/10.1007/3-540-34523-X_4

- Gourgoulhon, Éric. 2013. Special Relativity in General Frames. Graduate Texts in Physics Berlin: Springer Verlag. url: https://doi.org/10.1007/978-3-642-37276-6
- Greub, Werner H. 1975. *Linear Algebra*. Vol. 23 of *Graduate Texts in Mathematics* fourth ed. New York: Springer Verlag. url: https://doi.org/10.1007/978-1-4684-9446-4
- Jacobson, Nathan. 1985. *Basic Algebra I.* second ed. New York: W.H. Freeman and Co.
- Koczan, Grzegorz Marcin. 2023. Relative binary and ternary 4D velocities in the Special Relativity in terms of manifestly covariant Lorentz transformation. In Scientific Legacy of Professor Zbigniew Oziewicz. Vol. 75 of Series on Knots and Everything Singapore: World Scientific p. 169–206. url: https://doi.org/10.1142/9789811271151_0009
- Künzle, Hans-Peter. 1972. "Galilei and Lorentz structures on space-time: comparison of the corresponding geometry and physics." Annales de l'Institut Henri Poincaré (Section A) 17(4):337–362. url: http://www.numdam.org/item/?id=AIHPA_1972__17_4_ 337_0
- Laue, Max von. 1911. *Das Relativitätsprinzip*. Vol. 38 of *Die Wissenschaft* 1st ed. Braunschweig: Verlag von Friedrich Vieweg & Sohn.
- Macfarlane, Alan J. 1962. "On the Restricted Lorentz Group and Groups Homomorphically Related to It." *Journal of Mathematical Physics* 3(6):1116–1129. url: https://doi.org/10.1063/1.1703854
- Matolcsi, Tamás and A. Goher. 2001. "Spacetime without reference frames: An application to the velocity addition paradox." *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 32(1):83–99.

url: https://doi.org/10.1016/S1355-2198(00)00037-X

Matolcsi, Tamás and Máté Matolcsi. 2005. "Thomas rotation and Thomas precession." *International Journal or Theoretical Physics* 44(1):63–77. url: https://doi.org/10.1007/s10773-005-1437-y

- Mocanu, Constantin I. 1986. "Some difficulties within the framework of relativistic electrodynamics." *Archiv für Elektrotechnik* 69:97–110. url: https://doi.org/10.1007/BF01574845
- O'Neill, Barrett. 1983. Semi-Riemannian Geometry. With Applications to Relativity. Pure and Applied Mathematics New York: Academic Press.
- Oziewicz, Zbigniew. 2006. "The Lorentz boost-link is not unique. Relative velocity as a morphism in a connected groupoid category of null objects.". url: https://doi.org/10.48550/arXiv.math-ph/0608062
- Oziewicz, Zbigniew. 2007. "Relativity groupoid instead of relativity group." *International Journal of Geometric Methods in Modern Physics* 04(05):739–749. url: https://doi.org/10.1142/S0219887807002260
- Oziewicz, Zbigniew. 2011. "Ternary relative velocity; astonishing conflict of the Lorentz group with relativity.". url: https://doi.org/10.48550/arXiv.1104.0682
- Oziewicz, Zbigniew and William S. Page. 2011. "Concepts of relative velocity.". url: https://doi.org/10.48550/arXiv.1104.0684
- Robb, Alfred A. 1911. Optical Geometry of Motion: A New View of the Theory of Relativity. Cambridge: W. Heffer & Sons Ltd.
- Silberstein, Ludwik. 1914. *The Theory of Relativity*. London: MacMillan and Co. Limited. Based on lectures delivered at University College London in 1912-13.
- Streater, Raymond F. and Arthur Strong Wightman. 1964. PCT, Spin and Statistics, and All That. New York: W.A. Benjamin, Inc.
- Ungar, Abraham. 1989. "The relativistic velocity composition paradox and the Thomas rotation." *Foundations of Physics* 19(11):1385–1396. url: https://doi.org/10.1007/BF00732759
- Ungar, Abraham. 2002. Beyond the Einstein Addition Law and its Gyroscopic Thomas Precession. The Theory of Gyrogroups and Gyrovector Spaces. New York: Kluwer Academic Publishers.
- Ungar, Abraham Albert. 1988. "Thomas Rotation and the Parametrization of the Lorentz Transformation Group." *Foundations of Physics Letters* 1(1):57–89. url: https://doi.org/10.1007/BF00661317
- Ungar, Abraham Albert. 1997. "Thomas Precession: Its Underlying Gyrogroup Axioms and Their Use in Hyperbolic Geometry and Relativistic Physics." *Foundations of Physics* 27(6):881–951.
- Ungar, Abraham Albert. 2005. Analytic Hyperbolic Geometry: Mathematical Foundations and Applications. Singapore: World Scientific.

Urbantke, Helmuth. 1990. "Physical holonomy: Thomas precession, and Clifford algebra." *American Journal of Physics* 58(8):747–750. Erratum ibid. 59(12), 1991, 1150-1151.

url: https://doi.org/10.1119/1.16401

- Urbantke, Helmuth. 2003. "Lorentz Transformations from Reflections: Some Applications." *Foundations of Physics Letters* 16(2):111–117. url: https://doi.org/10.1023/A:1024162409610
- Weyl, Hermann. 1918. *Raum Zeit Materie*. 1st ed. Berlin: Verlag von Julius Springer.
- Weyl, Hermann. 1991. *Raum Zeit Materie*. 8th ed. Berlin: Springer Verlag. Based on the 1923 5th edition, with additional appendices, edited and annotated by Jürgen Ehlers.