

A NEW PROOF OF THE SHARP GORDON'S LEMMA: NO EIGENVALUES FOR SCHRÖDINGER OPERATORS WITH ALMOST REPETITION POTENTIALS

WENCAI LIU

Dedicated to Barry Simon on the occasion of his 80th birthday

ABSTRACT. Building on the work of Jitomirskaya-Simon and Jitomirskaya-Liu, who established the absence of eigenvalues for Schrödinger operators with almost reflective repetition potentials, we provide a new proof of the sharp Gordon's lemma, which asserts the absence of eigenvalues for Schrödinger operators with almost repetition potentials.

1. INTRODUCTION

In this paper, we study one-dimensional discrete Schrödinger operators given by:

$$(1) \quad (Hu)(k) = u(k+1) + u(k-1) + V(k)u(k), \quad k \in \mathbb{Z},$$

where $V = \{V(k)\}$ is a bounded potential.

We focus on two types of potentials: almost reflective repetition potentials and almost repetition potentials. These potentials are closely related to those studied in the context of quasi-periodic Schrödinger operators.

A potential $V = \{V(k)\}_{k \in \mathbb{Z}}$ is said to have γ -repetitions for some $\gamma > 0$ (a modification of the definition in [24]) if there exists an increasing sequence $k_n \rightarrow \infty$ or $k_n \rightarrow -\infty$ such that, for all $k \in \mathbb{Z}$:

$$(2) \quad |V(k_n + k) - V(k)| \leq e^{-\gamma|k_n|}.$$

Similarly, a potential $V = \{V(k)\}_{k \in \mathbb{Z}}$ is said to have γ -reflective repetitions if there exists a sequence $k_n \rightarrow \infty$ or $k_n \rightarrow -\infty$ such that, for all $k \in \mathbb{Z}$:

$$(3) \quad |V(k_n - k) - V(k)| \leq e^{-\gamma|k_n|}.$$

Potentials with repetitions are often referred to as *Gordon-type potentials*, while reflective repetition potentials are also known as *palindromic potentials*.

We define one-step transfer matrices

$$A_k(E) := \begin{pmatrix} E - V(k) & -1 \\ 1 & 0 \end{pmatrix}$$

and the multi-step transfer matrices by

$$A_{k,j}(E) := \begin{cases} A_{k-1}(E)A_{k-2}(E) \cdots A_j(E), & j < k; \\ A_{k,j} = I, & k = j; \\ A_{j,k}^{-1}(E), & j > k. \end{cases}$$

Define

$$(4) \quad \mathcal{L}(E) := \limsup_{k \rightarrow \infty} \sup_{j \in \mathbb{Z}} \frac{\ln \|A_{k+j,j}(E)\|}{k}.$$

It is clear that for bounded V , we have $\mathcal{L}(E) < \infty$ for every E .

The foundational work of Gordon [12] and Simon [27], commonly referred to as the Gordon-type argument, laid the groundwork for proving that for γ -repetition potentials, H has no eigenvalues in the regime

$$\{E \in \mathbb{C} : \gamma > 2\mathcal{L}(E)\}.$$

Although Gordon and Simon did not explicitly prove this exact result, it can be established by carefully analyzing their proofs. Gordon first presented his argument in the 1970s in Russian [12], but it remained largely unfamiliar to the broader mathematical community until Barry Simon provided a detailed discussion of the Gordon argument in his 1982 article [27] (also consult with [2]). For additional historical context, see [21].

Recently, Avila-You-Zhou [1] improved Gordon type arguments, extending the regime to

$$\{E \in \mathbb{C} : \gamma > \mathcal{L}(E)\}.$$

The absence of eigenvalues in the regime $\{E \in \mathbb{C} : \gamma > \mathcal{L}(E)\}$ is sharp, as demonstrated by the example of almost Mathieu operators (see the definition later) [1, 14, 18].

Over the past decades, Gordon-type arguments have been widely applied to the spectral theory of various models [5–7, 9, 10, 15–17, 23, 25, 26, 28, 29].

For γ -reflective repetition potentials, Jitomirskaya and Simon [22] proved that H has no eigenvalues in the regime

$$(5) \quad \{E \in \mathbb{C} : \gamma > 2\mathcal{L}(E)\}.$$

Recently, Jitomirskaya and the author improved the threshold, reducing the numerical number in (5) from 2 to 1 [19]. This result is also sharp, as evidenced by the almost Mathieu operator [19] (also see Ge-You-Zhou [11]). The Jitomirskaya-Simon argument has been commonly used to study Schrödinger operators with reflective repetition potentials, e.g., [4, 8, 13, 20].

In the present paper, we build on the approaches developed by Jitomirskaya-Simon [22] and Jitomirskaya-Liu [19] to study almost reflective repetition potentials. Our methods provide a new proof of the absence of eigenvalues for almost repetition potentials.

Theorem 1.1. *Assume $\gamma > \mathcal{L}(E)$. Then the eigen-equation $Hu = Eu$ has no $\ell^2(\mathbb{Z})$ solutions.*

Consider the (discrete) quasi-periodic Schrödinger operator on $\ell^2(\mathbb{Z})$:

$$(6) \quad (H_{v,\alpha,\theta}u)(k) = u(k+1) + u(k-1) + v(\theta + k\alpha)u(k), \quad k \in \mathbb{Z},$$

where $v : \mathbb{T} = \mathbb{R}/\mathbb{Z} \rightarrow \mathbb{R}$ is a continuous potential, α is the frequency, and θ is the phase.

Let $L(E)$ denote the Lyapunov exponent of the operator (6):

$$(7) \quad L(E) = \limsup_{j \rightarrow \infty} \frac{1}{j} \int_{\mathbb{T}} \ln \|T_{j,0}(E)\| d\theta = \lim_{j \rightarrow \infty} \frac{1}{j} \int_{\mathbb{T}} \ln \|T_{j,0}(E)\| d\theta.$$

For quasi-periodic Schrödinger operators with continuous potentials, it is known that $L(E) = \mathcal{L}(E)$ (independent of θ).

Define

$$\beta(\alpha) = \limsup_{k \rightarrow \infty} \frac{-\ln \|k\alpha\|_{\mathbb{T}}}{k},$$

where $\|x\|_{\mathbb{T}} = \text{dist}(x, \mathbb{Z})$.

Corollary 1.2. *Assume v is κ -Hölder continuous. Then H has no eigenvalues in the regime $\{E \in \mathbb{C} : L(E) < \kappa\beta(\alpha)\}$.*

By setting $v(\theta) = 2\lambda \cos(2\pi\theta)$ in (6), we obtain the almost Mathieu operator $H_{\lambda,\alpha,\theta}$. For the almost Mathieu operator, the Lyapunov exponent on the spectrum is given by $L(E) = \max\{\ln |\lambda|, 0\}$ [3].

As a consequence, Corollary 1.2 implies:

Corollary 1.3. *Assume α satisfies $\beta(\alpha) > 0$. For the almost Mathieu operator, $H_{\lambda,\alpha,\theta}$ has no eigenvalues when $|\lambda| < e^{\beta(\alpha)}$.*

Remark 1. *As mentioned earlier, Corollary 1.3 was previously proved using an improved Gordon-type argument [1].*

2. PROOF OF THEOREM 1.1

Without loss of generality, assume that $k_n \rightarrow \infty$. The proof of the case $k_n \rightarrow -\infty$ is similar. Since the spectrum of $H = \Delta + V$ is bounded, we can assume E is bounded. By the definition of $\mathcal{L}(E)$, one has that

$$(8) \quad \|T_{m,j}(E)\| \leq C e^{(\mathcal{L}(E)+\varepsilon)|m-j|},$$

where the constant C depends on ε and the potential V .

Lemma 2.1. *For all $m, j \in \mathbb{Z}$, we have*

$$\|T_{m,j}(E) - T_{m+k_n,j+k_n}(E)\| \leq C e^{-\gamma k_n} e^{(\mathcal{L}(E)+\varepsilon)|m-j|}.$$

Proof. By (8), the proof of Lemma 2.1 follows from the standard telescoping. See [28, Lemma 4.5] for example. \square

Proof of Theorem 1.1 . We prove it by contradiction. Suppose there exists E with $\mathcal{L}(E) < \gamma$ such that the eigen-equation $Hu = Eu$ has an $\ell^2(\mathbb{Z})$ solution u . Assume u is normalized, namely

$$\|u\|_{\ell^2} = \sum_{k \in \mathbb{Z}} |u(k)|^2 = 1.$$

For each $n \in \mathbb{Z}_+$, define sequences $\{u_n(k)\}_{k \in \mathbb{Z}}$ and $\{V_n(k)\}_{k \in \mathbb{Z}}$ as $u_n(k) = u(k_n + k)$ and $V_n(k) = V(k_n + k)$, $k \in \mathbb{Z}$. Then by (2), one has that

$$(9) \quad |V(k) - V_n(k)| \leq e^{-\gamma k_n}, \text{ for all } k \in \mathbb{Z}.$$

We also have

$$(10) \quad u(k+1) + u(k-1) + V(k)u(k) = Eu(k)$$

and

$$(11) \quad u_n(k+1) + u_n(k-1) + V_n(k)u_n(k) = Eu_n(k).$$

Let $W(k) = W(f, g) = f(k+1)g(k) - f(k)g(k+1)$ be the Wronskian.

By a standard calculation using (9), (10) and (11), we have

$$\begin{aligned} |W(u, u_n)(k) - W(u, u_n)(k-1)| &\leq |V(k) - V_n(k)| |u(k)u_n(k)| \\ &\leq e^{-\gamma k_n} |u(k)u_n(k)|. \end{aligned}$$

This implies for any $m \geq 0$ and k ,

$$\begin{aligned} |W(u, u_n)(k+m) - W(u, u_n)(k-1)| &\leq e^{-\gamma k_n} \sum_{j=0}^m |u(k+j)u_n(k+j)| \\ (12) \quad &\leq e^{-\gamma k_n}, \end{aligned}$$

where the second inequality holds by the Cauchy–Schwarz inequality and the fact that $\|u\|_{\ell^2} = \|u_n\|_{\ell^2} = 1$.

Notice that $\sum_k |W(u, u_n)(k)| \leq 2$. Therefore,

$$(13) \quad \lim_{k \rightarrow \infty} |W(u, u_n)(k)| = 0.$$

By (12) and (13), we must have that for all $k \in \mathbb{Z}$,

$$(14) \quad |W(u, u_n)(k)| \leq e^{-\gamma k_n}.$$

Letting $k = -1$ in (14), one has that

$$(15) \quad |W(u, u_n)(-1)| = \left| \det \begin{pmatrix} u(0) & u(k_n) \\ u(-1) & u(k_n - 1) \end{pmatrix} \right| \leq e^{-\gamma k_n}.$$

Write $\begin{pmatrix} u(k_n) \\ u(k_n - 1) \end{pmatrix}$ as the linear combination of $\begin{pmatrix} u(0) \\ u(-1) \end{pmatrix}$ and $\begin{pmatrix} -u(-1) \\ u(0) \end{pmatrix}$, namely

$$(16) \quad \begin{pmatrix} u(k_n) \\ u(k_n - 1) \end{pmatrix} = a_n \begin{pmatrix} u(0) \\ u(-1) \end{pmatrix} + b_n \begin{pmatrix} -u(-1) \\ u(0) \end{pmatrix}.$$

Since u is non-trivial, one has that

$$(17) \quad |u(0)|^2 + |u(-1)|^2 > 0.$$

By (15), (16) and (17), we have that

$$(18) \quad b_n = O(e^{-\gamma k_n}).$$

Since $u \in \ell^2(\mathbb{Z})$, one has that

$$(19) \quad a_n = o(1),$$

as $n \rightarrow \infty$. By the definition of transfer matrices, one has that

$$(20) \quad \begin{pmatrix} u(k) \\ u(k-1) \end{pmatrix} = T_{k,m}(E) \begin{pmatrix} u(m) \\ u(m-1) \end{pmatrix}.$$

Multiplying T_{0,k_n} on both sides of (16), we conclude that

$$(21) \quad \begin{pmatrix} u(0) \\ u(-1) \end{pmatrix} = a_n T_{0,k_n}(E) \begin{pmatrix} u(0) \\ u(-1) \end{pmatrix} + b_n T_{0,k_n}(E) \begin{pmatrix} -u(-1) \\ u(0) \end{pmatrix}$$

$$(22) \quad = a_n T_{0,k_n}(E) \begin{pmatrix} u(0) \\ u(-1) \end{pmatrix} + O(e^{-(\gamma - \mathcal{L}(E) - \varepsilon)k_n})$$

$$(23) \quad = a_n T_{-k_n,0}(E) \begin{pmatrix} u(0) \\ u(-1) \end{pmatrix} + O(e^{-(\gamma - \mathcal{L}(E) - \varepsilon)k_n})$$

$$(24) \quad = a_n \begin{pmatrix} u(-k_n) \\ u(-k_n - 1) \end{pmatrix} + O(e^{-(\gamma - \mathcal{L}(E) - \varepsilon)k_n}),$$

where (21) and (24) hold by (20), (22) holds by (8) and (18), and (23) holds by Lemma 2.1. The equality (24) can not hold since $\left\| \begin{pmatrix} u(-k_n) \\ u(-k_n - 1) \end{pmatrix} \right\| = o(1)$ and $a_n = o(1)$ as $n \rightarrow \infty$. We finish the proof. \square

ACKNOWLEDGMENTS

W. Liu was a 2024-2025 Simons fellow. W. Liu was supported by NSF DMS-2246031 and DMS-2052572.

REFERENCES

- [1] A. Avila, J. You, and Q. Zhou. Sharp phase transitions for the almost Mathieu operator. *Duke Math. J.*, 166(14):2697–2718, 2017.
- [2] J. Avron and B. Simon. Singular continuous spectrum for a class of almost periodic Jacobi matrices. *Bull. Amer. Math. Soc. (N.S.)*, 6(1):81–85, 1982.
- [3] J. Bourgain and S. Jitomirskaya. Continuity of the Lyapunov exponent for quasiperiodic operators with analytic potential. volume 108, pages 1203–1218. 2002. Dedicated to David Ruelle and Yasha Sinai on the occasion of their 65th birthdays.
- [4] C. Cedzich and J. Fillman. Absence of bound states for quantum walks and CMV matrices via reflections. *J. Spectr. Theory*, 14(4):1513–1536, 2024.
- [5] D. Damanik. Gordon-type arguments in the spectral theory of one-dimensional quasicrystals. In *Directions in mathematical quasicrystals*, volume 13 of *CRM Monogr. Ser.*, pages 277–305. Amer. Math. Soc., Providence, RI, 2000.
- [6] D. Damanik. Absence of eigenvalues for a class of Schrödinger operators on the strip. *Forum Math.*, 14(5):797–806, 2002.
- [7] D. Damanik. A version of Gordon’s theorem for multi-dimensional Schrödinger operators. *Trans. Amer. Math. Soc.*, 356(2):495–507, 2004.
- [8] D. Damanik and R. Killip. Reflection symmetries of almost periodic functions. *J. Funct. Anal.*, 178(2):251–257, 2000.
- [9] L. Fang, S. Guo, Y. Peng, and F. Wang. Quantitative version of Gordon’s lemma for Hamiltonian with finite range. *Linear Algebra Appl.*, 687:91–107, 2024.
- [10] J. Fillman. Purely singular continuous spectrum for Sturmian CMV matrices via strengthened Gordon lemmas. *Proc. Amer. Math. Soc.*, 145(1):225–239, 2017.
- [11] L. Ge, J. You, and Q. Zhou. Structured quantitative almost reducibility and its applications. *arXiv preprint arXiv:2407.05490*, 2024.
- [12] A. Y. Gordon. The point spectrum of the one-dimensional Schrödinger operator. *Uspekhi Matematicheskikh Nauk*, 31(4):257–258, 1976.
- [13] A. Hof, O. Knill, and B. Simon. Singular continuous spectrum for palindromic Schrödinger operators. *Comm. Math. Phys.*, 174(1):149–159, 1995.
- [14] S. Jitomirskaya and I. Kachkovskiy. L^2 -reducibility and localization for quasiperiodic operators. *Math. Res. Lett.*, 23(2):431–444, 2016.
- [15] S. Jitomirskaya and I. Kachkovskiy. Sharp arithmetic delocalization for quasiperiodic operators with potentials of semi-bounded variation. *arXiv preprint arXiv:2408.16935*, 2024.
- [16] S. Jitomirskaya and S. Kocić. Spectral theory of Schrödinger operators over circle diffeomorphisms. *Int. Math. Res. Not. IMRN*, (13):9810–9829, 2022.

- [17] S. Jitomirskaya and W. Liu. Arithmetic spectral transitions for the Maryland model. *Comm. Pure Appl. Math.*, 70(6):1025–1051, 2017.
- [18] S. Jitomirskaya and W. Liu. Universal hierarchical structure of quasiperiodic eigenfunctions. *Ann. of Math. (2)*, 187(3):721–776, 2018.
- [19] S. Jitomirskaya and W. Liu. Universal reflective-hierarchical structure of quasiperiodic eigenfunctions and sharp spectral transition in phase. *J. Eur. Math. Soc. (JEMS)*, 26(8):2797–2836, 2024.
- [20] S. Jitomirskaya, W. Liu, and L. Mi. Sharp palindromic criterion for semi-uniform dynamical localization. *arXiv preprint arXiv:2410.21700*, 2024.
- [21] S. Jitomirskaya, S. A. Molchanov, B. Simon, and B. R. Vainberg. Alexander Gordon. *J. Spectr. Theory*, 9(4):1157–1164, 2019.
- [22] S. Jitomirskaya and B. Simon. Operators with singular continuous spectrum. III. Almost periodic Schrödinger operators. *Comm. Math. Phys.*, 165(1):201–205, 1994.
- [23] S. Jitomirskaya and F. Yang. Singular continuous spectrum for singular potentials. *Comm. Math. Phys.*, 351(3):1127–1135, 2017.
- [24] S. Jitomirskaya and S. Zhang. Quantitative continuity of singular continuous spectral measures and arithmetic criteria for quasiperiodic Schrödinger operators. *J. Eur. Math. Soc. (JEMS)*, 24(5):1723–1767, 2022.
- [25] W. Liu. Small denominators and large numerators of quasiperiodic Schrödinger operators. *Peking Mathematical Journal to appear*.
- [26] W. Liu. Continuous quasiperiodic Schrödinger operators with Gordon type potentials. *J. Math. Phys.*, 59(6):063501, 6, 2018.
- [27] B. Simon. Almost periodic Schrödinger operators: a review. *Advances in Applied Mathematics*, 3(4):463–490, 1982.
- [28] B. Simon. Almost periodic Schrödinger operators. IV. The Maryland model. *Ann. Physics*, 159(1):157–183, 1985.
- [29] J. You, S. Zhang, and Q. Zhou. Point spectrum for quasi-periodic long range operators. *J. Spectr. Theory*, 4(4):769–781, 2014.

(W. Liu) DEPARTMENT OF MATHEMATICS, TEXAS A&M UNIVERSITY, COLLEGE STATION, TX 77843-3368, USA

Email address: liuwencai1226@gmail.com; wencail@tamu.edu