Extrinsic Mechanisms of Phonon Magnetic Moment

Rui Xue,^{1, 2} Zhenhua Qiao,^{1, 2, *} Yang Gao,^{1, 2, [†](#page-0-1)} and Qian Niu^{1, 2}

¹International Centre for Quantum Design of Functional Materials,

CAS Key Laboratory of Strongly-Coupled Quantum Matter Physics, and Department of Physics,

University of Science and Technology of China, Hefei, Anhui 230026, China

²Hefei National Laboratory, University of Science and Technology of China, Hefei 230088, China

(Dated: January 7, 2025)

We develop a general formalism of phonon magnetic moment by including the relaxation process. We then identify the skew-scattering and side-jump contributions to the phonon magnetic moment originating from the non-adiabaticity, both of which are related to the nonlocal phonon Berry curvature and are in close analogy to those in the electronic Hall effect. All contributions of the phonon magnetic moment are exemplified in a honeycomb lattice, showing that the extrinsic contribution can be as important as the intrinsic one and that the resulting phonon angular momentum varies significantly across the Brillouin zone. Our work offers a systematic framework of the phonon chirality and paves the way of tuning the phonon magnetic moment through the non-adiabaticity.

Recent years have seen a surge of interest in phonon chirality [\[1](#page-4-0)[–15\]](#page-4-1). As a unique degree of freedom characterizing the circular motion of phonons, phonon chirality can couple to the spin and valley degree of freedom of electrons and hence plays important roles in various quantum phenomena, such as exciton formation and relaxation [\[5,](#page-4-2) [15,](#page-4-1) [16\]](#page-4-3), magnetic dynamics [\[17,](#page-4-4) [18\]](#page-4-5), unconventional superconductivity [\[13\]](#page-4-6), etc.. As phonon modes can generally carry Born effective charges, their circular motion naturally leads to a nontrivial magnetic moment, which not only directly couples to the external magnetic field through the Zeeman effect [\[2,](#page-4-7) [4,](#page-4-8) [10\]](#page-4-9) but also is responsible for the phonon Hall effect [\[10,](#page-4-9) [19–](#page-4-10)[24\]](#page-4-11), a fundamental response of phonons. Under the adiabatic Born-Oppenheimer approximation, the phonon magnetic moment has been related to the Berry curvature of electrons in the parameter space characterizing the motion of nucleus, and hence favours topological systems [\[3,](#page-4-12) [4,](#page-4-8) [6–](#page-4-13) [9,](#page-4-14) [14\]](#page-4-15).

As successful as it is, the current formulation of the phonon magnetic moment does not take into account the relaxation process, which is essential in understanding the chiral response of electrons [\[25–](#page-4-16)[30\]](#page-4-17), such as the anomalous Hall effect, spin Hall effect, etc.. Its basis, i.e., the Born-Oppenheimer approximation, relies on the fact that the electron mass is much smaller than that of atoms, such that the characteristic energy scale of the electronic motion to be much larger than that of the atomic motion. However, treating the relaxation process introduces non-Hermiticity by the addition of extra energy scales to the phononic and electronic systems, originating from either externally controlled gain and loss or the internal correlation effects [\[27,](#page-4-18) [28,](#page-4-19) [31–](#page-4-20)[42\]](#page-5-0). As these energy scales do not necessarily fulfill the requirement of

the Born-Oppenheimer approximation, it is thus highly desirable to unravel their effect on the phonon magnetic moment.

In this work, we present a microscopic theory of phonon magnetic moment using the time-dependent density-matrix perturbation theory. The relaxation process is added through effective lifetime parameters to the phonon frequency and electronic Hamiltonian. A straightforward modification to the phonon magnetic moment is that the electron band gap is renormalized by the phonon frequency. As a result, the leading order correction is proportional to the ratio of the phonon frequency and the electron energy gap, as anticipated from the adiabatic approximation.

Strikingly, we also identify the skew-scattering and side-jump contributions to the phonon magnetic moment. Both vanish when the adiabaticity is restored in the relaxation process and depend on the ratio of different lifetimes otherwise. Microscopically, they originate from the molecular Berry curvature which characterizes the strength of the chiral-phonon-chiral-electron coupling. The skew-scattering and side-jump contribution is further due to the change of the magnetic moment and coordinate in the phonon-electron interaction, respectively, in analogy to those in the electronic Hall effect [\[25\]](#page-4-16). Our theory is exemplified in the honeycomb lattice, showing that the extrinsic contribution can be as important as the intrinsic one and that it can induce a sizable phonon angular momentum.

Microscopic theory.— The phonon magnetic moment m manifests as a Zeeman-type energy shift under a magnetic field \mathbf{B} : $\hat{H}_Z = -\mathbf{B} \cdot \mathbf{m}$. It is proportional to the circular atomic motion [\[3,](#page-4-12) [24\]](#page-4-11):

$$
m_i = \lambda_{ij} (\mathbf{u}^a \times \dot{\mathbf{u}}^a)_j , \qquad (1)
$$

where λ_{ij} is the response coefficient and u^a is the atomic displacement at lattice site a. Therefore, H_Z is the Raman term in the phonon Hamiltonian and is essential for the phonon Hall effect.

[∗]Correspondence author: qiao@ustc.edu.cn

[†]Correspondence author: ygao87@ustc.edu.cn

To derive **m** or equivalently λ_{ij} , we observe that the Raman term is quadratic in u and linear in B . We then consider the electronic Hamiltonian involving the electron-phonon coupling and the magnetic field, and expand it up to the order of interest. The result reads (for simplicity, we set $e = \hbar = 1$)

$$
\hat{H} = \hat{H}_0 + \hat{H}_B + \hat{H}_{ep} + \hat{H}_{mix},\tag{2}
$$

where H_0 is the bare electronic Hamiltonian, $\hat{H}_B =$ $\hat{v} \cdot A$ with A being the vector potential due to external magnetic field, $\hat{H}_{ep} = \hat{F}_i^a u_i^a + \hat{F}_{ij}^{ab} u_i^a u_j^b$ describes the electron-phonon coupling, $\hat{F}_i^a = \frac{\partial \hat{H}_0}{\partial R_i^a} (R_i^a)$ is the equilibrium atomic position at lattice site a), $\hat{F}^{ab}_{ij} =$ $(1/2) \partial^2 \hat{H}_0 / \partial R^a_i \partial R^b_j$, $\hat{H}_{mix} = \hat{F}^a_{i,j} u^a_i A_j + \hat{F}^{ab}_{ij,k} u^a_i u^b_j A_k$ describes the coupling between the electron-phonon interaction and the magnetic field, $\hat{F}_{i,j}^a = \partial \hat{v}_j / \partial R_i^a$, and $\hat{F}^{ab}_{ij,k} = (1/2)\partial^2 \hat{v}_k / \partial R_i^a \partial R_j^b$. Here and hereafter, Einstein summation convention is implied for repeated indices. Without loss of generality, we will consider the part of electron-phonon coupling with $a = b$, i.e., the onsite coupling. The generalization to $a \neq b$ is straightforward.

To proceed, we consider a single phonon mode: $u_i \rightarrow$ $u_i(\omega, \boldsymbol{q})e^{-i\omega t + i\boldsymbol{q}\cdot \boldsymbol{r}}$. Then $\boldsymbol{u} \times \dot{\boldsymbol{u}} \to \boldsymbol{u}(\omega, \boldsymbol{q}) \times \boldsymbol{u}(\omega, \boldsymbol{q})^*$. The magnetic field can be included by allowing the vector potential \boldsymbol{A} varies in the form of $\boldsymbol{A} = \boldsymbol{A}(\boldsymbol{q}_0)e^{i\boldsymbol{q}_0 \cdot \boldsymbol{r}}$ so that $\mathbf{B} = iq_0 \times \mathbf{A}$. The coefficient $\lambda_{ij}(\omega, \mathbf{q})$ can then be derived by integrating out the electronic degree of freedom in the full Hamiltonian in Eq. [\(2\)](#page-1-0). We note that the second terms in both \hat{H}_{ep} and \hat{H}_{mix} do not contribute to λ_{ij} , as they are symmetric with respect to u_i and u_j . For the other terms in Eq. [\(2\)](#page-1-0), we seek the following response functions

$$
\langle v_k e^{iq_0 r} \rangle = \theta_{kij}^1(\omega, \mathbf{q}, \mathbf{q}_0) u_i(\omega, \mathbf{q}) u_j(-\omega, -\mathbf{q} - \mathbf{q}_0),
$$

$$
\langle \hat{F}_i^a e^{i\mathbf{q} \cdot \mathbf{R}^a} \rangle = \theta_{kij}^2(\omega, \mathbf{q}, \mathbf{q}_0) u_j(-\omega, -\mathbf{q} - \mathbf{q}_0) A_k(\mathbf{q}_0),
$$

$$
\langle \hat{F}_{i,k}^a e^{i(\mathbf{q} + \mathbf{q}_0) \cdot \mathbf{R}^a} \rangle = \theta_{kij}^3(\omega, \mathbf{q}, \mathbf{q}_0) u_j(-\omega, -\mathbf{q} - \mathbf{q}_0).
$$
 (3)

Combining with the Zeeman energy shift and Eq. [\(1\)](#page-0-2), λ_{ij} can then be expressed as follows [\[43\]](#page-5-1):

$$
\lambda_{ij}(\omega, \boldsymbol{q}) = \frac{1}{4} \epsilon_{j\ell k} \epsilon_{ik_1k_2} \lim_{\boldsymbol{q}_0 \to 0} \partial_{q_{0\ell}} \partial_{\omega} \sum_{i=1}^3 [\theta^i_{kk_1k_2}(\omega, \boldsymbol{q}, \boldsymbol{q}_0) + \theta^i_{kk_2k_1}(-\omega, -\boldsymbol{q}, \boldsymbol{q}_0)]. \tag{4}
$$

The response functions in Eq. [\(3\)](#page-1-1) can be derived using the density-matrix perturbation theory. The density matrix $\hat{\rho}$ satisfies the following equation [\[36\]](#page-4-21):

$$
\dot{\hat{\rho}} = -i(\hat{\rho}\hat{H} - \hat{H}^{\dagger}\hat{\rho}).
$$
\n(5)

Originally $\hat{H} = \hat{H}^{\dagger}$ as in Eq. [\(3\)](#page-1-1). To include the relaxation process, we add non-Hermitian self-energy part to \hat{H} which manifests as an imaginary frequency in the phonon degree of freedom, i.e., $\omega \rightarrow \omega + i\eta_p$, and an imaginary self-energy Σ in the electron degree of freedom. Without loss of generality, we further assume that both η_p and Σ are constant, and that Σ is band-diagonal [\[27\]](#page-4-18), i.e., $\Sigma = i\eta_m \delta_{mn}$ with m being band index.

We then solve Eq. [\(5\)](#page-1-2) order-by-order by taking $u(\omega, \mathbf{q})$ and $\mathbf{A}(\mathbf{q}_0)$ as small quantities, yielding $\hat{\rho} = \hat{\rho}_0 + \hat{\rho}_1 + \hat{\rho}_2$ $\hat{\rho}_2 + \cdots$. The response function θ_{kij}^1 and θ_{kij}^2 can be

derived using $\hat{\rho}_2$ and θ_{kij}^3 using $\hat{\rho}_1$. The phonon magnetic moment, or λ_{ij} can then be obtained using Eq. [\(4\)](#page-1-3).

Interestinly, we identify two distinct types of terms in λ_{ij} :

$$
\lambda_{ij}(\omega, \mathbf{q}) = \lambda_{ij}^A(\omega, \mathbf{q}) + \lambda_{ij}^{NA}(\omega, \mathbf{q}). \tag{6}
$$

The first part λ_{ij}^A can be smoothly reduced to the phonon magnetic moment under the Born-Oppenheimer approximation in semiconductors and insulators [\[43\]](#page-5-1). Its leading order correction is in the form of ω/ε_0 where ε_0 is the energy scale of the electronic band, in accordance with the adiabatic approximation.

In sharp contrast, the second term λ_{ij}^{NA} originates from non-adiabaticity and reads as:

$$
\lambda_{ij}^{NA} = -\frac{1}{2} \sum_{n_1 n_3} \int d\mathbf{k} \left[\delta_{n_1}^{n_3} (m_j^e)_{n_1 \mathbf{k}} G_1(\omega, \mathbf{q}) (\Omega_i)_{n_1 n_1}^{n_2} (\mathbf{q}) + \epsilon_{j j_1 j_2} \delta_{n_1}^{n_3} (v_{j_2})_{n_1 \mathbf{k}} G_2(\omega, \mathbf{q}) T(\mathbf{q}) - (\omega, \mathbf{q} \to -\omega, -\mathbf{q}) \right], \tag{7}
$$

where
$$
\delta_{n_1}^{n_3} = \eta_{n_3} / (2\eta_p + \eta_{n_1}), \, \boldsymbol{m}_{n\boldsymbol{k}}^e = 1/2\mathrm{Im}\langle \boldsymbol{\partial} n\boldsymbol{k} | \times \hat{\boldsymbol{v}} | n\boldsymbol{k} \rangle
$$

is the electronic orbital magnetic moment in band n, $|n\mathbf{k}\rangle$

FIG. 1: Extrinsic contributions to the phonon magnetic moment. (a) Skew-scattering mechanism. (b) Side-jump Mechanism.

is the periodic part of the electronic Bloch wave function with energy $\varepsilon_{n\mathbf{k}}$ and distribution function $f_{n\mathbf{k}}$, $(v_{j_2})_{n_1\mathbf{k}} =$ $\langle n_1\bm{k}|\hat{v}_{j_2}|n_1\bm{k}\rangle,$

$$
G_{1}(\omega, \mathbf{q}) = \frac{\Delta f_{n_{1}\mathbf{k}, n_{2}\mathbf{k} - \mathbf{q}} \Delta \varepsilon_{n_{1}\mathbf{k}, n_{2}\mathbf{k} - \mathbf{q}}^{2}}{(-\omega + \Delta \varepsilon_{n_{1}\mathbf{k}, n_{2}\mathbf{k} - \mathbf{q}})^{3}},
$$

$$
(\Omega_{i})_{n_{1}n_{3}}^{n_{2}}(\mathbf{q}) = \epsilon_{ii_{1}i_{2}} \frac{\left(\frac{\partial \hat{H}}{\partial u_{i_{1},\mathbf{q}}}\right)_{n_{1}\mathbf{k}, n_{2}\mathbf{k} - \mathbf{q}}\left(\frac{\partial \hat{H}}{\partial u_{i_{2},\mathbf{q}}^{*}}\right)_{n_{2}\mathbf{k} - \mathbf{q}, n_{3}\mathbf{k}}}{(\varepsilon_{n_{1},\mathbf{k}} - \varepsilon_{n_{2},\mathbf{k} - \mathbf{q}})(\varepsilon_{n_{3},\mathbf{k}} - \varepsilon_{n_{2},\mathbf{k} - \mathbf{q}})},
$$
(8)

 $G_2 = G_1 \Delta \varepsilon_{n_3 k, n_2 k-q} / \Delta \varepsilon_{n_1 k, n_2 k-q}, \ \ \Delta f_{n_1 k, n_2 k-q} =$ $f_{n_1,k} - f_{n_2,k-q}, \ \ \Delta \varepsilon_{n_1k,n_2k-q} = \varepsilon_{n_1,k} - \varepsilon_{n_2,k-q},$ $(\partial \hat{H}/\partial u_{i_1,\boldsymbol{q}})_{n_1\boldsymbol{k},n_2\boldsymbol{k}-\boldsymbol{q}} \quad = \quad \langle n_1\boldsymbol{k}|\partial \hat{H}/\partial u_{i_1,\boldsymbol{q}}|n_2\boldsymbol{k} \; - \; \boldsymbol{q}\rangle,$ and $T(\boldsymbol{q}) = \text{Im} \sum_{n_3 \neq n_1}^{\bullet} (\Omega_i)_{n_1n_3}^{n_2}(\boldsymbol{q}) (A_{j_1})_{n_3 \boldsymbol{k}, n_1 \boldsymbol{k}}$ with $(A_{j_1})_{n_3\mathbf{k},n_1\mathbf{k}} = \langle n_3\mathbf{k}|i\partial_{j_2}|n_1\mathbf{k}\rangle$ being the interband Berry connection.

Equation [\(7\)](#page-1-4) is our main result and it does not have any counterpart in the adiabatic perturbation theory. It has several striking features. First, it is closely related to the relaxation process through the prefactor $\delta_{n_1}^{n_3}$. Interestingly, it relies not on the absolute strength of the relaxation but on the relative ratio of the electronic and phononic lifetime. In the limit that $\eta_p \gg \eta_{n_i}, \delta_{n_1}^{n_3} \to 0$, so that $\lambda_{ij}^{NA} \rightarrow 0$, and only λ_{ij}^{A} contributes. The nonadiabatic correction emerges with η_p either comparable with or dominated by certain η_{n_i} .

This observation sheds light on the nature of adiabaticity. On one hand, for λ_{ij}^A , the correction to the adiabatic result starts at the order of ω/ε_0 , which can be safely ignored if the phonon dynamics is much slower than the electron dynamics, so that the electron motion can follow the atomic one. On the other hand, the life time or the energy exchange rate of electrons shall be much slower than that of phonons, so that the distribution of electronic state is unaffected. Therefore, there are two conditions on the adiabaticity instead of the dynamic condition only. When the second condition is broken, λ_{ij}^{NA} emerges.

Secondly, Eq. [\(7\)](#page-1-4) depends on an essential geometric quantity $(\Omega_i)_{n1n3}^{n_2}(\boldsymbol{q})$, which is the interband molecular Berry curvature. For the diagonal element with n_1 = $n_3, \sum_{n_2 \neq n_1} (\Omega_i)_{n1n1}^{n_2}(\boldsymbol{q})$ recovers the full molecular Berry curvature introduced previously [\[3,](#page-4-12) [6,](#page-4-13) [9\]](#page-4-14). Specifically, for

two-band systems, $(\Omega_i)_{n_1n_1}^{n_2}(\mathbf{q})$ coincides with molecular Berry curvature.

The physical meaning of Eq. [\(7\)](#page-1-4) can be well understood with the help of the modern theory of electronic orbital magnetization M [\[44–](#page-5-2)[47\]](#page-5-3). We start from the following expression of M :

$$
\mathbf{M} = \int \frac{d\mathbf{k}}{8\pi^3} (\mathbf{m}_{n\mathbf{k}} f_{n\mathbf{k}} + \mathbf{v}_{n\mathbf{k}} \times \mathbf{A}_{n\mathbf{k},n\mathbf{k}} f_{n\mathbf{k}}). \tag{9}
$$

The first terms in both Eq. [\(7\)](#page-1-4) and Eq. [\(9\)](#page-2-0) contain the orbital magnetic moment m . Therefore, the first term in Eq. [\(7\)](#page-1-4) orignates from the correction to the distribution function due to the electron-phonon coupling. The appearance of the band-resolved Berry curvature is thus natural, as the transition rate induced by a circular phonon mode is proportional to $[u(\omega, \boldsymbol{q}) \times u^*(\omega, \boldsymbol{q})]$. $(\Omega_i)_{n1n1}^{n_2}(\boldsymbol{q}).$

More precisely, the first term in Eq. [\(7\)](#page-1-4) represents a skew-scattering contribution to the phonon magnetic moment. To see this, we note that in non-magnetic materials, there is equal number of states with positive and negative magnetic moment. Typical examples are transitional metal dichalcogenides, where m takes opposite signs in K and K' valley. On the other hand, the scattering of those electrons by chiral phonons depends on the band-resolved molecular Berry curvature, which shares the same symmetry property with the magnetic moment. Therefore, states with positive and negative magnetic moment are scattered differently by the chiral phonon, as illustrated in Fig. $1(a)$, leading to a net magnetic moment given by the first term in Eq. [\(7\)](#page-1-4). This is in close analogy to the skew-scattering mechanism of the anomalous Hall effect [\[25\]](#page-4-16), in which electrons with opposite spin polarization are scattered differently by impurities.

By similar logic, the second term in Eq. [\(7\)](#page-1-4) is a side-jump contribution to the phonon magnetic moment. First, it shares a similar structure with the second term in Eq. [\(9\)](#page-2-0). We then observe that $G_2(\omega, \mathbf{q})T(\mathbf{q})$ contains the off-diagonal part of both the electron-phonon coupling (i.e., $u(\omega, \mathbf{q}) \times u^*(\omega, \mathbf{q}) \cdot \Omega_{n1n3}^{n2}(\mathbf{q})$) and the position operator (i.e., A_{n_3k,n_1k}). It thus represents the shift of the electronic coordinate during the electron-chiralphonon coupling. If such coordinate shift is perpendicular to the group velocity of the electron, a net magnetic moment can then emerge, as illustrated by Fig. [1\(](#page-2-1)b). This closely resembles the side-jump mechanism of the anomalous Hall effect [\[25,](#page-4-16) [26\]](#page-4-22), where electrons change its center-of-mass position after scattered by impurities.

Lattice model. \leftarrow As a concrete example, we consider phonons in a honeycomb lattice. The electronic Hamiltonian contains the nearest-neighbour hopping and staggered potential: $\hat{H}_{el} = \sum_{\langle ij \rangle} t c_i^{\dagger} c_j + \sum_i \Delta c_i^{\dagger} c_i$. The electron-phonon coupling is added by allowing the hopping paramter t varies with the bond length $[48, 49]$ $[48, 49]$: $tc_i^{\dagger}c_j \rightarrow [t + \epsilon(u_i - u_j)]c_i^{\dagger}c_j$. In the phonon Hamiltonian, up to third nearest-neighbour elastic potentials are

FIG. 2: Adiabatic and non-adiabatic contributions to the phonon magnetic moment.(a) The isoenergy surface of electronic valence band; (b) λ_{ij}^A and λ_{ij}^{NA} along the high symmetry lines of the phonon Brillouin zone with $\eta_p = 0.04$ eV, $\eta_c = 0.04$ eV, $\eta_v = 0.04$ eV. (c) The skew-scattering and side-jump contribution in λ_{ij}^{NA} along the high symmetry lines of the phonon Brillouin zone with the same parameter of (b) , (c) the inset displays that an electron state near K can be scattered to a energy degenerate state near $K^{'}$ by a phonon with momentum K, vice versa. (d)-(g) The adiabatic term λ_{ij}^A and non-adiabatic term λ_{ij}^{NA} for the highest phonon band in phonon Brillouin zone for (d) $\eta_p = 0.02$ eV, $\eta_c = 0.002$ eV, $\eta_v = 0.004$ eV; (e) $\eta_p = 0.02$ eV, $\eta_c = 0.02$ eV, $\eta_v = 0.004$ eV; (f) $\eta_p = 0.02$ eV, $\eta_c = 0.004$ eV, $\eta_v = 0.02$ eV; (g) $\eta_p = 0.004$ eV, $\eta_c = 0.02$ eV, $\eta_v = 0.04$ eV, respectively. (h) The angular momentum of phonon in phonon Brillouin zone with the same parameter of (f).

considered. We also use real unit to make it clear that the phonon part has a much smaller energy scale than the electronic part. Detailed model description and parameter choices can be found in supplementary materials [\[43\]](#page-5-1).

Figure [2\(](#page-3-0)a) shows the electronic band structure, and near K/K' point there are gapped Dirac cones. We then use Eq. [\(4\)](#page-1-3) and [\(7\)](#page-1-4) to calculate λ_{ij} and λ_{ij}^{NA} , respectively. The adiabatic part λ_{ij}^A is obtained by their difference. The results along high-symmetry path in the phonon Brillouin zone are shown in Fig. [2\(](#page-3-0)b). We find that with relaxation process, both the adiabatic and non-adiabatic part of the phonon magnetic moment are peaked near the Γ and K piont, and that they can be generally comparable.

To further illuminate the origin of the non-adiabatic magnetic moment, we plot the skew-scattering and sidejump contribution in Fig. $2(c)$. We find that the side jump contribution almost vanishes, as it requires the breaking of particle-hole symmetry for two-band systems, which is prevserved in our model [\[43\]](#page-5-1). In comparison, the skew-scattering contribution shows a large peak around the phonon K point and as shown in the inset plot, it is due to the degeneracy of the part of the electronic band around K and K' point as they are connected by phonon momentum at K point. By expanding the lattice Hamitonian around K point, we can simiplify the skewscattering contribution which clearly shows its geometric

origin [\[43\]](#page-5-1)

$$
\lambda_{z,\mathbf{K}}^{SS,NA} = -\frac{1}{2} \left(\delta_c^v + \delta_v^c \right) m_0^z \Omega_0^z G_0 \,, \tag{10}
$$

where $G_0 = 1/(-\omega + 2\varepsilon_0) - (\omega \to -\omega)$, $\Omega_0^z = \frac{4\beta^2 v_f^2}{a_0^4}$ $rac{\Delta}{\varepsilon_0}$ is the molecular Berry curvature, ε_0 is the energy of conduction band, v_f is the Fermi velocity, a_0 is the atomic spacing and $\beta = -\frac{\partial ln t_0}{\partial ln a_0}$.

As different contributions to the phonon magnetic moment are sensitive to the relaxation process, we plot the distribution of λ_{ij} across the whole phonon Brillouin zone in different regimes in Fig. $2(d)-(g)$. We choose the phonon band with the highest energy. In Fig. $2(d)$, $\eta_p \gg \eta_c, \eta_v$, and the adiabaticity is restored. In this case, the adiabatic contribution is order-of-magnitude larger than the non-adiabatic one. The phonon magnetic moment is highly non-uniform, with a large peak at two valleys and small variation in other regions.

In Fig. [2\(](#page-3-0)e-f), the parameters are so chosen that only electrons in one band is adiabatic and those in the other one is not. The non-adiabatic phonon magnetic moment shows a C_3 symmetry, consistent with the point-group symmetry of the lattice. Moreover, although it is peaked at the valley, there are obvious variations in other regions of the Brillouin zone, even with sign changing. Due to the asymmetry between the valence and conduction band parameters, both the skew-scattering and side jump mechanism contribute, and they should correspond to the common part and the difference between (e) and (f), respectively. In Fig. $2(g)$, both bands are in the non-adiabatic regime and the resulting non-adiabatic magnetic moment increases.

In Fig. [2\(](#page-3-0)h), we show the phonon angular moment with the electronic conduction band in the adiabatic regime. This can be realized with hole-doping in the electronic degree of freedom. We find that since the phonon magnetic moment is highly nonuniform in the Brillouin zone, so is the angular momentum. Moreover, the angular momentum at different regions are roughly on the same order, showing the importance of the non-adiabatic contribution, as the adiabatic magnetic moment is mainly localized near the K and K' point. This inhomogeneity can manifest in the optical response of phonons.

In summary, we have derived the skew-scattering and side-jump contributions to phonon magnetization by including the relaxation process, both of which are connected to the nonlocal molecular Berry curvature. Using the honeycomb lattice as an exmple, we show that the non-adiabatic magnetic moment can be as important as the adiabatic one and that the resulting phonon angular momentum can vary drastically in the Brilloun zone.

This work was financially supported by the National Natural Science Foundation of China (Grants Nos. 12474158, 12234017, 12374164, and 12488101), Anhui Initiative in Quantum Information Technologies (AHY170000), and Innovation Program for Quantum Science and Technology (2021ZD0302800). We also thank the Supercomputing Center of University of Science and Technology of China for providing the high performance computing resources.

- [1] L. Zhang and Q. Niu, Physical Review Letters 112, 085503 (2014).
- [2] T. Nomura, X.-X. Zhang, S. Zherlitsyn, J. Wosnitza, Y. Tokura, N. Nagaosa, and S. Seki, Physical Review Letters 122, 145901 (2019).
- [3] T. Saito, K. Misaki, H. Ishizuka, and N. Nagaosa, Physical Review Letters 123, 255901 (2019).
- [4] B. Cheng, T. Schumann, Y. Wang, X. Zhang, D. Barbalas, S. Stemmer, and N. P. Armitage, Nano Letters 20, 5991 (2020).
- [5] M. He, P. Rivera, D. Van Tuan, N. P. Wilson, M. Yang, T. Taniguchi, K. Watanabe, J. Yan, D. G. Mandrus, H. Yu, et al., Nature Communications 11 (2020).
- [6] Y. Ren, C. Xiao, D. Saparov, and Q. Niu, Physical Review Letters 127, 186403 (2021).
- [7] L.-H. Hu, J. Yu, I. Garate, and C.-X. Liu, Physical Review Letters 127, 125901 (2021).
- [8] O. Bistoni, F. Mauri, and M. Calandra, Physical Review Letters 126, 225703 (2021).
- [9] D. Saparov, B. Xiong, Y. Ren, and Q. Niu, Physical Review B 105, 064303 (2022).
- [10] D. M. Juraschek, T. Neuman, and P. Narang, Physical Review Research 4, 013129 (2022).
- [11] A. Zabalo, C. E. Dreyer, and M. Stengel, Physical Review B 105, 094305 (2022).
- [12] H. Ueda, M. García-Fernández, S. Agrestini, C. P. Romao, J. van den Brink, N. A. Spaldin, K.-J. Zhou, and U. Staub, Nature 618, 946 (2023).
- [13] Y. Gao, Y. Pan, J. Zhou, and L. Zhang, Physical Review B 108, 064510 (2023).
- [14] J. Bonini, S. Ren, D. Vanderbilt, M. Stengel, C. E. Dreyer, and S. Coh, Physical Review Letters 130, 086701 (2023).
- [15] D. Lujan, J. Choe, S. Chaudhary, G. Ye, C. Nnokwe, M. Rodriguez-Vega, J. He, F. Y. Gao, T. N. Nunley, E. Baldini, et al., Proceedings of the National Academy of Sciences 121 (2024).
- [16] H.-Y. Chen, D. Sangalli, and M. Bernardi, Physical Review Letters 125, 107401 (2020).
- [17] X. Zhang, Y. Zhang, S. Okamoto, and D. Xiao, Physical Review Letters 123, 167202 (2019).
- [18] Q. Wang, M.-Q. Long, and Y.-P. Wang, Physical Review B 110, 024423 (2024).
- [19] C. Strohm, G. L. J. A. Rikken, and P. Wyder, Physical Review Letters 95, 155901 (2005).
- [20] L. Sheng, D. N. Sheng, and C. S. Ting, Physical Review Letters 96, 155901 (2006).
- [21] A. V. Inyushkin and A. N. Taldenkov, JETP Letters 86, 379 (2007).
- [22] Y. Kagan and L. A. Maksimov, Physical Review Letters 100, 145902 (2008).
- [23] J.-S. Wang and L. Zhang, Physical Review B 80, 012301 (2009) .
- [24] L. Zhang, J. Ren, J.-S. Wang, and B. Li, Physical Review Letters 105, 225901 (2010).
- [25] N. Nagaosa, J. Sinova, S. Onoda, A. H. MacDonald, and N. P. Ong, Reviews of Modern Physics 82, 1539 (2010).
- [26] N. A. Sinitsyn, Q. Niu, and A. H. MacDonald, Physical Review B 73, 075318 (2006).
- [27] D. Kaplan, T. Holder, and B. Yan, Physical Review Letters 125, 227401 (2020).
- [28] D. Kaplan, T. Holder, and B. Yan, Nature Communications 14 (2023).
- [29] R. Raimondi, P. Schwab, C. Gorini, and G. Vignale, Annalen der Physik 524 (2011).
- [30] C. Xiao, Y. Liu, Z. Yuan, S. A. Yang, and Q. Niu, Physical Review B 100, 085425 (2019).
- [31] H. Shen, B. Zhen, and L. Fu, Physical Review Letters 120, 146402 (2018).
- [32] Y. Chen and H. Zhai, Physical Review B 98, 245130 (2018).
- [33] S. Yao and Z. Wang, Physical Review Letters 121, 086803 (2018).
- [34] K. Yokomizo and S. Murakami, Physical Review Letters 123, 066404 (2019).
- [35] L. Pan, X. Chen, Y. Chen, and H. Zhai, Nature Physics 16, 767 (2020).
- [36] Y. Michishita and R. Peters, Physical Review Letters 124, 196401 (2020).
- [37] Y. Nagai, Y. Qi, H. Isobe, V. Kozii, and L. Fu, Physical Review Letters 125, 227204 (2020).
- [38] J.-H. Wang, Y.-L. Tao, and Y. Xu, Chinese Physics Letters 39, 010301 (2022).
- [39] H. Geng, J. Y. Wei, M. H. Zou, L. Sheng, W. Chen, and D. Y. Xing, Physical Review B 107, 035306 (2023).
- [40] T. Yu, J. Zou, B. Zeng, J. W. Rao, and K. Xia, Nonhermitian topological magnonics (2023).
- [41] V. Kozii and L. Fu, Physical Review B 109, 235139 (2024).
- [42] G. Yang, Y.-K. Li, Y. Fu, Z. Wang, and Y. Zhang, Physical Review B 109, 045110 (2024).
Supplemental material, URL_will_be_inserted_by_
- $[43]$ Supplemental material, [publisher](URL_will_be_inserted_by_publisher).
- [44] D. Xiao, J. Shi, and Q. Niu, Physical Review Letters 95, 137204 (2005).
- [45] T. Thonhauser, D. Ceresoli, D. Vanderbilt, and R. Resta, Physical Review Letters 95, 137205 (2005).
- [46] J. Shi, G. Vignale, D. Xiao, and Q. Niu, Physical Review Letters 99, 197202 (2007).
- [47] D. Xiao, M.-C. Chang, and Q. Niu, Reviews of Modern Physics 82, 1959 (2010).
- [48] T. Ando, Journal of the Physical Society of Japan 75, 124701 (2006).
- [49] L. Balents, SciPost Physics 7 (2019).