THEOREMS AND CONJECTURES ON AN ARITHMETIC SUM ASSOCIATED WITH THE CLASSICAL THETA FUNCTION θ_3

BRUCE C. BERNDT, RAGHAVENDRA BHAT, JEFFREY L. MEYER, LIKUN XIE, ALEXANDRU ZAHARESCU

ABSTRACT. Appearing in the modular transformation formula for the classical theta function $\theta_3(z)$ is the sum $S(h,k) := \sum_{j=1}^{k-1} (-1)^{j+1+\lfloor hj/k \rfloor}$, which is an analogue of the classical Dedekind sum s(h,k). We establish several properties for S(h,k) and $S(k) := \sum_{h=1}^{k-1} S(h,k)$. Several conjectures about the values of S(k) are given.

1. INTRODUCTION AND MOTIVATION

The arithmetical sum

$$S(h,k) := \sum_{j=1}^{k-1} (-1)^{j+1+[hj/k]},$$
(1.1)

where h and k are positive integers and [x] is the greatest integer less than or equal to x, appears in the transformation formula for the classical theta function $\theta_3(z)$. The sum S(h,k) is analogous to the classical Dedekind sum s(h,k), which appears in the transformation formula for the Dedekind eta-function $\eta(z)$. In contrast to Dedekind sums, S(h,k) has hardly been studied in the literature.

For positive integers h and k, define

$$S(k) := \sum_{h=1}^{k-1} S(h,k).$$
(1.2)

Extensive calculations lead to the following conjecture.

Conjecture 1.1. If k is an odd prime, then

S(k) > 0.

²⁰²⁰ Mathematics Subject Classification. Primary 11N56; Secondary 11N37, 11F20.

Key words and phrases. arithmetic sums, theta functions, analogue of Dedekind sum, asymptotic formula, Riemann zeta function.

A priori, since S(k) is the sum of $(k-1)^2$ summands, each equal to either +1 or -1, one would naturally expect that neither sign would dominate, i.e., S(k) would be positive roughly half of the time. But this intuition is apparently misleading.

Conjecture 1.1 appears to be very difficult to prove. The primary goal of this paper is to establish several theorems for S(k), and related sums, in an effort to prove Conjecture 1.1 and related conjectures. Throughout the sequel, we see that the difficulties we face in proving these conjectures arise from our lack of knowledge about the distribution of the residues jh modulo k, where h and k are co-prime and $1 \le j \le k - 1$.

The map which sends a pair (j, k) to the congruence class of jh modulo k describes a surface modulo k, and one may reinterpret the main questions discussed in this paper in terms of some delicate/subtle properties of this surface. We remark in passing that those level curves on this surface given by congruences of the form $jh \equiv C \pmod{k}$, where C and k are relatively prime, are modular hyperbolas modulo k. Modular hyperbolas come with their own interesting distribution problems, results, and conjectures. For a survey on modular hyperbolas, the reader is referred to I. E. Shparlinski [7].

2. TRANSFORMATION FORMULA

We begin by stating the transformation formula containing S(h, k).

Definition 2.1. Let

$$S := \frac{z+1}{z}$$
 and $T := -\frac{1}{z}$, $z \in \mathbb{H}$.

Then Γ_{θ} is that subgroup of transformations of the modular group $\Gamma(1)$ that is generated by S^2 and T.

Recall that the classical theta function $\theta_3(z)$ is defined by

$$\theta_3(z) := \sum_{n=-\infty}^{\infty} q^{n^2}, \quad q = e^{\pi i z}, \quad z \in \mathbb{H}.$$

We now offer a modular transformation formula for $\log\{\theta_3(z)\}$ [1, p. 339, Theorem 4.1], where here and in what follows, $\log w$ denotes the principal value of the logarithm.

Theorem 2.2. If c and d are co-prime integers of opposite parity with c > 0, and if

$$Vz = \frac{az+b}{cz+d} \in \mathbb{H},$$

then

$$\log\{\theta_3(Vz)\} = \log\{\theta_3(z)\} + \frac{1}{2}\log(cz+d) - \frac{1}{4}\pi i + \frac{1}{4}\pi i S(d,c)$$

where S(d, c) is defined by (1.1).

The sum S(d, c) satisfies a beautiful reciprocity theorem [1, p. 339, Theorem 4.2]. Let c and d be positive, co-prime integers of opposite parity. Then,

$$S(d,c) + S(c,d) = 1.$$
 (2.1)

(Note that the sum S(d,c) has one less term than the sum $\mathcal{S}(d,c)$ appearing in [1, p. 339].)

Let

$$((x)) := \begin{cases} x - [x] - \frac{1}{2}, & \text{if } x \notin \mathbb{Z}, \\ 0, & \text{if } x \in \mathbb{Z}. \end{cases}$$

Then, the Dedekind sum s(d, c) is defined by

$$s(d,c) := \sum_{j=1}^{c-1} \left(\left(\frac{dj}{c} \right) \right) \left(\left(\frac{j}{c} \right) \right).$$

As remarked above, Dedekind sums appear in the transformation formula for the Dedekind eta-function $\eta(z)$, which is defined by

$$\eta(z) := q^{1/24} \prod_{n=1}^{\infty} (1-q^n), \quad q = e^{2\pi i z}, \quad z \in \mathbb{H}.$$

The Dedekind eta-function satisfies the following transformation formula.

Theorem 2.3. [5, p. 145] If c and d are co-prime integers of opposite parity with c > 0, and if

$$Vz = \frac{az+b}{cz+d}, \quad z \in \mathbb{H},$$

then

$$\log\{\eta(Vz)\} = \log\{\eta(z)\} + \frac{1}{2}\log\left(\frac{cz+d}{i}\right) + \pi i s(-d,c).$$

Dedekind sums satisfy a famous reciprocity theorem [6, p. 4, Theorem 1]. If c and d are relatively prime, positive integers, then

$$s(c,d) + s(d,c) = -\frac{1}{4} + \frac{1}{12} \left(\frac{c}{d} + \frac{1}{cd} + \frac{d}{c} \right).$$
(2.2)

Thus, S(d,c) is an analogue of the Dedekind sum s(d,c), and the reciprocity theorem (2.1) is an analogue of (2.2).

3. Arithmetic Sums Related to S(h,k)

Recall that S(h, k) and S(k) are defined in (1.1) and (1.2), respectively. Furthermore, set

$$S(h, j, k) := (-1)^{j+1+\lfloor hj/k \rfloor}.$$
(3.1)

In this paper, we shall also study analogues wherein the number of summands is 'doubled.' To that end, define

$$T(k) := \sum_{h=1}^{2k-1} T(h,k), \qquad (3.2)$$

where

$$T(h,k) := \sum_{j=1}^{2k-1} (-1)^{j+1+[hj/k]}.$$
(3.3)

Also, set

$$T(h, j, k) := S(h, j, k) := (-1)^{j+1+[hj/k]}.$$
(3.4)

4. NUMERICAL CALCULATIONS AND CONJECTURES

The tables that follow give the first few values for S(k) and T(k).

As we remarked in Section 1, calculations led us to Conjecture 1.1. If we remove the hypothesis that k is prime, the conjecture is false. If k is prime, considerably more appears to be true. We strengthen Conjecture 1.1 below.

Conjecture 4.1. For each odd prime k > 5,

S(k) > k.

Stronger conjectures can be made. In particular, there are only 17 numbers less than 10000, namely, 2, 3, 5, 7, 11, 13, 17, 23, 29, 41, 53, 59, 83, 113, 149, 179, 233, for which S(k) is less than 2k. Note that the largest is only 233.

Conjecture 4.2. *For* k > 233*,*

S(k) > 2k.

Moreover, with 3119 being the largest, there are only 87 numbers less than 10000 that fail the inequality S(k) < 3k: 2, 3, 5, 7, 11, 13, 17, 19, 23, 29, 31, 37, 41, 43, 47, 53, 59, 61, 67, 71, 73, 79, 83, 89, 97, 101, 103, 107, 109, 113, 131, 137, 139, 149, 163, 167, 173, 179, 193, 197, 233, 239, 251, 257, 263, 269, 293, 317, 347, 349, 359, 383, 389, 419, 439, 443, 449, 479, 503, 509, 557, 563, 569, 593, 599, 683, 719, 743, 797, 809, 827, 839, 863, 1013, 1019, 1049, 1103, 1229, 1259, 1409, 1733, 1889, 1913, 2339, 2459, 2969, 3119. Thus, we make the following conjecture:

4

TABLE 1.

TABLE 2.

$\begin{array}{c ccccccccccccccccccccccccccccccccccc$					
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	\boldsymbol{k}	S(k)		\boldsymbol{k}	1
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	1	0	-	1	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	2	1		2	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	3	2	-	3	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	4	5		4	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	5	4		5	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	6	7	-	6	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	7	10		7	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	8	11		8	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	9	11	-	9	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	10	8		10	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	11	17	-	11	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	12	14		12	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	13	21		13	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	14	20	-	14	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	15	15		15	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	16	18	-	16	
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	17	39		17	-
$\begin{array}{c ccccccccccccccccccccccccccccccccccc$	18	24	-	18	-
20 38 20 -	19	21		19	-
	20	38		20	-

Conjecture 4.3. For k > 3119,

$$S(k) > 3k.$$

Our current calculations do not lead us to the conjecture that for k sufficiently large, S(k) > 4k. We make the even stronger conjecture below, but we have no evidence for it.

Conjecture 4.4. Let $n \in \mathbb{N}$. Then,

$$S(k) > nk$$

for all primes k sufficiently large, and depending on n.

We provide two graphs below. The first image depicts the first 10000 values of S(k). Note that there are several negative values of S(k). In the second graph below S(k) is depicted for only primes k.





We offer some observations about the negative values of S(k).

For $k \leq 10000$, there are 39 negative values of S(k) for which k is a multiple of 3, but not a multiple of 5, with the first being S(2079) = -1390 and the last being S(9933) = -448.

For $k \leq 10000$, there are 8 negative values of S(k) when k is a multiple of 5, but not a multiple of 3, with the first being S(5005) = -1332 and the last being S(8855) = -7950. For $k \leq 10000$, there are 104 negative values of S(k) when k is a multiple of both 5

and 3, with the first being S(945) = -296 and the last being S(9975) = -22450.

Hence, altogether, there are 151 negative values of S(k) for $k \leq 10000$. It is remarkable that all of these values of k are multiples of either 3 or 5 (or both). Contrary to what one might expect, there are many more values of k divisible by 15 than there are divisible by 3 or 5 only. Although our computations have extended only to k = 10000, we conjecture, with some trepidation, that if S(k) < 0, then k is a multiple of either 3, 5, or 15.

5. Elementary results

We provide some elementary results about S(h, k).

Proposition 5.1. For each positive integer k, S(k) and k have opposite parity.

Proof. Assume first that k is odd. Thus, k - 1 is even. For $1 \le h \le k$, S(h, k) is a sum of k - 1 values of either +1 or -1. Clearly, a sum over an even number of ± 1 's is even, i.e., S(h, k) is even. Thus, S(h, k) and k have opposite parity.

By a similar argument, if k is even, we easily see that S(h, k) is odd. Thus, again, S(h, k) and k have opposite parity.

Corollary 5.2. The only tuple for which both k and S(k) are prime is (3, 2).

Proposition 5.3. *When* h = k - 1*,* S(h, k) = h*.*

Proof. Recall that S(h,k) is defined in (1.1). If h = k - 1, then for each value of $j, 1 \le j \le k$, we see that $\left\lfloor \frac{hj}{k} \right\rfloor = \left\lfloor \frac{(k-1)j}{k} \right\rfloor = j - 1$. Hence,

$$S(k-1,k) = \sum_{j=1}^{k-1} 1 = k-1.$$

Lemma 5.4. Let gcd(h, k) = 1, with k odd. Then,

$$S(k) \equiv \begin{cases} 0, & \text{if } k \equiv 1 \pmod{4}, \\ 2, & \text{if } k \equiv 3 \pmod{4}. \end{cases}$$

Proof. We shall use the familiar equality

$$[x] + [-x] = -1,$$

when x is not an integer. Assume first that j is odd. Then,

$$S(k-h,j,k) = (-1)^{[(k-h)j/k]} = (-1)^{[j-hj/k]} = -(-1)^{[-hj/k]} = (-1)^{[hj/k]}.$$
 (5.1)

Hence, for odd j,

$$S(k-h, j, k) = S(h, j, k), \quad 1 \le h \le \frac{k-1}{2}, \ 1 \le j \le k-1.$$
(5.2)

If j is even, upon using the same argument as we did in (5.1), we conclude that (5.2) holds as well. By (1.1) and (5.2),

$$S(k) = 2\sum_{h=1}^{\frac{k-1}{2}} S(h,k).$$
(5.3)

If $k \equiv 1 \pmod{4}$, then

$$\sum_{h=1}^{\frac{k-1}{2}} S(h,k) \tag{5.4}$$

contains an even number of terms, and so by (5.3), $S(k) \equiv 0 \pmod{4}$. If $k \equiv 3 \pmod{4}$, then the sum (5.4) contains an odd number of terms, and so, by (5.3), $S(k) \equiv 2 \pmod{4}$. This completes the proof.

The following important theorem relates S(h,k) and T(h,k). Thus, if we prove a result about T(h,k), we can obtain a corresponding theorem for S(h,k).

Theorem 5.5. If h and k are positive, co-prime integers, then

$$T(h,k) = \left(1 + (-1)^{k+h}\right)S(h,k) + (-1)^{k+h+1}.$$

Proof. We have

$$T(h,k) = \sum_{j=1}^{2k-1} (-1)^{j+1+\lfloor hj/k \rfloor}$$

= $\sum_{j=1}^{k-1} (-1)^{j+1+\lfloor hj/k \rfloor} + (-1)^{k+1+\lfloor hk/k \rfloor} + \sum_{j=k+1}^{2k-1} (-1)^{j+1+\lfloor hj/k \rfloor}$
= $S(h,k) + (-1)^{k+h+1} + \sum_{j=1}^{k-1} (-1)^{k+j+1+\lfloor h(k+j)/k \rfloor}$
= $S(h,k) + (-1)^{k+h+1} + (-1)^{k+h} \sum_{j=1}^{k-1} (-1)^{j+1+\lfloor hj/k \rfloor}$
= $S(h,k) + (-1)^{k+h+1} + (-1)^{k+h} S(h,k)$
= $(1 + (-1)^{k+h}) S(h,k) + (-1)^{k+h+1}$.

Corollary 5.6. If h and k are positive, co-prime integers of opposite parity, then

$$T(h,k) = 1.$$

Proof. The corollary follows immediately from Theorem 5.5.

Lemma 5.7. If h and k are odd, co-prime, positive integers, then

$$S(h,k) = 0.$$

Proof. For $1 \le j \le \frac{k-1}{2}$, $(-1)^{k-j+1+[(k-j)h/k]} = (-1)^{-j+[h-hj/k]} = (-1)^{j+h+[-hj/k]}$ $= (-1)^{j+1+1+[hj/k]} = -(-1)^{j+1+[hj/k]}$. (5.5)

It follows from (1.1) and (5.5) that S(h,k) = 0.

Theorem 5.8. Let gcd(h, k) = 1 with k > 0, and let q be any positive integer. Then

$$S(qh,qk) = \begin{cases} S(h,k), & \text{if } h+k \text{ and } q \text{ are odd;} \\ 1, & \text{if } h+k \text{ is odd and } q \text{ is even;} \\ -(q-1), & \text{if } h \text{ and } k \text{ are odd.} \end{cases}$$

Proof. From the definition (1.1), we have

$$\begin{split} S(qh,qk) &= \sum_{j=1}^{qk-1} (-1)^{j+1+[hj/k]} \\ &= \sum_{\ell=0}^{q-1} \sum_{j=\ell k+1}^{(\ell+1)k-1} (-1)^{j+1+[hj/k]} + \sum_{\ell=1}^{q-1} (-1)^{\ell k+1+\ell h} \\ &= \sum_{\ell=0}^{q-1} \sum_{j=1}^{k-1} (-1)^{j+\ell k+1+[h(j+\ell k)/k]} - \sum_{\ell=1}^{q-1} (-1)^{\ell (h+k)} \\ &= \sum_{\ell=0}^{q-1} (-1)^{\ell (h+k)} \sum_{j=1}^{k-1} (-1)^{j+1+[hj/k]} - \sum_{\ell=1}^{q-1} (-1)^{\ell (h+k)} \\ &= S(h,k) \sum_{\ell=0}^{q-1} (-1)^{\ell (h+k)} - \sum_{\ell=1}^{q-1} (-1)^{\ell (h+k)}. \end{split}$$

The theorem now follows upon an analysis of the sums on ℓ for the different parities of h, k, and q.

6. Lower bounds for S(k)

In considering Conjecture 4.1, we seek lower bounds for S(k). We propose several such bounds in this section. Our first bounds are 'easy.'

Recall from Lemma 5.7 that S(h, k) = 0, when h and k are co-prime positive integers. Thus, we can write

$$S(k) = \sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \sum_{\substack{h=1\\j \text{ odd}}}^{k-1} (-1)^{j+1+[hj/k]}$$
$$= \sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \sum_{\substack{h=1\\h=1}}^{k-1} (-1)^{[hj/k]}$$
$$= \frac{(k-1)^2}{2} - 2 \sum_{\substack{j=1\\j \text{ odd}}}^{k-1} r(j,k),$$
(6.1)

where

$$r(j,k) = \#\left\{h : 1 \le h \le k-1; \left\lfloor\frac{hj}{k}\right\rfloor \text{ is odd }\right\}$$

where we have used the elementary evaluation [6, p. 32, Equation (41)], [2, p. 214, Equation (4.6)],

$$\sum_{k=1}^{n-1} \left[\frac{mk}{n} \right] = \frac{(m-1)(n-1)}{2},$$
(6.2)

where *m* and *n* are relatively prime. Observe that if we assume that each value of $\left\lfloor \frac{hj}{k} \right\rfloor$ is odd, or that each value of $\left\lfloor \frac{hj}{k} \right\rfloor$ is even, then we obtain the trivial bounds

$$\frac{(k-1)^2}{2} \le S(k) \le \frac{(k-1)^2}{2}.$$
(6.3)

Using the elementary relation,

$$[x] - 2\left[\frac{x}{2}\right] = \begin{cases} 0, & \text{if } [x] \text{ is even,} \\ 1, & \text{if } [x] \text{ is odd,} \end{cases}$$
(6.4)

we can write

$$r(j,k) = \sum_{h=1}^{k-1} \left(\left[\frac{hj}{k} \right] - 2 \left[\frac{hj}{2k} \right] \right)$$
$$= \frac{(j-1)(k-1)}{2} - 2 \sum_{h=1}^{k-1} \left[\frac{hj}{2k} \right].$$
(6.5)

If we write j = 2b + 1, then

$$\left[\frac{hb}{k}\right] \le \left[\frac{hj}{2k}\right] = \left[\frac{h(2b+1)}{2k}\right] \le \left[\frac{h(b+1)}{k}\right]$$

Using these trivial bounds for the floor function, we obtain the same trivial bounds (6.3).

If we rewrite (6.1) using fractional parts, then we have

$$S(k) = (k-1)^2 - 4 \sum_{\substack{j=1\\ j \text{ odd}}}^{k-1} \sum_{\substack{h=1\\ k=1}}^{k-1} \left\{ \frac{hj}{2k} \right\}.$$

We are not able to obtain a positive lower bound unless we can obtain an estimate for the number of (h, j) such that $\left[\frac{hj}{2k}\right] = \left[\frac{hb}{k}\right] + 1$. Also, the sum over h in (6.5) does not

have a closed form. As we related, there is a closed form (6.2) for the sum $\sum_{h=1}^{n-1} \left[\frac{hm}{n}\right]$, but no known formula for $\sum_{h=1}^{[n/2]} \left\{ \frac{hm}{n} \right\}$. The following theorem provides a more interesting, but still unsuccessful, approach

to proving the positivity of S(k) for each prime k.

Theorem 6.1. If k is a prime, then

$$\sum_{h=1}^{k-1} S(h,k) = -(k-1)^2 + 4\sum_{\ell=1}^{\frac{k-1}{2}} \sum_{h=1}^{\frac{k-1}{2}} \left(\left\{ \frac{2h\ell}{k} \right\} + \left\{ \frac{h(2\ell-1)}{k} - \frac{1}{2} \right\} \right)$$

Proof. Using Lemma 5.7, we may write

$$\sum_{h=1}^{k} S(h,k) = \sum_{h=1}^{\frac{k-1}{2}} S(2h,k) = \sum_{h=1}^{\frac{k-1}{2}} \sum_{j=1}^{k-1} (-1)^{j+1+\lfloor 2hj/k \rfloor}$$

$$= \sum_{j=1}^{k-1} \sum_{h=1}^{\frac{k-1}{2}} (-1)^{j+1+\lfloor 2hj/k \rfloor}$$

$$= -\sum_{j=1}^{k-1} \sum_{h=1}^{\frac{k-1}{2}} (-1)^{j+\lfloor 2hj/k \rfloor}$$

$$= -\sum_{j=1}^{k-1} \sum_{h=1}^{\frac{k-1}{2}} (-1)^{\lfloor (2hj+jk)/k \rfloor}$$

$$= 2\sum_{j=1}^{k-1} m(j,k) - \frac{(k-1)^2}{2},$$
(6.6)

where

$$m(j,k) = \#\left\{h: 1 \le h \le \frac{k-1}{2}; \left[\frac{(2h+k)j}{k}\right] \text{ is odd}\right\},$$
 (6.7)

and where in the last line of (6.6), we added and subtracted the number of odd values of $\left[\frac{2hj+jk}{k}\right]$.

Therefore, using (6.4), we have

$$m(j,k) = \sum_{h=1}^{\frac{k-1}{2}} \left(\left[\frac{(2h+k)j}{k} \right] - 2 \left[\frac{(2h+k)j}{2k} \right] \right)$$
$$= \sum_{h=1}^{\frac{k-1}{2}} \left(j + \left[\frac{2hj}{k} \right] - 2 \left[\frac{hj}{k} + \frac{j}{2} \right] \right).$$

Using (6.2) in the fourth equality below, we find that

$$\begin{split} \sum_{j=1}^{k-1} m(j,k) &= \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left(2\ell + \left[\frac{2h \cdot 2\ell}{k} \right] - 2 \left[\frac{h \cdot 2\ell}{k} + \frac{2\ell}{2} \right] \right) \right) \\ &+ \left(2\ell - 1 + \left[\frac{2h(2\ell - 1)}{k} \right] - 2 \left[\frac{h(2\ell - 1)}{k} + \frac{2\ell - 1}{2} \right] \right) \right) \\ &= \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{2h \cdot 2\ell}{k} \right] + \left[\frac{2h(2\ell - 1)}{k} \right] \right) \\ &- \sum_{\ell=1}^{2} \sum_{h=1}^{k-1} \left(2 \left[\frac{h \cdot 2\ell}{k} \right] + 2 \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] + 1 \right) \\ &= \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left[\frac{2h\ell}{k} \right] - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right) - \frac{(k-1)^2}{4} \\ &= \sum_{h=1}^{k-1} \sum_{h=1}^{k-1} \left[\frac{2h\ell}{k} \right] - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right) - \frac{(k-1)^2}{4} \\ &= \sum_{h=1}^{k-1} \sum_{h=1}^{k-1} (2h - 1) - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right) - \frac{(k-1)^2}{4} \\ &= \frac{k-1}{2} \sum_{h=1}^{k-1} (2h - 1) - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right) - \frac{(k-1)^2}{4} \\ &= \frac{k-1}{2} \cdot \frac{(k-1)^2}{4} - \frac{(k-1)^2}{4} - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right) \\ &= \frac{(k-1)^2(k-3)}{8} - 2 \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell - 1)}{k} - \frac{1}{2} \right] \right). \end{split}$$

It seems reasonable to restate this last identity in terms of fractional parts, rather than greatest integer functions. We therefore find that

$$\sum_{\ell=1}^{\frac{k-1}{2}} \sum_{h=1}^{\frac{k-1}{2}} \left(\left[\frac{h \cdot 2\ell}{k} \right] + \left[\frac{h(2\ell-1)}{k} - \frac{1}{2} \right] \right)$$
$$= \frac{(k-1)^3}{16} - \sum_{\ell=1}^{\frac{k-1}{2}} \sum_{h=1}^{\frac{k-1}{2}} \left(\left\{ \frac{2h\ell}{k} \right\} + \left\{ \frac{h(2\ell-1)}{k} - \frac{1}{2} \right\} \right)$$

Finally, when k is an odd prime, we have

$$\sum_{h=1}^{k-1} S(h,k) = \frac{(k-1)^2(k-5)}{4} - 4\left(\frac{(k-1)^3}{16} - \sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left\{\frac{2h\ell}{k}\right\} + \left\{\frac{h(2\ell-1)}{k} - \frac{1}{2}\right\}\right)\right)$$
$$= \frac{(k-1)^2(k-5)}{4} - \frac{(k-1)^3}{4} + 4\sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left\{\frac{2h\ell}{k}\right\} + \left\{\frac{h(2\ell-1)}{k} - \frac{1}{2}\right\}\right)$$
$$= -(k-1)^2 + 4\sum_{\ell=1}^{k-1} \sum_{h=1}^{k-1} \left(\left\{\frac{2h\ell}{k}\right\} + \left\{\frac{h(2\ell-1)}{k} - \frac{1}{2}\right\}\right), \tag{6.8}$$

and this completes the proof.

It is here that the difficulty of proving that S(k) > 0 becomes apparent. If one surmises that the fractional parts on average are equal to $\frac{1}{2}$, then, on average, the right side of (6.8) should be close to 0. However, the data do not show this. So there must be a skewed distribution of the fractional parts so that the expressions in the sum are somehow larger than 1 on average.

We may pair the fractional parts in the sum of (6.8) as follows to form integers. Denote

$$f(l,h) = \left\{\frac{2hl}{k}\right\} + \left\{\frac{h(2l-1)}{k} - \frac{1}{2}\right\}.$$

Proposition 6.2. For $m < \frac{k+1}{4}$,

$$g(m,h) := f(m,h) + f\left(\frac{k-1}{2} - m + 1, h\right)$$

is an integer.

Proof. Rewrite g(m, h) as:

$$g(m,h) = \left\{\frac{2mh}{k}\right\} + \left\{\frac{h}{k} - \frac{2mh}{k}\right\} + \left\{\frac{2mh}{k} - \frac{h}{k} - \frac{1}{2}\right\} + \left\{\frac{-2mh}{k} - \frac{1}{2}\right\}.$$

We separate several cases.

A.
$$\left\{\frac{h}{k}\right\} \leq \left\{\frac{2mh}{k}\right\}$$
. Then,
 $g(m,h) = \left\{\frac{2mh}{k}\right\} + 1 + \left\{\frac{h}{k}\right\} - \left\{\frac{2mh}{k}\right\} + \left\{\frac{2mh}{k} - \frac{h}{k} - \frac{1}{2}\right\} + 1 - \left\{\frac{2mh}{k} + \frac{1}{2}\right\}$
 $= 1 + \left\{\frac{h}{k}\right\} + \left\{\frac{2mh}{k} - \frac{h}{k} - \frac{1}{2}\right\} + 1 - \left\{\frac{2mh}{k} + \frac{1}{2}\right\}.$

We have the following subcases.

A1. If
$$\left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} \ge \frac{1}{2}$$
, then we must have $\left\{\frac{2mh}{k}\right\} > \frac{1}{2}$, and so
 $g(m,h) = 1 + \left\{\frac{h}{k}\right\} + \left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} - \frac{1}{2} + 1 - \left(\left\{\frac{2mh}{k}\right\} + \frac{1}{2} - 1\right) = 2.$
A2. If $\left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} < \frac{1}{2}$ and $\left\{\frac{2mh}{k}\right\} > \frac{1}{2}$, then
 $g(m,h) = 1 + \left\{\frac{h}{k}\right\} + 1 + \left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} - \frac{1}{2} + 1 - \left(\left\{\frac{2mh}{k}\right\} + \frac{1}{2} - 1\right) = 3.$
A3. If $\left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} < \frac{1}{2}$ and $\left\{\frac{2mh}{k}\right\} < \frac{1}{2}$, then
 $g(m,h) = 1 + \left\{\frac{h}{k}\right\} + 1 + \left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} - \frac{1}{2} + 1 - \left(\left\{\frac{2mh}{k}\right\} + \frac{1}{2}\right) = 2.$
B. $\left\{\frac{h}{k}\right\} > \left\{\frac{2mh}{k}\right\}$ Then, since $\left\{\frac{h}{k}\right\} < \frac{1}{2}$, we must have $\left\{\frac{2mh}{k}\right\} < \frac{1}{2}$; thus
 $g(m,h) = \left\{\frac{h}{k}\right\} + 1 + \left\{\frac{2mh}{k}\right\} - \left\{\frac{h}{k}\right\} - \frac{1}{2} + 1 - \left(\left\{\frac{2mh}{k}\right\} + \frac{1}{2}\right) = 1.$

Finally, when $\frac{k-1}{2}$ is odd, there is an unpaired term f(l,h) in the sum (6.8) for $l = \frac{k+1}{4}$. Thus,

$$f(l,h) = \left\{\frac{2h(k+1)}{4k}\right\} + \left\{\frac{h}{k}\frac{k-1}{2} - \frac{1}{2}\right\}$$
$$= \left\{\frac{h}{2} + \frac{h}{2k}\right\} + \left\{\frac{h}{2} - \frac{h}{2k} - \frac{1}{2}\right\}.$$
(6.9)

Now, $\left\{\frac{h}{2}\right\} = \frac{1}{2}$ or 0 and $\left\{\frac{h}{2k}\right\} < \frac{1}{2}$. Hence, from (6.9),

$$f(l,h) = \left\{\frac{h}{2} + \frac{h}{2k}\right\} + \left\{\frac{h}{2} - \frac{h}{2k} - \frac{1}{2}\right\}$$
$$= \left\{\frac{h}{2}\right\} + \left\{\frac{h}{2k}\right\} + 1 + \left\{\frac{h}{2}\right\} - \left\{\frac{h}{2k}\right\} - \frac{1}{2} = 2\left\{\frac{h}{2}\right\} + \frac{1}{2}.$$

If one can show that the number of pairs (m, h) that satisfy condition A2 is greater than the number of pairs that satisfy B, then we have a proof of the conjecture S(k) > 0for prime k.

Corollary 6.3. For primes k, the values of S(k) are equally distributed modulo 4. Specifically, there are only two possible cases for primes k, namely $S(k) \equiv 0 \pmod{4}$ and $S(k) \equiv 2 \pmod{4}$, and these two cases are distributed evenly.

Proof. By Theorem 6.1 and Proposition 6.2, when $\frac{k-1}{2}$ is odd, i.e., when $k \equiv 3 \pmod{4}$,

$$S(k) \equiv -(k-1)^2 + (k-1) \equiv 2 \pmod{4}.$$

When $\frac{k-1}{2}$ is even, i.e., when $k \equiv 1 \pmod{4}$,

$$S(k) \equiv -(k-1)^2 \equiv 0 \pmod{4}.$$

Therefore, by the prime number theorem in arithmetic progression, the number of primes congruent to 1 and 3 (mod 4) have asymptotically the same density. In other words, the values of S(k) among primes k are distributed evenly modulo 4.

Employing Riemann–Stieltjes sums and integrals, we establish a lower bound for S(k). Let

$$H(k) := \sum_{\substack{j=1\\ j \text{ odd}}}^{k-1} \frac{1}{j}.$$

Theorem 6.4. Let k denote any odd prime. Then,

$$S(k) \ge -\frac{k-1}{2} + kH(k) - \frac{(k-1)(k+1)}{4},$$

where, as $k \to \infty$,

$$H(k) = \frac{1}{2}\log\left(2k\right) + \frac{\gamma}{2} + O\left(\frac{1}{k}\right),$$

and where γ denotes Euler's constant.

Proof. Recall from (6.6) the identity

$$S(k) = \frac{(k-1)^2}{2} - 2\sum_{\substack{j=1\\ j \text{ odd}}}^{k-1} m(j,k),$$
(6.10)

where m(j,k) is defined in (6.7), i.e.,

$$m(j,k) = \sum_{h=1}^{k-1} \left(\left[\frac{hj}{k} \right] - 2 \left[\frac{hj}{2k} \right] \right).$$
(6.11)

Thus, by (6.11),

$$\lim_{k \to \infty} \frac{1}{k} m(j,k) = \lim_{k \to \infty} \frac{1}{k} \sum_{h=1}^{k} \left(\left[\frac{hj}{k} \right] - 2 \left[\frac{hj}{2k} \right] \right)$$
$$= \int_{0}^{1} \left([jx] - 2 \left[\frac{jx}{2} \right] \right) dx$$
$$= \frac{1}{j} \left(\frac{j-1}{2} \right).$$
(6.12)

To obtain the last equality above, first note that

$$[jx] - 2\left[\frac{jx}{2}\right] = \begin{cases} 1, & \text{if } k + \frac{1}{2} \le \frac{jx}{2} \le k + 1, \\ 0, & \text{otherwise.} \end{cases}$$
(6.13)

Next, subdivide the interval [0,1] into subintervals of length $\frac{1}{j}$. From (6.13), we see that the largest value of k is $\frac{j-1}{2}$. The value of the integral is therefore $\frac{1}{j} \times \frac{j-1}{2}$, and consequently (6.12) follows.

From the limit in (6.12), we are then motivated to define an error term $\epsilon_j(k)$ by

$$\epsilon_j(k) := \frac{1}{k} m(j,k) - \frac{1}{j} \left(\frac{j-1}{2}\right).$$
 (6.14)

We now derive an upper bound for $\epsilon_j(k)$. Define

$$f(j,k) := [jx] - 2\left[\frac{jx}{2}\right].$$
 (6.15)

Subdivide the interval [0, 1] into subintervals of length $\frac{1}{j}$. Consider the sequence of fractions

$$\frac{1}{k}, \frac{2}{k}, \dots, \frac{k-1}{k},$$

and set

$$a_j = \frac{n}{k}, \quad 1 \le n \le k - 1.$$

Consider a point

$$\frac{n}{j}, \quad 1 \le n \le j-1.$$

Let a_m be the closest fraction to $\frac{n}{j}$ to the left. Of course, a_{m+1} is the closest fraction to $\frac{n}{j}$ on the right. We note that $|a_{m+1} - a_m| = \frac{1}{k}$. From (6.13) and (6.15), we see that the contribution of each of these subintervals to f(j,k) is $\frac{1}{2} \times \frac{1}{k}$. Since there are j-1 of these contributions, we conclude that

$$\epsilon_j(k) \le \frac{j-1}{2k}.$$

Therefore, by (6.14),

$$m(j,k) \le \frac{k}{2} - \frac{k}{2j} + \frac{j-1}{2}.$$
 (6.16)

Finally, from (6.10) and (6.16), we conclude that

$$S(k) \ge \frac{(k-1)^2}{2} - 2\sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \left(\frac{k}{2} - \frac{k}{2j} + \frac{j-1}{2}\right)$$
$$= -\frac{k-1}{2} + k\sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \frac{1}{j} - \frac{(k-1)(k+1)}{4}.$$
(6.17)

From a familiar asymptotic formula for the harmonic sum, as $k \to \infty$,

$$\sum_{j=1}^{k} \frac{1}{j} = \log k + \gamma + O\left(\frac{1}{k}\right),$$

where γ denotes Euler's constant, we deduce that, as $k \to \infty$,

$$\sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \frac{1}{j} = \sum_{j=1}^{k-1} \frac{1}{j} - \frac{1}{2} \sum_{j=1}^{(k-1)/2} \frac{1}{j} = \frac{1}{2} \log \left(2(k-1) \right) + \frac{\gamma}{2} + O\left(\frac{1}{k-1}\right)$$
$$= \frac{1}{2} \log(2k) + \frac{\gamma}{2} + O\left(\frac{1}{k}\right).$$
(6.18)

Together, (6.17) and (6.18) complete the proof of Theorem 6.4.

18

This is slightly better than our previous lower bounds, but it is still not positive. As we can see from the proof above, a proper upper bound for $\epsilon_j(k)$ is crucial to gaining a better lower bound for S(k), but to get a smaller upper bound for $\epsilon_j(k)$, we need to obtain a better understanding of the distribution of fractional parts in (6.8). In other words, we need to have a suitable estimate for the sum

$$\sum_{\substack{j=1\\j \text{ odd}}}^{k-1} \sum_{h=1}^{k-1} \left\{ \frac{hj}{2k} \right\},\,$$

which depends on the residue class of hj modulo 2k. Observe that, since j is odd, gcd(j, 2k) = 1, and therefore $\{hj\}$ exhausts all of the nonzero residue classes modulo 2k. However, our range of h is only $1 \le h \le k - 1$.

7. Elementary Results on T(k), k odd

Recall that T(k) is defined by (3.2); we now provide a general formula for T(k) when k is odd. In particular, we can conveniently express T(k) by a single summation, in contrast to S(k), for which we have representations by only double sums. Moreover, our formula does not involve sums over powers of -1, but instead sums involving the Euler phi-function $\phi(n)$. Recall that

$$\phi(n) = \{m : 1 \le m < n; (m, n) = 1\}.$$

Theorem 7.1. For odd k,

$$T(k) = 2k - 1 - 2\sum_{d|k} d \cdot \phi\left(\frac{k}{d}\right).$$

We show that Theorem 7.1 can be recast using the following lemma.

Lemma 7.2. For odd k, we have

$$\sum_{d|k} d \cdot \phi\left(\frac{k}{d}\right) = \sum_{\substack{1 \le h \le 2k-1 \\ h \text{ odd}}} \gcd(k, h).$$

Proof. We have

$$\sum_{\substack{1 \le h \le 2k-1 \\ h \text{ odd}}} \gcd(h,k) = \sum_{d|k} d \cdot \#\{h : \gcd(h,k) = d, 1 \le h \le 2k-1, h \text{ odd} \}$$
$$= \sum_{d|k} d \cdot \#\left\{h : \gcd\left(\frac{h}{d}, \frac{k}{d}\right) = 1, 1 \le h \le 2k-1, h \text{ odd} \right\}$$

$$= \sum_{d|k} d \cdot \# \left\{ m : \gcd\left(m, \frac{k}{d}\right) = 1, 1 \le m \le \left[\frac{2k-1}{d}\right], m \text{ odd} \right\}$$
$$= \sum_{d|k} d \cdot \phi\left(\frac{k}{d}\right), \tag{7.1}$$

since $d \nmid j, k+1 \leq j \leq 2k-1$.

We will prove Theorem 7.1 with the help of a few useful lemmas.

Lemma 7.3. For k odd and h odd, we have

$$T(h,k) = \sum_{\substack{1 \leq j \leq 2k-1 \\ hj/k \in \mathbb{Z}}} (-1).$$

Proof. Consider any j with $\frac{hj}{k} \notin \mathbb{Z}$. Then,

$$T(h, 2k - j, k) = (-1)^{2k - j + 1 + [h(2k - j)/k]} = (-1)^{2k - j + 1 + 2h + [-hj/k]}$$
$$= (-1)^{-j - [hj/k]} = (-1)^{j + [hj/k]} = -T(h, j, k).$$

Thus, within the summands of T(h, k), each T(h, j, k) cancels out T(h, 2k - j, k), when $\frac{hj}{k} \notin \mathbb{Z}$. On the other hand, if $\frac{hj}{k}$ is an integer, it follows from an analogous calculation that

$$T(h, 2k - j, k) = T(h, j, k)$$

Therefore, we conclude that

$$T(h,k) = \sum_{1 \le h \le 2k-1} T(h,j,k) = \sum_{\substack{1 \le j \le 2k-1 \\ hj/k \in \mathbb{Z}}} (-1)^{j+1+hj/k}.$$

Since the parity of $\frac{hj}{k}$ matches the parity of j when $\frac{hj}{k} \in \mathbb{Z}$, since h and k are both odd, we see that

$$(-1)^{j+1+hj/k} = -1.$$

Thus,

$$T(h,k) = \sum_{\substack{1 \le j \le 2k-1 \\ hj/k \in \mathbb{Z}}} (-1)^{j+1+hj/k} = \sum_{\substack{1 \le j \le 2k-1 \\ hj/k \in \mathbb{Z}}} (-1).$$

Lemma 7.4. For k odd and h odd,

$$T(h,k) = 1 - 2 \cdot \gcd(h,k).$$

Proof. Let d = gcd(h, k). The least common multiple of k and h can be expressed as

$$\operatorname{lcm}(h,k) = \frac{h \cdot k}{d}.$$

Thus, for some $n \in \mathbb{N}$,

$$\frac{hj}{k} \in \mathbb{Z} \Leftrightarrow j = \frac{n \cdot k}{d}.$$

Since $1 \le j \le 2k - 1$, we have $1 \le n < 2d$. Thus, by Lemma 7.3,

$$T(h,k) = \sum_{1 \le n < 2d} T\left(h, \frac{nk}{d}, k\right) = \sum_{1 \le n < 2d} (-1) = -(2d-1) = 1 - 2 \cdot \gcd(k,h).$$

Proof of Theorem 7.1. From Lemmas 5.6 and 7.4, for odd k,

$$T(k) = \sum_{\substack{1 \le h \le 2k-1 \\ h \text{ even}}} 1 + \sum_{\substack{1 \le h \le 2k-1 \\ h \text{ odd}}} (1 - 2 \cdot \gcd(h, k))$$
$$= 2k - 1 - 2 \sum_{\substack{1 \le h \le 2k-1 \\ h \text{ odd}}} \gcd(h, k).$$

Theorem 7.1 now follows from Lemma 7.2.

As immediate consequences of Theorem 7.1, we obtain the following results.

Corollary 7.5. For each prime p,

$$T(p) = 1 - 2p.$$

Proof. From Theorem 7.1, we have

$$T(p) = 2p - 1 - 2\sum_{\substack{1 \le h \le 2p - 1\\h \text{ odd}}} \gcd(h, p).$$

Since p is prime, gcd(h, p) = 1 for all h that are not multiples of p. In the interval of interest, there are p odd numbers, and the only multiple of p is p itself. Thus,

$$T(p) = 2p - 1 - 2 \cdot (\gcd(p, p)) - 2 \cdot (p - 1) = 2p - 1 - 2p - 2p + 2 = 1 - 2p.$$

Corollary 7.6. If p and q are odd primes,

$$T(pq) = 4(p+q) - 6pq - 3.$$

Proof. From Theorem 7.1,

$$T(pq) = 2pq - 1 - 2 \sum_{\substack{1 \le h \le 2pq - 1\\h \text{ odd}}} \gcd(h, pq).$$
(7.2)

First, consider the interval [1, pq]. Other than pq, there are (q - 1) multiples of p, and there are (p - 1) multiples of q. Thus, in [1, pq], this contribution to (7.1) is

$$-2((q-1)p + (p-1)q + 1).$$
(7.3)

Because $gcd(j, pq) = gcd(pq - j, pq), 1 \le j \le pq - 1$, this contribution of the terms in [pq + 1, 2pq - 1] is, by (7.3),

$$-2((q-1)p + (p-1)q).$$
(7.4)

In conclusion, from (7.2)–(7.4),

$$T(pq) = 2pq - 1 - 4((q-1)p + (p-1)q) - 2 = -6pq + 4p + 4q - 3.$$

8. Elementary Results on T(k), k even

We provide a list of elementary theorems on T(k) when k is even. The proofs of the analogue of Theorem 7.1 along with the necessary lemmas are similar to those in Section 7.

Theorem 8.1. When k is even,

$$T(k) = 2k - 1 - \sum_{\substack{1 \le h \le 2k - 1\\h \equiv 2 \pmod{4}}} 2 \gcd(h, k).$$

Lemma 8.2. For k even and h odd,

$$T(h,k) = 1.$$

Lemma 8.3. If $k \equiv 2 \mod 4$ and $h \equiv 0 \mod 4$, then

T(h,k) = 1.

Lemma 8.4. If $k \equiv 2 \mod 4$ and $h \equiv 2 \mod 4$, then

$$T(h,k) = 1 - 2\gcd(h,k).$$

Lemma 8.5. If $k \equiv 0 \mod 4$ and $h \equiv 0 \mod 4$, then

$$\Gamma(h,k) = 1 - 2\gcd(h,k)$$

Corollary 8.6. For all natural numbers n,

$$T\left(2^n\right) = -1.$$

9. Asymptotic Formula for Partial Sums Related to T(k)

Recall from Theorem 7.1 that

$$T(k) = 2k - 1 - 2\sum_{d|k} d \cdot \phi\left(\frac{k}{d}\right).$$
(9.1)

Motivated by (9.1), we now derive an asymptotic formula for the sum on the right-hand side above. Set

$$a_n := \begin{cases} \sum_{d|n} d \cdot \phi\left(\frac{n}{d}\right), & \text{if } n \text{ is odd,} \\ 0, & \text{if } n \text{ is even.} \end{cases}$$
(9.2)

Now define a generating function for a_n by

$$F(s) := \sum_{n=1}^{\infty} \frac{a_n}{n^s}, \qquad \Re(s) > 2.$$
 (9.3)

Next, define

$$G(s) := \sum_{n=1}^{\infty} \frac{\sum_{d|n} d \cdot \phi\left(\frac{n}{d}\right)}{n^s} = \frac{\zeta(s-1)^2}{\zeta(s)} = \prod_{p \text{ prime}} \frac{1 - \frac{1}{p^s}}{\left(1 - \frac{1}{p^{s-1}}\right)^2},\tag{9.4}$$

where the second identity can be established by multiplying the quotient of zeta functions on the right-hand side. (See also [8, p. 6, Equation (1.2.12)].) It follows from (9.2)-(9.4) that

$$F(s) = \prod_{p \text{ odd prime}} \frac{1 - \frac{1}{p^s}}{\left(1 - \frac{1}{p^{s-1}}\right)^2} = \frac{\left(1 - \frac{1}{2^{s-1}}\right)^2}{1 - \frac{1}{2^s}} \cdot G(s) = \frac{\left(1 - \frac{1}{2^{s-1}}\right)^2}{1 - \frac{1}{2^s}} \cdot \frac{\zeta(s-1)^2}{\zeta(s)}.$$
 (9.5)

We now come to the most important result in this section.

Theorem 9.1. For each $0 < \epsilon < \frac{1}{10}$, we have

$$\sum_{n \le x}' a_n = \frac{x^2}{\pi^2} \log x + \frac{1}{6\pi^2} \left(-36\zeta'(2) + A \right) x^2 + O_\epsilon \left(x^{\frac{3}{2} + \epsilon} \log^2 x \right),$$

where

$$A = 12\gamma - 3 + 10\log 2,$$

and where γ denotes Euler's constant. Furthermore, the prime \prime on the summation sign on the left-hand side indicates that if x is an integer, then we count only $\frac{1}{2}a_x$. *Proof.* By Perron's summation formula [3, pp. 12–14],

$$\sum_{n \le x}' a_n = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} F(s) \frac{x^s}{s} ds = \frac{1}{2\pi i} \int_{c-iT}^{c+iT} F(s) \frac{x^s}{s} ds + E(T),$$
(9.6)

where F(s) is defined in (9.5) and

$$E(T) = \frac{1}{2\pi i} \left(\int_{c-i\infty}^{c-iT} + \int_{c+iT}^{c+i\infty} \right) F(s) \frac{x^s}{s} ds.$$
(9.7)

Let c > 2 and $\frac{3}{2} \le \sigma_0 < c$, where c and σ_0 will be determined later. Consider

$$I_T := \frac{1}{2\pi i} \int_{C_T} F(s) \frac{x^s}{s} ds = \frac{1}{2\pi i} \int_{C_T} \frac{\left(1 - \frac{1}{2^{s-1}}\right)^2}{1 - \frac{1}{2^s}} \cdot \frac{\zeta(s-1)^2}{\zeta(s)} \frac{x^s}{s} ds$$

where C_T is the positively oriented rectangle with vertices $\sigma_0 \pm iT$ and $c \pm iT$. We shall apply the residue theorem. On the interior of C_T , the integrand is analytic except for a double pole at s = 2, arising from $\zeta^2(s-1)$. Let $R_a(f(s))$ denote the residue of a function f(s) at s = a. Therefore, by the residue theorem,

$$I_T = R_2 := R_2 \left(\frac{\left(1 - \frac{1}{2^{s-1}}\right)^2}{1 - \frac{1}{2^s}} \cdot \frac{\zeta(s-1)^2}{\zeta(s)} \frac{x^s}{s} \right)$$
$$= \frac{-36x^2\zeta'(2) + 12\gamma\pi^2x^2 - 3\pi^2x^2 + 10\pi^2x^2\log(2) + 6\pi^2x^2\log(x)}{6\pi^4}, \qquad (9.8)$$

where the residue can be calculated either by hand or by Mathematica.

We shall now calculate I_T by a second method. Let I_1, I_2, I_3, I_4 , respectively, denote the integrals over

$$\begin{split} & [c-iT,c+iT],\\ & [c+iT,\sigma_0+iT],\\ & [\sigma_0+iT,\sigma_0-iT],\\ & [\sigma_0-iT,c-iT]. \end{split}$$

We first find a suitable bound for I_2 . Accordingly,

$$I_{2} = -\frac{1}{2\pi i} \int_{\sigma_{0}+iT}^{c+iT} \frac{\left(1-\frac{1}{2^{s-1}}\right)^{2}}{1-\frac{1}{2^{s}}} \cdot \frac{\zeta(s-1)^{2}}{\zeta(s)} \frac{x^{s}}{s} ds$$
$$\ll \frac{1}{T} \int_{\sigma_{0}}^{c} x^{u} |\zeta(u-1+iT)|^{2} du.$$
(9.9)

From [4, Theorem 5.3], we have

$$|\zeta(u-1+iT)| \ll T^{1-(\sigma_0-1)},$$

when $\sigma_0 \leq u < 2$, and

$$|\zeta(u-1+iT)| \ll \log T,$$

when $2 \le u \le 3$. Using these bounds in (9.9), we obtain

$$I_2 \ll \frac{1}{T^{2\sigma_0 - 3}} \int_{\sigma_0}^2 x^u \, du + \frac{\log^2 T}{T} \int_2^c x^u \, du \ll \frac{x^2}{T^{2\sigma_0 - 3} \log x} + \frac{\log^2 T \, x^c}{T \log x}.$$
 (9.10)

By the same argument, the bound above holds for ${\cal I}_4$ as well.

We now turn to bounding I_3 . To that end,

$$I_{3} = \frac{1}{2\pi i} \int_{\sigma_{0} - iT}^{\sigma_{0} + iT} \frac{\left(1 - \frac{1}{2^{s-1}}\right)^{2}}{1 - \frac{1}{2^{s}}} \cdot \frac{\zeta(s-1)^{2}}{\zeta(s)} \frac{x^{s}}{s} ds$$

$$\ll x^{\sigma_{0}} \int_{-T}^{T} \frac{|\zeta(\sigma_{0} - 1 + it)|^{2}}{|\sigma_{0} + it|} dt$$

$$\ll x^{\sigma_{0}} + x^{\sigma_{0}} \int_{1}^{T} \frac{|\zeta(\sigma_{0} - 1 + it)|^{2}}{t} dt.$$
(9.11)

Using a dyadic decomposition, we find from (9.11) that

$$I_3 \ll x^{\sigma_0} + x^{\sigma_0} \sum_{\nu=0}^{\lfloor \log_2 T \rfloor} \frac{1}{2^{\nu}} \int_{2^{\nu}}^{2^{\nu+1}} |\zeta(\sigma_0 - 1 + it)|^2 dt.$$
(9.12)

From [8, p. 118, Theorem 7.2(A)], we have

$$\int_{2^v}^{2^{v+1}} |\zeta(\sigma_0 - 1 + it)|^2 \, dt \ll v 2^v.$$

Using the bound above in (9.12), we deduce that

$$I_3 \ll x^{\sigma_0} + x^{\sigma_0} \sum_{v=0}^{\lfloor \log_2 T \rfloor} v \ll x^{\sigma_0} \log^2 T.$$
(9.13)

In summary, by (9.10) and (9.13),

$$I_2 + I_3 + I_4 \ll \frac{x^2}{T^{2\sigma_0 - 3}\log x} + \frac{\log^2 T \ x^c}{T\log x} + x^{\sigma_0}\log^2 T.$$
(9.14)

To optimize the bounds for I_2, I_3 , and I_4 , we seek suitable constants such that the maximal exponents of x in the estimates of I_2 and I_3 are equal. Fix $0 < \epsilon < \frac{1}{10}$, and let

$$a = \frac{1}{4\epsilon} - \frac{1}{2}; \tag{9.15}$$

then a > 2. If we take

$$c = 3, \qquad T = x^a, \tag{9.16}$$

then the maximal exponent of x in the estimate of I_3 from (9.9) or (9.14) is

$$\max\{2 - a(2\sigma_0 - 3), c - a\} = \frac{3}{2} + \epsilon,$$

and the exponent of x in the estimate of I_2 , from (9.13) or (9.10), is

$$\sigma_0 = \frac{3}{2} + \epsilon$$

Putting these together in (9.14), we conclude that

$$I_2 + I_3 + I_4 = O_{\epsilon}(x^{\frac{3}{2} + \epsilon} \log^2 x).$$
(9.17)

Combining (9.17) with (9.8), we further conclude that

$$I_1 = I_T - I_2 - I_3 - I_4 = R_2 + O_\epsilon(x^{\frac{3}{2} + \epsilon} \log^2 x).$$
(9.18)

Returning to (9.6), we see that it remains to bound E(T), which is given by (9.7). With d(n) denoting the number of positive divisors of n below, we find that

$$E(T) \leq \frac{x^{c}}{T} \sum_{n=1}^{\infty} \frac{\sum_{d|n} d\phi(\frac{n}{d})}{n^{c} |\log(\frac{x}{n})|} = \frac{x^{c}}{T} \sum_{n=1}^{\infty} \frac{\sum_{d|n} \frac{n}{d}\phi(d)}{n^{c} |\log(\frac{x}{n})|} = \frac{x^{c}}{T} \sum_{n=1}^{\infty} \frac{\sum_{d|n} \frac{\phi(d)}{d}}{n^{c-1} |\log(\frac{x}{n})|}$$
$$\leq \frac{x^{c}}{T} \sum_{n=1}^{\infty} \frac{d(n)}{n^{c-1} |\log(\frac{x}{n})|} \leq 2\frac{x^{c}}{T} \sum_{n=1}^{\infty} \frac{1}{n^{c-3/2} |\log(\frac{x}{n})|}, \tag{9.19}$$

where we used the easy bound $d(n) \leq 2n^{1/2}$.

We now separate two cases. First, let $E_1(T)$ denote those summands on the right side of (9.19) for which $n \leq \frac{3}{4}x$ or $n \geq \frac{5}{4}x$. For these terms, $|\log(\frac{x}{n})|$ has a positive lower bound, and hence,

$$E_1(T) \ll \frac{x^c}{T} \frac{1}{x^{c-5/2}} = \frac{x^{5/2}}{T}.$$
 (9.20)

Second, let $E_2(T)$ denote those summands in E(T) for which $\frac{3x}{4} < n < \frac{5x}{4}$. Observe that $\sum_{n \leq y} a_n$ does not change in each interval, N < y < N + 1, for any fixed positive

integer N. Therefore, we may choose $x = N_0 + 1/2$, for each positive integer N_0 . Then for $\frac{3}{4}x < n < \frac{5}{4}x$, by the mean value theorem,

$$\left|\log\left(\frac{x}{n}\right)\right| \ge \frac{|x-n|}{x} \ge \frac{1}{2x}.$$

Therefore, the contribution in this case is

$$E_2(T) \ll \frac{x^{c+1}}{T} \sum_{\frac{3}{4}x < n < \frac{5}{4}x} \frac{1}{n^{c-3/2}} \ll \frac{x^{7/2}}{T}.$$
 (9.21)

Combining (9.20) and (9.21) and also using (9.15) and (9.16), we deduce that

$$E(T) \ll x^{7/2-a} \ll x^{3/2}.$$
(9.22)

In summary, from (9.6), (9.18), and (9.22) we have shown that

$$\sum_{n \le x}' a_n = I_1 + E(T)$$

= $R_2 + O_{\epsilon}(x^{\frac{3}{2} + \epsilon} \log^2 x) + O(x^{3/2})$
= $R_2 + O_{\epsilon}(x^{\frac{3}{2} + \epsilon} \log^2 x),$

where R_2 is given by (9.8). This completes the proof of Theorem 9.1.

References

- B. C. Berndt, Analytic Eisenstein series, theta-functions, and series relations in the spirit of Ramanujan, J. Reine Angew. Math. 304 (1978), 332–365.
- [2] B. C. Berndt and U. Dieter, Sums involving the greatest integer function and Riemann-Stieltjes integration, J. Reine Angew. Math. 337 (1982), 208–220.
- [3] G. H. Hardy and M. Riesz, The General Theory of Dirichlet's Series, Stechert-Hafner, New York, 1964.
- [4] A. J. Hildebrand. Introduction to Analytic Number Theory, Math 531 Lecture Notes, Fall 2005. URL: https://api.semanticscholar.org/CorpusID:1155809. Version 2013.01.07 (2006).
- [5] H. Rademacher, Topics in Analytic Number Theory, Springer-Verlag, New York, 1973.
- [6] H. Rademacher and E. Grosswald, *Dedekind Sums*, Carus Math. Monograph #16, Math. Assoc. Amer., 1972.
- [7] I. E. Shparlinski, Modular hyperbolas, Jpn. J. Math. 7 (2012), 235-294.
- [8] E. C. Titchmarsh. The Theory of the Riemann Zeta Function, Clarendon Press, Oxford, 1986.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, IL 61801, USA

Email address: berndt@illinois.edu

Department of Mathematics, University of Illinois, 1409 West Green Street, Urbana, IL 61801, USA

Email address: rnbhat2@illinois.edu

Department of Mathematics, 215 Carnegie Library, Syracuse University, Syracuse, NY 13244, USA

Email address: jlmeye01@syr.edu

Department of Mathematics, University of Illinois, 1409 West Green Street, Urbana, IL 61801, USA

Email address: likunx2@illinois.edu

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS, 1409 WEST GREEN STREET, URBANA, IL 61801, USA; INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O. BOX 1-764, BUCHAREST RO-70700, ROMANIA

Email address: zaharesc@illinois.edu

28