

THE ARITHMETIC OF CRITICAL VALUES I: EQUICRITICAL QUARTIC POLYNOMIALS

FRANCESCO NACCARATO

ABSTRACT. A polynomial f of degree d and coefficients in an algebraically closed field k defines a morphism $f : \mathbb{P}_k^1 \rightarrow \mathbb{P}_k^1$ which, if $\text{char}(k) \nmid d$, is unramified outside a finite set of points in the image: the critical values of f . In this work we establish a rigorous framework for the study of their arithmetic, which we carry out for $d = 4$ and $k = \overline{\mathbb{Q}}$, uncovering a connection to the arithmetic of elliptic curves. Recent progress in the theory of Weyl sums has sparked some interest in finding pairs of polynomials having the same critical values for “nontrivial” reasons: building on our analysis, we provide a complete classification of such pairs in the case of quartics over number fields.

1. INTRODUCTION

Let k be an algebraically closed field, $d \geq 2$ an integer such that $\text{char}(k) \nmid d$, and let $f \in k[x]$ be a polynomial of degree d . The (finite) *critical values* of f , denoted by C_f , are those $y \in k$ such that the equation $f(x) = y$ has less than d distinct solutions, i.e. the branching points of the ramified cover $f : \mathbb{A}_k^1 \rightarrow \mathbb{A}_k^1$ induced by f . They are obtained by evaluating f at its *critical points* Z_f , which are the roots of f' . These notions account for multiplicity, so Z_f and C_f are in principle multisets, and they are clearly finite. By compactifying, we can extend the definition of critical values to rational functions and, more generally, to ramified covers of $\mathbb{P}^1 = \mathbb{P}_k^1$ (to which we will refer just as “covers”):

Definition 1.1. *Let X/k be a projective curve and let $\phi : X \rightarrow \mathbb{P}^1$ be a morphism of degree d . The critical values of ϕ are its branching points $\text{Branch}(\phi)$. Two morphisms $\phi_1, \phi_2 : X \rightarrow \mathbb{P}^1$ with the same critical values are called equicritical.*

Notice how this makes $\infty \in \mathbb{P}^1$ a critical value of all polynomials. There is a natural action of $\text{Aut}_k(X)$ on the set of covers $\phi : X \rightarrow \mathbb{P}^1$, given by $\gamma \cdot \phi = \phi \circ \gamma^{-1}$; critical values behave well under this action as we clearly have $\text{Branch}(\gamma \cdot \phi) = \text{Branch}(\phi)$. Of special interest for us is the case $X = \mathbb{P}^1$, with $\text{PGL}_2(k)$ acting via fractional linear transformations. Observe that the subset of covers consisting of polynomials, which will be our focus, is preserved under this action precisely by the subgroup $\text{Aff}(k)$ of affine transformations. Following Arnold [1], we define:

Definition 1.2. *Two polynomials $f, g \in k[x]$ are said to be **topologically equivalent** if there exists $\lambda \in \text{Aff}(k)$ such that $f = g \circ \lambda$.*

We refer to the equivalence classes of this relation as *topological types*, adopting the notation $[f]$ for the topological type of f ; if, moreover, $f \in K[x]$ for a subfield $K \subset k$, we also set $[f]_K = \{f \circ \lambda, \lambda \in \text{Aff}(K)\}$, referring to any such set as a *K -topological type*. So, polynomials of the same topological type have the same critical values. In the following, when we refer to two polynomials

as (in)equivalent, we mean with respect to topological equivalence. We also call a polynomial f *normalized* if it is monic and $f(0) = 0$ (the normalization with f' monic is usually chosen in the literature, see e.g. [3]: our choice is motivated by purely expository reasons and does not affect the nature of the results).

The study of critical values turns out to be crucial in understanding various phenomena, ranging from holomorphic dynamics [10] to the inverse Galois problem [17] to, somewhat surprisingly, exponential sums: while investigating polynomial value sets over finite fields and rings, Kowalski and Soundararajan formulated [18, 1.2] a Fourier-analogue to the Davenport pair problem (see e.g. [14, 2] for an overview of this classical problem), observing that two integral polynomials whose reductions modulo a prime p are equicritical (and have all critical points in \mathbb{F}_p) give rise to a *Fourier-Davenport pair* over $\mathbb{Z}/p^2\mathbb{Z}$. They then point out that there are inequivalent such pairs in any degree at least 4. It is then natural to ask (see §5.3 for the application) if there exist inequivalent polynomials with the same global critical values, and not just modulo some prime. More generally, we pose the following problem:

Question 1 (Equicriticality problem). Given a field $K \subset k$, classify all elements of the set

$$\text{EC}(K, d) = \{([f], [g]) : f, g \in K[x], \deg f = \deg g = d, [f] \neq [g] \text{ and } C_f = C_g\}.$$

Observe that $\text{EC}(K, d)$ is a constructible set. This work deals with the number field case in degree 4: we give an explicit description of $\text{EC}(K, 4)$ for all number fields K . In particular, we show that $\text{EC}(K, 4)$ has dimension 3.

Remark 1.2.1. One can slightly modify the definition of $\text{EC}(K, d)$ by replacing topological types with K -topological types, but not much changes in characteristic 0: see Remark 4.1.

We saw that critical values are naturally attached to a topological type, rather than just a single polynomial: indeed, we can speak of equicritical topological types in the same way. Observe that $\text{Aff}(k)$ acts on the set of topological types of polynomials over k by

$$(1) \quad [f]^\lambda = [\lambda(f)],$$

with the resulting topological type having finite critical values $\lambda(C_f)$. This yields a component-wise action of $\text{Aff}(K)$ on $\text{EC}(K, d)$ which, when $\text{char}(k) = 0$, is generically free (for complex quartics there is only one exception, see Corollary 4.1.2); so, in this case, varying $\lambda \in \text{Aff}(K)$ allows us to produce infinitely many pairs of distinct equicritical K -topological types starting from a single one. In other words, at least in characteristic 0, if $\text{EC}(K, d)$ is nonempty then it has dimension at least 2: it is an open question to understand how this dimension behaves as the degree varies.

Given an ordered pair $c = (f, g)$ of polynomials, let us denote by \bar{c} the pair (g, f) and by $[c]$ the pair $([f], [g])$ of their topological types, while, for $\lambda \in \text{Aff}(k)$, we let $\lambda(c)$ be the pair $(\lambda(f), \lambda(g))$. In light of the previous remarks, we give the following definition:

Definition 1.3. *Let $K \subset k$ be a field. A **basis** for $\text{EC}(K, d)$ is a collection $\mathcal{E} = (c_i, i \in I)$ of ordered pairs of polynomials over K such that $\{[c_i] : i \in I\}$ is a set of representatives for the orbits of the $\text{Aff}(K)$ -action on $\text{EC}(K, d)$. A **quasi-basis** is a collection that can be extended to a basis by adding finitely many pairs.*

Set $\rho = 1 + \sqrt{3}$, $C = -720\sqrt{3} - 1248$, $R = 362 + 209\sqrt{3}$ and, for an algebraic number κ of degree 2, let $\bar{\kappa}$ be the other root of its minimal polynomial. Moreover, let $\omega \in \overline{\mathbb{Q}}$ be a

primitive third root of unity and, for a number field K , define $K' = K \setminus \{\omega, \omega^2\}$ and $K'' = K' \setminus \{0, 1, -2, -2\omega, -2\omega^2, \rho, \bar{\rho}, \omega\rho, \omega^2\rho, \omega\bar{\rho}, \omega^2\bar{\rho}\}$. We can now state our main result:

Theorem 1. *Let K be a number field and, for $t \in K''$, let $v = \frac{t^4(t-1)^3}{t+2}$. Then, the collection $\mathcal{E}_K = (c_t = (f_t, g_t), t \in K' \cup \{\infty\})$ given by:*

$$c_t = \left(x^4 - 6t^3x^2 - 8t^3x, -\frac{(t-1)^3v}{3(t+2)^3}x^4 + 2vx^2 + \frac{8}{3}vx - 8t^4(t^2 + t + 1) \right) \text{ for } t \in K'',$$

$$c_0 = \left(x^4 - x, -\frac{1}{48}x^4 - \frac{1}{4}x^2 + \frac{1}{6}x - \frac{1}{2} \right) = \overline{c_{-2}},$$

$$c_1 = \left(x^4 + 6x^3 + 9x^2, -3x^4 - 6x^3 \right) = \overline{c_\infty}$$

and

- if $\sqrt{3} \in K$,

$$c_\rho = (f_\rho, -f_\rho + 2C), \quad c_{\bar{\rho}} = (f_{\bar{\rho}}, -f_{\bar{\rho}} + 2\bar{C})$$

where $f_\rho(x) = x^4 - 6\rho^3x^2 - 8\rho^3x$ and $f_{\bar{\rho}}(x) = x^4 - 6\bar{\rho}^3x^2 - 8\bar{\rho}^3x$;

- if $\omega \in K$, $c_{-2\omega} = (g_0, \omega g_0) = \overline{c_{-2\omega^2}}$;
- if $\sqrt{3}, \omega \in K$,

$$c_{\omega\rho} = (f_\rho, iR(f_{\bar{\rho}} - \bar{C})) = \overline{c_{\omega\bar{\rho}}}, \quad c_{\omega^2\rho} = (f_\rho, iR(-f_{\bar{\rho}} + \bar{C})) = \overline{c_{\omega^2\bar{\rho}}},$$

is a basis for $EC(K, 4)$.

Remark 1.1. While it will follow from our proof, we can immediately check that the polynomials in a pair appearing in Theorem 1 are inequivalent: indeed, since (for $t \neq 1, \infty$) the coefficients of x^3 are all 0, if f_t and g_t were related by a linear change of variable, it would have to be a homothety, which is easily ruled out by looking at the other coefficients.

Kristiansen [19] has shown that, for each multiset Y of real numbers, there is exactly one real polynomial f with $C_f = Y$, up to topological equivalence. There is no hope for similar results to hold over number fields, but a natural question one can ask is that of classifying the sets of critical values of degree d polynomials. We are able to answer this question in the generic case for $d = 4$:

Theorem 2. *Let K be a number field and let y_1, y_2, y_3 be distinct algebraic numbers such that the elliptic curve $E : z^2 = (y - y_1)(y - y_2)(y - y_3)$ satisfies $j(E) \neq 0, 1728$. Then, there exists a quartic $f \in K[x]$ with $C_f = \{y_1, y_2, y_3\}$ if and only if E is defined over K and $j(E) = \frac{(u+3)^3(u+27)}{u}$ for some $u \in K^*$.*

Remark 2.1. The statement of Theorem 2 suggests that it is useful to interpret the critical values of a quartic over K as the 2-torsion of an elliptic curve: we will see why this is the case in §3.1. The theorem actually provides a characterization of almost all curves that arise this way; still, many questions about these curves and their connection to critical values remain worth asking: for instance, one may be interested in classifying the integral j -invariants that are realized for $K = \mathbb{Q}$, in which case an elementary p -adic argument shows that only $u = \pm 3^k$, $0 \leq k \leq 6$, work. Additional questions and applications, especially in light of the constructions of §4.2, will be the subject of a separate upcoming work of the author.

The paper is structured as follows: in §2 we review the literature on critical values of polynomials, setting up the algebro-geometric picture of the problem; in doing so, we prove a generalization of a result of Arnold, which may bear some independent interest. A key ingredient in the proof of Theorem 1 is a thorough study of the (reduced) Hurwitz space \mathcal{H}_4 associated to quartic covers of \mathbb{P}^1 with monodromy of polynomial type: in §3, after introducing Hurwitz spaces, we construct the aforementioned reduced space for any degree $d \geq 3$, and we study its geometry and arithmetic. §4 is centered around \mathcal{H}_4 : we prove that it is a certain well-known modular curve, and use this to establish Theorem 2. §5 is devoted to the equicriticality problem for quartics over number fields: we first give a nonconstructive proof - which may adapt to similar questions - for the existence of a parametrization like that of Theorem 1, and then go on to compute the explicit equations for the polynomials appearing in the statement, by exploiting the rich structure arising from the link to the arithmetic of elliptic curves.

1.1. Acknowledgments. I am grateful to my advisor Emmanuel Kowalski, for recommending this topic and for the helpful advice he offered during the development of this work, to Davide Lombardo, for working out the details of the proof of Lemma 4.2 and for his suggestion of following an explicit approach to the construction of the Hurwitz space, and to Jordan Ellenberg, for sketching the ideas behind the fiber product formulation of the equicriticality problem and for pointing me to the paper of Rubin and Silverberg on elliptic curves with constant mop p representations.

The author was partially supported by the Swiss National Science Foundation (grant number 219220). This work is part of ongoing research within the scope of the author's PhD thesis.

2. CRITICAL VALUES OF POLYNOMIALS: ARITHMETIC AND GEOMETRY

From now on, we assume that either $\text{char}(k) = 0$ or $\text{char}(k) > d$. Given a normalized polynomial $f \in k[x]$, we attach to it the set of its critical points, as a (closed) point $\{x_i : x_i \in Z_f\} \in \mathbb{A}^{d-1}/S_{d-1}$. Since with any point $\hat{x} \in \mathbb{A}^{d-1}/S_{d-1}$ we can associate the normalized formal integral (always assumed to have 0 constant term) of $\prod_{\hat{x}=x_i}(x - x_i)$ multiplied by d , we get a bijection:

$$(2) \quad \begin{array}{ccc} \{\text{normalized polynomials } f \in k[x] \text{ of degree } d\} & \xleftarrow{1:1} & \mathbb{A}^{d-1}/S_{d-1} \\ f & \longmapsto & Z_f. \end{array}$$

If one does not care about the subfield of definition of the coefficients, the study of polynomials with given critical values can be carried out just for normalized polynomials: the leading coefficient of $f(ax + b)$ is a^d times that of f , and its constant term is $f(b)$. Therefore, in light of (2), the starting point for our analysis will be the relation between critical points and critical values. We will then recover information about fields of definition via the study of rational points on Hurwitz spaces.

If the normalized polynomial f corresponds to $\hat{x} = \{x_1, \dots, x_{d-1}\} \in \mathbb{A}^{d-1}/S_{d-1}$, its finite critical values y_1, \dots, y_{d-1} are given by:

$$(3) \quad y_j = \sum_{i=0}^{d-1} (-1)^i \frac{d}{d-i} e_i(\hat{x}) x_j^{d-i},$$

where $e_i(\hat{x})$ is the elementary symmetric function of degree i in $d - 1$ variables. This translates the study of critical values for polynomials of degree d to that of the covering $\theta_d : \mathbb{A}^{d-1} \rightarrow \mathbb{A}^{d-1}$ defined by (3), whose fiber above a point (y_1, \dots, y_{d-1}) of \mathbb{A}^{d-1} corresponds under (2) (and the quotient) to the set of normalized polynomials of degree d having as finite critical values its coordinates. An

elementary but important observation to make is that, for any subfield $K \subset k$, the finite critical values of $f \in K[x]$ are the (only) simple roots of a polynomial defined over K : this can be seen directly by letting $G_K = \text{Gal}(\overline{K}/K)$ act on both sides of (3), since θ_d is defined over K .

Let U_d be the affine open set $\{\prod_{1 \leq i < j \leq d-1} (y_i - y_j) \neq 0\} \subset \mathbb{A}^{d-1}$ and let $V_d \subset U_d$ be the closed subscheme given by $y_1 = 0$. Polynomials with finite critical values in U_d - i.e., distinct - are called *Morse polynomials*. Arnold [1, p.5, Theorem], following Davis [8], Thom [23] and Zdravkovska [25], shows that for $d \geq 3$ there are d^{d-3} topological types of complex Morse polynomials of degree d with given critical values. We generalize his result to our fields k which, we recall, are algebraically closed and have characteristic either 0 or larger than d :

Proposition 2.1. *Let $d \geq 3$. For any $(y_1, \dots, y_{d-1}) \in U_d$ we have $\#\{[f] : f \in k[x], C_f = \{y_1, \dots, y_{d-1}\}\} = d^{d-3}$.*

Remark 2.1.1. Proposition 2.1 shows that, in characteristic different from 2 and 3, the equicriticality problem is trivial for cubic polynomials: we have $EC(k, 3) = \emptyset$.

Arnold's proof reduces the question to that of counting trees with d numbered edges, which had already been solved by Cayley. Our strategy builds on Beardon, Carne and Ng's study [3, Lemma 2.4], over \mathbb{C} , of the Jacobian J_d of θ_d above nonzero distinct critical values, extending it to our fields k and to the case where one of the critical values is 0 (for a generalization of their study in the case of equal but nonzero critical values, see [12, Theorem B]). Let the morphism $\tilde{\theta}_d$ be given by (3) except for the equation for y_1 , where the right-hand side is divided by x_1 .

Lemma 2.2. (i) *The morphism θ_d restricted to $\theta_d^{-1}(U_d \setminus V_d)$ is étale;*
(ii) *the morphism $\tilde{\theta}_d$ restricted to $\tilde{\theta}_d^{-1}(V_d)$ is étale.*

Proof. (i) Let $Q \in U_d \setminus V_d$, $P \in \theta_d^{-1}(Q)$ and set $J = J_d$ for convenience. The part of the proof of Lemma 2.4 in [3] in which the invertibility of J at P is shown clearly holds for the fields we are considering, since it only relies on integral identities that hold formally in this case, and on a polynomial of degree $\leq d$ with $d+1$ roots being identically 0. Then, the dimension of the cotangent space of our variety at P is $(d-1) - \text{rk}(J(P)) = 0$, so in particular it must equal its local dimension at P , and the Jacobian criterion (which we can apply even in positive characteristic since $k = \bar{k}$) tells us that θ_d is smooth at P . Clearly the local dimension at P is an upper bound for the relative dimension at P and hence, since smooth morphisms of relative dimension 0 are étale, by arbitrariness of P and Q we get the desired claim;

(ii) Let now $Q = (y_i) \in V_d$, $P = (x_i) \in \tilde{\theta}_d^{-1}(Q)$ and set $\tilde{J} = \text{Jac}_{\tilde{\theta}_d}$: we want to show independence of the columns of $\tilde{J}(P)$, from which the claim will follow as in (i). Suppose on the contrary that there exist $\lambda_1, \dots, \lambda_{d-1} \in k$ such that $\sum_{j=1}^{d-1} \lambda_j \tilde{J}_{ij}(P) = 0$ for $1 \leq i \leq d-1$. Let us start by assuming $x_1 \neq 0$: then, respectively from the first equation in the proof of Lemma 2.4 in [3] and from the definition of $\tilde{\theta}_d$ (our different normalization for θ_d by a factor of d does not impact these equations), we get:

$$(4) \quad \int_0^{x_i} \sum_{j=1}^{d-1} \lambda_j \prod_{\substack{1 \leq k \leq d-1 \\ k \neq j}} (w - x_k) dw = 0, \quad 2 \leq i \leq d-1$$

and

$$(5) \quad x_1 \int_0^{x_1} \sum_{j=1}^{d-1} \lambda_j \prod_{\substack{1 \leq k \leq d-1 \\ k \neq j}} (w - x_k) dw - \lambda_1 \int_0^{x_1} \prod_{1 \leq k \leq d-1} (w - x_k) dw = 0.$$

Now, observe that the second integral in (5) multiplied by d is just the value at x_1 of the normalized polynomial with critical points x_1, \dots, x_{d-1} , so none other than $y_1 = 0$. Therefore, denoting by $f_\lambda(w)$ the common integrand left in both equations, we see that the polynomial $F_\lambda(x) = \int_0^x f_\lambda(w) dw$ has degree at most $d - 1$ and vanishes for $x = 0, x_1, \dots, x_{d-1}$, and hence vanishes identically. But then so must do its derivative f_λ ; on the other hand, for any $1 \leq j \leq d - 1$, we find $f_\lambda(x_j) = \lambda_j \prod_{\substack{1 \leq k \leq d-1 \\ k \neq j}} (x_j - x_k)$: since $P \in U_d$, this means $\lambda_j = 0$, and we are done.

If $x_1 = 0$, (4) still holds, so $F_\lambda(w) = \int_0^w f_\lambda(w) dw$ defined as above vanishes on $0, x_2, \dots, x_{d-1}$; therefore, if it does not vanish identically - in which case we conclude as above - we must have

$$(6) \quad F_\lambda(x) = Cx \prod_{2 \leq j \leq d-1} (x - x_j)$$

for some $C \in k^*$. On the other hand, a simple direct computation with (3) yields $\tilde{J}(P)_{(1,1)} = (-1)^{d-1} d^{\frac{\prod_{2 \leq j \leq d-1} x_j}{2}} \neq 0$, while $\tilde{J}(P)_{(1,j)} = 0$ for $2 \leq j \leq d - 1$ as x_1 appears in each monomial of these partial derivatives, so we immediately deduce $\lambda_1 = 0$. But then, we find:

$$0 \stackrel{(4)}{=} f_\lambda(x_1) \stackrel{x_1=0}{=} F'_\lambda(0) \stackrel{(6)}{=} (-1)^{d-2} C \prod_{2 \leq j \leq d-1} x_j \neq 0,$$

which gives the desired contradiction. □

We are now ready to prove Proposition 2.1:

Proof of Proposition 2.1. As θ_d is étale over $U_d \setminus V_d$ by Lemma 2.2 (i), it is unramified with finite fibers. Applying Bezout's Theorem, we find that the cardinality of the fiber above $Q \in U_d \setminus V_d$ equals d^{d-1} minus the number of points at infinity of (3); as its right-hand side is homogeneous, the points at infinity are found by letting it vanish for $1 \leq j \leq d - 1$, and hence correspond to the *nonzero* vectors of critical points of normalized polynomials having all finite critical values 0. But a polynomial h with $C_h = \{0\}$ (as a set) must vanish at all roots of h' , necessarily with higher multiplicity, which implies that h' - and hence h - have a single root, of multiplicity equal to the degree. Therefore, if h is normalized then it must be of the form $h(x) = x^d$, which shows that there are no intersection points at infinity: we have $|\theta_d^{-1}(Q)| = d^{d-1}$. Now, any normalized polynomial f in our fiber is equivalent to d^2 normalized polynomials in the same fiber, given by $f(\mu_d x + b)$ for μ_d any d -th root of unity in k and b any of the d distinct roots of f . Therefore, we are left with d^{d-3} topological types over each point of $U_d \setminus V_d$.

Let now $Q \in V_d$, and assume that the entry of Q that is 0 is the first one (we can safely do so, since all our equations are symmetric). If $f \in \theta_d^{-1}(Q)$, then one of the critical values of f is 0, so f has a common root with f' and hence a double root - exactly one, since the critical values are distinct. Therefore, in computing the number of topological types in each fiber, we need to divide its cardinality by $d(d - 1)$ instead of d^2 . On the other hand, in virtue of how $\tilde{\theta}_d$ is defined, the

preimage of Q under θ_d has the same number of elements (without multiplicity) as the preimage of Q under $\tilde{\theta}_d$, since the first equation of (3) has the form $f_1(x_1, \dots, x_{d-1}) = 0$ with $x_1^2 \mid f_1$. But Lemma 2.2 (ii) tells us that $\tilde{\theta}_d$ is étale over V_d , so by the same argument as above it has fibers of cardinality $(d-1)d^{d-2}$, and we are done. \square

3. THE HURWITZ SPACE OF MORSE POLYNOMIALS

Hurwitz spaces are, roughly speaking, moduli spaces of covers $X \rightarrow \mathbb{P}^1$ with given ramification properties. We assume for simplicity that the field of definition k of our covers is \mathbb{C} (the following constructions generalize to any algebraically closed field of characteristic 0, but the nice topological setting offered by Riemann surfaces is missing; see [4, 1.1] for an account), keeping in mind that compact Riemann surfaces come from algebraic curves by Serre's GAGA. So, let $\mathbb{P}^1 = \mathbb{P}_{\mathbb{C}}^1$ and consider all objects as base-changed to \mathbb{C} when necessary. Moreover, let \mathcal{U}_r be the moduli space for r distinct unordered points in \mathbb{P}^1 equipped with its natural algebraic structure of affine variety over \mathbb{Q} , whereby the field of definition of the point (y_1, \dots, y_r) is that of the polynomial $\prod_{y_i \neq \infty} (y - y_i)$.

3.1. Moduli spaces of G -covers. We begin by giving an outline of the constructions in Hurwitz theory that are relevant for our study; a more comprehensive introduction can be found, e.g., in the work of Fried and Volklein [16]. Let r and d be positive integers, let G be a subgroup of S_d and let $C = (C_1, \dots, C_r)$ an unordered tuple of conjugacy classes in G . We will refer to such a triple (r, G, C) as *datum*, where the embedding of G in S_d is implicit.

Definition 3.1. A *Nielsen class* for the triple (r, G, C) is an element of

$$Ni_r(G, C) = \{(g_1, \dots, g_r) \in G^r : \exists \sigma \in S_r : g_i \in C_{\sigma(i)} \ \forall i = 1, \dots, r, \ g_1 \cdot \dots \cdot g_r = 1 \text{ and } \langle \{g_i\} \rangle = G\}.$$

An *inner Nielsen class* is an element of $Ni_r(G, C)^{in} = Ni(G, C)/G$, where G acts component-wise by conjugation.

An *absolute Nielsen class* is an element of $Ni_r(G, C)^{abs} = Ni(G, C)/N_{S_d}(G)$, where the normalizer acts component-wise by conjugation.

Recall that a G -cover is a pair $(\phi : X \rightarrow \mathbb{P}^1, \delta)$ - usually denoted just by ϕ - where ϕ is a Galois cover and $\delta : \text{Aut}(\phi) \rightarrow G$ is a group isomorphism. Let ϕ have $r_\phi = r$ critical values Q_1, \dots, Q_r , let $Q_0 \in \mathbb{P}^1$ be any other point and, for $1 \leq i \leq r$, let U_i be a disk neighborhood of Q_i that does not contain any other critical value but contains Q_0 . Then, taking a generator ρ_i of $\pi_1(U_i \setminus \{Q_i\}, Q_0)$ and lifting it to X gives a deck transformation $g_i \in G$ of X , via the isomorphism δ . Let $\text{Cl}(g_i)$ be the respective conjugacy class in G and denote by $C(\phi)$ the unordered tuple $(\text{Cl}(g_1), \dots, \text{Cl}(g_r))$: we say that ϕ branches with local monodromy $C(\phi)$; this is a good definition since picking different generators for the fundamental groups only changes the induced deck transformations by inner automorphisms of $\text{Aut}(\phi)$. We have just described a way to attach to a G -cover ϕ a datum, namely $(r_\phi, \text{Aut}(\phi), C(\phi))$, and by choosing the ρ_i 's so that $\prod_{i=1}^r \rho_i = 1$ under the inclusions $\pi_1(U_i \setminus \{Q_i\}, Q_0) \hookrightarrow \pi_1(\mathbb{P}^1 \setminus \{Q_1, \dots, Q_r\}, Q_0)$, $1 \leq i \leq r$, we have also attached to ϕ the Nielsen class (g_1, \dots, g_r) .

As in the case of polynomials, it is more natural to attach a datum to a suitable equivalence class of G -covers: we generalize the definition of topological equivalence by saying that two G -covers $(\phi : X \rightarrow \mathbb{P}^1, \delta)$ and $(\psi : Y \rightarrow \mathbb{P}^1, \xi)$ are equivalent, or isomorphic, if there is an isomorphism of covering spaces $\gamma : X \rightarrow Y$ that commutes with the respective actions of G , i.e.

$\delta(g) = \xi(\gamma g \gamma^{-1}) \forall g \in \text{Aut}(\phi)$. We adopt the terminology *G-cover class* and the notation (ϕ) for the equivalence class of ϕ under this relation. Isomorphic covers clearly have the same critical values, while the additional condition on the G -actions implies that the datum associated to a G -cover is an invariant of its class. Hence, we have a (noncanonical, due to its dependence on the choice of generators ρ_i) map

$$(7) \quad \Phi : \left\{ \begin{array}{l} \text{classes of } G\text{-covers with datum} \\ (r, G, C) \text{ branching at } Q_1, \dots, Q_r \end{array} \right\} \longrightarrow \text{Ni}_r(G, C)^{\text{in}},$$

sending (ϕ) to the inner Nielsen class representative of (g_1, \dots, g_r) as defined above. The following result (see [13, 1.2] and [14, 2.2.1]) shows that Φ is a bijection:

Theorem 3 (Riemann Existence Theorem, inner version). *Given a datum (r, G, C) and r points Q_1, \dots, Q_r in \mathbb{P}^1 , $\text{Ni}(G, C)^{\text{in}}$ is in bijection with the set of classes of G -covers $\phi : X \rightarrow \mathbb{P}^1$ that branch over Q_1, \dots, Q_r with local monodromy C .*

Remark 3.1. Here the genus of X is uniquely determined by the datum: this is the Branch Cycle Lemma, see [15, 5.1].

Notice how in Theorem 3 the branching set is fixed, while in the situation that interests us it varies. The Riemann Existence Theorem for families (see again [13, 1]) enables us to construct Hurwitz spaces as parameter families of Nielsen classes over \mathcal{U}_r : in particular (see [4, Theorem 1.1]), the **inner Hurwitz space**

$$(8) \quad \Psi_r : H_r^{\text{in}}(G, C) \longrightarrow \mathcal{U}_r$$

has the property that for any $Q = (Q_1, \dots, Q_r) \in \mathcal{U}_r$, the fiber above Q is in bijection with the set of G -cover classes that branch at Q with local monodromy C . Moreover, it admits a unique algebraic model as an affine variety over a number field $k_0 = k_0(C)$ such that Ψ_r is defined over k_0 and, for any extension K/k_0 , any $\sigma \in G_K$ and any $(\phi) \in H_r^{\text{in}}(G, C)(\overline{\mathbb{Q}})$, we have:

$$(9) \quad \sigma(\phi) = (\sigma \phi).$$

As mentioned at the end of §1, our focus will mainly lie on a *reduced* Hurwitz space. Observe that there is an *outer* $\text{PGL}_2(\mathbb{C})$ -action on G -cover classes, given by $(\phi)^\alpha = (\alpha(\phi))$. We say that two G -cover classes in the same orbit are *$\text{PGL}_2(C)$ -equivalent*, and refer to an equivalence class as a *G/PGL_2 -cover class*. Then, given the same datum as above, the reduced Hurwitz space $H_r^{\text{red}}(G, C)$ is the affine variety of dimension $r - 3$ obtained as the quotient of $H_r^{\text{in}}(G, C)$ by this action. Indeed, as we already observed in (1) for polynomials and affine transformations, we have $\text{Branch}(\alpha(f)) = \alpha(\text{Branch}(f))$, and local monodromy around critical values is clearly preserved by the action. This gives a commutative diagram:

$$\begin{array}{ccc} H_r^{\text{in}}(G, C) & \xrightarrow{\Pi_r} & H_r^{\text{red}}(G, C) \\ \Psi_r \downarrow & & \downarrow \beta_r \\ \mathcal{U}_r & \xrightarrow{\pi_r} & \mathcal{I}_r \end{array}$$

with $\mathcal{G}_r = \mathcal{U}_r/\mathrm{PGL}_2(\mathbb{C})$. A classical result of Mumford and Fogarty [20, Theorem 1.1] shows that both quotient spaces are well-defined in the category of affine k_0 -varieties. In virtue of its construction, K -rational points on $H_r^{\mathrm{red}}(G, C)$ correspond with $\mathrm{PGL}_2(\mathbb{C})$ -orbits of G -cover classes (ϕ) (with ϕ taken to be defined over \overline{K}) such that for each $\sigma \in G_K$ there exist $\alpha_\sigma \in \mathrm{PGL}_2(\mathbb{C})$ and an isomorphism γ_σ of G -covers from $\alpha_\sigma(\sigma\phi)$ to ϕ .

If $r = 4$, then \mathcal{G}_4 is the usual moduli space of elliptic curves $Y(1)$, since the j -invariant of an elliptic curve is that of the quadruple of its 2-torsion points. A fiber of the morphism

$$\beta_4 : H_4^{\mathrm{red}}(G, C) \longrightarrow Y(1)$$

is then naturally identified with the set of G/PGL_2 -cover classes with monodromy C and representatives that branch over the 2-torsions of the elliptic curves with j -invariant at the base. We will refer to the latter as the *critical j -invariant* of a G - or G/PGL_2 -cover class.

By construction, β_4 only ramifies above critical j -invariants of quadruples with “large” stabilizer in $\mathrm{PGL}_2(\mathbb{C})$, i.e. the elliptic j -invariants $0, 1728$. It is then well known that $H_r^{\mathrm{red}}(G, C)$ must be a quotient of the upper half-plane by a finite index subgroup $\Gamma = \Gamma(G, C)$ of the modular group $\mathrm{PSL}_2(\mathbb{Z})$, with β_4 induced by the inclusion $\Gamma < \mathrm{PSL}_2(\mathbb{Z})$. In other words, we are dealing with a (not necessarily congruence) modular curve. See again [13, 3] for a general overview of reduced Hurwitz spaces of dimension 1 as modular curves.

3.2. Lifting points on Hurwitz spaces, and the case of polynomials. Given $x \in H_r^{\mathrm{red}}(G, C)$, we call any of its preimages in $H_r^{\mathrm{in}}(G, C)(K)$ a “lift” of x . The question of which points $x \in H_r^{\mathrm{red}}(G, C)(K)$ lift to $H_r^{\mathrm{in}}(G, C)(K)$ is a rich and open one, whose main application is in the domain of the inverse Galois problem: indeed, the IGP reduces to finding rational points on inner Hurwitz spaces, thanks to Hilbert’s Irreducibility Theorem. Looking for K -rational points on $H_r^{\mathrm{red}}(G, C)$ is easier due to it being a rational quotient of lower dimension, especially in the case $r = 4$ when it is a curve. However, these points rarely lift to points of $H_r^{\mathrm{in}}(G, C)(K)$; this behavior is largely controlled by two factors:

- (a) the lifting of points from $\mathcal{G}_r(K)$ to $\mathcal{U}_r(\overline{K})$;
- (b) the *base invariant* of the $G/\mathrm{PGL}_2(\mathbb{C})$ -cover class x , i.e. the conjugacy class in $\mathrm{PGL}_2(\mathbb{C})$ of the stabilizer of any of its lifts to $H_r^{\mathrm{in}}(G, C)$.

In [4], Cadoret gives a thorough account of the theory and proves some strong lifting results, especially for G/PGL_2 -cover classes with nontrivial base invariant: see for instance [4, Lemma 3.12, (2)]. In point (1) of the same result, the author observes that, for $r = 4$, a point in $\mathcal{G}_r(K)$ always lifts to a point in $\mathcal{U}_r(K)$, as this is equivalent to fields of moduli being fields of definition for elliptic curves. Notice that a stronger condition than x having trivial base invariant is the $\mathrm{PGL}_2(\mathbb{C})$ -stabilizer of the branching set of any (i.e. all) of its lifts being trivial; for our purposes, it is enough to prove the following lifting result:

Lemma 3.2. *For any datum (r, G, C) , any tower of extensions $K_0/K/k_0$ and any $x \in H_r^{\mathrm{red}}(G, C)(K)$, if:*

- (1) $\beta_r(x)$ lifts to a K_0 -rational point on \mathcal{U}_r ;
- (2) $\mathrm{Stab}_{\mathrm{PGL}_2(\mathbb{C})}(\mathrm{Branch}(\phi_0)) = \{I\}$ for a lift ϕ_0 of x ,

then there exists a G -cover $\phi : X \longrightarrow \mathbb{P}^1$ such that $(\phi) \in H_r^{\mathrm{in}}(G, C)(K_0)$ and $\Pi_r((\phi)) = x$.

Proof. Let $Q \in \mathcal{U}_r(K_0)$ be a lift of $\beta_r(x)$ and (ϕ) any G -cover class in $H_r^{\text{in}}(G, C)(\overline{\mathbb{Q}})$ such that $\Pi_r((\phi)) = x$ and $\Psi_r((\phi)) = Q$ (which exists as the $\text{PGL}_2(\mathbb{C})$ -action on G -cover classes and that on critical values are compatible). We know that for each $\sigma \in \text{Gal}(\overline{K_0}/K_0)$ there exist $\alpha_\sigma \in \text{PGL}_2(\mathbb{C})$ and an isomorphism of G -covers $\gamma_\sigma \in \text{Aut}(X)$ from $\alpha_\sigma(\sigma\phi)$ to ϕ . Since $\text{Branch}(\sigma\phi) = \sigma Q = Q$, α_σ must lie in its stabilizer, which is conjugate to that of ϕ_0 and hence trivial. Therefore, α_σ is the identity, and we obtain $\sigma\phi = \phi(\gamma_\sigma)$ and that γ_σ commutes with the G -actions of ϕ and $\sigma\phi$; in other words, $(\phi) \in H_r^{\text{in}}(G, C)(K_0)$ by (9) and the paragraph preceding it. \square

Let us now go back to polynomials. A Morse polynomial $f \in k[t]$ defines a cover $f : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ of the same degree d , branching at $C_f \cup \{\infty\}$ with local monodromy C_{pol} given by a transposition at each of the finite critical values and a d -cycle at ∞ (see [14, 2.2.2]). It is a classical result going back to Hilbert that any choice of representatives for the classes in C_{pol} whose product is the identity must generate the full symmetric group, so the Galois group of the cover $f(t) - y$ is S_d . In order to apply Hurwitz theory to our question, we set $r = d$, $G = S_d$ and $C = C_{\text{pol}}$. Then necessarily $X = \mathbb{P}^1$ (see Remark 3.1), and we get the inner Hurwitz space

$$\Psi_d^{\text{pol}} : H_d^{\text{in}}(S_d, C_{\text{pol}}) \rightarrow \mathcal{U}_d,$$

whose fiber above $\{y_1, \dots, y_d\} \in \mathcal{U}_d$ is in bijection with the set of classes of S_d -covers $\phi : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ branching over the y_i 's with local monodromy C_{pol} . We call these covers *polynomial S_d -covers*. Since $C_{\text{pol}}^m = C_{\text{pol}}$ for odd m , the results in [16, 1.1] imply that our inner Hurwitz space is defined over \mathbb{Q} .

While our polynomials are not Galois covers themselves, we have:

Lemma 3.3. *Polynomial S_d -cover classes with given branch locus are in bijection with isomorphism classes of mere covers with the same branch locus and local monodromy C_{pol} . Moreover, each such class is $\text{PGL}_2(\mathbb{C})$ -equivalent to the isomorphism class of a polynomial.*

Proof. A bijection analogous to (7) holds for absolute Nielsen classes, in this case with isomorphism classes of (mere) covers with given datum and branch locus (G being the automorphism group of the Galois closure): this gives rise [15, Proposition 6.3] to the *absolute Hurwitz space* relative to the datum. If $G = S_d$, inner and absolute Nielsen classes coincide in virtue of Definition 3.1, so in this case the inner and absolute Hurwitz spaces are isomorphic (see [2, 3.1.3] for more details).

Since $\text{PGL}_2(\mathbb{C})$ acts transitively on \mathbb{P}^1 , given $(\phi) \in H_d^{\text{in}}(S_d, C_{\text{pol}})$ (taken as a mere cover class) there is $\alpha \in \text{PGL}_2(\mathbb{C})$ such that the critical value where $(\phi)^\alpha$ has a d -cycle is ∞ . But a map with a d -cycle at ∞ has a pole of order d at its only preimage t_1 , so it is a polynomial in a local coordinate around $1/t_1$, and the second part of the claim follows. \square

In other words, Lemma 3.3 tells us that once we pass to the reduced space we are, up to equivalences, dealing just with polynomials. The choice of not focusing directly on absolute Hurwitz spaces is motivated by our, albeit short, treatment of the issue of lifting rational points, which the literature usually covers in the inner case, in virtue of the connection to the inverse Galois problem.

Notice the similarity between Ψ_d^{pol} and the map θ_d of Section 2: if we denote by ι_d the inclusion $U_d/S_{d-1} \hookrightarrow \mathcal{U}_d$ given by $\{y_1, \dots, y_{d-1}\} \mapsto \{y_1, \dots, y_{d-1}, \infty\}$, and by \mathcal{W}_d its image, then $\Psi_d^{\text{pol}}|_{\Psi_d^{-1}(\mathcal{W}_d)}$ and $\iota_d \circ \theta_d$ are both rational maps from an affine variety to \mathcal{W}_d whose fibers convey information about the polynomials branching at the base. Nevertheless, it is not the *same* information: Ψ_d

accounts for all classes of covers with polynomial monodromy, not just (normalized) polynomials, and more importantly, it carries data about the field of definition of the covers by (9) (see Theorem 4).

We refer to $\beta_d : H_d^{\text{red}}(S_d, C_{\text{pol}}) \rightarrow \mathcal{G}_d$ as the Hurwitz space of (Morse) polynomials of degree d . Let us compute its degree:

Proposition 3.4. *Let $d \geq 3$. Then β_d has degree d^{d-3} , and it is unramified above the $\text{PGL}_2(\mathbb{C})$ -orbits with generic stabilizer.*

Proof. By Lemma 3.3, $\deg(\beta_d)$ is equal to the number of isomorphism classes of polynomials in a generic fiber. Since these clearly biject with topological types, there are d^{d-3} of them above each $\{y_1, \dots, y_d\} \in \mathcal{U}_d$ by Proposition 2.1; therefore, it suffices to show that the stabilizer for the $\text{PGL}_2(\mathbb{C})$ -outer action of the class (f) of a polynomial is generically trivial.

For $d = 3$ there is nothing to prove (in particular, our Hurwitz space is a rational point). For $d > 4$, the stabilizer $\text{Stab}_{\text{PGL}_2(\mathbb{C})}(\text{Branch}(f))$ is itself generically trivial, so we are done. For $d = 4$, the former stabilizer is generically a Klein four-group, so each nontrivial element τ is a double transposition; in particular, the critical value ∞ is not fixed, so the 4-cycle in the local monodromy of $\tau(f)$ is not at ∞ , and therefore $\tau(f) \notin (f)$. \square

As we just observed, a point $x \in H_d^{\text{red}}(S_d, C_{\text{pol}})(\mathbb{C})$ always lifts to a complex polynomial f : from now on, we will use the expression “topological type” also for (f) , whenever this does not create ambiguity. In order to shed light on the equicriticality problem, we need to be able to relate the minimal field of definition of f to that of x ; our spaces have the following lifting property:

Theorem 4. *Let K be a number field;*

- (i) *any $x \in H_4^{\text{red}}(S_4, C_{\text{pol}})(K)$ lifts to a quartic defined over K , so $H_4^{\text{red}}(S_4, C_{\text{pol}})(K)$ is in bijection with the set of $\text{PGL}_2(\mathbb{C})$ -orbits of topological types of quartics over K ;*
- (ii) *if $d > 4$, any $x \in H_d^{\text{red}}(S_d, C_{\text{pol}})(K)$ in the generic fiber of β_d lifts to a polynomial defined over a K -extension of degree at most $d!$.*

Proof. (i) For the moment we only prove the claim in the case of nonelliptic $\beta_4(x)$; the remaining cases will be dealt with in §4. The datum $(4, S_4, C_{\text{pol}})$ satisfies condition (1) in Lemma 3.2 with $K_0 = K$ by the aforementioned Lemma 3.12 in [4], but the stabilizers are Klein groups, not trivial. Let us keep the notation of the proof of Lemma 3.2, but with $f := \phi$ taken as a polynomial; then, α_σ must lie in the stabilizer of Q , which is a Klein four-group by the nonellipticity hypothesis. Assume α_σ is not the identity; as we saw in the proof of Proposition 3.4, its action on the critical values is free, and hence ${}^\sigma f$ would have a 2-cycle at infinity, which is absurd. Therefore α_σ is the identity, so for any $\sigma \in G_K$ there exists $\gamma_\sigma \in \text{PGL}_2(\overline{\mathbb{Q}})$ such that

$$(10) \quad {}^\sigma f = f \circ \gamma_\sigma.$$

For mere covers of \mathbb{P}^1 without automorphisms, (10) implies that K is a field of definition for f (see the Introduction in [9]; it is *a priori* only a field of moduli, see 2.2 in the same work for the precise definitions), and our covers indeed have no automorphisms since these biject with the center of the Galois group of the cover (see [5, 1]), which in our case is $Z(S_d) = \{e\}$.

Therefore, there is some $\alpha \in \mathrm{PGL}_2(\overline{\mathbb{Q}})$ such that

$$(11) \quad f \circ \alpha \in K(t).$$

If α is affine we are done, so suppose $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$, $c \neq 0$. Then α can be written as $\beta\alpha'$ with β affine and α' having top-left entry 0. Therefore, up to replacing f with $f \circ \beta$, γ_σ with $\beta^{-1}\gamma_\sigma\beta$ and α with α' , which does not affect (10) and (11), we can take $a = 0$, so say $\alpha = \begin{pmatrix} 0 & 1 \\ c & d \end{pmatrix}$.

Let us now combine (10) and (11); for any $\sigma \in G_K$, we have

$$(12) \quad f = (f \circ \alpha) \circ \alpha^{-1} = \sigma(f \circ \alpha) \circ \alpha^{-1} = f \circ (\gamma_\sigma(\sigma\alpha)\alpha^{-1}),$$

which implies that the fractional linear transformation on the right-hand side is the identity, since it induces an automorphism of the cover $f(t) - y$. Therefore, since γ_σ is affine by (10),

$$\text{so must be } \sigma\alpha\alpha^{-1} = \begin{pmatrix} 1 & 0 \\ -d\frac{\sigma c}{c} + \sigma d & \frac{\sigma c}{c} \end{pmatrix}.$$

Looking at the bottom-left coefficient we get $\sigma(\frac{d}{c}) = \frac{d}{c} \forall \sigma \in G_K \implies l = \frac{d}{c} \in K$, and hence $f\left(\frac{c^{-1}}{t+l}\right) \in K(t)$ by (11). But now, as the denominator in the variable is K -rational itself, simply taking common denominator $(t+l)^4$ shows that the coefficients a_i of t^i in f satisfy $a_i c^{-i} \in K$, so we finally get:

$$f\left(\frac{t}{c}\right) \in K[t].$$

- (ii) Since the generic stabilizer of a point in \mathcal{U}_d is trivial for $d > 4$, Lemma 3.2 gives us a polynomial lift with field of moduli K_0 equal to the field of definition of any lift of $\beta_d(x)$, and by [4, Lemma 3.10] we can take $[K_0 : K] \leq d!$. The same argument as above gives a polynomial lift defined over K_0 . □

Remark 4.1. The paragraph after (10) shows that, in defining $\mathrm{EC}(K, d)$, we can generically just take K -topological types; that is, to obtain all rational polynomials equivalent to a given generic rational polynomial f , we can simply take its rational topological type: indeed, for a Morse polynomial f , if we have $f \circ \lambda \in K[t]$ for some $\lambda \in \mathrm{Aff}(\overline{K})$, then for any $\sigma \in G_K$ we have that $\lambda^{-1} \circ \sigma \lambda$ is an automorphism of the cover $f(t) - y$, and hence trivial. Therefore, λ is defined over K .

4. THE CASE OF QUARTICS: MODULARITY

4.1. The Hurwitz space of quartics as a modular curve. From now on, K will denote a number field. We consider the case $d = 4$: compactifying our reduced Hurwitz space, we get a rational modular curve

$$\beta_4 : \mathcal{H}_4 \longrightarrow X(1),$$

to which we refer simply as the Hurwitz space of quartics, since the information about those that are non-Morse is carried by the fiber above the cusp. We adopt the notation j_{CV} for our coordinate on the level 1 modular curve.

Let us focus on the geometry of our modular curve $\mathcal{H}_4 = X(\Gamma)$. Recall the formula for the genus of quotients of the upper half plane by modular subgroups [11, p.66 and 68]:

$$(13) \quad g(X(\Gamma)) = 1 + \frac{[\mathrm{PSL}_2(\mathbb{Z}) : \Gamma]}{12} - \frac{\epsilon_2}{4} - \frac{\epsilon_3}{3} - \frac{\epsilon_\infty}{2},$$

where the ϵ_i 's count points of period 2 and 3, and cusps, respectively. Since $[\mathrm{PSL}_2(\mathbb{Z}) : \Gamma] = \deg \beta_4 = 4$, the index term gives a contribution of $\frac{1}{3}$. As the genus is nonnegative, $\epsilon_\infty \geq 1$, and 3 is coprime with 4, surely $\epsilon_3 = 1$. Then necessarily $g(\mathcal{H}_4) = 0$. Notice how *a priori* we have the two possibilities (2, 1) and (0, 2) for $(\epsilon_2, \epsilon_\infty)$.

Having degree less than 7, our Γ is a congruence subgroup by a result of Wohlfhart [24, Theorem 5], so \mathcal{H}_4 is a congruence modular curve. By the list of genus 0 congruence subgroups in the Cummins-Pauli database [7], it is either $X_0(3)$ or $4A^0 \backslash \mathbb{H} = X(4A^0)$ as a curve over \mathbb{C} (and hence over $\overline{\mathbb{Q}}$). Observe how these groups realize the two aforementioned possibilities for the number of elliptic points above $j_{CV} = 1728$ and of cusps.

To determine whether \mathcal{H}_4 is $X_0(3)$ or $X(4A^0)$ we just need to compute ϵ_2 . Recall that quadruples $\widehat{y} = \{y_1, y_2, y_3, y_4\} \in \mathcal{U}_4$ with elliptic j -invariant have stabilizer larger than the usual Klein group (in [4, 4.2.2], Cadoret gives explicit descriptions for the Legendre representatives), in particular D_8 for $j_{CV} = 1728$: \mathcal{H}_4 has no elliptic point above $j_{CV} = 1728$, i.e. $\epsilon_2 = 0$, precisely if this stabilizer acts freely on topological types in the fiber. Indeed, the elliptic point above $j_{CV} = 0$ corresponds to (the orbit of) the topological type of $p_0(x) = x^4 - x$, which is fixed by multiplication by ω . The following result shows that our reduced Hurwitz space is the modular curve $X_0(3)$:

Lemma 4.1. *The outer $\mathrm{PGL}_2(\mathbb{C})$ -action on polynomial topological types with critical j -invariant 1728 is free. Therefore, $\mathcal{H}_4 \simeq X_0(3)$ over $\overline{\mathbb{Q}}$.*

Proof. Let f be a polynomial having critical values as in the hypotheses and let $S_f = \mathrm{Stab}_{\mathrm{PGL}_2(\mathbb{C})}((f))$. Since $j_{CV} = 1728$, we can suppose without loss of generality that $C_f = \{0, 1, -1\}$ by applying an affine transformation to the polynomial: the new stabilizer will simply be the conjugate under this transformation of the old one; moreover, we can set $f(0) = 0$ by a change of variable $x \mapsto x + c$, which does not change the topological type. Any element of S_f must fix ∞ as the only critical value of (f) with 4-cycle local monodromy: from the aforementioned description of the stabilizers for Legendre representatives, we get that the only affine nontrivial element in S_f is $z \mapsto -z$. Therefore, if the statement were false, we would have algebraic numbers a, b such that

$$(14) \quad -f(x) = f(ax + b),$$

which immediately gives $a = \zeta_8$ a primitive 8-th root of unity. As f and $-f$ have the same roots and 0 is one of them, iteratively substituting in the right-hand side of (14) gives:

$$(15) \quad 0 = f(0) = -f(b) = f(b(\zeta_8 + 1)) = -f(b(\zeta_8^2 + \zeta_8 + 1)) = f(b(\zeta_8^3 + \zeta_8^2 + \zeta_8 + 1)).$$

Since all expressions in ζ_8 appearing in (15) are nonzero (notice how this is false for ω , which is the corresponding value in the $j_{CV} = 0$ case) and pairwise distinct, and f has at most four roots, we get $b = 0$, but then $f(x) = x^4$ which does not have distinct critical values. \square

Remark 4.1.1. The modular curve $X_0(3)$ is already known to be a reduced Hurwitz space for a different datum, specifically that of certain *Lattès maps*, see [13, 4.2] and [6].

Notice how Proposition 3.4 together with Lemma 4.1 imply the following nice result:

Corollary 4.1.2. *The only topological type of quartic Morse polynomials over \mathbb{C} having nontrivial stabilizer for the outer action of $\text{Aff}(\mathbb{C})$ is $[p_0] = [x^4 - x]$.*

At this point, we have a rational curve \mathcal{H}_4 that fits into a diagram:

$$\begin{array}{ccc} \mathcal{H}_4 & \xrightarrow{\psi_4} & X_0(3) \\ \beta_4 \downarrow & & \swarrow \pi_3 \\ & & X(1) \end{array}$$

where we take π_3 (not to be confused with one of the maps in the bottom row of the diagram in §3.1) as a rational map making $X_0(3)$ into the moduli space of isomorphism classes $[(E, C)]$ of elliptic curves E with a cyclic 3-torsion subgroup C (so that $j(E) = \pi_3([(E, C)])$). *A priori*, the isomorphism ψ_4 is only defined over $\overline{\mathbb{Q}}$: one may ask whether it is forced to be defined over \mathbb{Q} . The answer is affirmative:

Lemma 4.2. *For any curve C/\mathbb{Q} with a rational map $\beta : C \rightarrow X(1)$ and an isomorphism $\psi : C \rightarrow X_0(3)$ sitting in a diagram as above, ψ is defined over \mathbb{Q} . In particular, $\mathcal{H}_4 \simeq_{\mathbb{Q}} X_0(3)$.*

Proof. As all objects and morphisms are defined over \mathbb{Q} except at most for ψ , if we act on the diagram by an element σ of $G_{\mathbb{Q}}$ we get another diagram with ${}^{\sigma}\psi = \sigma \circ \psi \circ \sigma^{-1}$ in place of ψ , from which we extract an automorphism ${}^{\sigma}\psi \circ \psi^{-1}$ of $X_0(3)$ over $X(1)$. Therefore, it suffices to show that the only such automorphism is the identity; but these automorphisms are classified by Galois theory: their set bijects with the quotient by $\Gamma_0(3)$ of its normalizer N in $\Gamma(3)$, since $X(3) \rightarrow X(1)$ is the Galois closure of $X_0(3) \rightarrow X(1)$. As the index $[\Gamma(3) : \Gamma_0(3)] = 12/4 = 3$ is prime, the normalizer must be $\Gamma_0(3)$ itself, and we are done. \square

Remark 4.2.1. As the proof shows, this is generally false if in the statement we replace the pair $(X_0(3), \pi_3)$ with (C_0, π) having nontrivial automorphisms: for example, let C_0 be an elliptic curve over \mathbb{Q} and π the usual 2-covering, and let C be a nontrivial quadratic twist; the isomorphism α is then only defined over a quadratic extension of \mathbb{Q} , the generator of whose Galois group sends α to $[-1]_{C_0} \circ \alpha$, with $[-1]_{C_0}$ the multiplication by -1 on C_0 .

Corollary 4.2.2. *We have $X_0(3) \simeq_{\mathbb{Q}} X_1(3)$ and hence, for any elliptic curve E/\mathbb{Q} with a cyclic subgroup of order 3 defined over \mathbb{Q} , there exist an elliptic curve E'/\mathbb{Q} with a rational point of (exact) order 3 and such that $j(E) = j(E')$.*

Proof. As $\Gamma_0(3) = \Gamma_1(3)$, we have an isomorphism $\psi : X_1(3) \rightarrow X_0(3)$ over \mathbb{C} , and hence over $\overline{\mathbb{Q}}$, that commutes with the projections to $X(1)$. The claim now follows from Lemma 4.2. \square

4.2. A direct method. Our Hurwitz space \mathcal{H}_4 has genus 0 and a certain algebraic structure $\mathcal{H}_4 \simeq_{\mathbb{Q}} \mathbb{P}^1$ coming from Hurwitz theory. Since a point on \mathcal{H}_4 represents a $\text{PGL}_2(\mathbb{C})$ -orbit of topological types, it is natural to ask if we can describe its coordinate explicitly as a function of the orbit. Recall that the morphisms θ_d of Section 2 map critical points to critical values: one may then guess that β_4 maps the j -invariant of the critical points of the orbit to that of the critical values. Indeed, this map is well-defined: the critical points of a topological type form a $\text{PGL}_2(\mathbb{C})$ -orbit, while $Z_{\alpha(f)} = Z_f$

for $\alpha \in \widetilde{\mathrm{PGL}}_2(\mathbb{C})$ (basically, their behavior under the two $\mathrm{PGL}_2(\mathbb{C})$ -actions we introduced is the mirror image of that of critical values); moreover, to any $j \in \mathbb{C}$ we can associate the $\mathrm{PGL}_2(\mathbb{C})$ -orbit of the (unique, by the above discussion) topological type of $\int_0^x (u - u_1)(u - u_2)(u - u_3) du$ where $\{u_1, u_2, u_3, \infty\}$ is any quadruple with j -invariant j . In summary, by sending an orbit to the j -invariant of its critical points, we get a bijection:

$$(16) \quad \{ \mathrm{PGL}_2(\mathbb{C})\text{-orbits of topological types of complex quartics} \} \xleftarrow{1:1} \mathbb{P}^1(\mathbb{C}).$$

Identify \mathcal{H}_4 with \mathbb{P}^1 via the j -invariant j of the critical points: the above correspondence yields a map $\tilde{\beta}_4 : \mathcal{H}_4 \rightarrow X(1)$, $j \mapsto j_{CV}$. Note that even if we know that such a map exists, it need not be β_4 *a priori*: nonetheless, we will now prove that $\tilde{\beta}_4$ is indeed an algebraic morphism defined over \mathbb{Q} by computing it explicitly, so necessarily $\tilde{\beta}_4 = \beta_4$ by the uniqueness property for algebraic models of Hurwitz spaces. In the following, the variable j will represent a point in \mathcal{H}_4 with the above identification.

Before we work out an explicit model for β_4 , observe how the above remarks show that *any* point $j \in \mathcal{H}_4(K) \setminus \{\infty\}$ lifts to a quartic defined over K , since we can just integrate the Weierstrass polynomial of any K -rational elliptic curve with j -invariant j : this completes the proof of Theorem 4.

Fix an isomorphism $u : \mathbb{P}^1 \rightarrow X_0(3)$ and let us take for π_3 the model given by $u \mapsto \frac{(u+3)^3(u+27)}{u}$ (see [22, p.23]). For a complex cubic curve given by a Weierstrass equation $E : y^2 = x^3 + Ax + B$, let¹:

$$(17) \quad f_E(x) = 12 \int_0^x (s^3 + As + B) ds - A^2.$$

Moreover, let $\Delta_E = -16(4A^3 + 27B^2)$ be the discriminant of E (note that we are allowing Δ_E to vanish), let $j = -1728 \frac{(4A)^3}{\Delta_E} \in \mathbb{P}^1(\mathbb{C})$ be its j -invariant and $j_{CV} \in \mathbb{P}^1(\mathbb{C})$ be that of the cubic $E^{CV} : z^2 = p_E(x)$, where $p_E(x)$ is the monic polynomial vanishing at the critical values of f_E . Since j and j_{CV} represent respectively the j -invariant of the critical points and of the critical values of a variable quartic, there is no ambiguity with our coordinates for \mathcal{H}_4 and $X(1)$, as the latter stand for the same things in our moduli problem. Then, (3) and a series of lengthy computations give:

$$(18) \quad j_{CV} = \frac{2^{20} A^3 (A^3 - 54B^2)^3}{B^2 \Delta_E^3} = 2^{-18} \frac{j(j - 1536)^3}{j - 1728} = \pi_3 \left(\frac{j}{64} - 27 \right),$$

so the second intermediate expression is our explicit model for β_4 and we find $\psi_4(j) = \frac{j}{64} - 27$ in the diagram in §4.1. Notice that (18) implies that the integrals of the Weierstrass polynomials of \mathbb{C} -isomorphic elliptic curves have the same critical j -invariant, which is not a surprise since it is equivalent to the right direction of (16). Moreover, it allows us to characterize the critical j -invariants of quartics over K : let

$$CJ(K) = \{v \in K \cup \{\infty\} : \exists f \in K[x] \text{ s.t. } \deg f = 4 \text{ and } j(C_f) = v\}.$$

Proposition 4.3. *We have:*

$$CJ(K) = \left\{ \frac{(u+3)^3(u+27)}{u}, u \in K \cup \{\infty\} \right\}.$$

¹our choice of linear normalization for the integral turns out to be convenient in computing the transformation in (25), but any other choice would work.

Proof. Theorem (4), (i) and Lemma 4.2 imply that

(19)

$$\text{CJ}(K) \cup \{\beta_4(\infty)\} = \{\infty\} \cup \left\{ \frac{(u+3)^3(u+27)}{u}, u \in K \cup \{\infty\} \right\} = \left\{ \frac{(u+3)^3(u+27)}{u}, u \in K \cup \{\infty\} \right\}$$

(the singleton in the intermediate expression is added *a priori* to make sure we are accounting for non-Morse quartics), while (18) tells us that $\beta_4(\infty) = \infty = \beta_4(1728) \subset \text{CJ}(K)$, so we are done. \square

Let us examine the fiber of β_4 above $j_{CV} = 0$: since π_3 vanishes only for $u = -3$ and $u = -27$ (with order respectively 3 and 1, in accordance with the ramification structure we previously found), (18) tells us that the values of j corresponding to these two preimages are 0 and 1536: in other words, the elliptic curves with j -invariant 0 and 1536 are precisely those with Weierstrass integrals having critical j -invariant 0. What about elliptic curves with $j = 1728$? Since $\beta_4(1728) = \infty$, they have non-Morse Weierstrass integrals: this is indeed immediate to check with $E : y^2 = x^3 - x$.

We are now almost ready to prove Theorem 2: we only need the following strengthening of Theorem 4, (i):

Lemma 4.4. *Let $j_{CV} \in K \setminus \{0, 1728\}$. For any $j \in \mathcal{H}_4(K)$ such that $\beta_4(j) = j_{CV}$ and any quadruple $\hat{y} = \{y_1, y_2, y_3, \infty\} \in \mathcal{U}_4(K)$ with $j(\hat{y}) = j_{CV}$, j lifts to a quartic $f \in K[X]$ with $C_f \cup \{\infty\} = \hat{y}$.*

Proof. Let f_0 be a lift of j as in the statement of Theorem 4 and set $\hat{w} = C_{f_0} \cup \{\infty\}$. As $j(\hat{w}) = j_{CV}$ by construction, there is $\lambda \in \text{PGL}_2(\overline{\mathbb{Q}})$ such that $\lambda(\hat{w}) = \hat{y}$ (as sets); up to composing with the double transposition in $\text{Stab}_{\text{PGL}_2(\overline{\mathbb{Q}})}(\hat{y})$ that swaps ∞ and $\lambda(\infty)$, we can assume λ to be affine. Therefore, $\lambda(f_0)$ is a quartic with critical values \hat{y} , but it may not be defined over K . Still, it has nonelliptic critical j -invariant and it satisfies equation (10) for any $\sigma \in G_K$, so the by argument after that equation its topological type contains a rational polynomial. \square

Proof of Theorem 2. The claim follows immediately from Lemma 4.4 and Proposition 4.3. \square

5. EQUICRITICAL QUARTICS AND ELLIPTIC CURVES

We have come to the question of inequivalent equicritical quartics: the goal of this section is to prove Theorem 1. Let C_4 be the non-diagonal irreducible component of $\mathcal{H}_4 \times_{X(1)} \mathcal{H}_4$ (there may be

a priori up to three such components, but the results of §5.1 will show that there is only one) and let C_4 be its normalization; we refer to a point of $C_4 \setminus C_4$ (or to a pair lifting it) as “exceptional”. A pair of inequivalent equicritical quartics (f, g) over a number field K clearly defines a K -point on C_4 . Moreover, Lemma 4.4 implies that, generically, such a point lifts to a pair of K -rational equicritical quartics. Therefore, our task is reduced, at least generically, to that of finding rational points on C_4 .

5.1. Elliptic curves with constant mod 3 representation. As $H_4^{\text{red}}(S_4, C_{\text{pol}})(K) \simeq_{\mathbb{Q}} Y_0(3)$ as covers of $Y(1)$ by Lemma 4.2, C_4 turns out to be \mathbb{Q} -isomorphic to the curve X_3 studied by Rubin and Silverberg [21] in the context of parametrizing elliptic families with given mod p Galois representation. Indeed, in virtue of Corollary 4.2.2 we have

$$H_4^{\text{red}}(S_4, C_{\text{pol}})(K) \times_{Y(1)} H_4^{\text{red}}(S_4, C_{\text{pol}})(K) \simeq Y_0(3) \times_{Y(1)} Y_1(3),$$

so removing the diagonal we precisely obtain the second equivalent definition for Y_3 given in [21, 1]. This is enough for us to give a proof of a nonconstructive version of Theorem 1:

Lemma 5.1. *There exists a quasi-basis for $EC(K, 4)$ generically parametrized by $\mathbb{P}^1(K)$.*

Proof. Observe that any collection of pairs of K -rational equicritical quartics $\{(f_P, g_P), P \in C_4(K)\}$ parametrized by $C_4(K)$ (where we take such a pair for points above $j_{CV} \in \{0, 1728, \infty\}$ only if it exists) makes up a quasi-basis for $EC(K, 4)$. Since the genus of X_3 is 0 and it has rational points (see [21, 1.1]), the claim follows. Alternatively, if one did not know about the modular characterization of the fiber product, the genus can be computed via Riemann-Hurwitz for the desingularization, and the rational point $(0, 1536)$ can be exhibited. Even without knowing the explicit form for β_4 we worked out in §4, but just the *geometry* of the rational map $\beta_4 : \mathcal{H}_4 \rightarrow X(1)$, one can exhibit the same rational point by observing that, since the fiber $\beta_4^{-1}(0)$ has two elements and one of them is rational (it corresponds to the orbit of (p_0)), the other one must be too. \square

The curve X_3 is itself a moduli space of elliptic curves, with the additional structure induced by the self-fiber product of $X_0(3)$: in the aforementioned work of Rubin and Silverberg, the authors give a certain rational model $t : \mathbb{P}^1 \rightarrow X_3$ of X_3 , and write down the respective affine equation for the universal elliptic curve over X_3 as

$$E_t : Y^2 = X^3 - 27t(t^3 + 8)X + 54(t^6 - 20t^3 - 8),$$

having j -invariant

$$(20) \quad j_t = j(E_t) = 27 \left(\frac{t(t^3 + 8)}{t^3 - 1} \right)^3.$$

We take as our model for X_3 the one obtained by compactifying the self-fiber product of $(X_0(3), \pi_3)$. Thanks to (18) and (20), we can explicitly parametrize $C_4(K)$: this amounts to finding two rational functions $x_1(t), x_2(t)$ such that the pairs $(x_1(t), x_2(t))$ as t ranges in $K \cup \{\infty\}$ parametrize $X_3(K)$, and then pulling them back via ψ_4 . Searching for rational functions such that

$$(21) \quad \pi_3(x(t)) = j_t$$

by comparing the right-hand side of (20) with π_3 , it is not particularly hard to figure out that

$$(22) \quad x_1(t) = \frac{27}{(t^3 - 1)} \text{ and } x_2(t) = \frac{3(t - 1)^3}{t^2 + t + 1} = x_1 \left(\frac{t + 2}{t - 1} \right)$$

work. We then have:

Proposition 5.2. *The multiset $\{(\psi_4^{-1}(x_1(t)), \psi_4^{-1}(x_2(t))), t \in K \cup \{\infty\}\}$ parametrizes $C_4(K)$.*

Proof. It suffices to show that $X_3(K)$ is parametrized by $\{(x_1(t), x_2(t)), t \in K \cup \{\infty\}\}$. One easily verifies that (21) holds for $x(t) = x_i(t)$, $i = 1, 2$. Now, since $x_1(s) = x_1(t) \iff t \in \langle \omega \rangle s$, (22) implies that $x_1(t) \neq x_2(t)$ generically, so $t \mapsto (x_1(t), x_2(t))$ defines a morphism $\chi : \mathbb{P}^1 \rightarrow X_3$: we just need to show that its degree is 1.

Again by (22), this amounts to verifying that $x_2(\omega^k t) \neq x_2(t)$ does not hold identically for $k = 1, 2$; rewriting $x_2(t)$ as $3 \frac{(t-1)^4}{t^3-1}$, we see that equality holds when either $t \in \{\omega, \omega^2\}$ or $(\omega^k t - 1)^4 = (t - 1)^4$. Since the polynomials appearing at the left and right-hand sides are distinct, the claim follows. Let us remark that, since $t \mapsto \frac{t+2}{t-1}$ is an involution, (22) implies that $\chi(t)$ and $\chi(\frac{t+2}{t-1})$ give the same unordered pair. \square

Remark 4.2. By solving $x_1(t) = x_2(t)$, we find that for $t \in \{\omega, \omega^2, -2\omega, -2\omega^2, 1 \pm \sqrt{3}\}$ we get points on C_4 from the compactified diagonal of $\mathcal{H}_4 \times_{X(1)} \mathcal{H}_4$, with “coordinate” respectively ∞ for the first

two values, 1536 for the second two and $1152 \pm 384\sqrt{3}$ for the last two (as expected, these are the four ramification points of β_4). In particular, C_4 has two exceptional points above each of $j = \infty$ and $j = 1536$, but there are more: solving $\chi(s) = \chi(t)$, we find that the values $t = \omega(1 \pm \sqrt{3})$ yield the remaining ones, satisfying respectively $\chi(t) = \chi(\omega t) = (1152 \pm 384\sqrt{3}, 1152 \mp 384\sqrt{3})$.

Proposition 5.2 and (22) give:

$$(23) \quad C_4(K) = \left\{ \left(1728 \frac{t^3}{t^3-1}, 1728 + 192 \frac{(t-1)^3}{t^2+t+1} \right), t \in K \cup \{\infty\} \right\},$$

where we treat pairs appearing with multiplicity as distinct points. Recall the notation $\rho = 1 + \sqrt{3}$, $\bar{\rho} = 1 - \sqrt{3}$, let $\tilde{K} = K \setminus \{\omega, \omega^2, -2\omega, -2\omega^2, \rho, \bar{\rho}\}$ and set $j_1(t) = 1728 \frac{t^3}{t^3-1}$, $j_2(t) = 1728 + 192 \frac{(t-1)^3}{t^2+t+1}$. Then (23) means precisely that, for $t \in K \cup \{\infty\}$, the integrals - normalized as in (17) - of the Weierstrass polynomials of any two rational elliptic curves E_t, F_t with $j(E_t) = j_1(t)$, $j(F_t) = j_2(t)$, are two quartics with critical j -invariant j_t , and they are inequivalent if $t \in \tilde{K} \cup \{\infty\}$.

In order to parametrize a basis for $\text{EC}(K, 4)$, we need to find an equicritical pair over K for each t (if there is one), that is, we need to explicitly go through the procedure described in the proof of Lemma 4.4 for nonelliptic $j_{CV} = j_t$, while taking care of the excluded values separately. Specifically, given $t \in \tilde{K} \setminus \{-2, 0, 1\}$, we:

- (1) pick elliptic curves E_t, F_t over K with $j(E_t) = j_1(t)$, $j(F_t) = j_2(t)$, yielding us quartic lifts $f^t = f_{E_t}$, $g^t = f_{F_t} \in K[X]$ of $j_1(t)$ and $j_2(t)$, with the notation of (17);
- (2) find the lift \tilde{g}^t of $j_2(t)$ such that $C_{\tilde{g}^t} = C_{f^t}$, by computing $\lambda_t(g^t)$ with $\lambda_t \in \text{Aff}(K)$ the transformation mapping $F_t^{CV}[2]$ to $E_t^{CV}[2]$ (we will not need to pick another representative of the topological type since, as we will see, the family λ_t is defined over $\mathbb{Q}(t)$).

The first step is easy: it is well known that the elliptic curve $E_j : y^2 = x^3 + \frac{3j}{1728-j}x + \frac{2j}{1728-j}$ has j -invariant j for all $j \neq 0, 1728$, so we can just take:

$$E_t : y^2 = x^3 - 3t^3x - 2t^3 \text{ and } F_t : y^2 = x^3 - 3\left(\frac{t+2}{t-1}\right)^3 x - 2\left(\frac{t+2}{t-1}\right)^3, \quad t \in \tilde{K} \setminus \{-2, 0, 1\},$$

corresponding to

$$(24)$$

$$f^t(x) = 3x^4 - 18t^3x^2 - 24t^3x - 9t^6 \text{ and } g^t(x) = 3x^4 - 18\left(\frac{t+2}{t-1}\right)^3 x^2 - 24\left(\frac{t+2}{t-1}\right)^3 x - 9\left(\frac{t+2}{t-1}\right)^6$$

The second step is in principle more complex, but a careful examination of the critical values for the first few negative integer values for t suggests for λ_t the form

$$(25) \quad z \mapsto -\frac{t^4(t-1)^6}{3(t+2)^4}z - 36t^4(t^2+t+1),$$

which we easily verify with **SageMath**. As anticipated, the λ_t 's turn out to be defined over K , so there is no need to look for a rational representative in $[g^t]$. Let us remark that it is always the case that the 2-torsions of two isomorphic elliptic curves E, F over K with nonelliptic j -invariant are related by an affine transformation over K : any $\overline{\mathbb{Q}}$ -isomorphism $\phi : E \rightarrow F$ descends to an

isomorphism $\lambda : E[2] \rightarrow F[2]$ of the 2-torsions as Galois modules, so in particular it is affine. If λ is not defined over K , there is $\sigma \in G_K$ such that $\lambda \circ \sigma \lambda^{-1} \neq I$, so in particular $\phi \circ \sigma \phi^{-1} \in \text{Aut}_{\mathbb{C}}(E) \setminus \langle \iota \rangle$ with ι the elliptic involution; in other words, E has nontrivial automorphisms and hence elliptic j -invariant.

Putting together Proposition 5.2, (24) and (25), we have proved:

Lemma 5.3. *The collection of pairs of Morse polynomials $\mathcal{E}'_K = (c_t = (f_t, g_t), t \in \tilde{K} \setminus \{-2, 0, 1\})$ with*

$$(26) \quad \begin{aligned} f_t &= \frac{1}{3}f^t + 3t^6 = x^4 - 6t^3x^2 - 8t^3x, \\ g_t &= -\frac{t^4(t-1)^6}{3(t+2)^4}x^4 + 2\frac{t^4(t-1)^3}{t+2}x^2 + 8\frac{t^4(t-1)^3}{3(t+2)}x + t^4(t+2)^2 - 12t^4(t^2+t+1) + 3t^6 \end{aligned}$$

is a quasi-basis for $EC(K, 4)$.

More specifically, substituting in (23) we see that the values $t = -2, 0, 1, \infty$ we left out correspond to the pairs $(1536, 0), (0, 1536), (\infty, 1728), (1728, \infty) \in C_4(K)$, while Remark 4.2 tells us that the other points in $C_4(K)$ which we might be missing representatives for are the eight exceptional ones, along with the two diagonal points $(1152 \pm 384\sqrt{3}, 1152 \pm 384\sqrt{3})$.

5.2. Elliptic fibers and cusps. Let us now deal with equicritical pairs lifting the remaining eight points in $C_4(K)$. We start by examining the situation above $j_{CV} \in \{0, 1728\}$:

- (1) $j_{CV} = 0$: let us deal with the points $(0, 1536)$ and $(1536, 0)$ first; we already found a representative (p_0) for the $\text{PGL}_2(\mathbb{C})$ -orbit of topological types corresponding to $j = 0$, and we can do the same for $j = 1536$ simply by integrating the Weierstrass polynomial of any elliptic curve with j -invariant 1536. Still, the affine transformation relating the critical values of the two representatives is not guaranteed to be rational (recall that this is why Lemma 4.4 fails for elliptic j_{CV}), but it happens to be so in some cases: choosing the right elliptic curve with the help of the **LMFDB** and **SageMath**, we obtain the two additional equicritical pairs $c_0 = (f_0, g_0)$ and $c_{-2} = (g_0, f_0)$ with

$$(27) \quad f_0 = p_0 = x^4 - x, \quad g_0 = -\frac{1}{48}x^4 - \frac{1}{3}x^2 + \frac{1}{6}x - \frac{1}{2}.$$

Let us now look at the two exceptional points in this fiber, both represented by the coordinates $(1536, 1536)$ in our parametrization: we know that, for any $\hat{y} \in \mathcal{U}_4(K)$ such that $j(\hat{y}) = j_{CV}$, there are three topological types with critical values \hat{y} that lift $j = 1536$. We have just constructed a rational one $[g_0]$; the other two will be given by $[\lambda(g_0)]$ and $[\lambda^{(2)}(g_0)]$ with λ the nontrivial affine transformation preserving C_{g_0} (we know by Corollary 4.1.2 that these are distinct topological types), which, since $C_{g_0} = C_{p_0}$, is simply the multiplication by ω . As $\mathbb{Q}(\omega)$ is clearly the minimal field of definition of any conjugate of λ , we get two additional pairs

$$(28) \quad c_{-2\omega} = (g_0, \omega g_0) = \overline{c_{-2\omega^2}}$$

if and only if $\omega \in K$ (and no other, since the pairs $(g_0, \omega^2 g_0)$ and $(\omega g_0, \omega^2 g_0)$ belong to the orbit of one those appearing in (28)).

- (2) $j_{CV} = 1728$: we consider the diagonal points $(1152 \pm 384\sqrt{3}, 1152 \pm 384\sqrt{3})$ first; they correspond to the values $t = \rho, \bar{\rho}$ in our parametrization, so the quartics $f_\rho, f_{\bar{\rho}}$ (following the notation of §5.1) are lifts for $j = 1152 + 384\sqrt{3}$ and $j = 1152 - 384\sqrt{3}$ respectively. As we just saw in the case $j_{CV} = 0$, to find the other lifts making up the pairs we need to apply to f_ρ and $f_{\bar{\rho}}$ the affine transformations preserving their respective critical value sets. **SageMath** shows that $f_\rho = x^4 - 6\rho^3x^2 - 8\rho^3x$ satisfies $C_{f_\rho} = (-v + C, C, v + C)$ for $v = 72\sqrt{362\sqrt{3} + 627}$ and $C = -720\sqrt{3} - 1248$. As f_ρ and $f_{\bar{\rho}}$ are Galois conjugates, so are C_{f_ρ} and $C_{f_{\bar{\rho}}}$, and hence we get the additional pairs

$$(29) \quad c_\rho = (f_\rho, -f_\rho + 2C), \quad c_{\bar{\rho}} = (f_{\bar{\rho}}, -f_{\bar{\rho}} + 2\bar{C})$$

if and only if $\sqrt{3} \in K$.

The previous paragraph enables us to also find pairs lifting the four exceptional points in this fiber, corresponding to $t = \omega\rho, \omega^2\rho, \omega\bar{\rho}, \omega^2\bar{\rho}$, as their orbits in $\text{EC}(K, 4)$ are necessarily obtained by taking one orbit from each of the pairs in (29). Again with the help of **SageMath**, we find that the linear map sending $C_{f_{\bar{\rho}}}$ to C_{f_ρ} is $z \mapsto iRz - iR\bar{C}$, with $R = 362 + 209\sqrt{3}$, yielding the four additional pairs

$$(30) \quad c_{\omega\rho} = (f_\rho, iRf_{\bar{\rho}} - iR\bar{C}) = \overline{c_{\omega\bar{\rho}}}, \quad c_{\omega^2\rho} = (f_\rho, iR(-f_{\bar{\rho}} + 2\bar{C}) - iR\bar{C}) = \overline{c_{\omega^2\bar{\rho}}}$$

if and only if $\omega, \sqrt{3} \in K$. Note that the other four possible combinations do not appear in the basis, as they all belong to the $\text{PGL}_2(\mathbb{C})$ -orbit of one of those in (30).

Let us now look above $j_{CV} = \infty$. As anticipated, the two cusps $j = \infty, 1728$ of \mathcal{H}_4 correspond to the $\text{PGL}_2(\mathbb{C})$ -orbits of topological types (f) of non-Morse quartics; indeed, there are two of these: if f branches over less than three points, then either f' has a double root - in which case f is of the form $A(x - a)^3(x - b) + B$, $(A, B) \in K^* \times K$, $a, b \in \overline{\mathbb{Q}}$ (*type 1*) - or f takes the same value at two of the distinct roots of f' , i.e. $f = A(x - a)^2(x - b)^2 + B$ for A, B, a, b as above (*type 2*) (we can assume $a \neq b$, as we otherwise get the $\text{Aff}(K)$ -orbit of a fourth power, which is, above each choice for the critical value B , the class of a unique topological type). When we mod out by the outer action of $\text{Aff}(\overline{\mathbb{Q}})$, *type 1* and *type 2* are clearly distinct topological types. Observe that if a multiset $\{w, w, x\}$ is fixed by an affine transformation, this must necessarily fix w as the element of multiplicity 2, and hence also x , so it must be the identity: therefore, there are no equicritical non-Morse quartics arising from the same outer $\text{Aff}(\overline{\mathbb{Q}})$ -orbit. In other words, the two exceptional points with coordinates (∞, ∞) in our parametrization, which correspond to $t = \omega, \omega^2$, do not lift to an inequivalent equicritical pair.

All we have to do now is find - if it exists - an equicritical pair (f_1, g_1) with f of type 1 and g of type 2. It is not hard to see that

$$(31) \quad (f_1, g_1) = (x^2(x + 3)^2, -3x^3(x + 2)),$$

with critical values $\{0, 0, \frac{81}{16}\}$, works, and this gives the last two ordered pairs $c_1 = (f_1, g_1)$ and $c_\infty = (g_1, f_1)$. Finally, Theorem 1 now follows from Lemma 5.3 together with (27), (28), (29), (30) and (31).

5.3. Application to Weyl sums mod p^2 , and one example. As we anticipated in §1, our main result has an application in the theory of exponential sums attached to polynomials. Specifically, given $a \in \mathbb{Z}$ and a prime power q , in [18] Kowalski and Soundararajan consider the mod q Weyl

sum

$$W_f(a, q) = \frac{1}{\sqrt{q}} \sum_{x \in \mathbb{Z}/q\mathbb{Z}} e\left(\frac{af(x)}{q}\right)$$

attached to $f \in \mathbb{Z}[x]$, investigating pairs (f, g) satisfying $W_f(a, q) = W_g(a, q)$ (or other weaker variants). For $q = p^2$ they observe [18, Remark 1.10] that, when $(a, p) = 1$, one has

$$W_f(a, q) = \sum_{\substack{f'(v)=0 \\ v \in \mathbb{F}_p}} e\left(\frac{af(v)}{p}\right),$$

and hence equicritical polynomials $f, g \in \mathbb{Z}[x]$ whose leading coefficients are invertible mod p satisfy $W_f(a, p^2) = W_g(a, p^2)$ for $(a, p) = 1$. Therefore, following its notation, Theorem 1 implies:

Corollary 5.3.1. *Let $p > 3$ be a prime. The integral polynomials $3(t+2)^4 f_t$ and $3(t+2)^4 g_t$ satisfy $W_f(a, p^2) = W_g(a, p^2)$ for any $(a, p) = 1$ and $t \in \mathbb{Z} \setminus \{2\}$ such that $p \nmid t(t-1)$.*

Finally, let us test Theorem 1 with an example: take, for instance, $t = 42$. Then, exploiting the fact that roots of quartics have exact expressions in terms of radicals, the simple implementation

```

from sage.all import QQbar, polygen, sqrt

def critical_values_exact(f):
    x = f.parent().gen()

    fp = f.derivative()
    critical_points = fp.roots(multiplicities=False)
    critical_values = []

    for cp in critical_points:
        cp_radical = cp.radical_expression()
        cv = f(cp)
        cv_radical = cv.radical_expression()
        critical_values.append(cv_radical)

    return critical_values

def main():
    x = polygen(QQbar)
    t = 42
    v = t**4*(t-1)**3/(t+2)
    f = x**4 - 6*t**3*x**2 - 8*t**3*x
    g = -(t-1)**3*v/(3*(t+2)**3)*x^4 + 2*v*x**2 + 8/3*v*x - 8*t**4*(t**2+t+1)

    print("Polynomial f(x):")
    print(f)
    print("Polynomial g(x):")
    print(g)

```

```

critical_values_f = critical_values_exact(f)
critical_values_g = critical_values_exact(g)

print("\nCritical values of f:")
for i, cv in enumerate(critical_values_f, 1):
    print(f"Critical value {i}:\n {cv}")
    print()

print("\nCritical values of g:")
for i, cv in enumerate(critical_values_g, 1):
    print(f"Critical value {i}:\n {cv}")
    print()

if __name__ == "__main__":
    main()

```

in **SageMath** returns the same critical values:

Polynomial f(x):
 $x^4 - 444528x^2 - 592704x$

Polynomial g(x):
 $-307935007631307/234256x^4 + 107230600008/11x^2 + 142974133344/11x - 44982677376$

Critical values of f:

Critical value 1:

$$-1/2*(482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} * (I*\sqrt{3} + 1) - 135597252859697147904*(-I*\sqrt{3} + 1) / (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} - 32934190464$$

Critical value 2:

$$(482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} + 271194505719394295808 / (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} - 32934190464$$

Critical value 3:

$$-1/2*(482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} * (-I*\sqrt{3} + 1) - 135597252859697147904*(I*\sqrt{3} + 1) / (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} - 32934190464$$

Critical values of g:

Critical value 1:

$$-1/2*(482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} * (I*\sqrt{3} + 1) - 135597252859697147904*(-I*\sqrt{3} + 1) / (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} - 32934190464$$

Critical value 2:

$$\begin{aligned}
 & -1/2*(482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)}* \\
 & (-I*\sqrt{3} + 1) - 135597252859697147904*(I*\sqrt{3} + 1)/ \\
 & (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} \\
 & - 32934190464
 \end{aligned}$$

Critical value 3:

$$\begin{aligned}
 & (482155633265041602566848512*I*\sqrt{74087} + 4464098178074162551805159964672)^{(1/3)} + \\
 & 271194505719394295808/(482155633265041602566848512*I*\sqrt{74087} + \\
 & 4464098178074162551805159964672)^{(1/3)} - 32934190464
 \end{aligned}$$

as expected.

REFERENCES

- [1] V.I. Arnold. “Topological classification of trigonometric polynomials and combinatorics of graphs with an equal number of vertices and edges”. In: *Functional Analysis and its Applications* 30 (1996), pp. 1–17.
- [2] P. Bailey and M. Fried. “Hurwitz monodromy, spin separation and higher levels of a Modular Tower”. In: *Proceedings of Symposia in Pure Mathematics* (2001).
- [3] A.F. Beardon, T.K. Carne, and T.W. Ng. “The critical values of a polynomial”. In: *Constr. Approx* 18 (2002), pp. 343–354.
- [4] A. Cadoret. “Lifting results for rational points on Hurwitz moduli spaces”. In: *Israel Journal of Math.* 164 (2008), pp. 19–59.
- [5] K. Coombes and D. Harbater. “Hurwitz families and arithmetic Galois groups”. In: *Duke Math. J.* 52(4) (1985), pp. 821–839.
- [6] A.F. Costa, M. Izquierdo, and G. Riera. “One-Dimensional Hurwitz Spaces, Modular Curves, and Real Forms of Belyi Meromorphic Functions”. In: *International Journal of Mathematics and Mathematical Sciences* (2008).
- [7] C.J. Cummins and S. Pauli. URL: <https://mathstats.uncg.edu/sites/pauli/congruence/csg0.html>.
- [8] C. Davis. “Extrema of a polynomial”. In: *Am. Math. Monthly* 64 (1957), pp. 679–680.
- [9] P. Dèbes and J-C. Douai. “Algebraic covers: field of moduli versus field of definition”. In: *Annales Scientifiques de l’École Normale Supérieure* 30(3) (1997), pp. 303–338.
- [10] L. DeMarco. “Critical orbits and arithmetic equidistribution”. In: *Proceedings of the International Congress of Mathematicians (ICM 2018)* (2019), pp. 1847–1866.
- [11] F. Diamond and Shurman J. *A first Course in Modular Forms*. Springer, 2005.
- [12] M. Dougherty and J Mccammond. “Critical points, critical values, and a determinant identity for complex polynomials”. In: *Proc. of the AMS* 148(12) (2020).
- [13] M. Fried. “Arithmetic of 3 and 4 branch point covers”. In: *Séminaire de Théorie des Nombres* 43 (1987-88), p. 77.
- [14] M. Fried. “Relating two genus 0 problems of John Thompson”. In: *Progress in Galois Theory. Developments in Mathematics* 12 (2005), pp. 51–85.
- [15] M. Fried. “Variables separated equations: strikingly different roles for the branch cycle lemma and the finite simple group classification”. In: *Science China Mathematics* 55 (2012), pp. 1–72.
- [16] M. Fried and H. Völklein. “The inverse Galois problem and rational points on moduli spaces”. In: *Mathematische Annalen* 290(1) (1991), pp. 771–800.
- [17] D. Harbater. “Galois coverings of the arithmetic line”. In: *Number Theory*. Springer, 2006, pp. 165–195.
- [18] E. Kowalski and K. Soundararajan. “Remembrances of polynomials values: Fourier’s way”. In: (in progress).
- [19] G.K. Kristiansen. “Characterization of polynomials by means of their stationary values”. In: *Arch. Math.* (1984), pp. 44–48.
- [20] D. Mumford and J. Fogarty. *Geometric Invariant Theory, 2nd enlarged ed.* Vol. 34. Ergebnisse der Mathematik und ihrer Grenzgebiete, vol. 34. Springer-Verlag, 1982.
- [21] K. Rubin and A. Silverberg. “Families of elliptic curves with constant mod p representations”. In: *Elliptic curves, modular forms and Fermat’s last theorem: proceedings of a conference held in the Institute of Mathematics of the Chinese University of Hong Kong*. International Press of Boston, 1995, pp. 148–161.
- [22] A. Sutherland and D. Zywina. “Modular curves of prime-power level with infinitely many rational points”. In: *Algebra and Number Theory* 11(5) (2017), pp. 1199–1233.

- [23] R. Thom. “L’équivalence d’une fonction différentiable et d’un polynôme”. In: *Topology* 3 (1965), pp. 297–307.
- [24] K. Wohlfahrt. “An extension of F. Klein’s level concept”. In: *Illinois J. Math* 8(4) (1964), pp. 529–535.
- [25] S. Zdravkovska. “The topological classification of polynomial mappings”. In: *Usp. Mat. Nauk* 25(4) (1970), pp. 179–180.