Markov branching process with infinite variance and non-homogeneous immigration with infinite mean

Kosto V. Mitov,

Faculty of Pharmacy, Medical University, Pleven, Bulgaria email: kmitov@yahoo.com Nikolay M. Yanev, Institute of Mathematics and Informatics, BAS, Sofia, Bulgaria, email: yanev@math.bas.bg

January 8, 2025

Abstract

The paper studies a class of critical Markov branching processes with infinite variance of the offspring distribution. The processes admit also an immigration component at the jump-points of a nonhomogeneous Poisson process, assuming that the mean number of immigrants is infinite and the intensity of the Poisson process converges to zero. The asymptotic behavior of the probability for non-visiting zero is obtained. Limiting distributions are proved, under suitable normalization of the sample paths, depending on the offspring distribution, on the distribution of the immigrants and on the intensity of the Poisson process.

Key words: Markov branching process, infinite variance, limit theorems, non-homogeneous immigration

2020 Mathematics Subject Classification: Primary 60J80; Secondary 60F05, 60J85, 62P10

1 Introduction

The paper deals with Markov branching processes with immigration in timemoments generated by Poisson measure with a local intensity r(t). We consider the critical case when the offspring mean is equal to one, but the offspring variance is infinite. The distribution of immigrants belongs to the class of stable laws with infinite mean and r(t) is a regularly varying function (r.v.f.) converging to zero. Then the considered branching processes are non-homogeneous in time.

Recall that the first branching process with immigration was formulated by Sevastyanov [22]. He investigated a single-type Markov branching process in which immigration occurs according to a time homogeneous Poisson process, and proved limiting distributions. Branching processes with time non-homogeneous immigration were first proposed by Durham [5] and Foster and Williamson [7]. Further results can be found in Badalbaev and Rahimov [2] and Rahimov [19]. See also a review paper of Rahimov [20]. A model with critical non-homogeneous migration was investigated by Yanev and Mitov [24]. Critical Sevastyanov branching processes with non-homogeneous immigration were studied in [12] and critical multitype Markov branching processes with non-homogeneous Poisson immigration were considered by Mitov et al. [14]. Notice that the limiting distributions in these models were obtained in the case of finite first and second offspring characteristics as well as those of the immigration components.

The asymptotic behaviour of branching processes is quite different in the case of finite or infinite offspring variance. Zolotarev [25] was first who obtained limiting distributions for Markov branching processes with infinite offspring variance. Pakes [15], [16] investigated respectively Bienaymé-Galton-Watson process and Markov branching process in the critical case with infinite offspring variance and finite mean of the immigrants, where for the continuous time case it is assumed that the immigration occurs at time-points of a homogeneous Poisson process. Imomov and Tukhtaev [11] considered critical Bienaymé-Galton-Watson process with infinite offspring variance and infinite mean of immigrants and extended also some of the results of Pakes [15]. Sagitov [21] studied multi-type Markov branching processes in the case of homogeneous Poisson immigration with infinite second moments of the offspring distributions and infinite means of the immigrants.

Branching processes with time non-homogeneous immigration find applications for investigating the dynamics of biological systems, particularly cellular populations (see, for example, [23, 9, 10]). In these applications, the stem cells often are considered as an immigration component.

We have to mention that some of the results obtained here are similar to some of the results obtained in the discrete time case by Rahimov [17, 18] for Bienaymé-Galton-Watson branching processes and this is not surprising. Let us note also that the methods of studying in the present work are based on the functional equations for the probability generating functions, stationary measures and some other methods which essentially differ from the methods used in [17, 18].

A detailed description of the considered model is presented in Section 2. Some preliminary results and basic assumptions are given in Section 3. The asymptotic behavior of the P_t (probability for non-visiting zero state at moment t) is investigated in Section 4 (Theorems 4.1-4.3). Thus Theorem 4.1, Theorem 4.2 and Theorem 4.3-(i) establish the conditions under which P_t converges to zero as $t \to \infty$ with various rates depending of the parameters of reproduction (γ), of immigration (α) and of Poisson measure (θ). Under the conditions of Theorem 4.3-(ii) P_t converges to a positive probability less than 1 which is exactly calculated. Finally, at the conditions of Theorem 4.3-(iii) we obtain that $P_t \to 1$ as $t \to \infty$.

Under the same basic conditions various types of limiting distributions are obtained in five theorems presented in Section 5. In fact we proved eight different type limiting distributions. Note that seven of them are under the condition for non-visiting zero state. Thus in Theorems 5.1-(i), 5.2-(ii) and 5.4-(ii) under suitable additional conditions we proved a stationary discrete time limiting distribution and we obtained an integral form for the corresponding probability generating function (p.g.f.) depending only from the local characteristics of the process (offspring p.g.f. f(s) and the p.g.f. of the immigrants g(s), see Remark 5.1). For the other limiting distributions under a normalization with a suitable r.v.f. we obtained the corresponding Laplace transforms. The most interesting results are given in Theorem 5.1-(iii) where we have two singular to each other conditional limiting distributions, where the first one is a non-proper stationary distribution with an atom at infinity and the second one under a suitable normalization has an atom at zero. Finally in Theorem 5.5-(iii) we obtained non conditional limiting distribution under a suitable normalizing r.v.f. and the limiting random variable is just stable with parameter α (from the p.g.f. of the immigrants). Comments with discussion of the results are given after the proofs of all theorems in Sections 4 and 5.

2 Description of the models

A Markov branching process can be described as follows. The particles of a given type evolve independently of each other, lives random time τ with distribution function (d.f.) $G(t) = \mathbf{P} \{\tau \leq t\} = 1 - e^{-\mu t}, t \geq 0, \mu > 0$, and at the end of its life the particle produces random number $\xi \geq 0$ of new particles of the same type. The number of particles Z(t) at moment $t \geq 0$ is known as Markov branching process (see [1], [8]). Denote by $h(s) = \mathbf{E} [s^{\xi}]$ the offspring p.g.f. and $F(t;s) = \mathbf{E}\left[s^{Z(t)}|Z(0) = 1\right], t \ge 0, s \in [0,1]$. It is well known that (see e.g. [1], [8])

$$\frac{\partial F(t;s)}{\partial t} = \mu[h(F(t;s)) - F(t;s)], \qquad (1)$$

Let (S_k, I_k) , k = 0, 1, 2, ..., be independent of Z(t), where $0 = S_0 < S_1 < S_2 < \cdots$ are jump points of an non-homogeneous Poisson process $\nu(t)$ and the random variables $\{I_k\}$ are independent, identically distributed (i.i.d.) with non-negative integer values. Denote by r(t) the intensity of $\nu(t)$ with a mean measure $R(t) = \int_0^t r(u) du$. Let $g(s) = \mathbf{E} [s^{I_k}]$ be the p.g.f. of the immigrants.

Assume that at every jump-point S_k , a random number I_k of new particles immigrate into the process Z(t) and they participate in the evolution as the other particles. Let us denote the new process by Y(t). It can be strictly defined as follows

$$Y(t) = \sum_{k=1}^{\nu(t)} \sum_{j=1}^{I_k} Z^{(k,j)} \left(t - S_k \right), \ t \ge 0,$$

where $\{Z^{(k,j)}(t)\}\$ are i.i.d. copies of Z(t). The process Y(t), $t \ge 0$, is called Markov branching process with non-homogeneous Poisson immigration (MBPNPI).

For $\Phi(t;s) := \mathbf{E} \left[s^{Y(t)} | Y(t) = 0 \right]$ we have the following equation

$$\Phi(t;s) = \exp\left\{-\int_0^t r(t-u)(1-g(F(u;s)))du\right\}, \ \Phi(t;0) = 1.$$
(2)

The proof is given in [23] and in the more general multitype case in [14].

For the intensity of the Poisson process, we assume additionally the following condition

$$r(t) = t^{-\theta} L_R(t), \quad \text{where} \quad \theta > 0, \tag{3}$$

and $L_R(.)$ is a slowly varying function (s.v.f.) at infinity.

3 Basic assumptions and preliminary results

For the branching mechanism we assume that the offspring p.g.f. f(s) has the following representation

$$f(s) = s + (1-s)^{\gamma+1} L\left(\frac{1}{1-s}\right), \quad s \in [0,1),$$
(4)

where $\gamma \in (0, 1]$ and L(.) is a s.v.f. at infinity. Thus, the process $Z(t), t \ge 0$, is critical. If $\gamma < 1$ the offspring variance in infinite.

Comment 3.1. If $\gamma = 1$ and $L(t) \rightarrow b$ then the offspring variance is finite. The results for this case follow directly from the corresponding results for the multitype Markov processes with non-homogeneous Poisson immigration studied in [14]. If $\gamma = 1$ the offspring variance can also be infinite, depending on the properties of the slowly varying function L(.).

It is known (see [8], Theorem 12.1) that a critical Markov branching process has an invariant measure whose p.g.f. U(s) is given by $U(s) = \int_0^s \frac{du}{f(u) - u}$, $0 \le s \le 1$. The Kolmogorov backward equation (1) can be written as follows $\int_s^{F(t;s)} \frac{du}{f(u) - u} = \mu t$. This leads to $U(F(t;s)) = U(s) + \mu t$. Denote by

$$V(x) = U\left(1 - \frac{1}{x}\right) = \int_0^{1 - 1/x} \frac{du}{f(u) - u} = \int_1^x \frac{u^{\gamma - 1}}{L(u)} du, x \ge 1.$$

So V(x) is regularly varying with exponent γ . Then its inverse W(y) is regularly varying with exponent $1/\gamma$. Let us note that V(x) and W(y) are increasing (see e.g. [15],[16]). Using the above relations we get for $s \in [0, 1)$

$$1/[1 - F(t;s)] = W\left(\mu t + V(1/(1-s))\right), \quad s \in [0,1).$$
(5)

Substituting s = 0 we have $1 - F(t; 0) = 1/W(\mu t)$. For the p.g.f. of the immigrants we will assume that

$$g(s) = 1 - (1 - s)^{\alpha} l\left(\frac{1}{1 - s}\right), \quad s \in [0, 1]$$
(6)

where $\alpha \in (0, 1]$ and l(x) is a s.v.f. at infinity.

Comment 3.2. If $\alpha \in (0,1)$ the mean number of immigrants is infinite. In the case when $\alpha = 1$ the mean number of immigrants can be infinite or finite depending on the s.v.f. l(.). If $\alpha = 1$, and $l(x) \rightarrow m \in (0,\infty)$, then $\mathbf{E}[I_k] = m$ is finite.

Let us denote $\Psi(x) = 1/[1 - g(1 - \frac{1}{x})] = \frac{x^{\alpha}}{l(x)}, x \ge 1$. The function $\Psi(.)$ is non decreasing in $[1, \infty)$ and let $\overleftarrow{\Psi}(x), x \ge 1$, be its inverse function which

is also non-decreasing in $[1, \infty)$. Then g(s) can be written in the following form

$$g(s) = 1 - 1/\Psi \left(1/(1-s) \right), \ s \in [0,1].$$

Further for convenience we will denote (see also (5))

$$q(t;s) := 1 - g(F(t;s)) = (1 - F(t;s))^{\alpha} l \left(1/[1 - F(t;s)] \right)$$

= $1/\Psi \left(W \left(\mu t + V \left(1/(1-s) \right) \right) \right).$ (7)

From (7) with s = 0 it follows that as $t \to \infty$,

$$q(t) := q(t;0) = \frac{1}{\Psi(W(\mu t))} \sim t^{-\alpha/\gamma} L_Q(t),$$
(8)

where $L_Q(.)$ is s.v.f. at infinity.

We will use the following notations

$$Q(t) := \int_0^t q(u) du, \ Q := \int_0^\infty q(u) du, \ \Delta(s) := \int_0^\infty q(u;s) du, s \in [0,1],$$

when the last two integrals converge.

Proposition 3.1. The following representations hold

$$\Delta(s) = \int_{s}^{1} \frac{1 - g(u)}{\mu(f(u) - u)} du, \ Q = \int_{0}^{1} \frac{1 - g(u)}{\mu(f(u) - u)} du.$$
(9)

Proof. Since q(t;s) is non-increasing in $s \in [0,1]$ then the convergence of $Q = \int_0^{\infty} q(t)dt$ leads to the uniform convergence of $\Delta(s) = \int_0^{\infty} q(t;s)dt$ on [0,1]. Then $\frac{d}{ds}\Delta(s) = -\int_0^{\infty} \frac{d}{dF}g(F(t;s))\frac{\partial}{\partial s}F(t;s)dt$. Note that by the forward Kolmogorov equation $\frac{\partial}{\partial t}F(t;s) = f^*(s)\frac{\partial}{\partial s}F(t;s)$, F(0;s) = s, where $f^*(s) = \mu(f(s) - s)$ is the infinitesimal generating function. Therefore $\frac{d}{ds}\Delta(s) = -\frac{1}{f^*(s)}\int_0^{\infty} \frac{d}{dF}g(F(t;s))\frac{\partial}{\partial t}F(t;s)dt = -\frac{1}{f^*(s)}\int_0^{\infty} d_tg(F(t;s))$

$$= -\frac{1}{f^*(s)} [g(F(\infty;s)) - g(F(0;s))] = -\frac{1}{f^*(s)} [1 - g(s)].$$

Hence by integrating from s to 1 and using that $\Delta(1) = 0$ one obtains the first equation in (9). On the other hand, by integrating form 0 to s one obtains $\Delta(s) - \Delta(0) = -\int_0^s \frac{1-g(u)}{f^*(u)} du$. Since $\Delta(0) = Q$ then the second equation in (9) follows.

Lemma 3.1. For $s \in [0,1)$ it is fulfilled that $q(t;s)/q(t) \to 1$ as $t \to \infty$.

Proof. From (7) we have

$$q(t;s) = \frac{1}{\Psi\left(W\left(\mu t + V\left(\frac{1}{1-s}\right)\right)\right)} = \frac{1}{\Psi\left(W\left(\mu t\left(1 + \frac{1}{\mu t}V\left(\frac{1}{1-s}\right)\right)\right)\right)}.$$

Since, s is fixed then for every t large enough $1 + \frac{1}{\mu t}V\left(\frac{1}{1-s}\right) \in [1,2]$. Therefore, by the uniform convergence of regularly varying functions on compact sets (see [4], Theorem 1.5.2), it follows that $\Psi\left(W\left(\mu t\left(1+V\left(1/(1-s)\right)/\mu t\right)\right)\right) \sim \Psi(W(\mu t)), t \to \infty$, which together with (7) and (8) completes the proof. \Box

Lemma 3.2. Let $s(t) = \exp(-\lambda/W(\mu t)), \lambda > 0$. Then for c > 0,

$$q(ct; s(t))/q(t) \to (c + \lambda^{-\gamma})^{-\alpha/\gamma}, \ t \to \infty.$$

Proof. Note first that $1-s(t) \sim \lambda/W(\mu t)$, $t \to \infty$. Since V and W are inverse to each other then $V\left([1-s(t)]^{-1}\right) \sim V(\lambda^{-1}W(\mu t)) \sim \lambda^{-\gamma}\mu t$, $t \to \infty$. Now from (7) we have as $t \to \infty$

$$q(ct; s(t)) = 1/\Psi \left(W \left(c\mu t + V \left([1 - s(t)]^{-1} \right) \right) \right)$$

$$\sim 1/\Psi \left(W \left(c\mu t + \lambda^{-\gamma} \mu t \right) \right) \sim 1/\Psi \left(W \left(\mu t \right) (c + \lambda^{-\gamma})^{1/\gamma} \right)$$

$$\sim \frac{1}{\Psi(W(\mu t))} \left(c + \lambda^{-\gamma} \right)^{-\alpha/\gamma} = q(t)(c + \lambda^{-\gamma})^{-\alpha/\gamma},$$

because of (8) and the fact that W(t) and $\Psi(t)$ are r.v.f. with exponents $1/\gamma$ and α respectively.

4 Asymptotic behavior of the probability for nonvisiting the state zero

Let us denote

$$P_t := \mathbf{P}\{Y(t) > 0\} = 1 - \Phi(t; 0) = 1 - \exp(-I(t)),$$
(10)

where
$$I(t) = \int_0^t r(t-u)q(u)du$$
 (see (7) with $s = 0$ and (8))

Theorem 4.1. Let conditions (3), (4) , and (6) hold. Assume also that in (3) $\theta \ge 1$ and $\alpha/\gamma \ge 1$. Then

$$\mathbf{P}\{Y(t) > 0\} \sim R(t)q(t) + Q(t)r(t), \quad t \to \infty.$$
(11)

Proof. Note that R(t) and Q(t) are s.v.f. at infinity, tq(t) = o(Q(t)) and tr(t) = o(R(t)) as $t \to \infty$ (see [4], Proposition 1.5.8 Eq (1.5.8)). Let $\delta \in (0, 1/2)$ be fixed. Then we have

$$I(t) = \int_0^t r(t-u)q(u)du = \int_0^{t\delta} + \int_{t\delta}^{t(1-\delta)} + \int_{t(1-\delta)}^t = I_1(t) + I_2(t) + I_3(t).$$

Note first that

$$I_1(t) \le r(t(1-\delta)) \int_0^{t\delta} q(u)du \sim r(t)(1-\delta)^{-1}Q(t),$$

$$I_1(t) \ge r(t) \int_0^{t\delta} q(u)du \sim r(t).Q(t), \ t \to \infty.$$

Then

$$1 \le \liminf_{t \to \infty} \frac{I_1(t)}{r(t).Q(t)} \le \limsup_{t \to \infty} \frac{I_1(t)}{r(t).Q(t)} \le (1-\delta)^{-\theta}.$$
 (12)

Since

$$I_{3}(t) \leq q(t(1-\delta)) \int_{0}^{t\delta} r(u)du \sim q(t)(1-\delta)^{-\alpha/\gamma} R(t),$$

$$I_{3}(t) \geq q(t) \int_{0}^{t\delta} r(u)du \sim q(t).R(t), \ t \to \infty,$$

then

$$1 \le \liminf_{t \to \infty} \frac{I_3(t)}{q(t).R(t)} \le \limsup_{t \to \infty} \frac{I_3(t)}{q(t).R(t)} \le (1-\delta)^{-\alpha/\gamma}.$$
 (13)

Finally, for $I_2(t)$ we have

$$0 \leq I_2(t) = \int_{t\delta}^{t(1-\delta)} r(t-u)q(u)du$$

$$\leq r(t\delta)q(t\delta)t(1-2\delta) \sim t.r(t).q(t)\delta^{-2}(1-2\delta), t \to \infty.$$

Since tr(t) = o(R(t)) and tq(t) = o(Q(t)), as $t \to \infty$ then

$$I_2(t) = o(q(t).R(t)), \quad I_2(t) = o(r(t).Q(t)), \ t \to \infty.$$
 (14)

Having in mind that $\delta \in (0, 1/2)$ was arbitrary, we conclude from (12), (13), and (14)

$$I(t) = I_1(t) + I_2(t) + I_3(t) \sim r(t) \cdot Q(t) + q(t) \cdot R(t) \to 0, \ t \to \infty,$$

which together with (10) and $1 - e^{-x} = x(1 + o(1)), x \to 0$ proves (11). \Box

Comment 4.1. If $\theta > \frac{\alpha}{\gamma} \ge 1$ then $R < \infty$ and $P_t \sim RL_Q(t)t^{-\alpha/\gamma}$. If $\frac{\alpha}{\gamma} > \theta \ge 1$ then $Q < \infty$ and $P_t \sim QL_R(t)t^{-\theta}$. If $\theta = \frac{\alpha}{\gamma} > 1$ then $P_t \sim [RL_Q(t) + QL_R(t)]t^{-\theta}$. If $\theta = \frac{\alpha}{\gamma} = 1$ then R(t) and Q(t) are s.v.f. and $P_t \sim [R(t)L_Q(t) + Q(t)L_R(t)]t^{-1}$.

Theorem 4.2. Assume the conditions(3), (4), and (6) hold.

- (i) If $\theta \ge 1$ and $0 < \alpha/\gamma < 1$ then $\mathbf{P}\{Y(t) > 0\} \sim R(t).q(t), t \to \infty$.
- (ii) If $0 < \theta < 1$ and $\alpha/\gamma \ge 1$ then $\mathbf{P}\{Y(t) > 0\} \sim r(t).Q(t), t \to \infty$.

Proof. (i) Under the conditions of this case $R(t) = \int_0^t r(u) du \uparrow R \leq \infty$, R(t) is a s.v.f. at infinity and $tr(t) = o(R(t)), t \to \infty$. Let $\delta \in (0, 1)$ be fixed.

Consider
$$I(t) = \int_0^t r(t-u)q(u)du = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t) + I_2(t)$$
. Then
 $I_2(t) = \int_{t\delta}^t r(t-u)q(u)du \le q(t\delta) \int_0^{t(1-\delta)} r(u)du \sim q(t)\delta^{-\alpha/\gamma}R(t)$
 $I_2(t) \ge q(t)R(t(1-\delta)) \sim q(t)R(t), t \to \infty.$
 $0 \le I_1(t) = \int_0^{t\delta} r(t-u)q(u)du \le r(t(1-\delta)) \int_0^{t\delta} q(u)du$
 $\sim r(t(1-\delta))\frac{t\delta q(t\delta)}{1-\alpha/\gamma} \sim \frac{tr(t)q(t)}{1-\alpha/\gamma}(1-\delta)^{-\theta}\delta^{1-\alpha/\gamma}.$

As we mentioned above $tr(t) = o(R(t)), t \to \infty$. Therefore $I_1(t) = o(I_2(t)))$ and

$$1 \le \liminf_{t \to \infty} I(t)/(q(t)R(t)) \le \limsup_{t \to \infty} I(t)/(q(t)R(t)) \le \delta^{-\alpha/\gamma}.$$

Since $\delta \in (0, 1)$ was arbitrary then we get that $I(t) \sim q(t)R(t) \to 0, t \to \infty$. By (10) and $1 - e^{-x} \sim x, x \to 0$, we complete the proof of case (i). The proof of case (ii) is similar, one has to change only the role of r(t) and q(t).

Comment 4.2. (i) If $\theta \ge 1$ then R(t) is a s.v.f. and $P_t \sim R(t)L_Q(t)t^{-\alpha/\gamma}$. (ii) If $\alpha/\gamma \ge 1$ then Q(t) is a s.v.f. and $P_t \sim Q(t)L_R(t)t^{-\theta}$.

Theorem 4.3. Assume conditions(3), (4), and (6) hold. Let additionally $0 < \theta < 1$ and $0 < \alpha/\gamma < 1$.

(i) If
$$\theta + \alpha/\gamma > 1$$
, or $\theta + \alpha/\gamma = 1$ but $L_R(t)L_Q(t) \to 0$ then

$$\mathbf{P}\{Y(t) > 0\} \sim t.r(t).q(t).\mathbb{B}(1 - \alpha/\gamma, 1 - \theta),$$

where $\mathbb{B}(.,.)$ is Euler's beta function.

(ii) If $\theta + \alpha/\gamma = 1$ but $L_R(t)L_Q(t) \to K \in (0,\infty)$ then $\mathbf{P}\{Y(t) > 0\} \to 1 - e^{-K\pi/\sin\pi\theta}, t \to \infty.$

(iii) If
$$\theta + \alpha/\gamma < 1$$
, or $\theta + \alpha/\gamma = 1$ but $L_R(t)L_Q(t) \to \infty$ then

 $\mathbf{P}\{Y(t) > 0\} \to 1, \ t \to \infty.$

Proof. Let $\delta \in (0, 1/2)$ be fixed. Consider

$$I(t) = \int_0^{t\delta} + \int_{t\delta}^{t(1-\delta)} + \int_{t(1-\delta)}^t = I_1(t) + I_2(t) + I_3(t).$$

Changing variables u = vt in $I_2(t)$ we obtain $I_2(t) = t \int_{\delta}^{(1-\delta)} r(t(1-v))q(tv)dv$. By the uniform convergence of r.v.f. on compact sets we have for $v \in [\delta, 1-\delta]$,

$$r(t(1-v)) \sim r(t)(1-v)^{-\theta}, \quad q(tv) \sim q(t)v^{-\alpha/\gamma}, t \to \infty.$$

Therefore, as $t \to \infty$,

$$I_2(t) = tr(t)q(t) \int_{\delta}^{(1-\delta)} \frac{r(t(1-v))q(tv)dv}{r(t)q(t)} \sim tr(t)q(t) \int_{\delta}^{(1-\delta)} (1-v)^{-\theta} v^{-\alpha/\gamma} dv.$$

Further one gets

$$0 \leq I_{1}(t) = \int_{0}^{t\delta} r(t-u)q(u)du \leq r(t(1-\delta)) \int_{0}^{t\delta} q(u)du$$

$$\sim r(t)(1-\delta)^{-\theta} \frac{t\delta q(t\delta)}{1-\alpha/\gamma} \sim \frac{tr(t)q(t)}{1-\alpha/\gamma} (1-\delta)^{-\theta} \delta^{1-\alpha/\gamma}, \quad (15)$$

$$0 \leq I_{3}(t) = \int_{t(1-\delta)}^{t} r(t-u)q(u)du \leq q(t(1-\delta)) \int_{0}^{t\delta} r(u)du$$

$$\sim q(t)(1-\delta)^{-\alpha/\gamma} \frac{t\delta r(t\delta)}{1-\theta} \sim \frac{tr(t)q(t)}{1-\theta} (1-\delta)^{-\alpha/\gamma} \delta^{1-\theta}, t \to \infty. \quad (16)$$

Using the estimates for $I_1(t)$, $I_2(t)$, and $I_3(t)$ we obtain that

$$\begin{split} & \int_{\delta}^{(1-\delta)} (1-v)^{-\theta} v^{-\alpha/\gamma} dv \\ & \leq \liminf_{t \to \infty} \frac{I_1(t) + I_2(t) + I_3(t)}{tr(t)q(t)} \leq \limsup_{t \to \infty} \frac{I_1(t) + I_2(t) + I_3(t)}{tr(t)q(t)} \\ & \leq \frac{(1-\delta)^{-\theta} \delta^{1-\alpha/\gamma}}{1-\alpha/\gamma} + \int_{\delta}^{(1-\delta)} (1-v)^{-\theta} v^{-\alpha/\gamma} dv + \frac{(1-\delta)^{-\theta} \delta^{1-\alpha/\gamma}}{1-\theta}. \end{split}$$

These inequalities and the fact that $\delta \in (0, 1/2)$ was arbitrary yield

$$\begin{split} I(t) &\sim tr(t)q(t)\mathbb{B}(1-\alpha/\gamma,1-\theta) \\ &\sim t^{1-\theta-\alpha/\gamma}L_R(t)L_Q(t)\mathbb{B}(1-\alpha/\gamma,1-\theta), \quad t\to\infty. \end{split}$$

(i) In this case $I(t) \to 0$ which together (10) and $1 - e^{-x} = x(1 + o(1))$, $x \to 0$, completes the proof.

(ii) Now $I(t) \to K\mathbb{B}(\theta, 1-\theta) = K\pi/\sin\pi\theta$ which proves this case.

(iii) Since $I(t) \to \infty$ then by (10) the proof of this case is completed. \Box

Comment 4.3. In case (i) if $\theta + \alpha/\gamma > 1$ then

$$P_t \sim L_R(t)L_Q(t)\mathbb{B}(1-\alpha/\gamma, 1-\theta)t^{-(\theta+\alpha/\gamma-1)} \to 0, t \to \infty.$$

Otherwise if $\theta + \alpha/\gamma = 1$ then $P_t \sim L_R(t)L_Q(t)\mathbb{B}(1-\alpha/\gamma, 1-\theta) \to 0, t \to \infty$, i.e. the probability of non-visiting zero develops like a s.v.f.

5 Limit theorems

Note that the conditional p.g.f. of Y(t)|Y(t) > 0 has the form

$$\mathbf{E}\left[s^{Y(t)}|Y(t)>0\right] = 1 - (1 - \Phi(t;s))/(1 - \Phi(t;0)).$$
(17)

and from (2) we have

$$1 - \Phi(t;s) = 1 - \exp(-I(t;s)), \tag{18}$$

$$I(t;s) := \int_0^t r(t-u)q(u;s)du,$$
(19)

where q(t; s) is defined in (7).

Theorem 5.1. Assume the conditions (3), (4), and (6) hold. Assume also that in (3) $\theta \ge 1$ such that $R = \int_0^\infty r(t)dt < \infty$ and $\alpha/\gamma \ge 1$ such that $Q = \int_0^\infty q(t)dt < \infty$.

(i) If
$$q(t) = o(r(t))$$
 then $\mathbf{E}[s^{Y(t)}|Y(t) > 0] \to 1 - \Delta(s)/Q, t \to \infty.$

(ii) If r(t) = o(q(t)) then $\mathbf{P} \{Y(t)/W(\mu t) \le x | Y(t) > 0\} \to D(\alpha, \gamma; x), x \ge 0$, where

$$\int_0^\infty e^{-\lambda x} \mathrm{d}D(\alpha, \gamma; x) = \hat{D}(\alpha, \gamma; \lambda) = 1 - \frac{\lambda^\alpha}{(1 + \lambda^\gamma)^{\alpha/\gamma}}, \quad \lambda > 0.$$

(iii) If
$$r(t)/q(t) \to d \in (0,\infty), t \to \infty$$
, then as $t \to \infty$,

$$\mathbf{E}[s^{Y(t)}|Y(t) > 0] \to \frac{dQ}{dQ + R} \left(1 - \Delta(s)/Q\right),\tag{20}$$

$$\mathbf{P}\left\{Y(t)/W(\mu t) \le x | Y(t) > 0\right\} \to \frac{dQ + RD(\alpha, \gamma; x)}{dQ + R}, \ x \ge 0.$$
(21)

Remark 5.1. From Proposition 3.1 one obtains for the limiting p.g.f.

$$\varphi(s) = 1 - \frac{\Delta(s)}{Q} = \left(\int_0^s \frac{1 - g(u)}{f(u) - u} du\right) / \left(\int_0^1 \frac{1 - g(u)}{f(u) - u} du\right).$$

Proof. Notice first that under the conditions of the theorem Eq.(11) gets the form

$$\mathbf{P}\{Y(t) > 0\} \sim R.q(t) + Q.r(t), \quad t \to \infty.$$
(22)

By Lemma 3.1 we conclude that $Q = \int_0^\infty q(t)dt < \infty$ yields $\Delta(s) = \int_0^\infty q(t;s)dt < \infty$ and by the dominated convergence theorem $\Delta(s) \to Q$ as $s \downarrow 0$. In this way for every fixed $s \in [0,1), \frac{q(t,s)}{\Delta(s)}$ is a density on $[0,\infty)$. Under the conditions of the theorem one has also that r(t)/R is a density on $[0,\infty)$. Therefore by (Theorem 1, [3]) for any fixed $s \in [0,1)$,

$$I(t;s) \sim \Delta(s).r(t) + q(t;s).R \to 0, \quad t \to \infty.$$
 (23)

By the relation $1 - e^{-x} \sim x$, $x \to 0$ we get that for any fixed $s \in [0, 1)$

$$1 - \Phi(t;s) \sim I(t;s), \quad t \to \infty.$$
(24)

(i) Using equations (22), (23), and (24) we get

$$\frac{1-\Phi(t;s)}{1-\Phi(t;0)} \sim \frac{\Delta(s)r(t)+q(t;s).R}{Q.r(t)+q(t).R} = \frac{\Delta(s)+q(t;s)\frac{R}{r(t)}}{Q+q(t)\frac{R}{r(t)}}, t \to \infty.$$

In this case we have $q(t)R/r(t) \to 0$, $q(t;s)R/r(t) \sim q(t)R/r(t) \to 0$. Therefore,

$$[1 - \Phi(t;s)]/[1 - \Phi(t;0)] \to \Delta(s)/Q, \ t \to \infty,$$

which together with (17) completes the proof of this case.

(ii) In this case $r(t)/q(t;s) \to 0$, $t \to \infty$, for every fixed $s \in [0,1)$. So,

$$\frac{1-\Phi(t;s)}{1-\Phi(t;0)} \sim \frac{q(t;s)}{q(t)} \frac{\Delta(s)\frac{r(t)}{q(t;s)} + R}{Q\frac{r(t)}{q(t)} + R} \to 1, \ t \to \infty.$$

In other words, almost all non-degenerate sample paths go to infinity. So we need an appropriate normalization in order to get a proper limit distribution. Let now $s(t) = \exp(-\lambda/W(\mu t))$. For $\delta \in (0, 1)$ fixed one has

$$I(t; s(t)) = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t; s(t)) + I_2(t; s(t)).$$

Having in mind that q(t; s) is non-increasing in $t \ge 0$ we have

$$\begin{aligned} q(t;s(t)) \int_0^{t\delta} r(u) du &\leq I_1(t;s(t)) = \int_0^{t\delta} r(u) q(t-u;s(t)) du \\ &\leq q(t(1-\delta);s(t)) \int_0^{t\delta} r(u) du. \end{aligned}$$

Using that $\lim_{t\to\infty}\int_0^{t\delta}r(u)du=R<\infty$ and applying Lemma 3.2 with c=1 and $c=1-\delta$ one obtains

$$\begin{split} q(t;s(t)) \int_0^{t\delta} r(u) du &\sim q(t) (1+\lambda^{-\gamma})^{-\alpha/\gamma} R, \\ q(t(1-\delta);s(t)) \int_0^{t\delta} r(u) du &\sim q(t) \left(1-\delta+\lambda^{-\gamma}\right)^{-\alpha/\gamma} R, \ t \to \infty, \end{split}$$

Hence

$$R(1+\lambda^{-\gamma})^{-\frac{\alpha}{\gamma}} \leq \liminf_{t \to \infty} I_1(t;s(t))/q(t) \leq \limsup_{t \to \infty} I_1(t;s(t))/q(t) \leq R(1-\delta+\lambda^{-\gamma})^{-\frac{\alpha}{\gamma}}$$

On the other hand for $I_2(t; s(t))$ we obtain

$$I_2(t;s(t)) = \int_{t\delta}^t r(u)q(t-u;s(t))du \le r(t\delta) \int_0^{t(1-\delta)} q(u;s(t))du$$
$$\le r(t\delta) \int_0^{t(1-\delta)} q(u)du \le r(t\delta)Q = o(q(t)), t \to \infty.$$

From the relations for $I_1(t; s(t))$ and $I_2(t; s(t))$ we get

$$\frac{R\lambda^{\alpha}}{(1+\lambda^{\gamma})^{\alpha/\gamma}} \leq \liminf_{t \to \infty} \frac{I(t;s(t))}{q(t)} \leq \limsup_{t \to \infty} \frac{I(t;s(t))}{q(t)} \leq \frac{R\lambda^{\alpha}}{(1-\delta+\lambda^{\gamma})^{\alpha/\gamma}}$$

Hence $I(t;s(t)) \sim q(t) \frac{R\lambda^{\alpha}}{(1+\lambda^{\gamma})^{\alpha/\gamma}} \to 0$, which gives that

$$1 - \Phi(t; s(t)) \sim q(t) R \lambda^{\alpha} (1 + \lambda^{\gamma})^{-\alpha/\gamma}, t \to \infty,$$

using the asymptotic $1 - e^{-x} \sim x$, $x \to 0$. This relation, (17), and (22) with r(t) = o(q(t)) yield

$$\lim_{t \to \infty} \mathbf{E} \left[e^{-\lambda Y(t)/W(\mu t)} | Y(t) > 0 \right] = \hat{D}_{\gamma}(\lambda) = 1 - \lambda^{\alpha} (1 + \lambda^{\gamma})^{-\alpha/\gamma}, \quad \lambda > 0,$$

which completes the proof of case (ii).

(iii) We have from (11) that

$$1 - \Phi(t;0) \sim (R + dQ)q(t), t \to \infty, \tag{25}$$

From equations (22), (23), and (24) it follows that

$$1 - \Phi(t;s) \sim (R + d\Delta(s)))q(t;s), t \to \infty.$$

This relation and (25) yield

$$[1 - \Phi(t;s)]/[1 - \Phi(t;0)] \rightarrow [R + d\Delta(s)]/[R + dQ],$$

which is equivalent to (20).

The obtained discrete limiting distribution is not proper. It has mass at infinity $\frac{dQ}{R+dQ}$. In other words there are sample paths that grow very fast and they have to be normalized by some factor in order to obtain a proper limiting distribution. Let $s(t) = \exp(-\lambda/W(\mu t))$. For $\delta \in (0, 1)$ one has

$$I(t;s(t)) = \int_0^{t/\delta} + \int_{t\delta}^t = I_1(t;s(t)) + I_2(t;s(t))$$

and hence

$$q(t;s(t))\int_0^{t\delta} r(u)du \le I_1(t;s(t)) \le q\left(t(1-\delta);s(t)\right)\int_0^{t\delta} r(u)du$$

Applying Lemma 3.2 with c = 1 and $c = 1 - \delta$ we obtain

$$\left(1+\frac{1}{\lambda^{\gamma}}\right)^{-\alpha/\gamma} \leq \liminf_{t \to \infty} \frac{I_1(t;s(t))}{Rq(t)} \leq \limsup_{t \to \infty} \frac{I_1(t;s(t))}{Rq(t)} \leq \left(1-\delta+\frac{1}{\lambda^{\gamma}}\right)^{-\alpha/\gamma}$$

Since q(t) is non-increasing in $s \in [0, 1)$ then one has that

$$I_2(t; s(t) \le r(t\delta) \int_0^{t(1-\delta)} q(u; s(t)) du \le r(t\delta)q(t)t(1-\delta) = o(q(t)),$$

having in mind that $tr(t) = o(1), t \to \infty$. Therefore

$$\left(1+\frac{1}{\lambda^{\gamma}}\right)^{-\alpha/\gamma} \leq \liminf_{t \to \infty} \frac{I(t;s(t))}{Rq(t)} \leq \limsup_{t \to \infty} \frac{I(t;s(t))}{Rq(t)} \leq \left(1-\delta+\frac{1}{\lambda^{\gamma}}\right)^{-\alpha/\gamma}$$

Since $\delta \in (0,1)$ was arbitrary we get $I(t;s(t)) \sim R.q(t) \cdot (1+1/\lambda^{\gamma})^{-\alpha/\gamma} \to 0, t \to \infty$. Therefore,

$$1 - \Phi((t; s(t)) \sim R.q(t). (1 + 1/\lambda^{\gamma})^{-\alpha/\gamma} \to 0, \ t \to \infty,$$

which leads to

$$\lim_{t \to \infty} \frac{1 - \Phi(t; s(t))}{1 - \Phi(t; 0)} = \frac{R}{R + dQ} \left(1 + \frac{1}{\lambda^{\gamma}} \right)^{-\alpha/\gamma}.$$

This relation and (25) prove (21).

Comment 5.1. (i) Since $q(t)/r(t) = t^{-(\alpha/\gamma - \theta)}L_Q(t)/L_R(t) \to 0$ then $\alpha/\gamma > \theta$ or $\alpha/\gamma = \theta$ but $L_Q(t)/L_R(t) \to 0$. Let us consider a particular case when in (4) $L(s) \equiv 1/(1+\gamma)$ and in (6) $l(s) \equiv 1$. Then for $\alpha > \gamma$ we obtain

$$\int_{0}^{s} \frac{1 - g(x)}{f(x) - x} dx = \frac{1 + \gamma}{\alpha - \gamma} (1 - (1 - s)^{\alpha - \gamma}).$$

Therefore $Q = \int_0^1 \frac{1-g(x)}{f(x)-x} dx = \frac{1+\gamma}{\alpha-\gamma}$ and

$$\varphi(s) = \frac{1}{Q} \int_0^s \frac{1 - g(x)}{f(x) - x} dx = 1 - (1 - s)^{\alpha - \gamma}.$$

Hence the limiting r.v. belongs to the normal domain of attraction of a stable law with parameter $\alpha - \gamma$. Note that for $\alpha = \gamma$ we have $\int_0^s \frac{1-g(x)}{f(x)-x} dx = (1+\gamma) \log \frac{1}{1-s}$ and therefore $Q = \infty$.

(ii) By the Tauberian theorem (Feller [6], Ch. XIII, (5.20)) one has $1 - D(\alpha, \gamma; x) \sim x^{-\alpha} / \Gamma(1-\alpha), x \to \infty$. Note that $\alpha/\gamma < \theta$ or $\alpha/\gamma = \theta$ but $L_R(t)/L_Q(t) \to 0$.

(iii) We obtained (with different normalization) two singular to each other conditional limiting distributions. The first one is a discrete nonproper distribution similar to the case (i) but now with an atom at infinity

with probability R/(dQ + R). The second one is similar to the case (ii) but it has now an atom at zero with probability dQ/(dQ + R). In other words, as $t \to \infty$ it follows for the sample paths that $Y(t) \sim \zeta_1$ with probability dQ/(dQ+R) where ζ_1 is a limiting r.v. in the first case and $Y(t) \sim \zeta_2 W(\mu t)$ with probability R/(dQ+R) where ζ_2 is the limiting r.v. in the second case. Note that $\alpha/\gamma = \theta$ and $L_R(t)/L_Q(t) \to d$.

Theorem 5.2. Assume the conditions (3), (4), and (6) hold.

(i) If
$$\theta = 1$$
 such that $R(t) \to \infty, t \to \infty$, and $\alpha/\gamma > 1$ then
 $\mathbf{E}[s^{Y(t)}|Y(t) > 0] \to 1 - \Delta(s)/Q.$

(ii) If $\theta > 1$ and $\alpha/\gamma = 1$ such that $Q(t) \to \infty, t \to \infty$, then $\mathbf{P} \{Y(t)/W(\mu t) \le x | Y(t) > 0\} \to D(\alpha, \gamma; x), x \ge 0,$

where $D(\alpha, \gamma; x), x \ge 0$ is defined in Theorem 5.1 (ii).

Proof. (i) In this case $q(t,s) = o(r(t)), t \to \infty$, for any fixed $s \in [0,1)$ and $1 - \Phi(t;s) \sim r(t).\Delta(s)$. Following the same way as in the proof of Theorem 5.1 (i) we can complete now the proof.

(ii) In this case $R = \int_0^\infty r(t)dt < \infty$ and $r(t) = o(q(t)), t \to \infty$. Using that $q(t;s)/q(t) \to 1, t \to \infty$ for any fixed $s \in [0,1)$ (see Lemma 3.1) we can obtain similarly as in the proof of Theorem 5.1 (ii) that $1 - \Phi(t;s) \sim R.q(t;s), t \to \infty$, for any fixed $s \in [0,1)$. Therefore $[1-\Phi(t;s)]/[1-\Phi(t;0)] \to 1, t \to \infty$, i.e. almost all non-degenerate sample paths go to infinity. Working in the same way as in the proof of Theorem 5.1 (ii) we are able to complete the proof.

Comment 5.2. Theorem 5.2 can be interpreted as an extension of the cases (i) and (ii) of Theorem 5.1, where both R and Q are finite. Now $Q(t) \to \infty$ in (i) and $R(t) \to \infty$ in (ii).

Theorem 5.3. Assume the conditions (3), (4), and (6) hold. If $\theta = 1$ and $\alpha = \gamma$ such that $R(t) \uparrow R \leq \infty$, $Q(t) \uparrow \infty$, and $\frac{r(t)Q(t)}{q(t)R(t)} \to d \geq 0, t \to \infty$ then

$$\mathbf{P}\left\{Y(t)/W(\mu t) \le x | Y(t) > 0\right\} \to D_{\gamma, d}(x), x \ge 0,$$

where $D_{\gamma}(x)$ has Laplace transform

$$\hat{D}_{\gamma,d}(\lambda) = \frac{d}{1+d} + \frac{1}{1+d} \cdot \frac{1}{1+\lambda^{\gamma}}, \lambda > 0.$$

Proof. Let $s(t) = \exp(-\lambda/W(\mu t))$ and for $\delta \in (0, 1)$ fixed one has

$$I(t; s(t)) = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t; s(t)) + I_2(t; s(t)).$$

Since q(t,s) is non-increasing in t we get that

$$q(t;s(t))\int_{0}^{t(1-\delta)} r(u)du \le I_{2}(t;s(t)) \le q(t\delta;s(t))\int_{0}^{t(1-\delta)} r(u)du.$$

Applying Lemma 3.2 with $c = \delta$ and c = 1 respectively, one obtains

$$q(t\delta;s(t)) \sim \frac{q(t)}{\delta + \lambda^{-\gamma}}, \quad q(t;s(t)) \sim \frac{q(t)}{1 + \lambda^{-\gamma}}, \ t \to \infty,$$

having in mind that $\alpha = \gamma$. Therefore

$$\frac{1}{1+\lambda^{-\gamma}} \le \liminf_{t \to \infty} \frac{I_2(t;s(t))}{q(t)R(t)} \le \limsup_{t \to \infty} \frac{I_2(t;s(t))}{q(t)R(t)} \le \frac{1}{\delta + \lambda^{-\gamma}}.$$
 (26)

Let us consider $I_1(t; s(t))$. Since q(u; s) is non-increasing in u then

$$q(u;s(t))) \le q(0;s(t)) = \frac{1}{\Psi\left(W\left(V\left(\frac{1}{1-s(t)}\right)\right)\right)} = \frac{1}{\Psi\left(\frac{1}{1-s(t)}\right)},$$

because W(.) and V(.) are inverse to each other. Therefore

$$I_1(t;s(t)) = \int_0^{t\delta} r(t-u)q(u;s(t))du \le r(t(1-\delta))\int_0^{t\delta} \frac{1}{\Psi\left(\frac{1}{1-s(t)}\right)}du$$
$$\le r(t(1-\delta))\frac{t\delta}{\Psi\left(\frac{1}{1-s(t)}\right)} \sim r(t(1-\delta))\frac{t\delta}{\Psi\left(\frac{W(\mu t)}{\lambda}\right)} \sim r(t(1-\delta))(t\delta)q(t)\lambda^{\alpha},$$

using the relation $1 - \exp(-\lambda/W(\mu t)) \sim \lambda/W(\mu t), t \to \infty$, and the properties of $\Psi(.)$. Therefore $I_1(t; s(t)) = O(r(t(1 - \delta))q(t)t), t \to \infty$. Having in mind that tr(t) = o(R(t)) we conclude that $I_1(t; s(t)) = o(q(t)R(t)), t \to \infty$. From here and (26) we get that

$$\frac{1}{1+\lambda^{-\gamma}} \leq \liminf_{t\to\infty} \frac{I(t;s(t))}{q(t)R(t)} \leq \limsup_{t\to\infty} \frac{I(t;s(t))}{q(t)R(t)} \leq \frac{1}{\delta+\lambda^{-\gamma}}.$$

Since $\delta \in (0,1)$ was arbitrary we conclude that

$$\lim_{t \to \infty} I(t; s(t)) / [q(t)R(t)] = 1 / (1 + \lambda^{-\gamma}).$$

Then $I(t; s(t)) \sim q(t)R(t)/[1 + \lambda^{-\gamma}] \to 0$, and $1 - \Phi(t; s(t)) \sim q(t)R(t)/[1 + \lambda^{-\gamma}], t \to \infty$. Therefore

$$\frac{1 - \Phi(t; s(t))}{1 - \Phi(t; 0)} \sim \frac{(1 + \lambda^{-\gamma})^{-1} q(t) R(t)}{q(t) \cdot R(t) + r(t) Q(t)} \to \frac{(1 + \lambda^{-\gamma})^{-1}}{1 + d}.$$

which completes the proof of the theorem.

Comment 5.3. The conditional limiting distribution has an atom at zero with probability $\frac{d}{1+d}$. Then applying a Tauberian theorem (as in Comment 5.1) one can obtain that $1 - D_{\gamma,d}(x) \sim x^{-\gamma} / [(1+d)\Gamma(1-\gamma))$, i.e. the limiting r.v. belongs to a normal domain of attraction of a stable law with parameter γ . Note that $\alpha = \gamma$ and $\frac{L_R(t)Q(t)}{L_Q(t)R(t)} \rightarrow d$.

Theorem 5.4. Assume the conditions (3), (4), and (6) hold. (i) If $\theta \ge 1$ and $0 < \alpha/\gamma < 1$ then

$$\mathbf{P}\left\{Y(t)/W(\mu t) \le x | Y(t) > 0\right\} \to D(\alpha, \gamma; x), x \ge 0.$$

where $D(\alpha, \gamma; x), x \ge 0$ is defined in Theorem 5.1(ii). (ii) If $0 < \theta < 1$ and $\alpha/\gamma \ge 1$ such that $Q = \int_0^\infty q(u) du < \infty$ then $\mathbf{E} \left[s^{Y(t)} | Y(t) > 0 \right] \to 1 - \Delta(s)/Q.$

Proof. (i) Under the conditions of this case one has $R(t) \to R \leq \infty$, R(t) is a s.v.f. and $tr(t) = o(R(t)), t \to \infty$. Let $s(t) = 1 - \exp(-\lambda/W(\mu t))$ and consider for any fixed $\delta \in (0, 1)$

$$I(t; s(t)) = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t; s(t)) + I_2(t; s(t)).$$

Hence

$$q(t;s(t))\int_{0}^{t(1-\delta)} r(u)du \le I_2(t;s(t)) \le q(t\delta;s(t))\int_{0}^{t(1-\delta)} r(u)du.$$
(27)

Applying Lemma 3.2 with $c = \delta$ and c = 1 respectively, one gets

$$q(t\delta; s(t)) \sim q(t)(\delta + \lambda^{-\gamma})^{-\alpha/\gamma}, \ q(t; s(t)) \sim q(t)(1 + \lambda^{-\gamma})^{-\alpha/\gamma}, t \to \infty.$$

These two relations and (27) provided that

$$(1+\lambda^{-\gamma})^{-\alpha/\gamma} \leq \liminf_{t \to \infty} \frac{I_2(t;s(t))}{q(t).R(t)} \leq \limsup_{t \to \infty} \frac{I_2(t;s(t))}{q(t).R(t)} \leq (\delta+\lambda^{-\gamma})^{-\alpha/\gamma}.$$

On the other hand we have

$$0 \le I_1(t; s(t)) = \int_0^{t\delta} r(t-u)q(u; s(t))du \le r(t(1-\delta)) \int_0^{t\delta} q(u)du$$

~ $r(t)(1-\delta)^{\theta} \frac{t\delta q(t\delta)}{1-\alpha/\gamma} \sim \frac{r(t)(1-\delta)^{\theta} tq(t)\delta^{1-\alpha/\gamma}}{1-\alpha/\gamma} = o(R(t).q(t)), t \to \infty,$

remember that $tr(t) = o(R(t)), t \to \infty$. Therefore,

$$(1+\lambda^{-\gamma})^{-\alpha/\gamma} \leq \liminf_{t \to \infty} \frac{I(t;s(t))}{q(t).R(t)} \leq \limsup_{t \to \infty} \frac{I(t;s(t))}{q(t).R(t)} \leq (\delta+\lambda^{-\gamma})^{-\alpha/\gamma}.$$

Since $\delta \in (0, 1)$ was arbitrary, we get

$$I(t;s(t)) \sim (1 + \lambda^{-\gamma})^{-\alpha/\gamma} q(t) \cdot R(t) \to 0, t \to \infty,$$

for any fixed $\lambda > 0$. Therefore

$$1 - \Phi(t; s(t)) \sim I(t; s(t)) \sim (1 + \lambda^{-\gamma})^{-\alpha/\gamma} (1 - \Phi(t; 0))$$

This relation and (17) compete the proof of this case.

(ii) Let $\delta \in (0, 1)$ be fixed and

$$I(t;s) = \int_0^t r(t-u)q(u;s)du = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t;s) + I_2(t;s).$$
(28)

Then

$$r(t)\int_0^{t\delta} q(u;s)du \le I_1(t;s) \le r(t(1-\delta))\int_0^{t\delta} q(u;s)du.$$

Since $Q < \infty$ then by Lemma 3.1 it follows that $\Delta(s) = \int_0^\infty q(u; s) du < \infty$ and by the dominated convergence theorem $\Delta(s) \to Q$, as $s \to 0$. Therefore,

$$\frac{\Delta(s)}{Q} \le \liminf_{t \to \infty} \frac{I_1(t;s)}{r(t)Q(t)} \limsup_{t \to \infty} \frac{I_1(t;s)}{r(t)Q(t)} \le (1-\delta)^{-\theta} \frac{\Delta(s)}{Q}.$$
 (29)

Having in mind that q(t; s) is non-increasing in $t \ge 0$ and $s \in [0, 1)$ we obtain

$$I_2(t;s) = \int_{t\delta}^t r(t-u)q(u;s)du \le q(t\delta) \int_0^t r(u)du \sim q(t\delta)\frac{tr(t)}{1-\theta},$$

because r(t) varies regularly with exponent $-\theta \in (-1, 0)$. Therefore

$$\frac{I_2(t;s)}{r(t)Q(t)} \le \frac{tq(t\delta)r(t)}{(1-\theta)r(t)Q(t)} \to 0, t \to \infty,$$
(30)

because in this case $tq(t) = o(Q(t)), t \to \infty$. From (28), (29), and (30) it follows that

$$\frac{\Delta(s)}{Q} \le \liminf_{t \to \infty} \frac{I(t;s)}{r(t)Q(t)} \limsup_{t \to \infty} \frac{I(t;s)}{r(t)Q(t)} \le (1-\delta)^{-\theta} \frac{\Delta(s)}{Q}, \quad s \in [0,1).$$

Since δ was arbitrary, we get that $I(t;s) \sim \Delta(s)r(t)Q(t)/Q \to 0, t \to \infty$. Therefore $1 - \Phi(t;s) \sim \Delta(s)r(t)Q(t)/Q \to 0, t \to \infty$, which together with Theorem 4.2(ii) and (17) completes the proof.

Comment 5.4. Note that the obtained limiting distributions are the same as in Theorem 5.2 (respectively Theorem 5.1 - (ii) and (i)) nevertheless that the conditions and methods of the proofs are different.

Theorem 5.5. Assume the conditions (3), (4), and (6) hold with $0 < \theta < 1$ and $0 < \alpha/\gamma < 1$. (i) If $tr(t)q(t) \rightarrow 0$ then

$$\mathbf{P}\left\{Y(t)/W(t) \le x | Y(t) > 0\right\} \to D_1(\theta, \alpha, \gamma; x), x \ge 0,$$

where $D_1(\theta, \alpha, \gamma; x)$ has Laplace transform

$$\hat{D}_1(\theta, \alpha, \gamma; \lambda) = 1 - \frac{\lambda^{\alpha}}{B(1-\theta, 1-\frac{\alpha}{\gamma})} \int_0^1 (1-u)^{-\theta} (u\lambda^{\gamma}+1)^{-\alpha/\gamma} du.$$

(ii) If $tr(t)q(t) \to K \in (0,\infty)$ then

$$\mathbf{P}\left\{Y(t)/W(t) \le x | Y(t) > 0\right\} \to D_2(\theta, \alpha, \gamma; x), x \ge 0,$$

where $D_2(\theta, \alpha, \gamma; x)$ has Laplace transform

$$\hat{D}_2(\theta, \alpha, \gamma; \lambda) = 1 - \frac{1 - \exp\left(-K\lambda^\alpha \int_0^1 (1-u)^{-\theta} (u\lambda^\gamma + 1)^{1-\theta} du\right)}{1 - \exp\left(-K \cdot B(\theta, 1-\theta)\right)}.$$

(iii) If $tr(t)q(t) \to \infty$ then

$$\mathbf{P}\left\{Y(t)/\overleftarrow{\Psi}(W(t)) \le x\right\} \to D_3(\alpha; x), x \ge 0,$$

where $D_3(\theta, \alpha, \gamma; x)$ has Laplace transform $\hat{D}_3(\alpha; \lambda) = e^{-\lambda^{\alpha}}, \ \lambda > 0.$

Proof. (i) Let us denote $s(t) = \exp(-\lambda/W(\mu t))$. Let $\delta \in (0, \frac{1}{2})$ be fixed. Consider

$$I(t;s(t)) = \int_0^{t\delta} + \int_{t\delta}^{t(1-\delta)} + \int_{t(1-\delta)}^t = I_1(t;s(t)) + I_2(t;s(t)) + I_3(t;s(t)).$$

By changing variables u = vt in $I_2(t; s(t))$ we obtain

$$I_2(t;s(t)) = \int_{t\delta}^{t(1-\delta)} r(t-u)q(u;s(t))du = t \int_{\delta}^{1-\delta} r(t(1-v))q(tv;s(t))dv,$$

Applying Lemma 3.2 we have $q(vt; s(t)) \sim q(t) (v + \lambda^{-\gamma})^{-\alpha/\gamma}, t \to \infty$. Therefore

$$I_2(t;s(t)) = tr(t)q(t) \int_{\delta}^{1-\delta} \frac{r(t(1-v))}{r(t)} \frac{q(tv;s(t))}{q(t)} dv$$

~ $tr(t)q(t) \int_{\delta}^{1-\delta} (1-v)^{-\theta} (v+\lambda^{-\gamma})^{-\alpha/\gamma} dv \to 0, \quad t \to \infty,$

by the uniform convergence of the r.v.f. on compact set $[\delta, 1 - \delta]$. Further

$$I_{1}(t;s(t)) = \int_{0}^{t\delta} r(t-u)q(u;s(t))du \leq \int_{0}^{t\delta} r(t-u)q(u)du$$

$$I_{3}(t;s(t)) = \int_{t(1-\delta)}^{t} r(t-u)q(u;s(t))du \leq \int_{0}^{t\delta} r(u)q(t-u)du,$$

because q(t; s) is non-increasing in $s \in [0, 1]$. By the last three relations, having also in mind (15) and (16), and the fact that δ was arbitrary, we conclude that

$$I(t;s(t) \sim tr(t)q(t) \int_0^1 (1-v)^{-\theta} (v+\lambda^{-\gamma})^{-\alpha/\gamma} dv \to 0, t \to \infty.$$

Therefore $1 - \Phi(t; s(t)) \sim tr(t)q(t) \int_0^1 (1-v)^{-\theta} (v+\lambda^{-\gamma})^{-\alpha/\gamma} dv, t \to \infty$, which together with (17) and Theorem 4.3(i) completes the proof of this case.

(ii) Setting as above $s(t) = \exp(-\lambda/W(\mu t))$ and having in mind that in this case $\theta + \alpha/\gamma = 1$, in the same way as in the previous case we obtain that as $t \to \infty$,

$$\begin{split} &I(t,s(t)) \sim tq(t)r(t)\lambda^{\alpha}\int_{0}^{1}(1-u)^{\theta}(u\lambda^{\gamma}+1)^{1-\theta}du\\ & \rightarrow \quad K\lambda^{\alpha}\int_{0}^{1}(1-u)^{\theta}(u\lambda^{\gamma}+1)^{1-\theta}du. \end{split}$$

Then (see (18) and (19)),

_

$$\Phi(t,s(t)) \to \exp\left(-K\lambda^{\alpha}\int_0^1 (1-u)^{\theta}(u\lambda^{\gamma}+1)^{1-\theta}du\right), \quad t \to \infty.$$

This limit together with Theorem 4.3(ii) and (17) completes the proof of the case (ii).

(iii) Denote by $s(t) = \exp(-\lambda/\overleftarrow{\Psi}(R(t)))$. Remember that $\overleftarrow{\Psi}(.)$ is the inverse function of $\Psi(.)$. For $\delta \in (0, 1)$ we consider

$$I(t;s(t)) = \int_0^t r(t-u)q(u;s(t))du = \int_0^{t\delta} + \int_{t\delta}^t = I_1(t;s(t)) + I_2(t;s(t))$$

Since q(t; s) is non-increasing in $t \ge 0$ we have

$$q(t;s(t))R(t(1-\delta)) \le I_2(t;s(t)) \le q(t\delta;s(t))R(t).$$

From (8) it follows that $1/q(t) = \Psi(W(\mu t))$. Then $\mu t \sim V\left(\overleftarrow{\Psi}(1/q(t))\right)$, as $t \to \infty$. We have also that $1 - s(t) \sim \lambda/\overleftarrow{\Psi}(R(t))$. Therefore,

$$\frac{\mu t}{V\left(1/(1-s(t))\right)} \sim \frac{V\left(\overleftarrow{\Psi}\left(1/q(t)\right)\right)}{V\left(\overleftarrow{\Psi}(R(t))/\lambda\right)} \sim \lambda^{-\gamma} \frac{V\left(\overleftarrow{\Psi}\left(1/q(t)\right)\right)}{V\left(\overleftarrow{\Psi}(R(t))\right)} \sim \frac{\lambda^{-\gamma}}{q(t)R(t)} \to 0$$

because $q(t)R(t) \sim q(t)\frac{tr(t)}{\theta+1} \to \infty$. From this relation (see also (7), using the uniform convergence of regularly varying functions we get that as $t \to \infty$,

$$q(t\delta; s(t))R(t) = \frac{R(t)}{\Psi\left(W\left(V\left(1/(1-s(t))\right)\left(\frac{\mu\delta t}{V(1/(1-s(t))}+1\right)\right)\right)} \\ \sim \frac{R(t)}{\Psi\left(W\left(V\left(1/(1-s(t))\right)\right)} \sim \frac{R(t)}{\Psi(1/(1-s(t)))} \sim \frac{R(t)}{\Psi\left(\overleftarrow{\Psi}(R(t))/\lambda\right)} \\ \sim \lambda^{\alpha} \frac{R(t)}{\Psi\left(\overleftarrow{\Psi}(R(t))\right)} \sim \lambda^{\alpha} R(t) \cdot \frac{1}{R(t)} \to \lambda^{\alpha}.$$

In the same way one has that $q(t\delta; s(t))R(t(1-\delta)) \to \lambda^{\alpha}(1-\delta)$ as $t \to \infty$. Having in mind that W(.) is increasing and V(x) > 0 for any x > 1, we get

$$\begin{split} I_1(t;s(t)) &= \int_0^{t\delta} r(t-u)q(u;s(t))du \le r(t) \int_0^{t\delta} q(u;s(t))du \\ &= r(t) \int_0^{t\delta} \frac{du}{\Psi\left(W\left(\mu u + V\left(1/(1-s(t))\right)\right)\right)} \le r(t) \int_0^{t\delta} \frac{du}{\Psi\left(W\left(V\left(1/(1-s(t))\right)\right))} \\ &= \frac{t\delta r(t)}{\Psi\left(1/(1-s(t))\right)} \sim \frac{t\delta r(t)}{\Psi\left(\overline{\Psi}(R(t))/\lambda\right)} \sim \delta(1+\theta)R(t) \frac{\lambda^{\alpha}}{R(t)} \to \delta(1+\theta)\lambda^{\alpha}. \end{split}$$

Therefore

$$(1-\delta)^{\theta+1}\lambda^{\alpha} \le \liminf_{t\to\infty} I(t;s(t)) \le \limsup_{t\to\infty} I(t;s(t)) \le \lambda^{\alpha} + \delta(1+\theta)\lambda^{\alpha}.$$

Since δ was arbitrary then $\lim_{t\to\infty} I(t; s(t)) = \lambda^{\alpha}$, which completes the proof.

Comment 5.5. (i) In this case $tr(t)q(t) = L_R(t)L_Q(t)t^{1-\theta-\alpha/\gamma} \to 0$, which means that $\theta+\alpha/\gamma < 1$ or $\theta+\alpha/\gamma = 1$ but $L_R(t)L_Q(t) \to 0$. The normalizing function is regularly varying with parameter $1/\gamma$ and we obtain a conditional limiting distribution $D_1(x)$. Then by the Tauberian theorem

$$1 - D_1(x) \sim x^{-\alpha} / [\Gamma(1 - \alpha)(1 - \theta)B(1 - \theta, 1 - \alpha/\gamma)], \ x \to \infty,$$

i.e. the limiting r.v. belongs to a normal domain of attraction of a stable law with parameter α .

(ii) In this case $tr(t)q(t) = L_R(t)L_Q(t) \to K \in (0,\infty)$, because $\theta + \alpha/\gamma = 1$. Then, with the same normalizing function as in the previous case we obtain similarly that

$$1 - D_2(x) \sim K x^{-\alpha} / \{ \Gamma(1 - \alpha)(1 - \theta) [1 - \exp(-K \cdot B(\theta, 1 - \theta))] \}, \quad x \to \infty,$$

(iii) In this case $\theta + \alpha/\gamma < 1$ or $\theta + \alpha/\gamma = 1$ but $L_R(t)L_Q(t) \to \infty$. The normalization is by a s.v.f. with parameter θ/α and surprisingly the non-conditional limiting distribution is just stable with parameter α .

6 Concluding remarks

As it was shown, the asymptotic behavior of the non-visiting zero probability and the limiting distributions depend of the relations between parameters of reproduction (γ), of immigration (α) and of the decreasing Poisson intensity (θ), as well as from the corresponding s.v.f. (in some cases). The probability for non-visiting zero state converges to zero (with different rates), or to positive constant in (0, 1) which is exact calculated, or finally to 1. We obtained eight different limiting distributions under the suitable normalization. An interesting case is given in Theorem 5.1-(iii) where we obtained two singular to each other conditional limiting distributions (with different normalizing functions). Another interesting situation is presented in Theorem 5.5-(iii) where we obtained non-conditional limiting distribution which is just stable with parameter α . Note that the intensity r(t) can be interpreted as a control function. The case when r(t) is increasing is quite different and the obtained results are accepted for publication in [13].

References

- [1] K. Athreya and P. Ney Branching Processes, Springer, Berlin, 1972.
- [2] I. S. Badalbaev and I. Rahimov, Non-homogeneous Flows of Branching Processes, FAN, Tashkent, 1993. (In Russian)
- [3] C. Bingham, C. Goldie and E. Omey, *Regularly varying densities*, Publ. Inst. Math. (Beograd) (N.S.) 80(94), (2006), pp. 47–57.
- [4] C. Bingham, C. Goldie and J. Teugels, *Regular Variation*, Cambridge University Press, Cambridge, 1987.
- S. Durham, A problem concerning generalized age-dependent branching processes with immigration, Annals of Mathematical Statistics. 42 (1971), pp. 1121–1123.
- [6] W. Feller, An Introduction to Probability Theory and its Applications, Vol.2, 2nd edn., John Wiley, New York, 1971.
- [7] J. Foster and J. Williamson, Limit theorems for the Galton-Watson process with time-dependent immigration, Z. Wahrsch. Verw. Geb. 20 (1976), pp. 227–235.
- [8] T. Harris, *The Theory of Branching Processes*, Dover Publications, New York, 1969.
- [9] O. Hyrien, S. Peslak, N. M. Yanev and J. Palis, Stochastic modeling of stress erythropoiesis using a two-type age-dependent branching process with immigration, Journal of Mathematical Biology. 70 (2015), pp. 1485–1521.
- [10] HYRIEN, O. AND YANEV, N. M. (2020) Age-dependent branching processes with non-homogeneous Poisson immigration as models of cell kinetics, In *Modeling and Inference in Biomedical Sciences: In Memory* of Andrei Yakovlev, eds. D. Oakes, W. J. Hall, A. Almudevar, Springer Nature, 21–46
- [11] A. A. Imomov and E. E. Tukhtaev, On asymptotic structure of critical Galton-Watson branching processes allowing immigration with infinite variance, Stochastic Models. 39 (2023), pp. 118–140.
- [12] K. V. Mitov and N. M. Yanev, Sevastyanov branching processes with non-homogeneous Poisson immigration, Proc. Steklov Inst. Math. 282 (2013), pp. 172–185.

- [13] K. V. Mitov and N. M. Yanev, Critical Markov branching process with infinite variance allowing Poisson immigration with increasing intensity, Stochastic Analysis and Applications. (2024) https://doi.org/10.1080/07362994.2024.2384575.
- [14] K. V. Mitov, N. M. Yanev and O. Hyrien, Multitype branching processes with inhomogeneous Poisson immigration, Adv. Appl. Prob. 51A (2019), pp. 211–228.
- [15] A. G. Pakes, Some new limit theorems for critical branching processes allowing immigration, Stochastic Processes and their Applications. 3 (1975), pp. 175–185.
- [16] A. G. Pakes, Critical Markov branching processes limit theorems allowing infinite variance, Adv. Appl. Prob. 42 (2010), pp. 460–488.
- [17] I. Rahimov, Critical branching processes with infinite variance and decreasing immigration, Theory Probab. Appl. 31(1) (1986), pp. 98–110.
- [18] I. Rahimov, Critical branching processes with infinite variance and increasing immigration, Math. Notes. 53(6), (1993), pp. 628–634.
- [19] I. Rahimov, Random Sums and Branching Stochastic Processes, Springer, New York, 1995.
- [20] I. Rahimov, Homogeneous Branching processes with non-homogeneous immigration, Stochastics and Quality Control. 36 (2021), pp. 165–183.
- [21] S. Sagitov, Critical branching processes with several particle types and immigration, Theory Probab. Appl. 27(2) (1982), pp. 369–374.
- [22] B. A. Sevastyanov, Limit theorems for branching random processes of special type, Theory Probab. Appl. 2(3),(1957), pp. 321–331.
- [23] A. Y. Yakovlev and N. M. Yanev, Branching stochastic processes with immigration in analysis of renewing cell populations, Math. Biosci. 203 (2006), pp. 37-63.
- [24] N. M. Yanev and K. V. Mitov, Critical branching processes with nonhomogeneous migration, Annals of Probability. 13 (1985), pp. 923–933.
- [25] V.M. Zolotarev, More exact statement of several limit theorems in the theory of branching processes, Theor. Prob. Appl. 2 (1958), pp. 256–266.