ON THE PERIODIC ORBITS OF C⁰-TYPICAL IMPULSIVE SEMIFLOWS

JAQUELINE SIQUEIRA, MARIA JOANA TORRES, AND PAULO VARANDAS

ABSTRACT. Impulsive semiflows modeled by a Lipschitz continuous vector field and continuous impulse functions, defined over an impulsive region, are piecewise Lipschitz continuous semiflows with piecewise smooth trajectories. In this paper we contribute to the topological description of typical impulsive semiflows, parameterized by both vector fields and impulses. We prove that C^0 generic Lipschitz vector fields generate impulsive semiflows with denseness of periodic orbits on the non-wandering set. Additionally, we show that C^0 -generic impulses generate impulsive semiflows with denseness of periodic orbits on the impulsive semiflows with denseness of periodic orbits on the impulsive non-wandering set.

1. INTRODUCTION

Classical dynamical systems theory has proven essential in explaining a broad array of phenomena. However, real-world systems often experience sudden, discontinuous shifts, a factor that traditional dynamical systems theory tends to overlook. The theory of impulsive semiflows models such abrupt changes on the dynamics caused by external input and provide a quite rich playground to the study of the geometric, ergodic and topological properties of continuous-time dynamical systems under perturbations of the system by a discrete-time impulse map. These systems, characterized by a continuous semiflow acting on a compact metric space (M, d), a subset $D \subset M$ where the semiflow experiences abrupt perturbations, and an impulsive function $I : D \rightarrow I(D) \subset M$ that models these perturbations, frequently exhibit discontinuities, adding a layer of complexity to their behavior. The study of the topological properties of impulsive systems has been extensively studied in the last two decades (see e.g. [6] and references therein).

In analogy to classical dynamical systems, impulsive semiflows are likely to behave quite differently depending on the regularity of the unperturbed flows and the space of impulses. In [15], we initiated the study of the geometric and topological features for C^1 -typical impulsive flows by proving that, given an initial smooth flow, there exists a C^1 -Baire generic subset of impulses (in the space of C^1 diffeomorphisms of D onto its image) such that the corresponding impulsive semiflow admits a dense set of periodic orbits on the part of the non-wandering set that intersects the impulsive region. The methods, relying on the extension of the concepts

Date: January 8, 2025.

²⁰¹⁰ Mathematics Subject Classification. Primary: 37A05, 37A35.

Key words and phrases. Impulsive semiflows; Periodic points; Non-wandering set.

of hyperbolicity and connecting lemmas to C^1 -impulsive semiflows (possibly with discontinuities), depend strongly on the C^1 -topology in the space of impulses.

In this paper we consider impulsive semiflows on which the underlying flow is generated by Lipschitz continuous vector fields and impulses are chosen as homeomorphisms onto its image. In this way, the space of impulsive semiflows is parameterized by a vector field $X \in \mathfrak{X}^{0,1}(M)$ and an impulse $I \in \mathscr{I}_{D,\hat{D}}$ (cf. Section 2 for details). Endowing the space of vector fields and impulses with the C^0 -topology, a first natural question is to ask whether a typical impulsive semiflow (usually piecewise continuous) admits invariant measures or periodic points. The denseness of periodic orbits in the non-wandering set, often known as general density theorem, was obtained first by Pugh for C^1 -generic diffeomorphisms [14], prior to the attempts to address this problem for C^0 -generic homeomorphisms [1, 8, 10, 13]. In case of homeomorphisms, the notion of permanent periodic point replaces hyperbolicity as the key ingredient to deduce a C^0 -generic lower semicontinuity of the closure of the space of periodic orbits. In case of impulsive semiflows, the results on the abundance of periodic orbits under C^0 -perturbations are of substantial different nature whenever one considers either the vector fields or the impulses. This is due to the fact that perturbations on the vector fields have a global impact while perturbations of impulses affect only the impulsive region, and will not affect nonwandering poits that do not meet the impulsive region (compare the statements of Theorems A and C and see the examples in Subsection 3.3).

This paper is organized as follows. In Section 2 we introduce the class of impulsive semiflows considered here, and the topologies in the space of vector fields and impulses. Section 3 is devoted to the statement of the main results, considering perturbations of the impulses (Subsection 3.1) and of the vector fields (Subsection 3.2). Some applications of the main results appear in Subsection 3.3. In Section 4 we introduce the concept of Poincaré maps for this class of impulsive semiflows and prove several C^0 -perturbation lemmas (including C^0 -closing lemmas and creation of periodic attractors) both for flows and impulses. We also prove that C^0 -generic vector fields generate impulsive semiflows whose periodic orbits are all permanent, and that a similar conclusion holds for C^0 -generic impulses, while considering periodic orbits that intersect the impulsive region. The proofs of the main results are summoned in Sections 5 and 6. Finally, in Section 7 we discuss some possible future research directions.

2. Setting

In this paper, impulsive semiflows are determined by a initial flow and impulsive maps. In the next subsections we introduce the precise framework assumed for these objects. 2.1. **Impulsive semiflows.** Let *M* be a *d*-dimensional Riemaniann closed manifold $(d \ge 2)$. We will denote by dist (\cdot, \cdot) the distance induced by the Riemannian structure. The space $\mathfrak{X}^{0,1}(M)$ of Lipschitz vector fields endowed with the C^0 -topology

$$||X - Y||_{C^0} := \max_{x \in M} ||X(x) - Y(x)|| < \varepsilon$$

is a Baire set, although not complete. In particular, any C^0 -Baire generic set (i.e. a set containing a countable intersection of C^0 -dense and C^0 -open sets) is C^0 -dense. We denote by $(X^t)_t$ the flow associated to $X \in \mathfrak{X}^{0,1}(M)$. Clearly, $X^t \colon M \to M$ is a lipeomorphism (i.e. a Lipschitz map with Lipschitz inverse) for all $t \in \mathbb{R}$.

Let $X \in \mathfrak{X}^{0,1}(M)$ and let $\varphi : \mathbb{R}_0^+ \times M \to M$ be the Lipschitz continuous flow generated by it, that is to say that φ satisfies $\varphi(0, \cdot) = id$ and

$$\varphi(s+t,x) = \varphi(s,\varphi(t,x))$$
 for all $s,t \in \mathbb{R}_0^+$ and $x \in M$.

We will often use the notation $\varphi_t(x) = \varphi(t, x)$, for every $t \in \mathbb{R}_0^+$ and $x \in M$.

Given a non-empty set $D \subset M$ we will refer to any continuous map $I : D \to M$ so that $I(D) \cap D = \emptyset$ as an *impulse*. Under the above conditions we say that (M, φ, D, I) is an *impulsive* dynamical system which generates an impulsive semiflow, possibly with discontinuities, as follows.

Let $\tau_1: M \to (0, +\infty]$ be the first hitting time function to the set D given by

$$\tau_1(x) = \begin{cases} \inf \{t > 0 \colon \varphi_t(x) \in D\}, & \text{if } \varphi_t(x) \in D \text{ for some } t > 0; \\ +\infty, & \text{otherwise,} \end{cases}$$
(1)

which is known to be upper semicontinuous [7]. The subsequent impulsive times are defined in terms of τ_1 applied to the point obtained after the impulse. More precisely, setting

$$\gamma_x(t) = \varphi_t(x), \quad \forall 0 \leq t < \tau_1(x) \text{ and } \gamma_x(\tau_1(x)) = I(\varphi_{\tau_1(x)}(x)),$$

the remainder of the impulsive trajectory and impulsive times $(\tau_n(x))_{n \ge 2}$ (possibly finitely many) of the point $x \in M$ are defined recursively by

$$\gamma_x(\tau_n(x)) = I(\varphi_{\tau_n(x)-\tau_{n-1}(x)}(\gamma_x(\tau_{n-1}(x))))$$

and

$$\tau_{n+1}(x) = \tau_n(x) + \tau_1(\gamma_x(\tau_n(x))).$$

In particular,

$$\gamma_x(t) = \varphi_{t-\tau_n(x)}(\gamma_x(\tau_n(x)))$$
 whenever $\tau_n(x) \le t < \tau_{n+1}(x)$. (2)

We observe that under the standing assumption that $dist_H(D, I(D)) > 0$, where $dist_H$ stands for the Hausdorff distance, it is easy to check that

$$\sup_{n \ge 1} \{\tau_n(x)\} = +\infty \quad \text{for every } x \in M$$

hence the impulsive trajectories are defined for all positive *t*.

The *impulsive semiflow* $\psi_{X,I}$ associated to (M, φ, D, I) is now defined as

$$\begin{aligned}
\psi_{X,I} : & \mathbb{R}_0^+ \times M \longrightarrow M \\
& (t,x) \longmapsto & \gamma_x(t), \\
& 3
\end{aligned}$$
(3)

where γ_x stands for the impulsive trajectory of x determined by (M, φ, D, I) and described by (2). The arguments in [[5], Proposition 2.1] guarantee that $\psi_{X,I}$ is indeed a piecewise continuous semiflow, whose trajectories are piecewise smooth. The main goal of this paper is to describe the set of periodic orbits of generic impulsive semiflows when one considers the dependence on the impulse I and vector field X.

The orbit of $x \in M$ by the impulsive semiflow $\psi_{X,I}$ is the set $O_x = \{\gamma_x(t) : t \ge 0\}$ 0}. Whenever necessary we shall write $O_{I,x}$ and $\gamma_{I,x}$ to emphasize the impulse *I*.

We say that a point $x \in M$ is *non-wandering* if, for every neighborhood U of x and any T > 0, there exists $t \ge T$ such that

$$(\psi_{X,I})_t^{-1}(U) \cap U \neq \emptyset.$$

We denote by $\Omega(\psi_{XI})$ the set of the non-wandering-points for the semiflow ψ_{XI} .

Throughout the paper, except if mentioned otherwise, it will be a standard assumption that φ is a Lipschitz continuous flow generated by a Lipschitz continuous vector field X on a compact manifold M of dimension $d \ge 3$ and D is a codimension one smooth submanifold of M, transversal to the flow direction, such that

$$\overline{D} \cap \operatorname{Sing}(\varphi) = \emptyset, \tag{H}$$

where D stands for the closure of D and Sing(φ) stands for the equilibrium points of φ . Every codimension one compact submanifold D transversal to the flow direction satisfies (H). In this setting the first hitting time function τ_1 , defined by (1), is piecewise Lipschitz continuous.

2.2. Space of impulsive maps. In this subsection we introduce a topology on the space of impulsive semiflows, parameterized by their impulses. Let φ be a Lipschitz continuous flow generated by a vector field X, $D, \hat{D} \subset M$ be smooth codimension one submanifolds transversal to the flow such that (H) holds, and consider the space

$$\mathscr{I}_{D,\hat{D}} = \operatorname{Homeo}(D,\hat{D})$$

endowed with the C⁰-distance. In the special case that D, \hat{D} are compact, we note that the condition dist_{*H*} $(D, \hat{D}) > 0$ can be simply rewritten as $D \cap \hat{D} = \emptyset$.

As φ is a Lipschitz continuous flow and $D \subset M$ satisfies (H), if $I \in \mathscr{I}_{D\hat{D}}$ there exists $\xi > 0$ so that for every distinct, $x_1, x_2 \in D \cup \hat{D}$ one has $\{\varphi_t(x_1) : |t| \leq 1\}$ $\xi \} \cap \{\varphi_t(x_2) \colon |t| \leq \xi\} = \emptyset$. Moreover, any discontinuity point in the interior of \hat{D} arises as a point mapped by the original flow φ on the boundary of the impulsive region D.

2.3. Space of Lipschitz continuous vector fields. One of our goals will be to understand the dependence of an impulsive semiflow on the underlying vector field. In order to do so, consider two smooth codimension one submanifolds $D, \hat{D} \subset M$ so that dist_{*H*}(*D*, \hat{D}) > 0. We consider the space of vector fields

$$\mathfrak{B}_{D,\hat{D}} = \left\{ X \in \mathfrak{X}^{0,1}(M) \colon D \pitchfork X \text{ and } \hat{D} \pitchfork X \right\},\tag{4}$$

endowed with the C^0 -topology.

3. STATEMENT OF THE MAIN RESULTS

This section is devoted to the statement of the main results, concerning the set of periodic orbits of typical impulsive semiflows. In Subsection 3.1 we assume that a vector field $X \in \mathfrak{X}^{0,1}(M)$ is given and we consider the dependence on the impulse *I*. In Subsection 3.2 we assume that an impulse *I* is fixed and we consider the dependence of the impulsive semiflow on the vector field *X*.

3.1. C^0 -Baire generic impulses. Throughout let *M* be a compact Riemannian manifold of dimension $d \ge 3$.

Theorem A. Let φ be a Lipschitz continuous flow generated by $X \in \mathfrak{X}^{0,1}(M)$ and $D, \hat{D} \subset M$ be smooth codimension one submanifolds transversal to the flow such that hypothesis (H) holds. There exists a C^0 -Baire generic subset $\mathscr{R}_X \subset \mathscr{I}_{D,\hat{D}}$ of impulses such that

$$Per(\psi_{X,I}) \cap \mathring{D} = \Omega(\psi_{X,I}) \cap \mathring{D}$$
 (5)

for every $I \in \mathscr{R}_X$, where $Per(\psi_{X,I})$ denotes the set of periodic orbits of $\psi_{X,I}$, and \mathring{D} stands for the interior of D.

Some comments are in order. Firstly, as $(\psi_{X,I})_t(x) = \varphi_t(x)$ for every $x \in D$ and for every small t > 0, periodic orbits under $\psi_{X,I}$ never intersect D. Secondly, the conclusion of Theorem A cannot be written using the landing region \hat{D} , as there exist C^1 -open sets of impulses for which the equality $\overline{Per(\psi_{X,I})} \cap \hat{D} = \Omega(\psi_{X,I}) \cap \hat{D}$ fails (see e.g. [15]). The next result is a direct consequence of the continuous dependence on initial conditions for the Lipschitz continuous flow φ and Theorem A, whose proof follows closely Corollary B in [15].

Corollary B. Let φ be a Lipschitz continuous flow generated by $X \in \mathfrak{X}^{0,1}(M)$ and $D, \hat{D} \subset M$ be smooth codimension one submanifolds transversal to the flow such that assumption (H) holds. The following hold:

- (1) if $I_0 \in \mathscr{I}_{D,\hat{D}}$ is such that $\Omega(\psi_{I_0}) \cap \partial D = \emptyset$ then there exist $\delta > 0$, an open neighborhood \mathcal{V} of I_0 and a Baire generic subset $\mathscr{R} \subset \mathcal{V}$ so that, for every $I \in \mathscr{R}$ one can write the non-wandering set $\Omega(\psi_{X,I})$ as a (possibly non-disjoint) union $\Omega(\psi_{X,I}) = \overline{Per(\psi_{X,I})} \cup \Omega_2(\varphi, D)$, where $\Omega_2(\varphi, D) \subset \Omega(\psi_{X,I})$ is a φ -invariant set which does not intersect a δ neighborhood of the cross-section D. Moreover, the set $\Omega(\psi_{X,I}) \setminus D$ is a $\psi_{X,I}$ -invariant subset of M.
- (2) if φ is minimal then there exists a Baire generic subset $\mathscr{R} \subset \mathscr{I}_{D,\hat{D}}$ so that, for every $I \in \mathscr{R}$, the set of periodic orbits is dense in $\Omega(\psi_{X,I})$. Moreover, the set $\Omega(\psi_{X,I}) \setminus D$ is a $\psi_{X,I}$ -invariant subset of M.

3.2. C^0 -Baire generic Lipschitz flows. Throughout let M be a compact Riemannian manifold of dimension $d \ge 3$. Our next results address the set of periodic orbits of impulsive semiflows obtained from a fixed impulsive map and typical Lipschitz continuous vector fields. **Theorem C.** Let D, $\hat{D} \subset M$ be smooth manifolds of codimension one and $I \in \mathscr{I}_{D,\hat{D}}$ be an impulsive map. There exists a C^0 -Baire generic subset $\mathscr{R}_I \subset \mathfrak{V}_{D,\hat{D}}$ so that,

$$Per(\psi_{X,I}) = \Omega(\psi_{X,I}), \text{ for every } X \in \mathscr{R}_I$$

Some comments are in order. First, it should be noticed that the impulsive semiflows are at most as regular as the impulsive map, hence rarely Lipschitz continuous. Second, the statement of Theorem C is substantially stronger than the conclusion of Theorem A, and this is due to the fact that perturbing the vector field one can obtain denseness of periodic orbits even among points in the non-wandering set that do not visit the impulsive region.

3.3. **Examples.** In this section we shall provide two simple examples of the main results in order to illustrate possible applications. The first application concerns the class of impulsive Anosov flows.

Example 3.1. (C^0 -generic impulsive Anosov flows) Let X be a C^1 -vector field on a compact three-dimensional compact manifold generating a transitive Anosov flow ϕ , let D, \hat{D} be codimension one smooth submanifolds of M transversal to X and $I \in \mathscr{I}_{D,\hat{D}}$. Let $\mathcal{U} \in \mathfrak{X}^{0,1}(M)$ be a small open neighborhood of X (in the C^0 topology) so that $\mathcal{U} \subset \mathfrak{B}_{D,\hat{D}}$. Even though it is not known whether all flows generated by vector fields in \mathcal{U} are equivalent, hence have a dense set of periodic orbits, by the proof of Theorem C, periodic points are all permanent and the set of periodic points is dense in the non-wandering set, for a C^0 -generic choice of $Y \in \mathcal{U}$. Furthermore, Theorem A implies that the set of (permanent) periodic orbits is dense in the impulsive non-wandering set for a C^0 -generic choice of the impulses $I \in \mathscr{I}_{D,\hat{D}}$.

Example 3.2. (Perturbations of minimal flows) Let φ be a minimal flow on \mathbb{T}^3 generated by a C^1 vector field $X \in \mathfrak{X}^1(\mathbb{T}^3)$ and $D, \hat{D} \subset M$ be smooth codimension one closed submanifolds transversal to the flow. The flow φ has no periodic orbits. Nevertheless, Theorem A ensures that for a C^0 -generic choice of the impulse $I \in \mathscr{I}_{D,\hat{D}}$, the impulsive semiflow $\psi_{X,I}$ has a dense set of periodic orbits in the impulsive non-wandering set.

The next example, adapted from [15] illustrates some of the subtleties between the perturbations in the space of impulses and the perturbations in the space of vector fields.

Example 3.3. Consider the impulsive semiflow constructed on the cylinder written as

$$S = \{(x, y, z) \in \mathbb{R}^3 : x^2 + y^2 = 1, z \in [-1, 1]\} \subset \mathbb{R}^3$$

and let φ be the flow generated by the vector field X(x, y, z) = (-y, x, 0) whose trajectories are either constant at the origin or circles spinning counterclockwise in each disk $S \cap \{z = c\}$, with $c \in [-1, 1]$. One can extend this flow to the cube $[-3, 3]^3$ in a way that no new periodic orbits are created. Consider the local crosssections D, \hat{D} in such a way that $D \cap S = \emptyset$ and \hat{D} intersects S (cf. Figure 1).




FIGURE 1. Representation of the impulsive flow on the section $\{z = 0\}$, where $D_0 = D \cap \{z = 0\}$ and $\hat{D}_0 = \hat{D} \cap \{z = 0\}$. Existence of non-permanent periodic orbits in $\Omega(\psi_{X,I}) \cap \hat{D}_0$.

Theorem A implies that the set of (permanent) periodic orbits is dense in the impulsive non-wandering set for a C^0 -generic choice of the impulses $I \in \mathscr{I}_{D,\hat{D}}$. Clearly, the periodic orbits that intersect \hat{D} are permanent as a function of I (perturbations of the impulse keep them unnaltered) but are not permanent as a function of the vector field X. When perturbations of the vector field are allowed, Theorem C guarantees that for a generic vector field $X \in \mathfrak{B}_{D,\hat{D}}$, all periodic orbits are permanent.

4. Preliminares

4.1. **Poincaré maps for impulsive semiflows.** Given a Lipschitz continuous flow generated by a vector field *X*, a periodic orbit γ and a local cross-section Σ passing through a point $p \in \gamma$, the Poincaré map $P : U \subset \Sigma \to \Sigma$ is a bi-Lipschitz local homeomorphism defined on some open neighborhood $U \subset \Sigma$ of *p* which is the first return map of points in *U* to Σ .

In case of impulsive semiflows, by some abuse of notation we shall denote as Poincaré map the ones that concatenate the impulse and the first hitting time map to *D*. More precisely, given $I \in \mathscr{I}_{D,\hat{D}}$ we shall consider the map (which we denote as a Poincaré map)

$$P_{X,I}: \begin{array}{ccc} \hat{D} & \to & \hat{D} \\ x & \mapsto & I \circ \varphi_{\tau(x)}(x) \end{array}$$
(6)

where

$$\tau(x) = \inf\{t > 0 \colon \varphi_t(x) \in D\}$$
(7)

for every $x \in \hat{D}$ and $\hat{\hat{D}} = \{x \in \hat{D} : \tau(x) < \infty\}$. Notice that $\tau = \tau_1 \mid_{\hat{D}}$. Consider the set

$$\widehat{I_*(D)} = \{ x \in \widehat{\hat{D}} : \varphi_{\tau(x)}(x) \notin \partial D \}$$

which consists of the points in \hat{D} which are mapped in the interior of D. It will be also convenient to consider the map $f_{X,I}: D \to D$ given by

$$f_{X,I} = I^{-1} \circ P_{X,I} \circ I, \tag{8}$$

which is defined on the impulsive region *D*, which is the support of all perturbations of the impulses.

In what follows we prove that, even though the Poincaré map in (6) is defined by expressions which take into account the impulse it is a local homeomorphism.

Proposition 4.1. Let φ be a Lipschitz continuous flow generated by $X \in \mathfrak{X}^{0,1}(M)$, D, \hat{D} be smooth submanifolds of codimension one transversal to X so that (H) holds and $\overline{D} \cap \overline{\hat{D}} = \emptyset$, and let $I \in \mathscr{I}_{D,\hat{D}}$ be an impulse. Given $x \in \widehat{I_*(D)}$ there exist a local cross-section $\hat{\Sigma} \subset \hat{D}$ and a neighborhood $\hat{V} \subset \hat{\Sigma}$ of x such that the Poincaré map $P_I|_{\hat{V}}$ is an homeomorphism onto its image.

Proof. By the assumption $\overline{D} \cap \overline{\hat{D}} = \emptyset$, there is an open neighborhood W of \hat{D} such that $\psi_{X,I}|_W = \varphi$. Let $x \in \widehat{I_*(D)}$ and consider a local cross-section $\hat{\Sigma} \subset \hat{D}$. The Lipschitz flowbox theorem for φ (cf. Theorem 4 in [3]) guarantees the existence of a flow-box (V_1, h_1) at x where $V_1 \subset W$ is an open neighborhood of x, and the chart map $h_1 : V_1 \to [0, 1] \times U_1 \subset [0, 1]^d$ sends trajectories of X in V_1 into trajectories of the vector field $X_{V_1}(z) := (1, 0, \dots, 0)$ for every $z \in [0, 1]^d$.

Choose $t_0 > 0$ small such that $\varphi_{t_0}(x)$ still lies in V_1 and consider the compact segment of orbit $\Gamma := \{\varphi_t(x) : t_0 \leq t \leq \tau(x)\}$. Since $\tau(x) < \infty$, there exists a long tubular Lipschitz flow-box chart (V_2, h_2) for the original flow φ , say $h_2 :$ $V_2 \to [0, 1] \times U_2 \subset [0, 1]^d$ such that $V_2 \supset \Gamma$. Let $\Sigma_1 = h_2^{-1}(\{0\} \times U_2)$ and $\Sigma_2 = h_2^{-1}(\{1\} \times U_2)$ be the components of the boundary of V_2 which are transversal to the flow φ . Denote by $\pi_1 : \hat{V} \subset \hat{\Sigma} \to \Sigma_1, \pi_2 : \Sigma_1 \to \Sigma_2$, the projections along the positive trajectories of φ given by the tubular flowbox theorems, where \hat{V} is a small open neighborhood of x in $\hat{\Sigma} \cap \widehat{I_*(D)}$. Similarly, let $\pi_3 : \Sigma_2 \to \Sigma$ be the projection along the negative trajectories of φ , where Σ is a small neighborhood of $\varphi_{\tau(x)}(x)$ in D. Each $\pi_i, i = 1, 2, 3$ is a Lipschitz continuous map. Since

$$P_I|_{\hat{V}} \equiv I \circ \pi_3 \circ \pi_2 \circ \pi_1$$

and the impulse *I* is an homeomorphism we have that $P_I|_{\hat{V}}$ is continuous. Using the invertibility of the impulse *I* and the flow φ , the same argument can be done for construct the inverse of the Poincaré map as before, proving that it is a local homeomorphism.

4.2. C^0 -perturbations lemmas. In our context impulsive semiflows are determined by Lipschitz continuous vector fields and continuous impulses.

 C^{0} -perturbations of impulses. We consider first the case where the Lipschitz continuous vector field is fixed, and the C^{0} -perturbations are considered in the space of impulses.

Lemma 4.1. (C^0 -closing lemma: impulses) Let M be a compact manifold of dimension $d \ge 3$, let D, \hat{D} be smooth submanifolds of codimension one and let $X \in \mathfrak{B}_{D,\hat{D}}$. Assume that $I \in \mathscr{I}_{D,\hat{D}}$ is an impulsive map and $p \in \Omega(\psi_{X,I})$ is a regular point so that $\overline{\gamma_p} \cap D \neq \emptyset$ and that $\gamma_p \cap \partial \hat{D} = \emptyset$. For every $\varepsilon > 0$ there exists $J \in \mathscr{I}_{D,\hat{D}}$ which is ε - C^0 -close to I and so that $\psi_{X,J}$ has a periodic orbit x so that $dist(p, x) < \varepsilon$.

Proof. We may assume that $p \in \Omega(\psi_{X,I})$ is not periodic for $\psi_{X,I}$, otherwise there is nothing to be done. Up to consider a point on the orbit of p, we may assume without loss of generality that $p \in \Omega(\psi_{X,I}) \cap \hat{D}$. Observe that p is a non-wandering point for the Poincaré map P_I given by (6). In particular, $q = I^{-1}(p)$ is a non-wandering point for $f_I : D \to D$ expressed by (8).

Given $\varepsilon > 0$, there exist $x \in \hat{D}$ and a least integer $n \ge 1$ such that $d(x, p) < \varepsilon$ and $d(x_n, p) < \varepsilon$, where $x_i = P_I^i(x)$. Noting that \hat{D} is of dimension $d - 1 \ge 2$, that $\{x_i : 1 \le i \le n\}$ is a finite collection of pairwise disjoint points and $d(x_n, p) < \varepsilon$, there exists an homeomorphism $\zeta : \hat{D} \to \hat{D}$ homotopic to the identity such that $\zeta(x_n) = x, \zeta(x_i) = x_i$ for every $1 \le i \le n - 1$ and $d_{C^0}(\zeta, id) < \varepsilon$. The impulse $J = \zeta \circ I$ is ε - C^0 -close to I and $x \in Per(\psi_{X,J})$. This finishes the proof of the lemma.

 C^{0} -perturbations of vector fields. Let us now obtain some C^{0} -perturbation lemmas for Lipschitz continuous vector fields, whenever the impulse is predetermined. First we recall the flowbox theorem for Lipschitz vector fields.

Lemma 4.2. [3, Theorem 4] Let $X \in \mathfrak{X}^{0,1}(M)$ be a Lipschitz continuous vector field, $(\varphi_t)_t$ is the flow generated by X and $(T^t)_t$ is the trivial flow on $\mathbb{R} \times \mathbb{R}^{d-1}$ defined by $T^t(s,z) = (s + t, z)$. If p is a regular point then there exist $\delta > 0$, a neighborhood U_{δ} of p and a lipeomorphism $h : U_{\delta} \to [0, \delta] \times B(\vec{0}, \delta)$ so that $h \circ \varphi_t(x) = T^t \circ h(x)$ for every $x \in U_{\delta}$ and $t \in \mathbb{R}$ so that $\varphi_t(x) \in \overline{U_{\delta}}$.

In the notation of the previous flowbox theorem, given $0 < r < \delta$ we will consider also the flowbow 'cylinders' $\mathcal{F}_{X,\delta}(B) := h^{-1}([0,\delta] \times B)$, for $B \subset B(\vec{0},\delta)$, and the local cross-section $\Sigma_p := h^{-1}(\{0\} \times B(\vec{0},\delta))$ to the flow $(\varphi_t)_t$ at p. Since the points in the impulsive region D do not belong to the orbits of the impulsive semiflow, the latter provides a local trivializing chart for regular points of an impulsive semiflow whose orbit closure does not intersect ∂D . The following perturbation lemma will also be instrumental.

Lemma 4.3. [2] Let $X \in \mathfrak{X}^{0,1}(M)$, let p be a regular point, Σ_p be a local crosssection through p and let h be the lipeomorphism given by Lemma 4.2. Assume there exists a C^{1} -arc $\gamma : [0, \delta] \to \Sigma_p$ with constant speed ν (hence of length $\nu\delta$) such that $\gamma(0) = p$ and $\gamma(\delta) = q$. Then there exists C > 0 so that for any $\eta > \nu\delta$, any open tubular neighborhood $B \subset \Sigma_p$ of diameter smaller than η containing γ and any $\varepsilon > 0$ there exists a vector field Y such that: (a) $Y \in \mathfrak{X}^{0,1}(M)$; (b) Y is $C\eta(1 + \varepsilon)$ - C^0 -close to X; (c) $Y^{\delta}(p) = X^{\delta}(q)$ and (d) Y = X outside $\mathcal{F}_{X,\delta}(B)$. The previous perturbation lemma is the key ingredient in the proof of the following C^0 -closing lemma for impulsive semiflows, when perturbing the Lipschitz vector fields.

Lemma 4.4. (C^0 -closing lemma: vector fields) Let $X \in \mathfrak{X}^{0,1}(M)$, D, \hat{D} be smooth submanifolds of codimension one transversal to the flow direction and $I \in \mathscr{I}_{D,\hat{D}}$ be an impulsive map. If $p \in \Omega(\psi_{X,I})$ is a regular point so that $\gamma_p \cap \partial \hat{D} = \emptyset$, then for every $\varepsilon > 0$ there exists $Y \in \mathfrak{X}^{0,1}(M)$ which is ε - C^0 -close to X, coincides with X outside of a small open neighborhood of p, and such that the impulsive semiflow ψ_Y has a periodic orbit \tilde{p} so that $d(p, \tilde{p}) < \varepsilon$.

Proof. In case *p* is periodic for $\psi_{X,I}$ we are done. Hence, we may assume that *p* is non-periodic. We subdivide the proof in two cases. First, if $p \in \Omega(\psi_{X,I}) \setminus D$ is a regular point, for any $0 < r < \delta$ and corresponding flowbow $\mathcal{F}_{X,\delta}(B(\vec{0}, r)) = h_1^{-1}([0, \delta] \times B(\vec{0}, r))$ associated to *p* there exists T > 0 such that $\gamma_T(\Sigma_p) \cap \Sigma_p \neq \emptyset$ where $\Sigma_p = h_1^{-1}(\{0\} \times B(\vec{0}, r))$. Reduce $\delta > 0$, if necessary, so that $\mathcal{F}_{X,\delta}(B(\vec{0}, r)) \cap D = \emptyset$. Consequently, just pick $0 < r < \delta$ small (it is enough that $4Cr < \varepsilon$) so that the perturbed vector field $Y \in \mathfrak{X}^{0,1}(M)$ guaranteed by Lemma 4.3 is ε - C^0 -close to the vector field X and ψ_Y admits a periodic orbit that intersects $\mathcal{F}_{X,\delta}(B(\vec{0}, r))$. Second, if $p \in \Omega(\psi_{X,I}) \cap D$ then there exists a small t > 0 so that $q = X_{-t}(p) \in \Omega(\psi_{X,I}) \setminus D$, which reduces this case to the previous context. This finishes the proof of the lemma.

4.3. **Permanent periodic orbits.** Given a continuous map f on a compact manifold M, we will recall the fixed point index used in [8, 10]. Let B be an open ball on M whose boundary ∂B is a embedded sphere and assume that either (i) $f(\overline{B}) \cap \overline{B} = \emptyset$, or (ii) $f(\overline{B}) \cup \overline{B}$ is contained in a single coordinate chart. If, in addition, f has no fixed points in ∂B then the *fixed point index* $\iota_f(B)$ is defined as follows:

(1) $\iota_f(B) = 0$, if $f(\overline{B}) \cap \overline{B} = \emptyset$; and

(2) $\iota_f(B) = \deg(\gamma)$, in the case that $f(\overline{B}) \cup \overline{B}$ is contained in a single coordinate chart, where $\deg(\gamma)$ denotes the Hopf degree of the map $\gamma : \partial B \simeq S^{n-1} \rightarrow S^{n-1}$ which is defined (after a change of coordinates) by

$$\gamma(x) = \frac{f(x) - x}{\|f(x) - x\|}.$$

We will make use of the following properties of the fixed-point index for continuous maps, in the order we shall use them (cf. [8]):

- (P1) if $f(B) \subset B$, $f|_B$ is a topological contraction and p is the unique attracting fixed point in B then $\iota_f(B) \neq 0$;
- (P2) if $\iota_f(B)$ is well defined then $\iota_f(B) = \iota_g(B)$ for every continuous map g sufficiently C^0 -close to f;
- (P3) if $\iota_f(B) \neq 0$ then *f* has a fixed point in *B*.

In this framework of impulsive semiflows we will use the fixed-point index for Poincaré maps. More precisely, given a periodic orbit γ , a point $p \in \gamma$ and a local

cross section Σ to the impulsive semiflow passing through p, the Poincaré return map is defined in an open neighborhood $\Sigma_0 \subset \Sigma$ of p by

$$P(x) = \begin{cases} \varphi_{\tau_{\Sigma}^{+}(x)}(x) &, \text{ if } \gamma \cap \hat{D} = \emptyset \\ (\varphi_{\tau_{\Sigma}^{+}(\cdot)} \circ P_{I}^{n} \circ \varphi_{-\tau_{\hat{D}}^{-}(\cdot)})(x) &, \text{ otherwise,} \end{cases}$$
(9)

where $\tau_S^{\pm} : S \to \mathbb{R}^+$ denotes the first hitting time map of the flow $(\varphi_{\pm t})_{t \in \mathbb{R}}$ to a cross-section *S* and $n = \#(\gamma \cap \hat{D})$. Clearly, P(p) = p if $\gamma \cap \hat{D} = \emptyset$ and $P^n(p) = p$, otherwise.

We will say that a periodic orbit γ for the impulsive semiflow $\psi_{X,I}$ is an *attracting periodic orbit* if there exists a local cross-section Σ and $p \in \Sigma \cap \gamma$ that is an attracting fixed point for the Poincaré map *P*.




FIGURE 2. Three types of periodic orbits for the impulsive semiflows

We will use the following notions of permanence adapted from topological dynamics.

Definition 4.1. Fix $X \in \mathfrak{X}^{0,1}(M)$ and $I \in \mathscr{I}_{D,\hat{D}}$ and a periodic orbit γ for the impulsive semiflow $\psi_{X,I}$. We say that:

- (a) γ is permanent with respect to the impulses (we write $\gamma \in \mathscr{P}_{impulsive}(\psi_{X,I})$ if for any impulse $J \in \mathscr{I}_{D,\hat{D}} C^0$ -arbitrarily close to I the impulsive semiflow $\psi_{X,J}$ has a periodic orbit near γ ;
- (b) γ is permanent with respect to vector fields (we write simply $\gamma \in \mathscr{P}_{fields}(\psi_{X,I})$ if for any vector field $Y \in \mathfrak{X}^{0,1}(M)$ that is C^0 -arbitrarily close to X the impulsive semiflow $\psi_{Y,I}$ has a periodic orbit near γ .

For notational simplicity, we shall denote by $\mathscr{P}(\psi_{X,I})$ the set of all permanent closed orbits of $\psi_{X,I}$ irregardeless, whenever there is no confusion on the perturbations one considers. We need some instrumental results.

Lemma 4.5. Let $X \in \mathfrak{X}^{0,1}(M)$ and $I \in \mathscr{I}_{D,\hat{D}}$. Assume that $p \in Per(\psi_{X,I})$ and $\gamma_p \cap \partial \hat{D} = \emptyset$. There exists a C^0 -small perturbation $Y \in \mathfrak{X}^1(M)$ of X such that p is an attracting periodic point for $\psi_{Y,I}$.

Proof. Fix $X \in \mathfrak{X}^{0,1}(M)$ and $I \in \mathscr{I}_{D,\hat{D}}$. Assume that $\gamma_p \cap \partial \hat{D} = \emptyset$ and $\gamma_p \cap \hat{D} \neq \emptyset$ (the case $\bar{\gamma}_p \cap D = \emptyset$ is simpler and left to the reader). Write $\gamma_p \cap \hat{D} = \{y_0, y_1, \dots, y_{k-1}\}$ and $x_j = I^{-1}(y_j)$ for $0 \leq j \leq k-1$.

As the orbit γ_p does not intersect $\partial \hat{D}$, I^{-1} is equicontinuous and the space of C^1 vector fields is C^0 -dense in the space of Lipschitz vector fields, given $\varepsilon > 0$, there exists $Y^1 \in \mathfrak{X}^1(M)$ that is C^0 -sufficiently close to X in such a way that the first k intersections of the forward orbit γ_p^1 of p, by the impulsive semiflow ψ_{Y^1} , with the cross-section \hat{D} consists of the points $\{\bar{y}_0, \bar{y}_1, \dots, \bar{y}_{k-1}\}$ and their pre-images $\bar{x}_i = I^{-1}(\bar{y}_i) \in D$ satisfy $f_{Y^1,I}^i(\bar{x}_i) = \bar{x}_{i+1}$ for every $0 \leq i < k-1$ and dist $_D(f_{Y^1,I}(\bar{x}_{k-1}), \bar{x}_0) < \varepsilon$. Hence, one can perform a ε - C^0 -small perturbation $Y^2 \in \mathfrak{X}^1(M)$ of Y^1 supported outsise of a neighborhood of p in such a way that $f_{Y^2,I}^k(\bar{x}_0) = \bar{x}_0$, hence $p \in \operatorname{Per}(\psi_{Y^2,I})$.

We proceed with the proof by choosing an appropriate flowbox for a compact piece of orbit of the flow $(Y_t^2)_{t \in \mathbb{R}}$ containing the point \bar{x}_0 . More precisely, by the flowbox theorem there exist $\delta > 0$, $U_0 \subset M$ and a trivializing diffeomorphism $h_0: U_0 \rightarrow [0, \delta] \times B(\vec{0}, \delta)$ so that $h_0(X_{-\delta}(\bar{x}_0)) = (0, \vec{0})$ and $h_0(\bar{x}_0) = (\delta, \vec{0})$, as in Figure 3 below. Diminish $\delta > 0$ if necessary in such a way that $x_i \notin U_0$ for every $1 \leq i \leq k - 1$.




FIGURE 3. Flowbox on the last lap of the periodic orbit

The strategy will be to make a C^0 -small perturbation of Y^2 to obtain a C^1 -vector field Y^3 whose Poincaré map is a contraction. In order to do so, let us first construct the perturbation in local flowbox coordinates.

Given $0 < \eta \ll \delta < 1$ consider the C^1 bump-function $\beta_\eta \colon B(\vec{0}, \delta) \to [0, \eta]$ such that $\beta_\eta \equiv \eta$ inside $B(\vec{0}, \delta/3)$ and $\beta_\eta \equiv 0$ outside $B(\vec{0}, \delta/2)$, and consider a C^1 -function $\alpha_\eta \colon [0, \delta] \to [0, \eta]$ so that $\alpha_\eta(s) = 0$ for every $s \in [0, \eta] \cup [\delta - \eta, \delta]$ and $\alpha_\eta(s) = \eta$ for every $s \in [2\eta, \delta - 2\eta]$. We define the C^1 -vector field $Z \in \mathfrak{X}^1([0, \delta] \times B(\vec{0}, \delta))$ by $Z = Z^0 + P^\eta$ where $Z^0 = (1, 0, \dots, 0)$ and

$$P^{\eta}(s,z) = \left(0, \, \alpha_{\eta}(s) \cdot \beta_{\eta}(z) \cdot z\right).$$

The vector fields Z^0 and P^{η} commute, $||Z - Z^0||_{C^0} \leq \eta^2$ and $||Z_{\delta} \circ Z^0_{-\delta}(\delta, x)|| \leq \eta^2 ||(\delta, x)||$ for every $x \in B(\vec{0}, \delta/3)$.

Finally, let $Y = Y^3 \in \mathfrak{X}^1(M)$ be the vector field defined by

$$Y(x) = Dh_0(x)^{-1}Z(h_0(x))$$

for each $x \in U_0$ and $Y = Y^2$ otherwise, and note that $||Y^3 - Y^2||_{C^0} \leq C\eta^2$, where $C = ||Dh_0(\cdot)^{-1}||_{C^0} > 0$. Moreover, on $h_0^{-1}([0, \delta] \times B(\vec{0}, \delta/3)) \subset U_0$ the flow $(Y_t^3)_{t \in \mathbb{R}}$ is the composition of a translation and a strong transversal contraction. Altogether, $f_{Y^3}^k(x_0) = x_0$ and, as the hitting times $\bar{\tau}_{Y^3}$ and $\bar{\tau}_{Y^2}$ coincide,

$$f_{Y^{3},I}^{k} = \varphi_{\bar{\tau}_{Y^{3}}}^{Y^{3}} \circ I \circ f_{Y^{2},I}^{k-1} = [\varphi_{\delta}^{Y^{3}} \circ \varphi_{-\delta}^{Y^{2}}] \circ f_{Y^{2},I}^{k}$$

on $D \cap \overline{U_0}$. Choosing $0 < \eta \ll 1$ sufficiently small we obtain that $f_{Y^3,I}^k$ is a uniform contraction on an open neighborhood of x_0 . The vector field $Y = Y^3 \in \mathfrak{X}^1(M)$ satisfies the requirements of the lemma.

The following two results guarantees that, for typical impulsive semiflows, all periodic points are permanent. The first result concerns impulsive semiflows parameterized by vector fields. We denote by $\text{Per}_t(\psi_{X,I})$ the set of periodic points with period smaller or equal than *t* of $\psi_{X,I}$.

Proposition 4.2. Fix $I \in \mathscr{I}_{D,\hat{D}}$. There exists a C^0 -residual subset $\mathscr{R}_I \subset \mathfrak{V}_{D,\hat{D}}$ such that $Per(\psi_{X,I}) = \mathscr{P}(\psi_{X,I})$ for any $X \in \mathscr{R}_I$.

Proof. The proof exploits ideas developed in [8], adapted with the many subtleties of impulsive semiflows, due to possible discontinuities caused by impulses. Fix $I \in \mathscr{I}_{D,\hat{D}}$ and let $(\mathscr{B}_i)_{i\geq 1}$ be a countable open basis of the topology on M, which is formed by balls whose boundaries are embedded spheres. For every $i, n \geq 1$, consider the pairwise disjoint subsets $\mathscr{F}_{i,n}, \mathscr{I}_{i,n}$ of $\mathfrak{B}_{D,\hat{D}}$ defined as follows:

- (1) $X \in \mathscr{F}_{i,n}$ if and only if $\overline{\mathscr{B}_i} \cap \bigcup_{0 < t \leq n} \operatorname{Per}_t(\psi_{X,I}) = \emptyset$;
- (2) $X \in \mathscr{I}_{i,n}$ if and only if there exists $\overline{\mathscr{B}_j} \subset \mathscr{B}_i$, diam $(\mathscr{B}_j) < \text{diam}(\mathscr{B}_i)$, and a local cross-section Σ so that
 - (a) $\partial(\mathcal{B}_j \cap \Sigma) \cap \bigcup_{0 < t \le n} \operatorname{Per}_t(\psi_{X,I}) = \emptyset;$
 - (b) there exists 0 < t < n so that $\mathcal{B}_i \cap \operatorname{Per}_t(\psi_{X,I}) \neq \emptyset$;
 - (c) $\iota_{P_{X,I}}(\mathcal{B}_j \cap \Sigma) \neq 0$, where *P* stands for the Poincaré map defined in (9) for the impulsive semiflow $\psi_{X,I}$.

Let us denote by $\mathring{\mathscr{F}}_{i,n}$ and $\mathring{\mathscr{J}}_{i,n}$ the C^0 -interior of the subsets $\mathscr{F}_{i,n}$ and $\mathscr{J}_{i,n}$ of $\mathfrak{B}_{D,\hat{D}}$, respectively. We need the following.

Claim: $\mathring{\mathscr{F}}_{i,n} \cup \mathring{\mathscr{I}}_{i,n}$ is a C^0 -open and dense subset of $\mathfrak{V}_{D,\hat{D}}$ for each $i, n \ge 1$.

Proof of the Claim. It is enough to show that every $X \notin \mathring{\mathscr{F}}_{i,n}$ can be C^0 -approximated by some vector field $Y \in \mathring{\mathscr{F}}_{i,n}$. Indeed, if $X \notin \mathring{\mathscr{F}}_{i,n}$ then for every $\varepsilon > 0$ there exists $Y_0 \in \mathfrak{B}_{D,\hat{D}}$ such that Y_0 is ε - C^0 -close to X and $\overline{\mathscr{B}}_i \cap \bigcup_{0 < t \leq n} \operatorname{Per}_t(\psi_{Y_0,I}) \neq \emptyset$. One can use Lemma 4.3 to obtain a small C^0 -perturbation Y_1 of the vector field

 Y_0 , supported on a finite number of flowboxes which intersect D and \hat{D} , in such a way that there exists a periodic orbit intersecting the interior of \mathcal{B}_i that does not intersect $\partial \hat{D}$. By Lemma 4.5 one can perform a small ε - C^0 -perturbation $Y_2 \in \mathfrak{V}_{D,\hat{D}}$ of the vector field Y_1 which exhibits an attracting periodic orbit which intersects the interior of \mathcal{B}_i .

We proceed to show that there exists an open subset $\mathcal{B}_j \subset \mathcal{B}_i$ so that items (2a)-(2c) above hold for every vector field *Y* that is C^0 -arbitrarily close to Y_2 (which guarantees that $Y_2 \in \mathscr{I}_{i,n}$). Indeed, by property (P1) and the existence of the periodic attracting orbit, $\iota_{P_{Y_2,I}}(\mathcal{B}_j \cap \Sigma) \neq 0$. Then the index $\iota_{P_{Y,I}}(\mathcal{B}_j \cap \Sigma) \neq 0$ for every *Y* that is C^0 -close to Y_2 , by property (P2). Finally, the existence of a periodic orbit in \mathcal{B}_j for impulsive semiflows $\psi_{Y,I}$, with $Y \in \mathfrak{X}^{0,1}(M)$ that is sufficiently close to Y_2 , is guaranteed by property (P3). Since $\varepsilon > 0$ was chosen arbitrarily then the claim follows.

Finally we prove that the C^0 -residual subset

$$\mathscr{R}_{I} := \bigcap_{i,n \ge 1} [\mathring{\mathscr{F}}_{i,n} \cup \mathring{\mathscr{I}}_{i,n}] \subset \mathfrak{V}_{D,\hat{D}}$$

satisfies the requirements of the proposition. Take $X \in \mathscr{R}_I$, a periodic orbit $\gamma \in Per(\psi_{X,I})$ of period a > 0 and any \mathscr{B}_i which intersects γ . As $X \in \mathring{\mathscr{F}}_{i,n} \cup \mathring{\mathscr{I}}_{i,n}$ and γ intersects \mathscr{B}_i then there exist $n \ge 1$ and \mathscr{B}_j with diam $(\mathscr{B}_j) < \text{diam}(\mathscr{B}_i)$ such that properties (2a)-(2c) hold. As the index is preserved by homotopy classes (recall property (P2)) then for any C^0 -small perturbation Y of X one has $\iota_{P_Y}(\mathscr{B}_j \cap \Sigma) \neq 0$, where P_Y stands for the Poincaré map defined in (9) for the impulsive semiflow $\psi_{Y,I}$. Finally, property (P3) implies that P_Y has a periodic orbit $\mathfrak{B}_j \cap \Sigma$, for every C^0 -small perturbation of X (hence $\psi_{Y,I}$ has a periodic orbit close to γ for every Y C^0 -close to X), which proves that γ is permanent. This completes the proof of the proposition.

The second result yields a similar consequence for impulsive semiflows parameterized by impulses.

Proposition 4.3. Fix $X \in \mathfrak{X}^{0,1}(M)$. There exists a C^0 -residual subset \mathscr{R}_X of $\mathscr{I}_{D,\hat{D}}$ such that

$$\overline{Per(\psi_{X,I})} \cap \mathring{D} = \overline{\mathscr{P}(\psi_{X,I})} \cap \mathring{D}, \quad for any \ I \in \mathscr{R}_X.$$

Proof. The argument is similar to the one of Proposition 4.2, with a decomposition formed on the space of impulses. For that reason we shall emphasize the main modifications. Fix $X \in \mathfrak{X}^{0,1}(M)$ and let $(\mathcal{B}_i)_{i\geq 1}$ be a countable open basis of the topology on \mathring{D} , which is formed by balls whose boundaries are embedded spheres and $\overline{\mathcal{B}}_i \subset \mathring{D}$ for every $i \geq 1$. For every $i, n \geq 1$, consider the pairwise disjoint subsets $\mathscr{E}_{i,n}, \mathscr{H}_{i,n}$ of $\mathscr{I}_{D,\hat{D}}$ defined as follows:

- (1) $I \in \mathscr{E}_{i,n}$ if and only if $\overline{\mathscr{B}_i} \cap \overline{\bigcup_{0 < t \leq n} \operatorname{Per}_t(\psi_{X,I})} = \emptyset;$
- (2) $I \in \mathscr{H}_{i,n}$ if and only if there exists $\overline{\mathscr{B}_j} \subset \mathscr{B}_i$, diam $(\mathscr{B}_j) < \text{diam}(\mathscr{B}_i)$ so that

- (a) $\partial \mathcal{B}_j \cap \overline{\bigcup_{0 < t \le n} \operatorname{Per}_t(\psi_{X,I})} = \emptyset;$
- (b) there exists 0 < t < n so that $\mathcal{B}_j \cap \overline{\operatorname{Per}_t(\psi_{X,I})} \neq \emptyset$;
- (c) $\iota_{f_{X,I}}(\mathcal{B}_j) \neq 0$, where $f_{X,I}$ is given by (8).

Let $\mathscr{E}_{i,n}$ and $\mathscr{H}_{i,n}$ be the C^0 -interior of the subsets $\mathscr{E}_{i,n}$ and $\mathscr{H}_{i,n}$ of $\mathscr{I}_{D,\hat{D}}$, respectively. We claim that $\mathscr{E}_{i,n} \cup \mathscr{H}_{i,n}$ is a C^0 -open and dense subset of $\mathscr{I}_{D,\hat{D}}$ for each $i, n \ge 1$. Indeed, if $I \notin \mathscr{E}_{i,n}$ then it can be C^0 -approximated by $I_0 \in \mathscr{I}_{D,\hat{D}}$ so that $\overline{\mathcal{B}}_i \cap \bigcup_{0 < t \le n} \operatorname{Per}_t(\psi_{X,I_0}) \neq \emptyset$. Performing a C^0 -arbitrary small perturbation $I_1 \in \mathscr{I}_{D,\hat{D}}$ of I_0 one guarantees that ψ_{X,I_1} exhibits an attracting periodic orbit which intersects the interior of \mathcal{B}_i (hence it is contained in \mathring{D}). Properties (P1)-(P3), applied to f_{X,I_1} ensure that $I_1 \in \mathscr{H}_{i,n}$.

Finally we are left to show that the C^0 -residual subset

$$\mathscr{R}_X := \bigcap_{i,n \ge 1} \left[\mathring{\mathscr{E}}_{i,n} \cup \mathscr{\mathscr{H}}_{i,n} \right] \subset \mathscr{I}_{D,\hat{D}}$$

fulfills the requirements of the proposition. Since this proof follows closely the argument used in the proof of the C^0 -genericity of \mathscr{R}_I (cf. proof of Proposition 4.2) we leave it as a simple exercise to the reader.

Remark 4.1. In [15] we proved that, given a smooth flow $(\varphi_t)_t$ for a C^1 -generic impulse the periodic points whose closure intersects D are all hyperbolic. This guarantees that a C^0 -dense subset of impulses have only permanent periodic orbits intersecting the closure of D. The similar argument could not be used in the proofs of Propositions 4.2 and 4.3 above, as the flow is assumed to be only Lipschitz continuous and impulses are only assumed to be homeomorphisms.

5. C^0 -generic impulses

This section is devoted to the proof of Theorems A on the properties of impulsive semiflows for C^0 typical impulses. Let φ be a Lipschitz continuous flow generated by $X \in \mathfrak{X}^{0,1}(M)$ and $D, \hat{D} \subset M$ be smooth codimension one submanifolds transversal to the flow such that hypothesis (H) holds.

We proceed to show that there exists a Baire generic subset $\mathscr{R}_X \subset \mathscr{I}_{D,\hat{D}}$ of impulses such that every $I \in \mathscr{R}_X$ satisfies

$$\overline{Per(\psi_{X,I})} \cap \mathring{D} = \Omega(\psi_{X,I}) \cap \mathring{D}$$
(10)

First, according to Proposition 4.3, there exists a C^0 -residual subset \mathscr{R}^0_X of $\mathscr{I}_{D,\hat{D}}$ such that $\overline{Per(\psi_{X,I})} \cap \mathring{D} = \overline{\mathscr{P}(\psi_{X,I})} \cap \mathring{D}$, for any $I \in \mathscr{R}^0_X$. This guarantees that the map

 $\Gamma: \mathscr{I}_{D,\hat{D}} \to 2^M$ given by $\Gamma(I) = \overline{\operatorname{Per}(\psi_{X,I})} \cap \mathring{D}$

is lower semicontinuous in \mathscr{R}^0_X . The continuity points of $\Gamma|_{\mathscr{R}^0_X}$ form a C^0 -residual subset $\mathscr{R}_X \subset \mathscr{R}^0_X$ [11].

Fix $I \in \mathscr{R}_X$. We claim that (10) holds. Assume otherwise, that there exists a point $p \in [\Omega(\psi_{X,I}) \cap \mathring{D}] \setminus [\overline{\operatorname{Per}(\psi_{X,I})} \cap \mathring{D}]$. A C^0 -small perturbation I_1 of I guarantees

that ψ_{X,I_1} admits a periodic point *x* close to $I_1(p)$ (recall Lemma 4.1). Under a small C^0 -perturbation I_2 of I_1 we can guarantee that *x* is an attractor for the Poincaré map $P_{X,I_2}: \hat{D} \to \hat{D}$ (recall (6)) hence its orbit γ is a permanent periodic orbit for ψ_{X,I_2} . By persistence of permanent periodic orbits and denseness of the C^0 -Baire generic subset \mathscr{R}_X there exists $I_3 \in \mathscr{R}_X$ that is C^0 -close to I_2 and so that ψ_{X,I_3} has a periodic orbit *q* arbitrarily close to *p*. This is a contradiction with the fact that all impulsive maps in \mathscr{R}_X are continuity points of the map $\Gamma \mid_{\mathscr{R}_2^0}$.

6. C^0 -generic Lipschitz flows

This section is devoted to the proof of Theorem C. Throughout this section let D be a smooth submanifold of codimension one and $I \in \mathscr{I}_{D,\hat{D}}$ be an impulse. The proof of Theorem C is built over the fact that there exists a C^0 -Baire generic subset $\mathscr{R}^0_I \subset \mathfrak{B}_{D,\hat{D}}$ such that

$$\operatorname{Per}(\psi_{X,I}) = \mathscr{P}(\psi_{X,I}), \text{ for any } X \in \mathscr{R}_I^0$$

which has been proved in Proposition 4.2. This guarantees that the map Γ defined by $\Gamma(X) = \overline{\operatorname{Per}(\psi_{X,I})}$ for $X \in \mathfrak{B}_{D,\hat{D}}$ is lower semicontinuous in \mathscr{R}_I^0 . In particular, the continuity points of $\Gamma|_{\mathscr{R}_I^0}$ form a residual subset $\mathscr{R}_I \subset \mathscr{R}_I^0$.

We claim that $\Omega(\psi_{X,I}) = \overline{\operatorname{Per}(\psi_{X,I})}$ for every $X \in \mathscr{R}_I$. Indeed, if this was not the case then choose $p \in \Omega(\psi_{X,I}) \setminus \overline{\operatorname{Per}(\psi_{X,I})}$. There exists a C^0 -small perturbation $Y_1 \in \mathfrak{V}_{D,\hat{D}}$ of X such that p is a periodic orbit for ψ_{Y_1} (recall Lemma 4.4). Under a small C^0 -perturbation Y_2 of Y_1 we can guarantee that p is a permanent periodic orbit for ψ_{Y_2} (recall Lemma 4.5). By persistence of permanent periodic orbits and denseness of the C^0 -Baire generic subset $X \in \mathscr{R}_I$. we can perform a C^0 -small perturbation Y_3 of Y_0 in such a way that $Y_3 \in \mathscr{R}_I$ has a periodic orbit q arbitrarily close to p. This is a contradiction with the fact that all vector fields in $X \in \mathscr{R}_I$. are continuity points of the map $\Gamma \mid_{\mathscr{R}_1^0}$. This finishes the proof.

7. FINAL COMMENTS

Volume preserving impulsive semiflows. A particular interesting framework, with many physical motivations, concerns the case of volume preserving dynamical systems. One can define a volume preserving impulsive semiflow as one generated by a volume preserving flow and an impulsive map $I : D \to \hat{D}$ such that $\text{Leb}_{\hat{D}}(\cdot) = \text{Leb}_D(I^{-1}(\cdot))$, where Leb_D and $\text{Leb}_{\hat{D}}$ denote the induced volume measures on the submanifolds D, \hat{D} of M. Assume that $(X_t)_{t\in\mathbb{R}}$ is a volume preserving Lipschitz continuous flow on a smooth compact manifold of dimension larger or equal than 3 and that D, \hat{D} are smooth cross-sections transversal to the flow. In particular, in view of [12] and the main results of this paper one expects the following questions to have an affirmative answer:

Question 1. If the flow $(X_I)_{I \in \mathbb{R}}$ is ergodic does a C^0 -generic impulse I generate an ergodic impulsive semiflow $\psi_{X,I}$? Moreover, if $I : D \to \hat{D}$ preserves volume does a C^0 -generic vector field $X \in \mathfrak{B}_{D,\hat{D}}$ generate an ergodic impulsive semiflow $\psi_{X,I}$?

We also expect that the main results of this paper (Theorems A and C) may find a parallel for C^0 -typical volume preserving impulsive semiflows.

Shadowing and gluing orbit properties. It is known that C^0 -generic homeomorphisms satisfy the shadowing property (see e.g. [1, 9]). The latter, in presence of the denseness of periodic orbits, implies on the gluing orbit property, which itself ensures a strong transitivity, denseness of ergodic measures on the simplex of invariant measures, among several strong recurrence results (cf. [2, 4]). Let us recall some ingredients. Recall that, given real numbers $\delta > 0$ and T > 1, we say that a pair of sequences $[x_i, t_i]_{i \in \mathbb{Z}}$, where $x_i \in M$, $t_i \in \mathbb{R}$, $1 \leq t_i \leq T$, is a (δ, T) -pseudo-orbit of $\psi_{X,I}$ if

$$d(\gamma_{x_i}(t_i), x_{i+1}) < \delta$$
 for all $i \in \mathbb{Z}$.

For the sequence $(t_i)_{i\in\mathbb{Z}}$ we write, $\sigma(n) = t_0 + t_1 + \ldots + t_{n-1}$ if n > 0, $\sigma(n) = -(t_n + \ldots + t_{-2} + t_{-1})$ if n < 0 and $\sigma(0) = 0$. Let $x_0 \star t$ denote a point on a (δ, T) -chain t units time from x_0 . More precisely, for $t \in \mathbb{R}$, $x_0 \star t = \gamma_{t-\sigma(i)}(x_i)$ if $\sigma(i) \leq t < \sigma(i+1)$. By Rep we denote the set of all increasing homeomorphisms $\zeta \colon \mathbb{R} \to \mathbb{R}$, called *reparametrizations*, satisfying $\zeta(0) = 0$. Fixing $\varepsilon > 0$, we define the set

$$\operatorname{Rep}(\varepsilon) = \left\{ \zeta \in \operatorname{Rep} : \left| \frac{\zeta(t) - \zeta(s)}{t - s} - 1 \right| < \varepsilon, \ s, t \in \mathbb{R} \right\},\$$

of the reparametrizations ε -close to the identity. The impulsive semiflow $\psi_{X,I}$ satisfies the *shadowing property* if, for any $\varepsilon > 0$ and T > 1 there exists $\delta = \delta(\varepsilon, T) > 0$ such that for any (δ, T) -pseudo-orbit $[x_i, t_i]_{i \in \mathbb{Z}}$ there is $\tilde{x} \in M$ and a reparametrization $\zeta \in \text{Rep}(\varepsilon)$ such that $d(\gamma_{\tilde{x}}(\zeta(t)), x_0 \star t) < \varepsilon$, for every $t \in \mathbb{R}$.

While the shadowing property turns out to be C^0 -generic in the space of continuous flows generated by Lipschitz continuous vector fields [2] this is no longer true for impulsive semiflows, as shown in the next example.

Example 7.1. Assume that the vector field $X \in \mathfrak{X}^1(\mathbb{S}^2)$ generates the usual northpole south-pole flow on the sphere \mathbb{S}^2 , that $D \subset \mathbb{S}^2$ denotes the meridian, $\hat{D} = \varphi_{-1}(D)$ and $\tau(x) = \sup\{|t|: \varphi_t(x) \notin D\}$. Then, given a small $\varepsilon > 0$, for every vector field $Y \in \mathfrak{X}^{0,1}(\mathbb{S}^2)$ that is C^0 -close to X, any impulse $I \in \mathscr{I}_{D,\hat{D}}$ and $\delta > 0$, it is simple to check that the following δ -pseudo-orbit $[x_i, t_i]_{i=1,2}$ is not ε -shadowed: it is enough to take any regular point x_1 in the interior of the upper hemisphere, $t_1 = \tau(x_1) - \frac{\delta}{10\|X\|_{\infty}}$, $x_2 = \varphi_{\tau(x_1) + \frac{\delta}{10\|X\|_{\infty}}}(x_1)$ and $t_2 = 1$. The absence of shadowing holds by C^0 -perturbations of the Lipschitz vector field and impulse I.

The previous simple example clarifies that, in order for some shadowing property to hold for typical impulsive semiflows, it may be necessary to consider special classes of pseudo-orbits. This justifies the following:

Question 2. Does a typical impulsive semiflow (parameterized by either the space of impulses or vector fields) satisfy a shadowing property for some relevant class of pseudo-orbits?

Acknowledgements. Part of this work was done during the visit of the third named author to UFRJ and PUC-Rio, whose research environment and research conditions are acknowledged. JS was funded by the grant E-26/010/002610/2019, Rio de Janeiro Research Foundation (FAPERJ), and by the Coordenação de Aperfeiçoamento de Pessoal de Nível Superior - Brasil (CAPES), Finance Code 001. MJT was partially financed by Portuguese Funds through FCT (Fundação para a Ciência e a Tecnologia) through the research Projects UIDB/00013/2020 (DOI: 10.54499/UIDB/00013/2020) and UIDP/00013/2020 (DOI: 10.54499/UIDP/00013/2020). PV was funded by Fundação para a Ciência e Tecnologia (FCT) - Portugal, through the grant CEECIND/03721/2017 of the Stimulus of Scientific Employment, Individual Support 2017 Call and CMUP (UID/MAT/00144/2013).

References

- E. Akin, M. Hurley and J. Kennedy, *Dynamics of topologically generic homeomorphisms*, Memoirs of the Amer. Math. Soc., volume 164, 2003. 1, 7
- [2] M. Bessa, M. J. Torres and P. Varandas, On the periodic orbits, shadowing and strong transitivity of continuous flows, Nonlinear Analysis, 175 (2018) 191–209. 4.3, 7
- [3] A. Boldt and C. Calcaterra, *Lipschitz Flow-box Theorem*, J. Math. Anal. Appl. 338 (2008) 1108–1115. 4.1, 4.2
- [4] T. Bomfim, M. J. Torres and P. Varandas. *Topological features of flows with the reparametrized gluing orbit property*. J. Diff. Eq., 262 (8) (2017), 4292–4313. 7
- [5] E. Bonotto, Flows of characteristic 0+ in impulsive semidynamical system J. Math. Anal. Appl., 332:1 (2007) 81–96. 2.1
- [6] E. Bonotto, M. Bortolan, T. Caraballo and R. Collegari, A survey on impulsive dynamical systems, Electron. J. Qual. Theory Differ. Equ., (2016), 1–27. 1
- [7] K. Ciesielski, On semicontinuity in impulsive dynamical systems, Bull. Pol. Acad. Sci. Math. 52 (2004), 71–80. 2.1
- [8] E. M. Coven, J. Madden and Z. Nitecki, A note on generic properties of continuous maps, Ergodic theory and dynamical systems, II, 97–101, Progr. Math., 21, Birkhäuser, Boston, Mass., 1982. 1, 4.3, 4.3, 4.3
- [9] P. Koscielniak, M. Mazur, P. Oprocha and P. Pilarczyk, Shadowing is generic-a continuous map case, Discrete Contin. Dyn. Syst 34:9 (2014) 3591–3609. 7
- [10] M. Hurley, On proofs of the C^0 general density theorem, Proc. Amer. Math. Soc., 124 (4), (1996), 1305–1309. 1, 4.3
- [11] K. Kuratowski, Topology, vol. 1, Academic Press, 1966. 5
- [12] J. Oxtoby and S. Ulam, Measure-preserving homeomorphisms and metrical transitivity, Ann. of Math., 42 (2) (1941), 874–920. 7
- [13] J. Palis, C. Pugh, M. Shub and D. Sullivan, *Genericity theorems in topological dynamics*, Lecture Notes in Math., 468, 241–250, Springer, Berlin, 1975. 1
- [14] C. Pugh, An improved closing lemma and a general density theorem, Amer. J. Math., 89:4 (1967) 1010–1021. 1
- [15] J. Siqueira, M. J. Torres and P. Varandas, Abundance of periodic orbits for typical impulsive flows, Preprint 2024. 1, 3.1, 3.3, 4.1

JAQUELINE SIQUEIRA, DEPARTAMENTO DE MATEMÁTICA, INSTITUTO DE MATEMÁTICA, UNIVERSIDADE FED-ERAL DO RIO DE JANEIRO, CAIXA POSTAL 68530, RIO DE JANEIRO, RJ 21945-970, BRAZIL *Email address*: jaqueline@im.ufrj.br

CMAT AND DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE DO MINHO, CAMPUS DE GUALTAR, 4700-057 BRAGA, PORTUGAL

Email address: jtorres@math.uminho.pt

DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DA BAHIA, AV. ADEMAR DE BARROS S/N, 40170-110 SALVADOR, BRAZIL & CMUP, UNIVERSITY OF PORTO -PORTUGAL

Email address: paulo.varandas@ufba.br