

# Breather gas and shielding for the focusing nonlinear Schrödinger equation with nonzero backgrounds

Weifang Weng<sup>1</sup>, Guoqiang Zhang<sup>2,3</sup>, Boris A. Malomed<sup>4</sup>, Zhenya Yan<sup>2,3,\*</sup>

<sup>1</sup>*School of Mathematical Sciences, University of Electronic Science and Technology of China, Chengdu, Sichuan, 611731, China*

<sup>2</sup>*KLMM, Academy of Mathematics and Systems Science, Chinese Academy of Sciences, Beijing 100190, China*

<sup>3</sup>*School of Mathematical Sciences, University of Chinese Academy of Sciences, Beijing 100049, China*

<sup>4</sup>*Department of Physical Electronics, School of Electrical Engineering, Faculty of Engineering, Tel Aviv University, Tel Aviv 69978, Israel*

**Abstract:** Breathers have been experimentally and theoretically found in many physical systems – in particular, in integrable nonlinear-wave models. A relevant problem is to study the *breather gas*, which is the limit, for  $N \rightarrow \infty$ , of  $N$ -breather solutions. In this paper, we investigate the breather gas in the framework of the focusing nonlinear Schrödinger (NLS) equation with nonzero boundary conditions, using the inverse scattering transform and Riemann-Hilbert problem. We address aggregate states in the form of  $N$ -breather solutions, when the respective discrete spectra are concentrated in specific domains. We show that the breather gas coagulates into a single-breather solution whose spectral eigenvalue is located at the center of the circle domain, and a multi-breather solution for the higher-degree quadrature concentration domain. These coagulation phenomena in the breather gas are called *breather shielding*. In particular, when the nonzero boundary conditions vanish, the breather gas reduces to an  $n$ -soliton solution. When the discrete eigenvalues are concentrated on a line, we derive the corresponding Riemann-Hilbert problem. When the discrete spectrum is uniformly distributed within an ellipse, it is equivalent to the case of the line domain. These results may be useful to design experiments with breathers in physical settings.

## 1 Introduction

In 1834, solitary waves were discovered by Russell [1], and Korteweg and de Veris established the KdV equation to describe this wave phenomenon in 1895 [2]. However, these significant results did not receive enough attention at that time. Until 1955, Fermi, Pasta and Ulam [3] numerically investigated the thermalization process of a solid, which was called the Fermi-Pasta-Ulam (FPU) problem, and broke new branches of nonlinear science (e.g., solitons and chaos), and numerically simulating scientific problems [4]. In 1965, Zabusky and Kruskal, motivated by the Fermi-Pasta-Ulam-Tsingou (FPUT) problem [3], coined the concept of ‘*solitons*’, as elastically interacting solitary-waves solutions of the KdV equation (continuum limit of FPUT problem) with periodic initial data [5]. In 1967, Gardner *et al* [6] discovered the inverse scattering transform (IST) to produce exact  $\mathcal{N}$ -soliton solutions of the KdV, starting from its spectral problem (alias the Lax pair [7]), as elaborated in detail in the classical work of Ablowitz *et al* [8]. Parallel to that, the integrability of the nonlinear Schrödinger (NLS) equations and solitons produced by them were discovered by Zakharov *et al* in 1971 [9]. The discovery was an incentive to predict NLS solitons in nonlinear optical fibers by Hasegawa *et al* [10, 11]. The predicted fundamental bright solitons and breathers (periodically oscillating  $\mathcal{N}$ th-order solitons, which are also exact solutions of the NLS equation with the self-focusing nonlinearity [12]) were created in optical fibers, for  $\mathcal{N} = 1, 2$  and 3 by Mollenauer *et al* [13]. The  $\mathcal{N}$ th-order breather may be considered as a bound state of  $\mathcal{N}$  fundamental solitons with unequal amplitudes, the ratios between which are  $1 : 3 : 5 : \dots : 2\mathcal{N} - 1$ . These bound states are fragile ones, in the sense that their binding energy is zero [12]. Nevertheless, they can be readily stabilized in fiber lasers [14–17], where breathers are basic operation modes [18].

Another fundamentally important realization of the NLS equation (alias the Gross-Pitaevskii equation) is provided by quasi-one-dimensional Bose-Einstein condensates (BECs) with attractive inter-atomic inter-

---

\*Email address: zyyan@mmrc.iss.ac.cn (Corresponding author)

actions [19]. Fundamental solitons in BECs were first observed in 2002 in condensates of  $^7\text{Li}$  atoms [20,21]. Breathers of orders 2 [22,23] and 3 [23] have been experimentally demonstrated more recently. Moreover, solitons also appear in many fields of nonlinear science [24,25].

The interaction among multi-soliton solutions is a topic of fundamental significance in the theory of nonlinear waves [14–17,19,24–36]. In 1971, Zakharov first proposed the concept of *soliton gas*, defined as the large- $N$  limit of the  $N$ -soliton solution of the KdV [37]. It is relevant to stress that, although collisions between solitons governed by integrable nonlinear wave equations are elastic, collisional effects in the soliton gas are not trivial, as the elastic collisions give rise to phase shifts of solitons [6,8,9,27] (an exception is the 2D KP-I equation, where collisions between weakly localized lump solitons yield zero phase shifts [38]). Afterwards, the concept was extended to investigate the fluid dynamics of soliton gases, breather gases, and dense soliton gases for other nonlinear wave equations [39–49]. Especially, El *et al* elaborated the spectral theory [50] and numerical experiment [51] of soliton and breather gases for the NLS equation. Suret *et al* developed the nonlinear spectral synthesis of the soliton gas in deep-water surface gravity waves [52]. There were some related soliton gas experiments in optics [53–55] and shallow water regime [56]. In particular, the concept of soliton and breather gases is relevant for the implementation in fiber lasers, where it is possible to create chains composed of large numbers of solitons and breathers [49]. Recently, Girotti *et al* first presented the soliton gas of the KdV and modified KdV equations, respectively, starting from  $N$ -soliton solutions via Riemann-Hilbert (RH) problems [57,58]. Bertola *et al* further proposed the effect of *soliton shielding*, alias “soliton coagulation”, in dense soliton gases governed by the NLS with zero backgrounds [59]. The effect implies that the field generated by a superposition of a large set of specially placed solitons may become tantamount to a few-soliton configuration. However, the “coagulation” was not studied for large sets of NLS breathers, rather than fundamental solitons via RH problems. Compared to zero boundary condition of NLS equation, the discrete spectrum with nonzero boundary condition exhibit more symmetry. It is worth studying whether there is a phenomenon of breather-shielding effect in this case.

In this paper, motivated by Ref. [59] for the soliton gas of the NLS equation with zero backgrounds, we would like to analyze the breather-shielding effect and breather gas (the  $N \rightarrow \infty$  limit of the  $N$ -breather solutions) of the focusing NLS equation with nonzero boundary conditions (BCs) of the Dirichlet type at infinity [9]:

$$\begin{cases} iq_t + q_{xx} + 2(|q|^2 - q_0^2)q = 0, & (x, t) \in \mathbb{R}^2, \\ \lim_{x \rightarrow \pm\infty} q(x, t) = q_{\pm} = \text{const}, & |q_{\pm}| = q_0 > 0. \end{cases} \quad (1)$$

Our starting point is the Zakharov-Shabat (ZS) scattering problem (i.e., the Lax pair) [9,26,60]

$$\begin{cases} \Psi_x = U\Psi, & U = \frac{i}{2} \left( k - \frac{q_0^2}{k} \right) \sigma_3 + Q, \\ \Psi_t = W\Psi, & W = -\frac{i}{2} \left( k^2 + \frac{q_0^4}{k^2} \right) \sigma_3 + i\sigma_3 (Q_x - Q^2), \end{cases} \quad (2)$$

where  $\Psi = \Psi(x, t; k)$  is a second-order matrix-valued Jost function,  $k$  is a complex spectral parameter, and the potential function matrix and  $\sigma_3$  are given by

$$Q = \begin{bmatrix} 0 & q(x, t) \\ -q^*(x, t) & 0 \end{bmatrix}, \quad \sigma_3 = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix}, \quad (3)$$

with  $*$  denoting the complex conjugate. Notice that Eq. (1) is the compatibility condition (or zero-curvature equation)  $U_t - W_x + [U, W] = 0$  of the Lax pair (2).

Different types (Kuznetsov-Ma-type [61,62], Akhmediev-type [63]) of breathers were found for the NLS equation. Moreover, their parameter limits could generate its new non-periodic rogue wave (RW) [64]. Recently, a powerful approach was proposed for applying IST and obtaining exact solutions to the focusing and defocusing NLS equations with the nonzero BCs, in terms of RH problems and their extensions, including the discrete case, multi-component systems, and nonlocal equations [65–70]. Deift *et al* proposed the

steepest-descent approximation for RH problems to explore the long-time asymptotics of the modified KdV equation [71]. Techniques based on IST and RH problems were developed in other directions [72–81]. In particular, Bilman *et al* combined the robust IST and Darboux transform to obtain RW solutions of the NLS equation [82]. The asymptotics of multi-soliton solutions to the NLS equation was addressed [83,84]. Later, a scale transform and RH technique were applied to an  $N$ -RW solution of the NLS equation to analyze its near- and far-field asymptotic behaviors [85]. Recently, Romero-Ros *et al* experimentally demonstrated the RW dynamics in a 3D coupled BECs [86].

The rest of this paper is arranged as follows. In Sec. 2, we simply recall the direct and inverse scattering transforms and the corresponding RH problem of the NLS equation with nonzero backgrounds. In Sec. 3, we analyze the breather gas, which is the limit of the  $N$ -breather solution at  $N \rightarrow \infty$ , via the modified RH problem. Moreover, we address aggregate states in the form of  $N$ -breather solutions, when the respective discrete spectra are concentrated in specific domains. We show that the breather gas coagulates into a single-breather solution whose spectral eigenvalue is located at the center of the domain for the circle domain, and a multi-breather solution for the higher-degree quadrature concentration domain. These coagulation phenomena in the breather gas are called *breather shielding*. In particular, when the nonzero boundary conditions vanish, the breather gas reduces to an  $n$ -soliton solution. When the discrete eigenvalues are concentrated on a line, we derive the corresponding Riemann-Hilbert problem. When the discrete spectrum is uniformly distributed within an ellipse, it is equivalent to the case of the line domain. Finally, we present some conclusions and discussions in Sec. 4.

## 2 Preliminaries

In this section, we recall the basic properties about the direct and inverse scattering problems and RH problem of the NLS equation with nonzero BCs given by Eq. (1) (see [60] for the details). As  $x \rightarrow \pm\infty$ , the ZS scattering problem (or Lax pair) (2) becomes the asymptotic form

$$\begin{cases} \Psi_x^{bg} = U^{bg}\Psi^{bg}, & U^{bg} = \frac{i}{2} \left( k - \frac{q_0^2}{k} \right) \sigma_3 + Q_{\pm}, \\ \Psi_t^{bg} = W^{bg}\Psi^{bg}, & W^{bg} = -\frac{i}{2} \left( k^2 + \frac{q_0^4}{k^2} \right) \sigma_3 - iq_0^2 \sigma_3, \end{cases} \quad (4)$$

which admits the solution

$$\Psi_{\pm}^{bg}(x, t; k) = \begin{cases} P_{\pm}(k) e^{i\vartheta(x, t; k)\sigma_3}, & k \neq 0, \pm iq_0, \\ \mathbb{I}_2 + (x - 2kt)(ik\sigma_3 + Q_{\pm}), & k = \pm iq_0, \end{cases} \quad (5)$$

where  $Q_{\pm} = \lim_{x \rightarrow \pm\infty} Q(x, t) = Q(x, t)|_{q=q_{\pm}}$  and

$$P_{\pm}(k) = \mathbb{I}_2 + \frac{i}{k}\sigma_3 Q_{\pm}, \quad \vartheta(x, t; k) = \frac{1}{2} \left( k + \frac{q_0^2}{k} \right) \left[ x - \left( k - \frac{q_0^2}{k} \right) t \right], \quad (6)$$

with  $\mathbb{I}_2$  is a  $2 \times 2$  unit matrix.

Let  $\Sigma = \mathbb{R} \cup C_0$ ,  $\widehat{\Sigma} = \mathbb{R} \setminus \{0\} \cup C_0$ ,  $\Sigma_0 := \Sigma \setminus \{\pm iq_0\}$  with  $C_0 = \{k \in \mathbb{C} : |k| = q_0\}$ . The continuous spectrum of  $X_{\pm} = \lim_{x \rightarrow \pm\infty} X$  is the set of all values of  $k$  satisfying  $k + q_0^2/k \in \mathbb{R}$ , i.e.,  $k \in \Sigma = \mathbb{R} \cup C_0$ , which are the jump contours for the related Riemann-Hilbert problem (see the inverse scattering problem). Let  $\mathbb{D}^+ \equiv \{k | \text{Im}(k)(1 - q_0^2/|k|^2) > 0\}$ ,  $\mathbb{D}^- \equiv \{k | \text{Im}(k)(1 - q_0^2/|k|^2) < 0\}$ . Thus, one can simultaneously determine the Jost and modified Jost solutions  $\Psi_{\pm}(x, t; k)$  and  $\mu_{\pm}(x, t; k)$  of the Lax pair (2) satisfying the boundary conditions

$$\begin{aligned} \Psi_{\pm}(x, t; k) &= P_{\pm}(k) e^{i\vartheta(x, t; k)\sigma_3} + o(1), \quad x \rightarrow \pm\infty, \\ \mu_{\pm}(x, t; k) &= \Psi_{\pm}(x, t; k) e^{-i\vartheta(x, t; k)\sigma_3} \rightarrow P_{\pm}(k), \quad x \rightarrow \pm\infty, \end{aligned} \quad (7)$$

where

$$\mu_{\pm}(k) = \begin{cases} P_{\pm}(k) \left\{ \mathbb{I}_2 + \int_{\pm\infty}^x \exp\left(\frac{i}{2}\left(k + \frac{q_0^2}{k}\right)(x - \xi)\widehat{\sigma}_3\right) [P_{\pm}^{-1}(k)[Q(\xi, t) - Q_{\pm}] \mu_{\pm}(\xi, t; k)] d\xi \right\}, \\ k \neq 0, \pm iq_0, q - q_{\pm} \in L^1(\mathbb{R}^{\pm}), \\ P_{\pm}(k) + \int_{\pm\infty}^x [I + (x - \xi)(Q_{\pm} \mp q_0 \sigma_3)] [Q(\xi, t) - Q_{\pm}] \mu_{\pm}(\xi, t; k) d\xi, \\ k = \pm iq_0, (1 + |x|)(q - q_{\pm}) \in L^1(\mathbb{R}^{\pm}) \end{cases} \quad (8)$$

with  $e^{\alpha\widehat{\sigma}_3} A := e^{\alpha\sigma_3} A e^{-\alpha\sigma_3}$ .

Let  $\Psi_{\pm}(x, t; k) = (\Psi_{\pm 1}(x, t; k), \Psi_{\pm 2}(x, t; k))$  and  $\mu_{\pm}(x, t; k) = (\mu_{\pm 1}(x, t; k), \mu_{\pm 2}(x, t; k))$ . Then for the given  $q - q_{\pm} \in L^1(\mathbb{R}^{\pm})$ , the Jost functions  $\Psi_{\pm 2}(x, t; k)$  and modified forms  $\mu_{\pm 2}(x, t; k)$  given by Eqs. (7) and (8) both possess the unique solutions in  $\Sigma_0$ . Moreover,  $\mu_{+1, -2}(x, t; k)$  and  $\Psi_{+1, -2}(x, t; k)$  ( $\mu_{-1, +2}(x, t; k)$  and  $\Psi_{-1, +2}(x, t; k)$ ) can be extended analytically to  $\mathbb{D}^+$  ( $\mathbb{D}^-$ ), and continuously to  $\mathbb{D}^+ \cup \Sigma_0$  ( $\mathbb{D}^- \cup \Sigma_0$ ). Since  $\Psi_{\pm}(x, t; k)$ ,  $k \neq 0, \pm iq_0$  are both fundamental matrix solutions of the Lax pair (2), thus there exists a constant scattering matrix  $S(k)$  between them

$$\Psi_+(x, t; k) = \Psi_-(x, t; k) S(k), \quad \mu_+(x, t; k) = \mu_-(x, t; k) e^{i\theta\widehat{\sigma}_3} S(k), \quad k \in \Sigma_0, \quad (9)$$

where  $S(k) = (s_{ij}(k))_{2 \times 2}$  with the scattering coefficients  $s_{ij}(k)$ 's with  $s_{11}(k)$  and  $s_{22}(k)$  in  $k \in \Sigma_0$  being extended analytically to  $\mathbb{D}^+$  and  $\mathbb{D}^-$ , and continuously to  $\mathbb{D}^+ \cup \Sigma_0$  and  $\mathbb{D}^- \cup \Sigma_0$ , respectively, and  $s_{12}(k)$  and  $s_{21}(k)$  not being analytically continued away from  $\Sigma_0$ . The reflection coefficients are defined as  $\rho(k) = \frac{s_{21}(k)}{s_{11}(k)}$ ,  $\widehat{\rho}(k) = \frac{s_{12}(k)}{s_{22}(k)}$ ,  $k \in \Sigma_0$ . The Jost solutions  $\Phi(x, t; z)$ ,  $\mu_{\pm}(x, t; z)$ , the scattering matrix  $S(z)$ , and reflection coefficients admit the following symmetries

$$\Phi_{\pm}(x, t; k) = \sigma_2 \Phi_{\pm}^*(x, t; k^*) \sigma_2 = \frac{i}{k} \Phi_{\pm} \left( x, t; -\frac{q_0^2}{k} \right) \sigma_3 Q_{\pm}, \quad \Phi_{\pm} = \Psi_{\pm}, \mu_{\pm}, \quad \sigma_2 = \text{antidiag}(-i, i), \quad (10)$$

$$S(k) = \sigma_2 S^*(k^*) \sigma_2 = (\sigma_3 Q_-)^{-1} S \left( -\frac{q_0^2}{k} \right) \sigma_3 Q_+, \quad \rho(k) = -\widehat{\rho}^*(k^*) = \frac{q_-^*}{q_-} \widehat{\rho} \left( -\frac{q_0^2}{k} \right).$$

Moreover, the asymptotic behaviors for modified Jost solutions and scattering matrix are found by

$$\begin{cases} \mu_{\pm}(x, t; k) = \mathbb{I}_2 + O\left(\frac{1}{k}\right), & S(k) = \mathbb{I}_2 + O\left(\frac{1}{k}\right), & k \rightarrow \infty, \\ \mu_{\pm}(x, t; k) = \frac{i}{k} \sigma_3 Q_{\pm} + O(1), & S(k) = \text{diag}\left(\frac{q_-}{q_+}, \frac{q_+}{q_-}\right) + O(k), & k \rightarrow 0. \end{cases} \quad (11)$$

The discrete spectrum of the focusing NLS equation with nonzero BCs (1) is the set

$$K = K^+ \cup K^-, \quad K^+ = \left\{ k_j, -q_0^2/k_j^* \right\}_{j=1}^N \subset \mathbb{D}^+, \quad K^- = \left\{ k_j^*, -q_0^2/k_j \right\}_{j=1}^N \subset \mathbb{D}^-, \quad (12)$$

where  $s_{11}(k_j) = 0$ ,  $s'_{11}(k_j) \neq 0$ .

We construct a piecewise meromorphic function:

$$M(x, t; k) = \begin{cases} \left( \frac{\mu_{+1}(x, t; k)}{s_{11}(k)}, \mu_{-2}(x, t; k) \right) = \left( \frac{\Psi_{+1}(x, t; k)}{s_{11}(k)}, \Psi_{-2}(x, t; k) \right) e^{-i\theta(x, t; k)\sigma_3}, & k \in \mathbb{D}^+, \\ \left( \mu_{-1}(x, t; k), \frac{\mu_{+2}(x, t; k)}{s_{22}(k)} \right) = \left( \Psi_{-1}(x, t; k), \frac{\Psi_{+2}(x, t; k)}{s_{22}(k)} \right) e^{-i\theta(x, t; k)\sigma_3}, & k \in \mathbb{D}^-. \end{cases} \quad (13)$$

Then, the matrix function  $M(x, t; k)$  satisfies the following RH problem:

**Riemann-Hilbert problem 1** Find a  $2 \times 2$  matrix  $M(x, t; k)$  that satisfies the following conditions:

- *Analyticity:*  $M(x, t; k)$  is meromorphic in  $\{k | k \in \mathbb{C} \setminus (\Sigma \cup K)\}$  and takes continuous boundary values on  $\Sigma$ ;
- *The jump condition:* the boundary values on the jump contour  $\Sigma$  are defined as

$$M_+(k) = M_-(k)J(k), \quad J(k) = e^{i\theta(x,t;k)\hat{\sigma}_3} \begin{pmatrix} 1 - \rho(k)\hat{\rho}(k) & -\hat{\rho}(k) \\ \rho(k) & 1 \end{pmatrix}, \quad k \in \Sigma, \quad (14)$$

- *Normalization:*

$$M(x, t; k) = \begin{cases} \mathbb{I}_2 + O(1/k) & k \rightarrow \infty, \\ \frac{i}{k} \sigma_3 Q_- + O(1), & k \rightarrow 0. \end{cases} \quad (15)$$

- *The residue conditions:*  $M(x, t; k)$  has simple poles at each point in  $K := \{k_j, -\frac{q_0^2}{k_j^*}, k_j^*, -\frac{q_0^2}{k_j}\}_{j=1}^N$  with

$$\begin{aligned} \operatorname{Res}_{k=k_j} M(x, t; k) &= \lim_{k \rightarrow k_j} M(x, t; k) \begin{bmatrix} 0 & 0 \\ c_j e^{-2i\theta(k_j)} & 0 \end{bmatrix}, \\ \operatorname{Res}_{k=-\frac{q_0^2}{k_j^*}} M(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{k_j^*}} M(x, t; k) \begin{bmatrix} 0 & 0 \\ -\frac{q_0^2 q_-^*}{k_j^{*2} q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})} & 0 \end{bmatrix}, \\ \operatorname{Res}_{k=k_j^*} M(x, t; k) &= \lim_{k \rightarrow k_j^*} M(x, t; k) \begin{bmatrix} 0 & -c_j^* e^{2i\theta(k_j^*)} \\ 0 & 0 \end{bmatrix}, \\ \operatorname{Res}_{k=-\frac{q_0^2}{k_j}} M(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{k_j}} M(x, t; k) \begin{bmatrix} 0 & \frac{q_0^2 q_-}{k_j^2 q_-} c_j e^{2i\theta(-\frac{q_0^2}{k_j})} \\ 0 & 0 \end{bmatrix}, \end{aligned} \quad (16)$$

with  $c_j$ 's being complex constants.

Then for the reflectionless case  $\rho(k) = 0$ , the  $N$ -breather solution  $q(x, t)$  of the focusing NLS equation with nonzero BCs is given by

$$q(x, t) = -i \lim_{k \rightarrow \infty} (kM(x, t; k))_{12}, \quad (17)$$

where  $M(x, t; k)$  is determined by Eqs. (16) and (15) as

$$M(x, t; k) = \mathbb{I} + \frac{i}{k} \sigma_3 Q_- + \sum_{j=1}^{2N} \left( \frac{\begin{bmatrix} u_j(x, t; \eta) & 0 \\ v_j(x, t; \eta) & 0 \end{bmatrix}}{k - \eta_j} + \frac{\begin{bmatrix} 0 & \hat{v}_j(x, t; \hat{\eta}) \\ 0 & \hat{u}_j(x, t; \hat{\eta}) \end{bmatrix}}{k - \hat{\eta}_j} \right), \quad (18)$$

with  $u_j = u_j(x, t; \eta)$ ,  $v_j = v_j(x, t; \eta)$ ,  $\hat{u}_j = \hat{u}_j(x, t; \hat{\eta})$ ,  $\hat{v}_j = \hat{v}_j(x, t; \hat{\eta})$  that can be found from Eq. (16),  $Q_- = \operatorname{antidiag}(q_-, -q_-^*)$ , and  $\eta_j = k_j$ ,  $\eta_{N+j} = -q_0^2/k_j^*$ ,  $\hat{\eta}_j = -q_0^2/\eta_j$ ,  $\hat{\eta}_{N+j} = -q_0^2/\eta_{N+j}$ , ( $j = 1, 2, \dots, N$ ). Note that the limiting  $M(k)$  satisfies, in general, a  $\bar{\partial}$  problem.

### 3 Breather gas: the limit of the $N$ -breather solution at $N \rightarrow \infty$

Below, we consider the ZS spectral problem for a reflectionless potential ( $\rho(k) = 0$ ,  $k \in \Sigma$ ) with simple poles, which corresponds to focusing NLS equation with nonzero BC. We define a closed curve  $\Gamma_{1+}(\Gamma_{2+})$  with a very small radius encircling poles  $\{k_j\}_{j=1}^N (\{-\frac{q_0^2}{k_j^*}\}_{j=1}^N)$  counterclockwise in the upper half plane  $\mathbb{D}_+$ , and a closed curve  $\Gamma_{1-}(\Gamma_{2-})$  with a very small radius encircling the poles  $\{k_j^*\}_{j=1}^N (\{-\frac{q_0^2}{k_j}\}_{j=1}^N)$  clockwise in the lower half plane  $\mathbb{D}_-$ .

Based on the RH problem 1, we consider following transformation:

$$M_1(x, t; k) = \begin{cases} M(x, t; k) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1+}, \\ M(x, t; k) \begin{bmatrix} 1 & 0 \\ \sum_{j=1}^N \frac{\frac{q_0^2 q_-^*}{k_j^{*2} q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{k + \frac{q_0^2}{k_j^*}} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2+}, \\ M(x, t; k) \begin{bmatrix} 1 & \sum_{j=1}^N \frac{c_j^* e^{2i\theta(k_j^*)}}{k - k_j^*} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1-}, \\ M(x, t; k) \begin{bmatrix} 1 & -\sum_{j=1}^N \frac{\frac{q_0^2 q_-}{k_j^2 q_-^*} c_j e^{2i\theta(-\frac{q_0^2}{k_j})}}{k + \frac{q_0^2}{k_j}} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2-}, \\ M(x, t; k), & \text{otherwise.} \end{cases} \quad (19)$$

Therefore, according to RH problem 1, we know that the matrix function  $M_1(x, t; k)$  satisfies the following RH problem:

**Riemann-Hilbert problem 2** Find a  $2 \times 2$  matrix function  $M_1(x, t; k)$  that meets the following conditions:

- *Analyticity:*  $M_1(x, t; k)$  is analytic in  $\mathbb{C} \setminus (\Gamma_{1\pm} \cup \Gamma_{2\pm})$  and takes continuous boundary values on  $\Gamma_{1\pm} \cup \Gamma_{2\pm}$ .
- *The jump condition:* The boundary values on the jump contour  $\Gamma_{1+} \cup \Gamma_{1-}$  are defined as

$$M_{1+}(x, t; k) = M_{1-}(x, t; k) V_1(x, t; k), \quad \lambda \in \Gamma_{1\pm} \cup \Gamma_{2\pm}, \quad (20)$$

where

$$V_1(x, t; k) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ \sum_{j=1}^N \frac{\frac{q_0^2 q_-^*}{k_j^{*2} q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{k + \frac{q_0^2}{k_j^*}} & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & -\sum_{j=1}^N \frac{c_j^* e^{2i\theta(k_j^*)}}{k - k_j^*} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & -\sum_{j=1}^N \frac{\frac{q_0^2 q_-}{k_j^2 q_-^*} c_j e^{2i\theta(-\frac{q_0^2}{k_j})}}{k + \frac{q_0^2}{k_j}} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (21)$$

- *Normalization:*

$$M_1(x, t; k) = \begin{cases} \mathbb{I}_2 + O(1/k) & k \rightarrow \infty, \\ \frac{i}{k} \sigma_3 Q_- + O(1), & k \rightarrow 0. \end{cases} \quad (22)$$

According to Eq. (17), we recover  $q(x, t)$  by means of the following formula:

$$q(x, t) = -i \lim_{k \rightarrow \infty} (kM_1(x, t; k))_{12}. \quad (23)$$

**Proof** Without loss of generality, it suffices to consider the case of  $k \in \Gamma_{1+}$ :

$$\begin{aligned} M_{1+}(x, t; k) &= M(x, t; k) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{-2i\vartheta(k_j)}}{k - k_j} & 1 \end{bmatrix} \\ &= M_{1-}(x, t; k) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{-2i\vartheta(k_j)}}{k - k_j} & 1 \end{bmatrix}. \end{aligned} \quad (24)$$

Then we have

$$V_1(x, t; k) = \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^N \frac{c_j e^{-2i\vartheta(k_j)}}{k - k_j} & 1 \end{bmatrix}, \quad k \in \Gamma_{1+}. \quad (25)$$

### 3.1 Quadrature domains

In this subsection, we address the limit of  $N \rightarrow \infty$ , under the additional assumptions:

- The discrete spectra  $k_j, j = 1, \dots, N$  with the norming constants  $c_j, j = 1, \dots, N$  fill uniformly compact domain  $\Omega_1$  which is strictly contained in the domain  $D_{\Gamma_1}$  bounded by  $\Gamma_1$ , that is,

$$\Omega_1 := \{k \mid |(k - s_1)^m - s_2| < s_3\}, \quad \Omega_1 \subset D_+, \quad (26)$$

where  $m \in \mathbb{N}_+, s_1 \in \mathbb{C}_+$  and  $|s_2|, s_3$  are sufficiently small (see Figure 1). Note that Bertola *et al* [59] has discussed the soliton shielding of the NLS equation with zero BC in the domain  $\Omega_1$ .

- The normalization constants  $c_j, j = 1, \dots, N$  have the following form:

$$c_j = \frac{|\Omega_1| r(k_j, k_j^*)}{N\pi}. \quad (27)$$

where  $|\Omega_1|$  means the area of the domain  $\Omega_1$  and  $r(k, k^*) := n(k^* - s_1^*)^{n-1} r_1(k)$  is a smooth function of variables  $k$  and  $k^*$  and function  $r_1(k)$  is subject to the symmetry relation  $r_1^*(k) = r_1(k^*)$ .




Figure 1: (a) The distribution of discrete spectrum  $K$ , the parameters being  $s_1 = 2i, s_2 = \frac{1}{4}, s_3 = \frac{1}{2}, m = 1, q_0 = 1$ . (b) The 1-breather solution with the same parameters and  $r(k) = 1$ .

**Proposition 1** For any open set  $B_+$  containing the domain  $\Omega_1$ , the following identities hold:

$$\begin{aligned}
\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} &= \iint_{\Omega_1} \frac{r(\lambda, \lambda^*) e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda^* \wedge d\lambda, \\
\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\frac{q_0^2 q_-^*}{k_j^{*2} q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{k + \frac{q_0^2}{k_j^*}} &= \iint_{\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} r^*(\lambda, \lambda^*) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* \wedge d\lambda, \\
\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j^* e^{2i\theta(k_j^*)}}{k - k_j^*} &= \iint_{\Omega_1} \frac{r^*(\lambda, \lambda^*) e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^* \wedge d\lambda, \\
\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\frac{q_0^2 q_-}{k_j^2 q_-^*} c_j e^{2i\theta(-\frac{q_0^2}{k_j})}}{k + \frac{q_0^2}{k_j}} &= \iint_{\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_-^*} r(\lambda, \lambda^*) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda^* \wedge d\lambda.
\end{aligned} \tag{28}$$

uniformly for all  $\mathbb{C} \setminus B_+$ . The boundary  $\partial\Omega_1$  has the counterclockwise orientation.

**Proof** Using Eq. (27), we have

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} = \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{|\Omega_1|}{N} \frac{r(k_j, k_j^*) e^{-2i\theta(k_j)}}{\pi(k - k_j)} = \iint_{\Omega_1} \frac{r(\lambda, \lambda^*) e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda^* \wedge d\lambda. \tag{29}$$

Thus the proof is completed.

**Proposition 2** The following identities hold:

$$\begin{aligned}
\iint_{\Omega_1} \frac{r(\lambda, \lambda^*) e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda^* \wedge d\lambda &= \int_{\partial\Omega_1} \frac{(\lambda^* - s_1^n) r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda, \\
\iint_{\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} r^*(\lambda, \lambda^*) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* \wedge d\lambda &= - \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} (\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^*, \\
\iint_{\Omega_1} \frac{r^*(\lambda, \lambda^*) e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^* \wedge d\lambda &= - \int_{\partial\Omega_1} \frac{(\lambda - s_1)^n r_1^*(\lambda) e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^*, \\
\iint_{\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_-^*} r(\lambda, \lambda^*) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda^* \wedge d\lambda &= \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_-^*} (\lambda^* - s_1^n) r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda,
\end{aligned} \tag{30}$$

uniformly for all  $\mathbb{C} \setminus \Omega_1$ . The boundary  $\partial\Omega_1$  has the counterclockwise orientation.



**Proof** Note that  $r(k, k^*) := nk^{*(n-1)}r_1(k)$ , using Green theorem, we have

$$\begin{aligned} \iint_{\Omega_1} \frac{r(\lambda, \lambda^*)e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda^* \wedge d\lambda &= \iint_{\Omega_1} \frac{\bar{\partial}((\lambda^* - s_1^*)^n r_1(\lambda) e^{-2i\theta(\lambda)})}{2\pi i(k-\lambda)} d\lambda^* \wedge d\lambda \\ &= \iint_{\Omega_1} \bar{\partial} \left( \frac{(\lambda^* - s_1^*)^n r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} \right) d\lambda^* \wedge d\lambda \\ &= \int_{\partial\Omega_1} \frac{(\lambda^* - s_1^*)^n r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda, \end{aligned}$$

and

$$\begin{aligned} \iint_{\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} r^*(\lambda, \lambda^*) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* \wedge d\lambda &= \iint_{\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} \partial((\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})})}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* \wedge d\lambda \\ &= - \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} (\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^*. \end{aligned}$$

Thus the proof is completed.

According to RH problems 1 and 2, we arrive at the following RH problem  $M_2(x, t; k) := \lim_{N \rightarrow \infty} M_1(x, t; k)$ :

**Riemann-Hilbert problem 3** Find a  $2 \times 2$  matrix function  $M_2(x, t; k)$  that meets the following conditions:

- *Analyticity:*  $M_2(x, t; k)$  is analytic in  $\mathbb{C} \setminus (\Gamma_{1\pm} \cup \Gamma_{2\pm})$  and takes continuous boundary values on  $\Gamma_{1\pm} \cup \Gamma_{2\pm}$ .
- *The jump condition:* The boundary values on the jump contour  $\Gamma_{1+} \cup \Gamma_{1-}$  are defined as

$$M_{2+}(x, t; k) = M_{2-}(x, t; k) V_2(x, t; k), \quad \lambda \in \Gamma_{1\pm} \cup \Gamma_{2\pm}, \quad (31)$$

where

$$V_2(x, t; k) = \begin{cases} \begin{bmatrix} 1 & 0 \\ - \int_{\partial\Omega_1} \frac{(\lambda^* - s_1^*)^n r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ - \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} (\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & \int_{\partial\Omega_1} \frac{(\lambda - s_1)^n r_1^*(\lambda) e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^* \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_-^*} (\lambda^* - s_1^*)^n r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (32)$$

- *The normalization:*

$$M_2(x, t; k) = \begin{cases} \mathbb{I}_2 + O(1/k) & k \rightarrow \infty, \\ \frac{i}{k} \sigma_3 Q_- + O(1), & k \rightarrow 0. \end{cases} \quad (33)$$

According to Eq. (23), we recover  $q(x, t)$  by means of the following formula:

$$q(x, t) = -i \lim_{k \rightarrow \infty} (kM_2(x, t; k))_{12}. \quad (34)$$

To restrict the values of  $n$  and  $m$ , we analyze the following three situations.

**Case I.** The single-breather solution. In this case, we choose  $n = m = 1$ . Then we arrive at the following Proposition.

**Proposition 3** Let  $\lambda_0 := s_1 + s_2$ , then the solution of the RH problem 3 is a single-breather solution  $q_1(x, t)$  with the discrete eigenvalue  $\lambda_0$  and normalization constants  $c_1 = s_3^2 r_1(\lambda_0)$ .

**Proof** The boundary of  $\Omega_1^*$ , which is the complex-conjugate domain of  $\Omega_1$ , defined by

$$k^* = s_1^* + \left( s_2^* + \frac{s_3^2}{(k - s_1)^m - s_2} \right)^{\frac{1}{m}} = s_1^* + \left( s_2^* + \frac{s_3^2}{k - \lambda_0} \right), \quad k \in \partial\Omega_1. \quad (35)$$

Substituting Eq. (35) into Eq. (32), we obtain

$$\begin{aligned} \int_{\partial\Omega_1} \frac{(\lambda^* - s_1^*)r_1(\lambda)e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda &= \frac{s_3^2 r_1(\lambda_0)e^{-2i\theta(\lambda_0)}}{k - \lambda_0}, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda_0^{*2} q_-} (\lambda - s_1)r_1^*(\lambda)e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* &= -\frac{\frac{q_0^2 q_-^*}{\lambda_0^{*2} q_-} s_3^2 r_1^*(\lambda_0)e^{-2i\theta(-\frac{q_0^2}{\lambda_0^*})}}{k + \frac{q_0^2}{\lambda_0^*}}, \\ \int_{\partial\Omega_1} \frac{(\lambda - s_1)r_1^*(\lambda)e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^* &= -\frac{s_3^2 r_1^*(\lambda_0)e^{2i\theta(\lambda_0^*)}}{k - \lambda_0^*}, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda_0^2 q_-^*} (\lambda^* - s_1^*)r_1(\lambda)e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda &= \frac{\frac{q_0^2 q_-}{\lambda_0^2 q_-^*} s_3^2 r_1(\lambda_0)e^{2i\theta(-\frac{q_0^2}{\lambda_0})}}{k + \frac{q_0^2}{\lambda_0}}, \end{aligned} \quad (36)$$

uniformly for all  $\mathbb{C} \setminus \Omega_1$ . Then Eqs. (32) can be rewritten as:

$$V_2(x, t; k)|_{m=n=1} = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\frac{s_3^2 r_1(\lambda_0)e^{-2i\theta(\lambda_0)}}{k - \lambda_0} & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ \frac{\frac{q_0^2 q_-^*}{\lambda_0^{*2} q_-} s_3^2 r_1^*(\lambda_0)e^{-2i\theta(-\frac{q_0^2}{\lambda_0^*})}}{k + \frac{q_0^2}{\lambda_0^*}} & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & -\frac{s_3^2 r_1^*(\lambda_0)e^{2i\theta(\lambda_0^*)}}{k - \lambda_0^*} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & \frac{\frac{q_0^2 q_-}{\lambda_0^2 q_-^*} s_3^2 r_1(\lambda_0)e^{2i\theta(-\frac{q_0^2}{\lambda_0})}}{k + \frac{q_0^2}{\lambda_0}} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (37)$$

Then, we consider following transformation:

$$\hat{M}_2(x, t; k) = \begin{cases} M_2(x, t; k) \begin{bmatrix} 1 & 0 \\ \frac{s_3^2 r_1(\lambda_0) e^{-2i\theta(\lambda_0)}}{k - \lambda_0} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1+}, \\ M_2(x, t; k) \begin{bmatrix} 1 & 0 \\ -\frac{\frac{q_0^2 q_-^*}{\lambda_0^2 q_-} s_3^2 r_1^*(\lambda_0) e^{-2i\theta(-\frac{q_0^2}{\lambda_0^*})}}{k + \frac{q_0^2}{\lambda_0^*}} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2+}, \\ M_2(x, t; k) \begin{bmatrix} 1 & -\frac{s_3^2 r_1^*(\lambda_0) e^{2i\theta(\lambda_0^*)}}{k - \lambda_0^*} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1-}, \\ M_2(x, t; k) \begin{bmatrix} 1 & \frac{\frac{q_0^2 q_-}{\lambda_0^2 q_-^*} s_3^2 r_1(\lambda_0) e^{2i\theta(-\frac{q_0^2}{\lambda_0})}}{k + \frac{q_0^2}{\lambda_0}} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2-}, \\ M_2(x, t; k), & \text{otherwise.} \end{cases} \quad (38)$$

Through the above transformations, we can obtain the residue condition for the matrix function  $\hat{M}_2(x, t; k)$ .  $\hat{M}_2(x, t; k)$  has simple pole at point  $k := \lambda_0$  with

$$\begin{aligned} \text{Res}_{k=\lambda_0} \hat{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_0} \hat{M}_2(x, t; k) \begin{bmatrix} 0 & 0 \\ s_3^2 r_1(\lambda_0) e^{-2i\theta(\lambda_0)} & 0 \end{bmatrix}, \\ \text{Res}_{k=-\frac{q_0^2}{\lambda_0^*}} \hat{M}_2(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{\lambda_0^*}} \hat{M}_2(x, t; k) \begin{bmatrix} 0 & 0 \\ -\frac{q_0^2 q_-^*}{\lambda_0^2 q_-} s_3^2 r_1^*(\lambda_0) e^{-2i\theta(-\frac{q_0^2}{\lambda_0^*})} & 0 \end{bmatrix}, \\ \text{Res}_{k=\lambda_0^*} \hat{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_0^*} \hat{M}_2(x, t; k) \begin{bmatrix} 0 & -s_3^2 r_1^*(\lambda_0) e^{2i\theta(\lambda_0^*)} \\ 0 & 0 \end{bmatrix}, \\ \text{Res}_{k=-\frac{q_0^2}{\lambda_0}} \hat{M}_2(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{\lambda_0}} \hat{M}_2(x, t; k) \begin{bmatrix} 0 & \frac{q_0^2 q_-}{\lambda_0^2 q_-^*} s_3^2 r_1(\lambda_0) e^{2i\theta(-\frac{q_0^2}{\lambda_0})} \\ 0 & 0 \end{bmatrix}, \end{aligned} \quad (39)$$

Therefore, the solution of RH problem 3 is the single breather  $q_1(x, t)$  with the eigenvalue  $\lambda_0$  with the normalization constants  $s_3^2 r_1(\lambda_0)$ .

In particular, we take  $q_0 = 1$ . When  $s_1 + s_2 \rightarrow i$ , we obtain the Peregrine's rational solution (rogue wave)  $q_{rw}(x, t)$  of the focusing NLS equation [64]:

$$q_{rw}(x, t) = 1 - \frac{4(4it + 1)}{4x^2 + 16t^2 + 1}.$$

**Case II.** The  $n$ -breather solution. In this case, we choose  $n = m$ . Then we arrive at the following Proposition.

**Proposition 4** If  $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$  is a solution to equation  $(k - s_1)^n = s_2$ , then the solution of the RH problem 3 is the  $n$ -breather solution  $q_n(x, t)$  with discrete eigenvalues  $\lambda_j, j = 1, \dots, n$  and the normalization constants

$$c_j = \frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}, j = 1, \dots, n.$$

**Proof** The boundary of  $\Omega_1^*$ , which is the complex-conjugate domain of  $\Omega_1$ , is defined by

$$k^* = s_1^* + \left( s_2^* + \frac{s_3^2}{(k - s_1)^m - s_2} \right)^{\frac{1}{n}}, \quad k \in \partial\Omega_1. \quad (40)$$

Substituting Eq. (40) into Eq. (32), we obtain

$$\begin{aligned} \int_{\partial\Omega_1} \frac{(\lambda^* - s_1^*)^n r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k - \lambda)} d\lambda &= \sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)}, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} (\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* &= - \sum_{j=1}^n \frac{\frac{q_0^2 q_-^*}{\lambda_j^{*2} q_-} s_3^2 r_1^*(\lambda_j) e^{-2i\theta(-\frac{q_0^2}{\lambda_j^*})}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k + \frac{q_0^2}{\lambda_j^*})}, \\ \int_{\partial\Omega_1} \frac{(\lambda - s_1)^n r_1^*(\lambda) e^{2i\theta(\lambda^*)}}{2\pi i(k - \lambda^*)} d\lambda^* &= - \sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)}, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_+^*} (\lambda^* - s_1^*)^n r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda &= \sum_{j=1}^n \frac{\frac{q_0^2 q_-}{\lambda_j^2 q_+^*} s_3^2 r_1(\lambda_j) e^{2i\theta(-\frac{q_0^2}{\lambda_j})}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k + \frac{q_0^2}{\lambda_j})}, \end{aligned} \quad (41)$$

Then Eqs. (32) can be rewritten as

$$V_2(x, t; k)|_{m=n} = \begin{cases} \begin{bmatrix} 1 & 0 \\ - \sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)} & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ \sum_{j=1}^n \frac{\frac{q_0^2 q_-^*}{\lambda_j^{*2} q_-} s_3^2 r_1^*(\lambda_j) e^{-2i\theta(-\frac{q_0^2}{\lambda_j^*})}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k + \frac{q_0^2}{\lambda_j^*})} & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & - \sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & \sum_{j=1}^n \frac{\frac{q_0^2 q_-}{\lambda_j^2 q_+^*} s_3^2 r_1(\lambda_j) e^{2i\theta(-\frac{q_0^2}{\lambda_j})}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k + \frac{q_0^2}{\lambda_j})} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (42)$$

Then, we consider following transformation:

$$\tilde{M}_2(x, t; k) = \begin{cases} M_2(x, t; k) \begin{bmatrix} 1 & 0 \\ \sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1+}, \\ M_2(x, t; k) \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^n \frac{\frac{q_0^2 q_-}{\lambda_j^{*2} q_-} s_3^2 r_1^*(\lambda_j) e^{-2i\theta(-\frac{q_0^2}{\lambda_j^*})}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k + \frac{q_0^2}{\lambda_j^*})} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2+}, \\ M_2(x, t; k) \begin{bmatrix} 1 & -\sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1-}, \\ M_2(x, t; k) \begin{bmatrix} 1 & \sum_{j=1}^n \frac{\frac{q_0^2 q_-}{\lambda_j^2 q_-} s_3^2 r_1(\lambda_j) e^{2i\theta(-\frac{q_0^2}{\lambda_j})}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k + \frac{q_0^2}{\lambda_j})} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2-}, \\ M_2(x, t; k), & \text{otherwise.} \end{cases} \quad (43)$$

Through the above transformations, we can obtain the residue condition for the matrix function  $\tilde{M}_2(x, t; k)$ .  $\tilde{M}_2(x, t; k)$  has simple poles at each point in  $\{\lambda_j\}_{j=1}^n$  with

$$\begin{aligned} \text{Res}_{k=\lambda_j} \tilde{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_j} \tilde{M}_2(x, t; k) \begin{bmatrix} 0 & 0 \\ \frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)} e^{-2i\theta(\lambda_j)} & 0 \end{bmatrix}, \\ \text{Res}_{k=-\frac{q_0^2}{\lambda_j^*}} \tilde{M}_2(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{\lambda_j^*}} \tilde{M}_2(x, t; k) \begin{bmatrix} 0 & 0 \\ -\frac{q_0^2 q_-}{\lambda_j^{*2} q_-} \frac{s_3^2 r_1^*(\lambda_j)}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)} e^{-2i\theta(-\frac{q_0^2}{\lambda_j^*})} & 0 \end{bmatrix}, \\ \text{Res}_{k=\lambda_j^*} \tilde{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_j^*} \tilde{M}_2(x, t; k) \begin{bmatrix} 0 & -\frac{s_3^2 r_1^*(\lambda_j)}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)} e^{2i\theta(\lambda_j^*)} \\ 0 & 0 \end{bmatrix}, \\ \text{Res}_{k=-\frac{q_0^2}{\lambda_j}} \tilde{M}_2(x, t; k) &= \lim_{k \rightarrow -\frac{q_0^2}{\lambda_j}} \tilde{M}_2(x, t; k) \begin{bmatrix} 0 & \frac{q_0^2 q_-}{\lambda_j^2 q_-} \frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)} e^{2i\theta(-\frac{q_0^2}{\lambda_j})} \\ 0 & 0 \end{bmatrix}, \end{aligned} \quad (44)$$

Therefore, the solution of the RH problem 3 is the  $n$ -breather state  $q_n(x, t)$ , with discrete eigenvalues  $\lambda_j$ ,  $j = 1, \dots, n$  and normalization constants  $\frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}$ ,  $j = 1, \dots, n$ .

**Case III.** The  $n$ -soliton solution. Note that the solution of RH problem 3 with  $q_0 \rightarrow 0$  can reduce to the known  $n$ -soliton solution  $q_n(x, t)$  [59], which can be also directly derived as the limit for  $q_0 \rightarrow 0$  of the obtained  $n$ -breather solution. Here we choose  $n = m$ . Then we arrive at the following Proposition.

**Proposition 5** If  $\{\lambda_1, \lambda_2, \dots, \lambda_n\}$  is the solution to equation  $(k - s_1)^n = s_2$  and  $q_0 \rightarrow 0$ , then the solution of the RH problem 3 is the  $n$ -soliton state  $q_n(x, t)$  with discrete eigenvalues  $\lambda_j$ ,  $j = 1, \dots, n$  and normalization constants

$$c_j = \frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}, j = 1, \dots, n.$$

**Proof** Using Eqs. (41), we obtain

$$\begin{aligned} \lim_{q_0 \rightarrow 0} \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} (\lambda - s_1)^n r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i (k + \frac{q_0^2}{\lambda^*})} d\lambda^* &= 0, \\ \lim_{q_0 \rightarrow 0} \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^2 q_-^*} (\lambda^* - s_1^*)^n r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i (k + \frac{q_0^2}{\lambda})} d\lambda &= 0. \end{aligned} \quad (45)$$

Substituting Eqs. (41) and (45) in Eq. (32), we obtain

$$V_2(x, t; k)|_{m=n, q_0 \rightarrow 0} = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)} & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & -\sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}. \end{cases} \quad (46)$$

Therefore, we can rewrite the RH problem for matrix function  $M_2(x, t; k)$  as follows: Find a  $2 \times 2$  matrix function  $M_2(x, t; k)$  that satisfies the particular conditions:

- *Analyticity:*  $M_2(x, t; k)$  is analytic in  $\mathbb{C} \setminus \Gamma_{1\pm}$  and takes continuous boundary values on  $\Gamma_{1\pm}$ .
- *The jump condition:* The boundary values on the jump contour  $\Gamma_{1+}$  are defined as

$$M_{2+}(x, t; k) = M_{2-}(x, t; k) \begin{cases} \begin{bmatrix} 1 & 0 \\ -\sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)} & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & -\sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)} \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}. \end{cases} \quad (47)$$

- *The normalization:*

$$M_2(x, t; k) = \mathbb{I}_2 + O(1/k), \quad k \rightarrow \infty. \quad (48)$$

Then, we consider following transformation:

$$\overline{M}_2(x, t; k) = \begin{cases} M_2(x, t; k) \begin{bmatrix} 1 & 0 \\ \sum_{j=1}^n \frac{s_3^2 r_1(\lambda_j) e^{-2i\theta(\lambda_j)}}{\prod_{k \neq j} (\lambda_j - \lambda_k)(k - \lambda_j)} & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1+}, \\ M_2(x, t; k) \begin{bmatrix} 1 & -\sum_{j=1}^n \frac{s_3^2 r_1^*(\lambda_j) e^{2i\theta(\lambda_j^*)}}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)(k - \lambda_j^*)} \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1-}, \\ M_2(x, t; k), & \text{otherwise.} \end{cases} \quad (49)$$

Through the above transformations, we can obtain the residue condition for the matrix function  $\overline{M}_2(x, t; k)$ .

$\overline{M}_2(x, t; k)$  has simple poles at each point in  $\{\lambda_j\}_{j=1}^n$  with:

$$\begin{aligned} \operatorname{Res}_{k=\lambda_j} \overline{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_j} \overline{M}_2(x, t; k) \begin{bmatrix} 0 & 0 \\ \frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)} e^{-2i\theta(\lambda_j)} & 0 \end{bmatrix}, \\ \operatorname{Res}_{k=\lambda_j^*} \overline{M}_2(x, t; k) &= \lim_{k \rightarrow \lambda_j^*} \overline{M}_2(x, t; k) \begin{bmatrix} 0 & -\frac{s_3^2 r_1^*(\lambda_j)}{\prod_{k \neq j} (\lambda_j^* - \lambda_k^*)} e^{2i\theta(\lambda_j^*)} \\ 0 & 0 \end{bmatrix}, \end{aligned} \quad (50)$$

Therefore, the solution of the RH problem 3 is the  $n$ -soliton state  $q_n(x, t)$  with discrete eigenvalues  $\lambda_j, j = 1, \dots, n$  and normalization constants  $\frac{s_3^2 r_1(\lambda_j)}{\prod_{k \neq j} (\lambda_j - \lambda_k)}, j = 1, \dots, n$ .

### 3.2 The line domain

In this section, we address the limit of  $N \rightarrow \infty$ , under the additional assumptions:

- Poles  $\{k_j\}_{j=1}^N$  are sampled from a smooth density function  $\rho(k)$  so that  $\int_a^{-ik_j} \rho(\lambda) d\lambda = \frac{j}{N}, j = 1, \dots, N$ .
- The coefficients  $\{c_j\}_{j=1}^N$  satisfy

$$c_j = \frac{i(b-a)r(k_j)}{N\pi}, \quad b > a > 0, \quad (51)$$

where  $r(k)$  is a real-valued, continuous and non-vanishing function of  $k \in (ia, ib)$ , subject to the symmetry relation,  $r(k^*) = r(k) = r(-\frac{q_0^2}{k}) = r(-\frac{q_0^2}{k^*})$ .

**Proposition 6** For any open set  $A_+(B_+)$  containing the interval  $[ia, ib]$  ( $[-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}]$ ), and any open set  $A_-(B_-)$  containing the interval  $[-ib, -ia]$  ( $[\frac{iq_0^2}{b}, \frac{iq_0^2}{a}]$ ), the following identities hold:

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} = \int_{ia}^{ib} \frac{r(w) e^{-2i\theta(w)}}{\pi(k-w)} dw, \quad (52)$$

uniformly for all  $\mathbb{C} \setminus A_+$ .

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\frac{q_0^2 q_-^*}{k_j^* q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{k + \frac{q_0^2}{k_j^*}} = \int_{-\frac{iq_0^2}{a}}^{-\frac{iq_0^2}{b}} \frac{q_-^* r(w) e^{-2i\theta(w)}}{q_- \pi(k-w)} dw, \quad (53)$$

uniformly for all  $\mathbb{C} \setminus B_+$ .

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j^* e^{2i\theta(k_j^*)}}{k - k_j^*} = - \int_{-ib}^{-ia} \frac{r(w) e^{2i\theta(w)}}{\pi(k-w)} dw, \quad (54)$$

uniformly for all  $\mathbb{C} \setminus A_-$ .

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\frac{q_0^2 q_-}{k_j^2 q_-^*} c_j e^{2i\theta(-\frac{q_0^2}{k_j})}}{k + \frac{q_0^2}{k_j}} = - \int_{\frac{iq_0^2}{b}}^{\frac{iq_0^2}{a}} \frac{q_- r(w) e^{2i\theta(w)}}{q_-^* \pi(k-w)} dw, \quad (55)$$

uniformly for all  $\mathbb{C} \setminus B_-$ . The open intervals  $(ia, ib), (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}), (\frac{iq_0^2}{b}, \frac{iq_0^2}{a})$  and  $(-ib, -ia)$  are both oriented upwards.

**Proof** Using Eq. (51), we have

$$\lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{c_j e^{-2i\theta(k_j)}}{k - k_j} = \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{i(b-a) r(k_j) e^{-2i\theta(k_j)}}{N \pi(k - k_j)} = \int_{ia}^{ib} \frac{r(w) e^{-2i\theta(w)}}{\pi(k - w)} dw,$$

and

$$\begin{aligned} \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{\frac{q_0^2 q_-^*}{k_j^{*2} q_-} c_j^* e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{k + \frac{q_0^2}{k_j^*}} &= \lim_{N \rightarrow \infty} \sum_{j=1}^N \frac{-i(b-a) \frac{q_0^2 q_-^*}{k_j^{*2} q_-} r(k_j) e^{-2i\theta(-\frac{q_0^2}{k_j^*})}}{N \pi(k + \frac{q_0^2}{k_j^*})} \\ &= - \int_{-ib}^{-ia} \frac{\frac{q_0^2 q_-^*}{\lambda^2 q_-} r(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda})}}{\pi(k + \frac{q_0^2}{\lambda})} d\lambda \\ &= \int_{-\frac{iq_0^2}{a}}^{-\frac{iq_0^2}{b}} \frac{q_-^* r(w) e^{-2i\theta(w)}}{q_- \pi(k - w)} dw, \end{aligned}$$

Thus the proof is completed.

At  $N \rightarrow \infty$ , according to Proposition 6, the jump matrix  $V_1(x, t; k)$ , defined by Eq. (21), can be rewritten as:

$$V_1(x, t; k)|_{N \rightarrow \infty} = \begin{cases} \begin{bmatrix} 1 & 0 \\ - \int_{ia}^{ib} \frac{r(w) e^{-2i\theta(w)}}{\pi(k - w)} dw & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ \int_{-\frac{iq_0^2}{a}}^{-\frac{iq_0^2}{b}} \frac{q_-^* r(w) e^{-2i\theta(w)}}{q_- \pi(k - w)} dw & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & \int_{-ib}^{-ia} \frac{r(w) e^{2i\theta(w)}}{\pi(k - w)} dw \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & - \int_{\frac{iq_0^2}{b}}^{\frac{iq_0^2}{a}} \frac{q_- r(w) e^{2i\theta(w)}}{q_-^* \pi(k - w)} dw \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (56)$$



Apply the following transformation:

$$M_3(x, t; k) = \begin{cases} M_1(x, t; k) \begin{bmatrix} 1 & 0 \\ \int_{ia}^{ib} \frac{r(w)e^{-2i\theta(w)}}{\pi(k-w)} dw & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1+}, \\ M_1(x, t; k) \begin{bmatrix} 1 & 0 \\ -\int_{-\frac{iq_0^2}{a}}^{-\frac{iq_0^2}{b}} \frac{q_-^* r(w)e^{-2i\theta(w)}}{q_- \pi(k-w)} dw & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2+}, \\ M_1(x, t; k) \begin{bmatrix} 1 & \int_{-ib}^{-ia} \frac{r(w)e^{2i\theta(w)}}{\pi(k-w)} dw \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{1-}, \\ M_1(x, t; k) \begin{bmatrix} 1 & -\int_{\frac{iq_0^2}{b}}^{\frac{iq_0^2}{a}} \frac{q_- r(w)e^{2i\theta(w)}}{q_-^* \pi(k-w)} dw \\ 0 & 1 \end{bmatrix}, & k \text{ within } \Gamma_{2-}, \\ M_1(x, t; k), & \text{otherwise.} \end{cases} \quad (57)$$

Then the matrix function  $M_3(x, t; k)$  satisfies the following RH problem:

**Riemann-Hilbert problem 4** Find a  $2 \times 2$  matrix function  $M_3(x, t; k)$  that satisfies the particular conditions:

- *Analyticity:*  $M_3(x, t; k)$  is analytic in  $\mathbb{C} \setminus ((ia, ib) \cup (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}) \cup (-ib, -ia) \cup (\frac{iq_0^2}{b}, \frac{iq_0^2}{a}))$  and takes continuous boundary values on  $(ia, ib) \cup (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}) \cup (-ib, -ia) \cup (\frac{iq_0^2}{b}, \frac{iq_0^2}{a})$  (Directions of these open intervals are all facing upwards).
- *The jump condition:* The boundary values on the jump contour  $(ia, ib) \cup (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}) \cup (-ib, -ia) \cup (\frac{iq_0^2}{b}, \frac{iq_0^2}{a})$  are defined as

$$M_{3+}(x, t; k) = M_{3-}(x, t; k)V_3(x, t; k), \quad \lambda \in (ia, ib) \cup (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}) \cup (-ib, -ia) \cup (\frac{iq_0^2}{b}, \frac{iq_0^2}{a}),$$

where

$$V_3(x, t; k) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -2ir(k)e^{-2i\theta(k)} & 1 \end{bmatrix}, & k \in (ia, ib), \\ \begin{bmatrix} 1 & 0 \\ \frac{2iq_-^* r(k)e^{-2i\theta(k)}}{q_-} & 1 \end{bmatrix}, & k \in (-\frac{iq_0^2}{a}, -\frac{iq_0^2}{b}), \\ \begin{bmatrix} 1 & -2ir(k)e^{2i\theta(k)} \\ 0 & 1 \end{bmatrix}, & k \in (-ib, -ia), \\ \begin{bmatrix} 1 & \frac{2iq_- r(k)e^{2i\theta(k)}}{q_-^*} \\ 0 & 1 \end{bmatrix}, & k \in (\frac{iq_0^2}{b}, \frac{iq_0^2}{a}). \end{cases} \quad (58)$$

- *The normalization:*

$$M_3(x, t; k) = \begin{cases} \mathbb{I}_2 + O(1/k) & k \rightarrow \infty, \\ \frac{i}{k} \sigma_3 Q_- + O(1), & k \rightarrow 0. \end{cases} \quad (59)$$

**Proof** Using the Plemelj formula, we have

$$\begin{aligned}
M_{3+}(x, t; k) &= M_{1+}(x, t; k) \begin{bmatrix} 1 & 0 \\ \int_{ia}^{ib} \frac{r(w)e^{-2i\theta(w)}}{\pi(k_+ - w)} dw & 1 \end{bmatrix} \\
&= M_{1-}(x, t; k) \begin{bmatrix} 1 & 0 \\ \int_{ia}^{ib} \frac{r(w)e^{-2i\theta(w)}}{\pi(k_- - w)} dw & 1 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ -2ir(k)e^{-2i\theta(k)} & 1 \end{bmatrix}, \quad k \in (ia, ib),
\end{aligned} \tag{60}$$

Then, we have

$$V_3(x, t; k) \begin{bmatrix} 1 & 0 \\ -2ir(k)e^{-2i\theta(k)} & 1 \end{bmatrix}, \quad k \in (ia, ib). \tag{61}$$

Using the same method, we can prove other cases as well.

According to Eq. (23), we recover  $q(x, t)$  by the following formula:

$$q(x, t) = -i \lim_{k \rightarrow \infty} (kM_3(x, t; k))_{12}. \tag{62}$$

Then the RH problem 4 for the matrix function  $M_3(x, t; k)$  represents the breather gas.

### 3.3 The elliptic domain

In this section, we consider the limit of  $N \rightarrow \infty$ , under the additional assumptions:

- Discrete eigenvalues  $k_j, j = 1, \dots, N$  with normalization constants  $c_j, j = 1, \dots, N$  fill a uniformly compact domain  $\Omega_2$  of the complex upper half plane  $\mathbb{C}_+$ , that is,

$$\Omega_2 := \left\{ k \in \mathbb{C} \mid \frac{\operatorname{Re}(k)^2}{b_2^2} + \frac{(2\operatorname{Im}(k) - a_1 - a_2)^2}{4b_1^2} < 1 \right\}, \quad \Omega_2 \subset D_+, \tag{63}$$

where  $ia_1$  and  $ia_2$  ( $a_2 > a_1$ ) are the focal points of the ellipse  $\partial\Omega_2$ ,  $b_1 = \sqrt{b_2^2 + (\frac{a_2 - a_1}{2})^2}$ , and  $b_2$  is sufficiently small so that  $\Omega_2$  lies in domain  $D_+$ .

- The normalization constants  $c_j, j = 1, \dots, N$  have the following form:

$$c_j = \frac{|\Omega_2| r_1(k_j)}{N\pi}. \tag{64}$$

where  $|\Omega_2|$  means the area of the domain  $\Omega_2$  and  $r_1(k)$  is an analytic functions in domain  $\Omega_2$ , subject to the symmetry condition  $r_1^*(k) = r_1(k^*)$ .

According to RH problems 1 and 2, we arrive at the following RH problem  $M_4(x, t; k) := \lim_{N \rightarrow \infty} M_1(x, t; k)$ :

**Riemann-Hilbert problem 5** Find a  $2 \times 2$  matrix function  $M_4(x, t; k)$  that meets the particular conditions:

- *Analyticity:*  $M_4(x, t; k)$  is analytic in  $\mathbb{C} \setminus (\Gamma_{1\pm} \cup \Gamma_{2\pm})$  and takes continuous boundary values on  $\Gamma_{1\pm} \cup \Gamma_{2\pm}$ .
- *The jump condition:* The boundary values on the jump contour  $\Gamma_{1+} \cup \Gamma_{1-}$  are defined as

$$M_{4+}(x, t; k) = M_{4-}(x, t; k)V_4(x, t; k), \quad \lambda \in \Gamma_{1\pm} \cup \Gamma_{2\pm}, \tag{65}$$

where

$$V_4(x, t; k) = \begin{cases} \begin{bmatrix} 1 & 0 \\ -\int_{\partial\Omega_2} \frac{\lambda^* r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda & 1 \end{bmatrix}, & k \in \Gamma_{1+}, \\ \begin{bmatrix} 1 & 0 \\ -\int_{\partial\Omega_2} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} \lambda r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* & 1 \end{bmatrix}, & k \in \Gamma_{2+}, \\ \begin{bmatrix} 1 & \int_{\partial\Omega_2} \frac{\lambda r_1^*(\lambda) e^{2i\theta(\lambda^*)}}{2\pi i(k-\lambda^*)} d\lambda^* \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{1-}, \\ \begin{bmatrix} 1 & \int_{\partial\Omega_2} \frac{\frac{q_0^2 q_-}{\lambda^{*2} q_-} \lambda^* r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda \\ 0 & 1 \end{bmatrix}, & k \in \Gamma_{2-}. \end{cases} \quad (66)$$

• The normalization:

$$M_4(x, t; k) = \begin{cases} \mathbb{I}_2 + O(1/k) & k \rightarrow \infty, \\ \frac{i}{k} \sigma_3 Q_- + O(1), & k \rightarrow 0. \end{cases} \quad (67)$$

**Proposition 7** The following identities hold:

$$\begin{aligned} \int_{\partial\Omega_1} \frac{\lambda^* r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda &= \int_{ia_1}^{ia_2} \frac{\Delta F(k) r_1(\lambda) e^{-2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-^*}{\lambda^{*2} q_-} \lambda r_1^*(\lambda) e^{-2i\theta(-\frac{q_0^2}{\lambda^*})}}{2\pi i(k + \frac{q_0^2}{\lambda^*})} d\lambda^* &= \int_{-\frac{iq_0^2}{a}}^{-\frac{iq_0^2}{b}} \frac{\frac{q_-^*}{q_-} \Delta F^*(-\frac{q_0^2}{w^*}) r_1(w) e^{-2i\theta(w)}}{2\pi i(k-w)} dw, \\ \int_{\partial\Omega_1} \frac{\lambda r_1^*(\lambda) e^{2i\theta(\lambda^*)}}{2\pi i(k-\lambda^*)} d\lambda^* &= -\int_{-ia_2}^{-ia_1} \frac{\Delta F^*(k) r_1(\lambda) e^{2i\theta(\lambda)}}{2\pi i(k-\lambda)} d\lambda, \\ \int_{\partial\Omega_1} \frac{\frac{q_0^2 q_-}{\lambda^{*2} q_-} \lambda^* r_1(\lambda) e^{2i\theta(-\frac{q_0^2}{\lambda})}}{2\pi i(k + \frac{q_0^2}{\lambda})} d\lambda &= -\int_{\frac{iq_0^2}{b}}^{\frac{iq_0^2}{a}} \frac{\frac{q_-}{q_-^*} \Delta F(-\frac{q_0^2}{w}) r_1(w) e^{2i\theta(w)}}{2\pi i(k-w)} dw, \end{aligned} \quad (68)$$

where  $\Delta F(k) = F_+(k) - F_-(k)$ , and the function  $F(k)$  is analytic in complex plane  $\mathbb{C}$  away from the segment  $[ia_1, ia_2]$ , with boundary values  $F_{\pm}(k)$ .

**Proof** The boundary of the complex-conjugate domain  $\Omega_2^*$  of  $\Omega_2$  is described by

$$k^* = \left(1 - \frac{8b_1^2}{(a_2 - a_1)^2}\right) \left(k - \frac{i(a_1 + a_2)}{2}\right) + \frac{8b_1 b_2}{(a_2 - a_1)^2} F(k) - \frac{i(a_1 + a_2)}{2}, \quad k \in \partial\Omega_2, \quad (69)$$

where  $F(k)^2 = (k - ia_1)(k - ia_2)$ . Using Eq. (69), one can obtain Eq. (68).

According to Proposition 7, we find that the RH problem 5 for matrix function  $M_4(x, t; k)$  is equivalent to the one in case of the line domain.

## 4 Conclusions and discussions

Based on the IST and RH problems, we have investigated a breather gas represented by the  $N_\infty$ -breather solution of the focusing NLS equation with nonzero BCs. In terms of scattering data of IST, the  $N$ -breather solutions are based on the set of discrete eigenvalues  $K \equiv \{k_j, -\frac{q_0^2}{k_j^*}, k_j^*, -\frac{q_0^2}{k_j}\}_{j=1}^N$  with normalization constants.

$\{c_j, -\frac{q_0^2 c_j^*}{k_j^{*2}}, -c_j^*, \frac{q_0^2 c_j}{k_j^2}\}_{j=1}^N$ . By concentrating the set of  $\{k_j\}_{j=1}^N$  in different domains, we have produced different types of breather gases which coagulate into the following effective forms: i) The concentration domain in the form of a disk condenses the gas into the single-breather solution with the spectral eigenvalue located at the disk's center; ii) The quadrater domain with  $m = n$  and  $q \rightarrow 0$  leads to the coagulation of the gas into the  $n$ -breather solution. These are examples of the breather-gas shielding. The discrete spectra concentrated in line domains imply solving the corresponding RH problems. The case of the discrete spectra lying on an ellipse is tantamount to the case of the line domain. **When discrete spectra are uniformly distributed within a specified region, the interaction among breathers manifests itself in the form of  $n$ -breathers, where parameter  $n$  is correlated with the region in question. The phenomenon of breather shielding can explain the distribution of the breathers when the discrete spectrum is densely distributed. The methodology presented here can be extended to other integrable equations and can also be employed to investigate the asymptotic behavior of breathers in different regions. For the phenomenon of breather shielding, we have developed here only the analytical framework. Verification of the findings by means of numerical methods is a subject for a separate work.**

The breather-gas shielding predicted by the present analysis can be observed in fiber optics, BEC, and other physical realizations of the NLS. The approach developed in this work can be extended to other integrable models – first of all, those based on ZS-type spectral problems. A challenging possibility is to extend the analysis of the shielding phenomenology to quantized NLS fields, in which breather states feature specific fluctuations [87].

## Acknowledgments

We thank Prof. Marco Bertola (Concordia University, Canada) for valuable suggestions and discussions. This work of W.W. was supported by the China Postdoctoral Science Foundation (No.043201025). The work of Z.Y. was supported by the National Natural Science Foundation of China (Nos.11925108 and 12471242). The work of B.A.M. is supported, in part, by Israel Science Foundation (No. 1695/22).

## Data availability statement

All data that support the findings of this study are included within the article.

## References

- [1] J. S. Russell, Report on waves, Report of the 14th Meeting of the British Association for the Advancement of Science, 311-390 (1844).
- [2] D. J. Korteweg and G. de Vries, On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves, Phil. Mag. 39, 422 (1895).
- [3] E. Fermi, J. Pasta, and S. Ulam, Studies of nonlinear problems. I. Los Alamos report LA-1940 (1955).
- [4] T. Dauxois and S. Ruffo, Fermi-Pasta-Ulam nonlinear lattice oscillations, Scholarpedia 3, 5538 (2008).
- [5] N. J. Zabusky and M. D. Kruskal, Interaction of "solitons" in a collisionless plasma and the recurrence of initial states, Phys. Rev. Lett. 15, 240 (1965).
- [6] C. S. Gardner, J. M. Greene, M. D. Kruskal, and R. M. Miura, Method for solving the Korteweg-de Vries equation, Phys. Rev. Lett. 19, 1095-1097 (1967).
- [7] P. D. Lax, Integrals of nonlinear equations of evolution and solitary waves, Comm. Pure Appl. Math. 21, 467-490 (1968).

- [8] M. J. Ablowitz, D. J. Kaup, A. C. Newell, and H. Segur, The inverse scattering transform. Fourier analysis for nonlinear problems, Stud. Appl. Math. 53, 249-315 (1974).
- [9] V. E. Zakharov and A. B. Shabat, Exact theory of two-dimensional self-focusing and one-dimensional self-modulation of waves in nonlinear media, Zh. Eksp. Teor. Fiz. 61, 118-134 (1971) [Sov. Phys. JETP 34, 62-69 (1972)].
- [10] A. Hasegawa and F. Tappert, Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. 1. Anomalous dispersion, Appl. Phys. Lett. 23, 142-144 (1973).
- [11] A. Hasegawa and F. Tappert, Transmission of stationary nonlinear optical pulses in dispersive dielectric fibers. 2. Normal dispersion, Appl. Phys. Lett. 23, 171-172 (1973).
- [12] J. Satsuma and N. Yajima, Initial value problems of one-dimensional self-modulation of nonlinear waves in dispersive media, Suppl. Prog. Theor. Phys. 55, 284-306 (1974).
- [13] L. F. Mollenauer, R. H. Stolen, and J. P. Gordon, Experimental observation of picosecond pulse narrowing and solitons in optical fibers, Phys. Rev. Lett. 45, 1095-1098 (1980).
- [14] A. Hasegawa and Y. Kodama, *Solitons in Optical Communications* (Oxford University Press, Oxford, 1995).
- [15] N. Akhmediev and A. Ankiewicz, *Solitons: Nonlinear Pulses and Beams* (Chapman and Hall, London, 1997).
- [16] G. Agrawal, *Nonlinear Fibre Optics* (5th ed.) (Academic Press, New York, 2014).
- [17] Y. S. Kivshar and G. Agrawal, *Optical Solitons: From Fibers to Photonic Crystals* (Academic Press, San Diego, CA, 2003).
- [18] J. S. Peng, S. Boscolo, Z. H. Zhao, and H. P. Zeng, Breathing dissipative solitons in mode-locked fiber lasers, Science Adv. 5, eaax1110 (2019).
- [19] L. P. Pitaevskii and S. Stringari, *Bose-Einstein Condensation* (Clarendon Press, New York, 2003).
- [20] K. E. Strecker, G. B. Partridge, A. G. Truscott, and R. G. Hulet, Formation and propagation of matter-wave soliton trains, Nature 417, 150-153 (2002).
- [21] L. Khaykovich, F. Schreck, G. Ferrari, T. Bourdel, J. Cubizolles, L. D. Carr, Y. Castin, and C. Salomon, Formation of a matter-wave bright soliton, Science 296, 1290-1293 (2002).
- [22] A. Di Carli, C. D. Colquhoun, G. Henderson, S. Flannigan, G.-L. Oppo, A. J. Daley, S. Kuhr, and E. Haller, Excitation modes of bright matter-wave solitons, Phys. Rev. Lett. 123, 123602 (2019).
- [23] D. Luo, Y. Jin, J. H. V. Nguyen, B. A. Malomed, O. V. Marchukov, V. A. Yurovsky, V. Dunjko, M. Olshani, and R. G. Hulet, Creation and characterization of matter-wave breathers, Phys. Rev. Lett. 125, 183902 (2020).
- [24] T. Dauxois and M. Peyrard, *Physics of Solitons* (Cambridge University Press, Cambridge, 2006).
- [25] A. R. Osborne, *Nonlinear Ocean Waves* (Academic Press, New York, 2009).
- [26] L. D. Faddeev and L. A. Takhtajan, *Hamiltonian Methods in the Theory of Solitons* (Springer, Berlin, 1987).
- [27] M. J. Ablowitz and P. A. Clarkson, *Solitons, Nonlinear Evolution Equations and Inverse Scattering* (Cambridge University Press, Cambridge, 1991).
- [28] J. Yang, *Nonlinear Waves in Integrable and Nonintegrable Systems* (SIAM, Philadelphia, 2010).
- [29] P. G. Kevrekidis, D. J. Frantzeskakis, and R. CarreteroGonzález, *The Defocusing Nonlinear Schrödinger Equation: From Dark Soliton to Vortices and Vortex Rings* (SIAM, Philadelphia, 2015).
- [30] F. Ye, D. Mihalache, B. Hu, and N.C. Panoiu, Subwavelength plasmonic lattice solitons in arrays of metallic nanowires, Phys. Rev. Lett. 104, 106802 (2010).
- [31] B. A. Malomed, D. Mihalache, F. Wise, and L. Torner, Spatiotemporal optical solitons J. Opt. B Quantum Semiclassical Opt. 7, R53-R72 (2005).
- [32] L.-C. Crasovan, Y. V. Kartashov, D. Mihalache, L. Torner, Y. S. Kivshar, and V. M. Pérez-García, Soliton "molecules": Robust clusters of spatiotemporal optical solitons, Phys. Rev. E 67, 046610 (2003).
- [33] A. Weller, J. P. Ronzheimer, C. Gross, J. Esteve, M. K. Oberthaler, D. J. Frantzeskakis, G. Theocharis, and P. G. Kevrekidis, Experimental observation of oscillating and interacting matter wave dark solitons, Phys. Rev. Lett. 101, 130401 (2008).
- [34] Y. V. Kartashov, B. A. Malomed, and L. Torner, Solitons in nonlinear lattices, Rev. Mod. Phys. 83, 247 (2011).
- [35] V. V. Konotop, J. Yang, and D. A. Zezyulin, Nonlinear waves in PT-symmetric systems, Rev. Mod. Phys. 88, 035002 (2016).
- [36] B. A. Malomed, *Multidimensional Solitons* (American Institute of Physics Publishers: Melville, NY, 2022).
- [37] V.E. Zakharov, Kinetic equation for solitons, Sov. Phys. JETP 33, 538-541 (1971).
- [38] S. V. Manakov, V. E. Zakharov, L. A. Bordag, A. R. Its, and V. B. Matveev, Two-dimensional solitons of the Kadomtsev-Petviashvili equation and their interaction, Phys. Lett. A 63, 205-206 (1977).
- [39] G. A. El, A.M. Kamchantov, Kinetic equation for a dense soliton gas, Phys. Rev. Lett. 95, 204101 (2005).
- [40] G. A. El, A. M. Kamchatnov, M. V. Pavlov, and S. A. Zykov, Kinetic equation for a soliton gas and its hydrodynamic reductions, J. Nonlinear Sci. 21, 151-191 (2011).
- [41] G. A. El, Critical density of a soliton gas, Chaos 26, 023105 (2016).
- [42] T. Congy, G. A. El, and G. Roberti, Soliton gas in bidirectional dispersive hydrodynamics, Phys. Rev. E 103, 042201 (2021).

- [43] E. G. Shurgalina and E. N. Pelinovsky, Nonlinear dynamics of a soliton gas: Modified Korteweg-de Vries equation framework, *Phys. Lett. A* 380, 2049-2053 (2016).
- [44] D. Dutykh and E. Pelinovsky, Numerical simulation of a solitonic gas in KdV and KdV-BBM equations, *Phys. Lett. A* 378, 3102 (2014).
- [45] B. Doyon, T. Yoshimura, and J.-S. Caux, Soliton gases and generalized hydrodynamics, *Phys. Rev. Lett.* 120, 045301 (2018).
- [46] A. Gelash, D. Agafontsev, V. Zakharov, G. A. El, S. Randoux, and P. Suret, Bound state soliton gas dynamics underlying the spontaneous modulational instability, *Phys. Rev. Lett.* 123, 234102 (2019).
- [47] A. Gelash, D. Agafontsev, P. Suret, and S. Randoux, Solitonic model of the condensate, *Phys. Rev. E* 104, 044213 (2021).
- [48] Gennady A El, Soliton gas in integrable dispersive hydrodynamics, *J. Stat. Mech.* 2021, 114001 (2021).
- [49] P. Suret, S. Randoux, A. Gelash, D. Agafontsev, B. Doyon, and G. El, Soliton Gas: Theory, numerics and experiments, *arXiv: 2304.06541* (2024).
- [50] G. A. El and A. Tovbis, Spectral theory of soliton and breather gases for the focusing nonlinear Schrödinger equation, *Phys. Rev. E* 101, 052207 (2020).
- [51] G. Roberti, G. A. El, A. Tovbis, F. Copie, P. Suret, and S. Randoux, Numerical spectral synthesis of breather gas for the focusing nonlinear Schrödinger equation, *Phys. Rev. E* 103, 042205 (2021).
- [52] P. Suret, A. Tikan, F. Bonnefoy, F. Copie, G. Ducrozet, A. Gelash, G. Prabhudesai, G. Michel, A. Cazaubiel, E. Falcon, G. A. El, and S. Randoux, Nonlinear spectral synthesis of soliton gas in deep-water surface gravity waves, *Phys. Rev. Lett.* 125, 264101 (2020).
- [53] A. Mussot, A. Kudlinski, M. Kolobov, E. Louvergneaux, M. Douay, and M. Taki, Observation of extreme temporal events in CW-pumped supercontinuum, *Opt. Exp.* 17, 17010 (2009).
- [54] D. R. Solli, C. Ropers, P. Koonath, and B. Jalali, Optical rogue waves, *Nature* 450, 1054-1057 (2007).
- [55] J. M. Dudley, F. Dias, M. Erkintalo, and G. Genty, Instabilities, breathers and rogue waves in optics, *Nat. Photon.* 8, 755 (2014).
- [56] I. Redor, E. Barthélemy, H. Michallet, M. Onorato, and N. Mordant, Experimental evidence of a hydrodynamic soliton gas, *Phys. Rev. Lett.* 122, 214502 (2019).
- [57] M. Girotti, T. Grava, R. Jenkins, and K. T. R. McLaughlin, Rigorous asymptotics of a KdV soliton gas, *Commun. Math. Phys.* 384, 733-784 (2021).
- [58] M. Girotti, T. Grava, R. Jenkins, K. T. R. McLaughlin, and A. Minakov, Soliton versus the gas: Fredholm determinants, analysis, and the rapid oscillations behind the kinetic equation, *Comm. Pure Appl. Math.* 76, 3233-3299 (2023).
- [59] M. Bertola, T. Grava, and G. Orsatti, Soliton shielding of the focusing nonlinear Schrödinger equation, *Phys. Rev. Lett.* 130, 127201 (2023).
- [60] G. Biondini and G. Kovačič, Inverse scattering transform for the focusing nonlinear Schrödinger equation with nonzero boundary conditions, *J. Math. Phys.* 55, 031506 (2014).
- [61] E. A. Kuznetsov, Solitons in parametrically unstable plasma, *Dokl. Akad. Nauk SSR* 22, 507 (1977).
- [62] Y. C. Ma, The perturbed plane-wave solutions of the cubic Schrödinger equation, *Stud. Appl. Math.* 60, 43 (1979).
- [63] N. Akhmediev and V. I. Korreev, Modulation instability and periodic solutions of the nonlinear Schrödinger equation *Theor. Math. Phys.* 69, 1089 (1986).
- [64] D. Peregrine, Water waves, nonlinear Schrödinger equations and their solutions, *J. Aust. Math. Soc. B: Appl. Math.* 25, 16 (1983).
- [65] M. J. Ablowitz, G. Biondini, and B. Prinari, Inverse scattering transform for the integrable discrete nonlinear Schrödinger equation with nonvanishing boundary conditions, *Inverse Prob.* 23, 1711-1758 (2007).
- [66] G. Biondini and B. Prinari, On the spectrum of the Dirac operator and the existence of discrete eigenvalues for the defocusing nonlinear Schrödinger equation, *Stud. Appl. Math.* 132, 138-159 (2014).
- [67] D. Kraus, G. Biondini, and G. Kovačič, The focusing Manakov system with nonzero boundary conditions, *Nonlinearity* 28, 3101-3151 (2015).
- [68] G. Biondini and D. Kraus, Inverse scattering transform for the defocusing Manakov system with nonzero boundary conditions, *SIAM J. Math. Anal.* 47, 706-757 (2015).
- [69] G. Biondini, D. K. Kraus, and B. Prinari, The three-component defocusing nonlinear Schrödinger equation with nonzero boundary conditions, *Commun. Math. Phys.* 348, 475-533 (2016).
- [70] B. Prinari, F. Demontis, S. Li, and T. P. Horikis, Inverse scattering transform and soliton solutions for square matrix nonlinear Schrödinger equations with non-zero boundary conditions, *Physica D* 368, 22-49 (2018).
- [71] P. Deift and X. Zhou, A steepest descent method for oscillatory Riemann-Hilbert problems, *Asymptotics for the MKdV equation*, *Ann. of Math.* 137, 295 (1993).
- [72] X. Zhou, The Riemann-Hilbert problem and inverse scattering, *SIAM J. Math. Anal.* 20, 966 (1989).
- [73] P. Deift and X. Zhou, Long-time asymptotics for solutions of the NLS equation with initial data in a weighted Sobolev space, *Comm. Pure Appl. Math.* 56, 1029-1077 (2003).

- [74] A. H. Vartanian, Long-time asymptotics of solutions to the Cauchy problem for the defocusing nonlinear Schrödinger equation with finite-density initial data. II. Dark solitons on continua. *Math. Phys. Anal. Geom.* 5, 319-413 (2002).
- [75] A. Boutet de Monvel, A. Its, and V. Kotlyarov, Long-time asymptotics for the focusing NLS equation with time-periodic boundary condition on the half-line, *Commun. Math. Phys.* 290, 479-522 (2009).
- [76] K. T. R. McLaughlin and P. D. Miller, The  $\bar{\partial}$ -steepest descent method for orthogonal polynomials on the real line with varying weights, *Int. Math. Res. Not.* 2008, 075 (2008).
- [77] M. Dieng, K. D. T. McLaughlin, Dispersive asymptotics for linear and integrable equations by the Dbar steepest descent method, *Nonlinear dispersive partial differential equations and inverse scattering*, 253-291, *Fields Inst. Comm.*, 83, Springer, New York, 2019
- [78] S. Cuccagna, R. Jenkins, On asymptotic stability of N-solitons of the defocusing nonlinear Schrödinger equation, *Comm. Math. Phys.* 343, 921-969 (2016).
- [79] M. Borghese, R. Jenkins, K.D.T-R McLaughlin, Long time asymptotics behavior of the focusing nonlinear Schrödinger equation, *Ann. I. H. Poincaré*, 35, 887-920 (2018).
- [80] R. Jenkins, J. Liu, P. Perry, C. Sulem, Global well-posedness for the derivative nonlinear Schrödinger equation, *Commun. Math. Phys.* 363, 1003-1049 (2018).
- [81] K. Grunert and G. Teschl, Long-time asymptotics for the Korteweg de Vries equation via nonlinear steepest descent, *Math. Phys. Anal. Geom.* 12, 287-324 (2009).
- [82] D. Bilman and P. D. Miller, A robust inverse scattering transform for the focusing nonlinear Schrödinger equation, *Comm. Pure Appl. Math.* 72, 1722 (2019).
- [83] D. Bilman and R. Buckingham, Large-order asymptotics for multiple-pole solitons of the focusing nonlinear Schrödinger equation. *J. Nonlinear Sci.* 29, 2185-2229 (2019).
- [84] D. Bilman, R. Buckingham, and D. Wang, Far-field asymptotics for multiple-pole solitons in the large-order limit, *J. Differ. Equ.* 297, 320 (2021).
- [85] D. Bilman, L. Ling, and P. D. Miller, Extreme superposition: rogue waves of infinite order and the Painlevé-III hierarchy, *Duke Math. J.* 169, 671-760 (2020).
- [86] A. Romero-Ros, G. C. Katsimiga, S. I. Mistakidis, S. Mossman, G. Biondini, P. Schmelcher, P. Engels, and P. G. Kevrekidis, Experimental realization of the Peregrine soliton in repulsive two-component Bose-Einstein condensates, *Phys. Rev. Lett.* 132, 033402 (2024).
- [87] O. V. Marchukov, B. A. Malomed, M. Olshani, J. Ruhl, V. Dunjko, R. G. Hulet, and V. A. Yurovsky, Quantum fluctuations of the center-of-mass and relative parameters of NLS breathers, *Phys. Rev. Lett.* 125, 050405 (2020).