

APPROXIMATIONS BY SPECIAL VALUES OF MULTIPLE COSINE AND SINE FUNCTIONS

SU HU AND MIN-SOO KIM

ABSTRACT. Kurokawa and Koyama's multiple cosine function $\mathcal{C}_r(x)$ and Kurokawa's multiple cosine function $S_r(x)$ are generalizations of the classical cosine and sine functions from their infinite product representations, respectively. For any fixed $x \in [0, \frac{1}{2})$, let

$$B = \left\{ \frac{\log \mathcal{C}_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\}$$

and

$$C = \left\{ \frac{\log S_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\}$$

be the sets of special values of $\mathcal{C}_r(x)$ and $S_r(x)$ at x , respectively.

In this paper, we will show that the real numbers can be strongly approximated by linear combinations of elements in B and C respectively, with rational coefficients. Furthermore, let

$$D = \left\{ \frac{\zeta_E(3)}{\pi^2}, \frac{\zeta_E(5)}{\pi^4}, \dots, \frac{\zeta_E(2k+1)}{\pi^{2k}}, \dots; \frac{\beta(4)}{\pi^3}, \frac{\beta(6)}{\pi^5}, \dots, \frac{\beta(2k+2)}{\pi^{2k+1}}, \dots \right\}$$

be the set of special values of Dirichlet's eta and beta functions. We will prove that the set D has a similar approximation property, where the coefficients are values of the derivatives of rational polynomials. Our approaches are inspired by recent works of Alkan [1] and Lupu-Wu [14] as applications of the trigonometric integrals.

1. INTRODUCTION

The Riemann zeta function is defined by

$$\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}, \quad (1.1)$$

for $\operatorname{Re}(s) > 1$. Let

$$A = \{\zeta(3), \zeta(5), \dots, \zeta(2k+1), \dots\}$$

be the set of odd zeta values. For $x \in \mathbb{R}$, let $\lfloor x \rfloor = \max\{m \in \mathbb{Z} \mid m \leq x\}$ and $\lceil x \rceil = \min\{m \in \mathbb{Z} \mid m \geq x\}$. We have the following formula connecting the trigonometric integral $\int_0^{2\pi} t^r \log\left(2 \sin \frac{t}{2}\right) dt$ and the elements in A :

$$\int_0^{2\pi} t^r \log\left(2 \sin \frac{t}{2}\right) dt = \sum_{k=1}^{\lfloor r/2 \rfloor} (-1)^k \frac{r!(2\pi)^{r-2k+1}}{(r-2k+1)!} \zeta(2k+1), \quad (1.2)$$

2010 *Mathematics Subject Classification.* 11M06, 41A50.

Key words and phrases. Multiple cosine and sine functions, Zeta value, Trigonometric integral, Approximation.

where $r = 2, 3, 4, \dots$ (see [1, p. 3748, (2.9)]). From the integral (1.2), Alkan [1, Theorem 1] proved that the real numbers can be strongly approximated by linear combinations of elements in A with coefficients in $\mathbb{Q}[\pi^s : 1 \leq s \leq r - 2k + 1]$, where $\mathbb{Q}[\pi^s : 1 \leq s \leq n]$ is the \mathbb{Q} -vector space with basis π^s for $1 \leq s \leq n$. His result is an analogy of the classical Diophantine approximation of Liouville numbers by rational numbers, that is, a real number α is named a Liouville number, if for every positive integer n , there exists a pair of integers (m, n) with $n > 1$ such that

$$0 < \left| \alpha - \frac{m}{n} \right| < \frac{1}{n^q}.$$

It is also a reminiscent of a recent result by Brown and Zudilin [2], which constructed infinitely many *effective* rational approximations $\frac{m}{n}$ to $\zeta(5)$ satisfying

$$0 < \left| \zeta(5) - \frac{m}{n} \right| < \frac{1}{n^{0.86}}.$$

Let

$$A' = \left\{ \frac{\zeta(3)}{\pi^3}, \frac{\zeta(5)}{\pi^5}, \dots, \frac{\zeta(2k+1)}{\pi^{2k+1}}, \dots \right\}.$$

Recently, applying the following trigonometric integral by Orr ([15], also see [14, Lemma 2.1]):

$$\int_0^{2\pi} t^r \cot(t) dt = \left(\frac{\pi}{2} \right)^r \left(\log 2 + \sum_{k=1}^{\lfloor r/2 \rfloor} \frac{r!(-1)^k(4^k - 1)}{(r - 2k)!(2\pi)^{2k}} \zeta(2k + 1) \right) \\ + \delta_{\lfloor r/2 \rfloor, r/2} \frac{r!(-1)^{r/2} \zeta(r + 1)}{\pi^r} \quad (1.3)$$

for $r = 1, 2, 3, \dots$, Lupu and Wu [14] improved Alkan's result to obtain the following approximation, which shows that the real numbers can be strongly approximated by linear combinations of elements in A' , where the coefficients are values of the derivatives of rational polynomials.

Theorem 1.1 ([14, Theorem 1.1]). *Let k_0 and q be positive integers, let α be a real number. For any given integer $n \geq 3$, there exists an integer $r \leq n$ and a rational polynomial $P_n(t) = t^{2k_0}(1 - t)^{2k_0} s_r(t)$ with degree $4k_0 + r$ depending on α, n, q and k_0 , satisfying*

$$\left| \alpha - \sum_{k=k_0}^{2k_0 + \lfloor \frac{r}{2} \rfloor} c_k \frac{\zeta(2k+1)}{\pi^{2k+1}} \right| \ll_{\alpha, q, k_0} \frac{1}{n^q}$$

when $r \geq 2$, where

$$c_k = (-1)^k 2 \left[P_n^{(2k)}(1) \left(1 - \frac{1}{4^k} \right) + P_n^{(2k)}(0) \right]$$

for all $k_0 \leq k \leq 2k_0 + \lfloor \frac{r}{2} \rfloor$.

In fact, there is a long history for investigating the connections between zeta values and the trigonometric integrals such as (1.2) and (1.3). Let

$$\lambda(s) = \sum_{n=0}^{\infty} \frac{1}{(2n+1)^s}, \quad \operatorname{Re}(s) > 1 \quad (1.4)$$

be Dirichlet's lambda function (see [4, p. 954, (1.9)]), which was studied by Euler under the notation $N(s)$ (see [16, p. 70]). Long time ago, Euler has obtained a formula of $\lambda(3)$:

$$\lambda(3) = 1 + \frac{1}{3^3} + \frac{1}{5^3} + \cdots = \frac{\pi^2}{\log 2} + 2 \int_0^{\frac{\pi}{2}} t \log(\sin t) dt \quad (1.5)$$

(see [16, p. 63]), which connects to the problem of finding an explicit formula for $\zeta(3)$ (see [5, section 1.2] for a historical overview). Generalizing (1.5), for any $0 \leq x < \pi$ and $r = 2, 3, 4, \dots$, Koyama and Kurokawa [6, Theorem 1] showed that

$$\int_0^x t^{r-2} \log(\sin t) dt = \frac{x^{r-1}}{r-1} \log(\sin x) - \frac{\pi^{r-1}}{r-1} \log \mathcal{S}_r\left(\frac{x}{\pi}\right). \quad (1.6)$$

Here $\mathcal{S}_r(x)$ is Kurokawa's multiple sine function of order $r = 2, 3, 4, \dots$ (see [7, 8, 9]):

$$\mathcal{S}_r(x) = \exp\left(\frac{x^{r-1}}{r-1}\right) \prod_{n=1}^{\infty} \left\{ P_r\left(\frac{x}{n}\right) P_r\left(-\frac{x}{n}\right)^{(-1)^{r-1}} \right\}^{n^{r-1}}, \quad (1.7)$$

where

$$P_r(x) = (1-x) \exp\left(x + \frac{x^2}{2} + \cdots + \frac{x^r}{r}\right). \quad (1.8)$$

By letting $r = 1$, we recover the infinite product representation of the classical sine function:

$$\mathcal{S}_1(x) = 2 \sin(\pi x) = 2\pi x \prod_{n=1}^{\infty} \left(1 - \frac{x^2}{n^2}\right)$$

and by letting $r = 2$, we get Hölder's double sine function $\mathcal{S}_2(x)$ (see [3]):

$$\mathcal{S}_2(x) = e^x \prod_{n=1}^{\infty} \left\{ \left(\frac{1 - \frac{x}{n}}{1 + \frac{x}{n}} \right)^n e^{2x} \right\}. \quad (1.9)$$

Recently, we have proved a cosine counterpart integral of (1.6):

$$\int_0^x t^{r-2} \log\left(\cos \frac{t}{2}\right) dt = \frac{x^{r-1}}{r-1} \log\left(\cos \frac{x}{2}\right) - \frac{(2\pi)^{r-1}}{r-1} \log \mathcal{C}_r\left(\frac{x}{2\pi}\right) \quad (1.10)$$

(see [5, Theorem 2.1]). Here $\mathcal{C}_r(x)$ is Kurokawa and Koyama's multiple cosine function or order $r = 2, 3, 4, \dots$:

$$\begin{aligned} \mathcal{C}_r(x) &= \prod_{n=-\infty, n:\text{odd}}^{\infty} P_r\left(\frac{x}{\frac{n}{2}}\right)^{\left(\frac{n}{2}\right)^{r-1}} \\ &= \prod_{n=1, n:\text{odd}}^{\infty} \left\{ P_r\left(\frac{x}{\frac{n}{2}}\right) P_r\left(-\frac{x}{\frac{n}{2}}\right)^{(-1)^{r-1}} \right\}^{\left(\frac{n}{2}\right)^{r-1}}. \end{aligned} \quad (1.11)$$

(see [11], [12] and [13]). By letting $r = 1$, we recover the infinite product representation of the classical cosine function:

$$\mathcal{C}_1(x) = 2 \cos(\pi x) = 2 \prod_{n=1, n:\text{odd}}^{\infty} \left(1 - \frac{x^2}{\left(\frac{n}{2}\right)^2}\right). \quad (1.12)$$

And by letting $r = 2, 3, 4$, we get the following examples, respectively:

$$\begin{aligned} \mathcal{C}_2(x) &= \prod_{n=1, n:\text{odd}}^{\infty} \left\{ \left(\frac{1 - \frac{x}{\frac{n}{2}}}{1 + \frac{x}{\frac{n}{2}}} \right)^{\frac{n}{2}} e^{2x} \right\}, \\ \mathcal{C}_3(x) &= \prod_{n=1, n:\text{odd}}^{\infty} \left\{ \left(1 - \frac{x^2}{\left(\frac{n}{2}\right)^2}\right)^{\left(\frac{n}{2}\right)^2} e^{x^2} \right\}, \\ \mathcal{C}_4(x) &= \prod_{n=1, n:\text{odd}}^{\infty} \left\{ \left(\frac{1 - \frac{x}{\frac{n}{2}}}{1 + \frac{x}{\frac{n}{2}}} \right)^{\left(\frac{n}{2}\right)^3} e^{\frac{n^2}{2}x + \frac{2}{3}x^3} \right\} \end{aligned} \quad (1.13)$$

(see [10, 11, 13]).

Furthermore, by applying the integrals

$$\int_0^x t^{r-2} \log(\sin t) dt \quad \text{and} \quad \int_0^x t^{r-2} \log\left(\cos \frac{t}{2}\right) dt,$$

we can show that the special values of multiple sine and cosine functions will be expressed by the zeta values. For example, [5, Theorem 2.4] gives

$$\begin{aligned} \log \mathcal{C}_r\left(\frac{1}{4}\right) &= \frac{\log 2}{2^{2r-1}} - \frac{(r-1)!}{(2\pi)^{r-1}} \sin\left(\frac{r\pi}{2}\right) \zeta_E(r) \\ &\quad - \frac{r-1}{2^{2(r-1)}} \sum_{k=0}^{\lfloor \frac{r-2}{2} \rfloor} (-1)^k (2k)! \binom{r-2}{2k} \left(\frac{2}{\pi}\right)^{2k+1} \\ &\quad \times \beta(2k+2) \\ &\quad - \frac{r-1}{2^{2r-1}} \sum_{k=1}^{\lceil \frac{r-2}{2} \rceil} \frac{(-1)^{k-1} (2k-1)!}{\pi^{2k}} \binom{r-2}{2k-1} \\ &\quad \times \zeta_E(2k+1) \end{aligned} \quad (1.14)$$

for $r = 2, 3, 4, \dots$. Here

$$\zeta_E(s) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^s}, \quad \operatorname{Re}(s) > 0 \quad (1.15)$$

is the alternating zeta function (also known as Dirichlet's eta or Euler's eta function) and

$$\beta(s) = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^s}, \quad \operatorname{Re}(s) > 0 \quad (1.16)$$

is the alternating form of Dirichlet's lambda function (1.4) (also known as Dirichlet's beta function), which was studied by Euler under the notation $L(s)$ (see [16, p. 70]). Let

$$G = \sum_{n=0}^{\infty} \frac{(-1)^n}{(2n+1)^2} = 0.915965594177219015 \dots$$

be the Catalan constant. As an example, if setting $r = 3$ in (1.14), then we get

$$\log \mathcal{C}_3 \left(\frac{1}{4} \right) = \frac{\log 2}{32} - \frac{G}{4\pi} + \frac{7\zeta_E(3)}{16\pi^2},$$

which is equivalent to a formula of $\zeta(3)$:

$$\zeta(3) = \frac{4\pi^2}{21} \log \left(\frac{e^{\frac{4G}{\pi}} \mathcal{C}_3 \left(\frac{1}{4} \right)^{16}}{\sqrt{2}} \right)$$

(see [5, Corollary 2.7]).

Now for any $0 \leq x < \frac{1}{2}$, let

$$B = \left\{ \frac{\log \mathcal{C}_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\}$$

and

$$C = \left\{ \frac{\log S_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\}$$

be the sets of special values of multiple cosine and sine functions at x , respectively. In this paper, by applying the trigonometric integrals (see Lemmas 2.1 and 4.1)

$$\int_0^x t^r \tan(\pi t) dt \quad \text{and} \quad \int_0^x t^r \cot \left(\frac{\pi t}{2} \right) dt,$$

we show that for any fixed $x \in [0, \frac{1}{2})$, the real numbers can be strongly approximated by linear combinations of elements in B and C respectively, with rational coefficients (see Sections 2 and 4). Furthermore, by considering the integral

$$\int_0^1 t^{r-2} \log \left(\cos \frac{\pi t}{4} \right) dt$$

(see Lemma 3.1), we prove that the set of zeta and beta values

$$D = \left\{ \frac{\zeta_E(3)}{\pi^2}, \frac{\zeta_E(5)}{\pi^4}, \dots, \frac{\zeta_E(2k+1)}{\pi^{2k}}, \dots; \frac{\beta(4)}{\pi^3}, \frac{\beta(6)}{\pi^5}, \dots, \frac{\beta(2k+2)}{\pi^{2k+1}}, \dots \right\}$$

has a similar approximation property, where the coefficients are values of the derivatives of rational polynomials (see Section 3).

2. THE INTEGRAL $\int_0^x t^r \tan(\pi t) dt$

For any fixed $x \in [0, \frac{1}{2})$, let

$$B = \left\{ \frac{\log \mathcal{C}_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\}.$$

In this section, we will show that the real numbers can be approximated by linear combinations of elements in B with rational coefficients. First, we need the following lemmas.

Lemma 2.1. *For $0 \leq x < \frac{1}{2}$ and $r = 1, 2, 3, \dots$, we have*

$$\int_0^x t^r \tan(\pi t) dt = -\frac{\log \mathcal{C}_{r+1}(x)}{\pi}. \quad (2.1)$$

Proof. By [5, Proposition 3.3], for $0 \leq x < 1$ and $r = 2, 3, 4, \dots$, we have

$$\begin{aligned} \log \mathcal{C}_r\left(\frac{x}{2}\right) &= -\frac{1}{2^r} \int_0^x \pi t^{r-1} \tan\left(\frac{\pi t}{2}\right) dt \\ &= -\pi \int_0^x \left(\frac{t}{2}\right)^{r-1} \tan\left(\frac{\pi t}{2}\right) d\left(\frac{t}{2}\right) \\ &= -\pi \int_0^{\frac{x}{2}} t^{r-1} \tan(\pi t) dt, \end{aligned} \quad (2.2)$$

which is equivalent to

$$\int_0^x t^r \tan(\pi t) dt = -\frac{\log \mathcal{C}_{r+1}(x)}{\pi}$$

for $0 \leq x < \frac{1}{2}$ and $r = 1, 2, 3, \dots$ □

Lemma 2.2. *For any $x \in [0, \frac{1}{2})$ and a positive integer k_0 . Let α be a real number. Then there exists a function $f_\alpha \in C^\infty[0, x]$ such that*

$$\alpha = \int_0^x f_\alpha(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt.$$

Proof. Fixed an $x \in [0, \frac{1}{2})$, define a functional on $C^\infty[0, x]$ by

$$T_{x, k_0}(f(t)) = \int_0^x f(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt, \quad (2.3)$$

for $f(t) \in C^\infty[0, x]$. Since

$$\int_0^x t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt = K_{x, k_0} > 0,$$

we have

$$\begin{aligned} |T_{x, k_0}(f(t))| &= \left| \int_0^x f(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt \right| \\ &\leq K_{x, k_0} \sup_{t \in [0, x]} |f(t)| \end{aligned} \quad (2.4)$$

for $f(t) \in C^\infty[0, x]$, thus $T_{x, k_0}(f(t))$ is a bounded linear functional under the supremum metric. Then fixed a small $\delta > 0$, define a smooth function

$$f_1(t) = \begin{cases} M, & t \in [\frac{x}{2} - \delta, \frac{x}{2} + \delta] \\ 0, & t \in [0, \frac{x}{2} - 2\delta] \cup [\frac{x}{2} + 2\delta, x]. \end{cases} \quad (2.5)$$

Since

$$\inf_{t \in [\frac{x}{2} - \delta, \frac{x}{2} + \delta]} \{t^{2k_0}(1-t)^{2k_0} \tan(\pi t)\} = L_{x, k_0} > 0,$$

we have

$$\begin{aligned} T_{x, k_0}(f_1(t)) &= \int_0^x f_1(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt \\ &\geq 2M\delta L_{x, k_0}. \end{aligned} \quad (2.6)$$

Similarly, define a smooth function

$$f_2(t) = \begin{cases} N, & t \in [\frac{x}{2} - \delta, \frac{x}{2} + \delta] \\ 0, & t \in [0, \frac{x}{2} - 2\delta] \cup [\frac{x}{2} + 2\delta, x], \end{cases} \quad (2.7)$$

we have

$$\begin{aligned} T_{x, k_0}(f_2(t)) &= \int_0^x f_2(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt \\ &\leq NK_{x, k_0}. \end{aligned} \quad (2.8)$$

For any $\alpha > 0$, taking $N > 0$ small enough such that

$$NK_{x, k_0} < \alpha \quad (2.9)$$

and taking $M > 0$ big enough such that

$$2M\delta L_{x, k_0} > \alpha. \quad (2.10)$$

For any $0 \leq \lambda \leq 1$, define

$$f^{(\lambda)}(t) = \lambda f_1(t) + (1 - \lambda) f_2(t).$$

Obviously, it is in $C^\infty[0, x]$. Thus for $\lambda \in [0, 1]$,

$$g(\lambda) = T_{x, k_0}(f^{(\lambda)}(t))$$

becomes a smooth function from $[0, 1]$ to \mathbb{R} . Since by (2.9) and (2.10) we have

$$T_{x, k_0}(f_2(t)) < \alpha < T_{x, k_0}(f_1(t)),$$

from the intermediate value theorem, there exists an $\lambda \in (0, 1)$ such that $f^{(\lambda)}(t)$ satisfying

$$\alpha = T_{x, k_0}(f^{(\lambda)}(t)) = \int_0^x f^{(\lambda)}(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt.$$

Denote $f_\alpha(t) = f^{(\lambda)}(t)$. If $\alpha < 0$, then $-\alpha > 0$ and from the above result there exists a $f_{-\alpha}(t) \in C^\infty[0, x]$ such that

$$-\alpha = T_{x, k_0}(f_{-\alpha}(t)).$$

Let $f_\alpha = -f_{-\alpha}$, then

$$T_{x, k_0}(f_\alpha(t)) = -T_{x, k_0}(f_{-\alpha}(t)) = \alpha$$

satisfying our requirement. If $\alpha = 0$, taking $f_\alpha(t) \equiv 0$ for $t \in [0, x]$, then we have

$$T_{x,k_0}(f_\alpha(t)) = 0 = \alpha.$$

We thus obtain Lemma 2.2. \square

Now we are at the position to prove the approximation property of the set B .

Theorem 2.3. *Fixed an $x \in [0, \frac{1}{2}]$. Let k_0 and q be positive integers, let α be a real number. For any given integer $n \geq 3$, there exists an integer $r \leq n$ and a rational polynomial $P_n(t)$ with degree $4k_0 + r$ depending on α, n, q, k_0 and x :*

$$P_n(t) = a_{2k_0}t^{2k_0} + \cdots + a_{4k_0+r}t^{4k_0+r},$$

satisfying

$$\left| \alpha - \sum_{k=2k_0}^{4k_0+r} c_k \frac{\log \mathcal{C}_{k+1}(x)}{\pi} \right| \ll_{\alpha, q, k_0, x} \frac{1}{n^q},$$

where

$$c_k = -a_k.$$

Proof. Let $n \geq 3$ be a given integer. For $f(t) \in C^\infty[0, x]$, by a classical theorem of Jackson (see [1, p. 3749, (2.16)]), there exists a rational polynomial of degree $r \leq n$, whose coefficients depend only on f, n, q and x :

$$s_r(t) = a_0 + a_1t + \cdots + a_rt^r,$$

such that

$$\|f - s_r\| = \sup_{t \in [0, x]} |f(t) - s_r(t)| \ll_{f, q, x} \frac{1}{n^q}. \quad (2.11)$$

And by Lemma 2.2, there is a function $f_\alpha(t) \in C^\infty[0, x]$ such that

$$\alpha = \int_0^x f_\alpha(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt.$$

Let

$$\begin{aligned} P_n(t) &= t^{2k_0} (1-t)^{2k_0} s_r(t) \\ &= a_{2k_0} t^{2k_0} + \cdots + a_{4k_0+r} t^{4k_0+r}. \end{aligned} \quad (2.12)$$

Then applying (2.11) to $f_\alpha(t)$, we get

$$\begin{aligned} & \left| \alpha - \int_0^x P_n(t) \tan(\pi t) dt \right| \\ &= \left| \int_0^x f_\alpha(t) t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt - \int_0^x P_n(t) \tan(\pi t) dt \right| \\ &\leq \int_0^x |f_\alpha(t) - s_r(t)| t^{2k_0} (1-t)^{2k_0} \tan(\pi t) dt \\ &\ll_{\alpha, q, k_0, x} \frac{1}{n^q}. \end{aligned} \quad (2.13)$$

Now we calculate the integral $\int_0^x P_n(t) \tan(\pi t) dt$. By Lemma 2.1, we have

$$\begin{aligned} \int_0^x P_n(t) \tan(\pi t) dt &= \sum_{k=2k_0}^{4k_0+r} a_k \int_0^x t^k \tan(\pi t) dt \\ &= \sum_{k=2k_0}^{4k_0+r} \left(-\frac{a_k}{\pi} \right) \log \mathcal{C}_{k+1}(x). \end{aligned} \quad (2.14)$$

Combing with (2.12) we get

$$\left| \alpha - \sum_{k=2k_0}^{4k_0+r} c_k \frac{\log \mathcal{C}_{k+1}(x)}{\pi} \right| \ll_{\alpha, q, k_0, x} \frac{1}{n^q},$$

where

$$c_k = -a_k.$$

□

3. THE INTEGRAL $\int_0^1 t^r \log \left(\cos \frac{\pi t}{4} \right) dt$

In [5, Theorem 2.3], we have calculated the integral $\int_0^x t^{r-2} \log \left(\cos \frac{t}{2} \right) dt$ for $r = 2, 3, 4, \dots$ as follows

$$\begin{aligned} \int_0^{\frac{\pi}{2}} t^{r-2} \log \left(\cos \frac{t}{2} \right) dt &= -\frac{\log 2}{r-1} \left(\frac{\pi}{2} \right)^{r-1} + (r-2)! \sin \left(\frac{r\pi}{2} \right) \zeta_E(r) \\ &\quad + \sum_{k=0}^{\lfloor \frac{r-2}{2} \rfloor} (-1)^k (2k)! \binom{r-2}{2k} \left(\frac{\pi}{2} \right)^{r-2k-2} \\ &\quad \times \beta(2k+2) \\ &\quad + \sum_{k=1}^{\lfloor \frac{r-2}{2} \rfloor} \frac{(-1)^{k-1} (2k-1)!}{2^{2k+1}} \binom{r-2}{2k-1} \left(\frac{\pi}{2} \right)^{r-2k-1} \\ &\quad \times \zeta_E(2k+1). \end{aligned} \quad (3.1)$$

Recall that $\zeta_E(s)$ and $\beta(s)$ are Dirichlet's eta and beta functions, respectively (see (1.15) and (1.16)).

Let

$$D = \left\{ \frac{\zeta_E(3)}{\pi^2}, \frac{\zeta_E(5)}{\pi^4}, \dots, \frac{\zeta_E(2k+1)}{\pi^{2k}}, \dots; \frac{\beta(4)}{\pi^3}, \frac{\beta(6)}{\pi^5}, \dots, \frac{\beta(2k+2)}{\pi^{2k+1}}, \dots \right\}$$

be the set of zeta and beta values. In this section, we will show that the real numbers can be strongly approximated by linear combinations of elements in D , where the coefficients are values of the derivatives of rational polynomials. First, we state the following integral representation which is the $\log \left(\cos \frac{\pi t}{4} \right)$ -counterpart of Lemma 2.1.

Lemma 3.1. *For $r = 0, 1, 2, \dots$, we have*

$$\begin{aligned} \int_0^1 t^r \log \left(\cos \frac{\pi t}{4} \right) &= -\frac{\log 2}{r+1} - r! \sin \left(\frac{r\pi}{2} \right) 2^{r+1} \frac{\zeta_E(r+2)}{\pi^{r+1}} \\ &+ \sum_{k=0}^{\lfloor \frac{r}{2} \rfloor} (-1)^k (2k)! \binom{r}{2k} 2^{2k+1} \frac{\beta(2k+2)}{\pi^{2k+1}} \\ &+ \frac{1}{2} \sum_{k=1}^{\lfloor \frac{r}{2} \rfloor} (-1)^k (2k-1)! \binom{r}{2k-1} \frac{\zeta_E(2k+1)}{\pi^{2k}}. \end{aligned} \quad (3.2)$$

Proof. By (3.1), for $r = 2, 3, 4, \dots$ we have

$$\begin{aligned} \int_0^1 t^{r-2} \log \left(\cos \frac{\pi t}{4} \right) dt &= \left(\frac{2}{\pi} \right)^{r-1} \int_0^{\frac{\pi}{2}} t^{r-2} \log \left(\cos \frac{t}{2} \right) dt \\ &= \left(\frac{2}{\pi} \right)^{r-1} \left(-\frac{\log 2}{r-1} \left(\frac{\pi}{2} \right)^{r-1} + (r-2)! \sin \left(\frac{r\pi}{2} \right) \zeta_E(r) \right. \\ &+ \sum_{k=0}^{\lfloor \frac{r-2}{2} \rfloor} (-1)^k (2k)! \binom{r-2}{2k} \left(\frac{\pi}{2} \right)^{r-2k-2} \beta(2k+2) \\ &+ \left. \sum_{k=1}^{\lfloor \frac{r-2}{2} \rfloor} \frac{(-1)^{k-1} (2k-1)!}{2^{2k+1}} \binom{r-2}{2k-1} \left(\frac{\pi}{2} \right)^{r-2k-1} \zeta_E(2k+1) \right) \\ &= -\frac{\log 2}{r-1} + (r-2)! \sin \left(\frac{r\pi}{2} \right) \left(\frac{2}{\pi} \right)^{r-1} \zeta_E(r) \\ &+ \sum_{k=0}^{\lfloor \frac{r-2}{2} \rfloor} (-1)^k (2k)! \binom{r-2}{2k} \left(\frac{2}{\pi} \right)^{2k+1} \beta(2k+2) \\ &+ \sum_{k=1}^{\lfloor \frac{r-2}{2} \rfloor} \frac{(-1)^{k-1} (2k-1)!}{2^{2k+1}} \binom{r-2}{2k-1} \left(\frac{2}{\pi} \right)^{2k} \zeta_E(2k+1), \end{aligned} \quad (3.3)$$

which is equivalent to

$$\begin{aligned} \int_0^1 t^r \log \left(\cos \frac{\pi t}{4} \right) &= -\frac{\log 2}{r+1} - r! \sin \left(\frac{r\pi}{2} \right) 2^{r+1} \frac{\zeta_E(r+2)}{\pi^{r+1}} \\ &+ \sum_{k=0}^{\lfloor \frac{r}{2} \rfloor} (-1)^k (2k)! \binom{r}{2k} 2^{2k+1} \frac{\beta(2k+2)}{\pi^{2k+1}} \\ &+ \frac{1}{2} \sum_{k=1}^{\lfloor \frac{r}{2} \rfloor} (-1)^k (2k-1)! \binom{r}{2k-1} \frac{\zeta_E(2k+1)}{\pi^{2k}} \end{aligned} \quad (3.4)$$

for $r = 0, 1, 2, \dots$ □

The above lemma implies the following result.

Lemma 3.2. *For any polynomial $P(t) \in \mathbb{C}[t]$ with $P(0) = 0$ and $\deg P = n$, we have*

$$\begin{aligned} \int_0^1 P(t) \log \left(\cos \frac{\pi t}{4} \right) dt &= -\frac{P(1)}{r+1} \log 2 + \sum_{k=0}^{\lfloor \frac{n}{2} \rfloor} (-1)^k P^{(2k)}(1) 2^{2k+1} \frac{\beta(2k+2)}{\pi^{2k+1}} \\ &\quad + \sum_{k=1}^{\lfloor \frac{n}{2} \rfloor} \left[\frac{1}{2} (-1)^k P^{(2k-1)}(1) + P^{(2k-1)}(0) 2^{2k} \right] \\ &\quad \times \frac{\zeta_E(2k+1)}{\pi^{2k}}. \end{aligned} \quad (3.5)$$

Then for any $x \in [0, 1)$, considering the following functional on $C^\infty[0, x]$:

$$C_{x, k_0}(f(t)) = \int_0^x f(t) t^{2k_0} (1-t)^{2k_0} \log \left(\cos \frac{\pi t}{4} \right) dt, \quad (3.6)$$

where $f(t) \in C^\infty[0, x]$. For a rational polynomial $P_n(t) = t^{2k_0} (1-t)^{2k_0} s_r(t)$, notice that $P_n^{(2k)}(0) = P_n^{(2k)}(1) = 0$ for $0 \leq k \leq k_0 - 1$, the same reasoning as Theorem 2.3 implies the following result.

Theorem 3.3. *Let k_0 and q be positive integers, let α be a real number. For any given integer $n \geq 3$, there exists an integer $r \leq n$ and a rational polynomial $P_n(t) = t^{2k_0} (1-t)^{2k_0} s_r(t)$ with degree $4k_0 + r$ depending on α, n, q and k_0 , satisfying*

$$\left| \alpha - \sum_{k=k_0}^{2k_0 + \lfloor \frac{r}{2} \rfloor} c_k^\beta \frac{\beta(2k+2)}{\pi^{2k+1}} - \sum_{k=k_0}^{2k_0 + \lceil \frac{r}{2} \rceil} c_k^{\zeta_E} \frac{\zeta_E(2k+1)}{\pi^{2k}} \right| \ll_{\alpha, q, k_0} \frac{1}{n^q}$$

when $r \geq 2$, where

$$c_k^\beta = (-1)^k P_n^{(2k)}(1) 2^{2k+1}$$

for all $k_0 \leq k \leq 2k_0 + \lfloor \frac{r}{2} \rfloor$, and

$$c_k^{\zeta_E} = \frac{1}{2} (-1)^k P_n^{(2k-1)}(1) + P_n^{(2k-1)}(0) 2^{2k}$$

all $k_0 \leq k \leq 2k_0 + \lceil \frac{r}{2} \rceil$.

4. THE INTEGRAL $\int_0^x t^r \cot \left(\frac{\pi t}{2} \right) dt$

In this section, we will consider the cotangent counterpart of section 2. For any fixed $x \in [0, \frac{1}{2})$, let

$$C = \left\{ \frac{\log S_r(x)}{\pi} \mid r = 1, 2, 3, \dots \right\},$$

we will show that the real numbers can be approximated by linear combinations of elements in C with rational coefficients. First, we state the following integral representation which is the cotangent counterpart of Lemma 2.1.

Lemma 4.1. For $0 \leq x < 1$ and $r = 1, 2, 3, \dots$, we have

$$\int_0^x t^r \cot\left(\frac{\pi t}{2}\right) dt = \frac{2^r}{\pi} \log S_r\left(\frac{x}{2}\right). \quad (4.1)$$

Proof. For $0 \leq x < 1$ and $r = 2, 3, 4, \dots$, by [6, Proposition 2], we have

$$\begin{aligned} \frac{\log S_r\left(\frac{x}{2}\right)}{\pi} &= \int_0^{\frac{x}{2}} t^{r-1} \cot(\pi t) dt \\ &= \int_0^x \left(\frac{t}{2}\right)^{r-1} \cot\left(\frac{\pi t}{2}\right) d\left(\frac{t}{2}\right) \\ &= \frac{1}{2^r} \int_0^x t^{r-1} \cot\left(\frac{\pi t}{2}\right) dt, \end{aligned} \quad (4.2)$$

which is equivalent to (4.1). \square

Then fixed an $x \in [0, 1)$, by considering the following functional on $C^\infty[0, x]$:

$$S_{x, k_0}(f(t)) = \int_0^x f(t) t^{2k_0} (1-t)^{2k_0} \cot\left(\frac{\pi t}{2}\right) dt, \quad (4.3)$$

for $f(t) \in C^\infty[0, x]$, the same reasoning as Theorem 2.3 implies the following result.

Theorem 4.2. Fixed an $x \in [0, 1)$. Let k_0 and q be positive integers, let α be a real number. For any given integer $n \geq 3$, there exists an integer $r \leq n$ and a rational polynomial $P_n(t)$ with degree $4k_0 + r$ depending on α, n, q and x :

$$P_n(t) = a_{2k_0} t^{2k_0} + \dots + a_{4k_0+r} t^{4k_0+r},$$

satisfying

$$\left| \alpha - \sum_{k=2k_0}^{4k_0+r} c_k \frac{\log S_{k+1}\left(\frac{x}{2}\right)}{\pi} \right| \ll_{\alpha, q, k_0, x} \frac{1}{n^q},$$

where

$$c_k = 2^k a_k.$$

The following integral concerning the case of $x = 1$ was obtained by Lupu and Wu [14] recently. It can be viewed as a cotangent counterpart of Lemma 3.2.

Lemma 4.3 ([14, Lemma 2.3]). For any polynomial $P(t) \in \mathbb{C}[t]$ with $P(0) = 0$ and $\deg P = n$, we have

$$\begin{aligned} \int_0^1 P(t) \cot\left(\frac{\pi t}{2}\right) dt &= \frac{2P(1)}{\pi} \log 2 + 2 \sum_{k=1}^{\lfloor n/2 \rfloor} (-1)^k P^{(2k)}(1) \cdot \left(1 - \frac{1}{2^{2k}}\right) \\ &\quad \times \frac{\zeta(2k+1)}{\pi^{2k+1}} + 2 \sum_{k=1}^{\lfloor n/2 \rfloor} (-1)^k P^{(2k)}(0) \frac{\zeta(2k+1)}{\pi^{2k+1}}. \end{aligned} \quad (4.4)$$

If considering the following functional on $C^\infty[0, 1]$:

$$\begin{aligned} S_{k_0}(f(t)) &= \int_0^1 f(t)t^{2k_0}(1-t)^{2k_0} \cot\left(\frac{\pi t}{2}\right) dt \\ &= \lim_{x \rightarrow 1} \int_0^x f(t)t^{2k_0}(1-t)^{2k_0} \cot\left(\frac{\pi t}{2}\right) dt, \end{aligned} \quad (4.5)$$

then Lemma 4.3 and the same reasoning as Theorem 3.3 recover Lupu and Wu's result (see Theorem 1.1 above).

REFERENCES

- [1] E. Alkan, *Special values of the Riemann zeta function capture all real numbers*, Proc. Amer. Math. Soc. **143** (2015), no. 9, 3743–3752.
- [2] F. Brown and W. Zudilin, *On cellular rational approximations to $\zeta(5)$* , preprint at <https://arxiv.org/abs/2210.03391>.
- [3] O. Hölder, *Ueber eine transcendente Function*, Göttingen Nachrichten **16** (1886), 514–522.
- [4] S. Hu and M.-S. Kim, *On Dirichlet's lambda function*, J. Math. Anal. Appl. **478** (2019), no. 2, 952–972.
- [5] S. Hu and M.-S. Kim, *Euler's integral, multiple cosine function and zeta values*, Forum Math. **37** (2025), no. 1, 287–308.
- [6] S.-Y. Koyama and N. Kurokawa, *Euler's integrals and multiple sine functions*, Proc. Amer. Math. Soc. **133** (2005), no. 5, 1257–1265.
- [7] N. Kurokawa, *Multiple sine functions and Selberg zeta functions*, Proc. Japan Acad. Ser. A Math. Sci. **67** (1991), no. 3, 61–64.
- [8] N. Kurokawa, *Gamma factors and Plancherel measures*, Proc. Japan Acad. Ser. A Math. Sci. **68** (1992), no. 9, 256–260.
- [9] N. Kurokawa, *Multiple zeta functions: an example*, Zeta functions in geometry (Tokyo, 1990), 219–226, Adv. Stud. Pure Math., **21**, Kinokuniya, Tokyo, 1992.
- [10] N. Kurokawa and S. Koyama, *Multiple sine functions*, Forum Math. **15** (2003), no. 6, 839–876.
- [11] N. Kurokawa and M. Wakayama, *Duplication formulas in triple trigonometry*, Proc. Japan Acad. Ser. A Math. Sci. **79** (2003), no. 8, 123–127.
- [12] N. Kurokawa and M. Wakayama, *Zeta regularized product expressions for multiple trigonometric functions*, Tokyo J. Math. **27** (2004), no. 2, 469–480.
- [13] N. Kurokawa and M. Wakayama, *Extremal values of double and triple trigonometric functions*, Kyushu J. Math. **58** (2004), no. 1, 141–166.
- [14] C. Lupu and D. Wu, *Approximations by special values of the Riemann zeta function via cotangent integrals*, J. Math. Anal. Appl. **545** (2025), no. 2, Article ID 129144, 8 pp.
- [15] D. Orr, *Generalized rational zeta series for $\zeta(2n)$ and $\zeta(2n+1)$* , Integral Transforms Spec. Funct. **28** (2017), no. 12, 966–987.
- [16] V.S. Varadarajan, *Euler through time: a new look at old themes*, American Mathematical Society, Providence, RI, 2006.

DEPARTMENT OF MATHEMATICS, SOUTH CHINA UNIVERSITY OF TECHNOLOGY,
GUANGZHOU, GUANGDONG 510640, CHINA
Email address: mahusu@scut.edu.cn

DEPARTMENT OF MATHEMATICS EDUCATION, KYUNGNAM UNIVERSITY, CHANG-
WON, GYEONGNAM 51767, REPUBLIC OF KOREA
Email address: mskim@kyungnam.ac.kr