ON THE NOWICKI CONJECTURE FOR THE FREE LIE ALGEBRA OF RANK 2

LUCIO CENTRONE, ȘEHMUS FINDIK, MANUELA DA SILVA SOUZA

ABSTRACT. We consider the free Lie algebra generated by two elements x and y, namely $\mathcal{L}(x, y)$ under the action of the Weitzenböck derivation δ sending y to x and x to 0 and we present its algebra of constants as the algebra generated by the integrals of the relations among the generators of a proper subalgebra of $\mathcal{L}(x, y)$.

1. INTRODUCTION

In this paper we deal with the study of the algebra of constants of some special derivations of the free Lie algebra in the variable x, y namely $L := \mathcal{L}(x, y)$.

Given a linear derivation it is not easy to describe its ring of constants even if we assume that its associated matrix (as an endomorphism) is nilpotent. Let K be a field of characteristic zero, $X_m = \{x_1, \ldots, x_m\}$ be a set of variables, and KX_m be the vector space with basis X_m . Now consider a non-zero nilpotent linear operator δ of KX_m . Certainly δ can be extended to a derivation of the polynomial algebra $K[X_m]$, the free algebra of rank m in the class of all commutative unitary algebras. Such derivations are called Weitzenböck due to the classical theorem of Weitzenböck [27], dating back to 1932, which states that the algebra $K[X_m]^{\delta} = \ker \delta$ of constants of the derivation δ in the algebra $K[X_m]$ is finitely generated. It is easy to see that any Weitzenböck derivation δ is locally nilpotent and the linear operator $\exp(\delta)$ which acts on the vector space KX_m is unipotent. Moreover the algebra $K[X_m]^{\delta}$ of constants of δ is equal to the algebra of invariants

$$K[X_m]^{\exp(\delta)} = \{ x \in K[X_m] \mid \exp(\delta)(x) = x \}.$$

The algebra of invariants of $K[X_m]$ under the action of $\exp(\delta)$ is equal to the algebra of invariants $K[X_m]^{UT_2(K)}$ of the unitriangular group $UT_2(K)$ consisting of all elements $\exp(c\delta)$, $c \in K$. Actually the study of the algebra of constants $K[X_m]^{\delta}$ inherits methods from the classical invariant theory. On this purpose it is worthy mentioning the book by Nowicki [21], Derksen and Kemper [10], and Sturmfels [26] concerning numerical aspects of algebras of constants and general invariant theory.

 $^{2010\} Mathematics\ Subject\ Classification.\ 17B01;\ 17B30;\ 17B40;\ 16R10;\ 16S15; 16W25;\ 13N15;\ 13A50.$

Key words and phrases. algebras of constants; Weitzenböck derivations, the Nowicki conjecture.

The first author is partially supported by FAPESP grant 2018/02108-7 and by CNPq.

The research of the second named author was partially supported by Çukurova University (BAP/10746) for Bilateral Scientific Cooperation between Turkey and Brazil.

Indeed the Jordan normal form of Weitzenböck derivations consists of Jordan cells with zero diagonals. Moreover Weitzenböck derivations are in one-to-one correspondence with the partition of m, up to a linear change of variables. Thus there is essentially a finite number of Weitzenböck derivations for a fixed m. In particular, let m = 2d, $d \ge 1$, and assume the Jordan form of δ contains the Jordan cells of size 2×2 only, i.e.,

$$J(\delta) = \begin{pmatrix} 0 & 1 & \cdots & 0 & 0 \\ 0 & 0 & \cdots & 0 & 0 \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ 0 & 0 & \cdots & 0 & 1 \\ 0 & 0 & \cdots & 0 & 0 \end{pmatrix}.$$

We assume that $\delta(x_{2j}) = x_{2j-1}$, $\delta(x_{2j-1}) = 0$, $j = 1, \ldots, d$. Nowicki conjectured in his book [21] (Section 6.9, page 76) that in this case the algebra $K[X_{2d}]^{\delta}$ is generated by $x_1, x_3, \ldots, x_{2d-1}$, and $x_{2i-1}x_{2j} - x_{2i}x_{2j-1}$, $1 \le i < j \le d$.

The conjecture was proved by several authors with different techniques. The first published proof appeared in 2004 by Khoury [17] in his PhD thesis, followed by his paper [18], where he makes use of a computational approach involving Gröbner basis techniques. Derksen and Panyushev applied ideas of classical invariant theory in order to prove the Novicki conjecture but their proof remained unpublished. Later in 2009, Drensky and Makar-Limanov [14] confirmed the conjecture by an elementary proof from undergraduate algebra, without involving invariant theory. Another simple short proof was given by Kuroda [19] in the same year. Bedratyuk [1] proved the Nowicki conjecture by reducing it to a well known problem of classical invariant theory.

The following problem arises naturally. Let δ be a Weitzenböck derivation of the free associative unitary algebra $F_m(\mathfrak{V})$ of rank m in a given variety \mathfrak{V} (or relatively free algebra of rank m of \mathfrak{V}), then give an explicit set of generators of the algebra of constants $F_m^{\delta}(\mathfrak{V})$. Recently, Dangovski et al. [8, 9] gave some results in this direction. They showed that the algebra of constants $F_m^{\delta}(\mathfrak{V})$ is not finitely generated as an algebra except for some cases: when \mathfrak{V} is the variety of metabelian Lie algebras or the variety of metabelian associative algebras. Although this, they gave a finite set of generators of $(F'_d(\mathfrak{V}))^{\delta}$ as a module over the algebra of constants of $K[U_d, V_d]$. Here $F'_d(\mathfrak{V})$ denotes the commutator ideal of $F_d(\mathfrak{V})$ and δ acts on U_d and V_d in the same way as on X_d .

In [13] Drensky and Gupta studied Weitzenböck derivations δ acting on $F_m(\mathfrak{V})$ proving that if the algebra $UT_2(K)$ of 2×2 upper left triangular matrices over a field K of characteristic zero belongs to a variety \mathfrak{V} , then $F_m^{\delta}(\mathfrak{V})$ is not finitely generated whereas if $UT_2(K)$ does not belong to \mathfrak{V} , then $F_m^{\delta}(\mathfrak{V})$ is finitely generated by a result of Drensky in [11].

The goal of this paper goes in this direction but in a Lie setting. We recall by the well known dichotomy a variety of Lie algebras either satisfies the Engel condition (hence it is nilpotent by [29]) or contains the metabelian variety \mathfrak{U}_2 consisting of all solvable Lie algebras of class 2 and is defined by the identity $[[x_1, x_2], [x_3, x_4]] = 0$. Since the finitely generated nilpotent Lie algebras are finite dimensional, the problem for the finite generation of the algebras of constants of relatively free nilpotent Lie algebras is solved trivially. In this paper we presented the algebra of constants of the free Lie algebra in two generators as the algebra generated by the integrals of the relations among the infinite generators of $\mathcal{L}(x, y) - \operatorname{span}_F\{y\}$.

The latter is a reliable 'representation' of $\mathcal{L}^{\delta}(x, y)$. We will call such generators pseudodeterminant due to some own algebraic properties and their similarities with the determinant-like generators used by Makar-Limanov and Drensky in their proof of Novicki's Conjecture for the free commutative algebra. A precious tool used in order to get this result is represented by some integration rules introduced in Section 3. We also make use of a linear basis of the free Lie algebra, the Hall basis in particular and we figure out how do the monomials which are constants are. It turns out they are monomials in the algebra generated by the pseudodeterminant. We finally give some exemple of an explicit set of generators for constants of small degree (up to degree 6).

2. Preliminaries

We start off with some basic facts.

Definition 2.1. If A is any algebra (associative or not) over a field K, a derivation of A is a K-linear map $\delta : A \to A$ so that $\delta(ab) = \delta(a)b + a\delta(b)$ for every $a, b \in A$.

Of course, every derivation of a free algebra is completely determined by the value it takes on its generators. Moreover we shall denote by

$$A^{\delta} := \{a \in A | \delta(a) = 0\}$$

the algebra of constants of A.

As mentioned before we are dealing with one particular locally nilpotent derivation of $\mathcal{L}(x, y)$. We recall a locally nilpotent derivation of an algebra A is a derivation δ so that $\delta^n(a) = 0$ for every $a \in A$ with n depending on a. Here we shall study the behaviour of $\mathcal{L}(x, y)^{\delta}$, where δ is the derivation of $\mathcal{L}(x, y)$ sending $y \mapsto x$ and $x \mapsto 0$. Notice that δ is a locally nilpotent derivation of $\mathcal{L}(x, y)$.

Let us fix some notations in our Lie setting.

Definition 2.2. A Lie algebra *L* is a vector space over the field *K* with a binary operation $[\cdot, \cdot] : L \to L$ so that

$$[a, a] = 0$$
 and $[[a, b], c] + [[c, a], b] + [[b, c], a] = 0$

for every $a, b, c \in L$.

The second of the relations above is called Jacobi's identity. For the sake of simplicity of notations we shall omit brackets in case of left-normed products, that is, we define $[a_1, a_2, \ldots, a_{n-1}, a_n] = [[[\ldots [a_1, a_2], \ldots], a_{n-1}], a_n]$ for all $a_1, \ldots, a_n \in L$. An immediate consequence of Jacobi's identity is the relation

(1)
$$[a, b, c] = [a, c, b] - [a, [c, b]]$$

for all $a, b, c \in L$. In particular [a, b, a, b] = [a, b, b, a].

As well as for the associative case, in the Lie setting we can construct a free object, called the free Lie algebra. If X is a set of indeterminates we are allowed to consider $\mathcal{L}(X)$ the free Lie algebra over the field K generated by the set X. In the sequel we shall use linear basis of the free Lie algebra in order to perform our computations. Indeed linear basis of the free Lie algebra have been studied by several authors using different techniques. We shall use one of them in particular which construction will be presented below.

Suppose X is linearly ordered, then using an induction on the degree of Lie monomials, we construct basic words starting from the "smallest" basic words:

Each $x_i \in X$ is a basic word of degree 1. Then let n > 1 and suppose that for every d < n all basic words of degree d are defined and ordered. Suppose also that provided w, u and v are basic words of degree d < n, then w = uv implies w > v. Hence the monomial q of degree n is basic if whenever $q = q_1q_2$, then q_1, q_2 are basic words and $q_1 > q_2$ and the decomposition $q = (q'q'')q_2$ implies $q'' \leq q_2$. Any monomial not respecting the previous axioms is to be considered not basic. We have the next result.

Theorem 2.3. The basic words in X form a basis of the free Lie algebra $\mathcal{L}(X)$.

Instead of basic words we shall use the expression Hall basis. We will denote by \mathcal{B} the Hall basis of the free Lie algebra $\mathcal{L}(x, y)$. The order used here is the deg-lexicographic one: $f \leq g$ if, and only if, $\deg(f) < \deg(g)$ or $\deg(f) = \deg(g)$ and $f \leq_{\text{lex}} g$ where \leq_{lex} is the lexicographic order and x < y. Here we use [x, y], $[x, y, x], [x, y, y] \in \mathcal{B}$ instead of [y, x], [y, x, x] and [y, x, y].

The first basis of a free Lie algebra was found by Hall in [16] whereas Shirshov in [24] and Lyndon in [20] constructed the basis of a free Lie algebra consisting of the so called *Lyndon-Shirshov words* (see for example the book of Bokut again). In a former work [23] Shirshov was able to develop the composition method for Lie algebras which was refined and rewritten in a modern language setting by Bokut in [3]. The smart contribution of Shirshov is highlighted in [24] where he suggested a path for choosing bases for free Lie algebras generalizing those of Hall's basis and of Lyndon-Shirshov's basis. Other examples of bases for free Lie algebras were found by Bokut in [4], by Reutenauer in [22], by Blessenohl and Laue in [2], by Bryant, Kovacs and Stöhr in [6], by Guilfoyle and Stöhr in [15] and by Chibrikov in [7] in which the author shows a right normed monomial basis for the free Lie algebra. For more details on the theory of free Lie algebras we remand to the book of Reutenauer [22].

We will also use a well celebrated result by Shirshov [25] and Witt [28] which separately obtained the same result.

Theorem 2.4. Every Lie subalgebra of a free Lie algebra is free. Moreover every finite dimensional Lie subalgebra of a free Lie algebra has dimension 1.

3. Novicki's conjecture for $\mathcal{L}(x, y)$

As already mentioned in the introduction, the algebra of constants $\mathcal{L}(x, y)$ is not finitely generated. Our target in this paper is trying to get a reformulation of Novicki's conjecture in terms of relations of the subalgebra $\mathcal{L}(x, y) - span_K\{y\}$. In what follows we shall construct an infinite set of generators of $\mathcal{L}(x, y)$ which ideal of relations will be used to describe $\mathcal{L}(x, y)^{\delta}$. Moreover every field is of characteristic 0.

We recall the first formulation of Novicki's conjecture stated the algebra of constants of $K[X,Y]^{\delta}$, where X and Y are two finite sets of the same number of variables, is generated by the variables X and the $u_{ij} = \begin{vmatrix} x_i & x_j \\ y_i & y_j \end{vmatrix}$. In [14] Drensky and Makar-Limanov found a uniformly looking explicit set of defining relations of the algebra of constants $K[X,Y]^{\delta}$ which corresponds to the reduced Gröbner basis of the related ideal of K[X,U], where $U = u_{ij}|1 \le i < j \le n$. The basic idea of the paper is the construction of objects which are similar to the *determinants* u_{ij} .

We shall denote $L := \mathcal{L}(x, y)$. We consider the following elements of L,

$$U_{A,B}^{(m,k)} := \left| \begin{array}{cc} \delta^m(A) & \delta^k(A) \\ \delta^m(B) & \delta^k(B) \end{array} \right| = [\delta^m(A), \delta^k(B)] - [\delta^k(A), \delta^m(B)],$$

where A and B are Hall monomials and we shall call $U_{A,B}^{(m,k)}$ pseudoterminant of degree (m, k) in the Hall monomials A and B.

The pseudodeterminants will turn out to be close to a generating set of L. First of all notice the following facts.

Remark 3.1. Notice that $\delta(U_{AB}^{(m,k-1)}) = U_{AB}^{(m,k-1)}$ if $\delta^{m+1}(A) = \delta^{m+1}(B) = 0$.

- (1) Remark that $U_{A,A} = 2[\delta(A), A]$. In particular if A and B Remark 3.2. are Hall monomials such that $\delta(A) = B$, then $U_{A,A} = 2[B, A]$. In particular $[x,y] = U_{y,y}.$
 - (2) If we swap two columns in $U_{A,B}$ we get $-U_{A,B}$ whereas if we swap two rows in $U_{A,B}$ we do not change any sign.

Let us introduce the main tool of the section.

Definition 3.3. Let $f \in U$. If there exists $f' \in U$ such that $\delta(f') = f$ we say f' is an integral of f.

Remark 3.4. Of course, if f', f'' are two integrals of f, then there exists $c \in U^{\delta}$ such that f' = f'' + c. Hence we are allowed to write $\int f = f' + c$. With abuse of notation, if g is an integral of f' we will write $\int_{-\infty}^{2} f = g + c$.

We may develop easily some integration rules for the intent we shall use later on. In particular we have:

Lemma 3.5.

 $\begin{array}{l} \textbf{mma 3.5.} & (1) \quad \int x = y + c \\ (2) \quad \int (U_{A,B}^{(m,k)}) = U_{A,B}^{(m,k-1)} + c, \ c \in U^{\delta}, \ if \ \delta^{m+1}(A) = \delta^{m+1}(B) = 0. \\ (3) \quad \int (U_{A,B}^{(m,k)}) = \sum_{s=0}^{l} (-1)^{s} U_{A,B}^{(m+s,k-(s+1))} + c, \ c \in U^{\delta}, \ if \ \delta^{m+l+1}(A) = 0, \\ \delta^{m+l+1}(B) = 0 \ and \ k - (s+1) \neq 0 \ for \ every \ s = 0, \dots, l. \\ (4) \quad \int (U_{A,B}^{(m+1,0)}) = \sum_{s=0}^{l} (-1)^{s} U_{A,B}^{(m-s,s)} + c, \ c \in U^{\delta}, \ if \ \delta^{l+1}(A) = 0, \ \delta^{l+1}(B) = 0 \end{array}$

(4)
$$\int (U_{A,B}^{(a)}(r)) = \sum_{s=0}^{r} (-1)^{s} U_{A,B}^{(a)}(r) + c, c \in U^{s}, \text{ if } \delta^{s+1}(A) = 0, \delta^{s+1}(B) = 0$$

and $m - s \neq 0$ for every $s = 0, \dots, l$.

Proof. The statements (1) and (2) are immediate. To prove the statement (3), it is sufficient to show $\delta(\sum_{s=0}^{l}(-1)^{s}U_{A,B}^{(m+s,k-(s+1))}) = U_{A,B}^{(m,k)}$ if $\delta^{m+l+1}(A) = 0$, $\delta^{m+l+1}(B) = 0$ and $k - (s+1) \neq 0$ for every $s = 0, \dots, l$. We have that

$$\begin{split} \delta(\sum_{s=0}^{l} U_{A,B}^{(m+s,k-(s+1))}) &= \sum_{s=0}^{l} (-1)^{s} ([\delta^{m+s+1}(A), \delta^{k-(s+1)}(B)] + [\delta^{m+s}(A), \delta^{k-s}(B)]) \\ &\sum_{s=0}^{l} (-1)^{s+1} ([\delta^{k-s}(A), \delta^{m+s}(B)] + [\delta^{k-(s+1)}, \delta^{m+s+1}(B)]) \\ &= (-1)^{l} [\delta^{m+l+1}(A), \delta^{k-(l+1)}] + [\delta^{m}(A), \delta^{k}(B)] \\ &- [\delta^{k}(A), \delta^{m}(B)] + (-1)^{l+1} [\delta^{k-(l+1)}(A), \delta^{m+l+1}(B)] \\ &= U_{A,B}^{(m,k)} \end{split}$$

since $\delta^{m+l+1}(A) = \delta^{m+l+1}(B) = 0$. The proof of statement (4) is analogous.

We do consider the following subalgebra of L,

$$U := \langle x, U_{A,B}^{(m,k)} | m, k \in \mathbb{N} \rangle,$$

then we get the next result.

Proposition 3.6. The algebra U is an algebra with derivation δ . Moreover it coincides with $L - \operatorname{span}_F\{y\}$.

Proof. For the first statement it is sufficient to test δ on each generator. By Remark 3.1 we get

$$\delta(U_{A,B}^{(m,k-1)}) = U_{A,B}^{(m,k)} \in U$$

if $\delta^{m+1}(A) = \delta^{m+1}(B) = 0$. By definition $\delta(x) = 0 \in U$ and this is sufficient to get $\delta(U) \subseteq U$. Hence U is an algebra with derivation δ .

For the other sentence we have to use an inductive argument on the multihomogeneous degree. Let $S := L - span_F\{y\}$ and take $f \in S$. We can suppose f being multihomogeneous in the variables x and y. If $\deg_x(f) = 1$, then $f = \alpha x$ where $\alpha \in K$ and we are done. If $\deg_y(f) = 1$, then $f = \alpha[y, x]$ where $\alpha \in K$ and $f \in U$ because $x = \delta(y)$. Suppose now the assertion true for multihomogeneous elements of S of degree $n - 1 \ge 2$ and take f of degree n. Of course by Theorem 2.4 S is free and has a Hall basis. We have $f = \sum_i \alpha_i M_i$, where the M_i 's are Hall monomials. Hence for every i we have $M_i = p_i q_i$, where p_i, q_i are Hall monomials such that $p_i \ge q_i$. For every i, if $\deg(q_i) \ge 2$, then $n > \deg(p_i) \ge \deg(q_i)$ and by induction p_i, q_i belong to U. It remains to tudy the case $\deg(q_i) = 1$. If $q_i = \alpha x$ we have nothing to prove. Indeed $q_i \ne \alpha y$ because q_i is a Hall monomial of S and $y \notin S$ and this concludes the proof. \Box

From the previous result if $f \neq \alpha y$, then $f \in U$. Hence f can be written as

(2)
$$f = \sum_{\overline{i}} \alpha_{\overline{i}} u_{i_1} \cdots u_{i_{k_{\overline{i}}}}$$

where $\alpha_{\overline{i}} \in K$ and the u_{i_j} 's are generators of U as above. Indeed we have f is a constant if and only if

(3)
$$0 = \delta(f) = \sum_{\overline{i}} \sum_{j=1}^{k_{\overline{i}}} \alpha_{\overline{i}} u_{i_1} \cdots u_{i_{j-1}} \delta(u_{i_j}) u_{i_{j+1}} \cdots u_{i_k},$$

By Remark 3.1 we get $\delta(u_{i_j})$ is again a generator of the algebra U so the formula (3) turns out to be a *relation* in the generators of U. Let \mathfrak{R} being the set of all relations as above, then Lemma 3.5 gives us an effective tool to find out constants if one of the relation among the generators of U is given. Of course we have:

Theorem 3.7. The algebra of constant of $\mathcal{L}(x, y)$ is the algebra

$$\mathcal{L}(x,y)^{\delta} = \langle \int g | g \in \mathfrak{R} \rangle$$

and $\int g$ can be effectively computed.

4. Some special cases

In the previous section we reformulated in a teorethical way Novicki's conjecture. Our intent now is to give the concrete answer to the Novicki's conjecture in some special cases. We shall see how do figure out Lie monomials which are constants and constants of small degrees. 4.1. The monomial case. First of all we investigate on the structure of Hall monomials which are constants. We hope this will share light on the structure of constants in general.

The next easy and well known result will be of help in our work and we would like to give a simple proof of it in order to highlight the strict relations between the Lie algebra structure of $\mathcal{L}(x, y)$ and its elements. In fact it is a direct consequence of the Shirshov-Witt Theorem.

Proposition 4.1. Let $f, g \in \mathcal{L}(x, y)$ such that [f, g] = 0, then $f = \alpha g$, where α belongs to the field.

Proof. Let us suppose $g \neq 0$ and do consider S the Lie algebra generated by f and g. By Theorem 2.4 S is free and all its products are 0. Hence the dimension of S as a vector space is less than or equal to 2. This means S is a finite dimensional free Lie algebra. By the second part of Theorem 2.4 and the fact that $g \neq 0$ we get the dimension of S is 1 and we are done.

Let us denote by U' the sub Lie algebra of L^{δ} generated by the $U_{A,A}$, where A is a Hall monomial.

Proposition 4.2. Let $M = [A, B] \neq 0$ be a monomial of L^{δ} where A, B are monomials in the Hall basis. Then $M \in U'$ if at least one between A and B belongs to U.

Proof. Suppose firstly A, B are constants. If $\deg(M) = 3$ then $M = \pm [x, y, x] = \pm [[x, y], x]$ and consequently $M \in U'$. Since A, B are monomials such that $\deg(A)$, $\deg(B) < \deg(M)$, the result follows by induction. Now suppose that A is a constant and B is not constant. Then by Proposition 4.1 $0 = \delta(M) = [A, \delta(B)]$ implies $\delta(B) = \pm A$ since $A \neq 0$ and $\delta(B) \neq 0$. Thus $M = \pm [\delta(B), B]$ and hence $0 = [\delta^2(B), B]$. Therefore $\delta^2(B) = 0$ since $B \neq 0$ and $0 < \deg_y(B) < \deg_y(\delta^2(B))$. Therefore $M = \pm [\delta(B), B]$ and the proof follows by item (1) of Remark 3.2.

4.2. Lie polynomials of small degree. In this subsection we shall calculate the generators of the constants of degree up to 6. In particular we will show that if $f = f(x, y) \in \mathcal{L}(x, y)$ is a polynomial of degree less than or equal to 6 and f is a constant then f belongs to the sub Lie algebra generated by x, [x, y], [x, y, y, [x, y, x]].

The next result is a technical lemma that will be useful to our intents.

Lemma 4.3. For all $a \ge 0$ and b > 0

$$[x, y, \underbrace{x, \dots, x}_{a}, y, \underbrace{x, \dots, x}_{b}] = [x, y, \underbrace{x, \dots, x}_{a+b}, y] + \sum_{i=0}^{b-1} \binom{b}{i} [x, y, \underbrace{x, \dots, x}_{a+i}, [y, \underbrace{x, \dots, x}_{b-i}]]$$

where $\binom{b}{i} = \frac{b!}{i!(b-i)!}$ is a binomial coefficient.

Proof. The proof will be performed by induction on b. If b = 1 the result follows from (1). We have that

$$\begin{split} [x, y, \underbrace{x, \dots, x}_{a}, y, \underbrace{x, \dots, x}_{b+1}] &= [x, y, \underbrace{x, \dots, x}_{a+1}, y, \underbrace{x, \dots, x}_{b}] + [x, y, \underbrace{x, \dots, x}_{a}, [y, x], \underbrace{x, \dots, x}_{b}] \\ &= [x, y, \underbrace{x, \dots, x}_{a+b+1}, y] + [x, y, \underbrace{x, \dots, x}_{a+b}, [y, x]] \\ &+ \sum_{i=0}^{b-1} {b \choose i} [x, y, \underbrace{x, \dots, x}_{a+1+i}, [y, \underbrace{x, \dots, x}_{b-i}]] \\ &+ \sum_{j=0}^{b-1} {b \choose j} [x, y, \underbrace{x, \dots, x}_{a+j}, [y, \underbrace{x, \dots, x}_{b+1)-j}] \\ &= [x, y, \underbrace{x, \dots, x}_{a+b+1}, y] + [x, y, \underbrace{x, \dots, x}_{a+j}, [y, x]] \\ &+ \sum_{j=0}^{b} {b \choose j-1} [x, y, \underbrace{x, \dots, x}_{a+j}, [y, \underbrace{x, \dots, x}_{b+1)-j}] \\ &+ \sum_{j=0}^{b-1} {b \choose j} [x, y, \underbrace{x, \dots, x}_{a+j}, [y, \underbrace{x, \dots, x}_{b+1)-j}] \\ &= [x, y, \underbrace{x, \dots, x}_{a+b+1}, y] + \sum_{j=0}^{b} {b \choose j} [x, y, \underbrace{x, \dots, x}_{a+j}, [y, \underbrace{x,$$

where the first equality holds by the induction hypothesis and the last by relation $\binom{b+1}{j} = \binom{b}{j-1} + \binom{b}{j}$ for all $1 \le j \le b-1$.

Corollary 4.4. If $f \neq 0$ is a multihomogeneous polynomial such that $deg_y(f) = 2$ then $f \in L^{\delta}$ if and only if f belongs to the subalgebra generated by x, [x, y].

Proof. Let f be a multihomogeneous polynomial, $\deg_y(f) = 2$ such that $f \in L^{\delta}$. Since left-normed brackets generate $\mathcal{L}(x, y)$ write

$$f = \alpha_0[x, y, y, \underbrace{x, \dots, x}_k] + \alpha_1[x, y, x, y, \underbrace{x, \dots, x}_{k-1}] + \dots + \alpha_k[x, y, \underbrace{x, \dots, x}_k, y]$$

we have $\delta(f) = 0$ implies that

$$(\alpha_0 + \dots + \alpha_k)[x, y, \underbrace{x, \dots, x}_{k+1}] = 0$$

and therefore $\alpha_0 + \cdots + \alpha_k = 0$. The result follows from previous lemma.

The next propositions prove that if $f = f(x, y) \in \mathcal{L}(x, y)$ is a polynomial of degree less than or equal to 6 and $f \in L^{\delta}$ then f belongs to the subalgebra generated by x, [x, y], [x, y, y, [x, y, x]].

Proposition 4.5. If $f \neq 0$ is a polynomial of degree less than or equal to 5 and $f \in L^{\delta}$ then f belongs to the subalgebra generated by x, [x, y].

Proof. Suppose without loss of generality that f is a multihomogeneous polynomial. In fact if $f = \sum_{\alpha} f_{\alpha}$ where f_{α} is a component multihomogeneous of f, $\delta(f) = \sum \delta(f_{\alpha}) = 0$ if, and only if, $\delta(f_{\alpha}) = 0$ if, and only if, $\delta(f_{\alpha}) = 0$. If $\deg_y(f) = 1$, f belongs to the subalgebra generated by x, [x, y] and consequently is a constant. If $\deg_y(f) = 2$ the result follows from previous corollary. Suppose f homogeneous of degree 4 such that $\deg_y(f) = 3$. In this case $f = \alpha[x, y, y, y]$ and consequently $\delta(f) = 2\alpha[x, y, x, y] = 0$ implies that $\alpha = 0$.

Now suppose f homogeneous of degree 5. In this case we have two possibilities: $\deg_y(f) = 3$ and 4. Let $\alpha[x, y, y, y, y] \in L^{\delta}$, that is, $\alpha[x, y, x, y, y] + \alpha[x, y, y, x, y] + \alpha[x, y, y, x, y] = 0$. Since [x, y, x, y, y] = [x, y, y, x, y] and [x, y, y, y, x] = [x, y, y, x, y] - [x, y, y, [x, y]], the last equation implies that $3\alpha[x, y, x, y, y] - \alpha[x, y, y, [x, y]] = 0$. Therefore $\alpha = 0$ since [x, y, x, y, y], [x, y, y, [x, y]] belong to Hall's basis of $\mathcal{L}(x, y)$. Let $\alpha[x, y, x, y, y] + \beta[x, y, y, y, x] \in L^{\delta}$. Therefore $(\alpha + 2\beta)[x, y, x, y, x] + \alpha[x, y, x, x, y] = (\alpha + 2\beta)([x, y, x, x, y] - [x, y, x, [x, y]) + \alpha[x, y, x, x, y] = 0$. Since [x, y, x, x, y], [x, y, x, x, y], [x, y, x, [x, y]] belong to Hall's basis we have that $\alpha = \beta = 0$.

Proposition 4.6. If $f \neq 0$ is a polynomial of degree 6 and $f \in L^{\delta}$ then f belongs to the subalgebra generated by x, [x, y] and [x, y, y, [x, y, x]].

Proof. There are three cases to check: $\deg_y(f) = 3, 4$ and 5. Suppose $\alpha[x, y, y, y, y, y] \in L^{\delta}$, i.e., $2\alpha[x, y, x, y, y, y] + \alpha[x, y, y, y, x, y] + \alpha[x, y, y, y, y, x] = 0$. Since

$$[x, y, y, y, x, y] = [x, y, x, y, y, y] - [x, y, y, y, [x, y]]$$

and

$$[x, y, y, y, y, x] = [x, y, x, y, y, y] - 2[x, y, y, y, [x, y]],$$

it follows that $\alpha = 0$. Suppose

(4)
$$f = \alpha[x, y, x, y, y, y] + \beta[x, y, y, y, x, y] + \gamma[x, y, y, y, y, x] \in L^{\delta},$$

that is, $\delta(f) = \alpha[x, y, x, x, y, y] + (\alpha + 2\beta)[x, y, x, y, x, y] + (\alpha + 2\gamma)[x, y, x, y, y, x] + (\beta + \gamma)[x, y, y, y, x, x] = 0$. Since $v_1 = [x, y, x, x, y, y]$, $v_2 = [x, y, x, y, [x, y]]$, $v_3 = [x, y, y, [x, y, x]]$ belong to Hall's basis and by simple calculations using (1) $[x, y, x, y, x, y] = v_1 - v_2 + v_3$, $[x, y, x, y, y, x] = v_1 - 2v_2 + v_3$, $[x, y, y, y, x, x] = v_1 - 3v_2$, we have the following system equations: $\alpha + \beta + \gamma = 0$ and $-3\alpha - 5\beta - 7\gamma = 0$ whose solution is $\beta = -2\alpha$ and $\gamma = \alpha$. Replacing this solution in (4) we get $f = -\alpha[x, y, y, [x, y, y]] = 0$. Finally consider

$$(5) \ f = \alpha[x, y, x, y, y, x] + \beta[x, y, y, x, x] + \gamma[x, y, x, x, y, y] + \epsilon[x, y, x, y, x, y] \in L^{o},$$

that is, $(\alpha + \gamma)[x, y, x, x, y, x] + (\alpha + 2\beta + \epsilon)[x, y, x, y, x, x] + (\gamma + \epsilon)[x, y, x, x, x, y] = 0$. Denote by $v_1 = [x, y, x, x, x, y]$ and $v_2 = [x, y, x, x, [x, y]]$. Since $[x, y, x, x, y, x] = v_1 - v_2$ and $[x, y, x, y, x, x] = v_1 - 2v_2$ follow that $\alpha + \beta + \gamma + \epsilon = 0$ and $-3\alpha - 4\beta - \gamma - 2\epsilon = 0$. Consequently $\gamma = \alpha + 2\beta$ and $\epsilon = -2\alpha - 3\beta$. Replacing this solution in (5) we obtain $f = \alpha([x, y, x, y, y, x] + [x, y, x, x, y, y] - 2[x, y, x, y, x, y]) = \beta([x, y, y, y, x, x] + 2[x, y.x, x, y, y] - 3[x, y, x, y, x, y]) \in L^{\delta}$. By simple calculations we conclude that

$$\begin{split} [x,y,x,y,y,x] + [x,y,x,x,y,y] - 2[x,y,x,y,x,y] &= -[x,y,y,[x,y,x]], \\ [x,y,y,y,x,x] + 2[x,y,x,x,y,y] - 3[x,y,x,y,x,y] &= -3[x,y,y,[x,y,x]] \\ \text{and this ends the proof.} \\ \Box$$

Indeed if A := [x, y, y], B := [x, y, x], then $\delta(A) = B$, then $M = 2U_{A,A} \in U'$. We are suspicious all constants Hall monomials satisfy the hipothesis of the previous proposition. In particular we can state the next conjecture.

Conjecture 4.7. Let M be a Hall monomial so that it is a constant under the derivation δ , then if M = [A, B], where A, B are Hall monomials again, then at least one between A and B is a constant.

We can also state a stronger conjecture supported by the results of this section.

Conjecture 4.8. The algebra of constants $\mathcal{L}(x, y)^{\delta}$ equals U'.

References

- L. Bedratyuk, A note about the Nowicki conjecture on Weitzenbock derivations, Serdica Math. J. 35 (2009), 311-316.
- [2] D. Blessenohl, H. Laue, A basis construction for free Lie algebras, Exp. Math. 11 (1993), 145-152.
- [3] L. A. Bokut, Unsolvability of the equality problem and subalgebras of finitely presented Lie algebras, Math. USSR Izv. 6 (1972), 1153-1199.
- [4] L. A. Bokut, A basis for free polynilpotent Lie algebras, Algebra Logika 3 (1963), 13-19.
- [5] L. A. Bokut, Y. Chen, Grobner-Shirshov bases for Lie algebras: after A. I. Shirshov, Southeast Asian Bull. Math. **31** (2007), 1057-1076.
- [6] R. M. Bryant, L. G. Kovacs, R. Stöhr, Invariant bases for free Lie rings, Quart J. Math. 53 (2002), 1-17.
- [7] E. S. Chibrikov, A right normed basis for free Lie algebras and Lyndon-Shirshov words, J. Algebra 302(2) (2006), 593-612.
- [8] R. Dangovski, V. Drensky, S. Fındık, Weitzenböck derivations of free metabelian Lie algebras, Linear Algebra and its Applications, 439 10 (2013), 3279-3296.
- R. Dangovski, V. Drensky, Ş. Fındık, Weitzenböck derivations of free associative algebras, Journal of Algebra and Its Applications, 16 03 (2017), 1750041/1-26.
- [10] H. Derksen, G. Kemper, Computational Invariant Theory, Encyclopaedia of Mathematical Sciences, Invariant Theory and Algebraic Transformation Groups 130, Springer-Verlag, Berlin, 2002.
- [11] V. Drensky, Invariants of unipotent transformations acting on noetherian relatively free algebras, Serdica Math. J. 30 (2004), 395–404.
- [12] V. Drensky, S. Findik, The Nowicki conjecture for free metabelian Lie algebras, preprint.
- [13] V. Drensky, C.K. Gupta, Constants of Weitzenböck derivations and invariants of unipotent transformations acting on relatively free algebras, J. Algebra 292 (2005), No. 2, 393-428.
- [14] V. Drensky, L. Makar-Limanov, The conjecture of Nowicki on Weitzenböck derivations of polynomial algebras, J. Algebra Appl. 8 (2009), 41-51.
- [15] S. Guilfoyle, R. Stöhr, Invariant bases for free Lie algebras, J. Algebra 204 (1998) 337-346.
- [16] M. Hall, A basis for free Lie rings and higher commutators in free groups, Proc. Amer. Math. Soc. 1 (1950), 575-581.
- [17] J. Khoury, Locally Nilpotent Derivations and Their Rings of Constants, Ph.D. Thesis, Univ. Ottawa, 2004.
- [18] J. Khoury, A Groebner basis approach to solve a conjecture of Nowicki, J. Symbolic Comput. 43 (2008), 908-922.
- [19] S. Kuroda, A simple proof of Nowicki's conjecture on the kernel of an elementary derivation, Tokyo J. Math. 32 (2009), 247-251.
- [20] R. C. Lyndon, On Burnsides problem I, Trans. Amer. Math. Soc. 77 (1954), 202-215.
- [21] A. Nowicki, Polynomial Derivations and Their Rings of Constants, Uniwersytet Mikolaja Kopernika, Torun, 1994. www-users.mat.umk.pl/~anow/ps-dvi/pol-der.pdf.
- [22] C. Reutenauer, Dimensions and characters of the derived series of the free Lie algebra, M. Lothaire (Ed.), Mots, Melanges Offerts a M.-P. Schutzenberger, Hermes, Paris, 1990, 171-184.
- [23] A. I. Shirshov, Some algorithmic problems for Lie algebras, Sibirsk. Mat. Z. 3 (1962) 292-296 (Russian); Translation: ACM SIGSAM Bull. 33 (2) (1999) 3-6.
- [24] A. I. Shirshov, On free Lie rings, Mat. Sb. 45 (2) (1958), 113-122 (Russian).

- [25] A. I. Shirshov, Subalgebras of free Lie algebras, Mat. Sb. 33 75 (1953) 441-452 (Russian).
- [26] B. Sturmfels, Algorithms in Invariant Theory. 2nd ed. Texts and Monographs in Symbolic Computation, Springer-Verlag, Wien, 2008.
- [27] R. Weitzenböck, Über die Invarianten von linearen Gruppen, Acta Math. 58 (1932), 231-293.
- [28] E. Witt, Die Unterringe der freien Liescher Ringe, Math. Z. 64 (1956) 195-216.
- [29] E. I. Zelmanov, On Engel Lie algebras (Russian), Sibirsk. Mat. Zh. 29(5) (1988), 112-117. Translation: Sib. Math. J. 29 (1988), 777-781.

IMECC, UNIVERSIDADE ESTADUAL DE CAMPINAS, RUA SERGIO BUARQUE DE HOLANDA 651,13083-859, CAMPINAS (SP), BRAZIL

Email address: centrone@ime.unicamp.br

DEPARTMENT OF MATHEMATICS, ÇUKUROVA UNIVERSITY, 01330 BALCALI, ADANA, TURKEY *Email address:* sfindik@cu.edu.tr

XXX

 $Email \ address: \verb"manuela.dss0gmail.com"$