# **Stochastic volatility model with long memory for water quantity-quality dynamics**

Hidekazu Yoshioka<sup>a,\*</sup> and Yumi Yoshioka<sup>b</sup>

<sup>a</sup>Japan Advanced Institute of Science and Technology, 1-1 Asahidai, Nomi 923-1292, Japan <sup>b</sup>Gifu University, 1-1 Yanagido, Gifu 501-1193, Japan

\*Corresponding author: yoshih@jaist.ac.jp, ORCID: 0000-0002-5293-3246

**Abstract:** Water quantity and quality are vital indices for assessing fluvial environments. These indices are highly variable over time and include sub-exponential memory, where the influences of past events persist over long durations. Moreover, water quantity and quality are interdependent, with the former affecting the latter. However, this relationship has not been thoroughly studied from the perspective of long-memory processes, which this paper aims to address. We propose applying a new stochastic volatility model, a system of infinite-dimensional stochastic differential equations, to describe dynamic asset prices in finance and economics. Although the stochastic volatility model was originally developed for phenomena unrelated to the water environment, its mathematical universality allows for an interdisciplinary reinterpretation: river discharge is analogous to volatility, and water quality to asset prices. Moreover, the model's infinitedimensional nature enables the analytical description of sub-exponential memory. The moments and autocorrelations of the model are then obtained analytically. We mathematically analyze the stochastic volatility model and investigate its applicability to the dynamics of water quantity and quality. Finally, we apply the model to real time-series data from a river in Japan, demonstrating that it effectively captures both the memory and the correlation of water quality indices to river discharge. This approach, grounded in infinite-dimensional stochastic differential equations, represents a novel contribution to the modelling and analysis of environmental systems where long memory processes play a role.

**Keywords:** Measure-valued processes, Stochastic volatility model, Long memory, River flow, Water quality.

#### **(Statements & Declarations)**

**Fundings** This study was supported by the Japan Society for the Promotion of Science (KAKENHI No. 22K14441) and the Japan Science and Technology Agency (PRESTO No. JPMJPR24KE).

**Competing interests** The authors have no relevant financial or non-financial interests to disclose.

**Data availability** The data will be made available upon reasonable request to the corresponding author.

**Acknowledgments** The authors would like to express their gratitude to Dr. Ikuo Takeda of Shimane University for providing a part of the water quality data used in this paper.

**Declaration of generative AI in scientific writing** The authors did not use generative AI for scientific writing of this manuscript.

#### **1. Introduction**

#### **1.1 Study background**

Assessing water quantity and quality in rivers is a core issue for fostering sustainable and harmonious coexistence between nature and humans. Examples of such efforts globally include, but are not limited to, hydrological connectivity in river systems (Durighetto et al., 2023; Zanetti et al., 2024)[1,2], surveys of international rivers under severe anthropogenic pressure (Ding et al., 2024)[3], flood hazard management in the context of climate change (Bahramloo et al., 2024)[4], investigations into potential threats to biodiversity loss in river environments (Rai et al., 2024)[5], fish habitat assessments in riverscapes (Hansen et al., 2024)[6], and water quality analyses in lakes (Gao et al., 2024)[7].

The physical, chemical, and biological dynamics of river environments are highly variable and subject to random fluctuations. Consequently, it is reasonable to conceptualize these dynamics as stochastic processes, characterized by continuous and discrete time-series data that fluctuate randomly, such as environmental variables including rainfall intensity and river discharge (Doukhan, 2018; Volpi et al., 2024)[8,9]. Particularly, stochastic differential equations (SDEs) (Øksendal and Sulem, 2019; Pascucci, 2024)[10,11] are well-suited for balancing mechanistic and phenomenological modeling. They have been widely employed in both scientific and engineering fields to develop simple models that capture the essence of the target phenomenon. SDEs play a central role in ecological modeling, where randomness significantly affects system stability (Mandal et al., 2024; Paul et al., 2025)[12,13]. They have also been effectively used to investigate disease propagation (Mohammad et al., 2024; Đorđević and Dahl, 2024)[14,15] and asset price dynamics and green financing in economics (Hambel et al., 2024; Zhang and Ren, 2024)[16,17]. Coastal sand dynamics (Ramakrishnan et al., 2024; Vinent et al., 2021)[18,19] and salt intrusion in tidal marshes (Dijkstra et al., 2023)[20] are additional examples of SDE applications.

Memory is an essential element in the stochastic process modeling of water quantity and quality in rivers, and it is effectively evaluated through autocorrelation. The rate at which autocorrelation decays characterizes the forgetting timescale of past events. Traditionally, this decay has been assumed to follow an exponential pattern, or Markovian dynamics, due to its analytical tractability in applications (Calvani and Perona, 2023; Latella et al., 2024; Olson et al., 2021; Yoshioka and Yoshioka, 2023a; Yoshioka and Yoshioka, 2023b)[21-25]. However, empirical hydrological data, including discharge, water quality, and hydrological signatures, often exhibit sub-exponential decay, indicative of longer memory and non-Markovian behavior (Fatni et al., 2024; Maftei et al., 2016; Guo et al., 2024a; Pizarro et al., 2024; Rahmani and Fattahi, 2024)[26-30]. Therefore, employing non-Markovian models with long memory is a reasonable strategy for addressing issues related to water quantity and quality.

Some non-Markovian processes can alternatively be expressed as a superposition (i.e., suitable integration) of Markovian processes, albeit at the cost of increased dimensionality, which may extend from finite to infinite dimensions (e.g., Abi Jaber, 2019; Cuchiero and Teichmann, 2020)[31,32]. This mathematical technique is called Markovian lifts or Markovian embedding, depending on the research domain, with slight differences in terminology. The shared principle underlying these techniques is to reinterpret a complex process as a collection or continuum of simpler processes. Markovian lifts have been employed in various fields, including physics (Kanazawa and Sornette, 2024; Wiśniewski and Spiechowicz, 2024)[33,34] and economics (Bondi et al., 2024; Damian and Frey, 2024; Dupret and Hainaut, 2024)[35- 37]. The Markov chain approximation of the Volterra process is an example of Markovian lifts (Yang et al., 2024)[38]. In computational fluid dynamics, the principle of decomposing long memory into a sum of shorter memories has been recently explored (Guo et al., 2024b)[39], suggesting the applicability of Markovian lifts across diverse research areas. Moreover, the use of numerous hypothetical reservoir cells has been shown to be effective in physics-informed machine learning models (He et al., 2024)[40].

Despite their mathematical complexity, Markovian lifts have practical implications in the context of autocorrelation. They decompose sub-exponential functions into integrals of exponential functions, with exponents distributed according to a probability measure (e.g., Fasen and Klüppelberg, 2005)[41]. This property has been exploited in modeling long-memory dynamics in water quantity and quality (Yoshioka and Yoshioka, 2024a; Yoshioka and Yoshioka, 2024b)[42,43]. However, these studies have focused on either water quantity or water quality in isolation. The coupled dynamics of water quantity and quality which interact in real river environments, remain unexplored from the perspective of long-memory modeling via Markovian lifts. Addressing this gap requires innovative approaches to integrate the two different dynamics while preserving the analytical tractability of Markovian lifts and accurately capturing their statistical characteristics.

Stochastic volatility models in asset dynamics have demonstrated that long-memory processes, based on a superposition approach, can be coupled with another stochastic differential equation (SDE) without compromising overall tractability (Barndorff-Nielsen and Stelzer, 2013)<sup>[44]</sup>. In such models, volatility is represented as the fluctuating diffusion coefficient of an SDE. The key to this approach lies in its affine nature (e.g., Duffie et al., 2003)[45], which allows statistics such as mean, variance, and autocovariance to be obtained in closed form. Stochastic volatility models with long-memory have been extensively studied, focusing on the roughness of their sample paths (Bayer et al., 2021; de Truchis et al., 2024; Wu et al., 2022)[46-48]. While such models have been applied to river discharge modeling (Wang et al., 2023)[49], they have yet to incorporate water quality dynamics.

Water quality models for river environments based on SDEs have addressed various aspects, including the influence of suspended sediment concentrations on fluvial processes (Jing et al., 2020)[50], dissolved oxygen and biological oxygen demand (Mansour, 2023)[51], antibiotic bacterial dynamics (Gothwal and Thatikonda, 2020)[52], nutrient loads with a focus on regime shifts in river environments (Park and Rao, 2014)[53], and phycocyanin as a pigment indicator of cyanobacteria (Carpenter and Brock, 2024)[54]. Theoretical studies have also explored topics such as the most probable pollution control paths (Chen et al., 2024)[55] and stochastic differential games for basin-wide carbon and pollution control (Song et al., 2023; Song et al., 2024)[56,57]. However, most existing SDE models for water quality dynamics are Markovian, and therefore do not adequately account for memory effects, which are known to influence a wide variety of water quality indices (WQIs) (Chong et al., 2023; Kovács et al., 2010; Rahmani and Fattahi, 2024; Rozental and Tambieva, 2021; Spezia et al., 2021; Yoshioka and Yoshioka, 2024b)[58-62,43]. Another critical issue is the concentration-discharge relationship, which expresses the concentration of WOIs as a function of discharge. This relationship has been studied for various indices (Oin et al., 2024; Tunqui Neira et al., 2021; Wichman et al., 2024; Wymore et al., 2023; Zhan et al., 2022)[63-67] and quantifies the correlation between water quality and quantity. Despite its importance, this relationship has not been adequately incorporated into existing SDE models, leaving a gap in the development of realistic stochastic models for coupled water quantity-quality dynamics.

These challenges motivated this study, which seeks to address the aforementioned gaps by conceptualizing water quantity and quality as analogous to asset prices and volatility, respectively. The aims and contributions of this study are detailed below.

#### **1.2 Aim and contribution**

The aims of this study are twofold: (i) the proposal and analysis of a new coupled water quantity-quality model within the framework of affine processes, and (ii) the application of the model to real-world data. The study's contributions to achieving these aims are detailed below.

## **(i) Modeling and analysis**

The proposed model is formulated as two coupled superposed processes representing river discharge and a WQI. It is based on the superposition approach introduced by Barndorff-Nielsen (2001)[68], in which a long-memory process is generated by superposing infinitely many "small", measure-valued Ornstein– Uhlenbeck processes with exponential memory decay (supOU). These small processes, characterized by different reversion timescales, are combined via stochastic integration using appropriate random measures. Initially, small processes are chosen as linear SDEs but are later generalized to affine processes, with nonlinear extensions being examined (Iglói and Terdik, 2003)[69]. The supOU process is particularly tractable, making it useful in applications such as option pricing models with memory (Leonenko et al., 2024)[70] and in exploring statistical properties such as large deviations and fractal properties (Grahovac et al., 2019; Grahovac and Kevei, 2025)[71,72]. By construction, the supOU process constitutes an infinitedimensional system of SDEs containing many small processes. The success of the superposition approach lies in the affine form of the coefficients of the infinitesimal generators associated with these small processes (e.g., Chevalier et al., 2022; Motte and Hainaut, 2024; Yoshioka et al., 2023)[73-75], which enable the derivation of the generalized Riccati equation governing the moment-generating function of the superposed process.

In this paper, we realize a stochastic volatility model for coupled water quantity and quality dynamics by first formulating a jump-driven supOU process to represent river discharge. The temporal evolution of the WQI is then modeled as the product of a deterministic seasonal function and the exponential of an SDE, with drift and diffusion modulated by the supOU process. This framework allows for the representation of both positive and negative correlations between river discharge and the WQI (**Figure 1**). We refer to this coupled model as the Ornstein–Uhlenbeck stochastic volatility (supOUSV) model. The supOUSV model naturally incorporates the concentration-discharge relationship without compromising analytical tractability. We derive a closed-form expression for the autocorrelation of the WQI, along with its tail estimates. Moreover, the approach facilitates the derivation of a generalized Riccati equation for the moment-generating function, ensuring boundedness of the solution. Generalized Riccati equations are pivotal for analyzing infinite-dimensional stochastic volatility models (Cox et al., 2022; Friesen and Karbach, 2024)[76,77].



**Figure 1.** Conceptual diagram of the proposed model.

## **(ii) Model application**

We apply the proposed supOUSV process to real river discharge and WQI data at a study site in the midstream section of the Hii River, a Class A river in Japan. Hourly river flow time-series data are publicly available for the site, and almost weekly sampled data of multiple WQIs have been collected over 30 years. The empirical data indicate that both discharge and WQIs exhibit long-memory characteristics. The model is fitted to the data based on least-squares and moment-matching methods, enabling computational investigation of the proposed model. The results show that the model effectively captures both the longmemory dynamics and the discharge dependence of WQIs. Additionally, we analyze the concentrationdischarge relationship, focusing on the hysteresis patterns of the WQI (Mazilamani et al., 2024)[78], as represented by the model. In summary, this study contributes to the formulation, analysis, and application of a novel mathematical model for long-memory processes, advancing the understanding of coupled water quantity and quality dynamics.

# **1.3 Organization of this paper**

The remainder of this paper is organized as follows. **Section 2** describes our proposed model. **Section 3** mathematically analyzes the model by focusing on statistics. **Section 4** describes the application of the model to river discharge and WQIs at a study site in Japan. **Section 5** concludes this paper and presents its perspectives. **Appendices** contain auxiliary results concerning the finite-dimensional version of the proposed model (**Section A. 1**), proofs of propositions (**Section A. 2**), and auxiliary results (**Sections A. 3– A. 4**).

#### **2. Stochastic volatility model**

# **2.1 Overview**

We work on a complete probability space  $(\Omega, \mathbb{F}, \mathbb{P})$  as assumed in stochastic analysis (e.g., Definition 1.1.7) in Pascucci (2024))[11] ( $\Omega$  is collection of events,  $\mathbb F$  is filtration, and  $\mathbb P$  is probability). The primary variables in the model are river discharge and WQI. River discharge is represented as a supOU process driven by a pure-jump subordinator as in previous studies (e.g., Yoshioka and Yoshioka, 2024a)[42] (**Figure 1**). The WQI, the focus of this study, is modeled as a positive scalar process expressed as the product of a seasonal part and an exponential superposition of the stochastic volatility SDE. The supOU process of river discharge affectsthe drift and diffusion terms of the stochastic volatility SDE, thereby creating a correlation between water quality and quantity. Since the details of supOU processes have been well-documented (e.g., Barndorff-Nielsen, 2001)[68], this section emphasizes the stochastic volatility SDE of the WQI.

### **2.2 River discharge**

River discharge (referred to as discharge in the sequel) in this study represents the volume of water passing through a cross-section of a river per unit of time. The discharge at a station located within the river is modeled as a nonnegative, continuous-time scalar stochastic process,  $Y = (Y_t)_{t \in \mathbb{R}}$ , where  $t \in \mathbb{R}$  is time. Assuming a stationary state (e.g., Botter, 2010)[79], the discharge is modeled as a supOU process

(Barndorff-Nielsen, 2001; Barndorff-Nielsen and Stelzer, 2013)[44,68]:  
\n
$$
Y_t = \int_{s=-\infty}^{s=t} \int_{z=0}^{z=\infty} \int_{r=0}^{r=\infty} ze^{-r(t-s)} N\left(ds, dz, dr\right), \ t \in \mathbb{R}.
$$
\n(1)

Here,  $z > 0$  is the jump size at each jump (i.e., flood event),  $r > 0$  is recession rate of a flood event and is distributed according to the probability measure  $\pi = \pi(\text{d}r)$ , and N is a Poisson random measure on  $\times (0, +\infty) \times (0, +\infty)$  with the compensated measure  $\pi(dr)\nu(dz)ds$ , where  $\nu$  is the Lévy measure of a pure-jump subordinator (i.e., a pure-jump Lévy process with positive jumps). The two measures  $\nu$  and  $\pi$ control the size-frequency relationship and the recession of jumps, respectively. Here, N is extended to as in Barndorff-Nielsen and Stelzer (2013)[44] to deal with the stationary process.

Physically, the supOU process represents a situation where each flood event, assumed to occur randomly in time, has its own recession rate. The heterogeneity in recession rates originates from the interplay between antecedent water storage in the catchment and rainfall intensity (e.g., Hameed et al., 2023; Lee et al., 2023)[80,81]. This heterogeneity was previously described as a finite sum of independent Ornstein–Uhlenbeck processes by Botter (2010)[79]. The supOU process generalizes this concept to infinite dimensions. The long memory of discharge arises from catchment heterogeneity, where both fast (surface water) and slow runoffs (groundwater) coexist. Consequently, discharge can be understood as an aggregated variable—namely, a superposition in this context—of runoffs with different timescales (Mudelsee, 2007; Di Dato et al., 2023)[82,83].

For technical reasons, we assume the following conditions for  $\nu$  and  $\pi$ :

$$
\int_0^{+\infty} z^k \nu(\mathrm{d}z) < +\infty \ \ (k = 1, 2, 3, \dots) \tag{2}
$$

and

$$
\int_0^{+\infty} r^{-1} \pi\big(\mathrm{d} r\big) < +\infty \,. \tag{3}
$$

Assumption (2) means that the jumps in the supOU process admit moments of arbitrary order. This condition is not restrictive for applications to discharge, e.g., a tempered stable model:  $v(dz) = a_1 e^{-a_2 z} z^{-(1+a_3)} dz$  with  $a_1, a_2 > 0$  and  $a_3 < 1$  (e.g., Yoshioka and Yoshioka, 2024a)[42]. The tempered stable model includes the gamma distribution ( $a<sub>3</sub> < 0$ ), exponential distribution ( $a<sub>3</sub> = 0$ ), and infinite-activity case  $(a_3 \in [0,1])$ ,  $\int_0^{+\infty} \nu(\text{d}z) = +\infty$  in a unified framework. The assumption (3) ensures that the distribution of the recession rate is not singular at the origin  $r = 0$ . This condition is satisfied for the gamma distribution  $\pi(\text{d}r) = \gamma_{\alpha,\beta}(r) \text{d}r = \frac{1}{\Gamma(\alpha)\beta^{\alpha}} r^{\alpha-1} e^{-r}$  $\pi(\mathbf{d}r) = \gamma_{\alpha,\beta}(r) \mathbf{d}r = \frac{1}{\Gamma(\alpha)\beta^{\alpha}} r^{\alpha-1} e^{-r/\beta} \mathbf{d}r$  with  $\alpha > 1$  and  $\beta > 0$ . Together,

assumptions (2) and (3) allow the stationary moments and autocorrelation of the supOU process (1) to be defined, as shown in **Section 3**.

#### **2.3 Water quality index**

## **2.3.1 Seasonal decomposition**

We propose an SDE model for the WQI of a river. The WQI  $C = (C_t)_{t \in \mathbb{R}}$ , representing the concentration of a solute, is assumed to be nonnegative and consists of seasonal and stationary parts (Yoshioka and Yoshioka, 2024b)[43]:

$$
C_t = \overline{C}e^{S_t + X_t}, \ t \in \mathbb{R} \,. \tag{4}
$$

Here,  $\overline{C} > 0$  is a constant;  $S = (S_t)_{t \in \mathbb{R}}$  is a deterministic, periodic, real-valued coefficient; and  $X = (X_t)_{t \in \mathbb{R}}$  is the logarithm of the stationary part that encodes the (hidden) long memory of WQI. Under this formulation, water quality investigations aim to identify the two processes *X* and *<sup>Y</sup>* , as well as the periodic part *<sup>S</sup>* .

We assume annual periodicity ( $T = 1$  year or  $T = 365.25$  days, adjusted for leap years) and expressed the seasonal part as follows:

$$
S_{t} = \sum_{i=1}^{n} A_{i} \sin\left(\frac{2\pi t}{T} + B_{i}\right), \ t \in \mathbb{R}
$$
 (5)

with some  $n \in \mathbb{N}$  and real parameters  $A_i$  and  $B_i$ . The memory of the WQI is assumed to be encoded in *<sup>X</sup>* , which is described using a stochastic volatility model. We therefore focus on mathematically modeling X. Without loss of generality, we set  $\mathbb{E}[X_t] = 0$  for all  $t \in \mathbb{R}$ , where  $\mathbb{E}$  is expectation.

#### **2.3.2 Classical stochastic volatility model**

The core of the proposed model is the SDE for the auxiliary scalar process  $X$ . A classical stochastic volatility model takes the form (e.g., Barndorff-Nielsen and Stelzer, 2013)<sup>[44]</sup>: for  $t \in \mathbb{R}$ ,

$$
dX_t = -RX_t dt + \sigma \sqrt{RY_t} dB_t
$$
\n(6)

or equivalently

$$
X_t = \sigma \int_{-\infty}^t \sqrt{RY_s} e^{-R(t-s)} dB_s , \qquad (7)
$$

where  $B = (B_t)_{t \in \mathbb{R}}$  is a (double-sided) standard Brownian motion independent of the Poisson random measure N,  $R > 0$  is the reversion rate, and  $\sigma > 0$  is the noise intensity. The last term in (6) is defined in the Itô's sense. The SDE in (6) is driven by a noise term with stochastic volatility  $\sigma \sqrt{RY_{t}}$ .

A key limitation of the SDE in (6) is that it results in an exponential autocorrelation. Indeed, from (7), for any  $h \ge 0$  and  $t \in \mathbb{R}$  (noting that  $\mathbb{E}[X_t] = 0$ ):

$$
\mathbb{E}\left[X_{t}X_{t+h}\right] = \mathbb{E}\left[\sigma\int_{-\infty}^{t}\sqrt{RY_{s}}e^{-R(t-s)}dB_{s}\sigma\int_{-\infty}^{t+h}\sqrt{RY_{u}}e^{-R(t+h-u)}dB_{u}\right]
$$
\n
$$
= \sigma^{2}Re^{-2Rt-Rh}\mathbb{E}\left[\int_{u=-\infty}^{u=t+h}\int_{s=-\infty}^{s-t}\sqrt{Y_{s}Y_{u}}e^{Rs+Ru}dB_{s}dB_{u}\right]
$$
\n
$$
= \sigma^{2}Re^{-2Rt-Rh}\int_{s=-\infty}^{s-t}\mathbb{E}\left[Y_{s}\right]e^{2Rs}ds
$$
\n
$$
= \frac{\sigma^{2}\overline{Y}}{2}e^{-Rh}
$$
\n(8)

Here, we used the Itô's isometry (e.g., Theorem 10.1.6 of Pascucci (2024))[11] and the stationarity of *<sup>Y</sup>* , with the average denoted by *<sup>Y</sup>* . The autocorrelation decays exponentially as shown in (8). Another drawback of (6) is that X does not correlate with Y. Indeed, for  $t \in \mathbb{R}$ ,

$$
\mathbb{E}\left[X_{t}Y_{t}\right] = \mathbb{E}\left[\sigma\int_{-\infty}^{t}\sqrt{RY_{s}}e^{-R(t-s)}dB_{s}Y_{t}\right]
$$
\n
$$
= \mathbb{E}\left[\sigma\int_{-\infty}^{t}\sqrt{RY_{s}}e^{-R(t-s)}dB_{s}\int_{u=-\infty}^{u=t}\int_{z=0}^{z=\infty}\int_{r=0}^{r=\infty}ze^{-r(t-u)}N\left(\mathrm{d}u,\mathrm{d}z,\mathrm{d}r\right)\right].
$$
\n(9)\n
$$
= 0
$$

This lack of correlation restricts the model's applicability, as it assumes no interaction between discharge and water quality. Such an assumption is simplistic for real-world scenarios, including the application study presented in **Section 4**.

# **2.3.3 SDE model**

The two drawbacks, exponential and hence short memory and no correlation to discharge, are overcome by the proposed supOUSV model. For  $t \in \mathbb{R}$ ,

$$
X_t = \underbrace{\int_{R=0}^{R=+\infty} Z_t \left( dR \right)}_{\text{Superposition to generate long memory}}
$$
(10)

with 
$$
Z = (Z_t (dR))_{t \in \mathbb{R}}
$$
 being a measure-valued small process governed by a formal SDE for  $R > 0$ :  
\n
$$
dZ_t (dR) = -R(Z_t (dR) - \mu (Y_t - \overline{Y}) \rho (dR)) dt + \underbrace{\sigma \sqrt{RY_t} B (dR, dt)}_{\text{Volatility-driven fluctuation}},
$$
\n(11)

also expressed as

$$
Z_{t}(\mathrm{d}R) = \int_{s=-\infty}^{s=t} e^{-R(t-s)} \left\{ R \mu \left( Y_{s} - \overline{Y} \right) \rho \left( \mathrm{d}R \right) \mathrm{d}s + \sigma \sqrt{RY_{s}} B(\mathrm{d}R, \mathrm{d}s) \right\}.
$$
 (12)

Here,  $B(dR, dt)$  is a space-time Gaussian random measure (e.g., Gomez et al., 2023)[84] with covariance  $[B(dR, dt)B(dR', dt')] = \delta(R - R')\delta(t - t')\rho(dR)\rho(dR')dtdt'$  is a probability measure  $\rho(dR)$  of the distributed reversion  $R > 0$ ,  $\mu \in \mathbb{R}$  modulates the dependence of the water quality on discharge, and  $\delta$  is the Dirac delta. We assume that the random measures  $B$  and  $N$  are mutually independent. Assuming a stationary state, the system (10) is rewritten as

$$
X_{t} = \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} \left\{ R \mu \left( Y_{s} - \overline{Y} \right) \rho \left( dR \right) ds + \sigma \sqrt{RY_{s}} B \left( dR, ds \right) \right\}.
$$
 (13)

The supOUSV model is a system containing (1) and (13), and the WQI is obtained from (4).

# **3. Mathematical analysis**

#### **3.1 Moments and autocorrelation**

# **3.1.1 supOU process**

The moments (and hence cumulants) and autocorrelation of the supOUSV model can be obtained explicitly under certain conditions. The cumulants and autocorrelation of the supOU process are as follows (e.g., Barndorff-Nielsen and Stelzer, 2013; Yoshioka, 2022)[44,85]. Here,  $\nabla$ ,  $\mathbb{S}$ , and  $\mathbb{K}$  are variance, (unnormalized, and hence not divided by using  $V$ ) skewness, and kurtosis, respectively, and AC<sub>*Y*</sub> (*h*) is the autocorrelation of Y with the time lag  $h \geq 0$ :

$$
\mathbb{E}\big[Y_{t}\big]=\overline{Y}=M_{1}\int_{0}^{+\infty}\frac{\pi\big(\mathrm{d}r\big)}{r},\ \ \mathbb{V}\big[Y_{t}\big]=\overline{V}=\frac{M_{2}}{2}\int_{0}^{+\infty}\frac{\pi\big(\mathrm{d}r\big)}{r},\tag{14}
$$

$$
\mathbb{S}\left[Y_{t}\right] = \frac{M_{3}}{3}\int_{0}^{+\infty} \frac{\pi\left(\mathrm{d}r\right)}{r}, \ \mathbb{K}\left[Y_{t}\right] = \frac{M_{4}}{4}\int_{0}^{+\infty} \frac{\pi\left(\mathrm{d}r\right)}{r}, \tag{15}
$$

and

$$
AC_{Y}(h) = \left(\int_0^{+\infty} \frac{\pi(dr)}{r}\right)^{-1} \int_0^{+\infty} \frac{e^{-rh}\pi(dr)}{r}.
$$
 (16)

If  $\pi$  has the gamma density  $\gamma_{\alpha_r,\beta_r}$  with  $\alpha_r > 1$  and  $\beta_r > 0$ , then the autocorrelation exhibits a subexponential decay:

$$
AC_{Y}(h) = \frac{1}{(1 + \beta_{r}h)^{\alpha_{r}-1}}.
$$
\n(17)

The process X is said to exhibit truly long memory if  $AC_Y(h)$  is not integrable ( $\alpha_r \in (1,2]$ ), and moderately long memory if it is ( $\alpha_r > 2$ ). In the sequel, we write  $\bar{V} = \mathbb{V}[Y_r]$ .

#### **3.1.2 supOUSV process**

Some cumulants and autocorrelations of *X* are derived in this paper. We have the following proposition concerning the average and variance of  $X$  and the covariance between  $X$  and  $Y$ , which can be utilized in applications such as the moment matching for identifying a model.

# *Proposition 1*

*For any*  $t \in \mathbb{R}$ , the followings hold true:

$$
\mathbb{E}\big[X_t\big]=0\,,\tag{18}
$$

$$
\mathbb{V}\left[X_{t}\right] = \underbrace{\frac{1}{2}\sigma^{2}\overline{Y}}_{\text{Variance due to diffusion}} + \mu^{2}\overline{V}\left(\int_{0}^{+\infty} \frac{\pi\left(\text{d}r\right)}{r}\right)^{-1} \int_{R=0}^{R=+\infty} \int_{P=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{1}{r} \frac{PR}{P+R}\left(\frac{1}{P+r}+\frac{1}{R+r}\right) \pi\left(\text{d}r\right) \rho\left(\text{d}R\right) \rho\left(\text{d}P\right) \text{ (19)} \\ \text{Variance due to discharge-dependent drift} \tag{19}
$$

*and*

$$
\mathbb{E}\left[X_{t}Y_{t}\right] = \mu\overline{V}\left(\int_{0}^{+\infty} \frac{\pi(\mathbf{d}r)}{r}\right)^{-1} \int_{R=0}^{R=\infty} \int_{r=0}^{r=\infty} \frac{R}{r(R+r)}\pi(\mathbf{d}r)\rho(\mathbf{d}R). \tag{20}
$$

The first term on the right-hand side of (19) represents the variance due to diffusion, which is proportional to  $\sigma^2$  and arises from stochastic volatility. By contrast, the second term in the right-hand side of (19) represents the variance due to the discharge-dependent drift, which is proportional to  $\mu^2$  and independent of the sign of  $\mu$ . The integrals in (19)-(20) are analytically intractable and is numerically discretized in this paper. According to (20), the sign of  $\mu$  determines the sign of the covariance, providing a potential method for distinguishing the supOUSV model in real data because it is able to deal with both positive and negative correlations between discharge and WQI by properly specifying  $\mu$ .

*Remark 1* The relationships (19)–(20) are useful for determining the associated Orlicz spaces when quantifying the impact of model uncertainty (Yoshioka and Yoshioka, 2024a)[42] in the measures  $\pi, \rho$  on statistics of X. A measure with stronger singularity (i.e., a more pronounced blow-up) at the origin corresponds to a smaller Orlicz space and is more sensitive to modeling errors.

We also obtain the closed-form expression for the autocorrelation  $AC_x(h)$  of X.

# *Proposition 2*

*For any*  $t \in \mathbb{R}$  and  $h \ge 0$ , the autocorrelation  $AC_X(h)$  is expressed as follows:

$$
AC_{X}(h) = \frac{1}{\mathbb{V}[X_{t}]} \left( \underbrace{\frac{\sigma^{2} \overline{Y}}{2} I_{1}(h)}_{\text{Correlation due to diffusion}} + \mu^{2} \overline{V} \left( \int_{0}^{+\infty} \frac{\pi (dr)}{r} \right)^{-1} (I_{2}(h) + I_{3}(h)) \right), \tag{21}
$$

*where*

$$
I_1(h) = \int_{R=0}^{R=+\infty} e^{-Rh} \rho(R) dR,
$$
 (22)

$$
I_2(h) = \int_{R=0}^{R=\infty} \int_{P=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r(P+R)} \left( \frac{1}{P+r} + \frac{1}{R+r} \right) e^{-Ph} \pi \left( dr \right) \rho \left( dR \right) \rho \left( dP \right), \tag{23}
$$

*and*

$$
I_{3}(h) = \int_{R=0}^{R=+\infty} \int_{P=0}^{P=+\infty} \int_{r=0}^{r=+\infty} \frac{RP}{r(R+r)(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(dr) \rho(dR) \rho(dP). \tag{24}
$$

The equation (21) demonstrates how drift and diffusion, and hence the two driving noises *<sup>N</sup>* and *B* contribute to the autocorrelation. The weighting between drift and diffusion is determined by the relative strengths of stochastic volatility and the correlation to discharge,  $I_k(h)$  ( $k = 1, 2, 3$ ), which may behave differently at large time lag  $h > 0$ , as studied below.

While **Proposition 2** formally represents  $AC_X(h)$ , it does not address the existence or decay rate, which is analyzed below. In the rest of **Section 3.1.2**, we assume the following gamma distributions to quantify the tail behavior of  $I_k$  in a tractable manner:

$$
\pi(dR) = \gamma_{\alpha_r, \beta_r}(R) dR \text{ and } \rho(dR) = \gamma_{\alpha_R, \beta_R}(R) dR \text{ with } \alpha_r - 1, \alpha_R, \beta_r, \beta_R > 0. \tag{25}
$$

We have the following estimate of  $AC_x(h)$  at large  $h > 0$ .

## *Proposition 3*

*Assume (25) with*

$$
(\alpha_r - 1)(\alpha_R + 2) > 1 \text{ if } \alpha_r \in (1, 2]. \tag{26}
$$

*Then, it follows that*  $\lim_{h\to\infty} AC_x(h) = 0$  *as*  $h\to\infty$ *. Moreover, if*  $\alpha_r > 2$ *, then* 

$$
AC_x(h) \le \sigma^2 O\big(h^{-\alpha_R}\big) + \mu^2 \Big\{O\big(h^{-\alpha_R}\big) + O\big(h^{-(\alpha_r - 2)}\big)\Big\} \,. \tag{27}
$$

*If*  $\alpha_r \in (1,2]$ , then

$$
AC_x(h) \le \sigma^2 O\big(h^{-\alpha_R}\big) + \mu^2 \big\{O\big(h^{-\alpha_R}\big) + O\big(h^{-\eta}\big)\big\} \tag{28}
$$

*with*

$$
\eta = \frac{(\alpha_r - 1)(\alpha_R + 2) - 1}{\alpha_R + 2} > 0.
$$
\n(29)

According to **Propositions 2** and 3, the term with the heaviest tail is  $I_3$  when  $\alpha_R$  is large at which the memory of river discharge (i.e., flood events) dominates. Specifically, the case  $\mu \neq 0$  arises due to the coupling of the two superposed processes and is a unique feature of the proposed model, and the tail of AC<sub>X</sub> (h) in this case possibly becomes the longer than that with  $\mu = 0$ . An engineering implication of **Proposition 3** is that autocorrelation may decay very slowly or even fail to converge to zero if the noise components of X and Y have sufficiently long memory, characterized by small  $\alpha_r, \alpha_R$ , as implied in (29).

# **3.2 Generalized Riccati equation**

We have investigated the process  $X$ , while what directly contributes to water quality is its exponential, i.e.,  $e^X$ . We therefore analyze when the exponential moment of the form  $M(q) = \mathbb{E}\left[e^{qX_t}\right]$  ( $q \in \mathbb{R}$ ) exists. We focus on cases  $q > 0$  because cases  $q \leq 0$  are trivial.

It is well known that, for a finite-dimensional system of affine SDEs, the moment-generating function is governed by a generalized Riccati equation as a nonlinear ordinary differential equation (e.g., Duffie et al., 2003)[45]. Because our system of SDEs is infinite-dimensional and consists of measure-valued processes, the direct derivation of the associated generalized Riccati equation may not be straightforward. However, starting with a finite-dimensional version of the supOUSV model, we can infer the form of the desired generalized Riccati equation. We present the discussion based on a finite-dimensional system in the **Section A. 1** of **Appendix** and only present the result here. In a stationary state, we have

$$
M(q) = \exp\left(\lim_{t \to +\infty} \phi_t\right),\tag{30}
$$

where

$$
\frac{\mathrm{d}\phi_t}{\mathrm{d}t} = \int_0^{+\infty} \int_0^{+\infty} \left( e^{\psi_t(R)z} - 1 \right) \nu \left( \mathrm{d}z \right) \pi \left( \mathrm{d}r \right) - \mu \int_0^{+\infty} P \omega_t \left( P \right) \rho \left( \mathrm{d}P \right), \ t > 0 \,, \tag{31}
$$

$$
\frac{\partial \omega_t(R)}{\partial t} = -R\omega_t(R), \ t, R > 0,
$$
\n(32)

and

$$
\frac{\partial \psi_t(r)}{\partial t} = -r\psi_t(r) + \mu q \int_0^{+\infty} \omega_t(P) P \rho(dP) + \frac{\sigma^2}{2} \int_0^{+\infty} {\{\omega_t(P)\}}^2 P \rho(dP), \ t, r > 0 \tag{33}
$$

subject to initial conditions:

$$
\omega_0(\cdot) = q \text{ and } \psi_0(\cdot) = \phi_0 = 0. \tag{34}
$$

By (33) and (34), we have

$$
\psi_t(r) = \int_{P=0}^{P=\infty} \int_{s=0}^{s=r} P e^{-r(t-s)} \left( \mu q e^{-Pt} + \frac{\sigma^2 q^2}{2} e^{-2Pt} \right) ds \rho(P) dP. \tag{35}
$$

The well-posedness of the differential equation part (33) is evident because it is a continuum of linear ordinary differential equations, and  $\psi_t(\cdot) \to 0$  as  $t \to 0$ . To analyze the generalized Riccati equation in more detail, we assume the tempered stable model  $v(dz) = a_1 e^{-a_2 z} z^{-(1+a_3)} dz$ . To ensure  $M(q)$  is welldefined, by (31) we need to have a bound  $\psi_t(r) \le a_2$  for all  $t \ge 0$  and  $r > 0$ . The following proposition gives a sufficient condition for  $M(q)$  to exist.

# *Proposition 4*

Assume that  $v(dz) = a_1e^{-a_2z}z^{-(1+a_3)}dz$  with  $a_1, a_2 > 0$  and  $a_3 < 1$ . To ensure  $M(q) < +\infty$ , it is necessary to *have* 

$$
\max\{\mu, 0\} q + \frac{\sigma^2 q^2}{4} \le a_2 e \ .
$$
 (36)

*Remark 2* Moment explosion phenomena have been reported using theoretical criteria for Markovian models (Deelstra et al., 2019)[86] and stochastic Volterra models (Gerhold et al., 2019)[87].

We set  $q_{\text{max}}$  as the maximum q satisfying (36), which is found as follows:

$$
q_{\text{max}} = \frac{2a_2e}{\mu + \sqrt{\mu^2 + \sigma^2 a_2 e}}.
$$
\n
$$
(37)
$$

From (4), the moments  $\mathbb{E}\left[C_t^k\right]$  of the concentration C exist only if  $k \leq q_{\text{max}}$ . Physically, higher-order moments are undefined for WQIs that fluctuate more (i.e., larger  $\sigma$ ) and are more positively correlated with the discharge Y (i.e., larger  $\mu$ ). These effects are due to the jump-driven nature of the discharge Y, where high value of *Y* may induce high excitations of *<sup>X</sup>* .

The condition (36) is specific to the tempered stable  $\nu$ , and becomes unnecessary if we use the generalized model  $v(dz) = a_1e^{-a_2z^{a_4}}z^{-(1+a_3)}dz$  with  $a_1, a_2 > 0$ ,  $a_3 < 1$ , and  $a_4 > 1$  (e.g., Grabchak, 2021)[88]. A drawback of the generalized tempered stable model is the increase in model parameters and the necessity for a more complicated numerical algorithm for simulating supOU processes. The generalized model is advantageous for stationary statistics but not for pathwise properties because moments *M<sup>k</sup>*  $(k = 1, 2, 3, \dots)$  are critical. The case study in **Section 4** shows that the tempered stable model fails in applications when  $q_{\text{max}} < 1$ ; in such cases, the average C does not exist. This issue can be efficiently avoided with a tempered stable  $\nu$  by redefining the supOU process from (1) as follows:

$$
Y_{t} = \int_{s=-\infty}^{s=t} \int_{z=0}^{z=\infty} \int_{r=0}^{r=-\infty} \frac{1}{z^{1+\varepsilon}} e^{-r(t-s)} N\big(\mathrm{d}s, \mathrm{d}z, \mathrm{d}r\big) , \ t \in \mathbb{R}
$$
 (38)

with a regularization parameter  $\varepsilon > 0$ . Accordingly, the generalized Riccati equation is modified as follows: (31) becomes:

$$
\frac{d\phi_t}{dt} = \int_0^{+\infty} \int_0^{+\infty} \left( \exp\left(\psi_t\left(r\right)z^{\frac{1}{1+\varepsilon}}\right) - 1 \right) \nu\left(\frac{dz}{\pi}\left(dr\right) - \mu \int_0^{+\infty} R\omega_t\left(R\right) \rho\left(\frac{dR}{dt}\right), \ t > 0, \tag{39}
$$

while (32)–(33) remain unchanged. Moments and autocorrelation functions of the processes X and Y are

also unchanged, except that  $M_k$  ( $k = 0,1,2,...$ ) must be formally replaced by  $z^k$  by  $z^{k/(1+\varepsilon)}$  in (2). An advantage of this regularization (38) is that it allows the use of simple numerical algorithms for simulating tempered stable subordinators (e.g., the rejection sampling method in Algorithm 0 of Kawai and Masuda (2011)[89]). Moreover, this regularization does not alter the overall structure of the proposed model but completely avoids the theoretical restriction in (36). Additionally, this regularization is qualitatively equivalent to employing a generalized tempered stable model in terms of moments, as shown by the following equality: for any  $p > 0$ ,

$$
\int_0^{+\infty} z^{\frac{p}{1+\varepsilon}} \frac{\exp(-c_2 z)}{z^{1+c_3}} dz = (1+\varepsilon) \int_0^{+\infty} z^p \frac{\exp(-c_2 z^{1+\varepsilon})}{z^{1+(1+\varepsilon)c_3}} dz , \qquad (40)
$$

where the right-hand side corresponds to the  $p$  th moment of a generalized tempered stable model. Thus, the two models are interchangeable in this sense.

# **4. Application**

## **4.1 Study site**

The study site is the Kisuki point in the midstream of the Hii River (a Class-A River in Japan), located in the eastern part of Shimane Prefecture (**Figure 2**). The watershed of the whole Hii River system covers an area of 2,540 (km<sup>2</sup>), with the main river stream extending 153 km. Approximately 500,000 people reside within the watershed (MLIT, 2024a)[90]. According to Takeda (2023)[91], the catchment area of the Kisuki point is 451 km<sup>2</sup> and is predominantly forest (83.5%), with farmland (8.7%) and residential areas (0.7%). Theu, the Kisuki point has a rural catchment. The mainstream Hii River also contains two cascading brackish lakes, Lake Shinji and Lake Nakaumi. These lakes serve as Ramsar sites, providing unique habitats for migratory birds and functioning as important wetlands (MLIT, 2024b)[92]. The biology (Kawaida et al., 2024)[93], hydrodynamics (Hafeez and Inoue, 2024)[94], and water quality (Kim et al., 2024)[95] of these brackish lakes have been studied. The water quantity and quality of the mainstream Hii River are critical in determining the characteristics of these lakes.

The water quality at Kisuki point has been investigated by Takeda (2023)[91], focusing on multiple indices, including total nitrogen (TN: August 20, 1991, to December 27, 2022), total phosphorus (TP: August 20, 1991, to December 27, 2022), and total organic carbon (TOC: April 5, 2005, to December 27, 2022). These indices were sampled and measured almost weekly from 1991 to 2022. Part of this dataset has been investigated by Yoshioka and Yoshioka (2024a)[42] using a superposition of diffusion processes without incorporating discharge data. This paper extends their analysis by including enhanced data from the 2022 survey, courtesy of Dr. Ikuo Takeda, the author of Takeda (2023)[91]. TN and TP are WQIs that determine environmental criteria for water use and eutrophication in aquatic systems (Atazadeh et al., 2024; Tian et al., 2024; Yuan and Paul, 2024)[96-98]. TOC is an effective trace element for carbon cycling, both locally in river environments and globally (Biedunkova and Kuznietsov, 2024; Starr et al., 2023)[99,100].

Another dataset used in this study is the weekly to monthly time series of dissolved silica (DSi,

amount of Silicon in water samples) measured in a 1-km river reach, including Kisuki point, from May 22, 2018, to December 1, 2022. Part of this dataset has been investigated in previous studies (Yoshioka and Yoshioka, 2023a; Yoshioka and Yoshioka, 2023b)[24,25]. The data are treated as representative of the Kisuki point, assuming longitudinal homogeneity of water quality over this short reach. DSi has been selected as a WQI because it does not correlate with flow discharge and serves as a conservative tracer for tracking hydrological pathways in different regions (Das et al., 2022; Hachgenei et al., 2024; Jewell et al., 2023)[101-103]. For DSi, the data from May 2, 2022, to December 1, 2022, are excluded from autocorrelation analysis due to a 150-day gap from the just before sampling date.

The WQIs are measured using the following methods: TN is determined by UV spectrophotometry after potassium peroxodisulfate decomposition; TP by molybdenum blue spectrophotometry after potassium peroxodisulfate decomposition; TOC using a TOC analyzer (TOC-Vcsn, Shimadzu); and DSi by ICP emission spectrometry (ICPE9000, Shimadzu). Hourly river discharge data for Kisuki point (MLIT,  $2024c$ )[104] from January 1, 2017, at 01:00 to December 31, 2022, at 24:00 are also used in this study. Time-series data for discharge and WQIs are presented in **Figures 3**-**4**. The average, variance, and skewness of the discharge and each WQI are summarized in **Table 1**. The discharge is positively skewed due to the balance between persistent baseflow and flood events. TN, TP, and TOC are also positively skewed, while DSi is negatively skewed, though the magnitude of skewness for DSi is an order smaller for DSi than for the other three indices.



**Figure 2.** Study site and its surrounding area.



**Figure 3.** Hourly time-series data of discharge at Kisuki point.



**Figure 4.** Time-series data of discharge and WQIs at Kisuki point: (a) TN, (b) TP, (c) TOC, and (d) DSi. Time shifts have been chosen for purely visualization purposes.

**Table 1.** Average, variance, and skewness of discharge-end WQI. Units of variables are discharge  $(m^3/s)$ ,

TN (mg/L), TP (mg/L), TOC (mg/L), and DSi (mg/L)

	Discharge	TN	TР	TOC.	DSi
Average	$1.701.E+01$	5.964 F-01	3.285 E-02	8.634.E-01	$5.869.E+00$
Variance	$8.308.E+02$	8.369 E-02	1.841.E-03	4.640 F-01	9.469 F-01
<b>Skewness</b>	1 406 E+01	1 056 E+01	$1574E+01$	$1.014.E+01$	$-7.141.E-01$

#### **4.2 Model identification**

The supOUSV model for each WQI is identified as follows: We assume gamma distributions as in (25) and a tempered stable type with regularization, as in (38). The supOU process of discharge, which is common to all indices, is first identified using the moment-matching methodology (e.g., Yoshioka, 2022; Yoshioka and Yoshioka, 2024a)[42,85]. Moment-matching and estimation have been widely employed for finite superposition SDEs (Wu and Hu, 2024)[105] and Volterra processes (Abi Jaber et al., 2024)[106]. Here, the autocorrelation is fitted by minimizing the least-squares error between the empirical and theoretical values from (17) to identify  $\pi$ , and the Lévy measure  $\nu$  is identified by minimizing the sum of relative errors in the average, variance, and skewness:

$$
\left(\frac{\text{Ave}_{\varepsilon}(Y) - \text{Ave}_{\varepsilon}(Y)}{\text{Ave}_{\varepsilon}(Y)}\right)^{2} + \left(\frac{\text{Var}_{\varepsilon}(Y) - \text{Var}_{\varepsilon}(Y)}{\text{Var}_{\varepsilon}(Y)}\right)^{2} + \left(\frac{\text{Skew}_{\varepsilon}(Y) - \text{Skew}_{\varepsilon}(Y)}{\text{Skew}_{\varepsilon}(Y)}\right)^{2},\tag{41}
$$

where Ave, Var, and Skew represent average, variance, and skewness of discharge *<sup>Y</sup>* , respectively. Subscripts "e" and "m" in (41) represent empirical and modeled values. Similar notations apply to *<sup>X</sup>* . The time lags for identifying autocorrelation are set at 30 days for discharge and 730 days for WQIs. We find that the identified models are not critically different when different lag durations are used (**Section A. 3**). We assume that the theoretical average of *X* equals zero for each concentration (consistent with **Table 5** presented later).

After identifying the supOU process, the supOUSV model for each WQI, that is, *X* is identified. The seasonality  $\bar{C}e^{S_t}$  is determined by least-squares fitting between  $\ln(\bar{C}e^{S_t})$  and observed concentrations In  $C<sub>t</sub>$ , ensuring that the residual component  $X<sub>t</sub>$  has a numerically close-to-zero mean. Empirically,  $n = 2$ is fixed to resolve the yearly variation of each WQI. Once seasonality component is identified,  $e^{X_t}$  is calculated as  $C_t / (\bar{C}e^{S_t})$ , and hence  $X_t$  by taking the logarithm. We need to identify the three quantities, which are the probability measure  $\rho$  and parameters  $\sigma$  and  $\mu$ . For DSi, which has a small correlation

between X and Y, we set  $\mu = 0$  and fitted  $\sigma$  using (19) as follows:  $\sigma = \sqrt{\frac{2 \text{Var}_{\epsilon}(X)}{N}}$  $\frac{e^{(-1)}}{e^{(-Y)}}$ 2Var Ave *X*  $\sigma = \sqrt{\frac{2 \times m_e (Y)}{\text{Ave } (Y)}}$ . The measure  $\rho$ 

is then fitted using a least-squares method based on (21) because, in this case, we have  $AC_{X}(h) = (1 + \alpha_{R}h)^{-\beta_{R}}$  ( $h \ge 0$ ).

For other indices (e.g., TN, TP, TOC), where the correlation between *X* and *Y* is not negligible  $O\big( 10^{-1} \big),$  the above strategy does not apply. Instead, we used a two-step strategy. The temporal model assumption  $\mu = 0$  is employed, and then we fit the auxiliary parameter  $\alpha_{R,a}, \beta_{R,a}$  ( $h \ge 0$ ). Next, using a trial-and-error method, we determine the parameter set  $(\alpha_R, \sigma, \mu)$ , as follows: Fix  $\beta_R = \beta_{R,a}$  and rewrite the autocorrelation in (21) as

$$
AC_x(h) = \frac{I_1(h) + w(I_2(h) + I_3(h))}{I_1(0) + w(I_2(0) + I_3(0))}, \quad h \ge 0
$$
\n(42)

with the weight

$$
w = \frac{2\mu^2 \text{Var}_e(Y)}{\sigma^2 \text{Ave}_e(Y)} \left( \int_0^{+\infty} \frac{\pi(\text{d}r)}{r} \right)^{-1}.
$$
 (43)

Here, we test several values of  $\alpha_R$  around  $\alpha_{R,a}$  (with an interval of 0.005 for TN and TP, and 0.05 for TOC). For each  $\alpha_R$ , we find w such that the least-squares error between empirical and theoretical autocorrelation (42) is minimized. Having obtained the optimal w for the given  $\alpha_R$ , we obtain  $\sigma$  from

$$
Var_e(X) = \frac{1}{2}\sigma^2 Ave_e(Y)\{I_1(0) + w(I_2(0) + I_3(0))\}
$$
\n(44)

or equivalently

$$
\sigma = \sqrt{\frac{2\text{Var}_{\text{e}}(X)}{\text{Ave}_{\text{e}}(Y)\{I_1(0) + w(I_2(0) + I_3(0))\}}},\tag{45}
$$

Finally, we find  $\mu$  from (43) as follows:

$$
\mu = \frac{\mathbb{E}\left[X_{\cdot}Y_{\cdot}\right]}{\left|\mathbb{E}\left[X_{\cdot}Y_{\cdot}\right]\right|} \sigma \sqrt{\frac{wAve_{e}\left(Y\right)}{2Var_{e}\left(Y\right)} \int_{0}^{+\infty} \frac{\pi\left(dr\right)}{r}}
$$
(46)

ensuring that the signs of  $\mu$  and  $\mathbb{E}[X_t, Y_t]$  are consistent (assuming that the latter is not zero). After obtaining  $\sigma$  and  $\mu$  for a given  $\alpha<sub>R</sub>$ , the relative difference between empirical and theoretical correlations of *<sup>X</sup>* and *<sup>Y</sup>* is evaluated based on (20). The best parameter set is then selected by minimizing this difference. A key advantage of this method is that the theoretical variance exactly matches the empirical variance due to (45).

This empirical fitting strategy was chosen in this study due to direct application of continuous optimization methods for identifying X is hindered by its complex and non-explicit dependence on  $\alpha_R$ ,  $\beta_R$ . As such, the identified parameters in this paper are suboptimal. The integrals in the autocorrelation (21) are evaluated numerically using quantile-based quadrature (e.g., Yoshioka et al., 2023; Yoshioka and Yoshioka, 2024a)[42,75] and a forward-in-time explicit discretization is applied to the finite-dimensional version of the supOUSV model in **Section A. 1**. The computational resolution, which are specified by the degree-of freedoms  $I_r$ ,  $I_R$  in Section A. 1, are set as 2,048. The time increment of 0.01 day is used. Sampling X of each WQI is conducted during a 1,000-year period after another 1,000-year period for burn-in with a deterministic initial condition.

**Remark 3** Fitting the autocorrelation of a WQI under the assumption  $\mu$  infers the decay rate with respect to the time lag h, i.e.,  $AC_x(h) \sim h^{-\alpha_R}$  for large  $h > 0$ .

## **4.3 Results and discussion**

**Tables 2–3** summarize the identified parameter values. For discharge, models with  $(\varepsilon = 0.1)$  and without  $(\varepsilon = 0)$  regularization are examined. From **Table 2**, it is evident that the model without regularization  $(\varepsilon = 0)$  fails for all indices except DSi, due to  $q_{\text{max}} < 1$ , which results in the nonexistence of the exponential moment of X in the mean. This highlights the necessity of the proposed regularization method for this case study. Sensitivity of the discharge model to different values of  $\varepsilon$  is explored in **Section A. 4** of **Appendix**, showing that the regularization introduced by  $\varepsilon$  does not qualitatively affect the computational results in this section. We therefore focus on the model with  $\varepsilon = 0.1$ . **Table 3** shows that the stochastic components of the four WQIs have long memories that would not be adequately captured by exponential ones. Particualrlly, those of TN and TP exhibit truly long memories.

In view of **Remark 1**, among the four WQIs, TN is the most vulnerable to modeling errors, followed by TP, TOC, and DSi. This is because smaller  $\alpha<sub>R</sub>$  implies less robustness. Moreover, evaluation of modeling uncertainty in TN should pay more attention than the other indices in this paper because the Orlicz space associated with TN is the smallest among the four indices in this paper (e.g., Proposition 1 of Yoshioka and Yoshioka (2024a))[42]. More specifically, assume that some modeling errors are included in the probability measure  $\rho$  and it is the Gamam type as considered in this section. Then, an integrable function against  $\rho$  should not have the singularity of the form  $O(R^{-\gamma})$  with  $\gamma \ge \alpha_R$  near the origin  $R = 0$ , or equivalently

$$
\int_0^{\kappa} R^{-\gamma} \rho \big( dR \big) = \begin{cases} \langle +\infty & (\gamma < \alpha_R) \\ +\infty & (\gamma \ge \alpha_R) \end{cases} \tag{47}
$$

for any small  $\kappa > 0$ . The theory of model uncertainty (Yoshioka and Yoshioka (2024a))[42] then suggests that the worst-case upper-bound of X is more sensitive to the index with a smaller  $\alpha_R$ ; namely, a process having a longer memory is more vulnerable to misspecification of this type. Consequently, TN having the smallest  $\alpha_R$  should be sampled more intensively to enable more accurate modeling at the study site.

**Table 4** compares the empirical and theoretical averages, variances, and skewnesses of discharge, showing strong agreement with relative errors smaller than 0.001%. **Table 5** shows the empirical statistics of *<sup>X</sup>* for each WQI, where the regularized model is slightly more accurate for cumulants. **Figure 5**  compares the empirical and theoretical autocorrelations of *<sup>Y</sup>* , computed from the identified models. Similarly, **Figure 6** compares the empirical and theoretical autocorrelations of X (with  $\mu \neq 0$  for TN, TP, and TOC). Table 6 compares the empirical and theoretical covariances between  $X$  and  $Y$ , suggesting that the identified model successfully captures the empirical correlations. Incorporating the discharge dependence in the SDE (11) through the parameter  $\mu \neq 0$  is essential to obtain this result.

The empirical and computed probability density functions (PDFs) of *<sup>X</sup>* for each WQI are compared in **Figure 7**. **Figure 8** shows the same comparison on a logarithmic scale to better visualize the PDF tails. **Figure 9** shows that the burn-in period of 1,000 years is sufficiently long so that fluctuations in the process *X* is close to being stationary. As demonstrated in **Figures 7**-**8**, the identified models capture the empirical results, accurately representing the height and location of the PDF mode. The underestimation of the mode is due to the bounded range of empirical *X* data, which is necessarily concentrated around the origin, while the model encompasses the entire range of *<sup>X</sup>* . Moreover, the model predicts the PDF tails,

which are not always clear in the empirical data. The positive correlation between X and discharge Y manifests as heavier right tails in the theoretical PDFs for TN, TP, and TOC, whereas such behavior is absent for DSi.

Finally, we investigate the so-called concentration-discharge curve at flood events to reveal the complex hysteresis relationship between discharge and concentration of a WQI. We note that the water quality data used in this study were sampled weekly, and hence the direct use of the empirical data fails in this task. **Figure 10** shows some flood events and TN concentration generated by the Monte Carlo simulation and their concentration-discharge curve, which was extracted from a one-year sample path with the average of *C* being 0.617 mg/L (empirical value is 0.596 mg/L) and the correlation between *Y* and *C* being 0.389 (empirical value is 0.441, and hence both are positive). In **Figure 10**, we plot the concentrationdischarge curve by using daily-averaged *Y* and *C* so the shape of the curve becomes better visible. The resulting concentration-discharge curve is clockwise (**Figure 10(b)**) or counterclockwise (**Figure 10(d)**) depending on flood events. Event-based concentration-discharge curve directions have been reported for total reactive phosphorus around a Great Lakes (Dialameh and Ghane, 2023)[107], TN for a rural catchment (Rodríguez-Blanco et al., 2023)[108], dissolved organic carbon in a subtropical karst catchment (Qin et al., 2024)[63], and suspended sediment in the upper Yangtze River (Xue et al., 2024)[109]. Our model is consistent with these observations.

Physically, a concentration-discharge curve of a flood event in a river results from a complex hysteric dependence of the WQI not only on discharge but also on residual concentrations in surrounding groundwater of the river (Speir et al., 2024)[110]. Moreover, the correlation between water quantity and quality should also play a role; the WQIs considered in this paper have a positive correlation to the discharge, while some index, such as chloride and calcium ion, have a negative correlation (Cairoli et al., 2024; Knapp et al., 2022)[111,112]. Distance from the pollutant source to the river (Roberts et al., 2023)[113] and climatic and vegetation factors in watershed also affect concentration-discharge curves (Zhu et al., 2023)[114]. Such mechanisms are not directly considered in the proposed model; nevertheless, as demonstrated above, the proposed model could capture both the memory and correlation to discharge WQIs from the standpoint of infinite-dimensional SDEs, which is a novel contribution. Moreover, the proposed model, despite it is a simple model that does not account for such detailed physics, can also generate the two directions of the concentration-discharge curve found in the literature. To the best of the authors' knowledge, such a report does not exist for SDE models including long memory ones. Encoding a physicsinformed event-based relationship into the proposed model requires the use of a higher-dimensional random measures in the SDEs, which is theoretically possible, but the issue is how to identify them from (not necessarily large) field data. This would be a challenging and interesting topic to be resolved in future.

	Not regularized ( $\varepsilon = 0$ )	Regularized ( $\varepsilon = 0.1$ )
$\alpha_{r}$ (-)	$2.143.E+00$	
$\beta_r$ (1/day)	$1.034.E+00$	
$a_1$ (m <sup>3c<sub>3</sub>+<math>\frac{3\varepsilon}{1+\varepsilon}</math></sup> /s <sup>c<sub>3</sub>+<math>\frac{\varepsilon}{1+\varepsilon}</math></sup> /day)	$1.266.E+00$	$1.124.E+00$
$a_2$ (s/m <sup>3</sup> )	1.960.E-03	8.920.E-04
$a_{2}(-)$	8.084.E-01	7.500.E-01

**Table 2.** Parameters for the model of discharge *<sup>Y</sup>* .

**Table 3.** Parameter values for process *X* of each WQI.

	TN	TP	<b>TOC</b>	DSi
$\bar{C}$	5.553.E-01	$2.642.E-02$	7.676.E-01	$5.745.E+00$
$A_{\rm i}$	7.104.E-02	$-2.752.E-01$	$-2.512.E-01$	$-1.133.E-01$
$A_{2}$	$6.562.E-02$	7.639.E-02	$6.511.E-02$	$-2.330.E-02$
B <sub>1</sub>	7.198.E-01	$1.344.E+00$	5.396.E-01	$1.801.E+00$
B <sub>2</sub>	7.185.E-01	8.146.E-01	$1.640.E+00$	4.360.E-01
$\alpha_{R}^{-}$ (-) ( $\mu$ = 0)	4.217.E-01	5.477.E-01	$2.334.E+00$	$2.510.E+00$
$\beta_{R}$ (1/day)	2.699.E-01	5.253.E-01	2.506.E-02	2.806.E-02
$\alpha_{R}(\cdot)$ ( $\mu \neq 0$ )	3.750.E-01	4.850.E-01	$2.650.E+00$	
$\sigma$ (-)	1.077.E-01	1.483.E-01	1.037.E-01	5.483.E-02
$\mu$ (-)	2.752.E-02	2.917.E-02	2.567.E-02	$0.000,E+00$
$q_{\text{max}}$ (-)	1.899.E-01	1.768.E-01	2.033.E-01	$2.663.E+00$

**Table 4.** Comparison between empirical and theoretical average, variance, and skewness of discharge *<sup>Y</sup>* .

	Empirical	Theoretical ( $\varepsilon = 0$ )	Theoretical ( $\varepsilon$ = 0.1)
Average $(m^3/s)$	$1.701.E+01$	$1.701.E+01$	$1.701.E+01$
Variance $(m^6/s^2)$	$8.308.E+02$	$8.308.E+02$	$8.308.E+02$
Skewness (-)	$1.406.E+01$	$1.406.E+01$	$1.406.E+01$

**Table 5.** Statistics of process X for each WQI. Covariance and correlation between X and Y are calculated using daily-averaged *<sup>Y</sup>* .

	TN	TР	TOC	DSi.
Average of $X$ (-)	5.890.E-07	3.634.E-06	2.628.E-07	$1.120.E-06$
Variance of $X$ (-)	1.373.E-01	2.794.E-01	1.459.E-01	2.556.E-02
Skewness of $X$ (-)	$-3.947.E-01$	$1.313.E+00$	$1.304.E+00$	$-9.859.E-01$
Covariance between X and Y $(m^3/s)$	$2.671.E+00$	$5.161.E+00$	$2.436.E+00$	8.105.E-02
Correlation between $X$ and $Y$ (-)	3.531.E-01	4.786.E-01	2.986.E-01	4.467.E-02

**Table 6.** Comparison between empirical and theoretical covariance  $(m^3/s)$  between X (-) and Y  $(m^3/s)$ .





**Figure 5.** Comparison between empirical (black) and theoretical (blue) autocorrelation (AC) of discharge *<sup>Y</sup>* .



**Figure 6.** Comparison between empirical (black) and theoretical (blue) autocorrelations (AC) of X: (a) TN, (b) TP, (c) TOC, and (d) DSi.



**Figure 7.** Comparison between empirical (red) and theoretical (blue) PDFs of  $X : (a) TN, (b) TP, (c) TOC,$ and (d) DSi.



**Figure 8.** Comparison between empirical (red) and theoretical (blue) PDFs of *X* in ordinary logarithmic scale: (a) TN, (b) TP, (c) TOC, and (d) DSi.



**Figure 9.** Convergence history of computed (a) average (Ave) and (b) variance (Var) of the process X in the supOUSV model in the burn-in period. The grey line in each figure panel corresponds to theoretical value.



Figure 10. concentration-discharge relationships at flood events in model: (a) discharge *Y* m<sup>3</sup>/s and TN concentration *<sup>C</sup>* mg/L during some simulated flood events in ordinary logarithmic scale and (b) corresponding concentration-discharge curve. Panels (c) and (d) show results for different flood events. Time is suitably normalized in panels (a) and (c), respectively.

#### **5. Conclusion**

We proposed a supOUSV model for coupled water quantity-quality dynamics in a river environment. The model demonstrated flexibility in representing the long-memory nature of discharge and that included in WQIs, as well as the ability to analytically obtain moments and autocorrelation, enabling the application of straightforward model identification methods. Analysis of the generalized Riccati equation provided the condition for defining each moment of the WQI for a specific case. The supOUSV model was applied to data collected at a study site in the midstream of the Hii River, clarifying its applicability and limitations for each WQI. The innovative integration of mathematical theory and computational techniques presented in this study would enhance the understanding of water environmental dynamics and their potential management strategies.

Several limitations of the supOUSV model are identified, which should be addressed in future research to develop a more accurate model with broader applicability. First, the effects of incorporating small processes with more generic forms have not been investigated and remain an open area of study. Another principal issue not addressed in this study is the biology and ecology of river environments, where water quantity and quality significantly impact aquatic organisms. The proposed model with the exponential nature of the seasonality decomposition allows for effectively separating the deterministic and stochastic parts of the positivity-preserving WQI dynamics, but its drawback is the possible loss of theoretical stability as indicated in **Proposition 3**. We will explore another decomposition method that does not lose the tractability of the model but is more stable. While space-time stochastic modeling (e.g., Nishimoto et al., 2024; Puica and Benth, 2023)[115,116] could potentially address these interactions, the lack of spatially detailed water quality data near the study site made such modeling technically infeasible at this stage. Addressing this issue will require an intensive and large-scale survey to collect comprehensive data. This effort will be crucial for advancing the understanding of stochastic water quality and quantity dynamics and expanding the scope of the proposed model.

## **Appendix**

## **A. 1 Finite-dimensional version**

We present a finite-dimensional version of the supOUSV model and derive its associated generalized Riccati equation to enhance understanding of the infinite-dimensional versions discussed in the main text. The finite-dimensional version is formulated following Yoshioka et al. (2023)[75], who discussed a discretized supOU-type process.

Let  $I_r, I_R \in \mathbb{N}$  be the degrees of freedom of the small processes of X, Y, respectively. Define nonnegative sequences  ${c_i}_{i \text{ s}}$  and  ${d_i}_{i \text{ s}}$  that weakly approximate the probability measures  $\pi$  and

 $\rho$ , respectively. We assume that  $\sum_{i=1}^{n} c_i = \sum_{i=1}^{n} c_i$  $\sum_{i=1}^{I_r} c_i = \sum_{i=1}^{I_R} d_i = 1$  $\sum_{i=1}^{i}$   $\sum_{j=1}^{i}$   $\sum_{j=1}^{i}$  $c = \sum d$  $\sum_{i=1}^{n} c_i = \sum_{j=1}^{n} d_j = 1$ . We prepare  $I_r$  independent Poisson random measures  $N^{(i)}$  ( $1 \le i \le I_r$ ) with compensated measures  $c_i v(\text{d}z) \text{d}s$ . Also, we prepare  $I_R$  independent Brownian

motions  $B^{(j)}$   $(1 \le j \le I_R)$ . All  $N^{(i)}$  and  $B^{(j)}$  are assumed to be mutually independent. Additionally, define discrete recession rates  $\{r_i\}_{1\le i\le I_r}$  and discrete reversion rates  $\{R_i\}_{1\le j\le I_R}$ , both strictly increasing.

Under these conditions, the finite-dimensional supOU process  $\hat{Y} = (\hat{Y}_t)_{t \in \mathbb{R}}$  is defined as follows:

$$
\hat{Y}_t = \sum_{i=1}^{I_r} y_t^{(i)} \tag{48}
$$

with

$$
y_t^{(i)} = \int_{s=-\infty}^{s=t} \int_{z=0}^{z=\infty} z e^{-r_t(t-s)} N^{(i)} \left( ds, dz, dr \right). \tag{49}
$$

Similarly, the discretized auxiliary process  $\hat{X} = (\hat{X}_t)_{t \in \mathbb{R}}$  is defined as follows:

$$
\hat{X}_t = \sum_{j=1}^{I_R} x_t^{(j)} \tag{50}
$$

with

$$
x_t^{(j)} = \int_{s=-\infty}^{s=t} e^{-R_j(t-s)} \left\{ d_j R_j \mu \left( \hat{Y}_s - \overline{Y} \right) \mathrm{d}s + \sigma \sqrt{R_j d_j \hat{Y}_s} \mathrm{d}B_s^{(j)} \right\} . \tag{51}
$$

The corresponding SDEs for (49) and (51) are given by

$$
dy_t^{(i)} = -r_i y_t^{(i)} dt + \int_{z=0}^{z=\infty} z N^{(i)} (dt, dz)
$$
 (52)

and

$$
\mathrm{d}x_t^{(j)} = -R_j \left\{ x_t^{(j)} - d_j \mu \left( \hat{Y}_t - \overline{Y} \right) \right\} \mathrm{d}t + \sigma \sqrt{R_j d_j \hat{Y}_t} \mathrm{d}B_t^{(j)}\,,\tag{53}
$$

respectively.

According to Duffie et al. (2003)[45], SDEs (52) and (53) are affine. Therefore, it is possible to obtain the generalized Riccati equation associated with the moment-generating function  $\hat{M}(q) = \mathbb{E}\left[e^{q\hat{X}_t}\right]$ 

in a stationary state. The generator *G* associated with SDEs (52) and (53) is given by the linear partial integro-differential operator acting on a generic smooth function  $\Phi = \Phi(x_1, x_2, ..., x_{I_k}, y_1, y_2, ..., y_{I_r})$  with arguments in  $(x_1, x_2, ..., x_{I_k}, y_1, y_2, ..., y_{I_r}) \in \mathbb{R}^{I_k + I_r}$  $X_1, X_2, \dots, X_{I_n}, Y_1, Y_2, \dots, Y_{I_n}$   $\in \mathbb{R}^{I_R + I_r}$  as follows:

$$
G\Phi = \sum_{i=1}^{I_r} \left\{ -r_i y_i \frac{\partial \Phi}{\partial y_i} + c_i \int_0^{+\infty} \Delta \Phi(z_i) v(\mathrm{d}z_i) \right\} + \sum_{j=1}^{I_R} \left\{ -R_j \left( x_j - \mu d_j \left( \sum_{i=1}^{I_r} y_i - \overline{Y} \right) \right) \frac{\partial \Phi}{\partial x_j} + \frac{1}{2} \sigma^2 R_j d_j \left( \sum_{i=1}^{I_r} y_i \right) \frac{\partial^2 \Phi}{\partial x_j^2} \right\} \tag{54}
$$

Here, the first and second lines of (54) correspond to the generators of the discretized supOU and auxiliary processes, respectively. To estimate the functional form of the conditional moment-generating function

$$
\Phi(t, x_1, x_2, ..., x_{I_R}, y_1, y_2, ..., y_{I_r}) = \mathbb{E}\left[e^{q\hat{X}_t} \Big| \Big( x_1, x_2, ..., x_{I_R}, y_1, y_2, ..., y_{I_r} \Big) \right]
$$
  
=  $\exp\left(\hat{\phi}_t + \sum_{i=1}^{I_r} \psi_t^{(i)} y_i + \sum_{j=1}^{I_r} \omega_t^{(j)} x_j \right)$ , (55)

we substitute (55) into the Kolmogorov equation:

$$
\frac{\partial \Phi}{\partial t} = G\Phi, \ t > 0, \ (x_1, x_2, ..., x_{I_k}, y_1, y_2, ..., y_{I_r}) \in \mathbb{R}^{I_k + I_r}, \tag{56}
$$

yielding (noting that  $\Phi > 0$ ):

$$
\frac{d}{dt} \left( \hat{\phi}_t + \sum_{i=1}^{I_r} \psi_i^{(i)} y_i + \sum_{j=1}^{I_r} \omega_i^{(j)} x_j \right) \n= \sum_{i=1}^{I_r} \left\{ -r_i y_i \psi_i^{(i)} + c_i \int_0^{+\infty} \left( e^{\psi_i^{(i)} z_i} - 1 \right) v \left( dz_i \right) \right\} \n+ \sum_{j=1}^{I_R} \left\{ -R_j x_j \omega_i^{(j)} - \mu R_j d_j \overline{Y} \omega_i^{(j)} + \mu R_j d_j \left( \sum_{i=1}^{I_r} y_i \right) \omega_i^{(j)} + \frac{1}{2} \sigma^2 R_j d_j \left( \sum_{i=1}^{I_r} y_i \right) \left( \omega_i^{(j)} \right)^2 \right\}
$$
\n(57)

and obtain the finite-dimensional generalized Riccati equation:

$$
\frac{d\hat{\phi}_t}{dt} = \sum_{i=1}^{I_r} c_i \int_0^{+\infty} \left( e^{w_t^{(i)}z} - 1 \right) \nu \left( dz \right) - \mu \overline{Y} \sum_{j=1}^{I_r} d_j R_j \omega_t^{(j)} \text{ , } t > 0 \text{ ,}
$$
 (58)

$$
\frac{d\omega_t^{(j)}}{dt} = -R_j \omega_t^{(j)}, \ 1 \le j \le I_R, \ t > 0,
$$
\n(59)

and

$$
\frac{d\psi_t^{(i)}}{dt} = -r_i\psi_t^{(i)} + \mu \sum_{j=1}^{I_R} d_j R_j \omega_t^{(j)} + \frac{1}{2} \sigma^2 \sum_{j=1}^{I_R} d_j R_j \left(\omega_t^{(j)}\right)^2, \ 1 \le i \le I_r, \ t > 0 \tag{60}
$$

subject to initial conditions:

$$
\omega_0^{(\cdot)} = q \text{ and } \psi_0^{(\cdot)} = \phi_0 = 0. \tag{61}
$$

Finally, letting  $I_r, I_R \to +\infty$  and  $t \to +\infty$  formally recovers (31)-(33).

# **A. 2 Proofs**

# *Proof of Proposition 1*

For the average in (18), by (13) we formally have

$$
\mathbb{E}\left[X_{t}\right] = \mathbb{E}\left[\int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} \left\{ R\mu\left(Y_{s}-\overline{Y}\right) \rho\left(dR\right) ds + \sigma \sqrt{RY_{s}} B\left(dR, ds\right) \right\}\right]
$$
\n
$$
= \mathbb{E}\left[\int_{r=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} R\mu\left(Y_{s}-\overline{Y}\right) \rho\left(dR\right) ds\right]
$$
\n
$$
+ \mathbb{E}\left[\int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} \sigma \sqrt{RY_{s}} B\left(dR, ds\right)\right]
$$
\n
$$
= \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} R\mu \mathbb{E}\left[Y_{s}-\overline{Y}\right] \rho\left(dR\right) ds
$$
\n
$$
= \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-R(t-s)} R\mu\left(\overline{Y}-\overline{Y}\right) \rho\left(dR\right) ds
$$
\n
$$
= 0
$$
\n(62)

because *Y* is stationary and independent of *<sup>B</sup>* .

For variance from (19), because *B* is a space-time Gaussian random measure and (18), and *Y* is a nonnegative stationary process, we have

$$
\mathbb{V}[X_{t}] = \mathbb{E}[\sum_{R=0}^{R^{2}}\int_{s=-\infty}^{s=r} e^{-R(t-s)} \{R\mu(Y_{s}-\bar{Y})\rho(dR)ds + \sigma\sqrt{RY_{s}}B(dR, ds)\}]^{2}]
$$
\n
$$
= \mathbb{E}\left[\int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-R(t-s)} \{R\mu(Y_{s}-\bar{Y})\rho(dR)ds + \sigma\sqrt{RY_{s}}B(dR, ds)\} \times \int_{R=\infty}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-P(t-s)} \{R\mu(Y_{s}-\bar{Y})\rho(dR)ds + \sigma\sqrt{RY_{s}}B(dR, ds)\} \times \int_{R=\infty}^{R=\infty}\int_{s=0}^{s=r} \int_{u=-\infty}^{u=r} e^{-P(t-u)} \{P\mu(Y_{u}-\bar{Y})\rho(dP)du + \sigma\sqrt{PY_{u}}B(dR, ds)B(dP, du)\} \right]
$$
\n
$$
= \sigma^{2} \mathbb{E}[\int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} \int_{p=0}^{p=\infty}\int_{u=-\infty}^{u=r} e^{-R(t-s)-P(t-u)} (Y_{s}-\bar{Y})\rho(dR)\rho(dP)dsdu]
$$
\n
$$
= \sigma^{2} \int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-2R(t-s)} \mathbb{E}[Y_{s}]R\rho(R)dkds
$$
\n
$$
+ \mu^{2} \int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-2R(t-s)} \mathbb{E}[Y_{s}]R\rho(R)dkds
$$
\n
$$
= \sigma^{2} \int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-2R(t-s)} \mathbb{E}[Y_{s}]R\rho(dR)ds
$$
\n
$$
+ \mu^{2} \int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-2R(t-s)} \mathbb{E}[Y_{s}]R\rho(dR)ds
$$
\n
$$
+ \mu^{2} \int_{R=0}^{R=\infty}\int_{s=-\infty}^{s=r} e^{-2R(t-s)} \mathbb{E}[Y_{s}]R\rho(d
$$

We have

$$
\int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-2R(t-s)} R \rho(dR) ds = \int_{R=0}^{R=\infty} \frac{1}{2R} R \rho(dR) dR = \frac{1}{2}.
$$
 (64)

Due to

$$
AC_{Y}\left(|s-u|\right) = \left(\int_0^{+\infty} \frac{\pi(\mathrm{d}r)}{r}\right)^{-1} \int_{r=0}^{r=+\infty} \frac{1}{r} e^{-r|s-u|} \pi(\mathrm{d}r), \tag{65}
$$

we also have

$$
\int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} \int_{p=0}^{P=\infty} \int_{u=-\infty}^{u=t} RPe^{-R(r-s)-P(t-u)} \text{AC}_{Y}\left(|s-u|\right) \rho(dR) \rho(dP) \text{dsdu}
$$
\n
$$
= \left(\int_{0}^{+\infty} \frac{\pi(dr)}{r}\right)^{-1} \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} \int_{p=0}^{P=\infty} \int_{u=-\infty}^{u=t} \int_{r=0}^{r=\infty} \frac{RP}{r} e^{-R(t-s)-P(t-u)-r|s-u|} \pi(dr) \rho(dR) \rho(dP) \text{dsdu} \quad . \quad (66)
$$
\n
$$
= \left(\int_{0}^{+\infty} \frac{\pi(dr)}{r}\right)^{-1} \int_{R=0}^{R=\infty} \int_{p=0}^{P=\infty} \int_{r=0}^{r=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r} \left(\int_{s=-\infty}^{s=t} \int_{u=-\infty}^{u=t} e^{-R(t-s)-P(t-u)-r|s-u|} \text{dsdu}\right) \pi(dr) \rho(dR) \rho(dP)
$$

$$
= \left(\int_{0}^{+\infty} \frac{h(x)}{h} \right) \int_{R=0}^{R=\infty} \int_{P=0}^{P=\infty} \int_{r=0}^{P=\infty} \frac{h^{R}}{h} \left(\int_{0}^{+\infty} \int_{-\infty}^{H} e^{-R(t-x)-P(t-u)-r|s-u} dS du\right) \pi(dr) \rho(dR) \rho(dP)
$$
\nThe inner integral in the last line of (66) is rewritten as follows:

\n
$$
\int_{s=-\infty}^{s=t} \int_{u=-\infty}^{u=t} e^{-R(t-s)-P(t-u)-r|s-u|} dS du = \int_{s=-\infty}^{s=0} \int_{u=-\infty}^{u=0} e^{Rs+Pu-r|s-u|} dS du
$$
\n
$$
= \int_{s=0}^{s=\infty} \int_{u=0}^{u=s} e^{-Rs-Pu-r|s-u|} dS du
$$
\n
$$
= \int_{s=0}^{s=\infty} \int_{u=s}^{u=s} e^{-Rs-Pu-r|s-u|} dS du
$$
\n
$$
= \int_{s=0}^{s=\infty} \int_{u=s}^{u=s} e^{-Rs-Pu-r|s-u|} dS du
$$
\n
$$
= \int_{s=0}^{s=\infty} \int_{u=s}^{u=s} e^{-Rs-Pu-r|s-u|} dS du + \int_{u=0}^{u=s} \int_{s=u}^{s=\infty} e^{-Rs-Pu-r|s-u|} dS du
$$
\n
$$
= \int_{s=0}^{s=\infty} e^{-Rs+rs} \int_{u=s}^{u=s} e^{-(P+r)s} du + \int_{u=0}^{u=s} e^{-Pu+n} \int_{s=u}^{s=u} e^{-(R+r)u} dS du.
$$
\n
$$
= \int_{s=0}^{s=\infty} e^{-Rs+rs} \frac{e^{-(P+r)s}}{P+r} dx + \int_{u=0}^{u=s} e^{-Pu+n} \frac{e^{-(R+r)u}}{R+r} du
$$
\n
$$
= \frac{1}{P+R} \left(\frac{1}{P+r} + \frac{1}{R+r}\right)
$$

By combining  $(63)$ – $(67)$ , we obtain (19).

$$
\int_{1}^{1} \sum_{n=-n}^{n} \int_{p=0}^{n} \int_{1}^{1} \int_{B=0}^{n} \int_{1}^{1} e^{-x} P(e^{-x}) P(-x) P(-x)} \Delta C_{\gamma} ([s - u]) \rho (dR) \rho (dr) dx
$$
\n
$$
\int_{0}^{1} \sum_{n}^{n} \frac{\pi (dr)}{r} \int_{1}^{2} \int_{1}^{2} e^{-x} \int_{1}^{2} e^{-
$$

and

$$
\int_{s=-\infty}^{s=t} e^{-R(t-s)-r(t-s)} ds = \int_{s=-\infty}^{s=0} e^{(R+r)s} ds = \frac{1}{R+r}.
$$
 (69)

Substituting (69) into (67), we obtain the expression for covariance (20).

# *Proof of Proposition 2*

By the isometry and zero-mean Gaussian property of B and  $\mathbb{E}[X_t] = 0$ , we have

$$
\begin{split}\n&\mathbb{V}[X_{t}]A C_{x}(h) \\
&= \mathbb{E}\Big[X_{t}X_{t+h}\Big] \\
&= \mathbb{E}\Big[\int_{R=0}^{R=+\infty} \int_{S=-\infty}^{s=t} e^{-R(t-s)} \left\{ R\mu (Y_{s}-\overline{Y}) \rho (dR) ds + \sigma \sqrt{RY_{s}} B(dR, ds) \right\} \\
&= \int_{X} \int_{P=0}^{R=+\infty} \int_{u=-\infty}^{s=t} e^{-P(t+h-s)} \left\{ P\mu (Y_{u}-\overline{Y}) \rho (dP) du + \sigma \sqrt{PY_{u}} B(dP, du) \right\} \Big] \\
&= \sigma^{2} \mathbb{E}\Big[\int_{R=0}^{R=+\infty} \int_{s=-\infty}^{s=t} \int_{P=0}^{P=\infty} \int_{u=-\infty}^{u=t+h} \sqrt{RP} e^{-R(t-s)-P(t+h-u)} \sqrt{Y_{s}Y_{u}} B(dR, ds) B(dP, du)\Big] \\
&+ \mu^{2} \mathbb{E}\Big[\int_{R=0}^{R=+\infty} \int_{s=-\infty}^{s=t} \int_{P=0}^{P=\infty} \int_{u=-\infty}^{u=t+h} R P e^{-R(t-s)-P(t+h-u)} (Y_{s}-\overline{Y}) (\gamma_{u}-\overline{Y}) \rho (dR) \rho (dP) ds du\Big] \\
&= \sigma^{2} \overline{Y} \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} e^{-2R(t-s)-Rh} R \rho (R) dR ds \\
&+ \mu^{2} \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} \int_{P=0}^{P=\infty} \int_{u=-\infty}^{u=t+h} R P e^{-R(t-s)-P(t+h-u)} \overline{V} A C_{Y} (|s-u|) \rho (dR) \rho (dP) ds du \\
&= \sigma^{2} \overline{Y} \int_{R=0}^{R=\infty} e^{-Rh} \left( \int_{s=-\infty}^{s=t} e^{-2R(t-s)} ds \right) R \rho (R) dR \\
&+ \mu^{2} \overline{V} \int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=t} \int_{P=0}^{P=\infty} \int_{u=-\infty}^{u=t+h} R P e^{-R(t-s)-P(t
$$

where we used (64).

For the second integral in (70), we proceed as follows:

$$
\int_{R=0}^{R=+\infty} \int_{s=-\infty}^{s=t} \int_{p=0}^{p=\infty} \int_{u=-\infty}^{u=t+h} RPe^{-R(t-s)-P(t+h-u)} \text{AC}_{Y}\left(|s-u|\right) \rho(dR) \rho(dP) \text{dsdu}
$$
\n
$$
= \left(\int_{0}^{+\infty} \frac{\pi(dr)}{r}\right)^{-1} \int_{R=0}^{R=+\infty} \int_{s=-\infty}^{s=t} \int_{p=0}^{p=\infty} \int_{u=-\infty}^{u=t+h} \int_{r=0}^{r=\infty} \frac{RP}{r} e^{-R(t-s)-P(t+h-u)-r|s-u|} \pi(dr) \rho(dR) \rho(dP) \text{dsdu}
$$
\n
$$
= \left(\int_{0}^{+\infty} \frac{\pi(dr)}{r}\right)^{-1} \int_{R=0}^{R=+\infty} \int_{p=0}^{p=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r} e^{-Ph} \left(\int_{s=-\infty}^{s=t} \int_{u=-\infty}^{u=t+h} e^{-R(t-s)-P(t-u)-r|s-u|} \text{dsdu}\right) \pi(dr) \rho(dR) \rho(dP) \tag{71}
$$

and

$$
\int_{s=-\infty}^{s=t} \int_{u=-\infty}^{u=t+h} e^{-R(t-s)-P(t-u)-r|s-u|} ds du = \int_{s=-\infty}^{s=0} \int_{u=-\infty}^{u=h} e^{-Rs+Pu-r|s-u|} ds du
$$
  
\n
$$
= \int_{s=0}^{s=\infty} \int_{u=-h}^{u=\infty} e^{-Rs-Pu-r|s-u|} ds du + \int_{s=0}^{s=\infty} \int_{u=-h}^{u=0} e^{-Rs-Pu-r|s-u|} ds du
$$
(72)

The first term on the last line of (72) is identical to (67). For the second term, we focus on the case where  $P \neq r$  in the integrand since the Lebesgue measure of the set  $\{P = r\}$  is zero in  $(0, +\infty)^2$  as assumed in **Proposition 2**:

$$
\int_{s=0}^{s=\infty} \int_{u=-h}^{u=0} e^{-Rs - Pu - r|s-u|} ds du = \int_{s=0}^{s=\infty} \int_{u=-h}^{u=0} e^{-Rs - Pu - r(s-u)} ds du
$$

$$
= \int_{s=0}^{s=\infty} e^{-(R+r)s} ds \int_{u=-h}^{u=0} e^{-(P-r)u} du
$$

$$
= \frac{1}{R+r} \frac{1}{P-r} \left( e^{(P-r)h} - 1 \right)
$$
(73)

Substituting  $(72)$  and  $(73)$  into  $(71)$ , we have

$$
\int_{R=0}^{R=\infty} \int_{s=-\infty}^{s=r} \int_{p=0}^{P=\infty} \int_{u=-\infty}^{u=r+h} RPe^{-R(r-s)-P(r+h-u)} A C_Y(|s-u|) \rho(dR) \rho(dP) ds du \n= \left( \int_0^{+\infty} \frac{\pi(dr)}{r} \right)^{-1} \int_{R=0}^{R=\infty} \int_{p=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r} \frac{1}{P+R} \left( \frac{1}{P+r} + \frac{1}{R+r} \right) e^{-Ph} \pi(dr) \rho(dR) \rho(dP) . \tag{74}
$$
\n
$$
+ \left( \int_0^{+\infty} \frac{\pi(dr)}{r} \right)^{-1} \int_{R=0}^{R=\infty} \int_{p=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r(R+r)(P-r)} \left( e^{-rh} - e^{-Ph} \right) \pi(dr) \rho(dR) \rho(dP)
$$

Finally, substituting (74) into (70) yields (21). Then, combining Equations (22), (23), and (24) directly follow from (21).

 $\Box$ 

# *Proof of Proposition 3*

Under the assumption of **Proposition 3**, we have

$$
I_1(h) = \frac{1}{(1 + \beta_R h)^{\alpha_R}} = O\left(h^{-\alpha_R}\right),\tag{75}
$$

$$
\int_{x=0}^{x=0} \int_{x=-k}^{x=0} e^{-Rx-p_{0}-r_{1}-r_{1}-r_{1}} dSd\mu = \int_{x=0}^{x=0} \int_{x=-k}^{x=0} e^{-(Rx+r_{1})} dSd\mu
$$
\n
$$
= \int_{x=0}^{x=0} e^{-(Rx+r_{1})} dS \int_{x=-k}^{x=0} e^{-(x-x+p_{1})} d\mu.
$$
\n2) and (73) into (71), we have  
\n
$$
\int_{x=-k}^{x=0} \int_{x=-0}^{x=0} e^{-(x-x+p_{1}+k-x)} AC_{x} (|s-u|) \rho (dR) \rho (dP) dSd\mu
$$
\n
$$
\int_{x=-k}^{x=0} \int_{x=0}^{x=0} e^{-(x-x+p_{1}+k-x)} AC_{x} (|s-u|) \rho (dR) \rho (dP) dSd\mu
$$
\n
$$
\int_{x=-k}^{x=0} \int_{x=0}^{x=0} \int_{y=0}^{x=0} \int_{y=0}^{x=0} \frac{RP}{r P + R} \left( \frac{1}{P+r} + \frac{1}{R+r} \right) e^{-px} \pi (dr) \rho (dR) \rho (dP).
$$
\n(74)  
\n
$$
\int_{x=0}^{x=0} \int_{x=0}^{x=0} \int_{y=0}^{x=0} \int_{y=0}^{x=0} \frac{RP}{r (R+r) (P-r)} (e^{-rx} - e^{-rx}) \pi (dr) \rho (dR) \rho (dP).
$$
\n1).  
\n3. If  $\int_{0}^{x=0} (1) \pi$  (7) yields (21). Then, combining Equations (22), (23), and (24) directly  
\n
$$
\int_{1}^{x=0} (h^{-2}e)^{x} \int_{y=0}^{x=0} \int_{y=0}^{x=0} \frac{RP}{r (P + R)} \left( \frac{1}{P+r} + \frac{1}{R+r} \right) e^{-px} \pi (dr) \rho (dR) \rho (dP)
$$
\n
$$
\leq \int_{R=0}^{x=0} \int_{y=0}^{x=0} \int_{y=0}^{x=0} \int_{y=0}^{x=0} \frac{RP}{r (P + R)}
$$

and

$$
I_{3}(h) = \int_{R=0}^{R=\infty} \int_{P=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{RP}{r(R+r)(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(dr) \rho(dr) \rho(dr)
$$
  
\n
$$
\leq \int_{R=0}^{R=\infty} \int_{P=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{P}{r(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(dr) \rho(dr) \rho(dr)
$$
  
\n
$$
= \int_{P=0}^{P=\infty} \int_{r=0}^{r=\infty} \frac{P}{r(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(dr) \rho(dr)
$$
 (77)

We note the following elementary inequality, which will be used in the proof: for any  $P > r > 0$ ,

$$
\frac{\partial}{\partial P} \left( \frac{e^{-rh} - e^{-Ph}}{P - r} \right) = \frac{(P - r)h + 1 - e^{(P - r)h}}{(P - r)^2} e^{-Ph} \le 0 \tag{78}
$$

due to  $e^x \geq 1 + x$  ( $x \in \mathbb{R}$ ), and the limit

$$
\lim_{P \to r} \frac{e^{-rh} - e^{-Ph}}{P - r} = h e^{-rh}, \ r > 0.
$$
\n(79)

The most technical aspect of this proof lies in evaluating the last integral in (77). We divide the domain of integration  $(0, +\infty) \times (0, +\infty)$  as follows: for a small  $\varepsilon > 0$  specified later,  $(0, +\infty)^2 = D_0 \cup D_1 \cup D_2$  with  $D_0 = \{(p, r) \in (0, +\infty)^2 : |p - r| \le \varepsilon\}$ ,  $D_1 = \{(p, r) \in (0, +\infty)^2 : p > r + \varepsilon\}$ , and  $D_2 = \{(p, r) \in (0, +\infty)^2 : r > p + \varepsilon\}$ . Then, from (78), for large  $h > 0$ , we have

$$
\int_{D_1} \frac{P}{r(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P) \le \int_{D_1} \frac{P}{r\epsilon} \left(e^{-rh} - e^{-(r+\epsilon)h}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
\n
$$
\le \frac{1 - e^{-\epsilon h}}{\epsilon} \int_{D_1} \frac{P}{r} e^{-rh} \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
\n
$$
\le \frac{1 - e^{-\epsilon h}}{\epsilon} \int_{P=0}^{P=\epsilon p} P \rho(\mathrm{d}P) \int_{r=0}^{r=\epsilon p} \frac{1}{r} e^{-rh} \pi(\mathrm{d}r)
$$
\n
$$
= \frac{1 - e^{-\epsilon h}}{\epsilon} O\left(h^{1-\alpha_r}\right)
$$
\n(80)

and

$$
\int_{D_2} \frac{P}{r(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P) \leq \int_{D_2} \frac{P}{r\epsilon} \left(e^{-Ph} - e^{-(P+\epsilon)h}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
  

$$
\leq \frac{1 - e^{-\epsilon h}}{\epsilon} \int_{D_2} \frac{P}{r} e^{-Ph} \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
  

$$
\leq \frac{1 - e^{-\epsilon h}}{\epsilon} \int_{P=0}^{P+\infty} P e^{-Ph} \rho(\mathrm{d}P) \int_{r=0}^{r=\infty} \frac{1}{r} \pi(\mathrm{d}r)
$$
  

$$
= \frac{1 - e^{-\epsilon h}}{\epsilon} O\left(h^{-1-\alpha_R}\right)
$$
 (81)

From (78)-(79), we also have

$$
\int_{D_3} \frac{P}{r(P-r)} \left(e^{-rh} - e^{-Ph}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P) \le \int_{D_3} \frac{P}{r} h \max\left\{e^{-rh}, e^{-Ph}\right\} \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
\n
$$
\le h \int_{D_3} \frac{P}{r} \left(e^{-rh} + e^{-Ph}\right) \pi(\mathrm{d}r) \rho(\mathrm{d}P) \qquad . \tag{82}
$$
\n
$$
\le h \int_{D_3} \frac{P}{r} e^{-rh} \pi(\mathrm{d}r) \rho(\mathrm{d}P) + h \int_{D_3} \frac{P}{r} e^{-Ph} \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$

Next, we estimate the last two integrals in (82):

$$
\int_{D_3} \frac{P}{r} e^{-rh} \pi(\mathrm{d}r) \rho(\mathrm{d}P) = \int_{r=0}^{r=\infty} \int_{P=\max\{0,r-\varepsilon\}}^{P=r+\varepsilon} \frac{P}{r} e^{-rh} \pi(\mathrm{d}r) \rho(\mathrm{d}P)
$$
  
\n
$$
= \int_{r=0}^{r=\infty} \frac{1}{r} e^{-rh} \left( \int_{P=\max\{0,r-\varepsilon\}}^{P=r+\varepsilon} P \rho(\mathrm{d}P) \right) \pi(\mathrm{d}r)
$$
  
\n
$$
= O\left(h^{1-\alpha_r}\right) \times O\left(\varepsilon^{\alpha_{R}+1}\right)
$$
  
\n
$$
= O\left(h^{1-\alpha_r}\varepsilon^{\alpha_{R}+1}\right)
$$
  
\n(83)

and

$$
\int_{D_{2}} \frac{P}{r} e^{-p_{k}} \pi(\mathbf{d}r) \rho(\mathbf{d}P) = \int_{r=0}^{r=r_{0}} \int_{r=\text{max}(0, r-s)}^{r=r_{0}+r} \frac{P}{r} e^{-p_{k}} \pi(\mathbf{d}r) \rho(\mathbf{d}P)
$$
\n
$$
= \int_{r=0}^{r=r_{0}} P e^{-p_{k}} \int_{r=\text{max}(0, r-s)}^{r=r_{0}+r} \frac{1}{r} \pi(\mathbf{d}r) \rho(\mathbf{d}P)
$$
\n
$$
= O\left(h^{-1-\alpha_{k}} \times O\left(e^{\alpha_{k}-1}\right)\right)
$$
\nobtain\n
$$
= O\left(h^{-1-\alpha_{k}} e^{\alpha_{k}-1}\right)
$$
\nobtain\n
$$
\frac{P}{r(P-r)} \left(e^{-r h} - e^{-r h}\right) \pi(\mathbf{d}r) \rho(\mathbf{d}P) \leq O\left(h^{-2\alpha_{k}} e^{\alpha_{k}+1}\right) + O\left(h^{-\alpha_{k}} e^{\alpha_{k}-1}\right).
$$
\n(85)\n(77), (80), (81), and (85), we arrive at the following estimate for large  $h > 0$ :\n
$$
\int \frac{1-e^{-\alpha_{k}}}{\varepsilon} O\left(h^{-\alpha_{k}}\right) + \frac{1-e^{-\alpha_{k}}}{\varepsilon} O\left(h^{-\alpha_{k}}\right) + O\left(h^{-2\alpha_{k}} e^{\alpha_{k}+1}\right) + O\left(h^{-2\alpha_{k}} e^{\alpha_{k}-1}\right).
$$
\n(86)\nchoose  $\varepsilon$  depending on  $h$  as follows:  $\varepsilon = h^{-\lambda}$  with some  $\lambda \in (0,1)$ . For large  $h > 0$ ,\n
$$
\frac{1-e^{-\alpha_{k}}}{\varepsilon} = h^{\lambda}\left(1-\exp\left(-h^{1-\lambda}\right)\right) = O\left(h^{\lambda})
$$
\n
$$
\varepsilon(0,1),
$$
\n
$$
I_{3}(h) \leq O\left(h^{\lambda-1-\alpha_{k}}\right) + O\left(h^{\lambda-1-\alpha_{k}}\right) + O\left(h^{\lambda-\alpha_{k}-\lambda(\alpha_{k}+1)}\right) + O\left(h^{-\
$$

From  $(82)$ - $(84)$ , we obtain

$$
\int_{D_3} \frac{P}{r(P-r)} \Big(e^{-rh} - e^{-Ph}\Big) \pi\Big(\mathrm{d}r\Big) \rho\Big(\mathrm{d}P\Big) \le O\Big(h^{2-\alpha_r} \varepsilon^{\alpha_r+1}\Big) + O\Big(h^{-\alpha_R} \varepsilon^{\alpha_r-1}\Big). \tag{85}
$$

Consequently, from (77), (80), (81), and (85), we arrive at the following estimate for large  $h > 0$ :

$$
I_{3}(h) \leq \frac{1-e^{-\varepsilon h}}{\varepsilon}O\left(h^{-1-\alpha_{k}}\right)+\frac{1-e^{-\varepsilon h}}{\varepsilon}O\left(h^{1-\alpha_{r}}\right)+O\left(h^{-\alpha_{r}}\varepsilon^{\alpha_{k}+1}\right)+O\left(h^{-\alpha_{k}}\varepsilon^{\alpha_{r}-1}\right).
$$
(86)

Now, we choose  $\varepsilon$  depending on h as follows:  $\varepsilon = h^{-\lambda}$  with some  $\lambda \in (0,1)$ . For large  $h > 0$ ,

we have

$$
\frac{1 - e^{-\varepsilon h}}{\varepsilon} = h^{\lambda} \left( 1 - \exp\left( - h^{1 - \lambda} \right) \right) = O\left( h^{\lambda} \right),\tag{87}
$$

and hence, since  $\lambda \in (0,1)$ ,

$$
I_{3}(h) \leq O\left(h^{\lambda-1-\alpha_{R}}\right)+O\left(h^{\lambda+1-\alpha_{r}}\right)+O\left(h^{2-\alpha_{r}-\lambda(\alpha_{R}+1)}\right)+O\left(h^{-\alpha_{R}-\lambda(\alpha_{r}-1)}\right)
$$
  
\n
$$
\leq O\left(h^{-\alpha_{R}}\right)+O\left(h^{\lambda+1-\alpha_{r}}\right)+O\left(h^{2-\alpha_{r}-\lambda(\alpha_{R}+1)}\right)+O\left(h^{-\alpha_{R}}\right)
$$
  
\n
$$
=O\left(h^{-\alpha_{R}}\right)+O\left(h^{\lambda+1-\alpha_{r}}\right)+O\left(h^{2-\alpha_{r}-\lambda(\alpha_{R}+1)}\right)
$$
\n
$$
(88)
$$

Due to  $\alpha_r > 1$ , there exists some  $\lambda = \lambda^* \in (0,1)$  such as  $\lambda^* + 1 - \alpha_r < 0$ . We consider two separate cases:  $\alpha_r > 2$  and  $\alpha_r \in (1,2]$ . First, if  $\alpha_r > 2$ , we have, automatically,

$$
2 - \alpha_r - \lambda^* (\alpha_R + 1) < 0 \,. \tag{89}
$$

By combining (75), (76), and (88), it follows that (without assuming (26))  $\lim_{h\to\infty} AC_x(h) = 0$  as  $h \to +\infty$ . Second, if  $\alpha_r \in (1,2]$ ,  $\lambda^*$  such that  $\lambda^* + 1 - \alpha_r < 0$  with (89) exists if

$$
\frac{2-\alpha_r}{\alpha_r+1} < \alpha_r - 1 \Leftrightarrow 2-\alpha_r < (\alpha_r - 1)(\alpha_r + 1) \Leftrightarrow 1 < (\alpha_r - 1)(\alpha_r + 2),\tag{90}
$$

which is equivalent to the inequality in (26). From (26), we conclude that  $\lim_{h\to\infty} AC_x(h) = 0$  as  $h \to +\infty$ because we can choose  $\lambda = \lambda^*$  to obtain

$$
I_{3}(h) \leq O\left(h^{-\alpha_{R}}\right) + O\left(h^{\lambda^{*}+1-\alpha_{r}}\right) + O\left(h^{2-\alpha_{r}-\lambda^{*}(\alpha_{R}+1)}\right). \tag{91}
$$

Finally, we close the proof by estimating  $AC_x(h)$ . First, if  $\alpha_r > 2$ , we have, by taking  $\lambda \to +0$ ,

$$
I_3(h) \le O\left(h^{-\alpha_k}\right) + O\left(h^{-(\alpha_r - 2)}\right) \tag{92}
$$

leading directly to (27). Second, if  $\alpha_r \in (1,2]$ , by equating the powers of h in the second and third terms in (88), we have

$$
I_3(h) \le O\left(h^{-\alpha_R}\right) + O\left(h^{-\eta}\right) \tag{93}
$$

with

$$
\eta = \frac{(\alpha_r - 1)(\alpha_R + 2) - 1}{\alpha_R + 2} > 0
$$
\n(94)

due to Equation (26). This leads directly to the estimate in (29).

□

## *Proof of Proposition 4*

We analyzed the upper bound of  $\psi_t$ . For this purpose, for each  $r, P > 0$  with  $P \neq r$ , consider the following function of  $t \geq 0$ :

$$
f_t = P \int_0^t e^{-r(t-s)} e^{-Ps} ds = \frac{P}{r - P} \left( e^{-Pt} - e^{-rt} \right).
$$
 (95)

We have  $f_0 = 0$ ,

$$
\frac{df_{t}}{dt} = \frac{P}{r - P} \left( -P e^{-Pt} + r e^{-rt} \right), \text{ and } \frac{d^{2} f_{t}}{dt^{2}} = \frac{P}{r - P} \left( P^{2} e^{-Pt} - r^{2} e^{-rt} \right). \tag{96}
$$

The function f is maximized at  $t = T$  such that the right-hand side of (96) vanishes. This T is obtained as  $T = \frac{1}{P-r} \ln \left( \frac{P}{r} \right)$  $=\frac{1}{P-r}\ln\left(\frac{P}{r}\right)$ . We then have the maximum value of the function f as follows:  $1-\frac{P}{\cdots}$  $\left(1-\left(\frac{P}{P}\right)^{\frac{P}{P-r}-\frac{r}{P-r}}\right)=\left(\frac{P}{P}\right)^{1-\frac{P/r}{P/r-1}}$  $\frac{p}{p}$  *(n)* $\frac{r}{p}$  *n (n)* $\frac{p}{p}$  *(n)* $\frac{p}{p}$  *r r n*) *r p*)  $f_r = \frac{P}{r - p} \left[ \left( \frac{P}{r}\right)^{-\frac{P}{p-r}} - \left( \frac{P}{r}\right)^{-\frac{r}{p-r}} \right] = \frac{P}{r - p} \left( \frac{P}{r}\right)^{-\frac{P}{p-r}} \left( 1 - \left( \frac{P}{r}\right)^{\frac{P}{p-r} - \frac{r}{p-r}} \right) = \left( \frac{P}{r}\right)^{1-\frac{P}{p}}$  $\frac{P}{r-P} \left( \left( \frac{P}{r} \right)^{-\frac{P}{P-r}} - \left( \frac{P}{r} \right)^{-\frac{r}{P-r}} \right) = \frac{P}{r-P} \left( \frac{P}{r} \right)^{\frac{P}{P-r}} \left( 1 - \left( \frac{P}{r} \right)^{\frac{P}{P-r} - \frac{r}{P-r}} \right) = \left( \frac{P}{r} \right)^{1-\frac{P/r}{P/r-1}}.$  $\frac{1}{r}$ . We then have the maximum value of the function  $f$  as follows:<br>=  $\frac{P}{r-P}\left(\left(\frac{P}{r}\right)^{\frac{p}{p-r}} - \left(\frac{P}{r}\right)^{\frac{r}{p-r}}\right) = \frac{P}{r-P}\left(\frac{P}{r}\right)^{\frac{p}{p-r}}\left(1 - \left(\frac{P}{r}\right)^{\frac{p}{p-r}-\frac{r}{p-r}}\right) = \left(\frac{P}{r}\right)^{1-\frac{p}{p/r-1}}$ .  $\frac{P}{-P}\left(\left(\frac{P}{r}\right)^{\frac{P}{P-r}} - \left(\frac{P}{r}\right)^{\frac{r}{P-r}}\right) = \frac{P}{r-P}\left(\frac{P}{r}\right)^{\frac{P}{P-r}}\left(1 - \left(\frac{P}{r}\right)^{\frac{P}{P-r}-\frac{r}{P-r}}\right) = \left(\frac{P}{r}\right)^{1-\frac{P/r}{P/r-1}}.$ . (97)

The right-hand side of (97) is expressed as  $g(u)$  with  $u = P/r > 0$ , where  $g(u)$  $g(u) = u^{\frac{1 - u}{u - 1}} = u^{\frac{-1}{u - 1}}$ . We have

$$
g(1) = e^{-1}
$$
 by  $g(u) = u^{\frac{1-u}{u-1}} = 1/(u-1+1)^{\frac{1}{u-1}}$  and taking the limit  $u \to 1$ . We have, for  $u \ne 1$ ,

$$
\frac{d}{du}\ln g\left(u\right) = \frac{d}{du}\left(\frac{1}{1-u}\ln u\right) = \frac{1}{\left(1-u\right)^2}\ln u - \frac{1}{u\left(1-u\right)} = \frac{1}{\left(1-u\right)^2}\left(\ln u + 1 - \frac{1}{u}\right). \tag{98}
$$

We also have

$$
\frac{d}{du}\left(\ln u + 1 - \frac{1}{u}\right) = \frac{1}{u} + \frac{1}{u^2} > 0 \text{ for } u > 0.
$$
 (99)

Moreover, it follows that

$$
\lim_{u \to +0} \left( \ln u + 1 - \frac{1}{u} \right) = -\infty, \quad \lim_{u \to +\infty} \left( \ln u + 1 - \frac{1}{u} \right) = +\infty, \text{ and } \ln 1 + 1 - \frac{1}{1} = 0. \tag{100}
$$

This implies that  $g(u)$  is maximized at  $u = 1$  with the maximum value  $g(1) = e^{-1}$ .

The function f with  $P = r$  is given by  $f_t = Pte^{-Pt}$ , which is maximized at  $t = P^{-1}$  as  $e^{-t}$ . Consequently, we have  $0 \le f_t \le e^{-1}$  for any  $P, r > 0$ . We apply this to the right-hand side of (35): for all  $t \geq 0$  and  $r > 0$ ,

$$
\psi_{t}(r) = \int_{P=0}^{P=\infty} \int_{s=0}^{s=t} P e^{-r(t-s)} \left( \mu q e^{-Pt} + \frac{\sigma^{2} q^{2}}{2} e^{-2Pt} \right) ds \rho(dP)
$$
\n
$$
= \mu q \int_{P=0}^{P=\infty} \int_{s=0}^{s=t} \left( P e^{-r(t-s)} e^{-Pt} \right) ds \rho(dP) + \frac{\sigma^{2} q^{2}}{4} \int_{P=0}^{P=\infty} \int_{s=0}^{s=t} \left( 2P e^{-r(t-s)} e^{-2Pt} \right) ds \rho(dP)
$$
\n
$$
\leq \max \{ \mu, 0 \} q \int_{P=0}^{P=\infty} \int_{s=0}^{s=t} \left( P e^{-r(t-s)} e^{-Pt} \right) ds \rho(dP) + \frac{\sigma^{2} q^{2}}{4} \int_{P=0}^{P=\infty} \int_{s=0}^{s=t} \left( 2P e^{-r(t-s)} e^{-2Pt} \right) ds \rho(dP)
$$
\n
$$
\leq \left( \max \{ \mu, 0 \} q + \frac{\sigma^{2} q^{2}}{4} \right) e^{-1}
$$
\n(101)

Finally, from (101), we arrive at the strict upper bound  $\psi_t(r) \le a_2$  for all  $t \ge 0$  and  $R > 0$ , provided condition in (36) is satisfied.



# A. **3** Identified values of  $\alpha_r$  ,  $\beta_r$  ,  $\alpha_R$  ,  $\beta_R$  for different time lag durations

For different time lag durations of autocorrelations, the identified values of the parameters  $\alpha_r, \beta_r$  in the probability measure  $\pi$  for the process Y and  $\alpha_r$ ,  $\beta_r$  in the probability measure  $\rho$  for the process X are summarized in **Table A1**. The results demonstrate that the exponential decay of the autocorrelation in SS, the truly long-memory nature in TN and TP, and the moderately long-memory nature in TOC and DSi remain consistent, irrespective of the lag duration.

Table A1. Parameter values for process X of each WQI. Numbers in "()" represent lag durations used for parameter identification.

	TN	TР	TOC.	<b>DSi</b>
$\alpha_{R}$ (-) (365 day)	3.410.E-01	$6.563.E-01$	$2.001.E+00$	$2.351.E+00$
$\beta_{R}$ (1/day) (365 day)	4.970.E-01	3.376.E-01	3.105.E-02	3.056.E-02
$\alpha_{R}$ (-) (730 day)	4.217.E-01	5.477.E-01	$2.334.E+00$	$3.510.E+00$
$\beta_R$ (1/day) (730day)	2.699.E-01	5.253.E-01	$2.506.E-02$	2.806.E-02
$\alpha_{R}$ (-) (1095 day)	$4.045.E-01$	5.673.E-01	$2.310.E+00$	$3.554.E+00$
$\beta_{R}$ (1/day) (1095 day)	$3.062.E-01$	4.766.E-01	$2.543.E-02$	2.741.E-02

## **A. 4 Sensitivity against**

**Table A2** shows the identified parameter values and theoretical statistics of the proposed model for river discharge under different  $\varepsilon$  values. This table serves as a supplementary version of **Table 2** in the main text. The results indicate that the average, variance, and skewness of the proposed model are not significantly affected by changes in  $\varepsilon$ , while kurtosis shows notable sensitivity. It is important to note that kurtosis is not included in the objective function in (41), and the autocorrelation of *X* depends on neither skewness nor kurtosis, as established in **Proposition 2**. Moreover, the measure  $\pi$ , which governs the autocorrelation, remains independent of  $\varepsilon$ . Consequently, the regularization introduced by  $\varepsilon$  does not qualitatively affect the computational results presented in the main text.

**Table A2.** Identified parameter values and theoretical statistics of model for river discharge against different values of  $\varepsilon$ .

	$\varepsilon = 0$	$\varepsilon = 0.01$	$\varepsilon = 0.1$	$\varepsilon = 0.2$	$\varepsilon = 0.4$
$c_1 \left( \text{ m}^{\frac{3c_3+\frac{3\varepsilon}{1+\varepsilon}}/s^{c_3+\frac{\varepsilon}{1+\varepsilon}}/ \text{day} \right)$	$1.266.E+00$	$1.251.E+00$	$1.124.E+00$	$1.009.E+00$	8.333.E-01
$c_{2}$ (s/m <sup>3</sup> )	1.960.E-03	1.811.E-03	8.920.E-04	$4.101.E-04$	8.871.E-05
$c_{3}(-)$	8.084.E-01	8.023.E-01	7.500.E-01	6.980.E-01	$6.109.E-01$
Average $(m^3/s)$	$1.701.E+01$	$1.701.E+01$	$1.701.E+01$	$1.700.E+01$	$1.700.E+01$
Variance $(m^6/s^2)$	$8.308.E+02$	$8.308.E+02$	$8.308.E+02$	$8.308.E+02$	$8.308.E+02$
Skewness (-)	$1.406.E+01$	$1.406.E+01$	$1.406.E+01$	$1.406.E+01$	$1.406.E+01$
Kurtosis (-)	$4.089.E+02$	$4.074.E+02$	$3.948.E+02$	$3.831.E+02$	$3.652.E+02$

#### **References**

- [1] Durighetto, N., Noto, S., Tauro, F., Grimaldi, S., & Botter, G. (2023). Integrating spatially-and temporallyheterogeneous data on river network dynamics using graph theory. Iscience, 26(8). <https://doi.org/10.1016/j.isci.2023.107417>
- [2] Zanetti, F., Botter, G., & Camporese, M. (2024). Stream network dynamics of non‐perennial rivers: Insights from integrated surface‐subsurface hydrological modeling of two virtual catchments. Water Resources Research, 60(2), e2023WR035631.<https://doi.org/10.1029/2023WR035631>
- [3] Ding, Z., Angarita, H., Albert Montesinos Cáceres, C., Lavado-Casimiro, W., Goldstein, J. A., Batista, N., ... & Schmitt, R. J. (2024). Sustainable land and irrigation management to limit loss of hydropower in the Andes-Amazon headwaters. Communications Earth & Environment, 5(1), 648. [https://doi.org/10.1038/s43247-024-](https://doi.org/10.1038/s43247-024-01738-4) [01738-4](https://doi.org/10.1038/s43247-024-01738-4)
- [4] Bahramloo, R., Wang, J., Sepehri, M., Faghfouri, A., Ghermezcheshmeh, B., Atapourfard, A., ... & Bazrafshan, E. (2024). Novel MCDA methods for flood hazard mapping: a case study in Hamadan, Iran. Stochastic Environmental Research and Risk Assessment, published online.<https://doi.org/10.1007/s00477-024-02836-7>
- [5] Rai, P., Kumar, J. Y., & Sen, A. (2024). Ganga, GAP, and lockdown: potential threats to the biodiversity of the river. Journal of Water and Climate Change, jwc2024322.<https://doi.org/10.2166/wcc.2024.322>
- [6] Hansen, H. H., Comoglio, C., Elings, J., Ericsson, P., Goethals, P., Gosselin, M. P., ... & Bergman, E. (2024). Fish habitat models for a future of novel riverscapes. BioScience, 74(9), 624-639. <https://doi.org/10.1093/biosci/biae081>
- [7] Gao, C., Liang, Z., Xin, P., Wang, C., Zhang, Y., & Chen, X. (2024). Identification of key water environmental factor contributions and spatiotemporal differential characteristics for eutrophication in Dianchi Lake. Environmental Monitoring and Assessment, 196(12), 1-18.<https://doi.org/10.1007/s10661-024-13425-0>
- [8] Doukhan, P. (2018). Stochastic models for time series. Springer, New York.
- [9] Volpi, E., Grimaldi, S., Aghakouchak, A., Castellarin, A., Chebana, F., Papalexiou, S. M., ... & Sharma, A. (2024). The legacy of STAHY: Milestones, achievements, challenges, and open problems in statistical hydrology. Hydrological Sciences Journal, 69(14), 1913-1949[. https://doi.org/10.1080/02626667.2024.2385686](https://doi.org/10.1080/02626667.2024.2385686)
- [10] Øksendal, B., & Sulem, A. (2019). Applied Stochastic Control of Jump Diffusions. Springer, Cham.
- [11] Pascucci, A. (2024). Probability Theory II: Stochastic Calculus. Springer, Cham.
- [12] Mandal, S., Sk, N., Tiwari, P. K., & Chattopadhyay, J. (2024). Bistability in modified Holling II response model with harvesting and Allee effect: Exploring transitions in a noisy environment. Chaos, Solitons & Fractals, 178, 114365[. https://doi.org/10.1016/j.chaos.2023.114365](https://doi.org/10.1016/j.chaos.2023.114365)
- [13] Paul, B., Sikdar, G. C., & Ghosh, U. (2025). Effect of fear and non-linear predator harvesting on a predator–prey system in presence of environmental variability. Mathematics and Computers in Simulation, 227, 442-460. <https://doi.org/10.1016/j.matcom.2024.08.021>
- [14] Mohammad, K. M., Tisha, M. S., & Kamrujjaman, M. (2024). Wiener and Lévy processes to prevent disease outbreaks: Predictable vs stochastic analysis. Partial Differential Equations in Applied Mathematics, 10, 100712. <https://doi.org/10.1016/j.padiff.2024.100712>
- [15] Đorđević, J., & Dahl, K. R. (2024). Stochastic optimal control of pre-exposure prophylaxis for HIV infection for a jump model. Journal of Mathematical Biology, 89(5), 55[. https://doi.org/10.1007/s00285-024-02151-3](https://doi.org/10.1007/s00285-024-02151-3)
- [16] Hambel, C., Kraft, H., & van der Ploeg, F. (2024). Asset diversification versus climate action. International Economic Review, 65(3), 1323-1355[. https://doi.org/10.1111/iere.12691](https://doi.org/10.1111/iere.12691)
- [17] Zhang, J., & Ren, L. (2024). A mean field game model of green economy. Digital Finance, published online. <https://doi.org/10.1007/s42521-024-00118-z>
- [18] Ramakrishnan, K. A., Rinaldo, T., Rodriguez-Iturbe, I., & Durán Vinent, O. (2024). Permanent loss of barrier island resilience due to a critical transition in dune ecosystems. Communications Earth & Environment, 5(1), 447[. https://doi.org/10.1038/s43247-024-01611-4](https://doi.org/10.1038/s43247-024-01611-4)
- [19] Vinent, O. D., Schaffer, B. E., & Rodriguez-Iturbe, I. (2021). Stochastic dynamics of barrier island elevation. Proceedings of the National Academy of Sciences, 118(1), e2013349118. [https://doi.org/10.1038/s43247-024-](https://doi.org/10.1038/s43247-024-01611-4) [01611-4](https://doi.org/10.1038/s43247-024-01611-4)
- [20] Dijkstra, H. A., Biemond, B., Lee, J., & de Swart, H. E. (2023). Explaining the Statistical Properties of Salt Intrusion in Estuaries Using a Stochastic Dynamical Modeling Approach. Water Resources Research, 59(7), e2023WR034454.<https://doi.org/10.1029/2023WR034454>
- [21] Calvani, G., & Perona, P. (2023). Splitting probabilities and mean first-passage times across multiple thresholds of jump-and-drift transition paths. Physical Review E, 108(4), 044105. <https://doi.org/10.1103/PhysRevE.108.044105>
- [22] Latella, M., Notti, D., Baldo, M., Giordan, D., & Camporeale, C. (2024). Short-term biogeomorphology of a gravel‐bed river: Integrating remote sensing with hydraulic modelling and field analysis. Earth Surface Processes and Landforms, 49(3), 1156-1178[. https://doi.org/10.1002/esp.5760](https://doi.org/10.1002/esp.5760)
- [23] Olson, R., Timmermann, A., Lee, J. Y., & An, S. I. (2021). A low order dynamical model for runoff predictability. Climate Dynamics, 56, 399-422.<https://doi.org/10.1007/s00382-020-05479-w>
- [24] Yoshioka, H., & Yoshioka, Y. (2023a). Stochastic streamflow and dissolved silica dynamics with application to the worst-case long-run evaluation of water environment. Optimization and Engineering, 24(3), 1577-1610. <https://doi.org/10.1007/s11081-022-09743-2>
- [25] Yoshioka, H., & Yoshioka, Y. (2023b). Dual stochastic descriptions of streamflow dynamics under model ambiguity through a Markovian embedding. Journal of Mathematics in Industry, 13(1), 7. <https://doi.org/10.1186/s13362-023-00135-4>
- [26] Fatoni, H., Kholis, A. N., Kurniawan, I. A., Haryono, E., Waskito, W. A., Cahyadi, A., ... & Adji, T. N. (2024). Use of discharge variability, aquifer memory effect, and recession curve analysis in several springs to characterize karst aquifers in the tropical area. Geosciences Journal, 28(6), 925-939. [https://doi.org/10.1007/s12303-024-](https://doi.org/10.1007/s12303-024-0034-0) [0034-0](https://doi.org/10.1007/s12303-024-0034-0)
- [27] Maftei, C., Barbulescu, A., & Carsteanu, A. A. (2016). Long-range dependence in the time series of Taita River discharges. Hydrological Sciences Journal, 61(9), 1740-1747[. https://doi.org/10.1080/02626667.2016.1171869](https://doi.org/10.1080/02626667.2016.1171869)
- [28] Guo, T., Song, S., Liu, X., Wang, H., & Zhang, G. (2024a). A comparison of several cyclo-stationary and fractionally differenced time-varying autoregressive models in runoff simulation and prediction. Journal of Hydrology, 131509[. https://doi.org/10.1016/j.jhydrol.2024.131509](https://doi.org/10.1016/j.jhydrol.2024.131509)
- [29] Pizarro, A., Acuña, P., Dimitriadis, P., Iliopoulou, T., & Koutsoyiannis, D. (2024). Past, present, and future of the Hurst-Kolmogorov dynamics in Stochastics: A bibliometric analysis of the last 50 years in water resources. Journal of Hydrology, 643, 132008.<https://doi.org/10.1016/j.jhydrol.2024.132008>
- [30] Rahmani, F., & Fattahi, M. H. (2024). Long-term evaluation of land use/land cover and hydrological drought patterns alteration consequences on river water quality. Environment, Development and Sustainability, 26(7), 19051-19068[. https://doi.org/10.1007/s10668-023-03302-0](https://doi.org/10.1007/s10668-023-03302-0)
- [31] Abi Jaber, E. (2019). Lifting the Heston model. Quantitative finance, 19(12), 1995-2013. <https://doi.org/10.1080/14697688.2019.1615113>
- [32] Cuchiero, C., & Teichmann, J. (2020). Generalized Feller processes and Markovian lifts of stochastic Volterra processes: the affine case. Journal of evolution equations, 20(4), 1301-1348[. https://doi.org/10.1007/s00028-020-](https://doi.org/10.1007/s00028-020-00557-2) [00557-2](https://doi.org/10.1007/s00028-020-00557-2)
- [33] Kanazawa, K., & Sornette, D. (2024). Standard form of master equations for general non-Markovian jump processes: The Laplace-space embedding framework and asymptotic solution. Physical Review Research, 6(2), 023270[. https://doi.org/10.1103/PhysRevResearch.6.023270](https://doi.org/10.1103/PhysRevResearch.6.023270)
- [34] Wiśniewski, M., & Spiechowicz, J. (2024). Dynamics of non-Markovian systems: Markovian embedding versus effective mass approach. Physical Review E, 110(5), 054117.<https://doi.org/10.1103/PhysRevE.110.054117>
- [35] Bondi, A., Pulido, S., & Scotti, S. (2024). The rough Hawkes Heston stochastic volatility model. Mathematical Finance, 34(4), 1197-1241[. https://doi.org/10.1111/mafi.12432](https://doi.org/10.1111/mafi.12432)
- [36] Damian, C., & Frey, R. (2024). Detecting rough volatility: a filtering approach. Quantitative Finance, 24(10), 1493-1508[. https://doi.org/10.1080/14697688.2024.2399284](https://doi.org/10.1080/14697688.2024.2399284)
- [37] Dupret, J. L., & Hainaut, D. (2022). A fractional Hawkes process for illiquidity modeling. Mathematics and Financial Economics, published online[. https://doi.org/10.1007/s11579-024-00379-7](https://doi.org/10.1007/s11579-024-00379-7)
- [38] Yang, W., Ma, J., & Cui, Z. (2024). A general valuation framework for rough stochastic local volatility models and applications. European Journal of Operational Research, published online. <https://doi.org/10.1016/j.ejor.2024.11.002>
- [39] Guo, X., Jiang, S., Xiong, Y., & Zhang, J. (2024b). Compressing the memory variables in constant-Q viscoelastic wave propagation via an improved sum-of-exponentials approximation. Journal of Computational Physics, 518, 113326[. https://doi.org/10.1016/j.jcp.2024.113326](https://doi.org/10.1016/j.jcp.2024.113326)
- [40] He, L., Shi, L., Song, W., Shen, J., Wang, L., Hu, X., Zha, Y. (2024). Synergizing Intuitive Physics and Big Data in Deep Learning: Can We Obtain Process Insights While Maintaining State-Of-The-Art Hydrological Prediction Capability? Water Resources Research, 60(12), e2024WR037582<https://doi.org/10.1029/2024WR037582>
- [41] Fasen, V., & Klüppelberg, C. (2005). Extremes of supOU processes. In Stochastic Analysis and Applications: The Abel Symposium 2005 (pp. 339-359). Berlin, Heidelberg: Springer, Berlin, Heidelberg. [https://doi.org/10.1007/978-3-540-70847-6\\_14](https://doi.org/10.1007/978-3-540-70847-6_14)
- [42] Yoshioka, H., & Yoshioka, Y. (2024a). Generalized divergences for statistical evaluation of uncertainty in long-memory processes. Chaos, Solitons & Fractals, 182, 114627[. https://doi.org/10.1016/j.chaos.2024.114627](https://doi.org/10.1016/j.chaos.2024.114627)
- [43] Yoshioka, H., & Yoshioka, Y. (2024b). Risk assessment of river water quality using long-memory processes subject to divergence or Wasserstein uncertainty. Stochastic Environmental Research and Risk Assessment, 38, 3007-3030[. https://doi.org/10.1007/s00477-024-02726-y](https://doi.org/10.1007/s00477-024-02726-y)
- [44] Barndorff-Nielsen, O. E., & Stelzer, R. (2013). The multivariate supOU stochastic volatility model. Mathematical Finance: An International Journal of Mathematics, Statistics and Financial Economics, 23(2), 275-296. <https://doi.org/10.1111/j.1467-9965.2011.00494.x>
- [45] Duffie, D., Filipović, D., & Schachermayer, W. (2003). Affine processes and applications in finance. The Annals of Applied Probability, 13(3), 984-1053[. https://doi.org/10.1214/aoap/1060202833](https://doi.org/10.1214/aoap/1060202833)
- [46] Bayer, C., Harang, F. A., & Pigato, P. (2021). Log-modulated rough stochastic volatility models. SIAM Journal on Financial Mathematics, 12(3), 1257-1284[. https://doi.org/10.1137/20M135902X](https://doi.org/10.1137/20M135902X)
- [47] de Truchis, G., Desgraupes, B., & Dumitrescu, E. I. (2024). Assessing volatility persistence in fractional Heston models with self-exciting jumps. Econometric Reviews, published online. <https://doi.org/10.1080/07474938.2024.2409475>
- [48] Wu, P., Muzy, J. F., & Bacry, E. (2022). From rough to multifractal volatility: The log S-fBM model. Physica A: Statistical Mechanics and its Applications, 604, 127919[. https://doi.org/10.1016/j.physa.2022.127919](https://doi.org/10.1016/j.physa.2022.127919)
- [49] Wang, H., Song, S., Zhang, G., Ayantobo, O. O., & Guo, T. (2023). Stochastic volatility modeling of daily streamflow time series. Water Resources Research, 59(1), e2021WR031662. <https://doi.org/10.1029/2021WR031662>
- [50] Jing, H., Zhong, D., Zhang, H., Shi, X., & Wang, Y. (2020). Accumulation phenomena in fluvial processes and the corresponding stochastic model. Journal of Geographical Sciences, 30, 1021-1040. <https://doi.org/10.1007/s11442-020-1768-x>
- [51] Mansour, M. B. (2023). On modelling water quality with stochastic differential equations. The ANZIAM Journal, 65(3), 273-284. <https://doi.org/10.1017/S1446181123000214>
- [52] Gothwal, R., & Thatikonda, S. (2020). Modeling transport of antibiotic resistant bacteria in aquatic environment using stochastic differential equations. Scientific reports, 10(1), 15081. [https://doi.org/10.1038/s41598-020-](https://doi.org/10.1038/s41598-020-72106-3) [72106-3](https://doi.org/10.1038/s41598-020-72106-3)
- [53] Park, J., & Rao, P. S. C. (2014). Regime shifts under forcing of non-stationary attractors: Conceptual model and case studies in hydrologic systems. Journal of contaminant hydrology, 169, 112-122. <https://doi.org/10.1016/j.jconhyd.2014.08.005>
- [54] Carpenter, S. R., & Brock, W. A. (2024). Stochastic dynamics of phycocyanin in years of contrasting phosphorus load. Ecosphere, 15(6), e4903[. https://doi.org/10.1002/ecs2.4903](https://doi.org/10.1002/ecs2.4903)
- [55] Chen, J., Gao, T., Li, Y., & Duan, J. (2023). Detecting the most probable high dimensional transition pathway based on optimal control theory. 127, 217-236.<https://doi.org/10.1016/j.apm.2023.12.008>
- [56] Song, J., & Wu, D. (2023). Modeling forest carbon sink trading with carbon credit using stochastic differential game. Environmental Science and Pollution Research, 30(26), 68934-68950. [https://doi.org/10.1007/s11356-](https://doi.org/10.1007/s11356-023-26974-7) [023-26974-7](https://doi.org/10.1007/s11356-023-26974-7)
- [57] Song, J., Wu, D., Bian, Y., & Dong, J. (2024). A decision support system based on stochastic differential game model in pollution control chain. IEEE Transactions on Systems, Man, and Cybernetics: Systems. 54(3), 1670- 1682[. https://doi.org/10.1109/TSMC.2023.3300469](https://doi.org/10.1109/TSMC.2023.3300469)
- [58] Chong, L., Zhong, J., Sun, Z., & Hu, C. (2023). Temporal variations and trends prediction of water quality during 2010–2019 in the middle Yangtze River, China. Environmental Science and Pollution Research, 30(11), 28745- 28758.<https://doi.org/10.1007/s11356-022-23968-9>
- [59] Kovács, J., Hatvani, I. G., Korponai, J., & Kovács, I. S. (2010). Morlet wavelet and autocorrelation analysis of long-term data series of the Kis-Balaton water protection system (KBWPS). Ecological Engineering, 36(10), 1469-1477[. https://doi.org/10.1016/j.ecoleng.2010.06.028](https://doi.org/10.1016/j.ecoleng.2010.06.028)
- [60] Rahmani, F., & Fattahi, M. H. (2024). Long-term evaluation of land use/land cover and hydrological drought patterns alteration consequences on river water quality. Environment, Development and Sustainability, 26(7), 19051-19068[. https://doi.org/10.1007/s10668-023-03302-0](https://doi.org/10.1007/s10668-023-03302-0)
- [61] Rozental, O. M., & Tambieva, D. A. (2021). Cyclic variations of quality characteristics in river water in an industrial region. Water Science and Technology, 83(4), 854-862[. https://doi.org/10.2166/wst.2020.597](https://doi.org/10.2166/wst.2020.597)
- [62] Spezia, L., Vinten, A., Paroli, R., & Stutter, M. (2021). An evolutionary Monte Carlo method for the analysis of turbidity high-frequency time series through Markov switching autoregressive models. Environmetrics, 32(8), e2695[. https://doi.org/10.1002/env.2695](https://doi.org/10.1002/env.2695)
- [63] Qin, C., Li, S. L., Wu, Y., Bass, A. M., Luo, W., Ding, H., ... & Zhang, P. (2024). High sensitivity of dissolved organic carbon transport during hydrological events in a small subtropical karst catchment. Science of The Total Environment, 946, 174090.<https://doi.org/10.1016/j.scitotenv.2024.174090>
- [64] Tunqui Neira, J. M., Andréassian, V., Tallec, G., & Mouchel, J. M. (2021). Multi-objective fitting of concentration-discharge relationships. Hydrological Processes, 35(11), e14428. <https://doi.org/10.1002/hyp.14428>
- [65] Wichman, G., Johnston, S. G., & Maher, D. T. (2024). Antimony Flux and Transport Dynamics in a Mining-Impacted River Is Linked to Catchment Hydrodynamics and Climate Oscillations. Hydrological Processes, 38(11), e15323[. https://doi.org/10.1002/hyp.15323](https://doi.org/10.1002/hyp.15323)
- [66] Wymore, A. S., Larsen, W., Kincaid, D. W., Underwood, K. L., Fazekas, H. M., McDowell, W. H., ... & Webster, A. J. (2023). Revisiting the origins of the power-law analysis for the assessment of concentration-discharge relationships. Water Resources Research, 59(8), e2023WR034910[. https://doi.org/10.1029/2023WR034910](https://doi.org/10.1029/2023WR034910)
- [67] Zhan, Q., Kong, X., & Rinke, K. (2022). High-frequency monitoring enables operational opportunities to reduce the dissolved organic carbon (DOC) load in Germany's largest drinking water reservoir. Inland Waters, 12(2), 245-260[. https://doi.org/10.1080/20442041.2021.1987796](https://doi.org/10.1080/20442041.2021.1987796)
- [68] Barndorff-Nielsen, O. E. (2001). Superposition of Ornstein–Uhlenbeck type processes. Theory of Probability & Its Applications, 45(2), 175-194.<https://doi.org/10.1137/S0040585X97978166>
- [69] Iglói, E., & Terdik, G. (2003). Superposition of diffusions with linear generator and its multifractal limit process. ESAIM: Probability and Statistics, 7, 23-88.<https://doi.org/10.1051/ps:2003008>
- [70] Leonenko, N., Liu, A., & Shchestyuk, N. (2024). Student models for a risky asset with dependence: Option pricing and Greeks. Austrian Journal of Statistics, 54(1), 138-165.<https://doi.org/10.17713/ajs.v54i1.1952>
- [71] Grahovac, D., & Kevei, P. (2025). Tail Behavior and Almost Sure Growth Rate of Superpositions of Ornstein– Uhlenbeck-type Processes. Journal of theoretical probability, 38(1), 1-15. [https://doi.org/10.1007/s10959-024-](https://doi.org/10.1007/s10959-024-01374-w) [01374-w](https://doi.org/10.1007/s10959-024-01374-w)
- [72] Grahovac, D., Leonenko, N. N., & Taqqu, M. S. (2019). Limit theorems, scaling of moments and intermittency

for integrated finite variance supOU processes. Stochastic processes and their applications, 129(12), 5113-5150. https://doi.org/10.1016/j.spa.2019.01.010

- [73] Chevalier, E., Pulido, S., & Zúñiga, E. (2022). American options in the Volterra Heston model. SIAM Journal on Financial Mathematics, 13(2), 426-458[. https://doi.org/10.1137/21M140674X](https://doi.org/10.1137/21M140674X)
- [74] Motte, E., & Hainaut, D. (2024). Partial hedging in rough volatility models. SIAM Journal on Financial Mathematics, 15(3), 601-652[. https://doi.org/10.1137/23M1583090](https://doi.org/10.1137/23M1583090)
- [75] Yoshioka, H., Tanaka, T., Yoshioka, Y., & Hashiguchi, A. (2023). Stochastic optimization of a mixed moving average process for controlling non-Markovian streamflow environments. Applied Mathematical Modelling, 116, 490-509[. https://doi.org/10.1016/j.apm.2022.11.009](https://doi.org/10.1016/j.apm.2022.11.009)
- [76] Cox, S., Karbach, S., & Khedher, A. (2022). An infinite-dimensional affine stochastic volatility model. Mathematical Finance, 32(3), 878-906[. https://doi.org/10.1111/mafi.12347](https://doi.org/10.1111/mafi.12347)
- [77] Friesen, M., & Karbach, S. (2024). Stationary covariance regime for affine stochastic covariance models in Hilbert spaces. Finance and Stochastics,28, 1077-1116.<https://doi.org/10.1007/s00780-024-00543-3>
- [78] Mazilamani, L. S., Walsh, R. P. D., Annammala, K. V., Bidin, K., Yusop, Z., Reynolds, G., & Nainar, A. (2024). Concentration-discharge hysteresis: current approaches and future directions for quantifying pollutant dynamics in storm events—with a particular focus on the tropics. Sustainable Water Resources Management, 10(4), 156. <https://doi.org/10.1007/s40899-024-01130-2>
- [79] Botter, G. (2010). Stochastic recession rates and the probabilistic structure of stream flows. Water Resources Research, 46(12).<https://doi.org/10.1029/2010WR009217>
- [80] Hameed, M., Nayak, M. A., & Ahanger, M. A. (2023). Event-Based Recession Analysis for Estimation of Basin-Wide Characteristic Drainage Timescale and Groundwater Storage Trends. Water Resources Research, 59(12), e2023WR035829.<https://doi.org/10.1029/2023WR035829>
- [81] Lee, J. Y., Yang, C. J., Peng, T. R., Lee, T. Y., & Huang, J. C. (2023). Landscape structures regulate the contrasting response of recession along rainfall amounts. Hydrology and Earth System Sciences, 27(23), 4279-4294. <https://doi.org/10.5194/hess-27-4279-2023>
- [82] Mudelsee, M. (2007). Long memory of rivers from spatial aggregation. Water Resources Research, 43(1). <https://doi.org/10.1029/2006WR005721>
- [83] Di Dato, M., Bellin, A., Cvetkovic, V., Dagan, G., Dietrich, P., Fiori, A., ... & Attinger, S. (2023). Baseflow statistics in aggregated catchments. Water Resources Research, 59(12), e2023WR035894. <https://doi.org/10.1029/2023WR035894>
- [84] Gomez, C., Iyer, G., Le, H., & Novikov, A. (2023). An oscillator driven by algebraically decorrelating noise. Communications in Mathematical Physics, 402(1), 231-284.<https://doi.org/10.1007/s00220-023-04744-3>
- [85] Yoshioka, H. (2022). Fitting a superposition of Ornstein–Uhlenbeck process to time series of discharge in a perennial river environment. ANZIAM Journal, 63, C84-C96.<https://doi.org/10.21914/anziamj.v63.16985>
- [86] Deelstra, G., Grasselli, M., & Van Weverberg, C. (2019). Explosion time for some Laplace transforms of the Wishart process. Stochastic Models, 35(1), 89-104[. https://doi.org/10.1080/15326349.2019.1578237](https://doi.org/10.1080/15326349.2019.1578237)
- [87] Gerhold, S., Gerstenecker, C., & Pinter, A. (2019). Moment explosions in the rough Heston model. Decisions in Economics and Finance, 42, 575-608[. https://doi.org/10.1007/s10203-019-00267-6](https://doi.org/10.1007/s10203-019-00267-6)
- [88] Grabchak, M. (2021). An exact method for simulating rapidly decreasing tempered stable distributions in the finite variation case. Statistics & Probability Letters, 170, 109015.<https://doi.org/10.1016/j.spl.2020.109015>
- [89] Kawai, R., & Masuda, H. (2011). On simulation of tempered stable random variates. Journal of Computational and Applied Mathematics, 235(8), 2873-2887[. https://doi.org/10.1016/j.cam.2010.12.014](https://doi.org/10.1016/j.cam.2010.12.014)
- [90] MLIT, Japan. Japan. (2024a). [https://www.mlit.go.jp/river/toukei\\_chousa/kasen/jiten/nihon\\_kawa/0713\\_hiikawa/0713\\_hiikawa\\_00.html.](https://www.mlit.go.jp/river/toukei_chousa/kasen/jiten/nihon_kawa/0713_hiikawa/0713_hiikawa_00.html) Last Accessed on November 24, 2024.
- [91] Takeda, I. (2023). Changes in river water quality for 30 years in a watershed characterized by population decline. Journal of Environmental Conservation Engineering, 52(1), 41-49, in Japanese with English abstract]. [https://doi.org/10.5956/jriet.52.1\\_41](https://doi.org/10.5956/jriet.52.1_41)
- [92] MLIT, Japan. (2024b).<https://www.mlit.go.jp/tagengo-db/en/H30-00978.html> Last accessed on November 24, 2024.
- [93] Kawaida, S., Horinouchi, M., Kurata, K., & Toda, K. (2024). Differences in benthic assemblage structures between vegetated and unvegetated habitats in a salt marsh in Lake Shinji, western Japan. Fisheries Science, published online[. https://doi.org/10.1007/s12562-024-01831-9](https://doi.org/10.1007/s12562-024-01831-9)
- [94] Hafeez, M. A., & Inoue, T. (2024). Three-dimensional hydrodynamic modelling of saltwater ingression and circulation in a Brackish Lake Shinji, Japan. Advances in Water Resources, 184, 104627. <https://doi.org/10.1016/j.advwatres.2024.104627>
- [95] Kim, S., Ando, T., Nakamura, Y., Hayashi, S., & Kawaida, S. (2024). Clustering evaluation of water quality for various classes of in-flow rivers in connected brackish lakes. Environmental Monitoring and Assessment, 196(6), 501[. https://doi.org/10.1007/s10661-024-12670-7](https://doi.org/10.1007/s10661-024-12670-7)
- [96] Atazadeh, E., Gell, P., Mills, K., Barton, A., & Newall, P. (2024). Ecological and statistical models to configure flow regime for environment benefit in highly engineered rivers: a case study in the MacKenzie River, Southeast Australia. Environmental Science and Pollution Research, 31(5), 7408-7427. [https://doi.org/10.1007/s11356-](https://doi.org/10.1007/s11356-023-31596-0) [023-31596-0](https://doi.org/10.1007/s11356-023-31596-0)
- [97] Tian, J., Wang, G., Xiang, D., Huang, S., & Li, W. (2024). Causality analysis and prediction of riverine algal blooms by combining empirical dynamic modeling and machine learning techniques. Water Resources Research, 60(5), e2023WR036334.<https://doi.org/10.1029/2023WR036334>
- [98] Yuan, L. L., & Paul, M. J. (2024). Predicting lake chlorophyll from stream phosphorus concentrations. JAWRA Journal of the American Water Resources Association. Published online. [https://doi.org/10.1111/1752-](https://doi.org/10.1111/1752-1688.13243) [1688.13243](https://doi.org/10.1111/1752-1688.13243)
- [99] Biedunkova, O., & Kuznietsov, P. (2024). Investigation of the formation and variability of dissolved inorganic carbon and dissolved organic carbon in the water of a small river (on the example of the Styr River, Ukraine). Environmental Monitoring and Assessment, 196(11), 1-17[. https://doi.org/10.1007/s10661-024-13309-3](https://doi.org/10.1007/s10661-024-13309-3)
- [100] Starr, S., Johnston, S. E., Sobolev, N., Perminova, I., Kellerman, A., Fiske, G., ... & Spencer, R. G. (2023). Characterizing uncertainty in Pan‐Arctic land‐ocean dissolved organic carbon flux: Insights from the Onega River, Russia. Journal of Geophysical Research: Biogeosciences, 128(5), e2022JG007073. <https://doi.org/10.1029/2022JG007073>
- [101] Das, A., Kumar, M., Jha, P. K., Kumari, R., Panday, D. P., Hdeib, R., ... & Deshpande, R. D. (2022). Isotopic and hydrogeochemical tracking of dissolved nutrient dynamics in the Brahmaputra River System: A source delineation perspective. Chemosphere, 307, 135757[. https://doi.org/10.1016/j.chemosphere.2022.135757](https://doi.org/10.1016/j.chemosphere.2022.135757)
- [102] Hachgenei, N., Nord, G., Spadini, L., Ginot, P., Voiron, C., & Duwig, C. (2024). Transit time tracing using wetness-adaptive StorAge Selection functions—application to a Mediterranean catchment. Journal of Hydrology, 131267[. https://doi.org/10.1016/j.jhydrol.2024.131267](https://doi.org/10.1016/j.jhydrol.2024.131267)
- [103] Jewell, K., Myers, K. D., Lipsi, M., Hossain, S., Datta, S., Cardenas, M. B., ... & Knappett, P. S. (2023). Redox trapping of arsenic in hyporheic zones modified by silicate weathering beneath floodplains. Applied Geochemistry, 159, 105831.<https://doi.org/10.1016/j.apgeochem.2023.105831>
- [104] MLIT, Japan. (2024c).<http://www1.river.go.jp/cgi-bin/SiteInfo.exe?ID=307041287705030> Last accessed on November 25, 2024.
- [105] Wu, Y. F., & Hu, J. Q. (2023). Method of moments estimation for the superposition of square-root diffusions. Probability in the Engineering and Informational Sciences. Published online. <https://doi.org/10.1017/S0269964824000147>
- [106] Abi Jaber, E., Illand, C., & Li, S. (2024). Joint SPX & VIX calibration with Gaussian polynomial volatility models: Deep pricing with quantization hints. Mathematical Finance. Published online. <https://doi.org/10.1111/mafi.12451>
- [107] Dialameh, B., & Ghane, E. (2023). Investigation of phosphorus transport dynamics using high-frequency monitoring at a subsurface-drained field in the Western Lake Erie Basin. Journal of Great Lakes Research, 49(4), 778-789[. https://doi.org/10.1016/j.jglr.2023.04.005](https://doi.org/10.1016/j.jglr.2023.04.005)
- [108] Rodríguez-Blanco, M. L., Taboada-Castro, M. T., & Taboada-Castro, M. M. (2023). Improving the understanding of N transport in a rural catchment under Atlantic climate conditions from the analysis of the concentration– discharge relationship derived from a high-frequency data set. Hydrology and Earth System Sciences, 27(6), 1243-1259[. https://doi.org/10.5194/hess-27-1243-2023](https://doi.org/10.5194/hess-27-1243-2023)
- [109] Xue, J., Chen, L., Zhang, W., Yuan, J., & Dong, B. (2024). A new method for analyzing storm event dischargeconcentration hysteresis: The upper Yangtze River as an example. Water Resources Research, 60(12), e2023WR035795.<https://doi.org/10.1029/2023WR035795>
- [110] Speir, S. L., Rose, L. A., Blaszczak, J. R., Kincaid, D. W., Fazekas, H. M., Webster, A. J., ... & Wymore, A. S. (2024). Catchment concentration–discharge relationships across temporal scales: A review. Wiley Interdisciplinary Reviews: Water, 11(2), e1702. <https://doi.org/10.1002/wat2.1702>
- [111] Cairoli, M., Souza, F., Stroomberg, G., Postma, G., Buydens, L., & Jansen, J. (2024). BaHys—A Bayesian modeling framework for long-term concentration-discharge hysteresis: A case study on chloride. Water Resources Research, 60(6), e2023WR035427.<https://doi.org/10.1029/2023WR035427>
- [112] Knapp, J. L., Li, L., & Musolff, A. (2022). Hydrologic connectivity and source heterogeneity control concentration–discharge relationships. Hydrological Processes, 36(9), e14683. <https://doi.org/10.1002/hyp.14683>
- [113] Roberts, M. E., Kim, D., Lu, J., & Hamilton, D. P. (2023). HARP: A suite of parameters to describe the hysteresis of streamflow and water quality constituents. Journal of Hydrology, 626, 130262. <https://doi.org/10.1016/j.jhydrol.2023.130262>
- [114] Zhu, M., Yu, X., Li, Z., Xu, X., & Ye, Z. (2023). Quantifying and interpreting the hysteresis patterns of monthly sediment concentration and water discharge in karst watersheds. Journal of Hydrology, 618, 129179. <https://doi.org/10.1016/j.jhydrol.2023.129179>
- [115] Nishimoto, M., Miyashita, T., & Fukasawa, K. (2024). Spatiotemporal smoothing of water quality in a complex riverine system with physical barriers. Science of The Total Environment, 948, 174843. <https://doi.org/10.1016/j.scitotenv.2024.174843>
- [116] Puica, M., & Benth, F. E. (2023). A spatio-temporal model for predicting wind speeds in Southern California. Communications in Statistics: Case Studies, Data Analysis and Applications, 9(3), 321-349. <https://doi.org/10.1080/23737484.2023.2217137>