Modular Features of Superstring Scattering Amplitudes: Generalised Eisenstein Series and Theta Lifts

Daniele Dorigoni^{1,2}, Michael B. Green³, Congkao Wen⁴

¹ Centre for Particle Theory & Department of Mathematical Sciences, Durham University, Lower Mountjoy, Stockton Road, Durham DH1 3LE, UK

² Max-Planck-Institut für Gravitationsphysik (Albert-Einstein-Institut), am Mühlenberg 1, Potsdam, 14476, Germany

³ Department of Applied Mathematics and Theoretical Physics Wilberforce Road, Cambridge CB3 0WA, UK

⁴ Centre for Theoretical Physics, Department of Physics and Astronomy, Queen Mary University of London, London, E1 4NS, UK

Abstract

In previous papers it has been shown that the coefficients of terms in the large-N expansion of a certain integrated four-point correlator of superconformal primary operators in $\mathcal{N} = 4$ supersymmetric Yang-Mills theory are rational sums of real-analytic Eisenstein series and "generalised Eisenstein series". The latter are novel modular functions first encountered in the context of graviton amplitudes in type IIB superstring theory. Similar modular functions, known as two-loop modular graph functions, are also encountered in the low-energy expansion of the integrand of genus-one closed superstring amplitudes. In this paper we further develop the mathematical structure of such generalised Eisenstein series emphasising, in particular, the occurrence of L-values of holomorphic cusp forms in their Fourier mode decomposition. We show that both the coefficients in the large-N expansion of this integrated correlator and two-loop modular graph functions admit a unifying description in terms of four-dimensional lattice sums generated by theta lifts of certain local Maass functions, which generalise the structure of real-analytic Eisenstein series. Through the theta lift representation, we demonstrate that elements belonging to these two families of non-holomorphic modular functions can be expressed as rational linear combinations of generalised Eisenstein series for which all the L-values of holomorphic cusp forms

Contents

| 1 | Overview and results 1.1 Introduction 1.2 Results 1.3 Outline | 3 3 4 6 |
|---|--|-----------------------------|
| 2 | Theta lift of local Maass functions2.1Theta lift: definitions and properties2.2Modular local polynomials and local Maass functions | 7 7 10 |
| 3 | Inhomogeneous Laplace Equation3.1Boundary terms3.2Alternative approach3.3General expression for $\mathcal{E}_{i,j}^w(\tau)$ | 14 15 19 21 |
| 4 | Structure of generalised Eisenstein series 4.1 Fourier mode expansion 4.2 L-values and generalised Eisenstein series 4.3 Absence of L-values in theta-lifted local Maass functions | 22 22 25 33 |
| 5 | Discussion 5.1 Cuspidal generalised Eisenstein series 5.2 Finite-N integrated correlators 5.3 S-dual modular forms | 39 39 39 40 |
| A | Non-holomorphic Eisenstein series | 41 |
| в | Modular local polynomials | 42 |
| С | Holomorphic cusp forms and their L-values | 44 |

1 Overview and results

1.1 Introduction

Two interesting and related classes of modular functions have arisen from the study of distinct aspects of closed superstring scattering amplitudes. Elements in these families can both be expressed as rational linear combinations of *Generalised Eisenstein Series* (GESs), which are defined to be modular invariant functions satisfying the inhomogeneous Laplace eigenvalue equation

$$(\Delta_{\tau} - s(s-1))\mathcal{E}(s; s_1, s_2; \tau) = E(s_1; \tau) E(s_2; \tau), \qquad (1.1)$$

where $s \in \mathbb{N}$ and $\tau \coloneqq \tau_1 + i\tau_2 \in \mathfrak{H}$ is a complex modulus, which takes values in a upper-half complex plane $\mathfrak{H} \coloneqq \{\tau \in \mathbb{C} \mid \operatorname{Im}(\tau) > 0\}$ and Δ_{τ} is the hyperbolic laplacian, $\Delta_{\tau} \coloneqq \tau_2^2(\partial_{\tau_1}^2 + \partial_{\tau_2}^2)$. The modular invariant function $E(s;\tau)$ appearing in the source term in (1.1) is a non-holomorphic Eisenstein series, which satisfies

$$(\Delta_{\tau} - s(s-1)) E(s;\tau) = 0.$$
(1.2)

The first class of GESs that we consider has $s_1, s_2 \in \mathbb{N}$ and it appears to be the relevant vector space of non-holomorphic modular invariant functions to describe *two-loop Modular Graph Functions* (MGFs) [1, 2, 3]. Interest in MGFs originated from the study of the low energy expansion of the integrand for genusone contributions to four-graviton scattering in either of the type II superstring theories [1, 4]. The name MGF originates from the fact that these functions are the values of connected Feynman graphs for a free scalar field theory living on a two-torus with complex structure τ . Modular invariance (SL(2, Z)-invariance) is a direct consequence of diffeomorphism invariance on the genus-one world-sheet torus. The mathematical structure of these functions generalises in an obvious manner to modular graph *forms* whose study has led to an impressive symbiosis with novel ideas in algebraic geometry [5, 6, 7]. The construction of MGFs may be extended to higher loops but in this paper we will focus on the two-loop cases since these are related to GESs. With a slight abuse of terminology, in what follows the acronym MGFs specifically refers to two-loop modular graph functions.

Two-loop MGFs are defined by the following lattice sums,

$$C_{a,b,c}(\tau) \coloneqq \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1+p_2+p_3=0}} \frac{(\tau_2/\pi)^{a+b+c}}{|p_1|^{2a}|p_2|^{2b}|p_3|^{2c}},$$
(1.3)

where $a, b, c \in \mathbb{N}$ and the sum is over three complex lattice momenta $p_i = m_i + n_i \tau \in \Lambda' = \mathbb{Z} + \tau \mathbb{Z} \setminus \{0\}$, with i = 1, 2, 3, subject to the conservation condition $p_1 + p_2 + p_3 = 0$. Notably, specific linear combinations [8] of MGFs satisfy inhomogeneous Laplace eigenvalue equations with source terms that are bilinear in nonholomorphic Eisenstein series of the form (1.1) with $s_1, s_2 \in \mathbb{N}$. However, for values of $s \ge 6$ modular-invariant solutions of (1.1) with integer indices cannot all be expressed as superpositions of MGFs. The construction of GESs requires additional contributions proportional to *L*-functions of holomorphic cusp forms [9, 10] with modular weight 2s. Conjecturally, MGFs select only rational combinations of this family of GES for which the *L*-value contributions precisely cancel.

A second class of GESs consists of solutions to (1.1) with sources given by non-holomorphic Eisenstein series with half-integer indices, i.e. with $s_1, s_2 \in \mathbb{N} + 1/2$. An element in this family of GESs arises in the low-energy expansion of four-graviton scattering amplitudes in type IIB superstring theory [11, 12]. In this case SL(2, \mathbb{Z}) covariance is known as S-duality and can be motivated by interpreting superstring theory in terms of eleven-dimensional M-theory compactified on a two-torus. The complex structure of the torus is here identified with the complex axion-dilaton field, $\tau = \tau_1 + i\tau_2$, where $\tau_2 = 1/g_s$ and g_s is the string coupling constant. For lack of better terminology, we refer to this second class of non-holomorphic modular invariant functions arising in the context scattering amplitudes in type IIB superstring theory as S-Dual Modular Functions (SMFs).

SMFs also appear in the study of correlation functions in $\mathcal{N} = 4$ supersymmetric Yang–Mills (SYM) in four dimensions. This four-dimensional quantum field theory exhibits Montonen–Olive duality [13], in which SL(2, Z) acts on the complex Yang–Mills coupling constant $\tau = \theta/2\pi + 4\pi i/g_{_{YM}}^2$. This fits beautifully with the AdS/CFT conjecture [14], where type IIB superstring theory in an $AdS_5 \times S^5$ background is identified with $\mathcal{N} = 4$ SYM with SU(N) gauge group. S-duality in the dual superstring theory here is interpreted as the holographic image of Montonen–Olive duality of $\mathcal{N} = 4$ SYM. It was shown in [15, 16], that certain integrated correlators of four superconformal primary operators in the stress tensor multiplet are computable using supersymmetric localisation for all N [17], and their properties depend sensitively on the choice of integration measures (see e.g. the review [18]). In the simplest example the large-N expansion of the correlator is a series of half-integer powers of 1/N and the coefficients are rational linear combinations of non-holomorphic Eisenstein series with half-integer indices [19]. The second example of integrated correlators has a large-N expansion with both half-integer and integer powers of 1/N [20, 21]. The half-integer powers again have coefficients that are rational linear combinations of non-holomorphic Eisenstein series with halfinteger indices, while the coefficients of integer powers of 1/N are given by finite rational sums of SMFs, which are of relevance to the observations in this paper.

SMFs satisfy inhomogeneous Laplace eigenvalue equations of the form (1.1) with source terms that are bilinear in non-holomorphic Eisenstein series, just as was the case with MGFs, but here the Eisenstein series in the sources have half-integer indices, $s_1, s_2 \in \mathbb{N} + 1/2$. Once again, for $s \ge 6$ not every GES solution to (1.1) with half-integer indices can be expressed as a superposition of SMFs. As before, the construction of this second family of GESs involves *L*-functions of holomorphic cusp forms [22] with modular weight 2s. SMFs are conjecturally equal to special linear combinations of GESs, where the *L*-values cancel as was the case for MGFs.

We will demonstrate that one of the main results of our paper establishes that both MGFs and SMFs emerge as specific instances of a broader class of non-holomorphic modular-invariant functions. These functions can be systematically constructed via a particular theta lift applied to a family of local modular Maass functions. This construction automatically leads to rational linear combinations of GES solutions to (1.1), where, as previously noted, the contributions from holomorphic cusp forms are entirely cancelled.

1.2 Results

As discussed in the introduction, both MGFs and SMFs can be expressed as specific rational linear combinations of GESs. We will see that these two families of non-holomorphic modular invariant functions are special cases of the theta-lifts defined by

$$\mathcal{E}_{i,j}^{w}(\tau) = \int_{(\mathbb{R}^+)^3} B_{i,j}^{w}(t) \, \Gamma'_{2,2}(\tau;t) \, \mathrm{d}^3 t \,, \tag{1.4}$$

where t represents three cartesian coordinates t_1, t_2, t_3 , while the indefinite theta-series $\Gamma'_{2,2}(\tau; t)$ is expressed as a four-dimensional lattice sum, as defined in (2.33). The lifted functions $B^w_{i,j}(t)$ are related to local Maass functions that will be defined in detail later in the paper, and the indices $i, j, w \in \mathbb{N}$.

We will show that the function $\mathcal{E}_{i,j}^{w}(\tau)$ is a specific sum of modular invariant solutions to the inhomogeneous Laplace equation (1.1). More concretely, we will prove that every $\mathcal{E}_{i,j}^{w}(\tau)$ defined in (1.4) can be expressed as a rational linear combination of GESs and non-holomorphic Eisenstein series,

$$\mathcal{E}_{i,j}^{w}(\tau) = \sum_{r=-\frac{(i+j-1)}{2}}^{\frac{(i+j-1)}{2}} b_{i,j}(w,r) \,\mathcal{E}(s; \frac{w}{2}+r, \frac{w}{2}-r; \tau) + d_{i,j}(w) \,E(w; \tau) \,, \tag{1.5}$$

where the eigenvalue s = 3i + j + 1 and the explicit expression of the general coefficients $b_{i,j}(w, r)$ is given by (3.53). All MGFs are obtained by considering $\mathcal{E}_{i,j}^w(\tau)$ when s = 3i + j + 1 and w have the same parity with $1 \leq s \leq w - 2$ and $w \geq 3$. Conversely, SMFs are generated by $\mathcal{E}_{i,j}^w(\tau)$ when s and w have opposite parity with $s \geq w + 1$ and $w \geq 3$.¹

We then analyse in more detail the properties of an individual GES, $\mathcal{E}(s; s_1, s_2; \tau)$ of the type that appears in the rational linear combination (1.5). Given the parity arguments mentioned above, we retrieve the claim made in the introduction that GESs relevant for MGFs have integer indices $s_1, s_2 \in \mathbb{N}$ with $s_1, s_2 \geq 2$, while for SMFs they have half-integral indices $s_1, s_2 \in \mathbb{N} + 1/2$.

In [9, 10] it was argued that the modular invariant solution to (1.1) with $s_1, s_2 \in \mathbb{N}$ and $s_1, s_2 \geq 2$ and eigenvalue s with the same parity as $s_1 + s_2$ must take the following form,

$$\mathcal{E}(s;s_1,s_2;\tau) = \mathcal{E}_{\mathbf{p}}(s;s_1,s_2;\tau) + \sum_{\Delta \in \mathcal{S}_{2s}} \lambda_{\Delta}(s;s_1,s_2) H_{\Delta}(\tau), \qquad (1.6)$$

where \mathcal{E}_{p} is a specific particular solution to (1.1) constructed explicitly in [9, 10]. Our main interest here is the second term, which is a solution to the homogeneous equation. Here $H_{\Delta}(\tau)$ denotes a particular Eichler integral (4.14) of the Hecke-normalised cusp form $\Delta \in \mathcal{S}_{2s}$ with \mathcal{S}_{2s} the vector space of holomorphic cusp forms of weight 2s. The coefficients $\lambda_{\Delta}(s; s_1, s_2)$ were determined in a multitude of cases in [9, 10], demonstrating that these contain the completed *L*-functions, $\Lambda(\Delta, t)$, of the corresponding holomorphic cusp form.

For the case $s_1, s_2 \in \mathbb{N}+1/2$ and eigenvalue $s \geq s_1+s_2+1$ of opposite parity, we use the method presented in [23] to construct a particular solution to (1.1) and show that whenever dim $S_{2s} \neq 0$ this does not correspond to the GES solution. In such cases, a suitable multiple of the solution of the homogeneous equation must be added to the particular solution to ensure that the complete expression is the modular invariant solution $\mathcal{E}(s; s_1, s_2; \tau)$. We show that again in this case, the GES solution to (1.1) with $s_1, s_2 \in \mathbb{N}+1/2$ and eigenvalue $s \geq s_1+s_2+1$ of opposite parity must take the same form (1.6). Using the results of [22] concerning certain convolutions of divisor sigma functions, we find an analytic expression for the coefficients of the solution of the homogeneous equation, which is given as

$$\frac{\lambda_{\Delta}(s;s_{1},s_{2}) =}{\Gamma(s_{1})^{\frac{s+s_{1}-s_{2}+2}{2}} 2^{5-4s} \Gamma(2s-1) \Gamma\left(\frac{s_{1}+s_{2}-s}{2}\right)}{\Gamma(s_{1}) \Gamma(s_{2}) \Gamma\left(\frac{s_{1}+s_{2}+s}{2}\right) \Gamma\left(\frac{s+s_{1}-s_{2}+1}{2}\right) \Gamma\left(\frac{s-s_{1}+s_{2}+1}{2}\right)} \frac{\Lambda(\Delta,s+s_{1}-s_{2}) \Lambda(\Delta,s+s_{1}+s_{2}-1)}{\langle \Delta,\Delta \rangle},$$
(1.7)

where $\langle \Delta, \Delta \rangle$ denotes the Petersson norm of the Hecke eigenform $\Delta \in \mathcal{S}_{2s}$.

Given the analytic dependence of the coefficient of the solution of the homogeneous equation (1.7) on the indices s_1, s_2 , we have verified that if we specialise (1.7) to the case of GESs with $s_1, s_2 \in \mathbb{N}$ and $s_1, s_2 \geq 2$ and eigenvalue s with the same parity as $s_1 + s_2$ we reproduce all the particular cases presented in $[9, 10]^2$. This leads to the following conjecture:

Conjecture 1.

The generalised Eisenstein series with $s_1, s_2 \in \mathbb{N}$ and $s_1, s_2 \geq 2$ and eigenvalue s with the same parity as $s_1 + s_2$ must take the form (1.6), where the particular solution has been constructed in [9, 10] and the coefficient of the solution of the homogeneous equation is given by (1.7).

This result is quite striking, since the particular solutions constructed in [9, 10] and [23] for the two different cases of GESs have in principle nothing to do with one another. Yet in both cases, the "modular

¹It is worth noting that many of our conclusions do apply to a wider range of weights w, for example SMFs with s < w. However, we will focus on modular functions with parameters satisfying these specific ranges since they are most relevant for superstring amplitudes and the integrated correlator.

²In these references, the coefficient of the solution of the homogeneous equation $\lambda_{\Delta}(s; s_1, s_2)$ was denoted by $a^+_{\Delta_{2s}, s_1, s_2}$.

form" content of the particular solution (which we stress is not modular invariant on its own) captures everything *but* the very same Eichler integral of holomorphic cusp forms.

Having clarified the form of the solution given in (1.6) for the families of GESs relevant to both MGFs and SMFs, we will revisit the general rational linear combination of GESs (1.5), which is produced by the theta lift (1.4). The explicit linear combination (1.5) together with the solution for the coefficients of the homogeneous equation (1.7) lead to a further conjecture:

Conjecture 2.

The modular invariant function $\mathcal{E}_{i,j}^w(\tau)$ defined in (1.4) is a rational linear combination of generalised Eisenstein series that admits a four-dimensional lattice sum representation, which receives no contributions from L-values of holomorphic cusp forms.

For MGFs that are given by rational linear combinations of GESs with $s_1, s_2 \in \mathbb{N}$ and $s_1, s_2 \geq 2$ and eigenvalue s with the same parity as $w = s_1 + s_2$, this is indeed a conjecture since we do not prove (1.7) for the coefficient of the solution of the homogeneous equation. However, for SMFs this conjecture can be proven since these are expressed in terms of GESs with $s_1, s_2 \in \mathbb{N} + 1/2$ and eigenvalue $s \geq s_1 + s_2 + 1$ of opposite parity for which the proof of (1.7) follows from [22].

In either case, assuming the general validity of (1.7), we will show that the *L*-function contribution to $\mathcal{E}_{i,j}^w(\tau)$ coming from the holomorphic cusp form $\Delta \in \mathcal{S}_{2s}$ with s = 3i + j + 1 is proportional to

$$\mathcal{K}_{i,j}(\Delta) = \int_0^{i\infty} \Delta(\tau) \, [\tau^2 (1-\tau)^2]^i (\tau^2 - \tau + 1)^j \mathrm{d}\tau \,. \tag{1.8}$$

The polynomials in τ appearing as part of the integrand in the above equation play a fundamental role in the construction of the local Maass functions, $B_{i,j}^w(t)$, whose theta lift (1.4) we will analyse. We will show that $\mathcal{K}_{i,j}(\Delta) = 0$ for all $i, j \in \mathbb{N}$ and for every holomorphic cusp form $\Delta \in S_{2s}$ with s = 3i + j + 1, thus providing strong evidences that the modular functions $\mathcal{E}_{i,j}^w(\tau)$ as defined in (1.4), or equivalently the particular rational linear combinations of GESs as in (1.5), are indeed free of holomorphic cusp forms.

Remarkably, precisely these particular linear combinations of GESs appear to be the only relevant ones for the study of important physical observables in superstring theory and its holographic dual CFT. This is true for MGFs for perturbative genus-one contributions to superstring amplitudes, as well as for SMFs arising in the low-energy expansion of superstring amplitudes and the large-N expansion of integrated correlators in $\mathcal{N} = 4$ SYM.

1.3 Outline

The paper is organised as follows. In section 2 we introduce the main characters of our story and define an infinite class of modular invariant non-holomorphic functions, constructed as theta lifts of certain modular local Maass forms. The defining property of these Maass forms is that they are constructed via a raising procedure from particular negative modular weight holomorphic polynomials, known as modular local polynomials. In appendix A we present some basic properties of Maass functions and non-holomorphic Eisenstein series, while in appendix B we discuss more technical details regarding modular local polynomials. We show in section 3 that the theta lift thus defined satisfies an inhomogeneous Laplace eigenvalue equation with source terms that are bilinear in non-holomorphic Eisenstein series. Hence, we derive a general expression for the theta lift as a rational linear combination of generalised Eisenstein series in section 4. In particular, we revisit two methods for constructing a particular solution to the differential equation (1.1) and show how to determine the homogeneous part so as to obtain a modular invariant solution. The solution of the homogeneous equation takes the form of an Eichler integral of a special holomorphic cusp form multiplied by the product of two of its completed L-values that we determine analytically. Some fundamental properties

of holomorphic cusp forms and their *L*-values are reviewed in appendix C. We conclude section 4 by showing that the sum of the holomorphic cusp form solutions to the homogeneous equation that contribute to $\mathcal{E}_{i,j}^w(\tau)$, i.e. to the sum of generalised Eisenstein series obtained from the theta lifts, cancels out completely. Finally, in section 5 we present some interesting open problems that we believe our analysis will help in solving.

2 Theta lift of local Maass functions

In this section we will define a particular example of a non-holomorphic indefinite theta-series that is a four-dimensional lattice sum, which will enter the subsequent analysis. This theta-series is related to the genus-one partition function for a closed string moving on a target space compactified on T^2 . We will use this theta-series to define a class of modular invariant functions via theta lift of certain local Maass functions. This theta lift provides an elegant and unified approach to both MGFs and SMFs and clarifies some intriguing properties shared by these two families of modular functions.

2.1 Theta lift: definitions and properties

In the study of superstring scattering amplitudes, being that MGFs or SMFs, we discover beautiful families of non-holomorphic modular invariant functions. The simplest examples of MGFs and SMFs are the nonholomorphic Eisenstein series, $E(s; \tau)$, with $s \in \mathbb{N}^+$ and $s \in \mathbb{N} + \frac{1}{2}$, respectively. Some properties of these modular functions, which enter into the source terms in (1.1), are reviewed in appendix A.

However, the space of modular invariant functions needed to discuss higher loop MGFs and higher order coefficients in the low-energy effective action of the scattering amplitudes in type IIB becomes significantly more complicated and goes beyond the standard world of non-holomorphic Eisenstein series or Maass functions. Importantly, both MGFs and the SMFs considered here are closely related, albeit for very different physical reasons, to a particular non-holomorphic modular function of three complex variables that can be expressed in terms of an indefinite theta-series.

The particular indefinite theta series of relevance arises quite naturally in string theory when considering a closed-string partition function for a genus-one world-sheet with modular parameter τ , embedded in a T^2 -compactified target space. The moduli space of the target space is the symmetric space:

$$[SO(2;\mathbb{R}) \times SO(2;\mathbb{R})] \setminus SO(2,2;\mathbb{R}) / SO(2,2;\mathbb{Z}), \qquad (2.1)$$

which describes the product of two hyperbolic planes, with each parameterised by a complex coordinate. These are the complex structure denoted as $\rho = \rho_1 + i\rho_2$, and the Kahler structure denoted by $\nu = B + iV$, where V is the volume of the torus and B is the value of an antisymmetric tensor on the torus.

We now define a lattice sum $\Gamma_{2,2} : \mathfrak{H}^3 \to \mathbb{R}$, which is a real-analytic function of the three upper-half plane variables $\tau, \rho, \nu \in \mathfrak{H} \coloneqq \{z \in \mathbb{C}, \text{Im } z > 0\},\$

$$\Gamma_{2,2}(\tau;\rho,\nu) \coloneqq \sum_{\underline{m},\underline{n}\in\mathbb{Z}^2} V \exp\left(-\frac{\pi}{\tau_2}(\underline{m}+\underline{n}\tau)G(\rho,V)(\underline{m}+\underline{n}\bar{\tau}) + 2\pi i B \det M\right),\tag{2.2}$$

where we have introduced the two 2×2 matrices

$$G(\rho, V) \coloneqq \frac{V}{\rho_2} \begin{pmatrix} 1 & \rho_1 \\ \rho_1 & |\rho|^2 \end{pmatrix}, \qquad M \coloneqq \begin{pmatrix} m_1 & n_1 \\ m_2 & n_2 \end{pmatrix} = (\underline{m}\,\underline{n}) .$$

$$(2.3)$$

with $m_1, m_2, n_1, n_2 \in \mathbb{Z}$. This function has the important symmetry properties,

$$\Gamma_{2,2}(\tau;\rho,\nu) = \Gamma_{2,2}(\rho;\tau,\nu), \qquad (2.4)$$

$$\Gamma_{2,2}(-\frac{1}{\tau};\rho,\nu) = \Gamma_{2,2}(\tau;\rho,\nu), \qquad \Gamma_{2,2}(\tau+1;\rho,\nu) = \Gamma_{2,2}(\tau;\rho,\nu), \qquad (2.5)$$

$$\Gamma_{2,2}(\tau;\rho,-\frac{1}{\nu}) = \Gamma_{2,2}(\tau;\rho,\nu), \qquad \Gamma_{2,2}(\tau;\rho,\nu+1) = \Gamma_{2,2}(\tau;\rho,\nu).$$
(2.6)

The first identity can be derived from (2.2) by exchanging the summation variables $n_1 \leftrightarrow m_2$, while the S-transformation for the ν variable $\nu \rightarrow -1/\nu$ can be derived upon performing a quadruple Poisson resummation over m_1, m_2, n_1, n_2 .

The Laplace operator on this symmetric space reduces to

$$\Delta_{SO(2,2)} = \Delta_{\rho} + \Delta_{\nu} = \rho_2^2 (\partial_{\rho_1}^2 + \partial_{\rho_2}^2) + V^2 (\partial_B^2 + \partial_V^2), \qquad (2.7)$$

and the lattice-sum $\Gamma_{2,2}$ satisfies

$$\Delta_{SO(2,2)}\Gamma_{2,2}(\tau;\rho,\nu) = 2\Delta_{\tau}\Gamma_{2,2}(\tau;\rho,\nu) = 2\tau_2^2(\partial_{\tau_1}^2 + \partial_{\tau_2}^2)\Gamma_{2,2}(\tau;\rho,\nu).$$
(2.8)

We note also that

$$\Delta_{\tau}\Gamma_{2,2}(\tau;\rho,\nu) = \Delta_{\rho}\Gamma_{2,2}(\tau;\rho,\nu) = \Delta_{\nu}\Gamma_{2,2}(\tau;\rho,\nu).$$
(2.9)

In string theory, one usually integrates over τ , the modular parameter of the genus-one worldsheet, for example, in obtaining a genus-one correction to a closed-string scattering amplitude for external states moving on a toroidally compactified target space. However, in the present discussion this is not the case, instead of integrating over τ we are interested in considering a particular theta lift that involves integration over ρ and $V = \text{Im } \nu$ after having set B = 0 in (2.2). More explicitly, we will consider

$$\mathcal{E}_{i,j}^{w}(\tau) = \int_{\mathbb{R}^{+} \times \mathcal{F}_{\rho}} V^{w} A_{i,j}(\rho) \Gamma_{2,2}(\tau;\rho,iV) \, 2 \frac{\mathrm{d}V}{V^{2}} \frac{\mathrm{d}^{2}\rho}{\rho_{2}^{2}} \,, \tag{2.10}$$

where, as will appear natural later, the domain of integration \mathcal{F}_{ρ} is the fundamental domain with respect to the congruence subgroup³ $\Gamma_0(2)$, i.e. $\mathcal{F}_{\rho} = \Gamma_0(2) \setminus \mathfrak{H}$. The parameter w is a non-negative integer, $w \in \mathbb{N}$. We will shortly provide some string theory arguments that justify the importance of the integral representation (2.10) in the discussion of MGFs as well as in the context of higher-derivative corrections to the low-energy expansion of type IIB string theory.

In the rest of this paper we consider the theta lift (2.10) where the lifted elements, $A_{i,j}(\rho)$, belong to a special class of Maass functions. The construction of the relevant family of functions $A_{i,j}(\rho)$ was given in unpublished notes by Don Zagier [24] that are reviewed and slightly expanded in section 5.4 of [25], and will be summarised later in section 2.2. The functions $A_{i,j}(\rho)$ are local Maass functions that transform as modular functions under $\Gamma_0(2)$ apart from on an 'exceptional set' of points forming the boundaries of \mathcal{F}_{ρ} (hence the denomination 'local'). These functions are constructed from special holomorphic polynomials with negative modular weight via a Maass raising operation so as to become non-holomorphic modular invariant functions. As a result $A_{i,j}(\rho)$ is a Laurent series in ρ_2 with coefficients that are polynomial in $(\rho_1^2 - \rho_1)$ and satisfies the Laplace equation

$$(\Delta_{\rho} - s(s-1))A_{i,j}(\rho) = 0, \qquad (2.11)$$

with s = 3i + j + 1.

³The congruence subgroup $\Gamma_0(2)$ is defined as $\Gamma_0(2) \coloneqq \{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}) \mid c \equiv 0 \mod 2 \}.$

Before moving on to discuss the construction of the functions $A_{i,j}(\rho)$ and the properties of their theta lifts (2.10) in more detail in section 2.2 we present a different parametrisation for the indefinite theta series $\Gamma_{2,2}$, which is related to the string-theory origins of MGFs [1] and SMFs [11, 12]. One property we immediately note is that (assuming convergence) the result of performing a quadruple Poisson resummation of the integers in the theta-series (2.2) leads to $\Gamma_{2,2}(\tau; \rho, iV) = \Gamma_{2,2}(\tau; \rho, iV^{-1})$ so that by changing variables $V \to 1/V$ in (2.10) we find that $\mathcal{E}_{i,j}^{w}(\tau)$ satisfies the functional relation

$$\mathcal{E}_{i,j}^{w}(\tau) = \mathcal{E}_{i,j}^{2-w}(\tau), \qquad (2.12)$$

which is very reminiscent of the functional equation (A.5) for the non-holomorphic Eisenstein series. We now discuss an alternative parameterisation of $\Gamma_{2,2}$, which will prove to be very useful.

An alternative parameterisation of $\Gamma_{2,2}$

This reparameterization maps the real integration parameters ρ_1 , ρ_2 and V in (2.10) into the three Schwinger parameters, t_1, t_2, t_3 of a two-loop vacuum Feynman diagram defined in the range $0 \le t_i \le \infty$ and given by the following equations⁴

$$\rho_1(t) = \frac{t_1}{t_1 + t_2}, \qquad \rho_2(t) = \frac{V(t)}{t_1 + t_2}, \qquad V(t) = \sqrt{t_1 t_2 + t_1 t_3 + t_2 t_3}, \tag{2.13}$$

where the measure in terms of the new variables is given by

$$2dVV^2 \frac{d^2\rho}{\rho_2^2} = dt_1 dt_2 dt_3.$$
(2.14)

The domain of integration $0 \le t_i \le \infty$ translates into the domain in (2.10) that is given by the real positive axis $0 \le V \le \infty$ and

$$0 \le \rho_1 \le 1$$
, $(\rho_1 - \frac{1}{2})^2 + \rho_2^2 \ge \frac{1}{4}$, (2.15)

which is precisely the fundamental domain of $\Gamma_0(2)$ illustrated by the grey domain in figure 1 (which is reproduced from [11]).

We will now express (2.2) in these new variables. With a slight abuse of notation, we denote $\Gamma_{2,2}(\tau;t) := \Gamma_{2,2}(\tau;\rho(t),iV(t))$, and after a simple redefinition of summation variables we arrive at

$$\Gamma_{2,2}(\tau;t) \coloneqq \Gamma_{2,2}(\tau;\rho(t),iV(t)) = \sqrt{t_1 t_2 + t_1 t_3 + t_2 t_3} \sum_{\substack{p_1,p_2,p_3 \in \Lambda\\p_1+p_2+p_3=0}} \exp\left(-\frac{\pi}{\tau_2} \sum_{j=1}^3 t_j |p_j|^2\right),$$
(2.16)

where the sum is over three complex lattice momenta $p_i = m_i + n_i \tau \in \Lambda = \mathbb{Z} + \tau \mathbb{Z}$, with i = 1, 2, 3, subject to the conservation condition $p_1 + p_2 + p_3 = 0$.

The integrand for the theta lift (2.10) can be re-expressed in terms of the t_i variables in the form

$$\mathcal{E}_{i,j}^{w}(\tau) = \int_{(\mathbb{R}^{+})^{3}} V(t)^{w-4} A_{i,j}(t) \Gamma_{2,2}(\tau;t) \,\mathrm{d}^{3}t \,.$$
(2.17)

The functions $A_{i,j}(t)$ must be symmetric functions of t_1 , t_2 , t_3 and the integrand in (2.17) is invariant under the symmetric group \mathfrak{S}_3 , a property that is not immediately apparent in (2.10). The manifest invariance of

⁴The notation f(t) denotes a function of the three t_i variables with i = 1, 2, 3.



Figure 1: In the ρ -plane, the domain $t_i > 0$ is mapped to the region shaded in grey, isomorphic to a fundamental domain of $\Gamma_0(2)$. Here the labels (ijk) in each sub-domain denote the ordering $t_i < t_j < t_k$.

the expression (2.17) under permutations of t_1 , t_2 and t_3 is exhibited by (2.10) in terms of the ρ variable, which is invariant under the six elements of transformation:

$$\rho \to \rho, \qquad \rho \to 1 - \rho^{-1}, \qquad \rho \to (1 - \rho)^{-1}, \qquad \rho \to 1 - \bar{\rho}, \qquad \rho \to \bar{\rho}^{-1}, \qquad \rho \to (1 - \bar{\rho}^{-1})^{-1}.$$
 (2.18)

The first three of these correspond to transpositions of pairs of t_i whereas the last three correspond to three-cycles with t_1 , t_2 , t_3 in different orders.

The Laplace identities (2.9) imply the relation

$$\Delta_{\tau}\Gamma_{2,2}(\tau;t) = \Delta_t\Gamma_{2,2}(\tau;t), \qquad (2.19)$$

where the laplacian in the t_i variables is given by

$$\Delta_t[F(t)] = -2\sum_{i=1}^3 \partial_i[t_i F(t)] + \sum_{i,j=1}^3 \partial_i \partial_j \{ [t_i t_j + (2\delta_{ij} - 1)V^2(t)]F(t) \}.$$
(2.20)

Importantly, we see that $\Delta_t[V(t)^{\alpha}F(t)] = V(t)^{\alpha}\Delta_t[F(t)]$ for all $\alpha \in \mathbb{R}$, with $V(t) = \sqrt{t_1t_2 + t_1t_3 + t_2t_3}$.

We will now discuss the particular functions $A_{i,j}(\rho)$ that are invariant under the group of symmetries (2.18), which will play a central role in the theta lift (2.10).

2.2 Modular local polynomials and local Maass functions

We present here the systematic construction of a basis of $\Gamma_0(2)$ -invariant functions $A_{i,j}(\rho)$, given by Don Zagier in unpublished notes that are expanded on in section 5.4 of [25]. The function $A_{1,0}(\rho)$ originally arose in the construction of the modular invariant coefficient of the higher-derivative interaction d^6R^4 in the four-graviton amplitude of type IIB superstring theory [12, 26]. This coefficient function is $\mathcal{E}(4; \frac{3}{2}, \frac{3}{2}; \tau)$ in the notation of (1.1)⁵.

The functions $A_{i,j}(\rho)$ here considered are particular examples of 'modular local Maass forms', constructed from special modular forms known as modular local polynomials [27, 28].

Definition [27]: For an SL(2, \mathbb{Z})-invariant nowhere dense set E, a function $\mathcal{P} : \mathfrak{H} \to \mathbb{C}$ is called a weight-k modular local polynomial with exceptional set E if (with k even and non-positive) \mathcal{P} is such that

(i) For every $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z})$ we have $\mathcal{P}(\rho)|_k \gamma = \mathcal{P}(\rho)$, where the usual $|_k \gamma$ action is defined as

$$\mathcal{P}(\rho)\big|_k \gamma = (c\rho + d)^{-k} \mathcal{P}\left(\frac{a\rho + b}{c\rho + d}\right); \qquad (2.21)$$

(ii) On each connected component $\mathcal{C} \subset \mathfrak{H} \setminus E$ we have $\mathcal{P}(\rho) = P_{\mathcal{C}}(\rho)$ where $P_{\mathcal{C}}$ is a polynomial in ρ .

Furthermore, as explained in [27] we must impose some regularity condition so that the limiting values exist when $\rho \to \rho^* \in E$. Importantly, while the modular local polynomials are finite on the exceptional set E their derivatives are discontinuous.

In [28] modular local polynomials were introduced for which the exceptional sets are

$$E_D \coloneqq \bigcup_{Q \in \mathcal{Q}_D} S_Q \,, \tag{2.22}$$

where Q_D denotes the set of binary quadratic forms with discriminant D, i.e. $Q = [a, b, c] \in Q_D$ is such that $Q(X, Y) = [a, b, c](X, Y) \coloneqq aX^2 + bXY + cY^2$ with $a, b, c \in \mathbb{Z}$ and discriminant $D = b^2 - 4ac$. The sets S_Q are defined by

$$S_Q \coloneqq \left\{ \rho \in \mathfrak{H} \, \middle| \, a |\rho|^2 + b \operatorname{Re}(\rho) + c = 0 \right\} \,. \tag{2.23}$$

For the case of interest the discriminant we want to consider is D = 1. As we show in appendix B that the connected component $\mathcal{C}_{\rho^*} \subset \mathfrak{H} \setminus E_1$, which contains $\rho^* \coloneqq e^{\frac{\pi i}{3}}$, i.e. $\rho \in \mathcal{C}_{\rho^*}$, coincides with a fundamental domain of $\Gamma_0(2)$, i.e.

$$\mathcal{C}_{\rho^*} = \Gamma_0(2) \backslash \mathfrak{H} \coloneqq \left\{ \rho \in \mathfrak{H} \left| 0 < \operatorname{Re}(\rho) < 1, \left| \rho - \frac{1}{2} \right|^2 > \frac{1}{4} \right\} \right.$$
(2.24)

We can then think of these modular local polynomials as the modular analogues of the periodic seesaw function. Once we define the polynomial on one of the connected components, say e.g. C_{ρ^*} , we extend it to a function on the whole upper-half plane \mathfrak{H} by simply patching together all of its images under $\Gamma_0(2)$.

The function $A_{i,j}(\rho)$ in the integrand of the theta lift (2.10) is modular invariant, so it has zero holomorphic and anti-holomorphic weight with respect to all $\gamma \in \Gamma_0(2)$. It can be constructed by successive applications of the Maass raising operator D_k to a modular polynomial $\mathcal{P}(\rho)$ of even negative weight, where $D_k \coloneqq \partial_{\rho} + k/(\rho - \bar{\rho})$ maps a modular form of weight (k, 0) to a modular form of weight (k + 2, 0). More explicitly, starting with a modular form of weight -2n we can raise it to a real-analytic modular function by acting with the iterated Maass derivative operator

$$D_{-2n}^{(n)} \coloneqq \frac{(-2i)^n n!}{(2n)!} D_{-2} \circ D_{-4} \circ \dots \circ D_{-2n+2} \circ D_{-2n}, \qquad (2.25)$$

⁵In [12] the function $A_{1,0}(\rho)$ was called $A(\tau)$ and the coefficient function was called $\mathcal{E}_{\frac{3}{2},\frac{3}{2}}(\Omega,\bar{\Omega})$. See also [26] for alternative early notation.

which can also be rewritten as

$$D_{-2n}^{(n)} = \frac{(-2i)^n n!}{(2n)!} \sum_{m=0}^n \binom{n}{m} \frac{(-n-m)_m}{(\rho-\bar{\rho})^m} \frac{\partial^{n-m}}{\partial \rho^{n-m}},$$
(2.26)

where $(x)_m := x(x+1)\dots(x+m-1)$ is the Pochhammer symbol. We note that the Maass operator D_k satisfies the property $\Delta_{k+2} \cdot D_k - D_k \Delta_k = -kD_k$, where $\Delta_k \coloneqq 4D_{k-2}\rho_2^2 \partial_{\bar{\rho}}$ is the Laplacian acting on weight k modular forms. As a consequence, given that a modular local polynomial, $\mathcal{P}(\rho)$, of weight k = -2n is a holomorphic function and therefore satisfies $\Delta_{-2n} \mathcal{P}(\rho) = 0$. It follows that $A(\rho) \coloneqq D_{-2n}^{(n)} \mathcal{P}(\rho)$ is a modular function satisfying the Laplace eigenvalue equation

$$[\Delta_{\rho} - n(n+1)] A(\rho) = 0, \qquad (2.27)$$

where $\Delta_{\rho} = \Delta_0 = \rho_2^2 (\partial_{\rho_1}^2 + \partial_{\rho_2}^2)$. As proved in [24, 25], the subspace of the modular local polynomials with exceptional set E_1 and weight k that have been lifted as described above to be modular invariant functions with respect to $\Gamma_0(2)$ as well as being invariant with respect to the group of automorphism of \mathfrak{H} generated by the involutions $\rho \to 1 - \bar{\rho}$ and $\rho \to 1/\bar{\rho}$, is spanned by the basis:

$$A_{i,j}(\rho) \coloneqq D_{-2n}^{(n)}[u^i v^j], \qquad (2.28)$$

where $i, j \in \mathbb{N}$ such that k = -2n = -2(3i + j), and

$$u \coloneqq \rho^2 (1-\rho)^2, \qquad v \coloneqq \rho^2 - \rho + 1.$$
 (2.29)

From (2.26) and (2.27), it is easy to show that the $A_{i,j}(\rho)$ are Laurent polynomials in ρ_2 with coefficients that are polynomial in ρ_1 , satisfying the Laplace equation

$$[\Delta_{\rho} - s(s-1)] A_{i,j}(\rho) = 0, \qquad (2.30)$$

for all $\rho \in \Gamma_0(2) \setminus \mathfrak{H}$, where s = 3i + j + 1. For example, we have

$$\begin{aligned} A_{0,1}(\rho) &= \rho_2 + \frac{\rho_1(\rho_1 - 1) + 1}{\rho_2} ,\\ A_{1,0}(\rho) &= \frac{\rho_2}{5} + \frac{\left[1 + 6\rho_1(\rho_1 - 1)\right]}{5\rho_2} + \frac{\rho_1^2(\rho_1 - 1)^2}{\rho_2^3} , \end{aligned} \tag{2.31} \\ A_{1,1}(\rho) &= \frac{\rho_2^2}{7} + \frac{3\left[4 + 15\rho_1(\rho_1 - 1)\right]}{35} + \frac{3\rho_1(\rho_1 - 1)\left[5\rho_1(\rho_1 - 1) + 3\right] + 1}{7\rho_2^2} + \frac{\rho_1^2(\rho_1 - 1)^2\left[\rho_1(\rho_1 - 1) + 1\right]}{\rho_2^4} . \end{aligned}$$

The particular theta lifts we want to consider are $\Gamma_{2,2}$ lifts of the modular-invariant local Maass forms $A_{i,j}(\rho)$, which are defined by (2.10). As will be clarified in the next section, the integral in (2.10) does not quite converge if one uses the definition (2.2) for $\Gamma_{2,2}$ as a complete lattice sum over \mathbb{Z}^4 . To obtain a convergent expression it is necessary to exclude the points $(m_1, n_1) = 0$, $(m_2, n_2) = 0$ and $(m_1 + m_2, n_1 + n_2) = 0$ from the lattice sum. In terms of the t_i variables the expression for the theta lift (2.17) becomes

$$\mathcal{E}_{i,j}^{w}(\tau) \coloneqq \int_{(\mathbb{R}^+)^3} [V(t)]^{w-4} A_{i,j}(\rho(t)) \, \Gamma'_{2,2}(\tau;t) \, \mathrm{d}^3 t \,, \tag{2.32}$$

where using (2.16) we have defined

$$\Gamma_{2,2}'(\tau;t) \coloneqq \sqrt{t_1 t_2 + t_1 t_3 + t_2 t_3} \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1 + p_2 + p_3 = 0}} \exp\left(-\frac{\pi}{\tau_2} \sum_{j=1}^3 t_j |p_j|^2\right),\tag{2.33}$$

and the sum runs over three lattice momenta $p_i = m_i + n_i \tau \in \Lambda' = \mathbb{Z} + \tau \mathbb{Z} \setminus \{0\}$, with i = 1, 2, 3, again subject to the conservation condition $p_1 + p_2 + p_3 = 0$. The symbol Λ' indicates that we exclude the point p = 0 from the lattice $p \in \Lambda = \mathbb{Z} + \tau \mathbb{Z}$, which is a requirement for the convergence of the integral (2.32). The objects of interest can then be rewritten in terms of the lattice sum integral

$$\mathcal{E}_{i,j}^{w}(\tau) = \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1 + p_2 + p_3 = 0}} \int_{(\mathbb{R}^+)^3} B_{i,j}^{w}(t) \exp\left(-\frac{\pi}{\tau_2} \sum_{j=1}^3 t_j |p_j|^2\right) \mathrm{d}^3 t \,, \tag{2.34}$$

where we have defined the auxiliary functions

$$B_{i,j}^{w}(t) \coloneqq [V(t)]^{w-3} A_{i,j}(\rho(t)) \,. \tag{2.35}$$

For example when expressed in terms of the t_i variables, the $A_{i,i}(\rho)$ given in (2.31) become

$$A_{0,1}(\rho(t)) = \sigma_2^{-\frac{1}{2}} \sigma_1,$$

$$A_{1,0}(\rho(t)) = \sigma_2^{-\frac{3}{2}} \left(\frac{1}{5}\sigma_1\sigma_2 - \sigma_3\right),$$

$$A_{1,1}(\rho(t)) = \sigma_2^{-2} \left(\frac{1}{7}\sigma_1^2\sigma_2 + \frac{2}{35}\sigma_2^2 - \sigma_1\sigma_3\right),$$
(2.36)

where we have expressed the results in terms of the basis for symmetric polynomials in three variables σ_1 , σ_2 , and σ_3 , defined as

$$\sigma_1 \coloneqq t_1 + t_2 + t_3, \qquad \sigma_2 \coloneqq V(t)^2 = t_1 t_2 + t_1 t_3 + t_2 t_3, \qquad \sigma_3 \coloneqq t_1 t_2 t_3. \tag{2.37}$$

It is important to realise that whereas the reflection identity (2.12) can be obtained by performing a quadruple Poisson resummation this procedure cannot be carried out for the restricted lattice sum (2.33). However, after correcting for the absence of the $(m_1, n_1) = (m_2, n_2) = (m_1 + m_2, n_1 + n_2) = (0, 0)$ terms in the sums the identity (2.12) still holds modulo the addition of terms proportional to non-holomorphic Eisenstein series, namely we find

$$\mathcal{E}_{i,j}^{w}(\tau) = \mathcal{E}_{i,j}^{2-w}(\tau) + \left(\text{non-holomorphic Eisenstein series}\right).$$
(2.38)

We can now better clarify why the theta lift considered here is a suitable framework for discussing both MGFs and SMFs. Firstly, it follows directly from (2.34) that $\mathcal{E}_{i,j}^w(\tau)$ is closely related to MGFs (1.3) introduced in [1]. If in equation (2.34) we replace the integrand $B_{i,j}^w(t)$ by the simple monomial $t_1^{a-1}t_2^{b-1}t_3^{c-1}$ with $a, b, c \in \mathbb{N}$ we deduce

$$\sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1+p_2+p_3=0}} \int_{(\mathbb{R}^+)^3} t_1^{a-1} t_2^{b-1} t_3^{c-1} \exp\left(-\frac{\pi}{\tau_2} \sum_{j=1}^3 t_j |p_j|^2\right) \mathrm{d}^3 t = \Gamma(a) \Gamma(b) \Gamma(c) C_{a,b,c}(\tau) \,.$$
(2.39)

Here the parameters t_i correspond to the Schwinger parameters for the two-loop graph associated with the MGF, $C_{a,b,c}(\tau)$.

Similarly in the context of SMFs the integral representation (2.34) originates by considering the fourgraviton two-loop scattering amplitude in eleven-dimensional supergravity compactified on a two-torus [12], where the t_i variables are related to the Schwinger parameters for the two-loop Feynman diagrams. While in [12] a prominent role was played by (2.35) specialised to the case of the local Maass function $A_{1,0}(\rho)$, in [21] it was shown that at large-N the expansion coefficient of a certain integrated correlator in $\mathcal{N} = 4$ SYM can be represented as a rational linear combination of theta lifts (2.34) involving more general $A_{i,j}(\rho)$. The precise connection between the theta lift (2.34) and both MGFs and SMFs will be derived by analysing the Laplace equation satisfied by $\mathcal{E}_{i,j}^w(\tau)$.

In the next section we will study the real-analytic modular invariant functions $\mathcal{E}_{i,j}^w(\tau)$ of the variable $\tau \in \mathfrak{H}$ and show that they satisfy an inhomogeneous Laplace eigenvalue equation with sources given by bilinears in non-holomorphic Eisenstein series. By varying the integers w, i, j, we will show that $\mathcal{E}_{i,j}^w(\tau)$ corresponds to either MGFs that appear in genus-one superstring amplitudes or to the SMFs that appear in the low energy effective action of the scattering amplitudes in type IIB superstring theory and in the context of the large-Nexpansion of integrated correlators in $\mathcal{N} = 4$ SYM.

3 Inhomogeneous Laplace Equation

In this section we analyse the properties of the theta-lifted modular local Maass functions defined in (2.32) by studying their behaviour under the action of the hyperbolic Laplace operator Δ_{τ} .

From their definition (2.28), it follows that the functions $A_{i,j}(\rho)$ are Laurent polynomials in ρ_2 with coefficients that are polynomial in ρ_1 . We recall that $A_{i,j}(\rho)$ satisfies the homogeneous Laplace eigenvalues equation (2.30) with respect to the Laplace-Beltrami operator $\Delta_{\rho} = \rho_2^2(\partial_{\rho_1}^2 + \partial_{\rho_2}^2)$

$$[\Delta_{\rho} - s(s-1)] A_{i,j}(\rho) = 0, \qquad (3.1)$$

where ρ is inside the domain $\rho \in \mathcal{C}_{\rho^*} = \Gamma_0(2) \setminus \mathfrak{H}$ and s = 3i + j + 1.

When translated into t-variables, the differential equation (3.1) implies that the function $B_{i,j}^w(t)$ defined in (2.35) satisfies another homogeneous Laplace eigenvalue equation

$$[\Delta_t - s(s-1)] B^w_{i,j}(t) = 0, \qquad (3.2)$$

where again s = 3i + j + 1 and the laplacian Δ_t is defined in (2.20). Furthermore it is straightforward to show that

$$\Delta_{\tau} \exp\left(-\frac{\pi}{\tau_2} \sum_{i=1}^{3} t_i |p_i|^2\right) = \Delta_t \exp\left(-\frac{\pi}{\tau_2} \sum_{i=1}^{3} t_i |p_i|^2\right),\tag{3.3}$$

so by acting with Δ_{τ} on (2.32) and combining (3.3) and (3.2) we deduce

$$\left[\Delta_{\tau} - s(s-1)\right] \mathcal{E}_{i,j}^{w}(\tau) = \text{b.t.}, \qquad (3.4)$$

where 'b.t.' denotes the boundary terms that arise from integrating the Laplacian Δ_t by parts.

We will shortly analyse these boundary terms and show that they produce the source terms for the inhomogeneous Laplace eigenvalue equation,

$$\left[\Delta_{\tau} - s(s-1)\right] \mathcal{E}_{i,j}^{w}(\tau) = \sum_{r=-\frac{(i+j-1)}{2}}^{\frac{(i+j-1)}{2}} b_{i,j}(w,r) E\left(\frac{w}{2} + r;\tau\right) E\left(\frac{w}{2} - r;\tau\right) + \tilde{d}_{i,j}(w) E(w;\tau),$$
(3.5)

for particular values of the coefficients $b_{i,j}(w,r)$ and $d_{i,j}(w)$. Note that the source terms in this equation have total "transcendental weight" $w = s_1 + s_2$. We also note that when the weight w and the eigenvalue s = 3i + j + 1 have the same parity the source terms contain bilinears in non-holomorphic Eisenstein series with integer indices. This is relevant to the discussion of MGFs for which $s \le w - 2$ so that the indices of the non-holomorphic Eisenstein series on the right hand-side of (3.5) are always integers greater or equal than two. Conversely, when the w and s have opposite parity the source terms contain bilinears in nonholomorphic Eisenstein series with half-integer indices. This is the case that is relevant to the discussion of SMFs, in which the eigenvalues are such that $s \ge w+1$. Hence the indices of the non-holomorphic Eisenstein series in the bilinear term in (3.5) may have different signs. Nonetheless, it is always possible to use the functional equation

$$\Gamma(s)E(s;\tau) = \Gamma(1-s)E(1-s;\tau), \qquad (3.6)$$

to rewrite (3.5) so that all Eisenstein series in the source term appear with positive indices, although in that case the functional equation does not respect uniform transcendentality in weight.

It is important to notice that for all the cases we are interested in the eigenvalue $s \neq w$ hence it is always possible to use (A.2) and invert the laplacian⁶ over the single Eisenstein series appearing as source to obtain

$$\mathcal{E}_{i,j}^{w}(\tau) = \sum_{r=-\frac{(i+j-1)}{2}}^{\frac{(i+j-1)}{2}} b_{i,j}(w,r) \mathcal{E}\left(s; \frac{w}{2}+r, \frac{w}{2}-r; \tau\right) + d_{i,j}(w) E(w; \tau),$$
(3.7)

where $d_{i,j}(w) \coloneqq \tilde{d}_{i,j}(w)/[w(w-1) - s(s-1)]$. We now describe how to compute the coefficients $b_{i,j}(w,r)$ and $d_{i,j}(w)$ starting from the definition (2.32) for the theta lift of local Maass functions.

3.1 Boundary terms

Applying the laplacian Δ_{τ} to (2.32) gives

$$\Delta_{\tau} \mathcal{E}_{i,j}^{w}(\tau) = \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1 + p_2 + p_3 = 0}} \int_0^\infty [V(t)]^{w-3} A_{i,j}(\rho(t)) \Delta_t \exp\left(-\frac{\pi}{\tau_2} \sum_{i=1}^3 t_i |p_i|^2\right) \mathrm{d}^3 t \,, \tag{3.8}$$

where we have used (3.3). We integrate (3.8) by parts and note that

$$\Delta_t[V(t)^{\alpha}F(t)] = V(t)^{\alpha}\Delta_t[F(t)], \qquad (3.9)$$

for all α , hence by using the definition (2.35) and the relation (2.30) it follows that

$$\Delta_t \Big[B_{i,j}^w(t) \Big] = [V(t)]^{w-3} \Big[\Delta_\rho A_{i,j}(\rho) \Big]_{\rho=\rho(t)} = s(s-1) B_{i,j}^w(t) , \qquad (3.10)$$

with s = 3i + j + 1. From this we we see that

$$\Delta_{\tau} \mathcal{E}_{i,j}^{w}(\tau) = \sum_{\substack{p_{1}, p_{2}, p_{3} \in \Lambda' \\ p_{1}+p_{2}+p_{3}=0}} \int_{0}^{\infty} B_{i,j}^{w}(t) \Delta_{t} \Big[\exp\left(-\frac{\pi}{\tau_{2}} \sum_{i=1}^{3} t_{i} |p_{i}|^{2}\right) \Big] \mathrm{d}^{3}t$$
$$= \sum_{\substack{p_{1}, p_{2}, p_{3} \in \Lambda' \\ p_{1}+p_{2}+p_{3}=0}} \int_{0}^{\infty} \Delta_{t} \Big[B_{i,j}^{w}(t) \Big] \exp\left(-\frac{\pi}{\tau_{2}} \sum_{i=1}^{3} t_{i} |p_{i}|^{2}\right) \mathrm{d}^{3}t + \mathrm{b.t.}$$
$$= s(s-1) \mathcal{E}_{i,j}^{w}(\tau) + \mathrm{b.t.}, \qquad (3.11)$$

⁶We can check from the lattice sum representation (2.34) that for the range of eigenvalues relevant for the SMFs the coefficient of the solution of the homogeneous equation $E(s;\tau)$ vanishes. Similarly, for the case relevant for discussing MGFs we can directly evaluate (2.34) to find a vanishing coefficient for the solution of the homogeneous equation $E(s;\tau)$, apart from the case s = 1 and w odd where this solution reduces to a simple rational multiple of $\zeta(w)$ which can be computed [29]. This implies that there is no issue in inverting the Laplace operator in (3.5).

where 'b.t.' represents boundary terms.

The boundary terms in (3.11) are easily collected by integrating Δ_t by parts using the definition (2.20). We start by noting that since all the momenta $p_1, p_2, p_3 = -p_1 - p_2$ are non-vanishing we only need to worry about boundary contributions coming from $t_i \to 0$ since the integral is exponentially suppressed along any direction $t_i \to \infty$. Using the fact that the integrand is invariant under permutations of (t_1, t_2, t_3) we arrive at the expression

$$b.t. = \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1 + p_2 + p_3 = 0}} 3 \int_0^\infty \left[t_1 t_2 (\partial_3 - \partial_1 - \partial_2) B_{i,j}^w(t_1, t_2, t_3) \right]_{t_3 = 0} \exp\left(-\frac{\pi t_1 |p_1|^2}{\tau_2} - \frac{\pi t_2 |p_2|^2}{\tau_2} \right) d^2 t + \sum_{p \in \Lambda'} \frac{6\pi |p|^2}{\tau_2} \int_0^\infty \left[t_1 t_2 B_{i,j}^w(t_1, t_2, t_3) \right]_{t_3 = 0} \exp\left(-\frac{\pi |p|^2 (t_1 + t_2)}{\tau_2} \right) d^2 t ,$$
(3.12)

where we have defined $p = m + n\tau \in \Lambda'$ with $(m, n) \in \mathbb{Z}^2 \setminus \{(0, 0)\}$. We now show that with the definition of $B_{i,j}^w(t)$ given earlier that the first term produces a bilinear in Eisenstein series, while the second term produces an expression linear in Eisenstein series.

Although we do not have a closed formula for the complete boundary terms associated with a general $B_{i,j}^w(t)$, we can make some general observations. It follows from (2.28) that when $A_{i,j}(\rho)$ is rewritten in terms of t_i via (2.13) it can be expressed in terms of the basis $\sigma_1, \sigma_2, \sigma_3$ for symmetric polynomials in three variables defined in (2.37). Using these variables, $A_{i,j}(\rho)$ takes the following general form

$$A_{i,j}(\rho(t)) = \sigma_2^{-\frac{s-1}{2}} \sum_{\substack{\alpha,\beta,\gamma \ge 0\\\alpha+2\beta+3\gamma=s-1}} c(\alpha,\beta,\gamma) \,\sigma_1^{\alpha} \sigma_2^{\beta} \sigma_3^{\gamma} \,, \tag{3.13}$$

where the coefficients $c(\alpha, \beta, \gamma) \in \mathbb{Q}$ implicitly depend on i, j (and s = 3i + j + 1). We note that when s is even the overall factor is an half-integer power of σ_2 while for s odd the power becomes integer. The condition on the sum that constrains the powers to satisfy $\alpha + 2\beta + 3\gamma = s - 1$ comes from the fact that the $A_{i,j}(\rho)$ are homogeneous functions of t_i since ρ_1 and ρ_2 defined in (2.13) are. From (2.13) we also deduce that the transformation $\rho \to \overline{\rho}$ is equivalent to $t_i \to -t_i$, or equivalently

$$\sigma_1 \to -\sigma_1, \qquad \sigma_2 \to \sigma_2, \qquad \sigma_3 \to -\sigma_3, \tag{3.14}$$

hence using (2.26) and (2.28) we see that

$$A_{i,j}(\bar{\rho}) = (-1)^{i+j} A_{i,j}(\rho) = (-1)^{s-1} A_{i,j}(\rho)$$

$$\implies A_{i,j}(\rho(-t)) = A_{i,j}(\bar{\rho}(t)) = (-1)^{s-1} A_{i,j}(\rho(t)).$$
(3.15)

Using the expansion (3.13) this implies

$$c(\alpha, \beta, \gamma) = 0$$
, for $\alpha + \gamma \not\equiv (s - 1) \mod 2$. (3.16)

For example we can derive formulae for infinite families of $A_{i,j}(\rho)$ at fixed low value of *i* and arbitrary *j*. From (2.28) we can then prove by induction over *j* that

$$\sigma_2^{\frac{j}{2}} A_{0,j}(\rho(t)) = \sigma_1^{j} {}_2F_1\left(\frac{1}{2} - \frac{j}{2}, -\frac{j}{2}; \frac{1}{2} - j | \frac{3\sigma_2}{\sigma_1^2}\right), \qquad (3.17)$$

and

We note that all the hypergeometric functions ${}_{2}F_{1}(a,b;c|z)$ that arise are actually polynomials in their argument z.

We can then focus on the integrand (2.35) and expand it as

$$B_{i,j}^{w}(t_1, t_2, t_3) = \sigma_2^{\frac{w-s-2}{2}} \sum_{\substack{\alpha, \beta, \gamma \ge 0\\ \alpha+2\beta+3\gamma=s-1}} c(\alpha, \beta, \gamma) \, \sigma_1^{\alpha} \sigma_2^{\beta} \sigma_3^{\gamma} \,. \tag{3.19}$$

From this equation it is rather easy to compute the boundary contribution (3.12) where only the terms with $\gamma = 0$ and $\gamma = 1$ contribute. Therefore, for the purpose of extracting the source terms in the differential equation we can restrict our analysis of (3.19) to terms constant or linear in σ_3 . In general, these terms take the form

$$B_{i,j}^{w}(t_{1},t_{2},t_{3}) = \sigma_{2}^{\frac{w-s-2}{2}} \left[\left(\sum_{n=0}^{\lfloor \frac{i+j}{2} \rfloor} c_{i,j}^{(0)}(n) \sigma_{1}^{i+j-2n} \sigma_{2}^{i+n} \right) + \sigma_{3} \left(\sum_{n=0}^{\lfloor \frac{i+j-1}{2} \rfloor} c_{i,j}^{(1)}(n) \sigma_{1}^{i+j-2n-1} \sigma_{2}^{i+n-1} \right) + O(\sigma_{3}^{2}) \right], \quad (3.20)$$

where we compactly denoted the coefficients of (3.19) as $c_{i,j}^{(\ell)}(n) \coloneqq c(i+j-2n-\ell,i+n-\ell,\ell)$.

Although the coefficients $c_{i,j}^{(\ell)}(n)$ are not known in closed form for arbitrary i, j we have found expressions for infinite families at fixed i or at fixed j. For example for the infinite family $B_{0,j}^w(t)$ from (3.17) we find

$$c_{0,j}^{(0)}(n) = \frac{\Gamma(j+1)}{\Gamma\left(j+\frac{1}{2}\right)} \times \frac{\left(-\frac{3}{4}\right)^n \Gamma\left(j-n+\frac{1}{2}\right)}{\Gamma(n+1)\Gamma(j-2n+1)},$$
(3.21)

with $c_{0,j}^{(\ell>0)}(n) = 0$. Similarly from (3.18) we deduce that for the family $B_{1,j}^w(t)$,

$$c_{1,j}^{(0)}(n) = \frac{\Gamma(j+1)}{\Gamma(j+\frac{\tau}{2})} \times \frac{(-\frac{3}{4})^n \Gamma(j-n+\frac{5}{2})}{18\Gamma(n+1)\Gamma(j-2n+2)} [j(9-6n) + 4(n-7)n+9],$$
(3.22)

$$c_{1,j}^{(1)}(n) = -\frac{\Gamma(j+1)}{\Gamma(j+\frac{\tau}{2})} \times \frac{(-\frac{3}{4})^n \Gamma(j-n+\frac{\tau}{2})}{\Gamma(n+1)\Gamma(j-2n+1)},$$
(3.23)

with $c_{1,j}^{(\ell>1)}(n) = 0$. While for $B_{i,0}^w(t)$ we have

$$c_{i,0}^{(0)}(n) = \frac{\sqrt{\pi}\,\Gamma(i+1)^2}{3^{3i}\Gamma\left(i+\frac{1}{6}\right)\Gamma\left(i+\frac{5}{6}\right)} \times \frac{2^{2n+1}\Gamma\left(2i-n+\frac{1}{2}\right)}{\Gamma(2n+1)\Gamma(i-2n+1)\Gamma(i-n+1)}\,,\tag{3.24}$$

$$c_{i,0}^{(1)}(n) = -\frac{\sqrt{\pi}\,\Gamma(i+1)^2}{3^{3i}\Gamma\left(i+\frac{1}{6}\right)\Gamma\left(i+\frac{5}{6}\right)} \times \frac{2^{2n+2}\Gamma\left(2i-n+\frac{3}{2}\right)}{\Gamma(2n+2)\Gamma(i-2n)\Gamma(i-n+1)},\tag{3.25}$$

where now in general $c_{i,j}^{(\ell>1)}(n) \neq 0$ for $i \geq 2$. We can then focus on the boundary terms produced by a single monomial contribution to $B_{i,j}^w(t_1, t_2, t_3)$, i.e. rather than the sum over n in (3.20) we consider the individual term,

$$B_{i,j}^{w}(t_1, t_2, t_3) \to c_{i,j}^{(0)}(n) \sigma_1^{i+j-2n} \sigma_2^{\frac{w-s-2}{2}+i+n} + c_{i,j}^{(1)}(n) \sigma_1^{i+j-2n-1} \sigma_2^{\frac{w-s-2}{2}+i+n-1} \sigma_3, \qquad (3.26)$$

and substitute it into (3.12) to find

$$b.t. = \sum_{\substack{p_1, p_2, p_3 \in \Lambda' \\ p_1 + p_2 + p_3 = 0}} 3(2n - i - j)c_{i,j}^{(0)}(n) \int_0^\infty (t_1 t_2)^{\frac{w-s}{2} + i + n} (t_1 + t_2)^{i + j - 2n - 1} e^{-\frac{\pi t_1 |p_1|^2}{\tau_2} - \frac{\pi t_2 |p_2|^2}{\tau_2}} d^2 t$$

$$+ \sum_{\substack{p_1, p_2, p_3 \in \Lambda' \\ p_1 + p_2 + p_3 = 0}} 3c_{i,j}^{(1)}(n) \int_0^\infty (t_1 t_2)^{\frac{w-s}{2} + i + n} (t_1 + t_2)^{i + j - 2n - 1} e^{-\frac{\pi t_1 |p_1|^2}{\tau_2} - \frac{\pi t_2 |p_2|^2}{\tau_2}} d^2 t$$

$$+ \sum_{p \in \Lambda'} \frac{6\pi |p|^2}{\tau_2} c_{i,j}^{(0)}(n) \int_0^\infty (t_1 t_2)^{\frac{w-s}{2} + i + n} (t_1 + t_2)^{i + j - 2n} e^{-\frac{\pi |p|^2 (t_1 + t_2)}{\tau_2}} d^2 t, \qquad (3.27)$$

once again in the above we have only focused a single monomial contribution to the boundary term.

We see from (3.20) that in the above expression the exponent α of each term $(t_1 + t_2)^{\alpha}$ is always a positive integer, hence we can use binomial expansion and integrate term by term, arriving at the sum of three boundary contributions,

$$b.t. = 3 \left[(2n - i - j) c_{i,j}^{(0)}(n) + c_{i,j}^{(1)}(n) \right] \sum_{m=0}^{i+j-2n-1} {i+j-2n-1 \choose m} \Gamma\left(\frac{w + (i+j-2m-2n-1)}{2}\right) \Gamma\left(\frac{w - (i+j-2m-2n-1)}{2}\right) \\ \times \left[E\left(\frac{w + (i+j-2m-2n-1)}{2}; \tau\right) E\left(\frac{w - (i+j-2m-2n-1)}{2}; \tau\right) - E(w; \tau) \right] \\ + 6c_{i,j}^{(0)}(n) E(w; \tau) \left[\sum_{m=0}^{i+j-2n} {i+j-2n \choose m} \Gamma\left(\frac{w + 1 + (i+j-2m-2n)}{2}\right) \Gamma\left(\frac{w + 1 - (i+j-2m-2n)}{2}\right) \right],$$
(3.28)

where we have used the identity

$$\sum_{\substack{p_1,p_2,p_3\in\Lambda'\\p_1+p_2+p_3=0}} \frac{(\tau_2/\pi)^{s_1+s_2}}{|p_1|^{2s_1}|p_2|^{2s_2}} = \sum_{p_1,p_2\in\Lambda'} \frac{(\tau_2/\pi)^{s_1+s_2}}{|p_1|^{2s_1}|p_2|^{2s_2}} - \sum_{p\in\Lambda'} \frac{(\tau_2/\pi)^{s_1+s_2}}{|p|^{2(s_1+s_2)}} = E(s_1;\tau)E(s_2;\tau) - E(s_1+s_2;\tau).$$
(3.29)

Note that we have substituted for the eigenvalue s = 3i + j + 1 in (3.28) and we can easily see that the indices of the bilinear in non-holomorphic Eisenstein series are such that $w \pm (i + j - 1) \equiv w - s \pmod{2}$. Hence, we deduce that when w and s have the same parity the bilinear source terms is the product of two non-holomorphic Eisenstein series with integer indices. Similarly, when w and s have opposite parity the bilinear non-holomorphic Eisenstein series have half-integer indices. Finally, the third line gives terms proportional to the non-holomorphic Eisenstein series $E(w; \tau)$. Thus we have reproduce the claim (3.5).

To obtain the complete boundary term we simply need to sum over the boundary terms in (3.28) arising from (3.20). After changing summation variables $(n,m) \to (n,r := n + m - \frac{1}{2}(i+j-1))$ and inverting the Laplace equation (3.11) (with the caveat explained in footnote 6) we arrive at

$$\mathcal{E}_{i,j}^{w}(\tau) = \sum_{\substack{r=-\frac{(i+j-1)}{2}}}^{\frac{(i+j-1)}{2}} b_{i,j}(w,r) \mathcal{E}(s; \frac{w}{2}+r, \frac{w}{2}-r; \tau) + d_{i,j}(w) E(w; \tau), \qquad (3.30)$$

where the coefficients $b_{i,j}(w,r)$ and $d_{i,j}(w)$ are given by

$$b_{i,j}(w,r) = \Gamma\left(\frac{w}{2}+r\right) \Gamma\left(\frac{w}{2}-r\right) \times \left[\sum_{n=0}^{\min\left(\frac{i+j-1}{2}-r,\frac{i+j-1}{2}+r\right)} 3\left[(2n-i-j)c_{i,j}^{(0)}(n)+c_{i,j}^{(1)}(n)\right] \left(\frac{i+j-2n-1}{\frac{1}{2}(i+j-1)+r-n}\right) \right], \quad (3.31)$$

$$d_{i,j}(w) = -\sum_{r=-\frac{(i+j-1)}{2}}^{\frac{(i+j-1)}{2}} \frac{b_{i,j}(w,r)}{[w(w-1)-s(s-1)]} + \sum_{r=-\frac{(i+j)}{2}}^{\frac{(i+j)}{2}} \left\{ \frac{6\Gamma\left(\frac{w+1}{2}+r\right)\Gamma\left(\frac{w+1}{2}-r\right)}{[w(w-1)-s(s-1)]} \times \left[\sum_{n=0}^{\min\left(\frac{i+j}{2}-r,\frac{i+j}{2}+r\right)} c_{i,j}^{(0)}(n) \left(\frac{i+j-2n}{\frac{1}{2}(i+j)+r-n}\right) \right] \right\}. \quad (3.32)$$

Note that the GES terms in (3.30) are invariant under the exchange $w \to 2 - w$ as one can check from the coefficients (3.31) combined with the reflection property for the GESs

$$\Gamma(s_1)\Gamma(s_2)\mathcal{E}(s;s_1,s_2;\tau) = \Gamma(1-s_1)\Gamma(1-s_2)\mathcal{E}(s;1-s_1,1-s_2;\tau), \qquad (3.33)$$

which can be inferred from its differential equation (1.1) combined with the functional identity (2.12). On the other hand, the single Eisenstein series term in (3.30) is not invariant as one can see from the coefficient (3.32) and the result is the modified functional equation (2.38) for $\mathcal{E}_{i,j}^w(\tau)$.

In the next sections 3.2 and 3.3 we provide an alternative way of determining the coefficients $b_{i,j}(w,r)$ of the GES part of (3.30) for general values of $i, j \in \mathbb{N}$. Importantly, we will see that this second method does not rely on knowing the general form for the coefficients $c_{i,j}^{(0)}(n)$ and $c_{i,j}^{(1)}(n)$ in (3.20).

3.2 Alternative approach

In this section we discuss an alternative method, originally presented in [26], for obtaining the source terms in the Laplace equation (3.11). Although this method does not fully capture the single Eisenstein series contribution in (3.28), it leads to a particularly efficient procedure for determining the explicit coefficients $b_{i,j}(w,r)$ for generic values of parameters, i, j, w, r.

For this calculation we make use of the formulation for the theta lift of modular local polynomials in terms of the original variables ρ and V. As discussed just below equation (2.17), the Maass functions $A_{i,j}(\rho)$ are invariant under the permutation group of three elements \mathfrak{S}_3 as is manifest in terms of the t_i variables. In particular, the $A_{i,j}(\rho)$ are invariant under $\rho \to 1 - \bar{\rho}$. It is important to note that the symmetry $\rho_1 \mapsto 1 - \rho_1$, implies that the derivative $\partial_{\rho_1} A_{i,j}(\rho)$ satisfies the following conditions on the boundary of the shaded domain $\mathcal{F}_{\rho} = \Gamma_0(2) \backslash \mathfrak{H}$ in figure 1,

$$\partial_{\rho_1} A_{i,j}(\rho)|_{\rho_1=0} = -\partial_{\rho_1} A_{i,j}(\rho)|_{\rho_1=1}.$$
(3.34)

It is convenient to use the symmetries of $A_{i,j}(\rho)$ to map the original domain \mathcal{F}_{ρ} into a three-fold covering of the fundamental domain of $\mathrm{SL}(2,\mathbb{Z})$ given by $\hat{\mathcal{F}}_{\rho} = \mathrm{SL}(2,\mathbb{Z})\backslash\mathfrak{H}$. In that case the two vertical boundary lines in \mathcal{F}_{ρ} , $\rho_1 = 0$ and $\rho_1 = 1$, are respectively mapped to the lines $\rho_1 = 0 - \epsilon$ and $\rho_1 = 0 + \epsilon$ in $\hat{\mathcal{F}}_{\rho}$, with $\epsilon \to 0^+$. If we now define $\hat{A}_{i,j}(\rho)$ as the mapping of the local Maass function $A_{i,j}(\rho)$ to $\hat{\mathcal{F}}_{\rho}$, we see that the relation (3.34) implies the existence of a discontinuity in $\partial_{\rho_1} \hat{A}(\rho)$ at $\rho_1 = 0$. It therefore follows from $\partial^2_{\rho_1} |\rho_1| = 2\delta(\rho_1)$ that inside the domain $\hat{\mathcal{F}}_{\rho}$ we have

$$\rho_2^2 \,\partial_{\rho_1}^2 \hat{A}_{i,j}(\rho) = 2\rho_2 \,c_{i,j}(\rho_2) \,\delta(\rho_1) \,, \tag{3.35}$$

where $c_{i,j}(\rho_2) = c_{i,j}(1/\rho_2)$ is a Laurent polynomial in ρ_2 that will play a key role in the following.

A crucial consequence of this fact is that on the fundamental domain $\hat{\mathcal{F}}_{\rho}$ the Laplace equation (2.30) is modified to

$$[\Delta_{\rho} - s(s-1)] \hat{A}_{i,j}(\rho) = 2\rho_2 c_{i,j}(\rho_2) \,\delta(\rho_1) \,. \tag{3.36}$$

The contribution of the boundary terms to the Laplace equation may then be determined by an argument similar to (3.8). After changing the integration domain in the theta lift (2.32) from \mathcal{F}_{ρ} to its triple cover $\hat{\mathcal{F}}_{\rho}$, acting with the τ Laplacian gives

$$\Delta_{\tau} \mathcal{E}^{w}_{i,j}(\tau) = 6 \int_{\mathbb{R}^{+} \times \hat{\mathcal{F}}_{\rho}} V^{w} \hat{A}_{i,j}(\rho) \,\Delta_{\rho} \Gamma'_{2,2}(\tau;\rho,iV) \,2 \frac{\mathrm{d}V}{V^{2}} \frac{\mathrm{d}^{2}\rho}{\rho_{2}^{2}} \,, \tag{3.37}$$

where (2.9) has been used. At this point integrating by parts and using (3.36) leads to

$$[\Delta_{\tau} - s(s-1)]\mathcal{E}^{w}_{i,j}(\tau) \sim 6 \int_{\mathbb{R}^{+} \times [1,\infty)} V^{w} c_{i,j}(\rho_{2}) \Gamma'_{2,2}(\tau; i\rho_{2}, iV) 2 \frac{\mathrm{d}V}{V^{2}} \frac{\mathrm{d}\rho_{2}}{\rho_{2}}, \qquad (3.38)$$

where we have discarded contributions on the right-hand side, which are linear combinations of nonholomorphic Eisenstein series.

By performing the change of variables $\rho_2 = \sqrt{y/x}$, $V = \sqrt{xy}$ equation (3.38) becomes (ignoring a subtlety in the limits of integration to be described below),

$$[\Delta_{\tau} - s(s-1)]\mathcal{E}_{i,j}^{w}(\tau) \sim 3\sum_{p_{1},p_{2}\in\Lambda'} \int_{0}^{\infty} x^{\frac{w-2}{2}} y^{\frac{w-2}{2}} c_{i,j}(\sqrt{y/x}) e^{-\pi x \frac{|p_{1}|^{2}}{\tau_{2}} - \pi y \frac{|p_{2}|^{2}}{\tau_{2}}} \,\mathrm{d}x \,\mathrm{d}y.$$
(3.39)

In writing this expression we have assumed that the integration over ρ_2 can be extended from the range $[1, \infty)$ to the semi-infinite line $[0, \infty)$ by exploiting symmetry of the integrand under $\rho_2 \rightarrow 1/\rho_2$. However, this ignores a subtlety that arises from the fact that the manifest \mathfrak{S}_3 symmetry of (2.33) has been lost in mapping the domain \mathcal{F}_{ρ} to $\hat{\mathcal{F}}_{\rho}$ and the symmetry $\rho_2 \rightarrow 1/\rho_2$ only exists if we remove the restriction $p_1 + p_2 \neq 0$. In other words, the expression (3.39) includes the $p_1 + p_2 = 0$ term but the $p_1 = 0$ and $p_2 = 0$ terms must be omitted in order to avoid obvious divergences in the expression for the source term. So in order to maintain \mathfrak{S}_3 symmetry we must subtract the $p_1 + p_2 = 0$ term from (3.39) even though its presence does not cause a divergence. From (3.29) it is easy to see that subtracting the $p_1 + p_2 = 0$ term adds a term proportional to a single non-holomorphic Eisenstein series, while leaving the bilinear source term unaffected.

From (2.28) the Laurent polynomial $c_{i,j}(\rho_2)$ takes the form

$$c_{i,j}(\rho_2) = \sum_{r=-\frac{(i+j-1)}{2}}^{2} s_{i,j}(r) \,\rho_2^{2r} \,.$$
(3.40)

Importantly, we see that each monomial in the above expression leads to a particular bilinear in Eisenstein series in (3.39),

$$\left[\Delta_{\tau} - s(s-1)\right] \mathcal{E}_{i,j}^{w}(\tau) \sim \sum_{r=-\frac{(i+j-1)}{2}}^{\frac{(i+j-1)}{2}} 3s_{i,j}(r) \Gamma\left(\frac{w}{2}+r\right) \Gamma\left(\frac{w}{2}-r\right) E(\frac{w}{2}+r;\tau) E(\frac{w}{2}-r;\tau) , \qquad (3.41)$$

where we have made use of the integral representation for the non-holomorphic Eisenstein series (A.6). Comparing this equation to (3.30) we deduce that the coefficients $b_{i,j}(w,r)$ given in (3.31) can be written in the simpler form

$$b_{i,j}(w,r) = 3s_{i,j}(r)\Gamma\left(\frac{w}{2}+r\right)\Gamma\left(\frac{w}{2}-r\right).$$
(3.42)

In the next section this alternative computation of the source term (3.41) will be used to derive a general expression valid for all $i, j \in \mathbb{N}$ of the coefficients $b_{i,j}(w, r)$ of the GES part of (3.30).

3.3 General expression for $\mathcal{E}_{i,j}^w(\tau)$

We now exploit the analysis of section 3.2 to deduce a general expression for the source term (3.36) generated by the modular local Maass function $A_{i,j}(\rho)$ defined in (2.28) and hence derive an explicit formula for the particular linear combinations of GESs produced by $\mathcal{E}_{i,j}^{w}(\tau)$. Thanks to (3.40) and (3.42), this will in turn allow us to determines the coefficients $b_{i,j}(w,r)$ of the GES part of (3.30).

The coefficient $s_{i,j}(r)$ appearing in (3.40) for a particular $A_{i,j}(\rho)$, can be obtained by expanding the corresponding modular local polynomial, $u^i v^j$, as a polynomial in ρ ,

$$u^{i}v^{j} = \rho^{2i}(1-\rho)^{2i}(\rho^{2}-\rho+1)^{j} = \sum_{k=0}^{2i}\sum_{\ell=0}^{j}\sum_{m=0}^{\ell}(-1)^{k+m+\ell}C_{i,j;k,\ell,m}\rho^{2i+k+\ell+m}, \qquad (3.43)$$

where $C_{i,j;k,\ell,m}$ are simply the binomial coefficients, given by

$$C_{i,j;k,\ell,m} \coloneqq \binom{2i}{k} \binom{j}{\ell} \binom{\ell}{m}.$$
(3.44)

The task of computing the source term (3.35) for a fixed $A_{i,j}(\rho)$ is then simplified to determining the source contribution that originates from a single monomial term $\rho^{2i+k+m+\ell}$. Firstly, we need to use the definition (2.26) to perform the Maass raising (2.28) and compute

$$D_{-2n}^{(n)}[\rho^q] = \frac{(-2i)^n n!}{(2n)!} \sum_{m=0}^n \binom{n}{m} \frac{(-n-m)_m}{(2i\rho_2)^m} \frac{\Gamma(q+1)}{\Gamma(q+m-n+1)} \rho^{q+m-n}, \qquad (3.45)$$

where the integer n is related to the eigenvalue s via n = s - 1 = 3i + j. We then use (3.35) to derive

$$\rho_{2}^{2} \partial_{\rho_{1}}^{2} \left(D_{-2n}^{(n)}[\rho^{q}] \right) = 2\rho_{2} \frac{(-1)^{n+\frac{q+3}{2}} \Gamma(n+1)\Gamma(q+1)\rho_{2}^{q-n}}{\Gamma(2n+1)} \Big[\sum_{m=0}^{n} \frac{(-1)^{m} 2^{n-m} \Gamma(m+n+1)}{\Gamma(m+1)\Gamma(n-m+1)\Gamma(m-n+q)} \Big] \delta(\rho_{1}) \\ = 2\rho_{2} \left[(1-(-1)^{q}) \frac{\Gamma\left(\frac{q}{2}+1\right) \Gamma\left(n-\frac{q}{2}+1\right)}{\sqrt{\pi} \Gamma\left(n+\frac{1}{2}\right)} \rho_{2}^{q-n} \right] \delta(\rho_{1}).$$

$$(3.46)$$

From the above equation we use (3.39) to derive the Laplace eigenvalue equation satisfied by the theta lift of the single Maass lifted monomial $D_{-2n}^{(n)}[\rho^q]$, which we denote as $\mathcal{E}_{s;q}(\tau)$,

$$[\Delta_{\tau} - s(s-1)]\mathcal{E}_{s;q}(\tau) = 3[1 - (-1)^q]\widetilde{C}_{s;s_1,s_2}E(s_1;\tau)E(s_2;\tau), \qquad (3.47)$$

where s_1, s_2 have been defined to be

$$s_1 = \frac{w+s-q-1}{2}, \qquad s_2 = \frac{w-s+q+1}{2},$$
 (3.48)

and the coefficient $\widetilde{C}_{s;s_1,s_2}$ is given by

$$\widetilde{C}_{s;s_1,s_2} \coloneqq \frac{\Gamma(s_1)\Gamma(s_2)}{\sqrt{\pi}\Gamma\left(s-\frac{1}{2}\right)} \Gamma\left(\frac{s+s_1-s_2+1}{2}\right) \Gamma\left(\frac{s-s_1+s_2+1}{2}\right) \,. \tag{3.49}$$

We see that $\mathcal{E}_{s;q}(\tau)$ is simply a multiple of a single GES with a non-trivial normalisation factor related to $\widetilde{C}_{s;s_1,s_2}$. We anticipate that these coefficients $\widetilde{C}_{s;s_1,s_2}$ will play a key role in section 4.

Finally, we combine the result (3.47) for the single monomial with the general expansion (3.43) to arrive at

$$\left[\Delta_{\tau} - s(s-1)\right] \mathcal{E}_{i,j}^{w}(\tau) = 3\sum_{k=0}^{2i} \sum_{\ell=0}^{j} \sum_{m=0}^{l} \left(1 - (-1)^{k+\ell+m}\right) C_{i,j;k,\ell,m} \widetilde{C}_{s;s_1,s_2} E(s_1;\tau) E(s_2;\tau),$$
(3.50)

with s = 3i + j + 1, and

$$s_1 = \frac{w}{2} + \frac{k + \ell + m - i - j}{2}, \qquad s_2 = \frac{w}{2} - \frac{k + \ell + m - i - j}{2}.$$
(3.51)

We therefore conclude that the theta lift (2.32) of the modular local Maass form $A_{i,j}(\rho)$ is given by

$$\mathcal{E}_{i,j}^{w}(\tau) = 3\sum_{k=0}^{2i}\sum_{\ell=0}^{j}\sum_{m=0}^{\ell} \left[1 - (-1)^{k+\ell+m}\right] C_{i,j;k,\ell,m} \widetilde{C}_{s;s_1,s_2} \mathcal{E}(s;s_1,s_2;\tau) + d_{i,j}(w) E(w;\tau) \,. \tag{3.52}$$

In the above equation we have also re-instated the boundary term contribution producing a multiple of the non-holomorphic Eisenstein series $E(w; \tau)$ and evaluated earlier in (3.32), although this simpler contribution will not play any role for the following discussion.

Using (3.51) it is easy to see the linear combination of GESs $\mathcal{E}(s; s_1, s_2; \tau)$ in (3.52) appears with a range of indices s_1, s_2 , which precisely matches the earlier statement (3.7). We conclude that for arbitrary $i, j \in \mathbb{N}$, the coefficients $b_{i,j}(w, r)$ of the generalised Eisenstein contribution to $\mathcal{E}_{i,j}^w(\tau)$ as presented in (3.7) are given by

$$b_{i,j}(w,r) = 3\sum_{k=0}^{2i} \sum_{\ell=0}^{j} \sum_{m=0}^{\ell} \delta(2r - (k+\ell+m-i-j)) \left[1 - (-1)^{k+\ell+m}\right] C_{i,j;k,\ell,m} \widetilde{C}_{s;s_1,s_2},$$
(3.53)

with $\delta(r)$ denoting Kronecker delta and s_1, s_2 given in (3.51).

Although the general coefficients (3.53) are not quite expressed in a fully explicit form, in section 4 we will see that they do hide rather beautiful mathematical structures.

4 Structure of generalised Eisenstein series

This section emphasises some interesting number theoretic aspects of GESs. In particular, we discuss the occurrence of the *L*-values associated with holomorphic cusp forms in the Fourier mode expansion of GESs. Surprisingly, due to some intriguing cancellations such *L*-values are absent from the modular functions $\mathcal{E}_{i,j}^w(\tau)$, which are constructed as theta lifts of local Maass functions (2.32) and correspond to MGFs and SMFs that are relevant to superstring scattering amplitudes.

4.1 Fourier mode expansion

It is useful to consider the structure of the Fourier mode expansion of GESs with respect to $\text{Re } \tau = \tau_1$. As in (1.1) we define a GES as the modular invariant solution to the inhomogeneous Laplace eigenvalue equation

$$(\Delta_{\tau} - s(s-1))\mathcal{E}(s; s_1, s_2; \tau) = E(s_1; \tau)E(s_2; \tau), \qquad (4.1)$$

subject to the condition that $\mathcal{E}(s; s_1, s_2; \tau)$ has moderate growth at the cusp $\tau_2 \to \infty$ and fixing the boundary condition that the coefficient of the solution of the homogeneous equation τ_2^s vanishes, thus implying the

growth condition $\mathcal{E}(s; s_1, s_2; \tau) = O(\tau_2^{s_1+s_2})$ as $\tau \to i \infty$. For $s \in \mathbb{N}$, which is the case of present interest, $\mathcal{E}(s; s_1, s_2; \tau)$ is the unique modular invariant solution to (4.1) subject to the above boundary conditions.

It follows from the Fourier mode decomposition of the non-holomorphic Eisenstein series (A.3), that the GESs $\mathcal{E}(s; s_1, s_2; \tau)$ can be decomposed in Fourier modes as

$$\mathcal{E}(s;s_1,s_2;\tau) = \sum_{n,m=0}^{\infty} \mathcal{E}^{(n,m)}(s;s_1,s_2;\tau_2) q^n \bar{q}^m \,, \tag{4.2}$$

where $q = e^{2\pi i \tau}$ and $\bar{q} = e^{-2\pi i \bar{\tau}}$. The mode number is given by k = n - m and modes with k > 0 represent the contributions of instantons while modes with k < 0 are contributions of anti-instantons. It is useful to separate the terms with k = 0 into purely perturbative contributions, coming from the m = n = 0 term, and instanton/anti-instanton contributions where m = n > 0.

The $\mathcal{E}^{(0,0)}$ term is a Laurent polynomial that is given by⁷

$$\mathcal{E}^{(0,0)}(s;s_1,s_2;\tau_2) = \frac{4\pi^{-2s_1-2s_2}\zeta(2s_1)\zeta(2s_2)}{\pi^{s_1+s_2}(s_1+s_2-s)(s_1+s_2+s-1)}(\pi\tau_2)^{s_1+s_2} + \frac{4\pi^{-\frac{1}{2}-2s_2}\Gamma(s_1-\frac{1}{2})\zeta(2s_1-1)\zeta(2s_2)}{(s_2-s_1-s+1)\Gamma(s_1)}(\pi\tau_2)^{1-s_1+s_2} + \frac{4\pi^{-\frac{1}{2}-2s_1}\Gamma(s_2-\frac{1}{2})\zeta(2s_2-1)\zeta(2s_1)}{(s_1-s_2+s)(s_1-s_2-s+1)\Gamma(s_2)}(\pi\tau_2)^{1-s_2+s_1} + \frac{4\Gamma(s_1-\frac{1}{2})\Gamma(s_2-\frac{1}{2})\zeta(2s_1-1)\zeta(2s_2-1)}{\pi(s_1+s_2-s-1)(s_1+s_2+s-2)\Gamma(s_1)\Gamma(s_2)}(\pi\tau_2)^{2-s_1-s_2} + \beta(s;s_1,s_2)(\pi\tau_2)^{1-s},$$
(4.3)

where the first four terms can be obtained by matching powers of τ_2 on the left-hand and right-hand sides of (4.1) and the last term satisfies the homogeneous equation and its coefficient is given by $[12, 26]^8$

$$\beta(s;s_1,s_2) \coloneqq \frac{4\pi^{s-1}}{\Gamma(s_1)\Gamma(s_2)} \frac{\zeta^*(s-s_1-s_2+1)\zeta^*(s+s_1-s_2)\zeta^*(s-s_1+s_2)\zeta^*(s+s_1+s_2-1)}{(1-2s)\,\zeta^*(2s)}, \quad (4.4)$$

with $\zeta^*(s) \coloneqq \zeta(s)\Gamma(s/2)/\pi^{s/2}$.

(0.0)

In addition to the purely perturbative terms, the zero-mode sector contains an infinite tower of instanton/antiinstanton contributions, (i.e. all $\mathcal{E}^{(n,n)}$ terms with n > 0), which have the form

$$\mathcal{E}^{(n,n)}(s;s_1,s_2;\tau_2) = \frac{n^{s_1+s_2-2}\sigma_{1-2s_1}(n)\sigma_{1-2s_2}(n)}{\Gamma(s_1)\Gamma(s_2)}\Phi_{s;s_1,s_2}(4\pi n\tau_2), \qquad (4.5)$$

where $\sigma_s(n) \coloneqq \sum_{d|n} d^s$ denotes the standard divisor sigma function. When $s_1, s_2 \in \mathbb{N}$ the expression $\Phi_{s;s_1,s_2}(4\pi n\tau_2)$ is a polynomial in inverse powers of τ_2 of degree $s_2 + s_1 - 2$. The first two perturbative orders in the (n, n) sector do not depend on the eigenvalue s and have the form

$$\Phi_{s;s_1,s_2}(y) = \frac{8}{y^2} + \frac{8[s_1(s_1-1) + s_2(s_2-1) - 4]}{y^3} + O(y^{-4}).$$
(4.6)

⁷There are some special cases for the parameters s_1, s_2 and s for which either the differential equation (4.1) does not admit a modular invariant solution, or the Laurent polynomial has to be slightly modified, see e.g. [30]. In particular, we note that for the case where either $s_1 = \frac{1}{2}$ or $s_2 = \frac{1}{2}$, the Laurent polynomial requires the addition of a logarithmic term $\log(\tau_2)$. None of these cases seem to be important for either MGFs or SMFs.

⁸In these references, this coefficient is determined by projecting the Laplace equation (4.1) on $E(s;\tau)$. Alternatively, it may be obtained from a Poincaré series representation as in [9].

Higher-order coefficients do depend on s and can be computed from the differential equation (4.1) as performed in [20, 23] or via resurgence analysis methods [31, 32, 33]. Note that similar instanton/anti-instanton contributions contribute to the general k^{th} Fourier mode associated with terms in (4.2),

$$\mathcal{E}^{(n,m)}(s;s_1,s_2;\tau_2) = \frac{n^{s_1-1}m^{s_2-1}\sigma_{1-2s_1}(n)\sigma_{1-2s_2}(m)}{\Gamma(s_1)\Gamma(s_2)} \left(\frac{1}{4nm(\pi\tau_2)^2} + O(\tau_2^{-3})\right) + (n\leftrightarrow m), \quad (4.7)$$

where $k = n - m \neq 0$ and both m, n are non-vanishing.

Analysis of the (n, 0) and (0, m) terms requires more care since in these sectors we have the freedom of adding solutions of the homogeneous equation to the differential equation (4.1) without spoiling either the boundary condition or the moderate growth condition at the cusp. From the differential equation (4.1) it is easy to see that the solutions of the homogeneous equation in the zero-mode sector are simply τ_2^s and τ_2^{1-s} . While the coefficient of the τ_2^s term has been set to zero thanks to our choice of boundary condition, the coefficient of τ_2^{1-s} must hence take the value (4.4) as a consequence of modular invariance. For the non-zero Fourier mode the story is more interesting since the solution of the homogeneous equation in the n^{th} Fourier mode sector (with $n \neq 0$) is proportional to $K_{s-\frac{1}{2}}(2\pi |n|\tau_2)$, which satisfies ⁹

$$(\Delta_{\tau} - s(s-1)) \left[e^{2\pi i n \tau_1} \sqrt{|n| \tau_2} K_{s-\frac{1}{2}}(2\pi |n| \tau_2) \right] = 0.$$
(4.8)

The modified Bessel function is exponentially suppressed at the cusp, i.e.

$$\sqrt{|n|\tau_2}K_{s-\frac{1}{2}}(2\pi|n|\tau_2) = e^{-2\pi|n|\tau_2} \left(\frac{1}{2} + \frac{s(s-1)}{8\pi|n|\tau_2} + O(\tau_2^{-2})\right), \qquad \tau_2 \gg 1.$$
(4.9)

Furthermore we note for future convenience that at the origin $\tau_2 \rightarrow 0$ this solution of the homogeneous equation behaves as

$$\sqrt{|n|\tau_2}K_{s-\frac{1}{2}}(2\pi|n|\tau_2) = \frac{(\pi|n|\tau_2)^{1-s}\Gamma(s-\frac{1}{2})}{2\sqrt{\pi}} \left(1 - \frac{2(\pi|n|\tau_2)^2}{2s-3} + O(\tau_2^4)\right), \qquad \tau_2 \to 0.$$
(4.10)

A key point is that a particular solution of the differential equation (4.1) is not necessarily modular invariant. To ensure that we have indeed constructed the unique modular invariant solution to (4.1) (given the boundary condition above) it may be necessary to add to the particular solution a suitable solution to the homogeneous equation

$$(\Delta_{\tau} - s(s-1)) H(\tau) = 0, \qquad (4.11)$$

which is given by the linear combination¹⁰

$$H(\tau) = \sum_{n=1}^{\infty} \alpha_n \sqrt{n\tau_2} K_{s-\frac{1}{2}} (2\pi n\tau_2) \left(e^{2\pi i n\tau_1} + e^{-2\pi i n\tau_1} \right), \tag{4.12}$$

with particular real coefficients α_n . We now show that this is precisely the case when constructing GESs relevant for MGFs and SMFs, and the homogeneous coefficients α_n are crucially related to the Fourier coefficients of certain holomorphic cusp forms.

⁹Note that there is a second, linearly independent, solution of the homogeneous equation that is proportional to the modified Bessel function of the first kind $I_{s-\frac{1}{2}}(2\pi|n|\tau_2)$. This solution to the homogeneous equation is discarded since it grows exponentially at the cusp $\tau_2 \gg 1$.

¹⁰Here we are interested in constructing modular invariant solutions to (4.1) that are even under the upper-half plane involution $\tau \to -\bar{\tau}$. This condition forces the solution of the homogeneous equation to take the form of (4.12) and the coefficients α_n to be real. In section 5 we comment on a different family of modular invariant functions introduced in [9] that are odd under this involution and are related to cuspidal MGFs.

4.2 L-values and generalised Eisenstein series

In this subsection we clarify how to determine the coefficients in the solution of the homogeneous equation (4.12) so as to obtain a modular invariant solution to (4.1). We first discuss how to construct a particular solution to the differential equation (4.1) and how to check whether or not this is modular invariant. Since the discussion is rather different for the case of GESs relevant to MGFs (where $s_1, s_2 \in \mathbb{N}^+$) and the case relevant to SMFs (where $s_1, s_2 \in \mathbb{N}^+$) and the case relevant to SMFs (where $s_1, s_2 \in \mathbb{N} + \frac{1}{2}$) we will discuss these two cases separately. However, the basic arguments that determine the solution of the homogeneous equation are the same for both families of GESs. We will present a conjectured closed-form expression for the coefficient of the solution of the homogeneous equation for the GESs relevant for MGFs (4.20) that reproduces all previously known results. Strikingly, we prove that the analytic continuation (4.39) of this conjectural expression to the case of GESs relevant for SMFs also produces the correct solution for the coefficient of the solution of the homogeneous equation, which will be determined thanks to some recent results on convolution identities for divisor sums [22]. We will highlight these important parallels amongst the two families when they arise in the following discussion.

Two-loop Modular Graph Functions

In [9], two related methods for constructing solutions to the differential equation (4.1) were presented for the case

$$s_1, s_2 \in \mathbb{N}$$
, with $s_1, s_2 \ge 2$, $s \in \{|s_1 - s_2| + 2, |s_1 - s_2| + 4, \dots, s_1 + s_2 - 4, s_1 + s_2 - 2\}$. (4.13)

A first approach is via Poincaré series where the key idea is to write one of the non-holomorphic Eisenstein series on the right hand side of (4.1), say $E(s_1; \tau)$, as a sum over images under the action of $B(\mathbb{Z})\backslash SL(2,\mathbb{Z})$ of the much simpler function $\tau_2^{s_1}$ where $B(\mathbb{Z}) := \{ \begin{pmatrix} 1 & n \\ 0 & 1 \end{pmatrix} | n \in \mathbb{Z} \}$ is the Borel stabiliser of the cusp $\tau = i\infty$. It is then possible to find a solution to this modified differential equation via standard methods. Upon summing this solution over all the images under $B(\mathbb{Z})\backslash SL(2,\mathbb{Z})$ we retrieve the modular invariant GES solution.

One of the advantages of this approach is that it immediately generates a modular invariant solution to (4.1). However, the drawback is that while it is possible [31, 32] to extract the zero-mode sector $\mathcal{E}^{(n,n)}$ (4.5) from the Poincaré series, it becomes much harder, if not impossible, to say anything about the instanton or anti-instanton sectors $\mathcal{E}^{(n,0)}$ and $\mathcal{E}^{(0,n)}$.

Interestingly, this Poincaré series approach suggests a different way of tackling the problem of obtaining a particular solution to (4.1) for the spectrum of sources and eigenvalues of present interest. In [9] it was realised that passing to the Poincaré seeds for the GESs motivates a certain notion of *depth* for the functional space under consideration. Building on [6, 7, 34], in [9] it was shown that a possible Poincaré representation of the GESs with spectrum (4.13) can be given in terms of depth-one iterated integrals of holomorphic Eisenstein series, and their complex conjugates, thus suggesting that GESs themselves are given by depthtwo iterated integrals of holomorphic Eisenstein series and their complex conjugates plus possibly lower depth terms.

While invariance under a T-transformation $\tau \to \tau + 1$ is manifest in the space of iterated Eisenstein integrals, invariance under a S-transformation $\tau \to -1/\tau$ is not and has to be checked case by case. Given a particular solution constructed in terms of iterated integrals of two holomorphic Eisenstein series, its Stransform can be evaluated and it can be checked whether or not the particular solution is modular invariant. The difference between an iterated integral of holomorphic Eisenstein series and its S-dual is controlled by a family of periods called *multiple modular values*. While at depth-one the only multiple modular values we find are standard Riemann zeta values, the same is not true at higher depth. In particular, as discussed in [9, 10] multiple modular values at depth-two display (amongst other things) L-values of holomorphic cusp forms. As a consequence, we see that for the spectrum (4.13) the particular solution to (4.1) constructed via depth-two iterated Eisenstein integrals fails to be modular invariant whenever the eigenvalue s is such that dim $S_{2s} \neq 0$ where S_{2s} is the vector space of holomorphic cusp forms with weight 2s. In appendix C we review some important properties of holomorphic cusp forms and their associated *L*-values.

As argued above, if the particular solution is not modular invariant we must add a suitable solution of the homogeneous equation of the form (4.12) to obtain the unique GES modular invariant solution to (4.1). In [9, 10] it was shown that the failure of modularity for the particular depth-two iterated Eisenstein integral solution can be cancelled by a suitable multiple of the non-modular invariant solution of the homogeneous equation

$$H_{\Delta}(\tau) \coloneqq \sum_{n=1}^{\infty} \frac{a_{\Delta}(n)}{n^s} \sqrt{n\tau_2} K_{s-\frac{1}{2}} (2\pi n\tau_2) \left(e^{2\pi i n\tau_1} + e^{-2\pi i n\tau_1} \right)$$
$$= (-1)^s \frac{\pi^s i}{\Gamma(s)} \tau_2^{1-s} \int_{\tau}^{i\infty} (\tau - \tau_1)^{s-1} (\bar{\tau} - \tau_1)^{s-1} \Delta(\tau) \, \mathrm{d}\tau_1 + \mathrm{c.c.} \,, \tag{4.14}$$

which can be thought of as a depth-one iterated integral of the Hecke-normalised, holomorphic cusp form $\Delta(\tau) = \sum_{n=1}^{\infty} a_{\Delta}(n)q^n \in S_{2s}$. Hecke normalisation implies $a_{\Delta}(1) = 1$ and we refer again to appendix C for important aspects on the theory of holomorphic cusp forms.

It should be stressed that the Eichler integral $H_{\Delta}(\tau)$ is not modular invariant on its own. However, the addition of a suitable multiple of $H_{\Delta}(\tau)$ to the particular solution $\mathcal{E}_{p}(s; s_{1}, s_{2}; \tau)$ is crucial for obtaining a modular invariant solution to (4.1). As a result the GESs can be expressed as

$$\mathcal{E}(s;s_1,s_2;\tau) = \mathcal{E}_{\mathbf{p}}(s;s_1,s_2;\tau) + \sum_{\Delta \in \mathcal{S}_{2s}} \lambda_{\Delta}(s;s_1,s_2) H_{\Delta}(\tau), \qquad (4.15)$$

where the sum is over a basis of Hecke-normalised cusp forms for S_{2s} . In this expression $\mathcal{E}_{p}(s; s_1, s_2; \tau)$ denotes the particular solution constructed in [9], which contains only iterated integrals of holomorphic Eisenstein series with depth less than or equal to two and their complex conjugates.

Recalling the formula for the dimension of the vector space of holomorphic cusp forms:

$$\dim \mathcal{S}_{2s} = \begin{cases} \left\lfloor \frac{2s}{12} \right\rfloor - 1 & 2s \equiv 2 \mod 12, \\ \left\lfloor \frac{2s}{12} \right\rfloor & \text{otherwise}, \end{cases}$$
(4.16)

we see that the first instance where the particular solution fails to coincide with the GESs happens at eigenvalue s = 6 for which we have $S_{12} = \operatorname{span}\{\Delta_{12}\}$ where $\Delta_{12} = \sum_{n=1}^{\infty} \tau(n)q^n$ is the discriminant modular form whose q-series coefficients are given by the Ramanujan tau-function $\tau(n)$ (not to be confused with the modular parameter τ). For the spectrum (4.13) the eigenvalue s = 6 is attained by considering sources with indices $(s_1, s_2) \in \{(2, 6), (3, 5), (4, 4)\}$ and we find that the corresponding GESs are given by [9, 10]

$$\mathcal{E}(6;2,6;\tau) = \mathcal{E}_{p}(6;2,6;\tau) + \frac{2}{17275} \frac{\Lambda(\Delta_{12},13)}{\Lambda(\Delta_{12},11)} H_{\Delta_{12}}(\tau) ,$$

$$\mathcal{E}(6;3,5;\tau) = \mathcal{E}_{p}(6;3,5;\tau) - \frac{5}{11056} \frac{\Lambda(\Delta_{12},13)}{\Lambda(\Delta_{12},11)} H_{\Delta_{12}}(\tau) ,$$

$$\mathcal{E}(6;4,4;\tau) = \mathcal{E}_{p}(6;4,4;\tau) + \frac{7}{10365} \frac{\Lambda(\Delta_{12},13)}{\Lambda(\Delta_{12},11)} H_{\Delta_{12}}(\tau) .$$

(4.17)

Here $\Lambda(\Delta, t)$ denotes the standard completed *L*-function, defined in (C.8), of the cusp form $\Delta \in \mathcal{S}_{2s}$.

While in [9] the coefficients $\lambda_{\Delta}(s; s_1, s_2)$ have not been determined in closed form for general parameters s, s_1, s_2 in (4.13), they appear to take the form

$$\lambda_{\Delta}(s;s_1,s_2) = \kappa_{\Delta,s,s_1,s_2} \frac{\Lambda(\Delta, w+s-1)}{\Lambda(\Delta, 2s-1)}, \qquad (4.18)$$

where $w = s_1 + s_2$. From the spectrum (4.13) it is easy to see that since $w = s_1 + s_2 \ge s + 2$ it follows that the ratio of *L*-values in the coefficients (4.18) always consists of an odd critical *L*-value in the denominator and an odd non-critical *L*-value in the numerator where, for a cusp form $\Delta \in S_{2s}$, the critical *L*-values are the numbers $\Lambda(\Delta, t)$ with $t \in \{1, \ldots, 2s - 1\}$. The general expression for the coefficients $\kappa_{\Delta,s,s_1,s_2}$ is not determined at this stage but they can be evaluated on a case by case basis by evaluating the multiple modular values associated with the particular solution part of (4.15) and they are generically in the number field associated with S_{2s} .¹¹ However, we will shortly propose a strongly motivated conjectural expression for the coefficients in (4.18) based on the expression (4.40) for these coefficients in the context of SMFs, which will be discussed in the next sub-section.

In order to make contact with the expression for λ_{Δ} that we will find for SMFs in the next sub-section it is convenient to re-express (4.18) so that the critical *L*-value in the denominator is converted to the Petersson norm $\langle \Delta, \Delta \rangle$ defined in (C.15). This makes use of a classic result by Rankin [35] that states that the Petersson norm of a Hecke eigenform Δ can be written in terms of the product of two critical *L*-values with opposite parity. For example the Petersson norm of the Ramanujan cusp form Δ_{12} , i.e. the unique Hecke normalised element of S_{12} , can be written as (see (9.1) of [35])

$$\langle \Delta_{12}, \Delta_{12} \rangle = \frac{691}{7680} \Lambda(\Delta_{12}, 8) \Lambda(\Delta_{12}, 11) \,. \tag{4.19}$$

The general expression for the norms of arbitrary cusp forms follows from Rankin's theorem stated explicitly in (C.16).

Furthermore, a beautiful result by Manin [36] known as the periods theorem and briefly reviewed in appendix C, proves that the ratio of any two even critical L-values or any two odd critical L-values are rational over the algebraic number field generated by the Fourier coefficients of the cusp forms. Hence by combining (C.16) and Manin's theorem we can rewrite (4.18) as

$$\lambda_{\Delta}(s;s_1,s_2) = \tilde{\kappa}_{\Delta,s,s_1,s_2} \frac{\Lambda(\Delta,s+s_1-s_2)\Lambda(\Delta,s+w-1)}{\langle \Delta,\Delta \rangle} \,. \tag{4.20}$$

According to Conjecture 1 presented in section 1.2 the expression for the coefficients of the solution of the homogeneous equation, $\lambda_{\Delta}(s; s_1, s_2)$, in the MGF case (with integer s_1 and s_2) are the same as those in the SMF case with (half-integer s_1 and s_2). An explicit expression for the latter will be derived in the next sub-section and is given in (4.39)-(4.40). As a result the coefficient $\tilde{\kappa}_{\Delta,s,s_1,s_2}$ which appears in (4.20) is given by

$$\tilde{\kappa}_{\Delta,s,s_1,s_2} = (-1)^{\frac{s+s_1-s_2+2}{2}} \frac{\Gamma(s)\Gamma\left(\frac{w-s}{2}\right)}{2^{2s-3}\Gamma\left(\frac{w+s}{2}\right)\tilde{C}_{s;s_1,s_2}},\tag{4.21}$$

with $w = s_1 + s_2$ and where $\widetilde{C}_{s;s_1,s_2}$ is the rational number defined in (3.49). The number field content for the coefficient $\lambda_{\Delta}(s;s_1,s_2)$ in the expression (4.20) is entirely carried by the Petersson norm $\langle \Delta, \Delta \rangle$ in the denominator, since $\tilde{\kappa}_{\Delta,s,s_1,s_2}$ in (4.21) is manifestly rational for the spectrum (4.13). Furthermore, for this spectrum $\lambda_{\Delta}(s;s_1,s_2)$ always contains the product of an even critical *L*-value and an odd non-critical *L*-value.

We have checked that our conjectural expression (4.20) does indeed reproduce all particular examples presented in the ancillary file of [9]. If we rewrite the particular examples (4.17) using (4.19) in order to

¹¹Whenever dim $S_{2s} = 1$ this number field actually coincides with the rational numbers. However, when dim $S_{2s} > 1$ that is no longer true. For example, with 2s = 24 we have dim $S_{24} = 2$ and the Fourier coefficients of the two Hecke eigenforms basis elements of S_{24} belong to the number field $\mathbb{Q}[\sqrt{144169}]$. by the Fourier coefficients a(n) of $\Delta(\tau) = \sum_{n=1}^{\infty} a(n)q^n$.

implement the above representation we can check that (4.20) is correct for s = 6 and we find

$$\mathcal{E}(6;2,6;\tau) = \mathcal{E}_{p}(6;2,6;\tau) + \frac{1}{184320} \frac{\Lambda(\Delta_{12},10)\Lambda(\Delta_{12},13)}{\langle\Delta_{12},\Delta_{12}\rangle} H_{\Delta_{12}}(\tau),$$

$$\mathcal{E}(6;3,5;\tau) = \mathcal{E}_{p}(6;3,5;\tau) - \frac{125}{32} \cdot \frac{1}{184320} \frac{\Lambda(\Delta_{12},10)\Lambda(\Delta_{12},13)}{\langle\Delta_{12},\Delta_{12}\rangle} H_{\Delta_{12}}(\tau),$$

$$\mathcal{E}(6;4,4;\tau) = \mathcal{E}_{p}(6;4,4;\tau) + \frac{35}{6} \cdot \frac{1}{184320} \frac{\Lambda(\Delta_{12},10)\Lambda(\Delta_{12},13)}{\langle\Delta_{12},\Delta_{12}\rangle} H_{\Delta_{12}}(\tau),$$

(4.22)

where we have expressed all the critical L-values in terms of $\Lambda(\Delta_{12}, 10)$ by using Manin's relations [36]. We will shortly see that the coefficients in front of the solutions of the homogeneous equation appearing in (4.22)are actually deeply connected with the particular combinations of GESs obtained from the theta-lifted local Maass functions discussed in section 3. However, before doing that we will now review how a similar interplay between the particular solution and the solution of the homogeneous equation related to holomorphic cusp forms is also present in the context of GESs associated with SMFs.

S-dual Modular Functions

We now consider the GES solution to (4.1) with spectrum¹²

$$s_1, s_2 \in \mathbb{N} + \frac{1}{2}, \qquad s \in \{s_1 + s_2 + 1, s_1 + s_2 + 3, \ldots\}.$$
 (4.23)

An algorithm for constructing a particular solution, $\mathcal{E}_{p}(s; s_1, s_2; \tau)$, to (4.1) for the special case $\mathcal{E}_{p}(4; \frac{3}{2}, \frac{3}{2}; \tau)$ in [37] (see also [20]), and later generalised in [23] to the spectrum (4.23). Particularly important for the present discussion is the form of the particular solution for $\mathcal{E}_{p}^{(n,0)}$ and $\mathcal{E}_{p}^{(0,n)}$ with $n \neq 0$ as defined in (4.2). Given the differential equation (4.1) and the Fourier mode expansion for the non-holomorphic Eisenstein

series (A.3) we see that $\mathcal{E}_p^{(n_1,n_2)}$ with $n_1 + n_2 \neq 0$ can be represented as

$$e^{-2\pi(n_1+n_2)\tau_2} \mathcal{E}_{\mathbf{p}}^{(n_1,n_2)}(s;s_1,s_2;\tau_2) = \hat{g}_{n_1,n_2}(\tau_2), \qquad (4.24)$$

where the functions $\hat{g}_{n_1,n_2}(\tau_2)$ are solutions to the second order differential equations

$$\left[\tau_2^2 \partial_2 - (2\pi n\tau_2)^2 - s(s-1) \right] \left(\hat{g}_{n,0}(\tau_2) + \hat{g}_{0,n}(\tau_2) \right) = \\ \left(\frac{2\zeta(2s_1)}{\pi^{s_1}} \tau_2^{s_1} + \frac{2\Gamma(s_1 - \frac{1}{2})}{\pi^{s_1 - \frac{1}{2}}\Gamma(s_1)} \zeta(2s_1 - 1) \tau_2^{1-s_1} \right) \frac{4\sqrt{\tau_2}}{\Gamma(s_2)} n^{s_2 - \frac{1}{2}} \sigma_{1-2s_2}(n) K_{s_2 - \frac{1}{2}}(2\pi n\tau_2) + (s_1 \leftrightarrow s_2) ,$$

$$(4.25)$$

and

$$\left[\tau_2^2 \partial_2 - (2\pi (n_1 + n_2)\tau_2)^2 - s(s-1) \right] \hat{g}_{n_1,n_2}(\tau_2) = \frac{16\tau_2}{\Gamma(s_1)\Gamma(s_2)} n^{s_1 + s_2 - 1} \sigma_{1-2s_1}(n_1) \sigma_{1-2s_2}(n_2) K_{s_1 - \frac{1}{2}}(2\pi n_1\tau_2) K_{s_2 - \frac{1}{2}}(2\pi n_2\tau_2) + (s_1 \leftrightarrow s_2) .$$

$$(4.26)$$

The terms $\hat{g}_{n,0}(\tau_2)$ and $\hat{g}_{0,n}(\tau_2)$ in (4.25) originate from contributions to the source term coming from the product of the *n*-instanton sector of $E(s_1; \tau)$ and the perturbative expansion of $E(s_2; \tau)$ (together with the terms with $s_1 \leftrightarrow s_2$) while $\hat{g}_{n_1,n_2}(\tau_2)$ in (4.26) comes from contributions from the product of the n_1 -instanton sector of $E(s_1; \tau)$ with the n_2 -instanton sector of $E(s_2; \tau)$.

¹²As mentioned in footnote 7, the case with either s_1 or s_2 equal to $\frac{1}{2}$ is slightly different from the other half-integer cases since the Fourier zero-mode sector does contain a term proportional to $\log(\tau_2)$. Although our present discussion encompasses this case as well, these particular modular invariant functions do not seem to be relevant when discussing SMFs.

Equation (4.25) can be solved by standard methods, while (4.26) can be brought into a nicer form by writing it as

$$\hat{g}_{n_1,n_2}(\tau_2) = \frac{16}{\Gamma(s_1)\Gamma(s_2)} n_1^{-s_1 + \frac{1}{2}} n_2^{-s_2 + \frac{1}{2}} \sigma_{2s_1 - 1}(n_1) \sigma_{2s_2 - 1}(n_2) G(2\pi n_1, 2\pi n_2, \tau_2), \qquad (4.27)$$

where we have used $\sigma_{-\alpha}(n) = n^{-\alpha} \sigma_{\alpha}(n)$. The auxiliary function $G(n_1, n_2, y)$ solves:

$$\left[y^2 \partial_y^2 - \left((n_1 + n_2)y\right)^2 - s(s-1)\right] G(n_1, n_2, y) = y K_{s_1 - \frac{1}{2}}(n_1 y) K_{s_2 - \frac{1}{2}}(n_2 y).$$
(4.28)

The results in [23] lead to a particular solution of the form:

$$G(n_1, n_2, y) = \sum_{i,j=0}^{1} \eta_{ij}(n_1, n_2, y) K_i(n_1 y) K_j(n_2 y), \qquad (4.29)$$

with $\eta_{ij}(n_1, n_2, y)$ a rational function depending on the parameters s_1, s_2 and s_2 . The key question now is whether this particular solution of (4.1) is modular invariant and is therefore a GES. In order to answer this question we recall an important lemma proved in [37].

Lemma. If $F(\tau)$ is an $SL(2,\mathbb{Z})$ invariant function on the upper half-plane such that at the cusp $\tau_2 \to \infty$, with $\tau_2 = Im \tau$, it satisfies the growth condition $F(\tau) = O(\tau_2^w)$ with w > 1, then each of its Fourier modes $F_n(\tau_2) = \int_0^1 F(\tau) e^{-2\pi i n \tau_1} d\tau_1$ satisfies the bound $F_n(\tau_2) = O(\tau_2^{1-w})$ in the limit $\tau_2 \to 0$. It is important to note that in the special case where the modular invariant function $F(\tau)$ satisfies the growth condition $F(\tau) = O(\tau_2)$, we can obtain the slightly weaker bound $F_n(\tau_2) = O(\tau_2^{-\epsilon})$ in the limit $\tau_2 \to 0$ for all $\epsilon > 0$.

The proof of this lemma is based on the fact that a cuspidal growth of order τ_2^w implies that the modular invariant function $F(\tau)$ must be bounded by $E(w;\tau)$ on the whole upper half-plane and since for small τ_2 we have $E(w;\tau) = O(\tau_2^{1-w})$, then the same bound must hold for $F_n(\tau)$.

We will now argue that thanks to the result of [22], the particular solution constructed in [23] and given by the solution of (4.27)-(4.29) violates the Lemma whenever the eigenvalue s is such that dim $S_{2s} \neq 0$ and hence it is not a modular invariant solution to (4.1). However, by adding a suitable multiple of exactly the same solution of the homogeneous equation (4.14) previously considered, we will be able to construct a modular invariant solution to (4.1) for the spectrum (4.23).

From the asymptotic expansion as $\tau_2 \to \infty$ presented in (4.3), we see that the GES $\mathcal{E}(s; s_1, s_2; \tau)$ grows at the cusp as $\tau_2^{s_1+s_2} = \tau_2^w$. Hence from the above Lemma we know that the behaviour of any of its Fourier modes at the origin is bounded by τ_2^{1-w} . In the zero-mode sector we immediately see that the perturbative solution of the homogeneous equation τ_2^{1-s} violates the growth condition at the origin for the spectrum (4.23) since $s \ge w + 1$. However, the zero-mode sector also contains the infinite tower (4.5) of instanton/anti-instanton terms. While these terms are are exponentially suppressed at the cusp $\tau_2 \gg 1$, it is possible to show either by direct calculation [37, 32] or via $\mathrm{SL}(2,\mathbb{Z})$ spectral theory [33] that near the origin $\tau_2 \to 0$ the sum of instanton/anti-instanton terms behaves as τ_2^{1-s} and cancels the unwanted power of τ_2^{1-s} coming from the perturbative part, thus proving that in the zero-mode sector the growth condition implied by the Lemma is indeed respected.

We now turn to consider the small- τ_2 behaviour of the non-zero mode part, which is given by

$$\int_{0}^{1} \mathcal{E}_{\mathbf{p}}(s; s_{1}, s_{2}; \tau) e^{-2\pi i n \tau_{1}} d\tau_{1} = \hat{g}_{n,0}(\tau_{2}) + \hat{g}_{0,n}(\tau_{2}) + \sum_{\substack{n_{1}+n_{2}=n\\n_{1},n_{2}\neq 0}} \hat{g}_{n_{1},n_{2}}(\tau_{2}) .$$
(4.30)

The small- τ_2 limits of $\hat{g}_{n,0}(\tau_2)$ and $\hat{g}_{0,n}(\tau_2)$ are easy to obtain from the differential equation (4.25), but the small- τ_2 limit of the infinite sum of the $\hat{g}_{n_1,n_2}(\tau_2)$ contributions is more complicated. Firstly, we notice from

(4.29) that since the modified Bessel functions $K_0(x)$, $K_1(x)$ decrease exponentially fast for large and positive x (4.9), the sum over $n_1 + n_2 = n$ appearing in (4.30) is absolutely convergent for all $n \neq 0$. Hence we can first expand $\hat{g}_{n_1,n_2}(\tau_2)$ for small- τ_2 before performing the sum over $n_1 + n_2 = n$. Using the results of [23] the expansion of the particular solution $\hat{g}_{n_1,n_2}(\tau_2)$ given by (4.27)-(4.29) in the limit $\tau_2 \to 0$ with $n_1 + n_2 = n$ and $n_1, n_2 \neq 0$ is given by

$$\hat{g}_{n_1,n_2}(\tau_2) \stackrel{\tau_2 \to 0}{\sim} R_{s;s_1,s_2} \tau_2^{1-s} \frac{1}{n^{s+s_1+s_2-1}} \Big[\sigma_{2s_1-1}(n_1) \sigma_{2s_2-1}(n_2) Q_{s-s_1-s_2}^{2s_1-1,2s_2-1} \Big(\frac{n_2-n_1}{n_2+n_1} \Big) \Big] + O(\tau_2^{3-s}) , \quad (4.31)$$

where $Q_d^{r_1,r_2}$ is a Jacobi function of the second kind and the coefficients $R_{s;s_1,s_2}$ are given by:

$$R_{s;s_1,s_2} \coloneqq 4 \, \frac{\Gamma\left(\frac{s-s_1-s_2+1}{2}\right) \Gamma\left(\frac{s+s_1+s_2-1}{2}\right)}{\pi^s \, \widetilde{C}_{s;s_1,s_2}} \,, \tag{4.32}$$

and the coefficients $\widetilde{C}_{s;s_1,s_2}$ have been defined in (3.49). Although we do not have a proof of this expression, we have checked its validity by computing the particular solution (4.29) using the methods of [23] for a multitude of values for the parameters s, s_1, s_2 in the spectrum (4.23).

Furthermore, although (4.31) only displays the leading power of the small- τ_2 expansion of $\hat{g}_{n_1,n_2}(\tau_2)$ we have checked that, for the range of parameters s, s_1, s_2 in the spectrum (4.23), the particular solution \hat{g}_{n_1,n_2} does produce all power-like terms $\tau_2^{-\ell}$ in the range $\ell \in \{w, w + 2, \ldots, s - 3, s - 1\}$ with $w = s_1 + s_2$ that violate the boundary condition at small- τ_2 . Importantly, these "unwanted" terms can be all written in the form (4.31) with exactly the same prefactor function of n_1, n_2 multiplying a polynomial in inverse powers of $(n\tau_2)$ that is identical to the small- τ_2 expansion of $\sqrt{n\tau_2} K_{s-1/2}(2\pi n\tau_2)$ given in (4.10).

To check whether the complete n^{th} Fourier mode (4.30) does in fact violate the Lemma we still need to perform the sum over $n_1 + n_2 = n$ in (4.31). To this end we utilise the following crucial theorem.

Theorem FKR. (Fedosova, Klinger-Logan, Radchenko [22]) For any $d \ge 1$ and $r_1, r_2 \in 2\mathbb{Z}_{>0}$ we have:

$$\sum_{\substack{n_1+n_2=n\\n_1,n_2\neq 0}} \sigma_{r_1}(n_1)\sigma_{r_2}(n_2)Q_d^{r_1,r_2}\left(\frac{n_2-n_1}{n_2+n_1}\right) = (-1)^d C_d^{r_1,r_2}(n)\sigma_{r_1}(n) - C_d^{r_2,r_1}(n)\sigma_{r_2}(n) + \frac{\tilde{a}(n)}{n^d}, \qquad (4.33)$$

where

$$C_d^{r_1,r_2}(n) = \frac{(r_2-1)!(r_1+d)!}{2(r_1+r_2+d)!}\zeta(r_2)n^{r_2} + \binom{d+r_2}{d}\frac{\zeta'(-r_2)}{2},$$
(4.34)

while $\tilde{\Delta} = \sum_{n=1}^{\infty} \tilde{a}(n)q^n \in \mathcal{S}_k$ is a holomorphic cusp form of weight k:

$$k = 2d + r_1 + r_2 + 2, (4.35)$$

given by $\tilde{\Delta}(\tau) = \sum_{\Delta \in S_k} \mu'_{\Delta} \Delta$ where Δ runs over normalised Hecke eigenforms in S_k and

$$\mu_{\Delta}' = \frac{\pi(-1)^{d+\frac{r_2}{2}+1}}{2^k} \binom{k-2}{d} \frac{\Lambda(\Delta, d+1)\Lambda(\Delta, r_1+d+1)}{\langle \Delta, \Delta \rangle} \,. \tag{4.36}$$

If we specialise this theorem to the present case (4.31) we have $r_1 = 2s_1 - 1$, $r_2 = 2s_2 - 1$ while $d = s - s_1 - s_2$. Hence in our case *L*-values of holomorphic cusp forms will appear whenever

$$k = 2d + r_1 + r_2 + 2 = 2(s - s_1 - s_2) + (2s_1 - 1) + (2s_2 - 1) + 2 = 2s,$$
(4.37)

with dim $S_k \neq 0$. We then apply this theorem to the infinite sum over $n_1 + n_2 = n$ of the small- τ_2 expansion (4.31) for the particular solution \hat{g}_{n_1,n_2} to deduce the schematic form:

$$\sum_{\substack{n_1+n_2=n\\n_1,n_2\neq 0}} \hat{g}_{n_1,n_2}(\tau_2) \overset{\tau_2\to 0}{\sim} \left[\frac{\zeta(2s_1)+\zeta(2s_2)}{n^{s+s_1+s_2-1}} + \frac{\zeta(2s_1-1)}{n^{s+s_2-s_1}} + \frac{\zeta(2s_2-1)}{n^{s+s_1-s_2}} + \frac{\tilde{a}(n)}{n^{2s-1}} \right] \tau_2^{1-s} + O(\tau_2^{3-s}).$$
(4.38)

Substituting this behaviour in the full *n*-instanton sector (4.30) we see that the terms proportional to Riemann zetas cancel against the same terms coming from the small- τ_2 expansion of $\hat{g}_{n,0} + \hat{g}_{0,n}$, in exactly the same fashion as for the zero-mode sector. However, when dim $S_{2s} \neq 0$ we see that the power behaved term proportional to $\tilde{a}(n)\tau_2^{1-s}$ does not vanish and therefore violates the growth condition coming from the Lemma. We deduce that whenever the eigenvalue *s* is such that dim $S_{2s} \neq 0$ the particular solution constructed using the methods of [23] cannot possibly be a GES.

However, although the small- τ_2 limit of the n^{th} Fourier mode contains power-like terms $\tau_2^{-\ell}$ in the range $\ell \in \{w, w+2, \ldots, s-3, s-1\}$ with $w = s_1 + s_2$, which all violate the Lemma, as mentioned below equation (4.31) they all multiply the same overall factor $\tilde{a}(n)/n^s$ and their relative coefficients coincide with the small- τ_2 expansion of $\sqrt{n\tau_2}K_{s-1/2}(2\pi n\tau_2)$. This tells us that if we add to the particular solution a suitable multiple of the solution of the homogeneous equation $H_{\tilde{\Delta}}(\tau)$ given in (4.14) for the holomorphic cusp form $\tilde{\Delta}$ defined in Theorem FKR, we can cancel all of the unwanted powers of τ_2 in (4.38) that do not satisfy the Lemma without spoiling the differential equation (4.1), i.e. for the spectrum (4.23) we are led to the solution

$$\mathcal{E}(s; s_1, s_2; \tau) = \mathcal{E}_{p}(s; s_1, s_2; \tau) + [\widetilde{C}_{s; s_1, s_2}]^{-1} \sum_{\Delta \in \mathcal{S}_{2s}} \lambda'_{\Delta}(s; s_1, s_2) H_{\Delta}(\tau) , \qquad (4.39)$$

where $C_{s;s_1,s_2}$ is given in (3.49) and the coefficients λ'_{Δ} are obtained from (4.36)

$$\lambda_{\Delta}'(s;s_1,s_2) = (-1)^{\frac{s+s_1-s_2+2}{2}} \frac{\Gamma(s)\Gamma\left(\frac{w-s}{2}\right)}{2^{2s-3}\Gamma\left(\frac{w+s}{2}\right)} \frac{\Lambda(\Delta,s+s_1-s_2)\Lambda(\Delta,s+w-1)}{\langle\Delta,\Delta\rangle}, \qquad (4.40)$$

with $w = s_1 + s_2$. Using the functional equation (C.8) for the completed *L*-function it is easy to show that λ'_{Δ} is symmetric under $s_1 \leftrightarrow s_2$. Importantly, when compared with the identical conjectural coefficients (4.18) for the case (4.13) relevant for MGFs, we see that λ'_{Δ} for the spectrum (4.23) is now always given by the product of two critical even *L*-values.

Crucially, the solution (4.39) produces the GES solution to (4.1) with spectrum (4.23). The proof of this statement follows again from the Lemma. Firstly, recall that the particular solution defined in (4.39) was constructed so as not to violate the growth condition at the origin. While this growth condition is a necessary condition for the modular invariance of $\mathcal{E}(s; s_1, s_2; \tau)$ in this case it turns out to be sufficient as well. This can be seen as follows.

Firstly, we can use $SL(2, \mathbb{Z})$ spectral methods to show that the modular invariant GES solution to (4.1) for the spectrum of parameters (4.23) does in fact exist [30]. Suppose now that the solution (4.39) is not modular invariant and therefore differs from the GES solution, even though it respects the growth condition at the origin for any of its Fourier modes. Since both (4.39) and the GES solution solve the same differential equation (4.1) they must differ by a solution to the homogeneous equation. However, it is not possible to add any such solution to (4.39) without spoiling the growth condition at the origin. In the zero-mode sector this would amount to either introducing a τ_2^s term or changing the coefficient (4.4) of the τ_2^{1-s} term, neither of which is allowed. In the non-zero mode sector, this would amount to adding a new term of the form (4.12). However, since we know that the solution (4.39) grows like $O(\tau_2^{1-w})$ at the origin, adding a new solution to the homogeneous equation to the homogeneous equation to the homogeneous equation to the limit $\tau_2 \to 0$, thus spoiling the necessary growth condition coming from the Lemma. We conclude that (4.39) must be the modular invariant GES solution to (4.1) for the SMF spectrum of parameters (4.23).

A similar argument regarding the necessity of having to add a solution of the homogeneous equation can be made for the GES relevant for the MGFs, with spectrum given in (4.13). Firstly, note that for a GES $\mathcal{E}(s; s_1, s_2; \tau)$ with parameters satisfying (4.13) we have that a term proportional to τ_2^{1-s} , originating from the particular solution constructed, does not violate the growth condition at the origin of the form $\tau_2^{1-s_1-s_2}$ dictated by the Lemma. However, we simply need to consider a more refined version of the Lemma where rather than $\mathcal{E}(s; s_1, s_2; \tau)$, we analyse $F = \mathcal{E}(s; s_1, s_2; \tau) + \alpha E(s_1 + s_2; \tau)$ where α is chosen in such a way as to cancel the leading $\tau_2^{s_1+s_2}$ cuspidal growth. The auxiliary function F has now the tamer growth at the cusp $\tau_2^{s_1+1-s_2}$ (without loss of generality we may assume $s_1 \geq s_2$), thus implying a behaviour at the origin bounded by $\tau_2^{s_2-s_1}$. If $s_1 > s_2$ we can furthermore construct yet another "improved" modular invariant function by considering $F' = F + \beta E(s_1 - s_2 + 1; \tau)$ with β chosen as to cancel the cuspidal behaviour $\tau_2^{s_1+1-s_2}$ so that $F' = O(\tau_2)$ at the cusp. We can then use the Lemma and bound F' by the Eisenstein series $E(1+\epsilon;\tau)$ for all $\epsilon > 0$. For the diagonal case, $s_1 = s_2$, we have $F = O(\tau_2)$ at the cusp and this is directly bounded by $E(1+\epsilon;\tau)$ for all $\epsilon > 0$. As a consequence, we deduce that every Fourier modes must diverge at the origin at most as $\tau_2^{-\epsilon}$. The particular solutions constructed in [9] via iterated integrals of holomorphic Eisenstein series do not always satisfy this bound, hence they are not in general modular invariant functions and a suitable multiple of the homogenous solution (4.15) must be added so as to obtain the GES solution to (1.1) with spectrum (4.13).

It should be stressed that there is no analogue of Theorem FKR for the GESs with integer s_1 and s_2 , i.e., with the spectrum (4.13), and so far the only way to fix the coefficient of the solution of the homogeneous equation so as to construct a modular invariant solution to the inhomogeneous Laplace equation has relied on the analysis of [9, 10]. In addition to the asymptotic argument just presented, we believe that there should be a more general version of Theorem FKR responsible for the validity of Conjecture 1.

Returning to the case of SMFs, we see that the first instance where L-values of holomorphic cusp forms appear in the GESs (4.39) is again for eigenvalue s = 6 where we know that S_{12} is a one-dimensional vector spaced spanned by Ramanujan cusp form Δ_{12} . From the spectrum (4.23) we see that an eigenvalue s = 6necessitates weight $w = s_1 + s_2 = 3$ or w = 5. In these cases the GESs are given by

$$\begin{aligned} \mathcal{E}(6; \frac{1}{2}, \frac{5}{2}; \tau) &= \mathcal{E}_{p}(6; \frac{1}{2}, \frac{5}{2}; \tau) - \frac{625}{6048} \cdot \frac{\Lambda(\Delta_{12}, 10)^{2}}{\pi\langle\Delta_{12}, \Delta_{12}\rangle} H_{\Delta_{12}}(\tau) \,, \\ \mathcal{E}(6; \frac{3}{2}, \frac{3}{2}; \tau) &= \mathcal{E}_{p}(6; \frac{3}{2}, \frac{3}{2}; \tau) + \frac{25}{72} \cdot \frac{\Lambda(\Delta_{12}, 10)^{2}}{\pi\langle\Delta_{12}, \Delta_{12}\rangle} H_{\Delta_{12}}(\tau) \,, \\ \mathcal{E}(6; \frac{1}{2}, \frac{9}{2}; \tau) &= \mathcal{E}_{p}(6; \frac{1}{2}, \frac{9}{2}; \tau) - \frac{32}{6615} \cdot \frac{\Lambda(\Delta_{12}, 10)^{2}}{\pi\langle\Delta_{12}, \Delta_{12}\rangle} H_{\Delta_{12}}(\tau) \,, \\ \mathcal{E}(6; \frac{3}{2}, \frac{7}{2}; \tau) &= \mathcal{E}_{p}(6; \frac{3}{2}, \frac{7}{2}; \tau) + \frac{10}{189} \cdot \frac{\Lambda(\Delta_{12}, 10)^{2}}{\pi\langle\Delta_{12}, \Delta_{12}\rangle} H_{\Delta_{12}}(\tau) \,, \\ \mathcal{E}(6; \frac{5}{2}, \frac{5}{2}; \tau) &= \mathcal{E}_{p}(6; \frac{5}{2}, \frac{5}{2}; \tau) - \frac{8}{81} \cdot \frac{\Lambda(\Delta_{12}, 10)^{2}}{\pi\langle\Delta_{12}, \Delta_{12}\rangle} H_{\Delta_{12}}(\tau) \,, \end{aligned}$$

where, again, we have used Manin's relations to express the results in terms of $\Lambda(\Delta_{12}, 10)$ only. As was the case with (4.22) the seemingly random rational numbers appearing in front of the solutions of the homogeneous equation will shortly prove to be rather special.

To summarise, in this section we have shown that for both spectra (4.13) and (4.23) (relevant to MGFs and SMFs, respectively) it is possible to find GES solutions of the form (4.15) and (4.39). In either case the modular invariant solution is obtained by first constructing a particular solution whose Fourier modes have asymptotic expansions at the cusp with coefficients that are quadratic forms in Riemann zeta values. Furthermore, in each case it is necessary to add a suitable multiple of the solution of the homogeneous equation that contributes only to the (0, n) and (n, 0) instanton sector and has an asymptotic expansion at the cusp with coefficients controlled by the q-series coefficients of holomorphic cusp forms and their associated critical and non-critical L-values.

4.3 Absence of *L*-values in theta-lifted local Maass functions

Although there is no known lattice representation for an individual GES we now use the results of section 3 to show that the lattice sum constructed as a theta lift of a local Maass function (2.32) produces special linear combinations of GESs for both MGFs and SMFs. Furthermore, these four-dimensional lattice sum representations produce precisely the rational linear combinations of GESs (4.15) and (4.39) for which the cuspidal contribution H_{Δ} cancels. Once again, we will present the cases of MGFs and SMFs separately.

Two-loop Modular Graph Functions

Using the results of section 3, we see that if we want to obtain linear combinations of GESs with parameters in the spectrum (4.13) relevant for MGFs we have to consider the theta lift (2.32), or equivalently (2.34), for $w \ge 4$ and $1 \le s \le w - 2$ with the same parity. As a consequence we see from (3.13) that the integrand (2.35) reduces to a symmetric polynomial in t_i of degree w - 3.

It is then straightforward to perform the integration (2.34) using the basic integral representation of a MGF given in (2.39). We deduce that the theta lift (2.32) of a local Maass function with $w \ge 4$ and $1 \le s \le w-2$ produces a rational linear combination of MGFs ($C_{a,b,c}(\tau)$ with w = a + b + c), that solves an inhomogeneous Laplace equation with eigenvalue s and source terms given by bilinears, $E(s_1; \tau)E(s_2; \tau)$, in non-holomorphic Eisenstein series with $w = s_1 + s_2$ and possibly a multiple of the single non-holomorphic Eisenstein series $E(w; \tau)$.

Moreover, the dimension of the vector eigenspace $\mathcal{V}_C(w, s)$ of MGFs of a given weight w = a + b + c and eigenvalue s was determined in [8] to be

$$\dim \mathcal{V}_C(w,s) = \begin{cases} \left\lfloor \frac{s+2}{3} \right\rfloor & \text{for } 1 \le s \le w-2 \text{ and } s, w \text{ of same parity,} \\ 0 & \text{otherwise.} \end{cases}$$
(4.42)

Note that this matches the total number of functions $B_{i,j}^w$ defined in (2.35) consistent with the spectrum (4.13) given that s = 3i + j + 1. Hence we see that the theta lift off local Maass functions with $w \ge 3$ and $1 \le s \le w - 2$ results in the vector space of $C_{a,b,c}(\tau)$ conveniently arranged in terms of eigenvectors of the Laplace operator.

However, a similar calculation of the number of independent GESs $\mathcal{E}(s; s_1, s_2; \tau)$ with spectrum (4.13) gives

$$\dim \mathcal{V}_{\mathcal{E}}(w,s) = \begin{cases} \left\lfloor \frac{s}{2} \right\rfloor & \text{for } 2 \le s \le w-2 \text{ and } s, w \text{ of the same parity,} \\ 0 & \text{otherwise,} \end{cases}$$
(4.43)

where $w = s_1 + s_2$. Note that $\mathcal{V}_C(w, s)$ includes the eigenvalue s = 1, which occurs when w is odd. In this case, the Laplace equation reduces in complexity and the corresponding eigenvector becomes a non-holomorphic Eisenstein series plus a constant. This special case with s = 1 corresponds to the theta lifts of $B_{0,0}^w$ for $w \ge 3$. For example, when w = 3 we have $B_{0,0}^3(t) = 1$ and

$$\mathcal{E}_{0,0}^{3}(\tau) = \sum_{\substack{p_1, p_2, p_3 \in \Lambda'\\ p_1 + p_2 + p_3 = 0}} \int_{(\mathbb{R}^+)^3} \exp\left(-\frac{\pi}{\tau_2} \sum_{j=1}^3 t_j |p_j|^2\right) \mathrm{d}^3 t = C_{1,1,1}(\tau) = E(3;\tau) + \zeta(3).$$
(4.44)

Comparing the dimensions (4.42) and (4.43) we see that there are in general more GESs $\mathcal{E}(s; s_1, s_2; \tau)$ with spectrum (4.13) than the $C_{a,b,c}(\tau)$. As noticed in [9] for $s \geq 2$ the difference in dimensions gives strikingly,

$$\dim \mathcal{V}_{\mathcal{E}}(w,s) - \dim \mathcal{V}_{C}(w,s) = \dim \mathcal{S}_{2s}.$$
(4.45)

Since dim $S_{2s} = 0$ for s < 6 and since s < w - 2 it follows that at weight w = 8 and at any weight $w \ge 10$, the number of independent GESs is strictly larger than that of $C_{a,b,c}(\tau)$. More precisely, the number of GESs at weight w and eigenvalue s in the MGF spectrum (4.13) that cannot be written in terms of MGFs equals the number of holomorphic cusp forms at modular weight 2s (which is given in (4.16)).

The first case for which this mismatch starts playing a role is s = 6, for which S_{12} is the one-dimensional vector space $S_{12} = \text{span}\{\Delta_{12}\}$, and necessarily $w \ge 8$. From the spectrum (4.13) we see that for (s, w) = (6, 8) there are three GESs, $\mathcal{E}(6; 2, 6; \tau), \mathcal{E}(6; 3, 5; \tau)$ and $\mathcal{E}(6; 4, 4; \tau)$, each of which can be written in terms of iterated integrals of holomorphic Eisenstein series of depth at most two and crucially a non-zero homogeneous contribution originating from an iterated integral of the Ramanujan cusp form Δ_{12} as seen in (4.17)-(4.22).

Conversely, there are only two theta-lifted local Maass functions (2.34) corresponding to MGFs with weight 8 and eigenvalue 6 given by $\mathcal{E}_{1,2}^8(\tau)$ and $\mathcal{E}_{0,5}^8(\tau)$. Thanks to the results of section 3, we have

$$\begin{aligned} \mathcal{E}_{1,2}^{8}(\tau) &= \sum_{\substack{p_1, p_2, p_3 \neq 0\\ p_1 + p_2 + p_3 = 0}} \int_0^\infty \frac{\left(7\sigma_1{}^3\sigma_2 - 63\sigma_1{}^2\sigma_3 + 3\sigma_1\sigma_2{}^2 + 21\sigma_2\sigma_3\right)}{63} \exp\left(-\frac{\pi}{\tau_2} \sum_{i=1}^3 t_i |p_i|^2\right) \mathrm{d}^3t \\ &= -\frac{24}{7} \left[112\,\mathcal{E}(6; 3, 5; \tau) + 75\,\mathcal{E}(6; 4, 4; \tau)\right] - \frac{436}{7}\,E(8; \tau) \\ &= \frac{8}{7} \left[14\,C_{1,2,5}(\tau) + 24\,C_{1,3,4}(\tau) - 2\,C_{2,2,4}(\tau) - C_{2,3,3}(\tau)\right], \end{aligned}$$
(4.46)

where the notation in the first line utilises the basis of symmetric polynomials in three variables $\sigma_1 = t_1 + t_2 + t_3$, $\sigma_2 = t_1 t_2 + t_1 t_3 + t_2 t_3$ and $\sigma_3 = t_1 t_2 t_3$. Similarly we find,

$$\mathcal{E}_{0,5}^{8}(\tau) = \sum_{\substack{p_1, p_2, p_3 \neq 0\\p_1+p_2+p_3=0}} \int_0^\infty \frac{\sigma_1 \left(21\sigma_1^4 - 70\sigma_1^2\sigma_2 + 45\sigma_2^2\right)}{21} \exp\left(-\frac{\pi}{\tau_2} \sum_{i=1}^3 t_i |p_i|^2\right) \mathrm{d}^3 t$$

$$= -\frac{180}{7} \left[140 \,\mathcal{E}(6; 2, 6; \tau) + 112 \,\mathcal{E}(6; 3, 5; \tau) + 51 \,\mathcal{E}(6; 4, 4; \tau)\right] + \frac{87270}{91} \,\mathcal{E}(8; \tau) \qquad (4.47)$$

$$= \frac{60}{7} \left[42 \,C_{1,1,6}(\tau) + 28 \,C_{1,2,5}(\tau) + 18 \,C_{1,3,4}(\tau) + 2 \,C_{2,2,4}(\tau) + C_{2,3,3}(\tau)\right].$$

At this point it is straightforward to substitute (4.22) in the above relations and check that indeed the solution of the homogeneous equation involving *L*-values of holomorphic cusp forms do drop out and both $\mathcal{E}_{1,2}^8(\tau)$ and $\mathcal{E}_{0,5}^8(\tau)$ can be written as linear combinations of iterated integrals involving solely holomorphic Eisenstein series (and their complex conjugate).

This phenomenon whereby the holomorphic cusp form contributions to GESs conspire to cancel out in linear combinations ultimately given by lattice sums corresponding to MGFs had been studied in [9, 10]. These magical cancellations provide the first hint of a beautiful algebraic structure at play behind the scenes. To better understand this story we need to consider a wider class of non-holomorphic modular covariant forms introduced by Francis Brown. The intriguing mathematical properties of MGFs stimulated Brown's construction of an infinite family of non-holomorphic modular forms dubbed equivariant iterated Eisenstein integrals [5, 6, 7] but the connection between MGFs and equivariant iterated Eisenstein integrals is indirect, see e.g. [38, 39].

In [6, 7] Brown has shown how to construct non-holomorphic forms as combinations of non-modular invariant iterated integrals of holomorphic Eisenstein series and their complex conjugates. However, not every equivariant iterated integral corresponds to a higher-loop MGF. Notably, in the Fourier expansion of modular graph forms we encounter only a very small subset of the vast number of multiple modular values

given by single-valued multiple zeta values and, in particular, we do not encounter completed L-values of holomorphic cusp forms, which can instead be present in a given equivariant iterated integral. However, if we consider the generating series of all modular graph forms constructed in [40, 41] we see that only special linear combinations of Brown's equivariant iterated integrals correspond to modular graph forms.

To extract a particular MGF from the generating series we simply need to extract the coefficient of certain non-commutative sign-post variables, which furnish a representation of an important mathematical object called Tsunogai's derivation algebra [42]. As a consequence of this fact, we find that MGFs are related only to specific linear combinations of equivariant iterated Eisenstein integrals, namely those that follow from the so-called Tsunogai's relations [42, 43], see also [7, 41, 10, 38, 44]. While the modular completion of iterated integrals of holomorphic Eisenstein series necessitates the addition of iterated integrals of holomorphic cusp forms [5, 7], such as for the depth-two examples (4.17), the linear combinations of equivariant iterated integrals selected by Tsunogai's relations and hence corresponding to MGFs are expressible in terms of iterated integrals involving solely holomorphic Eisenstein series (and their complex conjugates) and in addition the periods arising in this construction are conjecturally restricted solely to single-valued MZVs.

As a concrete consequence of this structure, we find that the theta lift of modular local polynomials corresponding to the spectrum (4.13) produces precisely the linear combinations of GESs in (4.46)-(4.47), or equivalently of depth-two equivariant iterated Eisenstein integrals for which the cusp form contributions have dropped out. We expect that a generalisation of (2.32) to a higher-dimensional lattice sum will produce higher-loop MGFs and hence the corresponding linear combinations of higher-depth iterated Eisenstein integrals selected by Tsunogai's relations.

S-dual Modular Functions

We now turn to consider the theta lift of local Maass functions giving rise to linear combinations of GESs with the spectrum (4.23), in which case s_1 and s_2 are half-integral. From the general structure (3.30), we see that in order to obtain linear combinations of GESs with this spectrum we need to consider the case where w and s have opposite parity. Furthermore, since in general $i + j \leq s - 1$ where s = 3i + j + 1, we see that the indices s_1, s_2 of the GESs $\mathcal{E}(s; s_1, s_2; \tau)$, which appear in (3.30) always lie between $(w - s + 2)/2 \leq s_1, s_2 \leq (w + s - 2)/2$. Hence in order to obtain the spectrum (4.23) where $s \geq w + 1$, some of these GESs might have negative indices. This is however not a problem since if s_1 is a negative half-integer the functional relation (3.33) can be used to consider instead $\mathcal{E}(s; 1 - s_1, s_2; \tau)$. As previously mentioned, this comes at the price of spoiling uniform transcendentality in (3.30).

We can now analyse the vector space of theta-lifted Maass functions at fixed weight w and eigenvalue s, as we did earlier for MGFs. From the analysis of the previous section we expect that the lowest eigenvalue for which some cusp form might appear to be s = 6, which from (4.23) requires either w = 3 or w = 5. This leads us to consider the theta lifts $\mathcal{E}_{1,2}^w(\tau), \mathcal{E}_{0,5}^w(\tau)$ with $w \in \{3,5\}$. It is now a matter of exploiting (3.30) combined with (3.31)-(3.32) to obtain

$$\mathcal{E}_{1,2}^{3}(\tau) = \sum_{\substack{p_1, p_2, p_3 \neq 0\\ p_1 + p_2 + p_3 = 0}} \int_0^\infty \frac{\left(7\sigma_1{}^3\sigma_2 - 63\sigma_1{}^2\sigma_3 + 3\sigma_1\sigma_2{}^2 + 21\sigma_2\sigma_3\right)}{63\sigma_2^{\frac{5}{2}}} \exp\left(-\frac{\pi}{\tau_2}\sum_{i=1}^3 t_i |p_i|^2\right) \mathrm{d}^3t$$
$$= -\frac{\pi}{14} \left[84 \,\mathcal{E}(6; \frac{1}{2}, \frac{5}{2}; \tau) + 25 \,\mathcal{E}(6; \frac{3}{2}, \frac{3}{2}; \tau)\right] - \frac{\pi}{2} \,E(3; \tau) \,, \tag{4.48}$$
$$\mathcal{E}_{0,5}^{3}(\tau) = \sum_{\substack{p_1, p_2, p_3 \neq 0\\ p_1 + p_2 + p_3 = 0}} \int_0^\infty d^3t \, \frac{\sigma_1 \left(21\sigma_1{}^4 - 70\sigma_1{}^2\sigma_2 + 45\sigma_2{}^2\right)}{21\,\sigma_2^{\frac{5}{2}}} \exp\left(-\frac{\pi}{\tau_2}\sum_{i=1}^3 t_i |p_i|^2\right)$$
$$= -\frac{15\pi}{56} \left[168 \,\mathcal{E}(6; \frac{1}{2}, \frac{5}{2}; \tau) + 34 \,\mathcal{E}(6; \frac{3}{2}, \frac{3}{2}; \tau) + 105 \,\mathcal{E}(6; \frac{3}{2}, \frac{7}{2}; \tau)\right] + \frac{225\pi}{32} \,E(3; \tau) \,, \tag{4.49}$$

with similar equations for $\mathcal{E}_{1,2}^5(\tau), \mathcal{E}_{0,5}^5(\tau)$, which can be expressed as

$$\mathcal{E}_{1,2}^{5}(\tau) = -\frac{15\pi}{56} \left[28 \,\mathcal{E}(6; \frac{3}{2}, \frac{7}{2}; \tau) + 15 \,\mathcal{E}(6; \frac{5}{2}, \frac{5}{2}; \tau) \right] - \frac{125\pi}{56} \,E(5; \tau) \,, \tag{4.50}$$

$$\mathcal{E}_{0,5}^{5}(\tau) = -\frac{45\pi}{112} \left[490 \,\mathcal{E}(6; \frac{1}{2}, \frac{9}{2}; \tau) + 140 \,\mathcal{E}(6; \frac{3}{2}, \frac{7}{2}; \tau) + 51 \,\mathcal{E}(6; \frac{5}{2}, \frac{5}{2}; \tau) \right] - \frac{16347\pi}{224} \,E(5; \tau) \,. \tag{4.51}$$

Note that the GESs on the right-hand side of (4.49) do not have uniform weight 3 but also contain $\mathcal{E}(6; \frac{3}{2}, \frac{\tau}{2}; \tau)$, which arises from inverting the negative index of $\mathcal{E}(6; -\frac{1}{2}, \frac{\tau}{2}; \tau)$ using the functional equation.

Crucially, if we substitute (4.41) into the linear combinations (4.48)-(4.51) we can easily check that the cuspidal contributions to the solution of the homogeneous equations do in fact cancel out. In other words we see that the linear combinations of GESs belonging to the spectrum (4.23) that are selected by the theta lift of local Maass functions are precisely the same as those for which the *L*-values of holomorphic cusp forms are absent. In contrast to the MGF case, in the present case there is no Tsunogai's derivation algebra, which could be responsible for such cancellations. However, it is entirely possible that upon studying higher-dimensional lattice versions of these particular theta lifts, which play the analogue of higher-loop MGFs in the previous section, such a structure might emerge thus justifying this seemingly magical cancellation of holomorphic cusp forms.

The results described in this section provide substantial evidence for the validity of Conjecture 2 presented in section 1.2, which implies that the modular invariant function $\mathcal{E}_{i,j}^w(\tau)$, given by the theta lift of $B_{i,j}^w(t)$ defined in (2.34), is a rational linear combination of GESs, which admits a four-dimensional lattice sum representation and receives no contributions from *L*-values of holomorphic cusp forms. This applies both to MGFs and to SMFs. It is worth stressing that for the case of SMFs, thanks to the result of [22] we have derived a general expression (4.39)-(4.40) for the *L*-value contribution to an arbitrary GES in the spectrum (4.23). Hence what we will now show amounts to a proof of Conjecture 2 specialised to the case of SMFs. However, for the MGFs the expression for the *L*-value contribution (4.20) to an arbitrary GES in the spectrum (4.13) is only conjectural.

Let us consider the theta lift of $B_{i,j}^w(t)$ as defined in (2.34). From the general expression (3.52) we know how to write $\mathcal{E}_{i,j}^w(\tau)$ as a rational linear combination of GESs. Thanks to (3.52) we then deduce that the *L*-value contribution to $\mathcal{E}_{i,j}^w(\tau)$ coming from a single Hecke eigenform $\Delta \in \mathcal{S}_{2s}$, which we denote by $\mathcal{E}_{i,j}^w(\tau)|_{\Delta}$, must take the form

$$\mathcal{E}_{i,j}^{w}(\tau)\Big|_{\Delta} = 3\sum_{k=0}^{2i}\sum_{\ell=0}^{j}\sum_{m=0}^{\ell}\left[1 - (-1)^{k+\ell+m}\right]C_{i,j;k,\ell,m}\,\widetilde{C}_{s;s_1,s_2}\,\mathcal{E}(s;s_1,s_2;\tau)\Big|_{\Delta},\tag{4.52}$$

with $s_1 = (w+k+\ell+m-i-j)/2$ and $s_2 = [w-(k+\ell+m-i-j)]/2$. For the MGF case, the GESs appearing in the linear combination on the right-hand side of this equation belong to the spectrum of parameters (4.13), while for the SMFs they belong to the spectrum (4.23). Hence, if we want to extract the *L*-value contribution to (4.52) specialised to the MGFs we must use the conjectural expressions (4.20), while for SMFs we have proved that the *L*-value contribution is given by (4.39)-(4.40).

Importantly, our Conjecture 1 states that the *L*-value contribution for GESs with spectrum (4.13) has the same functional dependence on the parameters s, s_1, s_2 as (4.39)-(4.40). Substituting (4.39)-(4.40), we deduce that for both MGFs and SMFs the *L*-value contribution (4.52) takes the form

$$\mathcal{E}_{i,j}^{w}(\tau)\Big|_{\Delta} = 3 \frac{\Gamma(s)\Gamma\left(\frac{w-s}{2}\right)}{2^{2s-4}\Gamma\left(\frac{w+s}{2}\right)} \frac{\Lambda(\Delta, s+w-1)}{\langle \Delta, \Delta \rangle} \mathcal{K}_{i,j}(\Delta) , \qquad (4.53)$$

where we have defined

$$\mathcal{K}_{i,j}(\Delta) \coloneqq \frac{1}{2} \sum_{k=0}^{2i} \sum_{\ell=0}^{j} \sum_{m=0}^{\ell} \left[(-1)^{k+\ell+m} - 1 \right] C_{i,j;k,\ell,m} \left(-1 \right)^{\frac{2i+k+\ell+m+1}{2}} \Lambda(\Delta_{2s}, 2i+k+\ell+m+1), \quad (4.54)$$

and again s = 3i + j + 1. It is important to note that the coefficients $\tilde{C}_{s;s_1,s_2}$ defined in (3.49) and appearing in (4.52) cancel out when we consider the cuspidal contributions (4.20) and (4.39).

We shall prove that $\mathcal{K}_{i,j}(\Delta)$ vanishes identically for all $i, j \in \mathbb{N}$ and for every Hecke eigenforms $\Delta \in \mathcal{S}_{2s}$ with s = 3i + j + 1, i.e.

$$\mathcal{K}_{i,j}(\Delta) = 0, \qquad \forall i, j \in \mathbb{N}, \quad \forall \Delta \in \mathcal{S}_{6i+2j+2}.$$
(4.55)

To prove this statement we proceed as follows. Firstly, we rewrite the L-values in (4.54) using the integral representation discussed in appendix C,

$$\Lambda(\Delta, \ell+1) = i^{-\ell-1} \int_0^{i\infty} \Delta(\tau) \tau^\ell \mathrm{d}\tau \,. \tag{4.56}$$

The sum of completed L-values (4.54) can then be written simply as

$$\mathcal{K}_{i,j}(\Delta) = \frac{1}{2} \int_0^{i\infty} \Delta(\tau) \Big[\sum_{k=0}^{2i} \sum_{\ell=0}^j \sum_{m=0}^\ell \big[(-1)^{k+\ell+m} - 1 \big] C_{i,j;k,\ell,m} \tau^{2i+k+\ell+m+1} \Big] d\tau$$
$$= \frac{1}{2} \int_0^{i\infty} \Delta(\tau) \, \tau^{2i} \left[(\tau - 1)^{2i} (\tau^2 - \tau + 1)^j - (\tau + 1)^{2i} (\tau^2 + \tau + 1)^j \right] d\tau \,, \tag{4.57}$$

where we have used the definition (3.44) for the coefficients $C_{i,j;k,\ell,m}$ originating from having expanded (3.43). The two terms in (4.57) are actually identical as one can easily check by changing the integration variable of the first term to $\tau = S \cdot \tau' = -1/\tau'$ and using the fact that Δ is a modular form of weight 2s = 2(3i + j + 1). We then deduce that (4.57) is identical to

$$\mathcal{K}_{i,j}(\Delta) = \int_0^{i\infty} \Delta(\tau) \, u^i v^j \,\mathrm{d}\tau \,, \tag{4.58}$$

where $u^i v^j$ is precisely the modular local polynomial defined in (2.29) (here a function of the variable τ) whose theta lift $\mathcal{E}^w_{i,i}(\tau)$ we are presently studying.

To prove that (4.58) vanishes identically for all $i, j \in \mathbb{N}$ and for every Hecke eigenforms $\Delta \in \mathcal{S}_{2s}$ with s = 3i + j we simply need to consider the action of the order-three generator of $SL(2,\mathbb{Z})$ given by $U = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$, which maps

$$U \cdot 0 = i\infty$$
, $U \cdot 1 = 0$, $U \cdot i\infty = 1$.

Since the modular local polynomial $u^i v^j$ transforms under U with modular weight -2(3i+j) = -2(s-1)and given that $\Delta \in S_{2s}$ we see that $\Delta(\tau) u^i v^j d\tau$ is invariant under U. We deduce that under the change of integration variables $\tau = U \cdot \tau'$ equation (4.58) becomes

$$\mathcal{K}_{i,j}(\Delta) = \left(\int_0^1 + \int_1^{i\infty} \Delta(\tau) \, u^i v^j \, \mathrm{d}\tau \right)$$
$$= \left(\int_1^{i\infty} + \int_{i\infty}^0 \Delta(\tau') \, u'^i v'^j \, \mathrm{d}\tau',$$
(4.59)

where with $u'^i v'^j$ simply means the same modular local polynomial that was defined in (2.29) in the variable τ' . Similarly, we rewrite (4.58) using the change of integration variables $\tau = U^2 \cdot \tau''$,

$$\mathcal{K}_{i,j}(\Delta) = \left(\int_0^1 + \int_1^{i\infty} \Delta(\tau) \, u^i v^j \, \mathrm{d}\tau\right)$$

$$= \left(\int_{i\infty}^{0} + \int_{0}^{1}\right) \Delta(\tau'') \, u''^{i} v''^{j} \mathrm{d}\tau'' \,, \tag{4.60}$$

Combining (4.59) with (4.60) we deduce that $\mathcal{K}_{i,j}(\Delta) = 0$ for all $i, j \in \mathbb{N}$ and for every Hecke eigenforms $\Delta \in \mathcal{S}_{2s}$ with s = 3i + j + 1. Finally, thanks to (4.53) we conclude that $\mathcal{E}_{i,j}^w(\tau)$ receives no contributions from *L*-values of holomorphic cusp forms contained in each individual GES.

Furthermore, a quick counting argument shows that any rational linear combination of GESs, $\mathcal{E}(s; s_1, s_2; \tau)$, at fixed weight $w = s_1 + s_2$ and eigenvalue s with parameters (s, s_1, s_2) in either (4.13) or (4.23) and for which the cusp form contribution cancels must be a linear combination of theta lifts $\mathcal{E}_{i,j}^w(\tau)$ with s = 3i + j + 1. For the spectrum of parameters (4.13) it is easy to see that at fixed eigenvalue s we have $\lfloor \frac{s}{2} \rfloor$ linearly independent GESs at fixed weight $w = s_1 + s_2$. Similarly, for the spectrum of parameters (4.23) at fixed eigenvalue s the largest number of GESs is found at weight w = s - 1 where there are again $\lfloor \frac{s}{2} \rfloor$ linearly independent GESs. In both cases we can write a general rational linear combination of GESs and impose cancellation of the *L*-values, using either the conjectural expression (4.20) or (4.40), and we obtain a system of dim \mathcal{S}_{2s} equations¹³ in $\lfloor \frac{s}{2} \rfloor$ variables. From the dimension formula (4.16) we see that this system of equations has exactly $\lfloor \frac{s-1}{3} \rfloor$ linearly independent solutions, which is precisely the number of $\mathcal{E}_{i,j}^w(\tau)$ with s = 3i + j + 1.

We conclude this section by noting that the cuspidal contribution defined in (4.54) for arbitrary $i, j \in \mathbb{N}$ as a finite sum over completed *L*-values can be simplified dramatically for the particular infinite families of $\mathcal{E}_{i,j}^w(\tau)$ considered in section 3.1. For example, if we consider the infinite family $\mathcal{E}_{i,0}^w$ with eigenvalue s = 3i+1and integer weight $w \leq s-1$ constructed using (3.24)-(3.25), we find that the vanishing of the corresponding cuspidal contribution $\mathcal{K}_{i,0}(\Delta_{6i+2})$ with $\Delta_{6i+2} \in \mathcal{S}_{6i+2}$ reduces to

$$\mathcal{K}_{i,0}(\Delta_{6i+2}) \propto \sum_{r=-\frac{i-1}{2}}^{\frac{i-1}{2}} (-1)^{r+\frac{i-1}{2}} \frac{\Lambda(\Delta_{6i+2}, (3i+1)+2r)}{\Gamma(i+1+2r)\Gamma(i+1-2r)} \,.$$
(4.61)

The first value of *i* for which the identity $\mathcal{K}_{i,0}(\Delta_{6i+2}) = 0$ becomes non-trivial is i = 3 for which we must consider the unique Hecke normalised cusp form of weight 6i + 2 = 20 and, after using the *L*-function reflection identity (C.8), equation (4.61) for i = 3 reduces to

$$-\frac{\Lambda(\Delta_{20}, 10)}{36} + \frac{\Lambda(\Delta_{20}, 12)}{60} = 0, \qquad (4.62)$$

which is a consequence of Manin's periods theorem [36]. It is easy to show that using the integral representation (4.56), equation (4.61) coincides with the general vanishing expression (4.57) specialised to j = 0.

If we repeat a similar argument for the infinite family $\mathcal{E}_{0,j}^w(\tau)$ with eigenvalue s = j + 1 and weight $w \leq s - 1$ and use (3.21) to compute the cuspidal contribution $\mathcal{K}_{0,j}(\Delta_{2j+2})$ we find

$$\mathcal{K}_{0,j}(\Delta_{2j+2}) \propto \sum_{r=-\frac{j-1}{2}}^{\frac{j-1}{2}} \frac{(-1)^r \,_2 F_1\left(-\frac{j}{2} - r + \frac{1}{2} - \frac{j}{2} + r + \frac{1}{2}; \frac{1}{2} - j|\frac{3}{4}\right) \Lambda(\Delta_{2j+2}, 2r+j+1)}{\Gamma(j-2r+1)\Gamma(j+2r+1)} \,. \tag{4.63}$$

Surprisingly, for this infinite family of linear combination of GESs, the cuspidal contribution (4.63) becomes a rather complicated linear combination of completed *L*-values, although it is the result of the integral representation (4.56) by specialising to i = 0. We have not performed an analytic derivation of this starting from (4.56), however, with the aid of Pari/GP [45] we have numerically verified that (4.63) is satisfied for all cusp forms of weight $2s \leq 100$ within the numerical precision of 10^{-50} utilised.

¹³Since we are working with rational linear combinations of GESs and the cusp form coefficients (4.20) and (4.40) take values in the number field associated with S_{2s} we deduce that a single linear equation with number field coefficients becomes a system of dim S_{2s} equations with rational coefficients.

5 Discussion

We conclude with comments on some possible extension of the current work.

5.1 Cuspidal generalised Eisenstein series

Within the context of the low-energy expansion for genus-one string scattering amplitudes, it is also necessary to consider a different class of odd MGFs [49, 41]. These odd MGFs are again modular invariant functions that can be represented in terms of lattice sums similar to (1.3). However, they are odd under the involution of the upper-half plane $\tau \to -\bar{\tau}$, thus necessarily leading to a cuspidal expansion at $\tau_2 \gg 1$, i.e. rather than having polynomial growth as for the GESs (4.3) their asymptotic expansion at the cusp must necessarily be $O(e^{2\pi i \tau})$. In [9, 10] it was realised that equation (1.1) can be modified so as to obtain a second vector space of cuspidal generalised Eisenstein series,

$$(\Delta_{\tau} - s(s-1))\mathcal{E}^{-}(s;s_1,s_2;\tau) = \frac{\nabla E(s_1;\tau)\overline{\nabla}E(s_2;\tau) - \nabla E(s_2;\tau)\overline{\nabla}E(s_1;\tau)}{2\tau_2^2},$$
(5.1)

with $\nabla := 2i\tau_2^2 \partial_{\tau}$. It is easy to see that the modular invariant solutions to the above equation must be odd under the involution $\mathcal{E}^-(s; s_1, s_2; -\bar{\tau}) = -\mathcal{E}^-(s; s_1, s_2; \tau)$.

In the same way as the GESs with spectrum of parameters (4.13) are related to MGFs, rational combinations of cuspidal GESs with spectrum of parameters

$$s_1, s_2 \in \mathbb{N}$$
, with $s_1, s_2 \ge 2$, $s \in \{|s_1 - s_2| + 1, |s_1 - s_2| + 3, \dots, s_1 + s_2 - 3, s_1 + s_2 - 1\}$, (5.2)

produce all odd MGFs. Importantly, it was shown in [9, 10] that whenever the eigenvalue s, which now has the opposite parity to the weight $w = s_1 + s_2$, is such that dim $S_{2s} \neq 0$, the function $\mathcal{E}^-(s; s_1, s_2; \tau)$ receives a non-vanishing contribution from an odd variant of the solution of the homogeneous equation (4.14). However, compared to (4.20) the coefficient of the solution to the homogeneous equation is proportional to the product of an even non-critical *L*-value times an odd critical one. Cancellation of *L*-values in the rational linear combinations of cuspidal GESs that produce odd MGFs is again a consequence of the same Tsunogai's derivation algebra as discussed in section 4.3 for MGFs.

Although these modular functions have not yet appeared in the physics literature, it is rather natural to ask whether in the odd SMFs case a generalisation of (5.1) to the case of half-integer indices $s_1, s_2 \in \mathbb{N} + 1/2$ and eigenvalue with the same parity as the weight $w = s_1 + s_2$ plays a role analogous to that of (4.23). Firstly, we believe that a suitable modification of Theorem FKR [22] ought to produce an *L*-value contribution which, in parallel to (4.40), should now result in the product of two critical odd values. We are not aware of an expression similar to (4.58) that is responsible for the cancellation of the odd critical *L*-values in the rational linear combinations of cuspidal GESs corresponding to odd MGFs or odd SMFs. Furthermore, even in the case of odd MGFs where a lattice sum representation is available, we do not know whether these objects can be constructed in terms of some modified theta lift akin to (1.4).

5.2 Finite-N integrated correlators

As mentioned in the introduction, it was conjectured in [21] that the correct vector space of modular invariant functions containing the coefficients of the large-N expansion of a certain integrated correlator, $\mathcal{H}_N(\tau)$, of four superconformal primary operators in the stress-tensor multiplet in $\mathcal{N} = 4$ SYM is given by the direct sum of non-holomorphic Eisenstein series with half-integer index and SMFs $\mathcal{E}_{i,j}^w(\tau)$ where w and s = 3i + j + 1have opposite parity. An interesting open question is how can this integrated correlator be represented at generic finite value of N. Inspired by a simpler case [50, 51] where the same four-point correlator is integrated with respect to a different measure, we may expect that in passing from large-N to finite N, the vector space of modular invariant functions changes to the direct sum of non-holomorphic Eisenstein series with integer index and MGFs $\mathcal{E}_{i,j}^w(\tau)$ where w and s = 3i + j + 1 have the same parity. At finite N we expect the integrated correlator to be given by a formal sum over infinitely many elements of this vector space.

For computational purposes, it might be useful to try and enlarge this vector space and use as a basis all GESs with spectrum (4.13). However, at any given N this physical observable can be computed from a well-known matrix model originating from supersymmetric localisation, which is believed to not produce L-values of holomorphic cusp forms. It is of interest to find explicitly the expression for the integrated correlator in terms of MGFs, thus immediately proving the absence of L-values, or as a sum of GESs with spectrum (4.13) and subsequently verify whether the cusp forms do indeed cancel in the infinite sum.

Guided by [50, 51], we conjecture that at finite N the representation of the integrated correlator $\mathcal{H}_N(\tau)$ as an infinite sum over MGFs and integer-index non holomorphic Eisenstein series should be equivalent to

$$\mathcal{H}_N(\tau) = \int_{(\mathbb{R}^+)^3} B_N(t) \,\Gamma_{2,2}(\tau; t) \,\mathrm{d}^3 t \,.$$
(5.3)

Two additional key properties motivate this conjecture. Firstly, the finite-N expression (5.3) when expanded at large-N ought to reproduce the results derived in [21]. In particular, the integrand function $B_N(t)$ originates from the resummation of the infinite series of MGFs with coefficients depending from N. Order by order at large-N, the expansion of $B_N(t)$ should produce a linear combination of integrands $B_{i,j}^w(t)$ of the form (2.35), which upon integrating over t produces the linear combination of SMFs found in [21]. This is exactly what happens in the simpler case [50, 51].

Secondly, in (5.3) we have used the complete theta-series (2.16) which includes terms in the lattice sum with vanishing momenta $p_i = 0$. While for a single $\mathcal{E}_{i,j}^w(\tau)$ the inclusion of these terms leads to an ill-defined singular integral in (2.34), for the resummed expression in (5.3) we expect the integral to converge even in the absence of the exponential suppression coming from (2.33). Furthermore, if we consider $\mathcal{E}_{i,j}^w(\tau)$ by extending the theta-series to the complete lattice sum as in (2.17), we can regularise the integral by introducing a cutoff, i.e. by integrating over $(t_1, t_2, t_3) \in [0, L]^3$ rather than over $(\mathbb{R}^+)^3$. One can then easily check that terms where one or three momenta vanish¹⁴ produce divergent terms as $L \gg 1$, which are either constant or proportional to integer index non-holomorphic Eisenstein series when w and s = 3i + j + 1 have the same parity, or half-integer index when w and s have opposite parity. We believe that this phenomenon is a signal of the fact that the large-N expansion of (5.3) produces both integer powers in 1/N terms, which are linear combinations of SMFs, as well as the half-integer powers in 1/N, which are rational linear combinations of non-holomorphic Eisenstein series with half-integer indices.

5.3 S-dual modular forms

The generic type IIB superstring amplitudes and correlators in $\mathcal{N} = 4$ SYM are not necessarily invariant under SL(2, \mathbb{Z}). In particular, these include U(1)-violating interactions, which transform covariantly with non-zero modular weight under SL(2, \mathbb{Z}). A natural extension of the cases studied in this paper involves the maximal U(1)-violating amplitudes [46] and their holographic dual correlators in $\mathcal{N} = 4$ SYM [47, 48].

Explicit examples discussed in [46] include the maximal U(1)-violating scattering amplitudes of four gravitons and h axion-dilatons excitations in type IIB superstring theory, which are described by modular forms with holomorphic and anti-holomorphic weights (h, -h), here denoted by $\mathcal{E}_h(s; s_1, h_1; s_2, h_2; \tau)$. In

¹⁴Due to momentum conservation $p_1 + p_2 + p_3 = 0$ in the lattice sum (2.16), if two momenta are zero then the third one is necessarily zero.

general, such modular forms transform as

$$f_h(\gamma \cdot \tau) = \left(\frac{c\tau + d}{c\overline{\tau} + d}\right)^h f_h(\tau), \quad \text{for} \quad \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \text{SL}(2, \mathbb{Z}).$$
(5.4)

It was argued in [46], the coefficients in the low-energy expansion of such amplitudes obey a Laplace eigenvalue equation of the form,

$$\left[\Delta_{(+)}^{(h)} - (s+h)(s-h-1)\right] \mathcal{E}_h(s;s_1,h_1;s_2,h_2;\tau) = E_{h_1}(s_1;\tau) E_{h_2}(s_2;\tau) , \qquad (5.5)$$

where $h_1 + h_2 = h$, and

$$\Delta_{(+)}^{(h)} = 4\overline{\mathcal{D}}_{-h-1}\mathcal{D}_h, \quad \text{with} \quad \mathcal{D}_h = i\left(\tau_2 \frac{\partial}{\partial_\tau} - i\frac{h}{2}\right), \quad (5.6)$$

The modular forms $E_h(s;\tau)$ in the source term of (5.5) are defined by

$$E_{h}(s;\tau) = \frac{2^{h}\Gamma(s)}{\Gamma(s+h)}\mathcal{D}_{h-1}\cdots\mathcal{D}_{0}E(s;\tau) = \sum_{(m,n)\neq(0,0)} \left(\frac{m+n\bar{\tau}}{m+n\tau}\right)^{h} \frac{(\tau_{2}/\pi)^{s}}{|m+n\tau|^{2s}}.$$
(5.7)

When h = 0, $E_0(s; \tau)$ reduces to the non-holomorphic Eisenstein series, $E(s; \tau)$.

The Laplace eigenvalue equation (5.5) is a very natural generalisation of (1.1) that was studied in this paper. An interesting case that is relevant to superstring amplitudes was considered in [46] corresponds to $s = 4, s_1 = s_2 = \frac{3}{2}$ and h > 0. In this case solutions to (5.5) can be related to the action of the covariant derivatives \mathcal{D}_w on $\mathcal{E}(4; \frac{3}{2}, \frac{3}{2}; \tau)$ with appropriate values of w. It is of interest to study the solution to equation (5.5) with more general parameters (with $s_1, s_2 \in \mathbb{N}$ and $s_1, s_2 \in \mathbb{N} + \frac{1}{2}$, respectively), and in particular examine the roles played by the *L*-values of holomorphic cusp forms and their potential cancellations.

Acknowledgements

We thank Guillaume Bossard, Ksenia Fedosova, Jens Funke, Abhiram Kidambi, Axel Kleinschmidt, Kim Klinger-Logan, Oliver Schlotterer for helpful discussions. We would also like to thank Axel Kleinschmidt and Oliver Schlotterer for providing extremely helpful comments on the draft. DD is indebted to the Albert Einstein Institute and in particular to Axel Kleinschmidt, Hermann Nicolai and Stefan Theisen for the hospitality and financial support. CW is supported by a Royal Society University Research Fellowship, URF\R\221015 and partly a STFC Consolidated Grant, ST\T000686\1 "Amplitudes, strings & duality".

A Non-holomorphic Eisenstein series

Non-holomorphic Eisenstein series constitute an important class of modular invariant functions that arise both in the study of one-loop MGFs as well as in the analysis of low-order coefficients in the low energy effective action of type IIB superstring theory.

They are special examples of Maass functions for the group $SL(2, \mathbb{Z})$ which we recall are modular invariant functions $f(\tau)$, i.e.

$$f(\gamma \cdot \tau) = f(\tau), \qquad \forall \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \mathrm{SL}(2, \mathbb{Z}), \qquad (A.1)$$

with $\gamma \cdot \tau \coloneqq \frac{a\tau+b}{c\tau+d}$, of moderate growth for large τ_2 , i.e. there exists $N \in \mathbb{N}$ such that $f(\tau) = O(\tau_2^N)$ as $\tau_2 \gg 1$, and importantly $f(\tau)$ satisfies the Laplace eigenvalue equation

$$[\Delta_{\tau} - s(s-1)]f(\tau) = 0, \qquad (A.2)$$

for some $s \in \mathbb{C}$. The non-holomorphic Eisenstein series, $E(s; \tau)$, is perhaps the simplest example of a Maass function and can be defined by the two-dimensional lattice sum,

$$E(s;\tau) = \sum_{(m,n)\neq(0,0)} \frac{(\tau_2/\pi)^s}{|m+n\tau|^{2s}}.$$
(A.3)

The function $E(s; \tau)$ satisfies the Laplace equation (A.2) and has a Fourier expansion given by

$$E(s;\tau) = \frac{2\zeta(2s)}{\pi^s} \tau_2^s + \frac{2\Gamma(s-\frac{1}{2})}{\pi^{s-\frac{1}{2}}\Gamma(s)} \zeta(2s-1)\tau_2^{1-s} + \frac{4\sqrt{\tau_2}}{\Gamma(s)} \sum_{k\neq 0} |k|^{s-\frac{1}{2}} \sigma_{1-2s}(k) K_{s-\frac{1}{2}}(2\pi |k| \tau_2) e^{2\pi i k \tau_1},$$
(A.4)

where $K_{s-\frac{1}{2}}(2\pi |k| \tau_2)$ is a modified Bessel function of the second kind and $\sigma_s(n) \coloneqq \sum_{d|n} d^s$ is the divisor sigma function. Although the lattice sum definition (A.3) only converges for $\operatorname{Re}(s) > 1$, it is possible to analytically continue $E(s;\tau)$ to a meromorphic function of $s \in \mathbb{C}$, with a simple pole located at s = 1. For later convenience we note that with the normalisation convention chosen the non-holomorphic Eisenstein series satisfies the functional equation

$$\Gamma(s)E(s;\tau) = \Gamma(1-s)E(1-s;\tau), \qquad (A.5)$$

and we also recall the integral representation

$$E(s;\tau) = \sum_{(m,n)\neq(0,0)} \frac{1}{\Gamma(s)} \int_0^\infty t^{s-1} e^{-\pi t \frac{|m+n\tau|^2}{\tau_2}} \mathrm{d}t\,,\tag{A.6}$$

valid for $\operatorname{Re}(s) > 1$.

B Modular local polynomials

In this appendix we provide more details regarding the modular local polynomials presented in section 2.2. Following [28] we are interested in analysing the exceptional set E_D , as defined in (2.22), where the modular local polynomials become non-differentiable. In particular, we now prove the claim made in the main text that for the case of discriminant D = 1, modular local polynomials with exceptional set E_1 are uniquely specified by their values on the fundamental domain of $\Gamma_0(2)$.

We start by noting that the exceptional set E_D defined in (2.22) is $SL(2,\mathbb{Z})$ invariant since for all $\gamma \in SL(2,\mathbb{Z})$ and for any $Q = [a, b, c] \in \mathcal{Q}_D$ (where $D = b^2 - 4ac$ is the discriminant) we have

$$S_Q|\gamma = \left\{ \rho \in \mathfrak{H} \mid a|\gamma \cdot \rho|^2 + b\operatorname{Re}(\gamma \cdot \rho) + c = 0 \right\} = S_{Q'},$$
(B.1)

with $Q' = [a', b', c'] \in \mathcal{Q}_D$ given by

$$a' = aI^{2} + bIK + cK^{2},$$

$$b' = 2aIJ + bIL + bJK + 2cKL,$$

$$c' = aJ^{2} + bJL + cL^{2},$$

(B.2)

and

$$\gamma = \begin{pmatrix} I & J \\ K & L \end{pmatrix} \in \operatorname{SL}(2, \mathbb{Z}).$$
(B.3)

From now on, we focus our attention to the case where the quadratic forms have discriminant D = 1 and note that the class [0, 1, c] with $c \in \mathbb{Z}$ belongs to \mathcal{Q}_1 with corresponding set

$$S_{[0,1,c]} = \left\{ \rho \in \mathfrak{H} \, \middle| \, \operatorname{Re}(\rho) + c = 0 \right\} \,. \tag{B.4}$$

We also see that $[-1, 1, 0] \in \mathcal{Q}_1$ with set

$$S_{[-1,1,0]} = \left\{ \rho \in \mathfrak{H} \, \Big| \, |\rho - \frac{1}{2}|^2 = \frac{1}{4} \right\} \,. \tag{B.5}$$

The subsets of E_1 given by $S_{[0,1,0]}$, $S_{[0,1,1]}$ and $S_{[-1,1,0]}$ coincide precisely with the boundary of the fundamental domain of $\Gamma_0(2)$, hence we claim that the connected component $\mathcal{C}_{\rho^*} \subset \mathfrak{H} \setminus E_1$ which contains $\rho^* := e^{\frac{\pi i}{3}}$ coincides with the fundamental domain of $\Gamma_0(2)$, i.e.

$$\mathcal{C}_{\rho^*} = \mathcal{F}_{\Gamma_0(2)} \coloneqq \left\{ \rho \in \mathfrak{H} \left| 0 < \operatorname{Re}(\rho) < 1, \left| \rho - \frac{1}{2} \right|^2 > \frac{1}{4} \right\} \right.$$
(B.6)

Proof: Firstly we show that ρ^* cannot possibly belong to the exceptional set, i.e. $\rho^* \neq E_1$. By contradiction, if $\rho^* \in E_1$ then there must exist $Q = [a, b, c] \in Q_1$ such that

$$a|\rho^*|^2 + b\operatorname{Re}(\rho^*) + c = 0 \Rightarrow b = -2(a+c),$$
 (B.7)

which would imply $D = b^2 - 4ac$ even thus contradicting the fact that D = 1.

Now we will show that the sets $S_{[0,1,0]}$, $S_{[0,1,-1]}$ and $S_{[-1,1,0]}$ form the boundaries of $\mathcal{F}_{\Gamma_0(2)}$, $\operatorname{Re}(\rho) = 0$, $\operatorname{Re}(\rho) = 1$ and $|\rho - \frac{1}{2}|^2 = \frac{1}{4}$, respectively. Suppose there exists $Q = [a, b, c] \in \mathcal{Q}_1$ such that $S_Q \cap \mathcal{F}_{\Gamma_0(2)} \neq \emptyset$, without loss of generality we can assume that $a \ge 0$, since

$$[a, b, c] \in \mathcal{Q}_1 \iff [-a, -b, -c] \in \mathcal{Q}_1, \qquad (B.8)$$

and furthermore

$$S_{[a,b,c]} \cap \mathcal{F}_{\Gamma_0(2)} \neq \emptyset \iff S_{[-a,-b,-c]} \cap \mathcal{F}_{\Gamma_0(2)} \neq \emptyset.$$
(B.9)

We can furthermore discard the case a = 0 since we already know that $S_{[0,1,c]}$ either does not intersect $\mathcal{F}_{\Gamma_0(2)}$ or produces the two vertical boundaries.

Let us suppose there exists $\tilde{\rho} = x + iy \in S_{[a,b,c]} \cap \mathcal{F}_{\Gamma_0(2)}$, i.e.

$$a(x^2 + y^2) + bx + c = 0, \qquad \rho^* \in \mathcal{F}_{\Gamma_0(2)}.$$
 (B.10)

Firstly, we see immediately that if c < 0 it is impossible to satisfy the discriminant condition $D = b^2 - 4ac = 1$ unless a = 0, b = 1 which we have already considered, hence we can assume $c \ge 0$.

Since $\tilde{\rho} \in \mathcal{F}_{\Gamma_0(2)}$ we know already that 0 < x < 1 and $x^2 + y^2 > x$ so we deduce

$$a(x^{2} + y^{2}) + bx + c > (a + b)x + c.$$
 (B.11)

If we assume that $(a + b) \ge 0$, since as discussed above $c \ge 0$, we find that the only possibility of satisfying (B.10) given the constraint (B.11) and D = 1 is to consider $S_{[1,-1,0]} = S_{[-1,1,0]}$ which correspond to the lower boundary of $\mathcal{F}_{\Gamma_0(2)}$. Let us consider then the case (a + b) < 0. Since $\tilde{\rho} \in \mathcal{F}_{\Gamma_0(2)}$ we must have x < 1 hence for (a + b) < 0 (B.11) becomes

$$a(x^{2} + y^{2}) + bx + c > a + b + c.$$
 (B.12)

To satisfy (B.10) given this constraint we must have $-b > a + c \ge 0$ hence the discriminant must satisfy

$$1 = D = b^2 - 4ac \ge (a - c)^2.$$
(B.13)

This implies either $a = c \in \mathbb{N}^+$ and hence $b = -\sqrt{4a^2 + 1} \notin \mathbb{Z}$ or $a = c \pm 1$ and hence $b = -2c \mp 1$ so that the constraint (B.12) becomes $a(x^2 + y^2) + bx + c > 0$ and again $\tilde{\rho}$ can only live at the boundary of $\mathcal{F}_{\Gamma_0(2)}$.

C Holomorphic cusp forms and their *L*-values

In this appendix we review some important facts about holomorphic cusp forms for $SL(2, \mathbb{Z})$. We start by defining M_k as the vector space of modular forms of weight $k \in \mathbb{Z}$ under $SL(2, \mathbb{Z})$, which are furthermore analytic in the upper half-plane \mathfrak{H} . It is well-known (see for example [52]) that when k odd, or if k < 0, or k = 2 then the only element of M_k is given by the zero-function while for k = 0 the only element of M_0 is the constant function. For k > 2 and even, M_k splits naturally as

$$M_k = \operatorname{span}\{G_k\} \oplus \mathcal{S}_k \,, \tag{C.1}$$

where G_k denotes the holomorphic Eisenstein series

$$G_k(\tau) = \sum_{n=0}^{\infty} \alpha_k(n) q^n , \qquad (C.2)$$

where $q = \exp(2\pi i \tau)$ and the Fourier coefficients $\alpha_k(n)$ are given by

$$\alpha_k(0) = 1, \qquad \alpha_k(1) = \alpha_k = -\frac{2k}{B_k}, \qquad \alpha_k(n) = \alpha_k \sigma_{k-1}(n), \qquad (C.3)$$

with B_k denoting the k^{th} Bernoulli number and $\sigma_s(n) = \sum_{d|n} d^s$ the divisor sigma function.

The second term in (C.1) is given by the vector space S_k of $SL(2, \mathbb{Z})$ holomorphic cusp forms. A function Δ analytic in the upper half-plane \mathfrak{H} belongs to S_k if it is a modular form of weight k and vanishes at the cusp $\tau \to i\infty$, i.e. its q-series Fourier expansion must not contain the constant term

$$\Delta(\tau) = \sum_{n=1}^{\infty} a(n)q^n \,. \tag{C.4}$$

We furthermore say that the cusp form $\Delta \in S_k$ is *Hecke normalised* if a(1) = 1. The dimension of the vector space of holomorphic cusp forms is given by

$$\dim \mathcal{S}_k = \begin{cases} \left\lfloor \frac{k}{12} \right\rfloor - 1 & k \equiv 2 \mod 12, \\ \left\lfloor \frac{k}{12} \right\rfloor & \text{otherwise.} \end{cases}$$
(C.5)

In [53] Hecke introduced an infinite family of linear operators $T_n : S_k \to S_k$ with $n \in \mathbb{N}$, now called Hecke operators, defined by

$$[T_n \Delta](\tau) = n^{k-1} \sum_{d|n} d^{-k} \sum_{b=0}^{d-1} \Delta\left(\frac{n\tau + bd}{d^2}\right) .$$
(C.6)

A preferred basis for S_k can be constructed by considering the basis of Hecke eigenforms which simultaneously diagonalise all T_n . Importantly, the eigenvalues of a Hecke eigenform $\Delta(\tau) = \sum_{n=1}^{\infty} a(n)q^n \in S_k$ are precisely given by its Fourier q-series coefficients, i.e. we have $[T_n\Delta](\tau) = a(n)\Delta(\tau)$ so that from (C.6) we see that a Hecke eigenform is automatically Hecke normalised.

Since the Hecke operators are multiplicative, i.e. $T_mT_n = T_{m\cdot n}$ if (m, n) = 1, it follows that the Fourier coefficients a(n) of every Hecke eigenform must be multiplicative as well, i.e. $a(m)a(n) = a(m \cdot n)$ for (m, n) = 1. This important property leads to the fact that in general the coefficients of a Hecke eigenform live in a number field extension of the rationals, which is a finite degree field extension of \mathbb{Q} by the roots of an associated defining polynomial of degree dim S_k . Whenever dim $S_k = 1$ we have that this number field actually coincides with the rational numbers. However, when dim $S_k > 1$ that is not true any longer. For example if we consider k = 24, for which dim $S_{24} = 2$, we find that the Fourier coefficients of the two unique

Hecke normalised basis elements of S_{24} belong to the number field $\mathbb{Q}[\sqrt{144169}]$ originating from the defining polynomial $x^2 - x - 36042$.

Given a Hecke eigenform $\Delta(\tau) = \sum_{n=1}^{\infty} a(n)q^n \in S_k$, an important quantity to consider is the associated *L*-function defined via the Dirichlet series

$$L(\Delta, s) \coloneqq \sum_{n=1}^{\infty} \frac{a(n)}{n^s}.$$
 (C.7)

In general this series converges for $\operatorname{Re}(s)$ large enough but it can be analytically continued to a meromorphic function of $s \in \mathbb{C}$. From $L(\Delta, s)$ we construct the completed *L*-function for a Hecke eigenform,

$$\Lambda(\Delta, s) \coloneqq (2\pi)^{-s} \Gamma(s) L(\Delta, s) , \qquad (C.8)$$

which satisfies the important functional equation

$$\Lambda(\Delta, s) = (-1)^{\frac{k}{2}} \Lambda(\Delta, k - s).$$
(C.9)

We remind the reader that the critical strip for the completed L-function $\Lambda(\Delta, s)$ of a cusp form $\Delta \in S_k$ is defined as the strip $\operatorname{Re} s \in (0, k)$ and the critical L-values of Δ are given by $\Lambda(\Delta, s)$ with $s \in (0, k) \cap \mathbb{N}$ while values outside of this range are called non-critical. A beautiful result by Manin [36], known as the periods theorem, proves that the ratio of any two even critical L-values or any two odd critical L-values are all rational over the algebraic number field generated by the Fourier coefficient of the cusp forms. This theorem effectively means that there are only two independent critical values for any given cusp forms, usually denoted as ω_{Δ}^{\pm} and called the periods of the cusp form. On the other hand, all non-critical L-values are conjecturally transcendentally independent from one another.

Manin's results rely heavily on the notion of period polynomials. For every Hecke-normalised cusp form $\Delta \in S_k$ we define its associated period polynomial

$$r_{\Delta}(X) \coloneqq \int_0^{i\infty} \Delta(\tau) (\tau - X)^{k-2} \mathrm{d}\tau \,, \tag{C.10}$$

which is a degree k-2 polynomial in the variable X. We can use the explicit q-series representation $\Delta(\tau) = \sum_{n=1}^{\infty} a(n)q^n$ with $q = \exp(2\pi i\tau)$ to show that

$$\Lambda(\Delta, \ell+1) = i^{-\ell-1} \int_0^{i\infty} \Delta(\tau) \tau^\ell d\tau \,. \tag{C.11}$$

Using (C.11) it is easy to prove that the period polynomial (C.10) takes the form,

$$r_{\Delta}(X) = \sum_{\ell=0}^{k-2} i^{\ell+1} (-X)^{k-2-\ell} \binom{k-2}{\ell} \Lambda(\Delta, \ell+1).$$
 (C.12)

We immediately note that the coefficients of this polynomial are the completed *L*-values for the cusp form Δ , and $r_{\Delta}(X)$ carries important information about the underlying modular group SL(2, \mathbb{Z}). In particular, considering the order-2 and 3 generators of SL(2, \mathbb{Z}) respectively given by $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ and $U = \begin{pmatrix} 1 & -1 \\ 1 & 0 \end{pmatrix}$, the period polynomial must satisfy the two cocycle conditions

$$r_{\Delta}|_{1}(X) + r_{\Delta}|_{S}(X) = r_{\Delta}|_{1}(X) + r_{\Delta}|_{U}(X) + r_{\Delta}|_{U^{2}}(X) = 0, \qquad (C.13)$$

where the $SL(2, \mathbb{Z})$ slash-action on r_{Δ} is given by

$$r_{\Delta}|_{\gamma}(X) \coloneqq (cX+d)^{k-2}r_{\Delta}\left(\frac{aX+b}{cX+d}\right), \qquad \gamma = \begin{pmatrix} a & b\\ c & d \end{pmatrix} \in \mathrm{SL}(2,\mathbb{Z}).$$
(C.14)

It is easy to check that for the critical *L*-values, the reflection identity $\Lambda(\Delta, s) = (-1)^{k/2} \Lambda(\Delta, k-s)$ is a direct consequence of the *S*-cocycle condition in (C.13). The aforementioned periods theorem of Manin [36] is a consequence of considering both *S* and *U* cocycle conditions in (C.13) as well as the action on $r_{\Delta}(X)$ induced by the Hecke operators (C.6) which are sensitive to the number field generated by the *q*-series coefficients of the Hecke normalised cusp forms in S_k given that their eigenvalues are the precisely Fourier coefficients of the Hecke eigenform considered.

Lastly, it is important that it is possible to put a norm, called the Petersson norm, on the vector space S_k , which is defined by

$$\langle \Delta, \Delta \rangle \coloneqq \int_{\mathcal{F}} |\Delta(\tau)|^2 \tau_2^{k-2} \,\mathrm{d}^2 \tau \,, \qquad \Delta \in \mathcal{S}_k \,,$$
 (C.15)

where the integral is over the fundamental domain $\mathcal{F} = \mathrm{SL}(2,\mathbb{Z}) \setminus \mathfrak{H}$. In section 4.2, we use a classic result by Rankin [35] which allows us to write the product of two critical *L*-values with opposite parity, i.e. one even and one odd, in terms of the Petersson norm. For the j^{th} Hecke normalised eigenform $\Delta^{(j)} \in \mathcal{S}_k$ with $j \in \{1, \ldots, \dim \mathcal{S}_k\}$, Rankin's theorem states

$$\langle \Delta^{(j)}, \Delta^{(j)} \rangle = -\frac{q r \Lambda(\Delta^{(j)}, k-1) \Lambda(\Delta^{(j)}, q)}{B_q B_r \kappa(j, q, r)}, \qquad (C.16)$$

where B_n is the n^{th} Bernoulli number and q and r are any even integers such that q+r = k and $4 \le r \le \frac{k}{2}-2$. The function $\kappa(j, q, r)$ takes value in the number field generated by the Fourier coefficients of the Hecke eigenforms of S_k and it is given by

$$\kappa(j,q,r) \coloneqq \sum_{n=1}^{\dim \mathcal{S}_k} \left(H^{-1} \right)_{jn} \left[\sum_{m=0}^n \alpha_q(m) \alpha_r(n-m) - \alpha_k(n) \right], \tag{C.17}$$

with $\alpha_k(n)$ given in (C.3). The $(\dim S_k) \times (\dim S_k)$ matrix H is defined by $H_{nj} \coloneqq a^{(j)}(n)$, i.e. the entry in the n^{th} row and j^{th} column of the matrix H is precisely the n^{th} q-series coefficient of the j^{th} Hecke eigenform, i.e. $\Delta^{(j)}(\tau) = \sum_{n=1}^{\infty} a^{(j)}(n)q^n \in S_k$. A particularly simple expression can be derived when we specialise (C.16) to the cases where $\dim S_k = 1$, i.e. $k \in \{12, 16, 20, 22, 26\}$. In this case the 1×1 matrix H reduces to 1 from the Hecke normalisation, and if we furthermore choose r = 4, q = k - 4 we find that (C.16) reduces to

$$\langle \Delta, \Delta \rangle = -\frac{15 \ 2^{5-k} (k-4) B_k \Lambda(\Delta, k-4) \Lambda(\Delta, k-1)}{\left[(k-4) B_k - (k+120 B_k) B_{k-4} \right]}.$$
 (C.18)

For example, for the Ramanujan cusp form Δ_{12} which is the unique Hecke normalised element of S_{12} , equation (C.18) reduces to (see (9.1) in [35])

$$\langle \Delta_{12}, \Delta_{12} \rangle = \frac{691\Lambda(\Delta_{12}, 8)\Lambda(\Delta_{12}, 11)}{7680} = \frac{10! \times 14511}{2^{24}\pi^{19}} L(\Delta_{12}, 8)L(\Delta_{12}, 11).$$
(C.19)

References

- M. B. Green and P. Vanhove, "The Low-energy expansion of the one loop type II superstring amplitude," Phys. Rev. D 61 (2000), 104011 [arXiv:hep-th/9910056 [hep-th]].
- [2] E. D'Hoker, M. B. Green, Ö. Gürdogan and P. Vanhove, "Modular Graph Functions," Commun. Num. Theor. Phys. 11, 165-218 (2017) [arXiv:1512.06779 [hep-th]].
- [3] E. D'Hoker and M. B. Green, "Identities between Modular Graph Forms," J. Number Theor. 189, 25-80 (2018) [arXiv:1603.00839 [hep-th]].

- [4] M. B. Green, J. G. Russo and P. Vanhove, "Low energy expansion of the four-particle genus-one amplitude in type II superstring theory," JHEP **02** (2008), 020 [arXiv:0801.0322 [hep-th]].
- [5] F. Brown, "Multiple modular values and the relative completion of the fundamental group of $\mathcal{M}_{1,1}$," arXiv:1407.5167 [math.NT].
- [6] F. Brown, "A class of non-holomorphic modular forms I," Res. Math. Sci. 5 (2018) no. 1, Paper No. 7, 40, arXiv:1707.01230 [math.NT].
- [7] F. Brown, "A class of nonholomorphic modular forms II: Equivariant iterated Eisenstein integrals," Forum Math. Sigma 8 (2020) Paper No. e31, 62, arXiv:1708.03354 [math.NT].
- [8] E. D'Hoker, M. B. Green and P. Vanhove, "On the modular structure of the genus-one Type II superstring low energy expansion," JHEP **08** (2015), 041 [arXiv:1502.06698 [hep-th]].
- [9] D. Dorigoni, A. Kleinschmidt and O. Schlotterer, "Poincaré series for modular graph forms at depth two. Part I. Seeds and Laplace systems," JHEP 01 (2022), 133 [arXiv:2109.05017 [hep-th]].
- [10] D. Dorigoni, A. Kleinschmidt and O. Schlotterer, "Poincaré series for modular graph forms at depth two. Part II. Iterated integrals of cusp forms," JHEP 01 (2022), 134 [arXiv:2109.05018 [hep-th]].
- [11] M. B. Green, H. h. Kwon and P. Vanhove, "Two loops in eleven-dimensions," Phys. Rev. D 61 (2000), 104010 [arXiv:hep-th/9910055 [hep-th]].
- [12] M. B. Green and P. Vanhove, "Duality and higher derivative terms in M theory," JHEP 01 (2006), 093 [arXiv:hep-th/0510027 [hep-th]].
- [13] C. Montonen and D. I. Olive, "Magnetic Monopoles as Gauge Particles?," Phys. Lett. B 72 (1977), 117-120
- [14] J. M. Maldacena, "The Large N limit of superconformal field theories and supergravity," Adv. Theor. Math. Phys. 2, 231-252 (1998) [arXiv:hep-th/9711200 [hep-th]].
- [15] D. J. Binder, S. M. Chester, S. S. Pufu and Y. Wang, "N = 4 Super-Yang-Mills correlators at strong coupling from string theory and localization," JHEP 12 (2019), 119 [arXiv:1902.06263 [hep-th]].
- [16] S. M. Chester and S. S. Pufu, "Far beyond the planar limit in strongly-coupled $\mathcal{N} = 4$ SYM," JHEP **01** (2021), 103 [arXiv:2003.08412 [hep-th]].
- [17] V. Pestun, "Localization of gauge theory on a four-sphere and supersymmetric Wilson loops," Commun. Math. Phys. 313, 71-129 (2012) [arXiv:0712.2824 [hep-th]].
- [18] D. Dorigoni, M. B. Green and C. Wen, "The SAGEX review on scattering amplitudes Chapter 10: Selected topics on modular covariance of type IIB string amplitudes and their supersymmetric Yang–Mills duals," J. Phys. A 55 (2022) no.44, 443011 [arXiv:2203.13021 [hep-th]].
- [19] S. M. Chester, M. B. Green, S. S. Pufu, Y. Wang and C. Wen, "Modular invariance in superstring theory from N = 4 super-Yang-Mills," JHEP 11, 016 (2020) [arXiv:1912.13365 [hep-th]].
- [20] S. M. Chester, M. B. Green, S. S. Pufu, Y. Wang and C. Wen, "New modular invariants in $\mathcal{N} = 4$ Super-Yang-Mills theory," JHEP **04** (2021), 212 [arXiv:2008.02713 [hep-th]].
- [21] L. F. Alday, S. M. Chester, D. Dorigoni, M. B. Green and C. Wen, "Relations between integrated correlators in $\mathcal{N} = 4$ supersymmetric Yang-Mills theory," JHEP 05 (2024), 044 [arXiv:2310.12322].

- [22] K. Fedosova, K. Klinger-Logan and D. Radchenko, "Convolution identities for divisor sums and modular forms," Proc. Natl. Acad. Sci., Vol. 121, No. 44 (2024) [arXiv:2312.00722 [math.NT]].
- [23] K. Fedosova and K. Klinger-Logan, "Whittaker Fourier type solutions to differential equations arising from string theory," [arXiv:2209.09319 [math.NT]].
- [24] D. Zagier, Unpublished notes 2008.
- [25] E. D'Hoker, M. B. Green and B. Pioline, "Asymptotics of the D⁸R⁴ genus-two string invariant," Commun. Num. Theor. Phys. 13 (2019), 351-462 [arXiv:1806.02691]
- [26] M. B. Green, J. G. Russo and P. Vanhove, "Modular properties of two-loop maximal supergravity and connections with string theory," JHEP 07 (2008), 126 [arXiv:0807.0389 [hep-th]].
- [27] K. Bringmann and B. Kane, "Modular local polynomials," Math. Res. Lett 23 (2016) 973 1405.0589 [math.NT]
- [28] K. Bringmann, B. Kane, and W. Kohnen, "Locally harmonic Maass forms and the kernel of the Shintani lift," Int. Math. Res. Not. IMRN (2015), no. 11, 3185–3224. [arXiv:1206.1100 [math.NT]]
- [29] E. D'Hoker and W. Duke, "Fourier series of modular graph functions," J. Number Theor. 192 (2018), 1-36 [arXiv:1708.07998 [math.NT]].
- [30] K. Klinger-Logan, "Differential equations in automorphic forms," Commun. Num. Theor. Phys. **12** (2018), 767-827
- [31] D. Dorigoni and A. Kleinschmidt, "Modular graph functions and asymptotic expansions of Poincaré series," Commun. Num. Theor. Phys. 13 (2019) no.3, 569-617 [arXiv:1903.09250 [hep-th]].
- [32] D. Dorigoni, A. Kleinschmidt and R. Treilis, "To the cusp and back: resurgent analysis for modular graph functions," JHEP 11 (2022), 048 [arXiv:2208.14087 [hep-th]].
- [33] D. Dorigoni and R. Treilis, "Two string theory flavours of generalised Eisenstein series," JHEP 11 (2023), 102 [arXiv:2307.07491 [hep-th]].
- [34] J. Broedel, O. Schlotterer and F. Zerbini, "From elliptic multiple zeta values to modular graph functions: open and closed strings at one loop," JHEP 01 (2019), 155 [arXiv:1803.00527 [hep-th]].
- [35] R. A. Rankin, "The Scalar Product of Modular Forms," Proceedings of the London Mathematical Society, Volume s3-2, Issue 1, 1952, Pages 198–217.
- [36] J. I. Manin, "Periods of parabolic forms and p-adic hecke series," Mathematics of the USSR-Sbornik 21 (1973) 371–393.
- [37] M. B. Green, S. D. Miller and P. Vanhove, "SL(2,ℤ)-invariance and D-instanton contributions to the D⁶R⁴ interaction," Commun. Num. Theor. Phys. **09** (2015), 307-344 [arXiv:1404.2192 [hep-th]].
- [38] D. Dorigoni, M. Doroudiani, J. Drewitt, M. Hidding, A. Kleinschmidt, N. Matthes, O. Schlotterer and B. Verbeek, "Modular graph forms from equivariant iterated Eisenstein integrals," JHEP 12 (2022), 162 [arXiv:2209.06772 [hep-th]].
- [39] D. Dorigoni, M. Doroudiani, J. Drewitt, M. Hidding, A. Kleinschmidt, O. Schlotterer, L. Schneps and B. Verbeek, "Non-holomorphic modular forms from zeta generators," JHEP 10 (2024), 053 [arXiv:2403.14816 [hep-th]].

- [40] J. E. Gerken, A. Kleinschmidt and O. Schlotterer, "All-order differential equations for one-loop closedstring integrals and modular graph forms," JHEP 01 (2020), 064 [arXiv:1911.03476 [hep-th]].
- [41] J. E. Gerken, A. Kleinschmidt and O. Schlotterer, "Generating series of all modular graph forms from iterated Eisenstein integrals," JHEP 07 (2020) no.07, 190 [arXiv:2004.05156 [hep-th]].
- [42] H. Tsunogai, "On some derivations of Lie algebras related to Galois representations," Publ. Res. Inst. Math. Sci., 31 (1995), 113–134.
- [43] A. Pollack, "Relations between derivations arising from modular forms", undegraduate thesis Duke University (2009), https://dukespace.lib.duke.edu/dspace/handle/10161/1281.
- [44] D. Dorigoni, M. Doroudiani, J. Drewitt, M. Hidding, A. Kleinschmidt, O. Schlotterer, L. Schneps and B. Verbeek, "Canonicalizing zeta generators: genus zero and genus one," [arXiv:2406.05099 [math.QA]].
- [45] The PARI Group, Univ. Bordeaux, PARI/GP version 2.13.1, 2020. http://pari.math.u-bordeaux.fr/.
- [46] M. B. Green and C. Wen, "Modular Forms and SL(2,ℤ)-covariance of type IIB superstring theory," JHEP 06, 087 (2019) [arXiv:1904.13394 [hep-th]].
- [47] M. B. Green and C. Wen, "Maximal U(1)_Y-violating n-point correlators in $\mathcal{N} = 4$ super-Yang-Mills theory," JHEP **02**, 042 (2021) [arXiv:2009.01211 [hep-th]].
- [48] D. Dorigoni, M. B. Green and C. Wen, "Exact expressions for *n*-point maximal $U(1)_Y$ -violating integrated correlators in $SU(N) \mathcal{N} = 4$ SYM," JHEP **11**, 132 (2021) [arXiv:2109.08086 [hep-th]].
- [49] E. D'Hoker and J. Kaidi, "Modular graph functions and odd cuspidal functions. Fourier and Poincaré series," JHEP 04 (2019), 136 [arXiv:1902.04180 [hep-th]].
- [50] D. Dorigoni, M. B. Green and C. Wen, "Novel Representation of an Integrated Correlator in N = 4 Supersymmetric Yang-Mills Theory," Phys. Rev. Lett. **126**, no.16, 161601 (2021) [arXiv:2102.08305 [hep-th]].
- [51] D. Dorigoni, M. B. Green and C. Wen, "Exact properties of an integrated correlator in $\mathcal{N} = 4$ SU(N) SYM," JHEP **05**, 089 (2021) [arXiv:2102.09537 [hep-th]].
- [52] T. M. Apostol, "Modular functions and Dirichlet series in number theory." Graduate texts in mathematics; 41. Springer-Verlag, New York, 1976.
- [53] E. Hecke, "Über Modulfunktionen und die Dirichletschen Reihen mit Eulerscher Produktentwicklung. I," Math. Ann. 114 (1937) no. 1, 1–28.