

# Decay spectroscopy of heavy and superheavy nuclei

Dieter Ackermann<sup>a,1</sup>

<sup>a</sup>*Grand Accélérateur National d'Ions Lourds – GANIL, CEA/DRE-CNRS/IN2P3, 55027, F-14076 Caen, France*

---

## Abstract

After more than half a century after the first predictions of the so-called "*island of stability of superheavy nuclei*", exploring the limits of nuclear stability at highest atomic numbers is still one of the most prominent challenges in low-energy nuclear physics. These exotic nuclear species reveal their character and details of some of their properties by their induced or spontaneous disintegration. The achievements in the field superheavy nuclei (SHN) research have been reported in a number of review papers throughout the decades. This review is an attempt to summarize the progress that has been made in recent years by employing the versatile tool park of Decay Spectroscopy After Separation (DSAS) for the heaviest isotopes from  $Z=99$  (einsteinium) to  $Z=118$  (oganesson). In addition, references to earlier reviews and the relevant papers, together with the major decay properties of all these isotopes are given.

**Keywords:** Superheavy Nuclei, decay spectroscopy, nuclear structure, limits of nuclear stability, heavy ion reactions

---

*Email address:* d.ackermann@ganil.fr (Dieter Ackermann)

# Contents

<b>1</b>	<b>Dedication</b>	<b>3</b>
<b>2</b>	<b>Introduction</b>	<b>3</b>
<b>3</b>	<b>Decay spectroscopy after separation - DSAS</b>	<b>6</b>
<b>4</b>	<b>Specific nuclear structure features</b>	<b>8</b>
4.1	Single-particle states, deformation and shell gaps . . . . .	10
4.2	<i>K</i> -isomerism . . . . .	10
4.3	Decay mode competition . . . . .	15
4.3.1	Competition between <i>SF</i> and $\alpha$ decay . . . . .	15
4.3.2	Competition <i>SF</i> and $\beta$ decay . . . . .	15
<b>5</b>	<b>Nuclear properties of the isotopes of the heaviest twenty elements</b>	<b>17</b>
5.1	Einsteinium - $Z=99$ . . . . .	17
5.2	Fermium - $Z=100$ . . . . .	22
5.3	Mendelevium - $Z=101$ . . . . .	24
5.4	Nobelium - $Z=102$ . . . . .	29
5.5	Lawrencium - $Z=103$ . . . . .	33
5.6	Rutherfordium - $Z=104$ . . . . .	35
5.7	Dubnium - $Z=105$ . . . . .	42
5.8	Seaborgium - $Z=106$ . . . . .	45
5.9	Bohrium - $Z=107$ . . . . .	46
5.10	Hassium - $Z=108$ . . . . .	47
5.11	Meitnerium - $Z=109$ . . . . .	47
5.12	Darmstadtium - $Z=110$ . . . . .	48
5.13	Roentgenium - $Z=111$ . . . . .	50
5.14	Corpernicium - $Z=112$ . . . . .	50
5.15	Nihonium - $Z=113$ . . . . .	51
5.16	Flerovium - $Z=114$ . . . . .	51
5.17	Moscovium - $Z=115$ . . . . .	52
5.18	Livermorium - $Z=116$ . . . . .	53
5.19	Tennessine - $Z=117$ . . . . .	54
5.20	Oganesson - $Z=118$ . . . . .	54
<b>6</b>	<b>Outlook</b>	<b>64</b>

## 1. Dedication

A substantial part of my understanding of the nature of these fascinating and unique objects at the upper right end of the Segré chart, the superheavy nuclei was formed in the common research activities that I pursued together with **Christophe Theisen**, working on our common publications and in our numerous discussions. Writing our review paper on "*Nuclear structure features of very heavy and superheavy nuclei—tracing quantum mechanics towards the ‘island of stability’*" [1] was more pleasure than effort thanks to a fruitful and rewarding exchange of ideas with him. With his infinite wisdom and profound knowledge of all aspects of nuclear structure and reaction mechanism in heavy ion collisions, not only for the heaviest nuclear species, attacking new frontiers having Christophe as a mate, was pleasure and joy. We had exciting visions and promising projects together, based and initiated often based on his ingenious ideas and foresight, some of which just commencing now when he went missing. Christophe left us in March 2024, way too early at the height of his impressive career, in the middle of his creative and productive life. Continuing on the path he has prescribed, following his visions, we will do our best to honor his legacy. We will miss him infinitely.



Figure 1.1: Christophe Theisen  
(Foto with permit from Véronique Theisen)

I dedicate this paper to my colleague and dearest friend **Christophe Theisen**.

## 2. Introduction

The quest to explore the limits of stability of nuclear matter towards the highest atomic number  $Z$  and the so-called "*island of stability*" of spherical superheavy nuclei (SHN) is at present as popular as it was in the mid-sixties when, as a consequence of the nuclear shell model, introduced by Maria Goeppert-Mayer [5, 6], and Otto Haxel, Hans Jensen and Hans Suess [7], the first predictions for the next closed proton and neutron shells beyond  $^{208}\text{Pb}$  were proposed by Meldner [8] and Sobiczewski [9]. Already at that time the occurrence of nuclear deformation and its major consequences for nuclear stability of these heavy nuclei had been recognized by Strutinsky [10, 11]. Beyond the desire and the reward for the discovery of a new chemical element or a new nuclear isotope, the nuclear structure features of these exotic species at the limit of nuclear stability and its quantum mechanical origin, as well as their chemical and ground state properties were the traditional driving forces for manifold activities throughout the last half-century.

The advancement of the field was periodically reported in a number of reviews for the various disciplines involved, like for the synthesis of new superheavy elements (SHE) the recent review by Oganessian, Sobiczewski and Ter-Akopian [12] or concerning the synthesis and search for even- $Z$  elements by Hofmann et al. [13]. The observation of three "promising"  $\alpha$  decay signals in the latter has been critically discussed in Ref. [14].

SHE research also has a long tradition as one of the major subjects in modern nuclear chemistry. The determination of characteristic chemical properties has the potential to identify the atomic number of a hitherto unknown species. In a comprehensive and detailed overview Türler and Pershina reviewed the state of the research on the chemical properties of transactinide elements ( $Z \geq 104$ ) up to flerovium ( $Z=114$ ) [15].

Beyond the chemical applications of the properties of the atom, also atomic physics' methods like precise mass measurements and laser spectroscopy have become more and more fashionable in the last two decades. "*The quest*




Figure 2.1: Excerpt of the chart of nuclides showing the heaviest nuclei observed ( $Z=96-118$  and  $N=137-177$ ). The colors indicate the decay mode and the subdivision in areas of different sizes the relative decay probabilities. Yellow denotes  $\alpha$ -decay, green spontaneous fission (SF), red  $\beta^+$  or electron capture (EC) and blue  $\beta^-$  decay. The scheme is adopted from [2]. For  $^{267}\text{Ds}$  a faint yellow has been chosen as the observation of its  $\alpha$  decay is uncertain (see Ref.s [3, 4]). For the  $\beta$  (EC) decay of  $^{266-268}\text{Db}$  a faint red has been chosen as the competition between SF and EC is being debated (see subsection 5.7).

for superheavy elements and the limit of the periodic table" is the title of a recent review of the state and perspectives of SHE discussing in addition to the advancement in SHE chemistry, also aspects of atomic and nuclear structure features of superheavy atoms and nuclei [16]. The recent progress in laser spectroscopy of actinides, referring to the potential of this type of investigation for the heaviest nuclei, has been summarized by Block, Laatiaoui and Raeder in 2021 [17], while the capabilities of laser spectroscopic methods for the investigation of nuclear structure have been detailed and emphasized by Campbell, Moore and Parson [18]. Ion traps and Multi-Reflection Time of Flight Mass Spectrometers (MRTOF-MS) contribute substantially to the investigation of nuclear ground state properties of actinides and SHN providing nuclear binding energies by high-precision mass measurements with mass resolution values of 30 keV to 100 keV [19, 20]. Titled "Smooth trends

in fermium charge radii and the impact of shell effects" by Warbinek et al reported on the most advanced results of systematic laser spectroscopy studies of the isotopic chains for nobelium and fermium, yielding fundamental findings concerning ground state properties like charge radii and nuclear structure features like nucleon density distributions and the shell structure around the deformed proton and neutron shell gaps for  $Z = 100$  and  $N = 152$  at  $^{253}\text{Fm}$  [21]. In a dedicated special issue of the journal Nuclear Physics A reviews involving basically all disciplines pursuing SHE/SHE research had been collected in 2015 [22]. The list of papers comprises experimental as well as theory contributions, including for atomic physics mass measurements, laser spectroscopy and atomic/electronic structure theory, for chemistry liquid and gas-phase methods, and for the nuclear physics partition synthesis and reaction studies, as well as nuclear structure research.

Focussing on Decay Spectroscopy After Separation (DSAS) of SHN, the present paper is dedicated to the nuclear structure of SHN. It is partly an update of a recent review, where Christophe Theisen and I were reporting on in-beam spectroscopy as well as on DSAS [1]. In this previous work we presented the history of the discovery and synthesis of chemical elements starting from the discovery of radium and polonium by Marie Curie [23, 24] to the synthesis of SHE during the first decade of this millennium with oganesson being the highest  $Z$  and tennessine [25] the up to now last element to be synthesized [26]. We discussed the theoretical background driving the search for new elements and their properties, introduced the various nuclear decay processes and presented in detail spectroscopic methods and at that time accumulated knowledge on SHN nuclear structure by DSAS and in-beam spectroscopy. We discussed selected nuclear structure theory approaches and reported on the advances in experimental instrumentation. Here I will only briefly refer to the more detailed discussion of the historical and theoretical background in our previous work, while I will focus essentially on the recent experimental achievements and future perspectives in instrumentation.

The status of the field of SHN is illustrated by the representation of the presently known isotopes in Fig. 2.1 as an update of Fig. 19 in Ref. [1], which shows the upper right part of the chart of nuclides using the color scheme for the decay mode indication of Ref. [2] with 258 isotopes for the elements from  $Z=96$  (curium) to  $Z=118$ , 13 more than in our earlier review.

One of the nuclei missing in Fig. 19 of Ref. [1] is  $^{267}\text{Ds}$ . As it is still included in the "Karlsruher Nuclidkarte" [2] and listed in various other sources, I included it also here. It is, however, presented in a faint yellow as the observation and assignment of its  $\alpha$  decay [3] is uncertain (see Ref. [4]).

Two more darmstadtium isotopes,  $^{275,276}\text{Ds}$  were observed at the new SHE-factory of Flerov Laboratory of Nuclear reactions (FLNR) of the Joint Institute for Nuclear Research (JINR) in Dubna, Russia [27] as  $4n$  and  $5n$  fusion-evaporation channels in the reaction

$^{48}\text{Ca} + ^{232}\text{Th}$  [28, 29]. Together with  $^{276}\text{Ds}$  also its  $\alpha$ -decay daughter  $^{272}\text{Hs}$  and  $^{268}\text{Sg}$  terminating the decay sequence by  $SF$ , were observed for the first time in this experiment.

For the  $\beta$  decay(  $EC$ ), of  $^{263,266,268}\text{Db}$  a faint red has been chosen as the competition between spontaneous fission ( $SF$ ) and  $EC$  is being debated (see subsections 4.2 and 5.7). Around the possible discovery of  $^{244}\text{Md}$  and its  $\alpha$  decay daughter  $^{240}\text{Es}$  a debate has been triggered by contradicting results from GSI, Darmstadt and LBNL, Berkeley. This is discussed in detail in section 5.3.

The discovery of  $^{250}\text{No}$  by Belozerov et al. in 2003 [30], Kallunkathariyil et al. at the gas-filled separator RITU of the University of Jyväskylä cyclotron laboratory (JYFL) reported in 2020 the detection of an additional  $K$ -isomeric state [31] (see subsection 5.4).

Three of the other 6 additional isotopes are located at the upper end of the region with  $^{290}\text{Fl}$  and  $^{286}\text{Cn}$  observed by Kaji et al. in 2017 [32], and with  $^{280}\text{Ds}$  another darmstadtium isotope as an additional  $SF$  termination, populated by the new  $\alpha$  decay branch of  $^{284}\text{Cn}$  of the  $^{288}\text{Fl}$   $\alpha$ -decay chain, identified by Samark-Roth et al. in 2021 [33].

The still missing three additions to the chart of nuclides all belong to the  $\alpha$ -decay chain of  $^{244}\text{Md}$ , discovered in 2020. The debate about its properties and different assignments will be discussed in subsection 5.3 [34, 35, 36]. Its daughter and granddaughter isotopes,  $^{240}\text{Es}$  and  $^{236}\text{Bk}$ , were reported earlier in 2017 by Konki et al. [37].

More details on these thirteen recently discovered isotopes will be given in the main section of this review in section 5. In this section also the major decay properties of the 200 known isotopes of the elements from einsteinium to oganesson are listed in Table 5.1 together with references to all relevant publications.

Referring to the various disciplines connected to the properties of the heaviest atoms and nuclei, I recall here the definitions we have formulated in our previous review for the categories superheavy elements: SHE, and superheavy nuclei: SHN [1]. Atomic physics and, in particular, nuclear chemistry are predominantly interested in atomic properties

of these heavy objects and mainly talk about SHE, defining them as the transactinides with  $Z \geq 104$ . For the community interested in their nuclear structure, the quantum mechanical relevance is important. That leads to the definition of SHN as those nuclides, around and beyond fermium, where the liquid drop fission barrier drops below the zero-point-motion energy of 0.5 MeV. These species owe their existence solely to quantum mechanical effects, which makes them an ideal laboratory to investigate and define the constraints of the underlying interaction, the strong nuclear force. The access to these interesting objects is strongly governed and limited by low and with increasing  $Z$  ever decreasing production cross-sections, demanding substantial efforts in method and instrumentation development which we addressed also in detail in Ref. [1]. In this review, based on this earlier work, I will start with a brief reminder on the spectroscopic methods employed for DSAS 3 and the specific nuclear structure features in section 4 on the decay properties of the heaviest nuclei. In the main part in section 5 I will after a brief overview, present the recent findings for the isotopes of all elements from  $Z=99$  (einsteinium) to  $Z=118$  (oganesson). This review will be concluded in section 6 presenting the technological progress pursued by new facilities, being presently constructed, coming online soon or being operational since recently.

### 3. Decay spectroscopy after separation - DSAS

The general production mechanism for most of the nuclei discussed here is heavy ion fusion-evaporation leading to so-called evaporation residues (ER) with partly extremely low cross-sections for which highly efficient separation schemes are needed to suppress the primary beam and products from unwanted reactions to provide the highest sensitivity for those rare events. A variety of different ion-optical separators is used at the various laboratories, being mainly of two types: i) vacuum separators like the velocity filters SHIP of GSI [39] and SHELTS of FLNR JINR [40], or mass spectrometers like the FMA at ANL in Lemont, IL, U.S.A.; ii) gas-filled separators like RITU [41] at the cyclotron laboratory of the University of Jyväskylä in Finland, or the many gas-filled separators like DGFRS [42] of FLNR JINR in Dubna, Russia, GARIS at RIKEN in Wako, Japan [43], BGS equipped with the FIONA mass separator at LBNL, Berkeley, CA, U.S.A. [44], TASCA at GSI/FAIR [45], and AGFA [46] at ANL, or in the near future also the combined separator and mass spectrometer installation S<sup>3</sup> of GANIL SPIRAL, presently being commissioned at GANIL SPIRAL2 (see also section 6). For a detailed discussion of ion-optical separation principles and devices see our previous review [1].

After being separated from the primary beam the decay properties of the ER are studied employing comprehensive particle and photon detectors. Alpha particles, electrons or fragments from SF are typically registered by silicon detector arrays with varying granularity, while germanium detectors collect  $\gamma$  and X-ray quanta emitted in the decay process of metastable states possibly populated in the ER, and from excited states of subsequent decay products. This method of decay spectroscopy after separation is schematically shown in Fig. 2.2.

The detection principle of  $\alpha$ -decay chains and genetic correlations, a powerful tool for particle identification for hitherto unknown species is shown in this figure which is




Figure 2.2: Decay spectroscopy after separation and genetic correlations: the recoiling nucleus  ${}^A_Z X$  is implanted in a position-sensitive detector at position  $(X_r, Y_r)$ . It subsequently decays via  $\alpha$  emission in its neighborhood at position  $(X_{\alpha 1}, Y_{\alpha 1})$  to the daughter nucleus  ${}^{A-4}_{Z-2} Y$  which itself decays to the granddaughter  ${}^{A-8}_{Z-4} Y$  at position  $(X_{\alpha 2}, Y_{\alpha 2})$ . In addition to the recoils and  $\alpha$  particles  $\gamma$ -, X-rays and CEs are detected in coincidence. The technique allows correlations in position and/or time of the recoil implantation and its subsequent decay due to the inclusive detection of the particles and photons involved. (Figure and caption are taken from ref. [38])

taken from ref. [38], where more details on this powerful and well-established tool to investigate rare activities are given, including single-event detection can be found [47].

While emitted  $\alpha$  particles from g.s.-to-g.s. decays and fission fragments, correlated to the ER, implanted in the focal plane silicon detector, or the previous decay in a decay sequence are most important for the identification of the decaying isotope,  $\gamma$  rays and conversion electrons (CE), together with *alpha* decay into excited decay-daughter states, yield detailed spectroscopic information on their low-lying nuclear structure. CEs have recently also proven to be a powerful tool for additional background suppression (see, e.g., ref. [48] as well as the detection *x*-rays which in addition can also serve for efficient Z-identification. The sensitivity of this correlation method down to lowest count rates, including single event detection, is the key for an efficient accumulation of nuclear structure data for the heaviest

nuclear species through the last decades. Highlights are, e.g., the discovery of the heaviest of the heaviest  $K$  isomers  ${}^{270m}\text{Ds}$  [49] and  ${}^{266}\text{Hs}$  [50], or the  $\alpha$ - $\gamma$  decay spectroscopy of odd-even nuclei like  ${}^{247}\text{Md}$  [51, 52],  ${}^{255}\text{Lr}$  [53, 54] or even-odd nuclei  ${}^{257}\text{Rf}$  [55, 56]. The important role the unpaired nucleon is playing for exotic nuclear structure features has been underlined in ref. [57], and, in particular, for the creation of high-spin/high- $K$  isomeric states in ref. [38] (see also subsections 4.1 and 4.2).

Similar to  $\alpha$  decay, highly excited states in heavy nuclei can also be populated by  $\beta$  decay, de-exciting by a series of internal transitions proceeding via CE and  $\gamma$  emission. However, as beta decay, proceeding by electron capture (EC) or  $e^+$  (, and  $e^-$ ) emission, is difficult to detect, *x*-rays and CEs, being emitted from the populated daughter states, play a major role for its identification, as in the case of the  ${}^{253}\text{Md}$   $\beta$  decay, revealing the  $11/2^-$  [725] isomer

in  $^{253}\text{Fm}$  [58], or the first attempt to identify  $EC$ -decay of an isomeric state in  $^{257}\text{Rf}$  [55].

Short lifetimes of nuclear states in heavy nuclei, and, in particular, in nuclear isomers pose a major challenge to the separation and detection set-ups. Despite optimized installation, like, e.g., in the case of the compact design of the gas-filled separator of ATLAS ANL, AGFA, typical flight paths of the order of  $\mu\text{s}$  lead to losses in the detection of  $\mu\text{s}$  isomers. Lifetimes of this order in addition call for special measures to separate the various electronic signals from  $ER$  implantation and the subsequent decays. Thank to the advancement in electronics development in recent years, the required time resolution is achieved by achieved by the application of flash-ADCs and pulse-shape analysis. So-called "digital" data acquisition systems [59], provide this functionality which is demonstrated by successful isomer identification in  $^{254}\text{Rf}$  [60, 61] and  $^{255}\text{Rf}$  [61], or  $^{250}\text{No}$  [31, 62] (see also subsection 5.4 for the latter) (see subsections 5.6 and 5.4). More details on the role of  $CE$  and pulse shape analysis is discussed in subsection 4.3.2 on the competition of  $SF$  and  $\beta$  decay. The advanced DSAS technologies have the potential to reveal the hitherto almost completely missing information on  $\beta$  decay properties of nuclides for all elements beyond dubnium (see Fig. 2.1).

## 4. Specific nuclear structure features

As mentioned in the introduction, the heaviest nuclei owe their existence to quantum mechanics and shell effects as the macroscopic barrier against fission which can be estimated using the liquid drop model (see e.g. [63], Appendix A2), vanishes with increasing atomic number  $Z$ . Within the liquid drop model the nuclear binding energy is described by the Bethe-Weizsäcker formula:

$$\begin{aligned} M(A, Z) = & a_V A - a_C Z^2/A^{1/3} - a_S A^{2/3} \\ & + a_A (N - Z)^2/A - \delta(A, Z) \end{aligned} \quad (4.1)$$

with

$$\delta(A, Z) = \begin{cases} +\delta_0 & N \text{ and } Z \text{ even} \\ 0 & A \text{ odd} \\ -\delta_0 & N \text{ and } Z \text{ odd} \end{cases} \quad (4.2)$$

and  $\delta_0 = a_P A^{-1/2}$ .

The  $a_i$  coefficients are typically obtained by fitting experimental mass values (see e.g. [64, 65, 66, 67]).

In our previous review [1] (section 1.2.1) we discussed the limits of stability according to the liquid drop model and summarized it in Fig. 4.1. As can be seen from the blue lines in Fig. 4.1, a definition of superheavy nuclei based on the vanishing liquid drop fission barrier does not result in a defined atomic number like for the chemist's definition of SHE with  $Z=104$ , but refers rather to an area between fermium ( $Z=100$ ) and rutherfordium ( $Z=104$ ) depending also on the neutron number (blue lines in Fig. 4.1). For more details refer to Ref. [1].

The nuclear structure features of nuclei in the region spanning from fermium to darmstadtium, in particular, single particle configurations at a given deformation, meta-stable states and the competition of decay modes ( $\alpha$ ,  $\beta$ ,  $SF$  and internal transitions ( $IT$ ) with the potential to explore the limits of stability in terms if highest  $Z$  and isospin (mainly towards low neutron numbers) are presently in the focus of decay spectroscopy after separation (DSAS) [68, 1, 38].




Figure 4.1: Nuclear chart and its limits of stability deduced from the liquid-drop model. The green lines correspond to the neutron  $S_n = 0$  and proton  $S_p = 0$  drip lines, with  $S_n$  and  $S_p$  denoting the neutron and proton separation energies, respectively. The blue lines  $B_f = 0$  and  $t_{1/2} = 10^{-14}$  s correspond to a vanishing fission barrier and a spontaneous fission half-life of  $10^{-14}$  s, respectively. The red line corresponds to beta-stable nuclei (Green line). Besides magic numbers shown with lines, the horizontal dotted line corresponds to  $Z = 104$ . (Figure and caption are taken from Ref. [1])

In the vicinity of low-level-density areas and large energy gaps in the plane spanned by single particle level (SPL) energies and deformation in a Nilsson diagram representation (see e.g. Ref. [69]), the properties of the single-particle states (SPLs) can be studied with reduced perturbation by mixing of other quantum states. The region of prolate-deformed nuclei around  $N=152$  and  $Z=100$  provides access to SPLs stemming from proton orbitals like e.g.  $f_{5/2}$  and  $f_{7/2}$ , and neutron orbitals like e.g.  $d_{3/2}$  and  $h_{11/2}$  which are predicted to form the spherical proton and neutron shell closures in the region of the so-called “island of stability” [69]. Investigating those SPLs can lead to findings that support conclusions on the development of shell gaps towards sphericity where those SPLs are degenerate at or-

bital energies. Recent findings, reported in section 5, in this respect, will be discussed in the next subsection 4.1.

As mentioned above, the unpaired nucleon plays a major role here as a probe for the development of specific quantum states with deformation and single particle energy and the possible occurrence of metastable states. A specific class of those, i.e.  $K$ -isomers, and the advances in recent studies as reported in section 5 will be discussed in subsection 4.2.

A comprehensive example for the intertwined decay properties leading to networks of communicating decay sequences built competing decay modes ( $\alpha$ -,  $\beta$ -decay and  $SF$ ), is shown for the example of the  $^{258}\text{Db}$  decay in Fig. 5.21.




Figure 4.2: Definition of the  $K$  quantum number as the total projection of the sum  $j_i$  of the spin of the nucleon  $s_i$  and the orbital angular momentum  $l_i$  of all excited 2-quasi-particle states onto the symmetry axis of the nucleus. (Figure and caption are taken from Ref. [1].)

#### 4.1. Single-particle states, deformation and shell gaps

One of the major problems for the assignment and interpretation of single particle states (SPLs) is the possible mixing with neighboring states in regions of high level density. This can be, at least partly, overcome in the vicinity of gaps in the level sequence like in the fermium/nobelium region with the  $Z=100$  and  $N=152$  shell gaps. The most promising and possible most conclusive studies are, therefore, carried for isotopes along these proton and neutron numbers, as shown in the comprehensive nuclear structure compilation of Asai, Lopez-Martens and Heßberger which, e.g., investigate isotonic trends for nuclei with neutron numbers 151 and 153. One addition to this systematic investigation are the new results for  $^{255}\text{No}$  reported by Bronis et al. [48] completing the isotonic trend of the  $11/2^-$ [725] state for the  $N=153$  isotones from curium to hassium (see subsection 5.4 and Fig. 5.14). Another striking feature of  $^{255}\text{No}$  is the presence of high- $K$  isomers in this nucleus as shown in Fig. 5.15, revealed consistently by two groups independently [48, 70]. The above mentioned  $11/2^-$ [725] state plays also a significant role in  $N=151$  isotones and has been revealed in  $^{253}\text{No}$  populated by the  $\alpha$  decay from the analog state in  $^{257}\text{Rf}$  where it is isomeric due to its low excitation energy as the

second state above the ground state (see subsections 5.4 and 5.6 and Ref.[56].) A nice example, where the experimental assignment provides information on the location of the next proton and neutron shell closures beyond  $^{208}\text{Pb}$  is the excitation energy  $E^*$  assignment of two SPLs in  $^{243}\text{Es}$  which are stemming from orbitals defining a possible proton shell gap at  $Z=114$  as proposed by theory. The experimental findings for the low-lying structure in  $^{243}\text{Es}$  contradict this prediction (see subsection 5.1).

#### 4.2. $K$ -isomerism

In a recent review, we discussed the role of meta-stable states in superheavy nuclei [38], where more details can be found. Here I restrict myself to one class of isomeric states which are particularly interesting for the region of heavy and superheavy nuclei, focussed on here. The nuclei which are accessible for more detailed spectroscopic studies in the region around  $Z=100$  and  $N=152$  exhibit a substantial degree of deformation with a typical quadrupole deformation of  $\beta_2 \approx 0.25$  [71, 72]. Therefore,  $K$ -isomers play a dominant role in this region with the  $K$  quantum number being the sum of the total spin projection on the symmetry axis of the deformed nucleus

$$K = \sum_i \Omega_i = \sum_i (\Lambda_i + \Sigma_i) \quad (4.3)$$

with  $\Lambda_i$  and  $\Sigma_i$  being the projection on the nucleus' symmetry axis of the orbital angular momentum  $l_i$  and the spin of the nucleon  $s_i$ . This definition is schematically shown in Fig. 4.2.

The heaviest nuclei for which a  $K$ -isomer has been observed are  $^{270}\text{Ds}$  and its  $\alpha$ -decay daughter  $^{266}\text{Hs}$ , where the delay due to isomerism is observed in  $\alpha$ -decay times with the specific feature that the isomeric decay proceeds slower than the decay of the ground state [49, 50]. An attempt to explain these findings by a phenomenological superfluid tunneling model succeeds in reproducing the experimental data when applying a reduction of the pairing gap by a factor of 0.6 [73, 74]. For most of the  $K$ -isomers shown in Fig. 4.3 and listed in Table 4.1 the prominent decay




Figure 4.3: Update of Fig. 43 in Ref. [1]: Summary of  $K$  isomers for the heaviest nuclei at and above  $Z=96$ . For detailed properties see table 5.1 and references given therein.

mode proceeds by internal transitions proceeding by  $\gamma$  emission or internal conversion. Using the conversion electron- $\gamma$  correlation is efficiently reducing background, leading to the observation of numerous high- $K$  states in the nuclei beyond fermium, like in the recently published cases for  $^{249,251}\text{Md}$  [75] and  $^{255}\text{No}$  [48], for details see subsections 5.3 and 5.4, respectively. For these odd- $Z$ , respectively odd- $N$  nuclei, the unpaired nucleon plays an important role here. Together with the high- $j$  orbitals accessible around  $Z=100$  and  $N=152$  as shown in Fig. 5.11 for  $^{249,251}\text{Md}$  it helps to generate high- $K$  values. Apart from isomers decaying by  $\alpha$  emission or internal transitions also delayed SF has been observed as for example for  $^{258}\text{Rf}$ , where the ground state and an excited state is populated by  $\beta$ -decay ( $EC$ ) of  $^{258}\text{Db}$ , while the nature of the meta-stability has still to be clarified here [76]. The scarce existing data for  $K$  isomers decaying by  $\alpha$ -decay - apart from  $^{270}\text{Ds}$  and  $^{266}\text{Hs}$  there is only  $^{178m^2}\text{Hf}$  - and SF is discussed in a recent review by R.M. Clark [74].

After initially being found in even-even nuclei heavy nuclei like, e.g.,  $^{254}\text{No}$ , more and more  $K$  isomers are reported in odd-even or even-odd nuclei. Indeed the unpaired nucleon and the vicinity of the proton and neutron shell gaps in the nobelium-fermium region with high orbitals accessible, favor the generation of high- $K$  values which are major ingredients for the formation of meta-stable states. Nice examples are the recently discovered 3-quasi-particle isomers in  $^{249,251}\text{Md}$  [75] (see subsection 5.3) and  $^{255}\text{No}$  [48, 70] including a suggested 5-quasi-particle excitation (see subsection 5.4).

In Table 4.1 and Fig. 4.3 the six even-odd nobelium and rutherfordium isotopes, and the three mendelevium and lawrencium isotopes, for which high- $K$  isomers have been reported are presented with their major properties. Up to date no high- $K$  isomer has been reported for an odd-odd nucleus with  $Z \geq 96$ .

Table 4.1: Extension of the table of known  $K$  isomers in heavy and SHN from curium to darmstadtium, including odd- $Z/N$  and odd-odd isotopes and single-particle excitations with respect to earlier listings [77, 50].  $IT$  denotes internal transitions that can proceed via  $\gamma$  emission or internal conversion. Note of caution: the configuration assignments in many cases are often based on systematics and model assumptions, rather than on safe experimental findings. For details of those assignments see the corresponding references.  
 (Table and caption taken from [38])

state	$K^\pi$	$T_{1/2}$	$E_x$	decay mode	configuration assignments	Ref.
$^{244m}\text{Cm}$	$6^+$	34 ms	1.040 MeV	$IT$	$5/2^+[622]_\nu \otimes 7/2^+[624]_\nu$	[78]
						[79]
$^{246m}\text{Cm}$	$8^-$	-	1.179 MeV	$IT$	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[80]
$^{248m}\text{Cf}$	$\geq 5$	$>140$ ns	0.9 MeV	(IT)	-	[81]
$^{248m}\text{Fm}$	$(6^+)$	$10.1(6)$ ms	-	$IT$	two-quasiparticle	[82]
						[83]
						[84]
$^{250m}\text{Fm}$	$8^-$	$1.92(5)$ s	1.195 MeV	$IT$	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[85]
$^{253m}\text{Fm}$	$11/2^-$	$0.56(6)$ $\mu$ s	$\approx 350$ MeV	$IT$	$11/2^-[725]_\nu$	[58]
$^{256m}\text{Fm}$	$7^-$	$70(5)$ ns	1.425 MeV	$IT, SF$	$7/2^+[633]_\pi \otimes 7/2^-[514]_\pi$	[86]
$^{249m}\text{Md}$	$(19/2^-)$	$2.4(3)$ ms	$\geq 0.910$ MeV	$IT$	$7/2^-[514]_\pi \otimes 5/2^+[622]_\pi$	[75]
					$\otimes 7/2^+[624]_\nu$	
$^{251m}\text{Md}$	$(23/2^+)$	$1.37(6)$ ms	$\geq 0.844$ MeV	$IT$	$7/2^-[514]_\pi \otimes 7/2^+[624]_\nu$	[75]
					$\otimes 9/2^-[734]_\nu$	
$^{250m}\text{No}^a$	$6^+$	$34.9^{+3.9}_{-3.2}$	$\approx 1.2$ MeV	$IT, (SF)$	$5/2^+[622]_\nu \otimes 7/2^+[624]_\nu$	[87]
						[31]
$^{251m}\text{No}$	-	$\approx 2$ $\mu$ s	$> 1.7$ MeV	$IT$	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[88]
						[89]
$^{252m}\text{No}$	$8^-$	$109(3)$ ms	1.254 MeV	$IT$	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[90]
						[91]
$^{253m}\text{No}$	$\geq 23/2$	$627(5)$ $\mu$ s	$> 1.44$ MeV	$IT$	$9/2^-[734]_\nu \otimes 7/2^+[624]_\nu$	[92]
					$\otimes 7/2^+[613]_\nu$	[93]
				and/or <sup>b</sup>	$9/2^+[624]_\pi \otimes 7/2^-[514]_\pi$	[94]
					$\otimes 9/2^-[734]_\nu$	[58]

$^{254m^1}\text{No}^c$	$8^-$	$265(2)$ ms	1.296 MeV	<i>IT</i>	two-quasiparticle	[95]
$^{254m^2}\text{No}^c$	$(16^+)$	$184(3)$ $\mu\text{s}$	$\approx 2.5$ MeV	<i>IT</i>	four-quasiparticle	[96]
						[97]
						[98]
$^{255m^1}\text{No}$	$11/2^-$	$109(9)$ $\mu\text{s}$	240-300 keV	<i>IT</i>	$11/2^-[725]_\nu$	[48]
	$11/2^-$	$86(6)$ $\mu\text{s}$	$\approx 200$ keV	<i>IT</i>	$11/2^-[725]_\nu$	[99]
$^{255m^2}\text{No}^d$	$19/2-23/2$	$77(6)$ $\mu\text{s}$	1.4-1.6 MeV	<i>IT</i>	$1/2^-[521]_\pi \otimes 9/2^+[624]_\pi$	[48]
	$21/2^+$	$2(1)$ $\mu\text{s}$	$\approx 1.3$ MeV	<i>IT</i>	$1/2^-[521]_\pi \otimes 9/2^+[624]_\pi$	[99]
					$\otimes 11/2^-[725]_\nu$	
$^{255m^3}\text{No}^d$	$\geq 19/2$	$\geq 1.2_{-0.4}^{+0.6}$ $\mu\text{s}$	$\geq 1.5$ MeV	<i>IT</i>	-	[48]
	$27/2^+$	$92(13)$ $\mu\text{s}$	$\geq 1.5$ MeV	<i>IT</i>	$7/2^-[514]_\pi \otimes 9/2^+[624]_\pi$	[99]
					$\otimes 11/2^-[725]_\nu$	
$^{255m^4}\text{No}$	-	$5(1)$ $\mu\text{s}$	$\geq 2.5$ MeV	<i>IT</i>	five-quasiparticle	[99]
$^{256m}\text{No}$	$5^-/7^-$	$7.8_{-2.6}^{+8.3}$ $\mu\text{s}$	$\geq 1.1$ MeV and/or <sup>e</sup>	<i>IT</i>	$11/2^-[725]_\nu \otimes 1/2^+[620]_\nu$	[100]
					$11/2^-[725]_\nu \otimes 3/2^+[622]_\nu$	
$^{255m^2}\text{Lr}$	$(15/2)$	10-100 ns	$> 1.6$ MeV	<i>IT</i>	$1/2^-[521]_\pi \otimes 7/2^-[514]_\pi$	[101]
					$\otimes 9/2^+[624]_\pi$	
$^{255m^3}\text{Lr}^f$	$(25/2)$	$\geq 1.70(3)$ ms	$\geq 1.6$ MeV	<i>IT</i>	$7/2^-[514]_\pi \otimes 7/2^+[624]_\nu$	[102]
					$\otimes 11/2^-[725]_\nu$	[54]
						[101]
$^{253m}\text{Rf}^g$	-	$0.66_{-18}^{+40}$ ms	$\geq 1.02$ MeV	<i>IT</i>	-	[89]
	-	$\approx 0.6$ $\mu\text{s}$	-	<i>IT</i>	-	[103]
$^{254m^1}\text{Rf}$	$8^-$	$4.7(1.1)$ $\mu\text{s}$	-	<i>IT, (SF)</i>	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[60]
$^{254m^2}\text{Rf}$	$16^+$	$247(73)$ $\mu\text{s}$	-	<i>IT, (SF)</i>	$7/2^+[624]_\nu \otimes 9/2^-[734]_\nu$	[60]
					$\otimes 7/2^-[514]_\nu \otimes 9/2^+[624]_\pi$	
$^{255m^2}\text{Rf}^h$	$19/2^+$	$29_{-5}^{+7}$ $\mu\text{s}$	1.103 MeV	<i>IT</i>	$9/2^-[734]_\nu \otimes 1/2^-[521]_\pi$	[104]
					$\otimes 9/2^+[624]_\pi$	[105]
$^{255m^3}\text{Rf}^h$	$25/2^+$	$49_{-10}^{+13}$ $\mu\text{s}$	1.303 MeV	<i>IT</i>	$9/2^-[734]_\nu \otimes 7/2^-[514]_\pi$	[104]
					$\otimes 9/2^+[624]_\pi$	[105]
$^{256m^1}\text{Rf}$	$6,7$	$25(2)$ $\mu\text{s}$	$\approx 1.12$ MeV	<i>IT</i>	-	[106]
$^{256m^2}\text{Rf}$	$10^+$	$17(2)$ $\mu\text{s}$	$\approx 1.4$ MeV	<i>IT</i>	$(9/2^-[734]_\nu \otimes 11/2^-[725]_\nu)$	[106]
$^{256m^3}\text{Rf}$	-	$27(5)$ $\mu\text{s}$	2.2 MeV	<i>IT</i>	-	[106]
$^{257m^1}\text{Rf}$	$11/2^-$	$4.1(4)$ s	$\approx 75$ keV	<i>IT</i>	$11/2^-[725]_\nu$	[107]
$^{257m^2}\text{Rf}$	$(21/2^+)$	$106(6)$ $\mu\text{s}$	1.151(11) MeV	<i>IT</i>	$1/2^-[521]_\pi \otimes 9/2^+[624]_\pi$	[108]
					$\otimes 11/2^-[725]_\nu$	[107]
						[109]
$^{266m}\text{Hs}$	-	$\approx 74$ ms	$\approx 1.2$ MeV	$\alpha$	-	[110]
						[50]

$^{270m}\text{Ds}$	$9^-, 10^-$	$\approx 6 \text{ ms}$	$\approx 1.13 \text{ MeV}$	$\alpha$	$11/2^-[725]_\nu \otimes 7/2^+[613]_\nu$	[49]
				or	$11/2^-[725]_\nu \otimes 9/2^+[615]_\nu$	[50]

<sup>a</sup> values taken from Ref. [31]

<sup>b</sup> see Ref. [58]

<sup>c</sup> the configurations for the two isomers are still under debate (see corresponding references)

<sup>d</sup> obviously the assignments for  $^{255m^2}\text{No}$  and  $^{255m^3}\text{No}$  are inverted in Ref.s [48] and [99]

<sup>e</sup> see Ref. [100]

<sup>f</sup> values taken from Ref. [101]

<sup>g</sup> note: half-lives differ by three orders of magnitude for the two literature values from [89] and [103]

configurations and the values for  $T_{1/2}$  and  $E^*$  are taken from Ref. [105]

### 4.3. Decay mode competition

The quantum character of nuclear states is often crucial for the decay-mode dependent transition probability due to different hindrance effects. The decay probability of internal transitions mainly depends, apart from the transition ( $\gamma$ ) energy, on multi-polarity, difference of initial and final state spin values and transition type (electric or magnetic). It can be estimated employing relations established for electric and magnetic transitions by V.F. Weisskopf [111]. The major driving parameter for  $\alpha$  and  $\beta$  decay is the energy difference between initial and final states while the decay into analog states of equal or similar spin and parity configurations is preferred. Parity change and larger spin differences lead to an increased reduction of the transition probability, a decay hindrance. For  $SF$  the fission barrier height is crucial. In the region of heavy and superheavy nuclei, these various modes are in strong competition due to similar decay probabilities for the various decay modes. As a consequence experimentally observed branching ratios provide valuable information on the underlying quantum character of the decaying states. In the following sections, recent observations of decay mode competition between  $SF$  and  $\alpha$  decay (section 4.3.1), as well as between  $SF$  and  $\beta$  decay 4.3.2 will be discussed.

#### 4.3.1. Competition between $SF$ and $\alpha$ decay

Various theory predictions suggest a fission hindrance generated on the more or less complex quantum character of a nuclear state. In a Nilsson-Strutinsky model approach, on the basis of an average Woods-Saxon potential and a monopole pairing residual interaction, Ćwiok et al. predicted that fission is hindered for nuclei with odd nucleon numbers, in particular, for high-spin Nilsson configurations [112]. This fact is supported by experimental observations for heavy nuclear systems.

One example was reported earlier by Antalic et al. regarding the decay properties of  $^{259}\text{Sg}$  [113]. For both its  $11/2^-[725]$  ground state,  $T_{1/2} = 411$  ms, and the first excited  $1/2^+[620]$ ,  $T_{1/2} = 254$  ms,  $\alpha$  decay is observed while

the measured  $SF$  half-life is measured with 235 ms, suggesting that only the state with the low-spin configuration ( $1/2^+$ ) is fissioning while fission from the high-spin negative-parity g.s. is strongly hindered.

The recent results for the decay from g.s. and isomer in  $^{247}\text{Md}$  support a similar conclusion, as shown recently by Heßberger et al. by the comparison of the measured  $SF-\alpha$  branching ratios which differ considerably for these two Nilsson states with similarly different configurations. While the  $1/2^-[521]$  isomer decays with a probability of 20(2)% by  $SF$ , the fission branch of its  $7/2^-[514]$  g.s. amounts to a fission probability of  $8.6 \times 10^{-3}(10)$  (see subsection 5.3).

For the new data on the  $^{273,275}\text{Ds}$  and their decay products obtained at the FLNR JINR SHE-factory [29] use similar arguments to produce scenarii, based on  $SF$  hindrance arguments to construct in a first approximation decay sequences for both decay chains with decay mode, spin and level order assignments (see Fig. 5.25 and section 5.12).

#### 4.3.2. Competition $SF$ and $\beta$ decay

As mentioned in the introduction of this section,  $\beta$  decay, in particular,  $EC$  decay is difficult to detect for heavy and superheavy nuclei which is reflected in the almost lack of information on  $\beta$ -decay properties for isotopes with  $Z > 105$ , beyond dubnium (see Fig. 2.1). The same figure illustrates, however, how competitive  $\beta$  decay is for the heavy nuclei up to dubnium. This is also clearly demonstrated by for the dubnium isotopes in a systematic analysis of  $\beta$ -decay half-lives as a function of *beta*-decay energies for heavy isotopes from neptunium to dubnium in Fig. 5.23 in section 5.7. It shows that the measured half-lives deduced from the fission events that were attributed to the odd-odd dubnium isotopes at the end of the decay chains originating from the decay of moscovium and tennessine isotopes,  $^{266,268}\text{Db}$ , are competitive with  $\beta$  decay half-lives deduced from their  $Q_\beta$  values, raising some doubt about the  $SF$  assignment (see subsection 5.7). Following this argumentation in Fig. 2.1 the possible partial  $SF$  decay is indicated by faint red colored triangles for these nuclides and, for similar reasons,




Figure 4.4: Evaporation residue trace examples where a small-energy signal and/or fission were observed. Left panels show traces in a full time range (3000 channels) and right panels show the expanded view of traces, where the small-energy electron signals occurred. Corresponding time differences of signals are given relative to the preceding event. a) ER followed by fission of  $^{254}\text{Rf}$  in the analog data branch. A small-energy signal was detected after  $6.1 \mu\text{s}$ . b) ER trace, in which both small-energy and fission signals associated with  $^{254}\text{Rf}$  were detected. c) ER trace similar to b) but with two small-energy signals. d) ER followed by  $\alpha$  decay of  $^{255}\text{Rf}$  in the analog data branch. See text for details. (Figure and caption are taken from [61])

also for  $^{263}\text{Db}$  (see subsection 5.7).

The difficulties in identifying  $\beta$  decay in a decay chain can successfully overcome with the state-of-the-art DSAS detection systems with their advanced time and energy resolution capabilities, corroborated by reduced thresholds (see section 3). As shown for the  $\beta$  decay of  $^{258}\text{Db}$  to  $^{258}\text{Rf}$ ,  $CE$ s play also here a decisive role which led to the identification of two decay activities, one of them populating an excited state in the daughter nucleus [76].

A impressive example for resolution of shortest times by advanced digital electronics is the trace analysis of a small electron signal on the tail of the large signal produced by the implantation of an ER. Fig. 4.4 shows the detection of  $CE$  signals on the tail of  $ER$  signals which were successfully

used by Khuyagbaatar et al. to identify the isomers  $^{254m}\text{Rf}$  and  $^{255m}\text{Rf}$  (see subsection 5.6).

Trace analysis can also be used to measure large fission fragment energies by the time-over-threshold method as demonstrated by Khuyagbaatar et al. for the  $^{254}\text{Rf}$  decay in Fig. 5.18 [61] together with the much lower  $\alpha$  or even electron energies with a single preamplifier configuration (see subsection 5.6). The there quoted sample time of 17 ns allows for the time separation of processes in sub- $\mu\text{s}$  regime.

## 5. Nuclear properties of the isotopes of the heaviest twenty elements

The continuous history of synthesis and discovery of the isotopes from einsteinium to oganesson starts in the mid-1950ies and the most recent isotopes were discovered in 2023. The lower panel of Fig. 5.1 illustrates the development in terms of year of discovery for all 199 nuclides, shown on the background of shell-correction energies calculated in the framework of the macroscopic-microscopic model introduced by Adam Sobiczewski [114, 115]. The year of discovery, identified as the year of first publication, has been verified with the data reported in [116], including updates from the "Discovery of Nuclides Project" at:

<https://people.nscl.msu.edu/~thoenes/isotopes/>.

The upper part of the same figure shows their half-lives which have been taken from the National Nuclear Data Center (NNDC) of Brookhaven National Laboratory (BNL) via its website: <https://www.nndc.bnl.gov/> and form recent publications as listed in Table 5.1.

In both panels shell closures, predicted by the various models, for protons ( $Z=114$  and 120) and neutrons ( $N=184$ ) are indicated, together with the shell gaps of deformed SHN at  $Z=108$  and  $N=152$  and 162. From the shown nuclear half-lives only, partly due to the large range spanning from  $10^{-7}$  s to  $10^9$  s, the location of these assignments is not obvious at first glance. They become more obvious from the combined theoretical (open circles) and experimental  $Q_\alpha$  values as a function of neutron number  $N$  presented in Fig. 5.2 with pronounced minima for a broad elemental range at the deformed neutron shell gaps. In addition large gaps between the isotopic lines in the vicinity of  $N=162$  are visible at  $Z=108$ . A similar less pronounced effect is present around fermium at  $N=152$ , which confirms the shell gap at  $Z=100$  as shown in the Nilsson representation of single-particle levels as a function of deformation in Ref.s [63] (Fig. 5.4) and [69] (Fig. 4) at quadrupole deformations around 0.25. This is supported by the trend of two-neutron separation energies in Fig. 5.3 which shows a

more pronounced slope change for fermium isotopes around  $N=152$  as compared to the neighboring even- $Z$  isotopic chains. Another indication for a possible shell gap at  $Z=100$  is provided by the trend of excitation energies  $E^*$  of the first  $2^+$  states of isotopes in the vicinity of  $N=152$  (see subsection 5.2 and Fig. 5.9).

In table 5.1 basic properties, like half-lives, ground-state spin and parities (if known), decay mode branching ratios,  $Q_\alpha$  values and year of discovery for the 199 known isotopes from einsteinium to oganesson are listed together with the relevant publications. In addition to the ground states, experimentally found isomeric states indicated as  $^{Am}Z$  are also listed with their excitation energies  $E^*$ , if known. The following subsections will report on recent results for the isotopes of the elements from einsteinium to oganesson. A comprehensive collection of papers since 2017 is organized in subsections for each isotope for which new data has been accumulated, with a more detailed discussion for specifically interesting cases. For the highest  $Z$  elements, seaborgium and beyond, many isotopes are produced as members of decay chains of heavier ERs, with recent re-measurements of some of those decay chains, leading mainly to a reduction of uncertainties of their decay properties listed in Table 5.1 with updated values with respect to data provided by the NNDC website. An updated overview of the decay modes is presented in Fig. 2.1 in the introduction.

### 5.1. Einsteinium - $Z=99$

For the element with atomic mass number 99, einsteinium, a total of 18 isotopes from  $^{240}\text{Es}$  to  $^{257}\text{Es}$  has been observed. For many odd- $Z$ /odd- $A$  einsteinium isotopes the assignment of the ground state spin and parity is still open, with a unambiguous assignment being hampered by the predicted small energy difference of the g.s. candidate configurations  $7/2^+[633]$  and the  $3/2^-$ [521]. This can be seen from Fig. 5.4 which shows the state of knowledge in a global decay scheme for odd- $Z$  odd- $A$  isotopes from einsteinium to dubnium as an update of the scheme shown in [1]. While the experimental activities concerning the heavier einsteinium isotopes date




Figure 5.1: Shell correction energies from Ref.s [114, 115] as function of proton and neutron numbers with (upper panel) experimental half-lives  $T_{1/2}$  as given in table 5.1 and (lower panel) years of discovery from Ref. [116], including updates from the *Discovery of Nuclides Project* at <https://people.nscl.msu.edu/~thoennes/isotopes/>, for the heaviest nuclei from einsteinium to oganesson. The predicted closed shells on the “island of stability” for spherical nuclei SHN for protons at  $Z=114$  and  $120$ , and neutrons at  $N=84$  are indicated by solid red lines. The subshell closures or shell gaps for deformed shell-stabilized nuclei for protons at  $Z=108$ , and for neutrons at  $N=152$  and  $162$  are indicated by dashed green lines.

further back in time, the three lightest ones received quite some attention lately. The lightest two isotopes are subject of a recent debate in the community, studying the nuclear properties of the heaviest nuclear species [34, 35, 36] which will be discussed in subsection 5.3 together with the mendelevium isotopes from which they were populated via  $\alpha$  decay.

### $^{240}\text{Es}$

$^{240}\text{Es}$  is the most recent addition to the einsteinium isotopic chain and was first reported in 2017 by Konki et al. [37], together with its  $\alpha$ -decay daughter nucleus  $^{236}\text{Bk}$ . For  $^{240}\text{Es}$   $\alpha$  decay and  $EC$  with branchings of  $b_\alpha = 0.7(1)$

and  $b_{EC} = 0.3(1)$  and a half-life of  $T_{1/2} = 6(2)$  s were established while for  $^{236}\text{Bk}$  only  $EC$  decay was observed, with a half-life of  $T_{1/2} = 22^{+13}_{-6}$  s. In both cases the  $EC$  decay leads to ( $\beta/EC$ -delayed) fission (ECDF) from excited states populated in their respective decay-daughter nuclei (see Fig. 5.5). These findings complete a systematic picture of einsteinium and berkelium ECDF probabilities, showing an increasing linear trend as a function of the difference between the  $EC$ -decay  $Q$ -value and the fission barrier  $Q_{EC} - B_{SF}$  towards decreasing neutron number for the light members of their respective isotopic chains, with  $^{240}\text{Es}$  being the lightest known einsteinium isotope. For a more detailed discussion of  $\beta$ -delayed fission see Ref. [126]




Figure 5.2:  $Q_\alpha$  values from Ref. [117] as a function of neutron number for isotopes from  $Z=94$  to  $118$ , indicated by their chemical Symbols. Full circles denote experimental and open circles theoretical values. Blue dotted lines mark the predicted neutron shell closure at  $N=184$  and the deformed neutron shell gaps at  $N=152$  and  $162$ , respectively.

and updates in Ref. [127].

### $^{241}\text{Es}$

The identification of  $^{241}\text{Es}$  was first reported in 1996, produced in the irradiation of  $^{209}\text{Bi}$  by  $^{40}\text{Ar}$  at the velocity filter SHIP at GSI [120] by  $\alpha$  decay of  $^{245}\text{Md}$ . In addition to  $\alpha$  decay,  $EC$  was observed. As  $^{241}\text{Es}$  is part of the discussion in the debate mentioned in the introduction of this subsection which will be further discussed in subsection 5.3, it is attracting some interest presently after it was first observed in 1996 together with  $^{242}\text{Es}$  as  $\alpha$ -decay products of  $^{245,246}\text{Md}$  [120].

### $^{242}\text{Es}$

In 2024 Khuyagbaatar et al. reported on a new measurement of the decay properties of  $^{242}\text{Es}$  [128]. In addition to 662  $\alpha$  decays they observed 26 fission events which they assigned to  $\beta$ -( $EC$ -)delayed fission (ECDF) which

is expected to occur for heavy odd-odd nuclei with an enhanced probability, due to the fission hindrance in for odd and odd-odd heavy nuclei. The isotopic ECDF systematics regarding ECDF probabilities in relation to  $Q_{EC}$  values and fission barrier heights for einsteinium in comparison to berkelium and americium isotopic chains is shown in Fig. 5.6.

### $^{243}\text{Es}$

In a new measurement, employing for the first time the direct production reaction  $^{197}\text{Au}(^{48}\text{Ca},2n)^{243}\text{Es}$ , the earlier observed decay properties of  $^{243}\text{Es}$  have been confirmed by Briselet et al. [133]. This was part of a series of measurements performed at the cyclotron accelerator laboratory of the University of Jyväskylä (JYFL), using also  $^{203,205}\text{Tl}$  targets to populate the mendelevium isotopes  $^{249,250}\text{Md}$ , which will be discussed in subsection 5.3.

A first tentative decay scheme for  $^{243}\text{Es}$  was proposed

*Fig. 8. Two-neutron separation energies  $N = 142$  to  $165$*




Figure 5.3: Two-neutron separation energy values  $S_{2n}$  of even-Z elements from Fr ( $Z=87$ ) to Rg ( $Z=111$ ) and  $N=142$  to  $165$ . (Figure taken from Ref. [117])

in Ref. [51] from  $^{247}\text{Md}$  to  $^{239}\text{Bk}$  which is expected to decay almost exclusively by  $\beta$  ( $EC$ ) decay. New experimental information is now also available from a recent  $^{247}\text{Md}$   $\alpha$ - $\gamma$  decay study (see also subsection 5.3), which provides important information for the above mentioned ground state spin and parity assignment in a recent paper [52]. Following the earlier work [51] strong arguments for the order of the  $7/2^+[633]$  and the  $3/2^-[521]$  states have now been presented and the  $^{243}\text{Es}$  ground-state could be assigned as  $3/2^-[521]$  establishing the hitherto uncertain order in competition with the close-lying  $7/2^+[633]$ . The excitation energy  $E^*$  of the  $7/2^+[633]$  above g.s. is given as 10 keV. With this a total of six low-lying Nilsson states have been assigned tentatively (see Fig. 5.7). The  $7/2^-[514]$  level at  $E^*=219$  keV as well as the  $5/2^-[523]$  are populated by  $8421$  keV and  $8406$   $\alpha$  transitions from the  $^{247}\text{Md}$   $7/2^-[514]$

g.s. An  $\alpha$  transition of 8402 keV populates the  $3/2^-[512]$  state at  $E^*\approx 350$  keV. Finally the  $1/2^-[521]$  which is populated by a 8720 keV  $\alpha$  decay from the  $^{247m}\text{Md}$   $1/2^-[521]$ , has been placed at  $E^*=68(11)$  keV above the  $3/2^-[521]$  g.s.

A particular conclusion is drawn by the authors considering the low energy difference of  $E^*$  of  $68(11)$  keV observed for the  $3/2^-[521]$  and  $1/2^-[521]$  Nilsson levels in  $^{243}\text{E}$ . These two states have their origin in the  $2f_{7/2}$  and  $2f_{5/2}$  spin-orbit partners which according to model predictions define the proton shell gap at  $Z=114$  (see, e.g., Ref. [69]). This low energy difference of 68 keV was found to be in contradiction to an energy gap of  $>1$  MeV of these two levels at sphericity, as predicted by some microscopic-macroscopic models (see, e.g., Ref. [69]). The authors regarded this as a hint for the next spherical proton shell location being at  $Z=120$  rather than at  $Z=114$ .




Figure 5.4: Decay scheme for odd- $Z$ /even- $N$  isotopes from einsteinium to dubnium in the vicinity of the  $N=152$  closed shell updated from Ref. [1]. Data are taken from [118, 119, 120, 121, 122, 123, 53, 124, 102, 54, 108, 51, 125, 55, 34, 35, 36, 52]. For the ground state assignment for  $^{243}\text{Es}$ , the presentation of Antalic et al. has been chosen who propose two possible scenarios [51]. For the shown case with the  $\pi7/2^+[633]$  Nilsson level being the ground state,  $\delta$  would be the excitation energy of the  $\pi3/2^-[521]$  level and  $\Delta=0$ . For the opposite assignment  $\Delta$  would be the excitation energy of the  $\pi7/2^+[633]$  state. For the other einsteinium ground state assignments the representation of [124] has been adopted here. For  $^{245}\text{Md}$  and  $^{241}\text{Es}$   $\alpha$  transitions are shown as reported in the original paper ([120]). For the complete transition data see Ref. [36].

### $^{244}\text{Es}$

In 2024 Pore et al. reported on new data for its  $\alpha$  (three events) and  $\beta$  decay (four events) [134] which are in line with its previous observation in Ref.s [135] and [122], respectively.

### $^{245}\text{Es}$

In the irradiation campaign of  $^{203,205}\text{Tl}$  at JYFL mentioned above, Briselet et al. confirmed earlier measurements of the  $\alpha$ -decay half-life of  $^{245}\text{Es}$  populated by  $\alpha$  decay of  $^{249}\text{Md}$  [133].

### $^{246}\text{Es}$

$^{246}\text{Es}$  is the last member in the  $^{258}\text{Db}$  decay shown in

Fig. 5.21. In a recent study a tentative level scheme was presented on the basis of  $\gamma$  rays observed in coincidence with  $^{250,250m}\text{Md}$   $\alpha$  decays [136] (see also subsections 5.3). Despite the scarce statistics four distinct transition could be tentatively assigned, two of them originating from the g.s. and two from the isomer. The  $^{258}\text{Db}$  decay is an instructive example for the competing decays ( $\alpha$ ,  $\beta(EC)$  and  $SF$ ) building a decay-network passing across the  $N=152$  neutron shell gap and in the vicinity of  $^{252}\text{Fm}$  with passages from odd-odd to even-even nuclei granted by  $EC$ . Some of these features will be discussed in more detail for nuclei which are members of this decay network in the following subsections.




Figure 5.5: Decay scheme proposed for  $^{240}\text{Es}$  as proposed in Fig. 5 in [37].

### $^{251}\text{Es}$

Total kinetic energy (TKE) and mass distributions have recently been reported by A. Pal et al. for the fission of  $^{251}\text{Es}$  at two excitation energies, 47.6 MeV and 52.4 MeV in a comparative investigation of three nuclei including also  $^{256}\text{Fm}$  and  $^{257}\text{Md}$  [137] (see also subsections 5.2 and 5.3).

## 5.2. Fermium - $Z=100$

As discussed in the introduction of this section there are indications for a shell gap for SPLs at  $Z=100$  at quadrupole deformations around  $\beta_2$  around 0.25. With the deformed neutron shell gap at  $N=152$ ,  $^{252}\text{Fm}$  occupies a prominent position among the 19 experimentally known fermium isotopes. Decay spectroscopy studies after separation of this nucleus are, however, hampered by its half-life of 25.39(4) h. Hence our knowledge on its low lying nuclear structure is limited to the first  $2^+$  state. With respect to literature data (Ref. [138]) Asai et al. established an improved value for the excitation energy  $E(2^+)$ , employing  $\alpha$  decay of  $^{256}\text{No}$  [139]. They observed two  $\alpha$  decay lines, a main transition at  $E_\alpha = 8446$  keV and a second transition at  $E_\alpha = 8402$  keV, with improved resolution as compared to Ref. [138], from which they deduced  $E(2^+)(2_1^+ = 42.1(13)$  keV. This is the lowest  $E(2^+)$  for the fermium isotopes in the vicinity of

$N=152$  as well as for the  $N=152$  isotones around fermium supporting the assumption of a pronounced double shell gap in  $N$  and  $Z$  for  $^{252}\text{Fm}$  as shown in Fig. 5.9.

Beyond the scope of this review, yet related via the accessible nuclear g.s. properties, are the recent advances in laser spectroscopy of the heaviest nuclei for which such studies were performed. The respective review and summary papers have been mentioned in the introduction. Nevertheless, it is worthwhile to mention some of those achievements for particular cases. As the nobelium isotopes, subject of pioneering efforts of precise mass measurements and laser spectroscopy, the fermium isotopes at the proton shell gap  $Z=100$  and passing the  $N=152$  neutron shell are part of those. Therefore, the recent results on laser spectroscopy for fermium isotopes as reported by Warbinek et al. [21], are shown in Fig. 5.8, taken from this publication. It shows the laser resonance spectra, illustrating nicely the isotopic shift from which nuclear charge radii were deduced, for eight fermium isotopes between  $^{245}\text{Fm}$  and  $^{257}\text{Fm}$ . Among the missing ones is the for the nuclear physics most interesting  $^{252}\text{Fm}$  sitting on the intersection of the two subshell closures. For further details I refer Ref. [21].

### $^{243,245}\text{Fm}$

In a recent investigation of the  $\alpha$  decaying isotopes  $^{243}\text{Fm}$  and  $^{245}\text{Fm}$  the literature values for decay time and energy could be improved [141], but apart of a tentatively suggested first excited  $1/2^+[631]$  state in  $^{243}\text{Fm}$  which is based on systematic considerations, nothing is known regarding the excitation structure of these two nuclei.

### $^{244}\text{Fm}$

For the decay of the new mendelevium isotope  $^{244}\text{Md}$  (see also section 5.3) Khuyagbaatar et al. discuss in Ref. [35] the possible assignment of eight fission events which might partly be attributed to  $^{244}\text{Fm}$  fission populated by  $^{244}\text{Md}$  EC decay (see also subsection 5.3). This will be further detailed in the next section on mendelevium isotopes in the context of the earlier mentioned debate around the




Figure 5.6: Upper panels: Probabilities of ECDF in Es, Bk, and Am isotopes. Solid and open circles mark the experimental [129, 37] and calculated [130]  $P_{ECDF}$ , respectively. A retardation factor  $P_R$  used for the description of the delayed fission probabilities according to Ref. [130] and corresponding to calculated values (open circles) is given for each element. The presently measured value for  $^{242}\text{Es}$  is shown by a solid square. Arrows indicate the present upper limit for  $^{238}\text{Bk}$  and the recently reported lower limit for  $^{230}\text{Am}$  [131]. Lower panels: Theoretical  $B_f$  and  $Q_{EC}$  values taken from Refs. [132, 65] are shown as a function of the neutron number of the EC-decaying odd-odd nucleus. Dashed vertical lines mark the crossing of the  $Q_{EC}$  and  $B_f$  curves. (Figure and caption are taken from Ref. [128].




Figure 5.7: Proposed tentative decay scheme of  $^{247}\text{Md}$ . Half-lives for  $^{247g}\text{s},^{247m}\text{Md}$  are from this work [52], half-life of  $^{243}\text{Es}$  is taken from [51]. Alpha transitions are represented by full lines,  $\gamma$  transitions by dashed lines. (Figure and caption are taken from Ref. [52]).

discovery of  $^{244}\text{Md}$ .

### $^{248}\text{Fm}$

In a recent experiment at the BGS of LBNL 22  $\alpha$  decays of  $^{248}\text{Fm}$  were detected, following the decay sequence of

$^{256}\text{Db}$  [134]. Two  $\alpha$  decay energies were assigned with 7.830(47) MeV and 7.950(47) MeV which differ from the literature. Nurminia et al. had given two  $^{248}\text{Fm}$   $\alpha$  decay lines with  $E_\alpha = 7.83(2)$  MeV and 7.87(2) which were measured by employing a rotating-drum recoil detection setup which was not sensitive to electron-summing [142] in contrast to the implantation detection used by Pore et al. [134] and in other studies (see 5.1). It is, however, unclear why the higher decay energy, differing by 120 keV from the lower one was not reported in earlier investigations. Half-lives could not be determined due to the population of  $^{248}\text{Fm}$  by the undetected  $EC$  decay of  $^{248}\text{Md}$  (see subsections 5.3, 5.7).

### $^{251}\text{Fm}$

For the  $N=151$  isotope  $^{251}\text{Fm}$  the low lying level structure was proposed on the basis of  $\alpha$ -decay energy differences by Asai et al. [143]. Employing combined  $CE$  and  $\gamma$  spectroscopy Rezynkina et al. [144] found the new transitions of 192 keV from the  $1/2^+$  state to the  $5/2^-$  23.7(11)  $\mu\text{s}$ -isomer




Figure 5.8: Laser resonances observed in studies of different fermium isotopes in on-line and off-line measurements. Error bars show statistical uncertainties (one standard deviation). The solid lines are fits to the data to extract the respective centroid wave number. The detected number of  $\alpha$  events for the on-line investigated isotopes are normalized to the respective primary beam charge integral in units of 0.01 C. A more detailed zoom into the spectrum of  $^{255}\text{Fm}$  is shown as an inset in Fig. 1 in the main text (of Ref. [21]). (Figure and caption are taken from [21])

and of 200 keV from the isomer to the  $9/2^-$  ground state, confirming the decay scheme established in the earlier investigations. The isomer decay time was deduced from  $CE$  time distribution. In a systematic analysis of the neighboring in  $N=151$  and  $150$  isotones, and in comparison with theory the authors discuss particle-phonon mixing and its important role in the  $Z=100$  and  $N=152$  region.

### $^{256}\text{Fm}$

Total Kinetic Energy (TKE) and mass distributions have recently been reported by A. Pal et al. for fission of  $^{256}\text{Fm}$

at an excitation energy of 47.6 MeV in a comparative investigation of three nuclei including also  $^{251}\text{Es}$  and  $^{257}\text{Md}$  [137] (see also subsections 5.1 and 5.3).

### 5.3. Mendelevium - $Z=101$

Up to now 17 mendelevium isotopes, from  $A=244$  to 260, have been observed and studied experimentally. Recently, the lightest ones are in the focus of the community due to a debated mass assignment for  $^{244,245}\text{Md}$ . A particular feature is the observation high- $K$  isomers for the odd-even  $^{249}\text{Md}$  and  $^{251}\text{No}$  illustrating the role high-spin orbitals in the vicinity of the  $Z=100$  and  $N=152$  shell gaps.

#### $^{244,245}\text{Md}$

Among the recent mendelevium isotopes studied are  $^{245}\text{Md}$ , first observed in 1996 at the velocity filter SHIP of GSI [120], and  $^{244}\text{Md}$  reported for the first time in 2020 [34, 35]. Two independent studies presented almost simultaneously results which due to conflicting assignments were initiating the above mentioned debate. One of these experiments, employing the same reaction,  $^{40}\text{Ar} + ^{209}\text{Bi}$ , as the first experiment at GSI, was performed at the Berkeley Gas-filled Separator (BGS) [146] of the Lawrence Berkeley National Laboratory (LBNL) in Berkley, California, U.S.A., using two complementary detection set-ups [34]. The decay properties were measured at the Berkeley Gas-filled Separator's (BGS) focal plane by means of a retractable  $32 \times 32$  double-sided silicon strip detector (DSSD), while a mass assignment was obtained in a separate measurement using the FIONA set-up, a mass spectrometer consisting of a gas-stopping cell, re-acceleration, a trochoidal spectrometer and particle detection [147]. Pore et al. [34] reported six decay chains which were all assigned to the  $\alpha$ -decay chain  $^{244}\text{Md} \rightarrow ^{240}\text{Es} \rightarrow ^{236}\text{Bk}$ . The proposed scenario included two  $\alpha$  transitions for  $^{244}\text{Md}$ , with  $\alpha$ -decay energies and half-lives of  $E_\alpha = 8663(23)$  keV and  $T_{1/2} = 0.4^{+0.4}_{-0.1}$  s, and  $E_\alpha = 8306(223)$  keV and  $T_{1/2} \approx 6$  s, respectively. In the original paper Pore et al. suggested the longer lived state to be




Figure 5.9: Systematic behavior of the energies of the first  $2^+$  states in even-even nuclei. Upper panel: For isotopes of plutonium to rutherfordium with neutron number from 146 to 156. Lower panel: For isotopes with neutron numbers from 146 to 154 from plutonium to nobelium. The figure including the data points for  $^{252}\text{Fm}$  from Ref. [139] was taken from Ref. [140].

the ground state, while in a later paper by the same authors the assignment was changed to the slower transition originating from the isomeric state [44]. One of the observed decay sequences was assigned to the decay of  $^{240}\text{Es}$  with the  $^{244}\text{Md}$   $\alpha$  decay missing. These assignments were motivated by the decay properties of  $^{240}\text{Es}$  from literature [37] and the mass determination using FIONA.

The second measurement was performed at the gas-filled separator TASCA of GSI, Darmstadt, Germany [35]. Khuyagbaatar et al. employed the reaction  $^{50}\text{Ti} + ^{197}\text{Au}$  at two beam energies (231.5 MeV and 239.8 MeV at center of target) leading to excitation energies  $E^*$  of 26.2 MeV and 32.7 MeV, expected to lead to the 2- and 3-neutron fusion-evaporation channels  $^{245,244}\text{Md}$ , respectively. They observed three decay chains at the lower beam energy and attributed them to the  $2n$  fusion-evaporation channel  $^{245}\text{Md}$ . The observed decay properties were in agreement with literature data as reported in Ref. [120]. At the higher beam en-

ergy a total of ten  $\alpha$ -decay sequences were extracted. Three of them were again in agreement with an assignment to  $^{245}\text{Md}$  while the seven others showed similar decay times as the ones assigned to  $^{245}\text{Md}$  but higher  $\alpha$ -decay energies of 8.73 MeV to 8.86 MeV. With a half-life of  $0.30^{+0.1}_{-0.09}$  s they were attributed to the new isotope  $^{244}\text{Md}$ . In particular, the decay properties of the subsequent decays reproduced the results assigned by Konki et al. to  $^{240}\text{Es}$  [37], while half-lives and  $Q_\alpha$  values are almost identical to  $^{241}\text{Es}$  (see Table 5.1).

These two results are, in their respective interpretation, in contradiction. While the  $A$  determination by FIONA suggests that an object with 244 amu was observed at the BGS, the reported energy values would suggest the observation of  $^{245}\text{Md}$  decay events. In a comment to these findings, Heßberger et al. discussed the decay properties of both experiments in the context of the literature values [36]. Fig. 5.10 which is taken from [36], shows a comparison of




Figure 5.10: Summary of decays attributed to  $^{245}\text{Md}$  in Ref. [120] (squares) as well as in Ref. [35] (triangles) together with data reported by Pore et al. [34] [circles: events attributed to  $^{245}\text{Md}$  by the present authors, diamonds: events attributed to  $^{245}\text{Fm}$  or (tentatively) to  $^{246}\text{Md}$ ]. The dashed lines are to guide the eyes: the red lines represent the  $\alpha$  energies given for  $^{245}\text{Md}$  (8640, 8680 keV) in Ref. [120] and the energy given for  $^{244}\text{Md}$  (8663 keV) in Ref. [34]; the blue lines represent the  $\alpha$  energy for  $^{241}\text{Es}$  (8113 keV) given in Ref. [120] and the highest daughter energy (P5) in Ref. [34]; the purple line represents the literature value of the  $\alpha$  energy of  $^{241}\text{Cf}$  (7335 keV) [145]. The orange hatched area marks the range of  $\alpha$  energies where the events attributed to  $^{244}\text{Md}$  in Ref. [35] were observed. (Figure and caption are taken from Ref. [36])

the  $\alpha$ -decay data for the original and the two new measurements and proposed re-assignments of the BGS decay chains to  $^{245}\text{Md}$  and other possible reaction channels (see figure caption). As an additional argument Hefberger et

al. analyzed the fusion-evaporation excitation function in comparison to calculations performed with the statistical model code HIVAP [148], suggesting that the beam energy used by Pore et al. would favor a  $4n$  fusion-evaporation channel corresponding to  $^{245}\text{Md}$ . The beam energy for the reaction  $^{40}\text{Ar} + ^{209}\text{Bi}$  chosen for the LBNL experiment was given as approximately 220 MeV in Ref. [34] corresponding to  $E^* \approx 45$  MeV.

To clarify this situation more statistics, but in particular, instead of the two-step procedure as applied in the case of FIONA, a simultaneous measurement of  $A$ ,  $E_\alpha$  and decay times would be extremely useful, using a suited set-up, like e.g. the Fragment Mass Analyzer FMA ANL or in future the separator-spectrometer set-up S<sup>3</sup> [149], presently under construction at GANIL/SPIRAL2.

In addition to  $\alpha$  decay, fission is a common and expected decay mode in that region of the chart of nuclides. Pore et al. did not report any fission events, partly due to instrumental insensitivity, while Khuyagbaatar et al. observed at total of six and eleven fission events at the two applied beam energies, respectively, part of which could be attributed to the fission activity of  $^{245}\text{Md}$  and to fission of its isomeric state. For eight fission events, however, the situation remains somewhat unclear.

Three possible scenarios are discussed none of which could be ruled out or definitely confirmed. The possibility of direct fission decay of  $^{244}\text{Fm}$  populated in a  $p2n$  fusion-evaporation reaction (emission of one proton and two neutrons from the excited compound nucleus (CN)) and fission from an excited state in  $^{244}\text{Md}$  are in contradiction to all earlier observations. For the first case the decay times of the observed fission decays ( $5_{-2}^{+3}$  ms) and the literature values of  $^{244}\text{Fm}$  (3.21(8) ms) would match. However,  $p$ -evaporation channels are known to be strongly suppressed in reactions of this kind, as recently shown for example by Lopez-Martens et al. irradiating a  $^{209}\text{Bi}$  target with  $^{50}\text{Ti}$  projectiles for the production  $^{256,258}\text{Rf}$  as  $p$  and  $p2n$  evaporation channels at the velocity filter SHELS of FLNR JINR, Dubna [150] (see also section 5.6 and 5.7). For odd nuclei fission is known to

be strongly hindered and the observation of fission from an excited state in the odd-odd nucleus  $^{244}\text{Md}$ , in particular, in comparison with the fission properties of neighboring nuclei is rather unlikely.

As a third scenario the authors discussed *EC*-delayed fission (ECDF) from a short-lived isomeric state for which often de-excitation by CE is observed. As a hint for the existence of such an isomeric state a low-energy signal was observed in the tail of one evaporation residue (ER) signal trace, registered by the digital electronics system used in this experiment, which was correlated to a  $^{244}\text{Md}$   $\alpha$  decay. In addition two fission-like signals of similar decay time were observed in the tail of ER-like events which could originate from such a short-lived state. The evidence is, according to Khuyagbaatar et al., still too weak to firmly assign this observation to  $^{244}\text{Md}$ - $^{244}\text{Fm}$  ECDF. They conclude that all three scenarios could contribute to the observed fission events of unclear origin, which needs certainly further confirmation, given the rather speculative assumptions and counter-indications.

### $^{247}\text{Md}$

In 1981  $^{247}\text{Md}$  was first observed at the velocity filter SHIP of GSI, Darmstadt [151], Germany. In a number of subsequent investigations employing  $\alpha$ - and  $\alpha$ - $\gamma$  decay spectroscopy more details of the low-lying nuclear structure of this odd- $Z$ /even- $N$  nucleus, including an isomeric state were revealed [152, 124, 51]. In a recent decay study at the same instrument [52] the excitation energy of the 231(30) ms isomer could be established with 153 keV. For the level structure deduced for the daughter nucleus  $^{243}\text{Es}$  from this decay study see subsection 5.1.

In addition to population of low-lying levels, Heßberger et al. could establish the effect of the quantum structure of the g.s. and isomer,  $^{247g.s.}\text{Md}$  and  $^{247m}\text{Md}$ , on the fission probability of those states. The relatively high spin of the Nilsson 7/2 $^-$ [517] g.s. configuration results in the low  $SF$  branching ratio  $b_{SF}=8.6\times 10^{-3} \pm 0.0010$  as compared to the isomeric 1/2 $^-$ [521] state with  $b_{SF}=0.20(2)$ . The

corresponding partial spontaneous fission half-lives were established as  $\approx 140$  s and  $\approx 1.2$  s for  $^{247g.s.}\text{Md}$  and  $^{247m}\text{Md}$ , respectively.

### $^{248}\text{Md}$

New data which are generally in agreement with previous observations on the  $\alpha$  and  $\beta$  decay of  $^{248}\text{Md}$  as a member of the decay chain of  $^{256}\text{Db}$  were recently accumulated at the BGS of LBNL [134]. Some ambiguities in the assignment of  $\alpha$  decays to either  $^{248}\text{Md}$  or its  $\beta$  decay daughter  $^{248}\text{Fm}$  from  $^{252}\text{No}$ , the  $^{252}\text{Lr}$   $\beta$  decay daughter, lead to different branching estimates. The authors of this investigation propose on the basis of two different assumptions for the  $^{252}\text{Lr}$  *EC* branch with 10% or 30%(see subsection 5.5, an  $\alpha$  branching for  $^{248}\text{Md}$  of 61(16)% or 68(22)%, respectively, which in any case overlap within the (large) error bars, and are both slightly higher then the previously adopted value of 58% (see Table 5.1).

### $^{249,251}\text{Md}$

For  $^{249}\text{Md}$  the possible existence of an isomeric state was first proposed by Heßberger et al. based on the  $\alpha$ -decay population scheme from two different states in  $^{253}\text{Lr}$  [122].

A recent series of experiments was conducted at the the gas-filled separator RITU of the cyclotron accelerator laboratory of the University of Jyväskylä. In this context Briselet and co-workers reported on in-beam spectroscopy results for  $^{249,251}\text{Md}$  collected with the RITU target area detection system for electron and  $\gamma$  spectroscopy SAGE [133, 153]. As mentioned in subsection 5.1, these mendelevium isotopes were produced employing targets of  $^{203,205}\text{Tl}$  in the reactions  $^{203,205}\text{Tl}(^{48}\text{Ca},2n)^{249,251}\text{Md}$ .

In continuation of this study, Goigoux et al. reported hitherto unobserved  $K$  isomers in  $^{249,251}\text{Md}$  [75] revealed in the same experiment series, using the combined silicon-germanium detector array GREAT in the RITU focal plane. They investigated the decay of these two isotopes by means of recoil- $\alpha$ -CE- $\gamma$  coincidences. For  $^{249}\text{Md}$  a fast activity of 2.4(3) ms with a lower excitation energy limit of  $E^* \geq$

910 keV could be established from recoil- $\alpha$ -e<sup>-</sup> correlations. For <sup>251</sup>Md a state of 1.37(6) s half-life and a lower excitation energy limit of  $E^* \geq 844$  keV was found applying the same method, however, with the need of the condition of a subsequent  $\alpha$  detection in addition to the recoil-e<sup>-</sup> correlation, due to the slower decay time overlapping partly with random-correlations.




Figure 5.11: Schematic single-particle configurations for the g.s. and isomeric state in <sup>249,251</sup>Md. (Figure and caption are taken from [75])

For the assignment of the new metastable states to  $K$ -isomers formed by 3-quasi-particle states the authors use a microscopic-macroscopic model as described by Muntian et al. [154] based for the macroscopic component on a Yukawa-plus-exponential folding function [155] and on a deformed Woods-Saxon single particle potential for the shell correction energies [156]. For both isotopes the resulting proton configuration is the same with the unpaired proton in the highest occupied SPL 7/2-[514]. With 148 and 150 neutrons, respectively, <sup>249</sup>Md and <sup>251</sup>Md are 2 and 4 neutrons short of the  $N=152$  shell gap. Thanks to this, they have access to the three SPLs below the shell gap with relatively high spin values (5/2, 7/2 and 9/2; see Fig. 5.11) which leads to the high  $K$ -values generated by the quasi-particle excitations. A schematic view of the single-particle configurations is given in Fig. 5.11.

Two approaches within the microscopic-macroscopic model framework, one using a level blocking scheme and a second one applying a quasi-particle method (for details see [75] and references therein), provide consistently arguments to extract energy and spin/parity arguments for

the assignment of the SPLs involved in the 3-quasi-particle configuration. For both nuclei the combination of the unpaired proton in its original g.s. level, breaking the pair in the last occupied neutron level and elevating one of the neutrons into the next higher level turns out to be the most likely configuration for the respective  $K$ -isomer. This leads to  $\pi\nu^2=19/2^-$  with  $\pi7/2^-[514]\otimes\nu5/2^+[622]\otimes\nu7/2^+[624]$  and  $\pi\nu^2=23/2^+$  with  $\pi7/2^-[514]\otimes\nu7/2^+[624]\otimes\nu9/2^-[734]$  for the new 3-quasiparticle  $K$  isomers in <sup>249</sup>Md and <sup>251</sup>Md, respectively.

### <sup>250</sup>Md

As <sup>246</sup>Es and <sup>254</sup>Lr (see subsections 5.1 and 5.5), also the <sup>258</sup>Db decays chain member <sup>250</sup>Md was subject of the investigations reported in Ref. [136]. Analyzing the decay times of partitions of the relatively broad distribution of  $\alpha$ -decay energies, the authors could establish, apart from the g.s. decay with a half-life of  $25^{+10}_{-5}$  s, a second activity which was attributed to an isomeric state with  $T_{1/2}=42.2(45)$  s at an excitation energy of  $E^*=123$  keV. The ratio of the relative population of these two states by  $\alpha$  decay is found to be different for correlations to g.s. and isomer decays from the earlier members in the decay chain, <sup>254</sup>Lr and <sup>258</sup>Db (see subsections 5.5 and 5.7). This is probably related to the quantum-mechanical character of the involved states with  $\alpha$  decay preferable connecting analog initial and final states. The accumulated statistics, however, is still too low to draw firm conclusions on spins and parities.

### <sup>257</sup>Md

TKE and mass distributions have been reported by A. Pal et al. for fission of <sup>257</sup>Md at three excitation energies, 37.5 MeV, 42.9 MeV and 47.6 MeV in a comparative investigation of three nuclei including also <sup>251</sup>Es and <sup>256</sup>Fm [137] (see also subsections 5.1 and 5.2). In contrast to the other two nuclei which show symmetric TKE distributions, <sup>257</sup>Md shows a skewness extending to higher TKE which is interpreted by the authors as an indication of a new so-called

"supershort" fission mode, originating from a compact, non-stretched configuration of the fissioning system at the scission point.

#### 5.4. Nobelium - $Z=102$

Being just a proton-pair above fermium, the nobelium isotopes and, in particular,  $^{254}\text{No}$  due to its relatively large production cross-section, are in the center of interest regarding decay spectroscopy as well as in-beam studies since more than two decades. In our review from 2017 [1] nobelium isotopes played a central role, in particular, regarding in-beam studies. We dedicated a section to what we called "*The  $^{254}\text{No}$  breakthrough*" which was a boost for the nuclear structure studies in the fermium/nobelium region of the Segré chart and beyond. In-beam spectroscopy resulted in the observation of rotational bands up to  $^{256}\text{Rf}$ , while DSAS efforts succeeded in revealing features like the  $K$  isomers in heavy nuclei up to  $^{270}\text{Ds}$  (see subsections 4.2 and 5.12). The relatively high production cross-section of  $^{254}\text{No}$  in the fusion-evaporation reaction  $^{208}\text{Pb}(^{48}\text{Ca}, 2n)^{254}\text{No}$  of  $\approx 2 \mu\text{b}$  made detailed in-beam studies of this nucleus possible. A natural link between in-beam spectroscopy and DSAS is provided by isomeric states which survive the flight path (typically a couple of  $\mu\text{s}$ ) through the separator, connecting the single particle features of a nucleus to its collective behavior. Among the twelve known nobelium isotopes from  $^{250}\text{No}$  to  $^{262}\text{No}$ , the  $K$ -isomers of  $^{252}\text{No}$  and  $^{254}\text{No}$  are textbook cases for this link, as demonstrated in a number of exemplary studies [90, 91, 96, 97, 98].

The pertinent role of nobelium isotopes for the nuclear structure of heavy nuclei triggered also the investigation of their fundamental properties. In a seminal work by Block et al. [160] the atomic masses of  $^{252,253,254}\text{No}$  had been precisely measured employing the Penning trap installation SHIPTRAP of GSI. As mentioned in the introduction of this review, an overview of mass measurements for the heaviest nuclei can be found in references [19, 20]. Raeder et al. [159] measured the isotopic shift of the laser-excited atomic  $^1P_1$  level in these isotopes from which the could

establish nuclear charge radii as in the case of fermium isotopes mentioned above (see subsection 5.2). In Fig. 5.12, where also the laser excitation spectra are shown for  $^{252,253,254}\text{No}$ , measured charge radii for isotopes of the even- $Z$  element from curium to nobelium are compared with the predictions of density functional models [158, 157].

##### $^{249}\text{No}$

In 2021 almost contemporaneously two groups reported the discovery of the lightest of the thirteen known nobelium isotopes,  $^{249}\text{No}$ . Khuyagbaatar et al. [103] observed three decay chains of three correlated  $\alpha$ -decays, starting from  $^{253}\text{Rf}$  for which hitherto only a fission branch was known. The third  $\alpha$ -decay energy in this sequence was found to be in good agreement with literature data for  $^{245}\text{Fm}$  (see subsection 5.2 and references in Table 5.1) and confirmed this assignment, resulting in the discovery of a new  $\alpha$ -decay branch for  $^{253}\text{Rf}$  and the new isotope  $^{249}\text{No}$ . The latter was directly produced in the reaction  $^{48}\text{Ca} + ^{204}\text{Pb}$  by Svirikhin et al. [162] who accumulated a total of about 220  $^{249}\text{No}$   $\alpha$  decays. The decay energies and times for this new isotope reported by the two groups are in good agreement.

##### $^{250}\text{No}$

For  $^{250}\text{No}$ , Kallunkathariyil et al. reported recently on half-life measurements for its ground and  $K$ -isomeric state performed at the gas-filled separator RITU of the University of Jyväskylä cyclotron laboratory [31]. On the basis of *ERCE-SF* coincidence measurements they found a half-life for the ground state of  $T_{1/2} = 3.8(3) \mu\text{s}$  and for the  $K$  isomer of  $T_{1/2} = 34.9^{+3.9}_{-3.2} \mu\text{s}$ . The rare occurrence of isomeric states being longer lived than the ground state, here for the neutron-deficient  $^{250}\text{No}$ , is also observed for the superheavy isotopes  $^{266}\text{Hs}$  and  $^{270}\text{Ds}$  (see subsection 4.2).

Apart from electron and time spectra, Kallunkathariyil et al. also show a spectrum of photons in coincidence with *CES* which they associate with the de-excitation of the isomer. It contains, apart from nobelium  $x$ -rays, also some strength in between 150 keV and 200 keV just above the no-




Figure 5.12: a) Measured (atomic) excitation spectra of the  $^1\text{P}_1$  level for the isotopes  $^{254}\text{No}$ ,  $^{253}\text{No}$ , and  $^{252}\text{No}$  with a best fit to the data (solid line). The dashed line represents the center of each resonance, while the solid vertical lines in the  $^{253}\text{No}$  spectrum indicate the position and strength of the individual hyperfine structure components with total angular momentum  $F=7/2$ ,  $9/2$ , and  $11/2$  at 3.99 GHz, 4.10 GHz, and 10.74 GHz, respectively. The inset shows a schematic ionization scheme. b) The change in the nuclear mean square charge radii  $\delta\langle r^2 \rangle$ , for  $^{252-254}\text{No}$  and even Z actinide nuclei starting from thorium, is plotted as a function of the neutron number with arbitrary offset. For each element, the DFT calculations with two Skyrme energy density functionals, UNEDF1 [157] (dashed line) and SV-min [158](solid line), are shown. The slope according to a schematic droplet model assuming a constant deformation for the actinide elements referenced to  $N=138$  is marked in gray. (Figures and captions are taken from [159])




Figure 5.13: Compilation of the data on average number of neutrons in spontaneous fissions of different heavy nuclei, including those for the  $^{254}\text{No}$  nucleus reported in [161]. The measurements carried out with the VASSILISSA and SHELS separators are denoted by diamonds. (Figure and caption are taken from [161])

belium  $K_\alpha$  and  $K_\beta$  x-ray lines. In addition some strength is being observed in the region 250-300 keV and around 400 keV. In an experiment performed recently at the SHELS

separator of FLNR JINR in Dubna [165], Kuznetsova et al. also reported on  $\gamma$  rays observed for the  $^{250}\text{No}$   $K$  isomer, this time in coincidence with the detected  $SF$  only.

With a total of 18,000 SF events, Kuznetsova et al. observe various  $\gamma$  lines in coincidence with  $SF$  which could correspond to a consistent scenario of the isomer decay into the g.s. rotational band. However, the accumulated statistics with  $\approx 5$  to 10 counts peak height for the observed transitions is too low to confirm possible assignments by more sensitive tests, like e.g.  $\gamma$ - $\gamma$  coincidences. A comparison of the results from these two independent studies is difficult as the coincidence criteria are different. Ideally, an experiment aiming at higher count rates and combining  $CE$ -coincidence and  $\gamma$  decay spectroscopy could help to shed light on the quantum nature and decay of this neutron-deficient nobelium isotope. Such an investigation could possibly help to reveal the mechanism responsible for the enhanced stability of the isomeric state with respect to the ground state which is observed by now in a number of cases.




Figure 5.14: (a) Experimental single particle level systematics of  $N=153$  isotones. Theoretical calculations of single particle levels based on self-consistent models (b) [163] and (c) [164] or microscopic-macroscopic models (d) [112] and (e) [72] are given for comparison. Dashed horizontal lines denote the range of the excitation energy for the  $11/2^-[725]$  level (260 – 320 keV) in  $^{255}\text{No}$  assigned in [48]. (Figure and caption are taken from [48])

### $^{251}\text{No}$

A new isomeric state in  $^{251}\text{No}$  was observed by Lopez-Martens et al. in the irradiation of an isotopically enriched  $^{204}\text{Pb}$  target by  $^{50}\text{Ti}$  projectiles [89]. The spin and parity assignment of  $1/2^+$  was supported by systematics involving the neighboring isotopes and, in particular, by the observation of isomeric states in  $^{253}\text{Rf}$  reported in the same paper and in Ref. [103] (see subsection 5.6).

### $^{252,254}\text{No}$

Despite the efforts and findings regarding  $^{254}\text{No}$  reported above, there are still open questions which are lively debated like the structure of the second  $K$ -isomer in  $^{254}\text{No}$ . Kuznetsova et al. [165] and in a recent paper by the same group, Isaev et al., [161] reported on the investigation of the fission properties of  $^{252,254}\text{No}$  at SHELs. While in the earlier brief report mainly decay times and TKE measurements are mentioned, the more recent work presents more detailed results including for the first time for these isotopes neutron

multiplicity distributions. A summary of experimentally obtained neutron multiplicities for isotopes from plutonium to rutherfordium are shown in Fig. 5.13.

For  $^{254}\text{No}$  Forge et al. accumulated additional decay data [166] in a measurement using the DSAS set-up GABRIELA [167, 168] at the focal plane of the velocity filter SHELS [169] of FLNR JINR, Dubna. They report on the observation of a strong transition from the  $8^-$  isomer,  $^{254m1}\text{No}$ , which was revealed by employing  $CE$  correlations. Suggesting  $E0$  as the most likely transition type, they discuss a possible shape co-existence and presence of a low-energy super-deformed state.

### $^{253}\text{No}$

From the isotopic shift, obtained by laser excitation of the atomic  $^1p_1$  state in  $^{252,253,254}\text{No}$  and the hyperfine splitting in  $^{253}\text{No}$  Raeder et al. could determine experimentally the magnetic moment

$$\frac{\mu}{\mu_N} = g_K \frac{I^2}{I+1} + g_R \frac{I}{I+1}, \quad (5.1)$$

for  $^{253}\text{No}$ , where  $g_K$  and  $g_R$  are the intrinsic single-particle and rotational  $g$ -factor, respectively. With the range for  $g_R$  of  $0.7\text{--}1 \times Z/A$  and  $g_K^{\text{exp}} = -0.22(5)$  the Nilsson configuration  $9/2^-[734]$  obtained from in-beam  $CE$  and  $\gamma$  spectroscopy [170, 171] was confirmed while the value  $g_K = -1.2$  from an earlier study [172] was ruled out.

The  $\alpha$  decay of  $^{257m}\text{Rf}$  with the Nilsson configuration  $11/2^-[725]$  is known to populate the ground state and excited states of  $^{253}\text{No}$  with two prominent transitions differing by  $\approx 750$  keV as observed by Heßberger et al. [55]. Two  $\alpha$ - $\gamma$  correlations observed in this study could, however, not be assigned to the internal electromagnetic transition which was expected to link the isomer to its homologue state in the decay daughter nucleus  $^{253}\text{No}$ . Accumulating substantially more statistics, Hauschild et al. detected an evident  $\gamma$  line with  $E_\gamma = 750$  keV correlated to the  $\alpha$  decay transition with the lower decay energy of  $E_\alpha = 8986(5)$  MeV [56], which was identified as the searched for  $11/2^-[725]$  Nilsson state. This state, originating from the  $j15/2$  neutron orbital, has been observed for all the  $N=153$  isotones from curium to hassium [68], completed by the recent result for  $^{255}\text{No}$  discussed below (see Fig. 5.14). For the  $N=151$  isotones which have  $9/2^-[734]$  as the g.s. Nilsson configuration, stemming also from the  $j15/2$  neutron orbital,  $^{253}\text{No}$  is the second  $N=151$  isotope after  $^{255}\text{Rf}$  for which this state is observed.

### $^{255}\text{No}$

An isomeric state had been tentatively assigned for  $^{255}\text{No}$  first in 2006 [173]. Re-analyzing these data, employing conversion electrons to construct  $\alpha$ - $CE$ - $\gamma$  correlations Bronis et al. proposed in 2022 a tentative level including three isomeric states [48]. The lowest lying isomeric state,  $^{255m1}\text{No}$ , they tentatively assigned to the  $\nu 11/2^-[725]$  Nilsson state, placed in an excitation energy range of 240 keV to 300 keV and decaying with a half-life of  $109(9)$   $\mu\text{s}$ . With this assignment the systematics of  $N=153$  isotones from  $^{249}\text{Cf}$  to  $^{269}\text{Sg}$  is completed with the  $\nu 11/2^-[725]$  state placed at an excitation energy in between its analog states in  $^{253}\text{Fm}$  and

in  $^{257}\text{Rf}$  (see Fig. 5.14). For the second isomer  $^{255m2}\text{No}$  with a measured half-life of  $77(6)$   $\mu\text{s}$ , a spin of  $21/2$  or  $23/2$  and an excitation energy range of 1400 keV to 1600 keV are estimated. The third isomer,  $^{255m3}\text{No}$ , is the shortest lived with  $1.2_{-0.4}^{+0.6}$  keV and is expected to have an excitation energy of  $E^* \geq 1500$  keV.

These assignments of isomeric states in  $^{255}\text{No}$  are in fair agreement with results obtained recently at the velocity separator SHELS of the FLNR JINR by Kessaci et al. [70]. Fig. 5.15 shows in addition to the three isomers reported by Bronis et al. a forth meta-stable state at higher excitation energy,  $E^* > 2.5$  MeV, interpreted as a 5-quasi-particle excitation. The lowest isomer is also here assigned to the same  $11/2^-[725]$  Nilsson level as a 1-quasi-particle excitation. For isomer II and III in their nomenclature, interpreted as 3-quasi-particle excitations, Kessaci et al. assigned spin values, based on the  $11/2^-[725]$  state coupled to  $5^-$  and  $8^-$ , i.e.  $21/2$  and  $27/2$ , respectively, which are close to the tentative suggestions made by Bronis et al.

### $^{256}\text{No}$

In an irradiation of  $^{238}\text{U}$  with a  $^{22}\text{Ne}$  beam Kessaci et al. populated by via the 4-neutron fusion-evaporation channel  $^{256}\text{No}$  as ER [100]. They observed 15 ER- $e^-$ - $\alpha$  correlations with an average decay time of  $7.8_{-2.6}^{+8.3}$   $\mu\text{s}$  which was interpreted as a high- $K$  isomer decay. A second scenario assuming a truncation of the time distribution for short times and applying the maximum likelihood estimate as proposed by K.-H. Schmidt et al. [174] resulted in a slow decay component of  $10.9_{-4.3}^{+21.7}$   $\mu\text{s}$  and a fast one of  $\approx 6$   $\mu\text{s}$ . A possible interpretation of the existence of two isomeric states in  $^{256}\text{No}$  has to be confirmed in a future experiment aiming at substantially higher statistics. In Table 5.1 the average decay-time value was adopted. The observation of  $\gamma$  and  $x$  rays in coincidence with the isomer decays led to a lower limit for its excitation energy of 1089 keV. An assignment of the isomer's quasi-particle excitation configuration with the relative spin and parities needs further experimental efforts.




Figure 5.15: Proposed decay scheme of  $^{255}\text{No}$ . One can see the four isomeric states observed denoted in red along with the intermediate structures (a)–(c) discussed. The observed  $\gamma$ -ray transitions are denoted in blue while the unobserved ones are denoted in gray and transitions deexciting the isomers are denoted in purple. (Figure and caption taken Ref. [70])

## 5.5. Lawrencium - $Z=103$

The heaviest of the fourteen known lawrencium isotopes is the end-point of the  $^{294}\text{Ts}$  ( $Z=117$ ) decay chain  $^{266}\text{Lr}$  [175, 176] and the nuclide with the lowest  $Z$  populated in a hot-fusion or  $^{48}\text{Ca}$ -induced reaction, leading to the heaviest nuclides with highest  $Z$  and  $N$  observed so far. The lawrencium isotopes with their unpaired proton provide a sensitive probe to investigate SPLs and  $K$ -isomerism, in particular, in the vicinity of the  $N=152$  deformed shell gap with  $^{255}\text{Lr}$ .

### $^{251}\text{Lr}$

In 2022 Huang et al. reported the discovery of the new isotope  $^{251}\text{Lr}$ , produced as the  $2n$  fusion-evaporation channel in the reaction  $^{50}\text{Ti} + ^{203}\text{Tl}$  at the gas-filled separator AGFA

of Argonne National Laboratory's linear accelerator facility ATLAS [177]. In addition to the ground state  $\alpha$  decay they observed an isomeric state decaying as well by  $\alpha$  emission. The spin and parity assignments, with  $7/2^+$  for the g.s. and  $1/2^-$  for the excited isomer, and the decay scheme for the new pair  $^{251}\text{Lr}$ - $^{247}\text{Md}$  are found to be similar to the neighboring isotopes  $^{253}\text{Lr}$ - $^{249}\text{Md}$  (also investigated in this work) and  $^{255}\text{Lr}$ - $^{251}\text{Md}$  (see Fig. 5.4).

### $^{252}\text{Lr}$

In an irradiation of  $^{206}\text{Pb}$  with  $^{51}\text{V}$  projectiles at the BGS of LBNL Pore et al. accumulated new data for  $^{252}\text{Lr}$ , observed as member of the  $^{256}\text{Db}$  decay chain [134]. The authors report on the detection of a previously unknown  $EC$  decay branch for  $^{252}\text{Lr}$ , based on one safe assignment

and additional four events which could not be assigned unambiguously. Pore et al. interpret this in terms of a probability range which leads them to propose a range for the  $\beta$  decay branching ratio of  $b_\beta = 10\%$  to 30%. Despite this uncertain assignment these values are reported in Table 5.1. However, as this assignment is uncertain which is in addition also affected by the low number of events (1 to 5), these numbers should be taken with caution.

### $^{253,254}\text{Lr}$

Ito et al. measured the g.s. masses of  $^{253,254}\text{Lr}$  together with other nuclides in the region around  $Z=100$  and  $N=152$  for the first time using a Multi-Reflection Time-of-Flight mass spectrometer (MRTToF MS) installed in the focal plane of the gas-filled separator GARIS II operated at the Nishina Accellerator Centre of RIKEN, Wako, Japan [178].




Figure 5.16: Possible ground state configuration for  $^{254}\text{Lr}$  in a single-particle level scheme for protons (left) and neutrons (right) extracted from Ref. [69].

### $^{254}\text{Lr}$

As  $^{246}\text{Es}$  and  $^{250}\text{Md}$  (see subsections 5.1 and 5.3)  $^{254}\text{Lr}$  is a member of the  $^{258}\text{No}$   $\alpha$ -decay chain. Vostinar et al. constructed a low-lying level scheme combining states populated by  $\alpha$  decay of the ground and isomeric state of  $^{258}\text{Db}$  (see subsection 5.7) as well as by the observed  $\alpha$ -decay transitions of  $^{254}\text{Lr}$  itself [136]. The observation of a previously proposed isomeric state could be confirmed by  $\alpha$ -decay curves for both activities with half-lives of 11.9(9) s for the g.s. and 20.3(42) s for the isomer, respectively. The literature value for the  $^{254}\text{Lr}$  lies with 18.4(18) s in between these values. For Table 5.1 the values of Ref. [136] have been adopted. Spin and parities for  $^{254g.s.}\text{Lr}$  and  $^{254m}\text{Lr}$

were tentatively assigned as  $J^\pi=4^+$  and  $1^-$ , respectively.

The excitation energy of the isomer with  $E^*=108$  keV has been confirmed by precision mass measurements performed at the Penning trap installation SHIPTRAP at GSI [179].

Fig. 5.16 shows a possible ground-state configuration for  $^{254}\text{Lr}$  with single-particle levels in the vicinity of  $Z=100$  and  $N=152$  extracted from Ref. [69] for a quadrupole deformation of  $\approx 0.25$ . Various scenarii for spin and parity assignments to  $^{254g.s.}\text{Lr}$  and  $^{254m}\text{Lr}$  can be assumed, some of which would put the assignment as proposed by Vostinar et al. in question. For a final conclusion concerning the properties of these states and for a comparison with the various theoretical model approaches additional data with higher statistics are mandatory.

### $^{257}\text{Lr}$

Alpha decay of  $^{257}\text{Lr}$  was observed in correlation to  $^{257}\text{Rf}$  and to  $CEs$  by Heßberger et al. [55] which was regarded to be a direct proof for  $EC$  decay of both, the ground state  $^{257g.s.}\text{Rf}$  and the isomer  $^{257m}\text{Rf}$  which were subject of a recent study by Hauschild et al. [56] (see subsection 5.6).

### $^{258}\text{Lr}$

Employing the fusion-evaporation reaction  $^{248}\text{Cm}(^{23}\text{Na},5n)^{266}\text{Bh}$  and the rotating wheel detection system MANON, Haba et al. collected additional  $\alpha$ -decay data for  $^{258}\text{Lr}$  as the granddaughter in the  $^{266}\text{Bh}$  decay chain (see subsection 5.9) [180].

### $^{264}\text{Lr}$

In a recent experiment with the new SHE-factory facility of FLNR JINR, Oganessian et al. assigned a new  $\alpha$ -decay activity to  $^{268}\text{Db}$  in the  $^{288}\text{Mc}$  decay chain [181, 182] (see also section 5.7) which lead to the discovery of  $^{264}\text{Lr}$ . A 4.9-h  $SF$  decay was assigned to the new isotope  $^{264}\text{Lr}$ . This activity was previously, before this new measurement at the FLNR JINR SHE-factory, not observed despite the collection of 104  $^{288}\text{Mc}$  decay chains by the various

experiments at the FLNR in 2003 (3 chains) [183], in 2010-2012 (31 chains) [184], at the gas-filled separator TASCA of GSI (22 chains) [185], at the Berkeley gas-filled separator BGS (48 chains) [186, 147]. The population of  $^{264}\text{Lr}$  was possible because here the authors investigated for the first time "α-like" events in between the  $^{272}\text{Mt}$  and the long- $T_{1/2}$  (28 h) SF events following it and found a new α activity for  $^{268}\text{Db}$  with an α branching of  $55^{+20}_{-15}\%$ , an α-decay energy of 7.6-8.0 MeV and a new  $^{268}\text{Db}$  half-life of  $T_{1/2}=16^{+6}_{-4}\text{h}$  (see subsection 5.7). The half-life for the new fissioning isotope  $^{264}\text{Lr}$  they quote with  $T_{1/2}=4.9^{+2.1}_{-1.3}\text{ h}$ .

### $^{266}\text{Lr}$

In 2019 Khuyagbaatar et al.[176] updated the previously published results [187] with more details on the decay sequence starting with  $^{294}\text{Ts}$  which they observed to be terminated by SF of  $^{266}\text{Lr}$ . The α decay of  $^{270}\text{Db}$  to  $^{266}\text{Lr}$  detected here in the two longer decay chains, was not observed in the earlier experiments at the DGFRS of FLNR JINR [188] (see also section 5.19).

## 5.6. Rutherfordium - $Z=104$

For rutherfordium isotopes spontaneous fission becomes more important and influenced by the transition from  $N=152$  deformed shell gap towards the next higher one at  $N=162$ . The detailed discussion of SF properties of heavy and superheavy nuclei by F.P. Heßberger [189] is addressing the various aspects of fission for nuclear structure and stability. Rf isotopes play a major role in this comprehensive review of the state-of-the-art of fission studies. Among the thirteen rutherfordium isotopes  $^{256}\text{Rf}$ , positioned on the  $N=152$  deformed shell gap, occupies a special position. Apart from being the heaviest nuclide for which a rotational band has been observed (up to a spin of  $20\ \hbar$ ) [190], three  $K$  isomers with half-lives around  $20\ \mu\text{s}$  were initially reported by Jeppesen et al. [106]. This has attracted until recently the attention of various collaborations, as we had also reported in our previous review [1].

### $^{253}\text{Rf}$

Recently two groups reported on new data collection for  $^{253}\text{Rf}$  [103, 89]. As mentioned in subsection 5.4, Khuyagbaatar et al. [103] found a new α decay branch followed also by the α decay of the new isotope  $^{249}\text{No}$  (see subsection 5.4). For  $^{253}\text{Rf}$  they observed two fission activities with fission half lives of  $12.8^{+7.0}_{-3.4}\text{ ms}$  and  $44^{+17}_{-10}\ \mu\text{s}$ , respectively. The longer-lived decay was assigned to the ground state as  $7/2^+[624]$  and the short one as the excited  $1/2^+[631]$  Nilsson state.

The motivation for the investigation of the fission properties of  $^{253}\text{Rf}$  was based on the fission hindrance, for which a hindrance factor  $F_H$  is introduced (see also [189]). This hindrance is assumed to be caused by the possibly complex quantum mechanical configuration due to an unpaired nucleon and it can be defined on the basis of the SF half-lives  $T_{SF}$  of the neighboring isotopes like follows ([103]):

$$F_H(K\pi, A) = \frac{T_{SF}(K\pi, A)}{T_{SF}(0^+, A-1)T_{SF}(0^+, A+1)^{1/2}}, \quad (5.2)$$

where  $\pi$  and  $K$  are the parity and the projection of the total angular momentum of the respective unpaired nucleon (single-particle state) on the symmetry axis of the nucleus.

This hindrance is, in particular, observed in the rutherfordium isotopic chain in long lifetimes for odd- $A$  isotopes. Regarding only even-even isotopes a specific behavior of SF half-lives is observed around the deformed neutron shell gap at  $N=152$ . While for lighter isotopic chains up to nobelium and lawrencium spontaneous fission can compete with α (and β) decay more towards the neutron-rich and neutron-deficient tails (see Fig. 2.1) resulting in a bell-shaped half-life distribution around  $N=152$ , for rutherfordium fission is a prevailing decay even for isotopes more in the centre of the chain with a much smaller retardation at  $N=152$ . For the heaviest rutherfordiums the effect of the next higher deformed shell gap at  $N=162$  becomes evident resulting in similar or even longer half-lives for  $^{260}\text{Rf}$  and  $^{262}\text{Rf}$ . In Fig. 5.17 (taken from [103]) the shown partial spontaneous fission half-lives  $T_{SF}$  for fermium to rutherfordium isotopes exhibit clearly this transition. With respect to the half-lives




Figure 5.17: Spontaneous fission half-lives of Fm, No, and Rf isotopes [189]. Theoretically predicted half-lives for Rf isotopes [115] are shown by rectangles. Dashed lines connect even-even isotopes. Vertical dotted lines show the fission hindrance factors for odd- $A$  nuclei. The dash-dotted line crossing to the isotopic border shows an abruptly falling tendency in half-lives of neutron-deficient Rf isotopes based on an empirically estimated half-life of 1 ps for  $^{252}\text{Rf}$ . This estimate is based on a 48  $\mu\text{s}$  half-life of  $^{253}\text{Rf}$  [189] and an assumed  $F_H=104$ . (Figure and caption are taken from Ref. [103]). See text and Ref. [103] for details.

of odd- $N$  rutherfordium isotopes the hindrance is clearly visible and seem to follow a stable trend. Employing formula 5.2 and with  $F_H=10^4$  estimated on the basis of spin and parities of lighter  $N=149$  isotopes, Khuyagbaatar et al. establish  $T_{SF}$  of the yet unknown  $^{252}\text{Rf}$  as 1 ps. In addition to the two fission activities a single short-lived event was interpreted as a fast  $CE$ , possibly originating from a second high-lying isomer as observed in the next lighter  $N=151$  isotope  $^{251}\text{No}$  with an excitation energy  $E^* > 1.7$  MeV.

In an independent study, Lopez-Martens et al. also observed two fission activities for  $^{253}\text{Rf}$  with half-lives 52.8(4.4)  $\mu\text{s}$  and 9.9(1.2) ms which are within error bars in agreement with the ones reported in Ref. [103]. In addition, they observed the electro-magnetic decay of a high-lying state ( $E^* > 1.02$  MeV) with a half-life of  $0.66^{+40}_{-18}$  ms which is proposed to be a  $K$ -isomer. This isomer decaying

exclusively to a state followed by the faster fission, led them to the conclusion of an inverted assignment with the short-lived fissioning state being the excited  $7/2^+[624]$  Nilsson level, and the slower fission originating from the  $1/2^+[631]$  ground state. Comparing the partial  $SF$  half-life for  $^{251}\text{No}$  (g.s.:  $7/2^+[624]$ ) with 571 s to the one of the same state in  $^{253}\text{Rf}$  and based on the exclusion of fission for  $^{251m}\text{No}$ , in conclusion from a measurement of the same group [162], they estimate a  $T_{SF}(^{252}\text{Rf})$  of 0.69 ps, being close to the value proposed by Khuyagbaatar et al. and therefore placing  $^{252}\text{Rf}$  at the very limit of nuclear stability.

### $^{254}\text{Rf}$

In our earlier review [1] we had mentioned  $^{254}\text{Rf}$  as an example for lifetime inversion leading to meta-stable stable states living longer than the ground state. Apart from the




Figure 5.18: Examples of different types of traces containing fission events: a) Single-signal trace corresponding to  $^{254}\text{Rf}$ . b and c) Double-signal traces where the implantation signals (noted by ER) of  $^{254}\text{Rf}$  were followed by fissions after  $2.9$  and  $31.6\ \mu\text{s}$ , respectively. (Figure and caption are taken from Ref. [61], where more details are given)

long-lived one with  $T_{1/2}=247(73)\ \mu\text{s}$  David et al. had observed, thanks to the use of advanced digital electronics, also a short lived  $K$  isomer with  $T_{1/2}=4.7(11)\ \mu\text{s}$  as compared to the g.s. with  $T_{1/2}=23(3)\ \mu\text{s}$ . Both isomers were identified by the detection of  $CEs$  emitted during the de-excitation cascade, and the subsequent g.s. fission. In a recent investigation of the products of the fusion-evaporation reactions  $^{206}\text{Pb}(\text{Ti},1n/2n)^{255/254}\text{Rf}$  Khuyagbaatar et al. [61] could confirm the short-lived activity in  $^{254}\text{Rf}$  employing pulse-shape analysis, made possible by a digital electronics data acquisition system. They were, however not sensitive to the slow decay. An interesting technical detail is the recovery of the  $SF$  energy from the large and saturated fission pulse utilizing the knowledge of the form of the electronic pulse. The so-called "time-over-threshold" method allows extending in this way the dynamic range to detect energies from a few hundred keV for the  $CEs$  to fission fragment signals

of  $\approx 50$ - $200$  MeV with the same preamplifier configuration. The method is illustrated in Fig. 5.18.

In an in-beam spectroscopy experiment, employing the  $\gamma$  detection array GAMMASPHERE combined with the Argonne Gas-filled Fragment Analyzer (AGFA), despite extremely low statistics, Seweryniak et al. succeeded in extracting some information on the first members of the ground state rotational band of  $^{264}\text{Rf}$  [191]. Using the existing knowledge for rotational bands of neighboring isotopes, in particular, moments of inertia as function of spin and rotational frequency, and the observation of a cluster of counts which was identified as the  $6^+$  to  $4^+$  transition, they succeeded in locating the position of  $\gamma$  transitions up  $14^+$  in the  $^{254}\text{Rf}$   $\gamma$  spectrum. A comparison of the moment of inertia development as a function of the rotational frequency  $\omega$  with the  $N=150$  ( $^{250}\text{Fm}$ ,  $^{252}\text{No}$ ) and  $N=152$  ( $^{254}\text{No}$ ,  $^{256}\text{Rf}$ ) isotones, a behavior similar

to that of the lighter  $N = 150$  isotones is found for  $^{254}\text{Rf}$ , confirming the effect of this deformed shell gap on the collective properties of nuclear matter.




Figure 5.19: Proposed decay scheme of the  $K$  isomers in  $^{255}\text{Rf}$ , populated in the reaction  $^{207}\text{Pb}(^{50}\text{Ti}, 2n)^{255}\text{Rf}$  (left) and in the  $\alpha$  decay of  $^{259}\text{Sg}$  (right). (Figure and caption are taken from Ref. [192])

### $^{255}\text{Rf}$

While in the context of the above mentioned investigation Khuyagbaatar et al., could reproduce within error bars also the literature values for the basic decay properties of  $^{255}\text{Rf}$ , the limited time range of  $30 \mu\text{s}$  (trace length; see Fig. 5.18) of their digital electronics resulted in a truncation of the decay-time spectrum for the detected  $CE$ s [61]. Only the qualitative statement could be made that the observed decay time was longer than  $30 \mu\text{s}$  which is consistent with the findings of Antalic et al. [113] of  $T_{1/2}(^{255m1}\text{Rf}) = 50(17) \mu\text{s}$ .

In a study of spontaneous fission for the three rutherfordium isotopes  $^{255,256,258}\text{Rf}$  Mosat et al. revisited isomeric structures in  $^{255}\text{Rf}$  employing  $ER - CE - \alpha/SF$  correlations [192]. In an energy-decay-time analysis they revealed two new  $K$ -isomers with  $T_{1/2}(^{255m2}\text{Rf}) = 15^{+6}_{-4} \mu\text{s}$  at  $E^* 0.9\text{-}1.2 \text{ MeV}$  and  $T_{1/2}(^{255m3}\text{Rf}) = 38^{+12}_{-7} \mu\text{s}$  at  $E^* 1.15\text{-}1.45 \text{ MeV}$ . To populate  $^{255}\text{Rf}$  and its low lying excited states they used the direct production reaction  $^{207}\text{Pb}(^{50}\text{Ti}, 2n)^{255}\text{Rf}$ . After  $\alpha$  decay of  $^{259}\text{Sg}$  these isomeric states were not observed [113]. This is illustrated Fig. 5.19 taken from [192]. An analysis of available SPLs was not conclusive as many different combinations could lead to appropriate high- $K$  configurations to explain the decay hindrance which

is causing the observed decay times. Advances in modern, self-consistent theory, in particular, an improvement of the SPL-energy predictions are needed to disentangle this puzzle. The accumulation of more and more experimental cases of  $K$ -isomeric structures with their direct link to SPL properties, building up networks in the SPL-energy vs. deformation plane of a Nilsson description could be a route to eventually also enable robust predictions for the region of spherical shell stabilized superheavy nuclei. In




Figure 5.20: TKE distributions of SF fragments from STOP-BOX coincidences (a) for SF of  $^{255}\text{Rf}$  obtained from  $ER - SF$  correlations (inset shows  $ER - SF$  time differences), (b) for SF of  $^{256}\text{Rf}$  obtained from  $ER - SF$  correlations (inset shows  $ER - SF$  time differences), and (c) for SF of  $^{258}\text{Rf}$  obtained from STOP-BOX coincidences (inset shows  $CE - SF$  time differences in cases when  $CE$  was registered). (Figure and caption are taken from Ref. [192])

addition to the investigation of the isomer decay, Mosat et al. investigated also the Total Kinetic Energy ( $TKE$ )

distribution of the *SF* decay of  $^{255}\text{Rf}$  employing the method presented by Nishio et al. in Ref. [193]. Despite the observed asymmetry in the *TKE* distribution, a double-Gaussian fit analysis was not conclusive regarding bi-modal fission which was observed in that region of the Segré chart and predicted for  $^{256,258}\text{Rf}$  [194] (see also the respective paragraphs below). A comparison of the *TKL* distribution for  $^{255}\text{Rf}$  with  $^{256,258}\text{Rf}$  is shown in Fig. 5.20, taken from [192].

### $^{256}\text{Rf}$

As mentioned in the introduction of this subsection the observation of possibly three *K*-isomeric states with half-lives of  $\approx 20\ \mu\text{s}$  has attracted the attention of the scientific community throughout the last decade.

Reported first by Jeppesen et al. in 2009 [106] and shortly after by Berryman et al., Robinson et al., and Rissanen et al. [107, 195, 109], it was revisited in 2020/2021 by Mosat et al. and Khuyagbaatar et al. [104, 103]. In particular, Khuyagbaatar et al. [103], employing trigger-less digital electronics and pulse-shape analysis (PSA) of registered traces confirm the existence of two isomeric states, despite the much lower statistics as compared to the results from Jeppesen et al. [106]. Apart from being a demonstration how effective digital electronics and PSA can be applied to short-lived decays, a major aim of this investigation was the evaluation of the *K*-isomer population probability. A value of  $\geq 28\%$  for the total population of the meta-stable states indicates, according to the authors, that the population of high-*K* isomers in higher-*Z* nuclei like Sg (*Z*=104) and Hs (*Z*=108) is to be expected.

Also for  $^{256}\text{Rf}$  Mosat et al. investigated the *SF* *TKE* distributions and compared it to  $^{255,258}\text{Rf}$  [192] (see respective paragraphs above and below).

Lopez-Martens et al. reported recently on the production of  $^{256}\text{Rf}$  as a *p2n*-evaporation channel in the reaction  $^{209}\text{Bi}(^{50}\text{Ti}, p2n)$  together with  $^{258}\text{Rf}$  produced in the same reaction at lower excitation energy in the *1p*-evaporation channel (see below).

### $^{257}\text{Rf}$

Mosat et al. confirmed in a study of rutherfordium isotopes ( $^{255,256,257}\text{Rf}$ ) the earlier findings for isomeric states in  $^{257}\text{Rf}$  (see also above) [104]. As mentioned in section 5.4, the decay from ground and isomeric states of  $^{257}\text{Rf}$  was employed by Hauschild et al. to reveal details of the low-lying states in  $^{253}\text{No}$  including the discovery of the searched-for  $11/2^-$  [725] Nilsson state.




Figure 5.21: Decay scheme of  $^{258}\text{Db}$  (color scheme and data taken from Ref. [2]).

### $^{258}\text{Rf}$

As shown in Fig. 5.21,  $^{258}\text{Rf}$  is produced by *EC* decay of  $^{258}\text{Db}$  and is mainly decaying by *SF*. By delayed *CE*- $\alpha$  coincidences and correlations to  $^{254}\text{No}$   $\alpha$  decay Heßberger et al. observed a weak  $\alpha$ -decay branch with  $b_\alpha=0.049(16)$  [76] which was about an order of magnitude lower than the previously reported value of  $b_\alpha=0.31$  by Gates et al. [196]. The  $^{258}\text{Rf}$  mass can be obtained by the here established  $Q_\alpha$  value from the precisely measured mass value of  $^{254}\text{No}$

at the Penning trap set-up SHIPTRAP [160]. On this basis the two-neutron separation energy trends in the vicinity of the deformed shell gap at  $N=152$  and towards  $N=162$  are also discussed in comparison with model predictions in the paper by Heßberger et al. [76].

In addition to the improved  $\alpha$  decay data, in this paper  $CE$ -coincident photons, followed by SF, were studied.

Fig. 5.22 shows  $EC-SF$  correlated photon spectra from the  $^{258}\text{Db}$  decay. Here enhanced rutherfordium  $x$ -ray lines were observed for photons in coincidence with  $CE$  while they were suppressed with at the same time enhanced  $\gamma$  transitions appearing in the spectrum in anti-coincidence with  $CE$ s. From  $EC-\gamma/x\text{-ray}-CE-SF$  correlations the existence and half-lives of two isomeric states in  $^{258}\text{Rf}$  could be established with  $T_{1/2}(^{258m1}\text{Rf}) = 2.4^{+2.4}_{-0.8}$  ms and  $T_{1/2}(^{258m2}\text{Rf}) = 15(10)$   $\mu\text{s}$ , respectively. In particular, the observation of  $x$ -rays of the decay daughter rutherfordium is interesting with respect to the identification of the atomic charge  $Z$  for cases where  $SF$  is terminating a decay chain of unknown nuclei (see subsection 5.7).

In an irradiation of  $^{209}\text{Bi}$  with  $^{50}\text{Ti}$  Lopez-Martens et al. found evidence for proton-evaporation channels ( $p$  and  $p2n$ ) leading to  $^{258}\text{Rf}$  (and  $^{256}\text{Rf}$ ) as ERs by the observation of short-lived  $SF$  events [150] (see also subsection 5.7). In an earlier measurement in 1985 [197], Heßberger reported the observation of one SF event with a decay time of 5 ms which was tentatively assigned to the reaction channel  $^{209}\text{Bi}(^{50}\text{Ti},p)^{258}\text{Rf}$  at a beam energy of 243 MeV. The obtained cross-sections in both experiments, being consistent with each, are suppressed with respect to neutron-evaporation channels by a factor of 30 to 100 at the given beam energies, with  $\approx 4$  pbarn [150] to  $\approx 18$  pbarn [197] for  $p$ -evaporation channels and  $\approx 4$  nbarn for the maximum cross-section of the neutron-evaporation channels.

The detailed structure of the  $SF$   $TKL$  distribution of  $^{258}\text{Rf}$  was compared to the one of the two lighter isotopes  $^{255,256}\text{Rf}$  by Mosat et al. [192] (see also above). The two-Gaussian analysis could possibly indicate bi-modal fission due to the observed skewness for the latter, while

the situation is less clear for the more symmetric  $TKL$  distribution observed in the case of  $^{258}\text{Rf}$  with only a small improvement of the  $\chi^2$  value for the double-Gaussian with respect to the single-Gaussian fit (see Fig. 5.20). The two possibilities of either single mode fission or the existence of two modes with similar distributions is supported by theory predictions [194].

### $^{261}\text{Rf}$

In an irradiation on  $^{238}\text{U}$  with  $^{40}\text{Ar}$  projectiles at the new FLNR JINR SHE-factory Oganessian et al. observed two  $\alpha$ -decay chains of type  $ER$ - $\alpha$ - $\alpha$ - $SF$  which they assigned to  $5n$  fusion-evaporation channel with the  $ER$   $^{273}\text{Ds}$ , terminated by the spontaneous fission of  $^{261m}\text{Rf}$  which is consistent with literature data [29]. For a discussion of the whole decay chain in which the population of isomers in all chain members is proposed by the authors of Ref. [29] see subsection 5.12.

### $^{262}\text{Rf}$

In a systematic study of the competition between fusion and quasi-fission,  $^{262}\text{Rf}$  was one of the nuclei investigated by Hinde et al. [198]. This work was mainly focussed on reaction dynamics being relevant for production and survival of very heavy nuclei. As nuclear structure features of the heavy collision participants, in particular nuclear deformation, play a decisive role, this reaction mechanism work with its findings is mentioned in this review on decay spectroscopy.

Investigating fission mass angular distributions (MAD) for the  $^{28}\text{Si} + ^{238}\text{U}$  and  $^{34}\text{S} + ^{232}\text{Th}$  leading to the excited compound nucleus  $^{266}\text{Sg}$ , and  $^{24}\text{S} + ^{238}\text{U}$  forming  $^{262}\text{Rf}$ , Hinde et al. have access to the reaction dynamics and can distinguish between fast and slow processes. From peaked MADs they deduce the existence of a fast quasifission (FQF) mode leading to an asymmetric mass distribution which can be distinguished from slow quasifission (SQF) which shows a symmetric mass distribution. These two quasifission modes compete with fusion of the colliding




Figure 5.22: Photon spectra observed in delayed coincidence with *SF* events; (a) spectrum of events, recorded between the beam bursts and preceding a *SF* event within  $\Delta t \leq 30$  ms (black line); the dotted line (pink) shows the spectrum of  $\gamma$  background down scaled to similar peak-heights for the background lines at  $E=139$  and  $197$  keV; (b) same as (a), but for photons registered in anti-coincidence with a *CE*; (c) spectrum of photons registered in coincidence with a *CE* preceding an *SF* event within  $\Delta t \leq 50$  ms. The dotted lines (purple) mark the energy positions of the  $K_{\alpha 1,2,\beta 1,2}$  x-rays of rutherfordium ( $Z = 104$ ). No attempts to identify the background lines were made. (Figure and caption are taken from Ref. [76])

system. Systematically analyzing the relative strength of symmetric and asymmetric fission modes and angular distributions as a function of kinetic energy of the colliding system as well as of the atomic charge of the projectile impinging on the deformed actinide targets, the authors discuss the competing processes.

### $^{263}\text{Rf}$

At the new FLNR JINR SHE-factory, in a similar investiga-

tion as mentioned above for the decay chain from  $^{273}\text{Ds}$  to  $^{261}\text{Rf}$ , Oganessian et al. observed  $^{263}\text{Rf}$ , terminating three of six decay chains assigned to the  $5n$  fusion-evaporation channel,  $^{275}\text{Ds}$ , of the reaction  $^{48}\text{Ca} + ^{232}\text{Th}$  [29]. This constitutes the discovery of this hitherto not known darmstadtium isotope. For this decay chain also the observation of new isomers are reported which will be discussed in more detail in section 5.12.

### $^{265}\text{Rf}$

Utyonkov et al. collected data for three new decay chains originating from  $^{285}\text{Fl}$  and terminating after five subsequent  $\alpha$  decays with *SF* of  $^{265}\text{Rf}$  [199] (see also corresponding subsections below). These new decay chains were mainly confirming literature data [200, 201] with an exception for the  $\alpha$  decay of  $^{269}\text{Sg}$  in the third of the new chains (see subsection 5.8).

### $^{266,267,268,270}\text{Rf}$

The decay chains originating from  $^{282}\text{Nh}$ ,  $^{287,288}\text{Mc}$  and  $^{294}\text{Ts}$  are terminated by *SF* which in the most recent publications by Oganessian et al. are attributed to the *SF* decay of  $^{266,267,268}\text{Db}$  [12]. However, the decay by *EC* of these dubnium isotopes cannot be ruled out, as given in earlier papers of the same group like in the discovery paper for  $^{282}\text{Nh}$  [202] or for  $^{266,268}\text{Db}$  decays in a review paper in the same year (2007) [203]. Similarly, for the endpoint of the  $^{294}\text{Ts}$  decay chain in the discovery paper from 2010  $^{270}\text{Db}$  is given in Ref. [25] from 2010, while in a review from 2011 the possibility of *EC* decay to  $^{270}\text{Rf}$  is considered in Fig. 5 of Ref. [204].

As discussed for  $^{258}\text{Db}(EC)^{258}\text{Rf}$  the possible population of excited states in the decay daughter could result in *x-ray* emission due to internal conversion (*IC*) which would be of special interest for atomic charge identification for those chains with no connection to known nuclides (see also subsection 5.7). Clear evidence for this process is, however, still missing and only  $^{267}\text{Rf}$  is confirmed as the end point of the  $\alpha$ -decay sequence starting with  $^{291}\text{Lv}$  [205]

(see Fig. 2.1). The possibility of *EC* decay for  $^{266,268}\text{Db}$  populating  $^{266,268}\text{Rf}$  is indicated in Fig. 2.1 by faint-red triangles (see also discussion in section 5.7 for  $^{266,267,268}\text{Db}$ ).

### $^{267}\text{Rf}$

The halflife of fissioning  $^{257}\text{Rf}$  has been updated with improved statistics obtained at the new FLNR JINR SHE-factory [181] by Oganessian et al. in the irradiation of  $^{242}\text{Pu}$  and  $^{238}\text{U}$  with  $^{48}\text{Ca}$  projectiles. For the plutonium target six decay chains terminated by *SF* of  $^{267}\text{Rf}$ , while in the  $^{238}\text{U}$  irradiation one additional decay chain of this type was observed [206] (see subsections 5.14 and 5.16).

## 5.7. Dubnium - $Z=105$

### $^{255}\text{Db}$

$^{255}\text{Db}$  is the lightest dubnium isotope for which reports of its observation exist. However, the reported properties differ significantly from each other. The earliest paper mentioning this isotope is a conference report from 1976 by G.N. Flerov [207], which just lists the branching ratio of 80%  $\alpha$  decay and 20% SF together with a half-life of 1.5 s in a table without any mentioning in the text. This is probably the same activity which is mentioned by Oganessian et al. in the same year in Ref. [208] where two activities with a half-lives of  $1.2^{+0.6}_{-0.3}$  s and  $>1$  s are given, produced in the same reactions  $^{51}\text{V} + ^{207,206}\text{Pb}$ , respectively. In 1986 in a laboratory report by the same group  $^{255}\text{Db}$  is mentioned with a half-life of 1.6 s [209].

A.-P. Leppänen reported in his Ph.D. thesis in 2005 [210] on two recoil-SF and one recoil- $\alpha$ -SF correlation events which he assigned to the decay of  $^{255}\text{Db}$  with a half-life of  $37^{+51}_{-14}$  ms. This half-life differs by orders of magnitude from the earlier reported values.

Finally, in 2024 Pore et al. reported on the detection of 55 fission events which they assign to  $^{255}\text{Db}$  produced as the  $2n$  fusion-evaporation channel in the reaction  $^{51}\text{V} + ^{206}\text{Pb}$  [134]. From this they deduce a half-life of  $T_{SF} = 2.7^{+4}_{-3}$  ms which is adopted here in Table 5.1.

### $^{256}\text{Db}$

In the above mentioned irradiation of a  $^{206}\text{Pb}$  target with  $^{51}\text{V}$  projectiles at the BGS of LBNL (see subsections 5.1, 5.2, 5.3 and 5.5) a total of 86 decay chains starting at  $^{256}\text{Db}$  were accumulated by Pore et al. [134]. To extract the decay properties of  $^{256}\text{Db}$  and its  $\alpha$  decay daughter  $^{252}\text{Lr}$  the authors considered only 43 events with  $\alpha$  decay energies above a certain threshold ( $E_\alpha > 8.75$  MeV), which is still substantially more than previously collected by Heßberger et al. [122] and Nelson et al. [211]. The decay properties are in agreement with the earlier findings, leading to a better precision of the measured values.

The authors also discuss the possible population and decay of isomeric states in the daughter nucleus  $^{252}\text{Lr}$ . To search for such states they investigate 43 selected events in a restricted  $E_\alpha$  window, using  $\alpha_1(^{256}\text{Db})-\alpha_2(^{252}\text{Lr})$  time and energy correlation plots which do not show any sign of the presence of an isomeric state. Yet, the population and decay of isomeric states at finite excitation energy can proceed via transitions with lower  $Q_\alpha$  values. The non-considered  $\alpha_1-\alpha_2$  correlation with  $E_{\alpha 2}$  below the threshold of 8.75 MeV chosen by the authors could indeed originate from such decay routes. This is, however, not discussed in the paper [134] and, lacking the timing information, remains subject of speculation. The only statement regarding those events given, is that all 86 decay chains were used "To determine the properties of all subsequent daughters in the decay chain, ...".

### $^{256-258}\text{Db}$

In an experiment to investigate proton evaporation channels Lopez-Martens et al. used the reaction  $^{50}\text{Ti} + ^{209}\text{Bi} \rightarrow ^{259}\text{Db}^*$  to populate  $xn$  and  $pxn$  evaporation channels (see also section 5.6). The data obtained for the three observed  $xn$  ERs  $^{256-258}\text{Db}$  were consistent with the earlier findings of Heßberger et al. in Ref. [122]. Some aspects of this measurement are also reported By Kuznetsova et al. and Yeremin et al. in Ref.s [165, 212].

## $^{258}\text{Db}$

Fig. 5.21 shows the decay scheme of  $^{258}\text{Db}$ , illustrating how the various decay paths via  $\alpha$  and  $\beta$  decay evolve across the  $N=152$  shell gap in the vicinity of the  $Z=100$  isotope  $^{252}\text{Fm}$ . Together with its  $\alpha$ -decay daughter nuclei  $^{246}\text{Es}$  (section 5.1),  $^{250}\text{Md}$  (section 5.3) and  $^{254}\text{Lr}$  (section 5.5),  $^{258}\text{Db}$  is subject of a detailed decay study reported in two article focusing separately on it's  $\beta$ - (Heßberger et al. [76]) and  $\alpha$ -decay branch (Vostinar et al. [136]), respectively.

Being an odd-odd nucleus in a region with access to high- $J$  orbitals, complex  $J - \pi$  configurations of its low-lying excitation states with consequences for decay-mode competition ( $\alpha$ -,  $\beta$ -, SF- and intrinsic  $\gamma$ -decay),  $^{258}\text{Db}$  represents the starting point of a complex decay network (as shown in Fig. 5.21). The dependence of the various decay modes as well as decay properties like half-lives and decay energies provide a rich source of information on the microscopic structure features of the involved nuclei.

The earlier of the two papers reports on the  $^{258}\text{Db}$  EC decay into excited states and the ground state of  $^{258}\text{Rf}$  [76] of the two activities known in  $^{258}\text{Db}$  from earlier  $\alpha$ -spectroscopy results [197, 213], supporting their existence. The half-lives deduced for these two states from the measured  $\beta$  decay were  $T_{1/2} = 4.3(5)$  s and  $1.9(5)$  s. The findings for  $^{258}\text{Rf}$  are summarized in the previous subsection 5.6.

The second paper reports on a more detailed analysis of the collected  $\alpha$  decays for  $^{258}\text{Db}$  producing a rather complex spectrum with various components contributing to the two activities. From  $\alpha$ - $\gamma$  coincidences on the basis of decay energy systematics a tentative level scheme of the low-lying structure in  $^{254}\text{Lr}$  and possible spin and parities in the two states could be inferred. In addition the authors propose the assignment as ground state for  $^{258}\text{Db}(2)$  (the nomenclature of Ref. [197] was preserved) with  $J^\pi = (0-)$  and an improved half-life value of  $T_{1/2} = 2.17(36)$  s.  $^{258}\text{Db}(1)$  was then assigned as an isomeric state with  $E^* = 51$  keV,  $J^\pi = (5+)$  or  $(10-)$  and  $T_{1/2} = 4.41(21)$  s.

## $^{262}\text{Db}$

In 2020 Haba et al. reported on new data for the  $\alpha$  decay of  $^{262}\text{Db}$  as the daughter of  $^{266}\text{Bh}$  produced in the reaction  $^{248}\text{Cm}(^{23}\text{Na},5n)^{266}\text{Bh}$ . The experiment was performed using the rotational wheel detection set-up MANON [180] (see also section 5.8).  $^{262}\text{Db}$  was first observed in the decay chain of  $^{278}\text{Nh}$  by Morita et al. [214, 215, 216, 217] and later directly produced in the reaction  $^{248}\text{Cm}(^{19}\text{F},5n)^{262}\text{Db}$  [218].

## $^{265}\text{Db}$

D. Fernandez et al. reported in 2023 on fission studies of the compound nucleus  $^{265}\text{Db}$  produced in inverse kinematics, irradiating  $^{238}\text{U}$  with  $^{27}\text{Al}$  ions [219]. Employing the magnetic spectrometer VAMOS++ [220], the isotopic fission yield distribution was investigated for the highly excited nuclear system. Apart from the presenting the experimental fission yield distributions, determined by quasi-fission, the authors suggest the observed even-odd staggering for the heavy fragments as a probe to investigate the interplay of quasi-fission and fusion-fission. No properties of the ground state of  $^{265}\text{Db}$  or its decay are known.

## $^{266-268}\text{Db}$

The three isotopes  $^{266,267,268}\text{Db}$  were assigned to SF events, observed as the termination of decay chains of nihonium and moscovium isotopes by Oganessian et al. [183] while in Ref. [202] the possibility of  $\beta$  decay(EC) is considered for  $^{266}\text{Db}$ , as the detection system used for those measurements was not sensitive to  $\beta$  decay (see discussion above). In Ref. [223] the authors discuss the probability of  $\beta$  decay of the heaviest isotopes competing with SF observed as chain terminations. In Fig. 5.23 the comparison of the observed half-lives assigned to SF (colored symbols) with measured partial EC/ $\beta^+$  half-lives shows that for the dubnium isotopes  $^{266,268,270}\text{Db}$  the assigned  $T_{SF}$  values are similar to the  $T_\beta$  value trend of the lighter odd- $Z$  nuclei, indicating a finite probability of  $\beta$  decay to occur. Part of the observed SF decays would then belong to the rutherfordium daughter isotopes, at least for  $^{266}\text{Db}$  and  $^{268}\text{Db}$ .




Figure 5.23: Partial EC/ $\beta^+$ -decay half-lives vs. decay energy for known odd-odd isotopes of Np-Db (open symbols). The systematics [221] for odd-odd and odd-A isotopes are shown by lines. Experimental half-lives of terminal isotopes for which spontaneous fission was detected in the decay chains produced in  $^{48}\text{Ca}$ -induced reactions with  $^{237}\text{Np}$ ,  $^{243}\text{Am}$ , and  $^{249}\text{Bk}$  targets are given by colored solid squares. EC/ $\beta^+$ -decay energies for  $^{266,268,270}\text{Db}$  and  $^{281}\text{Rg}$  were calculated from Ref. [222]. (Figure and caption are taken from Ref. [223])

For  $^{266}\text{Db}$  one event in addition to the one previously observed in the  $^{282}\text{Nh}$  chain [203], was reported in Ref. [182] with a half-life shorter by a factor of  $\approx 25$ . This is, however, still consistent with a single decay activity according to the Poisson statistics analysis in the prescription of Ref. [225], leading to a combined half-life of  $11^{+21}_{-4}$  min (see Table 5.1).

From the shorter half-lives observed for the two isotopes  $^{267}\text{Db}$  and  $^{281}\text{Rg}$  the authors deduce that SF would be more likely the dominating decay mode. They also discuss decay time systematics across the  $N=162$  shell gap. A striking observation are the long half-lives of up to  $16^{+6}_{-4}$  h for  $^{268}\text{Db}$  as given in a recent paper on the first results obtained at the new SHE-factory of FLNR JINR [181]. Here the authors report on 55 new decay chains obtained for  $^{288}\text{Mc}$  and six new chains for  $^{289}\text{Mc}$ .

In addition to the known decay features, they observed also  $\alpha$  decay of  $^{268}\text{Db}$  leading to the discovery of the new isotope  $^{264}\text{Lr}$  (see subsection 5.5). This new assignment leads to changes in the decay branching ratios and consequently partial decay half-lives.

A possible EC-decay of the odd-odd isotope  $^{267}\text{Db}$  with

$T_{1/2} = 1.2^{+58}_{-6}$  h to the daughter nucleus  $^{267}\text{Rf}$  with a very similar half-life of  $T_{1/2} = 1.3^{+23}_{-5}$  h is more unlikely, as supported also by the EC/ $\beta^+$ -decay half-life systematics in Fig. 5.23. The close-by lawrencium isotopes  $^{260,261,262}\text{Lr}$  exhibit a similar decay mode pattern with  $\beta$  decay being observed for the two odd-odd isotopes while the even-odd isotope in between decays exclusively by fission (see Fig 2.1). The half-life trends are also similar with 0.4 h, 1.3 h and 16 h for  $^{266,267,268}\text{Db}$ , and 180 s, 39 min and 4 h for  $^{260,261,262}\text{Lr}$ , respectively. To disentangle  $\beta$  decay (EC) from SF, correlations by CE emission and the coincident  $x$ -rays, originating from possibly populated excited states in the daughter nucleus can be employed, as applied in the case of  $^{258}\text{Db}$  and its  $\beta$ -decay daughter  $^{258}\text{Rf}$  by Heßberger et al. in ref. [76]. In addition, the correlated  $x$ -rays emitted with the CE can provide the identification of the atomic charge  $Z$  of the decaying nuclide which would yield the still missing direct experimental prove for nuclides of elements exclusively produced in hot fusion, for  $^{48}\text{Ca}$  induced reactions with actinide targets. This concerns all elements from flerovium to oganesson, the five heaviest elements with  $Z=114$  to 118.

In the second irradiation of  $^{243}\text{Am}$  with  $^{48}\text{Ca}$  projectiles during this first series of experiments at the SHE-factory with 52 new  $^{288}\text{Mc}$  decay chains collected, the same  $\alpha$  activity was found.

The SF half-life systematics, including the new data, is shown in Fig. 5.24 taken from Ref. [182], reporting on this second experiment. Extending the investigation to higher beam energies, the  $4n$  ( $^{287}\text{Mc}$ ) and, for the first time the  $5n$  fusion-evaporation channel with the new isotope  $^{286}\text{Mc}$  were observed.

### $^{270}\text{Db}$

In the irradiation of  $^{249}\text{Bk}$  with  $^{48}\text{Ca}$ , Khuyagbaatar et al. investigated the decay sequence starting with  $^{294}\text{Ts}$  which was, as mentioned above (subsection 5.5), terminated by SF of  $^{266}\text{Lr}$  [176]. The  $\alpha$  decay of  $^{270}\text{Db}$ , detected in the two longer decay chains was not observed in the earlier experiments at the DGFRS of FLNR JINR [188] (see also




Figure 5.24: Common logarithm of partial spontaneous fission half-life [ $\log_{10}(T_{SF}[s])$ ] vs neutron number for isotopes of elements with  $Z = 100\text{--}114$ . The area of half-lives of even-even isotopes with  $Z = 104\text{--}110$ ,  $N = 160\text{--}164$  and  $Z = 110\text{--}114$ ,  $N = 164\text{--}177$  predicted in [224] is highlighted in gray. Partial half-lives of nuclei produced in the  $^{48}\text{Ca}$ -induced reactions with actinide targets are shown by filled symbols. (Figure and caption are taken from Ref. [182])

section 5.19).

## 5.8. Seaborgium - $Z=106$

For the heaviest nuclei, the nuclear structure information collected is, due to the low production cross-section often limited to basic decay properties. E.g., as mentioned before, starting with the seaborgium isotopes no information on the  $\beta$ -decay properties is available for the higher- $Z$  nuclides.

Seaborgium is also the only even- $Z$  element for  $Z$  from 100 (fermium) to 110 (darmstadtium), for which no  $K$ -isomer has been reported yet. However, new data, including the synthesis of the new isotope  $^{268}\text{Sg}$ , has been recorded for some of its isotopes as the endpoints of decay chains originating from isotopes of heavier element, mainly due to the first experiments performed at the new SHE-factory of FLNR JINR, as for many of the lighter superheavy elements.

### $^{264}\text{Sg}$

For  $^{264}\text{Sg}$  Galkina et al. investigated fission properties of this nucleus produced in the fusion reaction  $^{32}\text{S}+^{232}\text{Th}$  at the three different excitation energies  $E^*=45$  MeV, 59 MeV, and 75 MeV, around the Coulomb barrier [226]. While a substantial part of the reaction products is found to be due to quasifission, an analysis of the Total Kinetic Energy (TKE), based on the fission mode classification by Brosa et al. [227], leads the authors to the conclusion that a large contribution to symmetric a mass split of up to 80% results from compound-nucleus fission.

### $^{265}\text{Sg}$

As a  $\alpha$ -decaying member of the two  $^{273}\text{Ds}$  decay chains, produced at the FLNR JINR SHE-factory in the  $5n$  fusion-evaporation channel of the reaction  $^{40}\text{Ar}+^{238}\text{U}$ , additional

two decay events were collected for  $^{265}\text{Sg}$  with respect to earlier results [29]. For More details of this experiment see subsection 5.12.

### $^{267}\text{Sg}$

Also for  $^{267}\text{Sg}$  new data has been reported from an experiment at the new FLNR JINR SHE-factory [29]. A more detailed discussion on measured properties for  $^{275}\text{Ds}$  and its decay products, will be given in section 5.12. Three of the six  $^{275}\text{Ds}$  decay chains collected in this experiment exhibit  $\alpha$  decay of  $^{267}\text{Sg}$  with relatively long decay times, ranging from 760 s to 970 s, while for the three fission events decay times vary from  $\approx 1$  s (2 events) to 360 s. The authors propose an isomeric state decaying by fission and the ground state decaying by  $\alpha$  emission with a half-life of about 10 min (see Table 5.1), exceeding the detection time range of 300 s of the earlier chemistry experiment [228, 229] where this nucleus was observed before without an indication for an isomeric state.

### $^{268}\text{Sg}$

Irradiating a  $^{232}\text{Th}$  target with  $^{48}\text{Ca}$  projectiles, Oganessian et al. observed in one of the series of experiments carried out at the new FLNR SHE-factory three decay chains, consisting of two subsequent  $\alpha$  decays followed by a  $SF$  which they assigned to start from the  $4n$  fusion-evaporation residue  $^{276}\text{Ds}$  populated in this as the  $4-n$  channel [28]. All three nuclei in this chain, including as its termination  $^{268}\text{Sg}$ , were hitherto unknown (see subsection 5.12).

### $^{269}\text{Sg}$

In 2018 Utyonkov et al. reported on a second investigation of the reaction  $^{240}\text{Pu}(^{48}\text{Ca},3n)^{285}\text{Fl}$ , with  $^{269}\text{Sg}$  being populated in the decay chain, at a beam energy of  $E_{beam}=250$  MeV [199] with respect to the earlier measurement at  $E_{beam}=245$  MeV (see subsection 5.16).

### $^{271}\text{Sg}$

$^{271}\text{Sg}$  is the last but one member of the  $^{297}\text{Fl}$  decay chain

and exhibits  $\alpha$ -decay as well as  $SF$ . It was recently again observed at the new SHE-factory of FLNR JINR in an irradiation of  $^{242}\text{Pu}$  [206] (see subsection 5.16). As  $^{267}\text{Rf}$  and  $^{275}\text{Hs}$ , discussed above and below, also  $^{271}\text{Sg}$  is produced following the  $\alpha$ -decay of  $^{279}\text{Ds}$  with seven and two new decay chains observed in irradiation of  $^{242}\text{Pu}$  and of  $^{238}\text{U}$ , respectively [206] (see subsections 5.16 and 5.14). The values in Table 5.1 have been updated accordingly, taking the values from [206] which include all previously accumulated data.

## 5.9. Bohrium - $Z=107$

Apart from a re-evaluation of an earlier measurement for  $^{274}\text{Bh}$ , for the bohrium isotopes new data were collected exclusively in experiments at the new SHE-factory of FLNR JINR with the respective bohrium isotopes being members of the decay chains of heavier nuclei.

### $^{270}\text{Bh}$

In the first campaign of experiments at the Dubna SHE-factory Oganessian et al. reported in 2022 on the discovery of  $^{286}\text{Mc}$  of which  $^{270}\text{Bh}$  is the last  $\alpha$  decaying member [182] (see subsection 5.17), while it was first observed in the  $\alpha$ -decay sequence of  $^{282}\text{Nh}$  in 2007 [202]. In addition to the earlier observed event a second  $^{270}\text{Bh}$   $\alpha$  decay was reported in Ref. [182].

### $^{271}\text{Bh}$

Similar as for  $^{270}\text{Bh}$  Oganessian et al. collected also for  $^{271}\text{Bh}$  new data in the decay products following  $^{287}\text{Mc}$  (see subsection 5.17) which they reported in 2022 [182], again doubling the number of observed events from two to four.

### $^{272}\text{Bh}$

Until the last  $^{288}\text{Mc}$  synthesis experiment performed at the FLNR SHE-factory [182]  $^{272}\text{Bh}$  was the last  $\alpha$  decaying member of the decay chain, before the  $^{266}\text{Db}$   $\alpha$  decay was extracted from the new data (see subsection 5.7). In an earlier re-investigation, performed at the Berkeley Gas-filled Separator (BGS), Gates et al. had used the trochoidal

mass spectrometer to establish the atomic mass number A of  $^{288}\text{Mc}$  [147], where they also detected  $^{272}\text{Bh}$  as the last  $\alpha$  decay of the two observed chains (see subsection 5.17).

### $^{274}\text{Bh}$

In 2019 Khuyagbaater et al. published more details with respect to the first publication in 2015 [187] on the reaction  $^{48}\text{Ca} + ^{249}\text{Bk}$  to synthesize  $^{294}\text{Ts}$  [176].  $^{274}\text{Bh}$  is a member of the decay chain (see subsection 5.19).

## 5.10. Hassium - $Z=108$

For the new hassium data the same is valid as stated in the introduction of the previous subsection 5.9 for the bohrium isotopes.

### $^{269}\text{Hs}$

As the  $\alpha$ -decay daughter of  $^{273}\text{Ds}$ , of  $^{269}\text{Hs}$ , its  $\alpha$  decay was discussed in a recent paper of an experiment at the new FLNR JINR SHE-factory [29]. Despite the fact that only for one chain an indication of this decay by an escape- $\alpha$  signal was observed, the authors discuss the presence of an isomeric state with spin 1/2 and a half-life of  $2.8_{-1.3}^{+3.6}$  s above the ground state with spin 9/2 and a half-life of  $13_{-4}^{+10}$  s (see Table 5.1) on the basis of existing earlier data for the  $^{277}\text{Cn}$  decay chain [230, 231] and its direct production [228, 229]. For more details on the properties the whole decay chain see subsection 5.12.

### $^{271}\text{Hs}$

A similar scenario as for  $^{269}\text{Hs}$  has been proposed for  $^{271}\text{Hs}$ , again observed at the FLNR JINR SHE-factory in six decay chains following the  $\alpha$  decay of  $^{275}\text{Ds}$  which was newly synthesized as the 5n fusion-evaporation channel of the reaction  $^{48}\text{Ca} + ^{232}\text{Th}$  [29]. The authors propose again an isomeric state with low spin (3/2) and  $T_{1/2} = 7.1_{-2.5}^{+8.4}$  s above a ground state with high spin (13/2) and  $T_{1/2} = 46_{-16}^{56}$  s (see Table 5.1). More details on the interpretation of the decay chain and the population of isomeric states by  $\alpha$  decay is discussed in section 5.12.

### $^{272}\text{Hs}$

Irradiating a  $^{232}\text{Th}$  target with  $^{48}\text{Ca}$  projectiles Oganessian et al. observed three decay chains consisting of two subsequent  $\alpha$  decays, followed by a *SF*, which they assigned to start from the 4n fusion-evaporation residue  $^{276}\text{Ds}$  populated in this reaction. All three nuclei in this chain, including as its termination  $^{268}\text{Sg}$ , were hitherto unknown (see subsections 5.8 and 5.12).

### $^{273}\text{Hs}$

As mentioned earlier for the lighter isotopes, Utyonkov et al. collected data for three new decay chains originating from  $^{285}\text{Fl}$  and terminating after five subsequent  $\alpha$  decays, of which  $^{273}\text{Hs}$  is the 4<sup>th</sup> member, with *SF* of  $^{265}\text{Rf}$  [199] (see also corresponding subsections below and above).

### $^{275}\text{Hs}$

Also for  $^{275}\text{Hs}$ , as amember of the decay chains of  $^{283}\text{Cn}$  and  $^{287}\text{Fl}$ , new  $\alpha$ -decay events were observed at the new SHE-factory of FLNR JINR in the irradiation of  $^{242}\text{Pu}$  and of  $^{238}\text{U}$  targets with  $^{48}\text{Ca}$  projectiles [206] (see subsections 5.14 and 5.16).

### $^{277,278}\text{Hs}$

In 2019 Khuyagbaatar et al. observed four decay chains produced in the reaction  $^{48}\text{Ca} + ^{249}\text{Bk}$  of which they assigned two each as starting from  $^{294}\text{Ts}$  and  $^{293}\text{Ts}$ , respectively. For the latter, they assumed the possibility of  $^{277}\text{Mt}$   $\beta$  decay leading to  $^{277}\text{Hs}$ , similar to what was discussed for  $^{266,267,268}\text{Db}$  in section 5.7.

## 5.11. Meitnerium - $Z=109$

Also for meitnerium new or updated data have been reported for five meitnerium isotopes as members of decay chains originating from heavier nuclei, for measurements performed by Oganessian et al. at the new Dubna SHE-factory, as well as for earlier experiments reevaluated by Khuyabaatar et al.




Figure 5.25: A proposed experimental and calculated decay scheme for  $^{275}\text{Ds}$  based on present and previously published data [228, 229]. Alpha or  $\gamma$  transitions to approximate one-quasiparticle states and SF decays are shown by red (solid), blue (dashed), and green (dotted) lines, respectively. Alpha-decay energies ( $Q_\alpha$ ) and half-lives are provided for states decaying by  $\alpha$  decay ( $T_\alpha$ ) or spontaneous fission ( $T_{SF}$ ). Hindrance factors  $HF = T_{1/2}^{\text{exp}}/T_{1/2}^{\text{calc}}$  were derived from experimental and calculated half-lives according to Ref.s [205, 232, 233, 234, 235] (see discussion in [29]). Population probabilities  $P$  of energy levels for  $^{271}\text{Hs}$  are based on experimental data [228, 229]. and this[29]. (Figure and caption are taken from Ref. [29])

$^{274,275,276}\text{Mt}$

For the three meitnerium isotopes  $^{274,275,276}\text{Mt}$  as members of the decay chains of the respective moscovium isotopes, additional data were accumulated at the new SHE-factory at FLNR JINR in 2022 [182] (see subsection 5.17). In this experiment for  $^{274}\text{Mt}$  no  $\alpha$  decay energy, in addition to the previously observed one event was measured, and only indirect information on its decay time was deduced.

$^{277,278}\text{Mt}$

In 2019 Khuyagbaatar et al. [176] reported, in addition to the earlier published  $^{294}\text{Ts}$  decay data [175], hitherto unpublished data for  $^{293}\text{Ts}$ .  $^{278}\text{Mt}$  is an  $\alpha$ -decaying member of the  $^{294}\text{Ts}$  decay chain and  $^{277}\text{Mt}$  terminates the  $^{293}\text{Ts}$

decay chain, while the authors suggest, in line with what was discusses in subsection 4.3.2, alternatively EC to  $^{277}\text{Hs}$  which then decays by SF [176] (see subsection 5.19).

## 5.12. Darmstadtium - $Z=110$

The possibly most interesting new data is the synthesis new darmstadtium isotopes  $^{275}\text{Ds}$  adding another stepping stone in between the island populated mainly by reactions of  $^{48}\text{Ca}$  projectiles on actinide target nuclei. A complete connection to the "mainland" with the heaviest nuclei produced in cold fusion reactions, resulting in ER at low excitation energy would strengthen the up to now missing direct  $Z$ -identification.

### $^{273}\text{Ds}$

Similar to the interpretation of the six decay chains observed for  $^{275}\text{Ds}$  (see below) and obviously encouraged by the apparent consistency, Oganessian et al. constructed a similar decay scenario for the decay sequence starting from  $^{273}\text{Ds}$ . To this end, they used two decay chains of type *ER*- $\alpha$ - $\alpha$ -*SF* which they observed in the irradiation of  $^{238}\text{U}$  with  $^{40}\text{Ar}$  projectiles at the FLNR JINR SHE-factory [29], combined with available literature data. In Fig. 5 and Table 3 of Ref. [29] they propose new isomers in all four members,  $^{273}\text{Ds}$ ,  $^{269}\text{Hs}$ ,  $^{265}\text{Sg}$  and  $^{261}\text{Rf}$ , of the decay sequence (see also sections 5.6, 5.8 and 5.10). This scenario also suggests that *SF* occurs for  $^{261}\text{Rf}$  only for the low-spin 3/2 isomeric state and is suppressed for the high-spin (11/2) ground state (see also the discussion below for the  $^{275}\text{Ds}$  decay chain and  $^{267}\text{Sg}$ ).

### $^{275}\text{Ds}$

The hitherto unknown isotope  $^{275}\text{Ds}$  was synthesized for the first time in a recent experiment at the FLNR JINR SHE-factory in the reaction  $^{48}\text{Ca} + ^{232}\text{Th}$  [29]. A total of six decay chains were accumulated of which three were of type *ER*- $\alpha$ - $\alpha$ -*SF*, ending in fission of  $^{267}\text{Sg}$ . The same number of decay chains was recorded showing the decay sequence *ER*- $\alpha$ - $\alpha$ -*SF*, terminated by *SF* of  $^{263}\text{Rf}$ . (see also sections 5.6, 5.8 and 5.10).

For the second decay chain member,  $^{271}\text{Hs}$ , two different decay times correlated to two different  $\alpha$  decay energies were observed. For  $^{267}\text{Sg}$  the three observed *SF* events exhibited decay times similar to what had been observed in chemistry experiments before [228, 229], while the decay times of the three  $\alpha$  decays of this isotope were detected with an average decay time of about 10 minutes, a time interval much longer than the maximum time range of 300 s accessible for the chemistry studies reported by Dvorak et al. in 2006 and 2008 [228, 229].

These findings were interpreted on the basis of model predictions for  $Q_\alpha$  values, decay times, the competition between  $\alpha$  decay and *SF* and spin values of ground and excited

states, referring to model calculations within the Di-Nuclear System (DNS) and the Two-Center Shell Model (TCSM) approach [236, 237, 238].

As a result the possible decay sequence from  $^{275}\text{Ds}$  to  $^{263}\text{Rf}$ , shown in Fig. 5.25 (taken from [29]), has been constructed. Here the authors follow the assumption that  $\alpha$  decay between analog spin states, i.e. Fermi transitions with  $\Delta S = 0$ , are favored and *SF* from high-spin states is hindered. This leads to new isomeric states in  $^{271}\text{Hs}$  and  $^{267}\text{Sg}$ , respectively, while in this scenario the isomer in  $^{267}\text{Sg}$  disintegrates by *SF* only and the ground state decays exclusively by  $\alpha$  emission. This situation is similar to other cases, apart from  $^{261}\text{Rf}$  in the previous paragraph, like, e.g.,  $^{259}\text{Sg}$  [113] or  $^{247}\text{Md}$  [52], where *SF* is also observed to decay only from the excited low-spin state (see also the discussion in subsection 4.3.1).

To conclusively address the scenarii proposed for  $^{253,257}\text{Ds}$  here, apart from a larger body of data, a more self-consistent model approach, than DNS or TCSM, explaining these intriguing observations based on firmer grounds would be highly desirable.

### $^{276}\text{Ds}$

Before the discovery and investigation of  $^{275}\text{Ds}$ ,  $^{276}\text{Ds}$  had already been synthesized at the FLNR JINR SHE-factory as the  $4n$  fusion-evaporation channel of the same reaction  $^{48}\text{Ca} + ^{232}\text{Th}$  at lower beam energies [29]. The in total used three beam energies allowed the construction of an excitation function for an excitation energy range ranging from  $E^* = 35$  MeV to 50 MeV with the maximum located at 40 MeV.

In total seven events consisting of four *ER-SF* and three *ER*- $\alpha$ - $\alpha$ -*SF* decay sequences. The isotopes  $^{272}\text{Hs}$  and  $^{268}\text{Sg}$  were also observed for the first time. The authors support the isotope assignment comparing properties like  $\alpha$  decay energies and partial  $\alpha$  and *SF* half-lives to neighboring isotopes. In particular, the systematic behavior of partial *SF* half-life as shown in Fig. 5.24 (Fig. 3 in Ref. [182]) and Table II in Ref. [182] show the onset of

a rising  $\log_{10} T_{SF}$  value towards lower neutron numbers with  $\log_{10} T_{SF} = -0.357$  s for  $^{276}\text{Ds}$  (166 neutrons) as compared to the value for  $^{279}\text{Ds}$  (169 neutrons) of -0.658 s approaching the value for  $^{279}\text{Rg}$  (168 neutrons) of -0.148 s, lying, however, a bit below the general trend. An update of Fig. 5.24 was not provided by Oganessian et al. in Ref. [29].

### $^{277}\text{Ds}$

Utyonkov et al. collected additional data for  $^{277}\text{Ds}$ , being an  $\alpha$ -decaying member of the  $^{285}\text{Fl}$  decay chain [199] (see subsection 5.16).

### $^{279}\text{Ds}$

Also for  $^{279}\text{Ds}$  new  $\alpha$ -decay events were observed at the new SHE-factory of FLNR JINR in the irradiation of  $^{242}\text{Pu}$  and of  $^{238}\text{U}$ , as member of the decay chains of  $^{283}\text{Cn}$  and  $^{287}\text{Fl}$  [206] (see subsections 5.14 and 5.16). The authors claim that the  $SF$  branch of  $^{279}\text{Ds}$  was observed in terms of ER- $SF$  correlations, where one or two  $\alpha$  decays for  $^{238}\text{U}$  and  $^{242}\text{Pu}$ , respectively, were missing due to the limited  $\alpha$ -particle detection efficiency.

## 5.13. Roentgenium - $Z=111$

For three roentgenium isotopes new or updated data were collected, being members of decay sequences of heavier nuclides.

### $^{279}\text{Rg}$

Being the third member of the  $^{287}\text{Mc}$  decay chain,  $^{279}\text{Rg}$  was also observed in the recent experiment at the FLNR JINR SHE-factory [182]. Apart from two additional  $\alpha$ -decay events, one  $SF$  was observed for the first time for this roentgenium isotope, leading to an  $\alpha$  decay branch of  $87^{+5}_{-19}\%$  (see Table 5.1). Following the argumentation of the authors, the  $SF$  assignment is supported by a missing  $\alpha$  decay being very unlikely, and by decay time systematics. The partial  $SF$  half-life resulting from the observation of the new  $SF$  branch in  $^{279}\text{Rg}$  is included in the half-life systematics in Fig. 5.24.

### $^{281,282}\text{Rg}$

In 2019 Khuyagbaatar et al. reported the  $^{294}\text{Ts}$  decay hitherto unpublished data for  $^{293}\text{Ts}$ .  $^{281,282}\text{Rg}$  are the first  $\alpha$ -decaying members of the decay chains of these nuclei [176] (see subsection 5.19).

## 5.14. Copernicium - $Z=112$

For the last but one element for which an isotope was produced by cold fusion, new data were accumulated for all five isotopes synthesized in hot fusion reactions.

### $^{281}\text{Cn}$

In 2018 Utyonkov et al. reported additional data for  $^{281}\text{Cn}$ , being an  $\alpha$ -decaying member of the  $^{285}\text{Fl}$  decay chain [199] (see subsection 5.16).

### $^{282,283,287,284}\text{Cn}$

As mentioned repeatedly above, Oganessian et al. irradiated at the new SHE-factory of FLNR JINR targets of  $^{242}\text{Pu}$  and of  $^{238}\text{U}$  with  $^{48}\text{Ca}$ , each at two beam energies [206] (see also section 5.16).

$^{282}\text{Cn}$  with its 100%  $SF$  branch, was only observed as  $\alpha$ -decay daughter of the  $4n$  ER  $^{286}\text{Fl}$ , for the plutonium target. Fourteen new  $SF$  decays were accumulated. In the same experiment,  $^{283}\text{Cn}$  was produced directly as the  $3n$  ER in the reaction with the uranium target, and as the first member of  $\alpha$ -decay chain of  $^{287}\text{Fl}$ . For the direct production 16 new decay chains were observed, improving the uncertainties of experimental decay properties, including decay energies, half-lives and branching ratios as shown in Table 5.1.

In an investigation of flerovium decay chains new data have been collected in two experimental runs at the GSI gas-filled separator TASCA for  $^{282,284}\text{Cn}$  [33, 239] and  $^{285}\text{Cn}$  [240] (see also section 5.16).

Terminating the  $^{290}\text{Lv}$  of the decay chain by  $SF$   $^{282}\text{Cn}$  was observed in a very recent experiment at the BGS of LBNL [241] (see subsection 5.18).

## 5.15. Nihonium - $Z=113$

### $^{282}\text{Nh}$

The lightest nihonium isotope, produced directly in a hot fusion reaction, i.e.  $^{237}\text{Np}(^{48}\text{Ca},3n)^{282}\text{Nh}$ , was recently also observed as the  $\alpha$ -decay daughter in the decay chain of  $^{286}\text{Mc}$  [181] (see subsection 5.17).

### $^{285,286}\text{Nh}$

$^{289,290}\text{Mc}$  are the first  $\alpha$ -decaying members of the decay chains of  $^{294}\text{Ts}$  and  $^{293}\text{Ts}$  for which Khuyagbaatar et al published updated and new data in 2019 [176] (see subsection 5.19).

## 5.16. Flerovium - $Z=114$

Flerovium is the first element that was produced in hot fusion only. Flerovium isotopes were also among the first superheavy nuclei being investigated at the new Dubna SHE-factory. But also at GSI and, as a decay product of the first livermorium synthesized at LBNL, flerovium isotopes were investigated recently.

### $^{284,285}\text{Fl}$

Results on the decay properties of the lightest flerovium isotopes  $^{284,285}\text{Fl}$  produced as the  $4n$  and  $3n$  fusion-evaporation channel, respectively, in the reaction  $^{48}\text{Ca}+^{240}\text{Pu}$ , were first reported by Utyonkov et al. in 2015 [200]. While the lighter of the two decays by  $SF$ , the  $ER$   $^{285}\text{Fl}$  is followed by five subsequent  $\alpha$  decays and the  $SF$  of  $^{264}\text{RF}$  terminating the decay chain.

In a new experiment, in order to favor the population of the  $4n$  fusion-evaporation channel,  $^{285}\text{Fl}$ , Utyonkov et al. chose the higher beam energy of the two, used in the first experiment, with  $E_{beam}=250$  MeV [199]. The  $3n$  channel was indeed not observed at this energy in the earlier measurement, but with the higher beam dose applied for the new run three  $^{285}\text{Fl}$  decay chains were observed which show consistent properties with respect to the three events

observed earlier at the lower beam energy. In addition three  $ER-SF$  correlations were detected and assigned to  $^{284}\text{Fl}$ .

### $^{286,287,288,289}\text{Fl}$

The irradiation of  $^{242}\text{Pu}$  and  $^{238}\text{U}$  targets with  $^{48}\text{Ca}$  beam, at the new SHE-factory of FLNR JINR at two beam energies provided new decay data for  $^{287}\text{Fl}$  and  $^{286}\text{Fl}$ , produced as  $3n$  and  $4n$  evaporation channels, respectively [206]. The various members of both decay chains with up to six subsequent  $\alpha$  decays for  $^{287}\text{Fl}$  terminated by fission of  $^{267}\text{Rf}$  for longest chains, and short decay sequences of  $\alpha-SF$  and  $SF$  of the ER for  $^{286}\text{Fl}$  have been presented in the respective subsections (see subsections 5.6, 5.8, 5.10, 5.12 and 5.14).

Concluding on specific decay properties observed in the  $^{287,286}\text{Fl}$  and  $^{283}\text{Cc}$  decay chains, in Ref. [206] the possible occurrence of isomeric states is discussed. For  $^{279}\text{Ds}$ , e.g., based on the observation of difference in the newly measured experimental  $Q_\alpha$  value of  $9.531(15)$  MeV, as compared to the theoretical value of  $10.11(12)$  MeV [117], Oganessian et al. suggest that  $SF$  occurs from an excited, low-spin isomeric state and is hindered for the supposedly high-spin ground state. A theoretical scenario supporting this interpretation had been suggested by Kuzmina et al. in Ref. [242].

The competition between  $SF$  and  $\alpha$  decay, with fission being hindered by the complex quantum structure of particular states, has been discussed in subsection 4.3.1, and for various examples through this review like, e.g., the case of  $^{247}\text{Md}$  in subsection 5.3 [52]. Similarly in Ref. [206] the possibly existence of isomeric states in  $^{286}\text{Fl}$  and  $^{282}\text{Cn}$  is discussed on the basis of three alpha decays which exhibit a by  $\approx 200$  keV lower  $\alpha$ -decay energy, as compared to the main activity. These findings show the potential offered by facilities like the SHE-factory and others providing high-intensity beams (see section 6), like here with up to 6 particle  $\mu\text{A}$  of  $^{48}\text{Ca}$  beam intensity. With these capabilities, it will be possible extending the study of the heaviest SHN towards more detailed spectroscopy, providing access to more complex nu-




Figure 5.26: (a) Summary of the decay chains of  $^{286,288}\text{Fl}$  (cf. Supplemental Material (of Ref. [33]) and Table II). (b),(c) Digitized preamplifier pulses of events associated with decay chains 08 and 02. Numbers in the panels are calibrated energies in mega-electron volts. Correlation times are given between recoil implantation (orange squares),  $\alpha$  decays (yellow squares), and fission (green squares). The 0.36(1)-MeV event in an upstream DSSD (blue) is in prompt coincidence with the 9.60(1)-MeV event in the implantation DSSD. Black triangles in the lower right corner of a square indicate detection during beam-off periods. (d) Suggested decay sequence of chain 02 through excited states in  $^{282}\text{Cn}$ . Tentative levels and transitions are dashed. (Figure and caption are taken from [33])

clear structure features.

The spontaneous fission decay of  $^{286}\text{Fl}$  has also been seen in the decay chain of  $^{294}\text{Og}$  detected by Brewer et al. in their first attempt to synthesize oganesson isotopes employing a target composed of a mixture of californium isotopes [243] (see subsection 5.20).

$^{286}\text{Fl}$  has very recently also been detected at the BGS of LBNL as  $\alpha$  decaying member of the  $^{290}\text{Lv}$  decay chain [241] (see subsection 5.18).

In 2021 and 2023 Såmark-Roth et al. reported on experiments employing the fusion-evaporation reactions  $^{48}\text{Ca} + ^{242,244}\text{Pu}$  to study the decay sequences of flerovium isotopes [33, 239]. In two experimental runs at the gas-filled separator TASCA of GSI, they collected a total of 29 decay chains in the  $3n$  and  $4n$  fusion-evaporation channels for the isotopes  $^{286}\text{Fl}$  (2 chains),  $^{288}\text{Fl}$  (12 chains) and  $^{289}\text{Fl}$  (15 chains). For details see Table 1 in Ref. [239]. The smoothly increasing trend of  $Q_\alpha$  values for the isotope series  $^{284}\text{Cn}$ - $^{288}\text{Fl}$ - $^{292}\text{Lv}$  is, according to the authors, indicative for the absence of a possible shell gap at  $Z=114$  (at  $N=174$ ) as predicted by some theoretical models as the next proton shell gap beyond  $^{208}\text{Pb}$  [33].

A specific feature observed in one of the decay chains for  $^{286}\text{Fl}$  is the detection of a 360-keV conversion electron,

marking the population and de-excitation of an excited state in  $^{282}\text{Cn}$  (see Fig. 5.26). On the basis of triaxial beyond-mean-field calculations (TBMF) [244, 245] and employing GEANT4 simulations to simulate the decay pattern involving this 360 keV  $CE$  four scenarios are discussed. All of them suggest an excited state populated by the  $^{286}\text{Fl}$   $\alpha$  decay having spin and parity of  $0^+$  (scenario i-iii) or  $2^+$  (scenario iv) [239]. For decay chain 8 (see supplemental material of Ref. [33]) which is assigned to  $^{288}\text{Fl}$  as  $ER$ , a second  $\alpha$  is observed instead of  $SF$  as seen for  $^{284}\text{Cn}$  in all other chains. This sequence is then terminated by  $SF$  of  $^{280}\text{Ds}$ . The TBMF calculations which are applied in two versions, one based on a Gogny, the other on a UNIDEF-Skyrme interaction, underline the importance of triaxiality for the ground and excited states, explicitly  $0^+$  states, of livermorium and flerovium isotopes and their decay products.

The details of the fifteen decay chains observed for  $^{289}\text{Fl}$  are discussed in detail in Ref. [240].

## 5.17. Moscovium - $Z=115$

Moscovium isotopes, in particular,  $^{288}\text{Mc}$  produced in  $^{48}\text{Ca}$ -induced reactions have proven to be good candidates for the investigation of the heaviest SHN, due to their large

production cross-sections of up to the order of 15 pbarn. From the nuclear structure point of view, they are promising candidates as well, considering the long  $\alpha$ -decay chains reaching down to lawrencium isotopes ( $Z=103$ ).

### $^{286}\text{Mc}$

The lightest moscovium isotope was first produced recently at the SHE-factory of JINR/FLNR employing the new gas-filled separator DGFRS-2 [246] as  $5n$  fusion-evaporation channel of the reaction  $^{48}\text{Ca} + ^{243}\text{Am}$ , by the observation of one decay chain ending with SF of  $^{266}\text{Db}$  [182].

### $^{287}\text{Mc}$

In Ref. [181] the authors report also on new data for the decay of  $^{287}\text{Mc}$ , for which four new chains were observed including the observation of SF for  $^{279}\text{Rg}$  in one of them.

### $^{288,289}\text{Mc}$

In the two experiments at the SHE-factory of FLNR JINR irradiating  $^{243}\text{Am}$  targets with  $^{48}\text{Ca}$  projectiles, Ref.s [181, 182], 52 and 55 new decay chains, respectively, were observed for  $^{288}\text{Mc}$  produced as the  $3n$  fusion-evaporation channel. This adds up to a total production of about 210 decay sequences observed for this nucleus, including measurements at GSI and LBNL [247, 186, 248, 147]. The new data confirm the earlier observation. The sequence of nuclei populated by subsequent  $\alpha$  decay down to  $^{268}\text{Db}$  which terminates the sequence, according to present assignments, by SF (see subsections 5.7 and, in particular, 4.3.2 for the discussion of a possible SF-EC decay competition).

In addition to those longer decay chains, a total of 24 shorter ones of type ER- $\alpha$ -SF or ER- $\alpha$ - $\alpha$ -SF have been observed in the various experiments. The origin and interpretation of these chains, including scenarii which propose  $\beta$  decay ( $\beta^+/\text{EC}$ ) of  $^{288}\text{Mc}$  and its daughter nucleus  $^{284}\text{Nh}$  [248] or the assignment of  $^{289}\text{Mc}$  as the decaying ER [182]) are even after the new findings at the SHE-factory still under debate. Despite this situation and valid counterarguments opposing the assignment by Oganessian et al, these latter

interpretation has been adopted in Table 5.1.

As a possible solution the authors of Ref. [182] propose the employment of instrumentation providing mass identification like the trochoidal mass spectrometer FIONA of LBNL [147] or the MASHA separator at FLNR JINR [249]. The separator-spectrometer setup S<sup>3</sup> [250], approaching the completion of its installation, together with highest beam intensities from the LINAC of GANIL SPIRAL2 and its upgrade, the new injector NEWGAIN [251], being presently constructed, would be an ideal tool to tackle this problem (see section 6).

### $^{289,290}\text{Mc}$

In 2019 Khuyagbaatar et al. reported the  $^{294}\text{Ts}$  decay hitherto unpublished data for  $^{293}\text{Ts}$ .  $^{289,290}\text{Mc}$  are the first  $\alpha$ -decaying members of the decay chains of these nuclei [176] (see subsection 5.19).

## 5.18. Livermorium - $Z=116$

Also for second heaviest even- $Z$  element new data has been accumulated for its lightest isotope  $^{290}\text{Lv}$ , as a member of the  $^{294}\text{Og}$  decay chain as well as in direct production.

### $^{290}\text{Lv}$

In an interesting approach,  $^{290}\text{Lv}$  was detected as a member of the  $^{294}\text{Og}$  decay chain by Brewer et al. in their first attempt to synthesize oganesson isotopes, employing a target composed of a mixture of californium isotopes [243] (see subsection 5.20).

Very recently  $^{290}\text{Lv}$  has also been synthesized at the BGS of LBNL. Irradiating a  $^{244}\text{Pu}$  target with a  $^{50}\text{Ti}$  beam, produced with the 88-inch cyclotron of LBNL, Gates et al. observed two decay chains of this nucleus with a production cross-section of 0.44 pbarn [241]. For both events the same decay sequence was observed, consisting of  $\alpha$  decay of  $^{290}\text{Lv}$  and  $^{286}\text{Fl}$ , terminated by SF of  $^{282}\text{Cn}$ .

## 5.19. Tennessine - $Z=117$

For last discovered element with the second-highest atomic number only one activity has been reported recently, studying both known tennessine isotopes.

### $^{293,294}\text{Ts}$

In 2015 Khuyagbaatar et al. had reported two long decay chains, observed at the GSI gas-filled separator TASCA in an irradiation of  $^{249}\text{Bk}$  with  $^{48}\text{Ca}$  beam, consisting of seven consecutive  $\alpha$  decays following the ER  $^{294}\text{Ts}$  and being terminated by  $SF$  of  $^{266}\text{Lr}$  [187]. These two chains are by one  $\alpha$  decay longer than four decay sequences observed at the DGFRS of FLNR JINR, which end with  $SF$  of  $^{270}\text{Db}$  [188].

In a paper in 2019, more details of this experiment are discussed and two shorter decay chains from the same irradiation assigned to  $^{293}\text{Ts}$  are presented [176]. These chains running down to  $^{277}\text{Mt}$  are in agreement with two out of sixteen decay sequences observed also at the DGFRS of FLRN JINR [188], where the additional 14 decay chains are by one member shorter and end with  $SF$  of  $^{273}\text{Rg}$ .

In line with what was discussed earlier in this review, e.g. for the dubnium isotopes  $^{266-268}\text{Db}$ , Khuyagbaatar et al. consider the possibility that  $^{277}\text{Mt}$  undergoes  $EC$  decay and the observed  $SF$  belongs to  $^{277}\text{Hs}$  which would be in agreement with the known properties of this nucleus.

## 5.20. Oganesson - $Z=118$

Finally for the only one known isotope of the heaviest element oganesson,  $^{294}\text{Og}$ , only the attempt to use a composite target, holding a cocktail of different californium isotopes was published recently.

### $^{294}\text{Og}$

The nucleus with the highest atomic charge  $Z=118$  and together with  $^{294}\text{Ts}$  the highest mass number  $A=294$ ,  $^{294}\text{Og}$  was observed four times [223], before Brewer et al. followed various theoretical predictions for synthesis cross-section of up to  $\approx 80$  pbarn of oganesson isotopes with mass numbers

from 292 to 298, employing the californium isotopes  $^{249}\text{Cf}$  to  $^{252}\text{Cf}$  as target material.

In a first attempt they used a mixed target consisting of a combination of  $^{249}\text{Cf}$  and  $^{251}\text{Cf}$ . For the beam energy two values were chosen with 252 MeV favoring the  $3n$  fusion-evaporation channel and 258 MeV populating most likely the  $4n$  channel. One decay chain was observed, with three subsequent decays assigned to the sequence of a  $^{294}\text{Og}$   $\alpha$  decay, followed by  $\alpha$  decay of  $^{290}\text{Lv}$  and terminated by  $SF$  of  $^{286}\text{Fl}$ , increasing the total number of observed  $^{294}\text{Og}$  nuclei to five.

Table 5.1: List of basic decay properties, including year of discovery and relevant literature references, for all hitherto experimentally known isotopes ( ${}^A Z$ ) and isomeric states ( ${}^{Am} Z$  from  $Z = 99$  to 118. Most of the data were retrieved from the various services offered by the National Nuclear Data Center (NNDC) of Brookhaven National Laboratory (BNL) via its website: <https://www.nndc.bnl.gov/> (see also Ref. [252]).  $Q_\alpha$  values are taken from [117]. Exceptions are indicated in the footnotes to this table. Half-lives  $T_{1/2}$ , and g.s. spins and parities,  $J_{gs}^\pi$  (where available), are given also for isomeric states. For excitation energies of isomeric states please refer to the quoted literature or the to the respective (sub)section in this review. Symmetric error bars are given in brackets for the last digits of the corresponding value, while for asymmetric ones they are quoted in the value units.  $Q_\alpha$  values marked by # are not purely experimental but are estimated on the basis of the values of neighboring nuclei (see explanation in [117]). For isomers the excitation energy  $E^*$  of the meta-stable state is shown instead of  $Q_\alpha$ , if known. The year of discovery has been verified with the data reported in [116], including updates from Discovery of Nuclides Project at <https://people.nscl.msu.edu/~thoennes/isotopes/>. For references that were originally published in Russian language their corresponding publications are given which provide the translation in English.

Isotope	$T_{1/2}$	$J_{gs}^\pi$	decay	branching	$Q_\alpha(g.s.)$	discovery	references
					mode		
					[%]	[keV]	year
${}^{240}\text{Es}$	6.7(13) s		$\alpha, EC, SF$	70/30/?	8260(60)	2017	[37, 35, 34]
${}^{241}\text{Es}^a$	$4.3^{+2.4}_{-1.2}$ s		$\alpha, EC$		8259(17)	1996	[120, 35, 36]
${}^{242}\text{Es}$	17.8(16) s		$\alpha, EC$	57/43	8160(20)	1994	[253, 120, 254, 51, 128]
${}^{243}\text{Es}$	22(1) s	(3/2 $^-$ )	$\alpha, EC, SF$	61/39/<1	8072(10)	1973	[255, 118, 253, 124, 51, 133] [35, 52]
${}^{244}\text{Es}$	37(4) s		$\alpha, EC$	4/96	7940(100) #	1973	[255, 256, 122, 135, 257]
${}^{245}\text{Es}$	1.12(6) min	(3/2 $^-$ )	$\alpha, EC$	54/46	7909(3)	1967	[258, 259, 118, 124, 196, 213] [178, 133]
${}^{246}\text{Es}$	7.5(5) min		$\alpha, EC$	9.9/90.1	7640(100) #	1954	[260, 258, 256, 118, 261, 124] [196, 178, 136]
${}^{247}\text{Es}$	4.55(26) min	(7/2 $^+$ )	$\alpha, EC$	7/93	7464(20)	1967	[258, 118, 124, 53]
${}^{248}\text{Es}$	24(2) min	(2 $^-, 0^+$ )	$\alpha, EC$	0.3/99.7	7160(50) #	1956	[262, 263, 256, 264, 118, 261]
${}^{249}\text{Es}$	102.2(6) min	7/2 $^+$	$\alpha, EC$	0.57/99.43	6940(30) #	1956	[265, 263, 266, 267, 264, 268] [118, 124, 125]
${}^{250}\text{Es}$	8.6(1) min	(6 $^+$ )	$\alpha, EC$	0>97/<3	6830(120) #	1956	[265, 263, 267, 264, 268, 269]
${}^{250m}\text{Es}$	2.25(5) min	1( $^-$ )	$EC$	100		1970	[263, 270]
${}^{251}\text{Es}$	33(1) h	3/2 $^-$	$\alpha, EC$	99.50/0.50	6597.1(10)	1956	[265, 263, 267, 271, 264, 268] [272, 137]
${}^{252}\text{Es}$	417.7(19) d	(5 $^-$ )	$\alpha, EC$	78/22	6738.6(5)	1956	[265, 273, 274, 268, 119, 272] [124, 272, 275]
${}^{253}\text{Es}$	20.47(2) d	7/2 $^+$	$\alpha, EC$	100/8.8E-6	6739.24(5)	1954	[276, 277, 278, 279, 264, 268] [119, 280, 281, 282, 283, 284] [275, 285]
${}^{254}\text{Es}$	275.7(5) d	(7 $^+$ )	$\alpha,$ $\beta^-, SF$	100 1.7E-4/<3.0E-6	6617.2(5)	1954	[286, 277, 287, 288, 289, 268] [279, 290, 291, 292, 293, 119] [294, 284, 275, 295, 285]
${}^{254m}\text{Es}$	39.3(5) h	2 $^+$	$\beta^-, IT$ $\alpha, EC, SF$	98/3 0.32/0.08/0.05	84.2(25)		[278, 296, 297, 275, 295]
${}^{255}\text{Es}$	39.8(12) d	(7/2 $^+$ )	$\alpha, EC, SF$	92/8/4.1E-3	6436.3(13)	1954	[286, 298, 291, 299, 268, 275] [285]
${}^{256}\text{Es}$	25.4(12) min	(1 $^+, 0^-$ )	$\beta^-$	100	6230(220) #	1955	[300, 301, 302, 299, 303]
${}^{256m}\text{Es}$	7.6 h	(8 $^+$ )	$\beta^-$	100		1976	[304, 86, 275, 295]
${}^{257}\text{Es}$	7.7(2) d		$\beta^-, SF$	100/?	6050(200) #	1987	[305]
${}^{241}\text{Fm}$	0.73(6) ms		$\alpha, EC, SF$	<14/<12/>78	8860(320) #	2008	[257]

$^{242}\text{Fm}$	0.8(2) ms	$0^+$	$SF$	100	8700(500) #	1975	[306, 307, 257]
$^{243}\text{Fm}$	231(9) ms	$(7/2^-)$	$\alpha, EC, SF$	$91/<10/9$	8690(50)	1981	[308, 309, 257, 141]
$^{244}\text{Fm}$	3.21(8) ms	$0^+$	$\alpha, EC, SF$	$<1/<2/>97$	8550(200) #	1967	[142, 310, 311, 312, 307, 257] [313, 189, 141, 103, 314]
$^{245}\text{Fm}$	5.4(6) ms		$\alpha, EC, SF$	$88.5/11.5/<0.3$	8305(25)	1967	[142, 307, 315, 316, 141, 162] [103, 89, 317]
$^{246}\text{Fm}$	1.53(4) s	$0^+$	$\alpha, EC, SF$	$93.6/6.4/\leq 1.3$	8379(5)	1966	[318, 319, 142, 310, 320, 120] [321, 51, 322, 323, 324, 316] [62, 317, 21]
$^{247}\text{Fm}$	31(1) s	$(7/2^+)$	$\alpha, EC$	64/36	8258(10)	1967	[319, 325, 326, 88, 327, 328]
$^{247m}\text{Fm}$	5.1(2) s	$(1/2^+)$	$\alpha, IT$	88/12	45(7)	1967	[319, 326, 88, 327]
$^{248}\text{Fm}$	34.5(12) s	$0^+$	$\alpha, EC, SF$	0.99/0.1/?	7995(8)	1958	[329, 318, 142, 259, 330, 307] [331, 332, 333, 82, 334, 91] [134, 21]
$^{249}\text{Fm}$	1.6(1) m	$(7/2^+)$	$\alpha, EC$	15.6/84.4	7709(6)	1959	[335, 318, 330, 326, 336, 93] [213, 108, 125, 109, 337, 21]
$^{250}\text{Fm}$	30.9(10) m	$0^+$	$\alpha, EC, SF$	$100/?/6.9E-3$	7557(8)	1954	[338, 339, 318, 340, 341, 264] [342, 343, 344, 331, 345, 346] [347, 85, 196, 348, 98, 91] [170, 144, 349, 48, 21]
$^{250m}\text{Fm}$	1.90(5) s		$IT$	100		1973	[350, 342]
$^{251}\text{Fm}$	5.30(8) h	$(9/2^-)$	$\alpha, EC$	1.80/98.20	7424.5(10)	1957	[339, 297, 271, 264, 351, 88] [143, 144, 178, 295]
$^{252}\text{Fm}$	25.38(12) h	$0^+$	$\alpha, SF$	$100/2.3E-3$	7557(8)	1956	[352, 353, 354, 355, 264, 321] [275, 139, 295]
$^{253}\text{Fm}$	3.00(12) d	$(1/2^+)$	$\alpha, EC$	$1.7E-4/<3.0E-6$	7197.9 (10)	1959	[356, 357, 358, 264, 359, 94] [58, 275, 295]
$^{254}\text{Fm}$	3.240(2) h	$0^+$	$\alpha, SF$	99.94/0.06	7307.3(10)	1954	[360, 286, 278, 291, 264, 321] [275, 295, 21]
$^{255}\text{Fm}$	27.07(7) h	$7/2^+$	$\alpha, SF$	$1.7E-4/<3.0E-6$	7240.6(5)	1954	[286, 278, 289, 361, 272, 307] [362, 284, 275, 363, 295]
$^{256}\text{Fm}$	157.1(12) m	$0^+$	$\alpha, SF$	91.90/8.10	7025.3(19)	1954	[300, 364, 365, 366, 367, 302] [368, 264, 86, 369, 272, 275] [295, 137]
$^{256m}\text{Fm}$	70(5) ns	$7^-$	$IT, SF$		1425	1989	[86]
$^{257}\text{Fm}$	100.5(2) d	$(9/2^+)$	$\alpha, SF$	99.79/0.21	6863.7(9)	1964	[370, 365, 298, 371, 372, 119] [307, 275]
$^{258}\text{Fm}$	370(140) $\mu$ s	$0^+$	$SF$	100	6660(200) #	1971	[373, 374, 375, 376, 377, 378] [379, 119]
$^{259}\text{Fm}$	1.5(2) s		$SF$	100	6470(200) #	1980	[380, 376, 307]
$^{244}\text{Md}^a$	$0.30_{-0.09}^{+0.19}$ s		$\alpha, EC$	$\approx 100/<14$	8950(80)	2020	[34, 35, 36]
$^{245}\text{Md}$	0.90(25) ms	$(7/2^-)$	$\alpha, SF$	?/?	9010(120) #	1996	[120, 35, 36]
$^{245m}\text{Md}$	$0.35_{-0.16}^{+0.23}$ s	$1/2^-$	$\alpha, EC$	?/?	$\approx 300$	1996	[120, 307, 35, 36]
$^{246m1}\text{Md}^b$	0.9(2) s		$\alpha, EC$	$100/?/?$	8890(40)	1996	[120, 51]
$^{246m2}\text{Md}^b$	4.4(8) s		$\alpha, EC$	$<23/>77/?$		1996	[120, 51]
$^{247}\text{Md}$	1.3(1) s	$(7/2^-)$	$\alpha, SF$	99.14/0.86	8764(10)	1981	[308, 152, 124, 51, 52]
$^{247m}\text{Md}$	230(3) ms	$1/2^-$	$IT, SF$	79/21	153 <sup>c</sup>	1994	[152, 124, 51, 52]
$^{248}\text{Md}$	7.0(30) s		$\alpha, EC$	20/80	8500(30)	1973	[259, 256, 122, 211]
$^{249}\text{Md}$	25.6(9) s	$(7/2^-)$	$\alpha, EC$	$75/<25$	8441(18) #	1973	[259, 197, 122, 381, 178, 133] [153, 75]
$^{249m1}\text{Md}$	1.9(9) s	$1/2^-$	$\alpha$	100		2001	[122, 382]
$^{249m2}\text{Md}$	2.4(3) ms		$IT$	100	$\geq 910$	2001	[75]
$^{250}\text{Md}$	53.4(45) s		$\alpha, EC$	7/93	8155(28)	1973	[259, 256, 197, 178, 54, 381] [136]
$^{250m}\text{Md}$	42.4(45) s		$\alpha$	100	123	2019	[136]
$^{251}\text{Md}$	4.27(26) m	$(7/2^-)$	$\alpha, EC$	$<10/>90$	7963(4)	1973	[259, 331, 53, 383, 178, 153] [75]
$^{251m}\text{Md}$	1.37(6) s	$(23/2^+)$	$IT$	100	$\geq 844$	2001	[75]

$^{252}\text{Md}$	2.3(8) m	(1/2 <sup>+</sup> )	$EC$	100	7740(110)#+	1966	[353, 259, 178]
$^{253}\text{Md}$	$6_{-3}^{+12}$ m	(7/2 <sup>-</sup> )	$\alpha, EC$	$\approx 0.7/\approx 99.3$	7573(8)	1992	[384, 124, 58, 125]
$^{254}\text{Md}$	10(3) m		$EC$	$\leq 100$	7800(140)#+	1970	[385]
$^{254m}\text{Md}$	28(8) m	(3 <sup>-</sup> )	$EC$	$\leq 100$	50	1970	[385]
$^{255}\text{Md}$	27(2) m	(7/2 <sup>-</sup> )	$\alpha, EC$	7/93	7905.6(17)	1958	[364, 365, 385, 386, 272, 307] [295]
$^{256}\text{Md}$	77.7(15) m	(1 <sup>-</sup> )	$\alpha, EC, SF$	9.5/90.5/<3	7740(110)#+	1955	[387, 364, 365, 385, 386, 119] [272, 295]
$^{257}\text{Md}$	5.52(5) h	(7/2 <sup>-</sup> )	$\alpha, EC, SF$	15/85/<1	7557.1(9)	1965	[365, 385, 386, 119, 307, 137] [295]
$^{258}\text{Md}$	51.50(29) d	(8 <sup>-</sup> )	$\alpha, EC, SF$	100	7271.3(19)	1968	[388, 385, 375, 377, 119, 272] [295]
$^{258m}\text{Md}$	57.5(11) m	(1 <sup>-</sup> )	$\alpha, EC, SF$	,1.2/>85/<15		1968	[388, 385, 375, 377, 119, 272]
$^{259}\text{Md}$	1.61(6) h		$\alpha, SF$	<1.3/100	7050(100)#+	1982	[389, 379, 369, 119, 307]
$^{260}\text{Md}$	31.8(5) d		$\alpha, EC, \beta^-, SF$	<25/<23/<10/>42	6940(300)#+	1986	[379, 390, 295]
$^{249}\text{No}^d$	38.1(25)ms		$\alpha$	100	9276(20)	2021	[162, 103, 89]
$^{250}\text{No}$	4.6(2) $\mu\text{s}$		$\alpha, SF$	<2/100	8950(200)#+	2003 <sup>e</sup>	[306, 391, 30, 87, 392]
$^{250m}\text{No}^f$	$43.0_{-15}^{+22}$ $\mu\text{s}$		$SF$	100		2006	[30, 393, 394, 87, 31, 165] [161]
$^{251}\text{No}$	0.8(1) s	(7/2 <sup>+</sup> )	$\alpha, EC, SF$	90/>0/<1.4E-3	8752(4)	1967	[395, 307, 122, 391, 30, 326] [328, 89]
$^{251m1}\text{No}$	1.00(4) s	(1/2 <sup>+</sup> )	$\alpha,$	100	106(6)	2001	[122, 88, 327, 83, 113, 89]
$^{251m2}\text{No}$	2 $\mu\text{s}$		$\gamma,$	100	$\geq 1700$	2001	[88, 327, 83, 113, 89]
$^{252}\text{No}$	2.44(4) s	0 <sup>+</sup>	$\alpha, EC, SF$	65.3/1.7/33	8549(5)	1967	[396, 395, 397, 398, 399, 330] [400, 401, 391, 402, 30, 331] [333, 334, 403, 159, 165, 161]
$^{252m}\text{No}$	109(3) ms	(8 <sup>-</sup> )	$IT,$	100	1254.6(15)	1989	[342, 90, 91]
$^{253}\text{No}$	1.61(5) m	9/2 <sup>-</sup>	$\alpha, EC$	55/45	8415(4)	1967	[396, 395, 404, 405, 326, 108] [406, 125, 109, 55, 407, 159] [56]
$^{253m}\text{No}$	31.1(12) $\mu\text{s}$	5/2 <sup>+</sup>	$\alpha, EC, IT$			1973	[408, 93, 94, 58, 56]
$^{254}\text{No}$	51.2(4) s	0 <sup>+</sup>	$\alpha, EC, SF$	90/10/0.17	8226(8)	1966	[329, 409, 410, 411, 350, 197] [412, 342, 413, 343, 331, 95] [414, 415, 416, 417, 418, 178] [159, 165, 161, 166]
$^{254m1}\text{No}$	265(1) ms	(8 <sup>-</sup> )	$IT, \alpha, SF$	99.98/ $\leq 0.01/0.02$	1296.4(11)	1973	[350, 95, 96, 97, 98, 166]
$^{254m2}\text{No}$	184(3) $\mu\text{s}$	(16 <sup>+</sup> )	$IT, SF$	100/ $\leq 0.012$	2314.8+ $\epsilon$	2006	[95, 96, 97, 98, 166]
$^{255}\text{No}$	3.52(18) m	(1/2 <sup>+</sup> )	$\alpha, EC$	30/70	8428.2(30)	1967	[419, 420, 395, 421, 422, 423] [424, 425, 173, 426, 143, 48]
$^{255m1}\text{No}$	109(9) $\mu\text{s}$	(11/2 <sup>-</sup> )	$\gamma$	100	230-240	2006	[173, 48]
$^{255m2}\text{No}$	77(6) $\mu\text{s}$	(11/2 <sup>-</sup> )	$\gamma$	100	1400-1600	2006	[48]
$^{255m3}\text{No}$	$1.2_{-0.4}^{+0.6}$ $\mu\text{s}$	(11/2 <sup>-</sup> )	$\gamma$	100	$\leq 1500$	2006	[48]
$^{256}\text{No}$	2.93(7) s	0 <sup>+</sup>	$\alpha, SF$	99.47/0.53	8582(5)	1963	[427, 428, 395, 429, 138, 139] [100]
$^{256m}\text{No}^g$	$7.8_{-2.6}^{+8.3}$ $\mu\text{s}$		$\alpha$		1089	2021	[100]
$^{257}\text{No}$	24.5(5) s	(7/2 <sup>+</sup> )	$\alpha, EC, SF$	85/15/<1.5	8477(6)	1967	[395, 421, 422, 307, 430, 424] [431, 359, 432, 433]
$^{258}\text{No}$	$1.23_{-11}^{+12}$ ms	0 <sup>+</sup>	$SF$	$\leq 100$	8150(100)#+	1969	[434, 377, 379, 435, 436] [437, 438]
$^{259}\text{No}$	58(5) m	(9/2 <sup>+</sup> )	$\alpha, EC, SF$	75/25/<10	7854(5)	1973	[439, 389, 307, 440, 295]
$^{260}\text{No}$	107(8) ms	0 <sup>+</sup>	$SF$	100	7700(200)#+	1969	[441, 442, 295, 438]
$^{262}\text{No}$	$\approx 5$ ms	0 <sup>+</sup>	$SF$	100	7250(300)#+	1989	[379, 307]
$^{251}\text{Lr}$	$24.4_{-4.5}^{+7.0}$ ms	(7/2 <sup>-</sup> )	$\alpha$	100	9210(19)	2001	[177]
$^{251m}\text{Lr}$	$42_{-14}^{+42}$ ms	(1/2 <sup>-</sup> )	$\alpha$	100	117 (9402(19))	2001	[177]
$^{252}\text{Lr}^h$	$0.41_{-0.05}^{+0.07}$ s	0 <sup>+</sup>	$\alpha, EC, SF$	$\approx 70-90/?/\approx 10-30$	9164(17)	2001	[208, 122, 211, 134]
$^{253}\text{Lr}$	0.65(4) s	(7/2 <sup>-</sup> )	$\alpha, EC, SF$	99/1	8918(20)	1985	[197, 443, 444, 307, 345, 178]
$^{253m}\text{Lr}$	1.44(10) s	(1/2 <sup>-</sup> )	$\alpha, SF$	88/12		2001	[122, 345]

$^{254}\text{Lr}$	11.9(9) s <sup>8</sup>	(4 <sup>+</sup> )	$\alpha, EC, SF$	71.70/28.30/<0.1	8822(8)	1981	[151, 445, 197, 443, 307, 121] [345, 54, 381, 213, 178, 136]
$^{254m}\text{Lr}$	20.3(42) s <sup>i</sup>	(1 <sup>-</sup> )	$\alpha$	100	108	2019	[136]
$^{255}\text{Lr}$	31.1(13) s	(1/2 <sup>-</sup> )	$\alpha, EC, SF$	99.70/0.30/≤0.1	8556(7)	1971	[446, 447, 448, 208, 307, 121] [449, 54, 113]
$^{255m1}\text{Lr}$	2.54(4) s	(7/2 <sup>-</sup> )	$\alpha, IT$	40/60	38(10)	2006	[53, 102, 54, 101, 383, 113]
$^{255m2}\text{Lr}$	<1 $\mu\text{s}$	(15/2 <sup>+</sup> )			740.0+ $\epsilon$ (5)	2009	[101]
$^{255m3}\text{Lr}$	1.70(3) ms	(25/2 <sup>+</sup> )	$\alpha, IT$	0.15/99.85	1408+ $\epsilon$ (10)	2008	[102, 54, 101, 383]
$^{256}\text{Lr}$	27.9(10) s		$\alpha, EC$	85/15	8850(120)	1965	[450, 451, 447, 448, 54, 452] [418, 453]
$^{257}\text{Lr}$	5.9(4) s		$\alpha, EC, SF$	100/<15/≤0.03	9070(30)	1971	[454, 447, 455, 448, 404, 307] [123, 108, 55]
$^{258}\text{Lr}$	3.86 <sup>+24</sup> <sub>-423</sub> s		$\alpha, SF$	97.4/2.6	8904(19)	1971	[447, 455, 448, 456, 457, 458] [216, 217, 218, 180]
$^{259}\text{Lr}$	6.2(3) s		$\alpha, SF$	78/22	8580(70)≠	1971	[447, 457, 459, 307, 460, 180]
$^{260}\text{Lr}$	180(30) s		$\alpha, EC$	8>60/<40	8400(140)≠	1971	[447]
$^{261}\text{Lr}$	39(12) m		$SF$	100	8140(200)≠	1987	[461, 462, 463]
$^{262}\text{Lr}$	≈4 h		$\alpha, EC, SF$	?/?<10	7990(200)≠	1987	[461, 462, 295]
$^{264}\text{Lr}^j$	4.8 <sup>+2.2</sup> <sub>-1.3</sub> h		$SF$	100	7400(300)≠	2022	[181, 182]
$^{266}\text{Lr}$	11 <sup>+21</sup> <sub>-5</sub> h		$SF$	100	7570(300)≠	2014	[175, 295, 176]
$^{253}\text{Rf}^k$	12.8 <sup>+7.0</sup> <sub>-3.4</sub> ms	(7/2 <sup>+</sup> )	$\alpha, SF$	12.5/87.5	9170(30)	1997	[404, 444, 307, 103, 89]
$^{253m}\text{Rf}$	48 <sup>+17</sup> <sub>-10</sub> $\mu\text{s}$		$\alpha, SF$	?/≤100	9430(300)≠	1997	[404, 444, 307, 103, 89]
$^{254}\text{Rf}$	22(8) $\mu\text{s}$	0 <sup>+</sup>	$\alpha, SF$	<1.5/100	9210(200)≠	1997	[464, 306, 404, 444, 465, 60] [61, 191]
$^{254m1}\text{Rf}$	4.7(11) $\mu\text{s}$	(8 <sup>-</sup> )	$IT, SF$	≈100/<10		2015	[60, 61]
$^{254m2}\text{Rf}$	247(73) $\mu\text{s}$	(16 <sup>+</sup> )	$IT, SF$	≈100/<40		2015	[60, 61]
$^{255}\text{Rf}$	1.67(2) s	9/2 <sup>-</sup>	$\alpha, EC, SF$	49/<0.08/51	9055(4)	1975	[466, 464, 467, 468, 469, 470] [471, 325, 312, 472, 404, 473] [307, 122, 474, 465, 143, 113] [113, 61, 192, 134]
$^{255m1}\text{Rf}$	50(17) $\mu\text{s}$	5/2 <sup>+</sup>	$IT$	100	≈135 keV	1975	[113, 61, 192]
$^{255m2}\text{Rf}^l$	15 <sup>+6</sup> <sub>-4</sub> $\mu\text{s}$		$IT$	100	900-1200 keV	1975	[61, 192]
$^{255m3}\text{Rf}^l$	38 <sup>+12</sup> <sub>-7</sub> $\mu\text{s}$		$IT$	100	1150-1450 keV	1975	[192]
$^{256}\text{Rf}$	6.66(6) ms	0 <sup>+</sup>	$\alpha, SF$	0.31/99.69	8926(15)	1975	[464, 468, 469, 475, 441, 404] [465, 190, 150, 212, 192, 165] [349, 103, 476]
$^{256m1}\text{Rf}$	25(2) $\mu\text{s}$	(5 <sup>-</sup> )	$IT, SF$	?/?	≈1120	2009	[106, 107, 195, 109, 103]
$^{256m2}\text{Rf}$	17(2) $\mu\text{s}$	(8 <sup>-</sup> )	$IT, SF$	?/?	≈1400	2009	[106, 107, 195, 109, 103]
$^{256m3}\text{Rf}$	27(5) $\mu\text{s}$		$IT, SF$	?/?	>2200	2009	[106, 107, 195, 109, 103]
$^{257}\text{Rf}$	4.5(2) s	(1/2 <sup>+</sup> )	$\alpha, EC, SF$	89.30/9.4/1.30	9083(8)	1969	[340, 341, 408, 477, 445, 469] [441, 404, 444, 307, 465, 94] [415, 109, 55, 104, 56]
$^{257m1}\text{Rf}$	4.1(4) s	11/2 <sup>-</sup>	$\alpha, EC, SF$	<88/11/<1.4	≈75	2009	[106, 108, 107, 478, 94, 83] [109, 55, 104, 56]
$^{257m2}\text{Rf}$	106(6) $\mu\text{s}$	(21/2 <sup>+</sup> )	$IT$		1151(11)	2009	[108, 107, 109]
$^{258}\text{Rf}$	9.871(68) ms	0 <sup>+</sup>	$\alpha, SF$	95.1/4.9	9196(8)	1969	[340, 445, 469, 441, 413, 400] [196, 150, 212, 165, 192]
$^{258m1}\text{Rf}$	2.4 <sup>+2.4</sup> <sub>-0.8</sub> ms		$\alpha, IT, SF$	?/?/?		2016	[76]
$^{258m2}\text{Rf}$	15(10) $\mu\text{s}$		$\alpha, IT, SF$	?/?/?		2016	[76]
$^{259}\text{Rf}$	2.5(3) s		$\alpha, EC, SF$	85/15/<3	9130(70)≠	1969	[340, 341, 479, 480, 481, 441] [482, 430, 196, 483, 425, 484]
$^{260}\text{Rf}$	21.7(8) ms	0 <sup>+</sup>	$\alpha, SF$	35/65	8900(200)≠	1985	[485, 397, 486, 487, 488, 489] [441, 379, 490]
$^{261}\text{Rf}$	70(3) s	(11/2)	$\alpha$	100	8650(70)	1970	[491, 492, 493, 494, 495, 496] [430, 497, 498, 230, 499, 500] [332, 501, 231, 29]
$^{261m}\text{Rf}$	1.9(4) s	(1/2)	$\alpha, SF$	18/82	234+ $\epsilon$ (57)	2006	[228, 229, 196, 502, 432, 503] [433, 332, 501, 231, 29]

$^{262}\text{Rf}$	47 ms	$0^+$	$SF$	100	8490(200)♯	1985	[504, 441, 435, 436, 496, 430]
$^{263}\text{Rf}^m$	$5.1^{+4.6}_{-1.7}$ s		$\alpha, SF$	100	8250(150)♯	1985	[198]
$^{265}\text{Rf}$	$1^{+1.2}_{-0.3}$ m		$SF$	$\approx 100$	7810(300)♯	2010	[201, 200, 199]
$^{267}\text{Rf}^n$	$48^{+23}_{-12}$ h		$SF$	100	7810(300)♯	2004	[205, 203, 204, 206]
$^{255}\text{Db}^h$	$2.7^{+0.4}_{-0.3}$ ms		$\alpha, SF$	33/66	9716(30)	2005	[398, 210, 134]
$^{256}\text{Db}$	$1.9^{+0.4}_{-0.3}$ s		$\alpha, EC, SF$	$\approx 70/\approx 30/\approx 0.02$	9340(30)	1999	[444, 507, 122, 211, 150, 212]
							[165, 134]
$^{257}\text{Db}$	$2.3(2)$ s	$(9/2^+)$	$\alpha, SF$	$\geq 94/\leq 6$	9206(20)	1985	[208, 197, 508, 444, 507, 381]
							[213, 509, 178, 150, 212, 165]
							[510, 511, 512]
$^{257m}\text{Db}$	0.67(6) s	$(1/2^-)$	$\alpha, SF$	$\geq 87/\leq 13$	250	1999	[444, 507, 122, 326, 381, 213]
							[509, 512]
$^{258}\text{Db}$	2.08(29) s	$(0^-)$	$\alpha, SF$	64/36	9437(100)	1981	[151, 445, 443, 121, 213, 76]
							[136, 150, 212, 192, 165, 513]
							[512]
$^{258m}\text{Db}$	4.37(16) s		$\alpha, EC$	77/23		1989	[197, 213, 76, 136]
$^{259}\text{Db}$	0.51(16) s	$(9/2^+)$	$\alpha$	100	9620(50)	2001	[121, 514, 213, 327, 113]
$^{260}\text{Db}$	1.52(13) s		$\alpha, EC, SF$	90/10	9500(40)♯	1970	[515, 516, 517, 518, 487, 519]
							[230, 424, 123, 520, 483]
$^{261}\text{Db}$	$4.1^{+1.4}_{-0.8}$ s		$\alpha, SF$	73/27	9220(100)♯	1970	[455, 518, 521, 522, 123, 523]
$^{262}\text{Db}$	34(4) s		$\alpha, EC, SF$	$46/\leq 5/54$	9050(100)♯	1971	[521, 524, 456, 525, 526, 527]
							[528, 529, 530, 531, 214, 532]
							[501, 216, 217, 218, 180]
$^{263}\text{Db}$	$27^{+10}_{-7}$ s		$\alpha, EC, SF$	42/3/55	8830(150)♯	1992	[533, 528, 505, 458, 534, 535]
							[530, 506, 180]
$^{265}\text{Db}$			$SF$	?/?/?	8830(150)♯	-	[219]
$^{266}\text{Db}^m$	$11^{+21}_{-4}$ min		$EC, SF$	?/100	8210(200)♯	2007	[202, 203, 474, 204, 184, 223]
							[12, 181, 182]
$^{267}\text{Db}^j$	$1.4^{+1.0}_{-0.4}$ h		$EC, SF$	?/100	7920(300)♯	2004	[536, 537, 203, 204, 182]
$^{268}\text{Db}^j$	$16^{+6}_{-4}$ h		$\alpha, EC, SF$	51/?/49	7700-8100	2004	[536, 537, 538, 539, 203, 540]
							[204, 541, 184, 185, 186, 223]
							[247, 248, 12, 181, 182, 542]
$^{270}\text{Db}$	$15^{+10}_{-4}$ h		$EC, SF$	?/100	8310(200)♯	2004	[536, 537, 538, 539, 203, 540]
							[541, 204, 175, 176]
$^{258}\text{Sg}$	$2.7^{+0.5}_{-0.4}$ ms	$0^+$	$\alpha, SF$	$<20/100$	9670(300)♯	1997	[404, 444, 327]
$^{259}\text{Sg}$	0.40(6) s	$(1/2^+)$	$\alpha, EC, SF^p$	$97/3/\leq 13$	9765(8)	1985	[543, 467, 475, 307, 327, 213]
							[544, 113]
$^{260}\text{Sg}$	4.83(37) ms	$0^+$	$\alpha, SF$	71/29	9901(10)	1984	[467, 545, 475, 546, 471, 213]
							[547]
$^{261}\text{Sg}$	0.18(5) s		$\alpha, EC, SF$	98.1/1.3/0.6	9714(15)	1984	[548, 445, 475, 471, 549, 473]
							[550, 551, 444, 552, 213, 107]
							[415, 547]
$^{262}\text{Sg}$	11.4(22) ms	$0^+$	$\alpha, SF$	6/94	9600(15)	2001	[49, 484, 332, 484, 553, 554]
							[50, 1]
$^{263}\text{Sg}$	$0.90^{+0.17}_{-0.16}$ s		$\alpha, SF$	87/13	9400(60)	1974	[480, 525, 482, 555, 473, 550]
							[556, 551, 557, 483, 520, 424]
							[425, 484, 332, 558, 559]
$^{263m}\text{Sg}$	0.43(11) s		$\alpha, IT, SF$	100/?/?		1995	[473, 550, 551, 444, 557, 483]
$^{264}\text{Sg}$	$60^{+26}_{-20}$ ms	$0^+$	$\alpha, SF$	$<20/\geq 80$	9210(200)♯	2006	[560, 556, 484, 332, 561, 226]
$^{265}\text{Sg}^m$	$14.4^{+3.7}_{-2.5}$ s	$(3/2)$	$\alpha, SF$	$50/\leq 50$	8970(50)♯	1974	[435, 492, 562, 495, 563, 496]
							[307, 564, 230, 499, 500, 228]
							[332, 501, 502, 565, 503, 566]
							[231, 29]
$^{265m}\text{Sg}$	$8.8^{+1.8}_{-1.3}$ s	$(11/2)$	$\alpha, SF$	$>49/\leq 51$		1974	[496, 230, 499, 228, 332, 501]
							[502, 561, 503, 231, 29]
$^{266}\text{Sg}$	$0.34^{+0.12}_{-0.09}$ s	$0^+$	$SF$	100	8800(100)♯	2006	[435, 492, 495, 563, 496, 498]

							[564, 499, 500, 228, 229, 204] [566, 567, 198]
$^{267}\text{Sg}^m$	$9.8_{-4.5}^{+11.3}$ s	$(9/2)^l$	$\alpha, SF$	100?	$8400(20)$ $^l$	2008	[229, 561, 566, 29]
$^{267m}\text{Sg}^m$	$100_{-39}^{+92}$ s	$(1/2)$	$SF$	100?	$8400(20)$ $^l$	2024	[29]
$^{268}\text{Sg}^o$	$13_{-4}^{+17}$ s		$SF$	100	$8300(300)\#$	2023	[28]
$^{269}\text{Sg}$	$3.1_{-1.1}^{+3.7}$ min		$\alpha$	100	$8580(70)$	2010	[201, 558, 200, 199]
$^{271}\text{Sg}^n$	$31_{-7}^{+13}$ s		$\alpha, SF$	73/27	$8.629(16)\#$	2004	[205, 204, 206]
 $^{260}\text{Bh}$	$35_{-9}^{+19}$ ms		$\alpha, EC, SF$	$\approx 100 / <18 / <18$	10400(60)	2008	[211]
$^{261}\text{Bh}$	$11.7_{-24}^{+37}$ ms		$\alpha, SF$	$100 / <10$	10500(70)	1989	[208, 151, 443, 568, 509, 178] [512]
$^{262}\text{Bh}$	$88_{-10}^{+11}$ ms		$\alpha$	$<100$	10319(15)	1981	[151, 445, 569, 443, 570, 307] [345, 568, 213, 348, 178, 512]
$^{264}\text{Bh}$	$0.99_{-0.18}^{+0.23}$ s		$\alpha, SF$	85/15	$9767(20)^q$	1995	[519, 230, 123, 571, 520, 424]
$^{265}\text{Bh}$	$0.94_{-0.31}^{+0.70}$ s		$\alpha$	100	$9660(210)\#$	2004	[123]
$^{266}\text{Bh}$	$10_{-1.6}^{+2.7}$ s		$\alpha$	100	$9430(80)\#$	2000	[530, 214, 572, 215, 216, 573] [217, 559, 180]
$^{267}\text{Bh}$	$16_{-4}^{+7}$ s		$\alpha$	100	$9230(200)\#$	2000	[530, 574, 575, 180]
$^{270}\text{Bh}^j$	$2.4_{-0.9}^{+4.4}$ min		$\alpha$	$\approx 100$	9060(100)	2007	[202, 474, 204, 223, 182]
$^{271}\text{Bh}^j$	$2.9_{-0.9}^{+2.2}$ s		$\alpha$	100	9420(90)	2004	[537, 203, 184, 185, 223, 182]
$^{272}\text{Bh}^j$	$8.8(0.7)$ s		$\alpha$	100	9300(50)	2004	[183, 538, 539, 576, 541, 184] [185, 577, 186, 223, 247, 248] [578, 12, 147, 181, 182]
$^{274}\text{Bh}$	$44_{-13}^{+34}$ s		$\alpha, SF$	100/?	8940(60)	2010	[25, 576, 204, 579, 184, 185] [176]
 $^{263}\text{Hs}$	$0.74_{-0.21}^{+0.48}$ ms		$\alpha, SF$	100/?	10730(80)	2009	[328]
$^{264}\text{Hs}$	$0.63_{-0.16}^{+0.34}$ ms	$0^+$	$\alpha, SF$	80/20	10591(20)	1986	[546, 471, 547]
$^{265}\text{Hs}$	$1.9_{-0.1}^{+0.2}$ ms		$\alpha, SF$	$100 / <1$	10470(15)	1984	[548, 468, 471, 549, 570, 444] [213, 547]
$^{265m}\text{Hs}$	$0.75_{-0.12}^{+0.17}$ ms		$\alpha, SF$		$\geq 300$	2009	[549, 444, 213]
$^{266}\text{Hs}$	$0.74_{-0.21}^{+0.48}$ ms	$0^+$	$\alpha, SF$	76/24	10346(16)	2001	[49, 554, 110, 580, 50, 581] [1, 38]
$^{266m}\text{Hs}$	$74_{-36}^{+354}$ ms		$\alpha$	$\approx 100$	$\approx 1200$	2012	[110, 580, 50, 581, 1, 38]
$^{267}\text{Hs}$	$53_{-8}^{+12}$ ms		$\alpha$	100	10038(13)	1995	[555, 473, 550, 551, 483, 424] [425, 558, 582, 559]
$^{267m}\text{Hs}$	$0.8_{-0.4}^{+4.0}$ s	$(3/2^+)$	$\alpha$	100		2004	[483, 425]
$^{268}\text{Hs}$	$0.4_{-0.2}^{+1.8}$ s	$0^+$	$\alpha$	100	9623(16)	2010	[558, 561]
$^{269}\text{Hs}^m$	$13_{-4}^{+10}$ s	$(9/2)$	$\alpha$	100	$9340(40)\#$	1996	[494, 562, 564, 499, 583, 500] [584, 561, 566, 231, 29]
$^{269m}\text{Hs}^m$	$2.8_{-1.3}^{+13.6}$ s	$(1/2)$	$\alpha$	100		2024	[29]
$^{270}\text{Hs}$	$7.6_{-2.2}^{+4.9}$ s	$0^+$	$\alpha, SF$	$>50 / <50$	9070(40)	2006	[564, 499, 583, 228, 229, 584] [561, 204, 566, 567]
$^{271}\text{Hs}^m$	$7.1_{-2.5}^{+8.4}$ s	$(11/2)$	$\alpha$		$9480(20)\#$	2008	[229, 584, 561, 15, 29]
$^{271m}\text{Hs}^m$	$46_{-16}^{+56}$ s	$(3/2)$	$\alpha$			2024	[29]
$^{272}\text{Hs}^o$	$0.16_{-0.06}^{+0.19}$ s		$\alpha$	100	9772(21)	2023	[28]
$^{273}\text{Hs}$	$0.51_{-0.13}^{+0.30}$ s	$r$	$\alpha$	$\approx 100$	9650(60)	2010	[201, 200, 199]
$^{275}\text{Hs}^n$	$0.15_{-0.06}^{+0.27}$ s		$\alpha, SF$	$>89 / <11$	9450(50)	2004	[205, 206]
$^{277}\text{Hs}$	$3_{-1}^{+15}$ ms		$SF$	100	$9030(200)\#$	2010	[585, 334, 586, 176, 240]
$^{277m}\text{Hs}$	$0.6_{-0.3}^{+2.7}$ ms		$SF$	100		2012	[586]
 $^{266}\text{Mt}$	2.1(7) ms		$\alpha$	$\$ \leq 100$	10996(25)	1982	[587, 588, 569, 589, 570, 444] [348, 178]
$^{268}\text{Mt}$	$21_{-5}^{+8}$ ms		$\alpha$	100	$10770(150)\#$	1995	[519, 230, 483, 571, 424]
$^{270}\text{Mt}$	$0.48_{-0.18}^{+0.66}$ s		$\alpha$	100	10181(50)	2004	[214, 215, 590, 216, 217, 559]
$^{274}\text{Mt}^j$	$0.64_{-0.23}^{+0.76}$ s		$\alpha, SF$	$\approx 100 / ?$	$10600(230)\#$	2007	[202, 203, 474, 576, 184, 223] [12, 182]
$^{275}\text{Mt}^j$	$20_{-6}^{+13}$ ms		$\alpha$	100	10480(50)	2004	[537, 539, 203, 474, 576, 184] [223, 12, 182]

$^{276}\text{Mt}^j$	$0.62_{-0.04}^{+0.06}$ s	$\alpha$	100	10100(10)	2004	[183, 538, 539, 576, 541, 184] [185, 577, 186, 223, 247, 248] [578, 12, 147, 182]
$^{277}\text{Mt}$	$3.9_{-1.3}^{+3.6}$ ms	$SF$	100	9900(100)≠	2013	[188, 223, 12, 176]
$^{278}\text{Mt}$	$4.5_{-1.3}^{+3.5}$ s	$\alpha, SF$	100/?	9580(30)	2010	[25, 576, 204, 579, 175, 223] [12, 176]
$^{267}\text{Ds}$	$2.8_{-0.1}^{+13.3}$ $\mu\text{s}$ ?	$\alpha$ ?	100 (?)	11780(50)	1995	[3, 4]
$^{269}\text{Ds}$	$170_{-60}^{+160}$ $\mu\text{s}^s$	$\alpha$	100	11510(30)	1995	[549, 230]
$^{270}\text{Ds}$	$0.20_{-0.04}^{+0.07}$ ms	$0^+$	$\alpha, SF$	100/<20	2001	[49, 230, 553, 554, 50, 591] [581, 1]
$^{270m}\text{Ds}$	$3.9_{-0.8}^{+1.3}$ ms	(9-,10-)	$\alpha, IT$	>70/<30	≈1130	2001 [49, 230, 553, 554, 50, 591] [581, 1]
$^{271}\text{Ds}$	$1.63_{-0.02}^{+0.44}$ ms		$\alpha$	100	10870(18)	1995 [519, 473, 550, 551, 4, 230] [483, 425, 424, 520]
$^{271m}\text{Ds}$	$69_{-21}^{+56}$ ms		$\alpha, IT$	>0/?		2004 [425]
$^{273}\text{Ds}^m$	$30_{-15}^{+140}$ ms	(11/2)	$\alpha$	100	11090(20)	1996 [562, 494, 551, 592, 29]
$^{273m}\text{Ds}^m$	$0.18_{-0.05}^{+0.11}$ ms	(1/2)	$\alpha$	100		2024 [29]
$^{275}\text{Ds}^m$	$0.43_{-0.12}^{+0.29}$ ms		$\alpha$	100	11370(20)	1996 [29]
$^{276}\text{Ds}^o$	$0.15_{-0.04}^{+0.10}$ s		$\alpha, SF$	43/57	10904(28)	2023 [28]
$^{277}\text{Ds}$	$3.5_{-0.9}^{+2.1}$ ms		$\alpha$	100	10710(40) <sup>r</sup>	2010 [201, 200]
$^{279}\text{Ds}^n$	$186_{-17}^{+21}$ ms		$\alpha, SF$	≈13/87	9827(15)≠	2004 [183, 205, 593, 206]
$^{280}\text{Ds}$	$0.36_{-0.16}^{+1.72}$ s		$SF$	100	9710(200)≠	2021 [33]
$^{281}\text{Ds}$	$12_{-2}^{+4}$ s		$\alpha, SF$	6/94	9470(210)≠	2004 [183, 205, 240]
$^{272}\text{Rg}$	$3.8_{-0.8}^{+1.4}$ ms		$\alpha$	100	11197(13)	1995 [549, 473, 550, 551, 444, 483] [520, 571]
$^{274}\text{Rg}$	$12_{-5}^{+17}$ ms		$\alpha$	100	11480(90)	2004 [214, 215, 590, 594, 217, 559]
$^{278}\text{Rg}^i$	$4.6_{-1.6}^{+5.5}$ ms		$\alpha, SF$	≈100/?	10850(100)	2007 [202, 203, 474, 204, 541, 184] [223, 12, 182]
$^{279}\text{Rg}^j$	$90_{-25}^{+60}$ ms		$\alpha, SF$	87/13	10530(170)	2004 [536, 183, 185, 248, 181, 182]
$^{280}\text{Rg}^j$	$3.9(3)$ s		$\alpha$	100	10149(10)	2004 [183, 538, 539, 576, 541, 184] [185, 577, 186, 223, 247, 248] [578, 12, 147, 182]
$^{281}\text{Rg}^j$	$11_{-1}^{+3}$ s		$\alpha, SF$	13/≈100	9900(400)≠	2010 [25, 576, 204, 579, 223, 12] [176, 182]
$^{282}\text{Rg}$	$100_{-30}^{+70}$ s		$\alpha, SF$	≈100/?	9550(100)≠	2010 [25, 576, 204, 579, 188, 175] [223, 12, 176]
$^{277}\text{Cn}$	$0.61_{-0.18}^{+0.46}$ ms		$\alpha$	100	11620(60)	1996 [494, 551, 230, 592, 501, 231] [559, 581]
$^{281}\text{Cn}$	$0.18_{-0.05}^{+0.10}$ s		$\alpha$	100	10430(60)	2010 [201, 200, 199]
$^{282}\text{Cn}^n$	$0.83_{-0.13}^{+0.18}$ ms		$SF$	100	10150(200)≠	2004 [183, 205, 595, 26, 596, 201] [204, 579, 223, 12, 33, 239] [206, 241]
$^{283}\text{Cn}^n$	$3.81_{-0.36}^{+0.45}$ s		$\alpha, SF$	≥90/leq10	9667(15)	2004 [183, 205, 593, 26, 596, 204] [223, 12, 176] [206]
$^{284}\text{Cn}$	$121_{-15}^{+20}$ ms		$\alpha, SF$	2/198	9670(150)≠	2004 [183, 205, 585, 597, 334, 204] [586, 598, 599, 223, 32, 12] [33]
$^{285}\text{Cn}$	$33_{-6}^{+10}$ s		$\alpha$	100	9390(120)≠	2004 [183, 205, 600, 601, 204, 223] [12, 240]
$^{278}\text{Nh}$	$1.4_{-0.5}^{+1.9}$ ms		$\alpha$	100	11990(80)	2004 [214, 215, 590, 594, 217, 559]
$^{282}\text{Nh}^j$	$61_{-22}^{+73}$ ms		$\alpha$	100	10780(80)	2007 [202, 602, 204, 184, 223, 12] [182]
$^{283}\text{Nh}^j$	$123_{-35}^{+80}$ ms		$\alpha$	100	10420(110)≠	2004 [536, 537, 185, 223, 248, 12] [182]

$^{284}\text{Nh}^j$	$0.90_{-0.06}^{+0.07}$ s	$\alpha$	100	10280(40)	2004	[537, 538, 539, 576, 541, 184] [185, 577, 186, 223, 247, 248] [578, 12, 147, 182]
$^{285}\text{Nh}^j$	$2.1_{-0.3}^{+0.6}$ s	$\alpha, SF$	100/?	10010(40)	2010	[25, 576, 204, 579, 223, 12] [176, 182]
$^{286}\text{Nh}$	$9.5_{-2.7}^{+6.3}$ s	$\alpha, SF$	$\approx 100/?$	9790(50)	2010	[25, 576, 204, 579, 188, 175] [223, 12, 176]
$^{284}\text{Fl}$	$2.5_{-0.8}^{+1.8}$ ms	$\alpha, SF$	<25/100	10700(300)≠	2015	[200, 12, 199]
$^{285}\text{Fl}$	$0.10_{-0.03}^{+0.06}$ s	$\alpha$	100	10560(70)	2010	[201, 200, 12, 199]
$^{286}\text{Fl}^n$	$105_{-13}^{+17}$ s	$\alpha, SF$	40/60	1036035(10)	2004	[183, 205, 595, 26, 596, 201] [204, 579, 223, 12, 33, 239] [206, 241]
$^{287}\text{Fl}^n$	$360_{-36}^{+45}$ ms	$\alpha$	>70	10157(15)	2004	[183, 205, 26, 596, 204, 223] [12, 206]
$^{288}\text{Fl}$	$0.64_{-0.10}^{+0.14}$ s	$\alpha$	100	10076(12)	2004	[183, 205, 595, 585, 597, 334] [204, 586, 598, 599, 223, 32] [12, 33, 239]
$^{289}\text{Fl}$	$2.4_{-0.5}^{+0.8}$ s	$\alpha$	100	9950(70)	2004	[183, 205, 600, 601, 204, 223] [12, 240]
$^{286}\text{Mc}^j$	$20_{-9}^{+98}$ ms	$\alpha$	100	10860(20)	2022	[182]
$^{287}\text{Mc}^j$	$38_{-10}^{+22}$ ms	$\alpha$	100	10760(70)	2004	[536, 537, 185, 223, 248, 603] [12, 181, 182]
$^{288}\text{Mc}^j$	$193_{-13}^{+15}$ ms	$\alpha$	100	10280(40)	2004	[537, 538, 539, 604, 576, 541] [184, 185, 577, 186, 223, 247] [248, 12, 578, 12, 147, 181] [182]
$^{289}\text{Mc}^j$	$250_{-35}^{+51}$ ms	$\alpha, SF$	100/?	10490(50)	2010	[25, 576, 204, 579, 223, 12] [223, 12, 176, 182]
$^{290}\text{Mc}$	$0.65_{-0.20}^{+0.49}$ ms	$\alpha$	100	10410(40)	2010	[25, 576, 204, 579, 188, 175] [176]
$^{290}\text{Lv}$	$8.3_{-1.9}^{+3.5}$ ms	$\alpha$	100	11000(60)	2004	[183, 205, 595, 26, 204, 579] [223, 12, 241]
$^{291}\text{Lv}$	$19_{-6}^{+17}$ ms	$\alpha$	100	10890(90)	2004	[183, 205, 26, 204, 223, 12]
$^{292}\text{Lv}$	$12.8_{-3.3}^{+7.4}$ ms	$\alpha$	100	10791(12)	2004	[183, 205, 595, 204, 586, 598] [223, 32, 12]
$^{293}\text{Lv}$	$95_{-27}^{+63}$ ms	$\alpha$	100	10680(60)	2004	[183, 205, 595, 204, 586, 598] [223, 32, 12]
$^{293}\text{Ts}$	$21_{-4}^{+7}$ ms	$\alpha$	100	11320(50)	2010	[25, 576, 204, 579, 188, 223] [603, 12, 176]
$^{294}\text{Ts}$	$51_{-16}^{+38}$ ms	$\alpha$	100	11180(40)	2010	[25, 576, 204, 579, 188, 223] [603, 12, 176]
$^{294}\text{Og}$	$0.58_{-0.18}^{+0.44}$ ms	$\alpha$	100	11870(30)	2006	[26, 204, 203, 474, 204, 579] [223, 12, 243]

<sup>a</sup> values for  $T_{1/2}$  and branching are taken from [35] (see subsection 5.3 and discussion in Ref. [36])

<sup>b</sup> nomenclature taken from Ref. [51]. The g.s.  $Q_\alpha$  values are given while isomer level energies are not known.

<sup>c</sup> value taken from Ref. [52]

<sup>d</sup> values are taken from [162] while a conservative error for  $Q_\alpha$  which was not given, was deduced from Fig. 1 in [162]

<sup>e</sup> first discovery report [306] not confirmed and disproved by new results [391, 30, 87]

<sup>f</sup> earlier assignment to  $^{249}\text{No}$  [30, 393, 394] (see subsection 5.4 and discussion in Ref. [87])

<sup>g</sup> values are taken from Ref. [100]

<sup>h</sup> values are taken from Ref. [134]

<sup>i</sup> values are taken from Ref. [136]

<sup>j</sup> values are taken from Ref. [182]

<sup>k</sup> values are taken from Ref. [103], confirming the original tentative assignment of an  $11_{-3}^{+6}$  ms activity in Ref. [404] (see subsection 5.6)

<sup>l</sup> values are taken from Ref. [192]

<sup>m</sup> values are taken from Ref. [29]

<sup>n</sup> values are taken from Ref. [206]

<sup>o</sup> values are taken from Ref. [28]

<sup>p</sup> for  $\alpha$ -SF competition – see discussion in section 5.8

<sup>q</sup>  $Q_\alpha$  value deduced from Ref. [519] which is consistent with the theoretical value of 9860(150) given in Ref. [117]

<sup>r</sup> value as reported in Ref. [200]

<sup>s</sup> value as reported in Ref. [47]

## 6. Outlook

The long history of the search for the heaviest nuclear species, almost coinciding with the birth of nuclear physics through the discovery of radioactivity by Henri Becquerel in 1896, has provided us with many fascinating discoveries and exciting insights into fundamental properties of nuclear matter. Throughout these last almost 130 years, not only 32 new elements have been found, seven discovered in the gap between bismuth and uranium and 26 newly synthesized from neptunium to oganesson, but more importantly many fundamental properties of nuclear matter revealed in the study of this exotic species, advancing science as a whole. Here the decay processes,  $\alpha$ -,  $\beta$ -, *gamma*-, *IC*-decay, and *SF*, first exciting new features by themselves have developed into powerful tools to understand basic properties like nuclear excitation, deformation and shapes, isomerism and the stability of nuclear matter itself. The first nuclear isomer has been found in a heavy nucleus,  $^{234}\text{Pa}$ , by chemists which explains the notion borrowed from chemistry [1]. The findings in (heavy) nuclei lead to many applications in medicine and applied science science.

SHE and SHN research, throughout the years and decades, always reached a point when progress seemed to come to a hold. A step in technological development and advancement of the experimental tool, introducing accelerators, detection techniques and efficient separation methods, often lead then to a quantum leap in the progress of our understanding of the heaviest nuclei, but not only.

At present, with the advance of acceleration technology, promising and partly delivering already the capability of an order of magnitude higher beam intensity for the low cross-section processes governing SHE/SHN research. New separators developed further for efficient separation, mass identification and handling of those high beam intensities, including target technology, are starting operation now. The advancement of detection set-ups with digital signal processing and the introduction of new methods, borrowed from neighboring fields, like the classic connection to nuclear chemistry, but also the recent introduction of meth-

ods coming from atomic physics like precise mass measurements in Penning traps and MR-ToF mass spectrometers, or laser spectroscopy, have the potential to push fundamental SHN/SHE research to new horizons. Atomic and nuclear properties like binding energies, ionization potentials and atomic excitation levels, as well as the single-particle and collective nuclear excitations will foster the progress towards the eventual localization for the next proton and neutron shell closures, with the potential to live our dream of revealing the final understanding of the strong interaction, with a direct and detailed insight into the foundation of fundamental physics.

New installations are partly already available at accelerator facilities worldwide. The quest for the discovery of new elements following traditional pathways is in the focus of some SHE research teams like the group working at the e.g. at the gas-filled separators DGFRS [42] of FLNR JINR in Dubna, Russia, or GARIS at RIKEN in Wako, Japan [43]. The introduction of novel techniques like ion traps for high-precision mass measurements and laser spectroscopy like the velocity filter SHIP of GSI/FAIR, Darmstadt, Germany, are pursued by others.

The basic nuclear structure features are in the focus pursuing research programs focussed on nuclear spectroscopy, in-beam and, in particular, decay spectroscopy after separation (DSAS) at installations like the mass spectrometer FMA [605] and the gas-filled separator AGFA [46] at ANL in Lemont, IL, U.S.A., RITU [41] at the cyclotron laboratory of the University of Jyväskylä in Finland, the gas-filled separator BGS equipped with the FIONA mass separator at LBNL, Berkeley, CA, U.S.A. [44], the gas-filled separator TASCA at GSI/FAIR [45] and the velocity filter SHELS at FLNR JINR, Dubna, Russia [40].

For the major working horses, the accelerators, the next-generation high-intensity stable beam facilities are already or are coming online soon, like the SHE-factory of FLNR JINR with the new gas-filled separator DGFRS2-2 [246] with its early achievements mentioned in this review, the linear accelerator RILAC with its upgraded performance in

terms of beam intensities at RIKEN, Tokyo, Japan, or the HELIAC project of GSI/FAIR which is still in an early stage with the first components being tested. The next facility in line to actively attack the challenge of SHN research is the SPIRAL2 LINAC at GANIL which has started operation of its first phase being limited for the highest intensities to projectile masses  $A \lesssim 40$ . The complete capabilities of this installation will be available with the second phase of the project, the new injector NEWGAIN [251] will then provide the highest intensities for all ions and together with the separator-spectrometer set-up S<sup>3</sup> [250] will be one of the worldwide most competitive facilities for the investigation of SHN. Combined with the comprehensive separation and detection installation S<sup>3</sup>, be equipped with the detection array for Spectroscopy and Identification of Rare Isotopes Using S<sup>3</sup> (SIRIUS) as well as the S<sup>3</sup> Low Energy Branch (S<sup>3</sup> LEB) [606] offering tools for laser spectroscopy, mass measurement and the setup for laser/trap-assisted DSAS SEASON, it will be ready to advance SHE/SHN research to the next level.

In conclusion, building on the experience of more than a century, with the installations starting operation in these days worldwide SHE/SHN research faces possibly the next quantum leap towards a fundamental understanding of nuclear matter and the nature of the strong force.

## Acknowledgments

Over the decades I profited enormously from the vast knowledge and experience which my teachers, supervisors and collaborators shared generously with me. The recent years saw some of them go, leaving us with partly unrecoverable which we will have a hard time to fill. After Sigurd Hofmann had left us in 2022, this year, 2024, saw Gottfried Münzenberg and Peter Armbruster go, all of them supporting me and my scientific development, starting from my early days as a diploma student in Darmstadt. To Christophe Theisen I owe a lot and I mentioned it already in the dedication at the beginning of this review. There are numerous scientists, students, engineers, technicians and administrators without whom our work and our achievements, and in particular mine, would not be possible. For this review, I would like to express my deepest gratitude to three colleagues who shaped my understanding of our common research in many discussions and collaborative efforts, leading often to acknowledged publications. Fritz Heßberger, my mentor from my first insecure steps on, and throughout my whole career, Stanislav Antalic, my collaborator in DSAS projects and together with Fitz co-author of our last review on SHN isomers, and Christelle Stodel, my local colleague and daily advisor at GANIL, with her patience and friendly understanding, and her help with some of the graphics in this review. Finally, I would like to thank my wife, Elena Litvinova for her patient support, understanding and advice when it comes to the real, i.e. the theory questions.

## References

- [1] D. Ackermann, C. Theisen, Nuclear structure features of very heavy and superheavy nuclei—tracing quantum mechanics towards the ‘island of stability’, *Physica Scripta* 92 (8) (2017) 083002. [doi:10.1088/1402-4896/aa7921](https://doi.org/10.1088/1402-4896/aa7921). URL <https://dx.doi.org/10.1088/1402-4896/aa7921>
- [2] Soti, Zsolt, Dreher, Raymond, Magill, Joseph, Karl-sruhe nuclide chart – a tradition in progress for nuclear data, *EPJ Nuclear Sci. Technol.* 10 (2024) 7. [doi:10.1051/epjn/2024009](https://doi.org/10.1051/epjn/2024009) URL <https://doi.org/10.1051/epjn/2024009>
- [3] A. Ghiorso, D. Lee, L. P. Somerville, W. Loveland, J. M. Nitschke, W. Ghiorso, G. T. Seaborg, P. Wilmarth, R. Leres, A. Wydler, M. Nurmia, K. Gregorich, K. Czerwinski, R. Gaylord, T. Hamilton, N. J. Hannink, D. C. Hoffman, C. Jarzynski, C. Kacher, B. Kadkhodayan, S. Kreek, M. Lane, A. Lyon, M. A. McMahan, M. Neu, T. Sikkeland, W. J. Swiatecki, A. Türler, J. T. Walton, S. Yashita, Evidence for the possible synthesis of element 110 produced by the  $^{59}\text{Co} + ^{209}\text{Bi}$  reaction, *Phys. Rev. C* 51 (1995) R2293–R2297. [doi:10.1103/PhysRevC.51.R2293](https://doi.org/10.1103/PhysRevC.51.R2293). URL <https://link.aps.org/doi/10.1103/PhysRevC.51.R2293>
- [4] P. J. Karol, H. Nakahara, B. W. Petley, E. Vogt, On the discovery of the elements 110-112 (iupac technical report):, *Pure and Applied Chemistry* 73 (6) (2001) 959–967. [doi:doi:10.1351/pac200173060959](https://doi.org/10.1351/pac200173060959). URL <https://doi.org/10.1351/pac200173060959>
- [5] M. Goeppert Mayer, On Closed Shells in Nuclei. II, *Phys. Rev.* 75 (1949) 1969–1970. [doi:10.1103/PhysRev.75.1969](https://doi.org/10.1103/PhysRev.75.1969). URL <https://link.aps.org/doi/10.1103/PhysRev.75.1969>
- [6] M. Goeppert Mayer, Nuclear configurations in the spin-orbit coupling model. i. empirical evidence, *Phys. Rev.* 78 (1950) 16–21. [doi:10.1103/PhysRev.78.16](https://doi.org/10.1103/PhysRev.78.16). URL <http://link.aps.org/doi/10.1103/PhysRev.78.16>
- [7] O. Haxel, J. H. D. Jensen, H. E. Suess, On the "magic numbers" in nuclear structure, *Phys. Rev.* 75 (1949) 1766–1766. [doi:10.1103/PhysRev.75.1766.2](https://doi.org/10.1103/PhysRev.75.1766.2). URL <http://link.aps.org/doi/10.1103/PhysRev.75.1766.2>
- [8] H. Meldner, Predictions of new magic regions and masses for super-heavy nuclei from calculations with realistic shell model single particle hamiltonians, *Ark. Fys.* 36 (0) (1967) 593.
- [9] A. Sobiczewski, F. A. Gareev, B. N. Kalinkin, Closed shells for  $Z > 82$  and  $N > 126$  in a diffuse potential well, *Phys. Lett.* 22 (4) (1966) 500–502. URL <https://www.sciencedirect.com/science/article/abs/pii/0031916366912431>
- [10] V. Strutinsky, Microscopic calculations of nucleon shell effects in the deformation energy of nuclei, in: Nucleides far off the stability line, Proceedings of the Lysekil Symposium, 1966, Vol. 36, 1966, p. 629.
- [11] V. Strutinsky, Shell effects in nuclear masses and deformation energies, *Nuclear Physics A* 95 (2) (1967) 420–442. [doi:10.1016/0375-9474\(67\)90510-6](https://doi.org/10.1016/0375-9474(67)90510-6). URL [https://dx.doi.org/10.1016/0375-9474\(67\)90510-6](https://dx.doi.org/10.1016/0375-9474(67)90510-6)
- [12] Y. T. Oganessian, A. Sobiczewski, G. M. Ter-Akopian, Superheavy nuclei: from predictions to discovery, *Physica Scripta* 92 (2) (2017) 023003. [doi:10.1088/1402-4896/aa53c1](https://doi.org/10.1088/1402-4896/aa53c1). URL <https://dx.doi.org/10.1088/1402-4896/aa53c1>
- [13] S. Hofmann, S. Heinz, R. Mann, J. Maurer, G. Münenberg, S. Antalic, W. Barth, H. G. Burkhard, L. Dahl, K. Eberhardt, R. Grzywacz,

- J. H. Hamilton, R. A. Henderson, J. M. Kenneally, B. Kindler, I. Kojouharov, R. Lang, B. Lommel, K. Miernik, D. Miller, K. J. Moody, K. Morita, K. Nishio, A. G. Popeko, J. B. Roberto, J. Runke, K. P. Rykaczewski, S. Saro, C. Scheidenberger, H. J. Schött, D. A. Shaughnessy, M. A. Stoyer, P. Thörle-Pospiech, K. Tinschert, N. Trautmann, J. Uusitalo, A. V. Yeremin, [Review of even element super-heavy nuclei and search for element 120](#), The European Physical Journal A 52 (6) (2016) 1–34. doi:[10.1140/epja/i2016-16180-4](https://doi.org/10.1140/epja/i2016-16180-4).  
URL <http://dx.doi.org/10.1140/epja/i2016-16180-4>
- [14] F. P. Heßberger, D. Ackermann, [Some critical remarks on a sequence of events interpreted to possibly originate from a decay chain of an element 120 isotope](#), The European Physical Journal A 53 (6) (2017) 123. doi:[10.1140/epja/i2017-12307-5](https://doi.org/10.1140/epja/i2017-12307-5).  
URL <https://doi.org/10.1140/epja/i2017-12307-5>
- [15] A. Türler, V. Pershina, Advances in the production and chemistry of the heaviest elements, Chem. Rev. 13 (2013) 1237.
- [16] O. R. Smits, C. E. Düllmann, P. Indelicato, W. Nazarewicz, P. Schwerdtfeger, [The quest for superheavy elements and the limit of the periodic table](#), Nature Reviews Physics 6 (2) (2024) 86–98. doi:[10.1038/s42254-023-00668-y](https://doi.org/10.1038/s42254-023-00668-y).  
URL <https://doi.org/10.1038/s42254-023-00668-y>
- [17] M. Block, M. Laatiaoui, S. Raeder, [Recent progress in laser spectroscopy of the actinides](#), Progress in Particle and Nuclear Physics 116 (2021) 103834. doi:<https://doi.org/10.1016/j.ppnp.2020.103834>.  
URL <https://www.sciencedirect.com/science/article/pii/S0146641020300818>
- [18] P. Campbell, I. Moore, M. Pearson, [Laser spectroscopy for nuclear structure physics](#), Progress in Particle and Nuclear Physics 86 (2016) 127–180. doi:<https://doi.org/10.1016/j.ppnp.2015.09.003>.  
URL <https://www.sciencedirect.com/science/article/pii/S0146641015000915>
- [19] M. Block, Precise ground state properties of the heaviest elements for studies of their atomic and nuclear structure, Radiochimica Acta 107 (7) (2019) 603–613. doi:[doi:10.1515/ract-2019-0002](https://doi.org/10.1515/ract-2019-0002).  
URL <https://doi.org/10.1515/ract-2019-0002>
- [20] M. Block, Direct mass measurements and ionization potential measurements of the actinides:, Radiochim. Acta 107 (9-11) (2019) 821–831. doi:[doi:10.1515/ract-2019-3143](https://doi.org/10.1515/ract-2019-3143).  
URL <https://doi.org/10.1515/ract-2019-3143>
- [21] J. Warbinek, E. Rickert, S. Raeder, T. Albrecht-Schönzart, B. Andelic, J. Auler, B. Bally, M. Bender, S. Berndt, M. Block, A. Brizard, P. Chauveau, B. Cheal, P. Chhetri, A. Claessens, A. de Roubin, C. Devlin, H. Dorrer, C. E. Düllmann, J. Ezold, R. Ferrer, V. Gadelshin, A. Gaiser, F. Giacoppo, S. Goriely, M. J. Gutiérrez, A. Harvey, R. Hasse, R. Heinke, F.-P. Heßberger, S. Hilaire, M. Kaja, O. Kaleja, T. Kieck, E. Kim, N. Kneip, U. Köster, S. Kraemer, M. Laatiaoui, J. Lantis, N. Lecesne, A. T. Loria Basto, A. K. Mistry, C. Mokry, I. Moore, T. Murböck, D. Münzberg, W. Nazarewicz, T. Niemeyer, S. Nothelfer, S. Péru, A. Raggio, P.-G. Reinhard, D. Renisch, E. Rey-Herme, J. Romans, E. Romero Romero, J. Runke, W. Ryssens, H. Savajols, F. Schneider, J. Sperling, M. Stemmler, D. Studer, P. Thörle-Pospiech, N. Trautmann, M. Urquiza-González, K. van Beek, S. Van Cleve, P. Van Duppen, M. Vandebrouck, E. Verstraeten, T. Walther, F. Weber, K. Wendt, [Smooth trends in fermium charge radii and the impact of shell effects](#), Nature 634 (8036) (2024) 1075–1079. doi:[10.1038/s41586-024-08062-z](https://doi.org/10.1038/s41586-024-08062-z).

- URL <https://doi.org/10.1038/s41586-024-08062-z>
- [22] C. E. Düllmann, R.-D. Herzberg, W. Nazarewicz, Y. Oganessian (Eds.), *Special Issue on Superheavy Elements*, Vol. 944, Elsevier, B.V., 2015.
- URL <https://www.sciencedirect.com/journal/nuclear-physics-a/vol/944/suppl/C>
- [23] M. Curie, P. Curie, Sur une substance nouvelle radioactive contenue dans la pechblende, Comptes Rendus de l'Académie des Sciences 127 (1898) 175–178.
- [24] P. Curie, M. Curie-Skłodowska, G. B'ement, Sur une nouvelle substance fortement radioactive contenue dans la pechblende, Comptes Rendus de l'Académie des Sciences 127 (1898) 1215.
- [25] Y. T. Oganessian, F. S. Abdullin, P. D. Bailey, D. E. Benker, M. E. Bennett, S. N. Dmitriev, J. G. Ezold, J. H. Hamilton, R. A. Henderson, M. G. Itkis, Y. V. Lobanov, A. N. Mezentsev, K. J. Moody, S. L. Nelson, A. N. Polyakov, C. E. Porter, A. V. Ramayya, F. D. Riley, J. B. Roberto, M. A. Ryabinin, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, V. G. Subbotin, R. Sudowe, A. M. Sukhov, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, P. A. Wilk, *Synthesis of a new element with atomic number  $z = 117$* , Phys. Rev. Lett. 104 (2010) 142502. doi:10.1103/PhysRevLett.104.142502.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.104.142502>
- [26] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, R. N. Sagaidak, I. V. Shirokovsky, Y. S. Tsyanov, A. A. Voinov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, K. Subotic, V. I. Zagrebaev, G. K. Vostokin, M. G. Itkis, K. J. Moody, J. B. Patin, D. A. Shaughnessy, M. A. Stoyer, N. J. Stoyer, P. A. Wilk, J. M. Kenneally, J. H. Landrum, J. F. Wild, R. W. Lougheed, *Synthesis of the isotopes of elements 118 and 116 in the  $^{249}\text{Cf}$  and  $^{245}\text{Cm} + ^{48}\text{Ca}$  fusion reactions*, Phys. Rev. C 74 (2006) 044602. doi:10.1103/PhysRevC.74.044602.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.74.044602>
- [27] G. G. Gulbekian, S. N. Dmitriev, M. G. Itkis, Y. T. Oganessian, B. N. Gikal, I. V. Kalagin, V. A. Semin, S. L. Bogomolov, V. A. Buzmakov, I. A. Ivanenko, N. Y. Kazarinov, N. F. Osipov, S. V. Pashenko, V. A. Sokolov, N. N. Pchelkin, S. V. Prokhorov, M. V. Khabarov, K. B. Gikal, *Start-up of the dc-280 cyclotron, the basic facility of the factory of superheavy elements of the laboratory of nuclear reactions at the joint institute for nuclear research*, Physics of Particles and Nuclei Letters 16 (6) (2019) 866–875. doi:10.1134/S1547477119060177.
- URL <https://doi.org/10.1134/S1547477119060177>
- [28] Y. T. Oganessian, V. K. Utyonkov, M. V. Shumeiko, F. S. Abdullin, S. N. Dmitriev, D. Ibadullayev, M. G. Itkis, N. D. Kovrzhnykh, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibakin, A. N. Polyakov, A. G. Popeko, I. S. Rogov, R. N. Sagaidak, L. Schlattauer, V. D. Shubin, D. I. Solovyev, Y. S. Tsyanov, A. A. Voinov, V. G. Subbotin, N. S. Bublikova, M. G. Voronyuk, A. V. Sabelnikov, A. Y. Bodrov, Z. G. Gan, Z. Y. Zhang, M. H. Huang, H. B. Yang, *New isotope  $^{276}\text{Ds}$  and its decay products  $^{272}\text{Hs}$  and  $^{268}\text{Sg}$  from the  $^{232}\text{Th} + ^{48}\text{Ca}$  reaction*, Phys. Rev. C 108 (2023) 024611. doi:10.1103/PhysRevC.108.024611.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.108.024611>
- [29] Y. T. Oganessian, V. K. Utyonkov, M. V. Shumeiko, F. S. Abdullin, G. G. Adamian, S. N. Dmitriev, D. Ibadullayev, M. G. Itkis, N. D. Kovrzhnykh, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibakin, A. N. Polyakov, A. G. Popeko, I. S. Rogov,

- R. N. Sagaidak, L. Schlattauer, V. D. Shubin, D. I. Solovyev, Y. S. Tsyganov, A. A. Voinov, V. G. Subbotin, N. S. Bublikova, M. G. Voronyuk, A. V. Sabelnikov, A. Y. Bodrov, N. V. Aksenov, A. V. Khalkin, Z. G. Gan, Z. Y. Zhang, M. H. Huang, H. B. Yang, [Synthesis and decay properties of isotopes of element 110:  \$^{273}\text{Ds}\$  and  \$^{275}\text{Ds}\$](#) , Phys. Rev. C 109 (2024) 054307. doi:[10.1103/PhysRevC.109.054307](https://doi.org/10.1103/PhysRevC.109.054307). URL <https://link.aps.org/doi/10.1103/PhysRevC.109.054307>
- [30] A. V. Belozerov, M. L. Chelnokov, V. I. Chepigin, T. P. Drobina, V. A. Gorshkov, A. P. Kabachenko, O. N. Malyshev, I. M. Merkin, Y. T. Oganessian, A. G. Popeko, R. N. Sagaidak, A. I. Svirikhin, A. V. Yeremin, G. Berek, I. Brida, Š. Šáro, [Spontaneous-fission decay properties and production cross-sections for the neutron-deficient nobelium isotopes formed in the 44, 48ca + 204, 206, 208pb reactions](#), The European Physical Journal A - Hadrons and Nuclei 16 (4) (2003) 447–456. doi:[10.1140/epja/i2002-10109-6](https://doi.org/10.1140/epja/i2002-10109-6). URL <https://doi.org/10.1140/epja/i2002-10109-6>
- [31] J. Kallunkathariyil, B. Sulignano, P. T. Greenlees, J. Khuyagbaatar, C. Theisen, K. Auranen, H. Badran, F. Bisso, P. Brionnet, R. Briselet, A. Drouart, Z. Favier, T. Goigoux, T. Grahn, K. Hauschild, A. Herzan, F. P. Heßberger, U. Jakobsson, R. Julin, S. Juutinen, J. Konki, M. Leino, A. Lightfoot, J. Pakarinen, P. Papadakis, J. Partanen, P. Peura, P. Rahkila, K. Rezynkina, P. Ruotsalainen, M. Sandzelius, J. Saren, C. Scholey, M. Siciliano, J. Sorri, S. Stolze, A. I. Svirikhin, J. Uusitalo, M. Vandebrouck, A. Ward, C. Wraith, M. Zielińska, [Stability of the heaviest elements:  \$k\$  isomer in  \$^{250}\text{No}\$](#) , Phys. Rev. C 101 (2020) 011301. doi:[10.1103/PhysRevC.101.011301](https://doi.org/10.1103/PhysRevC.101.011301). URL <https://link.aps.org/doi/10.1103/PhysRevC.101.011301>
- [32] D. Kaji, K. Morita, K. Morimoto, H. Haba, M. Asai, K. Fujita, Z. Gan, H. Geissel, H. Hasebe, S. Hofmann, M. Huang, Y. Komori, L. Ma, J. Maurer, M. Murakami, M. Takeyama, F. Tokanai, T. Tanaka, Y. Wakabayashi, T. Yamaguchi, S. Yamaki, A. Yoshida, [Study of the reaction  \$^{48}\text{Ca} + ^{248}\text{Cm} \rightarrow ^{296}\text{Lv}^\*\$  at riken-garis](#), Journal of the Physical Society of Japan 86 (3) (2017) 034201. arXiv: <https://doi.org/10.7566/JPSJ.86.034201>, doi: [10.7566/JPSJ.86.034201](https://doi.org/10.7566/JPSJ.86.034201). URL <https://doi.org/10.7566/JPSJ.86.034201>
- [33] A. Samark-Roth, D. M. Cox, D. Rudolph, L. G. Sarmiento, B. G. Carlsson, J. L. Egido, P. Golubev, J. Heery, A. Yakushev, S. Aberg, H. M. Albers, M. Albertsson, M. Block, H. Brand, T. Calverley, R. Cantemir, R. M. Clark, C. E. Düllmann, J. Eberth, C. Fahlander, U. Forsberg, J. M. Gates, F. Giacoppo, M. Götz, S. Götz, R.-D. Herzberg, Y. Hrabar, E. Jäger, D. Judson, J. Khuyagbaatar, B. Kindler, I. Kojouharov, J. V. Kratz, J. Krier, N. Kurz, L. Lens, J. Ljungberg, B. Lommel, J. Louko, C.-C. Meyer, A. Mistry, C. Mokry, P. Papadakis, E. Parr, J. L. Pore, I. Ragnarsson, J. Runke, M. Schädel, H. Schaffner, B. Schausten, D. A. Shaughnessy, P. Thörle-Pospiech, N. Trautmann, J. Uusitalo, [Spectroscopy along flerovium decay chains: Discovery of  \$^{280}\text{Ds}\$  and an excited state in  \$^{282}\text{Cn}\$](#) , Phys. Rev. Lett. 126 (2021) 032503. doi:[10.1103/PhysRevLett.126.032503](https://doi.org/10.1103/PhysRevLett.126.032503). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.032503>
- [34] J. L. Pore, J. M. Gates, R. Orford, C. M. Campbell, R. M. Clark, H. L. Crawford, N. E. Esker, P. Fallon, J. A. Gooding, J. T. Kwarsick, A. O. Macchiavelli, C. Morse, D. Rudolph, A. Samark-Roth, C. Santamaria, R. S. Shah, M. A. Stoyer, [Identification of the new isotope  \$^{244}\text{Md}\$](#) , Phys. Rev. Lett. 124 (2020) 252502. doi:[10.1103/PhysRevLett.124.252502](https://doi.org/10.1103/PhysRevLett.124.252502).

- URL <https://link.aps.org/doi/10.1103/PhysRevLett.124.252502>
- [35] J. Khuyagbaatar, H. M. Albers, M. Block, H. Brand, R. A. Cantemir, A. Di Nitto, C. E. Düllmann, M. Götz, S. Götz, F. P. Heßberger, E. Jäger, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, B. Lommel, L. Lens, A. Mistry, B. Schausten, J. Uusitalo, A. Yakushev, **Search for electron-capture delayed fission in the new isotope  $^{244}\text{Md}$** , Phys. Rev. Lett. 125 (2020) 142504. doi:[10.1103/PhysRevLett.125.142504](https://doi.org/10.1103/PhysRevLett.125.142504).
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.142504>
- [36] F. P. Heßberger, M. Block, C. E. Düllmann, A. Yakushev, M. Leino, J. Uusitalo, **Some remarks on the discovery of  $^{244}\text{Md}$** , Phys. Rev. Lett. 126 (2021) 182501. doi:[10.1103/PhysRevLett.126.182501](https://doi.org/10.1103/PhysRevLett.126.182501).
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.182501>
- [37] J. Konki, J. Khuyagbaatar, J. Uusitalo, P. Greenlees, K. Auranen, H. Badran, M. Block, R. Briselet, D. Cox, M. Dasgupta, A. Di Nitto, C. Düllmann, T. Grahn, K. Hauschild, A. Herzán, R.-D. Herzberg, F. Heßberger, D. Hinde, R. Julin, S. Juutinen, E. Jäger, B. Kindler, J. Krier, M. Leino, B. Lommel, A. Lopez-Martens, D. Luong, M. Mallaburn, K. Nishio, J. Pakarinen, P. Papadakis, J. Partanen, P. Peura, P. Rahkila, K. Rezynkina, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, S. Stolze, B. Sulignano, C. Theisen, A. Ward, A. Yakushev, V. Yakusheva, **Towards saturation of the electron-capture delayed fission probability: The new isotopes  $^{240}\text{es}$  and  $^{236}\text{bk}$** , Physics Letters B 764 (2017) 265–270. doi:<https://doi.org/10.1016/j.physletb.2016.11.038>.
- URL <https://www.sciencedirect.com/science/article/pii/S0370269316307018>
- [38] D. Ackermann, S. Antalic, F. P. Heßberger, **Iso-**mers in superheavy nuclei, The European Physical Journal Special Topics 233 (5) (2024) 1017–1036. doi:[10.1140/epjs/s11734-024-01150-1](https://doi.org/10.1140/epjs/s11734-024-01150-1).
- URL <https://doi.org/10.1140/epjs/s11734-024-01150-1>
- [39] G. Münzenberg, W. Faust, S. Hofmann, P. Armbruster, K. Güttner, H. Ewald., **The velocity filter ship and a separator of unslowed heavy ion fusion products.**, Nucl. Instr. and Meth. 161 (1979) 65.
- [40] A. Yeremin, O. Malyshev, A. Popeko, V. Chepigin, A. Svirikhin, A. Lopez-Martens, K. Hauschild, O. Dorvaux, B. Gall, J. Gehlot, **First experimental tests of the kinematic separator shells (separator for heavy element spectroscopy).**, EPJ Web of Conferences 86 (2015) 00065. doi:[10.1051/epjconf/20158600065](https://doi.org/10.1051/epjconf/20158600065).
- URL <http://dx.doi.org/10.1051/epjconf/20158600065>
- [41] M. Leino, J. Äystö, T. Enqvist, P. Heikkinen, A. Jokinen, M. Nurmi, A. Ostrowski, W. Trzaska, J. Uusitalo, K. Eskola, P. Armbruster, V. Ninov, **Gas-filled recoil separator for studies of heavy elements**, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 99 (1) (1995) 653–656, application of Accelerators in Research and Industry '94. doi:[https://doi.org/10.1016/0168-583X\(94\)00573-7](https://doi.org/10.1016/0168-583X(94)00573-7).
- URL <https://www.sciencedirect.com/science/article/pii/0168583X94005737>
- [42] Y. S. Tsyganov, **The dubna gas-filled recoil separator: status and developments**, Journal of Physics G: Nuclear and Particle Physics 25 (4) (1999) 937. doi:[10.1088/0954-3899/25/4/074](https://doi.org/10.1088/0954-3899/25/4/074).
- URL <https://dx.doi.org/10.1088/0954-3899/25/4/074>
- [43] K. Morita, A. Yoshida, T. Inamura, M. Koizumi, T. Nomura, M. Fujioka, T. Shinozuka, H. Miyatake, K. Sueki, H. Kudo, Y. Nagai, T. Toriyama,

- K. Yoshimura, Y. Hatsukawa, Riken isotope separator on-line garis/igisol, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 70 (1) (1992) 220–225. doi:[https://doi.org/10.1016/0168-583X\(92\)95935-K](https://doi.org/10.1016/0168-583X(92)95935-K). URL <https://www.sciencedirect.com/science/article/pii/0168583X9295935K>
- [44] J. M. Gates, J. L. Pore, Studies of heavy and super heavy elements with fiona: the broad impact of mass-number identifications., Eur. Phys. J. A 58 (10) (2022) 1 – 21. URL <https://search-ebscohost-com.in2p3.bib.cnrs.fr/login.aspx?direct=true&db=a9h&AN=160254331&lang=fr&site=ehost-live>
- [45] A. Semchenkov, W. Brückle, E. Jäger, E. Schimpf, M. Schädel, C. Mühle, F. Klos, A. TÄrlirler, A. Yakuhev, A. Belov, T. Belyakova, M. Kaparkova, V. Kukhtin, E. Lamzin, S. Sytchevsky, The transactinide separator and chemistry apparatus TASCA at GSI - optimization of ion-optical structures and magnet designs, Nucl. Instr. and Meth. B 266 (2008) 4153.
- [46] D. Seweryniak, Status and plans for recoil separators for experiments with intense stable beams from ATLAS, Nucl. Inst. Meth. B 317, Part B (2013) 274 – 276, XVIth International Conference on ElectroMagnetic Isotope Separators and Techniques Related to their Applications, December 27, 2012 at Matsue, Japan. doi:<http://dx.doi.org/10.1016/j.nimb.2013.07.027>. URL <https://www.sciencedirect.com/science/article/pii/S0168583X13008215>
- [47] S. Hofmann, G. Münzenberg, The discovery of the heaviest elements, Rev. Mod. Phys. 72 (2000) 733–767. doi:[10.1103/RevModPhys.72.733](https://doi.org/10.1103/RevModPhys.72.733). URL <https://link.aps.org/doi/10.1103/RevModPhys.72.733>
- [48] A. Bronis, F. P. Heßberger, S. Antalic, B. Andel, D. Ackermann, S. Heinz, S. Hofmann, J. Khuyagbaatar, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, A. G. Popeko, B. Streicher, B. Sulignano, J. Uusitalo, M. Venhart, A. V. Yeremin, Decay studies of new isomeric states in  $^{255}\text{No}$ , Phys. Rev. C 106 (2022) 014602. doi:[10.1103/PhysRevC.106.014602](https://doi.org/10.1103/PhysRevC.106.014602). URL <https://link.aps.org/doi/10.1103/PhysRevC.106.014602>
- [49] S. Hofmann, F.P. Heßberger, D. Ackermann, S. Antalic, P. Cagarda, S. Ćwiok, B. Kindler, J. Kojouharova, B. Lommel, R. Mann, G. Münzenberg, A.G. Popeko, S. Saro, H.J. Schött, A.V. Yeremin, The new isotope  $^{270}\text{No}$  and its decay products  $^{266}\text{Hs}$  and  $^{262}\text{Sg}$ , Eur. Phys. J. A 10 (01) (2001) 5–10. doi:[10.1007/s100500170137](https://doi.org/10.1007/s100500170137). URL <https://doi.org/10.1007/s100500170137>
- [50] D. Ackermann, Nuclear spectroscopy in nuclei with  $Z \geq 110$ , Nucl. Phys. A 944 (2015) 376 – 387. doi:<http://dx.doi.org/10.1016/j.nuclphysa.2015.09.002>. URL <https://www.sciencedirect.com/science/article/pii/S0375947415002018>
- [51] S. Antalic, F. P. Heßberger, S. Hofmann, D. Ackermann, S. Heinz, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, Š. Šáro, Studies of neutron-deficient mendelevium isotopes at SHIP, Eur. Phys. J. A 43 (1) (2010) 35–44. doi:[10.1140/epja/i2009-10896-0](https://doi.org/10.1140/epja/i2009-10896-0). URL <http://dx.doi.org/10.1140/epja/i2009-10896-0>
- [52] F. P. Heßberger, S. Antalic, F. Giacoppo, B. Andel, D. Ackermann, M. Block, S. Heinz, J. Khuyagbaatar, I. Kojouharov, M. Venhart, Alpha-gamma decay studies of  $^{247}\text{Md}$ , Eur. Phys. J. A 58 (1) (2022) 11. doi:[10.1140/epja/s10050-022-00663-4](https://doi.org/10.1140/epja/s10050-022-00663-4). URL <https://doi.org/10.1140/epja/s10050-022-00663-4>

- [53] A. Chatillon, C. Theisen, P. T. Greenlees, G. Auger, J. E. Bastin, E. Bouchez, B. Bouriquet, J. M. Casandjian, R. Cee, E. Clément, R. Dayras, G. de France, R. de Tourel, S. Eeckhaudt, A. Görgen, T. Grahn, S. Grévy, K. Hauschild, R. D. Herzberg, P. J. C. Ikin, G. D. Jones, P. Jones, R. Julin, S. Juutinen, H. Kettunen, A. Korichi, W. Korten, Y. Le Coz, M. Leino, A. Lopez-Martens, S. M. Lukyanov, Y. E. Penionzhkevich, J. Perkowski, A. Pritchard, P. Rahkila, M. Rejmund, J. Saren, C. Scholey, S. Siem, M. G. Saint-Laurent, C. Simenel, Y. G. Sobolev, C. Stodel, J. Uusitalo, A. Villari, M. Bender, P. Bonche, P. H. Heenen, **Spectroscopy and single-particle structure of the odd- $z$  heavy elements 255 $\text{lr}$ , 251 $\text{md}$  and 247 $\text{es}$** , The European Physical Journal A - Hadrons and Nuclei 30 (2) (2006) 397–411. [doi:10.1140/epja/i2006-10134-5](https://doi.org/10.1140/epja/i2006-10134-5). URL <https://doi.org/10.1140/epja/i2006-10134-5>
- [54] S. Antalic, F. P. Heßberger, S. Hofmann, D. Ackermann, S. Heinz, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, Š. Šáro, B. Streicher, B. Sulignano, M. Venhart, **Decay studies of neutron-deficient lawrencium isotopes**, The European Physical Journal A 38 (2) (2008) 219–226. [doi:10.1140/epja/i2008-10665-7](https://doi.org/10.1140/epja/i2008-10665-7). URL <https://doi.org/10.1140/epja/i2008-10665-7>
- [55] F. P. Heßberger, S. Antalic, A. K. Mistry, D. Ackermann, B. Andel, M. Block, Z. Kalaninova, B. Kindler, I. Kojouharov, M. Laatiaoui, B. Lommel, J. Piot, M. Vostinar, **Alpha- and ec-decay measurements of  $^{257}\text{rf}$** , Eur. Phys. J. A 52 (7) (2016) 192. [doi:10.1140/epja/i2016-16192-0](https://doi.org/10.1140/epja/i2016-16192-0). URL <https://doi.org/10.1140/epja/i2016-16192-0>
- [56] K. Hauschild, A. Lopez-Martens, R. Chakma, M. L. Chelnokov, V. I. Chepigin, A. V. Isaev, I. N. Izosimov, D. E. Katrasev, A. A. Kuznetsova, O. N. Malyshev, A. G. Popeko, Y. A. Popov, E. A. Sokol, A. I. Svirikhin, M. S. Tezekbayeva, A. V. Yeremin, Z. Asfari, O. Dorvaux, B. J. P. Gall, K. Kessaci, D. Ackermann, J. Piot, P. Mosat, B. Andel, **Alpha-decay spectroscopy of  $^{257}\text{rf}$** , The European Physical Journal A 58 (1) (2022) 6. [doi:10.1140/epja/s10050-021-00657-8](https://doi.org/10.1140/epja/s10050-021-00657-8). URL <https://doi.org/10.1140/epja/s10050-021-00657-8>
- [57] D. Ackermann, **Nuclear isomers in the heaviest nuclei – the odd nucleon as a sensitive probe of low-lying nuclear structure**, Acta Phys. Pol. B Proc. Suppl. 16 (4-A9) (2023). [doi:https://doi.org/10.5506/APhysPolBSupp.16.4-A9](https://doi.org/10.5506/APhysPolBSupp.16.4-A9). URL <https://www.actaphys.uj.edu.pl/S/16/4-A9/pdf>
- [58] S. Antalic, F. P. Heßberger, D. Ackermann, S. Heinz, S. Hofmann, Z. Kalaninová, B. Kindler, J. Khuyagbaatar, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, Š. Šáro, B. Streicher, B. Sulignano, M. Venhart, **Isomeric states in  $^{253}\text{No}$  and  $^{253}\text{Fm}$** , The European Physical Journal A 47 (5) (2011) 62. [doi:10.1140/epja/i2011-11062-y](https://doi.org/10.1140/epja/i2011-11062-y). URL <https://doi.org/10.1140/epja/i2011-11062-y>
- [59] V. T. Jordanov, G. F. Knoll, **Digital synthesis of pulse shapes in real time for high resolution radiation spectroscopy**, Nucl. Instr. and Meth. A 345 (2) (1994) 337–345. [doi:https://doi.org/10.1016/0168-9002\(94\)91011-1](https://doi.org/10.1016/0168-9002(94)91011-1). URL <https://www.sciencedirect.com/science/article/pii/0168900294910111>
- [60] H. M. David, J. Chen, D. Seweryniak, F. G. Kondev, J. M. Gates, K. E. Gregorich, I. Ahmad, M. Albers, M. Alcorta, B. B. Back, B. Baartman, P. F. Bertone, L. A. Bernstein, C. M. Campbell, M. P. Carpenter, C. J. Chiara, R. M. Clark,

- M. Cromaz, D. T. Doherty, G. D. Dracoulis, N. E. Esker, P. Fallon, O. R. Gothe, J. P. Greene, P. T. Greenlees, D. J. Hartley, K. Hauschild, C. R. Hoffman, S. S. Hota, R. V. F. Janssens, T. L. Khoo, J. Konki, J. T. Kwartsick, T. Lauritsen, A. O. Macchiavelli, P. R. Mudder, C. Nair, Y. Qiu, J. Risnanen, A. M. Rogers, P. Ruotsalainen, G. Savard, S. Stolze, A. Wiens, S. Zhu, Decay and fission hindrance of two- and four-quasiparticle  $k$  isomers in  $^{254}\text{Rf}$ , Phys. Rev. Lett. 115 (2015) 132502. doi:[10.1103/PhysRevLett.115.132502](https://doi.org/10.1103/PhysRevLett.115.132502). URL <https://link.aps.org/doi/10.1103/PhysRevLett.115.132502>
- [61] J. Khuyagbaatar, A. Mistry, D. Ackermann, L.-L. Andersson, M. Block, H. Brand, C. Düllmann, J. Even, F. Heßberger, J. Hoffmann, A. Hübiner, E. Jäger, B. Kindler, J. Krier, N. Kurz, B. Lommel, B. Schausten, J. Steiner, A. Yakushev, V. Yakusheva, The identification and confirmation of isomeric states in  $^{254}\text{rf}$  and  $^{255}\text{rf}$  through conversion electron detection, Nuclear Physics A 994 (2020) 121662. doi:<https://doi.org/10.1016/j.nuclphysa.2019.121662>. URL <https://www.sciencedirect.com/science/article/pii/S0375947419302374>
- [62] J. Khuyagbaatar, H. Brand, C. E. Düllmann, F. P. Heßberger, E. Jäger, B. Kindler, J. Krier, N. Kurz, B. Lommel, Y. Nechiporenko, Y. N. Novikov, B. Schausten, A. Yakushev, Search for fission from a long-lived isomer in  $^{250}\text{No}$  and evidence of a second isomer, Phys. Rev. C 106 (2022) 024309. doi:[10.1103/PhysRevC.106.024309](https://doi.org/10.1103/PhysRevC.106.024309). URL <https://link.aps.org/doi/10.1103/PhysRevC.106.024309>
- [63] A. Bohr, B. R. Mottelson, Nuclear Structure Volume II, W.A. Benjamin, Inc., Massachusetts; World Scientific 1998, 1999, 2008, 1975.
- [64] W. Myers, W. Swiatecki, Nuclear masses and deformations, Nucl. Phys. 81 (1966) 1.
- [65] P. Möller, J. Nix, W. Myers, W. Swiatecki, Nuclear ground-state masses and deformations, At. Data and Nucl. Data Tables 59 (1995) 185.
- [66] K. Pomorski, J. Dudek, Nuclear liquid-drop model and surface-curvature effects, Phys. Rev. C 67 (2003) 044316. doi:[10.1103/PhysRevC.67.044316](https://doi.org/10.1103/PhysRevC.67.044316). URL <http://link.aps.org/doi/10.1103/PhysRevC.67.044316>
- [67] K. Pomorski, Fission-barrier heights in some newest liquid-drop models, Physica Scripta 2013 (T154) (2013) 014023. URL <http://stacks.iop.org/1402-4896/2013/i=T154/a=014023>
- [68] M. Asai, F. P. Heßberger, A. Lopez-Martens, Nuclear structure of elements with  $100 \leq Z \leq 109$  from alpha spectroscopy, Nucl. Phys. A 944 (2015) 308 – 332, special Issue on Superheavy Elements. doi:<http://dx.doi.org/10.1016/j.nuclphysa.2015.06.011>. URL <http://www.sciencedirect.com/science/article/pii/S0375947415001402>
- [69] R. R. Chasman, I. Ahmad, A. M. Friedman, J. R. Erskine, Survey of single-particle states in the mass region  $A > 228$ , Rev. Mod. Phys. 49 (1977) 833–891. doi:[10.1103/RevModPhys.49.833](https://doi.org/10.1103/RevModPhys.49.833). URL <http://link.aps.org/doi/10.1103/RevModPhys.49.833>
- [70] K. Kessaci, B. J. P. Gall, O. Dorvaux, M. Forge, A. Lopez-Martens, R. Chakma, K. Hauschild, M. L. Chelnokov, V. I. Chepigin, A. V. Isaev, A. A. Kuznetsova, O. N. Malyshev, R. Mukhin, J. Piot, A. G. Popeko, Y. A. Popov, E. A. Sokol, A. I. Svirikhin, A. V. Yeremin, Cascade of high- $k$  isomers in  $^{255}_{102}\text{No}_{153}$ , Phys. Rev. C 110 (2024) 054310. doi:[10.1103/PhysRevC.110.054310](https://doi.org/10.1103/PhysRevC.110.054310).

- URL <https://link.aps.org/doi/10.1103/PhysRevC.110.054310>
- [71] A. Parkhomenko, A. Sobiczewski, **Proton one quasi-particle states of heaviest nuclei**, Acta Phys. Pol. B 35 (10) (2004) 2447.
- URL <https://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=35&page=2447>
- [72] A. Parkhomenko, A. Sobiczewski, **Phenomenological formula for  $\alpha$ -decay half-lives of heaviest nuclei**, Acta Phys. Pol. B 36 (10) (2005) 3095–3108.
- URL <https://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=36&page=3115>
- [73] R. M. Clark, D. Rudolph,  $\alpha$  decay of high-k isomers in  $^{270}\text{Ds}$  and  $^{266}\text{Hs}$  in a superfluid tunneling model, Phys. Rev. C 97 (2018) 024333. doi:[10.1103/PhysRevC.97.024333](https://doi.org/10.1103/PhysRevC.97.024333).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.97.024333>
- [74] Clark, R. M., **Hindrances to the alpha decay and fission of high-k isomers**, Eur. Phys. J. Spec. Top. 233 (5) (2024) 1007–1015. doi:[10.1140/epjs/s11734-024-01138-x](https://doi.org/10.1140/epjs/s11734-024-01138-x).
- URL <https://doi.org/10.1140/epjs/s11734-024-01138-x>
- [75] T. Goigoux, C. Theisen, B. Sulignano, M. Airiau, K. Auranen, H. Badran, R. Briselet, T. Calverley, D. Cox, F. Déchery, F. D. Bisso, A. Drouart, Z. Favier, B. Gall, T. Grahn, P. T. Greenlees, K. Hauschild, A. Herzáñ, R. D. Herzberg, U. Jakobsson, R. Julin, S. Juutinen, J. Konki, M. Leino, A. Lightfoot, A. Lopez-Martens, A. Mistry, P. Nieminen, J. Pakarinen, P. Papadakis, J. Partanen, P. Peura, P. Rahkila, E. Rey-Herme, J. Rubert, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, S. Stolze, J. Uusitalo, M. Vandebrouck, A. Ward, M. Zielińska, P. Jachimowicz, M. Kowal, J. Skalski, **First observation of high-k isomeric states in  $^{249}\text{Md}$  and  $^{251}\text{Md}$** , Eur. Phys. J. A 57 (12) (2021) 321. doi:[10.1140/epja/s10050-021-00631-4](https://doi.org/10.1140/epja/s10050-021-00631-4).
- URL <https://doi.org/10.1140/epja/s10050-021-00631-4>
- [76] F. P. Heßberger, S. Antalic, D. Ackermann, B. Andel, M. Block, Z. Kalaninova, B. Kindler, I. Kojouharov, M. Laatiaoui, B. Lommel, A. K. Mistry, J. Piot, M. Vostinar, **Investigation of electron capture decay of  $^{258}\text{Db}$  and  $\alpha$  decay of  $^{258}\text{Rf}$** , Eur. Phys. J. A 52 (11) (2016) 328. doi:[10.1140/epja/i2016-16328-2](https://doi.org/10.1140/epja/i2016-16328-2).
- URL <http://dx.doi.org/10.1140/epja/i2016-16328-2>
- [77] R.-D. Herzberg, P. Greenlees, **In-beam and decay spectroscopy of transfermium nuclei**, Progress in Particle and Nuclear Physics 61 (2) (2008) 674–720. doi:<https://doi.org/10.1016/j.ppnp.2008.05.003>.
- URL <https://www.sciencedirect.com/science/article/pii/S0146641008000409>
- [78] R. W. Hoff, T. von Egidy, R. W. Lougheed, D. H. White, H. G. Börner, K. Schreckenbach, G. Barreau, D. D. Warner, **Levels of  $^{244}\text{Cm}$  populated by the beta decay of 10-h  $^{244}\text{Am}^g$  and 26-min  $^{244}\text{Am}^m$** , Phys. Rev. C 29 (1984) 618–622. doi:[10.1103/PhysRevC.29.618](https://doi.org/10.1103/PhysRevC.29.618).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.29.618>
- [79] P. G. Hansen, K. Wilsky, C. V. K. Baba, S. E. Vandenbosch, **Decay of an isomeric state in  $\text{cm}^{244}$** , Nuclear Physics 45 (0) (1963) 410 – 416.
- URL <https://orbit.dtu.dk/en/publications/decay-of-an-isomeric-state-in-cm244>
- [80] L. G. Multhauf, K. G. Tirsell, R. A. Meyer, **Collective excitations in  $^{246}\text{Cm}$  and the decay of  $^{246}\text{Am}^m$** , Phys. Rev. C 13 (1976) 771–789. doi:[10.1103/PhysRevC.13.771](https://doi.org/10.1103/PhysRevC.13.771).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.13.771>
- [81] R. Orlandi, H. Makii, K. Nishio, K. Hirose, M. Asai,

- K. Tsukada, T. K. Sato, Y. Ito, F. Suzuki, Y. Nagame, A. N. Andreyev, E. Ideguchi, N. Aoi, T. T. Pham, S. Q. Yan, Y. P. Shen, B. Gao, G. Li, *New  $k$  isomers in  $^{248}\text{Cf}$* , Phys. Rev. C 106 (2022) 064301. [doi:10.1103/PhysRevC.106.064301](https://doi.org/10.1103/PhysRevC.106.064301).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.106.064301>
- [82] S. Ketelhut, Rotational Structures and High-K Isomerism in  $^{248,250}\text{Fm}$ , Ph.D. thesis, University of Jyväskylä (2010).
- [83] R.-D. Herzberg, D. M. Cox, *Spectroscopy of actinide and transactinide nuclei*, Radiochimica Acta 99 (7-8) (2011) 441–457 [cited 2024-02-21]. [doi:doi:10.1524/ract.2011.1858](https://doi.org/10.1524/ract.2011.1858).  
 URL <https://doi.org/10.1524/ract.2011.1858>
- [84] H. L. Liu, P. M. Walker, F. R. Xu, *Favored configurations for four-quasiparticle  $k$  isomerism in the heaviest nuclei*, Phys. Rev. C 89 (2014) 044304. [doi:10.1103/PhysRevC.89.044304](https://doi.org/10.1103/PhysRevC.89.044304).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.89.044304>
- [85] P. T. Greenlees, R.-D. Herzberg, S. Ketelhut, P. A. Butler, P. Chowdhury, T. Grahn, C. Gray-Jones, G. D. Jones, P. Jones, R. Julin, S. Juutinen, T.-L. Khoo, M. Leino, S. Moon, M. Nyman, J. Pakarinen, P. Rahkila, D. Rostron, J. Sarén, C. Scholey, J. Sorri, S. K. Tandel, J. Uusitalo, M. Venhart, *High- $k$  structure in  $^{250}\text{fm}$  and the deformed shell gaps at  $n = 152$  and  $z = 100$* , Phys. Rev. C 78 (2008) 021303. [doi:10.1103/PhysRevC.78.021303](https://doi.org/10.1103/PhysRevC.78.021303).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.78.021303>
- [86] H. L. Hall, K. E. Gregorich, R. A. Henderson, D. M. Lee, D. C. Hoffman, M. E. Bunker, M. M. Fowler, P. Lysaght, J. W. Starner, J. B. Wilhelmy,  *$\beta$ -delayed fission from  $^{256}\text{Es}^m$  and the level scheme of  $^{256}\text{Fm}$* , Phys. Rev. C 39 (1989) 1866–1875.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.39.1866>
- [87] D. Peterson, B. B. Back, R. V. F. Janssens, T. L. Khoo, C. J. Lister, D. Seweryniak, I. Ahmad, M. P. Carpenter, C. N. Davids, A. A. Hecht, C. L. Jiang, T. Lauritsen, X. Wang, S. Zhu, F. G. Kondev, A. Heinz, J. Qian, R. Winkler, P. Chowdhury, S. K. Tandel, U. S. Tandel, *Decay modes of  $^{250}\text{No}$* , Phys. Rev. C 74 (2006) 014316. [doi:10.1103/PhysRevC.74.014316](https://doi.org/10.1103/PhysRevC.74.014316).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.74.014316>
- [88] F. P. Heßberger, S. Hofmann, D. Ackermann, S. Antalic, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, A. G. Popeko, B. Sulignano, S. Saro, B. Streicher, M. Venhart, A. V. Yeremin, *Alpha-gamma decay studies of  $^{255}\text{rf}$  and  $^{251}\text{no}$  and  $^{247}\text{fm}$* , The European Physical Journal A - Hadrons and Nuclei 30 (3) (2006) 561–569. [doi:10.1140/epja/i2006-10137-2](https://doi.org/10.1140/epja/i2006-10137-2).  
 URL <https://doi.org/10.1140/epja/i2006-10137-2>
- [89] A. Lopez-Martens, K. Hauschild, A. I. Svirikhin, Z. Asfari, M. L. Chelnokov, V. I. Chepigin, O. Dorvaux, M. Forge, B. Gall, A. V. Isaev, I. N. Izosimov, K. Kessaci, A. A. Kuznetsova, O. N. Malyshev, R. S. Mukhin, A. G. Popeko, Y. A. Popov, B. Sailaubekov, E. A. Sokol, M. S. Tezekbayeva, A. V. Yeremin, *Fission properties of  $^{253}\text{Rf}$  and the stability of neutron-deficient rf isotopes*, Phys. Rev. C 105 (2022) L021306. [doi:10.1103/PhysRevC.105.L021306](https://doi.org/10.1103/PhysRevC.105.L021306).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.105.L021306>
- [90] B. Sulignano, S. Heinz, F. P. Heßberger, S. Hofmann, D. Ackermann, S. Antalic, B. Kindler, I. Kojouharov, P. Kuusiniemi, B. Lommel, R. Mann, K. Nishio, A. G. Popeko, S. Saro, B. Streicher, M. Venhart, A. V. Yeremin, *Identification of a  $k$  isomer in  $^{252}\text{no}$* ,

- The European Physical Journal A 33 (4) (2007) 327–331. doi:10.1140/epja/i2007-10469-3.  
 URL <https://doi.org/10.1140/epja/i2007-10469-3>
- [91] B. Sulignano, C. Theisen, J.-P. Delaroche, M. Girod, J. Ljungvall, D. Ackermann, S. Antalic, O. Dorvaux, A. Drouart, B. Gall, A. Görgen, P. T. Greenlees, K. Hauschild, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, W. Korten, M. Leino, A. Lopez-Martens, M. Nyman, A. Obertelli, J. Pakarinen, P. Papadakis, E. Parr, P. Peura, J. Piot, P. Rahkila, D. Rostron, P. Ruotsalainen, J. Sarén, C. Scholey, J. Sorri, J. Uusitalo, M. Venhart, M. Zielińska, Investigation of high- $k$  states in  $^{252}\text{no}$ , Phys. Rev. C 86 (2012) 044318. doi:10.1103/PhysRevC.86.044318.  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.86.044318>
- [92] B. Štreicher, Synthesis and spectroscopic properties of transfermium isotopes with  $z = 105$ , 106 and 107, Ph.D. thesis, Comenius University Bratislava (2006).  
 URL <http://sok11249.dnp.fmph.uniba.sk/sok11249/?cont=prace&id=44>
- [93] A. Lopez-Martens, K. Hauschild, A. V. Yeremin, O. Dorvaux, A. V. Belozerov, C. Briançon, M. L. Chelnokov, V. I. Chepigin, D. Curien, P. Désesquelles, B. Gall, V. A. Gorshkov, M. Guttormsen, F. Hanappe, A. P. Kabachenko, F. Khalfallah, A. Korichi, A. C. Larsen, O. N. Malyshev, A. Minkova, Y. T. Oganessian, A. G. Popeko, M. Rousseau, N. Rowley, R. N. Sagaidak, S. Sharo, A. V. Shutov, S. Siem, L. Stuttgé, A. I. Svirikhin, N. U. H. Syed, C. Theisen, Isomeric states in  $^{253}\text{no}$ , The European Physical Journal A 32 (3) (2007) 245–250. doi:10.1140/epja/i2007-10391-8.  
 URL <https://doi.org/10.1140/epja/i2007-10391-8>
- [94] B. Streicher, F. P. Heßberger, S. Antalic, S. Hofmann, D. Ackermann, S. Heinz, B. Kindler, J. Khuyagbaatar, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, Š. Šáro, B. Sulignano, J. Uusitalo, M. Venhart, Alpha-gamma decay studies of  $^{261}\text{sg}$  and  $^{257}\text{rf}$ , The European Physical Journal A 45 (3) (2010) 275–286. doi:10.1140/epja/i2010-11005-2.  
 URL <https://doi.org/10.1140/epja/i2010-11005-2>
- [95] R. D. Herzberg, P. T. Greenlees, P. A. Butler, G. D. Jones, M. Venhart, I. G. Darby, S. Eeckhaudt, K. Eskola, T. Grahn, C. Gray-Jones, F. P. Hessberger, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, W. Korten, M. Leino, A. P. Leppänen, S. Moon, M. Nyman, R. D. Page, J. Pakarinen, A. Pritchard, P. Rahkila, J. Sarén, C. Scholey, A. Steer, Y. Sun, C. Theisen, J. Uusitalo, Nuclear isomers in superheavy elements as stepping stones towards the island of stability, Nature 442 (7105) (2006) 896–899. doi:10.1038/nature05069.  
 URL <https://doi.org/10.1038/nature05069>
- [96] S. K. Tandel, T. L. Khoo, D. Seweryniak, G. Mukherjee, I. Ahmad, B. Back, R. Blinstrup, M. Carpenter, J. Chapman, P. Chowdhury, C. N. Davids, A. A. Hecht, A. Heinz, P. Ikin, R. V. F. Janssens, F. Kondev, T. Lauritsen, C. J. Lister, E. F. Moore, D. Peterson, P. Reiter, U. S. Tandel, X. Wang, S. Zhu, K isomers in  $^{254}\text{no}$ : Probing single-particle energies and pairing strength in the heaviest nuclei, Phys. Rev. Lett. 97 (2006) 082502. doi:10.1103/PhysRevLett.97.082502.  
 URL <http://link.aps.org/doi/10.1103/PhysRevLett.97.082502>
- [97] R. Clark, K. Gregorich, J. Berryman, M. Ali, J. Allmond, C. Beausang, M. Cromaz, M. Deleplanque, I. Dragojević, J. Dvorak, P. Ellison, P. Fallon, M. Garcia, J. Gates, S. Gros, H. Jeppesen, D. Kaji, I. Lee, A. Macchiavelli, K. Morimoto, H. Nitsche, S. Paschalis, M. Petri, L. Stavse-

- tra, F. Stephens, H. Watanabe, M. Wiedeking, **High- $k$  multi-quasiparticle states in  $^{254}\text{No}$** , Physics Letters B 690 (1) (2010) 19–24. doi:<https://doi.org/10.1016/j.physletb.2010.04.079>. URL <https://www.sciencedirect.com/science/article/pii/S0370269310005757>
- [98] F. P. Heßberger, S. Antalic, B. Sulignano, D. Ackermann, S. Heinz, S. Hofmann, B. Kindler, J. Khuyagbaatar, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, A. G. Popeko, Š. Šáro, B. Streicher, J. Uusitalo, M. Venhart, A. V. Yeremin, **Decay studies of K isomers in  $^{254}\text{No}$** , The European Physical Journal A 43 (1) (2010) 55. doi:<https://doi.org/10.1140/epja/i2009-10899-9>. URL <https://doi.org/10.1140/epja/i2009-10899-9>
- [99] K. Kessaci, **Synthèse de nouveaux éléments superlourds et spectroscopie des nobélia 255 et 256**, Ph.D. thesis, l’Université de Strasbourg (2022). URL <https://www.theses.fr/s269617>
- [100] K. Kessaci, B. J. P. Gall, O. Dorvaux, A. Lopez-Martens, R. Chakma, K. Hauschild, M. L. Chelnokov, V. I. Chepigin, M. Forge, A. V. Isaev, I. N. Izosimov, D. E. Katrasev, A. A. Kuznetsova, O. N. Malyshev, R. Mukhin, J. Piot, A. G. Popeko, Y. A. Popov, E. A. Sokol, A. I. Svirikhin, M. S. Tezekbayev, A. V. Yeremin, **Evidence of high- $k$  isomerism in  $^{256}\text{No}_{154}$** , Phys. Rev. C 104 (2021) 044609. doi:<https://doi.org/10.1103/PhysRevC.104.044609>. URL <https://link.aps.org/doi/10.1103/PhysRevC.104.044609>
- [101] H. B. Jeppesen, R. M. Clark, K. E. Gregorich, A. V. Afanasjev, M. N. Ali, J. M. Allmond, C. W. Beau-sang, M. Cromaz, M. A. Deleplanque, I. Dragojević, J. Dvorak, P. A. Ellison, P. Fallon, M. A. Garcia, J. M. Gates, S. Gros, I. Y. Lee, A. O. Macchiavelli, S. L. Nelson, H. Nitsche, L. Stavsetra, F. S. Stephens, M. Wiedeking, **High- $k$  multi-quasiparticle states and rotational bands in  $^{255}\text{Lr}$** , Phys. Rev. C 80 (2009) 034324. doi:<https://doi.org/10.1103/PhysRevC.80.034324>. URL <https://link.aps.org/doi/10.1103/PhysRevC.80.034324>
- [102] K. Hauschild, A. Lopez-Martens, A. V. Yeremin, O. Dorvaux, S. Antalic, A. V. Belozerov, C. Brianc̄on, M. L. Chelnokov, V. I. Chepigin, D. Curien, B. Gall, A. Görgen, V. A. Gorshkov, M. Guttormsen, F. Hanappe, A. P. Kabachenko, F. Khalfallah, A. C. Larsen, O. N. Malyshev, A. Minkova, A. G. Popeko, M. Rousseau, N. Rowley, S. Saro, A. V. Shutov, S. Siem, L. Stuttḡe, A. I. Svirikhin, N. U. H. Syed, C. Theisen, M. Venhart, **High- $k$ ,  $t_{1/2} = 1.4(1)$  ms, isomeric state in  $^{255}\text{Lr}$** , Phys. Rev. C 78 (2008) 021302. doi:<https://doi.org/10.1103/PhysRevC.78.021302>. URL <https://link.aps.org/doi/10.1103/PhysRevC.78.021302>
- [103] J. Khuyagbaatar, H. Brand, R. A. Cantemir, C. E. Düllmann, F. P. Heßberger, E. Jäger, B. Kindler, J. Krier, N. Kurz, B. Lommel, B. Schausten, A. Yakushev, **Spontaneous fission instability of the neutron-deficient no and rf isotopes: The new isotope  $^{249}\text{No}$** , Phys. Rev. C 104 (2021) L031303. doi:<https://doi.org/10.1103/PhysRevC.104.L031303>. URL <https://link.aps.org/doi/10.1103/PhysRevC.104.L031303>
- [104] P. Mosat, S. Antalic, F. P. Heßberger, D. Ackermann, B. Andel, M. S. Block, Z. Kalaninova, B. Kindler, M. Laatiaoui, B. Lommel, A. K. Mistry, J. Piot, M. Vostinar, **Isomeric states in  $^{255}\text{Rf}$ ,  $^{256}\text{Rf}$  and  $^{257}\text{Rf}$** , Acta Phys. Pol. B 51 (3) (2020) 849–854. doi:<https://doi.org/10.1103/PhysRevC.101.034310>. URL <https://www.actaphys.uj.edu.pl/R/51/3/849>
- [105] R. Chakma, A. Lopez-Martens, K. Hauschild, A. V. Yeremin, O. N. Malyshev, A. G. Popeko, Y. A. Popov, A. I. Svirikhin, V. I. Chepigin, E. A. Sokol, A. V. Isaev, A. A. Kuznetsova, M. L.

- Chelnokov, M. S. Tezekbayeva, I. N. Izosimov, O. Dorvaux, B. Gall, Z. Asfari, [Investigation of isomeric states in  \$^{255}\text{Rf}\$](#) , Phys. Rev. C 107 (2023) 014326. doi:[10.1103/PhysRevC.107.014326](https://doi.org/10.1103/PhysRevC.107.014326). URL <https://link.aps.org/doi/10.1103/PhysRevC.107.014326>
- [106] H. B. Jeppesen, I. Dragojević, R. M. Clark, K. E. Gregorich, M. N. Ali, J. M. Allmond, C. W. Beausang, D. L. Bleuel, M. Cromaz, M. A. Deleplanque, P. A. Ellison, P. Fallon, M. A. Garcia, J. M. Gates, J. P. Greene, S. Gros, I. Y. Lee, H. L. Liu, A. O. Macchiavelli, S. L. Nelson, H. Nitsche, J. R. Pavan, L. Stavsetra, F. S. Stephens, M. Wiedeking, R. Wyss, F. R. Xu, [Multi-quasiparticle states in  \$^{256}\text{Rf}\$](#) , Phys. Rev. C 79 (2009) 031303. doi:[10.1103/PhysRevC.79.031303](https://doi.org/10.1103/PhysRevC.79.031303). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.031303>
- [107] J. S. Berryman, R. M. Clark, K. E. Gregorich, J. M. Allmond, D. L. Bleuel, M. Cromaz, I. Dragojević, J. Dvorak, P. A. Ellison, P. Fallon, M. A. Garcia, S. Gros, I. Y. Lee, A. O. Macchiavelli, H. Nitsche, S. Paschalis, M. Petri, J. Qian, M. A. Stoyer, M. Wiedeking, [Electromagnetic decays of excited states in  \$^{261}\text{Sg}\$  \( \$z = 106\$ \) and  \$^{257}\text{Rf}\$  \( \$z = 104\$ \)](#), Phys. Rev. C 81 (2010) 064325. doi:[10.1103/PhysRevC.81.064325](https://doi.org/10.1103/PhysRevC.81.064325). URL <https://link.aps.org/doi/10.1103/PhysRevC.81.064325>
- [108] J. Qian, A. Heinz, T. L. Khoo, R. V. F. Janssens, D. Peterson, D. Seweryniak, I. Ahmad, M. Asai, B. B. Back, M. P. Carpenter, A. B. Garnsworthy, J. P. Greene, A. A. Hecht, C. L. Jiang, F. G. Kondev, T. Lauritsen, C. J. Lister, A. Robinson, G. Savard, R. Scott, R. Vondrasek, X. Wang, R. Winkler, S. Zhu, [Spectroscopy of  \$^{257}\text{Rf}\$](#) , Phys. Rev. C 79 (2009) 064319. doi:[10.1103/PhysRevC.79.064319](https://doi.org/10.1103/PhysRevC.79.064319). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.064319>
- [109] J. Rissanen, R. M. Clark, K. E. Gregorich, J. M. Gates, C. M. Campbell, H. L. Crawford, M. Cromaz, N. E. Esker, P. Fallon, U. Forsberg, O. Gothe, I.-Y. Lee, H. L. Liu, A. O. Macchiavelli, P. Mudder, H. Nitsche, G. Pang, A. Rice, D. Rudolph, M. A. Stoyer, A. Wiens, F. R. Xu, [Decay of the high- \$k\$  isomeric state to a rotational band in  \$^{257}\text{rf}\$](#) , Phys. Rev. C 88 (2013) 044313. doi:[10.1103/PhysRevC.88.044313](https://doi.org/10.1103/PhysRevC.88.044313). URL <https://link.aps.org/doi/10.1103/PhysRevC.88.044313>
- [110] D. Ackermann, F. P. Heßberger, S. Antalic, M. Block, H.-G. Burkhard, V. F. Comas, P. T. Greenlees, S. Heinz, S. Hofmann, S. Ketelhut, J. Khuyagbaatar, B. Kindler, I. Kojouharov, M. Mazzocco, M. Leino, B. Lommel, R. Mann, J. Maurer, A. G. Popeko, J. Sorri, J. Uusitalo, A. V. Yeremin, [Discovery of a K-isomer in  \$^{266}\text{Hs}\$](#) , GSI Sci. Rep. 2011 (2012) 208.
- [111] V. F. Weisskopf, [Radiative transition probabilities in nuclei](#), Phys. Rev. 83 (1951) 1073–1073. doi:[10.1103/PhysRev.83.1073](https://doi.org/10.1103/PhysRev.83.1073). URL <https://link.aps.org/doi/10.1103/PhysRev.83.1073>
- [112] S. Ćwiok, S. Hofmann, W. Nazarewicz, [Shell structure of the heaviest elements](#), Nucl. Phys. A 573 (3) (1994) 356 – 394. doi:[http://dx.doi.org/10.1016/0375-9474\(94\)90349-2](http://dx.doi.org/10.1016/0375-9474(94)90349-2). URL <http://www.sciencedirect.com/science/article/pii/0375947494903492>
- [113] S. Antalic, F. P. Heßberger, D. Ackermann, S. Heinz, S. Hofmann, B. Kindler, J. Khuyagbaatar, B. Lommel, R. Mann, [Nuclear isomers in  \$^{259}\text{Sg}\$  and  \$^{255}\text{rf}\$](#) , Eur. Phys. J. A 51 (4) (2015) 41. doi:[10.1140/epja/i2015-15041-0](https://doi.org/10.1140/epja/i2015-15041-0). URL <https://doi.org/10.1140/epja/i2015-15041-0>
- [114] R. Smolańczuk, A. Sobiczewski, [Shell effects in the properties of heavy and superheavy nuclei](#), in:

- Y. Oganessian, R. Kalpakchieva, W. von Oertzen (Eds.), Proceedings of the XV. Nuclear Physics Divisional Conference on Low Energy Nuclear Dynamics, World Scientific, Singapore, St.Petersburg, Russia, 1995, p. 313.
- [115] R. Smończuk, J. Skalski, A. Sobiczewski, Spontaneous-fission half-lives of deformed super-heavy nuclei, Phys. Rev. C 52 (1995) 1871.
- [116] M. Thoennessen, The Discovery of Isotopes - A Complete Compilation, 1st Edition, Springer International Publishing, 2016. doi:[10.1007/978-3-319-31763-2](https://doi.org/10.1007/978-3-319-31763-2).
- [117] M. Wang, W. Huang, F. Kondev, G. Audi, S. Naimi, The AME 2020 atomic mass evaluation (II). tables, graphs and references, Chinese Physics C 45 (3) (2021) 030003. doi:[10.1088/1674-1137/abddaf](https://doi.org/10.1088/1674-1137/abddaf)  
URL <https://doi.org/10.1088/1674-1137/abddaf>
- [118] H. Yuichi, O. Tsutomu, S. Keisuke, N. Hiromichi, K. Isao, M. Masaaki, S. Nobuo, H. L. Hall, R. A. Henderson, C. M. Gannet, J. A. Leyba, R. B. Chadwick, K. E. Gregorich, D. Lee, M. J. Nurmia, D. C. Hoffman, Alpha decay properties of light einsteinium isotopes, Nucl. Phys. A 500 (1) (1989) 90–110. doi:[https://doi.org/10.1016/0375-9474\(89\)90131-0](https://doi.org/10.1016/0375-9474(89)90131-0).  
URL <https://www.sciencedirect.com/science/article/pii/0375947489901310>
- [119] K. Moody, R. Lougheed, J. Wild, R. Dougan, E. Hulet, R. Hoff, C. Henderson, R. Dupzyk, R. Hahn, K. Sümmerer, G. O’Kelley, G. Bethune, Decay properties of heavy mendelevium isotopes, Nuclear Physics A 563 (1) (1993) 21 – 73. doi:[http://dx.doi.org/10.1016/0375-9474\(93\)90010-U](https://dx.doi.org/10.1016/0375-9474(93)90010-U).  
URL [http://www.sciencedirect.com/science/article/pii/037594749390010U](https://www.sciencedirect.com/science/article/pii/037594749390010U)
- [120] V. Ninov, F. P. Heßberger, S. Hofmann, H. Folger, G. Münzenberg, P. Armbruster, A. V. Yeremin, A. G. Popeko, M. Leino, S. Saro, Identification of new mendelevium and einsteinium isotopes in bombardments of  $^{209}\text{Bi}$  with  $^{40}\text{Ar}$ , Zeitschrift für Physik A Hadrons and Nuclei 356 (1) (1996) 11–12. doi:[10.1007/s002180050141](https://doi.org/10.1007/s002180050141).  
URL <https://doi.org/10.1007/s002180050141>
- [121] Z.-G. Gan, Z. Qin, H.-M. Fan, X. Lei, Y. Xu, J. He, H. Liu, X. Wu, J. Guo, X. Zhou, et al., A new alpha-particle-emitting isotope  $^{259}\text{Db}$ , The European Physical Journal A-Hadrons and Nuclei 10 (1) (2001) 21–25.
- [122] F.P. Heßberger, S. Hofmann, D. Ackermann, V. Ninov, M. Leino, G. Münzenberg, S. Saro, A. Lavrentev, A.G. Popeko, A.V. Yeremin, Ch. Stodel, Decay properties of neutron-deficient isotopes  $^{256,257}\text{db}$ ,  $^{255}\text{rf}$ ,  $^{252,253}\text{lr}$ , Eur. Phys. J. A 12 (1) (2001) 57–67. doi:[10.1007/s100500170039](https://doi.org/10.1007/s100500170039).  
URL <https://doi.org/10.1007/s100500170039>
- [123] Z.-G. Gan, J. Guo, X. Wu, Z. Qin, H.-M. Fan, X. Lei, H. Liu, B. Guo, H. Xu, R. Chen, et al., New isotope  $^{265}\text{Bh}$ , The European Physical Journal A-Hadrons and Nuclei 20 (3) (2004) 385–387.
- [124] F. P. Heßberger, S. Antalic, B. Streicher, S. Hofmann, D. Ackermann, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, S. Saro, B. Sulignano, Energy systematics of low-lying Nilsson levels in odd-mass einsteinium isotopes, The European Physical Journal A - Hadrons and Nuclei 26 (2) (2005) 233–239. doi:[10.1140/epja/i2005-10171-6](https://doi.org/10.1140/epja/i2005-10171-6).  
URL <https://doi.org/10.1140/epja/i2005-10171-6>
- [125] F. P. Heßberger, S. Antalic, D. Ackermann, Z. Kalaninovà, S. Heinz, S. Hofmann, B. Streicher, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, Š. Šáro, B. Sulignano, M. Venhart, Alpha-gamma decay studies of  $^{253}\text{No}$  and its

- daughter products  $^{253}\text{Md}$  and  $^{249}\text{Fm}$ , Eur. Phys. J. A 48: (2012) 75.
- [126] A. N. Andreyev, M. Huyse, P. Van Duppen, **Colloquium: Beta-delayed fission of atomic nuclei**, Rev. Mod. Phys. 85 (4) (2013) 1541–1559. doi:10.1103/RevModPhys.85.1541. URL <https://link.aps.org/doi/10.1103/RevModPhys.85.1541>
- [127] A. N. Andreyev, K. Nishio, K.-H. Schmidt, Nuclear fission: a review of experimental advances and phenomenology., Rep Prog Phys 81 (1) (2018) 016301. doi:10.1088/1361-6633/aa82eb.
- [128] J. Khuyagbaatar, R. A. Cantemir, C. E. Düllmann, E. Jäger, B. Kindler, J. Krier, N. Kurz, B. Lommel, B. Schausten, A. Yakushev, **Decay properties of the neutron-deficient isotope  $^{242}\text{Es}$** , Phys. Rev. C 109 (2024) 034311. doi:10.1103/PhysRevC.109.034311. URL <https://link.aps.org/doi/10.1103/PhysRevC.109.034311>
- [129] A. N. Andreyev, M. Huyse, P. Van Duppen, **Colloquium: Beta-delayed fission of atomic nuclei**, Rev. Mod. Phys. 85 (2013) 1541–1559. doi:10.1103/RevModPhys.85.1541. URL <https://link.aps.org/doi/10.1103/RevModPhys.85.1541>
- [130] J. Khuyagbaatar,  **$\beta$ -delayed fission in the heaviest nuclei**, The European Physical Journal A 55 (8) (2019) 134. doi:10.1140/epja/i2019-12824-1. URL <https://doi.org/10.1140/epja/i2019-12824-1>
- [131] G. L. Wilson, M. Takeyama, A. N. Andreyev, B. Andel, S. Antalic, W. N. Catford, L. Ghys, H. Haba, F. P. Heßberger, M. Huang, D. Kaji, Z. Kalaninova, K. Morimoto, K. Morita, M. Murakami, K. Nishio, R. Orlandi, A. G. Smith, K. Tanaka, Y. Wakabayashi, S. Yamaki,  **$\beta$ -delayed fission of  $^{230}\text{Am}$** , Phys. Rev. C 96 (2017) 044315. doi:10.1103/PhysRevC.96.044315. URL <https://link.aps.org/doi/10.1103/PhysRevC.96.044315>
- [132] P. Möller, A. J. Sierk, T. Ichikawa, A. Iwamoto, R. Bengtsson, H. Uhrenholt, S. Åberg, **Heavy-element fission barriers**, Phys. Rev. C 79 (2009) 064304. doi:10.1103/PhysRevC.79.064304. URL <http://link.aps.org/doi/10.1103/PhysRevC.79.064304>
- [133] R. Briselet, C. Theisen, M. Vandebruck, A. Marchix, M. Airiau, K. Auranen, H. Badran, D. Boilley, T. Calverley, D. Cox, F. Déchery, F. Defranchi Bisso, A. Drouart, B. Gall, T. Goigoux, T. Grahn, P. T. Greenlees, K. Hauschild, A. Herzan, R. D. Herzberg, U. Jakobsson, R. Julin, S. Juutinen, J. Konki, M. Leino, A. Lightfoot, A. Lopez-Martens, A. Mistry, P. Nieminen, J. Pakarinen, P. Papadakis, J. Partanen, P. Peura, P. Rahkila, J. Rubert, P. Ruotsalainen, M. Sandzelius, J. Saren, C. Scholey, J. Sorri, S. Stolze, B. Sulignano, J. Uusitalo, A. Ward, M. Zielińska, **Production cross section and decay study of  $^{243}\text{Es}$  and  $^{249}\text{Md}$** , Phys. Rev. C 99 (2019) 024614. doi:10.1103/PhysRevC.99.024614. URL <https://link.aps.org/doi/10.1103/PhysRevC.99.024614>
- [134] J. L. Pore, J. M. Gates, R. Orford, F. H. Garcia, H. L. Crawford, P. Fallon, J. A. Gooding, M. Kireeff Covo, M. McCarthy, M. A. Stoyer, **Properties of the  $^{256}\text{Db}$  decay chain**, Phys. Rev. C 110 (2024) 044313. doi:10.1103/PhysRevC.110.044313. URL <https://link.aps.org/doi/10.1103/PhysRevC.110.044313>
- [135] D. A. Shaughnessy, K. E. Gregorich, J. L. Adams, M. R. Lane, C. A. Laue, D. M. Lee, C. A. McGrath, V. Ninov, J. B. Patin, D. A. Strellis, E. R. Sylvester, P. A. Wilk, D. C. Hoffman, **Electron-capture delayed fission properties of  $^{244}\text{Es}$** , Phys. Rev. C 65 (2002)

024612. doi:[10.1103/PhysRevC.65.024612](https://doi.org/10.1103/PhysRevC.65.024612).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.65.024612>
- [136] M. Vostinar, F. P. Heßberger, D. Ackermann, B. Andel, S. Antalic, M. Block, C. Droese, J. Even, S. Heinz, Z. Kalaninova, I. Kojouharov, M. Laatiaoui, A. K. Mistry, J. Piot, H. Savajols, Alpha-gamma decay studies of  $^{258}\text{db}$  and its (grand)daughter nuclei  $^{254}\text{lr}$  and  $^{250}\text{md}$ , Eur. Phys. J. A 55 (2) (2019) 17. doi:[10.1140/epja/i2019-12701-y](https://doi.org/10.1140/epja/i2019-12701-y).  
 URL <https://doi.org/10.1140/epja/i2019-12701-y>
- [137] A. Pal, S. Santra, P. C. Rout, R. Gandhi, A. Baishya, T. Santhosh, R. Tripathi, T. N. Nag, Observation of a fission mode with very short elongation for the neutron-rich  $^{257}\text{Md}$  nucleus at high excitation energy, Phys. Rev. C 104 (2021) L031602. doi:[10.1103/PhysRevC.104.L031602](https://doi.org/10.1103/PhysRevC.104.L031602).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.104.L031602>
- [138] D. C. Hoffman, D. M. Lee, K. E. Gregorich, M. J. Nurmia, R. B. Chadwick, K. B. Chen, K. R. Czerwinski, C. M. Gannett, H. L. Hall, R. A. Henderson, B. Kadkhodayan, S. A. Kreek, J. D. Leyba, Spontaneous fission properties of 2.9-s  $^{256}\text{No}$ , Phys. Rev. C 41 (1990) 631–639. doi:[10.1103/PhysRevC.41.631](https://doi.org/10.1103/PhysRevC.41.631).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.41.631>
- [139] M. Asai, K. Tsukada, Y. Kasamatsu, Y. Ishii, A. Toyoshima, T. K. Sato, Y. Nagame, I. Nishinaka, H. Haba, T. Ishii, Measurement of the first excited  $2^+$  state in  $^{252}\text{fm}$ , Jaea-tokai tandem annual report 2013 , page 9, JAEA-Tokai (2016). doi:DOI: [10.11484/jaea-review-2016-025](https://doi.org/10.11484/jaea-review-2016-025).
- [140] C. Theisen, P. Greenlees, T.-L. Khoo, P. Chowdhury, T. Ishii, In-beam spectroscopy of heavy elements, Nucl. Phys. A 944 (2015) 333–375, special Issue on Superheavy Elements. doi:<http://dx.doi.org/10.1016/j.nuclphysa.2015.07.014>.  
 URL <http://www.sciencedirect.com/science/article/pii/S0375947415001621>
- [141] J. Khuyagbaatar, F. P. Heßberger, S. Hofmann, D. Ackermann, H. G. Burkhard, S. Heinz, B. Kindler, I. Kojouharov, B. Lommel, R. Mann, J. Maurer, K. Nishio,  $\alpha$  decay of  $^{243}\text{Fm}_{143}$  and  $^{245}\text{Fm}_{145}$ , and of their daughter nuclei, Phys. Rev. C 102 (2020) 044312. doi:[10.1103/PhysRevC.102.044312](https://doi.org/10.1103/PhysRevC.102.044312).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.102.044312>
- [142] M. Nurmia, T. Sikkeland, R. Silva, A. Ghiorso, Spontaneous fission of light fermium isotopes; new nuclides  $^{244}\text{fm}$  and  $^{245}\text{fm}$ , Physics Letters B 26 (2) (1967) 78–80. doi:[https://doi.org/10.1016/0370-2693\(67\)90552-7](https://doi.org/10.1016/0370-2693(67)90552-7).  
 URL <https://www.sciencedirect.com/science/article/pii/0370269367905527>
- [143] M. Asai, K. Tsukada, H. Haba, Y. Ishii, T. Ichikawa, A. Toyoshima, T. Ishii, Y. Nagame, I. Nishinaka, Y. Kojima, K. Sueki, Neutron one-quasiparticle states in  $^{251}\text{fm}_{151}$  populated via the  $\alpha$  decay of  $^{255}\text{No}$ , Phys. Rev. C 83 (2011) 014315. doi:[10.1103/PhysRevC.83.014315](https://doi.org/10.1103/PhysRevC.83.014315).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.83.014315>
- [144] K. Rezynkina, A. Lopez-Martens, K. Hauschild, I. Deloncle, S. Péru, P. Brionnet, M. L. Chelnokov, V. I. Chepigin, O. Dorvaux, F. Déchery, H. Faure, B. Gall, A. V. Isaev, I. N. Izosimov, D. E. Katrasev, A. N. Kuznetsov, A. A. Kuznetsova, O. N. Malyshev, A. G. Popeko, Y. A. Popov, E. A. Sokol, A. I. Svirikhin, A. V. Yeremin, Influence of octupole vibration on the low-lying structure of  $^{251}\text{Fm}$  and other heavy  $n = 151$  isotones, Phys. Rev. C 97 (2018) 054332. doi:[10.1103/PhysRevC.97.054332](https://doi.org/10.1103/PhysRevC.97.054332).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.97.054332>

- [145] R. B. Firestone, V. S. Shirley, S. F. Chu, C. M. Baglin, J. Zipkin (Eds.), R. B. Firestone et al., Table of Isotopes, 8th Edition, John Wiley & Sons, New York,, 1996.
- [146] K. Gregorich, [Simulation of recoil trajectories in gas-filled magnetic separators](#), Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 711 (2013) 47–59. doi:<https://doi.org/10.1016/j.nima.2013.01.020>. URL <https://www.sciencedirect.com/science/article/pii/S0168900213000673>
- [147] J. M. Gates, G. K. Pang, J. L. Pore, K. E. Gregorich, J. T. Kwarsick, G. Savard, N. E. Esker, M. Kireeff Covo, M. J. Mogannam, J. C. Batchelder, D. L. Bleuel, R. M. Clark, H. L. Crawford, P. Fallon, K. K. Hubbard, A. M. Hurst, I. T. Kolaja, A. O. Macchivelli, C. Morse, R. Orford, L. Phair, M. A. Stoyer, First direct measurements of superheavy-element mass numbers, Phys. Rev. Lett. 121 (2018) 222501. doi:[10.1103/PhysRevLett.121.222501](https://doi.org/10.1103/PhysRevLett.121.222501). URL <https://link.aps.org/doi/10.1103/PhysRevLett.121.222501>
- [148] W. Reisdorf, Analysis of fissionability data at high excitation energies, Z. Phys. A 300 (2) (1981) 227–238. doi:[10.1007/BF01412298](https://doi.org/10.1007/BF01412298). URL <https://doi.org/10.1007/BF01412298>
- [149] F. Déchery, H. Savajols, M. Authier, A. Drouart, J. Nolen, D. Ackermann, A. Amthor, B. Bastin, A. Berryhill, D. Boutin, L. Caceres, M. Coffey, O. Delferrière, O. Dorvaux, B. Gall, K. Hauschild, A. Hue, B. Jacquot, N. Karkour, B. Laune, F. Le Blanc, N. Lecesne, A. Lopez-Martens, F. Lutton, S. Manikonda, R. Meinke, G. Olivier, J. Payet, J. Piot, O. Pochon, V. Prince, M. Souli, G. Stelzer, C. Stodel, M.-H. Stodel, B. Sulignano, E. Traykov, D. Uriot, The super separator spectrometer s3 and the associated detection systems: Sirius & leb-reglis3, Nucl. Instr. and Meth. B 376 (2016) 125–130, proceedings of the XVIIth International Conference on Electromagnetic Isotope Separators and Related Topics (EMIS2015), Grand Rapids, MI, U.S.A., 11-15 May 2015. doi:<https://doi.org/10.1016/j.nimb.2016.02.036>. URL <https://www.sciencedirect.com/science/article/pii/S0168583X1600166X>
- [150] A. Lopez-Martens, A. Yeremin, M. Tezekbayeva, Z. Asfari, P. Brionnet, O. Dorvaux, B. Gall, K. Hauschild, D. Ackermann, L. Caceres, M. Chelnokov, V. Chepigin, M. Gustova, A. Isaev, A. Karpov, A. Kuznetsova, J. Piot, O. Malyshев, A. Popeko, Y. Popov, K. Rezynkina, H. Savajols, A. Svirikhin, E. Sokol, P. Steineger, Measurement of proton-evaporation rates in fusion reactions leading to transfermium nuclei, Physics Letters B 795 (2019) 271–276. doi:<https://doi.org/10.1016/j.physletb.2019.06.010>. URL <https://www.sciencedirect.com/science/article/pii/S0370269319303880>
- [151] G. Münzenberg, S. Hofmann, F. P. Heßberger, W. Reisdorf, K. H. Schmidt, J. H. R. Schneider, P. Armbruster, C. C. Sahm, B. Thuma, Identification of element 107 by  $\alpha$  correlation chains, Z. Phys. A 300 (1) (1981) 107–108. doi:[10.1007/BF01412623](https://doi.org/10.1007/BF01412623). URL <https://doi.org/10.1007/BF01412623>
- [152] S. Hofmann, V. Ninov, F. Heßberger, H. Folger, G. Münzenberg, H. Schött, P. Armbruster, A. Andreyev, A. Popeko, A. Yeremin, M. Leino, R. Janik, Š. Šáro, M. Veselsky, Alpha decay and fission of new neutron deficient mendelevium isotopes, GSI Scientific Report 1993 GSI 94-1 (1994), GSI (1994).
- [153] R. Briselet, C. Theisen, B. Sulignano, M. Airiau, K. Auranen, D. M. Cox, F. Déchery, A. Drouart, Z. Favier, B. Gall, T. Goigoux, T. Grahn, P. T. Greenlees, K. Hauschild, A. Herzan, R.-D. Herzberg, U. Jakobsson, R. Julin, S. Juutinen, J. Konki, M. Leino, A. Lopez-Martens, A. Mistry, P. Nieminen,

- J. Pakarinen, P. Papadakis, P. Peura, E. Rey-Herme, P. Rahkila, J. Rubert, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, S. Stolze, J. Uusitalo, M. Vandebrouck, A. Ward, M. Zielińska, B. Bally, M. Bender, W. Ryssens, In-beam  $\gamma$ -ray and electron spectroscopy of  $^{249,251}\text{Md}$ , Phys. Rev. C 102 (2020) 014307. doi:[10.1103/PhysRevC.102.014307](https://doi.org/10.1103/PhysRevC.102.014307).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.102.014307>
- [154] I. Muntian, Z. Patyk, A. Sobczewski, Are super-heavy nuclei around  $^{270}\text{Hs}$  really deformed?, Physics Letters B 500 (3–4) (2001) 241 – 246. doi:[http://dx.doi.org/10.1016/S0370-2693\(01\)00090-9](http://dx.doi.org/10.1016/S0370-2693(01)00090-9).  
URL <http://www.sciencedirect.com/science/article/pii/S0370269301000909>
- [155] H. J. Krappe, J. R. Nix, A. J. Sierk, Unified nuclear potential for heavy-ion elastic scattering, fusion, fission, and ground-state masses and deformations, Phys. Rev. C 20 (1979) 992–1013. doi:[10.1103/PhysRevC.20.992](https://doi.org/10.1103/PhysRevC.20.992).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.20.992>
- [156] S. Cwiok, J. Dudek, W. Nazarewicz, J. Skalski, T. Werner, Single-particle energies, wave functions, quadrupole moments and g-factors in an axially deformed Woods-Saxon potential with applications to the two-centre-type nuclear problems, Computer Physics Communications 46 (3) (1987) 379–399. doi:[https://doi.org/10.1016/0010-4655\(87\)90093-2](https://doi.org/10.1016/0010-4655(87)90093-2).  
URL <https://www.sciencedirect.com/science/article/pii/0010465587900932>
- [157] M. Kortelainen, J. McDonnell, W. Nazarewicz, P.-G. Reinhard, J. Sarich, N. Schunck, M. V. Stoitsov, S. M. Wild, Nuclear energy density optimization: Large deformations, Phys. Rev. C 85 (2012) 024304. doi:[10.1103/PhysRevC.85.024304](https://doi.org/10.1103/PhysRevC.85.024304).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.85.024304>
- [158] P. Klüpfel, P.-G. Reinhard, T. J. Bürvenich, J. A. Maruhn, Variations on a theme by Skyrme: A systematic study of adjustments of model parameters, Phys. Rev. C 79 (2009) 034310. doi:[10.1103/PhysRevC.79.034310](https://doi.org/10.1103/PhysRevC.79.034310).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.79.034310>
- [159] S. Raeder, D. Ackermann, H. Backe, R. Beerwerth, J. C. Berengut, M. Block, A. Borschevsky, B. Cheal, P. Chhetri, C. E. Düllmann, V. A. Dzuba, E. Eliav, J. Even, R. Ferrer, V. V. Flambaum, S. Fritzsche, F. Giacoppo, S. Götz, F. P. Heßberger, M. Huyse, U. Kaldor, O. Kaleja, J. Khuyagbaatar, P. Kunz, M. Laatiaoui, F. Lautenschläger, W. Lauth, A. K. Mistry, E. Minaya Ramirez, W. Nazarewicz, S. G. Porsev, M. S. Safronova, U. I. Safronova, B. Schuetrumpf, P. Van Duppen, T. Walther, C. Wraith, A. Yakushev, Probing sizes and shapes of nobelium isotopes by laser spectroscopy, Phys. Rev. Lett. 120 (2018) 232503. doi:[10.1103/PhysRevLett.120.232503](https://doi.org/10.1103/PhysRevLett.120.232503).  
URL <https://link.aps.org/doi/10.1103/PhysRevLett.120.232503>
- [160] M. Block, D. Ackermann, K. Blaum, C. Droese, M. Dworschak, S. Eliseev, T. Fleckenstein, E. Haettner, F. Herfurth, F. P. Heßberger, S. Hofmann, J. Ketelaer, J. Ketter, H.-J. Kluge, G. Marx, M. Mazzocco, Y. N. Novikov, W. R. Plaß, A. Popeko, S. Rahaman, D. Rodríguez, C. Scheidenberger, L. Schweikhart, P. G. Thirolf, G. K. Vorobyev, C. Weber, Direct mass measurements above uranium bridge the gap to the island of stability, Nature 463 (7282) (2010) 785–788.
- [161] A. V. Isaev, A. V. Andreev, M. L. Chelnokov, V. I. Chepigin, I. N. Izosimov, A. A. Kuznetsova, O. N. Malyshev, R. S. Mukhin, A. G. Popeko, Y. A. Popov, T. M. Shneidman, E. A. Sokol, A. I. Svirikhin, M. S. Tezekbayeva, A. V. Yeremin, N. I. Zamyatin,

- P. Brionnet, O. Dorvaux, B. Gall, K. Kessaci, A. Sel-lam, K. Hauschild, A. Lopez-Martens, S. Antalic, P. Mosat, Comparative study of spontaneous-fission characteristics of  $^{252}\text{No}$  and  $^{254}\text{No}$  isotopes, Physics of Particles and Nuclei Letters 18 (4) (2021) 449–456. doi:10.1134/S1547477121040087.
- URL <https://doi.org/10.1134/S1547477121040087>
- [162] A. I. Svirikhin, A. V. Yeremin, N. I. Zamyatin, I. N. Izosimov, A. V. Isaev, A. A. Kuznetsova, O. N. Malyshev, R. S. Mukhin, A. G. Popeko, Y. A. Popov, E. A. Sokol, B. Sailaubekov, M. S. Tezekbayeva, M. L. Chelnokov, V. I. Chepigin, B. Andel, S. Antalic, A. Bronis, P. Mosat, B. Gall, O. Dorvaux, A. Lopez-Martens, K. Hauschild, The new  $^{249}\text{No}$  isotope, Physics of Particles and Nuclei Letters 18 (4) (2021) 445–448. doi:10.1134/S1547477121040154.
- URL <https://doi.org/10.1134/S1547477121040154>
- [163] Z.-H. Zhang, X.-T. He, J.-Y. Zeng, E.-G. Zhao, S.-G. Zhou, Systematic investigation of the rotational bands in nuclei with  $Z \approx 100$  using a particle-number conserving method based on a cranked shell model, Phys. Rev. C 85 (2012) 014324. doi:10.1103/PhysRevC.85.014324.
- URL <http://link.aps.org/doi/10.1103/PhysRevC.85.014324>
- [164] M. Bender, P. Bonche, T. Duguet, P.-H. Heenen, Skyrme mean-field study of rotational bands in transfermium isotopes, Nuclear Physics A 723 (3) (2003) 354 – 364. doi:[http://dx.doi.org/10.1016/S0375-9474\(03\)01081-9](http://dx.doi.org/10.1016/S0375-9474(03)01081-9).
- URL <http://www.sciencedirect.com/science/article/pii/S0375947403010819>
- [165] A. A. Kuznetsova, A. V. Yeremin, A. Lopez-Martens, K. Hauschild, A. G. Popeko, O. N. Malyshev, V. I. Chepigin, A. I. Svirikhin, A. V. Isaev, Y. A. Popov, M. L. Chelnokov, O. Dorvaux, B. Gall, M. S. Tezekbayeva, Detailed study of the radioactive decay properties of no, rf, and db isotopes, Bulletin of the Russian Academy of Sciences: Physics 84 (8) (2020) 932–937. doi:10.3103/S1062873820080225.
- URL <https://doi.org/10.3103/S1062873820080225>
- [166] M. Forge, Étude des états isomériques des noyaux dits superlourds : cas du  $^{254}\text{No}$ , Theses, Université de Strasbourg (Sep. 2023).
- URL <https://theses.hal.science/tel-04397507>
- [167] K. Hauschild, A. V. Yeremin, O. Dorvaux, A. Lopez-Martens, A. V. Belozerov, C. Briancon, M. L. Chelnokov, V. I. Chepigin, S. Garcia-Santamaria, V. A. Gorshkov, F. Hanappe, A. P. Kabachenko, A. Korichi, O. N. Malyshev, Y. T. Oganessian, A. G. Popeko, N. Rowley, A. V. Shutov, L. Stuttge, A. I. Svirikhin, GABRIELA: A new detector array for  $\gamma$ -ray and conversion electron spectroscopy of transfermium elements, Nucl. Instr. and Meth. in Phys. Research A 560 (2006) 388.
- [168] R. Chakma, K. Hauschild, A. Lopez-Martens, A. V. Yeremin, O. N. Malyshev, A. G. Popeko, Y. A. Popov, A. I. Svirikhin, V. I. Chepigin, O. Dorvaux, B. Gall, K. Kessaci, Gamma and conversion electron spectroscopy using gabriela, The European Physical Journal A 56 (10) (2020) 245. doi:10.1140/epja/s10050-020-00242-5.
- URL <https://doi.org/10.1140/epja/s10050-020-00242-5>
- [169] A. Popeko, On-line separators for the dubna superheavy element factory, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 376 (2016) 144 – 149, proceedings of the {XVIIth} International Conference on Electromagnetic Isotope Separators and Related Topics (EMIS2015), Grand Rapids, MI, U.S.A., 11-15 May 2015. doi:<http://dx.doi.org/10.1016/j.nimb.2016.02.025>.

- URL <http://www.sciencedirect.com/science/article/pii/S0168583X16001531>
- [170] A. Mistry, Z. Zhang, F. Heßberger, D. Ackermann, B. Andel, S. Antalica, M. Block, P. Chhetri, F. Dechery, C. Droese, C. Düllmann, F. Giacoppo, J. Hoffmann, O. Kaleja, J. Khuyagbaatar, N. Kurz, M. Laatiaoui, J. Maurer, P. Mosat, J. Piot, S. Raeder, M. Vostinar, A. Y. and, Decay spectroscopy of heavy isotopes at ship using the compass focal plane detection set-up, *Acta Phys. Pol. B* 49 (2018) 613–620.
- [171] R. D. Herzberg, S. Moon, S. Eeckhaudt, P. T. Greenlees, P. A. Butler, T. Page, A. V. Afanasjev, N. Amzal, J. E. Bastin, F. Becker, M. Bender, B. Bruyneel, J. F. C. Cocks, I. G. Darby, O. Dorvaux, K. Eskola, J. Gerl, T. Grahn, C. Gray-Jones, N. J. Hammond, K. Hauschild, P. H. Heenen, K. Helariutta, A. Herzberg, F. Hessberger, M. Houry, A. Hürstel, R. D. Humphreys, G. D. Jones, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, T. L. Khoo, W. Korten, P. Kuusiniemi, Y. LeCoz, M. Leino, A. P. Leppänen, C. J. Lister, R. Lucas, M. Muikku, P. Nieminen, M. Nyman, R. D. Page, J. Pakarinen, A. Pritchard, P. Rahkila, P. Reiter, M. Sandzelius, J. Saren, C. Schlegel, C. Scholey, C. Theisen, W. H. Trzaska, J. Uusitalo, A. Wiens, H. J. Wollersheim, Structure of rotational bands in  $^{253}\text{No}$ , *The European Physical Journal A* 42 (3) (2009) 333–337. doi:[10.1140/epja/i2009-10855-9](https://doi.org/10.1140/epja/i2009-10855-9). URL <http://dx.doi.org/10.1140/epja/i2009-10855-9>
- [172] P. Reiter, T. L. Khoo, I. Ahmad, A. V. Afanasjev, A. Heinz, T. Lauritsen, C. J. Lister, D. Seweryniak, P. Bhattacharyya, P. A. Butler, M. P. Carpenter, A. J. Chewter, J. A. Cizewski, C. N. Davids, J. P. Greene, P. T. Greenlees, K. Helariutta, R.-D. Herzberg, R. V. F. Janssens, G. D. Jones, R. Julin, H. Kankaanpää, H. Kettunen, F. G. Kondev, P. Kuusiniemi, M. Leino, S. Siem, A. A. Sonzogni, J. Uusitalo, I. Wiedenhöver, Structure of the odd- $a$ , shell-stabilized nucleus  $^{253}_{102}\text{No}$ , *Phys. Rev. Lett.* 95 (2005) 032501. doi:[10.1103/PhysRevLett.95.032501](https://doi.org/10.1103/PhysRevLett.95.032501).
- URL <http://link.aps.org/doi/10.1103/PhysRevLett.95.032501>
- [173] F. P. Heßberger, S. Hofmann, D. Ackermann, S. Antalic, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, K. Nishio, A. G. Popeko, B. Sulignano, S. Saro, B. Streicher, M. Venhart, A. V. Yeremin, Alpha-gamma decay studies of  $^{255}\text{No}$ , *Eur. Phys. J. A* 29 (2) (2006) 165–173.
- [174] K. H. Schmidt, C. C. Sahm, K. Pielenz, H. G. Clerc, Some remarks on the error analysis in the case of poor statistics, *Z. f. Physik A* 316 (1) (1984) 19–26. doi:[10.1007/BF01415656](https://doi.org/10.1007/BF01415656). URL <http://dx.doi.org/10.1007/BF01415656>
- [175] J. Khuyagbaatar, A. Yakushev, C. E. Düllmann, D. Ackermann, L.-L. Andersson, M. Asai, M. Block, R. A. Boll, H. Brand, D. M. Cox, M. Dasgupta, X. Derkx, A. Di Nitto, K. Eberhardt, J. Even, M. Evers, C. Fahlander, U. Forsberg, J. M. Gates, N. Gharibyan, P. Golubev, K. E. Gregorich, J. H. Hamilton, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. J. Hinde, J. Hoffmann, R. Hollinger, A. Hübner, E. Jäger, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, M. Laatiaoui, S. Lahiri, R. Lang, B. Lommel, M. Maiti, K. Miernik, S. Minami, A. Mistry, C. Mokry, H. Nitsche, J. P. Omtvedt, G. K. Pang, P. Papadakis, D. Renisch, J. Roberto, D. Rudolph, J. Runke, K. P. Rykaczewski, L. G. Sarmiento, M. Schädel, B. Schausten, A. Semchenkov, D. A. Shaughnessy, P. Steinegger, J. Steiner, E. E. Terešatov, P. Thörle-Pospiech, K. Tinschert, T. Torres De Heidenreich, N. Trautmann, A. Türler, J. Uusitalo, D. E. Ward, M. Wegrzecki, N. Wiehl, S. M. Van Cleve, V. Yakusheva,  $^{48}\text{Ca} + ^{249}\text{Bk}$  fusion reaction leading to element  $Z = 117$ : Long-lived  $\alpha$ -decaying  $^{270}\text{Db}$  and discovery of  $^{266}\text{Lr}$ , *Phys. Rev. Lett.* 112 (2014)

172501. doi:[10.1103/PhysRevLett.112.172501](https://doi.org/10.1103/PhysRevLett.112.172501).  
 URL <http://link.aps.org/doi/10.1103/PhysRevLett.112.172501>
- [176] J. Khuyagbaatar, A. Yakushev, C. E. Düllmann, D. Ackermann, L.-L. Andersson, M. Asai, M. Block, R. A. Boll, H. Brand, D. M. Cox, M. Dasgupta, X. Derkx, A. Di Nitto, K. Eberhardt, J. Even, M. Evers, C. Fahlander, U. Forsberg, J. M. Gates, N. Gharibyan, P. Golubev, K. E. Gregorich, J. H. Hamilton, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. J. Hinde, J. Hoffmann, R. Hollinger, A. Hübner, E. Jäger, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, M. Laatiaoui, S. Lahiri, R. Lang, B. Lommel, M. Maiti, K. Miernik, S. Minami, A. Mistry, C. Mokry, H. Nitsche, J. P. Omtvedt, G. K. Pang, P. Papadakis, D. Renisch, J. Roberto, D. Rudolph, J. Runke, K. P. Rykaczewski, L. G. Sarmiento, M. Schädel, B. Schausten, A. Semchenkov, D. A. Shaughnessy, P. Steinegger, J. Steiner, E. E. Tereshatov, P. Thörle-Pospiech, K. Tinschert, T. Torres De Heidenreich, N. Trautmann, A. Türler, J. Uusitalo, D. E. Ward, M. Wegrzejczi, N. Wiehl, S. M. Van Cleve, V. Yakusheva, **Fusion reaction  $^{48}\text{Ca} + ^{249}\text{Bk}$  leading to formation of the element ts ( $z = 117$ )**, Phys. Rev. C 99 (2019) 054306. doi:[10.1103/PhysRevC.99.054306](https://doi.org/10.1103/PhysRevC.99.054306).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.99.054306>
- [177] H. Huang, W. Zhang, A. Andreyev, Z. Liu, D. Seweryniak, Z. Li, C. Guo, A. Barzakh, P. Van Duppen, B. Andel, S. Antalic, M. Block, A. Bronis, M. Carpenter, P. Copp, J. Cubiss, B. Ding, D. Doherty, Z. Favier, F. Giacoppo, T. Huang, B. Kindler, F. Kondev, T. Lauritsen, J. Li, G. Li, B. Lommel, H. Lu, M. Al Monthery, P. Mošať, Y. Niu, C. Raiison, W. Reviol, G. Savard, S. Stolze, G. Wilson, H. Wu, Z. Wang, F. Xu, Q. Zeng, X. Yu, F. Zeng, X. Zhou, **First observation of the decay of the  $13/2^+$  isomer in  $^{183}\text{hg}$  and  $b(m2)$  systematics** of neutron transitions across the nuclear chart, Physics Letters B 833 (2022) 137345. doi:<https://doi.org/10.1016/j.physletb.2022.137345>.  
 URL <https://www.sciencedirect.com/science/article/pii/S0370269322004798>
- [178] Y. Ito, P. Schury, M. Wada, F. Arai, H. Haba, Y. Hirayama, S. Ishizawa, D. Kaji, S. Kimura, H. Koura, M. MacCormick, H. Miyatake, J. Y. Moon, K. Morimoto, K. Morita, M. Mukai, I. Murray, T. Niwase, K. Okada, A. Ozawa, M. Rosenbusch, A. Takamine, T. Tanaka, Y. X. Watanabe, H. Wollnik, S. Yamaki, **First direct mass measurements of nuclides around  $z = 100$  with a multireflection time-of-flight mass spectrograph**, Phys. Rev. Lett. 120 (2018) 152501. doi:[10.1103/PhysRevLett.120.152501](https://doi.org/10.1103/PhysRevLett.120.152501).  
 URL <https://link.aps.org/doi/10.1103/PhysRevLett.120.152501>
- [179] F. Giacoppo, M. Block, Private communication (2020).
- [180] H. Haba, F. Fan, D. Kaji, Y. Kasamatsu, H. Kikunaga, Y. Komori, N. Kondo, H. Kudo, K. Morimoto, K. Morita, M. Murakami, K. Nishio, J. P. Omtvedt, K. Ooe, Z. Qin, D. Sato, N. Sato, T. K. Sato, Y. Shigekawa, A. Shinohara, M. Takeyama, T. Tanaka, A. Toyoshima, K. Tsukada, Y. Wakabayashi, Y. Wang, S. Wulff, S. Yamaki, S. Yano, Y. Yasuda, T. Yokokita, **Production of  $^{266}\text{Bh}$  in the  $^{248}\text{Cm}(^{23}\text{Na}, 5n)^{266}\text{Bh}$  reaction and its decay properties**, Phys. Rev. C 102 (2020) 024625. doi:[10.1103/PhysRevC.102.024625](https://doi.org/10.1103/PhysRevC.102.024625).  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.102.024625>
- [181] Y. T. Oganessian, V. K. Utyonkov, N. D. Kovrizhnykh, F. S. Abdullin, S. N. Dmitriev, D. Ibadullayev, M. G. Itkis, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibaikin, A. N. Polyakov, A. G. Popeko, R. N. Sagaidak, L. Schlattauer, I. V. Shirokovski, V. D. Shubin, M. V. Shumeiko, D. I. Solovyev, Y. S. Tsyganov,

- A. A. Voinov, V. G. Subbotin, A. Y. Bodrov, A. V. Sabel'nikov, A. V. Khalkin, V. B. Zlokazov, K. P. Rykaczewski, T. T. King, J. B. Roberto, N. T. Brewer, R. K. Grzywacz, Z. G. Gan, Z. Y. Zhang, M. H. Huang, H. B. Yang, [First experiment at the super heavy element factory: High cross section of  \$^{288}\text{Mc}\$  in the  \$^{243}\text{Am} + ^{48}\text{Ca}\$  reaction and identification of the new isotope  \$^{264}\text{Lr}\$](#) , Phys. Rev. C 106 (2022) L031301. [doi:10.1103/PhysRevC.106.L031301](https://doi.org/10.1103/PhysRevC.106.L031301)  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.106.L031301>
- [182] Y. T. Oganessian, V. K. Utyonkov, N. D. Kovrizhnykh, F. S. Abdullin, S. N. Dmitriev, A. A. Dzhioev, D. Ibadullayev, M. G. Itkis, A. V. Karpov, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibakin, A. N. Polyakov, A. G. Popeko, I. S. Rogov, R. N. Sagaidak, L. Schlattauer, V. D. Shubin, M. V. Shumeiko, D. I. Solovyev, Y. S. Tsyanov, A. A. Voinov, V. G. Subbotin, A. Y. Bodrov, A. V. Sabel'nikov, A. V. Khalkin, K. P. Rykaczewski, T. T. King, J. B. Roberto, N. T. Brewer, R. K. Grzywacz, Z. G. Gan, Z. Y. Zhang, M. H. Huang, H. B. Yang, [New isotope  \$^{286}\text{Mc}\$  produced in the  \$^{243}\text{Am} + ^{48}\text{Ca}\$  reaction](#), Phys. Rev. C 106 (2022) 064306. [doi:10.1103/PhysRevC.106.064306](https://doi.org/10.1103/PhysRevC.106.064306)  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.106.064306>
- [183] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyanov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, J. B. Patin, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, R. W. Lougheed, [Measurements of cross sections for the fusion-evaporation reactions  \$^{244}\text{Pu}\(^{48}\text{Ca}, xn\)^{292-x}114\$  and  \$^{245}\text{Cm}\(^{48}\text{Ca}, xn\)^{293-x}116\$](#) , Phys. Rev. C 69 (2004) 054607. [doi:10.1103/PhysRevC.69.054607](https://doi.org/10.1103/PhysRevC.69.054607)
- URL <https://link.aps.org/doi/10.1103/PhysRevC.69.054607>
- [184] Y. T. Oganessian, F. S. Abdullin, S. N. Dmitriev, J. M. Gostic, J. H. Hamilton, R. A. Henderson, M. G. Itkis, K. J. Moody, A. N. Polyakov, A. V. Ramayya, J. B. Roberto, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, N. J. Stoyer, V. G. Subbotin, A. M. Sukhov, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vos-tokin, [Investigation of the  \$^{243}\text{Am} + ^{48}\text{Ca}\$  reaction products previously observed in the experiments on elements 113 and 115 and 117](#), Phys. Rev. C 87 (2013) 014302. [doi:10.1103/PhysRevC.87.014302](https://doi.org/10.1103/PhysRevC.87.014302).
- [185] D. Rudolph, U. Forsberg, P. Golubev, L. Sarmiento, A. Yakushev, L.-L. Andersson, A. D. Nitto, C. E. Düllmann, J. Gates, K. Gregorich, C. Gross, F. A. Heßberger, R.-D. Herzberg, J. Khuyagbaatar, J. V. Kratz, K. Rykaczewski, M. Schädel, S. Aberg, D. Ackermann, M. Block, H. Brand, B. Carlsson, D. Cox, X. Derkx, K. Eberhardt, J. Even, C. Fahlander, J. Gerl, E. Jäger, B. Kindler, J. Krier, I. Kojouharov, N. Kurz, B. Lommel, A. Mistry, C. Mokry, H. Nitsche, J. P. Omtvedt, P. Papadakis, I. Ragnarsson, J. Runke, H. Schaffner, B. Schausten, P. Thöle-Pospiech, T. Torres, T. Traut, N. Trautmann, A. Türler, A. Ward, D. E. Ward, N. Wiehl, [Spectroscopy of element 115 decay chains](#), Phys. Rev. Lett. 111 (2013) 112502.
- [186] J. M. Gates, K. E. Gregorich, O. R. Gothe, E. C. Uribe, G. K. Pang, D. L. Bleuel, M. Block, R. M. Clark, C. M. Campbell, H. L. Crawford, M. Cromaz, A. Di Nitto, C. E. Düllmann, N. E. Esker, C. Fahlander, P. Fallon, R. M. Farjadi, U. Forsberg, J. Khuyagbaatar, W. Loveland, A. O. Macchiavelli, E. M. May, P. R. Mudder, D. T. Olive, A. C. Rice, J. Rissanen, D. Rudolph, L. G. Sarmiento, J. A. Shusterman, M. A. Stoyer, A. Wiens, A. Yakushev, H. Nitsche, [Decay spectroscopy of element 115](#)

- daughters:  $^{280}\text{Rg} \rightarrow ^{276}\text{Mt}$  and  $^{276}\text{Mt} \rightarrow ^{272}\text{Bh}$ , Phys. Rev. C 92 (2015) 021301.  
doi:[10.1140/epja/i2017-12260-3](https://doi.org/10.1140/epja/i2017-12260-3).
- URL <https://doi.org/10.1140/epja/i2017-12260-3>
- [187] J. Khuyagbaatar, A. Yakushev, C. E. Düllmann, D. Ackermann, L.-L. Andersson, M. Block, H. Brand, D. M. Cox, J. Even, U. Forsberg, P. Golubev, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, J. Hoffmann, A. Hübner, E. Jäger, J. Jeppsson, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, B. Lommel, M. Maiti, S. Minami, A. K. Mistry, C. M. Mrosek, I. Pysmenetska, D. Rudolph, L. G. Sarmiento, H. Schaffner, M. Schädel, B. Schausten, J. Steiner, T. T. De Heidenreich, J. Uusitalo, M. Wegrzecki, N. Wiehl, V. Yakusheva, New short-lived isotope  $^{221}\text{U}$  and the mass surface near  $n = 126$ , Phys. Rev. Lett. 115 (2015) 242502. doi:[10.1103/PhysRevLett.115.242502](https://doi.org/10.1103/PhysRevLett.115.242502)
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.115.242502>
- [188] Y. T. Oganessian, F. S. Abdullin, C. Alexander, J. Binder, R. A. Boll, S. N. Dmitriev, J. Ezold, K. Felker, J. M. Gostic, R. K. Grzywacz, J. H. Hamilton, R. A. Henderson, M. G. Itkis, K. Miernik, D. Miller, K. J. Moody, A. N. Polyakov, A. V. Ramayya, J. B. Roberto, M. A. Ryabinin, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. V. Shumeiko, M. A. Stoyer, N. J. Stoyer, V. G. Subbotin, A. M. Sukhov, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, Experimental studies of the  $^{249}\text{Bk} + ^{48}\text{Ca}$  reaction including decay properties and excitation function for isotopes of element 117, and discovery of the new isotope  $^{277}\text{Mt}$ , Phys. Rev. C 87 (2013) 054621. doi:[10.1103/PhysRevC.87.054621](https://doi.org/10.1103/PhysRevC.87.054621)
- URL <https://link.aps.org/doi/10.1103/PhysRevC.87.054621>
- [189] F. P. Heßberger, Spontaneous fission properties of superheavy elements, The European Physical Journal A 53 (4) (2017) 75.
- [190] P. T. Greenlees, J. Rubert, J. Piot, B. J. P. Gall, L. L. Andersson, M. Asai, Z. Asfari, D. M. Cox, F. Dechery, O. Dorvaux, T. Grahn, K. Hauschild, G. Henning, , A. Herzan, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, T.-L. Khoo, M. Leino, J. Ljungvall, A. Lopez-Martens, R. Lozeva, P. Nieminen, J. Pakarinen, P. Papadakis, E. Parr, P. Peura, P. Rahkila, S. Rinta-Antila, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, D. Seweryniak, J. Sorri, B. Sulignano, C. Theisen, J. Uusitalo, , M. Venhart, Shell-structure and pairing interaction in super-heavy nuclei: Rotational properties of the  $Z=104$  nucleus  $^{256}\text{Rf}$ , Phys. Rev. Lett. 109 (2012) 012501. doi:[10.1103/PhysRevLett.109.012501](https://doi.org/10.1103/PhysRevLett.109.012501)
- URL <http://link.aps.org/doi/10.1103/PhysRevLett.109.012501>
- [191] D. Seweryniak, T. Huang, K. Auranen, A. D. Ayangeakaa, B. B. Back, M. P. Carpenter, P. Chowdhury, R. M. Clark, P. A. Copp, Z. Favier, K. Hauschild, X.-T. He, T. L. Khoo, F. G. Kondev, A. Korichi, T. Lauritsen, J. Li, C. Morse, D. H. Potterveld, G. Savard, S. Stolze, J. Wu, J. Zhang, Y.-F. Xu, Nuclear rotation at the fission limit in  $^{254}\text{Rf}$ , Phys. Rev. C 107 (2023) L061302. doi:[10.1103/PhysRevC.107.L061302](https://doi.org/10.1103/PhysRevC.107.L061302)
- URL <https://link.aps.org/doi/10.1103/PhysRevC.107.L061302>
- [192] P. Mosat, F. P. Heßberger, S. Antalic, D. Ackermann, B. Andel, M. Block, S. Hofmann, Z. Kalaninova, B. Kindler, M. Laatiaoui, B. Lommel, A. K. Mistry, J. Piot, M. Vostinar, K isomerism in  $^{255}\text{Rf}$  and total kinetic energy measurements for spontaneous fission of  $^{255,256,258}\text{Rf}$ , Phys. Rev. C 101 (2020) 034310. doi:[10.1103/PhysRevC.101.034310](https://doi.org/10.1103/PhysRevC.101.034310).

- URL <https://link.aps.org/doi/10.1103/PhysRevC.101.034310>
- [193] K. Nishio, S. Hofmann, F. P. Heßberger, D. Ackermann, S. Antalic, V. F. Comas, Z. Gan, S. Heinz, J. A. Heredia, H. Ikezoe, J. Khuyagbaatar, B. Kindler, I. Kojouharov, P. Kuusiniemi, B. Lommel, M. Mazzocco, S. Mitsuoka, Y. Nagame, T. Ohtsuki, A. G. Popeko, S. Saro, H. J. Schött, B. Sulignano, A. Svirikhin, K. Tsukada, K. Tsuruta, A. V. Yeremin, Measurement of evaporation residue cross-sections of the reaction  $^{30}\text{Si} + ^{238}\text{U}$  at subbarrier energies, AIP Conference Proceedings 891 (1) (2007) 71–79. [arXiv:https://aip.scitation.org/doi/pdf/10.1063/1.2713502](https://arxiv.org/abs/https://aip.scitation.org/doi/pdf/10.1063/1.2713502), doi:10.1063/1.2713502.
- URL <https://aip.scitation.org/doi/abs/10.1063/1.2713502>
- [194] N. Carjan, F. Ivanyuk, Y. Oganessian, G. Ter-Akopian, Fission of transactinide elements described in terms of generalized cassinian ovals: Fragment mass and total kinetic energy distributions, Nuclear Physics A 942 (2015) 97–109. doi:<https://doi.org/10.1016/j.nuclphysa.2015.07.019>.
- URL <https://www.sciencedirect.com/science/article/pii/S0375947415001815>
- [195] A. P. Robinson, T. L. Khoo, D. Seweryniak, I. Ahmad, M. Asai, B. B. Back, M. P. Carpenter, P. Chowdhury, C. N. Davids, J. Greene, P. T. Greenlees, K. Hauschild, A. Heinz, R.-D. Herzberg, R. V. F. Janssens, D. G. Jenkins, G. D. Jones, S. Ketelhut, F. G. Kondev, T. Lauritsen, C. J. Lister, A. Lopez-Martens, P. Marley, E. McCutchan, P. Papadakis, D. Peterson, J. Qian, D. Rostron, U. Shirwadkar, I. Stefanescu, S. K. Tandel, X. Wang, S. Zhu, Search for a 2-quasiparticle high- $k$  isomer in  $^{256}\text{Rf}$ , Phys. Rev. C 83 (2011) 064311. doi:10.1103/PhysRevC.83.064311.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.83.064311>
- [196] J. Gates, M. Garcia, K. Gregorich, C. Düllmann, I. Dragojević, J. Dvorak, R. Eichler, C. Folden, W. Loveland, S. Nelson, G. Pang, L. Stavsetra, R. Sudowe, A. Türler, H. Nitsche, Synthesis of rutherfordium isotopes in the  $^{238}\text{U}(^{26}\text{Mg}, xn)^{264-x}\text{Rf}$  reaction and study of their decay properties, Phys. Rev. C 77 (2008) 034603.
- [197] F. P. Heßberger, G. Münzenberg, S. Hofmann, Y. K. Agarwal, K. Poppensieker, W. Reisdorf, K. H. Schmidt, J. R. H. Schneider, W. F. W. Schneider, H. J. Schött, P. Armbruster, B. Thuma, C. C. Sahm, D. Vermeulen, The new isotopes  $^{258}\text{I}_{05}, ^{257}\text{I}_{05}, ^{254}\text{Lr}$  and  $^{253}\text{Lr}$ , Z. Phys. A 322 (4) (1985) 557–566. doi:10.1007/BF01415134.
- URL <https://doi.org/10.1007/BF01415134>
- [198] D. J. Hinde, D. Y. Jeung, E. Prasad, A. Wakhle, M. Dasgupta, M. Evers, D. H. Luong, R. du Rietz, C. Simenel, E. C. Simpson, E. Williams, Sub-barrier quasifission in heavy element formation reactions with deformed actinide target nuclei, Phys. Rev. C 97 (2018) 024616. doi:10.1103/PhysRevC.97.024616.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.97.024616>
- [199] V. K. Utyonkov, N. T. Brewer, Y. T. Oganessian, K. P. Rykaczewski, F. S. Abdullin, S. N. Dmitriev, R. K. Grzywacz, M. G. Itkis, K. Miernik, A. N. Polyakov, J. B. Roberto, R. N. Sagaidak, I. V. Shirokovsky, M. V. Shumeiko, Y. S. Tsyanov, A. A. Voinov, V. G. Subbotin, A. M. Sukhov, A. V. Karpov, A. G. Popeko, A. V. Sabel'nikov, A. I. Svirikhin, G. K. Vostokin, J. H. Hamilton, N. D. Kovrizhnykh, L. Schlattauer, M. A. Stoyer, Z. Gan, W. X. Huang, L. Ma, Neutron-deficient superheavy nuclei obtained in the  $^{240}\text{Pu} + ^{48}\text{Ca}$  reaction, Phys. Rev. C 97 (2018) 014320. doi:10.1103/PhysRevC.97.014320.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.97.014320>
- [200] V. K. Utyonkov, N. T. Brewer, Y. T. Oganessian,

- K. P. Rykaczewski, F. S. Abdullin, S. N. Dmitriev, R. K. Grzywacz, M. G. Itkis, K. Miernik, A. N. Polyakov, J. B. Roberto, R. N. Sagaidak, I. V. Shirokovsky, M. V. Shumeiko, Y. S. Tsyganov, A. A. Voinov, V. G. Subbotin, A. M. Sukhov, A. V. Sabel'nikov, G. K. Vostokin, J. H. Hamilton, M. A. Stoyer, S. Y. Strauss, [Experiments on the synthesis of superheavy nuclei  \$^{284}\text{Fl}\$  and  \$^{285}\text{Fl}\$  in the  \$^{239,240}\text{Pu} + ^{48}\text{Ca}\$  reactions](#), Phys. Rev. C 92 (2015) 034609. doi:[10.1103/PhysRevC.92.034609](https://doi.org/10.1103/PhysRevC.92.034609). URL <https://link.aps.org/doi/10.1103/PhysRevC.92.034609>
- [201] P. A. Ellison, K. E. Gregorich, J. S. Berryman, D. L. Bleuel, R. M. Clark, I. Dragojević, J. Dvorak, P. Fallon, C. Fineman-Sotomayor, J. M. Gates, O. R. Gothe, I. Y. Lee, W. D. Loveland, J. P. McLaughlin, S. Paschalis, M. Petri, J. Qian, L. Stavsetra, M. Wiedeking, H. Nitsche, [New superheavy element isotopes:  \$^{242}\text{Pu}\(^{48}\text{Ca}, 5n\)^{285}\text{Fl}\$](#) , Phys. Rev. Lett. 105 (2010) 182701. doi:[10.1103/PhysRevLett.105.182701](https://doi.org/10.1103/PhysRevLett.105.182701). URL <https://link.aps.org/doi/10.1103/PhysRevLett.105.182701>
- [202] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, R. N. Sagaidak, I. V. Shirokovsky, Y. S. Tsyganov, A. A. Voinov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, V. G. Subbotin, A. M. Sukhov, K. Subotic, V. I. Zagrebaev, G. K. Vostokin, M. G. Itkis, R. A. Henderson, J. M. Kenneally, J. H. Landrum, K. J. Moody, D. A. Shaughnessy, M. A. Stoyer, N. J. Stoyer, P. A. Wilk, [Synthesis of the isotope  \$^{282}\text{Fl}\$  in the  \$^{237}\text{Np} + ^{48}\text{Ca}\$  fusion reaction](#), Phys. Rev. C 76 (2007) 011601. doi:[10.1103/PhysRevC.76.011601](https://doi.org/10.1103/PhysRevC.76.011601). URL <https://link.aps.org/doi/10.1103/PhysRevC.76.011601>
- [203] Y. T. Oganessian, Heaviest nuclei from  $^{48}\text{Ca}$ -induced reactions, J. Phys. G: Nucl. Part. Phys. 34 (2007) R165.
- [204] Y. T. Oganessian, [Synthesis of the heaviest elements in  \$^{48}\text{Ca}\$ -induced reactions](#), Radiochimica Acta 99 (7-8) (2011) 429–439. doi:[doi:10.1524/ract.2011.1860](https://doi.org/10.1524/ract.2011.1860). URL <https://doi.org/10.1524/ract.2011.1860>
- [205] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyganov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, J. B. Patin, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, P. A. Wilk, R. W. Lougheed, R. I. Il'kaev, S. P. Vesnovskii, [Measurements of cross sections and decay properties of the isotopes of elements 112, 114, and 116 produced in the fusion reactions  \$^{233,238}\text{U}\$ ,  \$^{242}\text{Pu}\$ , and  \$^{248}\text{Cm} + ^{48}\text{Ca}\$](#) , Phys. Rev. C 70 (2004) 064609. doi:[10.1103/PhysRevC.70.064609](https://doi.org/10.1103/PhysRevC.70.064609). URL <https://link.aps.org/doi/10.1103/PhysRevC.70.064609>
- [206] Y. T. Oganessian, V. K. Utyonkov, D. Ibadullayev, F. S. Abdullin, S. N. Dmitriev, M. G. Itkis, A. V. Karpov, N. D. Kovrizhnykh, D. A. Kuznetsov, O. V. Petrushkin, A. V. Podshibaikin, A. N. Polyakov, A. G. Popeko, R. N. Sagaidak, L. Schlattauer, V. D. Shubin, M. V. Shumeiko, D. I. Solovyev, Y. S. Tsyganov, A. A. Voinov, V. G. Subbotin, A. Y. Bodrov, A. V. Sabel'nikov, A. Lindner, K. P. Rykaczewski, T. T. King, J. B. Roberto, N. T. Brewer, R. K. Grzywacz, Z. G. Gan, Z. Y. Zhang, M. H. Huang, H. B. Yang, [Investigation of  \$^{48}\text{Ca}\$ -induced reactions with  \$^{242}\text{Pu}\$  and  \$^{238}\text{U}\$  targets at the jinr superheavy element factory](#), Phys. Rev. C 106 (2022) 024612. doi:[10.1103/PhysRevC.106.024612](https://doi.org/10.1103/PhysRevC.106.024612).

- URL <https://link.aps.org/doi/10.1103/PhysRevC.106.024612>
- [207] G.N.Flerov, The dubna programme on heavy and superheavy elements, in: Proceedings of the 3rd INTERNATIONAL CONFERENCE ON NUCLEI FAR FROM STABILITY, Vol. 76-13, CERN, Geneva, 1976, pp. 542–547.
- [208] Y. Oganessian, A. Demin, N. Danilov, G. Flerov, M. Ivanov, A. Iljinov, N. Kolesnikov, B. Markov, V. Plotko, S. Tretyakova, On spontaneous fission of neutron-deficient isotopes of elements 103, 105 and 107, Nuclear Physics A 273 (2) (1976) 505–522. doi:[https://doi.org/10.1016/0375-9474\(76\)90607-2](https://doi.org/10.1016/0375-9474(76)90607-2)
- URL <https://www.sciencedirect.com/science/article/pii/0375947476906072>
- [209] Y. T. Oganessian, Certain aspects of fusion and fission of heavy nuclear system, JINR report JINR-P7-83-644 (1983) 55.
- [210] A.-P. Leppänen, Alpha-decay and decay-tagging studies of heavy elements using the ritu separator, Ph.d thesis, University of Jyväskylä (2005).
- [211] S. L. Nelson, K. E. Gregorich, I. Dragojević, M. A. Garcia, J. M. Gates, R. Sudowe, H. Nitsche, Lightest isotope of bh produced via the  $^{209}\text{Bi}(^{52}\text{Cr}, n)^{260}\text{Bh}$  reaction, Phys. Rev. Lett. 100 (2008) 022501. doi:[10.1103/PhysRevLett.100.022501](https://doi.org/10.1103/PhysRevLett.100.022501)
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.100.022501>
- [212] A. V. Yeremin, M. S. Tezekbayeva, A. G. Popeko, O. N. Malyshev, A. Lopez-Martens, K. Hauschild, O. Dorvaux, B. Gall, A. V. Isaev, A. V. Karpov, A. N. Kuznetsov, A. A. Kuznetsova, Y. A. Popov, A. I. Svirikhin, E. A. Sokol, M. L. Chelnokov, V. I. Chepigin, P. Moshat, Measurement of the production cross sections for isotopes of transfermium elements in complete fusion reactions with the subsequent evaporation of a proton and neutrons, Physics of Particles and Nuclei Letters 16 (3) (2019) 224–228. doi:[10.1134/S1547477119030099](https://doi.org/10.1134/S1547477119030099).
- URL <https://doi.org/10.1134/S1547477119030099>
- [213] F. P. Heßberger, S. Hofmann, B. Streicher, B. Sulignano, S. Antalic, D. Ackermann, S. Heinz, B. Kindler, I. Kojouharov, P. Kuusiniemi, M. Leino, B. Lommel, R. Mann, A. G. Popeko, Š. Šáro, J. Uusitalo, A. V. Yeremin, Decay properties of neutron-deficient isotopes of elements from z = 101 to z = 108, Eur. Phys. J. A 41 (2) (2009) 145–153. doi:[10.1140/epja/i2009-10826-2](https://doi.org/10.1140/epja/i2009-10826-2)
- URL <https://doi.org/10.1140/epja/i2009-10826-2>
- [214] K. Morita, K. Morimoto, D. Kaji, T. Akiyama, S.-i. Goto, H. Haba, E. Ideguchi, R. Kamungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, H. Xu, T. Yamaguchi, A. Yoneda, A. Yoshida, Y. Zhao, Experiment on the synthesis of element 113 in the reaction  $^{209}\text{Bi}(^{70}\text{Zn}, n)^{278}\text{113}$ , Journal of the Physical Society of Japan 73 (10) (2004) 2593–2596. arXiv:<https://doi.org/10.1143/JPSJ.73.2593>.
- URL <https://doi.org/10.1143/JPSJ.73.2593>
- [215] K. Morita, K. Morimoto, D. Kaji, T. Akiyama, S.-i. Goto, H. Haba, E. Ideguchi, K. Katori, H. Koura, H. Kikunaga, H. Kudo, T. Ohnishi, A. Ozawa, N. Sato, T. Suda, K. Sueki, F. Tokanai, T. Yamaguchi, A. Yoneda, A. Yoshida, Observation of second decay chain from  $^{278}\text{113}$ , Journal of the Physical Society of Japan 76 (4) (2007) 045001. arXiv:<https://doi.org/10.1143/JPSJ.76.045001>, doi:[10.1143/JPSJ.76.045001](https://doi.org/10.1143/JPSJ.76.045001).
- URL <https://doi.org/10.1143/JPSJ.76.045001>
- [216] K. Morita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, Y. Kudou, N. Sato, T. Sumita, A. Yoneda, T. Ichikawa, Y. Fujimori, S.-i. Goto, E. Ideguchi,

- Y. Kasamatsu, K. Katori, Y. Komori, H. Koura, H. Kudo, K. Ooe, A. Ozawa, F. Tokanai, K. Tsukada, T. Yamaguchi, A. Yoshida, Decay properties of  $^{266}\text{bh}$  and  $^{262}\text{db}$  produced in the  $^{248}\text{Cm} + ^{23}\text{na}$  reaction, Journal of the Physical Society of Japan 78 (6) (2009) 064201. [arXiv:<https://arxiv.org/abs/10.1143/JPSJ.78.064201>](https://arxiv.org/abs/10.1143/JPSJ.78.064201)
- URL <https://doi.org/10.1143/JPSJ.78.064201>
- [217] K. Morita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, Y. Kudou, T. Sumita, Y. Wakabayashi, A. Yoneda, K. Tanaka, S. Yamaki, R. Sakai, T. Akiyama, S. ichi Goto, H. Hasebe, M. Huang, T. Huang, E. Ideguchi, Y. Kasamatsu, K. Katori, Y. Kariya, H. Kikunaga, H. Koura, H. Kudo, A. Mashiko, K. Mayama, S. ichi Mitsuoka, T. Moriya, M. Murakami, H. Murayama, S. Namai, A. Ozawa, N. Sato, K. Sueki, M. Takeyama, F. Tokanai, T. Yamaguchi, A. Yoshida, New result in the production and decay of an isotope and  $^{278}\text{113}$  and of the 113th element, J. Phys. Soc. Jpn. 81 (2012) 103201.
- [218] H. Haba, M. Huang, D. Kaji, J. Kanaya, Y. Kudou, K. Morimoto, K. Morita, M. Murakami, K. Ozeki, R. Sakai, T. Sumita, Y. Wakabayashi, A. Yoneda, Y. Kasamatsu, Y. Kikutani, Y. Komori, K. Nakamura, A. Shinohara, H. Kikunaga, H. Kudo, K. Nishio, A. Toyoshima, K. Tsukada, Production of  $^{262}\text{Db}$  in the  $^{248}\text{Cm}(^{19}\text{F},5n)^{262}\text{Db}$  reaction and decay properties of  $^{262}\text{Db}$  and  $^{258}\text{Lr}$ , Phys. Rev. C 89 (2014) 024618. [doi:10.1103/PhysRevC.89.024618](https://doi.org/10.1103/PhysRevC.89.024618)
- URL <http://link.aps.org/doi/10.1103/PhysRevC.89.024618>
- [219] Fernández, D., Caamaño, M., Ramos, D., Lemasson, A., Rejmund, M., Álvarez-Pol, H., Audouin, L., Frankland, J. D., Fernández-Domínguez, B., Galiana-Baldó, E., Piot, J., Schmitt, C., Ackermann, D., Biswas, S., Clement, E., Durand, D., Farget, F., Fregeau, M. O., Galaviz, D., Heinz, A., Henriques, A., Jacquot, B., Jurado, B., Kim, Y. H., Morfouace, P., Ralet, D., Roger, T., Teubig, P., Tsekhanovich, I., Experimental study of high-energy fission and quasi-fission dynamics with fusion-induced fission reactions at vamos++, EPJ Web of Conf. 284 (2023) 04009. [doi:10.1051/epjconf/202328404009](https://doi.org/10.1051/epjconf/202328404009)
- URL <https://doi.org/10.1051/epjconf/202328404009>
- [220] A. Lemasson, M. Rejmund, Fast trajectory reconstruction techniques for the large acceptance magnetic spectrometer vamos++, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 1054 (2023) 168407. [doi:<https://doi.org/10.1016/j.nima.2023.168407>](https://doi.org/10.1016/j.nima.2023.168407)
- URL <https://www.sciencedirect.com/science/article/pii/S0168900223003972>
- [221] N. Kolesnikov, A. Demin, E. Cherepanov, n.n., Communication D4-80-572, Joint Institute for Nuclear Research (JINR, Dubna) (1980).
- [222] I. Muntian, Z. Patyk, A. Sobiczewski, Calculated masses of heaviest nuclei, Phys. of Atomic Nuclei 66 (6) (2003) 1015.
- [223] Y. Oganessian, V. Utyonkov, Superheavy nuclei from  $^{48}\text{Ca}$ -induced reactions, Nucl. Phys. A 944 (2015) 62 – 98. [doi:\[http://dx.doi.org/10.1016/j.nuclphysa.2015.07.003\]\(https://doi.org/10.1016/j.nuclphysa.2015.07.003\)](https://doi.org/10.1016/j.nuclphysa.2015.07.003)
- URL <https://www.sciencedirect.com/science/article/pii/S0375947415001517>
- [224] Z. Łojewski, A. Staszczak, Role of pairing degrees of freedom and higher multipolarity deformations in spontaneous fission process, Nuclear Physics A 657 (2) (1999) 134–157. [doi:\[https://doi.org/10.1016/S0375-9474\\(99\\)00328-0\]\(https://doi.org/10.1016/S0375-9474\(99\)00328-0\)](https://doi.org/10.1016/S0375-9474(99)00328-0)
- URL <https://www.sciencedirect.com/science/article/pii/S0375947499003280>
- [225] K. H. Schmidt, A new test for random events of an exponential distribution, Eur. J. Phys. A 8 (1) (2000)

- 141–145. doi:[10.1007/s100500070129](https://doi.org/10.1007/s100500070129). URL <https://doi.org/10.1007/s100500070129>
- [226] E. I. Galkina, E. M. Kozulin, G. N. Knyazheva, I. M. Itkis, A. A. Bogachev, I. N. Diatlov, M. Cheralu, D. Kumar, N. I. Kozulina, K. V. Novikov, A. N. Pan, I. V. Pchelintsev, I. V. Vorobiev, W. H. Trzaska, S. Heinz, B. Lommel, E. Vardaci, S. Spinosa, A. Di Nitto, A. Pulcini, C. Borcea, I. Harca, **Investigating mass–energy distributions of fragments produced in the  $^{32}\text{s} + ^{232}\text{Th} \rightarrow ^{264}\text{Sg}$  reaction at energies below and near the coulomb barrier**, Bulletin of the Russian Academy of Sciences: Physics 85 (10) (2021) 1085–1089. doi:[10.3103/S1062873821100154](https://doi.org/10.3103/S1062873821100154). URL <https://doi.org/10.3103/S1062873821100154>
- [227] U. Brosa, S. Grossmann, A. Müller, **Nuclear scission**, Physics Reports 197 (4) (1990) 167–262. doi:[https://doi.org/10.1016/0370-1573\(90\)90114-H](https://doi.org/10.1016/0370-1573(90)90114-H). URL <https://www.sciencedirect.com/science/article/pii/037015739090114H>
- [228] J. Dvorak, W. Brüchle, M. Chelnokov, R. Dressler, C. E. Düllmann, K. Eberhardt, V. Gorshkov, E. Jäger, R. Krücken, A. Kuznetsov, Y. Nagame, F. Nebel, Z. Novackova, Z. Qin, M. Schädel, B. Schausten, E. Schimpf, A. Semchenkov, P. Thörle, A. Türler, M. Wegrzecki, B. Wierczinski, A. Yakushev, A. Yeremin, **Doubly magic nucleus  $^{270}\text{Hs}_{162}$** , Phys. Rev. Lett. 97 (2006) 242501. doi:[10.1103/PhysRevLett.97.242501](https://doi.org/10.1103/PhysRevLett.97.242501). URL <https://link.aps.org/doi/10.1103/PhysRevLett.97.242501>
- [229] J. Dvorak, W. Brüchle, M. Chelnokov, C. E. Düllmann, Z. Dvorakova, K. Eberhardt, E. Jäger, R. Krücken, A. Kuznetsov, Y. Nagame, F. Nebel, K. Nishio, R. Perego, Z. Qin, M. Schädel, B. Schausten, E. Schimpf, R. Schuber, A. Semchenkov, P. Thörle, A. Türler, M. Wegrzecki, B. Wierczinski, A. Yakushev, A. Yeremin, **Observa-**tion of the  $3n$  evaporation channel in the complete hot-fusion reaction  $^{26}\text{Mg} + ^{248}\text{Cm}$  leading to the new superheavy nuclide  $^{271}\text{Hs}$ , Phys. Rev. Lett. 100 (2008) 132503. doi:[10.1103/PhysRevLett.100.132503](https://doi.org/10.1103/PhysRevLett.100.132503). URL <https://link.aps.org/doi/10.1103/PhysRevLett.100.132503>
- [230] S. Hofmann, F. P. Heßberger, D. Ackermann, G. Münzenberg, S. Antalic, P. Cagarda, B. Kindler, J. Kojouharova, M. Leino, B. Lommel, R. Mann, A. G. Popeko, S. Reshitko, S. Šaro, J. Usitalo, A. V. Yeremin, **New results on elements 111 and 112**, The European Physical Journal A - Hadrons and Nuclei 14 (2) (2002) 147–157. doi:[10.1140/epja/i2001-10119-x](https://doi.org/10.1140/epja/i2001-10119-x). URL <https://doi.org/10.1140/epja/i2001-10119-x>
- [231] T. Sumita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, R. Sakai, A. Yoneda, A. Yoshida, H. Hasebe, K. Katori, N. Sato, Y. Wakabayashi, S.-i. Mitsuoka, S.-i. Goto, M. Murakami, Y. Kariya, F. Tokanai, K. Mayama, M. Takeyama, T. Moriya, E. Ideguchi, T. Yamaguchi, H. Kikunaga, J. Chiba, K. Morita, **New result on the production of  $^{277}\text{Cn}$  by the  $^{208}\text{Pb} + ^{70}\text{Zn}$  reaction**, Journal of the Physical Society of Japan 82 (2) (2013) 024202. doi:[10.7566/JPSJ.82.024202](https://doi.org/10.7566/JPSJ.82.024202). URL <https://doi.org/10.7566/JPSJ.82.024202>
- [232] C. Qi, F. R. Xu, R. J. Liotta, R. Wyss, M. Y. Zhang, C. Asawatangtrakuldee, D. Hu, **Microscopic mechanism of charged-particle radioactivity and generalization of the geiger-nuttall law**, Phys. Rev. C 80 (2009) 044326. doi:[10.1103/PhysRevC.80.044326](https://doi.org/10.1103/PhysRevC.80.044326). URL <https://link.aps.org/doi/10.1103/PhysRevC.80.044326>
- [233] Y.-Y. Xu, D.-X. Zhu, X. Chen, X.-J. Wu, B. He, X.-H. Li, **A unified formula for  $\alpha$  decay half-lives**, The European Physical Journal A 58 (9) (2022) 163. doi:[10.1140/epja/s10050-022-00812-9](https://doi.org/10.1140/epja/s10050-022-00812-9).

- URL <https://doi.org/10.1140/epja/s10050-022-00812-9>
- [234] M. Ismail, A. Y. Ellithi, A. Adel, M. A. Abbas, An improved unified formula for  $\alpha$ -decay and cluster radioactivity of heavy and superheavy nuclei, The European Physical Journal A 58 (11) (2022) 225. doi:[10.1140/epja/s10050-022-00882-9](https://doi.org/10.1140/epja/s10050-022-00882-9).
- URL <https://doi.org/10.1140/epja/s10050-022-00882-9>
- [235] S. Luo, Y.-Y. Xu, D.-X. Zhu, B. He, P.-C. Chu, X.-H. Li, Improved geiger–nuttall law for  $\alpha$ -decay half-lives of heavy and superheavy nuclei, The European Physical Journal A 58 (12) (2022) 244. doi:[10.1140/epja/s10050-022-00898-1](https://doi.org/10.1140/epja/s10050-022-00898-1).
- URL <https://doi.org/10.1140/epja/s10050-022-00898-1>
- [236] I. S. Rogov, G. G. Adamian, N. V. Antonenko, Dynamics of a dinuclear system in charge-asymmetry coordinates:  $\alpha$  decay, cluster radioactivity, and spontaneous fission, Phys. Rev. C 100 (2019) 024606. doi:[10.1103/PhysRevC.100.024606](https://doi.org/10.1103/PhysRevC.100.024606).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.100.024606>
- [237] I. S. Rogov, G. G. Adamian, N. V. Antonenko, Cluster approach to spontaneous fission of even-even isotopes of u, pu, cm, cf, fm, no, rf, sg, and hs, Phys. Rev. C 104 (2021) 034618. doi:[10.1103/PhysRevC.104.034618](https://doi.org/10.1103/PhysRevC.104.034618).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.104.034618>
- [238] G. G. Adamian, N. V. Antonenko, A. N. Bezbakh, R. V. Jolos, Effect of properties of superheavy nuclei on their production and decay, Physics of Particles and Nuclei 47 (3) (2016) 387–455. doi:[10.1134/S1063779616030023](https://doi.org/10.1134/S1063779616030023).
- URL <https://doi.org/10.1134/S1063779616030023>
- [239] A. Samark-Roth, D. M. Cox, D. Rudolph, L. G. Sarmiento, M. Albertsson, B. G. Carlsson, J. L. Egido, P. Golubev, J. Heery, A. Yakushev, S. Aberg, H. M. Albers, M. Block, H. Brand, T. Calverley, R. Cantemir, R. M. Clark, C. E. Düllmann, J. Eberth, C. Fahlander, U. Forsberg, J. M. Gates, F. Giacoppo, M. Götz, S. Götz, R.-D. Herzberg, Y. Hrabar, E. Jäger, D. Judson, J. Khuyagbaatar, B. Kindler, I. Kojouharov, J. V. Kratz, J. Krier, N. Kurz, L. Lens, J. Ljungberg, B. Lommel, J. Louko, C.-C. Meyer, A. Mistry, C. Mokry, P. Papadakis, E. Parr, J. L. Pore, I. Ragnarsson, J. Runke, M. Schädel, H. Schaffner, B. Schausten, D. A. Shaughnessy, P. Thörle-Pospiech, N. Trautmann, J. Uusitalo, Spectroscopy along flerovium decay chains. iii. details on experiment, analysis,  $^{282}\text{Cn}$ , and spontaneous fission branches, Phys. Rev. C 107 (2023) 024301. doi:[10.1103/PhysRevC.107.024301](https://doi.org/10.1103/PhysRevC.107.024301).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.107.024301>
- [240] D. M. Cox, A. Samark-Roth, D. Rudolph, L. G. Sarmiento, R. M. Clark, J. L. Egido, P. Golubev, J. Heery, A. Yakushev, S. Aberg, H. M. Albers, M. Albertsson, M. Block, H. Brand, T. Calverley, R. Cantemir, B. G. Carlsson, C. E. Düllmann, J. Eberth, C. Fahlander, U. Forsberg, J. M. Gates, F. Giacoppo, M. Götz, S. Götz, R.-D. Herzberg, Y. Hrabar, E. Jäger, D. Judson, J. Khuyagbaatar, B. Kindler, I. Kojouharov, J. V. Kratz, J. Krier, N. Kurz, L. Lens, J. Ljungberg, B. Lommel, J. Louko, C.-C. Meyer, A. Mistry, C. Mokry, P. Papadakis, E. Parr, J. L. Pore, I. Ragnarsson, J. Runke, M. Schädel, H. Schaffner, B. Schausten, D. A. Shaughnessy, P. Thörle-Pospiech, N. Trautmann, J. Uusitalo, Spectroscopy along flerovium decay chains. ii. fine structure in odd- $a$   $^{289}\text{Fl}$ , Phys. Rev. C 107 (2023) L021301. doi:[10.1103/PhysRevC.107.L021301](https://doi.org/10.1103/PhysRevC.107.L021301).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.107.L021301>

- [241] J. M. Gates, R. Orford, D. Rudolph, C. Appleton, B. M. Barrios, J. Y. Benitez, M. Bordeau, W. Botha, C. M. Campbell, J. Chadderton, A. T. Chemey, R. M. Clark, H. L. Crawford, J. D. Despotopoulos, O. Dorvaux, N. E. Esker, P. Fallon, C. M. Folden, B. J. P. Gall, F. H. Garcia, P. Golubev, J. A. Gooding, M. Grebo, K. E. Gregorich, M. Guerrero, R. A. Henderson, R.-D. Herzberg, Y. Hrabar, T. T. King, M. Kireeff Covo, A. S. Kirkland, R. Krücken, E. Leisenschneider, E. M. Lykiardopoulou, M. McCarthy, J. A. Mildon, C. Müller-Gatermann, L. Phair, J. L. Pore, E. Rice, K. P. Rykaczewski, B. N. Sammis, L. G. Sarmiento, D. Seweryniak, D. K. Sharp, A. Sinjari, P. Steinagger, M. A. Stoyer, J. M. Szornel, K. Thomas, D. S. Todd, P. Vo, V. Watson, P. T. Woody, **Toward the discovery of new elements: Production of livermorium ( $z = 116$ ) with  $^{50}\text{Ti}$** , Phys. Rev. Lett. 133 (2024) 172502. doi:[10.1103/PhysRevLett.133.172502](https://doi.org/10.1103/PhysRevLett.133.172502). URL <https://link.aps.org/doi/10.1103/PhysRevLett.133.172502>
- [242] A. N. Kuzmina, G. G. Adamian, N. V. Antonenko, **Role of quasiparticle structure in  $\alpha$  decays of the heaviest nuclei**, Phys. Rev. C 85 (2012) 027308. doi:[10.1103/PhysRevC.85.027308](https://doi.org/10.1103/PhysRevC.85.027308). URL <https://link.aps.org/doi/10.1103/PhysRevC.85.027308>
- [243] N. T. Brewer, V. K. Utyonkov, K. P. Rykaczewski, Y. T. Oganessian, F. S. Abdullin, R. A. Boll, D. J. Dean, S. N. Dmitriev, J. G. Ezold, L. K. Felker, R. K. Grzywacz, M. G. Itkis, N. D. Kovrzhnykh, D. C. McInturff, K. Miernik, G. D. Owen, A. N. Polyakov, A. G. Popeko, J. B. Roberto, A. V. Sabel'nikov, R. N. Sagaidak, I. V. Shirokovsky, M. V. Shumeiko, N. J. Sims, E. H. Smith, V. G. Subbotin, A. M. Sukhov, A. I. Svirikhin, Y. S. Tsyanov, S. M. Van Cleve, A. A. Voinov, G. K. Vostokin, C. S. White, J. H. Hamilton, M. A. Stoyer, **Search for the heaviest atomic nuclei among the products from reactions of mixed-cf with a  $^{48}\text{Ca}$  beam**, Phys. Rev. C 98 (2018) 024317. doi:[10.1103/PhysRevC.98.024317](https://doi.org/10.1103/PhysRevC.98.024317). URL <https://link.aps.org/doi/10.1103/PhysRevC.98.024317>
- [244] J. L. Egido, A. Jungclaus, **Predominance of tri-axial shapes in transitional super-heavy nuclei: Ground-state deformation and shape coexistence along the flerovium ( $z = 114$ ) chain of isotopes**, Phys. Rev. Lett. 125 (2020) 192504. doi:[10.1103/PhysRevLett.125.192504](https://doi.org/10.1103/PhysRevLett.125.192504). URL <https://link.aps.org/doi/10.1103/PhysRevLett.125.192504>
- [245] J. L. Egido, A. Jungclaus, **Low-energy nuclear excitations along the  $\alpha$ -decay chains of superheavy  $^{292}\text{Lv}$  and  $^{294}\text{Og}$** , Phys. Rev. Lett. 126 (2021) 192501. doi:[10.1103/PhysRevLett.126.192501](https://doi.org/10.1103/PhysRevLett.126.192501). URL <https://link.aps.org/doi/10.1103/PhysRevLett.126.192501>
- [246] Y. Oganessian, V. Utyonkov, A. Popeko, D. Solov'yev, F. Abdullin, S. Dmitriev, D. Ibadullayev, M. Itkis, N. Kovrzhnykh, D. Kuznetsov, O. Petrushkin, A. Podshibyakin, A. Polyakov, R. Sagaidak, L. Schlattauer, I. Shirokovsky, V. Shubin, M. Shumeiko, Y. Tsyanov, A. Voinov, V. Subbotin, V. Bekhterev, N. Belykh, O. Chernyshev, K. Gikal, G. Ivanov, A. Khalkin, V. Konstantinov, N. Osipov, S. Paschenko, A. Protasov, V. Semin, V. Sorokoumov, K. Sychev, V. Verevochkin, B. Yakovlev, S. Antoine, W. Beeckman, P. Jehanno, M. Yavor, A. Shcherbakov, K. Rykaczewski, T. King, J. Roberto, N. Brewer, R. Grzywacz, Z. Gan, Z. Zhang, M. Huang, H. Yang, **Dgfrs-2—a gas-filled recoil separator for the dubna super heavy element factory**, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 1033 (2022) 166640. doi:<https://doi.org/10.1016/j.nima.2022.166640>.

- URL <https://www.sciencedirect.com/science/article/pii/S0168900222002078>
- [247] D. Rudolph, U. Forsberg, P. Golubev, L. Sarmiento, A. Yakushev, L.-L. Andersson, A. Di Nitto, C. E. Düllmann, J. M. Gates, K. E. Gregorich, C. J. Gross, R.-D. Herzberg, F. P. Heßberger, J. Khuyagbaatar, J. V. Kratz, K. Rykaczewski, M. Schädel, S. Aberg, D. Ackermann, M. Block, H. Brand, B. G. Carlsson, D. Cox, X. Derkx, K. Eberhardt, J. Even, C. Fahlander, J. Gerl, E. Jäger, B. Kindler, J. Krier, I. Kojouharov, N. Kurz, B. Lommel, A. Mistry, C. Mokry, H. Nitsche, J. P. Omtvedt, P. Papadakis, I. Ragnarsson, J. Runke, H. Schaffner, B. Schausten, P. Thörle-Pospiech, T. Torres, T. Traut, N. Trautmann, A. Türler, A. Ward, D. E. Ward, N. Wiehl, Selected spectroscopic results on element 115 decay chains, *J. Rad. Nuc. Chem.* 303 (2) (2015) 1185–1190.
- [248] U. Forsberg, D. Rudolph, L.-L. Andersson, A. Di Nitto, C. Düllmann, C. Fahlander, J. Gates, P. Golubev, K. Gregorich, C. Gross, R.-D. Herzberg, F. Heßberger, J. Khuyagbaatar, J. Kratz, K. Rykaczewski, L. Sarmiento, M. Schädel, A. Yakushev, S. Aberg, D. Ackermann, M. Block, H. Brand, B. Carlsson, D. Cox, X. Derkx, J. Dobaczewski, K. Eberhardt, J. Even, J. Gerl, E. Jäger, B. Kindler, J. Krier, I. Kojouharov, N. Kurz, B. Lommel, A. Mistry, C. Mokry, W. Nazarewicz, H. Nitsche, J. Omtvedt, P. Papadakis, I. Ragnarsson, J. Runke, H. Schaffner, B. Schausten, Y. Shi, P. Thörle-Pospiech, T. Torres, T. Traut, N. Trautmann, A. Türler, A. Ward, D. Ward, N. Wiehl, Recoil- $\alpha$ -fission and recoil- $\alpha$ - $\alpha$ -fission events observed in the reaction  $^{48}\text{Ca} + ^{243}\text{Am}$ , *Nuclear Physics A* 953 (2016) 117–138. doi:<https://doi.org/10.1016/j.nuclphysa.2016.04.025>. URL <https://www.sciencedirect.com/science/article/pii/S0375947416300768>
- [249] A. M. Rodin, A. V. Belozerov, D. V. Vanin, V. Y. Vedeneyev, A. V. Gulyaev, A. V. Gulyaeva, S. N. Dmitriev, M. G. Itkis, J. Kliman, N. A. Kondratiev, L. Krupa, Y. T. Oganessian, V. S. Salamatin, I. Siváček, S. V. Stepantsov, E. V. Chernysheva, S. A. Yuchimchuk, *Masha separator on the heavy ion beam for determining masses and nuclear physical properties of isotopes of heavy and superheavy elements, Instruments and Experimental Techniques* 57 (4) (2014) 386–393. doi:[10.1134/S0020441214030208](https://doi.org/10.1134/S0020441214030208).
- URL <https://doi.org/10.1134/S0020441214030208>
- [250] F. Déchery, A. Drouart, H. Savajols, J. Nolen, M. Authier, A. M. Amthor, D. Boutin, O. Delferrière, B. Gall, A. Hue, B. Laune, F. Blanc, S. Manikonda, J. Payet, M. H. Stodel, E. Traykov, D. Uriot, *Toward the drip lines and the superheavy island of stability with the Super Separator Spectrometer S3*, *Eur. Phys. J. A* 51 (6) (2015) 1–16. doi:[10.1140/epja/i2015-15066-3](https://doi.org/10.1140/epja/i2015-15066-3).
- URL <http://dx.doi.org/10.1140/epja/i2015-15066-3>
- [251] D. Ackermann, B. Blank, L. Caceres, M. Caamaño, G. De France, B. Gall, S. Grévy, C. Grygiel, E. Lamour, X. Ledoux, V. Manea, C. Peaucelle, J. Rangama, N. Redon, I. Stefan, C. Stodel, B. Sulignano, C. Theisen, *NEWGAIN White Book - Science Requirements - Work Package Physics*, Technical report, GANIL - document link: <https://www.ganil-spiral2.eu/scientists/ganil-spiral-2-facilities/accelerators/newgain/> (Jun. 2021).
- URL <https://hal.archives-ouvertes.fr/hal-03280595>
- [252] F. Kondev, M. Wang, W. Huang, S. Naimi, G. Audi, *The NUBASE2020 evaluation of nuclear physics properties*, *Chinese Physics C* 45 (3) (2021) 030001. doi:[10.1088/1674-1137/abddae](https://doi.org/10.1088/1674-1137/abddae).
- URL <https://doi.org/10.1088/1674-1137/abddae>

- [253] Y. Lazarev, Y. Oganessian, Z. Szeglowki, V. Utyonkov, Y. Kharitonov, O. Constantinescu, D. T. Lién, I. Shirokovsky, S. Tretyakova, [Cross sections of the \(hi,  \$\alpha\$ n\) channel in the cold-fusion-type reactions  \$^{209}\text{Bi}+^{40}\text{Ar}\$  and  \$^{208}\text{Pb}+^{37}\text{Cl}\$](#) , Nuclear Physics A 580 (1) (1994) 113–132. [doi:\[https://doi.org/10.1016/0375-9474\\(94\\)90818-4\]\(https://doi.org/10.1016/0375-9474\(94\)90818-4\)](https://doi.org/10.1016/0375-9474(94)90818-4). URL <https://www.sciencedirect.com/science/article/pii/0375947494908184>
- [254] D. A. Shaughnessy, J. L. Adams, K. E. Gregorich, M. R. Lane, C. A. Laue, D. M. Lee, C. A. McGrath, J. B. Patin, D. A. Strellis, E. R. Sylvester, P. A. Wilk, D. C. Hoffman, [Electron-capture delayed fission properties of  \$^{242}\text{Es}\$](#) , Phys. Rev. C 61 (2000) 044609. [doi:<https://doi.org/10.1103/PhysRevC.61.044609>](https://doi.org/10.1103/PhysRevC.61.044609). URL <https://link.aps.org/doi/10.1103/PhysRevC.61.044609>
- [255] P. Eskola, K. Eskola, M. Nurmia, A. Ghiorso, [Two new isotopes of einsteinium,  \$^{243}\text{es}\$  and  \$^{244}\text{es}\$](#) , Phys. Fenn. 8 (1973) 357.
- [256] Y. P. Gangrsky, M. B. Miller, L. V. Mikhailov, I. F. Kharisov, [Study of delayed fission of bk, es, md isotopes](#), Yad. Fiz. - Sov. J. Nucl. Phys. 31 (1980) 306 – 162.
- [257] J. Khuyagbaatar, S. Hofmann, F. P. Heßberger, D. Ackermann, H. G. Burkhard, S. Heinz, B. Kindler, I. Kojouharov, B. Lommel, R. Mann, J. Maurer, K. Nishio, Y. Novikov, [Spontaneous fission of neutron-deficient fermium isotopes and the new nucleus  \$^{241}\text{fm}\$](#) , The European Physical Journal A 37 (2) (2008) 177–183. [doi:<https://doi.org/10.1140/epja/i2008-10608-4>](https://doi.org/10.1140/epja/i2008-10608-4). URL <https://doi.org/10.1140/epja/i2008-10608-4>
- [258] V. L. Mikheev, V. I. Ilyushchenko, M. B. Miller, [Synthesis of einsteinium isotopes in reactions with nitrogen ions](#), Yadern. Fiz. 5 (1967) 49.
- [259] P. Eskola, [Studies of mendelevium isotopes with mass numbers 248 through 252](#), Phys. Rev. C 7 (1973) 280–289. [doi:<https://doi.org/10.1103/PhysRevC.7.280>](https://doi.org/10.1103/PhysRevC.7.280). URL <https://link.aps.org/doi/10.1103/PhysRevC.7.280>
- [260] A. Ghiorso, S. G. Thompson, G. H. Higgins, B. G. Harvey, G. T. Seaborg, [Evidence for subshell at  \$N = 152\$](#) , Phys. Rev. 95 (1954) 293–295. [doi:<https://doi.org/10.1103/PhysRev.95.293>](https://doi.org/10.1103/PhysRev.95.293). URL <http://link.aps.org/doi/10.1103/PhysRev.95.293>
- [261] D. A. Shaughnessy, K. E. Gregorich, M. R. Lane, C. A. Laue, D. M. Lee, C. A. McGrath, D. A. Strellis, E. R. Sylvester, P. A. Wilk, D. C. Hoffman, [Electron-capture delayed fission probabilities of  \$^{248}\text{Es}\$  and  \$^{246}\text{Es}\$](#) , Phys. Rev. C 63 (2001) 037603. [doi:<https://doi.org/10.1103/PhysRevC.63.037603>](https://doi.org/10.1103/PhysRevC.63.037603). URL <https://link.aps.org/doi/10.1103/PhysRevC.63.037603>
- [262] A. Chetham-Strode, L. W. Holm, [New isotope einsteinium-248](#), Phys. Rev. 104 (1956) 1314–1314. [doi:<https://doi.org/10.1103/PhysRev.104.1314>](https://doi.org/10.1103/PhysRev.104.1314). URL <https://link.aps.org/doi/10.1103/PhysRev.104.1314>
- [263] I. Ahmad, R. K. Sjoblom, R. F. Barnes, E. P. Horwitz, P. R. Fields, [Nuclear decay properties of light einsteinium isotopes](#), Nuclear Physics A 140 (1) (1970) 141–153. [doi:\[https://doi.org/10.1016/0375-9474\\(70\\)90890-0\]\(https://doi.org/10.1016/0375-9474\(70\)90890-0\)](https://doi.org/10.1016/0375-9474(70)90890-0). URL <https://www.sciencedirect.com/science/article/pii/0375947470908900>
- [264] R. M. Chasteler, R. A. Henderson, D. Lee, K. E. Gregorich, M. J. Nurmia, R. B. Welch, D. C. Hoffman, [Excitation functions for production of heavy actinides from interactions of  \$^{16}\text{O}\$  with  \$^{249}\text{Cf}\$](#) , Phys. Rev. C 36 (1987) 1820–1825. [doi:<https://doi.org/10.1103/PhysRevC.36.1820>](https://doi.org/10.1103/PhysRevC.36.1820). URL <https://link.aps.org/doi/10.1103/PhysRevC.36.1820>

- [265] B. G. Harvey, A. Chetham-Strode, A. Ghiorso, G. R. Choppin, S. G. Thompson, New isotopes of einsteinium, Phys. Rev. 104 (1956) 1315–1319. [doi:10.1103/PhysRev.104.1315](https://doi.org/10.1103/PhysRev.104.1315). URL <https://link.aps.org/doi/10.1103/PhysRev.104.1315>
- [266] I. Ahmad, R. K. Sjoblom, P. R. Fields, One- and three-quasiparticle states in  $^{249}\text{Cf}$  populated by the electron capture decay of 1.70-h  $^{249}\text{Es}$ , Phys. Rev. C 14 (1976) 218–228. [doi:10.1103/PhysRevC.14.218](https://doi.org/10.1103/PhysRevC.14.218). URL <https://link.aps.org/doi/10.1103/PhysRevC.14.218>
- [267] A. Gavron, H. C. Britt, P. D. Goldstone, R. Schoenmackers, J. Weber, J. B. Wilhelmy,  $\frac{\Gamma_n}{\Gamma_f}$  in heavy actinides, Phys. Rev. C 15 (1977) 2238–2240. [doi:10.1103/PhysRevC.15.2238](https://doi.org/10.1103/PhysRevC.15.2238). URL <https://link.aps.org/doi/10.1103/PhysRevC.15.2238>
- [268] K. E. Gregorich, K. J. Moody, D. Lee, W. K. Kot, R. B. Welch, P. A. Wilmarth, G. T. Seaborg, Actinide production in  $^{136}\text{Xe}$  bombardments of  $^{249}\text{Cf}$ , Phys. Rev. C 35 (1987) 2117–2124. [doi:10.1103/PhysRevC.35.2117](https://doi.org/10.1103/PhysRevC.35.2117). URL <https://link.aps.org/doi/10.1103/PhysRevC.35.2117>
- [269] R. B. Welch, K. J. Moody, K. E. Gregorich, D. Lee, G. T. Seaborg, Dependence of actinide production on the mass number of the projectile:  $\text{Xe} + ^{248}\text{cm}$ , Phys. Rev. C 35 (1987) 204–212. [doi:10.1103/PhysRevC.35.204](https://doi.org/10.1103/PhysRevC.35.204). URL <https://link.aps.org/doi/10.1103/PhysRevC.35.204>
- [270] M. S. Freedman, I. Ahmad, F. T. Porter, R. K. Sjoblom, R. F. Barnes, J. Lerner, P. R. Fields, Two-quasiparticle states in  $^{250}\text{Cf}$  populated by electron capture decay of 8.6-h  $^{250}\text{Es}$  isomer, Phys. Rev. C 15 (1977) 760–776. [doi:10.1103/PhysRevC.15.760](https://doi.org/10.1103/PhysRevC.15.760). URL <https://link.aps.org/doi/10.1103/PhysRevC.15.760>
- [271] I. Ahmad, R. K. Sjoblom, A. M. Friedman, S. W. Yates, Proton states in the  $Z = 99$  nucleus  $^{251}\text{Es}$  excited by  $^{251}\text{Fm}$  electron capture decay and  $^{250}\text{Cf}(\alpha, t)$  reaction, Phys. Rev. C 17 (1978) 2163–2175. [doi:10.1103/PhysRevC.17.2163](https://doi.org/10.1103/PhysRevC.17.2163). URL <http://link.aps.org/doi/10.1103/PhysRevC.17.2163>
- [272] I. Ahmad, R. R. Chasman, P. R. Fields, Spectroscopic studies beyond the  $n = 152$  neutron gap: Decay of  $^{255}_{101}\text{Md}$  and  $^{256}_{101}\text{Md}$ , Phys. Rev. C 61 (2000) 044301. [doi:10.1103/PhysRevC.61.044301](https://doi.org/10.1103/PhysRevC.61.044301). URL <http://link.aps.org/doi/10.1103/PhysRevC.61.044301>
- [273] P. Fields, I. Ahmad, R. Barnes, R. Sjoblom, W. McHarris, Nuclear spectroscopic studies of  $^{252}\text{Es}$ , Nuclear Physics A 208 (2) (1973) 269 – 286. [doi:\[http://dx.doi.org/10.1016/0375-9474\\(73\\)90375-8\]\(http://dx.doi.org/10.1016/0375-9474\(73\)90375-8\)](https://doi.org/10.1016/0375-9474(73)90375-8). URL <http://www.sciencedirect.com/science/article/pii/0375947473903758>
- [274] I. Ahmad, F. Wagner, Half-life of the longest-lived einsteinium isotope-252es, Journal of Inorganic and Nuclear Chemistry 39 (9) (1977) 1509–1511. [doi:\[https://doi.org/10.1016/0022-1902\\(77\\)80089-4\]\(https://doi.org/10.1016/0022-1902\(77\)80089-4\)](https://doi.org/10.1016/0022-1902(77)80089-4). URL <http://www.sciencedirect.com/science/article/pii/0022190277800894>
- [275] J. V. Kratz, M. Schädel, H. W. Gäggeler, Re-examining the heavy-ion reactions  $^{238}\text{U} + ^{238}\text{U}$  and  $^{238}\text{U} + ^{248}\text{Cm}$  and actinide production close to the barrier, Phys. Rev. C 88 (2013) 054615. [doi:10.1103/PhysRevC.88.054615](https://doi.org/10.1103/PhysRevC.88.054615). URL <http://link.aps.org/doi/10.1103/PhysRevC.88.054615>
- [276] S. G. Thompson, A. Ghiorso, B. G. Harvey, G. R. Choppin, Transcurium isotopes produced in the

- neutron irradiation of plutonium, Phys. Rev. 93 (1954) 908–908. doi:[10.1103/PhysRev.93.908](https://doi.org/10.1103/PhysRev.93.908). URL <https://link.aps.org/doi/10.1103/PhysRev.93.908>
- [277] P. R. Fields, M. H. Studier, J. F. Mech, H. Diamond, A. M. Friedman, L. B. Magnusson, J. R. Huizenga, Additional properties of isotopes of elements 99 and 100, Phys. Rev. 94 (1954) 209–210. doi:[10.1103/PhysRev.94.209](https://doi.org/10.1103/PhysRev.94.209). URL <https://link.aps.org/doi/10.1103/PhysRev.94.209>
- [278] M. Jones, R. P. Schuman, J. P. Butler, G. Cowper, T. A. Eastwood, H. G. Jackson, Isotopes of einsteinium and fermium produced by neutron irradiation of plutonium, Phys. Rev. 102 (1956) 203–207. doi:[10.1103/PhysRev.102.203](https://doi.org/10.1103/PhysRev.102.203). URL <https://link.aps.org/doi/10.1103/PhysRev.102.203>
- [279] D. Metta, H. Diamond, R. F. Barnes, J. Milsted, J. Gray, D. J. Henderson, C. M. Stevens, Nuclear constants of nine transplutonium nuclides, Journal of Inorganic and Nuclear Chemistry 27 (1) (1965) 33–39. doi:[https://doi.org/10.1016/0022-1902\(65\)80187-7](https://doi.org/10.1016/0022-1902(65)80187-7). URL <https://www.sciencedirect.com/science/article/pii/0022190265801877>
- [280] N. Severijns, A. A. Belyaev, A. L. Erzinkyan, P.-D. Eversheim, V. T. Filimonov, V. V. Golovko, G. M. Gurevich, P. Herzog, I. S. Kraev, A. A. Lukhanin, V. I. Noga, V. P. Parfenova, T. Phalet, A. V. Rusakov, Y. G. Toporov, C. Tramm, V. N. Vyachin, D. Zákoucký, E. Zotov, Angular distribution of  $\alpha$  particles from oriented  $^{253,254}\text{Es}$  and  $^{255}\text{Fm}$  nuclei, Phys. Rev. C 71 (2005) 044324. doi:[10.1103/PhysRevC.71.044324](https://doi.org/10.1103/PhysRevC.71.044324). URL <https://link.aps.org/doi/10.1103/PhysRevC.71.044324>
- [281] I. Ahmad, F. G. Kondev, E. F. Moore, M. P. Carpenter, R. R. Chasman, J. P. Greene, R. V. F. Janssens, T. Lauritsen, C. J. Lister, D. Seweryniak, R. W. Hoff, J. E. Evans, R. W. Lougheed, C. E. Porter, L. K. Felker, Energy levels of  $^{249}\text{Bk}$  populated in the  $\alpha$  decay of  $^{253}\text{Es}$  and  $\beta^-$  decay of  $^{249}\text{Cm}$ , Phys. Rev. C 71 (2005) 054305. doi:[10.1103/PhysRevC.71.054305](https://doi.org/10.1103/PhysRevC.71.054305). URL <https://link.aps.org/doi/10.1103/PhysRevC.71.054305>
- [282] N. Severijns, A. A. Belyaev, A. L. Erzinkyan, P.-D. Eversheim, V. T. Filimonov, V. V. Golovko, G. M. Gurevich, P. Herzog, I. S. Kraev, A. A. Lukhanin, V. I. Noga, V. P. Parfenova, T. Phalet, A. V. Rusakov, Y. G. Toporov, C. Tramm, V. N. Vyachin, F. Wauters, D. Zákoucký, E. Zotov,  $\alpha$ -decay half-life of  $^{253}\text{Es}$  in metallic fe at temperatures between 4 k and 50 mk, Phys. Rev. C 76 (2007) 024304. doi:[10.1103/PhysRevC.76.024304](https://doi.org/10.1103/PhysRevC.76.024304). URL <https://link.aps.org/doi/10.1103/PhysRevC.76.024304>
- [283] N. Severijns, A. A. Belyaev, A. L. Erzinkyan, P. D. Eversheim, V. T. Filimonov, V. V. Golovko, G. M. Gurevich, P. Herzog, I. S. Kraev, A. A. Lukhanin, V. I. Noga, V. P. Parfenova, T. Phalet, A. V. Rusakov, M. Tandecki, Y. G. Toporov, C. Tramm, E. Traykov, S. V. Gorp, V. N. Vyachin, F. Wauters, D. Zákoucký, E. Zotov, Hyperfine field of einsteinium in iron and nuclear magnetic moment of  $^{254}\text{Es}$ , Phys. Rev. C 79 (2009) 064322. doi:[10.1103/PhysRevC.79.064322](https://doi.org/10.1103/PhysRevC.79.064322). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.064322>
- [284] F. G. Kondev, I. Ahmad, M. P. Carpenter, C. J. Chiara, J. P. Greene, R. V. F. Janssens, M. A. Kellett, T. L. Khoo, T. Lauritsen, C. J. Lister, E. F. Moore, A. L. Nichols, D. Seweryniak, S. Zhu, Studies of nuclear structure and decay properties of actinide nuclei, AIP Conference Proceedings 1090 (1) (2009) 199–206. arXiv:[https://aip.scitation.org/doi/10.1063/1.3149322](https://arxiv.org/abs/https://aip.scitation.org/doi/10.1063/1.3149322)

- pdf/10.1063/1.3087013, doi:10.1063/1.3087013.  
 URL <https://aip.scitation.org/doi/abs/10.1063/1.3087013>
- [285] S. Nothelfer, T. E. Albrecht-Schönzart, M. Block, P. Chhetri, C. E. Düllmann, J. G. Ezold, V. Gadelshin, A. Gaiser, F. Giacoppo, R. Heinke, T. Kieck, N. Kneip, M. Laatiaoui, C. Mokry, S. Raeder, J. Runke, F. Schneider, J. M. Sperling, D. Studer, P. Thörle-Pospiech, N. Trautmann, F. Weber, K. Wendt, Nuclear structure investigations of  $^{253-255}\text{Es}$  by laser spectroscopy, Phys. Rev. C 105 (2022) L021302. doi:10.1103/PhysRevC.105.L021302.  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.105.L021302>
- [286] G. R. Choppin, S. G. Thompson, A. Ghiorso, B. G. Harvey, Nuclear properties of some isotopes of californium, elements 99 and 100, Phys. Rev. 94 (1954) 1080–1081. doi:10.1103/PhysRev.94.1080.  
 URL <https://link.aps.org/doi/10.1103/PhysRev.94.1080>
- [287] B. G. Harvey, S. G. Thompson, G. R. Choppin, A. Ghiorso, Nuclide 99 $^{254}$ , Phys. Rev. 99 (1955) 337–337. doi:10.1103/PhysRev.99.337.  
 URL <https://link.aps.org/doi/10.1103/PhysRev.99.337>
- [288] R. P. Schuman, T. A. Eastwood, H. G. Jackson, J. P. Butler, The half-life, neutron capture and fission cross-sections of long lived einsteinium-254, Journal of Inorganic and Nuclear Chemistry 6 (1) (1958) 1–2. doi:[https://doi.org/10.1016/0022-1902\(58\)80092-5](https://doi.org/10.1016/0022-1902(58)80092-5).  
 URL <https://www.sciencedirect.com/science/article/pii/0022190258800925>
- [289] L. Phillips, R. C. Gatti, R. Brandt, S. G. Thompson, Spontaneous fission half lives of  $^{254}\text{cf}$ ,  $^{255}\text{fm}$ , and  $^{250}\text{cf}$ , Journal of Inorganic and Nuclear Chemistry 25 (9) (1963) 1085–1087. doi:[https://doi.org/10.1016/0022-1902\(63\)80125-6](https://doi.org/10.1016/0022-1902(63)80125-6).
- [290] W. McHarris, F. S. Stephens, F. Asaro, I. Perlman, Decay scheme of einsteinium-254, Phys. Rev. 144 (1966) 1031–1045. doi:10.1103/PhysRev.144.1031.  
 URL <https://link.aps.org/doi/10.1103/PhysRev.144.1031>
- [291] P. R. Fields, H. Diamond, A. M. Friedman, J. Milsted, J. L. Lerner, R. F. Barnes, R. K. Sjöblom, D. N. Metta, E. P. Horwitz, Some new properties of  $^{254}\text{es}$  and  $^{255}\text{es}$ , Nuclear Physics A 96 (2) (1967) 440–448. doi:[https://doi.org/10.1016/0375-9474\(67\)90725-7](https://doi.org/10.1016/0375-9474(67)90725-7).  
 URL <https://www.sciencedirect.com/science/article/pii/0375947467907257>
- [292] I. Ahmad, J. P. Unik, Half-life of  $^{254}\text{es}$ , Journal of Inorganic and Nuclear Chemistry 37 (9) (1975) 1851–1853. doi:[https://doi.org/10.1016/0022-1902\(75\)80900-6](https://doi.org/10.1016/0022-1902(75)80900-6).  
 URL <https://www.sciencedirect.com/science/article/pii/0022190275809006>
- [293] L. S. Goodman, H. Diamond, H. E. Stanton, Nuclear and atomic moments and hyperfine-structure parameters of  $^{253}\text{Es}$  and  $^{254m}\text{Es}$ , Phys. Rev. A 11 (1975) 499–504. doi:10.1103/PhysRevA.11.499.  
 URL <https://link.aps.org/doi/10.1103/PhysRevA.11.499>
- [294] I. Ahmad, F. G. Kondev, Z. M. Koenig, W. C. McHarris, S. W. Yates, Two-quasiparticle states in  $^{250}\text{Bk}$  studied by decay scheme and transfer reaction spectroscopy, Phys. Rev. C 77 (2008) 054302. doi:10.1103/PhysRevC.77.054302.  
 URL <https://link.aps.org/doi/10.1103/PhysRevC.77.054302>
- [295] H. M. Devaraja, S. Heinz, O. Beliuskina, S. Hofmann, C. Hornung, G. Münenberg, D. Ackermann,

- M. Gupta, Y. K. Gambhir, R. A. Henderson, F. P. Heßberger, A. V. Yeremin, B. Kindler, B. Lommel, J. Maurer, K. J. Moody, K. Nishio, A. G. Popeko, M. A. Stoyer, D. A. Shaughnessy, Population of nuclides with  $z \geq 98$  in multi-nucleon transfer reactions of  $^{48}\text{Ca} + ^{248}\text{Cm}$ , The European Physical Journal A 55 (2) (2019) 25. [doi:10.1140/epja/i2019-12696-3](https://doi.org/10.1140/epja/i2019-12696-3).  
URL <https://doi.org/10.1140/epja/i2019-12696-3>
- [296] J. Unik, P. Day, S. Vandenbosch, The decay scheme of  $39.3\text{ h }^{254}\text{Es}$ , Nuclear Physics 36 (1962) 284–304. [doi:https://doi.org/10.1016/0029-5582\(62\)90452-2](https://doi.org/10.1016/0029-5582(62)90452-2).  
URL <https://www.sciencedirect.com/science/article/pii/0029558262904522>
- [297] I. Ahmad, J. Milsted, R. K. Sjoblom, J. Lerner, P. R. Fields, Alpha decay of  $^{251}\text{Fm}$ , Phys. Rev. C 8 (1973) 737–744. [doi:10.1103/PhysRevC.8.737](https://doi.org/10.1103/PhysRevC.8.737).  
URL <http://link.aps.org/doi/10.1103/PhysRevC.8.737>
- [298] C. R. Group, Nuclear decay properties of heavy nuclides produced in thermonuclear explosions-par and barbel events, Phys. Rev. 148 (1966) 1192–1198. [doi:10.1103/PhysRev.148.1192](https://doi.org/10.1103/PhysRev.148.1192).  
URL <https://link.aps.org/doi/10.1103/PhysRev.148.1192>
- [299] H. C. Britt, D. C. Hoffman, J. van der Plicht, J. B. Wilhelmy, E. Cheifetz, R. J. Dupzyk, R. W. Lougheed, Fission of  $^{255,256}\text{Es}$ ,  $^{255-257}\text{Fm}$ , and  $^{258}\text{Md}$  at moderate excitation energies, Phys. Rev. C 30 (1984) 559–565. [doi:10.1103/PhysRevC.30.559](https://doi.org/10.1103/PhysRevC.30.559).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.30.559>
- [300] G. R. Choppin, B. G. Harvey, S. G. Thompson, A. Ghiorso, Nuclear properties of  $^{100}\text{Es}$ , Phys. Rev. 98 (1955) 1519–1520. [doi:10.1103/PhysRev.98.1519](https://doi.org/10.1103/PhysRev.98.1519)
- URL <https://link.aps.org/doi/10.1103/PhysRev.98.1519>
- [301] H. C. Britt, E. Cheifetz, D. C. Hoffman, J. B. Wilhelmy, R. J. Dupzyk, R. W. Lougheed, Fission barriers for  $^{255}\text{Es}$ ,  $^{256}\text{Es}$ , and  $^{255}\text{Fm}$ , Phys. Rev. C 21 (1980) 761–763. [doi:10.1103/PhysRevC.21.761](https://doi.org/10.1103/PhysRevC.21.761).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.21.761>
- [302] R. Lougheed, E. Hulet, J. Wild, B. Qualheim, J. Evans, R. Dupzyk, Two new isotopes with  $n = 157$ :  $^{256}\text{es}$  and  $^{255}\text{cf}$ , Jour. Inorg. Nucl. Chem. 43 (10) (1981) 2239–2241. [doi:https://doi.org/10.1016/0022-1902\(81\)80239-4](https://doi.org/10.1016/0022-1902(81)80239-4).  
URL <https://www.sciencedirect.com/science/article/pii/0022190281802394>
- [303] L. Ghys, A. N. Andreyev, S. Antalic, M. Huyse, P. Van Duppen, Empirical description of  $\beta$ -delayed fission partial half-lives, Phys. Rev. C 91 (2015) 044314. [doi:10.1103/PhysRevC.91.044314](https://doi.org/10.1103/PhysRevC.91.044314).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.91.044314>
- [304] R. Lougheed, J. Landrum, D. Hoffman, W. Daniels, J. Wilhelmy, M. Bunker, J. Starner, S. Jackson, Discovery of a 7.6-hour high-spin isomer of einsteinium-256, in: 3rd INTERNATIONAL CONFERENCE ON NUCLEI FAR FROM STABILITY - Proceedings, 1976, pp. 563–564.
- [305] Y. Popov, G. Timofeev, V.B.Mishenev, V.N.Kovantsev, A.A.Elesin, Determining the half-lives of  $^{253}\text{es}$ ,  $^{254}\text{es}$ ,  $^{254}\text{mes}$ ,  $^{255}\text{es}$ ,  $^{257}\text{es}$ ,  $^{256}\text{fm}$ , Sov. J. Radiochem. 29 (1987) 431.
- [306] G. Ter-Akopyan, A. Iljinov, Y. Oganessian, O. Orlova, G. Popeko, S. Tretyakova, V. Chepigin, B. Shilov, G. Flerov, Synthesis of the new neutron-deficient isotopes  $^{250}\text{102}$ ,  $^{242}\text{fm}$ , and  $^{254}\text{ku}$ , Nucl. Phys. A 255 (2) (1975) 509–522. [doi:https://doi.org/10.1016/0375-9474\(75\)90696-X](https://doi.org/10.1016/0375-9474(75)90696-X).

- URL <https://www.sciencedirect.com/science/article/pii/037594747590696X>
- [307] N. Holden, D. Hoffman, Spontaneous fission half-lives for ground-state nuclide, Pure Appl. Chem. 72 (8) (2000) 1525–1562.
- [308] G. Münzenberg, S. Hofmann, W. Faust, F. P. Heßberger, W. Reisdorf, K. H. Schmidt, T. Kitahara, P. Armbruster, K. Güttner, B. Thuma, D. Vermeulen, The new isotopes  $^{247}\text{md}$ ,  $^{243}\text{fm}$ ,  $^{239}\text{cf}$ , and investigation of the evaporation residues from fusion of  $^{206}\text{pb}$ ,  $^{208}\text{pb}$ , and  $^{209}\text{bi}$  with  $^{40}\text{ar}$ , Zeitschrift für Physik A Atoms and Nuclei 302 (1) (1981) 7–14. doi:[10.1007/BF01425097](https://doi.org/10.1007/BF01425097).  
URL <https://doi.org/10.1007/BF01425097>
- [309] H. Gäggeler, T. Sikkeland, G. Wirth, W. Brüchle, W. Bögl, G. Franz, G. Herrmann, J. V. Kratz, M. Schädel, K. Sümmerer, W. Weber, Probing sub-barrier fusion and extra-push by measuring fermium evaporation residues in different heavy ion reactions, Zeitschrift für Physik A Atoms and Nuclei 316 (3) (1984) 291–307. doi:[10.1007/BF01439902](https://doi.org/10.1007/BF01439902).  
URL <https://doi.org/10.1007/BF01439902>
- [310] Y. Oganessian, A. Iljinov, A. Demin, S. Tretyakova, Experiments on the production of fermium neutron-deficient isotopes and new possibilities of synthesizing elements with  $z > 100$ , Nuclear Physics A 239 (2) (1975) 353–364. doi:[https://doi.org/10.1016/0375-9474\(75\)90456-X](https://doi.org/10.1016/0375-9474(75)90456-X).  
URL <https://www.sciencedirect.com/science/article/pii/037594747590456X>
- [311] H. Gaeggeler, A. S. Iljinov, G. S. Popeko, W. Seidel, G. M. Ter-Akopian, S. P. Tretyakova, A study of fusion reactions between  $^{206}$ ,  $^{207}\text{pb}$  nuclei and  $^{40}\text{ar}$  ions near the coulomb barrier, Zeitschrift für Physik A Atoms and Nuclei 289 (4) (1979) 415–420. doi:[10.1007/BF01409394](https://doi.org/10.1007/BF01409394).  
URL <https://doi.org/10.1007/BF01409394>
- [312] A. S. Iljinov, Y. T. Oganessian, E. A. Cherepanov, A. G. Demin, S. P. Tretyakova, V. K. Utyonkov, I. V. Shirokovsky, An energy shift between the fusion barrier and interaction barrier in heavy nuclear systems, Il Nuovo Cimento A (1965-1970) 101 (2) (1989) 225–240. doi:[10.1007/BF02813994](https://doi.org/10.1007/BF02813994).  
URL <https://doi.org/10.1007/BF02813994>
- [313] Svirikhin, Alexandr, Andreev, Alexander, Dushin, Victor, Chelnokov, Maxim, Chepigin, Victor, Gupta, Mohini, Isaev, Andrey, Izosimov, Igor, Katrasev, Denis, Kuznetsov, Alexey, Malyshev, Oleg, Minkova, Ani, Mullins, Simon, Popeko, Andrey, Sokol, Evgeniy, Yeremin, Alexander, The investigation of properties of short-lived sf isotopes ( $z > 100$ ) at the focal plane of vassilissa separator, EPJ Web of Conferences 62 (2013) 03005. doi:[10.1051/epjconf/20136203005](https://doi.org/10.1051/epjconf/20136203005).  
URL <https://doi.org/10.1051/epjconf/20136203005>
- [314] R. S. Mukhin, A. V. Isaev, I. I. Ulanova, A. I. Svirikhin, M. A. Bychkov, M. L. Chelnokov, V. I. Chepigin, H. M. Devaraja, O. Dorvaux, B. Gall, K. Hauschild, I. N. Izosimov, A. A. Kuznetsova, A. Lopez-Martens, O. N. Malyshev, A. G. Popeko, Y. A. Popov, B. Sailaubekov, E. A. Sokol, M. S. Tezekbayeva, N. I. Zamyatin, A. V. Yeremin, Prompt neutron multiplicity from spontaneous fission of  $^{244}\text{fm}$ , The European Physical Journal A 60 (11) (2024) 223. doi:[10.1140/epja/s10050-024-01441-0](https://doi.org/10.1140/epja/s10050-024-01441-0).  
URL <https://doi.org/10.1140/epja/s10050-024-01441-0>
- [315] H. Haba, T. Akiyama, D. Kaji, H. Kikunaga, T. Kurabayashi, K. Morimoto, K. Morita, K. Ooe, N. Sato, A. Shinohara, T. Takabe, Y. Tashiro, A. Toyoshima, A. Yoneda, T. Yoshimura, Startup of superheavy element chemistry at riken, The European Physical Journal D 45 (1) (2007) 81–86. doi:[10.1140/epjd/e2007-00135-4](https://doi.org/10.1140/epjd/e2007-00135-4).

- URL <https://doi.org/10.1140/epjd/e2007-00135-4>
- [316] K. Rezynkina, K. Hauschild, A. L.-M. O. Dorvaux, B. Gall, F. Déchery, H. Faure, A. Yeremin, M. Chelnokov, V. Chepigin, A. I. Izosimov, D. Katrasev, A. K. Kuznetsova, O. Malyshev, A. Popeko, Y. P. Sokol, A. SvirikhinFLNR, J. Piot, J. Ruber, First experimental tests of shels: a new heavy ion separator at the jinr, *Acta Phys. Pol. B* 46 (2015) 623.
- [317] M. S. Tezekbayeva, A. V. Yeremin, A. I. Svirikhin, A. Lopez-Martens, M. L. Chelnokov, V. I. Chepigin, A. V. Isaev, I. N. Izosimov, A. V. Karpov, A. A. Kuznetsova, O. N. Malyshev, R. S. Mukhin, A. G. Popeko, Y. A. Popov, V. A. Rachkov, B. S. Sailaubekov, E. A. Sokol, K. Hauschild, H. Jacob, R. Chakma, O. Dorvaux, M. Forge, B. Gall, K. Kessaci, B. Andel, S. Antalic, A. Broenis, P. Mosat, *Study of the production and decay properties of neutron-deficient nobelium isotopes*, *The European Physical Journal A* 58 (3) (2022) 52. doi:[10.1140/epja/s10050-022-00707-9](https://doi.org/10.1140/epja/s10050-022-00707-9).
- URL <https://doi.org/10.1140/epja/s10050-022-00707-9>
- [318] G.N.Akapev, A.G.Demina, V.A.Druin, E.G.Imaev, I.V.Kolesov, Y.V.Lobanov, L.P.Pashchenko, Synthesis of several isotopes of fermium and determination of their radioactive properties, *Soviet J.At.Energy* 21 (1966) 908.
- [319] G.N.Flerov, S.M.Polikanov, V.L.Mikheev, V.I.Ilyushchenko, M.B.Miller, V.A.Shchegolev, Synthesis of isotopes of fermium with mass numbers 247 and 246, *Sov.At.Energy* 22 (1967) 434.
- [320] D. Bogdanov, M. Ivanov, G. Popeko, A. Rodin, G. Ter-Akopian, V. Vakatov, A. Voronin, *The spontaneous fission of 224fm and 246fm*, *Physics Letters B* 113 (3) (1982) 213–214. doi:[https://doi.org/10.1016/0370-2693\(82\)90823-1](https://doi.org/10.1016/0370-2693(82)90823-1).
- URL <https://www.sciencedirect.com/science/article/pii/0370269382908231>
- [321] R. N. Sagaidak, V. I. Chepigin, A. P. Kabachenko, J. Roháč, Y. T. Oganessian, A. G. Popeko, A. V. Yeremin, A. D'Arrigo, G. Fazio, G. Giardina, M. Herman, R. Ruggeri, R. Sturiale, *Fission-evaporation competition in excited uranium and fermium nuclei*, *Journal of Physics G: Nuclear and Particle Physics* 24 (3) (1998) 611–625. doi:[10.1088/0954-3899/24/3/012](https://doi.org/10.1088/0954-3899/24/3/012).
- URL <https://doi.org/10.1088/0954-3899/24/3/012>
- [322] A. I. Svirikhin, V. N. Dushin, M. L. Chelnokov, V. I. Chepigin, I. N. Izosimov, D. E. Katrasev, O. N. Malyshev, A. Minkova, A. G. Popeko, E. A. Sokol, A. V. Yeremin, *Neutron multiplicity at spontaneous fission of 246fm*, *The European Physical Journal A* 44 (3) (2010) 393–396. doi:[10.1140/epja/i2010-10968-0](https://doi.org/10.1140/epja/i2010-10968-0).
- URL <https://doi.org/10.1140/epja/i2010-10968-0>
- [323] M. Venhart, F. P. Heßberger, D. Ackermann, S. Antalic, C. Gray-Jones, P. T. Greenlees, S. Heinz, R. D. Herzberg, S. Hofmann, S. Ketelhut, B. Kindler, I. Kojouharov, M. Leino, B. Lommel, R. Mann, P. Papadakis, D. Rostron, D. Rudolph, Š. Šáro, B. Sulignano, *Decay study of 246fm at ship*, *The European Physical Journal A* 47 (2) (2011) 20. doi:[10.1140/epja/i2011-11020-9](https://doi.org/10.1140/epja/i2011-11020-9).
- URL <https://doi.org/10.1140/epja/i2011-11020-9>
- [324] J. Piot, B. J.-P. Gall, O. Dorvaux, P. T. Greenlees, N. Rowley, L. L. Andersson, D. M. Cox, F. Dechery, T. Grahn, K. Hauschild, G. Henning, A. Herzan, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, P. Jones, R. Julin, S. Juutinen, S. Ketelhut, T.-L. Khoo, M. Leino, J. Ljungvall, A. Lopez-Martens, P. Nieminen, J. Pakarinen, P. Papadakis, E. Parr, P. Peura, P. Rahkila, S. Rinta-Antila, J. Rubert,

- P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, D. Seweryniak, J. Sorri, B. Sulignano, J. Uusitalo, In-beam spectroscopy with intense ion beams: Evidence for a rotational structure in  $^{246}\text{Fm}$ , Phys. Rev. C 85 (2012) 041301. doi:[10.1103/PhysRevC.85.041301](https://doi.org/10.1103/PhysRevC.85.041301). URL <http://link.aps.org/doi/10.1103/PhysRevC.85.041301>
- [325] F. Heßberger, S. Hofmann, G. Münzenberg, K.-H. Schmidt, P. Armbruster, R. Hingmann, Influence of energy summing of  $\alpha$ -particles and conversion electrons on the shapes of  $\alpha$ -spectra in the region of the heaviest nuclei, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 274 (3) (1989) 522–527. doi:[https://doi.org/10.1016/0168-9002\(89\)90184-8](https://doi.org/10.1016/0168-9002(89)90184-8). URL <https://www.sciencedirect.com/science/article/pii/0168900289901848>
- [326] F. P. Heßberger, S. Hofmann, D. Ackermann, P. Cagarda, R. D. Herzberg, I. Kojouharov, P. Kuusiniemi, M. Leino, R. Mann, Alpha-gamma decay studies of  $^{251}$ ,  $^{253}\text{no}$  and their daughter products  $^{247}$ ,  $^{249}\text{fm}$ , The European Physical Journal A - Hadrons and Nuclei 22 (3) (2004) 417–427. doi:[10.1140/epja/i2003-10238-4](https://doi.org/10.1140/epja/i2003-10238-4). URL <https://doi.org/10.1140/epja/i2003-10238-4>
- [327] C. M. Folden III, I. Dragojević, C. E. Düllmann, R. Eichler, M. A. Garcia, J. M. Gates, S. L. Nelson, R. Sudowe, K. E. Gregorich, D. C. Hoffman, H. Nitsche, Measurement of the  $^{208}\text{Pb}(^{52}\text{Cr}, n)^{259}\text{Sg}$  excitation function, Phys. Rev. C 79 (2009) 027602. doi:[10.1103/PhysRevC.79.027602](https://doi.org/10.1103/PhysRevC.79.027602). URL <http://link.aps.org/doi/10.1103/PhysRevC.79.027602>
- [328] I. Dragojević, K. E. Gregorich, C. E. Düllmann, J. Dvorak, P. A. Ellison, J. M. Gates, S. L. Nelson, L. Stavsetra, H. Nitsche, New isotope  $^{263}\text{Hs}$ , Phys. Rev. C 79 (2009) 011602. doi:[10.1103/PhysRevC.79.011602](https://doi.org/10.1103/PhysRevC.79.011602). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.011602>
- [329] A. Ghiorso, T. Sikkeland, J. R. Walton, G. T. Seaborg, Element no. 102, Phys. Rev. Lett. 1 (1958) 18–21. doi:[10.1103/PhysRevLett.1.18](https://doi.org/10.1103/PhysRevLett.1.18). URL <https://link.aps.org/doi/10.1103/PhysRevLett.1.18>
- [330] A. N. Andreyev, D. D. Bogdanov, V. I. Chepigin, A. P. Kabachenko, O. N. Malyshev, R. N. Sagaidak, L. I. Salamatin, G. M. Ter-Akopian, A. V. Yeremin, Cross sections of 102 element isotopes formation in the reactions of  $^{22}\text{ne} + ^{236}\text{u}$  and  $^{26}\text{mg} + ^{232}\text{th}$ , Zeitschrift für Physik A Hadrons and Nuclei 345 (4) (1993) 389–394. doi:[10.1007/BF01282899](https://doi.org/10.1007/BF01282899). URL <https://doi.org/10.1007/BF01282899>
- [331] P. T. Greenlees, N. Amzal, J. E. Bastin, E. Bouchez, P. A. Butler, A. Chatillon, O. Dorvaux, S. Eeckhaudt, K. Eskola, B. Gall, J. Gerl, T. Grahn, A. Görgen, N. J. Hammond, K. Hauschild, R. D. Herzberg, F. P. Heßberger, R. D. Humphreys, A. Hürstel, D. G. Jenkins, G. D. Jones, P. Jones, R. Julin, S. Juutinen, H. Kankaanpää, A. Keenan, H. Kettunen, F. Khalfallah, T. L. Khoo, W. Korten, P. Kuusiniemi, Y. Le Coz, M. Leino, A. P. Leppänen, M. Muikku, P. Nieminen, J. Pakarinen, P. Rahkila, P. Reiter, M. Rousseau, C. Scholey, C. Theisen, J. Uusitalo, J. Wilson, H. J. Wollersheim, In-beam and decay spectroscopy of trans fermium elements, The European Physical Journal A - Hadrons and Nuclei 25 (1) (2005) 599–604. doi:[10.1140/epjad/i2005-06-026-0](https://doi.org/10.1140/epjad/i2005-06-026-0). URL <https://doi.org/10.1140/epjad/i2005-06-026-0>
- [332] K. Nishio, S. Hofmann, F. P. Heßberger, D. Ackermann, S. Antalic, V. F. Comas, Z. Gan, S. Heinz, J. A. Heredia, H. Ikezoe, J. Khuyagbaatar, B. Kindler, I. Kojouharov, P. Kuusiniemi, B. Lommel, R. Mann,

- M. Mazzocco, S. Mitsuoka, Y. Nagame, T. Ohtsuki, A. G. Popeko, S. Saro, H. J. Schött, B. Sulignano, A. Svirikhin, K. Tsukada, K. Tsuruta, A. V. Yeremin, [Measurement of evaporation residue cross-sections of the reaction  \$^{30}\text{Si} + ^{238}\text{U}\$  at sub-barrier energies](#), The European Physical Journal A - Hadrons and Nuclei 29 (3) (2006) 281–287. doi:[10.1140/epja/i2006-10091-y](https://doi.org/10.1140/epja/i2006-10091-y).
- URL <https://doi.org/10.1140/epja/i2006-10091-y>
- [333] A. P. Leppänen, J. Uusitalo, P. T. Greenlees, R. D. Herzberg, N. Amzal, F. Becker, P. A. Butler, A. J. C. Chewter, J. F. C. Cocks, O. Dorvaux, S. Eeckhaudt, K. Eskola, J. Gerl, T. Grahn, N. J. Hammond, K. Hauschild, K. Helariutta, F. P. Heßberger, M. Houry, G. D. Jones, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, T. L. Khoo, W. Korten, P. Kuusiniemi, Y. Le Coz, M. Leino, C. J. Lister, R. Lucas, M. Muikku, P. Nieminen, M. Nyman, R. D. Page, J. Pakarinen, P. Rahkila, P. Reiter, J. Sarén, C. Schlegel, C. Scholey, O. Stezowski, C. Theisen, W. H. Trzaska, H. J. Wollersheim, [Recoil-fission tagging of the transfermium nucleus  \$^{252}\text{No}\$](#) , The European Physical Journal A - Hadrons and Nuclei 28 (3) (2006) 301–306. doi:[10.1140/epja/i2006-10056-2](https://doi.org/10.1140/epja/i2006-10056-2).
- URL <https://doi.org/10.1140/epja/i2006-10056-2>
- [334] J. M. Gates, C. E. Düllmann, M. Schädel, A. Yakushev, A. Türler, K. Eberhardt, J. V. Kratz, D. Ackermann, L.-L. Andersson, M. Block, W. Brüchle, J. Dvorak, H. G. Essel, P. A. Ellison, J. Even, U. Forsberg, J. Gellanki, A. Gorshkov, R. Graeger, K. E. Gregorich, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. Hild, A. Hübler, E. Jäger, J. Khuyagbaatar, B. Kindler, J. Krier, N. Kurz, S. Lahiri, D. Liebe, B. Lommel, M. Maiti, H. Nitsche, J. P. Omtvedt, E. Parr, D. Rudolph, J. Runke, H. Schaffner, B. Schausten, E. Schimpf, A. Semchenkov, J. Steiner, P. Thörle-Pospiech, J. Uusitalo, M. Wegrzecki, N. Wiehl, [First superheavy element experiments at the gsi recoil separator tasca: The production and decay of element 114 in the  \$^{244}\text{Pu}\(^{48}\text{Ca},3\text{-}4n\)\$  reaction](#), Phys. Rev. C 83 (2011) 054618. doi:[10.1103/PhysRevC.83.054618](https://doi.org/10.1103/PhysRevC.83.054618).
- URL <http://link.aps.org/doi/10.1103/PhysRevC.83.054618>
- [335] V. Perelygin, E. Donets, G. Flerov, Experiments in the production of a new fermium isotope, Soviet Phys. JETP 10 (1960) 1106.
- [336] A. Lopez-Martens, K. Hauschild, A. V. Yeremin, A. V. Belozerov, C. Briançon, M. L. Chelnokov, V. I. Chepigin, D. Curien, O. Dorvaux, B. Gall, V. A. Gorshkov, M. Guttormsen, F. Hanappe, A. P. Kabachenko, F. Khalfallah, A. Korichi, A. C. Larsen, O. N. Malyshev, A. Minkova, Y. T. Oganessian, A. G. Popeko, M. Rousseau, N. Rowley, R. N. Sagaidak, S. Sharo, A. V. Shutov, S. Siem, A. I. Svirikhin, N. U. H. Syed, C. Theisen, [Detailed spectroscopy of  \$^{249}\text{Fm}\$](#) , Phys. Rev. C 74 (2006) 044303. doi:[10.1103/PhysRevC.74.044303](https://doi.org/10.1103/PhysRevC.74.044303).
- URL <https://doi.org/10.1103/PhysRevC.74.044303>
- [337] D. Ackermann, A. Mistry, F. Heßberger, B. Andel, S. Antalic, M. Block, L. Caceres, P. Chhetri, F. Dechery, C. Droese, C. Düllmann, F. Giacoppo, J. Hoffmann, O. Kaleja, O. Kamalou, N. Kurz, J. Khuyagbaatar, R. Mändl, T. Murböck, M. Laatiaoui, J. Maurer, P. Mosat, J. Piot, S. Raeder, H. Savajols, O. Sorlin, C. Stodel, J. Thomas, M. Vostinar, P. Wieczorek, A. Yakushev, Z. Zhang, [Compass—a compact decay spectroscopy set-up](#), Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 907 (2018) 81–89, advances in Instrumentation and Experimental Methods (Special Issue in Honour of Kai Siegbahn). doi:

- <https://doi.org/10.1016/j.nima.2018.01.096>. URL <https://www.sciencedirect.com/science/article/pii/S0168900218301438>
- [338] H. Atterling, W. Forsling, L. W. Holm, L. Melander, B. Aström, Element 100 produced by means of cyclotron-accelerated oxygen ions, Phys. Rev. 95 (1954) 585–586. doi:[10.1103/PhysRev.95.585.2](https://doi.org/10.1103/PhysRev.95.585.2). URL <https://link.aps.org/doi/10.1103/PhysRev.95.585.2>
- [339] S. Amiel, A. Chetham-Strode, G. R. Choppin, A. Ghiorso, B. G. Harvey, L. W. Holm, S. G. Thompson, Production and properties of the nuclides fermium-250, 251, and 252, Phys. Rev. 106 (1957) 553–554. doi:[10.1103/PhysRev.106.553](https://doi.org/10.1103/PhysRev.106.553). URL <https://link.aps.org/doi/10.1103/PhysRev.106.553>
- [340] A. Ghiorso, M. Nurmia, J. Harris, K. Eskola, P. Eskola, Positive identification of two alpha-particle-emitting isotopes of element 104, Phys. Rev. Lett. 22 (1969) 1317–1320. doi:[10.1103/PhysRevLett.22.1317](https://doi.org/10.1103/PhysRevLett.22.1317). URL <https://link.aps.org/doi/10.1103/PhysRevLett.22.1317>
- [341] A. Ghiorso, M. Nurmia, J. Harris, K. Eskola, P. Eskola, Defence of the berkeley work on alpha-emitting isotopes of element 104, Nature 229 (5287) (1971) 603–607. doi:[10.1038/229603a0](https://doi.org/10.1038/229603a0). URL <https://doi.org/10.1038/229603a0>
- [342] Y. A. Lazarev, Y. V. Lobanov, R. N. Sagaidak, V. K. Utyonkov, M. Hussonnois, Y. P. Kharitonov, I. V. Shirokovsky, S. P. Tretyakova, Y. T. Oganessian, Study of the stability of the ground states and K-isomeric states of  $^{250}\text{fm}$  and  $^{254}\text{I02}$  against spontaneous fission, Physica Scripta 39 (4) (1989) 422–435. doi:[10.1088/0031-8949/39/4/002](https://doi.org/10.1088/0031-8949/39/4/002). URL <https://doi.org/10.1088/0031-8949/39/4/002>
- [343] M. Leino, H. Kankaanpää, R.-D. Herzberg, A. Chewter, F. P. Heßberger, Y. Le Coz, F. Becker, P. Butler, J. Cocks, O. Dorvaux, K. Eskola, J. Gerl, P. Greenlees, K. Helariutta, M. Houry, G. Jones, P. Jones, R. Julin, S. Juutinen, H. Kettunen, T. Khoo, A. Kleinböhl, W. Korten, P. Kuusiniemi, R. Lucas, M. Muikku, P. Nieminen, R. Page, P. Rahkila, P. Reiter, A. Savelius, C. Schlegel, C. Theisen, W. Trzaska, H.-J. Wollersheim, In-beam study of  $^{254}\text{No}$ , The European Physical Journal A - Hadrons and Nuclei 6 (1) (1999) 63–69. doi:[10.1007/s100500050318](https://doi.org/10.1007/s100500050318). URL <http://dx.doi.org/10.1007/s100500050318>
- [344] K. Nishio, H. Ikezoe, Y. Nagame, M. Asai, K. Tsukada, S. Mitsuoka, K. Tsuruta, K. Satou, C. J. Lin, T. Ohsawa, Evidence of complete fusion in the sub-barrier  $^{16}\text{O} + ^{238}\text{U}$  reaction, Phys. Rev. Lett. 93 (2004) 162701. doi:[10.1103/PhysRevLett.93.162701](https://doi.org/10.1103/PhysRevLett.93.162701). URL <https://link.aps.org/doi/10.1103/PhysRevLett.93.162701>
- [345] C. M. Folden III, S. L. Nelson, C. E. Düllmann, J. M. Schwantes, R. Sudowe, P. M. Zielinski, K. E. Gregorich, H. Nitsche, D. C. Hoffman, Excitation function for the production of  $^{262}\text{Bh}$  ( $z = 107$ ) in the odd-z-projectile reaction  $^{208}\text{Pb}(^{55}\text{Mn}, n)$ , Phys. Rev. C 73 (2006) 014611. doi:[10.1103/PhysRevC.73.014611](https://doi.org/10.1103/PhysRevC.73.014611). URL <https://link.aps.org/doi/10.1103/PhysRevC.73.014611>
- [346] J. E. Bastin, R.-D. Herzberg, P. A. Butler, G. D. Jones, R. D. Page, D. G. Jenkins, N. Amzal, P. M. T. Brew, N. J. Hammond, R. D. Humphreys, P. J. C. Ikin, T. Page, P. T. Greenlees, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, A. Keenan, H. Kettunen, P. Kuusiniemi, M. Leino, A. P. Leppänen, M. Muikku, P. Nieminen, P. Rahkila, C. Scholey, J. Uusitalo, E. Bouchez, A. Chatillon, A. Hürstel, W. Korten, Y. L. Coz, C. Theisen, D. Ackermann, J. Gerl, K. Helariutta, F. P. Hessberger, C. Schlegel,

- H. J. Wollersheim, M. Lach, A. Maj, W. Meczynski, J. Styczen, T. L. Khoo, C. J. Lister, A. V. Afanasjev, H. J. Maier, P. Reiter, P. Bednarczyc, K. Eskola, K. Hauschild, In-beam gamma ray and conversion electron study of  $^{250}\text{Fm}$ , Phys. Rev. C 73 (2006) 024308. doi:[10.1103/PhysRevC.73.024308](https://doi.org/10.1103/PhysRevC.73.024308). URL <http://link.aps.org/doi/10.1103/PhysRevC.73.024308>
- [347] A. Robinson, T. Khoo, I. Ahmad, S. Tandel, F. Kondev, T. Nakasukasa, D. Seweryniak, M. Asai, B. Back, M. Carpenter, P. Choudhury, C. Davids, S. Eeckhaudt, J. Greene, P. Greenlees, S. Gros, A. Heinz, R.-D. Herzberg, R. Janssens, G. Jones, T. Lauritsen, C. Lister, D. Peterson, J. Qian, U. Tandel, X. Wang, S. Zhu,  $K^\pi = 8^-$  isomers and  $k^\pi = 2^-$  octupole vibrations in  $n = 150$  shell-stabilized isotones, Phys. Rev. C 78 (2008) 034308.
- [348] S. L. Nelson, K. E. Gregorich, I. Dragojević, J. Dvořák, P. A. Ellison, M. A. Garcia, J. M. Gates, L. Stavsetra, M. N. Ali, H. Nitsche, Comparison of complementary reactions in the production of mt, Phys. Rev. C 79 (2009) 027605. doi:[10.1103/PhysRevC.79.027605](https://doi.org/10.1103/PhysRevC.79.027605). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.027605>
- [349] M. S. Tezekbayeva, A. V. Yeremin, O. N. Malyshev, A. V. Isaev, R. S. Mukhin, A. A. Kuznetsova, A. G. Popeko, Y. A. Popov, A. I. Svirikhin, E. A. Sokol, M. L. Chelnokov, V. I. Chepigin, A. Lopez-Martens, K. Hauschild, O. Dorvaux, B. Gall, B. S. Sailaubekov, Spectroscopy of the decay properties of transfermium isotopes in dubna, Bulletin of the Russian Academy of Sciences: Physics 85 (10) (2021) 1167–1171. doi:[10.3103/S1062873821100257](https://doi.org/10.3103/S1062873821100257). URL <https://doi.org/10.3103/S1062873821100257>
- [350] A. Ghiorso, K. Eskola, P. Eskola, M. Nurmia, Isomeric states in  $^{250}\text{Fm}$  and  $^{254}\text{No}$ , Phys. Rev. C 7 (1973) 2032–2036. doi:[10.1103/PhysRevC.7.2032](https://doi.org/10.1103/PhysRevC.7.2032). URL <https://link.aps.org/doi/10.1103/PhysRevC.7.2032>
- [351] A. G. Popeko, A. V. Belozerov, C. Briançon, V. I. Chepigin, O. Dorvaux, K. Hauschild, A. P. Kabachenko, A. Korichi, A. Lopez-Martens, O. N. Malyshev, Y. T. Oganessian, Š. Šáro, A. V. Shutov, A. I. Svirikhin, A. V. Yeremin, Gabriela setup for nuclear spectroscopy of the transfermium element isotopes at the vassilissa separator, Physics of Atomic Nuclei 69 (7) (2006) 1183–1187. doi:[10.1134/S1063778806070143](https://doi.org/10.1134/S1063778806070143). URL <https://doi.org/10.1134/S1063778806070143>
- [352] A. M. Friedman, J. E. Gindler, R. F. Barnes, R. Sjoblom, P. R. Fields, Properties of fm $^{252}$ , Phys. Rev. 102 (1956) 585–586. doi:[10.1103/PhysRev.102.585](https://doi.org/10.1103/PhysRev.102.585). URL <https://link.aps.org/doi/10.1103/PhysRev.102.585>
- [353] E. Donets, V. Shchegolev, V. Ermakov, Reactions involving evaporation of several neutrons on bombardment of u238 by accelerated ions of o18, f19, and ne22, Soviet J. Nucl. Phys. 2 (1966) 723.
- [354] J. Chwaszczecka, K. Gavrilov, W. Kusch, M. Szymczak, Some decay properties of 251fm and 252fm, Physics Letters B 25 (5) (1967) 331–332. doi:[https://doi.org/10.1016/0370-2693\(67\)90085-8](https://doi.org/10.1016/0370-2693(67)90085-8). URL <https://www.sciencedirect.com/science/article/pii/0370269367900858>
- [355] I. Ahmad, J. Lerner, Alpha-decay properties of 247cf, 248cf, 252fm and 254fm, Nuclear Physics A 413 (3) (1984) 423–431. doi:[https://doi.org/10.1016/0375-9474\(84\)90419-6](https://doi.org/10.1016/0375-9474(84)90419-6). URL <https://www.sciencedirect.com/science/article/pii/0375947484904196>
- [356] S. Amiel, Properties of fermium-253, Phys. Rev. 105

- (1957) 1412–1413. doi:[10.1103/PhysRev.105.1412](https://doi.org/10.1103/PhysRev.105.1412).  
URL <https://link.aps.org/doi/10.1103/PhysRev.105.1412>
- [357] T. Sikkeland, S. Amiel, S. Thompson, Spallation reactions of californium-252 with helium ions, Journal of Inorganic and Nuclear Chemistry 11 (4) (1959) 261–263. doi:[https://doi.org/10.1016/0022-1902\(59\)80039-7](https://doi.org/10.1016/0022-1902(59)80039-7).  
URL <https://www.sciencedirect.com/science/article/pii/0022190259800397>
- [358] I. Ahmad, A. M. Friedman, R. F. Barnes, R. K. Sjöblom, J. Milsted, P. R. Fields, Decay of  $\text{Fm}^{253}$ , Phys. Rev. 164 (1967) 1537–1544. doi:[10.1103/PhysRev.164.1537](https://doi.org/10.1103/PhysRev.164.1537).  
URL <https://link.aps.org/doi/10.1103/PhysRev.164.1537>
- [359] M. Asai, K. Tsukada, M. Sakama, S. Ichikawa, T. Ishii, Y. Nagame, I. Nishinaka, K. Akiyama, A. Osa, Y. Oura, K. Sueki, M. Shibata, Experimental identification of spin-parities and single-particle configurations in  $^{257}\text{No}$  and its  $\alpha$ -decay daughter  $^{253}\text{Fm}$ , Phys. Rev. Lett. 95 (2005) 102502. doi:[10.1103/PhysRevLett.95.102502](https://doi.org/10.1103/PhysRevLett.95.102502).  
URL <https://link.aps.org/doi/10.1103/PhysRevLett.95.102502>
- [360] B. G. Harvey, S. G. Thompson, A. Ghiorso, G. R. Choppin, Further production of transcurium nuclides by neutron irradiation, Phys. Rev. 93 (1954) 1129–1129. doi:[10.1103/PhysRev.93.1129](https://doi.org/10.1103/PhysRev.93.1129).  
URL <https://link.aps.org/doi/10.1103/PhysRev.93.1129>
- [361] F. Asaro, S. Björnholm, I. Perlman, Decay scheme of  $\text{fm}^{255}$ , Phys. Rev. 133 (1964) B291–B300. doi:[10.1103/PhysRev.133.B291](https://doi.org/10.1103/PhysRev.133.B291).  
URL <https://link.aps.org/doi/10.1103/PhysRev.133.B291>
- [362] Backe, H, Dretzke, A, Haire, RG, Kunz, P, Lauth, W, Sewtz, M, Trautmann, N, et al., Laser spectroscopic investigation of the element fermium ( $z=100$ ), Hyperfine interactions 162 (1-4) (2005) 3–14.  
URL <http://link.springer.com/article/10.1007/s10751-005-9209-x>
- [363] M. Asai, K. Tsukada, M. Sakama, H. Haba, T. Ichikawa, Y. Ishii, A. Toyoshima, T. Ishii, Y. Nagame, I. Nishinaka, Y. Nagame, Y. Kasamatsu, M. Shibata, Y. Kojima, H. Hayashi, Ground-state configuration of the  $N=157$  nucleus  $^{259}\text{No}$ , Phys. Rev. C 87 (2013) 014332.
- [364] L. Phillips, R. Gatti, A. Chesne, L. Muga, S. Thompson, Discovery of a new mendelevium isotope, Phys. Rev. Lett. 1 (1958) 215–217. doi:[10.1103/PhysRevLett.1.215](https://doi.org/10.1103/PhysRevLett.1.215).  
URL <https://link.aps.org/doi/10.1103/PhysRevLett.1.215>
- [365] T. Sikkeland, A. Ghiorso, R. Latimer, A. E. Larsh, Decay properties of the nuclides fermium-256 and -257 and mendelevium-255, -256, and -257, Phys. Rev. 140 (1965) B277–B282. doi:[10.1103/PhysRev.140.B277](https://doi.org/10.1103/PhysRev.140.B277).  
URL <https://link.aps.org/doi/10.1103/PhysRev.140.B277>
- [366] R. Hoff, J. Evans, E. Hulet, R. Dupzyk, B. Qualheim, Nuclear decay properties of  $^{256}\text{fm}$ , Nuclear Physics A 115 (1) (1968) 225–233. doi:[https://doi.org/10.1016/0375-9474\(68\)90657-X](https://doi.org/10.1016/0375-9474(68)90657-X).  
URL <https://www.sciencedirect.com/science/article/pii/037594746890657X>
- [367] K. F. Flynn, E. P. Horwitz, C. A. A. Bloomquist, R. F. Barnes, R. K. Sjöblom, P. R. Fields, L. E. Glendenin, Distribution of mass in the spontaneous fission of  $^{256}\text{Fm}$ , Phys. Rev. C 5 (1972) 1725–1729. doi:[10.1103/PhysRevC.5.1725](https://doi.org/10.1103/PhysRevC.5.1725).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.5.1725>

- [368] J. F. Wild, P. A. Baisden, R. J. Dougan, E. K. Hulet, R. W. Lougheed, J. H. Landrum, Light-charged-particle emission in the spontaneous fission of  $^{250}\text{Cf}$ ,  $^{256}\text{Fm}$ , and  $^{257}\text{Fm}$ , Phys. Rev. C 32 (1985) 488–495. doi:10.1103/PhysRevC.32.488.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.32.488>
- [369] E. A. Sokol, G. M. Ter-Akopian, A. I. Krupman, V. P. Katkov, L. F. Nikonova, N. V. Eremin, Experiments on the spontaneous fission gamma photons from  $^{248}\text{Cm}$ ,  $^{252}\text{Cf}$ ,  $^{256}\text{Fm}$ , and  $^{259}\text{Md}$ , Soviet Atomic Energy 71 (5) (1991) 906–909. doi:10.1007/BF01124209.
- URL <https://doi.org/10.1007/BF01124209>
- [370] E. K. Hulet, R. W. Hoff, J. E. Evans, R. W. Lougheed, 79-day fermium isotope of mass 257, Phys. Rev. Lett. 13 (1964) 343–345. doi:10.1103/PhysRevLett.13.343.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.13.343>
- [371] F. Asaro, I. Perlman, Nuclear spectroscopic studies of fm $^{257}$ , Phys. Rev. 158 (1967) 1073–1077. doi:10.1103/PhysRev.158.1073.
- URL <https://link.aps.org/doi/10.1103/PhysRev.158.1073>
- [372] J. Wild, E. Hulet, R. Lougheed, Some nuclear properties of fermium-257, Journal of Inorganic and Nuclear Chemistry 35 (4) (1973) 1063–1067. doi:[https://doi.org/10.1016/0022-1902\(73\)80176-9](https://doi.org/10.1016/0022-1902(73)80176-9).
- URL <https://www.sciencedirect.com/science/article/pii/0022190273801769>
- [373] E. K. Hulet, J. F. Wild, R. W. Lougheed, J. E. Evans, B. J. Qualheim, M. Nurmi, A. Ghiorso, Spontaneous-fission half-life of  $^{258}\text{Fm}$  and nuclear instability, Phys. Rev. Lett. 26 (1971) 523–526. doi:10.1103/PhysRevLett.26.523.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.26.523>
- [374] W. John, E. K. Hulet, R. W. Lougheed, J. J. Wesolowski, Symmetric fission observed in thermal-neutron-induced and spontaneous fission of  $^{257}\text{Fm}$ , Phys. Rev. Lett. 27 (1971) 45–48. doi:10.1103/PhysRevLett.27.45.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.27.45>
- [375] D. C. Hoffman, J. B. Wilhelmy, J. Weber, W. R. Daniels, E. K. Hulet, R. W. Lougheed, J. H. Landrum, J. F. Wild, R. J. Dupzyk, 12.3-min  $^{256}\text{Cf}$  and 43-min  $^{258}\text{Md}$  and systematics of the spontaneous fission properties of heavy nuclides, Phys. Rev. C 21 (1980) 972–981. doi:10.1103/PhysRevC.21.972.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.21.972>
- [376] D. C. Hoffman, D. Lee, A. Ghiorso, M. J. Nurmi, K. Aleklett, M. Leino, Fission properties of the 1.5-s spontaneous fission activity produced in bombardment of  $^{248}\text{Cm}$  with  $^{18}\text{O}$ , Phys. Rev. C 24 (1981) 495–499. doi:10.1103/PhysRevC.24.495.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.24.495>
- [377] E. K. Hulet, J. F. Wild, R. J. Dougan, R. W. Lougheed, J. H. Landrum, A. D. Dougan, M. Schadel, R. L. Hahn, P. A. Baisden, C. M. Henderson, R. J. Dupzyk, K. Sümmerer, G. R. Bethune, Bimodal symmetric fission observed in the heaviest elements, Phys. Rev. Lett. 56 (1986) 313–316. doi:10.1103/PhysRevLett.56.313.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.56.313>
- [378] E. K. Hulet, R. W. Lougheed, J. F. Wild, R. J. Dougan, K. J. Moody, R. L. Hahn, C. M. Henderson, R. J. Dupzyk, G. R. Bethune, Z and a identification of 370- $\mu\text{s}$   $^{258}\text{Fm}$ , Phys. Rev. C 34 (1986) 1394–1396. doi:10.1103/PhysRevC.34.1394.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.34.1394>

- [379] E. K. Hulet, J. F. Wild, R. J. Dougan, R. W. Lougheed, J. H. Landrum, A. D. Dougan, P. A. Baisden, C. M. Henderson, R. J. Dupzyk, R. L. Hahn, M. Schädel, K. Sümmerer, G. R. Bethune, Spontaneous fission properties of  $^{258}\text{Fm}$ ,  $^{259}\text{Md}$ ,  $^{260}\text{Md}$ ,  $^{258}\text{No}$ , and  $^{260}\text{No}$ : Bimodal fission, Phys. Rev. C 40 (1989) 770–784. doi:[10.1103/PhysRevC.40.770](https://doi.org/10.1103/PhysRevC.40.770). URL <https://link.aps.org/doi/10.1103/PhysRevC.40.770>
- [380] E. K. Hulet, R. W. Lougheed, J. H. Landrum, J. F. Wild, D. C. Hoffman, J. Weber, J. B. Wilhelmy, Spontaneous fission of  $^{259}\text{Fm}$ , Phys. Rev. C 21 (1980) 966–971. doi:[10.1103/PhysRevC.21.966](https://doi.org/10.1103/PhysRevC.21.966). URL <https://link.aps.org/doi/10.1103/PhysRevC.21.966>
- [381] J. M. Gates, S. L. Nelson, K. E. Gregorich, I. Dragojević, C. E. Düllmann, P. A. Ellison, C. M. Folden III, M. A. Garcia, L. Stavsetra, R. Sudowe, D. C. Hoffman, H. Nitsche, Comparison of reactions for the production of  $^{258,257}\text{Db}$ :  $^{208}\text{Pb}(\text{V}, xn)$  and  $^{209}\text{Bi}(\text{Ti}, xn)$ , Phys. Rev. C 78 (2008) 034604. doi:[10.1103/PhysRevC.78.034604](https://doi.org/10.1103/PhysRevC.78.034604). URL <https://link.aps.org/doi/10.1103/PhysRevC.78.034604>
- [382] P. Kuusiniemi, Nuclear structure investigations of heavy nuclei and the decay of she, AIP Conference Proceedings 764 (1) (2005) 231–236. arXiv:<https://arxiv.org/abs/https://aip.scitation.org/doi/pdf/10.1063/1.1905316>, doi:[10.1063/1.1905316](https://doi.org/10.1063/1.1905316). URL <https://aip.scitation.org/doi/abs/10.1063/1.1905316>
- [383] S. Ketelhut, P. T. Greenlees, D. Ackermann, S. Antalic, E. Clément, I. G. Darby, O. Dorvaux, A. Drouart, S. Eeckhaudt, B. J. P. Gall, A. Görgen, T. Grahn, C. Gray-Jones, K. Hauschild, R.-D. Herzberg, F. P. Heßberger, U. Jakobsson, G. D. Jones, P. Jones, R. Julin, S. Juutinen, T.-L. Khoo, W. Korten, M. Leino, A.-P. Leppänen, J. Ljungvall, S. Moon, M. Nyman, A. Obertelli, J. Pakarinen, E. Parr, P. Papadakis, P. Peura, J. Piot, A. Pritchard, P. Rahkila, D. Rostron, P. Ruotsalainen, M. Sandzelius, J. Sarén, C. Scholey, J. Sorri, A. Steer, B. Sulignano, C. Theisen, J. Uusitalo, M. Venhart, M. Zielinska, M. Bender, P.-H. Heenen,  $\gamma$ -ray spectroscopy at the limits: First observation of rotational bands in  $^{255}\text{Lr}$ , Phys. Rev. Lett. 102 (2009) 212501. doi:[10.1103/PhysRevLett.102.212501](https://doi.org/10.1103/PhysRevLett.102.212501). URL <http://link.aps.org/doi/10.1103/PhysRevLett.102.212501>
- [384] B. Kadkhodayan, R. A. Henderson, H. L. Hall, J. D. Leyba, K. R. Czerwinski, S. A. Kreek, N. J. Hannink, K. E. Gregorich, D. M. Lee, M. J. Nurmia, D. C. Hoffman, Identification of  $^{253}\text{md}$ , Radiochim. Acta 56 (1992) 1.
- [385] P. Fields, I. Ahmad, R. Barnes, R. Sjöblom, E. Horowitz, Nuclear properties of  $^{254}\text{md}$ ,  $^{255}\text{md}$ ,  $^{256}\text{md}$ ,  $^{257}\text{md}$  and  $^{258}\text{md}$ , Nuclear Physics A 154 (2) (1970) 407–416.
- [386] R. Hoff, E. Hulet, R. Dupzyk, R. Lougheed, J. Evans, Alpha decay of  $^{255}\text{md}$ ,  $^{256}\text{md}$  and  $^{257}\text{md}$ , Nuclear Physics A 169 (3) (1971) 641–650. doi:[https://doi.org/10.1016/0375-9474\(71\)90708-1](https://doi.org/10.1016/0375-9474(71)90708-1). URL <https://www.sciencedirect.com/science/article/pii/0375947471907081>
- [387] A. Ghiorso, B. G. Harvey, G. R. Choppin, S. G. Thompson, G. T. Seaborg, New element mendelevium, atomic number 101, Phys. Rev. 98 (1955) 1518–1519. doi:[10.1103/PhysRev.98.1518](https://doi.org/10.1103/PhysRev.98.1518). URL <https://link.aps.org/doi/10.1103/PhysRev.98.1518>
- [388] O. R. E.K.Hulet, A.Ghiorso, R.W.Lougheed, J.E.Evans, J.D.Brady, R.W.Hoff, R.E.Stone, B.J.Qualheim, Discovery of  $^{258}\text{md}$ , Bull.Am.Phys.Soc. 13 (4) (1968) 604.
- [389] J. F. Wild, E. K. Hulet, R. W. Lougheed, P. A.

- Baisden, J. H. Landrum, R. J. Dougan, M. G. Mustafa, [Unusually low fragment energies in the symmetric fission of  \$^{259}\text{Md}\$](#) , Phys. Rev. C 26 (1982) 1531–1543. doi:[10.1103/PhysRevC.26.1531](https://doi.org/10.1103/PhysRevC.26.1531). URL <https://link.aps.org/doi/10.1103/PhysRevC.26.1531>
- [390] R.W.Lougheed, E.K.Hulet, R. Dougan, J.F.Wild, R.J.Dupzyk, C. Henderson, K. Moody, R. Hahn, K. Sümmerer, G. Bethune, The discovery of  $^{260}\text{md}$  and the decay properties of  $^{258}\text{fm}$ ,  $^{258}\text{m}$ ,  $^{258}\text{gmd}$  and  $^{259}\text{md}$ , J.Less-Common Met. 122 (1986) 461.
- [391] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyganov, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, K. Subotic, O. V. Ivanov, A. N. Voinov, V. I. Zagrebaev, K. J. Moody, J. F. Wild, N. J. Stoyer, M. A. Stoyer, R. W. Lougheed, Measurements of cross sections for the fusion-evaporation reactions  $^{204,206,207,208}\text{Pb} + ^{48}\text{Ca}$  and  $^{207}\text{Pb} + ^{34}\text{S}$ : decay properties of the even-even nuclides  $^{238}\text{Cf}$  and  $^{250}\text{No}$ , Phys. Rev. C 64 (2001) 054606. doi:[10.1103/PhysRevC.64.054606](https://doi.org/10.1103/PhysRevC.64.054606). URL <https://link.aps.org/doi/10.1103/PhysRevC.64.054606>
- [392] G. N. Knyazheva, M. G. Itkis, S. V. Khlebnikov, E. M. Kozulin, V. G. Lyapin, V. A. Rubchenya, W. Trzaska, The influence of the entrance channel on the formation and decay of the compound nucleus  $^{250}\text{no}$ , Physics of Particles and Nuclei Letters 5 (1) (2008) 21–28. doi:[10.1134/S1547477108010044](https://doi.org/10.1134/S1547477108010044). URL <https://doi.org/10.1134/S1547477108010044>
- [393] A. Popeko, O. Malyshev, A. Yeremin, A. Belozerov, M. Chelnokov, V. Chepigin, V. Gorshkov, M. Itkis, A. Kabachenko, Y. Oganessian, R. Sagaidak, A. Shutov, A. Svirikhin, In-beam separation and mass determination of superheavy nuclei. part i, Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 510 (3) (2003) 371–376. doi:[https://doi.org/10.1016/S0168-9002\(03\)01823-0](https://doi.org/10.1016/S0168-9002(03)01823-0). URL <https://www.sciencedirect.com/science/article/pii/S0168900203018230>
- [394] A. V. Yeremin, A. V. Belozerov, M. L. Chelnokov, V. I. Chepigin, V. A. Gorshkov, A. P. Kabachenko, O. N. Malyshev, Y. T. Oganessian, A. G. Popeko, R. N. Sagaidak, A. I. Svirikhin, S. Hofmann, G. Berek, I. Brida, S. Saro, The upgrade of the kinematic separator VASSILISSA—experimental results and plans, Physics of Atomic Nuclei 66 (6) (2003) 1042–1052. doi:[10.1134/1.1586416](https://doi.org/10.1134/1.1586416). URL <http://dx.doi.org/10.1134/1.1586416>
- [395] A. Ghiorso, T. Sikkeland, M. J. Nurmia, Isotopes of element 102 with mass 251 to 258, Phys. Rev. Lett. 18 (1967) 401–404. doi:[10.1103/PhysRevLett.18.401](https://doi.org/10.1103/PhysRevLett.18.401). URL <https://link.aps.org/doi/10.1103/PhysRevLett.18.401>
- [396] V.L.Mikheev, V.I.Ilyushchenko, M.B.Miller, S.M.Polikanov, G.N.Flerov, Y. Kharitonov, Synthesis of isotopes of element 102 with mass numbers 254, 253, and 252, Soviet J.At.Energy 22 (1967) 93.
- [397] Y. Oganesyan, Y. Lobanov, S. Tretyakova, Y. Lazarev, I. Kolesov, K. Gavrilov, V. Plotko, Y. Poluboyarinov, Identification of the elements 102 and 104 by means of the collimation method, Soviet J.At.Energy 28 (1970) 502.
- [398] G. Flerov, Y. Oganessian, A. Pleve, N. Pronin, Y. Tretyakov, Acceleration of  $^{48}\text{ca}$  ions and new possibilities of synthesizing superheavy elements, Nuclear Physics A 267 (1976) 359–364. doi:[https://doi.org/10.1016/0375-9474\(76\)90666-7](https://doi.org/10.1016/0375-9474(76)90666-7). URL <https://www.sciencedirect.com/science/article/pii/0375947476906667>
- [399] C. E. Bemis, R. L. Ferguson, F. Plasil, R. J. Silva, F. Pleasonton, R. L. Hahn, Fragment-mass and

- kinetic-energy distributions from the spontaneous fission of  $^{252}\text{No}$ , Phys. Rev. C 15 (1977) 705–712. doi:[10.1103/PhysRevC.15.705](https://doi.org/10.1103/PhysRevC.15.705).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.15.705>
- [400] J. Wild, E. Hulet, R. Lougheed, K. Moody, B. Bandong, R. Dougan, A. Veeck, Spontaneous fission properties of  $^{252,254}\text{No}$  and  $^{256,258}\text{No}$ [104] and the disappearance of the outer fission barrier, Journal of Alloys and Compounds 213–214 (1994) 86 – 92, international Conference on Actinides. doi:[https://doi.org/10.1016/0925-8388\(94\)90885-0](https://doi.org/10.1016/0925-8388(94)90885-0).
- URL <http://www.sciencedirect.com/science/article/pii/0925838894908850>
- [401] R. Julin, In-beam spectroscopy of heavy actinides, Nuclear Physics A 685 (1) (2001) 221–232, nucleus-Nucleus Collisions 2000. doi:[https://doi.org/10.1016/S0375-9474\(01\)00542-5](https://doi.org/10.1016/S0375-9474(01)00542-5).
- URL <https://www.sciencedirect.com/science/article/pii/S0375947401005425>
- [402] R.-D. Herzberg, N. Amzal, F. Becker, P. A. Butler, A. J. C. Chewter, J. F. C. Cocks, O. Dorvaux, K. Eskola, J. Gerl, P. T. Greenlees, N. J. Hammond, K. Hauschild, K. Helariutta, F. Heßberger, M. Houry, G. D. Jones, P. M. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, T. L. Khoo, W. Korten, P. Kuusiniemi, Y. L. Coz, M. Leino, C. J. Lister, R. Lucas, M. Muikku, P. Nieminen, R. D. Page, P. Rahkila, P. Reiter, C. Schlegel, C. Scholey, O. Stezowski, C. Theisen, W. H. Trzaska, J. Uusitalo, H. J. Wollersheim, Spectroscopy of transfermium nuclei:  $^{252}_{102}\text{No}$ , Phys. Rev. C 65 (2001) 014303. doi:[10.1103/PhysRevC.65.014303](https://doi.org/10.1103/PhysRevC.65.014303).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.65.014303>
- [403] A. I. Svirikhin, A. V. Andreev, V. N. Dushin, M. L. Chelnokov, V. I. Chepigin, M. Gupta, A. V. Isaev, I. N. Izosimov, D. E. Katrasev, A. N. Kuznetsov, O. N. Malyshev, S. Mullins, A. G. Popeko, E. A. Sokol, A. V. Yeremin, The emission of prompt neutrons from the spontaneous fission of  $^{252}\text{No}$  and  $^{244}\text{Fm}$ , The European Physical Journal A 48 (9) (2012) 121. doi:[10.1140/epja/i2012-12121-7](https://doi.org/10.1140/epja/i2012-12121-7).
- URL <https://doi.org/10.1140/epja/i2012-12121-7>
- [404] F. P. Heßberger, S. Hofmann, V. Ninov, P. Armbruster, H. Folger, G. Münzenberg, H. J. Schött, A. G. Popeko, A. V. Yeremin, A. N. Andreyev, S. Saro, Spontaneous fission and alpha-decay properties of neutron deficient isotopes  $^{257-253}\text{No}$  and  $^{258}\text{Fm}$ , Zeitschrift für Physik A Hadrons and Nuclei 359 (4) (1997) 415–425. doi:[10.1007/s002180050422](https://doi.org/10.1007/s002180050422).
- URL <http://dx.doi.org/10.1007/s002180050422>
- [405] R.-D. Herzberg, N. Amzal, J. Bastin, F. Becker, P. Brew, P. Butler, A. Chewter, J. Cocks, O. Dorvaux, K. Eskola, J. Gerl, P. Greenlees, N. Hammond, K. Hauschild, K. Helariutta, F. Heßberger, M. Houry, A. Hürstel, R. Humphreys, G. Jones, P. Jones, R. Julin, S. Juutinen, H. Kankaanpää, H. Kettunen, T. Khoo, W. Korten, P. Kuusiniemi, Y. Le Coz, M. Leino, A. Leppänen, C.J. Lister, R. Lucas, M. Muikku, P. Nieminen, R. Page, T. Page, P. Rahkila, P. Reiter, C. Schlegel, C. Scholey, G. Sletten, O. Stezowski, C. Theisen, W. Trzaska, J. Uusitalo, H. Wollersheim, In-beam spectroscopy of  $^{253,254}\text{No}$ , The European Physical Journal A - Hadrons and Nuclei 15 (1) (2002) 205–208. doi:[10.1140/epja/i2001-10255-3](https://doi.org/10.1140/epja/i2001-10255-3).
- URL <http://dx.doi.org/10.1140/epja/i2001-10255-3>
- [406] A. Lopez-Martens, T. Wiborg-Hagen, K. Hauschild, M. Chelnkov, V. Chepigin, D. Curien, O. Dorvaux, G. Drafta, B. Gall, A. Görgen, M. Guttormsen, A. Isaev, I. Izosimov, A. Kabachenko, D. Katrasev, T. Kutsarova, A. Kuznetsov, A. Larsen, O. Malyshev, A. Minkova, S. Mullins, H. Nyhus, D. Pantel-

- ica, J. Piot, A. Popeko, S. Saro, N. Scintee, S. Siem, N. Syed, E. Sokol, A. Svirikin, A. Yeremin, Spectroscopy of  $^{253}\text{no}$  and its daughters, Nucl. Phys. A 852 (2011) 15.
- [407] A. K. Mistry, R. D. Herzberg, P. T. Greenlees, P. Papadakis, K. Auranen, P. A. Butler, D. M. Cox, A. B. Garnsworthy, T. Grahn, K. Hauschild, U. Jakobsson, D. T. Joss, R. Julin, S. Ketelhut, J. Konki, M. Leino, A. Lopez-Martens, R. D. Page, J. Pakarinen, P. Peura, P. Rahkila, M. Sandzelius, C. Scholey, J. Simpson, D. Seddon, J. Sorri, S. Stolze, J. Thornhill, J. Uusitalo, D. Wells, In-beam study of  $^{253}\text{no}$  using the sage spectrometer, The European Physical Journal A 53 (2) (2017) 24. doi:10.1140/epja/i2017-12215-8.  
URL <https://doi.org/10.1140/epja/i2017-12215-8>
- [408] C. E. Bemis, R. J. Silva, D. C. Hensley, O. L. Keller, J. R. Tarrant, L. D. Hunt, P. F. Dittner, R. L. Hahn, C. D. Goodman, X-ray identification of element 104, Phys. Rev. Lett. 31 (1973) 647–650.
- [409] V. E.D.Donets, V.A.Shchegolev, The properties of the isotope 102 $^{254}$ , Soviet J.At.Energy 20 (1966) 257.
- [410] B. A. Zager, M. B. Miller, V. L. Mikheev, S. M. Polikanov, A. M. Sukhov, G. N. Flerov, L. P. Chelnokov, The properties of the isotope 102 $^{254}$ , At.Energ.USSR 20 (1966) 230.
- [411] G. N. Flerov, Synthesis and investigation of element 102 (review), Soviet Atomic Energy 24 (1) (1968) 3–15. doi:10.1007/BF01133459.  
URL <https://doi.org/10.1007/BF01133459>
- [412] A. Türler, H. W. Gäggeler, D. T. Jost, P. Armbruster, W. Brüchle, H. Folger, F. P. Heßberger, S. Hofmann, G. Münzenberg, V. Ninov, M. Schädel, K. Sümmerer, J. V. Kratz, U. Scherer, Determination of the partial electron capture- and spontaneous-fission half-lives of  $^{254}\text{no}$ , Zeitschrift für Physik A Atomic Nuclei 331 (3) (1988) 363–364. doi:10.1007/BF01355610. URL <https://doi.org/10.1007/BF01355610>
- [413] E.K.Hulet, Role of the second barrier upon mass division in the spontaneous fission of the heaviest elements, Phys.Atomic Nuclei 57 (1994) 1099.
- [414] M. Block, D. Ackermann, K. Blaum, C. Droese, M. Dworschak, M. Eibach, S. Eliseev, T. Fleckenstein, E. Haettner, F. Herfurth, F. P. Heßberger, S. Hofmann, J. Ketelaer, J. Ketter, H. J. Kluge, G. Marx, M. Mazzocco, Y. N. Novikov, W. R. Plaß, A. Popeko, S. Rahaman, D. Rodríguez, C. Scheidenberger, L. Schweikhard, P. G. Thirolf, G. K. Vorobyev, C. Weber, Penning trap mass measurements of trans fermium elements with shiptrap, Hyperfine Interactions 196 (1) (2010) 225–231. doi:10.1007/s10751-010-0161-z.  
URL <https://doi.org/10.1007/s10751-010-0161-z>
- [415] J. S. Berryman, R. M. Clark, K. E. Gregorich, J. M. Allmond, D. L. Bleuel, R. J. Cooper, M. Cromaz, M. A. Deleplanque, I. Dragojević, J. Dvorak, P. A. Ellison, P. Fallon, M. A. Garcia, J. M. Gates, S. Gros, O. Gothe, H. B. Jeppesen, D. Kaji, I. Y. Lee, A. O. Macchiavelli, K. Morimoto, H. Nitsche, S. Paschalis, M. Petri, L. Stavsetra, F. S. Stephens, M. A. Stoyer, T. J. Ross, H. Watanabe, M. Wiedeking, Nuclear spectroscopy of the heaviest elements: studies of  $^{254}\text{no}$ ,  $^{257}\text{rf}$ , and  $^{261}\text{sg}$ , Journal of Physics: Conference Series 312 (9) (2011) 092017. doi:10.1088/1742-6596/312/9/092017.  
URL <https://dx.doi.org/10.1088/1742-6596/312/9/092017>
- [416] E. Parr, R. D. Herzberg, S. Antalic, P. T. Greenlees, F. P. Heßberger, S. Ketelhut, D. Rostron, New approaches to assign configurations using low-statistic  $\gamma$ -ray spectra, The European Physical Journal A 48 (10) (2012) 134. doi:10.1140/epja/i2012-12134-2.

- URL <https://doi.org/10.1140/epja/i2012-12134-2>
- [417] Dworschak, Michael, Block, Michael, Ackermann, D, Audi, Georges, Blaum, Klaus, Droese, C, Eliseev, Sergey, Fleckenstein, T, Haettner, E, Herfurth, Frank, et al., **Penning trap mass measurements on nobelium isotopes**, Physical Review C 81 (6) (2010) 064312.  
URL <http://journals.aps.org/prc/abstract/10.1103/PhysRevC.81.064312>
- [418] Ramirez, E. Minaya, Ackermann, D, Blaum, K, Block, M, Droese, C, Düllmann, C. E, Dworschak, M, Eibach, M, Eliseev, S, Haettner, E, et al., **Direct mapping of nuclear shell effects in the heaviest elements**, Science 337 (6099) (2012) 1207–1210.  
URL <http://www.sciencemag.org/content/337/6099/1207.short>
- [419] V. A. Druin, G. N. Akapev, A. G. Demin, Y. V. Lobanov, B. V. Fefilov, G. N. Flerov, L. P. Chelnokov, Nuclear properties of the isotopes of element 102 with mass numbers 255 and 256, At.Energ.USSR 22 (1967) 127.
- [420] G. N. Flerov, S. M. Polikanov, V. L. Mikheev, V. I. Ilyushchenko, V. F. Kushniruk, M. B. Miller, A. M. Sukhov, V. A. Shchegolev, Synthesis of isotopes of element 102 mass numbers 256 and 255 in reactions with oxygen ions, Soviet J.Nucl.Phys. 5 (1967) 848.
- [421] G. Flerov, G. Akapev, A. Demin, V. Druin, Y. Lobanov, B. Fefilov, On the properties of some isotopes of elements 102 and 103, Soviet J.Nucl.Phys. 7 (1968) 588.
- [422] P. Eskola, K. Eskola, M. Nurmi, A. Ghiorso, **Alpha decay of  $^{255}\text{No}$  and  $^{257}\text{No}$** , Phys. Rev. C 2 (1970) 1058–1062. doi:[10.1103/PhysRevC.2.1058](https://doi.org/10.1103/PhysRevC.2.1058).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.2.1058>
- [423] R. Silva, C. Bemis, Jr., P. Dittner, D. Hensley, Measurement of the electron-capture branch of the decay of 225no, Tech. Rep. ORNL-5137 1975 page 70, ORNL (1976).
- [424] K. Morimoto, K. Morita, D. Kaji, S. Goto, H. Haba, E. Ideguchi, R. Kanungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, J. Peter, T. Suda, K. Sueki, I. Tanihata, F. Tokanai, H. Xu, A. Yeremin, A. Yoneda, A. Yoshida, T.-L. Zhao, T. Zheng, **Status of super heavy element research using garis at riken**, Nuclear Physics A 738 (2004) 129–135, proceedings of the 8th International Conference on Clustering Aspects of Nuclear Structure and Dynamics. doi:<https://doi.org/10.1016/j.nuclphysa.2004.04.021>.  
URL <https://www.sciencedirect.com/science/article/pii/S0375947404005585>
- [425] K. Morita, K. Morimoto, D. Kaji, H. Haba, E. Ideguchi, R. Kanungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, I. Tanihata, H. Xu, A. V. Yeremin, A. Yoneda, A. Yoshida, Y. L. Zhao, T. Zheng, **Production and decay of the isotope  $^{271}\text{ds}$  ( $z = 110$ )**, The European Physical Journal A - Hadrons and Nuclei 21 (2) (2004) 257–263. doi:[10.1140/epja/i2003-10205-1](https://doi.org/10.1140/epja/i2003-10205-1).  
URL <https://doi.org/10.1140/epja/i2003-10205-1>
- [426] H. Haba, T. Akiyama, D. Kaji, H. Kikunaga, K. Morimoto, K. Morita, T. Nanri, K. Ooe, N. Sato, A. Shinohara, D. Suzuki, T. Takabe, I. Yamazaki, A. Yokoyama, A. Yoneda, Production of 255no via the  $^{238}\text{U}(22\text{ne}, 5\text{n})^{255}\text{no}$  reaction using the gas-jet transport system coupled to garis, Tech. rep., RIKEN Accelerator Progress Report 2007 (2008).
- [427] E.D.Donets, V.A.Shchegolev, V.A.Ermakov, Synthesis of a new isotope of element 102, Tech. rep., JINR-P-1383 (1963).
- [428] E. Donets, V. Shchegolev, V. Ermakov, Synthesis of the element 102 of mass number 256, Soviet J.At.Energy 16 (1964) 233.

- [429] G. Flerov, A. Demin, V. Druin, Y. Lobanov, V. Mikheev, S. Polikanov, V. Shchegolev, Properties of  $^{102}\text{C}^{256}$ , Soviet J.Nucl.Phys. 7 (1968) 168.
- [430] Y. A. Lazarev, Y. V. Lobanov, Y. T. Oganessian, V. K. Utyonkov, F. S. Abdullin, A. N. Polyakov, J. Rigol, I. V. Shirokovsky, Y. S. Tsyganov, S. Iliev, V. G. Subbotin, A. M. Sukhov, G. V. Buklanov, A. N. Mezentsev, K. Subotic, K. J. Moody, N. J. Stoyer, J. F. Wild, R. W. Lougheed, Decay properties of  $^{257}\text{No}$ ,  $^{261}\text{Rf}$ , and  $^{262}\text{Rf}$ , Phys. Rev. C 62 (2000) 064307. doi:[10.1103/PhysRevC.62.064307](https://doi.org/10.1103/PhysRevC.62.064307). URL <https://link.aps.org/doi/10.1103/PhysRevC.62.064307>
- [431] S. Hofmann, On beyond uranium: journey to the end of the periodic table, CRC Press, 2002.
- [432] H. Haba, D. Kaji, H. Kikunaga, Y. Kudou, K. Morimoto, K. Morita, K. Ozeki, T. Sumita, A. Yoneda, Y. Kasamatsu, Y. Komori, K. Ooe, A. Shinohara, Production and decay properties of the 1.9-s isomeric state in  $^{261}\text{rf}$ , Phys. Rev. C 83 (2011) 034602.
- [433] M. Murakami, S. Goto, H. Murayama, T. Kojima, H. Kudo, D. Kaji, K. Morimoto, H. Haba, Y. Kudou, T. Sumita, R. Sakai, A. Yoneda, K. Morita, Y. Kasamatsu, H. Kikunaga, T. K. Sato, Excitation functions for production of rf isotopes in the  $^{248}\text{cm} + ^{18}\text{o}$  reaction, Phys. Rev. C 88 (2013) 024618. doi:[10.1103/PhysRevC.88.024618](https://doi.org/10.1103/PhysRevC.88.024618). URL <https://link.aps.org/doi/10.1103/PhysRevC.88.024618>
- [434] M. Nurmia, K. Eskola, P. Eskola, A. Ghiorso, New nuclide  $^{258}\text{no}$ , Ucr1-18667 page 63, California University - Lawrence Radiation Laboratory - UCRL (1969).
- [435] Y. A. Lazarev, Y. V. Lobanov, Y. T. Oganessian, V. K. Utyonkov, F. S. Abdullin, G. V. Buklanov, B. N. Gikal, S. Iliev, A. N. Mezentsev, A. N. Polyakov, I. M. Sedykh, I. V. Shirokovsky, V. G. Subbotin, A. M. Sukhov, Y. S. Tsyganov, V. E. Zhuchko, R. W. Lougheed, K. J. Moody, J. F. Wild, E. K. Hulet, J. H. McQuaid, Discovery of enhanced nuclear stability near the deformed shells  $n = 162$  and  $z = 108$ , Phys. Rev. Lett. 73 (1994) 624–627. doi:[10.1103/PhysRevLett.73.624](https://doi.org/10.1103/PhysRevLett.73.624). URL <https://link.aps.org/doi/10.1103/PhysRevLett.73.624>
- [436] M. R. Lane, K. E. Gregorich, D. M. Lee, M. F. Mohar, M. Hsu, C. D. Kacher, B. Kadkhodayan, M. P. Neu, N. J. Stoyer, E. R. Sylvester, J. C. Yang, D. C. Hoffman, Spontaneous fission properties of  $^{262}_{104}\text{Rf}$ , Phys. Rev. C 53 (1996) 2893–2899. doi:[10.1103/PhysRevC.53.2893](https://doi.org/10.1103/PhysRevC.53.2893). URL <https://link.aps.org/doi/10.1103/PhysRevC.53.2893>
- [437] D. Peterson, W. Loveland, O. Batenkov, M. Majorov, A. Veshikov, K. Aleklett, C. Rouki, Direct observation of first-chance fission of  $^{258}\text{No}$ , Phys. Rev. C 79 (2009) 044607. doi:[10.1103/PhysRevC.79.044607](https://doi.org/10.1103/PhysRevC.79.044607). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.044607>
- [438] T. Banerjee, D. J. Hinde, D. Y. Jeung, K. Banerjee, M. Dasgupta, A. C. Beriman, L. T. Bezzina, H. M. Albers, C. E. Düllmann, J. Khuyagbaatar, B. Kindler, B. Lommel, E. C. Simpson, C. Sengupta, B. M. A. Swinton-Bland, T. Tanaka, A. Yakushev, K. Eberhardt, C. Mokry, J. Runke, P. Thörle-Pospiech, N. Trautmann, Systematic evidence for quasifission in  $^9\text{be-}$ ,  $^{12}\text{c-}$  and  $^{16}\text{o-induced}$  reactions forming  $^{258,260}\text{no}$ , Phys. Rev. C 102 (2020) 024603. doi:[10.1103/PhysRevC.102.024603](https://doi.org/10.1103/PhysRevC.102.024603). URL <https://link.aps.org/doi/10.1103/PhysRevC.102.024603>
- [439] R. Silva, P. Dittner, M. Mallory, O. Keller, K. Eskola, P. Eskola, M. Nurmia, A. Ghiorso, The new nuclide nobelium-259, Nuclear Physics A 216 (1) (1973) 97–108. doi:[https://doi.org/10.1016/0375-9474\(73\)90520-4](https://doi.org/10.1016/0375-9474(73)90520-4).

- URL <https://www.sciencedirect.com/science/article/pii/0375947473905204>
- [440] M. Asai, K. Tsukada, M. Sakama, Y. Ishii, A. Toyoshima, T. Ishii, I. Nishinaka, Y. Nagame, Y. Kasamatsu, M. Shibata, H. Hayashi, H. Haba, Y. Kojima, Alpha-gamma coincidence spectroscopy of  $^{259}\text{no}$ , Jaea-review 2008-054 page 40, JAEA-Tokai Tandem Annual Report (2008).
- [441] L. P. Somerville, M. J. Nurmia, J. M. Nitschke, A. Ghiorso, E. K. Hulet, R. W. Lougheed, Spontaneous fission of rutherfordium isotopes, Phys. Rev. C 31 (1985) 1801–1815. doi:10.1103/PhysRevC.31.1801.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.31.1801>
- [442] N. S.A.Karamyan, V.N.Bugrov, Fragment angular distribution and duration of nuclear fission reaction, Soviet J.Nucl.Phys. 43 (1986) 339.
- [443] G. Münzenberg, P. Armbruster, S. Hofmann, F. P. Heßberger, H. Folger, J. G. Keller, V. Ninov, K. Popensieker, A. B. Quint, W. Reisdorf, K. H. Schmidt, J. R. H. Schneider, H. J. Schött, K. Sümmerer, I. Zychor, M. E. Leino, D. Ackermann, U. Gollerthan, E. Hanelt, W. Morawek, D. Vermeulen, Y. Fujita, T. Schwab, Element 107, Zeitschrift für Physik A Atomic Nuclei 333 (2) (1989) 163–175. doi:10.1007/BF01565147.
- URL <https://doi.org/10.1007/BF01565147>
- [444] F. P. Hessberger, GSI experiments in the region of heaviest elements, Journal of Physics G: Nuclear and Particle Physics 25 (4) (1999) 877–879. doi:10.1088/0954-3899/25/4/059.
- URL <https://doi.org/10.1088/0954-3899/25/4/059>
- [445] Y. T. Oganessian, M. Hussonnois, A. G. Demin, Y. U. P. Kharitonov, H. Bruchertseifer, O. Constantinescu, Y. S. Korotkin, S. P. Tretyakova, V. K. Utyonkov, I. V. Shirokovsky, J. Estevez, Experimental studies of the formation and radioactive decay of isotopes with  $z = 104\text{--}109$ , Radiochimica Acta 37 (3) (1984) 113–120. doi:doi:10.1524/ract.1984.37.3.113.
- URL <https://doi.org/10.1524/ract.1984.37.3.113>
- [446] V. Druin, Radioactive properties of isotopes of element 103, Soviet J. Nucl. Phys. 12 (2) (1971) 146.
- [447] K. Eskola, P. Eskola, M. Nurmia, A. Ghiorso, Studies of lawrencium isotopes with mass numbers 255 through 260, Phys. Rev. C 4 (1971) 632–642. doi:10.1103/PhysRevC.4.632.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.4.632>
- [448] C. B. Jr., D. Hensley, P. Dittner, R. Hahn, R. Silva, J. Tarrant, L.D.Hunt, X-ray identification and decay properties of isotopes of lawrencium, Tech. Rep. ORNL-5137 1975 page 73, ORNL (1976).
- [449] F. Heßberger, S. Hofmann, D. Ackermann, S. Antalic, P. Cagarda, I. Kojouharov, P. Kuusiniemi, R. Mann, S. Šaro, Nilsson levels in odd mass odd  $z$  nuclei in the region  $z = (99\text{--}105)$ , gsi scientific report 2004, Tech. Rep. GSI 2004-1 (2004) p. 3, GSI (2004).
- [450] E. Donets, V. Shchegolev, V. Ermakov, Synthesis of the isotope of element 103 (lawrencium), Soviet J.At.Energy (At.Energ.USSR) 19(19) (1966(1965)) 995(109).
- [451] G. Flerov, Y. Korotkin, V. Mikheev, M. Miller, S. Polikanov, V. Stchegolev, On the nuclear properties of the isotopes 256103 and 257103, Nuclear Physics A 106 (2) (1967) 476–480. doi:[https://doi.org/10.1016/0375-9474\(67\)90892-5](https://doi.org/10.1016/0375-9474(67)90892-5).
- URL <https://www.sciencedirect.com/science/article/pii/0375947467908925>
- [452] N. Sato, M. Asai, K. Tsukada, T. Sato, A. Toyoshima, Z. Li, T. Kikuchi, S. Ichikawa, Y. Nagame, Production

- of  $^{256}\text{lr}$  with the  $^{249,250,251}\text{cf}(^{11}\text{b}, \text{xn})$  and  $^{243}\text{am}(^{18}\text{o}, 5\text{n})$  reactions, Jaea-review 2010-056 page 52, JAEA-Tokai Tandem Annual Report (2010).
- [453] N. Sato, T. Sato, M. Asai, A. Toyoshima, K. Tsukada, Z. Li, K. Nishio, Y. Nagame, M. Schädel, H. Haba, S. Ichikawa, H. Kikunaga, Production of  $^{256}\text{lr}$  in the  $^{249, 250, 251}\text{cf} + ^{11}\text{b}$ ,  $^{243}\text{am} + ^{18}\text{o}$ , and  $^{248}\text{cm} + ^{14}\text{n}$  reactions, Radiochim. Acta 102 (204) 211.
- [454] G. Flerov, Y. Oganesyan, Y. Lobanov, Y. Lazarev, S. Tretyakova, I. Kolesov, V. Plotko, Synthesis of element 105, Sov. At. Energy 29 (1970) 967.
- [455] G. Flerov, Y. Oganesian, Y. Lobanov, Y. Lasarev, S. Tretiakova, I. Kolesov, V. Plotko, On the synthesis of element 105, Nuclear Physics A 160 (1) (1971) 181–192. doi:[https://doi.org/10.1016/0375-9474\(70\)90182-X](https://doi.org/10.1016/0375-9474(70)90182-X). URL <https://www.sciencedirect.com/science/article/pii/037594747090182X>
- [456] C. B. Jr., R. Ferguson, R. Silva, F. Plasil, G. O'Kelley, R. Hahn, D. Hensley, E. Hulet, R.W. Lougheed, Production and decay of  $^{262}\text{I}05$  and spontaneous fission properties of  $^{262}\text{I}05$ - $^{262}\text{I}04$ , Bull. Am. Phys. Soc. 22, No. 4, 611, HE14 (1977) 22 (1977) 611.
- [457] K. E. Gregorich, H. L. Hall, R. A. Henderson, J. D. Leyba, K. R. Czerwinski, S. A. Kreek, B. A. Kadkhodayan, M. J. Nurmia, D. M. Lee, D. C. Hoffman, Fission branch in  $^{259}\text{Lr}$  and confirmation of  $^{258}\text{Lr}$  and  $^{259}\text{Lr}$  mass assignments, Phys. Rev. C 45 (1992) 1058–1063. doi:<https://doi.org/10.1103/PhysRevC.45.1058>. URL <https://link.aps.org/doi/10.1103/PhysRevC.45.1058>
- [458] M. Schädel, W. Brächte, E. Schimpf, H. P. Zimmermann, M. K. Gober, J. V. Kratz, N. Trautmann, K. E. Gregorich, A. Türler, K. R. Czerwinski, N. J. Hannink, B. Kadkhodayan, D. M. Lee, M. J. Nurmia, D. C. Hoffman, Chemical properties of element 105 in aqueous solution: Cation exchange separations with  $\alpha$ -hydroxyisobutyric acid, Radiochimica Acta 57 (2-3) (1992) 85–92. doi:[doi:10.1524/ract.1992.57.23.85](https://doi.org/10.1524/ract.1992.57.23.85). URL <https://doi.org/10.1524/ract.1992.57.23.85>
- [459] T. M. Hamilton, K. E. Gregorich, D. M. Lee, K. R. Czerwinski, N. J. Hannink, C. D. Kacher, B. Kadkhodayan, S. A. Kreek, M. J. Nurmia, M. R. Lane, M. P. Neu, A. Türler, D. C. Hoffman, Spontaneous fission properties of  $^{259}\text{Lr}$ , Phys. Rev. C 46 (1992) 1873–1879. doi:[10.1103/PhysRevC.46.1873](https://doi.org/10.1103/PhysRevC.46.1873). URL <https://link.aps.org/doi/10.1103/PhysRevC.46.1873>
- [460] M. Asai, H. Haba, Y. Kasamatsu, N. Sato, K. Tsukada, T. Sato,  $\alpha$ -fine structure spectroscopy for  $^{257}\text{lr}$  and  $^{259}\text{lr}$ , Riken accelerator progress report 2009 page ix, RIKEN (2010).
- [461] R. L. et al., Llnl rep. ucar 100062-87 pages 2-4, Tech. rep., Lawrence Livermore National Laboratory (1987).
- [462] E. Hulet, Results from heavy-ion bombardments of  $^{254}\text{es}$ , Ucrl-100763, California University - Lawrence Radiation Laboratory - UCRL (1989).
- [463] R. Henderson, K. Gregorich, H. Hall, J. Leyba, K. Czerwinski, B. Kadkhodayan, S. Kreek, N. Hannink, M. Nurmia, D. Lee, D. Hoffman,  $^{261}\text{lr}$  and  $^{262}\text{lr}$  studies, lbl-30798, page 65, Tech. rep., Lawrence Berkeley National Laboratory (1991).
- [464] Y. Oganessian, A. Demin, A. Iljinov, S. Tretyakova, A. Pleve, Y. Penionzhkevich, M. Ivanov, Y. Tretyakov, Experiments on the synthesis of neutron-deficient kurchatovium isotopes in reactions induced by  $^{50}\text{Ti}$  ions, Nuclear Physics A 239 (1) (1975) 157–171. doi:[https://doi.org/10.1016/0375-9474\(75\)91140-9](https://doi.org/10.1016/0375-9474(75)91140-9). URL <https://www.sciencedirect.com/science/article/pii/0375947475911409>

- [465] I. Dragojević, K. E. Gregorich, C. E. Düllmann, M. A. Garcia, J. M. Gates, S. L. Nelson, L. Stavsetra, R. Sudowe, H. Nitsche, [Influence of projectile neutron number in the  \$^{208}\text{Pb}\(^{48}\text{Ti}, n\)^{255}\text{Rf}\$  and  \$^{208}\text{Pb}\(^{50}\text{Ti}, n\)^{257}\text{Rf}\$  reactions](#), Phys. Rev. C 78 (2008) 024605. doi:[10.1103/PhysRevC.78.024605](https://doi.org/10.1103/PhysRevC.78.024605). URL <https://link.aps.org/doi/10.1103/PhysRevC.78.024605>
- [466] Y. Oganesyan, Y. Penionzhkevich, S. K. Kvan, A. Shamsutdinov, N. Maltseva, I. Chuburkova, Study of many-nucleon transfer reactions induced by heavy ions in nuclei of medium mass, Sov.J.Nucl.Phys. 11 (1970) 276.
- [467] A. G. Demin, S. P. Tretyakova, V. K. Utyonkov, I. V. Shirokovsky, [On the properties of the element 106 isotopes produced in the reactions pb+54cr](#), Zeitschrift für Physik A Atoms and Nuclei 315 (2) (1984) 197–200. doi:[10.1007/BF01419379](https://doi.org/10.1007/BF01419379). URL <https://doi.org/10.1007/BF01419379>
- [468] Y. T. Oganessian, A. G. Demin, M. Hussonnois, S. P. Tretyakova, Y. P. Kharitonov, V. K. Utyonkov, I. V. Shirokovsky, O. Constantinescu, H. Bruchertseifer, Y. S. Korotkin, [On the stability of the nuclei of element 108 with  \$a=263-265\$](#) , Zeitschrift für Physik A Atoms and Nuclei 319 (2) (1984) 215–217. doi:[10.1007/BF01415635](https://doi.org/10.1007/BF01415635). URL <https://doi.org/10.1007/BF01415635>
- [469] F. P. Heßberger, G. Münzenberg, S. Hofmann, W. Reisdorf, K. H. Schmidt, H. J. Schött, P. Armbruster, R. Hingmann, B. Thuma, D. Vermeulen, [Study of evaporation residues produced in reactions of  \$^{207,208}\text{pb}\$  with  \$^{50}\text{ti}\$](#) , Zeitschrift für Physik A Atoms and Nuclei 321 (2) (1985) 317–327. doi:[10.1007/BF01493453](https://doi.org/10.1007/BF01493453). URL <https://doi.org/10.1007/BF01493453>
- [470] F. Heßberger, G. Münzenberg, S. Hofmann, P. Armbruster, Y. Agarwal, W. Reisdorf, K. Poppensieker, K.-H. Schmidt, J. Schneider, W. Schneider, H. Schött, C.-C. Sahm, D. Vermeulen, B. Thuma, [Experiments on the synthesis of neutron-deficient isotopes with atomic numbers in the region from  \$Z = 102\$  to  \$Z = 105\$](#) , Journal of the Less Common Metals 122 (1986) 445–451, proceedings of Actinides 85 - Part 2. doi:[https://doi.org/10.1016/0022-5088\(86\)90440-6](https://doi.org/10.1016/0022-5088(86)90440-6). URL <https://www.sciencedirect.com/science/article/pii/0022508886904406>
- [471] G. Münzenberg, P. Armbruster, G. Berthes, H. Folger, F. P. Heßberger, S. Hofmann, J. Keller, K. Poppensieker, A. B. Quint, W. Reisdorf, K. H. Schmidt, H. J. Schött, K. Sümmeler, I. Zychor, M. E. Leino, R. Hingmann, U. Gollerthan, E. Hanelt, [Observation of the isotopes  \$^{264}\text{108}\$  and  \$^{265}\text{108}\$](#) , Zeitschrift für Physik A Atomic Nuclei 328 (1) (1987) 49–59. doi:[10.1007/BF01295182](https://doi.org/10.1007/BF01295182). URL <https://doi.org/10.1007/BF01295182>
- [472] G. Münzenberg, F. Hessberger, W. Faust, H. Folger, S. Hofmann, K.-H. Schmidt, H.-J. Schött, P. Armbruster, [On-line measurements of target properties in experiments for the investigation of the heaviest known elements](#), Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment 282 (1) (1989) 28–35. doi:[https://doi.org/10.1016/0168-9002\(89\)90105-8](https://doi.org/10.1016/0168-9002(89)90105-8). URL <https://www.sciencedirect.com/science/article/pii/0168900289901058>
- [473] S. Hofmann, [Heavy and superheavy nuclei](#), Zeitschrift für Physik A Hadrons and Nuclei 358 (2) (1997) 125–129. doi:[10.1007/s002180050286](https://doi.org/10.1007/s002180050286). URL <https://doi.org/10.1007/s002180050286>
- [474] Y. Oganessian, [Synthesis and decay properties of heaviest nuclei with 48ca-induced reactions](#), Nuclear Physics A 787 (1) (2007) 343–352, proceedings of the Ninth International Conference on Nucleus-Nucleus Collisions. doi:<https://doi.org/10.1016/j.nuclphysa.2006.12.055>.

- URL <https://www.sciencedirect.com/science/article/pii/S0375947406010347>
- [475] G. Münzenberg, S. Hofmann, H. Folger, F. P. Heßberger, J. Keller, K. Poppensieker, B. Quint, W. Reisdorf, K. H. Schmidt, H. J. Schött, P. Armbruster, M. E. Leino, R. Hingmann, *The isotopes 259106, 260106, and 261106*, Zeitschrift für Physik A Atoms and Nuclei 322 (2) (1985) 227–235. doi:10.1007/BF01411887.
- URL <https://doi.org/10.1007/BF01411887>
- [476] A. Isaev, R. Mukhin, A. Andreev, Z. Asfari, M. Chelnokov, V. Chepigin, H. Devaraja, O. Dorvaux, B. Gall, K. Hauschild, I. Izosimov, A. Kuznetsova, A. Lopez-Martens, O. Malyshев, A. Popeko, Y. Popov, A. Rahmatinejad, B. Sailaubekov, T. Shneidman, E. Sokol, A. Svirikhin, M. Tezekbayeva, A. Yeremin, N. Zamyatin, Structure of the prompt neutron multiplicity distribution in the spontaneous fission of  $^{256}\text{rf}$ , Physics Letters B 843 (2023) 138008. doi:https://doi.org/10.1016/j.physletb.2023.138008.
- URL <https://www.sciencedirect.com/science/article/pii/S0370269323003428>
- [477] C. Bemis, R. Silva, D. Hensley, O. K. Jr., J. Tarabant, L. Hunt, P. Dittner, R. Hahn, C. Goodman, X-ray identification and decay properties of isotopes of lawrencium, Tech. Rep. ORNL-4976 1973 page 37, ORNL (1974).
- [478] D. Seweryniak, T. Khoo, I. Ahmad, F. Kondev, A. Robinson, S. Tandel, M. Asai, B. Back, M. Carpenter, P. Chowdhury, C. Davids, S. Eeckhaudt, J. Greene, P. Greenlees, S. Gros, K. Hauschild, A. Heinz, R.-D. Herzberg, R. Janssens, D. Jenkins, G. Jones, S. Ketelhut, T. Lauritsen, C. Lister, A. Lopez-Martens, P. Marley, E. McCutchan, T. Nakatsukasa, P. Papadakis, D. Peterson, J. Qian, D. Rostron, I. Stefanescu, U. Tandel, X. Wang, S. Zhu, Bridging the nuclear structure gap between stable and super heavy nuclei, Nuclear Physics A 834 (1) (2010) 357c–361c, the 10th International Conference on Nucleus-Nucleus Collisions (NN2009). doi:<https://doi.org/10.1016/j.nuclphysa.2010.01.039>.
- URL <https://www.sciencedirect.com/science/article/pii/S0375947410000400>
- [479] V. B. Druin, Y. V. Lobanov, D. M. Nadkarni, Y. P. Kharitonov, Y. S. Korotkin, S. P. Tretyakova, V. I. Krashonkin, Nonobservance of spontaneous fission in kurchatovium at berkeley, At.Energ. 35 (1973) 279.
- [480] A. Ghiorso, J. M. Nitschke, J. R. Alonso, C. T. Alonso, M. Nurmia, G. T. Seaborg, E. K. Hulet, R. W. Lougheed, Element 106, Phys. Rev. Lett. 33 (1974) 1490–1493. doi:10.1103/PhysRevLett.33.1490.
- URL <http://link.aps.org/doi/10.1103/PhysRevLett.33.1490>
- [481] C. E. Bemis, P. F. Dittner, R. L. Ferguson, D. C. Hensley, F. Plasil, F. Pleasonton, Spontaneous-fission branching in the decay of  $^{259}\text{104}$ , Phys. Rev. C 23 (1981) 555–558. doi:10.1103/PhysRevC.23.555.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.23.555>
- [482] K. E. Gregorich, M. R. Lane, M. F. Mohar, D. M. Lee, C. D. Kacher, E. R. Sylwester, D. C. Hoffman, First confirmation of the discovery of element 106, Phys. Rev. Lett. 72 (1994) 1423–1426. doi:10.1103/PhysRevLett.72.1423.
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.72.1423>
- [483] C. M. Folden, K. E. Gregorich, C. E. Düllmann, H. Mahmud, G. K. Pang, J. M. Schwantes, R. Sudowe, P. M. Zielinski, H. Nitsche, D. C. Hoffman, Development of an odd-z-projectile reaction for heavy element synthesis:  $^{208}\text{Pb}(^{64}\text{Ni}, n)^{271}\text{Ds}$  and  $^{208}\text{Pb}(^{65}\text{Cu}, n)^{272}\text{111}$ , Phys. Rev. Lett. 93 (2004) 212702. doi:10.1103/PhysRevLett.93.212702.

- URL <https://link.aps.org/doi/10.1103/PhysRevLett.93.212702>
- [484] K. E. Gregorich, J. M. Gates, C. E. Düllmann, R. Sudowe, S. L. Nelson, M. A. Garcia, I. Dragojević, C. M. Folden III, S. H. Neumann, D. C. Hoffman, H. Nitsche, New isotope  $^{264}\text{Sg}$  and decay properties of  $^{262-264}\text{Sg}$ , Phys. Rev. C 74 (2006) 044611. doi:10.1103/PhysRevC.74.044611.
- URL <https://link.aps.org/doi/10.1103/PhysRevC.74.044611>
- [485] G. Flerov, Y. Oganesyan, Y. Lobanov, V. Kuznetsov, V. Druin, V. Perelygin, K. Gavrilov, S. Tretiakova, V. Plotko, Synthesis and physical identification of the isotope of element 104 with mass number 260, Physics Letters 13 (1) (1964) 73–75. doi:[https://doi.org/10.1016/0031-9163\(64\)90313-0](https://doi.org/10.1016/0031-9163(64)90313-0).
- URL <https://www.sciencedirect.com/science/article/pii/0031916364903130>
- [486] V. Druin, Y. Korotkin, Y. Lobanov, Y. Poluboyarinov, R. Sagaidak, G. Soloveva, S. Tretyakova, Y. Kharitonov, Synthesis of heavy isotopes of  $\text{ku}$  in bombardment of  $\text{cm}$  by  $\text{o}$  ions, Sov.J.Nucl.Phys. (Yad.Fiz.) 24(24) (1977(1976)) 131(254).
- [487] C. E. Bemis, P. F. Dittner, R. J. Silva, R. L. Hahn, J. R. Tarrant, L. D. Hunt, D. C. Hensley, Production,  $L$  x-ray identification, and decay of the nuclide  $^{260}\text{105}$ , Phys. Rev. C 16 (1977) 1146–1158. doi:10.1103/PhysRevC.16.1146.
- URL <http://link.aps.org/doi/10.1103/PhysRevC.16.1146>
- [488] V. A. Druin, B. Bochev, Y. S. Korotkin, V. N. Kosyakov, Y. V. Lobanov, E. A. Minin, Y. V. Poluboyarinov, A. G. Rykov, R. N. Sagaidak, S. P. Tretyakova, Y. P. Kharitonov, Synthesis of  $^{260}\text{ku}$  in the irradiation of  $^{249}\text{bk}$  by  $^{15}\text{n}$  ions, At.Energ. 43 (1977) 155.
- [489] J. Nitschke, M. Fowler, A. Ghiorso, R. Leber, M. Leino, M. Nurmia, L. Somerville, K. Williams, E. Hulet, J. Landrum, R. Lougheed, J. Wild, C. Bemis, R. Silva, P. Eskola, Search for an 80 ms spontaneous fission activity in bombardments of  $^{249}\text{bk}$  with  $^{15}\text{n}$ , Nuclear Physics A 352 (1) (1981) 138–146. doi:[https://doi.org/10.1016/0375-9474\(81\)90564-9](https://doi.org/10.1016/0375-9474(81)90564-9).
- URL <https://www.sciencedirect.com/science/article/pii/0375947481905649>
- [490] D. C. Hoffman, M. R. Lane, Spontaneous fission, Radiochim.Acta 70/71 (1995) 135.
- [491] A. Ghiorso, M. Nurmia, K. Eskola, P. Eskola,  $^{261}\text{rf}$ ; new isotope of element 104, Physics Letters B 32 (2) (1970) 95–98. doi:[https://doi.org/10.1016/0370-2693\(70\)90595-2](https://doi.org/10.1016/0370-2693(70)90595-2).
- URL <https://www.sciencedirect.com/science/article/pii/0370269370905952>
- [492] R. Lougheed, K. Moody, J. Wild, E. Hulet, J. McQuaid, Y. Lazarev, Y. Lobanov, Y. Oganessian, V. Utyonkov, F. Abdullin, G. Buklanov, B. Gikal, S. Iliev, A. Mezentsev, A. Polyakov, I. Sedykh, I. Shirokovsky, V. Subbotin, A. Sukhov, Y. Tsyanov, V. Zhuchko, Observation of enhanced nuclear stability near the 162 neutron shell, Journal of Alloys and Compounds 213-214 (1994) 61–66, international Conference on Actinides. doi:[https://doi.org/10.1016/0925-8388\(94\)90881-8](https://doi.org/10.1016/0925-8388(94)90881-8).
- URL <https://www.sciencedirect.com/science/article/pii/0925838894908818>
- [493] B. Kadkhodayan, A. Turler, K. E. Gregorich, P. A. Baisden, K. R. Czerwinski, B. Eichler, H. W. Gaggeler, T. M. Hamilton, D. T. Jost, C. D. Kacher, A. Kovacs, S. A. Creek, M. R. Lane, M. F. Mohar, M. P. Neu, N. J. Stoyer, E. R. Sylwester, D. M. Lee, M. J. Nurmia, G. T. Seaborg, D. C. Hoffman, On-line gas chromatographic studies of chlorides of rutherfordium and homologs zr and hf, Radiochim.Acta 72 (1996) 169.

- [494] S. Hofmann, V. Ninov, F. P. Heßberger, P. Armbruster, H. Folger, G. Münzenberg, H. J. Schött, A. G. Popeko, A. V. Yeremin, S. Saro, R. Janik, M. Leino, [The new element 112](#), Zeitschrift für Physik A Hadrons and Nuclei 354 (3) (1996) 229–230. [doi:10.1007/BF02769517](#).  
URL <https://doi.org/10.1007/BF02769517>
- [495] M. Schadel, W. Bruchle, B. Schausten, E. Schimpf, E. Jager, G. Wirth, R. Gunther, J. V. Kratz, W. Paulus, A. Seibert, P. Thorle, N. Trautmann, S. Zauner, D. Schumann, M. Andrassy, R. Misiak, K. E. Gregorich, D. C. Hoffman, D. M. Lee, E. R. Sylvester, Y. Nagame, Y. Oura, First aqueous chemistry with seaborgium (element 106), Radiochim.Acta 77 (1997) 149.
- [496] A. Türler, R. Dressler, B. Eichler, H. W. Gäggeler, D. T. Jost, M. Schädel, W. Brüchle, K. E. Gregorich, N. Trautmann, S. Taut, Decay properties of  $^{265}\text{Sg}$ ( $z = 106$ ) and  $^{266}\text{Sg}$ ( $z = 106$ ), Phys. Rev. C 57 (1998) 1648–1655. [doi:10.1103/PhysRevC.57.1648](#).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.57.1648>
- [497] E. R. Sylvester, K. E. Gregorich, D. M. Lee, B. Kadkhodayan, A. Turler, J. L. Adams, C. D. Kacher, M. R. Lane, C. A. Laue, C. A. McGrath, D. A. Shaughnessy, D. A. Strellis, P. A. Wilk, D. C. Hoffman, On-line gas chromatographic studies of rf, zr, and hf bromides, Radiochim.Acta 88 (2000) 837.
- [498] S. Hübener, S. Taut, A. Vahle, R. Dressler, B. Eichler, H. W. Gäggeler, D. T. Jost, D. Piguet, A. Türler, A. ler, W. Brüchle, Physico-chemical characterization of seaborgium as oxide hydroxide:, Radiochimica Acta 89 (11-12) (2001) 737–742. [doi:doi:10.1524/ract.2001.89.11-12.737](#).  
URL <https://doi.org/10.1524/ract.2001.89.11-12.737>
- [499] A. Türler, C. Düllmann, H. Gäggeler, U. Kirbach, A. Yakushev, M. Schädel, W. Brüchle, R. Dressler, K. Eberhardt, B. Eichler, R. Eichler, T. Ginter, F. Glaus, K. Gregorich, D. Hoffman, E. Jäger, D. Jost, D. Lee, H. Nitsche, J. Patin, V. Pershina, D. Piguet, Z. Qin, B. Schausten, E. Schimpf, H.-J. Schött, S. Soverna, R. Sudowe, P. Thörle, S. Timokhin, N. Trautmann, A. Vahle, G. Wirth, P. Zielinski, On the decay properties of  $^{269}\text{Hs}$  and indications for the new nuclide  $^{270}\text{Hs}$ , The European Physical Journal A - Hadrons and Nuclei 17 (4) (2003) 505–508. [doi:10.1140/epja/i2002-10163-0](#).  
URL <http://dx.doi.org/10.1140/epja/i2002-10163-0>
- [500] A. von Zweidorf, W. Brüchle, S. Bürger, H. Hummrich, J. V. Kratz, B. Kuczewski, G. Langrock, U. Rieth, M. Schädel, N. Trautmann, K. Tsukada, N. Wiehl, Evidence for the formation of sodium has-sate(viii):, Radiochimica Acta 92 (12) (2004) 855–861. [doi:doi:10.1524/ract.92.12.855.55112](#).  
URL <https://doi.org/10.1524/ract.92.12.855.55112>
- [501] K. Morita, K. Morimoto, D. Kaji, T. Akiyama, S.-i. Goto, H. Haba, E. Ideguchi, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, F. Tokanai, T. Yamaguchi, A. Yoneda, A. Yoshida, Experiment on synthesis of an isotope 277112 by  $^{208}\text{pb}+^{70}\text{zn}$  reaction, Journal of the Physical Society of Japan 76 (4) (2007) 043201. [arXiv:https://doi.org/10.1143/JPSJ.76.043201](#), [doi:10.1143/JPSJ.76.043201](#), [76.043201](#).  
URL <https://doi.org/10.1143/JPSJ.76.043201>
- [502] C. E. Düllmann, A. Türler,  $^{248}\text{cm}(^{22}\text{ne},\text{xn})^{270-x}\text{sg}$  reaction and the decay properties of  $^{265}\text{sg}$  re-examined, Phys. Rev. C 77 (2008) 064320. [doi:10.1103/PhysRevC.77.064320](#).  
URL <https://link.aps.org/doi/10.1103/PhysRevC.77.064320>
- [503] H. Haba, D. Kaji, Y. Kudou, K. Morimoto, K. Morita, K. Ozeki, R. Sakai, T. Sumita, A. Yoneda,

- Y. Kasamatsu, Y. Komori, A. Shinohara, Kikunaga, H. Kudo, K. Nishio, K. Ooe, N. Sato, K. Tsukada, Production of  $^{265}\text{sg}$  in the  $^{248}\text{cm}(^{22}\text{ne and}5\text{n})^{265}\text{sg}$  reaction and decay properties of two isomeric states in  $^{265}\text{sg}$ , Phys. Rev. C 85 (2012) 024611.
- [504] J. M. Nitschke, L. P. Somerville, M. J. Nurmia, R. C. Eggers, A. Ghiorso, Spontaneous fission activities in the bombardment of  $^{248}\text{cm}$  with oxygen ions, Tech. Rep. LBL-8151,P84, Lawrence Berkeley National Laboratory (1978).
- [505] J. V. Kratz, M. K. Gober, H. P. Zimmermann, M. Schädel, W. Brüchle, E. Schimpf, K. E. Gregorich, A. Türler, N. J. Hannink, K. R. Czerwinski, B. Kadkhodayan, D. M. Lee, M. J. Nurmia, D. C. Hoffman, H. Gäggeler, D. Jost, J. Kovacs, U. W. Scherer, A. Weber, New nuclide  $^{263}\text{Ha}$ , Phys. Rev. C 45 (1992) 1064–1069. doi:[10.1103/PhysRevC.45.1064](https://doi.org/10.1103/PhysRevC.45.1064). URL <https://link.aps.org/doi/10.1103/PhysRevC.45.1064>
- [506] J. V. Kratz, A. Nähler, U. Rieth, A. Kronenberg, B. Kuczewski, E. Strub, W. Brüchle, M. Schädel, B. Schausten, A. Türler, H. W. Gäggeler, D. T. Jost, K. E. Gregorich, H. Nitsche, C. Laue, R. Sudowe, P. A. Wilk, An ec-branch in the decay of 27-s  $^{263}\text{db}$ : Evidence for the isotope  $^{263}\text{rf}$ , Radiochimica Acta 91 (1) (2003) 59–62. doi:[doi:10.1524/ract.91.1.59.19010](https://doi.org/10.1524/ract.91.1.59.19010). URL <https://doi.org/10.1524/ract.91.1.59.19010>
- [507] F. P. Hessberger, Experiments on the synthesis of new superheavy elements, Acta Phys.Slovaca 49 (1999) 43.
- [508] A. N. Andreyev, D. D. Bogdanov, V. I. Chepigin, A. P. Kabachenko, O. N. Malyshev, Y. T. Oganessian, R. N. Sagaidak, G. M. Ter-Akopian, A. V. Yeremin, F. P. Hessberger, S. Hofmann, V. Ninov, M. Florek, S. Saro, M. Veselsky, Investigation of the fusion reaction  $^{27}\text{al} + ^{236}\text{u} \rightarrow ^{263}105$  at excitation energies of 57 mev and 65 mev, Z.Phys. A344 (1992) 225.
- [509] F. P. Heßberger, S. Antalic, D. Ackermann, S. Heinz, S. Hofmann, J. Khuyagbaatar, B. Kindler, I. Koujouharov, B. Lommel, R. Mann, Alpha-decay properties of  $^{261}\text{bh}$ , The European Physical Journal A 43 (2) (2010) 175–180. doi:[10.1140/epja/i2010-10913-3](https://doi.org/10.1140/epja/i2010-10913-3). URL <https://doi.org/10.1140/epja/i2010-10913-3>
- [510] P. Schury, T. Niwase, M. Wada, P. Brionnet, S. Chen, T. Hashimoto, H. Haba, Y. Hirayama, D. S. Hou, S. Iimura, H. Ishiyama, S. Ishizawa, Y. Ito, D. Kaji, S. Kimura, H. Koura, J. J. Liu, H. Miyatake, J.-Y. Moon, K. Morimoto, K. Morita, D. Nagae, M. Rosenbusch, A. Takamine, Y. X. Watanabe, H. Wollnik, W. Xian, S. X. Yan, First high-precision direct determination of the atomic mass of a superheavy nuclide, Phys. Rev. C 104 (2021) L021304. doi:[10.1103/PhysRevC.104.L021304](https://doi.org/10.1103/PhysRevC.104.L021304). URL <https://link.aps.org/doi/10.1103/PhysRevC.104.L021304>
- [511] T. Niwase, P. Schury, M. Wada, the SHE-Mass Collaboration, Accurate event assignment from the decay-correlated mass measurement of the superheavy nuclide  $^{257}\text{db}$ , J.Nucl.Radiochem.Sci. 23 (2023) 1.
- [512] Z. Zhao, Z. G. Gan, Z. Y. Zhang, J. G. Wang, M. H. Huang, L. Ma, H. B. Yang, M. M. Zhang, C. L. Yang, S. Y. Xu, X. Y. Huang, Z. C. Li, L. C. Sun, X. L. Wu, Y. S. Wang, Y. L. Tian, Y. H. Qiang, J. Y. Wang, W. X. Huang, Y. He, L. T. Sun, V. K. Utyonkov, A. A. Voinov, Y. S. Tsyganov, A. N. Polyakov, Reinvestigation of the decay properties of  $^{261,262}\text{Bh}$  at the gas-filled recoil separator shans2, Phys. Rev. C 109 (2024) 034314. doi:[10.1103/PhysRevC.109.034314](https://doi.org/10.1103/PhysRevC.109.034314). URL <https://link.aps.org/doi/10.1103/PhysRevC.109.034314>
- [513] O. Kaleja, B. Andelić, O. Bezrodnova, K. Blaum, M. Block, S. Chenmarev, P. Chhetri, C. Droese, C. E. Düllmann, M. Eibach, S. Eliseev, J. Even, P. Filianin, F. Giacoppo, S. Götz, Y. Gusev, M. J. Gutiérrez,

- F. P. Heßberger, N. Kalantar-Nayestanaki, J. J. W. van de Laar, M. Laatiaoui, S. Lohse, N. Martynova, E. Minaya Ramirez, A. K. Mistry, T. Murböck, Y. Novikov, S. Raeder, D. Rodríguez, F. Schneider, L. Schweikhard, P. G. Thirolf, A. Yakushev, [Direct high-precision mass spectrometry of superheavy elements with shiptrap](#), Phys. Rev. C 106 (2022) 054325. doi:[10.1103/PhysRevC.106.054325](https://doi.org/10.1103/PhysRevC.106.054325). URL <https://link.aps.org/doi/10.1103/PhysRevC.106.054325>
- [514] J. S. Guo, Z. Qin, Z. G. Gan, H. M. Fan, Y. B. Xu, J. J. He, X. G. Lei, X. L. Wu, H. Y. Liu, X. H. Zhou, S. G. Yuan, G. M. Jin, Preparation of thick americium targets and synthesis of  $^{259}\text{db}$ , J.Nucl.Radiochem.Sci. 3 (2002) No 1, 183.
- [515] G. N. Flerov, V. A. Druin, A. G. Demin, Y. V. Lobanov, N. K. Skobelev, G. N. Akapev, B. V. Feofilov, I. V. Kolesov, K. A. Gavrilov, Y. P. Kharitonov, L. P. Chelnokov, Experiments in search of alpha-radioactive isotopes of element 105, Tech. Rep. JINR P7 3808, JINR (1968).
- [516] G. N. Flerov, Y. T. Oganesyan, Y. V. Lobanov, Y. A. Lazarev, V. I. Kuznetsov, S. P. Tretiakova, Investigation of the regularities of the production of spontaneously fissionable isotope of element 105, Tech. Rep. JINR-P7-5108, JINR (1970).
- [517] A. Ghiorso, M. Nurmia, K. Eskola, J. Harris, P. Eskola, [New element hahnium, atomic number 105](#), Phys. Rev. Lett. 24 (1970) 1498–1503. doi:[10.1103/PhysRevLett.24.1498](https://doi.org/10.1103/PhysRevLett.24.1498). URL <https://link.aps.org/doi/10.1103/PhysRevLett.24.1498>
- [518] V. A. Druin, A. G. Demin, Y. P. Kharitonov, G. N. Akapev, V. I. Rud, G. Y. Sung-Ching-Yang, L. P. Chelnokov, K. A. Gavrilov, Study of  $\alpha$  decay of 105 element isotopes, Sov.J.Nucl.Phys. 13 (1971) 139.
- [519] S. Hofmann, V. Ninov, F. P. Heßberger, P. Armbruster, H. Folger, G. Münzenberg, H. J. Schött, A. G. Popeko, A. V. Yeremin, A. N. Andreyev, v. Šaro, R. Janik, M. Leino, [The new element 111](#), Zeitschrift für Physik A Hadrons and Nuclei 350 (4) (1995) 281–282. doi:[10.1007/BF01291182](https://doi.org/10.1007/BF01291182). URL <http://dx.doi.org/10.1007/BF01291182>
- [520] K. Morita, K. Morimoto, D. Kaji, S. Goto, H. Haba, E. Ideguchi, R. Kanungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, J. Peter, T. Suda, K. Sueki, I. Tanihata, F. Tokanai, H. Xu, A. Yeremin, A. Yoneda, A. Yoshida, Y.-L. Zhao, T. Zheng, [Status of heavy element research using garis at riken](#), Nuclear Physics A 734 (2004) 101–108. doi:<https://doi.org/10.1016/j.nuclphysa.2004.01.019>. URL <https://www.sciencedirect.com/science/article/pii/S0375947404000247>
- [521] A. Ghiorso, M. Nurmia, K. Eskola, P. Eskola, [Two new alpha-particle emitting isotopes of element 105,  \$^{261}\text{Ha}\$  and  \$^{262}\text{Ha}\$](#) , Phys. Rev. C 4 (1971) 1850–1855. doi:[10.1103/PhysRevC.4.1850](https://doi.org/10.1103/PhysRevC.4.1850). URL <https://link.aps.org/doi/10.1103/PhysRevC.4.1850>
- [522] M. R. Lane, K. E. Gregorich, D. M. Lee, B. Wierszczinski, C. A. McGrath, M. B. Hendricks, D. A. Shaughnessy, D. A. Strellis, E. R. Sylvester, P. A. Wilk, D. C. Hoffman, [Production cross sections of  \$^{261}\text{Ha}\$  from the  \$^{250}\text{Cf}\(\text{ \$^{15}\text{N}\$ , 4n}\)\$  and  \$^{243}\text{Am}\(\text{ \$^{22}\text{Ne}\$ , 4n}\)\$  reactions](#), Phys. Rev. C 58 (1998) 3413–3418. doi:[10.1103/PhysRevC.58.3413](https://doi.org/10.1103/PhysRevC.58.3413). URL <https://link.aps.org/doi/10.1103/PhysRevC.58.3413>
- [523] M. Asai, K. Tsukada, N. Sato, T. K. Sato, A. Toyoshima, S. Miyashita, K. Ooe, Y. Kaneya, T. Ishii, Y. Nagame, M. Schadel, Alpha-gamma coincidence measurement for  $^{261}\text{db}$ , Tech. Rep. JAEA-Review 2013-057,P11, JAEA, Tokai, Japan (2013).
- [524] C. E. Bemis, R. L. Ferguson, F. Plasil, R. J. Silva, G. D. O’Kelley, M. L. Kiefer, R. L. Hahn, D. C. Hens-

- ley, E. K. Hulet, R. W. Lougheed, **Mass asymmetry and total-kinetic-energy release in the spontaneous fission of  $^{262}$ [105]**, Phys. Rev. Lett. 39 (1977) 1246–1249. doi:[10.1103/PhysRevLett.39.1246](https://doi.org/10.1103/PhysRevLett.39.1246).
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.39.1246>
- [525] V. Druin, B. Bochev, Y. Lobanov, R. Sagaidak, Y. Kharitonov, S. Tretyakova, G. Gulbekyan, G. Bulkanov, E. Erin, V. Kosyakov, A. Rykov, Spontaneous fission of the heavy isotopes of nielsbohrium ( $z=105$ ) and element 106, Sov. J. Nucl. Phys. 29 (5) (1979) 591–595.
- [526] K. E. Gregorich, R. A. Henderson, D. M. Lee, M. J. Nurmia, R. M. Chasteler, H. L. Hall, D. A. Bennett, C. M. Gannett, R. B. Chadwick, J. D. Leyba, D. C. Hoffman, G. Herrmann, Aqueous chemistry of element 105, Radiochim.Acta 43 (1988) 223.
- [527] J. V. Kratz, H. P. Zimmermann, U. W. Scherer, M. Schädel, W. Brückle, K. E. Gregorich, C. M. Gannett, H. L. Hall, R. A. Henderson, D. M. Lee, J. D. Leyba, M. J. Nurmia, D. C. Hoffman, H. Gäggeler, D. Jost, U. Baltensperger, N. Q. Ya, A. Türler, C. Lienert, Chemical properties of element 105 in aqueous solution: Halide complex formation and anion exchange into triisooctyl amine, Radiochim.Acta 48 (1989) 121.
- [528] M. K. Gober, J. V. Kratz, H. P. Zimmermann, M. Schädel, W. Brückle, E. Schimpf, K. E. Gregorich, A. Türler, N. J. Hannink, K. R. Czerwinski, B. Kadkhodayan, D. M. Lee, M. J. Nurmia, D. C. Hoffman, H. Gäggeler, D. Jost, J. Kovacz, U. W. Scherer, A. Weber, Chemical properties of element 105 in aqueous solution: Extractions into diisobutylcarbinol, Radiochim.Acta 57 (1992) 77.
- [529] R. Dressler, B. Eichler, D. T. Jost, D. Piguet, A. Türler, C. Düllmann, R. Eichler, H. W. Gäggeler, M. Gärtner, M. Schädel, S. Taut, A. B. Yakushev, **Production of  $^{262}\text{Db}$ ( $z = 105$ ) in the reaction  $^{248}\text{Cm}(\text{F}, \text{n})$** , Phys. Rev. C 59 (1999) 3433–3436. doi:[10.1103/PhysRevC.59.3433](https://doi.org/10.1103/PhysRevC.59.3433).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.59.3433>
- [530] P. A. Wilk, K. E. Gregorich, A. Türler, C. A. Laue, R. Eichler, V. Ninov, J. L. Adams, U. W. Kirbach, M. R. Lane, D. M. Lee, J. B. Patin, D. A. Shaughnessy, D. A. Strellis, H. Nitsche, D. C. Hoffman, **Evidence for new isotopes of element 107:  $^{266}\text{bh}$  and  $^{267}\text{bh}$** , Phys. Rev. Lett. 85 (2000) 2697–2700. doi:[10.1103/PhysRevLett.85.2697](https://doi.org/10.1103/PhysRevLett.85.2697).
- URL <https://link.aps.org/doi/10.1103/PhysRevLett.85.2697>
- [531] Y. Nagame, M. Asai, H. Haba, K. Tsukada, I. Nishinaka, S. Goto, A. Toyoshima, K. Akiyama, M. Sakama, Y. L. Zhao, S. Ichikawa, H. Nakahara, Heavy element nuclear chemistry at JAERI, Physics of Atomic Nuclei 66 (2003) 1131–1136.
- [532] Z. Qin, X. l. Wu, H. j. Ding, W. Wu, W. x. Huang, X. g. Lei, Y. b. Xu, X. h. Yuan, B. Guo, W. f. Yang, Z. g. Gan, H. m. Fan, J. S. Guo, H. s. Xu, G. q. Xiao, Alpha-decay properties of  $^{266}\text{bh}$ , Yuan.Wul.Ping. 23 (2006) 400.
- [533] K. E. Gregorich, R. Leres, D. Lee, D. C. Hoffman, Possible detection of new isotope 263-105, Tech. Rep. LBL-22820,P54, Lawrence Berkeley National Laboratory (1987).
- [534] H. P. Zimmermann, M. K. Gober, J. V. Kratz, M. Schädel, W. Brückle, E. Schimpf, K. E. Gregorich, A. Türler, K. R. Czerwinski, N. J. Hannink, B. Kadkhodayan, D. M. Lee, M. J. Nurmia, D. C. Hoffman, H. Gäggeler, D. Jost, J. Kovacz, U. W. Scherer, A. Weber, Chemical properties of element 105 in aqueous solution: Back extraction from triisooctyl amine into 0.5 m hcl, Radiochim.Acta 60 (1993) 11.
- [535] K. Gregorich, C. Kacher, M. Mohar, D. Lee, M. Lane, E. Sylvester, D. Hoffman, M. Schädel, W. Brückle,

- B. Schausten, E. Schimpf, J. Kratz, R. Gunther, U. Becker, N. Trautmann, A search for the isotope 263 of element 104 - a possible ec-branch in 263ha, Tech. Rep. GSI-95-1 (1995) p. 14, GSI (1994). URL <https://repository.gsi.de/record/53537/files/GSI-Report-1995-1.pdf>
- [536] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyganov, G. G. Gulbekian, S. L. Bogomolov, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, J. B. Patin, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, R. W. Lougheed, Experiments on the synthesis of element 115 in the reaction  $^{243}\text{am}(^{48}\text{ca}, \text{xn})^{291-x}115$ , Tech. Rep. JINR-E7-2003-178, JINR (2003).
- [537] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyganov, G. G. Gulbekian, S. L. Bogomolov, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, J. B. Patin, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, R. W. Lougheed, Experiments on the synthesis of element 115 in the reaction  $^{243}\text{Am}(^{48}\text{Ca}, \text{xn})^{291-x}115$ , Phys. Rev. C 69 (2004) 021601. doi:[10.1103/PhysRevC.69.021601](https://doi.org/10.1103/PhysRevC.69.021601). URL <https://link.aps.org/doi/10.1103/PhysRevC.69.021601>
- [538] S. N. Dmitriev, Y. T. Oganessyan, V. K. Utyonkov, S. V. Shishkin, A. V. Yeremin, Y. V. Lobanov, Y. S. Tsyganov, V. I. Chepygin, E. A. Sokol, G. K. Vostokin, N. V. Aksенов, M. Hussonnois, M. G. Itkis, H. W. Gäggeler, D. Schumann, H. Bruchertseifer, R. Eichler, D. A. Shaughnessy, P. A. Wilk, J. M. Kenneally, M. A. Stoyer, J. F. Wild, Chemical identification of dubnium as a decay product of element 115 produced in the reaction  $^{48}\text{ca} + ^{243}\text{am}$ , Mendeleev Communications 15 (1) (2005) 1–4. doi:<https://doi.org/10.1070/MC2005v015n01ABEH002077>. URL <https://www.sciencedirect.com/science/article/pii/S0959943605701577>
- [539] Y. T. Oganessian, V. K. Utyonkov, S. N. Dmitriev, Y. V. Lobanov, M. G. Itkis, A. N. Polyakov, Y. S. Tsyganov, A. N. Mezentsev, A. V. Yeremin, A. A. Voinov, E. A. Sokol, G. G. Gulbekian, S. L. Bogomolov, S. Iliev, V. G. Subbotin, A. M. Sukhov, G. V. Buklanov, S. V. Shishkin, V. I. Chepygin, G. K. Vostokin, N. V. Aksenov, M. Hussonnois, K. Subotic, V. I. Zagrebaev, K. J. Moody, J. B. Patin, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, P. A. Wilk, R. W. Lougheed, H. W. Gäggeler, D. Schumann, H. Bruchertseifer, R. Eichler, Synthesis of elements 115 and 113 in the reaction  $^{243}\text{Am} + ^{48}\text{Ca}$ , Phys. Rev. C 72 (2005) 034611. doi:[10.1103/PhysRevC.72.034611](https://doi.org/10.1103/PhysRevC.72.034611). URL <http://link.aps.org/doi/10.1103/PhysRevC.72.034611>
- [540] N. Stoyer, J. Landrum, P. Wilk, K. Moody, J. Kenneally, D. Shaughnessy, M. Stoyer, J. Wild, R. Lougheed, S. Dmitriev, Y. Oganessian, S. Shishkin, N. Aksenov, E. Tereshatov, G. Bozhikov, G. Vostokin, V. Utyonkov, A. Yeremin, Chemical identification of a long-lived isotope of dubnium, a descendant of element 115, Nuclear Physics A 787 (1) (2007) 388–395, proceedings of the Ninth International Conference on Nucleus-Nucleus Collisions. doi:<https://doi.org/10.1016/j.nuclphysa.2006.12.060>. URL <https://www.sciencedirect.com/science/article/pii/S0375947406010396>
- [541] Y. Oganessian, F. Abdullin, S. Dmitriev, J. Gostic, J. Hamilton, R. Henderson, M. Itkis, K. Moody, A. Polyakov, A. Ramayya, J. Roberto, K. Rykaczewski, R. Sagaidak, D. Shaughnessy, I. Shirokovsky, M. Stoyer, V. Subbotin, A. Sukhov,

- Y. Tsyganov, V. Utyonkov, A. Voinov, G. Vostokin, New insights into the  $^{243}\text{am} + ^{48}\text{ca}$  reaction products previously observed in the experiments on elements 113 and 115 and and 117, Physical Review Letters 108 (2012) 022502.
- [542] A. A. Kuznetsova, First tests of the grand gas-filled separator at the superheavy element factory (flerov laboratory of nuclear reactions, joint institute for nuclear research), Bulletin of the Russian Academy of Sciences: Physics 87 (8) (2023) 1105–1111. doi:10.3103/S1062873823702921.
- URL <https://doi.org/10.3103/S1062873823702921>
- [543] Y. T. Oganessian, Y. P. Tretyakov, A. S. Ilinov, Synthesis of neutron-deficient isotopes of fermium, kurchatovium and element with atomic number 106, Tech. Rep. JINR-D7-8099, JINR (1974).
- [544] S. Antalic, F. Heßberger, D. Ackermann, M. Block, S. Heinz, S. Hofmann, Z. Kalaninová, B. Kindler, M. Leino, B. Lommel, R. Mann, K. Nishio, Š. Šá, B. Sulignano, Nuclear structure of heavy  $n = 153$  isotones, Acta Phys. Pol. B 44 (3) (2013) 387–390.
- URL <https://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=44&page=387>
- [545] Y. T. Oganessian, M. Hussonnois, A. G. Demin, Y. P. Kharitonov, H. Bruchertseifer, O. Constantinescu, Y. S. Korotkin, S. P. Tretyakova, V. K. Utyonkov, I. V. Shirokovsky, J. Estevez, Experimental studies of the formation and radioactive decay of isotopes with  $z = 104\text{--}109$ , Radiochim.Acta 37 (1984) 113.
- [546] G. Münzenberg, P. Armbruster, G. Berthes, H. Folger, F. P. Heßberger, S. Hofmann, K. Poppensieker, W. Reisdorf, B. Quint, K. H. Schmidt, H. J. Schött, K. Sümmerer, I. Zychor, M. E. Leino, U. Gollerthan, E. Hanelt, Evidence for 264108, the heaviest known even-even isotope, Zeitschrift für Physik A Atomic Nuclei 324 (4) (1986) 489–490. doi:10.1007/BF01290935.
- URL <https://doi.org/10.1007/BF01290935>
- [547] N. Sato, H. Haba, T. Ichikawa, D. Kaji, Y. Kudou, K. Morimoto, K. Morita, K. Ozeki, T. Sumita, A. Yoneda, E. Ideguchi, H. Koura, A. Ozawa, T. Shinozuka, T. Yamaguchi, A. Yoshida, Production and decay properties of 264hs and 265hs, Journal of the Physical Society of Japan 80 (9) (2011) 094201. arXiv:<https://doi.org/10.1143/JPSJ.80.094201>. doi:10.1143/JPSJ.80.094201.
- URL <https://doi.org/10.1143/JPSJ.80.094201>
- [548] G. Münzenberg, P. Armbruster, H. Folger, P. F. Heßberger, S. Hofmann, J. Keller, K. Poppensieker, W. Reisdorf, K. H. Schmidt, H. J. Schött, M. E. Leino, R. Hingmann, The identification of element 108, Zeitschrift für Physik A Atoms and Nuclei 317 (2) (1984) 235–236. doi:10.1007/BF01421260.
- URL <https://doi.org/10.1007/BF01421260>
- [549] S. Hofmann, V. Ninov, F. P. Heßberger, P. Armbruster, H. Folger, G. Münzenberg, H. J. Schött, A. G. Popeko, A. V. Yeremin, A. N. Andreyev, S. Saro, R. Janik, M. Leino, Production and decay of 269110, Zeitschrift für Physik A Hadrons and Nuclei 350 (4) (1995) 277–280. doi:10.1007/BF01291181.
- URL <http://dx.doi.org/10.1007/BF01291181>
- [550] S. Hofmann, Production of heavy and super-heavy nuclei, Nuclear Physics A 616 (1) (1997) 370–379, radioactive Nuclear Beams. doi:[https://doi.org/10.1016/S0375-9474\(97\)00121-8](https://doi.org/10.1016/S0375-9474(97)00121-8).
- URL <https://www.sciencedirect.com/science/article/pii/S0375947497001218>
- [551] S. Hofmann, New elements - approaching  $z=114$ , Report Progress Physics 61 (1998) 639–689.
- [552] B. Štreicher, S. Antalic, v. Šaro, F. Heßberger, S. Hofmann, D. Ackermann, B. Kindler, I. Kojouharov, B. Lommel, R. Mann, B. Sulignano, P. Kuusiniemi, Alpha-gamma decay studies of 261sg, Acta Phys.

- Pol. B 38 (4) (2007) Ackermann, B. Kindler I. Kojouharov, B. Lommel, R. Mann, B. Sulignano.  
URL <https://www.actaphys.uj.edu.pl/fulltext?series=Reg&vol=38&page=1561>
- [553] D. Ackermann, F. P. Heßberger, S. Antalic, M. Block, H.-G. Burkhard, V. F. Comas, P. Greenlees, S. Heinz, J. A. Heredia, S. Hofmann, S. Ketelhut, J. Khuyagbaatar, B. Kindler, I. Kojouharov, M. Mazzocco, M. Leino, B. Lommel, R. Mann, J. Maurer, A. G. Popeko, J. Sorri, J. Uusitalo, A. V. Yeremin, Investigation of  $^{270}\text{Ds}$  and its decay products, GSI Sci. Rep. 2010 (2011) 200.
- [554] D. Ackermann, Superheavy elements at gsi — investigating exotic nuclear matter, Acta Phys. Pol. B 42 (2011) 577.  
URL <https://www.actaphys.uj.edu.pl/R/42/3/577/pdf>
- [555] Y. A. Lazarev, Y. V. Lobanov, Y. T. Oganessian, Y. S. Tsyganova, V. K. Utyonkov, F. S. Abdullin, S. Iliev, A. N. Polyakov, J. Rigol, I. V. Shirokovsky, V. G. Subbotin, A. M. Sukhov, G. V. Buklanov, B. N. Gikal, V. B. Kutner, A. N. Mezentsev, I. M. Sedykh, D. V. Vakatov, R. W. Lougheed, J. F. Wild, K. J. Moody, E. K. Hulet, New nuclide  $^{267}\text{108}$  produced by the  $^{238}\text{U} + ^{34}\text{S}$  reaction, Phys. Rev. Lett. 75 (1995) 1903–1906. doi:10.1103/PhysRevLett.75.1903.  
URL <https://link.aps.org/doi/10.1103/PhysRevLett.75.1903>
- [556] H. Ikezoe, T. Ikuta, S. Mitsuoka, Y. Nagame, I. Nishinaka, K. Tsukada, T. Ohtsuki, T. Kuzumaki, J. Lu, First evidence for a new spontaneous fission decay produced in the reaction  $^{30}\text{Si} + ^{238}\text{U}$ , The European Physical Journal A - Hadrons and Nuclei 2 (4) (1998) 379–382. doi:10.1007/s100500050134.  
URL <https://doi.org/10.1007/s100500050134>
- [557] T. N. Ginter, K. E. Gregorich, W. Loveland, D. M. Lee, U. W. Kirbach, R. Sudowe, C. M. Folden, J. B. Patin, N. Seward, P. A. Wilk, P. M. Zielinski, K. Aleklett, R. Eichler, H. Nitsche, D. C. Hoffman, Confirmation of production of element 110 by the  $^{208}\text{Pb}(^{64}\text{Ni}, n)$  reaction, Phys. Rev. C 67 (2003) 064609. doi:10.1103/PhysRevC.67.064609.  
URL <https://link.aps.org/doi/10.1103/PhysRevC.67.064609>
- [558] K. Nishio, S. Hofmann, F. P. Heßberger, D. Ackermann, S. Antalic, Y. Aritomo, V. F. Comas, C. E. Düllmann, A. Gorshkov, R. Graeger, K. Hagino, S. Heinz, J. A. Heredia, K. Hirose, H. Ikezoe, J. Khuyagbaatar, B. Kindler, I. Kojouharov, B. Lommel, R. Mann, S. Mitsuoka, Y. Nagame, I. Nishinaka, T. Ohtsuki, A. G. Popeko, S. Saro, M. Schädel, A. Türler, Y. Watanabe, A. Yakushev, A. V. Yeremin, Nuclear orientation in the reaction  $^{34}\text{S} + ^{238}\text{U}$  and synthesis of the new isotope  $^{268}\text{Hs}$ , Phys. Rev. C 82 (2010) 024611. doi:10.1103/PhysRevC.82.024611.  
URL <https://link.aps.org/doi/10.1103/PhysRevC.82.024611>
- [559] K. Morita, She research at riken/garis, Nuclear Physics A 944 (2015) 30–61, special Issue on Superheavy Elements. doi:<https://doi.org/10.1016/j.nuclphysa.2015.10.007>.  
URL <https://www.sciencedirect.com/science/article/pii/S0375947415002304>
- [560] B. B. Back, H. G. Clerc, R. R. Betts, B. G. Glagola, B. D. Wilkins, Observation of anisotropy in the fission decay of nuclei with vanishing fission barrier, Phys. Rev. Lett. 46 (1981) 1068–1071. doi:10.1103/PhysRevLett.46.1068.  
URL <https://link.aps.org/doi/10.1103/PhysRevLett.46.1068>
- [561] R. Graeger, D. Ackermann, M. Chelnokov, V. Chepigin, C. E. Düllmann, J. Dvorak, J. Even, A. Gorshkov, F. P. Heßberger, D. Hild, A. Hübler, E. Jäger, J. Khuyagbaatar, B. Kindler, J. V. Kratz, J. Krier, A. Kuznetsov, B. Lommel, K. Nishio, H. Nitsche, J. P. Omtvedt, O. Petrushkin, D. Rudolph, J. Runke,

- F. Samadani, M. Schädel, B. Schausten, A. Türler, A. Yakushev, Q. Zhi, **Experimental study of the  $^{238}\text{U}(^{36}\text{S},3 - 5n)^{269-271}\text{Hs}$  reaction leading to the observation of  $^{270}\text{Hs}$** , Phys. Rev. C 81 (2010) 061601. doi:[10.1103/PhysRevC.81.061601](https://doi.org/10.1103/PhysRevC.81.061601).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.81.061601>
- [562] Y. A. Lazarev, Y. V. Lobanov, Y. T. Oganessian, V. K. Utyonkov, F. S. Abdullin, A. N. Polyakov, J. Rigol, I. V. Shirokovsky, Y. S. Tsyganov, S. Iliev, V. G. Subbotin, A. M. Sukhov, G. V. Buklanov, B. N. Gikal, V. B. Kutner, A. N. Mezentsev, K. Subotic, J. F. Wild, R. W. Lougheed, K. J. Moody,  $\alpha$  decay of  $^{273}\text{No}$ : Shell closure at n=162, Phys. Rev. C 54 (1996) 620–625. doi:[10.1103/PhysRevC.54.620](https://doi.org/10.1103/PhysRevC.54.620).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.54.620>
- [563] M. Schädel, W. Brüchle, R. Dressler, B. Eichler, H. W. Gäggeler, R. Günther, K. E. Gregorich, D. C. Hoffman, S. Hübener, D. T. Jost, J. V. Kratz, W. Paulus, D. Schumann, S. Timokhin, N. Trautmann, A. Türler, G. Wirth, A. Yakushev, **Chemical properties of element 106 (seaborgium)**, Nature 388 (6637) (1997) 55–57. doi:[10.1038/40375](https://doi.org/10.1038/40375).
- URL <https://doi.org/10.1038/40375>
- [564] C. E. Düllmann, W. Brüchle, R. Dressler, K. Eberhardt, B. Eichler, R. Eichler, H. W. Gäggeler, T. N. Ginter, F. Glauß, K. E. Gregorich, D. C. Hoffman, E. Jäger, D. T. Jost, U. W. Kirbach, D. M. Lee, H. Nitsche, J. B. Patin, V. Pershina, D. Piguet, Z. Qin, M. Schädel, B. Schausten, E. Schimpf, H. J. Schött, S. Soverna, R. Sudowe, P. Thörle, S. N. Timokhin, N. Trautmann, A. Türler, A. Vahle, G. Wirth, A. B. Yakushev, P. M. Zielinski, **Chemical investigation of hassium (element 108)**, Nature 418 (6900) (2002) 859–862. doi:[10.1038/nature00980](https://doi.org/10.1038/nature00980).
- URL <https://doi.org/10.1038/nature00980>
- [565] J. Dvorak, W. Brüchle, C. E. Düllmann, Z. Dvořáková, K. Eberhardt, R. Eichler, E. Jäger, Y. Nagame, Z. Qin, M. Schädel, B. Schausten, E. Schimpf, R. Schuber, A. Semchenkov, P. Thörle, A. Türler, M. Wegrzecki, A. Yakushev, **Cross section limits for the  $^{248}\text{Cm}(^{25}\text{Mg}, 4n-5n)^{268,269}\text{Hs}$  reactions**, Phys. Rev. C 79 (2009) 037602. doi:[10.1103/PhysRevC.79.037602](https://doi.org/10.1103/PhysRevC.79.037602).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.79.037602>
- [566] A. Türler, **Nuclear structure and reaction studies near doubly magic  $^{270}\text{Hs}$** , Radiochim. Acta 100 (2) (2012) 75–83. doi:[doi:10.1524/ract.2011.1874](https://doi.org/10.1524/ract.2011.1874).
- URL <https://doi.org/10.1524/ract.2011.1874>
- [567] Y. T. Oganessian, V. K. Utyonkov, F. S. Abdullin, S. N. Dmitriev, R. Graeger, R. A. Henderson, M. G. Itkis, Y. V. Lobanov, A. N. Mezentsev, K. J. Moody, S. L. Nelson, A. N. Polyakov, M. A. Ryabinin, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, N. J. Stoyer, V. G. Subbotin, K. Subotic, A. M. Sukhov, Y. S. Tsyganov, A. Türler, A. A. Voinov, G. K. Vostokin, P. A. Wilk, A. Yakushev, **Synthesis and study of decay properties of the doubly magic nucleus  $^{270}\text{Hs}$  in the  $^{226}\text{Ra} + ^{48}\text{Ca}$  reaction**, Phys. Rev. C 87 (2013) 034605. doi:[10.1103/PhysRevC.87.034605](https://doi.org/10.1103/PhysRevC.87.034605).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.87.034605>
- [568] S. L. Nelson, C. M. Folden III, K. E. Gregorich, I. Dragojević, C. E. Düllmann, R. Eichler, M. A. Garcia, J. M. Gates, R. Sudowe, H. Nitsche, **Comparison of complementary reactions for the production of  $^{261,262}\text{Bh}$** , Phys. Rev. C 78 (2008) 024606. doi:[10.1103/PhysRevC.78.024606](https://doi.org/10.1103/PhysRevC.78.024606).
- URL <https://link.aps.org/doi/10.1103/PhysRevC.78.024606>
- [569] G. Münzenberg, S. Hofmann, F. P. Heßberger, H. Folger, V. Ninov, K. Poppensieker, A. B. Quint, W. Reisdorf, H. J. Schött, K. Sümmerer, P. Armbruster,

- M. E. Leino, D. Ackermann, U. Gollerthan, E. Hanelt, W. Morawek, Y. Fujita, T. Schwab, A. Türler, [New results on element 109](#), Zeitschrift für Physik A Atomic Nuclei 330 (4) (1988) 435–436. [doi:10.1007/BF01290131](#).  
 URL <https://doi.org/10.1007/BF01290131>
- [570] S. Hofmann, F. P. Heßberger, V. Ninov, P. Armbruster, G. Münzenberg, C. Stodel, A. G. Popeko, A. V. Yeremin, S. Saro, M. Leino, [Excitation function for the production of 265108 and 266109](#), Zeitschrift für Physik A Hadrons and Nuclei 358 (4) (1997) 377–378. [doi:10.1007/s002180050343](#).  
 URL <https://doi.org/10.1007/s002180050343>
- [571] K. Morita, K. Morimoto, D. Kaji, H. Haba, E. Ideguchi, J. C. Peter, R. Kanungo, K. Katori, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, I. Tanihata, H. Xu, A. V. Yeremin, A. Yoneda, A. Yoshida, Y.-L. Zhao, T. Zheng, S. Goto, F. Tokanai, [Production and decay properties of 272111 and its daughter nuclei](#), Journal of the Physical Society of Japan 73 (7) (2004) 1738–1744. [arXiv:https://doi.org/10.1143/JPSJ.73.1738](#), [doi:10.1143/JPSJ.73.1738](#).  
 URL <https://doi.org/10.1143/JPSJ.73.1738>
- [572] K. Morimoto, K. Morita, D. Kaji, A. Yoneda, H. Haba, T. Akiyama, N. Sato, S. Goto, E. Ideguchi, K. Katori, H. Kikunaga, H. Koura, H. Kudo, T. Ohnishi, A. Ozawa, T. Suda, K. Sueki, F. Tokanai, T. Yamaguchi, A. Yoshida, [Experiments on synthesis of the heaviest elements at riken](#), AIP Conference Proceedings 853 (1) (2006) 253–258. [arXiv:https://aip.scitation.org/doi/pdf/10.1063/1.2338386](#), [doi:10.1063/1.2338386](#).  
 URL <https://aip.scitation.org/doi/abs/10.1063/1.2338386>
- [573] K. Morita, K. Morimoto, D. Kaji, H. Haba, K. Ozeki, Y. Kudou, N. Sato, T. Sumita, A. Yoneda, T. Ichikawa, Y. Fujimori, S. Goto, E. Ideguchi, Y. Kasamatsu, K. Katori, Y. Komori, H. Koura, H. Kudo, K. Ooe, A. Ozawa, F. Tokanai, K. Tsukada, T. Yamaguchi, A. Yoshida, [Decay properties of 266bh and 262db produced in the 248cm+23na reaction—further confirmation of the 278113 decay chain—](#), AIP Conference Proceedings 1238 (1) (2010) 331–336. [arXiv:https://aip.scitation.org/doi/pdf/10.1063/1.3455961](#), [doi:10.1063/1.3455961](#).  
 URL <https://aip.scitation.org/doi/abs/10.1063/1.3455961>
- [574] R. Eichler, W. Brüchle, R. Dressler, C. E. Düllmann, B. Eichler, H. W. Gäggeler, K. E. Gregorich, D. C. Hoffman, S. Hübener, D. T. Jost, U. W. Kirbach, C. A. Laue, V. M. Lavanchy, H. Nitsche, J. B. Patin, D. Piguet, M. Schädel, D. A. Shaughnessy, D. A. Strellis, S. Taut, L. Tobler, Y. S. Tsyanov, A. Türler, A. Vahle, P. A. Wilk, A. B. Yakushev, [Chemical characterization of bohrium \(element 107\)](#), Nature 407 (6800) (2000) 63–65. [doi:10.1038/35024044](#).  
 URL <https://doi.org/10.1038/35024044>
- [575] A. Türler, [Heavy-element chemistry –status and perspectives](#), The European Physical Journal A 15 (1) (2002) 271–274. [doi:10.1140/epja/i2001-10266-0](#).  
 URL <https://doi.org/10.1140/epja/i2001-10266-0>
- [576] Y. T. Oganessian, F. S. Abdullin, P. D. Bailey, D. E. Benker, M. E. Bennett, S. N. Dmitriev, J. G. Ezold, J. H. Hamilton, R. A. Henderson, M. G. Itkis, Y. V. Lobanov, A. N. Mezentsev, K. J. Moody, S. L. Nelson, A. N. Polyakov, C. E. Porter, A. V. Ramayya, F. D. Riley, J. B. Roberto, M. A. Ryabinin, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. A. Stoyer, V. G. Subbotin, R. Sudowe, A. M. Sukhov, R. Taylor, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, P. A. Wilk, [Eleven new heaviest isotopes of elements z = 105 to z = 117 identified among the products of  \$^{24}\text{bk} + ^{48}\text{ca}\$  reactions](#), Physical Review C 83 (2011) 054315.

- [577] S. N. Dmitriev, N. V. Aksenov, Y. V. Albin, G. A. Bozhikov, M. L. Chelnokov, V. I. Chepygin, R. Eichler, A. V. Isaev, D. E. Katrasev, V. Y. Lebedev, O. N. Malyshev, O. V. Petrushkin, L. S. Porobanuk, M. A. Ryabinin, A. V. Sabel'nikov, E. A. Sokol, A. V. Svirikhin, G. Y. Starodub, I. Usoltsev, G. K. Vostokin, A. V. Yeremin, *Pioneering experiments on the chemical properties of element 113*, Mendeleev Communications 24 (5) (2014) 253–256. doi:<https://doi.org/10.1016/j.mencom.2014.09.001>. URL <https://www.sciencedirect.com/science/article/pii/S0959943614001035>
- [578] N. V. Aksenov, P. Steinegger, F. S. Abdullin, Y. V. Albin, G. A. Bozhikov, V. I. Chepigin, R. Eichler, V. Y. Lebedev, A. S. Madumarov, O. N. Malyshev, O. V. Petrushkin, A. N. Polyakov, Y. A. Popov, A. V. Sabel'nikov, R. N. Sagaidak, I. V. Shirokovsky, M. V. Shumeiko, G. Y. Starodub, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, A. V. Yeremin, S. N. Dmitriev, *On the volatility of nihonium (nh, z = 113)*, The European Physical Journal A 53 (7) (2017) 158. doi:<https://doi.org/10.1140/epja/i2017-12348-8>. URL <https://doi.org/10.1140/epja/i2017-12348-8>
- [579] Y. T. Oganessian, F. S. Abdullin, C. Alexander, J. Binder, R. A. Boll, S. N. Dmitriev, J. Ezold, K. Felker, J. M. Gostic, R. K. Grzywacz, J. H. Hamilton, R. A. Henderson, M. G. Itkis, K. Miernik, D. Miller, K. J. Moody, A. N. Polyakov, A. V. Ramayya, J. B. Roberto, M. A. Ryabinin, K. P. Rykaczewski, R. N. Sagaidak, D. A. Shaughnessy, I. V. Shirokovsky, M. V. Shumeiko, M. A. Stoyer, N. J. Stoyer, V. G. Subbotin, A. M. Sukhov, Y. S. Tsyanov, V. K. Utyonkov, A. A. Voinov, G. K. Vostokin, *Production and decay of the heaviest nuclei  $^{293,294}117$  and  $^{294}118$* , Phys. Rev. Lett. 109 (2012) 162501. doi:[10.1103/PhysRevLett.109.162501](https://doi.org/10.1103/PhysRevLett.109.162501).
- [580] D. Ackermann, *Superheavy Elements – A Probe for Nuclear Matter at the Extremes*, Progress of Theoretical Physics Supplement 196 (2012) 255–262. arXiv:<https://academic.oup.com/ptps/article-pdf/doi/10.1143/PTPS.196.255/5342867/196-255.pdf>, doi:[10.1143/PTPS.196.255](https://doi.org/10.1143/PTPS.196.255). URL <https://doi.org/10.1143/PTPS.196.255>
- [581] G. Münzenberg, *From bohrium to copernicium and beyond {SHE} research at {SHIP}*, Nuclear Physics A 944 (2015) 5 – 29, special Issue on Superheavy Elements. doi:<http://dx.doi.org/10.1016/j.nuclphysa.2015.06.008>. URL <https://www.sciencedirect.com/science/article/pii/S0375947415001372>
- [582] and, , , , , , , , , , , and, *Observation of the superheavy nuclide  $^{271}\text{ds}$* , Chinese Physics Letters 29 (1) (2012) 012502. doi:[10.1088/0256-307X/29/1/012502](https://doi.org/10.1088/0256-307X/29/1/012502). URL <https://dx.doi.org/10.1088/0256-307X/29/1/012502>
- [583] H. Gäggeler, W. Brüchle, C. Düllmann, R. Dressler, K. Eberhardt, B. Eichler, R. Eichler, C. Folden, T. Ginter, F. Glaus, K. Gregorich, F. Haenssler, D. Hoffman, E. Ja"ger, D. Jost, U. Kirbach, J. Kratz, H. Nitsche, J. Patin, V. Pershina, D. Piguet, Z. Qin, U. Rieth, M. Schädel, E. Schimpf, B. Schausten, S. Soverna, R. Sudowe, P. Tho"rle, N. Trautmann, A. Tu"rler, A. Vahle, P. Wilk, G. Wirth, A. Yakushev, A. von Zweidorff, *Chemical and nuclear studies of hassium and element 112*, Nuclear Physics A 734 (2004) 208–212. doi:<https://doi.org/10.1016/j.nuclphysa.2004.01.036>. URL <https://www.sciencedirect.com/science/article/pii/S0375947404000417>
- [584] Y. T. Oganessian, S. N. Dmitriev, A. V. Yeremin,

- N. V. Aksenov, G. A. Bozhikov, V. I. Chepigin, M. L. Chelnokov, V. Y. Lebedev, O. N. Malyshev, O. V. Petrushkin, S. V. Shishkin, A. I. Svirikhin, E. E. Tereshatov, G. K. Vostokin, [Attempt to produce the isotopes of element 108 in the fusion reaction  \$^{136}\text{Xe} + ^{136}\text{Xe}\$](#) , Phys. Rev. C 79 (2009) 024608. doi:[10.1103/PhysRevC.79.024608](https://doi.org/10.1103/PhysRevC.79.024608).
- URL <http://link.aps.org/doi/10.1103/PhysRevC.79.024608>
- [585] C. E. Düllmann, M. Schädel, A. Yakushev, A. Türler, K. Eberhardt, J. V. Kratz, D. Ackermann, L.-L. Andersson, M. Block, W. Brüchle, J. Dvorak, H. G. Essel, P. A. Ellison, J. Even, J. M. Gates, A. Gorshkov, R. Graeger, K. E. Gregorich, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. Hild, A. Hübner, E. Jäger, J. Khuyagbaatar, B. Kindler, J. Krier, N. Kurz, S. Lahiri, D. Liebe, B. Lommel, M. Maiti, H. Nitsche, J. P. Omtvedt, E. Parr, D. Rudolph, J. Runke, B. Schausten, E. Schimpf, A. Semchenkov, J. Steiner, P. Thörle-Pospiech, J. Uusitalo, M. Wegrzecki, N. Wiehl, [Production and decay of element 114: High cross sections and the new nucleus  \$^{277}\text{Hs}\$](#) , Phys. Rev. Lett. 104 (2010) 252701. doi:[10.1103/PhysRevLett.104.252701](https://doi.org/10.1103/PhysRevLett.104.252701).
- URL <http://link.aps.org/doi/10.1103/PhysRevLett.104.252701>
- [586] S. Hofmann, S. Heinz, R. Mann, J. Maurer, J. Khuyagbaatar, D. Ackermann, S. Antalic, W. Barth, M. Block, H. Burkhard, V. Comas, L. Dahl, K. Eberhardt, J. Gostic, R. Henderson, J. Heredia, F. P. Heßberger, J. Kenneally, B. Kindler, I. Kojouharov, J. Kratz, R. Lang, M. Leino, B. Lommel, K. Moody, G. Münzenberg, S. Nelson, K. Nishio, A. Popelo, J. Runke, S. Saro, D. Shaughnessy, M. Stoyer, P. Thörle-Pospiech, K. Tinschert, N. Trautmann, J. Uusitalo, P. Wilk, A. Yeremin, The reaction  $^{48}\text{Ca} + ^{248}\text{Cr} \rightarrow ^{296}\text{116}^*$  studied at the gsi-ship, European Physics Journal A 48 (2012) 62.
- [587] G. Münzenberg, P. Armbruster, F. P. Heßberger, S. Hofmann, K. Poppensieker, W. Reisdorf, J. H. R. Schneider, W. F. W. Schneider, K. H. Schmidt, C. C. Sahm, D. Vermeulen, [Observation of one correlated  \$\alpha\$ -decay in the reaction  \$^{58}\text{Fe}\$  on  \$^{209}\text{Bi} \rightarrow ^{267}\text{109}\$](#) , Zeitschrift für Physik A Atoms and Nuclei 309 (1) (1982) 89–90. doi:[10.1007/BF01420157](https://doi.org/10.1007/BF01420157).
- URL <https://doi.org/10.1007/BF01420157>
- [588] G. Münzenberg, W. Reisdorf, S. Hofmann, Y. K. Agarwal, F. P. Heßberger, K. Poppensieker, J. R. H. Schneider, W. F. W. Schneider, K. H. Schmidt, H. J. Schött, P. Armbruster, C. C. Sahm, D. Vermeulen, [Evidence for element 109 from one correlated decay sequence following the fusion of  \$^{58}\text{Fe}\$  with  \$^{209}\text{Bi}\$](#) , Zeitschrift für Physik A Atoms and Nuclei 315 (2) (1984) 145–158. doi:[10.1007/BF01419373](https://doi.org/10.1007/BF01419373).
- URL <https://doi.org/10.1007/BF01419373>
- [589] G. Münzenberg, [Heavy element production and limits to fusion](#), Nuclear Physics A 502 (1989) 571–590. doi:[https://doi.org/10.1016/0375-9474\(89\)90690-8](https://doi.org/10.1016/0375-9474(89)90690-8).
- URL <https://www.sciencedirect.com/science/article/pii/0375947489906908>
- [590] K. Morita, [Experiments on searching for the heaviest elements](#), Nuclear Physics A 805 (1) (2008) 172c–179c, iNPC 2007. doi:<https://doi.org/10.1016/j.nuclphysa.2008.02.238>.
- URL <https://www.sciencedirect.com/science/article/pii/S0375947408003394>
- [591] F. Kondev, G. Dracoulis, T. Kibédi, [Configurations and hindered decays of isomers in deformed nuclei with  \$A > 100\$](#) , Atomic Data and Nuclear Data Tables 103–104 (2015) 50 – 105. doi:<http://dx.doi.org/10.1016/j.adt.2015.01.001>.
- URL <https://www.sciencedirect.com/science/article/pii/S0092640X15000029>
- [592] P. J. Karol, H. Nakahara, B. W. Petley, E. Vogt, [On the claims for discovery of elements 110, 111, 112,](#)

- 114, 116, and 118 (iupac technical report);, Pure and Applied Chemistry 75 (10) (2003) 1601–1611. doi: [doi:10.1351/pac200375101601](https://doi.org/10.1351/pac200375101601). URL <https://doi.org/10.1351/pac200375101601>
- [593] K. E. Gregorich, W. Loveland, D. Peterson, P. M. Zielinski, S. L. Nelson, Y. H. Chung, C. E. Düllmann, C. M. Folden III, K. Aleklett, R. Eichler, D. C. Hoffman, J. P. Omtvedt, G. K. Pang, J. M. Schwantes, S. Soverná, P. Sprunger, R. Sudowe, R. E. Wilson, H. Nitsche, Attempt to confirm superheavy element production in the  $^{48}\text{Ca} + ^{238}\text{U}$  reaction, Phys. Rev. C 72 (2005) 014605. doi: [doi:10.1103/PhysRevC.72.014605](https://doi.org/10.1103/PhysRevC.72.014605). URL <https://link.aps.org/doi/10.1103/PhysRevC.72.014605>
- [594] K. Morita, Superheavy research at riken, International Journal of Modern Physics E 18 (10) (2009) 2175–2178. arXiv:<https://doi.org/10.1142/S0218301309014500>, doi: [doi:10.1142/S0218301309014500](https://doi.org/10.1142/S0218301309014500). URL <https://doi.org/10.1142/S0218301309014500>
- [595] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tszyganov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, J. B. Patin, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, P. A. Wilk, R. W. Lougheed, New elements from dubna, The European Physical Journal A - Hadrons and Nuclei 25 (1) (2005) 589–594. doi: [doi:10.1140/epjad/i2005-06-134-9](https://doi.org/10.1140/epjad/i2005-06-134-9). URL <https://doi.org/10.1140/epjad/i2005-06-134-9>
- [596] L. Stavsetra, K. E. Gregorich, J. Dvorak, P. A. Ellison, I. Dragojević, M. A. Garcia, H. Nitsche, Independent verification of element 114 production in the  $^{48}\text{Ca} + ^{242}\text{Pu}$  reaction, Phys. Rev. Lett. 103 (2009) 132502. doi: [doi:10.1103/PhysRevLett.103.132502](https://doi.org/10.1103/PhysRevLett.103.132502). URL <https://link.aps.org/doi/10.1103/PhysRevLett.103.132502>
- [597] R. Eichler, N. V. Aksenov, Y. V. Albin, A. V. Belozarov, G. A. Bozhikov, V. I. Chepigin, S. N. Dmitriev, R. Dressler, H. W. Gäggeler, V. A. Gorshkov, G. Henderson, et al., Indication for a volatile element 114:, rca - Radiochimica Acta 98 (3) (2010) 133–139. doi: [doi:10.1524/ract.2010.1705](https://doi.org/10.1524/ract.2010.1705). URL <https://doi.org/10.1524/ract.2010.1705>
- [598] K. Morita, K. Morimoto, D. Kaji, H. Haba, Y. Wakabayashi, M. Takeyama, S. Yamaki, K. Tanaka, H. Hasebe, M. Huang, J. Kanaya, M. Murakami, A. Yoneda, A. Yoshida, T. Yamaguchi, T. Y. F. Tokanai, Z. Gan, L. Ma, H. Geissel, S. Hofmann, J. Maurer, K. Fujita, Y. Narikiyo, T. Tanaka, S. Yamamoto, M. Asai, K. Katori, Measurement of the  $^{248}\text{cm} + ^{48}\text{ca}$  fusion reaction products at riken garis, Riken accelerator progress report 2013, p.xi, RIKEN (2014).
- [599] A. Yakushev, J. M. Gates, A. Türler, M. Schädel, C. E. Düllmann, D. Ackermann, L.-L. Andersson, M. Block, W. Brüchle, J. Dvorak, K. Eberhardt, H. G. Essel, J. Even, U. Forsberg, A. Gorshkov, R. Graeger, K. E. Gregorich, W. Hartmann, R.-D. Herzberg, F. P. Heßberger, D. Hild, A. Hübner, E. Jäger, J. Khuyagbaatar, B. Kindler, J. V. Kratz, J. Krier, N. Kurz, B. Lommel, L. J. Niewisch, H. Nitsche, J. P. Omtvedt, E. Parr, Z. Qin, D. Rudolph, J. Runke, B. Schausten, E. Schimpf, A. Semchenkov, J. Steiner, P. Thörle-Pospiech, J. Uusitalo, M. Wegrzecki, N. Wiehl, Superheavy element flerovium (element 114) is a volatile metal, Inorganic Chemistry 53 (3) (2014) 1624–1629. doi: [doi:10.1021/ic4026766](https://doi.org/10.1021/ic4026766). URL <https://doi.org/10.1021/ic4026766>
- [600] Y. Oganessian, V. Utyonkov, Y. Lobanov, F. Abdullin, A. Polyakov, I. Shirokovsky, Y. Tszyganov,

- G. Gulbekian, S. Bogomolov, B. Gikal, A. Mezentsev, S. Iliev, V. Subbotin, A. Sukhov, A. Voinov, G. Buklanov, K. Subotic, V. Zagrebaev, M. Itkis, J. Patin, K. Moody, J. Wild, M. Stoyer, N. Stoyer, D. Shaughnessy, J. Kenneally, R. Lougheed, Heavy element research at dubna, Nuclear Physics A 734 (2004) 109–123. doi:<https://doi.org/10.1016/j.nuclphysa.2004.01.020>. URL <https://www.sciencedirect.com/science/article/pii/S0375947404000259>
- [601] Y. T. Oganessian, V. K. Utyonkov, Y. V. Lobanov, F. S. Abdullin, A. N. Polyakov, I. V. Shirokovsky, Y. S. Tsyanov, G. G. Gulbekian, S. L. Bogomolov, B. N. Gikal, A. N. Mezentsev, S. Iliev, V. G. Subbotin, A. M. Sukhov, A. A. Voinov, G. V. Buklanov, K. Subotic, V. I. Zagrebaev, M. G. Itkis, K. J. Moody, J. F. Wild, M. A. Stoyer, N. J. Stoyer, D. A. Shaughnessy, J. M. Kenneally, J. B. Patin, R. W. Lougheed, Synthesis of Superheavy Nuclei in  $^{48}\text{Ca}$ -Induced Reactions, Progress of Theoretical Physics Supplement 154 (2004) 406–416. arXiv:<https://academic.oup.com/ptps/article-pdf/doi/10.1143/PTPS.154.406/5224956/154-406.pdf>, doi:[10.1143/PTPS.154.406](https://doi.org/10.1143/PTPS.154.406). URL <https://doi.org/10.1143/PTPS.154.406>
- [602] Y. Oganessian, Heavy element research at flnr (dubna), The European Physical Journal A 42 (3) (2009) 361. doi:[10.1140/epja/i2009-10861-y](https://doi.org/10.1140/epja/i2009-10861-y). URL <https://doi.org/10.1140/epja/i2009-10861-y>
- [603] U. Forsberg, D. Rudolph, C. Fahlander, P. Golubev, L. Sarmiento, S. Aberg, M. Block, C. Düllmann, F. Heßberger, J. Kratz, A. Yakushev, A new assessment of the alleged link between element 115 and element 117 decay chains, Physics Letters B 760 (2016) 293–296. doi:<https://doi.org/10.1016/j.physletb.2016.07.008>.
- URL <https://www.sciencedirect.com/science/article/pii/S0370269316303495>
- [604] R. Dressler, R. Eichler, D. Schumann, S. Shishkin, Long-term  $\alpha$ - and spontaneous fission measurement of a Rf/Db sample chemically prepared in a  $^{48}\text{Ca}$  on  $^{243}\text{Am}$  experiment, Phys. Rev. C 79 (2009) 054605. doi:[10.1103/PhysRevC.79.054605](https://doi.org/10.1103/PhysRevC.79.054605). URL <https://link.aps.org/doi/10.1103/PhysRevC.79.054605>
- [605] C. N. Davids, J. D. Larson, The argonne fragment mass analyzer, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 40-41 (1989) 1224–1228. doi:[https://doi.org/10.1016/0168-583X\(89\)90624-1](https://doi.org/10.1016/0168-583X(89)90624-1). URL <https://www.sciencedirect.com/science/article/pii/0168583X89906241>
- [606] A. Ajayakumar, J. Romans, M. Authier, Y. Balasmeh, A. Brizard, F. Boumard, L. Caceres, J.-F. Cam, A. Claessens, S. Damoy, P. Delahaye, P. Desrues, W. Dong, A. Drouart, P. Duchesne, R. Ferrer, X. Fléchard, S. Fransoo, P. Gangnant, S. Geldhof, R. P. de Groote, F. Ivandikov, N. Lebesne, R. Leroy, J. Lory, F. Lutton, V. Manea, Y. Merrer, I. Moore, A. Ortiz-Cortes, B. Osmond, J. Piot, O. Pochon, S. Raeder, A. de Roubin, H. Savajols, D. Studer, E. Traykov, J. Uusitalo, C. Vandamme, P. Van den Bergh, P. Van Duppen, K. Wendt, In-gas-jet laser spectroscopy with s3-leb, Nuclear Instruments and Methods in Physics Research Section B: Beam Interactions with Materials and Atoms 539 (2023) 102–107. doi:<https://doi.org/10.1016/j.nimb.2023.03.020>. URL <https://www.sciencedirect.com/science/article/pii/S0168583X23001003>