CYCLIC COVERS OF AN ALGEBRAIC CURVE FROM AN ADELIC VIEWPOINT

LUIS MANUEL NAVAS VICENTE[®] AND FRANCISCO J. PLAZA MARTÍN[®]

ABSTRACT. We propose an algebraic method for the classification of branched Galois covers of a curve X focused on studying Galois ring extensions of its geometric adele ring \mathbb{A}_X . As an application, we deal with cyclic covers; namely, we determine when a given cyclic ring extension of \mathbb{A}_X comes from a corresponding cover of curves $Y \to X$, which is reminiscent of a Grunwald-Wang problem, and also determine when two covers yield isomorphic ring extensions, which is known in the literature as an equivalence problem. This completely algebraic method permits us to recover ramification, certain analytic data such as rotation numbers, and enumeration formulas for covers.

1. INTRODUCTION

The classification of finite extensions of the function field Σ of an algebraic curve X is a long standing problem with roots in the foundational work of Riemann, Klein, Hurwitz, and many others. The standard techniques for dealing with this question ultimately rest upon analytic methods in the theory of Riemann surfaces, appearing often in the guise of complex algebraic geometry.

Here we present a purely algebraic alternative approach based on the use of algebraic extensions of the geometric adele ring \mathbb{A}_X of an algebraic curve X over an algebraically closed base field \mathbb{k} . The motivation is the following fact: a finite Galois cover $Y \to X$ with group G naturally endows \mathbb{A}_Y with an \mathbb{A}_X -algebra structure together with an action of G, which makes \mathbb{A}_Y a Galois ring extension of \mathbb{A}_X with group G in the sense of Chase-Harrison-Rosenberg (CHR) [4].

The main object in our toolkit is the Harrison set $\mathbb{H}(R, G)$, defined for any commutative ring R and finite group G (see §1.B). It consists of isomorphism classes of Galois ring extensions with fixed Galois group G. For abelian groups, $\mathbb{H}(R, G)$ can be itself endowed with a group structure.

Although there is a geometric interpretation of the Harrison set in the language of torsors and the algebraic fundamental group, we choose to work with the Harrison group $\mathbb{H}(\mathbb{A}_X, G)$ rather than *G*-torsors over $\operatorname{Spec}(\mathbb{A}_X)$, since describing the full spectrum requires ultrafilters, which unnecessarily complicates matters. In our method we can avoid having to deal with the complete spectrum ([16, Lemma 3.13, Theorem 3.14]). For the interested reader, the survey [18] discusses the connections between the algebraic foundation of CHR and subsequent generalizations or alternative points of view, including the Galois-Grothendieck theory.

²⁰²⁰ Mathematics Subject Classification. 14H30 (Primary) 13B05 14H05, 11R56 (Secondary). Key words and phrases. Coverings of curves, algebraic curves and function fields, Galois theory of commutative rings, geometric adele ring, Kummer theory.

Author 1 ORCID: 0000-0002-5742-8679 Author 2 ORCID: 0000-0002-5532-7567.

Research of both authors supported by grant PID2023-150787NB-I00, and of the first author also by grant PID2021-124332NB-C22, both from the Ministerio de Ciencia e Innovación (Government of Spain).

The authors have no relevant financial or non-financial interests to disclose. No datasets are involved in any way in this paper, which only involves research in pure mathematics.

Using these algebraic methods, we classify the cyclic covers of X of prime order different from char(k). As must be the case, we recover the bijection between algebraic curves which are finite G-Galois covers branched over n points and conjugacy classes of n-tuples (g_1, \ldots, g_n) generating G and satisfying $\prod_i g_i = 1$. The essential tool for this is the Kummer sequence for ring extensions, described for example in [8, Theorem 5.4] and, from the point of view of étale cohomology in [14, Chapter III, Proposition 4.11]. For readers accustomed to the latter, we can show that $\mathbb{H}(R, \mathcal{C}_n) \simeq H^1_{\text{et}}(\operatorname{Spec} R, \mathcal{C}_n)$ when n is coprime with char(R), thus establishing an equivalence with our approach (but see the remark above regarding $\operatorname{Spec}(\mathbb{A}_X)$).

The paper is structured around three main themes: the first is determining when, for a prime p, given a \mathcal{C}_p -Galois ring extension of \mathbb{A}_X , there exists a \mathcal{C}_p -Galois cover of curves $Y \to X$ (§2.A) which gives rise to it in the sense explained below. We shall refer to this as the *existence problem*.

The second deals with what is known as an *equivalence problem*, in this case, for function fields of algebraic curves over algebraically closed fields (see [20] for a current survey of many instances of this question), namely, when two such covers induce isomorphic extensions of \mathbb{A}_X (§2.B).

The third, and final question we will deal with, is how geometric data (such as ramification, analytic rotation numbers, enumeration of covers, etc.) can be recovered from the constructions by completely algebraic means ($\S3$ and $\S4$).

The first two problems are solved in §2, by relating the Kummer theory of the function field Σ with that of its adele ring \mathbb{A}_X . To be precise, the map induced by tensoring with \mathbb{A}_X induces a group homomorphism $\mathbb{H}(\Sigma, \mathcal{C}_p) \to \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)$. The determination of its kernel and image by the exact sequence (2.11) in Theorem 2.10 solves the existence problem and also provides the key to answering the equivalence problem, which admits various forms (Corollaries 2.19, 2.23, and 2.25).

Motivated by the fact that the notion of ramification locus appears naturally in the correspondence between finite extensions of the function field Σ and covers of the curve X, we are led to consider a refinement of (2.11) in §3. For a finite nonempty subset \mathfrak{R} of closed points of X representing the ramification locus, we study the Harrison group of the ring corresponding to the affine curve $X \setminus \mathfrak{R}$. The main results are encapsulated in the commutative cube (3.12) of Theorem 3.11 and applied in §3.B to obtain various filtrations of Harrison groups by ramification.

The third question is taken up in the last section of the paper (§4), which presents various situations illustrating how our algebraic approach indeed recovers interesting geometric information. In §4.A we show how our study of the Harrison group corresponds with the classification of *p*-cyclic Galois covers in [1] and [5]. In §4.B we recover the enumeration formulas for *p*-cyclic covers with specified ramification given in [10, 12, 13]. In §4.C we show how our methods allow one to define so-called rotation data purely algebraically in terms of the local Kummer symbols at ramified points. We prove that for $\Bbbk = \mathbb{C}$, our definition coincides with the usual analytic notion (see for example [7]).

We believe that the approach presented here based on studying Galois extensions of the adeles, and in general, of finite \mathbb{A}_X -algebras, provides an interesting alternative algebraic perspective on several geometric problems, and, although limited here to cyclic extensions as an illustration, can be adapted to more general situations.

1.A. Geometric adeles. Let X be a projective, irreducible, non-singular curve over an algebraically closed field k. Let Σ be the function field of X. The notation $x \in X$ will be used to denote a closed point. $A_x, K_x, \mathfrak{m}_x, v_x$, will denote respectively the completion of the valuation ring at $x \in X$, its quotient field, maximal ideal,

and the completed valuation. Since \mathbb{k} is algebraically closed, the residue field is $A_x/\mathfrak{m}_x = \mathbb{k}$.

Recall that the adele ring \mathbb{A}_X is the restricted direct product

$$\mathbb{A}_X \coloneqq \prod_{x \in X}' (K_x, A_x) = \{ (\alpha_x)_{x \in X} : \alpha_x \in A_x \text{ for almost all } x \in X \}$$

equipped with the restricted product topology. \mathbb{A}_X is the direct limit over finite subsets F of (closed) points of X containing an arbitrary fixed F_0 ,

$$\mathbb{A}_X \simeq \varinjlim_{F \supseteq F_0} \mathbb{A}_{X,F}, \qquad \mathbb{A}_{X,F} \coloneqq \prod_{x \in F} K_x \times \prod_{x \in X \setminus F} A_x.$$
(1.1)

The limit topology is generated by the neighborhood basis at 0 consisting of the sets $\prod_x \mathfrak{m}_x^{n_x}$, where $(n_x)_{x \in X}$ runs over collections of non-negative integers n_x such that $n_x = 0$ for almost all x. It coincides with the restricted product topology. A third characterization is in terms of the k-vector subspaces commensurable with $\mathbb{A}_X^+ := \prod_{x \in X} A_x$ as considered by Tate.

The idele group \mathbb{I}_X of invertible elements of \mathbb{A}_X is the restricted product of K_x^* with respect to the unit groups A_x^* and also the direct limit $\varinjlim \mathbb{I}_{X,F}$ where $\mathbb{I}_{X,F} := \mathbb{A}_{X,F}^*$.

1.B. Galois theory of rings. Let us give a brief overview of the Galois theory of commutative ring extensions. Standard sources for this include [1, 4, 8]. A far-reaching survey of this topic and how it has evolved may be found in [18]. We have extracted the following definitions and results from our previous paper [16] and copied them here for the reader's convenience.

We begin by recalling the definition of a Galois extension of commutative rings (taken from [4, Theorem 1.3], which generalizes the classical case of fields.

Definition 1.2 (Galois extension of rings). A Galois extension of a commutative ring R consists of a pair (S, G), where S is a commutative ring extension of R and G is a finite group which acts faithfully on S by R-algebra automorphisms, with invariants $S^G = R$, and satisfying one of the following equivalent conditions (we only list the ones which we will use later on):

- **1.** S is a separable R-algebra and the elements of G are pairwise strongly distinct.
- **2.** The map $h: S \otimes S \to \text{Maps}(G, S)$ given by $h(s \otimes t)(g) = sg(t)$ is an S-algebra isomorphism.
- **3.** If $g \in G$ with $g \neq 1$, for any maximal ideal \mathfrak{m} of S there is some $s \in S$ such that $g(s) \not\equiv s \mod \mathfrak{m}$.

In this case we say that S is a Galois ring extension of R with Galois group G, or simply a G-Galois extension of R.

Recall that the notion of strongly distinct for a pair of morphisms of commutative rings $f, g: S \to T$ means that for every nonzero idempotent $e \in T$ there exist $s \in S$ such that $f(s)e \neq g(s)e$.

We will denote Galois extensions S of the commutative ring R with group G acting via R-algebra automorphisms as pairs (S, G).

Definition 1.3 (Kummerian ring). Let n be a natural number. A commutative ring R is n-Kummerian if n is prime to char(R) and its unit group R^* contains a distinguished n-cyclic subgroup μ_n .

A field K with char(K) prime to n containing the n-th roots of unity $\mu_n = \mu_n(K^*)$, is n-Kummerian and this is the only possible choice of subgroup.

In our context, having fixed an algebraically closed field k, choosing $\mu_n \coloneqq \mu_n(\mathbb{k}^*)$, the group of *n*th roots of unity in k, induces the structure of an *n*-Kummerian ring on any k-algebra R of characteristic prime to n. This is compatible with k-algebra morphisms.

Note that for the adele ring, $\mathbb{k} \subseteq \Sigma$ is diagonally embedded in \mathbb{A}_X and we also have copies of \mathbb{k} in each completion K_x . This is an example of how there may be infinitely many choices of subgroups $\mu_n \subseteq \mathbb{R}^*$, highlighting the need to specify one.

A particular type of cyclic Galois ring extensions of a Kummerian ring may be constructed as follows.

Definition 1.4 ((G, χ)-Kummer extensions). For a fixed n-Kummerian base ring R, a (G, χ)-Kummer extension of R is a triple ($R\{u^{1/n}\}, G, \chi$), where

$$R\{u^{1/n}\} \coloneqq R[T]/(T^n - u),$$

with $u \in R^*$, and G is an n-cyclic group which acts on S via the character $\chi: G \to \mu_n \subseteq R^*$ by

$$g(T) \coloneqq \chi(g)T. \tag{1.5}$$

That this in fact is a Galois extension is shown in [8, p.20].

Definition 1.6 (Equivariant Isomorphism). Let R be a commutative ring and G a fixed finite group. For i = 1, 2, let S_i be a ring extension of R with a faithful action of G by R-automorphisms of S_i . We say that the pairs (S_1, G) and (S_2, G) are equivariantly isomorphic, or simply G-isomorphic, via φ , if φ is an R-algebra isomorphism $\varphi : S_1 \xrightarrow{\sim} S_2$ such that $\varphi \circ g = g \circ \varphi$ for all $g \in G$.

It is clear that a G-isomorphism preserves the G-Galois property of a ring extension. Harrison showed how to classify the set of Galois extensions of a given ring R and group G with a fixed action, via the following object.

Definition 1.7 (The Harrison group [4, p. 67]). Given a group G, the set of all G-isomorphism classes of G-Galois ring extensions S over a fixed base ring R with a fixed faithful action of G is called the Harrison set of (R, G) and denoted by $\mathbb{H}(R, G)$. When G is a finite abelian group, $\mathbb{H}(R, G)$ can be endowed with a group structure. In this case it is called the Harrison group.

We review some of the basic facts regarding group actions on modules over a ring R. Let R be an n-Kummerian ring with distinguished subgroup μ_n , and San R-module. Suppose G is a finite abelian group of order n acting on S via Rmodule automorphisms. Its dual group \hat{G} will be identified with $\text{Hom}(G, \mu_n)$ and its elements referred to simply as characters of G. We may consider the decomposition of S with respect to the action of \hat{G} , namely, for $\chi \in \hat{G}$, we define the χ -eigenspace (or isotypical component)

$$S^{\chi} \coloneqq \{ \alpha \in S : g(\alpha) = \chi(g) \alpha \ \forall g \in G \}.$$

Projection onto the χ -eigenspace is given by $\alpha_{\chi} = e_{\chi}\alpha$, where e_{χ} is the corresponding idempotent in the group algebra,

$$e_{\chi} \coloneqq \frac{1}{|G|} \sum_{g \in G} \chi(g^{-1})g \in R[G].$$

We then have the decomposition

$$S = \bigoplus_{\chi \in \widehat{G}} S^{\chi}.$$
 (1.8)

If G is cyclic of prime order and the action is nontrivial, then each R-module S^{χ} is nontrivial.

With regard to the decomposition (1.8), the product of two G-Galois extensions S_i/R for i = 1, 2 is given by

$$S_1 \cdot S_2 \coloneqq \bigoplus_{\chi} (S_1^{\chi} \otimes S_2^{\chi}), \tag{1.9}$$

where G acts on the summand $S_1^{\chi} \otimes S_2^{\chi}$ via $g(s_1 \otimes s_2) \coloneqq \chi(g)(s_1 \otimes s_2)$. One checks that this product factors through equivariant equivalence and thus defines the group law on the Harrison group $\mathbb{H}(R, G)$.

The neutral element with respect to this product is the so-called trivial G-Galois extension, defined by $R^{(G)} := \bigoplus_{\chi} R$, considered as the set of maps from G to R under the standard sum and product, and with the action of G given by $g((r_{\chi})_{\chi}) = (r_{g^*\chi})_{\chi}$ where g^* denotes composition with multiplication by g.

For simplicity, as in the following result, we will restrict the rank n to be a prime number p, different from the characteristic of k.

Definition 1.10 (*G*-primitive element). Let *R* be a *p*-Kummerian ring and *S* an *R*-algebra on which a *p*-cyclic group *G* acts via *R*-automorphisms. Given a nontrivial character $\chi : G \to \mu_p \subseteq R^*$, an element $\alpha \in S$ is called (G, χ) -primitive if:

- $1, \alpha, \ldots, \alpha^{p-1}$ is an *R*-module basis of *S*.
- $g(\alpha) = \chi(g)\alpha$ for $g \in G$, i.e. $\alpha \in S^{\chi}$.

In this case we also say that α is G-primitive with character χ .

The subject of primitive elements in ring extensions is in itself a topic which has been studied by various authors ([15, 17]). In this regard, we cite the following result, which is [16, Proposition 3.21].

Proposition 1.11. Let S be a separable algebra over a p-Kummerian ring R on which the p-cyclic group G acts via R-automorphisms with $S^G = R$. Fix a nontrivial character $\chi: G \to \mu_p \subseteq R^*$. For an element $\alpha \in S^{\chi}$, the following are equivalent:

- **1.** α is (G, χ) -primitive, i.e. $1, \alpha, \alpha^2, \ldots, \alpha^{p-1}$ is an *R*-module basis of *S*.
- **2.** $\alpha^{p} \in R^{*}$.
- **3.** α is invertible in S.

If this is the case, then:

- **4.** α generates S^{χ} as an *R*-module, and *S* is a free *R*-module of rank *p*.
- 5. The characteristic polynomial of α is $C_{\alpha}(T) = T^p \alpha^p \in R[T]$. It is separable and generates $\operatorname{Ann}(\alpha)$.
- **6.** S is equivariantly isomorphic to the (G, χ) -Kummer extension (Definition 1.4) $(R\{u^{1/p}\}, G, \chi)$ for $u = \alpha^p$.

The basic result in the Kummer theory of ring extensions is the following, which is [8, Theorem 5.4].

Proposition 1.12 (The general Kummer sequence). For an n-Kummerian ring, R, fixing a nontrivial character $\chi : C_n \to \mu_n \subseteq R^*$ yields an exact sequence of groups:

$$1 \to \overset{R^*}{\nearrow}_{R^{*n}} \xrightarrow{i_{(R,\chi)}} \mathbb{H}(R,\mathcal{C}_n) \to \operatorname{Pic}(R)[n] \to 1.$$
(1.13)

Here $i = i_{(R,\chi)}$ sends $u \in R^*$ to the (\mathcal{C}_n, χ) -Kummer extension $R\{u^{1/n}\}$, and the second map is projection onto the χ -eigenspace. This sequence is called the Kummer sequence.

It follows from the Kummer sequence that if R is n-Kummerian and $\operatorname{Pic}(R)$ has trivial n-torsion, then

$$\mathbb{H}(R,\mathcal{C}_n) \simeq \overset{R^*}{\swarrow}_{R^{*n}} \tag{1.14}$$

and thus every isomorphism class of C_n -Galois extensions of R contains a representative which is (C_n, χ) -Kummer. In particular this holds for the function field $R = \Sigma$ and, more interestingly, also for the geometric adele ring $R = \mathbb{A}_X$ ([16, Theorem 3.11]).

Remark 1.15. As in [16, Lemma 3.41], one sees that $i_{(R,\chi)}$ is equivariant with respect to the action of $\operatorname{Aut}(\mathcal{C}_n) \simeq (\mathbb{Z}/(n))^*$ by

$$b \in (\mathbb{Z}/(n))^* \longmapsto (u \mapsto u^b) \in \operatorname{Aut}(R^*/R^{*n})$$
(1.16)

 $b \in (\mathbb{Z}/(n)) \longrightarrow (u \mapsto u) \in \operatorname{Aut}(R/R)$ (1.10) and the group isomorphism $\chi^* : \operatorname{Aut}(\mu_n) \simeq (\mathbb{Z}/(n))^* \xrightarrow{\sim} \operatorname{Aut}(\mathcal{C}_n)$ induced by $\chi : \mathcal{C}_n \xrightarrow{\sim} \mu_n \subseteq R^*.$

Example 1.17. Let p be a prime different from char k. For a choice of nontrivial character $\chi : \mathcal{C}_p \to \mu_p(\mathbb{k}^*)$, the Kummer sequence yields an isomorphism $\Sigma^*/\Sigma^{*p} \xrightarrow{\sim} \mathbb{H}(\Sigma, \mathcal{C}_p)$ given by sending $f \in \Sigma^*$ to the Galois ring extension

$$\Omega_f \coloneqq \Sigma[T] / (T^p - f) \tag{1.18}$$

with action via $g(T) = \chi(g)T$, which is a (\mathcal{C}_p, χ) -Kummer extension. Thus, without loss of generality, we may work with pairs of the form $(\Omega_f, \mathcal{C}_p)$ when considering general representatives of elements in $\mathbb{H}(\Sigma, \mathcal{C}_p)$.

By Capelli's Theorem, $f \notin \Sigma^{*p}$ if and only if Ω_f is a p-Kummer field extension of Σ . In this case C_p is identified with the classical Galois group $\operatorname{Gal}(\Omega_f/\Sigma)$. Recall that by the Kummer theory of fields, every p-cyclic field extension of Σ is of this form (see for example Birch's paper in [3, Ch. III, §2]).

If $f \in \Sigma^{*p}$, then Ω_f is the trivial extension $\Omega_f \simeq \prod^p \Sigma$, and C_p is identified with a subgroup of the symmetric group S_p generated by a p-cycle, permuting the copies of Σ .

We shall also need the following construction of cyclic Kummer extensions due to Borevich.

Theorem 1.19 ([1, §8, Theorem 2]). For an n-Kummerian ring R and fixed nontrivial character $\chi : C_n \to \mu_n \subseteq R^*$, there is a one-to-one correspondence between n-cyclic Kummer extensions S of R and pairs (\mathfrak{L}, φ) , where $\mathfrak{L} \in \operatorname{Pic}(R)[n]$ and φ is an isomorphism $\varphi : \mathfrak{L}^{\otimes n} \xrightarrow{\sim} R$.

The construction sends an *n*-cyclic extension S/R to the pair consisting of the eigenspace S^{χ} together with the map $\varphi : (S^{\chi})^{\otimes n} \to S^{\chi^n} = S^{\mathcal{C}_p} = R$, which is easily seen to be an isomorphism. The inverse sends a pair (\mathfrak{L}, φ) to the quotient of $\bigoplus_{i \geq 0} \mathfrak{L}^{\otimes i}$ by the ideal generated by $\varphi(\ell^{\otimes n}) - \ell^{\otimes n}$, where ℓ runs over the elements of \mathfrak{L} .

In fact this correspondence induces a group isomorphism of $\mathbb{H}(R, \mathcal{C}_p)$ to the set of pairs (\mathfrak{L}, φ) with group law given by $(\mathfrak{L}_1, \varphi_1) \cdot (\mathfrak{L}_2, \varphi_2) = (\mathfrak{L}_1 \otimes \mathfrak{L}_2, \varphi_1 \otimes \varphi_2)$ ([1, §11, Theorem 1]).

Definition 1.20 (Conjugacy of G-ring extensions). Let R be a commutative ring. For i = 1, 2, let S_i be a ring extension of R with a faithful action of a group G_i by R-automorphisms of S_i . We say that (S_1, G_1) and (S_2, G_2) are conjugate via (φ, τ) if $\varphi : S_1 \xrightarrow{\sim} S_2$ is an R-algebra isomorphism and $\tau : G_1 \xrightarrow{\sim} G_2$ is a group isomorphism, such that $\varphi \circ g = \tau(g) \circ \varphi$ for all $g \in G_1$. We will denote this relation by $(S_1, G_1) \sim (S_2, G_2)$.

Note that the group isomorphism τ is in fact determined by the *R*-algebra isomorphism φ , namely $\tau(g) = g^{\varphi} := \varphi \circ g \circ \varphi^{-1}$, although it is convenient to denote it as part of a pair (φ, τ) as we are doing here.

Remark 1.21. If one looks at the definition, it is immediately clear that being a Galois extension S/R is preserved by conjugation.

Definition 1.22. Suppose (S,G) is a ring extension of R with G a group acting faithfully by R-automorphisms os S. Given an automorphism $\tau \in \operatorname{Aut}(G)$ of G, we define the twist of (S,G) by τ , denoted by $(S,G)^{\tau}$, to be the same ring extension S/R but with the action of G now given by $g(s) \coloneqq \tau(g)s$.

We will make use of the bifunctoriality of the Harrison group, as expressed in the following result taken from [8, Proposition 3.10, Corollary 3.11].

Proposition 1.23. Given a homomorphism of commutative rings $R_1 \to R_2$ and a homomorphism $\tau : G_1 \to G_2$ of finite abelian groups, we have a commutative diagram

$$\begin{aligned}
& \mathbb{H}(R_1, G_1) \xrightarrow{\tau^*} \mathbb{H}(R_1, G_2) \\
& \mathbb{H}(R_2, G_1) \xrightarrow{\tau^*} \mathbb{H}(R_2, G_2)
\end{aligned}$$
(1.24)

of group homomorphisms of Harrison groups.

In particular, conjugacy classes of *n*-cyclic Galois ring extensions correspond bijectively to the quotient set of C_n -isomorphism classes of C_n -Galois extensions modulo Aut (C_n) , i.e.

$$\begin{cases} \text{Conjugacy classes of } n\text{-cyclic} \\ \text{Galois extensions } (S,G) \text{ of } R \end{cases} \xrightarrow{1:1} \mathbb{H}(R,\mathcal{C}_n) / \operatorname{Aut}(\mathcal{C}_n).$$
(1.25)

1.C. Covers as Galois ring extensions. Since our main goal is to characterize Galois covers of algebraic curves in terms of Galois ring extensions of the geometric adele ring \mathbb{A}_X , we need to begin by giving a precise notion of Galois cover of a curve. Note that we include covers which are not necessarily irreducible.

Definition 1.26 (Galois cover). A cover of projective, non-singular algebraic curves, $\pi: Y \to X$, where X is irreducible and π is separable, is said to be a Galois cover if the quotient of Y by the action of Gal(Y/X) is X.

Define two covers $Y' \to X$, $Y \to X$ to be equivalent when there is a birational map between them as schemes over X (i.e., there are dense open subschemes U', U of Y', Y respectively and an isomorphism of U' and U over X).

Given X as in §1.A, let $Cov(X, C_p)$ denote the set of isomorphism classes of pcyclic Galois covers $Y \to X$, where we restrict to Y having connected components which coincide with its irreducible components.

Note that with this definition, ramification is allowed (in accordance, for example, with the definition of finite Galois branched cover in [21, p.125]). Then each equivalence class has a distinguished representative that satisfies the additional property that its irreducible and connected components coincide.

An *irreducible* cover $Y \to X$, determines a finite separable extension Ω/Σ of function fields. Conversely, given such an Ω , the Zariski-Riemann variety Y of Ω , whose set of closed points is the set of discrete valuations on Ω , determines a cover of X. This is the classical equivalence between the category of finitely generated field extensions of \Bbbk of transcendence degree 1 and the category of nonsingular projective irreducible curves and nonconstant morphisms (e.g. [19, Theorem 0BY1]).

A general cover, which may not be irreducible, determines a finite separable *ring* extension Ω of Σ , which is isomorphic to a finite product of field extensions. In this case Y is the disjoint union of the corresponding Zariski-Riemann varieties of these fields and Ω is the total quotient ring of any affine dense open subscheme of Y.

Example 1.27. Consider $X = \mathbb{P}^1$ with homogeneous coordinates $[x_0, x_1]$ and the following two covers. First, $Y = \coprod^p X$ with the action of \mathcal{C}_p by cyclic permutation. Second, the cover $Z \subseteq \mathbb{P}^2$ given by the zeros of the homogeneous ideal

generated by $x_1^p - x_2^p$ where $[x_0, x_1, x_2]$ are homogeneous coordinates on \mathbb{P}^2 , with action $[x_0, x_1, x_2] \rightarrow [x_0, x_1, \zeta x_2]$ where ζ is a pth root of unity. Observe that Z is connected and birational to Y. The conditions imposed above are satisfied by Y but not by Z.

The following is an algebraic version of [3, (14.2)], taking into account the nonirreducible case. The reader may check that the proof (discarding the topological aspects) is analogous to the case of a global field considered there.

Theorem 1.28. Let Ω be a finite separable ring extension of Σ and let \mathbb{A}_Y denote the ring of adeles of Ω . Then

$$\mathbb{A}_X \otimes_{\Sigma} \Omega \simeq \mathbb{A}_Y \tag{1.29}$$

as \mathbb{A}_X -algebras. Furthermore, \mathbb{A}_Y^+ is the integral closure of \mathbb{A}_X^+ in $\mathbb{A}_X \otimes_{\Sigma} \Omega$.

A Galois cover $Y \to X$ gives a finite Galois extension of rings Ω/Σ of group $\operatorname{Gal}(Y/X)$. In this case, the isomorphism (1.29) is equivariant with respect to the action of $\operatorname{Gal}(Y/X)$, and

$$\mathbb{A}_{Y}^{\operatorname{Gal}(Y/X)} = \mathbb{A}_{X}.$$
(1.30)

The following result expresses the basic correspondence between the Kummer theory of the function field Σ , the Harrison group of isomorphism classes of C_{p} -Galois ring extensions of Σ , and covers of the algebraic curve X as we have defined above.

Theorem 1.31. Let X/\Bbbk be a projective, irreducible, non-singular curve over an algebraically closed field \Bbbk with char(\Bbbk) $\neq p$ and Σ its function field. There are canonical identifications

$$(\Sigma^*/\Sigma^{*p})/(\mathbb{Z}/(p))^* \simeq \mathbb{H}(\Sigma, \mathcal{C}_p)/_{\operatorname{Aut}(\mathcal{C}_p)} \simeq \operatorname{Cov}(X, \mathcal{C}_p),$$
 (1.32)

where $(\mathbb{Z}/(p))^*$ acts on Σ^*/Σ^{*p} as in (1.16).

Proof. Recall that throughout we fix a character $\chi : \mathcal{C}_p \to \mu_p(\mathbb{k}^*)$ that serves to determine the action used in defining the Harrison group.

The first correspondence in (1.32) is a direct consequence of (1.14) and equivariance with respect to the action of $\operatorname{Aut}(\mathcal{C}_p)$ (see Remark 1.15). Thus we focus on the second correspondence between extensions and covers.

Given an element of $\mathbb{H}(\Sigma, \mathcal{C}_p)$, i.e. a class of \mathcal{C}_p -ring extensions of Σ , we may choose a representative which is a (\mathcal{C}_p, χ) -Kummer extension, namely, of the form $\Omega_f = \Sigma[T]/(T^p - f)$ as in (1.18), with action via $g(T) = \chi(g)T$. As we discussed above, Ω_f is a finite separable ring, isomorphic to a finite product of field extensions of Σ . Let Y be the disjoint union of the Zariski-Riemann varieties of these extensions, endowed with the \mathcal{C}_p -action induced by a given isomorphism. This determines a cover $\pi: Y \to X$ as in Definition 1.26.

Note that f^b , where $b \in (\mathbb{Z}/(p))^*$, gives an equivalent representative in $\mathbb{H}(\Sigma, \mathcal{C}_p)/\operatorname{Aut}(\mathcal{C}_p)$ and does not change Y, i.e. it yields the same cover.

Conversely, given a *p*-cyclic Galois cover $\pi : Y \to X$, the localization of the morphism of \mathcal{O}_X -algebras $\mathcal{O}_X \to \pi_* \mathcal{O}_Y$ at the generic point of X yields a *p*-cyclic Galois ring extension $\Sigma \to \Omega$ and thus a class in $\mathbb{H}(\Sigma, \mathcal{C}_p)/\operatorname{Aut}(\mathcal{C}_p)$.

Remark 1.33. A fundamental reason for specifying the conditions on Galois covers as in Definition 1.26 is that, under the correspondence (1.32), the neutral element of $\mathbb{H}(\Sigma, C_p)$ indeed maps to the trivial cover in that sense (see also Example 1.27).

Given a cover $\pi : Y \to X$, the ramification locus $\operatorname{Ram}(\pi)$ is the image of the support of the sheaf of relative differentials $\omega_{Y/X}$. Equivalently, it consists of the

points $x \in X$ such that $\#\pi^{-1}(x) < \deg(\pi)$. For a finite subset of closed points $\mathfrak{R} \subseteq X$, it is natural to consider

$$\operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p) \coloneqq \{ \pi \in \operatorname{Cov}(X, \mathcal{C}_p) : \operatorname{Ram}(\pi) \subseteq \mathfrak{R} \},$$
(1.34)

so that

$$\operatorname{Cov}(X, \mathcal{C}_p) = \bigcup_{\mathfrak{R}} \operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p).$$
(1.35)

We will study the relation between (1.35) and (1.32) in §3.B.

Remark 1.36. As in the classical topological theory of Riemann surfaces, in the geometric case an element of $\text{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p)$ corresponds to a member $\pi \in \text{Cov}(X \setminus \mathfrak{R}, \mathcal{C}_p)$ which is unramified, i.e. $\text{Ram}(\pi) = \emptyset$ (see [21, Theorem 4.6.4]).

2. Relating the Kummer theories of \mathbb{A}_X and Σ

In [16] we extensively studied the structure of the Harrison group $\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)$, classifying the *p*-cyclic Galois extensions of the ring of adeles of the curve X. Now, by means of the Harrison group of its function field Σ , namely $\mathbb{H}(\Sigma, \mathcal{C}_p)$, we can now relate this to the classification of *p*-cyclic covers of X.

By Proposition 1.23, tensoring $\mathbb{A}_X \otimes_{\Sigma}$ – yields a canonical map

$$\mathbb{H}(\Sigma, \mathcal{C}_p) \to \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \tag{2.1}$$

which is equivariant under the action of the automorphism group $\operatorname{Aut}(\mathcal{C}_p)$. Our main goal in this section is to study (2.1), since it encapsulates the relation between the classical Galois theory of field extensions of Σ with that of its adele ring \mathbb{A}_X .

2.A. The fundamental exact sequence. The valuation vector of an idele $u \in \mathbb{I}_X$ is defined as

$$\upsilon(u) \coloneqq (\upsilon_x(u_x) \bmod p)_x \in \bigoplus_{x \in X} \mathbb{Z}/(p).$$
(2.2)

Clearly the map υ is a group homomorphism and, since \Bbbk is assumed to be algebraically closed, we have

$$\mathbb{I}_X^p = \ker \upsilon : \mathbb{I}_X \to \bigoplus_x \mathbb{Z}/(p)$$
(2.3)

(see [16, (3.29)]). We summarize several important facts in the following.

Proposition 2.4. There is commutative diagram of groups

where the bottom arrow is the map (2.1) which centers our attention, and the vertical arrows are the first terms in the Kummer sequences (1.13) of Σ and \mathbb{A}_X , respectively. In addition, (2.5) is equivariant with respect to the action of $\operatorname{Aut}(\mathcal{C}_p) \simeq (\mathbb{Z}/(p))^*$ as in (1.16) for $R = \Sigma$ or $R = \mathbb{A}_X$, and the action on valuation vectors given by multiplication, i.e.

$$b \in (\mathbb{Z}/(p))^* \longmapsto ((v_x)_x) \mapsto ((bv_x)_x)$$
(2.6)

Definition 2.7. Given an extension $(\mathbb{B}, \mathcal{C}_p)$ of \mathbb{A}_X , the valuation vector $v(\mathbb{B}, G, \chi)$ of the triple (\mathbb{B}, G, χ) is the image of the \mathcal{C}_p -isomorphism class of $(\mathbb{B}, \mathcal{C}_p)$ under $v \circ i_{(\mathbb{A}_X, \chi)}^{-1}$ as in (2.5).

The valuation vector of an extension is an invariant which is explicitly given by

$$\upsilon(\mathbb{B}, G, \chi) = \upsilon(\alpha^p) = (\upsilon_x(\alpha^p) \bmod p)_x \in \bigoplus_{x \in X} \mathbb{Z}/(p),$$
(2.8)

where α is a (\mathcal{C}_p, χ) -primitive element (Definition 1.10 and Proposition 1.11). The existence of the latter is guaranteed by [16, Theorem 3.22], where it is also shown that the definition does not depend on the choice of α .

Note that for a (\mathcal{C}_p, χ) -extension $\mathbb{A}_X \{ t^{1/p} \} = \mathbb{A}_X[T]/(T^p - t)$ where $t \in \mathbb{I}_X$, the class of T is a (\mathcal{C}_p, χ) -primitive element, and thus

$$\upsilon(\mathbb{A}_X\{\mathfrak{t}^{1/p}\}, \mathcal{C}_p, \chi) = \upsilon(\mathfrak{t}).$$
(2.9)

In order to proceed with our characterization of $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$, we need an auxiliary result, which is interesting in itself, and points out the role played by the geometry of the curve.

Theorem 2.10 (The fundamental exact sequence for $\mathbb{H}(\Sigma, C_p)$ and $\mathbb{H}(\mathbb{A}_X, C_p)$). Let X/\mathbb{k} be a curve satisfying our initial hypotheses, namely, a projective, irreducible, non-singular curve over an algebraically closed field \mathbb{k} with char(\mathbb{k}) $\neq p$. Fix a nontrivial character $\chi : C_p \to \mu_p \subseteq \mathbb{k}^*$ and consider the Harrison groups $\mathbb{H}(\Sigma, C_p)$ and $\mathbb{H}(\mathbb{A}_X, C_p)$ of C_p -isomorphism classes of p-cyclic Galois ring extensions of the function field Σ and the adele ring \mathbb{A}_X , respectively, where in both cases the C_p -action is via χ . Then there is a canonical exact sequence of groups

$$0 \to \mathbb{I}_X^p \cap \Sigma^*_{\Sigma^{*p}} \to \mathbb{H}(\Sigma, \mathcal{C}_p) \to \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \to \mathbb{Z}_{(p)} \to 0.$$
(2.11)

Proof. Our assumptions imply that the Picard scheme $\operatorname{Pic}^{0}(X)$ exists and is a projective k-scheme isomorphic to the Jacobian variety $\operatorname{Jac}(X)$.

Consider the commutative diagram (of groups)

where $\operatorname{Div}(X) = \bigoplus_{x \in X} \mathbb{Z}$ denotes the group of divisors on X, the map d sends a function $f \in \Sigma^*$ to its divisor, i.e. $\sum_x v_x(f)x$, and the vertical maps are raising to the *p*th power in the first arrow and multiplication by p in the other two. The Snake Lemma yields the exact sequence

$$0 \to \operatorname{Pic}(X)[p] \to \overset{\Sigma^*}{\longrightarrow} \overset{\bar{d}}{\longrightarrow} \overset{\operatorname{Div}(X)}{\longrightarrow} p(\operatorname{Div}(X)) \to \overset{\operatorname{Pic}(X)}{\longrightarrow} p(\operatorname{Pic}(X)) \to 0.$$

$$(2.12)$$

where $\operatorname{Pic}(X)[p]$ denotes the *p*-torsion subgroup of the Picard variety of X, which is easily seen to coincide with $\operatorname{Jac}(X)[p]$. The map \overline{d} sends a function $f \in \Sigma^*$ to its divisor modulo *p*, i.e, $\sum_x v_x(f)x \mod p$. By (2.3), we have

$$\operatorname{Div}(X)_{p(\operatorname{Div}(X))} \simeq \mathbb{I}_{X/\mathbb{I}_{X}^{p}} \simeq \bigoplus_{x \in X} \mathbb{Z}_{p},$$
 (2.13)

and therefore

$$\operatorname{Jac}(X)[p] = \operatorname{ker}(\bar{d}) = \left\{ \bar{f} \in {}^{\sum^{*}} / {}_{\sum^{*p}} : v_{x}(f) = 0 \mod p \quad \forall x \in X \right\}$$
$$= {}^{\mathbb{I}_{X}^{p}} \cap {}^{\sum^{*}} / {}_{\sum^{*p}}.$$
$$(2.14)$$

Finally, consider the commutative diagram corresponding to the degree map and multiplication by p,



Now, the group $\operatorname{Jac}(X)(\Bbbk)$ of points of $\operatorname{Jac}(X)$ with values in \Bbbk is divisible [19, Proposition 03RP], hence the first arrow is surjective and hence we obtain an isomorphism $\operatorname{Pic}(X)/p(\operatorname{Pic}(X)) \simeq \mathbb{Z}/(p)$. Thus using (2.5) we obtain (2.11).

Corollary 2.15. Given a positive integer n and a divisor $D \in \text{Div}(X)$ with degree equal to a multiple of n, there always exists a function $f \in \Sigma^*$ such that $D \equiv (f) \mod n$.

Proof. As we have seen in the proof of the theorem, $\operatorname{Jac}(X)(\Bbbk)$ is a divisible group. Thus, given a divisor $D \in \operatorname{Div}(X)$ of degree $d \in n\mathbb{Z}$ and any point x_0 , the composition

$$\operatorname{Jac}(X)(\Bbbk) \xrightarrow{[n]} \operatorname{Jac}(X)(\Bbbk) \xrightarrow{+D-dx_0} \operatorname{Jac}(X)(\Bbbk)$$

is surjective. Hence there is some $D' \in \text{Div}(X)^0$ such that $D - dx_0 + nD' \sim 0$, where \sim denotes linear equivalence; that is, $D - dx_0 + nD' = (f)$ for some function f as desired.

2.B. The geometric adelic equivalence problem. In this section we deal with the problem of characterizing when two *p*-cyclic Galois extensions of Σ have the same image under the map (2.1).

This problem is analogous to the so-called equivalence problem for global fields. The latter has a long history, dating back to the 19th century with Kronecker and Hurwitz. It deals in general with the question of when an invariant associated to a global field might classify it up to isomorphism. For example, one may consider its Dedekind or Weil zeta function, the Galois group, or its adele ring, leading to various related notions of equivalence and counterexamples where non-isomorphic fields turn out to have the same invariant. The first example, concerning the Dedekind zeta function, dates back to Gassmann. Komatsu and Perlis considered the Galois group and the adele ring. The problem constitutes an active area of research to this day, see for example [20] for a survey of these classical results and the current state of the art.

Definition 2.16. Denote by $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$ the subgroup of $\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)$ which is the image of $\mathbb{H}(\Sigma, \mathcal{C}_p)$ under (2.1).

We may describe $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$ as follows. Given a rational function $f \in \Sigma^*$, consider

$$\mathbb{A}_X\{\mathbb{f}^{1/p}\} = \mathbb{A}_X[T]/(T^p - \mathbb{f}), \qquad (2.17)$$

where $f = (f_x)_x$ is the idele such that $f_x \in K_x$ is the germ of f at x and C_p acts on the class of T via χ . Note that the class of T is a (C_p, χ) -primitive element. By Theorem 1.28, $\mathbb{A}_X\{f^{1/p}\} \simeq \mathbb{A}_Y$. This motivates the following definition.

Definition 2.18 (rational (C_p, χ) -Kummer extension of \mathbb{A}_X). A *p*-cyclic extension of \mathbb{A}_X which is C_p -isomorphic to one of the form (2.17) will be called a rational (C_p, χ) -Kummer extension of \mathbb{A}_X .

Thus $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$ consists of the \mathcal{C}_p -isomorphism classes of rational (\mathcal{C}_p, χ) -extensions of \mathbb{A}_X . This subgroup can be characterized as a kernel and a cokernel as follows.

Corollary 2.19. The exact sequence (2.11) splits into the following two short exact sequences:

$$0 \to \mathbb{H}_{\mathrm{ratl}}(\mathbb{A}_X, \mathcal{C}_p) \to \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \simeq \bigoplus_{x \in X} \mathbb{Z}_{(p)} \to \mathbb{Z}_{(p)} \to 0$$
(2.20)

$$0 \to \mathbb{I}_X^p \cap \Sigma^*_{\Sigma^{*p}} \to \mathbb{H}(\Sigma, \mathcal{C}_p) \simeq \Sigma^*_{\Sigma^{*p}} \to \mathbb{H}_{\mathrm{ratl}}(\mathbb{A}_X, \mathcal{C}_p) \to 0.$$
(2.21)

Remark 2.22. As we mentioned above, the maps between Harrison groups are equivariant under the action of $Aut(C_p)$ and hence induce actions on the cokernel in (2.20) and on the kernel in (2.21), which is isomorphic to Jac(X)[p] by (2.14).

The sequence (2.20) determines when a given adelic extension comes from a corresponding extension of the function field. This may be regarded as an adelic analog of the Grunwald-Wang problem, in the form given in [11].

Corollary 2.23. In terms of the valuation vector associated to a C_p -extension (\mathbb{B}, C_p) of \mathbb{A}_X (Definition 2.7), the previous result translates to the following set of equivalences:

- **1.** The class of $(\mathbb{B}, \mathcal{C}_p)$ belongs to the subgroup $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$.
- **2.** The valuation vector $v(\mathbb{B}, \mathcal{C}_p, \chi) = (v_x) \in \bigoplus_{x \in X} \mathbb{Z}/(p)$ satisfies

$$\sum_{x} v_x \equiv 0 \mod p. \tag{2.24}$$

3. $v(\mathbb{B}, \mathcal{C}_p, \chi) = v(f)$ for some function $f \in \Sigma^*$. In this case, $(\mathbb{B}, \mathcal{C}_p)$ is \mathcal{C}_p isomorphic to $\mathbb{A}_X\{\mathbb{f}^{1/p}\} \simeq \mathbb{A}_X \otimes_{\Sigma} \Omega_f$ as in (1.18) and (2.17).

From (2.21) we see that it is possible to have non-isomorphic field extensions of Σ whose adele rings are (topologically)¹ isomorphic as \mathbb{A}_X -algebras. An example is given below. This can be considered the analog for an algebraically closed base field of the known equivalence results for global function fields, which were originally studied by Tate [22] and Turner [23].

Corollary 2.25 (Geometric equivalence problem). There is an exact sequence

$$0 \to \operatorname{Jac}(X)[p] \to \mathbb{H}(\Sigma, \mathcal{C}_p) \to \bigoplus_{x \in X}^{0} \mathbb{Z}_{(p)} \to 0, \qquad (2.26)$$

where the notation indicates zero-sum modulo p tuples.

In particular, when g > 0, this shows that the geometric adele ring is not enough in general to classify Σ -extensions.

Proof. This follows from the exact sequence (2.21), recalling (2.14), and Corollary 2.23.

We illustrate the negative answer to the geometric equivalence problem with the following explicit example.

Example 2.27. Let \Bbbk be an algebraically closed field of characteristic different from 2,3 and set p = 2. Let X be the normalization of the completion of the affine plane curve of equation

$$v^{2} = (u-1)(u+1)u \tag{2.28}$$

as a closed subscheme of the affine plane Spec $\mathbb{k}[u, v]$. A routine computation shows that X is an irreducible, non singular, elliptic curve of equation $x_0x_2^2 = (x_1 - x_0)(x_1 + x_0)x_1$ in homogeneous coordinates of the projective plane $\operatorname{Proj} \mathbb{k}[x_0, x_1, x_2]$. Let $0 \in X$ denote the point $X \cap (x_0)_0$ with coordinates (0, 1, 0). Using the Abel map

¹We do not discuss topologies on \mathbb{A}_X -algebras here; see [16] for details.

with base point \mathfrak{O} , the law group of $\operatorname{Jac}(X)$ given by addition of divisors (or tensor product of invertible sheaves) can be transported to X and, in this way, \mathfrak{O} turns out to be the neutral element of the composition law on X. The reader may recognize in this construction the addition law in plane cubic defined geometrically by union of points and intersection of lines. Since X has genus 1, $\operatorname{Jac}(X)[2]$ consists of $p^{2g} = 4$ points which corresponds via \mathcal{A} to \mathfrak{O} and the three affine points of X whose tangent is parallel to the line $x_1 = 0$. That is

$$Jac(X)[2] = \{\mathcal{O}_X, \mathcal{O}_X((1, -1, 0) - (0, 1, 0)), \\ \mathcal{O}_X((1, 0, 0) - (0, 1, 0)), \mathcal{O}_X((1, 1, 0) - (0, 1, 0))\}.$$

Now, observe that Σ , the function field of X, is the given by the localization $(\mathbb{k}[u, v]/(v^2 - (u-1)(u+1)u))_{(0)}$ so that any function in Σ can be obtained as a rational function on u, v. Let us consider the rational function $f := \frac{u-1}{u} \in \Sigma^*$. The divisor of f is given by

$$(f) = \left(\frac{u-1}{u}\right) = \left(\frac{x_1 - x_0}{x_1}\right) = (x_1 - x_0)_0 - (x_1)_0 = 2(1, 1, 0) - 2(1, 0, 0)$$

so that $v_x(f) = 0 \mod 2$ for all $x \in X$. Note that $f \notin \Sigma^{*2}$, since if $f = h^2$ for some $h \in \Sigma^*$ then h would be a rational function with exactly one simple pole and one simple zero, which would imply that the curve X has genus 0. Proceeding similarly with the function $g := \frac{u+1}{u-1}$, one concludes that

$$\mathbb{I}_X^2 \cap \Sigma^* / _{\Sigma^{*2}} = \{1, f, g, fg\}.$$

Hence, Theorem 2.10 implies that we have the following four non-isomorphic 2-cyclic Galois extensions of Σ :

$$\Sigma[T]_{(T^2-1)}, \quad \Sigma[T]_{(T^2-f)}, \quad \Sigma[T]_{(T^2-g)}, \quad \Sigma[T]_{(T^2-fg)},$$

The latter three are nontrivial and, when tensored by \mathbb{A}_X , all yield trivial 2-cyclic Galois ring extensions of \mathbb{A}_X . Thus we have an example of four distinct elements of $\mathbb{H}(\Sigma, C_2)$ which map to the identity element in $\mathbb{H}(\mathbb{A}_X, C_2)$, in particular, they become isomorphic after tensoring with \mathbb{A}_X .

3. The role of ramification

3.A. The fundamental cube. As we outlined in the introduction, we will now consider a finite nonempty subset \mathfrak{R} of closed points of X, which will represent the ramification locus. By studying the Harrison group of the ring of the affine curve $X \setminus \mathfrak{R}$, we obtain a commutative cube (3.12) which refines (2.11).

In [16, §2] we introduced a notion of ramification for an adelic algebra $\mathbb{A}_X\{\mathfrak{t}^{1/n}\} = \mathbb{A}_X[T]/(T^n - \mathfrak{t})$ for an idele $\mathfrak{t} \in \mathbb{I}_X$ and *n* coprime to char(\Bbbk), which we now briefly describe. First, the *ramification locus* of \mathfrak{t} is

$$Ram(t) := \{ x \in X : (n, v_x(t_x)) \neq n \} = \{ x \in X : v_x(t_x) \not\equiv 0 \mod n \},$$
(3.1)

and the corresponding *ramification index* at x is

$$e_x \coloneqq \frac{n}{(n, v_x(t_x))}.$$

The vector $\mathbf{\mathfrak{e}} = (e_x)$ of integers is the ramification profile of $\mathbf{\mathfrak{t}}$. Observe that $\operatorname{Ram}(\mathbf{\mathfrak{t}}) = \{x \in X : e_x > 1\}$ is a finite set. By [16, Theorem 2.19], isomorphism classes of \mathbb{A}_X -algebras of the form $\mathbb{A}_X\{\mathbf{\mathfrak{t}}^{1/n}\}$ are classified by their ramification profile, namely, $\mathbb{A}_X\{\mathbf{\mathfrak{t}}_1^{1/n}\}$ and $\mathbb{A}_X\{\mathbf{\mathfrak{t}}_2^{1/n}\}$ are isomorphic if and only if $\mathbf{\mathfrak{e}}_1 = \mathbf{\mathfrak{e}}_2$.

Recalling the notation in §1.A, when n = p is prime it is straightforward to check (see [16, Lemma 2.13]) that $x \notin \text{Ram}(t)$ if and only if the K_x -algebra

 $K_x\{t_x^{1/p}\}$ is isomorphic to p copies of K_x . Since the latter is the neutral element of $\mathbb{H}(K_x, \mathcal{C}_p)$, this is also equivalent to $\mathbb{A}_X\{t^{1/p}\}$ lying in the kernel of the canonical map $\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \to \mathbb{H}(K_x, \mathcal{C}_p)$.

Since every C_p -extension of \mathbb{A}_X is isomorphic to a (C_p, χ) -Kummer extension, we have the following stratification by ramification detailed in [16, §3.E, Corollary 3.75]:

$$\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \xrightarrow{\text{Aut}(\mathcal{C}_p)} \stackrel{\text{1:1}}{\longleftrightarrow} \prod_{\mathfrak{R} \subseteq X} \left\{ \begin{array}{c} \text{Conjugacy classes of } p \text{-cyclic Galois} \\ \text{extensions } (\mathbb{B}, G) \text{ of } \mathbb{A}_X \text{ ramified at } \mathfrak{R} \end{array} \right\}$$
(3.2)

where \Re ranges over finite subsets of closed points of X.

In the following we recall the notation from §1.A. For a finite subset $\mathfrak{R} \subseteq X$ (possibly empty)

$$\mathbb{A}_{X,\mathfrak{R}} := \prod_{x \in \mathfrak{R}} K_x \times \prod_{x \in X \setminus \mathfrak{R}} A_x, \quad \mathbb{I}_{X,\mathfrak{R}} := \mathbb{A}^*_{X,\mathfrak{R}}.$$

In addition, we will also denote

$$A_{\mathfrak{R}} \coloneqq H^0(X \setminus \mathfrak{R}, \mathcal{O}_X). \tag{3.3}$$

Theorem 3.4. Let X/\Bbbk be a projective, irreducible, non-singular curve over an algebraically closed field \Bbbk with char(\Bbbk) $\neq p$, and $\varnothing \subsetneq \mathfrak{R} \subset X$ be a finite nonempty subset (of closed points). The Harrison group $\mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p)$ is characterized as

$$\mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p) \simeq \{ f \in \Sigma^* / (\Sigma^*)^p : v_x(f) \equiv 0 \mod p \ \forall x \notin \mathfrak{R} \}.$$
(3.5)

Proof. Consider the exact sequence of sheaves

$$0 \to \mathcal{O}_X^* \to \Sigma^* \to \Sigma^* / \mathcal{O}_X^* \to 0 \tag{3.6}$$

and restrict it to the open subscheme $X \setminus \mathfrak{R}$. From its long exact sequence of cohomology one obtains

$$0 \to \Sigma^*/A^*_{\mathfrak{R}} \to \operatorname{Div}(X \setminus \mathfrak{R}) \to \operatorname{Pic}(X \setminus \mathfrak{R}) \to 0.$$

Observe that, since we assume \mathfrak{R} to be nonempty, $X \setminus \mathfrak{R}$ is affine, equal to $\operatorname{Spec}(A_{\mathfrak{R}})$. This yields the following analog to (2.12):

$$0 \to \operatorname{Pic}(A_{\mathfrak{R}})[p] \to \overset{\Sigma^*}{/}_{A_{\mathfrak{R}}^*(\Sigma^*)^p} \xrightarrow{\Phi} \overset{\mathbb{I}_X}{\longrightarrow} \overset{\mathbb{I}_X}{/}_{\mathbb{I}_{X,\mathfrak{R}}\mathbb{I}_X^p} = \overset{\operatorname{Div}(A_{\mathfrak{R}})}{/}_{p(\operatorname{Div}(A_{\mathfrak{R}}))} \to 0 \quad (3.7)$$

On the other hand the Kummer sequences (1.13) yield

where Ψ is induced by $\Sigma \otimes_{A_{\Re}}$ – and injectivity of the left vertical arrow is clear. The Snake Lemma yields

$$0 \to \ker \Psi \to \operatorname{Pic} A_{\mathfrak{R}}[p] \xrightarrow{\delta} \Sigma^* / A_{\mathfrak{R}}^*(\Sigma^*)^p \to \operatorname{coker} \Psi \to 0,$$

where δ denotes the connecting morphism. Observe that δ coincides with the map on the l.h.s. of (3.7). From the injectivity of the latter, we conclude that ker $\Psi = 0$ and

$$\operatorname{coker} \delta = \operatorname{coker} \Psi = \operatorname{Im} \Phi = {}^{\mathbb{I}_X} / {}_{\mathbb{I}_X, \mathfrak{R}} {}^{p}_X.$$
(3.9)

Hence, one has the commutative diagram

which after another application of the Snake Lemma, using (3.9) yields

$$0 \to \mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p) \xrightarrow{\Psi} \overset{\Sigma^*}{\frown} (\Sigma^*)^p \to \overset{\mathbb{I}_X}{\longrightarrow} \overset{\mathbb{I}_X}{\longrightarrow} \mathbb{I}_{X, \mathfrak{R}} \mathbb{I}_X^p \to 0, \qquad (3.10)$$

from which (3.5) follows immediately.

Theorem 3.11. Let X/\Bbbk be a curve satisfying our initial hypotheses, namely, a projective, irreducible, non-singular curve over an algebraically closed field \Bbbk with char(\Bbbk) \neq p. For each finite nonempty subset $\Re \subset X$, we have the following commutative cube:



where the notation is as above. Moreover, the front face of this cube is a cartesian square:

$$\mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p) \simeq \mathbb{H}(\Sigma, \mathcal{C}_p) \underset{\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)}{\times} \mathbb{H}(\mathbb{A}_{X, \mathfrak{R}}, \mathcal{C}_p),$$
(3.13)

Furthermore, (3.12) is equivariant with respect to the action of $\operatorname{Aut}(\mathcal{C}_p)$ on the various objects, as described in Proposition 1.23 and (1.16).

Proof. The diagonal maps come from the Kummer sequence (1.13), while the horizontal and vertical maps arise from the functoriality of $\mathbb{H}(-, \mathcal{C}_p)$ in (1.24) and the following (cartesian) diagram:

The three diagonal isomorphisms follow from the triviality of the respective Picard groups of $\mathbb{A}_{X,\mathfrak{R}}$ and \mathbb{A}_X , shown in [16, Theorem 3.11] (and the field case for Σ). It is easy to check the injectivity of the vertical maps in back. That of the front right vertical map now follows from the one in back.

Finally, the injectivity of $\mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p) \to \mathbb{H}(\Sigma, \mathcal{C}_p)$ is part of (3.10), which was the injectivity of the map denoted by Ψ in the proof of Theorem 3.4. Combining (3.10)

with the front and right faces of (3.12) we obtain

from which (3.13) follows.

Corollary 3.15. There is an isomorphism from the direct limit over finite nonempty subsets $\mathfrak{R} \subset X$ of the top floor of (3.12) to the bottom floor.

Proof. The statement is straightforward for the right face of the cube, and for the left face, it means that

$$\mathbb{H}(\Sigma, \mathcal{C}_p) \simeq \varinjlim_{\mathfrak{R}} \mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p), \quad \overset{\Sigma^*}{\searrow}_{(\Sigma^*)^p} \simeq \varinjlim_{\mathfrak{R}} \overset{A^*_{\mathfrak{R}}}{\swarrow}_{(A^*_{\mathfrak{R}})^p}.$$

The first follows from (3.10), and the second is then a consequence of this and (3.8).

3.B. Algebraic vs. geometric ramification. The results of the previous section now let us to take ramification into account in the Harrison groups of C_p -extensions of both Σ and \mathbb{A}_X , allowing us to filter by ramification. We end by establishing the concordance between the algebraic and geometric notions of ramification (Proposition 3.35), which underlies the various geometric applications explored in §4.

Keeping in mind (2.5), Definition 2.7 and (2.9), and that any class of C_p -extensions of \mathbb{A}_X is represented by a (C_p, χ) -Kummer extension (see (1.14)), where the character χ has been fixed beforehand, the following notions are well-defined.

Definition 3.16 (Ramification for C_p -extensions of \mathbb{A}_X). Given a class of C_p -Galois ring extensions $\mathbb{B} \in \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)$, we define the ramification locus of \mathbb{B} and the corresponding valuations and indices by

$$\operatorname{Ram}(\mathbb{B}) \coloneqq \operatorname{Ram}(\mathfrak{t}), \quad v_x(\mathbb{B}) \coloneqq v_x(\mathfrak{t}), \quad e_x(\mathbb{B}) \coloneqq e_x, \tag{3.17}$$

where $t \in \mathbb{I}_X$ is such that \mathbb{B} is represented by the (\mathcal{C}_p, χ) -Kummer extension $\mathbb{A}_X\{t^{1/p}\}$.

Corollary 3.18 (Ramification filtration by subgroups). *Ramification provides us with a natural filtration by subgroups:*

$$\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \simeq \varinjlim_{\mathfrak{R}} \mathbb{H}(\mathbb{A}_{X, \mathfrak{R}}, \mathcal{C}_p) \simeq \varinjlim_{\mathfrak{R}} \left\{ \mathbb{B} \in \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) : \operatorname{Ram}(\mathbb{B}) \subseteq \mathfrak{R} \right\}.$$
(3.19)

For a given finite subset $\mathfrak{R} \subseteq X$,

$$\# \{ \mathbb{B} \in \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) : \operatorname{Ram}(\mathbb{B}) \subseteq \mathfrak{R} \} = p^{\#\mathfrak{R}}.$$
(3.20)

Proof. For a finite subset $\mathfrak{R} \subset X$, we conclude from the isomorphisms on the right face of (3.12), that the following diagram is commutative.

where, by Corollary 3.15, the bottom row is the limit of the top row. Thus, by observing that the following three conditions are equivalent,

1. $\mathbb{B} \in \mathbb{H}(\mathbb{A}_{X,\mathfrak{R}}, \mathcal{C}_p),$ 2. $\operatorname{Ram}(\mathbb{B}) \subseteq \mathfrak{R},$ **3.** $v_x(\mathbb{B}) \equiv 0 \mod p \text{ for } x \notin \mathfrak{R},$

we obtain (3.19) and (3.20).

Corollary 3.22 (Stratification by ramification). *There is a natural stratification of sets, indexed by ramification:*

$$\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) \simeq \bigsqcup_{\mathfrak{R}} \{\mathbb{B} \in \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) : \operatorname{Ram}(\mathbb{B}) = \mathfrak{R}\} \simeq \bigsqcup_{\mathfrak{R}} \bigoplus_{x \in \mathfrak{R}} \left(\mathbb{Z}_{(p)}\right)^*.$$
(3.23)

For a given finite subset $\mathfrak{R} \subseteq X$,

$$\# \{ \mathbb{B} \in \mathbb{H}(\mathbb{A}_X, \mathcal{C}_p) : \operatorname{Ram}(\mathbb{B}) = \mathfrak{R} \} = (p-1)^{\#\mathfrak{R}}.$$
(3.24)

Proof. This follows analogously to the above by observing that the following three conditions are also equivalent:

- 1. $\mathbb{B} \in \mathbb{H}(\mathbb{A}_{X,\mathfrak{R}}, \mathcal{C}_p) \setminus \bigcup_{\mathfrak{R}' \subsetneq \mathfrak{R}} \mathbb{H}(\mathbb{A}_{X,\mathfrak{R}'}, \mathcal{C}_p),$ 2. $\operatorname{Ram}(\mathbb{B}) = \mathfrak{R},$
- **3.** $v_x(\mathbb{B}) \equiv 0 \mod p$ if and only if $x \notin \mathfrak{R}$.

In view of (3.21), for a finite (possibly empty) subset $\mathfrak{R} \subset X$, we may express the conditions $\operatorname{Ram}(\mathbb{B}) \subseteq \mathfrak{R}$ and $\operatorname{Ram}(\mathbb{B}) = \mathfrak{R}$ respectively by

$$(\upsilon_x(\mathbb{B}))_{x\in X} \in \bigoplus_{x\in\mathfrak{R}} \mathbb{Z}/(p), \qquad (\upsilon_x(\mathbb{B}))_{x\in X} \in \bigoplus_{x\in\mathfrak{R}} (\mathbb{Z}/(p))^*,$$

if no confusion arises.

We will also need to define a corresponding notion of ramification for C_p -Galois ring extensions of the function field Σ .

Definition 3.25 (Ramification for C_p -extensions of Σ). Given a class of C_p -Galois ring extensions $\Omega \in \mathbb{H}(\Sigma, C_p)$, we define the ramification locus $\operatorname{Ram}(\Omega)$ of Ω via the map (2.1) as $\operatorname{Ram}(\mathbb{A}_X \otimes_{\Sigma} \Omega)$.

Remark 3.26. For $f \in \Sigma^*$, consider, as in (1.18), the C_p -extension of Σ given by $\Omega_f = \Sigma[T]/(T^p - f)$ with C_p -action by $g(T) = \chi(g)T$ where $\chi : C_p \to \mu_p(\mathbb{k}^*)$ is a fixed character. Then, from the definition and (2.9), we have

$$\operatorname{Ram}(\Omega_f) = \operatorname{Ram}(\mathbb{A}_X \otimes_{\Sigma} \Omega_f) = \operatorname{Ram}(\mathbb{A}_X \{\mathbb{f}^{1/p}\})$$

=
$$\operatorname{Ram}(\mathbb{f}) = \{x \in X : v_x(f_x) \not\equiv 0 \bmod p\},$$
(3.27)

where $f = (f_x)_x \in I_X$ is the idele of germs as in (2.17).

Definition 3.28. The subset of classes of C_p -Galois extensions of Σ with ramification contained in \mathfrak{R} will be denoted by

$$\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) \coloneqq \{\Omega \in \mathbb{H}(\Sigma, \mathcal{C}_p) : \operatorname{Ram}(\Omega) \subseteq \mathfrak{R}\}.$$

With a slight abuse of notation (recalling that the front face of (3.12) is a cartesian square), note that $\operatorname{Ram}(\Omega)$ is the intersection of the finite subsets $\mathfrak{R} \subset X$ such that $\Omega \in \mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p)$.

Proposition 3.29.

$$\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) = \begin{cases} \mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p), & \text{if } \mathfrak{R} \text{ is nonempty,} \\ \operatorname{Jac}(X)[p], & \text{if } \mathfrak{R} = \varnothing. \end{cases}$$
(3.30)

Proof. If \mathfrak{R} is nonempty, this follows from Corollary 3.15 and (3.13), while if $\mathfrak{R} = \emptyset$ then it follows from Corollary 2.25.

Remark 3.31. A posteriori, it follows that $\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p)$ is in fact a subgroup of the Harrison group $\mathbb{H}(\Sigma, \mathcal{C}_p)$, and there is a filtration by ramification

$$\mathbb{H}(\Sigma, \mathcal{C}_p) \simeq \varinjlim_{\mathfrak{R}} \mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) \simeq \varinjlim_{\mathfrak{R}} \left\{ \Omega \in \mathbb{H}(\Sigma, \mathcal{C}_p) : \operatorname{Ram}(\Omega) \subseteq \mathfrak{R} \right\}.$$
(3.32)

analogous to (3.19).

Proposition 3.33. For a given finite subset $\mathfrak{R} \subseteq X$, we have the following refinement of (2.26) filtered by ramification:

$$0 \to \operatorname{Jac}(X)[p] \to \mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) \to \bigoplus_{x \in \mathfrak{R}} 0^{\mathbb{Z}}(p) \to 0.$$
(3.34)

Proof. This follows by combining (2.26), (3.30) and (3.32).

We can now relate the geometric notion of ramification of a cover of an algebraic curve with the corresponding algebraic definition that we have given for extensions of its function field.

Proposition 3.35. If a p-cyclic Σ -ring extension (Ω, C_p) corresponds to the cover $\pi: Y \to X$ as in (1.32), then

$$\operatorname{Ram}(\Omega) = \operatorname{Ram}(\pi). \tag{3.36}$$

In particular, for a finite subset $\mathfrak{R} \subseteq X$,

$$\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) \xrightarrow{}_{\operatorname{Aut}(\mathcal{C}_p)} \xrightarrow{\simeq} \operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p)$$
(3.37)

with notation as in (1.34).

Proof. With notation as in Example 1.17 and the proof of Theorem 1.31, it suffices to check that for $f \in \Sigma^*$, we have $\operatorname{Ram}(\Omega_f) = \operatorname{Ram}(\pi)$. Recall that, by (3.27), $\operatorname{Ram}(\Omega_f) = \{x : v_x(f_x) \neq 0 \mod p\}$. Noting that in general $(\omega_{Y/X})_y \simeq (\omega_{Y/\Bbbk})/(\mathfrak{m}_y^{e-1}\omega_{X/\Bbbk})$ where e is the ramification index at $x = \pi(y)$, there are two cases to consider:

- If $v_x(f_x) \neq 0 \mod p$, then $T^p f_x$ is irreducible in $A_x[T]$, hence the class of T is a local parameter, provided that $v_x(f_x) > 0$. In this case, by uniqueness of extensions of valuations, we have $v_x(f_x) = pv_y(T)$ where y is the unique point in $\pi^{-1}(x)$. Thus $x \in \operatorname{Ram}(\pi)$. If $v_x(f_x) < 0$ then the argument is similar with T^{-1} as local parameter.
- If $v_x(f_x) \equiv 0 \mod p$, and $v_x(f_x) \geq 0$, then $T^p f_x$ splits completely as $\prod_{i=0}^{p-1} (T \zeta^i f_x^{1/p})$ where ζ is a primitive *p*th root of unity and $f_x^{1/p}$ is any choice of *p*th root in A_x . Now $\pi^{-1}(x) = \{y_1, \ldots, y_p\}$ and for each *i* we have $v_x(f_x^{1/p}) = v_{y_i}(T)$. Hence $x \notin \operatorname{Ram}(\pi)$. The case where $v_x(f_x) \leq 0$ is similar.

Finally, (3.37) follows from the previous results and (3.13).

Corollary 3.38. Let p be a prime number. Let X be a smooth, irreducible, nonsingular curve over an algebraically closed field \Bbbk of characteristic different from p. Let $\mathfrak{R} \subset X$ be a finite subset of closed points containing at least two elements. Then there exist:

- **1.** An irreducible p-cyclic Galois cover of X whose ramification locus is exactly \mathfrak{R} .
- **2.** An irreducible polynomial $T^p f \in \Sigma[T]$ which is ramified exactly at the points of \mathfrak{R} and whose Galois group is \mathcal{C}_p .

Proof. This follows from the previous result and Corollary 3.22. We need \Re to have at least two elements so that we can have non-zero valuations which sum to 0.

4. Geometric applications

The previous sections, concerned with C_p -Galois ring extensions of the function field Σ and the corresponding adele ring \mathbb{A}_X , allow us to recover some classical geometric results regarding *p*-cyclic Galois covers of an algebraic curve. Since we employ purely algebraic techniques, this is done without any underlying appeal to the analytical and topological theory of Riemann surfaces, as is the case in the standard approach.

4.A. Classification. We can give a second proof of Proposition 3.35, relating algebraic and geometric ramification, by a direct argument of independent interest, since it shows the concordance of two points of view, namely Borevich's correspondence given in Theorem 1.19 (and sketched following its statement) and that of Cornalba in [5, Lemma 1].

We refine these constructions to take into account a specified ramification locus \mathfrak{R} , via (3.37), with notation as in §3.B.

Proposition 4.1 (Proposition 3.35 via Borevich and Cornalba). For a finite subset $\mathfrak{R} \subseteq X$,

$$\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) / \operatorname{Aut}(\mathcal{C}_p) \xrightarrow{\sim} \operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p).$$

$$(4.2)$$

Proof. Cornalba [5] showed that there is a bijection of sets

$$\operatorname{Cov}(X, \mathcal{C}_p) \stackrel{1:1}{\longleftrightarrow} \left\{ \begin{array}{l} (\mathcal{L}, D) \text{ where } D \in \operatorname{Div}(X) \text{ is an effective} \\ \operatorname{divisor and } \mathcal{L} \in \operatorname{Pic}(X) \text{ is a line bundle} \\ \operatorname{with} \mathcal{L}^p \simeq \mathcal{O}_X(D) \end{array} \right\} \middle/ \sim, \qquad (4.3)$$

where two pairs are defined to be equivalent, denoted by $(\mathcal{L}_1, D_1) \sim (\mathcal{L}_2, D_2)$ with $D_i = \sum_{x \in X} a_x^i x$, if there exists $1 \le b < p$ such that

- $ba_x^1 \equiv a_x^2 \mod p$ for all $x \in X$, $\mathcal{L}_1^b \simeq \mathcal{L}_2(\sum c_x x)$ with $ba_x^1 = c_x p + a_x^2$ for each $x \in X$.

In this correspondence, Y is not required to be irreducible. Indeed, the trivial cover $Y = \prod^{p} X$ corresponds to the equivalence class of the pair $(\mathcal{O}_X, 0)$. In addition, a given cover $\pi \in \text{Cov}(X, \mathcal{C}_p)$ corresponding to a pair (\mathcal{L}, D) has ramification locus $\operatorname{Ram}(\pi)$ equal to $\operatorname{supp}(D)$.

Fix a character $\chi : \mathcal{C}_p \to \mu_p(\mathbb{k}^*)$, that determines the actions used in defining the different Harrison groups involved, as well as a finite subset $\mathfrak{R} \subseteq X$.

Let Ω be a \mathcal{C}_p -ring extension of Σ with $\mathfrak{R} \supset \operatorname{Ram}(\Omega)$, i.e. whose class lies in $\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p)$. By (3.30) and (3.32), \mathfrak{R} may be assumed nonempty and $\Omega \in \mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p)$. The unramified case, where $\Re = \emptyset$, is easily seen to follow from the general result.

Under Borevich's correspondence, there is a pair (\mathcal{L}, φ) associated to Ω , where $\mathcal{L} \in \operatorname{Pic}(A_{\mathfrak{R}})[p]$ and $\varphi : \mathcal{L}^{\otimes p} \xrightarrow{\sim} A_{\mathfrak{R}}$. The exact sequence

$$\bigoplus_{x_i \in \mathfrak{R}} \mathbb{Z}x_i \to \operatorname{Pic}(X) \to \operatorname{Pic}(X \setminus \mathfrak{R}) \to 0$$

shows that there exists $\mathfrak{L} \in \operatorname{Pic}(X)$ such that $\mathfrak{L}|_{X \setminus \mathfrak{R}} \simeq \mathcal{L}$. Then, applying Lemma 30.10.6 [19, Lemma 0FD0] to $\varphi^{-1} : \mathcal{O}_{X \setminus \mathfrak{R}} \xrightarrow{\sim} \mathcal{L}^{\otimes (-p)}$ shows that there exists an effective divisor \overline{D} with support contained in \mathfrak{R} and a morphism of \mathcal{O}_X -modules

$$\bar{\varphi}^{-1}: \mathcal{O}_X(-\bar{D}) \to \mathfrak{L}^{\otimes (-p)}$$

whose restriction to $X \setminus \mathfrak{R}$ coincides with φ^{-1} . Choosing a suitable \overline{D} , it can be assumed that $\bar{\varphi}^{-1}$ is an isomorphism. Write $\bar{D} = pE + D$ for divisors E, D with support contained in \mathfrak{R} and such that $D = \sum_{x_i \in \mathfrak{R}} a_i x_i$ with $0 \le a_i \le p-1$. Then, $\bar{\varphi}$ yields

$$(\mathfrak{L}(-E))^{\otimes p} \simeq \mathcal{O}_X(D).$$

It is straightforward to check that the pair $(\mathfrak{L}(-E), D)$ is the data associated by Cornalba to the cover corresponding to the extension Ω and that its branch locus is $\{x_i \in \mathfrak{R} : a_i \neq 0\} = \operatorname{supp}(D) \subseteq \mathfrak{R}$. Thus we obtain an element of $\operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p)$. If one chooses another character, the resulting pair is equivalent in the sense of (4.3).

Conversely, let (\mathcal{L}, D) be associated to an element of $\operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p)$ as in (4.3); thus $\mathfrak{R} \supseteq \operatorname{supp}(D)$. Fix an isomorphism $\varphi : \mathcal{L}^{\otimes p} \xrightarrow{\sim} \mathcal{O}_X(D)$, which is unique up to \Bbbk^* . Restricting to the open subset $X \setminus \mathfrak{R}$, let $\mathfrak{L} \coloneqq H^0(X \setminus \mathfrak{R}, \mathcal{L}) \in \operatorname{Pic}(A_{\mathfrak{R}})$ and $\varphi_{\mathfrak{R}} \coloneqq \varphi|_{X \setminus \mathfrak{R}} : \mathfrak{L}^{\otimes p} \xrightarrow{\sim} A_{\mathfrak{R}}$. Since $X \setminus \mathfrak{R}$ is affine with ring $A_{\mathfrak{R}}$, we obtain a Borevich pair $(\mathfrak{L}, \varphi_{\mathfrak{R}})$, which then corresponds (via the choice of character χ) to an element of $\mathbb{H}(A_{\mathfrak{R}}, \mathcal{C}_p) = \mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p)$.

Note that a different choice of character yields a conjugate extension, not necessarily an isomorphic one. Thus we obtain an element of $\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p) / \operatorname{Aut}(\mathcal{C}_p)$.

Theorem 1.31 gave a geometric description of the Harrison group $\mathbb{H}(\Sigma, \mathcal{C}_p)$ and now, with the previous discussion in mind, we obtain the following geometrical interpretation of its image $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$ inside $\mathbb{H}(\mathbb{A}_X, \mathcal{C}_p)$.

Theorem 4.4. There is a canonical bijection of sets

$$\operatorname{Cov}(X, \mathcal{C}_p)/\operatorname{Jac}(X)[p] \xrightarrow{1:1} \operatorname{\mathbb{H}_{ratl}}(\mathbb{A}_X, \mathcal{C}_p)/\operatorname{Aut}(\mathcal{C}_p).$$
(4.5)

Proof. First observe that Jac(X)[p] acts freely on the r.h.s. of (4.3) since $L \in Jac(X)[p]$ acts by sending a pair (\mathcal{L}, D) to $(D, \mathcal{L} \otimes L)$ and, thus, we conclude that

$$\operatorname{Cov}(X, \mathcal{C}_p)/\operatorname{Jac}(X)[p] \xrightarrow{1:1} \begin{cases} D \in \operatorname{Div}(X) \text{ such that} \\ \deg(D) \equiv 0 \mod p \end{cases} \right\} / \mathcal{Cov}(X, \mathcal{C}_p)/\operatorname{Jac}(X)[p]$$

where $D_1 = \sum_x a_x^1 x$ and $D_2 = \sum_x a_x^2 x$ are equivalent if there exists $b \in (\mathbb{Z}/(p))^*$ such that $ba_x^1 \equiv a_x^2 \mod p$ for all x.

On the other hand, recalling (2.13) and (2.20), it is clear that

$$\mathbb{H}_{\mathrm{ratl}}(\mathbb{A}_X, \mathcal{C}_p) \simeq \begin{cases} D \in \mathrm{Div}(X) \text{ with coefficients in } \mathbb{Z}/(p) \\ \mathrm{such that } \deg(D) \equiv 0 \mod p \end{cases} \end{cases}$$

Finally, from (2.14) one has an isomorphism $\operatorname{Jac}(X)[p] \simeq \mathbb{I}_X^p \cap \Sigma^* / \Sigma^{*p}$, and now the statement is deduced from the exact sequence (2.21), recalling Remark 2.22.

4.B. Enumeration.

Theorem 4.6. Let X/\Bbbk be a projective, irreducible, non-singular curve of genus g over an algebraically closed field \Bbbk with char(\Bbbk) $\neq p$. For a finite subset $\Re \subseteq X$ with r points,

1. The number of nontrivial unramified p-cyclic covers $\pi \in Cov(X, \mathcal{C}_p)$ is

$$\frac{p^{2g} - 1}{p - 1}.\tag{4.7}$$

2. The number of nontrivial p-cyclic covers $\pi \in Cov(X, C_p)$ with ramification contained in \mathfrak{R} , assuming \mathfrak{R} nonempty, is

$$\frac{p^{2g+r-1}-1}{p-1}.$$
(4.8)

3. The number of nontrivial p-cyclic covers $\pi \in Cov(X, C_p)$ with ramification equal to \mathfrak{R} , assuming \mathfrak{R} nonempty, is

$$p^{2g-1}((p-1)^{r-1} + (-1)^r). (4.9)$$

Proof.

- **1.** By (3.30), we have $\mathbb{H}_{\emptyset}(\Sigma, \mathcal{C}_p) = \operatorname{Jac}(X)[p]$, which has p^{2g} elements. Note that the isotropy subgroup of a class in $\mathbb{H}_{\emptyset}(\Sigma, \mathcal{C}_p)$ under the action of $\operatorname{Aut}(\mathcal{C}_p) \xrightarrow{\sim} (\mathbb{Z}/(p))^*$ is either trivial or all of $(\mathbb{Z}/(p))^*$. Observing that the former only holds for the trivial cover and recalling (1.32) yields (4.7).
- **2.** Combining (2.26), Corollary 3.18, and (3.36), $\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p)$ has p^{2g+r-1} elements. Reasoning as before, we obtain (4.8).
- **3.** Considering again (2.26) and the stratification by ramification (3.23), the number of elements in $\mathbb{H}(\Sigma, \mathcal{C}_p)$ with ramification exactly equal to \mathfrak{R} is p^{2g} times the number of solutions to the congruence

$$v_1 + \dots + v_r \equiv 0 \mod p, \qquad v_1, \dots, v_r \in (\mathbb{Z}/(p))^*,$$

which is given by

$$\frac{1}{p}((p-1)^r + (-1)^r(p-1)).$$
(4.10)

To see this, for a subset $A \subseteq \mathfrak{R}$, let h(A) be the number of tuples $v : \mathfrak{R} \to \mathbb{Z}/(p)$ with support equal to A, i.e. $v(x) \not\equiv 0 \mod p$ if and only if $x \in A$, and satisfying $\sum_{x \in \mathfrak{R}} v(x) \equiv 0 \mod p$. We want to compute $h(\mathfrak{R})$. Set $H(A) = \sum_{B \subseteq A} h(A)$, which is the number of zero-sum tuples with support contained in A. This is easily seen to equal $p^{\#A-1}$ when A is nonempty and 1 when $A = \emptyset$. Note that this only depends on the cardinality of A. By Möbius inversion, h(A) = $\sum_{B \subseteq A} (-1)^{\#A-\#B} H(B)$. Rearranging by cardinality we obtain $h(\mathfrak{R}) = (-1)^r +$ $\sum_{k=1}^r {r \choose k} (-1)^{r-k} p^{k-1}$, which simplifies to the given formula.

In this enumeration we are not counting the class of the trivial extension, since it is unramified and we assume \Re is nonempty. Noting that $\operatorname{Aut}(\mathcal{C}_p)$ acts freely on every class with prescribed nonempty ramification \Re yields (4.9) after dividing by p-1.

Alternatively, (4.9) may be deduced from (4.7) and (4.8) by an analogous process of Möbius inversion, which leads to

$$h(\mathfrak{R}) = (-1)^r \frac{p^{2g} - 1}{p - 1} + \sum_{k=1}^r \binom{r}{k} (-1)^{r-k} \frac{p^{2g+k-1} - 1}{p - 1},$$

and simplifying. In either case (2.26) is the starting point.

Note that (4.7) is [12, Theorem 3] and is also found on [9, p.490] and the first formula of [10, Theorem 7], while (4.9) is on [9, p. 500] and is the second formula in [10, Theorem 7]. These papers use different methods, although they ultimately rely on the properties of the fundamental group of a Riemann surface, as is common in this kind of problem. Thus we think it is interesting to point out that our approach, avoiding the dependence on the classical topological or analytic structure, allows us to generalize several aspects of *p*-cyclic Galois covers, such as classification and enumeration, to any algebraically closed base field of any characteristic (different from *p*) via the Galois theory of the adele ring. We may also deal with the general abelian case using these tools.

The reader may compare our methods to the references above, for example (4.10) is [13, Lemma 3] although we have arrived at this congruence via quite a different path.

Corollary 4.11. There exists a p-cyclic Galois cover of X with exactly $r \ge 0$ ramification points, except in the following two cases: g = 0 and r = 0, or $g \ge 0$ and r = 1.

Proof. This is immediate from the formulas in Theorem 4.6.

Recall that the Hurwitz existence problem is the question of whether a certain set of numerical data associated to a curve X, foremost among these the Riemann-Hurwitz relation, is in fact realizable by a cover $Y \to X$. This problem has been extensively studied, beginning with Hurwitz himself. It is known that for genus g >0 the answer is affirmative (major advances were made in [6] and many remaining cases dealt with later on), while the case g = 0 reduces to 3-point branched covers of the Riemann sphere and remains a difficult question. In particular when g > 0and r = 1, although as we see above, there are no p-cyclic Galois covers, other covers do exist as predicted by these general results. For g = 0 and r = 2, (4.9) says there is a unique p-cyclic cover. For r = 3 there are p - 2 such covers.

4.C. Rotation data. In this section we give an algebraic definition of rotation numbers (Definition 4.17), and using the Kummer pairing at ramification points, we prove that for $\mathbb{k} = \mathbb{C}$ it coincides with the classical one for the case of a compact connected Riemann surface.

To achieve this, we begin by considering the following classical construction of a p-cyclic cover of the projective line, which is standard in the literature (see e.g. [2, §5] and especially [7, §1]).

Set $\mathbb{k} = \mathbb{C}$, $X = \mathbb{P}^1$, p a prime number, and $\mathfrak{R} := \{x_1, \ldots, x_r\}$ a finite set of distinct points of X. Let Y be the normalization of

$$y^p = f(x) \coloneqq (x - x_1)^{v_1} \cdots (x - x_r)^{v_r},$$
 (4.12)

where $\pi : Y \to X$ maps (x, y) to x, and the exponents v_i satisfy $0 < v_i < p$ and $\sum_i v_i \equiv 0 \mod p$ (the latter is equivalent to being unramified at ∞). Observe that the Riemann-Hurwitz formula implies that $r \ge 1$ in the above expression for f(x).

Fix a nontrivial character $\chi : \mathcal{C}_p \to \mu_p \subseteq \mathbb{C}^*$ and let $\zeta = \chi(1) \in \mu_p$, which is a primitive *p*th root of unity. This defines an action of \mathcal{C}_p on *Y* where 1 acts via the automorphism

$$\tau: (x, y) \mapsto (x, \zeta y). \tag{4.13}$$

Thus τ is a generator of $\operatorname{Gal}(Y/X)$, and allows us to identify \mathcal{C}_p with $\operatorname{Gal}(Y/X)$.

If we assume that $gcd(v_1, \ldots, v_r) = 1$, then Y is irreducible and $\mathcal{O}_X \to \pi_* \mathcal{O}_Y$ at the generic point is

$$\Sigma \coloneqq \mathbb{C}(x) \hookrightarrow \Omega_f = \Sigma[y]/(y^p - f(x))$$

as considered in Example 1.17. This is what we termed a (\mathcal{C}_p, χ) -Kummer extension (Definition 1.4).

Thus $\pi: Y \to X$ is a *p*-cyclic cover with ramification locus

$$\operatorname{Ram}(\pi) = \mathfrak{R} = \{x_1, \dots, x_r\},\$$

having ramification index at x_i equal to $p/(p, v_i) = p$. Thus π defines a class in $\operatorname{Cov}_{\mathfrak{R}}(X, \mathcal{C}_p)$ and the class of Ω_f is an element of $\mathbb{H}_{\mathfrak{R}}(\Sigma, \mathcal{C}_p)$, related via the correspondence in Proposition 3.35.

Let us now change our point of view to the geometric adele ring. Under the canonical map (2.1), the corresponding adelic algebra $\mathbb{A}_Y = \mathbb{A}_X\{\mathbb{f}^{1/p}\}$ is a rational (\mathcal{C}_p, χ) -Kummer adelic extension of \mathbb{A}_X (Definition 2.18), hence lying by definition in $\mathbb{H}_{ratl}(\mathbb{A}_X, \mathcal{C}_p)$. The characteristic polynomial of y is equal to $C_y(T) = T^p - \mathbb{f}$, and thus the associated valuation vector of this extension (Definition 2.7) is

$$(v_x)_x \coloneqq \upsilon(\mathbb{A}_Y, \mathcal{C}_p, \chi) = (\upsilon_x(\mathbb{f}))_x = \begin{cases} v_i \mod p, & x = x_i \in \mathfrak{R}, \\ 0 \mod p, & x \notin \mathfrak{R}, \end{cases}$$

where we take as χ -primitive element the class of T modulo $T^p - f$. This invariant satisfies

$$\sum_{x} v_x \equiv 0 \bmod p$$

in accordance with Corollary 2.23. Recall that these invariants classify (\mathcal{C}_p, χ) adelic extensions and hence *p*-cyclic covers.

In the classical theory of Riemann surfaces, one considers the so-called *rota*tion numbers. These are defined in terms of the automorphism τ given by the action (4.13), namely, choosing $\zeta = e^{2\pi i/p}$, in a neighborhood of a branch point $x = (x_i, 0)$ one sees that τ rotates a small disk around this point by an angle $2\pi\rho_x/p$; the integer $\rho_x \mod p$ is the corresponding rotation number. A simple calculation shows they are the inverses modulo p of the valuations v_x :

$$\rho_x v_x \equiv 1 \mod p. \tag{4.14}$$

The usual approach based on complex analytic and topological methods (for example [7]), classifies the *p*-cyclic branched covers of the punctured Riemann sphere in terms of the rotation data $(\rho_x)_{x \in \Re}$.

On the other hand, in §2.B, we have already classified *p*-covers in arbitrary characteristic prime to p in a purely algebraic manner, reflected explicitly in the valuation vector $(v_x)_x$. To complete the algebraic picture, we will show how both the valuations and the rotation data arise naturally from the local Kummer symbols at each ramified point $x \in \mathfrak{R}$. For this we need to recall some notions introduced in [16].

For $x \in \mathfrak{R}$, by Kummer theory, there is a unique (up to isomorphism) abelian extension E_x of K_x of exponent p, generated by adjoining the pth root of any non pth power in K_x . This is the algebraic analog of the classical Puiseux series expansion in the theory of Riemann surfaces, since for example $K_x \simeq \Bbbk((z_x))$ where z_x is a uniformizer at x, and the local valuation v_x

$$\ker(K_x^* \xrightarrow{v_x} \mathbb{Z} \to \mathbb{Z}/(p)) = \{\lambda : v_x(\lambda) \equiv 0 \bmod p\} = K_x^{*p}, \tag{4.15}$$

induces an isomorphism $K_x^*/K_x^{*p} \simeq \mathbb{Z}/(p)$. Now, we have the Kummer perfect pairing for E_x ,

$$\langle g, \lambda \rangle_x = \frac{g(\lambda^{1/p})}{\lambda^{1/p}} : \operatorname{Gal}(E_x/K_x) \times \overset{K^*_x}{/} K^{*p}_x \to \mu_p,$$

$$(4.16)$$

where $\lambda^{1/p}$ is any *p*th root of λ in E_x .

Now, as we saw in §1.C, the isomorphism $\mathbb{A}_Y \simeq \mathbb{A}_X \otimes_{\Sigma} \Omega$ (1.29) is equivariant under the action of $\operatorname{Gal}(Y/X)$. Thus an automorphism τ of Y/X can be restricted to each fiber of the cover $\pi : Y \to X$. In particular, for a ramification point $x \in \mathfrak{R}$, τ induces an element $\tau_x \in \operatorname{Gal}(E_x/K_x)$ in each local Galois group.

The explicit computations with local Kummer symbols carried out in [16, §3.E] lead to the following algebraic definition of rotation data, valid over any algebraically closed base field k of characteristic prime to p.

Definition 4.17 (Algebraic rotation data). The algebraic rotation data for an automorphism τ of a p-cyclic Galois cover $\pi: Y \to X$ with ramification locus \mathfrak{R} is defined by

$$\left(\log_{\zeta}\langle \tau_x, z_x \rangle_x\right)_{x \in \mathfrak{R}} \in \prod_{x \in \mathfrak{R}} \mathbb{Z}/(p),$$
(4.18)

where \log_{ζ} is the discrete logarithm associated to a fixed primitive pth root of unity ζ .

Proposition 4.19. The algebraic rotation data for a nontrivial automorphism τ of the superelliptic curve (4.12) coincide with the classical analytic rotation data, under the assumption that $\chi(\tau) = \zeta$, where we identify C_p with $\operatorname{Gal}(Y/X)$ as above.

Proof. This follows from [16, Proposition 3.79], which together with our choices of χ , τ and ζ show that the numbers $\log_{\zeta} \langle \tau_x, z_x \rangle_x$ are indeed the inverses modulo p of the valuations $v_x = v_x(\mathbf{f})$ and thus by (4.14) are equal to the rotation numbers.

Remark 4.20. Note that (4.18) depends not only on the automorphism τ but also on the choice of primitive pth root of unity ζ , and that the relation $\chi(\tau) = \zeta$ is needed for the equivalence in Proposition 4.19. In general it would hold only modulo a constant multiple in $(\mathbb{Z}/(p))^*$.

Thus, the equivalence class of \mathfrak{R} -tuples as in (4.18) modulo the action of $(\mathbb{Z}/(p))^*$ by multiplication (see for example [16, Theorem 3.71]) is independent of these choices.

References

- [1] Borevich, A. Z., Kummer extensions of rings, J. Math. Sci. 11, 514-534 (1979).
- [2] Broughton, A; Shaska, T. and Wootton, A, On automorphisms of algebraic curves, in "Algebraic curves and their applications", 175–212, Contemp. Math., 724, Amer. Math. Soc., Providence, RI, (2019).
- [3] Cassels, J.W.S.; Fröhlich, A., Algebraic Number Theory (Proc. Instructional Conf., Brighton, 1965) Thompson, Washington, D.C.
- [4] Chase, S. U.; Harrison, D.K.; Rosenberg, A., Galois Theory and Cohomology of Commutative Rings, Memoirs of the AMS (52), American Mathematical Society, Providence, R.I. (1965).
- [5] Cornalba, M., On the locus of curves with automorphisms, Ann. Mat. Pura Appl. (4) 149 (1987), 135-151.
- [6] Edmonds, A. L.; Kulkarni, R. S.; Stong, R. E., Realizability of Branched Coverings of Surfaces Trans. AMS 282, No. 2 (1984), 773–790.
- [7] González Díez, G., Loci of Curves Which are Prime Galois Coverings of P¹, Proc. London Math. Soc. Volume (3) 62, 469–489.
- [8] Greither, C., Cyclic Galois Extensions of Commutative Rings, LNM 1534, Springer (1992).
- [9] Jones, G. A., Enumeration of Homomorphisms and Surface-Coverings, The Quarterly Journal of Mathematics, 46(4), (1995) 485–507.
- [10] Kwak, J.; Lee, J.; Mednykh, A., Enumerating Branched Surface Coverings from Unbranched Ones, LMS Journal of Computation and Mathematics 6 (2003), 89–104.
- [11] Lorenz, F.; Roquette, P., The theorem of Grunwald-Wang in the setting of valuation theory, Valuation theory and its applications, Vol. II (Saskatoon, SK, 1999) Fields Inst. Commun. 33 175–212 (2003)
- [12] Mednyh, A. D., On Unramified Coverings of Compact Riemann Surfaces, Soviet Math. Dokl. 20, No. 1 (1979).
- [13] Mednyh, A. D., Determination of the Number of Nonequivalent Coverings Over a Compact Riemann Surface, Soviet Math. Dokl. 19, No. 2 (1978).
- [14] Milne, J.S., Étale Cohomology, PMS-33, Princeton University Press (1980).
- [15] Nagahara, T., Separable polynomials over a commutative ring II, Math. J. Okayama Univ. 15 (1971/72), 149–162.
- [16] Navas Vicente, L.M.; Plaza Martín, F.J.; Serrano Holgado, Á., Kummer theory over the geometric adeles of a curve, arXiv:2310.13443 [math.AG] (2023).
- [17] Paques, A., On the primitive element and normal basis theorems, Comm. Algebra 16, no. 3 (1988), 443–455.
- [18] Paques, A., Galois theories: a survey, Advances in mathematics and applications, Springer (2018) 247–273.
- [19] The Stacks project, https://stacks.math.columbia.edu.
- [20] Sutherland, A., Stronger arithmetic equivalence, Discrete Analysis 2021:23, 23 pp.
- [21] Szamuely, T., Galois Groups and Fundamental Groups, Cambridge Studies in Advanced Mathematics (117), Cambridge University Press (2009).
- [22] Tate, J., Endomorphisms of abelian varieties over finite fields, Inv. Math. 2 (1966), 134– 144.
- [23] Turner, S., Adele rings of global field of positive characteristic, Bol. Soc. Bras. Mat. 9 (1978), 89–95.

Email address: navas@usal.es

Email address: fplaza@usal.es

Departamento de Matemáticas and IUFFyM, Universidad de Salamanca, Plaza de la Merced 1-4, 37008 Salamanca. Spain.