

Precision Higgs Constraints in U(1) Extensions of the Standard Model with a Light Z'-Boson

Zoltán Péli*

*Institute for Theoretical Physics, ELTE Eötvös Loránd University,
Pázmány Péter sétány 1/A, 1117 Budapest, Hungary*

Zoltán Trócsányi†

*Institute for Theoretical Physics, ELTE Eötvös Loránd University
and HUN-REN ELTE Theoretical Physics Research Group,
Pázmány Péter sétány 1/A, 1117 Budapest, Hungary, also at
University of Debrecen, Bem tér 18/A, 4026 Debrecen, Hungary*

Abstract

Anomaly free $U(1)$ extensions of the standard model (SM) predict a new neutral gauge boson Z' . When the Z' obtains its mass from the spontaneous breaking of the new $U(1)$ symmetry by a new complex scalar field, the model also predicts a second real scalar s and the searches for the new scalar and the new gauge boson become intertwined. We present the computation of production cross sections and decay widths of such a scalar s in models with a light Z' boson, when the decay $h \rightarrow Z'Z'$ may have a sizeable branching ratio. We show how Higgs signal strength measurement in this channel can provide stricter exclusion bounds on the parameters of the model than those obtained from the total signal strength for Higgs boson production.

I. INTRODUCTION

While the discovery of the Higgs boson [1, 2] has established the existence of a scalar elementary particle, the thorough understanding of the role of scalar fields in Nature remains elusive. So far all experimental results are in agreement with the structure of the scalar sector of the standard model (SM) [3], although the scalar potential has not yet been fully confirmed experimentally, which allows for the existence of an extended scalar sector. Indeed, there is a vigorous experimental search for new scalar particles at the LHC [4]. The more complex such an extended sector, the more new particles should exist and the more difficult the search strategies.

The $U(1)$ extensions of the SM have the potential to explain several beyond the standard model (BSM) phenomena at the cost of predicting the existence of Z' , a new neutral gauge boson. In the simplest scenario Z' acquires its mass from the spontaneous breaking of a new scalar field χ [5], hence, the model also predicts a new scalar particle s . Z' bosons appear in a wide variety of models, for a comprehensive review on them see Ref. [6]. Experiments searched extensively for new scalar particles, as well as a new Z' boson (for an incomplete list we refer to Refs. [7–10]). In such $U(1)$ extensions the existence of s and Z' are interconnected. For instance, if Z' is sufficiently light, then the Higgs boson and the new scalar can decay into a Z' pair as shown on Figure. 1. The channel $h \rightarrow Z'Z'$ can

* zoltan.peli@ttk.elte.hu

† zoltan.trocsanyi@cern.ch

significantly alter the decay properties of the Higgs boson, which consequently can be used to constrain the free parameters of the model.




FIG. 1. Tree-level diagram of the Higgs boson decaying into two Z' bosons.

An example of such a simple $U(1)$ extension of the standard model, with the potential to explain all established experimental observations that cannot be interpreted within the SM, is the superweak extension of the standard model (SWSM) [11]. It is designed to explain (i) the non-vanishing masses of neutrinos, (ii) the nature of dark matter, (iii) the origin of baryon asymmetry, and (iv) the role of scalar fields in the Universe.

The SWSM extends the gauge group of the SM $G = SU(3)_c \otimes SU(2)_L \otimes U(1)_Y$, to $G \otimes U(1)_z$. The corresponding neutral gauge field becomes massive by introducing a complex scalar field χ with non-zero vacuum expectation value (VEV). The gauge and gravity anomalies are required to cancel, which is achieved by introducing three generations of right-handed neutrinos – dubbed heavy neutral lepton, (HNL) below – with properly chosen z charges z_N . If we also allow the presence of Dirac-type Yukawa terms for the HNLs, then two independent z charges remain. We choose the z charge z_ϕ of the Brout-Englert-Higgs (BEH) field and z_N to be the free, model dependent parameters. All new fields are neutral under the standard model gauge interactions. The SWSM fixes the z charge of the BEH field to $z_\phi = 1$, and those of the HNLs to $z_N = \frac{1}{2}$ by allowing also Majorana-type Yukawa term in the Lagrangian for the HNLs and normalizing the new gauge coupling g_z such that the new scalar has z charge $z_\chi = -1$.

In this work we compute benchmark points for production and decay of a new scalar s for models with a light Z' boson. If the Z' is light the decay $h \rightarrow Z'Z'$ may have a sizeable branching ratio with dominant contribution independent of z_ϕ and z_N . We show that exclusion limits on new singlet scalar particles obtained from signal strength measurements can be improved by taking into account the process $h \rightarrow Z'Z'$. We provide a *Mathematica* notebook that can be used flexibly to compute benchmark points relevant for scalar searches at the LHC. While the specific model we have in mind is the SWSM, yet our analysis is

valid for an arbitrary assignment of z charges.

Experiments often search for the so-called dark photon, which appears in the $U(1)$ extensions with $z_\phi = z_N = 0$. These results can be translated to $U(1)$ extensions with arbitrary z charge assignment [12]. The prospects of discovering a dark photon were studied extensively [13] when one also takes into account the channels $h \rightarrow ZZ'$ and $h \rightarrow Z'Z'$. Our approach is different in the sense that we investigate the prospects of discovering a new scalar s when the model also predicts a Z' boson with unconstrained z_ϕ and z_N charges.

II. SCALAR SECTOR OF THE SWSM

In this section we collect the details of the model only to the extent used in the present analyses.

In the scalar sector, in addition to the $SU(2)_L$ -doublet Brout-Englert-Higgs field

$$\phi = \begin{pmatrix} \phi^+ \\ \phi^0 \end{pmatrix} = \frac{1}{\sqrt{2}} \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}, \quad (\text{II.1})$$

the model contains a complex scalar SM singlet χ . The Lagrangian of the scalar fields contains the potential energy

$$V(\phi, \chi) = -\mu_\phi^2 |\phi|^2 - \mu_\chi^2 |\chi|^2 + (|\phi|^2, |\chi|^2) \begin{pmatrix} \lambda_\phi & \frac{\lambda}{2} \\ \frac{\lambda}{2} & \lambda_\chi \end{pmatrix} \begin{pmatrix} |\phi|^2 \\ |\chi|^2 \end{pmatrix} \subset -\mathcal{L} \quad (\text{II.2})$$

where $|\phi|^2 = |\phi^+|^2 + |\phi^0|^2$. After spontaneous symmetry breaking, we parametrize the scalar fields as

$$\phi = \frac{1}{\sqrt{2}} \begin{pmatrix} -i\sqrt{2}\sigma^+ \\ v + h' + i\sigma_\phi \end{pmatrix}, \quad \chi = \frac{1}{\sqrt{2}}(w + s' + i\sigma_\chi) \quad (\text{II.3})$$

where v and w are the vacuum expectation values of ϕ and χ . The fields h' and s' are real scalars, σ^+ is a charged, while σ_ϕ and σ_χ are neutral Goldstone bosons that are gauge eigenstates.

The gauge and mass eigenstates are related by the rotations

$$\begin{pmatrix} h \\ s \end{pmatrix} = \mathbf{Z}_S \begin{pmatrix} h' \\ s' \end{pmatrix}, \quad \begin{pmatrix} \sigma_Z \\ \sigma_{Z'} \end{pmatrix} = \mathbf{Z}_G \begin{pmatrix} \sigma_\phi \\ \sigma_\chi \end{pmatrix}, \quad (\text{II.4})$$

with

$$\mathbf{Z}_X = \begin{pmatrix} \cos \theta_X & -\sin \theta_X \\ \sin \theta_X & \cos \theta_X \end{pmatrix} \quad (\text{II.5})$$

where we denoted the mass eigenstates with h , s and σ_Z , $\sigma_{Z'}$. The angles θ_S and θ_G are the scalar and Goldstone mixing angles that can be determined by the diagonalization of the mass matrix of the real scalars and that of the neutral Goldstone bosons. In the following, we are going to use the abbreviations $c_X = \cos \theta_X$ and $s_X = \sin \theta_X$ for mixing angles.

A. Scalar couplings

The vertices that involve the scalars are related to the corresponding vertices in the SM by simple proportionality factors involving the scalar mixing angle θ_S . Hence, if possible, we present the Feynman rules expressed using the corresponding SM rule.

- Scalar-fermion couplings:

$$\Gamma_{hf\bar{f}} = c_S \Gamma_{hf\bar{f}}^{\text{SM}}, \quad \Gamma_{sf\bar{f}} = s_S \Gamma_{hf\bar{f}}^{\text{SM}}. \quad (\text{II.6})$$

- Scalar-vector boson couplings: $i\Gamma_{SVV}g^{\mu\nu}$ where

$$\Gamma_{hWW} = c_S \Gamma_{hWW}^{\text{SM}}, \quad \Gamma_{sWW} = s_S \Gamma_{hWW}^{\text{SM}} \quad (\text{II.7})$$

and

$$\begin{aligned} \Gamma_{hZZ} &= 2M_Z^2 \left(c_S c_Z^2 - \frac{s_S}{\tan \beta} s_Z^2 \right), & \Gamma_{sZZ} &= 2M_Z^2 \left(s_S c_Z^2 + \frac{c_S}{\tan \beta} s_Z^2 \right), \\ \Gamma_{hZZ'} &= 2M_Z M_{Z'} s_Z c_Z \left(c_S + \frac{s_S}{\tan \beta} \right), & \Gamma_{sZZ'} &= 2M_Z M_{Z'} s_Z c_Z \left(s_S - \frac{c_S}{\tan \beta} \right), \\ \Gamma_{hZ'Z'} &= 2M_{Z'}^2 \left(c_S s_Z^2 - \frac{s_S}{\tan \beta} c_Z^2 \right), & \Gamma_{sZ'Z'} &= 2M_{Z'}^2 \left(s_S s_Z^2 + \frac{c_S}{\tan \beta} c_Z^2 \right) \end{aligned} \quad (\text{II.8})$$

where $\tan \beta = w/v$ is the ratio of the VeVs. In the SWSM region of the parameter space the neutral boson mixing angle θ_Z is smaller than $\mathcal{O}(10^{-3})$ [14], hence we consider the leading contributions to the vertex factors Γ_{SVV} in the limit $\theta_Z \rightarrow 0$ ($c_Z = 1$, $s_Z = 0$).

- Scalar-scalar couplings:

$$\begin{aligned} \Gamma_{hss} &= + \frac{M_H^2 + 2M_s^2}{v} s_S c_S \left(\frac{c_S}{\tan \beta} - s_S \right), \\ \Gamma_{shh} &= - \frac{2M_H^2 + M_s^2}{v} s_S c_S \left(\frac{s_S}{\tan \beta} + c_S \right). \end{aligned} \quad (\text{II.9})$$

In these vertex factors the replacement $h \rightarrow s$ can be achieved by ($c_S \rightarrow s_S$, $s_S \rightarrow -c_S$) and the replacement of one $Z \rightarrow Z'$ by *one factor* of ($c_Z \rightarrow s_Z$, $s_Z \rightarrow -c_Z$). Thus, it is sufficient to define explicitly Γ_{hZZ} and Γ_{hss} .

III. PRODUCTION OF SCALAR PARTICLES IN THE LHC

The production cross sections of a scalar particle that mixes with the SM scalar, which we shall refer to as *Higgs-like new scalar*, can be computed based on Ref. [15]. There are several production channels for the Higgs boson at the LHC, and similarly for the new scalar as well. These channels involve (i) gluon-gluon fusion (ggF), (ii) associated production of the scalar with a vector boson (VS), (iii) vector boson fusion (VBF), and (iv) associated production of the scalar with a $t\bar{t}$ pair (ttS).

In the SWSM the new scalar is directly coupled to heavy quarks, so the dominant production channel is the gluon-gluon fusion, as in the SM. The difference compared to the $\Gamma_{hf\bar{f}}$ vertex is only a proportionality factor obtained from the scalar mixing angle θ_S given in Eq. (II.6), hence the gluon fusion cross sections for the production of the two scalar particles are proportional,

$$\sigma(gg \rightarrow h) = c_S^2 \sigma^{\text{SM}}(gg \rightarrow h), \quad \sigma(gg \rightarrow s) = s_S^2 \sigma^{\text{SM}}(gg \rightarrow h)|_{M_h \rightarrow M_s}. \quad (\text{III.1})$$

The associated production of the scalar boson with a vector boson V involves the couplings (II.8) between the scalars and the vector bosons, which provide factors of c_S^2 and s_S^2 as compared to the SM. In addition, there is a contribution due to the SZ' channel, but it is negligibly small compared to the SW or SZ channels for small Z' masses, $\xi = M_{Z'}/M_Z \ll 1$, relevant in the SWSM parameter space. The V-A couplings of the Z boson to the quarks also receives BSM corrections, but these are well measured quantities and the deviation from the SM must be small. Hence, the Vh production cross section receives only an overall factor c_S^2 as compared to the SM model prediction,

$$\sigma(pp \rightarrow Vh) = c_S^2 \sigma^{\text{SM}}(pp \rightarrow Vh). \quad (\text{III.2})$$

Thus, measuring $\sigma(pp \rightarrow Vh)$ and comparing it to the SM prediction for this process constrains c_S .

By far the most complicated process to compute its cross section is the vector boson fusion. However, the radiative corrections are known to be small [15], so we consider this channel only at leading order (LO) in perturbation theory. The partonic process is $qq \rightarrow qq + (V^*V^* \rightarrow h)$. The squared matrix element of this process at LO is proportional to $G_F^3 M_V^8$, which means that the VBF process is also heavily suppressed, when $V = Z'$ and

$\xi \ll 1$. Thus, only $V = Z$ and W contributes with suppression factors c_S^2 for the Higgs production and s_S^2 for the new Higgs-like scalar production,

$$\sigma(qq \rightarrow qqh) = c_S^2 \sigma^{\text{SM}}(qq \rightarrow qqh), \quad \sigma(qq \rightarrow qqS) = s_S^2 \sigma^{\text{SM}}(qq \rightarrow qqh)|_{M_h \rightarrow M_s}. \quad (\text{III.3})$$

The scalar boson can also be produced in association with heavy quarks. This has the smallest contribution to the total production rate even when the heavy quark is the t-quark. The Higgs-like new scalar is directly coupled to heavy quarks just like in the Higgs boson, and the suppression factor in the production cross section is $\tan^2(\theta_S)$ just like in gluon-gluon fusion,

$$\sigma(pp \rightarrow t\bar{t}S) = s_S^2 \sigma^{\text{SM}}(pp \rightarrow t\bar{t}h)|_{M_h \rightarrow M_s}. \quad (\text{III.4})$$

Based on its small contribution to the total production cross section, we neglect this production channel in our analysis.

We see that all important production cross sections are proportional to the cross section of Higgs boson production in the SM with the relevant scalar mass value. Then it is sufficient to precisely compute the Higgs boson production cross section for different values of the Higgs mass. One can use automated software for that purpose, or alternatively one can save data using plots from Ref. [16], which we did here to obtain the relevant K -factors.

IV. DECAYS OF SCALARS

A. Total width of the Higgs boson

The SM theoretical prediction for the Higgs boson width is $\Gamma_h^{\text{SM}} = 4.07$ MeV, with a relative uncertainty of 4% [3]. The experimental measurements on the other hand are $\Gamma_h^{\text{ATLAS}} = 4.5_{-2.5}^{+3.3}$ MeV [17] and $\Gamma_h^{\text{CMS}} = 3.2_{-1.7}^{+2.4}$ MeV [18], display a much larger uncertainty than the SM theoretical prediction allowing for several BSM models to remain compatible with observations.

In the superweak and other $U(1)_z$ extensions with a light Z' boson ($M_{Z'} \ll M_Z$) the decay $h \rightarrow Z'Z'$ is allowed with partial width

$$\Gamma(h \rightarrow Z'Z') = \frac{G_F M_h^3}{16\sqrt{2}\pi} \left(\frac{s_S}{\tan\beta} \right)^2 + \mathcal{O}\left(\frac{M_{Z'}^2}{M_Z^2}\right), \quad (\text{IV.1})$$

and the total width of the Higgs boson is

$$\Gamma_h = c_S^2 \Gamma_h^{\text{SM}} + \Gamma(h \rightarrow Z'Z'), \quad (\text{IV.2})$$

which can be used to constrain the parameters θ_S and $\tan\beta$. We see that the total decay width can be smaller in the extended model than that in the SM depending on the relative effect of the scalar mixing angle and the partial decay width of the Higgs boson into the new neutral gauge boson pair.

B. Decay channels

A Higgs-like new scalar has similar decay channels as the Higgs-boson as listed.

Fermionic: The decays into SM fermions are only affected by an overall factor of s_S^2 as compared to the SM. If the mass of the Higgs-like scalar is above the $2m_N$ threshold, it can also decay into a pair of HNLs, with partial with

$$\Gamma(S \rightarrow N_i N_i) = \frac{G_f M_s m_{N_i}^2}{8\sqrt{2}\pi} \left(1 - \frac{4m_{N_i}^2}{M_s^2}\right)^{3/2} \left(\frac{c_S}{\tan\beta}\right)^2. \quad (\text{IV.3})$$

The Higgs boson can also decay into an HNL pair, provided the process is allowed kinematically. The corresponding partial decay width can be obtained from formula (IV.3) with the replacements $M_s \rightarrow M_h$ and $c_S \rightarrow s_S$.

Loop induced: This category includes the decays into $\gamma\gamma$, gg and $Z\gamma$ as well as $Z'\gamma$. In the important decay channels $s \rightarrow \gamma\gamma$ and gg , the decay widths are only multiplied by an overall s_S^2 factor,

$$\Gamma(s \rightarrow \gamma\gamma, Z\gamma, gg) = s_S^2 \Gamma^{\text{SM}}(h \rightarrow \gamma\gamma, Z\gamma, gg)|_{M_h \rightarrow M_s}. \quad (\text{IV.4})$$

The $Z'\gamma$ channel is also interesting because the LHC cast exclusion limits specifically on this channel (with branching ratio $\text{Br}(Z'\gamma) < 2\%$), but the prediction for this branching fraction is much smaller than the experimental limit. In the case of a light Z' and a modest $Z - Z'$ mixing, its contributions are negligibly small.

Decays into a pair of heavy vector bosons: The decays of the Higgs-like scalar into a charged W -pair is only affected by an overall factor of s_S^2 as compared to the SM prediction. Decays into heavy neutral bosons are controlled by the $SZ^{(\prime)}Z^{(\prime)}$ vertex. The vertex SZZ' is naturally suppressed, but the $SZ'Z'$ may be large.

Decays into scalars: In case $M_s > 2M_h$, the channel $s \rightarrow hh$ opens up, while the opposite decay, $h \rightarrow ss$ is excluded by the results for Γ_h^{exp} compared to the SM prediction.

The decay properties of a Higgs-like new scalar depend on all free parameters of the model. An indirect restriction on the parameter space can be derived from the experimental constraints on the Higgs-boson width Γ_h .

A more refined analysis reveals that for $\xi \ll 1$, such as the case in the SWSM, the free parameters reduce to a set of four parameters [19]:

$$M_s, s_S, \tan \beta, \text{ and } m_{N_i}, \quad (\text{IV.5})$$

where $\tan \beta$ absorbs the free parameters of the gauge sector in the combination

$$\tan \beta \propto \frac{M_{Z'}}{|s_Z| M_Z}. \quad (\text{IV.6})$$

The exact formula contains an additional factor depending on the z -charge of the BEH field and the kinetic mixing between the U(1) gauge fields [14], which is not important in our analysis. We note that for the B-L U(1) extension the proportionality factor is zero, so the scalar and gauge sector parameters are not related in that model.

V. IMPLEMENTATION

We computed the Higgs boson production cross section $\sigma(pp \rightarrow h + X)$ at $\sqrt{s} = 13.6$ TeV center of mass energies in proton proton collisions for several values of M_h in the range [100 GeV, 1 TeV]. Producing a Higgs-like scalar means that this cross section is multiplied with s_S^2 . In principle, the Z' boson affects these production rates in a nontrivial way, but these effects are negligibly small for $M_{Z'} \ll M_Z$, which is relevant for the SWSM.

We used the NNPDF3.0 set for the parton density functions (PDF), the LO set for LO predictions and the NLO set for the computations at the next-to-leading order (NLO) accuracy. We use the running strong coupling from the chosen PDF set. We take the values of the other input parameters m_b^{pole} , m_t^{pole} , G_f , M_Z and M_W from Ref. [3].

We compute the production cross section from ggF process at NLO QCD with NNLO QCD and NLO EW corrections included as a K-factor. The Vh process is implemented at NLO QCD and finally we compute the VBF at LO as the NLO corrections are known to be small with $K_{\text{NLO}}^{\text{VBF}} \sim 1.1$.

We include a *Mathematica* notebook `swsm_scalar.nb` which contains a precomputed set of the scalar boson productions cross sections and precisely computes the decay rates of the Higgs boson and the Higgs-like scalar.

VI. SIGNAL STRENGTHS

If a light neutral vector boson Z' exists, the Higgs boson can decay into a $Z'Z'$ pair, which affects the signal strengths in other channels. The signal strength measured at the LHC is

$$\mu_{\text{exp.}} = \frac{(\sigma \text{ Br})_{\text{obs}}}{(\sigma \text{ Br})_{\text{SM}}}, \quad (\text{VI.1})$$

while the theoretical value in the presence of new physics is

$$\mu_{\text{th.}} = \frac{(\sigma \text{ Br})_{\text{BSM}}}{(\sigma \text{ Br})_{\text{SM}}} \quad (\text{VI.2})$$

where the BSM subscript refers to the prediction in the full BSM model, including the SM contributions. The most precisely measured channel is $h \rightarrow ZZ^*$ [3]. In the superweak model and in general when the Higgs boson cannot decay into an ss or HNL pair, the predicted signal strength in the ZZ channel is

$$\mu_{ZZ} = \frac{c_S^4 \Gamma_h^{\text{SM}}}{c_S^2 \Gamma_h^{\text{SM}} + \Gamma(h \rightarrow Z'Z')} \quad (\text{VI.3})$$

because $\sigma_h^{\text{SWSM}} = c_S^2 \sigma_h^{\text{SM}}$ and $\Gamma^{\text{SWSM}}(h \rightarrow ZZ) = c_S^2 \Gamma^{\text{SM}}(h \rightarrow ZZ)$. Using the PDG24 [3] values ($\Gamma_h \simeq 0.00407$ GeV, $M_h \simeq 125$ GeV), we obtain

$$\mu_{ZZ} = \frac{c_S^4}{c_S^2 + 78.74 (c_S / \tan \beta)^2}, \quad (\text{VI.4})$$

which approaches c_S^2 for large values of $\tan \beta$. In that case the total signal strength provides a considerably more severe limit. The latter sums over all production and decay channels, hence

$$\mu_{\text{tot}} = \frac{\sigma_h^{\text{BSM}}}{\sigma_h^{\text{SM}}} = c_S^2, \quad (\text{VI.5})$$

in which case the experimental values are [20, 21]

$$\mu_{\text{tot}}^{\text{ATLAS}} = 1.05 \pm 0.06, \quad \mu_{\text{tot}}^{\text{CMS}} = 1.02 \pm 0.06. \quad (\text{VI.6})$$

Then, the total signal strength yields the bounds independent of $\tan \beta$

$$\theta_{S,\text{tot}}^{\text{ATL}} = 0.27, \quad \theta_{S,\text{tot}}^{\text{CMS}} = 0.32. \quad (\text{VI.7})$$




FIG. 2. Limit on the sine of the scalar mixing angle θ_S as a function of $\tan\beta$. The red (cyan) region is excluded at 95% C.L. by the ATLAS (CMS) measurement of the total and $h \rightarrow ZZ^*$ signal strengths. The blue disks with numbers i correspond to benchmark points BP i .

VII. BENCHMARKS AND EXCLUSION BOUNDS

For sufficiently small values of $\tan\beta$ the bound obtained from μ_{ZZ} tends to be more constraining than that obtained from μ_{tot} . The experimentally measured values for the ZZ^* signal strength are [20, 21]

$$\mu_{ZZ}^{\text{ATLAS}} = 1.04 \pm 0.09, \quad \mu_{ZZ}^{\text{CMS}} = 0.97 \pm 0.12, \quad (\text{VII.1})$$

which includes all the productions channels. Figure 2 shows limits obtained from the total signal strength measurements as well as those from the ZZ^* channel for $\tan\beta \leq 20$. The colored region on the $s_S - \tan\beta$ plane is excluded at 95% confidence level (C.L.). The blue discs with numbers i correspond to benchmark points BP i we propose based on the exclusion bounds obtained from the signal strength measurements. Explicitly, the three benchmark points are:

$$\begin{aligned} \text{BP1} : \theta_S &= 0.175, \tan\beta = 2, M_S = 500 \text{ GeV}, \\ \text{BP2} : \theta_S &= 0.175, \tan\beta = 10, M_S = 500 \text{ GeV}, \\ \text{BP3} : \theta_S &= 0.10, \tan\beta = 10, M_S = 1000 \text{ GeV}. \end{aligned} \quad (\text{VII.2})$$

where BP1 is allowed by μ_{tot} , but excluded by μ_{ZZ} , while BP2 and BP3 are still allowed. BP1 is chosen such that finding a scalar corresponding to BP1 would exclude a large class of BSM models predicting a light Z' boson. Using Eq. (IV.6) the gauge sector parameters are

$$\begin{aligned} \text{BP1} : M_{Z'} &= 18 \text{ MeV}, & s_Z &= 10^{-4}, \\ \text{BP2} : M_{Z'} &= 91 \text{ MeV}, & s_Z &= 10^{-4}, \\ \text{BP3} : M_{Z'} &= 91 \text{ MeV}, & s_Z &= 10^{-4}. \end{aligned} \tag{VII.3}$$

The Higgs-portal coupling λ can be expressed via the relation

$$\lambda = \frac{M_S^2 - M_H^2}{v^2} \frac{s_S c_S}{\tan \beta}, \tag{VII.4}$$

yielding

$$\lambda_{\text{BP1}} = 0.33, \quad \lambda_{\text{BP2}} = 0.07 \quad \text{and} \quad \lambda_{\text{BP3}} = 0.16. \tag{VII.5}$$

The SWSM model contains three families HNLs N_i . We consider the masses of these particles $M_{N_1} = \mathcal{O}(M_{Z'})$ and $M_{N_2} = M_{N_3} = 100 \text{ GeV}$ for all benchmark points. Then, the HNLs do not contribute to the decays of the Higgs boson, and can explain dark matter abundance [22]. We present the productions cross sections and branching ratios corresponding to BP1, BP2 and BP3 in Table I. Further points can be obtained similarly using the *Mathematica* code available on request.

VIII. SUMMARY

In our analysis we focused on $U(1)$ extensions of the SM where the new gauge boson becomes massive via the spontaneous breaking of the new $U(1)$ symmetry by a complex scalar field, such as in the superweak extension of the SM. We have shown that in the case of a light Z' , the new decay channel $h \rightarrow Z'Z'$ alters the signal strengths in all decay channels of the Higgs boson. Here we focused on the ZZ^* channel, which is measured experimentally with high precision. We obtained bounds on the free parameters of the model: the Higgs – new scalar mixing angle θ_S and the ratio $\tan \beta$ of the VeV of the new scalar to that of the BEH field. We presented three benchmark points – one (BP1) already excluded by the signal strength measurement in the ZZ^* decay channel of the Higgs boson and two other ones (BP2, BP3) still allowed. Finding a new scalar s corresponding to BP1 would mean

TABLE I. The productions cross sections and branching fractions corresponding to the benchmark points defined in Eq. (VII.2).

	h			s		
–	BP1	BP2	BP3	BP1	BP2	BP3
σ_{prod} [pb]	53	53	54	0.16	0.16	0.001
Γ [GeV]	$6.4 \cdot 10^{-3}$	$4.0 \cdot 10^{-3}$	$4.0 \cdot 10^{-3}$	8.3	2.9	8.5
$\text{Br}(hh)$	–	–	–	0.09	0.23	0.20
$\text{Br}(W^+W^-)$	0.13	0.21	0.21	0.13	0.37	0.37
$\text{Br}(ZZ)$	0.016	0.026	0.026	0.06	0.18	0.18
$\text{Br}(Z'Z')$	0.38	0.024	0.008	0.6	0.07	0.19
$\text{Br}(b\bar{b})$	0.36	0.57	0.58	$< 10^{-4}$	$< 10^{-4}$	$< 10^{-4}$
$\text{Br}(t\bar{t})$	–	–	–	0.05	0.14	0.05
$2 \times \text{Br}(NN)$	–	–	–	0.07	0.009	0.07

that no new light Z' boson exists as it would violate the exclusion bound derived from the effect of the decay $h \rightarrow Z'Z'$. Further benchmark points can be generated easily using the *Mathematica* notebook available on request.

ACKNOWLEDGMENTS

This research was funded by Excellence Programme of the Hungarian Ministry of Culture and Innovation grant number TKP2021-NKTA-64, by the National Research, Development and Innovation Office grant number NKFI-150794, and by the Hungarian Scientific Research Fund grant number PD-146527.

-
- [1] Aad G. *et al.* (ATLAS) Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC. *Phys. Lett.* **2012**, *B716*, 1–29 (*Preprint arXiv:1207.7214*)

- [2] Chatrchyan S. *et al.* (CMS) Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC. *Phys. Lett.* **2012**, *B716*, 30–61 (*Preprint* arXiv:1207.7235)
- [3] Navas S. *et al.* (Particle Data Group) Review of Particle Physics. *Phys. Rev. D* **2024**, *110*, 030001
- [4] Hayrapetyan A. *et al.* (CMS) Searches for Higgs boson production through decays of heavy resonances. **2024** (*Preprint* arXiv: 2403.16926)
- [5] Appelquist T., Dobrescu B. A., Hopper A. R., Nonexotic Neutral Gauge Bosons. *Phys. Rev. D* **2003**, *68*, 035012 (*Preprint* arXiv: 0212073)
- [6] Langacker P., The Physics of Heavy Z' Gauge Bosons, *em Rev. Mod. Phys.* **2009** *81*, 1199-1228 (*Preprint* arXiv: 0801.1345)
- [7] Robens T. Constraining extended scalar sectors at current and future colliders – an update. **2023** *Springer Proc. Phys* *292*, 141-152 (*Preprint* arXiv:2209.15544).
- [8] Aad G. *et al.* (ATLAS) Search for high-mass dilepton resonances using 139 fb⁻¹ of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector. *Phys. Lett. B*, **2019**, *796*, 68–87 (*Preprint* arXiv:1903.06248)
- [9] Sirunyan A.M. *et al.* (CMS) Search for resonant and nonresonant new phenomena in high-mass dilepton final states at $\sqrt{s} = 13$ TeV. *JHEP* **2021**, *07*, 208 (*Preprint* arXiv:2103.02708)
- [10] Banerjee D. *et al.* (NA64) Dark matter search in missing energy events with NA64. *Phys. Rev. Lett.* **2019**, *123*, 121801 (*Preprint* arXiv:1906.00176)
- [11] Trócsányi Z. Super-weak force and neutrino masses. *Symmetry* **2020**, *12*, 107 (*Preprint* arXiv:1812.11189)
- [12] Ilten P., Soreq Y., Williams M., and Xue W. Serendipity in dark photon searches. *JHEP* **2018**, *06*, 004 (*Preprint* arXiv:1801.04847)
- [13] Curtin D., Essig R., Gori S. and Shelton J. Illuminating Dark Photons with High-Energy Colliders. *JHEP* **2015**, *02* 157 (*Preprint* arXiv:1412.0018)
- [14] Péli Z. and Trócsányi Z. Exclusion bounds for neutral gauge bosons. *Phys. Rev. D* **2024**, *110*, 015027 (*Preprint* arXiv:2402.14786)
- [15] Djouadi A. The Anatomy of Electro-Weak Symmetry Breaking. I: The Higgs boson in the Standard Model. *Phys. Rept.* **2008**, *457*, 1–216 (*Preprint* arXiv:hep-ph/0503172)
- [16] [ATLAS wiki](#).

- [17] Aad G. *et al.* (ATLAS) Evidence of off-shell Higgs boson production from ZZ leptonic decay channels and constraints on its total width with the ATLAS detector. *Phys. Lett. B* **2023**, *846*, 138223 (*Preprint* arXiv:2304.01532)
- [18] Tumasyan A. *et al.* (CMS) Measurement of the Higgs boson width and evidence of its off-shell contributions to ZZ production. *Nature Phys.* **2022**, *18*, 1329–1334 (*Preprint* arXiv:2202.06923)
- [19] Péli Z. and Trócsányi Z. Precise prediction for the mass of the W boson in gauged U(1) extensions of the standard model. *Phys. Rev. D* **2023**, *108*, L031704 (*Preprint* arXiv:2305.11931)
- [20] Aad G. *et al.* (ATLAS) A detailed map of Higgs boson interactions by the ATLAS experiment ten years after the discovery. *Nature* **2022**, *607*, 52–59 [Erratum: *Nature* 612, E24 (2022)] (*Preprint* arXiv:2207.00092)
- [21] Tumasyan A. *et al.* (CMS) A portrait of the Higgs boson by the CMS experiment ten years after the discovery. *Nature* **2022**, *607*, 60–68 [Erratum: *Nature* 623, (2023)] (*Preprint* arXiv:2207.00043)
- [22] Iwamoto S., Seller K. and Trócsányi Z. Sterile neutrino dark matter in a U(1) extension of the standard model. *JCAP* **2022**, *01*, 035 (*Preprint* arXiv:2104.11248)