

LAZUTKIN COORDINATES OF THE MAXIMAL SYMMETRIC PERIODIC ORBITS ON THE ELLIPSE

KLAUDIUSZ CZUDEK

ABSTRACT. In De Simoi J., Kaloshin V., Wei Q. “Dynamical spectral rigidity among \mathbb{Z}_2 -symmetric strictly convex domains close to a circle (Appendix B coauthored with H. Hezari)”. Ann. of Math. 186.1 (2017), pp. 277–314 deformational spectral rigidity of \mathbb{Z}_2 symmetric domains close to the circle has been shown. One of the steps of the proof was to express the maximal symmetric periodic orbits in Lazutkin parametrization. Here using the action-angle variables we find the second order approximation of Lazutkin coordinates of the maximal symmetric periodic orbits on the ellipses.

1. INTRODUCTION

Let Ω be a C^1 domain in the plane, and let us consider billiard inside Ω , i.e. a particle moving along straight lines that bounces off the boundary with the rule the angle of incidence equals the angle of reflection. Having that the meaning of *periodic orbit* is clear. *The rotation number* of a periodic orbit equals p/q , where q is its period and p is the winding number, i.e. the number of times that the associated polygon wraps around the boundary $\partial\Omega$. *The length spectrum* is defined as

$$\mathcal{L}(\Omega) = \mathbb{N}\{\text{length of all periodic orbits in } \Omega\} \cup \mathbb{N}\{|\partial\Omega|\}.$$

A one-parameter family $(\Omega_t)_{t \in [-1,1]}$ of C^1 domains is called *isospectral* if $\mathcal{L}(\Omega_t) = \mathcal{L}(\Omega_0)$ for every $t \in [-1, 1]$. A domain Ω is *dynamically spectrally rigid* if every isospectral one-parameter family of domains with $\Omega_0 = \Omega$ is trivial (i.e. Ω_t is Ω_0 up to isometry).

It is an important open question if every sufficiently smooth domain Ω is dynamically spectrally rigid. Important progress towards the solution has been made by De Simoi, Kaloshin, Wei [4] where the authors show that every sufficiently smooth domain with axial symmetry, sufficiently close to the circle, is dynamically spectrally rigid if we admit only one-parameter families with axial symmetry (see also the introduction of [4] for more detailed exposition of known results).

The proof of the main result of [4] doesn’t incorporate all the periodic orbits in Ω but just some orbits of a specific type. Namely, for any axially symmetric Ω and $q \geq 2$ there exists a symmetric periodic orbit of rotation number $1/q$ passing through the marked point of $\partial\Omega$ and having maximal length among other symmetric periodic orbits of rotation number $1/q$ passing through the marked point. We call it the *maximal symmetric q -periodic orbit* (Lemma 3.4 [4]).

2020 *Mathematics Subject Classification.* Primary 37D40, Secondary 35P05, 35R30, 57J53.

Key words and phrases. convex billiards, Lazutkin coordinates, length spectrum, ellipse.

Another important ingredient of [4] is the Lazutkin parametrization. Let Ω be a C^1 domain. The Lazutkin parametrization of Ω is the parametrization of the form

$$(1) \quad x(s) = C_\Pi \int_0^s \rho(s')^{-2/3} ds', \quad C_\Pi = \left[\int_{\partial\Omega} \rho(s')^{-2/3} ds' \right]^{-1},$$

where ρ is the radius of curvature and s is the arc-length parametrization. Lazutkin weight is a function

$$\mu(x) = \frac{1}{2C_\Pi\rho(x)^{1/3}}.$$

It has been proven (Lemma 5.1 in [4]) that for every $\Omega \in \mathcal{S}^r$, $r \geq 8$, there exist C^{r-4} functions α (odd) and β (even) such that for any marked symmetric maximal q -periodic orbit $(x_q^0, \dots, x_q^{q-1})$

$$(2) \quad x_q^k = \frac{k}{q} + \frac{\alpha(k/q)}{q^2} + O(q^{-4}).$$

Moreover, if ϑ_q^k represents the angle of reflection of the trajectory at the k -th collision then

$$(3) \quad \vartheta_q^k = \frac{\mu(x_q^k)}{q} \left(1 + \frac{\beta(k/q)}{q^2} + O(q^{-4}) \right).$$

Theorem 1. Let us consider the ellipse

$$(4) \quad \mathcal{E}_e = \left\{ (x, y) \in \mathbb{R}^2 : x^2 + \frac{y^2}{b^2} = 1 \right\},$$

where $e = \sqrt{1 - b^2} \in (0, 1)$ represents the eccentricity. Then on $\Omega = \mathcal{E}_e$ the functions α and β defined by (2) and (3) take the forms

$$\alpha(t) = \frac{-eK(e)(1 - e^2)}{2\sqrt{1 - e^2 \operatorname{sn}^2(K(e)(1 - 4t), e)}} \left[(1 - 4t)K'(e)\operatorname{dn}(K(e)(1 - 4t), e) + \partial_2 \operatorname{am}(K(e)(1 - 4t), e) \right].$$

and

$$\begin{aligned} \beta(t) = \alpha'(t) - \frac{8K(e)^2 e^2}{3} & \left(\operatorname{cn}^2(\xi(t), e) - \operatorname{sn}^2(\xi(t), e) \right) + \frac{16K(e)^2 e^4 \operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)}{\operatorname{dn}^2(\xi(t), e)} \\ & + \frac{16K(e)}{1 - e^2} \hat{E}_1(t) \operatorname{dn}^5(\xi(t), e), \end{aligned}$$

where

$$\begin{aligned} \hat{E}_1(t) = & \frac{1}{24} \frac{e^4 \operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)}{[\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e)]^2} \\ & - \frac{1}{24} \frac{\operatorname{sn}^2(\xi(t), e) + b^2 \operatorname{cn}^2(\xi(t), e)}{\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e)} \\ & - \left[e^4 \frac{\operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)}{(\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e))^2} - \frac{1}{3} \right] \\ & \cdot \left[\frac{e^4 \operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)}{(\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e))^2} - \frac{\operatorname{sn}^2(\xi(t), e) + b^2 \operatorname{cn}^2(\xi(t), e)}{4(\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e))} + \frac{1}{3} \right] \\ & - \left[\frac{\operatorname{sn}^2(\xi(t), e) + b^2 \operatorname{cn}^2(\xi(t), e)}{4(\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e))} - \frac{1}{3} \right]^2 \\ & + e^4 \frac{\operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)}{\operatorname{cn}^2(\xi(t), e) + b^2 \operatorname{sn}^2(\xi(t), e)} \left[\frac{\operatorname{sn}^2(\xi(t), e) + b^2 \operatorname{cn}^2(\xi(t), e)}{4 \operatorname{sn}^2(\xi(t), e) \operatorname{cn}^2(\xi(t), e)} - \frac{1}{3} \right]. \end{aligned}$$

Here K is the complete elliptic integral of the first kind, dn is the Jacobi delta function, am is the Jacobi amplitude¹, ∂_2 represents the derivative with respect to the second variable and $\xi(t) = K(e)(1 - 4t)$.

2. NOTATION AND BASIC FACTS

The elliptic parametrization of (4) is given by

$$(5) \quad \gamma(\varphi) = (\sin \varphi, b \cos \varphi), \varphi \in [0, 2\pi).$$

The family of confocal ellipses C_λ , $\lambda \in (0, b)$, is given by

$$(6) \quad C_\lambda = \left\{ (x, y) \in \mathbb{R}^2 : \frac{x^2}{1 - \lambda^2} + \frac{y^2}{b^2 - \lambda^2} \right\}.$$

By $F(\varphi, k)$ and $K(k)$ we denote, respectively, the incomplete and complete Jacobi elliptic integral of the first kind

$$F(\varphi, k) = \int_0^\varphi \frac{d\tau}{\sqrt{1 - k^2 \sin^2 \tau}}, \quad K(k) = F\left(\frac{\pi}{2}, k\right).$$

Jacobi amplitude is defined by the relation $\operatorname{am}(u, k) = \varphi$ if and only if $F(\varphi, k) = u$. Further, the elliptic sine, cosine and delta amplitude are defined as

$$\operatorname{sn}(u, k) = \sin \operatorname{am}(u, k), \quad \operatorname{cn}(u, k) = \cos \operatorname{am}(u, k), \quad \operatorname{dn}(u, k) = \partial_u \operatorname{am}(u, k).$$

Let

$$(7) \quad k_\lambda^2 = \frac{e^2}{1 - \lambda^2}, \quad \delta_\lambda = F(\arcsin(\lambda/b), k_\lambda), \quad \omega_\lambda = \frac{\delta_\lambda}{2K(k_\lambda)}.$$

Let P_1, P_2, \dots be the collision points (with the boundary \mathcal{E}_e) of a certain billiard trajectory inside \mathcal{E}_e . Let us denote by $P_m P_{m+1}$ the segment joining P_m and P_{m+1} . It is a well-known fact (see e.g. [3]) that if $P_1 P_2$ is tangent to one of the confocal ellipses C_λ then $P_m P_{m+1}$ is

¹The definitions are given in Section 2.

tangent to C_λ for every $m \geq 1$. The next Proposition introduces a parametrization that is useful to study billiard trajectories whose all segments are tangent to C_λ .

Proposition 2 ([2]). Let $\lambda \in (0, b)$ and let us denote

$$S_\lambda(\varphi) = \frac{1}{4} - \frac{F(\frac{\pi}{2} - \varphi, k_\lambda)}{4K(k_\lambda)}.$$

Let $\varphi_0, \varphi_1, \dots$ be the collision points of a certain billiard trajectory inside \mathcal{E}_e whose all segments are tangent to C_λ . Then $S_\lambda(\varphi_k) = S_\lambda(\varphi_0) + k\omega_\lambda$ for every $k \geq 0$.

One the ellipse it is convenient to express the Lazutkin parametrization in terms of the elliptic parametrization on \mathcal{E}_e . Then it takes the form

$$(8) \quad x(\varphi) = \frac{1}{4} - \frac{F(\frac{\pi}{2} - \varphi, e)}{4K(e)}, \quad x(\varphi) \in [0, 1].$$

Observe (8) is the limit of the action angle variables $S_\lambda(\varphi)$ as $\lambda \rightarrow 0$. The Lazutkin weight takes the form (by 16.9.1 [1])

$$(9) \quad \mu(x) = \frac{2K(e)\sqrt{1-e^2}}{(\operatorname{cn}^2(K(e)(1-4x), e) + b^2\operatorname{sn}^2(K(e)(1-4x, e)))^{1/2}} = \frac{2K(e)\sqrt{1-e^2}}{|\operatorname{dn}(K(e)(1-4x, e))|}.$$

As explained in Introduction we are going to find the Lazutkin coordinates of orbits with the rotation number of the form $1/q$. Thus it is reasonable to define λ_q as the unique number with $\omega_{\lambda_q} = \frac{1}{q}$. Then by Proposition 2

$$(10) \quad S_{\lambda_q}(\varphi_q^k) = \frac{k}{q},$$

where φ_q^k is the elliptic coordinate of the k -th collision point if the maximal symmetric periodic orbit with the rotation number $1/q$.

3. CALCULATION OF α

3.1. The Taylor expansion of ω_λ and λ_q . Although it is we do not stress that in the notation many of the expressions below depend on the eccentricity e that should be treated as a fixed parameter. Taylor expansion of F at $\varphi = 0, k = e$ is

$$(11) \quad F(\varphi, k) = \varphi + \frac{1}{6}\varphi^3e^2 + \frac{1}{3}e\varphi^3(k - e) + O(\|(\varphi, k - e)\|^5).$$

We need also the expansions

$$(12) \quad \arcsin\left(\frac{\lambda}{\sqrt{1-e^2}}\right) = \frac{\lambda}{(1-e^2)^{1/2}} + \frac{\lambda^3}{6(1-e^2)^{3/2}} + O(\lambda^5) \quad \text{at } \lambda = 0$$

and

$$(13) \quad k_\lambda = \frac{e}{(1-\lambda^2)^{1/2}} = e\left(1 + \frac{\lambda^2}{2} + O(\lambda^4)\right) \quad \text{at } \lambda = 0.$$

Plugging (13) into the Taylor expansion of $K(k)$ at $k = e$ gives

$$(14) \quad K(k_\lambda) = K(e) + \frac{e}{2}K'(e)\lambda^2 + O(\lambda^4) \quad \text{at } \lambda = 0.$$

The Taylor series

$$\frac{1}{1+x} = 1 - x + x^2 - x^3 + O(x^4)$$

yields

$$(15) \quad \frac{1}{2K(k_\lambda)} = \frac{1}{2K(e)} - \frac{eK'(e)}{4K(e)^2} \lambda^2 + O(\lambda^4) \quad \text{at } \lambda = 0.$$

Plugging (12) and (13) to (11) we get (see (7))

$$(16) \quad \delta_\lambda = \frac{\lambda}{(1-e^2)^{1/2}} + \frac{\lambda^3(e^2+1)}{6(1-e^2)^{3/2}} + O(\lambda^5) \quad \text{at } \lambda = 0.$$

Multiplying (16) by (15) we arrive at

$$(17) \quad \omega_\lambda = \frac{\lambda}{2K(e)\sqrt{1-e^2}} \left[1 + \lambda^2 \left(\frac{-eK'(e)}{2K(e)} + \frac{e^2+1}{6(1-e^2)} \right) + O(\lambda^4) \right] \quad \text{at } \lambda = 0.$$

The application of the formula 3.6.25 [1] (p. 16) yields

$$(18) \quad \lambda_q = \frac{2K(e)\sqrt{1-e^2}}{q} \left(1 + \frac{1}{q^2} (2K(e)\sqrt{1-e^2})^2 \left(\frac{eK'(e)}{2K(e)} - \frac{e^2+1}{6(1-e^2)} \right) + O(q^{-4}) \right)$$

3.2. Elliptic coordinates of the maximal symmetric periodic orbit. Inverting the formula $\varphi \rightarrow S_\lambda(\varphi)$ from Proposition 2 yields

$$\varphi_q^k = \operatorname{am} \left(K(k_{\lambda_q}) \left(1 - \frac{4k}{q} \right), k_{\lambda_q} \right).$$

If $q \rightarrow \infty$, then $\lambda_q \rightarrow 0$ and $k_{\lambda_q} \rightarrow e$. Therefore using the Taylor expansion of Jacobi amplitude and (18) gives

$$\begin{aligned} \varphi_q^k &= \operatorname{am} \left(K(e) \left(1 - \frac{4k}{q} \right), e \right) \\ &\quad + \frac{1}{q^2} \left[2eK'(e)K(e)^2(1-e^2) \operatorname{dn} \left(K(e) \left(1 - \frac{4k}{q} \right), e \right) \left(1 - \frac{4k}{q} \right) \right. \\ &\quad \left. + 2eK(e^2)(1-e^2) \partial_2 \operatorname{am} \left(K(e) \left(1 - \frac{4k}{q} \right), e \right) \right] + O(q^{-4}), \end{aligned}$$

where ∂_2 stands for the partial derivative with respect to the second variable.

3.3. The Lazutkin coordinates of the maximal symmetric periodic orbit. By (8)

$$x_q^k = \frac{1}{4} - \frac{F(\frac{\pi}{2} - \varphi_q^k, e)}{4K(e)}$$

Using again the Taylor expansion of $\varphi \mapsto F(\varphi, e)$ at $\varphi = \operatorname{am} \left(K(e) \left(1 - \frac{4k}{q} \right), e \right)$

$$\partial_1 F|_{(\operatorname{am}(K(e)(1-4t), e))} = \frac{1}{\sqrt{1 - e^2 \operatorname{sn}^2(K(e)(1-4t), e))}}, \quad t \in [0, 1],$$

gives

$$(19) \quad x_q^k = \frac{k}{q} + \frac{\alpha(k/q)}{q^2} + O(q^{-4}),$$

where

$$(20) \quad \alpha(t) = \frac{-eK(e)(1-e^2)}{2\sqrt{1-e^2\operatorname{sn}^2(K(e)(1-4t), e))}} \left[(1-4t)K'(e)\operatorname{dn}(K(e)(1-4t), e) + \partial_2 \operatorname{am}(K(e)(1-4t), e) \right].$$

4. THE CALCULATION OF β

In order to find β we the expansion of $\frac{\sin \vartheta_q^k}{\mu(x_q^k)}$ in terms of $1/q$, $q \rightarrow \infty$, and then use the relation (B.1) from Appendix B [4]. To this end let us denote

$$L(\varphi, \varphi') = \|\gamma(\varphi) - \gamma(\varphi')\|,$$

where $\|\cdot\|$ is the Euclidean distance on \mathbb{R}^2 , γ is from (5). The function L is the generating function of the billiard map inside \mathcal{E}_e , thus

$$(21) \quad \begin{aligned} \sin \vartheta_q^k &= \sqrt{1 - \left[\frac{\partial_\varphi L(\varphi, \varphi_q^{k+1})|_{\varphi=\varphi_q^k}}{|\gamma'(\varphi_q^k)|} \right]^2} \\ &= \sqrt{1 - \frac{[(\cos \varphi_q^{k+1} - \cos \varphi_q^k) \sin \varphi_q^k - b^2(\sin \varphi_q^{k+1} - \sin \varphi_q^k) \cos \varphi_q^k]^2}{(\sin^2 \varphi_q^k + b^2 \cos^2 \varphi_q^k) \cdot ((\cos \varphi_q^{k+1} - \cos \varphi_q^k)^2 + b^2(\sin \varphi_q^{k+1} - \sin \varphi_q^k)^2)}} \end{aligned}$$

Let

$$(22) \quad t := \varphi_q^k, \quad d := \varphi_q^{k+1} - \varphi_q^k.$$

Then

$$(23) \quad \sin \varphi_q^{k+1} - \sin \varphi_q^k = d \cos t - \frac{d^2}{2} \sin t - \frac{d^3}{6} \cos t + \frac{d^4}{24} \sin t + \frac{d^5}{120} \cos t - \frac{d^6}{720} \sin t + O(d^7),$$

$$(24) \quad \cos \varphi_q^{k+1} - \cos \varphi_q^k = -d \sin t - \frac{d^2}{2} \cos t + \frac{d^3}{6} \sin t + \frac{d^4}{24} \cos t - \frac{d^5}{120} \sin t - \frac{d^6}{720} \cos t + O(d^7),$$

Let

$$(25) \quad \hat{A} = \sin^2 t + b^2 \cos^2 t, \quad \hat{B} = \sin t \cos t, \quad \hat{D} = \cos^2 t + b^2 \sin^2 t.$$

Then the denominator in the fraction appearing in (21) is

$$d^2 \hat{A}^2 + d^3 \hat{A} \hat{B} e^2 + d^4 \left(\frac{\hat{A} \hat{D}}{4} - \frac{\hat{A}^2}{3} \right) - \frac{d^5}{4} \hat{A} \hat{B} e^2 + d^6 \left(\frac{2}{45} \hat{A}^2 - \frac{1}{24} \hat{A} \hat{D} \right) + O(d^7).$$

In a similar manner the numerator is

$$d^2 \hat{A}^2 + d^3 \hat{A} \hat{B} e^2 + d^4 \left(\frac{1}{4} \hat{B}^2 e^4 - \frac{1}{3} \hat{A}^2 \right) - \frac{d^5}{4} \hat{A} \hat{B} e^2 + d^6 \left(\frac{2}{45} \hat{A}^2 - \frac{1}{24} \hat{B}^2 e^4 \right) + O(d^7).$$

After dividing the numerator and denominator by $d^2 \hat{A}^2$ and putting

$$(26) \quad M_1 = \frac{\hat{B}}{\hat{A}} e^2, \quad M_2 = \frac{\hat{D}}{4\hat{A}} - \frac{1}{3}, \quad M_3 = \frac{2}{45} - \frac{1}{24} \frac{\hat{D}}{\hat{A}}, \quad N_2 = \frac{\hat{B}^2}{4\hat{A}^2} e^4 - \frac{1}{3}, \quad N_3 = \frac{2}{45} - \frac{1}{24} \frac{\hat{B}^2}{\hat{A}^2} e^4$$

the denominator can be written down as

$$1 + dM_1 + d^2 M_2 - \frac{d^3}{4} M_1 + d^4 M_3 + O(d^5),$$

while the numerator

$$1 + dM_1 + d^2N_2 - \frac{d^3}{4}M_1 + d^4N_3 + O(d^5).$$

Using the Taylor expansion of $1/(1+x)$ the inverse of the denominator can be written down as

$$1 - dM_1 + d^2(M_1^2 - M_2) + d^3\left(\frac{M_1}{4} + 2M_1M_2 - M_1^3\right) + d^4\left(-M_3 + M_2^2 - \frac{M_1^2}{2} - 3M_1^2M_2 + M_1^4\right) + O(d^5).$$

The product of that with the numerator gives the expansion of the fraction appearing under the square root in (21) in terms of d and therefore of the entire expression under the square root. In order to obtain the expansion of $\sin \vartheta_q^k$ we factor out $d^2(M_2 - N_2)$ and use the Taylor expansion of $\sqrt{1+x}$. After that we arrive at

$$(27) \quad \sin \vartheta_q^k = \sqrt{M_2 - N_2} \left(d + d^2M_1 + d^3 \frac{\hat{E}_1}{M_2 - N_2} \right) + O(d^4),$$

where

$$(28) \quad \hat{E}_1 = M_3 - N_3 - N_2M_1^2 + N_2M_2 - M_2^2 + M_1^2M_2.$$

Now, let us express $d = \varphi_q^{k+1} - \varphi_q^k$ in terms of $1/q$. We have

$$x_q^{k+1} = \frac{k}{q} + \frac{1}{q} + \frac{\alpha(k/q)}{q^2} + \frac{\alpha'(k/q)}{q^3} + O(q^{-4}),$$

and therefore

$$x_q^{k+1} - x_q^k = \frac{1}{q} + \frac{\alpha'(k/q)}{q^3} + O(q^{-4}).$$

Let $y_q^k = K(e)(1 - 4x_q^k)$ and $z_q^k = K(e)(1 - 4k/q)$. By inverting (8) we arrive at

$$\begin{aligned} d = \varphi_q^{k+1} - \varphi_q^k &= 4K(e)\operatorname{dn}(y_q^k, e) \left[x_q^{k+1} - x_q^k + 2K(e)e^2(x_q^{k+1} - x_q^k)^2 \frac{\operatorname{sn}(y_q^k, e)\operatorname{cn}(y_q^k, e)}{\operatorname{dn}(y_q^k, e)} \right. \\ &\quad \left. - \frac{8}{3}K(e)^2e^2(x_q^{k+1} - x_q^k)^3 \left(\operatorname{cn}^2(y_q^k, e) - \operatorname{sn}^2(y_q^k, e) \right) \right] + O((x_q^{k+1} - x_q^k)^4) \\ &= 4K(e)\operatorname{dn}(y_q^k, e) \left[\frac{1}{q} + \frac{2K(e)e^2}{q^2} \frac{\operatorname{sn}(z_q^k, e)\operatorname{cn}(z_q^k, e)}{\operatorname{dn}(z_q^k, e)} \right. \\ &\quad \left. + \frac{1}{q^3} \left(\alpha'(k/q) - \frac{8K(e)^2e^2}{3} \left(\operatorname{cn}^2(z_q^k, e) - \operatorname{sn}^2(z_q^k, e) \right) \right) \right] + O(q^{-4}) \end{aligned}$$

Since

$$4K(e)\operatorname{dn}(y_q^k, e) = 4K(e)\operatorname{dn}(z_q^k, e) + \frac{16K(e)^2e^2\alpha(k/q)}{q^2} \operatorname{sn}(z_q^k, e)\operatorname{cn}(z_q^k, e) + O(q^{-4})$$

we have

$$d^2 = 4K(e)\operatorname{dn}(y_q^k, e) \left(\frac{4K(e)\operatorname{dn}(z_q^k, e)}{q^2} + \frac{16K(e)^2e^2}{q^3} \operatorname{sn}(z_q^k, e)\operatorname{cn}(z_q^k, e) + O(q^{-4}) \right)$$

and

$$d^3 = 4K(e)\operatorname{dn}(y_q^k, e) \left(\frac{4K(e)\operatorname{dn}(z_q^k, e)}{q^3} + O(q^{-4}) \right).$$

Moreover,

$$M_1 = \frac{e^2 \operatorname{sn}(K(e)(1 - 4k/q), e) \operatorname{cn}(K(e)(1 - 4k/q), e)}{\operatorname{dn}^2(K(e)(1 - 4k/q), e)} + O(q^{-2}).$$

By the definition of M_2 , N_2 , (8) and 16.9.1 in [1] we have

$$\begin{aligned} \sqrt{M_2 - N_2} &= \frac{\sqrt{1 - e^2}}{2(\sin^2 \varphi_q^k + b^2 \cos^2 \varphi_q^k)} = \frac{\sqrt{1 - e^2}}{2(\operatorname{cn}^2(K(e)(1 - 4x_q^k), e) + b^2 \operatorname{sn}^2(K(e)(1 - 4x_q^k), e))} \\ &= \frac{\sqrt{1 - e^2}}{2\operatorname{dn}^2(K(e)(1 - 4x_q^k), e)}. \end{aligned}$$

Going back to (27) we factor out

$$\frac{4K(e)\operatorname{dn}(K(e)(1 - 4x_q^k), e)}{q} \sqrt{M_2 - N_2} = \frac{\mu(x_q^k)}{q}$$

By (B.1) in [4] the term that we are looking for equals

$$\begin{aligned} \beta(k/q) &= \alpha'(k/q) - \frac{8K(e)^2 e^2}{3} \left(\operatorname{cn}^2(K(e)(1 - 4k/q), e) - \operatorname{sn}^2(K(e)(1 - 4k/q), e) \right) \\ &\quad + \frac{16K(e)^2 e^4 \operatorname{sn}^2(K(e)(1 - 4k/q), e) \operatorname{cn}^2(K(e)(1 - 4k/q), e)}{\operatorname{dn}^2(K(e)(1 - 4k/q), e)} \\ &\quad + \frac{16K(e)}{1 - e^2} \hat{E}_1(k/q) \operatorname{dn}^5(K(e)(1 - 4k/q), e), \end{aligned}$$

where

$$\begin{aligned} \hat{E}_1 &= \frac{1}{24} \frac{e^4 \operatorname{sn}^2(z_q^k, e) \operatorname{cn}^2(z_q^k, e)}{[\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e)]^2} \\ &\quad - \frac{1}{24} \frac{\operatorname{sn}^2(z_q^k, e) + b^2 \operatorname{cn}^2(z_q^k, e)}{\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e)} \\ &\quad - \left[e^4 \frac{\operatorname{sn}^2(z_q^k, e) \operatorname{cn}^2(z_q^k, e)}{(\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e))^2} - \frac{1}{3} \right] \\ &\quad \cdot \left[\frac{e^4 \operatorname{sn}^2(z_q^k, e) \operatorname{cn}^2(z_q^k, e)}{(\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e))^2} - \frac{\operatorname{sn}^2(z_q^k, e) + b^2 \operatorname{cn}^2(z_q^k, e)}{4(\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e))} + \frac{1}{3} \right] \\ &\quad - \left[\frac{\operatorname{sn}^2(z_q^k, e) + b^2 \operatorname{cn}^2(z_q^k, e)}{4(\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e))} - \frac{1}{3} \right]^2 \\ &\quad + e^4 \frac{\operatorname{sn}^2(z_q^k, e) \operatorname{cn}^2(z_q^k, e)}{\operatorname{cn}^2(z_q^k, e) + b^2 \operatorname{sn}^2(z_q^k, e)} \left[\frac{\operatorname{sn}^2(z_q^k, e) + b^2 \operatorname{cn}^2(z_q^k, e)}{4(\operatorname{sn}^2(z_q^k, e) \operatorname{cn}^2(z_q^k, e))} - \frac{1}{3} \right]. \end{aligned}$$

5. ACKNOWLEDGMENTS

The author thanks Vadim Kaloshin, Corentin Fierobe and Jacopo De Simoi for useful discussions.

REFERENCES

- [1] Milton Abramowitz and Irene A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume No. 55 of *National Bureau of Standards Applied Mathematics Series*. U. S. Government Printing Office, Washington, DC, 1964. For sale by the Superintendent of Documents.
- [2] Shau-Jin Chang and Richard Friedberg. Elliptical billiards and Poncelet's theorem. *J. Math. Phys.*, 29(7):1537–1550, 1988.
- [3] I. P. Cornfeld, S. V. Fomin, and Ya. G. Sinai. *Ergodic theory*, volume 245 of *Grundlehren der mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1982. Translated from the Russian by A. B. Sosinskiĭ.
- [4] Jacopo de Simoi, Vadim Kaloshin, and Qiaoling Wei. Dynamical spectral rigidity among \mathbb{Z}_2 -symmetric strictly convex domains close to a circle. *Ann. of Math. (2)*, 186(1):277–314, 2017. Appendix B coauthored with H. Hezari.

KLAUDIUSZ CZUDEK, INSTITUTE OF SCIENCE AND TECHNOLOGY AUSTRIA (ISTA), AM CAMPUS 1,
3400 KLOSTERNEUBURG

INSTITUTE OF APPLIED MATHEMATICS, FACULTY OF PHYSICS AND APPLIED MATHEMATICS, GDAŃSK
UNIVERSITY OF TECHNOLOGY, UL. GABRIELA NARUTOWICZA 11/12, 80-223 GDAŃSK, POLAND

Email address: klaudiusz.czudek@gmail.com