

ALPs coupled to gluons in the GeV mass range - data-driven and consistent

Maksym Ovchynnikov^{1,*} and Andrii Zaporozhchenko^{2,†}¹*Theoretical Physics Department, CERN, 1211 Geneva 23, Switzerland*²*Taras Shevchenko National University of Kyiv, 64 Volodymyrs'ka str., Kyiv 01601, Ukraine*

(Dated: March 10, 2025)

We present a new framework for describing GeV-scale axion-like particles (ALPs), discussing in-depth the case of purely gluonic coupling. Unlike previous approaches, our treatment simultaneously (i) enforces invariance under unphysical chiral rotations converting the ALP-gluon coupling to the quark sector, and (ii) incorporates heavier meson resonances relevant at the GeV scale. In particular, for the first time, we discuss the mixing of ALPs with pseudoscalar and axial-vector excitations and describe the ALP production in the quark fragmentation, which is one of the main ALP production mechanisms. We argue, however, that the lack of maturity of the light meson spectroscopy unavoidably leads to significant uncertainties in the phenomenology of ALP coupled to quarks and gluons in the GeV mass range. This is essential to take into account when studying the potential of future lifetime frontier experiments to search for ALPs.

QCD axion is a hypothetical pseudoscalar particle introduced to solve the strong CP problem [1–3]. The Lagrangian describing its interactions is

$$\mathcal{L}_a = c_G \frac{\alpha_s}{4\pi} \frac{a}{f_a} G^{\mu\nu} \tilde{G}_{\mu\nu}, \quad (1)$$

where a is the axion, c_G/f_a is its interaction constant, α_s is the QCD running coupling, $G_{\mu\nu}$ is the gluon field strength, $\tilde{G}^{\mu\nu}$ is its dual.

The a particle emerges as a Goldstone boson, and its mass m_a gets dynamically generated during the QCD crossover. Therefore, m_a and f_a are intimately related, which leads to severe limitations on the model. The obvious generalization is to assume their independence; corresponding particles are called Axion-Like Particles, or ALPs. ALPs naturally appear in various extensions of the Standard Model [4–7] and may have relation to various cosmological phenomena, being, for example, a portal to light dark matter [8–11]. Generic ALPs with arbitrary coupling pattern, including fermionic couplings, and mass in the GeV range are the subject of extensive studies [12–24]. In particular, the gluonic ALP model (1) has been considered by the PBC initiative as one of the benchmark models to compare the reach of various experiments [25, 26]. Therefore, they are often considered for searches at existing [23, 27] and future lifetime frontier experiments [15, 25, 26, 28], such as the recently approved SHiP [29–31]. At the latter, ALPs may be copiously produced in very intense proton-target collisions and then carefully reconstruct their decays in a displaced decay volume, potentially differentiating between the a particles with different coupling patterns [32].

Thus, it is essential to understand the phenomenology of the GeV-scale ALPs in these experiments. Existing studies, however, provide ambiguous phenomenology descriptions. In particular, some works [12, 16–18, 28] restrict the low-energy ALP interactions by the sector of

pseudoscalar mesons only. It does not allow the description of the ALP interactions in the mass range $m_a \gtrsim m_\sigma$, where various excitations may heavily change the results. On the other hand, Ref. [13] has extended the mesonic interactions by including some of these resonances. However, the resulting description suffers from the dependence of the results on unphysical chiral rotation, which converts the gluon coupling to the quark sector. The corresponding rotation-dependent description of the ALP production and decay modes have been subsequently implemented in various event generators [21, 33, 34], which are widely used by the community to calculate exclusion regions of different past and future experiments [26].

In this Letter, we, for the first time, provide the chiral rotation-independent treatment of the interactions of the GeV-scale ALPs relevant to the Lifetime Frontier experiments. Our main results can be summarized in three points. First, we include the mixing of ALPs with axial-vector mesons and heavier pseudoscalar excitations $\pi^0(1300)$, $\eta(1295)$, $\eta(1440)$, which may significantly enhance the production or decays of ALPs with masses above 1 GeV. Second, we provide self-consistent treatment of the ALP production in the quark fragmentation chain, which appears to be one of the dominant modes. Finally, our last point is that the state-of-the-art “data-driven” descriptions of effective meson interactions, necessary to derive the ALP phenomenology, have ambiguities and incompleteness, preventing us from having a fully robust and comprehensive model of ALP-meson interactions. All these points are not limited to the gluonic ALP (1) – they are equally applicable to the ALPs with arbitrary hadronic coupling patterns.

The Supplemental material discusses these points in-depth, while the main text highlights them.

Phenomenology of ALPs. If the ALP mass is in the GeV range, the description of the hadronic interactions in terms of quarks and gluons breaks down; instead, one needs to know how the ALPs couple to the operators with various bound states, such as mesons and baryons. The generic algorithm to study these interactions is the following [13, 16]. The first step is to make a matching be-

* maksym.ovchynnikov@cern.ch

† andriizaporozhchenko@knu.ua




FIG. 1. How the invariance of a generic observable with the ALPs a on the unphysical chiral rotation parametrized by the matrix κ_q is reached. In the matrix element of an arbitrary process, the direct κ_q -dependent piece of the Lagrangian (the left diagram) cancels the κ_q -dependent part coming from the mixing between a and pseudoscalar mesons $P^0 = \pi^0, \eta, \eta'$ parametrized by the mixing angle $\theta_{P^0 a}$ (the right diagram). The same cancellation holds for the mixing of the ALPs with heavy pseudoscalar and axial-vector mesons.

tween the ALP interactions with quarks and gluons and a modified Chiral Perturbation Theory (ChPT) describing the interaction of ALPs with the lightest pseudoscalar mesons $P = \pi^0, \eta, \eta', \dots$, forming an $U(3)$ nonet. To do this, one may perform the chiral rotation of the quark field:

$$q \rightarrow \exp \left[-i\gamma_5 c_G \kappa_q \frac{a}{f_a} \right] q, \quad (2)$$

where κ_q is a matrix satisfying the condition $\text{Tr}[\kappa_q] = 1$, chosen to be diagonal. This rotation converts the gluon coupling into the derivative coupling to the quark axial-vector current (and also modifies the quark mass term), which can be translated to ChPT.

The resulting ChPT+ALP Lagrangian has quadratic non-diagonal terms between a and neutral flavorless mesons $P^0 = \pi^0, \eta, \eta'$, i.e., there is a $P^0 - a$ mixing. It can be diagonalized by the linear transformation

$$P^0 \approx P_{\text{mass}}^0 + \theta_{P^0 a}(m_a)a + \dots, \quad (3)$$

where the modulus of the *mixing angle* $|\theta_{P^0 a}(m_a)| \ll 1$ everywhere except for the domain $m_a \simeq m_{P^0}$, where it gets resonantly enhanced.

To have consistency, the crucial next step is to perform backward matching – ensure that the hadronic interactions at the level of mesons smoothly repeat those described by the gluon Lagrangian (1). E.g., the hadronic decay width of ALPs calculated exclusively using the interactions with mesons must saturate the perturbative QCD prediction obtained by calculating the width into two gluons.

The pure ChPT is insufficient for making such a matching, because for $m_a \gtrsim m_{P^0}$ various other mesons (such as vector, scalar, and tensor resonances) may potentially contribute to the observables (see, e.g., [35]). Therefore, one has to include the interactions of a with these particles as well. An essential property of the resulting description is that it *must* predict the observables that are independent on the chiral rotation κ_q (modulo the unambiguous contribution $\kappa_u + \kappa_d + \kappa_s = 1$). This is realized, at the level of the matrix element of a particular observable, by summing the κ_q -dependent contributions from

ALP-meson operators coming directly from the chiral rotation and those appearing because of the diagonalization of the ALP-meson quadratic Lagrangian (see Fig. 1).

To the best of our knowledge, there are no studies that provide the approach fulfilling both of these requirements. On the one hand, Refs. [17, 18, 28] formulated the ALP interactions in a κ_q -invariant way. However, they only considered pure ChPT and the anomalous term coupling ALPs to photons. On the other hand, Ref. [13] added the phenomenological interactions with vector, scalar, and tensor mesons, and performed the matching of the ALP decay modes (see also Refs. [22, 36], which the ALPs coupled to quarks). However, the added interactions produced κ_q -dependent results: no direct κ_q terms were added, which means no cancellation. In addition, they miss the interaction of ALPs with axial-vector (such as $a_1, f_1(1285)$, etc.) and heavy pseudoscalar ($\pi^0(1300)$, etc.) excitations, which have mixing with ALPs and therefore are essential for describing the interactions of heavy ALPs with mass $m_a > 1$ GeV.

Our approach. The method we develop combines the self-consistent, κ_q -independent approach with the data-driven description of interactions of various mesons. We include the light pseudoscalar and scalar mesons, vector, and tensor mesons, similarly to how it is done in Refs. [13, 36], and then extend the meson sector by adding the axial-vector and heavy pseudoscalar mesons. The latter terms are essential for describing the ALP interactions, as they induce additional ALP-meson mixings.

Here and below, we consider the three-flavor setup (i.e., keeping η, η') and calculate all the quantities in the order $\mathcal{O}(\delta)$, where $\delta \equiv (m_d - m_u)/(m_d + m_u)$ is the isospin parameter, and $\mathcal{O}(\epsilon)$, where $\epsilon \equiv f_\pi/f_a \ll 1$ is the ALP dimensionless coupling, with $f_\pi \approx 93$ MeV being the pion decay constant. Finally, we assume that the Lagrangian (1) is defined at some scale $\Lambda > \Lambda_{\text{electroweak}}$.¹ Its effective analog at low scales $Q = 2$ GeV $\ll \Lambda$, relevant for the ALP interactions, would also contain additional couplings to quarks $c_q(Q, \Lambda)\partial^\mu a/f_a \bar{q}\gamma_\mu \gamma_5 q$ generated by the renormalization group flow.² We obtain them with the help of the approach of Refs. [17, 22].

To find the mixing between the ALPs and mesons, we first diagonalize the quadratic part of the ALP-mesonic interactions. This way, we find the mixing angles (Eq. (3)) $\theta_{P^0 a}$, $\theta_{A^0 a}$, and $\theta_{P_h^0 a}$, where P_h^0, A^0 are the corresponding neutral heavy pseudoscalar and axial-vector mesons. We utilize the description of their interactions from Ref. [37], which assumes $P_h^0 = \pi^0(1300), \eta(1295), \eta(1440)$ and $A^0 = a_1^0, f_1(1285), f_1(1415)$. Full expressions for the mixing angles, as well as the discussion of the mixing with axial-vector mesons, may be found in Supplementary Material,

¹ Throughout the text, we will consider the common choice $\Lambda = 1$ TeV [25, 26].

² Yet, the induced couplings are suppressed, at the level $\mathcal{O}(10^{-2})$, and weakly sensitive to the value of Λ .

while here, we only provide the κ -dependent part:

$$\theta_{\pi^0 a} \ni c_G \epsilon(\kappa_u - \kappa_d), \theta_{\pi^0(1300)a} \ni -c_G \epsilon(\kappa_u - \kappa_d) \quad (4)$$

$$\theta_{\eta^0 a} \ni -2\sqrt{\frac{2}{3}} c_G \epsilon(\kappa_u + \kappa_d), \theta_{\eta^0(1295)a} \ni -c_G \epsilon(\kappa_u + \kappa_d), \quad (5)$$

$$\theta_{\eta' a} \ni \frac{1}{\sqrt{3}} c_G \epsilon(\kappa_u + \kappa_d), \theta_{\eta(1440)a} \ni \sqrt{2} c_G \epsilon(\kappa_u + \kappa_d), \quad (6)$$

where the rest terms are κ_q -independent summands; in these expressions, we have utilized the relation $\text{Tr}[\kappa_q] = 1$, expressing $\kappa_s = 1 - \kappa_u - \kappa_d$. The ALP- P_h^0 mixing is very sensitive to the presence of some operators dropped in Ref. [37]. We will come back to this point later. Also, while the mixing with pseudoscalars exhibits resonant enhancement around their masses, this is not the case for the axial-vector mesons; for this reason, we drop them from the discussion.

The linear transformation (3) applied to the whole Lagrangian eliminates the non-diagonal terms and induces additional ALP- P interactions. We have ensured that the resulting pure ChPT+ALP theory, receiving contributions from both the mixing angles and the direct κ_q -dependent terms, is invariant on κ_q . Namely, when computing the matrix element of processes with ALPs, the κ_q dependence either vanishes identically after inserting Eqs. (4)-(6), or disappears after utilizing the conservation of 4-momentum at the level of individual vertices (if the same single vertex receives contributions from various operators).

The next step is to add the interactions with various excitations. The Standard Model analogs – Lagrangians of the interaction of these particles with pseudoscalar mesons – are discussed in, correspondingly, Refs. [38, 39] (vector mesons and anomalous photon vertex), [40] (light scalar sector), [39] (tensor mesons), and [37, 41, 42] (vector, axial-vector, and heavy pseudoscalar excitations). On top of that, we will also need the $s \rightarrow d$ operator induced by the octet operator from [19] to describe the decays of kaons into ALPs.

These interactions are given either in terms of the manifestly $U(3)$ covariant objects $\Sigma = \exp[2i\mathcal{P}/f_\pi]$ or in terms of just \mathcal{P} , where $\mathcal{P} = \sum_P t_P P$ is the matrix of the pseudoscalar mesons nonet. To ensure the κ_q -invariant results, we replace the Σ matrix with its transformed version

$$\Sigma \rightarrow \exp\left[-ic_G \kappa_q \frac{a}{f_a}\right] \cdot \Sigma \cdot \exp\left[-ic_G \kappa_q \frac{a}{f_a}\right], \quad (7)$$

and then expand the resulting Lagrangian in the powers of \mathcal{P} and a .³ As for the interactions written in terms of

\mathcal{P} , let us utilize the Baker-Campbell-Hausdorff formula:

$$\begin{aligned} & \exp\left[-ic_G \kappa_q \frac{a}{f_a}\right] \cdot \Sigma \cdot \exp\left[-ic_G \kappa_q \frac{a}{f_a}\right] \\ & \rightarrow \exp\left[i\frac{2}{f_\pi} (\mathcal{P}(x) - \epsilon c_G \kappa_q a) + \mathcal{O}(\epsilon^2)\right] \end{aligned} \quad (8)$$

To maintain the κ_q invariance, we have to replace (note the sign change)

$$\mathcal{P} \rightarrow \mathcal{P} + \epsilon c_G \kappa_q a \quad (9)$$

The resulting Lagrangian, using the modified \mathcal{P}, Σ matrices and the diagonalization (3), provides κ_q -invariant description of all ALP interactions in the $\mathcal{O}(\epsilon)$ limit. It is very complex though, including thousands of terms. In addition, we have to keep as many parameters as possible in the symbolic form rather than inserting their values. Given the computational complexity and to make our results usable and easily modifiable, we have implemented the whole machinery in the improved version of a Mathematica notebook studying the ALP phenomenology from Ref. [22].⁴

At this point, it is important to note that using the “rotated” matrices (7), (9) not only ensures the κ_q independence of the observables but also induces the additional terms with the unambiguous part $\text{Tr}[\kappa_q] = 1$; therefore, it may lead to the cancellation or enhancement of the κ_q -independent piece. We will return to this point later.

ALP production modes. The widely adopted approach from [21] describes the ALP production in deep inelastic scatterings and decays of light mesons by the product of the fluxes π^0, η, η' mesons times the corresponding squared mixing angles $|\theta_{P^0 a}|^2$. It has been consequently incorporated in the event generators with LLPs [26, 33, 34]. Such a description is not only κ_q -dependent – it also wrongly describes how the flux of ALPs depends on the ALP mass. In addition, it does not account for the theoretical uncertainty coming from different production modes.⁵

Within our approach, we split the production into components, considering 2- and 3-body decays of light mesons ($\eta, \eta', K_S, K^+, \rho^0, \omega$), the proton bremsstrahlung, the Drell-Yan process, quark fragmentation, and also decays of B mesons. Decays of ρ^0 and ω within the 3-flavor case are considered for the first time. For the description of the bremsstrahlung, we follow the quasi-real approximation [45–47]. We adopt the ALP-nucleon form factor, needed for accounting for the non-point-like structure of the proton, from [28]. We also incorporate the intrinsic theoretical uncertainty of the quasi-real approximation

³ A similar procedure has been adopted for the $s \rightarrow d$ and anomalous interactions with photons in [18, 28], and for the Wess-Zumino-Witten terms in [24, 28].

⁴ Available on [GitHub](https://github.com/maksymovchynnikov/ALPs-phenomenology) [10.5281/zenodo.14616404](https://doi.org/10.5281/zenodo.14616404).

⁵ Ref. [28] has partially disentangled different production mechanisms, leaving, however, unexplored theoretical uncertainties and some important production modes.




FIG. 2. Phenomenology of ALPs coupled to gluons. Here and below, we assume $c_G = 1$ in Eq. (1). *Top panel*: probabilities of various processes producing ALPs (the yield of ALPs per proton-on-target) in the collisions of a 400 GeV proton beam colliding with the Molybdenum target, corresponding to the SHiP experiment [43]. The bands denote theoretical uncertainties, which we derive following Refs. [22, 44]. *Bottom panel*: the total hadronic decay width. To illustrate the impact of our study, we show two results. The first one (the solid blue line) is obtained using the κ_q -independent approach of our work. The second one (the blue dashed line) is obtained by introducing interactions with various mesons in a manifestly κ_q -variant way (i.e., do not utilize the transformations (7), (9)), and using κ_q given by Eq. (10). The dashed red line shows the hadronic decay width obtained with the help of the perturbative QCD calculations. The vertical gray bands denote the vicinities of π^0, η, η' masses where the description of the ALP phenomenology based on the mixing with mesons breaks down.

– the hard scale defining the allowed ranges of the proton virtuality, see Refs. [47, 48]. For the Drell-Yan process, we follow Ref. [22]; in particular, we accommodate the theoretical uncertainties by varying factorization and renormalization scales of the hard process $GG \rightarrow a$.

Qualitatively, the production in the fragmentation chain occurs when, because of the mixing, each P^0 appearing at the end of the fragmentation is “replaced” by the ALP with a tiny probability. This process is attractive because of a relatively small uncertainty compared

to the proton bremsstrahlung. We incorporate [49] the production in the quark fragmentation via the mixing with light P^0 s in PYTHIA8 [50]. Including the contribution from the mixing with heavier resonances is left for future work; therefore, our results on this mode are conservative.

The probabilities of all these processes for the beam and target setup corresponding to the SHiP experiment are shown in Fig. 2 (the top panel). We note a large uncertainty in the proton bremsstrahlung (a common feature for the other LLPs, see Ref. [44]) and the Drell-Yan process, which may reach 1-2 orders in magnitude. The production in the fragmentation chain is one of the dominant modes.

ALP decay modes. To calculate the ALP decay width, we use the combined Lagrangian describing the interaction of ALPs with various mesons and consider exclusive decay modes, including $a \rightarrow 2\gamma, 3\pi, 2\pi\gamma, KK\pi, \eta^{(\prime)}\pi\pi, 4\pi/2\rho^0, 2\omega$. Our results apply to any interaction pattern of ALPs. They generalize and extend the past studies [13, 17, 22, 36], providing the κ_q -invariant description of the decay modes and incorporating new interactions with axial-vector and pseudoscalar excitations. In addition, we improve the treatment of the sector of light scalar mesons, re-analyzing the results of Refs. [40, 51] by using the definition of η, η' mesons in terms of isoscalar and singlet states η_0, η_8 that is consistent with the one adopted in the rest interaction sector.

The total width is shown in Fig. 2 (the bottom panel). To stress the importance of κ_q invariance, we show two curves: the one obtained using our approach, and another one obtained as if adding the scalar, vector, and tensor interactions without accounting for the replacements (7), (9). Then, we fix the κ_q dependence using the common choice

$$\kappa_q = \text{diag}(m_u^{-1}, m_d^{-1}, m_s^{-1}) / \sum_q m_q^{-1} \quad (10)$$

The widths heavily differ on the whole mass range; the difference may reach a few orders of magnitude.

In particular, in the domain $m_a \gtrsim 1$ GeV, the κ_q -independent width is smaller, which prevents making a naive matching with the perturbative width into gluons. This smallness is caused by the mutual cancellation between the contributions coming from the diagonalization (3) and the rotation (7) in some cases, which reflects the genuine property of the gluonic interaction (1). In particular, it leads to suppression of the contribution of the $P_h^0 a$ -mixing (such suppression is not present for the ALPs having large coupling to quarks).

This brings us to the following important message: the status of the ALP phenomenology is intimately linked to the robustness of the effective description of Standard Model interactions of mesons. Unfortunately, there is no unique description accommodating all the resonances simultaneously, especially in the domain of meson masses $M \gtrsim 1$ GeV; this is partially caused by the lack of robust experimental data for these resonances [42], mak-

ing it complicated even to conclude whether the observed mesons are pure 2-quark states, or also include an admixture of a 4-quark state. The interaction pattern of included mesons is also ambiguous. In particular, the study P_h interactions [37], motivated by simplicity, dropped some operators that may significantly enhance the production and decay rates of the ALPs (we discuss this point in detail in Supplementary Material). On the other hand, accommodating these terms is not straightforward, requiring re-fitting the resulting interactions to match the experimental data.

An important point is that, thanks to the implementation of the ALP phenomenology in the `Mathematica` notebook, our approach is very flexible and can be easily generalized by adding additional ALP interactions.

Conclusions. To summarize, we have systematically explored the phenomenology of the ALPs in the GeV mass range, considering as an example the ALPs coupled to gluons. Our study has two main outcomes. First, our

approach is simultaneously free from unphysical chiral rotation and includes interactions with various meson resonances. In particular, we have incorporated previously missing ALP production modes, as well as their mixing with axial-vector and heavy pseudoscalar excitations. Second, we have highlighted significant unavoidable uncertainties in the ALP phenomenology, linking it to the (not mature enough) state of the effective description of interactions of the mesons with masses $M < 2$ GeV. Our analysis, incorporating missing interactions and production modes, may serve as the new state-of-the-art study of the ALP phenomenology, with the possibility of easily incorporating new interactions once the meson spectroscopy improves. The revised ALP phenomenology will be implemented in the event rate sampler `SensCalc` [34], allowing the community to recast the parameter space of the ALPs probed by past and future experiments.

Acknowledgements. The authors thank Francesco Giacosa for useful discussions on the extended linear sigma model.

-
- [1] R. D. Peccei and H. R. Quinn, “CP Conservation in the Presence of Instantons,” *Phys. Rev. Lett.* **38** (1977) 1440–1443.
- [2] S. Weinberg, “A New Light Boson?,” *Phys. Rev. Lett.* **40** (1978) 223–226.
- [3] F. Wilczek, “Problem of Strong P and T Invariance in the Presence of Instantons,” *Phys. Rev. Lett.* **40** (1978) 279–282.
- [4] R. Essig *et al.*, “Working Group Report: New Light Weakly Coupled Particles,” in *Snowmass 2013: Snowmass on the Mississippi*. 10, 2013. [arXiv:1311.0029 \[hep-ph\]](#).
- [5] D. J. E. Marsh, “Axion Cosmology,” *Phys. Rept.* **643** (2016) 1–79, [arXiv:1510.07633 \[astro-ph.CO\]](#).
- [6] P. W. Graham, I. G. Irastorza, S. K. Lamoreaux, A. Lindner, and K. A. van Bibber, “Experimental Searches for the Axion and Axion-Like Particles,” *Ann. Rev. Nucl. Part. Sci.* **65** (2015) 485–514, [arXiv:1602.00039 \[hep-ex\]](#).
- [7] I. G. Irastorza and J. Redondo, “New experimental approaches in the search for axion-like particles,” *Prog. Part. Nucl. Phys.* **102** (2018) 89–159, [arXiv:1801.08127 \[hep-ph\]](#).
- [8] Y. Nomura and J. Thaler, “Dark Matter through the Axion Portal,” *Phys. Rev. D* **79** (2009) 075008, [arXiv:0810.5397 \[hep-ph\]](#).
- [9] M. J. Dolan, F. Kahlhoefer, C. McCabe, and K. Schmidt-Hoberg, “A taste of dark matter: Flavour constraints on pseudoscalar mediators,” *JHEP* **03** (2015) 171, [arXiv:1412.5174 \[hep-ph\]](#). [Erratum: *JHEP* **07**, 103 (2015)].
- [10] Y. Hochberg, E. Kuflik, R. McGehee, H. Murayama, and K. Schutz, “Strongly interacting massive particles through the axion portal,” *Phys. Rev. D* **98** (2018) no. 11, 115031, [arXiv:1806.10139 \[hep-ph\]](#).
- [11] P. J. Fitzpatrick, Y. Hochberg, E. Kuflik, R. Ovadia, and Y. Soreq, “Dark matter through the axion-gluon portal,” *Phys. Rev. D* **108** (2023) no. 7, 075003, [arXiv:2306.03128 \[hep-ph\]](#).
- [12] M. Bauer, M. Neubert, and A. Thamm, “Collider Probes of Axion-Like Particles,” *JHEP* **12** (2017) 044, [arXiv:1708.00443 \[hep-ph\]](#).
- [13] D. Aloni, Y. Soreq, and M. Williams, “Coupling QCD-Scale Axionlike Particles to Gluons,” *Phys. Rev. Lett.* **123** (2019) no. 3, 031803, [arXiv:1811.03474 \[hep-ph\]](#).
- [14] C. Cornella, P. Paradisi, and O. Sumensari, “Hunting for ALPs with Lepton Flavor Violation,” *JHEP* **01** (2020) 158, [arXiv:1911.06279 \[hep-ph\]](#).
- [15] G. Aielli *et al.*, “Expression of interest for the CODEX-b detector,” *Eur. Phys. J. C* **80** (2020) no. 12, 1177, [arXiv:1911.00481 \[hep-ex\]](#).
- [16] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, “The Low-Energy Effective Theory of Axions and ALPs,” *JHEP* **04** (2021) 063, [arXiv:2012.12272 \[hep-ph\]](#).
- [17] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, “Flavor probes of axion-like particles,” *JHEP* **09** (2022) 056, [arXiv:2110.10698 \[hep-ph\]](#).
- [18] M. Bauer, M. Neubert, S. Renner, M. Schnubel, and A. Thamm, “Consistent Treatment of Axions in the Weak Chiral Lagrangian,” *Phys. Rev. Lett.* **127** (2021) no. 8, 081803, [arXiv:2102.13112 \[hep-ph\]](#).
- [19] S. Gori, G. Perez, and K. Tobioka, “KOTO vs. NA62 Dark Scalar Searches,” *JHEP* **08** (2020) 110, [arXiv:2005.05170 \[hep-ph\]](#).
- [20] S. Chakraborty, M. Kraus, V. Loladze, T. Okui, and K. Tobioka, “Heavy QCD axion in $b \rightarrow s$ transition: Enhanced limits and projections,” *Phys. Rev. D* **104** (2021) no. 5, 055036, [arXiv:2102.04474 \[hep-ph\]](#).
- [21] J. Jerhot, B. Döbrich, F. Ertas, F. Kahlhoefer, and T. Spadaro, “ALPINIST: Axion-Like Particles In Numerous Interactions Simulated and Tabulated,” *JHEP* **07** (2022) 094, [arXiv:2201.05170 \[hep-ph\]](#).
- [22] G. Dalla Valle Garcia, F. Kahlhoefer, M. Ovchinnikov, and A. Zaporozhchenko, “Phenomenology of axionlike

- particles with universal fermion couplings revisited,” *Phys. Rev. D* **109** (2024) no. 5, 055042, [arXiv:2310.03524 \[hep-ph\]](#).
- [23] Y. Afik, B. Döbrich, J. Jerhot, Y. Soreq, and K. Tobioka, “Probing long-lived axions at the KOTO experiment,” *Phys. Rev. D* **108** (2023) no. 5, 055007, [arXiv:2303.01521 \[hep-ph\]](#).
- [24] Y. Bai, T.-K. Chen, J. Liu, and X. Ma, “Wess-Zumino-Witten Interactions of Axions,” [arXiv:2406.11948 \[hep-ph\]](#).
- [25] J. Beacham *et al.*, “Physics Beyond Colliders at CERN: Beyond the Standard Model Working Group Report,” *J. Phys. G* **47** (2020) no. 1, 010501, [arXiv:1901.09966 \[hep-ex\]](#).
- [26] C. Antel *et al.*, “Feebly Interacting Particles: FIPs 2022 workshop report,” [arXiv:2305.01715 \[hep-ph\]](#).
- [27] NA62 Collaboration, E. Cortina Gil *et al.*, “Search for dark photon decays to $\mu^+\mu^-$ at NA62,” *JHEP* **09** (2023) 035, [arXiv:2303.08666 \[hep-ex\]](#).
- [28] N. Blinov, E. Kowalczyk, and M. Wynne, “Axion-like particle searches at DarkQuest,” *JHEP* **02** (2022) 036, [arXiv:2112.09814 \[hep-ph\]](#).
- [29] S. Alekhin *et al.*, “A facility to Search for Hidden Particles at the CERN SPS: the SHiP physics case,” *Rept. Prog. Phys.* **79** (2016) no. 12, 124201, [arXiv:1504.04855 \[hep-ph\]](#).
- [30] SHiP Collaboration, R. e. a. Albanese, “BDF/SHiP at the ECN3 high-intensity beam facility,” tech. rep., CERN, Geneva, 2023. <https://cds.cern.ch/record/2878604>.
- [31] C. Ahdida *et al.*, “Post-LS3 Experimental Options in ECN3,” [arXiv:2310.17726 \[hep-ex\]](#).
- [32] O. Mikulenko, K. Bondarenko, A. Boyarsky, and O. Ruchayskiy, “New physics at the Intensity Frontier: how much can we learn and how?,” [arXiv:2312.00659 \[hep-ph\]](#).
- [33] F. Kling and S. Trojanowski, “Forward experiment sensitivity estimator for the LHC and future hadron colliders,” *Phys. Rev. D* **104** (2021) no. 3, 035012, [arXiv:2105.07077 \[hep-ph\]](#).
- [34] M. Ovchinnikov, J.-L. Tastet, O. Mikulenko, and K. Bondarenko, “Sensitivities to feebly interacting particles: public and unified calculations,” [arXiv:2305.13383 \[hep-ph\]](#).
- [35] P. Ilten, Y. Soreq, M. Williams, and W. Xue, “Serendipity in dark photon searches,” *JHEP* **06** (2018) 004, [arXiv:1801.04847 \[hep-ph\]](#).
- [36] H.-C. Cheng, L. Li, and E. Salvioni, “A theory of dark pions,” *JHEP* **01** (2022) 122, [arXiv:2110.10691 \[hep-ph\]](#).
- [37] D. Parganlija and F. Giacosa, “Excited Scalar and Pseudoscalar Mesons in the Extended Linear Sigma Model,” *Eur. Phys. J. C* **77** (2017) no. 7, 450, [arXiv:1612.09218 \[hep-ph\]](#).
- [38] T. Fujiwara, T. Kugo, H. Terao, S. Uehara, and K. Yamawaki, “Nonabelian Anomaly and Vector Mesons as Dynamical Gauge Bosons of Hidden Local Symmetries,” *Prog. Theor. Phys.* **73** (1985) 926.
- [39] F.-K. Guo, B. Kubis, and A. Wirzba, “Anomalous decays of η' and η into four pions,” *Phys. Rev. D* **85** (2012) 014014, [arXiv:1111.5949 \[hep-ph\]](#).
- [40] A. H. Fariborz and J. Schechter, “ η' —> η pi pi decay as a probe of a possible lowest lying scalar nonet,” *Phys. Rev. D* **60** (1999) 034002, [arXiv:hep-ph/9902238](#).
- [41] D. Parganlija, P. Kovacs, G. Wolf, F. Giacosa, and D. H. Rischke, “Meson vacuum phenomenology in a three-flavor linear sigma model with (axial-)vector mesons,” *Phys. Rev. D* **87** (2013) no. 1, 014011, [arXiv:1208.0585 \[hep-ph\]](#).
- [42] F. Giacosa, P. Kovács, and S. Jafarzade, “Ordinary and exotic mesons in the extended Linear Sigma Model,” [arXiv:2407.18348 \[hep-ph\]](#).
- [43] SHiP Collaboration, O. Aberle *et al.*, “BDF/SHiP at the ECN3 high-intensity beam facility,” tech. rep., CERN, Geneva, 2022. <http://cds.cern.ch/record/2839677>.
- [44] Y. Kyselov and M. Ovchinnikov, “Searches for long-lived dark photons at proton accelerator experiments,” *Phys. Rev. D* **111** (2025) no. 1, 015030, [arXiv:2409.11096 \[hep-ph\]](#).
- [45] G. Altarelli and G. Parisi, “Asymptotic Freedom in Parton Language,” *Nucl. Phys. B* **126** (1977) 298–318.
- [46] I. Boiarska, K. Bondarenko, A. Boyarsky, V. Gorkavenko, M. Ovchinnikov, and A. Sokolenko, “Phenomenology of GeV-scale scalar portal,” *JHEP* **11** (2019) 162, [arXiv:1904.10447 \[hep-ph\]](#).
- [47] S. Foroughi-Abari and A. Ritz, “Dark sector production via proton bremsstrahlung,” *Phys. Rev. D* **105** (2022) no. 9, 095045, [arXiv:2108.05900 \[hep-ph\]](#).
- [48] S. Foroughi-Abari, P. Reimitz, and A. Ritz, “A Closer Look at Dark Vector Splitting Functions in Proton Bremsstrahlung,” [arXiv:2409.09123 \[hep-ph\]](#).
- [49] Y. Kyselov, S. Mrenna, and M. Ovchinnikov, “New physics mixing with mesons: production in the fragmentation chain,” [arXiv:to appear \[hep-ph\]](#).
- [50] C. Bierlich *et al.*, “A comprehensive guide to the physics and usage of PYTHIA 8.3,” *SciPost Phys. Codeb.* **2022** (2022) 8, [arXiv:2203.11601 \[hep-ph\]](#).
- [51] D. Black, A. H. Fariborz, F. Sannino, and J. Schechter, “Putative light scalar nonet,” *Phys. Rev. D* **59** (1999) 074026, [arXiv:hep-ph/9808415](#).
- [52] Particle Data Group Collaboration, R. L. Workman *et al.*, “Review of Particle Physics,” *PTEP* **2022** (2022) 083C01.
- [53] V. Cirigliano, G. Ecker, H. Neufeld, A. Pich, and J. Portoles, “Kaon Decays in the Standard Model,” *Rev. Mod. Phys.* **84** (2012) 399, [arXiv:1107.6001 \[hep-ph\]](#).
- [54] Particle Data Group Collaboration, R. L. Workman and Others, “Review of Particle Physics,” *PTEP* **2022** (2022) 083C01.

SUPPLEMENTAL MATERIAL

In this Supplemental material, we describe the ALP interactions with various mesons and nucleons following the self-consistent approach presented in the paper. It is organized as follows.

Sec. A is devoted to the addition of ALPs to the Chiral Perturbation Theory (ChPT), describing light pseudoscalar meson nonet, and incorporating interactions with various other mesons. There, Subsec. A1 describes the Chiral Perturbation Theory. In Subsec. A2, we add ALPs, define the procedure of the diagonalization of the quadratic part of the ALP-meson Lagrangian, calculate the mixing angles, and outline our strategy to incorporate other mesons in the integration Lagrangian. In Subsec. A3, we discuss the addition of light scalar, vector, and tensor mesons. Subsec. A4 describes the ALP-nucleon interaction. Subsec. A5 is devoted to the description of the ALP interaction with axial-vector and heavy pseudoscalar mesons. Subsec. A6 is devoted to cross-checks of our approach. Finally, Subsec. A7 highlights the limitations of our approach, relating it to the currently immature status of incorporating the whole meson spectroscopy in the language of effective field theory.

Sec. B uses the results of the previous section to describe different ALP production mechanisms, including decays of light mesons and the production in the quark fragmentation. There, we also outline the problem of the widely adopted description [21], approximating the flux of ALPs by the flux of mesons P^0 times the squared mixing angle $|\theta_{P^0 a}|^2$.

Most of the results of these sections are obtained (unless stated otherwise) using our *Mathematica* notebook where we carefully implement the ALP phenomenology.⁶

Appendix A: Meson spectroscopy and ALP interactions

1. Pure ChPT

The pure ChPT Lagrangian we will use in our studies is

$$\mathcal{L}_{\text{ChPT, min}} = \frac{f_\pi^2}{2} B_0 \text{Tr} [\Sigma \hat{m}_q^\dagger + \hat{m}_q \Sigma^\dagger] + \mathcal{L}_{\text{anomaly}} + \frac{f_\pi^2}{4} \text{Tr} [D_\mu \Sigma D^\mu \Sigma^\dagger] \quad (\text{A1})$$

Here:

- $B_0 = m_{\pi^0}^2 / (m_u + m_d)$ and $f_\pi = 93$ MeV is the pion decay constant.
- Σ is the matrix of the pseudoscalar mesons:

$$\Sigma = \exp \left[\frac{2i\mathcal{P}}{f_\pi} \right], \quad \mathcal{P} = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{3}} + \frac{\eta'}{\sqrt{6}} & \pi^+ & K^+ \\ \pi^- & -\frac{\pi^0}{\sqrt{2}} + \frac{\eta}{\sqrt{3}} + \frac{\eta'}{\sqrt{6}} & K^0 \\ K^- & \bar{K}^0 & -\frac{\eta}{\sqrt{3}} + 2\frac{\eta'}{\sqrt{6}} \end{pmatrix} \quad (\text{A2})$$

while $D_\mu \Sigma = \partial_\mu \Sigma + ieA_\mu \Sigma$ is the covariant derivative. Here, we have fixed the $\eta - \eta'$ mixing angle as $\theta_{\eta\eta'} = \arcsin(-1/3)$, which provides a reasonable agreement with the experimental data while allowing us to provide the results in a simple analytic form [13].

- $\hat{m}_q = \text{diag}(m_u, m_d, m_s)$ is the matrix of quark masses.
- $\mathcal{L}_{\text{anomaly}}$ is the QCD anomaly-breaking term of the $U_A(1)$ symmetry:

$$\mathcal{L}_{\text{anomaly}} = -m_0^2 \eta_0^2 / 2, \quad (\text{A3})$$

with $\eta_0 = \cos(\theta_{\eta\eta'})\eta' - \sin(\theta_{\eta\eta'})\eta$ being its Goldstone. The coefficient m_0 is fixed in a way such that after summing the ChPT mass term (coming from the first summand in Eq. (A1)) and the m_0 term, there is no $\eta - \eta'$ mixing for the given $\theta_{\eta\eta'}$:

$$m_0^2 = \frac{3 m_{\pi^0}^2 (2m_s - m_u - m_d)}{2(m_u + m_d)} \quad (\text{A4})$$

⁶ Available on [GitHub: maksymovchynnikov/ALPs-phenomenology](https://github.com/maksymovchynnikov/ALPs-phenomenology) and

[10.5281/zenodo.14616404](https://zenodo.org/record/14616404).

Calculating the masses of η and η' , π^0 , and using the explicit form the isospin breaking parameter $\delta = (m_d - m_u)/(m_d + m_u)$, one may get the following consistency relations between quark and meson masses which we will use below:

$$m_u = m_d \frac{1 - \delta}{1 + \delta}, \quad m_d = (1 + \delta) \frac{m_{\pi^0}^2}{2m_\eta^2 - m_{\pi^0}^2} m_s, \quad m_{\eta'}^2 = 4m_\eta^2 - 3m_{\pi^0}^2, \quad m_{K^+}^2 = m_\eta^2 - \frac{\delta m_{\pi^0}^2}{2} \quad (\text{A5})$$

Here and below, we will only keep the terms being quartic in fields. For these purposes, for example, in $\text{Tr}[D_\mu \Sigma D^\mu \Sigma]$, we have to expand $\Sigma = \sum_{n=0}^4 \frac{1}{n!} \left(\frac{2i\mathcal{P}}{f_\pi}\right)^n$ and then drop all the terms with the dimensionality higher than 4.

2. Adding ALPs to ChPT and strategy of adding other excitations

Let us start with the interaction Lagrangian of the ALPs in terms of quarks and gluons, assuming the absence of sizeable flavor-changing operators:

$$\mathcal{L}_a = c_G \frac{a}{f_a} \frac{\alpha_s}{4\pi} G_{\mu\nu}^a \tilde{G}^{\mu\nu,a} + \frac{\partial_\mu a}{f_a} \sum_q c_q \bar{q} \gamma^\mu \gamma_5 q \quad (\text{A6})$$

Similarly to the studies [16, 17], we assume that this Lagrangian is defined at some scale $\Lambda \gg \Lambda_{\text{electroweak}}$. The scales of interest are $\Lambda_Q \simeq m_a$, and the Lagrangian (A6) non-trivially evolves down to this scale: the values of the couplings $c_q(\Lambda_Q) \neq c_q(\Lambda)$. In particular, even if $c_q(\Lambda) = 0$ (this is the case we concentrate on), we end up with non-zero $c_q(\Lambda_Q)$; for the choice $\Lambda = 1$ TeV commonly used in the literature, $c_q \simeq 10^{-2}$ [17, 22].

The analog of the Lagrangian (A6) at the scale Λ_Q can be translated to the effective Lagrangian in terms of mesonic degrees of freedom after performing the rotation

$$q \rightarrow \exp \left[-\frac{i c_G a}{f_a} \hat{\kappa}_q \gamma_5 \right] q, \quad \text{Tr}[\kappa_q] = 1 \quad (\text{A7})$$

eliminating the direct coupling to gluons. We parametrize the κ_q matrix by

$$\hat{\kappa}_q = \text{diag}(\kappa_u, \kappa_d, 1 - \kappa_u - \kappa_d) \quad (\text{A8})$$

The corresponding Lagrangian in terms of the mesons is [16]

$$\begin{aligned} \mathcal{L}_{\text{ChPT,min}} = & \frac{1}{2} (\partial_\mu a)^2 - \frac{m_a^2}{2} a^2 + \frac{f_\pi^2}{2} B_0 \text{Tr} [\Sigma \hat{m}_q^\dagger + \hat{m}_q \Sigma^\dagger] + \mathcal{L}_{\text{anomaly}} + \frac{f_\pi^2}{4} \text{Tr} [D_\mu \Sigma D^\mu \Sigma^\dagger] \\ & + \frac{f_\pi^2}{2} \frac{\partial_\mu a}{f} \text{Tr}[(\hat{c}_q + c_G \kappa_q)(\Sigma D^\mu \Sigma^\dagger - \Sigma^\dagger D^\mu \Sigma)] \quad (\text{A9}) \end{aligned}$$

Here:

- $\hat{c}_q = \text{diag}(c_u, c_d, c_s)$ is the direct ALP coupling to quarks. To shorten the expressions, below, we set them to zero, although keeping in all numerical calculations.
- \hat{m}_q becomes modified by the chiral transformation:

$$\hat{m}_q = \exp \left[-i c_G \frac{a}{f} \kappa_q \right] m_q \exp \left[-i c_G \frac{a}{f} \kappa_q \right], \quad (\text{A10})$$

The part of the Lagrangian (A9) quadratic in the ALP field a and flavorless pseudoscalar mesons $P^0 = \eta, \eta', \pi^0$ has the form

$$\mathcal{L}_{P^0 a, \text{quad}} = \frac{1}{2} K_{ij} \partial_\mu X_i \partial^\mu X_j - \frac{1}{2} M_{ij} X_i X_j, \quad (\text{A11})$$

where $X = (a, \pi^0, \eta, \eta')$, and the mass and kinetic matrices are

$$\hat{K} = \begin{pmatrix} 1 & \epsilon c_G (\kappa_u - \kappa_d) & \sqrt{\frac{2}{3}} \epsilon c_G (2\kappa_d + 2\kappa_u - 1) & -\frac{\epsilon c_G (\kappa_d + \kappa_u - 2)}{\sqrt{3}} \\ \epsilon c_G (\kappa_u - \kappa_d) & 1 & 0 & 0 \\ \sqrt{\frac{2}{3}} \epsilon c_G (2\kappa_d + 2\kappa_u - 1) & 0 & 1 & 0 \\ -\frac{\epsilon c_G (\kappa_d + \kappa_u - 2)}{\sqrt{3}} & 0 & 0 & 1 \end{pmatrix}, \quad (\text{A12})$$

$$\hat{M} = \begin{pmatrix} m_a^2 & M_{a\pi^0} & M_{a\eta} & M_{a\eta'} \\ M_{a\pi^0} & m_\pi^2 & -\sqrt{\frac{2}{3}}\delta m_{\pi^0}^2 & -\frac{\delta m_{\pi^0}^2}{\sqrt{3}} \\ M_{a\eta} & -\sqrt{\frac{2}{3}}\delta m_{\pi^0}^2 & m_\eta^2 & 0 \\ M_{a\eta'} & -\frac{\delta m_{\pi^0}^2}{\sqrt{3}} & 0 & m_{\eta'}^2 \end{pmatrix}, \quad (\text{A13})$$

where

$$M_{a\pi^0} = -\epsilon c_G m_{\pi^0}^2 ((\delta + 1)\kappa_d + (\delta - 1)\kappa_u), \quad (\text{A14})$$

$$M_{a\eta} = \sqrt{\frac{2}{3}}\epsilon c_G (m_{\pi^0}^2 (\delta\kappa_d - \delta\kappa_u + 1) + 2m_\eta^2 (\kappa_d + \kappa_u - 1)), \quad (\text{A15})$$

$$M_{a\eta'} = \frac{\epsilon c_G (m_{\pi^0}^2 ((\delta + 3)\kappa_d - (\delta - 3)\kappa_u - 2) - 4m_\eta^2 (\kappa_d + \kappa_u - 1))}{\sqrt{3}} \quad (\text{A16})$$

In Eq. (A13), we used the relation for the u and d quark masses in terms of δ and m_s , see Eq. (A5).

A generic transformation simultaneously diagonalizing these matrices in the $\mathcal{O}(\delta)$ order is [13]

$$a = a_{\text{phys}} - \sum_{P^0=\pi^0,\eta,\eta'} h(P^0, m_{P^0}) P_{\text{phys}}^0, \quad (\text{A17})$$

$$P^0 = P_{\text{phys}}^0 - \sum_{P^{0'} \neq P^0} \frac{M_{P^0 P^{0'}}}{m_{P^0}^2 - m_{P^{0'}}^2} + h(P^0, m_a) a_{\text{phys}}, \quad (\text{A18})$$

where

$$h(P^0, m_X) = \frac{1}{m_a^2 - m_{P^0}^2} \left[M_{aP^0} - m_X^2 K_{aP^0} + \sum_{P^{0'} \neq P^0} M_{P^0 P^{0'}} \frac{M_{aP^{0'}} - m_X^2 K_{aP^{0'}}}{m_X^2 - m_{P^{0'}}^2} \right] \quad (\text{A19})$$

Below, we will drop the index ‘‘phys’’.

Introducing the parameter $\epsilon = f_\pi/f_a$, the mixing angles (the coefficients in the expansion of P^0 in front of a) become

$$\theta_{\pi^0 a} = \frac{\frac{1}{3}\epsilon c_G \delta m_{\pi^0}^2 \left(\frac{(2m_a^2 - m_{\eta'}^2 - m_{\pi^0}^2)}{(m_a^2 - m_{\eta'}^2)} - \frac{2(m_a^2 - 2m_\eta^2 + m_{\pi^0}^2)}{(m_a^2 - m_\eta^2)} \right)}{m_a^2 - m_{\pi^0}^2} + \epsilon c_G (\kappa_d - \kappa_u), \quad (\text{A20})$$

$$\theta_{\eta a} = \frac{\sqrt{\frac{2}{3}}\epsilon c_G (m_a^2 + m_{\pi^0}^2 - 2m_\eta^2)}{m_a^2 - m_\eta^2} - 2\sqrt{\frac{2}{3}}\epsilon c_G (\kappa_d + \kappa_u), \quad (\text{A21})$$

$$\theta_{\eta' a} = -\frac{\epsilon c_G (2m_a^2 - m_{\eta'}^2 - m_{\pi^0}^2)}{\sqrt{3}(m_a^2 - m_{\eta'}^2)} + \frac{\epsilon c_G (\kappa_d + \kappa_u)}{\sqrt{3}} \quad (\text{A22})$$

If utilizing the relations (A5), the quadratic part of the Lagrangian becomes diagonalized up to terms $\mathcal{O}(\delta)$ inclusive. We note the presence of the pole terms from η, η' in the $\pi^0 a$ mixing and, vice versa, $\theta_{\eta a/\eta' a}$ include pole from π^0 . It is caused by the $\pi^0 - \eta/\eta'$ mixing induced by the mass matrix (A13).

Let us now outline our strategy to add other mesons to the ALP interaction. For this, we will utilize the transformations of the Σ and \mathcal{P} matrices under the quark chiral rotation (A7) driven by κ_q :

$$\Sigma \rightarrow \exp \left[-i c_G \kappa_q \frac{a}{f_a} \right] \Sigma \exp \left[-i c_G \kappa_q \frac{a}{f_a} \right], \quad (\text{A23})$$

$$\mathcal{P} \rightarrow \mathcal{P} + \epsilon c_G \kappa_q a \quad (\text{A24})$$

κ_q -invariant interactions of ALPs with the other mesons can be obtained with the help of the Lagrangian written in terms of the objects (A23), (A24), subsequently inserted linear transformation (A18), and keeping only $\mathcal{O}(\epsilon, \delta, \epsilon \cdot \delta)$ terms.

Having such a Lagrangian, we are ready to calculate the ALP decay widths. To do this, we first compute the matrix element of the given process and ensure that it is κ_q -independent. Then, following Ref. [13], we adopt the phenomenological suppression factors $F(m)$ for the whole matrix element, accounting for the QCD sum rules. We consider the limit $\mathcal{O}(\delta)$ everywhere except for the process $a \rightarrow 3\pi$, for which the squared matrix element is already $\propto \delta^2$.

Below, we describe how we add various excitations: lightest scalar, vector, tensor, axial-vector, and heavy pseudoscalar.

Regarding the sector of scalar, vector, and tensor mesons (Sec. A 3 for the detailed description), we mainly follow Appendix A of Ref. [36] (already utilized in our previous study [22]). It improves the treatment of these interactions in Ref. [13] but studies a different ALP model without the coupling to gluons. The improvements include: imposing unitarity constraints restricting the contribution of certain excitations to various kinematic phase space of ALP decays; including proper polarization sum of vector and tensor mesons when treating them as intermediate states; and the coupling of the tensor mesons to the stress-energy tensor in terms of pseudoscalar mesons rather than just the minimal coupling via the metrics. However, there is an important difference between our present analysis and Refs. [22, 36] concerning the sector of scalar mesons in our present analysis; it is discussed in Sec. A 3 a.

The addition of axial-vector and heavy pseudoscalar mesons is the new result that we discuss in Sec. A 5. It is based on the so-called Extended Linear Sigma Model from Refs. [37, 41, 42].

3. Adding scalar, vector, and tensor resonances

a. Scalar mesons

We add the interactions with the lightest scalar mesons following Refs. [40, 51], which introduced $SU(3)$ invariant description in terms of the matrices Σ and the scalar nonet matrix

$$S = \begin{pmatrix} a_0^0 - \sigma \sin(\theta_s) + f_0(980) \cos(\theta_s) & \sqrt{2}a_0^+ & \sqrt{2}\kappa^+ \\ \sqrt{2}a_0^- & -a_0^0 - \sigma \sin(\theta_s) + f_0(980) \cos(\theta_s) & \sqrt{2}\kappa \\ \sqrt{2}\kappa^- & \sqrt{2}\bar{\kappa} & \sqrt{2}\sigma \cos(\theta_s) + \sqrt{2}f_0(980) \sin(\theta_s) \end{pmatrix}, \quad (\text{A25})$$

with $\theta_s = -21^\circ$ being the scalar mixing angle, and κ is also known as $K_0^*(700)$. The interaction coefficients have been fixed to describe the $\pi\pi$ and πK scattering data, and also to reproduce the decay $\eta' \rightarrow \eta\pi\pi$.

These studies defined η, η' in terms of the isoscalar and singlet particles η_8, η_0 in a different way compared to the ALP studies. Namely, the mixing angle in [40] differs from the angle $\theta_{\eta\eta'} = \arcsin(-1/3)$, commonly used when describing the mixing of ALPs with these mesons. Thus, incorporating the interactions from [40, 51] into the ALP phenomenology “out of the box” would lead to inconsistency.

To fix the problem, we start with Eq. (B.4) of [51]. It is written in terms of phenomenological constants A, B, C, D and the matrix of the pseudoscalar nonet Σ (denoted as ξ there). As usual, we replace it with Eq. (A23), and perform the expansion in terms of \mathcal{P} . The parameters A, B, C, D are fitted to the observational data – π, K scattering for A, B , and η' decays for C, D . Due to the different definition of η, η' we use, in our expansion, the η, η' couplings depend differently on C, D than in the expansion (A.1) of Ref. [40], so we cannot just use their reported values – it would destroy the predictions on $\eta' \rightarrow \eta\pi\pi$ decay. Instead, we recompute the couplings of scalar mesons to η, η' in our expansion, and set their values to recover the width of the process $\eta' \rightarrow \eta 2\pi$, consistent with the experimental value [52].

We also utilize these couplings when adding the S -wave amplitude to the ALP decays into $KK\pi$ using the BaBar fit from [36].

b. Vector mesons

The Hidden Local Symmetry approach of including the interaction of vector mesons adds the following Lagrangian [38, 39]:

$$\mathcal{L}_{\text{vec+an}} = -\frac{3g^2}{8\pi^2 f_\pi} \epsilon^{\mu\nu\alpha\beta} \text{Tr}[\mathcal{P}(x) \partial_\mu V_\nu(x) \partial_\alpha V_\beta(x)] + \frac{7}{60\pi^2 f_\pi^5} \epsilon^{\mu\nu\alpha\beta} \text{Tr}[\mathcal{P}(x) \partial_\mu \mathcal{P} \partial_\nu \mathcal{P} \partial_\alpha \mathcal{P} \partial_\beta \mathcal{P}] \quad (\text{A26})$$

$$+ 2f_\pi^2 \text{Tr} \left| gV_\mu - eA_\mu Q - \frac{i}{2f_\pi^2} [\mathcal{P}, \partial_\mu \mathcal{P}] \right|^2 \quad (\text{A27})$$

Here, $g \approx m_\rho/\sqrt{2}f_\pi$, $Q = \text{diag}(2/3, -1/3, -1/3)$ is the quark charge matrix, V_μ is the matrix of vector mesons,

$$V_\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{\rho^0+\omega}{\sqrt{2}} & \rho^+ & K^{*+} \\ \rho^- & \frac{-\rho^0+\omega}{\sqrt{2}} & K^{*0} \\ K^{*-} & K^{*0} & \phi \end{pmatrix} \quad (\text{A28})$$

with K^* being associated with $K^*(892)$, and A_μ is the EM field. As discussed, to define the ALP interactions, we start with the corresponding Lagrangian, replace the \mathcal{P} matrix with Eq. (9), and perform the diagonalization (3).

The Lagrangian induces the mixing between A_μ and ρ^0, ω, ϕ , which effectively generates, e.g., electromagnetic decays of P^0 . This way, it is responsible for all the vertices of interactions, in particular for the κ_q cancellation. In particular, it includes the effect of the chiral rotation (A7) coming from the non-invariance of the fermion path integration measure, i.e., reproduces the chiral anomaly at the language of the quark bound states rather than quarks themselves.

Let us show how the κ_q dependence vanishes calculating the coupling of the ALPs to two photons. Expanding Eq. (A27), for the $a\gamma\gamma$ piece, we get

$$\mathcal{L}_{a\gamma\gamma} = \frac{\alpha_{\text{EM}}\epsilon}{\pi f_\pi} c_{\gamma\gamma}^{\text{eff}} F_{\mu\nu} \tilde{F}^{\mu\nu}, \quad (\text{A29})$$

where

$$c_{\gamma\gamma}^{\text{eff}} = \frac{1}{9} (6c_G(3\kappa_u + 1) - 4\sqrt{6}\theta_{\eta a} - 7\sqrt{3}\theta_{\eta' a} - 9\theta_{\pi^0 a}) \quad (\text{A30})$$

The κ_q dependence vanishes after inserting the expressions (A20)-(A22). The first term has the same structure as the term $\propto \text{Tr}[\kappa_q Q^2]$, which is explicitly added in the literature [18] because of the path integral measure non-invariance under the chiral rotation.

We perform two cross-checks. First, in the limit of the “ P^0 -like” ALP (i.e., setting $\theta_{P^0 a} = \delta_{P^0 P^0}$ and $c_G = 0$), the Lagrangian (A29) predicts the decay widths of π^0, η, η' mesons matching the observed data with $\simeq 10\%$ accuracy, typical for the model of Hidden Local Symmetry [35, 38].

Second, let us consider the 2-flavor limit. This is achieved by setting $\theta_{\eta/\eta' a} \rightarrow 0$, dropping the π^0 - η/η' mixing terms in Eq. (A21), and assuming $\kappa_u = 1 - \kappa_d$ (this is the relation between κ_u, κ_d in the case of the 2-flavor setup). Keeping now explicitly the ALP derivative coupling to quarks $\mathcal{L} \ni \partial_\mu a / f_a \sum_q c_q \bar{q} \gamma^\mu \gamma_5 q$, for the expression (A30) we obtain

$$c_{\gamma\gamma}^{\text{eff}} = c_G \left(-\delta \frac{m_{\pi^0}^2}{m_{\pi^0}^2 - m_a^2} - \frac{5}{3} \right) - \frac{m_a^2 (c_u - c_d)}{m_{\pi^0}^2 - m_a^2}, \quad (\text{A31})$$

which exactly matches Eq. (92) in Ref. [16] (after taking into account the relation $c_{qq} \equiv 2c_q$ there).

c. $s \rightarrow d$ transition operator

Next, let us proceed to the process $K \rightarrow a\pi$, generated by the $s \rightarrow d$ transition. The Standard Model analog of this process, $K \rightarrow \pi\pi$, receives contributions from two operators [53] classified by the transformation properties of the chiral operators – $SU(3)$ octet and 27-plet. The coupling constant in front of the latter is severely suppressed, $G_{27}/G_8 \approx 0.05$. Despite this, the SM decay is driven by G_{27} , as the G_8 contribution is proportional to the tiny factor $(m_{\pi^+}^2 - m_{\pi^0}^2)/m_K^2$. However, this is no longer the case for ALPs, and the octet typically makes the main contribution (see, e.g., [19]).

We implement the octet operator; the implementation of the 27-plet may be, in principle, done analogously. The matrix element of the process has the form

$$\begin{aligned} \mathcal{M}_{K^+ \rightarrow \pi^+ a} = & \frac{1}{3} i f_\pi G_8 [6\epsilon c_G (-m_a^2(2\kappa_d + \kappa_u) + m_{K^+}^2(\kappa_d + \kappa_u) + m_{\pi^+}^2 \kappa_d + m_a^2 - m_{\pi^+}^2) \\ & + \sqrt{3}\theta_{a\eta'}(m_a^2 + 2m_{K^+}^2 - 3m_{\pi^+}^2) + 2\sqrt{6}\theta_{\eta a}(m_{K^+}^2 - m_a^2) + 3\theta_{\pi^0 a}(m_a^2 - m_{\pi^+}^2)] \end{aligned} \quad (\text{A32})$$

Note that, unlike the work [18], we do not decouple the η' meson, which leads to the qualitative difference in the scaling of the matrix element with the ALP mass. Namely, after inserting the explicit form of the mixing angles (which identically cancels the κ_q dependence) and working in the limit $\delta = 0$, we get

$$\mathcal{M}_{K^+ \rightarrow \pi^+ a} \approx 8i\epsilon c_G f_\pi G_8 \frac{(m_{K^+}^2 - m_{\pi^+}^2)^2}{m_a^2 - m_{\eta'}^2} \quad (\text{A33})$$

d. *Tensor mesons*

We add the following Lagrangian of the interaction of P s with the tensor meson f_2 [39]:

$$\mathcal{L}_T = -g_T \frac{f_\pi^2}{4} \text{Tr} \left[\left(\partial^\mu \Sigma^\dagger \partial^\nu \Sigma - \frac{1}{2} g^{\mu\nu} \partial^\alpha \Sigma^\dagger \partial_\alpha \Sigma \right) \mathbf{f}_2 \right] f_{2\mu\nu}, \quad (\text{A34})$$

where $\mathbf{f}_2 = \text{diag}(1/2, 1/2, 0)$ is the SU(3) generator of the tensor meson. The coupling $g_T = 13.1 \text{ GeV}^{-1}$ recovers the decay widths of the f_2 meson [36].

Here, we use the replacement (A23) to achieve the κ_q independence. As an illustration, consider the vertex $f_2 a \eta$:

$$V_{f_2 a \eta}^{\mu\nu} = \frac{1}{6} \epsilon g_T \left(-g^{\mu\nu} (p_a \cdot p_\eta) + p_a^\mu p_\eta^\nu + p_a^\nu p_\eta^\mu \right) \left(\sqrt{2} \theta_{a\eta'} + \sqrt{6} c_G (\kappa_d + \kappa_u) + 2\theta_{\eta a} \right) \quad (\text{A35})$$

Inserting the explicit expressions (A21), (A22), for the c_G -dependent piece of the coefficient in front of the tensor structure, we get

$$c_G \frac{(m_\eta^2 - m_{\pi^0}^2) (m_a^2 - 2m_\eta^2 + m_{\pi^0}^2)}{(m_a^2 - m_\eta^2) (m_a^2 - 4m_\eta^2 + 3m_{\pi^0}^2)} \quad (\text{A36})$$

4. Interaction with nucleons

We follow Ref. [28] and consider the low momentum transfer ALP-nucleon interaction in the form

$$\mathcal{L}_{app} = g_{app} \partial_\mu a \bar{p} \gamma^\mu \gamma_5 p \quad (\text{A37})$$

Here,

$$g_{app} = \frac{(4D_s + 3D - F) \theta_{a\eta'}}{2\sqrt{3}} + \epsilon c_G (\kappa_d (F - D) + 2F \kappa_u) + \frac{\theta_{\eta a} (D_s + 2F)}{\sqrt{6}} + \frac{1}{2} (D + F) \theta_{\pi^0 a} \quad (\text{A38})$$

is the ALP-nucleon coupling, with D_s, F, D being phenomenological constants. Using Eqs. (A20)-(A22) and also the relation (A5), the explicit form of g_{app} becomes

$$g_{app} = \frac{\epsilon c_G}{f_\pi} \frac{(m_{\pi^0}^2 - m_a^2) (m_a^2 (3D_s + 2D) - 2m_\eta^2 (2D_s + D + F) + m_{\pi^0}^2 (D_s + 2F))}{(m_a^2 - m_\eta^2) (m_a^2 - m_{\eta'}^2)} + \delta \frac{\epsilon c_G}{f_\pi} \frac{m_{\pi^0}^2 (D + F) (m_{\pi^0}^2 - m_\eta^2) (m_a^2 - 2m_\eta^2 + m_{\pi^0}^2)}{(m_{\pi^0}^2 - m_a^2) (m_a^2 - m_\eta^2) (m_a^2 - m_{\eta'}^2)} \quad (\text{A39})$$

5. Adding axial-vector and heavy pseudoscalar excitations

In the mass range $m_a \gtrsim 1 \text{ GeV}$, axial-vector mesons A_μ and heavy pseudoscalar excitations P_h are essential to describe the ALP interactions. The reason is that ALPs mix with them. For excited pseudoscalar mesons, the mixing structure is similar to the light pseudoscalar case discussed in Sec. A2: the chiral rotation modifies the meson kinetic and mass terms, whereas the derivative coupling translates to the analog of the last term of Eq. (A9). The axial-vector sector generically adds the mixing terms of the type $L \ni M_{AP} A_\mu \partial^\mu P$, where M_{AP} is the mixing coefficient.

It turns out that the $A - a$ mixing is not as important as the mixing with pseudoscalar mesons: unlike the latter, it does not get resonantly enhanced at the vicinity of the A mass. To understand this feature, let us treat the $A_\mu \partial^\mu a$ term as the interaction term and use it as an internal vertex of some matrix element \mathcal{M} of the ALP production or decay. It has the form

$$\mathcal{M} \propto M_{A^0 a} p_a^\mu D_{\mu\nu}^{A^0} \tilde{\mathcal{M}}^\nu = M_{A^0 a} \frac{\left(1 - \frac{m_a^2}{m_{A^0}^2}\right)}{m_a^2 - m_{A^0}^2 - i\Gamma_{A^0} m_{A^0}} p^\nu \tilde{\mathcal{M}}_\nu \equiv \theta_{aA^0}^{\text{eff}} p^\nu \tilde{\mathcal{M}}_\nu, \quad (\text{A40})$$

where $\theta_{aA_0}^{\text{eff}}$ is the mixing angle. We see that it vanishes at $m_a = m_{A_0}$, and hence lacks the enhancement. Therefore, compared to the mixing with the heavy pseudoscalar mesons, it is expected to be subdominant. In addition, incorporating them in a consistent way with the other mesons that are already included in our approach is a non-trivial task. For these two reasons, we will drop their interactions with ALPs. We return to this question in Sec. A 7.

Nevertheless, as we will see, the mixing between axial-vectors and light pseudoscalar mesons are important to describe the interactions of pseudoscalar excitations with vector and light pseudoscalar mesons. Therefore, we briefly describe their phenomenology in Sec. A 5 a.

Our treatment of their interactions follows Refs. [37, 41, 42]. It utilizes the description of different vector, axial-vector, heavy scalar, and pseudoscalar excitations using the $SU(3)$ symmetry, assuming that masses originate from a condensate of the heavy scalar nonet, and fixing the interaction couplings by matching the theoretical prediction of various decay widths with experimentally observed data. This model is called the *Extended Linear Sigma Model*, or **ELSM**. The crucial point is that we turn off the contributions from axial-vector and heavy pseudoscalar sectors for the scale $m \lesssim 1$ GeV, which is explained by the incompleteness of the ELSM in the meson spectrum mass range $M < 1$ GeV (see Sec. A 7).

The mixing of ALPs with P_h contributes to all “ P_h -like” decay modes, including $3\pi, KK\pi, \eta 2\pi$, and many others.

a. Mixing of axial-vector and pseudoscalar mesons

In this section, we follow the approach of including the axial-vector mesons presented in Ref. [41].

We start with the Lagrangian describing the sector of axial-vector A , vector V , and pseudoscalar mesons P :

$$\begin{aligned} \mathcal{L}_0 = & \text{Tr}[(D_\mu \Phi)^\dagger (D_\mu \Phi)] - m_0^2 \left(\frac{G}{G_0}\right)^2 \text{Tr}(\Phi^\dagger \Phi) - \lambda_1 [\text{Tr}(\Phi^\dagger \Phi)]^2 - \lambda_2 \text{Tr}(\Phi^\dagger \Phi)^2 \\ & - \frac{1}{4} \text{Tr}(L_{\mu\nu}^2 + R_{\mu\nu}^2) + \text{Tr} \left[\left(\left(\frac{G}{G_0}\right)^2 \frac{m_1^2}{2} + \Delta \right) (L_\mu^2 + R_\mu^2) \right] + \text{Tr}[H(\Phi + \Phi^\dagger)] + \text{Tr}(\Phi^\dagger \Phi E_0 + \Phi \Phi^\dagger E_0) \\ & + c_1 (\det \Phi - \det \Phi^\dagger)^2 + i \frac{g_2}{2} (\text{Tr}\{L_{\mu\nu}[L^\mu, L^\nu]\} + \text{Tr}\{R_{\mu\nu}[R^\mu, R^\nu]\}) \\ & + \frac{h_1}{2} \text{Tr}(\Phi^\dagger \Phi) \text{Tr}(L_\mu^2 + R_\mu^2) + h_2 \text{Tr}[|L_\mu \Phi|^2 + |\Phi R_\mu|^2] + 2h_3 \text{Tr}(L_\mu \Phi R^\mu \Phi^\dagger) \\ & + g_3 [\text{Tr}(L_\mu L_\nu L^\mu L^\nu) + \text{Tr}(R_\mu R_\nu R^\mu R^\nu)] + g_4 [\text{Tr}(L_\mu L^\mu L_\nu L^\nu) + \text{Tr}(R_\mu R^\mu R_\nu R^\nu)] \\ & + g_5 \text{Tr}(L_\mu L^\mu) \text{Tr}(R_\nu R^\nu) + g_6 [\text{Tr}(L_\mu L^\mu) \text{Tr}(L_\nu L^\nu) + \text{Tr}(R_\mu R^\mu) \text{Tr}(R_\nu R^\nu)] . \end{aligned} \quad (\text{A41})$$

Here,

$$\Phi(x) = \langle \Phi \rangle + S_h + i\mathcal{P}(x) \quad (\text{A42})$$

is the (heavy scalar)-pseudoscalar nonet, with S_h being the heavy scalar nonet, and

$$\langle \Phi \rangle = \frac{1}{\sqrt{2}} \text{diag} \left(\frac{\phi_N}{\sqrt{2}}, \frac{\phi_N}{\sqrt{2}}, \phi_S \right) \quad (\text{A43})$$

being the scalar condensate giving mass to the fields.

$$L_\mu = V_\mu + A_\mu, \quad R_\mu = V_\mu - A_\mu, \quad (\text{A44})$$

are, correspondingly, left and right axial and vector nonets, with V_μ defined in Eq. (A28), and A_μ the axial-vector nonet:

$$A_\mu = \frac{1}{\sqrt{2}} \begin{pmatrix} \frac{f_{1N} + a_1^0}{\sqrt{2}} & a_1^+ & K_1^+ \\ a_1^- & \frac{f_{1N} - a_1^0}{\sqrt{2}} & K_1^0 \\ K_1^- & K_1^0 & f_{1S} \end{pmatrix} \quad (\text{A45})$$

f_{1N} is associated to $f_1(1285)$, f_{1S} to $f_1(1415)$, and K_1 to $K_1(1270)$. $L_{\mu\nu} = \partial_\mu L_\nu - \partial_\nu R_\mu$ (and similarly $R_{\mu\nu}$) is the field strength. Finally, $D_\mu \Phi = \partial_\mu \Phi - ig_1(L_\mu \Phi - \Phi R_\mu)$ is the covariant derivative, with g_1 being an interaction coupling.

The relevant values of the couplings entering Eq. (A41) are summarized in Table III of Ref. [41]; the rest are set to zero. Finally, the glueball G is set to its VEV value G_0 , whereas $E_0 = \text{diag}(0, 0, \epsilon_S)$.

We are interested in the mixing terms between A and P particles. Let us therefore turn off vector and heavy scalar fields S_h .⁷ Thus, the matrix $\Phi(x)$ has the form $\Phi(x) = \langle \Phi \rangle + iP(x)$. The quadratic part of the resulting interaction is

$$\mathcal{L}_{VA}^{(2)} = \sum_{A,X=\{P,a\}} \frac{M_{AX}}{2} A_\mu \partial^\mu X \quad (\text{A46})$$

With this Lagrangian, we reproduce the Lagrangian of the pure A - P mixing from Ref. [41].

It is possible to eliminate the mixing terms by performing the shift

$$A^\mu \rightarrow A^\mu + M_{A^0P} Z_P \partial^\mu P, \quad P \rightarrow Z_P P \quad (\text{A47})$$

where Z_P is the renormalization constant, which ensures that the kinetic term for P has the canonical form after the shift. In the ELSM, Z_P enters the expressions of the P masses in terms of the parameters of the model (A41), as well as the P s' interactions with other mesons.

Our approach of treating this mixing is to assume that the Lagrangian (A41) and the shift (A47) generate the ChPT mass term (A1). I.e., in the sector of pseudoscalar mesons only, the ELSM predictions exactly match the ChPT description. However, we will utilize it in the next section, where we describe the interactions of heavy pseudoscalar nonet.

For completeness, however, we provide the mixing coefficients between the neutral A^0 and ALPs which can be obtained if including them in the same fashion as in the other interactions:

$$M_{a_1^0 a} = g_1 \phi_N (c_G (\kappa_d - \kappa_u) - \theta_{\pi^0 a}), \quad (\text{A48})$$

$$M_{f_1(1285) a} = -\frac{1}{3} g_1 \phi_N \left(\sqrt{3} \theta_{\eta' a} + 3c_G (\kappa_d + \kappa_u) + \sqrt{6} \theta_{\eta a} \right), \quad (\text{A49})$$

$$M_{f_1(1415) a} = \frac{1}{3} g_1 \phi_S \left(-2\sqrt{3} \theta_{\eta' a} + 6c_G (\kappa_d + \kappa_u - 1) + \sqrt{6} \theta_{\eta a} \right) \quad (\text{A50})$$

As it should be, it is κ_q -independent; it also does not vanish in the limit $c_{u,d,s} \rightarrow 0$.

b. Heavy pseudoscalar sector

To describe the interactions of ALPs with heavy pseudoscalar mesons, we follow Ref. [37], which incorporated them in the ELSM. The starting Lagrangian is

$$\begin{aligned} \mathcal{L}_{P_h} = & \text{Tr}[(D_\mu \Phi_h)^\dagger (D_\mu \Phi_h)] + \alpha \text{Tr}[(D_\mu \Phi_h)^\dagger (D_\mu \Phi) + (D_\mu \Phi)^\dagger (D_\mu \Phi_h)] - (m_0^*)^2 \left(\frac{G}{G_0} \right)^2 \text{Tr}(\Phi_h^\dagger \Phi_h) \\ & - \lambda_0 \left(\frac{G}{G_0} \right)^2 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) - \lambda_1^* \text{Tr}(\Phi_h^\dagger \Phi_h) \text{Tr}(\Phi^\dagger \Phi) - \lambda_2^* \text{Tr}(\Phi_h^\dagger \Phi_h \Phi^\dagger \Phi + \Phi_h \Phi_h^\dagger \Phi \Phi^\dagger) \\ & - \kappa_1 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(\Phi^\dagger \Phi) - \kappa_2 [\text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h)]^2 - \kappa_3 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(\Phi_h^\dagger \Phi_h) - \kappa_4 [\text{Tr}(\Phi_h^\dagger \Phi_h)]^2 \\ & - \xi_1 \text{Tr}(\Phi_h^\dagger \Phi \Phi^\dagger \Phi + \Phi_h \Phi^\dagger \Phi \Phi^\dagger) - \xi_2 \text{Tr}(\Phi_h^\dagger \Phi \Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h \Phi^\dagger \Phi_h) - \xi_3 \text{Tr}(\Phi^\dagger \Phi_h \Phi_h^\dagger \Phi_h + \Phi \Phi_h^\dagger \Phi_h \Phi^\dagger) - \xi_4 \text{Tr}(\Phi_h^\dagger \Phi_h)^2 \\ & + \text{Tr}(\Phi_h^\dagger \Phi_h E_1 + \Phi_h \Phi_h^\dagger E_1) + c_1^* [(\det \Phi - \det \Phi_h^\dagger)^2 + (\det \Phi^\dagger - \det \Phi_h)^2] + c_{1E}^* (\det \Phi_h - \det \Phi_h^\dagger)^2 \\ & + \frac{h_1^*}{2} \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(L_\mu^2 + R_\mu^2) + \frac{h_{1E}^*}{2} \text{Tr}(\Phi_h^\dagger \Phi_h) \text{Tr}(L_\mu^2 + R_\mu^2) \\ & + h_2^* \text{Tr}(\Phi_h^\dagger L_\mu L^\mu \Phi + \Phi^\dagger L_\mu L^\mu \Phi_h + R_\mu \Phi_h^\dagger \Phi R^\mu + R_\mu \Phi^\dagger \Phi_h R^\mu) + h_{2E}^* \text{Tr}[|L_\mu \Phi_h|^2 + |\Phi_h R_\mu|^2] \\ & + 2h_3^* \text{Tr}(L_\mu \Phi_h R^\mu \Phi^\dagger + L_\mu \Phi R^\mu \Phi_h^\dagger) + 2h_{3E}^* \text{Tr}(L_\mu \Phi_h R^\mu \Phi_h^\dagger). \end{aligned} \quad (\text{A51})$$

Here,

$$\Phi_h = S_{hh} + \frac{i}{\sqrt{2}} \begin{pmatrix} \frac{\eta_N + \pi_E^0}{\sqrt{2}} & \pi_E^+ & K_E^+ \\ \pi_E^- & \frac{\eta_N - \pi_E^0}{\sqrt{2}} & K_E^0 \\ K_E^- & \bar{K}_E^0 & \eta_N \end{pmatrix}, \quad (\text{A52})$$

⁷ We have a posteriori checked that their large mass, absence of mixing with ALPs, and presence of the lighter scalar nonet makes

their contribution to the processes with ALPs subdominant.

is the excited scalar-pseudoscalar nonet, with S_{hh} denoting the second heavy scalar fields. Here, π_E is associated to $\pi(1300)$, K_E to $K(1460)$, η_N to $\eta(1295)$ and η_S to $\eta(1440)$. The association assumes that the mentioned states are 2-quark bound states, although there is ambiguity. As in the axial-vector sector, the covariant derivative has the form $D_\mu\Phi_h = \partial_\mu\Phi_h - g_{1E}(L_\mu\Phi_h - \Phi_h R_\mu)$, and $G = G_0$.

Following Ref. [37], we drop the terms proportional to $\alpha, \lambda_0^*, \lambda_1^*, \xi_1, h_1^*, \kappa_{1-4}$. It implies, in particular, the absence of the mixing between the heavy and light pseudoscalar mesons. We comment on its consequences in Sec. A 7. We also set $S_{hh} = 0$, as similarly to the nonet S_h , they are not expected to contribute sizeably to the interactions.

The interactions of Φ_h with vector and light pseudoscalar mesons are obtained with the help of the transformation (A47); it induces the vertices $\Phi_h\Phi V$, $\Phi_h\Phi\Phi\Phi$, dominating the decays of Φ_h and hence ALPs, due to the mixing with heavy pseudoscalars.

After inserting the explicit form of Φ matrix, Eq. (A42), one arrives at the following mass and kinetic matrices between the ALP and the neutral mesons $P_h^0 = \pi^0(1300), \eta(1295), \eta(1440)$, similar to Eqs. (A12), (A13):

$$\hat{M} = \begin{pmatrix} m_a^2 & M_{a\pi^0(1300)} & M_{a\eta(1295)} & M_{a\eta(1440)} \\ M_{a\pi^0(1300)} & m_{\pi^0(1300)}^2 & 0 & 0 \\ M_{a\eta(1295)} & 0 & m_{\eta(1295)}^2 & 0 \\ M_{a\eta(1440)} & 0 & 0 & m_{\eta(1440)}^2 \end{pmatrix} \quad (\text{A53})$$

with

$$M_{a\pi^0(1300)} = -c_G \frac{f_\pi}{2f_a} (\kappa_d - \kappa_u) (2m_0^{*2} + (\lambda_2^* - \xi_2) \phi_N^2), \quad (\text{A54})$$

$$M_{a\eta(1295)} = c_G \frac{f_\pi}{2f_a} (\kappa_d + \kappa_u) (2m_0^{*2} + (\lambda_2^* - \xi_2) \phi_N^2), \quad (\text{A55})$$

$$M_{a\eta(1440)} = -c_G \frac{f_\pi}{f_a} \sqrt{2} (\kappa_d + \kappa_u - 1) (m_0^{*2} + (\lambda_2^* - \xi_2) \phi_S^2 - 2\epsilon_S^E), \quad (\text{A56})$$

and

$$\hat{K} = \begin{pmatrix} 1 & \frac{f_\pi(c_G(\kappa_u - \kappa_d) - c_d + c_u)}{f_a} & \frac{f_\pi(c_G(\kappa_d + \kappa_u) + c_d + c_u)}{f_a} & \frac{\sqrt{2}f_\pi(c_s - c_G(\kappa_d + \kappa_u - 1))}{f_a} \\ \frac{f_\pi(c_G(\kappa_u - \kappa_d) - c_d + c_u)}{f_a} & 1 & 0 & 0 \\ \frac{f_\pi(c_G(\kappa_d + \kappa_u) + c_d + c_u)}{f_a} & 0 & 1 & 0 \\ \frac{\sqrt{2}f_\pi(c_s - c_G(\kappa_d + \kappa_u - 1))}{f_a} & 0 & 0 & 1 \end{pmatrix} \quad (\text{A57})$$

Unlike the case of the light pseudoscalar sector, there is no mixing between various mesons from the beginning – the mixing is solely between P_h^0 and a . Utilizing the relation between the parameters of the Lagrangian and the masses of the heavy pseudoscalars (with the help of our *Mathematica* notebook, we recovered Eqs. (21)-(27) in Ref. [37]), for the resulting mixing angles, we get

$$\theta_{\pi^0(1300)a} = -\frac{f_\pi}{f_a} \left(c_G (\kappa_u - \kappa_d) + \frac{m_a^2(c_u - c_d)}{m_a^2 - m_{\pi^0(1300)}^2} \right), \quad (\text{A58})$$

$$\theta_{\eta(1295)a} = -\frac{f_\pi}{f_a} \left(c_G (\kappa_d + \kappa_u) + \frac{m_a^2(c_d + c_u)}{m_a^2 - m_{\eta(1295)}^2} \right), \quad (\text{A59})$$

$$\theta_{\eta(1440)a} = -\sqrt{2} \frac{f_\pi}{f_a} \left(c_G (1 - \kappa_d - \kappa_u) + \frac{m_a^2 c_s}{m_a^2 - m_{\eta(1440)}^2} \right) \quad (\text{A60})$$

Unlike the light pseudoscalars, heavy pseudoscalar mesons have a non-negligible decay width, which smears the resonant enhancement. To include this effect in practice, we ensure that the κ_q -dependence vanishes in the zero-width limit, explicitly set $\kappa_q = 0$, and replace the denominators of the mixing angles: $m_a^2 - m_{P_h^0}^2 \rightarrow m_a^2 - m_{P_h^0}^2 - i\Gamma_{P_h^0} m_{P_h^0}$.

c. *How the c_G -mediated contribution vanishes*

Now, let us consider the contribution of the Lagrangian (A51) to some processes. Choosing, for instance, the process $a(\rightarrow \eta^*(1440)) \rightarrow KK\pi$, we get for the corresponding Lagrangian

$$\mathcal{L}_a = \frac{if_\pi}{2f_a} a(x) \partial^\mu K^-(x) K_{\mu^+}^*(x) h_3^* \theta_{K_1^-(1270)K^-} \left[\sqrt{2} (\phi_S (\theta_{\eta(1295)a} + \theta_{\pi^0(1300)a}) - \phi_N \theta_{\eta(1440)a}) + 2c_G (\phi_N (\kappa_d + \kappa_u - 1) + \sqrt{2}\phi_S \kappa_u) \right] \quad (\text{A61})$$

After using Eqs. (A58)-(A60), for the expression in the brackets, we get

$$\frac{2c_s \phi_N}{m_a^2 - m_{\eta(1440)}^2} - \frac{\sqrt{2}\phi_S (c_d(m_{\eta(1295)}^2 - m_{\pi^0(1300)}^2) + c_u(2m_a^2 - m_{\eta(1295)}^2 - m_{\pi^0(1300)}^2))}{(m_a^2 - m_{\eta(1295)}^2)(m_a^2 - m_{\pi^0(1300)}^2)} \quad (\text{A62})$$

I.e., whereas the $c_{u,d,s}$ terms obviously survive, the dependence on c_G vanishes identically. It happens because of the simple structure of the interaction operators and the trivial structure of heavy pseudoscalars utilized in Ref. [37]: unlike the case light pseudoscalars, $\eta(1295)$, $\eta(1440)$ exactly match the isoscalar and singlet components of the nonet; also, there is no analog of the isospin breaking.

The same cancellation happens also for any other interaction induced by the Lagrangian (A51). Therefore, if dropping the terms proportional to $\alpha, \lambda_0^*, \lambda_1^*, \xi_1, h_1^*, \kappa_{1-4}$ (as it is done in [37]), the ALPs coupled to gluons can only experience these interactions via tiny couplings $c_{u,d,s}$ induced because of the renormalization flow of the ALP gluonic operator from the scale Λ .

6. Cross-checks of our approach

We have validated our combined approach in various independent ways:

1. Comparing symbolic expressions for various quantities with the literature. For instance, assuming the 2-flavor limit, we recovered the mass matrix (A13), the ALP-photon coupling, and the ALP decay width into three pions with the works [12, 16] (see also Sec. A3b). Next, we have compared the expressions for the pure ChPT contribution to the matrix elements of ALP decays to Ref. [13] and mostly found the exact agreement. The only exception concerns the κ_q -dependent piece, which is absent for the $a \rightarrow \eta\pi\pi$ matrix element in [13]. For the added interactions with scalar, vector, and tensor mesons (where we followed Ref. [36]), we have found the agreement with [36] except for the sector of the scalar mesons, modified by our treatment (discussed in Sec. A3a).
2. Ensuring κ_q independence of various vertices and matrix elements. When computing different quantities, we insert an explicit form of the mixing angles and calculate the κ_q -dependent pieces. They always vanish.
3. Comparing the widths of the Standard Model processes with their measured values. We consider the processes $\eta \rightarrow \pi^+\pi^-\gamma$, $\eta' \rightarrow \eta 2\pi$, $\eta' \rightarrow 4\pi$, $\pi^0/\eta/\eta' \rightarrow 2\gamma$, which in the model we use goes via the mixing of vector mesons with photons [38] and intermediate vector and scalar excitations [39, 40]. We find good agreement within 10%. This deviation is subdominant compared to the other uncertainty sources (highlighted in Sec. A7) and also uncertainties coming from the experimental setup (which may include a cascade enhancement of the ALP production in the thick target – already be an $\mathcal{O}(1)$ effect) and lack of knowledge about the meson's spectra). Regarding the newly added axial-vector and heavy pseudoscalar sectors, we have recovered the results of Refs. [37, 41].
4. Reproducing the results of Ref. [22], which studied the ALPs universally coupled to fermions, after turning off the contributions from axial-vector and heavy pseudoscalar mesons. For this setup, the hadronic ALP widths differ within a factor of 1.5. The discrepancy is caused by the improvement of the sector of the interactions with the scalar mesons made in the given paper (see Sec. A3a).

We finalize the discussion by comparing our calculation of the ALP decay width with the results of Ref. [13]. The comparison is shown in Fig. 3. The result of Ref. [13] is similar to our calculations obtained within the κ_q -dependent approach (which does not utilize the rotations (A24), (A23)). The discrepancy arises mainly because of the differences in the treatment of light resonances, which we described in Sec. A2. In the same figure (right panel), we compare the total decay widths of ALPs that are universally coupled to fermions and those coupling solely to gluons. In the fermionic case, the effects of the heavy pseudoscalar resonances are much stronger (see Eq. (A60)), leading to a significant enhancement of the decay width.




FIG. 3. *Left panel*: the comparison of the ALP decay widths obtained in this work and in Ref. [13]. The meaning of the blue lines and the dashed red line is the same as in Fig. 2 of the main text. *Right panel*: the comparison of the total decay widths of the ALPs that have the universal coupling to all fermions (see Refs. [22, 25, 26] for the discussion on the model) and those having solely gluonic coupling at the scale $\Lambda = 1$ TeV. For the fermionic coupling case, the jump at $m_a = 2m_\mu$ is caused by the turning on of the di-muon decay, which dominates until the mass $m_a \simeq m_{\eta'}$.

7. Limitations of our approach

Our approach to ALP phenomenology, summarized in Sec. A, has limitations intimately related to the maturity of the frameworks describing effective interactions of mesons. There are two major problems.

The first problem arises because there is no unique comprehensive framework describing all the meson excitations with masses below 2 GeV [42]. For instance, ELSM is fitted to describe the main decays of some of the pseudoscalar, scalar, vector, and tensor mesons. On the other hand, it does not consistently incorporate the sectors of light scalar nonet and pseudovector fields, as well as anomalous interactions with vector mesons. The latter may be essential for decays of light pseudoscalar mesons [39, 40]. Partially because of this, it does not fit the experimental data on various subdominant decays of mesons which are essential for describing the ALP decay pattern, such as $\eta \rightarrow \pi^+\pi^-\gamma$ or $\eta' \rightarrow 4\pi$. Because of this, we use the hybrid framework, utilizing the ELSM to describe the sector of heavy pseudoscalars.

Next, some heavy pseudoscalar excitations, such as $\pi(1800)$, are not implemented as well. Because of the resonantly enhanced mixing of such mesons with the ALPs, it would severely influence the ALP phenomenology.

One of the main reasons for this is the lack of robust experimental data on these resonances [52], leading to huge uncertainties in decay widths and masses of some resonances. It leads to a complexity of interpreting various meson states assigned in Sec. A 5 as 2-quark bound states may be 4-quark bound states or an admixture between 2-quark and 4-quark bound states [42]; the latter would directly change the interaction operators of mesons and hence the ALP phenomenology.

An independent problem concerns the ambiguity of the phenomenological Lagrangians. For example, consider again

the interaction Lagrangian of heavy pseudoscalar mesons P_h as defined in Ref. [37]:

$$\begin{aligned}
\mathcal{L}_{P_h} = & \text{Tr}[(D_\mu \Phi_h)^\dagger (D_\mu \Phi_h)] + \alpha \text{Tr}[(D_\mu \Phi_h)^\dagger (D_\mu \Phi) + (D_\mu \Phi)^\dagger (D_\mu \Phi_h)] - (m_0^*)^2 \left(\frac{G}{G_0}\right)^2 \text{Tr}(\Phi_h^\dagger \Phi_h) \\
& - \lambda_0 \left(\frac{G}{G_0}\right)^2 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) - \lambda_1^* \text{Tr}(\Phi_h^\dagger \Phi_h) \text{Tr}(\Phi^\dagger \Phi) - \lambda_2^* \text{Tr}(\Phi_h^\dagger \Phi_h \Phi^\dagger \Phi + \Phi_h \Phi_h^\dagger \Phi \Phi^\dagger) \\
& - \kappa_1 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(\Phi^\dagger \Phi) - \kappa_2 [\text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h)]^2 - \kappa_3 \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(\Phi_h^\dagger \Phi_h) - \kappa_4 [\text{Tr}(\Phi_h^\dagger \Phi_h)]^2 \\
& - \xi_1 \text{Tr}(\Phi_h^\dagger \Phi \Phi^\dagger \Phi + \Phi_h \Phi^\dagger \Phi \Phi^\dagger) - \xi_2 \text{Tr}(\Phi_h^\dagger \Phi \Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h \Phi^\dagger \Phi_h) - \xi_3 \text{Tr}(\Phi^\dagger \Phi_h \Phi_h^\dagger \Phi_h + \Phi \Phi_h^\dagger \Phi_h \Phi) - \xi_4 \text{Tr}(\Phi_h^\dagger \Phi_h)^2 \\
& + \text{Tr}(\Phi_h^\dagger \Phi_h E_1 + \Phi_h \Phi_h^\dagger E_1) + c_1^* [(\det \Phi - \det \Phi_h^\dagger)^2 + (\det \Phi^\dagger - \det \Phi_h)^2] + c_{1E}^* (\det \Phi_h - \det \Phi_h^\dagger)^2 \\
& + \frac{h_1^*}{2} \text{Tr}(\Phi_h^\dagger \Phi + \Phi^\dagger \Phi_h) \text{Tr}(L_\mu^2 + R_\mu^2) + \frac{h_{1E}^*}{2} \text{Tr}(\Phi_h^\dagger \Phi_h) \text{Tr}(L_\mu^2 + R_\mu^2) \\
& + h_2^* \text{Tr}(\Phi_h^\dagger L_\mu L^\mu \Phi + \Phi^\dagger L_\mu L^\mu \Phi_h + R_\mu \Phi_h^\dagger \Phi R^\mu + R_\mu \Phi^\dagger \Phi_h R^\mu) + h_{2E}^* \text{Tr}[|L_\mu \Phi_h|^2 + |\Phi_h R_\mu|^2] \\
& + 2h_3^* \text{Tr}(L_\mu \Phi_h R^\mu \Phi^\dagger + L_\mu \Phi R^\mu \Phi_h^\dagger) + 2h_{3E}^* \text{Tr}(L_\mu \Phi_h R^\mu \Phi_h^\dagger). \tag{A63}
\end{aligned}$$

When fitting the coefficients of this Lagrangian to the data, Ref. [37] dropped for simplicity the terms inducing the mixing between light and heavy pseudoscalars. As we have discussed in Sec. A 5 b, in this case, κ_q invariance also leads to the cancellation of the κ_q -independent pieces in the interaction between ALPs and P_h . On the other hand, the dropped terms may be essential for the ALPs.

Indeed, let us consider the second term, which induces such mixing, and perform the usual diagonalization (A18). Among the other terms, it includes the ALP-meson mixing which does not vanish in the limit $c_{u,d,s} \rightarrow 0$:

$$\begin{aligned}
\mathcal{L}_\alpha \subset \alpha \frac{f_\pi}{f_a} c_G a(x) \left[\eta_{1440}(x) \frac{\sqrt{2} (m_a^2 - m_{\pi^0}^2) (m_\eta^2 - m_{\eta'}^2)}{3 (m_a^2 - m_\eta^2) (m_a^2 - m_{\eta'}^2)} - \right. \\
\left. - \eta_{1295}(x) \frac{(m_a^2 (m_{\eta'}^2 + 2m_\eta^2 - 3m_{\pi^0}^2) + 2m_{\pi^0}^2 m_{\eta'}^2 + m_\eta^2 (m_{\pi^0}^2 - 3m_{\eta'}^2))}{3 (m_a^2 - m_\eta^2) (m_a^2 - m_{\eta'}^2)} \right] \tag{A64}
\end{aligned}$$

Treating it for the moment as the additional mixing, we see that the c_G -dependent piece involving the ALP- P_h^0 mixing no longer vanishes (there is no corresponding ‘‘counterterm’’ from the rotation (A24), (A23)), and may sizeably contribute to the ALP decay widths. However, including them would require a complete re-analysis of the whole ELSM fit (as additional mixing between P_h and P may be non-negligible), which goes beyond the scope of our study.

Therefore, we conclude that the robustness of the description of the ALP phenomenology is significantly limited. Nevertheless, incorporating the interactions in `Mathematica` notebook, we provide a state-of-the-art analysis that can be easily improved in the future once the meson spectroscopy becomes more mature.

Appendix B: Production of ALPs

Below, we briefly discuss our method to describe the production modes of ALPs (more details may be found in Ref. [49]).

The commonly adopted approach to describe ALP production [21, 26] was to approximate the flux of ALPs by the formula

$$\frac{d^2 N_a}{d\phi_a dE_a} = \left(\frac{d^2 N_a}{d\phi_a dE_a} \right)_{\text{direct+light decays}} + \sum_{B, X_{s/d}} N_B \cdot \text{Br}(B \rightarrow X_{s/d} a) \frac{df_a^{(B)}}{d\phi_a dE_a} \tag{B1}$$

Here, ϕ, E stay for the polar angle and energy correspondingly.

$$\left(\frac{d^2 N_a}{d\phi_a dE_a} \right)_{\text{direct+light decays}} = \sum_{P^0} |\theta_{P^0 a}|^2 \frac{d^2 N_{P^0}}{d\phi_{P^0} dE_{P^0}} \Big|_{\phi_{P^0}, E_{P^0} \rightarrow \phi_a, E_a} \tag{B2}$$

describes the contribution of the ALPs produced by deep inelastic scatterings and decays of light mesons, with $\phi_{P^0}, E_{P^0} \rightarrow \phi_a, E_a$ being some transformation relating meson’s kinematic to the ALP kinematics. The second summand describes the production of the ALPs by decays of B mesons, with N_B being the total number of B mesons

of the given type produced in collisions, $\text{Br}(B \rightarrow X_{s/d} a)$ the branching ratio of the decay into an ALP and a hadronic state containing an s/d quark, and $f_a^{(B)}$ the distribution in polar angle and energy of the ALPs produced in this decay (it is normalized by one).

The term (B2) literally means “the flux of ALPs is the summed fluxes of mother mesons times the squared mixing angle”.

This approximation has two problems. First, it explicitly depends on unphysical κ_q parameters entering $\theta_{P^0 a}$. Second, it wrongly describes the mass dependence of the ALP flux, both in terms of the integrated value and the kinematics (the proposed ϕ, E transformation is ambiguous and leads to unphysical bumps in the angular distribution of ALPs at low center-of-mass frame collision energy). A clear illustration is to consider the ALPs from the mixing with pions; while $\simeq 60 - 70\%$ of pions are produced by decays of heavier mesons, this is not the case of the ALPs with $m_a \gg m_{\pi^0}$.

As we have discussed in the main paper, we explicitly decompose this production into separate modes: the ALP production in the proton bremsstrahlung (the initial state radiation process), quark fragmentation (mostly the final state radiation process relevant for masses $m_a \lesssim 2$ GeV), gluon fusion/Drell-Yan process (the final state radiation relevant for heavier ALPs), and decays of light mesons (see also Ref. [44] for the definition of the processes). The descriptions of the proton bremsstrahlung and gluon fusion processes may be found in Refs. [22, 28, 44], whereas below, we discuss the production in decays of mesons and the quark fragmentation.

1. Decays of mesons

Let us now present the method to calculate the branching ratios of various mesons, denoted by h , into ALPs. First, using the Lagrangian with the ALP interactions, we compute the matrix element of the process, \mathcal{M} . Then, we calculate the decay widths using the standard formulas [54]:

$$\Gamma_{h \rightarrow a}^{2\text{-body}} = \frac{|\mathcal{M}|^2 |\mathbf{p}|}{8\pi m_h^2} \quad (\text{B3})$$

for 2-body decays, where $|\mathbf{p}|$ is the momentum of the decay product at the rest frame of the decaying h , and

$$\Gamma_{h \rightarrow a}^{3\text{-body}} = F_s \frac{1}{(2\pi)^3 8m_h} \int dE_1 dE_3 |\mathcal{M}|^2, \quad (\text{B4})$$

for 3-body decays, where $E_{1,3}$ are the energies of the particles 1,3 in the process $h \rightarrow 1 + 2 + 3$ at h 's rest frame, and F_s is the symmetry factor, being $1/n!$ if there are n identical particles in the final state and 1 otherwise. We will consider the following decay processes:

$$\rho^\pm \rightarrow \pi^\pm a, \quad \eta \rightarrow \pi^+ \pi^- a, \quad \eta \rightarrow 2\pi^0 a, \quad K_S \rightarrow \pi^0 a, \quad \omega \rightarrow \pi^+ \pi^- a, \quad \omega \rightarrow a\gamma \quad (\text{B5})$$

As a cross-check, considering the “pion”-like ALP, we have checked that together with the production via fragmentation, decays of mesons accumulate $\simeq 90\%$ of the SM neutral pion production flux. The processes that we drop are, e.g., $\Lambda \rightarrow pa$ and $K^\pm \rightarrow a\pi^\pm$. The first process has too limited phase space and produces very soft ALPs (as most of the momentum is carried away by the outgoing proton). In their turn, K^\pm are long-lived and typically do not have time to decay before interacting with the infrastructure surrounding the collision point (or target for beam dump experiments). Given the complexity of describing its decays, and also the suppression of the corresponding event yield, it can be safely neglected.

2. Fragmentation

Qualitatively, the production in the quark fragmentation effectively replaces a meson P^0 occurring at the last stage of the fragmentation chain with the ALP a .

To account for the proper ALP mass dependence of the flux, we incorporate the production of ALPs via mixing with $P^0 = \pi^0, \eta, \eta'$ in PYTHIA8 [50]. The production of the ALPs from the mixing with the given meson P^0 is described by P^0 's fragmentation function, carefully tuned to the data on the meson fluxes available at different facilities and having at most $\mathcal{O}(1)$ uncertainty. More details can be found in Ref. [49].

The mixing with heavier mesons (axial-vector mesons A and heavy pseudoscalars P_h^0) cannot be properly accommodated, as PYTHIA8 misses these particles in the spectrum. Even if including them, it would be unrealistic to

properly accommodate the contribution of these mesons because of the complexity of properly tuning PYTHIA8 setup to describe. Therefore, our estimate of the production in this channel is conservative.

The rate of the replacement $P^0 \rightarrow a$ cannot be given by the mixing angle $\theta_{P^0 a}$, as it is clearly κ_q -dependent and misses contributions from the direct operator (see Fig. 1). Instead, we replace it with the ‘‘effective’’ rate $\Theta_{P^0 a}$, which we define from some typical processes involving the $P^0 a$ mixing and then extrapolate on the whole fragmentation process. Schematically, this rate has the form

$$\Theta_{P^0 a} = \theta_{P^0 a} + \dots, \quad (\text{B6})$$

where \dots stays for the direct operator contribution and other mixings.

Specifically, to define Θ for the mixing with $P^0 = \pi^0, \eta, \eta'$, we consider, correspondingly, the following processes:

$$\pi^0 \pi^0 \rightarrow \pi^0 a, \quad \pi^0 \eta \rightarrow \pi^0 a, \quad \pi^0 \eta' \rightarrow \pi^0 a \quad (\text{B7})$$

The specific processes have been chosen because they are the simplest scattering processes and because the $P^0 a$ mixing directly contributes to them. Namely, their matrix elements have the form

$$\mathcal{M}_{\pi^0 P^0 \rightarrow \pi^0 a} = \Theta_{\pi^0 a} \cdot \tilde{\mathcal{M}}, \quad (\text{B8})$$

Explicitly, we get

$$\begin{aligned} \Theta_{\pi^0 a} = \theta_{\pi^0 a} + c_G \left[\kappa_u - \kappa_d - \delta (\kappa_d + \kappa_u) \left(\sqrt{3} \theta_{\pi^0 \eta'} + \sqrt{6} \theta_{\pi^0 \eta} + 1 \right) \right] \\ - \frac{1}{3} \delta \left(3 \theta_{\pi^0 \eta'} + 3 \sqrt{2} \theta_{\pi^0 \eta} + \sqrt{3} \right) \theta_{\eta' a} - \frac{1}{3} \delta \theta_{\eta a} \left(3 \sqrt{2} \theta_{\pi^0 \eta'} + 6 \theta_{\pi^0 \eta} + \sqrt{6} \right), \end{aligned} \quad (\text{B9})$$

$$\begin{aligned} \Theta_{\eta a} = \theta_{\eta a} + \frac{1}{2} c_G \left[\kappa_d \left(\delta \left(2 \sqrt{2} \theta_{\pi \eta'} + \theta_{\pi \eta} + \sqrt{6} \right) + \sqrt{6} \right) + \kappa_u \left(\sqrt{6} - \delta \left(2 \sqrt{2} \theta_{\pi \eta'} + \theta_{\pi \eta} + \sqrt{6} \right) \right) \right] \\ - \frac{1}{2} \delta \theta_{\pi^0 a} \left(2 \sqrt{2} \theta_{\pi \eta'} + \theta_{\pi \eta} + \sqrt{6} \right) + \frac{\theta_{\eta' a}}{\sqrt{2}} \end{aligned} \quad (\text{B10})$$

$$\begin{aligned} \Theta_{\eta' a} = \theta_{\eta' a} + c_G \left[\kappa_d \left(\delta \left(-\theta_{\pi^0 \eta'} + 2 \sqrt{2} \theta_{\pi^0 \eta} + \sqrt{3} \right) + \sqrt{3} \right) + \kappa_u \left(\delta \left(\theta_{\pi^0 \eta'} - 2 \sqrt{2} \theta_{\pi^0 \eta} - \sqrt{3} \right) + \sqrt{3} \right) \right] + \\ + \delta \cdot \theta_{\pi^0 a} \left(\theta_{\pi^0 \eta'} - 2 \sqrt{2} \theta_{\pi^0 \eta} - \sqrt{3} \right) + \sqrt{2} \theta_{\eta a} \end{aligned} \quad (\text{B11})$$

Here,

$$\theta_{\pi^0 \eta} = -\frac{\sqrt{\frac{2}{3}} m_{\pi^0}^2}{m_{\eta}^2 - m_{\pi^0}^2}, \quad \theta_{\pi^0 \eta'} = -\frac{m_{\pi^0}^2}{\sqrt{3} (m_{\eta'}^2 - m_{\pi^0}^2)} \quad (\text{B12})$$

are the mixing angles between π^0 and η/η' (modulus δ) originating from the pure ChPT (see the mass matrix (A13)).

Inserting them and the ALP- P^0 mixing angles (A20)-(A22) in Eqs. (B9)-(B11), it can be seen that the κ_q dependence drops out.

Note that using $\Theta_{P^0 a}$ instead of $\theta_{P^0 a}$ is also needed in the case of ALPs with the dominant coupling to quarks.