RANDOM WALKS WITH SQUARE-ROOT BOUNDARIES: THE CASE OF EXACT BOUNDARIES $g(t) = c\sqrt{t+b} - a$

DENIS DENISOV, ALEXANDER SAKHANENKO, SARA TERVEER, AND VITALI WACHTEL

ABSTRACT. Let S(n) be a real valued random walk with i.i.d. increments which have zero mean and finite variance. We are interested in the asymptotic properties of the stopping time $T(g) := \inf\{n \ge 1 : S(n) \le g(n)\}$, where g(t) is a boundary function. In the present paper we deal with the parametric family of boundaries $\{g_{a,b}(t) = c\sqrt{t+b} - a, b \ge 0, a > c\sqrt{b}\}$. First, assuming that sufficiently many moments of increments of the walk are finite, we construct a positive space-time harmonic function W(a, b). Then we show that there exist p(c) > 0 and a constant $\varkappa(c)$ such that $\mathbf{P}(T_{g_{a,b}} > n) \sim \varkappa(c) \frac{W(a,b)}{np(c)/2}$ as $n \to \infty$.

1. INTRODUCTION, MAIN RESULTS AND DISCUSSION

Consider a one-dimensional random walk

$$S(0) = 0, \quad S(n) = X_1 + X_2 + \ldots + X_n, \ n \ge 1,$$

where X_1, X_2, \ldots are independent copies of a random variable X. We shall always assume that the random variable X has zero mean and unit variance, $\mathbf{E}X = 0$ and $\mathbf{E}X^2 = 1$. The main purpose of this paper is to study the behaviour of $\{S(n)\}$ killed at crossing a square-root boundary. More precisely, for a function g(t) we define the stopping time

$$T(g) := \inf\{n \ge 1 : S(n) \le g(n)\}.$$

We shall always assume that g(0) < 0 and that there exists a constant $c \in \mathbb{R}$ such that

$$g(t) \sim c\sqrt{t} \quad \text{as } t \to \infty.$$
 (1)

A quite important role in the study of the stopping time T_g with general g satisfying (1) is played by the following parametric family of boundary functions. For $b \ge 0$ and $a > c\sqrt{b}$ we set

$$g_{a,b}(t) = c\sqrt{t+b} - a, \ t \ge 0.$$
 (2)

In this case we set

$$T_{a,b} := T(g_{a,b}) = \inf\{n \ge 1 : a + S(n) \le c\sqrt{n+b}\}.$$

For this special family of boundaries we can interpret a as a starting position of the walk and b as a time shift. Using this interpretation of parameters a and b we

²⁰²⁰ Mathematics Subject Classification. Primary 60G50; Secondary 60G40, 60F17.

Key words and phrases. Random walk, exit time, harmonic function, conditioned process. S. Terveer was partially supported by the Deutsche Forschungsgemeinschaft (DFG, German

Research Foundation) – TRR 352 "Mathematics of Many-Body Quantum Systems and Their Collective Phenomena" – Project-ID 470903074.

V. Wachtel was partially supported by DFG.

may define a more general family of stopping times. For every boundary function q we set

$$T_{a,b}(g) := \inf\{n \ge 1 : a + S(n) \le g(n+b)\}, \quad a > g(b).$$

Thus, $T_{a,b} = T_{a,b}(c\sqrt{\cdot})$ and, consequently, we deal with the exact boundary $c\sqrt{t}$ modulo temporal and spatial shifts.

First-passage problems related to stopping times T(g) and $T_{a,b}$ can be seen as a one-sided analogue of the problem with two boundaries:

$$T(g_1, g_2) := \inf\{n \ge N : S(n) \notin (g_1(n), g_2(n))\},\$$

where $N \geq 1$, $g_i(t) \sim c_i \sqrt{t}$ with $c_1 \leq 0 \leq c_2$, $c_1 < c_2$. (The additional parameter N is needed here, since the interval $(g_1(n), g_2(n))$ can be empty for small values of n.) Apparently, the study of such two-sided first-passage problems has been initiated by Blackwell and Freedman [1]. They have considered first time when the absolute values of a symmetric simple random walk exceeds $c\sqrt{n}$ and have shown that this random time has finite expectation if and only if c < 1. Motivated by this surprising observation Breiman [2] has investigated the tail behaviour of T_{g_1,g_2} in the case when $-g_1(t) = g_2(t) = c\sqrt{t}$. Assuming that the third moment of X is finite he has shown that

$$\mathbf{P}(T(g_1, g_2) > n) \sim \alpha n^{-\gamma(c)}$$

where $\gamma(c)$ does not depend on the distribution of the walk. Breiman's result was later improved by Greenwood and Perkins [9], who have shown that if $\mathbf{E}X^2 \log(1 + |X|)$ is finite then there exist a positive number $p(c_1, c_2)$ and a slowly varying function L_{q_1, q_2} such that

$$\mathbf{P}(T(g_1, g_2) > n) \sim \frac{L_{g_1, g_2}(n)}{n^{p(c_1, c_2)/2}}$$
 as $n \to \infty$.

Moreover, they have proven a limit theorem for S(n) conditioned on $\{T_{q_1,q_2} > n\}$.

The corresponding picture in the case of one-sided boundaries is not as complete as in the two-sided case. Greenwood and Perkins [9] have shown that if $\mathbf{E}X^2 \log(1+|X|) < \infty$ and if $g(t) \sim c\sqrt{t}$ with some $c \leq 0$, then there exist p(c) > 0 and a slowly varying function L_g such that

$$\mathbf{P}(T(g) > n) \sim \frac{L_g(n)}{n^{p(c)/2}}.$$
(3)

As in the two-sided case, the authors prove also a limit theorem for S(n) conditioned on $\{T(g) > n\}$. Later it was shown in [6] that (3) remains valid without extra moment assumption in the case c = 0. Moreover, that paper contains sufficient and necessary conditions on boundaries g(t) under which g(t) is asymptotically constant. The case c = 0, i.e. $g(t) = o(\sqrt{t})$ is probably the most studied case of first-passage times over moving boundaries, and we refer to [6] for a rather detailed overview of existing results.

To the best of our knowledge, there is no version of (3) in the case when (1) holds with some c > 0. Furthermore, as far as we know, there was no progress after [9] in studying exit times T(g) and related problems in the case $c \neq 0$ despite the fact that walks with square-root boundaries play a significant role in various statistical and learning settings; see, for example, a recent paper Harvey et al [10].

The main purpose of the present paper is to develop a technique which will allow one to study the tail behaviour of $T_{a,b}$ for boundaries $g_{a,b}(t)$ defined in (2) with negative as well as positive values of c. The case of general boundaries g(t) satisfying (1) will be studied in the follow-up paper [4].

We shall take a route via potential theory for killed random walks and construct appropriate harmonic functions. In the most classical case of constant boundaries a positive harmonic function describes, on one hand, the dependence of the distribution of the exit time on the initial state. On the other hand, harmonic functions allow one to perform the corresponding Doob h-transform and to introduce the conditioned process. In the case of square-root boundaries one expects that the same role will be played by the so-called space-time harmonic functions. To give reasons for that we describe next some known results for the Brownian motion.

For a Brownian motion B(t) killed at crossing $g_{a,b}$ such functions have been constructed by Novikov [12]. Using these functions he obtained explicit expressions for moments of exit times. We now define that functions, since they will play a crucial role in our analysis of discrete time random walks. For every $p \in \mathbb{R}$ we set

$$\psi_p(x) := e^{x^2/4} D_p(x) \text{ and } V_p(x,t) := t^{p/2} \psi_p\left(\frac{x}{\sqrt{t}}\right), \quad t > 0,$$
 (4)

where $D_p(z)$ is the parabolic cylinder function. We also set

$$V_p(x,0) = x^p \quad \text{for} \quad x > 0. \tag{5}$$

Novikov has shown that the processes $V_p(a + B(t), b + t)$ are martingales.

Let p(c) be the minimal positive root of the mapping $p \mapsto \psi_p(c)$. This implies that $V_{p(c)}(c\sqrt{t},t) = 0$ and $V_{p(c)}(x,t) > 0$ for all $x > c\sqrt{t}$. Combining this with the mentioned above martingale property of $V_{p(c)}$, we conclude that

$$V_{p(c)}(a,b) = \mathbf{E}[V_{p(c)}(a+B(t),b+t);T_{a,b}^{(bm)} > t], \quad t > 0,$$
(6)

where

$$T_{a,b}^{(bm)} = \inf\{t > 0 : a + B(t) \le c\sqrt{b+t}\}.$$

In other words, $V_{p(c)}$ is a positive space-time harmonic function for B(t) killed at $T_{a,b}^{(bm)}$. Equivalently, the process $V_{p(c)}(a + B(t), b + t) \mathbb{1}_{\{T_{a,b}^{(bm)} > t\}}$ is a positive martingale. We note also that the space-time harmonicity is equivalent to the 'standard' harmonicity for the two-dimensional process (B(t), t) in the sense

$$\frac{\partial V_{p(c)}(x,t)}{\partial t} + \frac{1}{2} \frac{\partial^2 V_{p(c)}(x,t)}{\partial x^2} = 0,$$

$$V_{p(c)}(c\sqrt{t},t) = 0,$$

see (14) below. The importance of the space-time harmonic function $V_{p(c)}$ becomes clear from the relation

$$\mathbf{P}(T_{a,b}^{(bm)} > t) \sim \varkappa(c) \frac{V_{p(c)}(a,b)}{t^{p(c)/2}} \quad \text{as } t \to \infty,$$
(7)

where $\varkappa(c) > 0$. This relation has been shown by Uchiyama [15].

Our first result deals with space-time harmonic functions for discrete time random walks killed at down-crossing of the boundary $g_{a,b}$.

Theorem 1. Assume that $\mathbf{E}|X|^{2+\delta} < \infty$ for some $\delta > 0$ and $\mathbf{E}|X|^{p(c)} < \infty$, where p(c) is the minimal positive root of $p \mapsto \psi_p(c)$. Then the function

$$W(a,b) := \lim_{n \to \infty} \mathbf{E}[V_{p(c)}(a + S(n), b + n); T_{a,b} > n]$$

is well-defined and satisfies

$$W(a,b) = \mathbf{E}[W(a+S(n),b+n);T_{a,b} > n], \quad n \ge 1.$$
(8)

Moreover, the mapping $a \mapsto W(a, b)$ is monotone increasing and

$$W(x,t) \sim V_{p(c)}(x,t)$$
 as $x, t \to \infty$ and $\frac{x}{\sqrt{t}} \ge c + \varepsilon$ with $\varepsilon > 0$.

It is immediate from the definition of the stopping time $T_{a,b}$ that (8) is equivalent to

$$W(a,b) = \mathbf{E}\left[W(a+S(n),b+n);\min_{k\leq n}\left(a+S(k)-c\sqrt{n+b}\right) > 0\right].$$

Thus, also for the function W, we can interpret the variable a as a spatial (starting) point and b as a time shift for the boundary $c\sqrt{t}$. Equation (8) is a discrete-time analogue of (6) and it can be viewed as a standard discrete harmonicity for 2-dimensional process (S(n), n) killed at $T_{a,b}$.

The space-time harmonic function is intrinsically connected to the stopping time $T_{a,b}$. We underline this connection by the following result.

Corollary 2. Under the conditions of Theorem 1,

$$\mathbf{P}(T_{a,b} > n) \le M \frac{W(a,b)}{n^{p(c)/2}}, \quad n \ge b,$$

with some constant M which is independent of a and b. (This constant may depend on c and on the distribution of $X_{1.}$)

To prove Theorem 1 we first construct a family of positive supermartingales for S(n) killed at $T_{a,b}$. To this end we modify appropriately functions $V_p(x,t)$ for p < p(c). (To understand, why V_p with p < p(c) is a good starting point, we look again at the Brownian motion. Recalling that $V_p(a + B(t), b + t)$ is a martingale and that $V_p(c\sqrt{t},t) > 0$ for every p < p(c) and using the optional stopping theorem, we infer that $V_p(a + B(t), b + t) 1\{T_{a,b}^{(bm)} > t\}$ is a supermartingale.) Applying the Optimal Stopping Theorem to that martingales, we derive suboptimal upper bounds for the tail of $T_{a,b}$, see Lemma 19 below. Having that bound we construct W(a, b)by applying the strategy from [7], where harmonic functions for multidimensional walks in cones have been constructed.

Construction of supermartingales in the proof of Theorem 1 is lengthy but uses only quite elementary calculations. A disadvantage of this approach is a slightly stronger moment condition $\mathbf{E}|X|^{2+\delta} < \infty$. This moment condition will be relaxed in the follow-up paper [4].

We now state our result on the tail behaviour of the stopping time T_q .

Theorem 3. Assume that all the conditions of Theorem 1 are valid. Then, for all fixed a, b such that $a > c\sqrt{b}$,

$$\mathbf{P}(T_{a,b} > n) \sim \varkappa(c) \frac{W(a,b)}{n^{p(c)/2}},\tag{9}$$

where $\varkappa(c)$ is the same constant as in (7).

Comparing this with (3) we see that our result covers boundaries $g_{a,b}$ for all possible values of c. Furthermore, we show that a slowly varying function L_g can be replaced by a 'constant' W(a, b). A certain weakness of our result consists in

the moment condition $\mathbf{E}|X|^{2+\delta}$, which is relevant in the case $p(c) \leq 2$. Greenwood and Perkins impose a milder assumption $\mathbf{E}X^2 \log(1+|X|) < \infty$.

We conclude the introduction by discussing the moment assumption $\mathbf{E}|X|^{p(c)} < \infty$. We assume that c is such that p(c) > 2. The following, rather standard, example shows that that moment condition is the minimal one for such values of c. Assume that

$$\mathbf{P}(X > t) \ge \theta_0 t^{-\beta}, \quad t \ge 1 \tag{10}$$

with some $\theta_0 > 0$ and some $\beta \in (2, p(c))$. Clearly, $\mathbf{E}|X|^{p(c)}$ is then infinite. It is clear that

$$\mathbf{P}(T_{a,b} > n) \ge \mathbf{P}(X_1 > A\sqrt{n})\mathbf{P}\left(\min_{k < n} S(k) > \max_{k \le n} g_{a,b}(k) - A\sqrt{n}\right)$$

for every A > 0. It follows from (2) that

$$\frac{1}{\sqrt{n}}\max_{k\le n}g_{a,b}(k)\to c$$

Therefore, by the functional central limit theorem,

$$\mathbf{P}\left(\min_{k< n} S(k) > \max_{k\leq n} g_{a,b}(k) - A\sqrt{n}\right) \to \mathbf{P}\left(\min_{s\leq 1} B(s) > c - A\right)$$

and the probability on the right hand side is positive for all A > c. Taking A = c+1, we infer that there exists a constant θ_1 such that

$$\mathbf{P}(T_{a,b} > n) \ge \theta_1 n^{-\beta/2}.$$

Thus, the statement of Theorem 3 can not hold for walks satisfying (10). As we have mentioned before, this extra moment condition is not needed in the case of two-sided boundaries, which can be explained as follows. The second half of the boundary prevents appearance of big jumps as it was described above. Thus, one can think of a random walk with truncated increments.

2. Properties of the functions V_p

We start by collecting some simple but quite useful analytical properties of the functions $V_p(x, t)$.

The function $\psi_p(x) = e^{x^2/4} D_p(x)$ is defined for all real x and p. For p < 0 this function admits the following integral representation

$$\psi_p(x) = \frac{1}{\Gamma(-p)} \int_0^{+\infty} \exp\left(-xs - \frac{s^2}{2}\right) s^{-p-1} ds > 0.$$

This implies immediately that ψ_p has no real roots in the case p < 0. It is also known that ψ_p satisfies the following recurrence relation:

$$\psi_p(x) = x\psi_{p-1}(x) + (1-p)\psi_{p-2}(x). \tag{11}$$

This relation allows one to get an integral representation for ψ_p also in the case $p \ge 0$.

The derivative of ψ_p satisfies

$$\psi'_{p}(x) = p\psi_{p-1}(x). \tag{12}$$

Recalling definitions (5) and (4) of functions V_p and using (12), we obtain

$$\frac{\partial}{\partial x}V_p(x,t) = pV_{p-1}(x,t).$$
(13)

Combining (11) and (12), one gets easily the relation

$$\frac{\partial}{\partial t}V_p(x,t) = -\frac{1}{2}\frac{\partial^2}{\partial x^2}V_p(x,t) = -\frac{1}{2}p(p-1)V_{p-2}(x,t).$$
(14)

It is also known that

$$\psi_p(x) \sim x^p \quad \text{as } x \to \infty.$$
 (15)

All properties presented above may be found in [14], see also appendices in [13] and [15].

The following properties of V_p are proven in [3].

Lemma 4. It holds that

$$V_p(x,t)$$
 is increasing in $x \in (c\sqrt{t},\infty)$ for all $p \in [0,p(c)]$ (16)

and

$$V_{p(c)}(c\sqrt{t},t) = 0 < V_{p(c)}(x,t) \quad for \ all \quad x > c\sqrt{t}.$$
 (17)

In addition, for all $p \in \mathbb{R}$,

$$\sup_{x>z\sqrt{t}} |V_p(x,t)/x^p - 1| \to 0 \quad as \quad z \to \infty.$$
(18)

Furthermore, for each $\gamma > 0$ there exists a finite positive constant $C_0(\gamma) = C_0(\gamma, c)$ such that

$$C_0(\gamma)(x - c\sqrt{t})^{p(c)} \ge V_{p(c)}(x, t) \ge (x - c\sqrt{t})^{p(c)}/C_0(\gamma)$$
(19)

for all x, t, c and γ satisfying conditions:

$$x - c\sqrt{t} > \gamma\sqrt{t}$$
 with $t \ge 0.$ (20)

Since $c \mapsto p(c)$ is monotone increasing, we can invert this mapping. The inverse mapping will be denoted by $p \to c(p)$. It is then clear that c(p) is the largest zero of the function $\psi_p(z)$.

Lemma 5. For every p > 0 there exists a constant C = C(p) > 0 such that for $x > c(p)\sqrt{t}$ one has

$$C^{-1}V_{p-1}(x,t)(x-c(p)\sqrt{t}) \le V_p(x,t) \le CV_{p-1}(x,t)(x-c(p)\sqrt{t}).$$
(21)

If p > 1 there exists a constant A(p) such that

$$\frac{1}{A(p)}x^{p-1}(x-c(p)\sqrt{t}) \le V_p(x,t) \le A(p)x^{p-1}(x-c(p)\sqrt{t})$$

for all $x \ge c(p)\sqrt{t}$.

Proof. Since $\psi_p(x) \sim x^p, x \to \infty$ for all p, there exists sufficiently large A such that (21) holds for $x > A\sqrt{t}$.

Consider now the case $x \in (c(p)\sqrt{t}, A\sqrt{t})$. By the mean value theorem,

$$V_p(x,t) = V_p(x,t) - V_p(c(p)\sqrt{t},t)$$

= $\frac{\partial}{\partial\theta}V_p(\theta,t)(x-c(p)\sqrt{t}) = pV_{p-1}(\theta,t)(x-c(p)\sqrt{t})$

for some $\theta \in (c(p)\sqrt{t}, x)$.

By the definition of V_{p-1} ,

$$V_{p-1}(y,t) = t^{(p-1)/2} \psi_{p-1}(y/\sqrt{t})$$

Noting that $1/C(p, A) < \psi_{p-1}(u) < C(p, A)$ for all $u \in [c(p), A]$, we conclude that the ratio $V_{p-1}(y, t)/V_{p-1}(x, t)$ is bounded below and above by some positive constants, which depend on A and p only. This completes the proof of (21).

To prove the second claim we notice that c(p-1) < c(p). This allows us to apply (19) to the function V_{p-1} . This gives us the estimates

$$\frac{1}{A_1(p)}(x-c(p-1)\sqrt{t})^{p-1} \le V_{p-1}(x,t) \le A_1(p)(x-c(p-1)\sqrt{t})^{p-1}$$

for $x > c(p)\sqrt{t}$. For all $x \ge c(p)\sqrt{t}$ one has

$$x - \frac{c(p-1)^{+}}{c(p)}x \le x - c(p-1)\sqrt{t} \le x + \frac{c(p-1)^{-}}{c(p)}x,$$

where as usual $x^+ = \max(x, 0)$ and $x^- = \max(-x, 0)$. Using these estimates we obtain

$$\frac{1}{A_2(p)}x^{p-1} \le V_{p-1}(x,t) \le A_2(p)x^{p-1}.$$

Thus, the proof is complete.

Lemma 6. For every q < 0 and every T > 0 there exists a constant A > 0 such that

$$V_q(x,t) \le A(x+T\sqrt{t})^q$$

for all $x > -T\sqrt{t}$.

Proof. By the definition,

$$V_q(x,t) = t^{q/2} \psi_q\left(\frac{x}{\sqrt{t}}\right)$$

and

$$\psi_q(z) = \frac{1}{\Gamma(-q)} \int_0^\infty \exp\left(-zs - \frac{s^2}{2}\right) s^{-q-1} ds.$$

It is then clear that

$$\frac{d}{dz}\psi_q(z) = -\frac{1}{\Gamma(-q)}\int_0^\infty \exp\left(-zs - \frac{s^2}{2}\right)s^{-q}ds < 0.$$

Thus, $\psi_q(z)$ is decreasing. Recall that $\psi_q(z) \sim z^q$ as $z \to \infty$. Therefore, there are constants R and C_0 such that

$$\psi_q(z) \le C_0 \left(1_{[-T,R)}(z) + z^q 1_{[R,\infty)}(z) \right)$$

and consequently,

$$V_q(x,t) \le C_0 \left(t^{q/2} \mathbb{1}_{[-T\sqrt{t}, R\sqrt{t})}(x) + x^q \mathbb{1}_{[R\sqrt{t}, \infty)}(x) \right).$$
(22)

If $x \leq R\sqrt{t}$ then $(x + T\sqrt{t})^q \geq (R + T)^q \cdot t^{q/2}$. Furthermore, for $t > R\sqrt{t}$ we have $(x + T\sqrt{t})^q \geq (1 + \frac{T}{R})^q x^q$. Combining this with (22), we get the desired estimate.

Lemma 7. For every $p \in (0,1)$ there exists A(p) such that

$$V_p(x,t) \le A(p)(x-c(p)\sqrt{t})^p$$

for all $x \ge c(p)\sqrt{t}$, t > 0.

Proof. Using (21) and the fact that $V_{p-1}(x,t)$ is decreasing in x we obtain

$$V_p(x,t) \le A(p)V_{p-1}(c(p)\sqrt{t},t) \cdot (x-c(p)\sqrt{t}).$$

Applying Lemma 6 to V_{p-1} , we get

$$V_p(x,t) \le A_0(p)t^{(p-1)/2}(x-c(p)\sqrt{t}) = A_0(p)(x-c(p)\sqrt{t})^p \left(\frac{x-c(p)\sqrt{t}}{\sqrt{t}}\right)^{1-p}.$$

Noting that $\left(\frac{x-c(p)\sqrt{t}}{\sqrt{t}}\right)^{1-p} \leq 1$ for $x \in [c(p)\sqrt{t}, (c(p)+1)\sqrt{t}]$ we complete the proof.

3. Constructions of supermartingales

Our constructions of supermartingales for discrete time random walks will use functions V_p which are martingales for the Brownian motion. Set

$$\overline{V}_p(x,t) = \begin{cases} V_p(x,t), & x \ge c(p)\sqrt{t}, \\ 0, & x < c(p)\sqrt{t}. \end{cases}$$

In this section we will use an approach which is similar to the method suggested by McConnell [11], who has constructed supermartingales and derived bounds for exit times from cones by multidimensional walks.

We first bound the mean drift of the process $\overline{V}_p(a + S(n), b + n)$ for p > 1 and then for $p \in (0, 1)$.

Lemma 8 (p > 1). Assume that $\mathbf{E}|X|^p < \infty$ and $\mathbf{E}|X|^{2+\delta} < \infty$ for some $\delta > 0$. If p > 2 then there exists a constant C such that, for all $x > c(p)\sqrt{t}$,

$$\left|\mathbf{E}\overline{V}_{p}(x+X,t+1) - \overline{V}_{p}(x,t)\right| \le C\left(\frac{(1+x)^{p-1}}{(1+x-c(p)\sqrt{t})^{p-1}} + (1+x)^{p-3}\right).$$

If $p \in (1,2]$ then there exists a constant C such that, for all $x > c(p)\sqrt{t}$,

$$\left|\mathbf{E}\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t)\right| \le C\left(\frac{(1+x)^{p-1}}{(1+x-c(p)\sqrt{t})^{1+\delta}}\right)$$

Proof. Both statements are obvious for $x \in (c(p)\sqrt{t}, c(p)\sqrt{t+1}]$. From now on we assume that $x > c(p)\sqrt{t+1}$.

First we notice that

$$V_p(x+z,t+1) - V_p(x,t)$$

= $[V_p(x+z,t+1) - V_p(x+z,t)] + [V_p(x+z,t) - V_p(x,t)].$

Applying now the Taylor formula, we get the equalities

$$\begin{aligned} V_p(x+z,t+1) - V_p(x+z,t) \\ &= \frac{\partial}{\partial t} V_p(x+z,t) + \frac{1}{2} \frac{\partial^2}{\partial t^2} V_p(x+z,t+\phi) \\ &= \frac{\partial}{\partial t} V_p(x,t) + z \frac{\partial^2}{\partial x \partial t} V_p(x+\psi z,t) + \frac{1}{2} \frac{\partial^2}{\partial t^2} V_p(x+z,t+\phi) \end{aligned}$$

for some $\phi, \psi \in (0, 1)$. Similarly,

$$\begin{split} V_p(x+z,t) &- V_p(x,t) \\ &= z \frac{\partial}{\partial x} V_p(x,t) + z^2 \frac{1}{2} \frac{\partial^2}{\partial x^2} V_p(x,t) + z^3 \frac{1}{6} \frac{\partial^3}{\partial x^3} V_p(x+\theta z,t) \end{split}$$

$$V_{p}(x+z,t+1) - V_{p}(x,t)$$

$$= zpV_{p-1}(x,t) + \frac{z^{2}}{2}p(p-1)V_{p-2}(x,t) - \frac{1}{2}p(p-1)V_{p-2}(x,t) + R_{p}(x,t;z), \quad (23)$$

where

$$R_{p}(x,t;z) = \frac{z^{3}}{6}p(p-1)(p-2)V_{p-3}(x+\theta z,t) -\frac{z}{2}p(p-1)(p-2)V_{p-3}(x+\psi z,t) +\frac{1}{8}p(p-1)(p-2)(p-3)V_{p-4}(x,t+\phi), \text{ for } \phi, \psi, \theta \in (0,1).$$

Using the equality (23), we have

$$\begin{split} \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| &\leq (x-c(p)\sqrt{t+1})/2 \right] \\ &= \mathbf{E} \left[V_p(x+X,t+1) - V_p(x,t); |X| &\leq (x-c(p)\sqrt{t+1})/2 \right] \\ &= -\frac{1}{2} p(p-1) V_{p-2}(x,t) \mathbf{P}(|X| &\leq (x-c(p)\sqrt{t+1})/2) \\ &+ p V_{p-1}(x,t) \mathbf{E}[X; |X| &\leq (x-c(p)\sqrt{t+1})/2] \\ &+ \frac{1}{2} p(p-1) V_{p-2}(x,t) \mathbf{E}[X^2; |X| &\leq (x-c(p)\sqrt{t+1})/2] \\ &+ \mathbf{E}[R_p(x,t;X); |X| &\leq (x-c(p)\sqrt{t+1})/2]. \end{split}$$

Recall that p > 1 is equivalent to c(p) > 0. It follows from (19) that $V_{p-1}(x,t) = O(x^{p-1})$ and $V_{p-2}(x,t) = O(x^{p-2})$ uniformly in $x > c(p)\sqrt{t}$.

To bound $\mathbf{E}[R_p(x,t;X); |X| \le (x-c(p)\sqrt{t})/2]$, we notice that, by (19), $V_{p-3}(x+\theta z,t) = O((1+x)^{p-3})$ and $V_{p-4}(x,t+\phi) = O((1+x)^{p-4})$ for $x > c(p)\sqrt{t+1}$. These estimates imply that

$$|\mathbf{E}[R_p(x,t;X);|X| \le (x-c(p)\sqrt{t+1})/2]| \le C\left((1+x)^{p-4} + (1+x)^{p-3}\mathbf{E}[|X|^3;|X| \le (x-c(p)\sqrt{t+1})/2]\right).$$

Recalling that $\mathbf{E}X = 0$ and $\mathbf{E}X^2 = 1$ and using once again (14), we infer that

$$\mathbf{E}\left[\overline{V}_{p}(x+X,t+1)-\overline{V}_{p}(x,t);|X| \leq (x-c(p)\sqrt{t+1})/2\right] \\
\leq C\left(x^{p-2}\mathbf{P}(|X| > (x-c(p)\sqrt{t+1})/2) \\
+ x^{p-1}\mathbf{E}[X;|X| > (x-c(p)\sqrt{t+1})/2] \\
+ x^{p-2}\mathbf{E}[X^{2};|X| > (x-c(p)\sqrt{t+1})/2] \\
+ (1+x)^{p-4} + (1+x)^{p-3}\mathbf{E}[|X|^{3};|X| \leq (x-c(p)\sqrt{t+1})/2]\right).$$
(24)

It is immediate from (14) that $\overline{V}_p(x,t)$ increases in x and decreases in t. This observation implies that $\overline{V}_p(x+X,t+1) \leq \overline{V}_p(x,t+1) \leq \overline{V}_p(x,t)$ for all $X < -(x-c(p)\sqrt{t+1})/2$. This implies that

$$\begin{split} & \left| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| > (x-c(p)\sqrt{t+1})/2 \right] \right| \\ & \leq \mathbf{E} \left[\overline{V}_p(x+X,t+1); X > (x-c(p)\sqrt{t+1})/2 \right] \\ & \quad + \overline{V}_p(x,t) \mathbf{P}(|X| > (x-c(p)\sqrt{t+1})/2). \end{split}$$

Using Lemma 5,

$$\begin{aligned} \left| \mathbf{E} \left[\overline{V}_{p}(x+X,t+1) - \overline{V}_{p}(x,t); |X| > (x-c(p)\sqrt{t+1})/2 \right] \right| \\ &\leq A(p) \mathbf{E} \left[(x+X-c(p)\sqrt{t}) \cdot (x+X)^{p-1}; X > (x-c(p)\sqrt{t+1})/2 \right] \\ &\quad + A(p)(x-c(p)\sqrt{t})x^{p-1} \mathbf{P}(|X| > (x-c(p)\sqrt{t+1})/2) \\ &\leq 3 \cdot 2^{p-1}A(p) \mathbf{E} \left[|X|(x^{p-1}+|X|^{p-1}); X > (x-c(p)\sqrt{t+1})/2 \right] \\ &\quad + 2A(p)(x-c(p)\sqrt{t})x^{p-1} \mathbf{P}(|X| > (x-c(p)\sqrt{t+1})/2) \\ &\leq C \Big(\mathbf{E} \left[|X|^{p}; X > (x-c(p)\sqrt{t+1})/2 \right] + x^{p-1} \mathbf{E} \left[|X|; X > (x-c(p)\sqrt{t+1})/2 \right] \\ &\quad + (x-c(p)\sqrt{t})x^{p-1} \mathbf{P}(|X| > (x-c(p)\sqrt{t+1})/2) \Big). \end{aligned}$$

Assume that p > 2. Due to our moment assumption $\mathbf{E}|X|^p < \infty$,

$$\mathbf{P}(|X| > (x - c(p)\sqrt{t+1})/2) \le C(1 + x - c(p)\sqrt{t})^{-p},$$
(26)

$$\mathbf{E}[|X|; |X| > (x - c(p)\sqrt{t+1})/2] \le C(1 + x - c(p)\sqrt{t})^{1-p},$$
(27)

$$\mathbf{E}[X^2; |X| > (x - c(p)\sqrt{t+1})/2] \le (1 + x - c(p)\sqrt{t})^{2-p},$$
(28)

and

$$\mathbf{E}[|X|^{p}; X > (x - c(p)\sqrt{t+1})/2] \le C.$$
(29)

If $p \ge 3$ then $\mathbf{E}[|X|^3; |X| \le (x - c(p)\sqrt{t})/2]$ is bounded. Plugging in (26) – (29) into (24) and (25), we thus obtain

$$\begin{split} \left| \mathbf{E} \overline{V}_p(x + X, t + 1) - \overline{V}_p(x, t) \right| \\ &\leq \left| \mathbf{E} \left[\overline{V}_p(x + X, t + 1) - \overline{V}_p(x, t); |X| \leq (x - c(p)\sqrt{t + 1})/2 \right] \right| \\ &+ \left| \mathbf{E} \left[\overline{V}_p(x + X, t + 1) - \overline{V}_p(x, t); |X| > (x - c(p)\sqrt{t + 1})/2 \right] \right| \\ &\leq C \Big(\frac{x^{p-2}}{(1 + x - c(p)\sqrt{t})^p} + \frac{x^{p-1}}{(1 + x - c(p)\sqrt{t})^{p-1}} + \frac{x^{p-2}}{(1 + x - c(p)\sqrt{t})^{p-2}} \\ &+ (1 + x)^{p-4} + (1 + x)^{p-3} + 1 + \frac{(x - c(p)\sqrt{t})x^{p-1}}{(1 + x - c(p)\sqrt{t})^p} \Big) \\ &\leq C \left(\frac{(1 + x)^{p-4}}{(1 + x - c(p)\sqrt{t})^{p-1}} + (1 + x)^{p-3} \right), \end{split}$$

which is the desired bound for $p \ge 3$.

If $p \in (2,3)$ then

$$\mathbf{E}[|X|^3; |X| \le (x - c(p)\sqrt{t+1})/2] \le (x - c(p)\sqrt{t+1})^{3-p}\mathbf{E}|X|^p \le C(1 + x - c(p)\sqrt{t})^{3-p}.$$

Applying this together with (26) - (29) to (24) and (25), we find for $p \in (2,3)$

$$\begin{split} \mathbf{E}V_p(x+X,t+1) &- V_p(x,t) \big| \\ &\leq \big| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| \leq (x-c(p)\sqrt{t+1})/2 \right] \big| \\ &+ \big| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| > (x-c(p)\sqrt{t+1})/2 \right] \big| \\ &\leq C \frac{x^{p-1}}{(1+x-c(p)\sqrt{t})^{p-1}}. \end{split}$$

This completes the proof of the first claim.

If $1 then we assume that <math>\mathbf{E}|X|^{2+\delta} < \infty$ for some $\delta > 0$. In this case, instead of (26) – (28), we have

$$\mathbf{P}(|X| > (x - c(p)\sqrt{t+1})/2) \le C(1 + x - c(p)\sqrt{t})^{-(2+\delta)},$$
(30)

$$\mathbf{E}[|X|^r; X > (x - c(p)\sqrt{t+1})/2] \le C(1 + x - c(p)\sqrt{t})^{r-2-\delta} \quad \text{for all } r \le 2 \quad (31)$$

and

$$\mathbf{E}[|X|^{3}; X \le (x - c(p)\sqrt{t+1})/2] \le (x - c(p)\sqrt{t})^{1-\delta} \mathbf{E}\left[|X|^{2+\delta}\right] \le C(1 + x - c(p)\sqrt{t})^{1-\delta}.$$
(32)

Entering (30) - (32) into (24) and (25), we thus obtain for $p \in (1, 2]$

$$\begin{split} \mathbf{E}\overline{V}_p(x+X,t+1) &- \overline{V}_p(x,t) \big| \\ &\leq \big| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| \leq (x-c(p)\sqrt{t+1})/2 \right] \big| \\ &+ \big| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); |X| > (x-c(p)\sqrt{t+1})/2 \right] \big| \\ &\leq C \frac{(1+x)^{p-1}}{(1+x-c(p)\sqrt{t})^{1+\delta}}, \end{split}$$

which is the second claim.

Lemma 9 $(p \in (0,1))$. Assume that $\mathbf{E}|X|^{2+\delta} < \infty$ for some $\delta > 0$. If $p \in (0,1)$ then, as $x - c(p)\sqrt{t} \to \infty$,

$$\left|\mathbf{E}\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t)\right| = o\left((1+x-c(p)\sqrt{t})^{p-2-\delta}\right).$$

Proof. We shall use the same decomposition as in the proof of Lemma 8. Recalling that $\overline{V}_p(x,t)$ is decreasing in t and increasing in x, we obtain

$$\begin{split} E_1 &:= \left| \mathbf{E} \left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); X > (x-c(p)\sqrt{t})/2 \right] \right| \\ &\leq \mathbf{E} \left[\overline{V}_p(x+X,t); X > (x-c(p)\sqrt{t})/2 \right] + \overline{V}_p(x,t) \mathbf{P}(X > (x-c(p)\sqrt{t})/2) \\ &\leq 2 \mathbf{E} \left[\overline{V}_p(x+X,t); X > (x-c(p)\sqrt{t})/2 \right] \\ &\leq 2 A(p) \mathbf{E} \left[(x+X-c(p)\sqrt{t})^p; X > (x-c(p)\sqrt{t})/2 \right] \\ &\leq 6 A(p) \mathbf{E} \left[|X|^p; X > (x-c(p)\sqrt{t})/2 \right], \end{split}$$

where the third inequality follows from Lemma 7. Applying now the Chebyshev inequality, we conclude that

$$E_1 = o\left((1 + x - c(p)\sqrt{t})^{p-2-\delta} \right).$$
(33)

On the event $X < -(x-c(p)\sqrt{t})/2$ we have, due to Lemma 7, $\overline{V}_p(x+X,t+1) \leq C(x-c(p)\sqrt{t})^p$ and $\overline{V}_p(x,t) \leq C(x-c(p)\sqrt{t})^p$. Therefore,

$$\mathbf{E}\left[\overline{V}_p(x+X,t+1) - \overline{V}_p(x,t); X < -(x-c(p)\sqrt{t})/2\right]$$
$$= o\left((1+x-c(p)\sqrt{t})^{p-2-\delta}\right). \quad (34)$$

In the event $|X| \leq (x-c(p)\sqrt{t})/2$ we shall use the decomposition in (24). Combining Lemma 6 with the Chebyshev inequality, we have

$$V_{p-2}(x,t)\mathbf{P}(|X| > (x-c(p)\sqrt{t})/2) \le C(1+x-c(p)\sqrt{t})^{p-2}\mathbf{P}(|X| > (x-c(p)\sqrt{t})/2)$$
$$= o\left((1+x-c(p)\sqrt{t})^{p-4-\delta}\right),$$
$$V_{p-1}(x,t)\mathbf{E}\left[X; |X| > (x-c(p)\sqrt{t})/2\right] = o\left((1+x-c(p)\sqrt{t})^{p-2-\delta}\right)$$

and

$$V_{p-2}(x,t)\mathbf{E}\left[X^2; |X| > (x-c(p)\sqrt{t})/2\right] = o\left((1+x-c(p)\sqrt{t})^{p-2-\delta}\right).$$

Similarly,

$$\mathbf{E}\left[R_p(x,t,X); |X| \le (x - c(p)\sqrt{t})/2\right] = o\left((1 + x - c(p)\sqrt{t})^{p-2-\delta}\right).$$

This completes the proof.

We have bounded the absolute value of the mean drift. To construct a supermartingale we will correct \overline{V}_p with a term of order $x^{p-\delta}$ with some sufficiently small δ . In the next lemma we estimate the drift of this term.

Lemma 10 (p > 1). Assume that $\mathbf{E}|X|^p < \infty$ and $\mathbf{E}|X|^{2+\delta} < \infty$ for some $\delta > 0$.

(i) If p > 3, then for all $x > c(p)\sqrt{t}$ there is constant C > 0 depending only on p and δ such that

$$\begin{split} \mathbf{E}\left[|x+X|^{p-\delta}-x^{p-\delta}\right] \\ \geq \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^p} + x^{p-\delta-3}\right). \end{split}$$

(ii) If $p \in (2,3]$, then for all $x > c(p)\sqrt{t}$ there is a constant C > 0 depending only on p and δ such that

$$\mathbf{E}\left[|x+X|^{p-\delta} - x^{p-\delta}\right] \ge \frac{(p-\delta)(p-\delta-1)}{2} x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^p} + (x-c(p)\sqrt{t})^{p-2\delta-2}\right).$$

(iii) If $p \in (1, 2]$, then for all $x > c(p)\sqrt{t}$ there is a constant C > 0 depending only on p and δ such that

$$\mathbf{E}\left[|x+X|^{p-\delta}-x^{p-\delta}\right] \ge \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^{2+\delta}} + (x-c(p)\sqrt{t})^{p-2\delta-2}\right).$$

Proof. We split the expectation into three parts

$$\mathbf{E} \left[|x + X|^{p-\delta} - x^{p-\delta} \right] \\
= \mathbf{E} \left[|x + X|^{p-\delta} - x^{p-\delta}; |X| \le (x - c(p)\sqrt{t})/2 \right] \\
+ \mathbf{E} \left[|x + X|^{p-\delta} - x^{p-\delta}; X > (x - c(p)\sqrt{t})/2 \right] \\
+ \mathbf{E} \left[|x + X|^{p-\delta} - x^{p-\delta}; X < -(x - c(p)\sqrt{t})/2 \right] \\
=: A_1 + A_2 + A_3.$$
(35)

We find immediately that $A_2 \ge 0$. For A_3 , notice that $|x + X|^p \ge 0$ and thus due to (26) (if p > 2) and (30) (if 1), there is a constant <math>C > 0 such that

$$A_3 \ge -x^{p-\delta} \mathbf{P}(X < -(x - c(p)\sqrt{t})/2) \ge -C \frac{x^{p-\delta}}{(x - c(p)\sqrt{t})^p}.$$
 (36)

Finally, for A_1 , notice that due to $X \ge -(x - c(p)\sqrt{t})/2$, $x + X \ge 0$ so we can remove the absolutes. By Taylor's formula, there is a $\theta = \theta(x, X) \in [0, 1]$ such that

$$A_{1} = \mathbf{E} \left[|x + X|^{p-\delta} - x^{p-\delta}; |X| \le (x - c(p)\sqrt{t})/2 \right]$$

=
$$\mathbf{E} \left[(p-\delta)x^{p-\delta-1} \cdot X + \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} \cdot X^{2} + \frac{(p-\delta)(p-\delta-1)(p-\delta-2)}{6}(x+\theta X)^{p-\delta-3} \cdot X^{3}; |X| \le (x - c(p)\sqrt{t})/2 \right].$$
 (37)

Consider the terms in this expression separately. Firstly, due to $\mathbf{E}X = 0$ and (27) and (31), there is a constant C > 0 such that

$$\mathbf{E}\left[(p-\delta)x^{p-\delta-1}\cdot X; |X| \le (x-c(p)\sqrt{t})/2\right] \\
= -(p-\delta)x^{p-\delta-1}\mathbf{E}\left[X; |X| > (x-c(p)\sqrt{t})/2\right] \\
\ge -C\frac{x^{p-\delta-1}}{(x-c(p)\sqrt{t})^{p-1}} \ge -C\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^{p}}.$$
(38)

Secondly, due to $\mathbf{E} X^2 = 1$ and (28), there is a constant C > 0 such that in the case p > 2

$$\mathbf{E}\left[\frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} \cdot X^{2}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
= \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} \left(1 - \mathbf{E}\left[X^{2}; |X| > (x-c(p)\sqrt{t})/2\right]\right) \\
\geq \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\frac{x^{p-\delta-2}}{(x-c(p)\sqrt{t})^{p-2}} \\
\geq \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^{p}},$$
(39)

while for 1 using (31), there is a constant <math>C > 0 such that

$$\mathbf{E}\left[\frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} \cdot X^{2}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
= \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} \left(1 - \mathbf{E}\left[X^{2}; |X| > (x-c(p)\sqrt{t})/2\right]\right) \\
\geq \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\frac{x^{p-\delta-2}}{(x-c(p)\sqrt{t})^{\delta}} \\
\geq \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^{2+\delta}}.$$
(40)

Thirdly, we notice that if $|X| \leq (x - c(p)\sqrt{t})/2$, then for every $\theta \in [0, 1]$

$$\frac{3}{2}x \ge \frac{3}{2}x - \frac{1}{2}c(p)\sqrt{t} \ge x + \theta X \ge \frac{1}{2}(x + c(p)\sqrt{t}) \ge 0.$$

For p>3, the third moment of X is finite, so there is a constant C>0 depending only on p and δ such that

$$\mathbf{E}\left[\frac{(p-\delta)(p-\delta-1)(p-\delta-2)}{6}(x+\theta X)^{p-\delta-3} \cdot X^{3}; |X| \le (x-c(p)\sqrt{t})/2\right] \\
\ge -\frac{(p-\delta)(p-\delta-1)(p-\delta-2)}{6}\left(\frac{3}{2}x\right)^{p-\delta-3} \mathbf{E}\left[|X|^{3}; |X| \le (x-c(p)\sqrt{t})/2\right] \\
\ge -Cx^{p-\delta-3}.$$
(41)

Combining all results (35) – (39) and (41), we obtain that for p > 3 there is a constant C > 0 depending only on p and δ such that

$$\mathbf{E}[|x+X|^p - x^p] \ge \frac{(p-\delta)(p-\delta-1)}{2} x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^p} + x^{p-\delta-3}\right),$$

If $p \in (2,3]$, then we notice that $p - \delta - 3 < 0$. So if $|X| \le (x - c(p)\sqrt{t})/2$, then

$$0 \le (x + \theta X)^{p - \delta - 3} \le \left(x - (x - c(p)\sqrt{t})/2 \right)^{p - \delta - 3} \le C \left(x - c(p)\sqrt{t} \right)^{p - \delta - 3}$$

and using that $\mathbf{E}|X|^{2+\delta} < \infty$, we obtain

$$\mathbf{E}\left[\frac{(p-\delta)(p-\delta-1)(p-\delta-2)}{6}(x+\theta X)^{p-\delta-3} \cdot X^{3}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
\geq -C\left((x-c(p)\sqrt{t})/2\right)^{p-\delta-3} \mathbf{E}\left[|X|^{3}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
\geq -C\left(x-c(p)\sqrt{t}\right)^{p-\delta-3} \mathbf{E}\left[|X|^{3}\left(\frac{(x-c(p)\sqrt{t})/2}{|X|}\right)^{1-\delta}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
\geq -C\left(x-c(p)\sqrt{t}\right)^{p-2\delta-2} \mathbf{E}\left[|X|^{2+\delta}; |X| \leq (x-c(p)\sqrt{t})/2\right] \\
\geq -C\left(x-c(p)\sqrt{t}\right)^{p-2\delta-2}.$$
(42)

Combining all results (35) – (39) and (42), for $p \in (2,3]$ we obtain that there is a constant C > 0 depending only on p and δ such that

$$\mathbf{E}[|x+X|^{p} - x^{p}] \ge \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(1+x-c(p)\sqrt{t})^{p}} + (x-c(p)\sqrt{t})^{p-2\delta-2}\right).$$

If $p \in (1, 2]$, then we notice that $p - \delta - 3 . So analogous to the case <math>p \in (2, 3]$, there is some constant C > 0 such depending only on p and δ such that

$$\mathbf{E}\left[\frac{(p-\delta)(p-\delta-1)(p-\delta-2)}{6}(x+\theta X)^{p-\delta-3}\cdot X^3; |X| \le (x-c(p)\sqrt{t})/2\right]$$
$$\ge -C\left(x-c(p)\sqrt{t}\right)^{p-2\delta-2}.$$
(43)

Combining all results (35) - (38), (40) and (43), for $p \in (1, 2]$ we obtain that there is a constant C > 0 depending only on p and δ such that

$$\mathbf{E}[|x+X|^{p} - x^{p}] \ge \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2} - C\left(\frac{x^{p-\delta}}{(x-c(p)\sqrt{t})^{2+\delta}} + (x-c(p)\sqrt{t})^{p-2\delta-2}\right),$$

thereby completing the proof.

Remark 11. If $t \to \infty$ and $x \ge (c(p) + \gamma)\sqrt{t}$ for some $\gamma > 0$, then we have

$$\mathbf{E}\left[|x+X|^{p-\delta} - x^{p-\delta}\right] = \frac{(p-\delta)(p-\delta-1)}{2}x^{p-\delta-2}(1+o(1)),$$

as the dominating term was obtained only through equalities.

By Lemma 4, we notice that if $x \ge (c(p) + \gamma)\sqrt{t}$ for some $\gamma > 0$, then

$$V_p(x,t) - C \cdot x^{p-\delta} \ge \frac{1}{A(p,\gamma)} x^p - C x^{p-\delta}$$

For every constant C we can therefore find a constant R(C) such that for every $R \geq R(C)$ one has

$$V_p(x+R,t) - C \cdot |x+R|^{p-\delta} > 0$$
(44)

for all $x \ge (c(p) + \gamma)\sqrt{t}$.

With this in mind, we define for $k \in \mathbb{N}$ and $x \in \mathbb{R}$

$$h(x,t) := \left(\overline{V}_p(x+R,t) - C \cdot |x+R|^{p-\delta}\right)^+.$$

Lemma 12. Let p > 1. For every $\gamma > 0$ there exist constants C and R such that if $x \ge (c(p) + \gamma)\sqrt{t}$ then

$$\mathbf{E}\left[h(x+X,t+1) - h(x,t)\right] \le 0.$$

 \diamond

Proof. There exists a constant C > 0 depending only on p and δ such that

$$\mathbf{E}\left[|x+R+X|^{p-\delta}; X < -(x-c(p)\sqrt{t})/2\right] \\
\leq 2^{p}\left((x+R)^{p-\delta}\mathbf{P}\left(|X| > (x-c(p)\sqrt{t})/2\right) + \mathbf{E}\left[|X|^{p-\delta}; |X| > (x-c(p)\sqrt{t})/2\right]\right) \\
\leq 2^{2p}\left((x+R)^{p-\delta}\frac{\mathbf{E}[|X|^{p}]}{(x-c(p)\sqrt{t})^{p}} + \frac{\mathbf{E}\left[|X|^{p}\right]}{(x-c(p)\sqrt{t})^{\delta}}\right) \\
\leq C\left(\frac{(x+R)^{p-\delta}}{(x-c(p)\sqrt{t})^{p}} + \frac{1}{(x-c(p)\sqrt{t})^{\delta}}\right) \\
\leq C\frac{(x+R)^{p-\delta}}{(x-c(p)\sqrt{t})^{p}}.$$
(45)

Using (44) we infer that

$$\{\overline{V}_p(x+R+X,t+1) < C|x+R+X|^{p-\delta}\} \subseteq \{X < -(x-c(p)\sqrt{t})/2\}$$

for all $R \ge R(C)$, so

$$\begin{split} &\mathbf{E}\left[h(x+X,t+1)-h(x,t)\right] \\ &= \mathbf{E}\left[\overline{V}_p(x+R+X,t+1)-\overline{V}_p(x+R,t)-C\left(|x+R+X|^{p-\delta}-|x+R|^{p-\delta}\right)\right] \\ &- \mathbf{E}\left[\overline{V}_p(x+R+X,t+1)\right. \\ &- C|x+R+X|^{p-\delta}; \overline{V}_p(x+R+X,t+1) < C|x+R+X|^{p-\delta}\right] \\ &\leq \mathbf{E}\left[\overline{V}_p(x+R+X,t+1)-\overline{V}_p(x+R,t)\right] \\ &- C\cdot \mathbf{E}\left[\left(|x+R+X|^{p-\delta}-|x+R|^{p-\delta}\right)\right] \\ &+ \mathbf{E}\left[C|x+R+X|^{p-\delta}; X < -(x-c(p)\sqrt{t})/2\right]. \end{split}$$

Using Lemmas 8 and 10 together with (45), we find that for p > 3,

$$\mathbf{E} \left[h(x+X,t+1) - h(x,t) \right] \le -\frac{(p-\delta)(p-\delta-1)}{2} (x+R)^{p-\delta-2} + C \left((x+R)^{p-3} + \frac{(x+R)^{p-\delta}}{(x-c(p)\sqrt{t})^p} \right).$$

For $p \in (2, 3]$,

$$\mathbf{E}\left[h(x+X,t+1) - h(x,t)\right] \le -\frac{(p-\delta)(p-\delta-1)}{2}(x+R)^{p-\delta-2} + C\left(\frac{(x+R)^{p-1}}{(x-c(p)\sqrt{t})^{p-1}} + (x+R-c(p)\sqrt{t})^{p-2\delta-2}\right).$$

And if $p \leq 2$,

$$\mathbf{E} \left[h(x+X,t+1) - h(x,t) \right] \le -\frac{(p-\delta)(p-\delta-1)}{2} (x+R)^{p-\delta-2} + C \left(\frac{(x+R)^{p-\delta}}{(x-c(p)\sqrt{t})^{2+\delta}} + (x+R-c(p)\sqrt{t})^{p-2\delta-2} \right).$$

In all three cases, the entire term is dominated by $-\frac{(p-\delta)(p-\delta-1)}{2}(x+R)^{p-\delta-2}$ for sufficiently large x (all further terms are of lower order; indeed by Remark 11, we

obtain an asymptotic equality). Due to the negative sign, this term is non-positive, hence

$$\mathbf{E}\left[h(x+X,t+1)-h(x,t)\right] \le 0.$$

Lemma 13. For every c > 0 and every $1 < p_1 < p(c)$, there exist constants C and R such that

$$M_{n} := h(a + S(T_{a,b} \wedge n), b + T_{a,b} \wedge n)$$

= $\left(\overline{V}_{p_{1}}(R + a + S(T_{a,b} \wedge n), b + T_{a,b} \wedge n) - C \cdot |a + R + S(n)|^{p_{1} - \delta}\right)^{+}$

is a supermartingale.

Proof. Consider

$$\mathbf{E} [M_{n+1} - M_n \mid \mathcal{F}_n] = \mathbf{E} [(M_{n+1} - M_n) \mathbf{1}_{\{T_{a,b} \le n\}} \mid \mathcal{F}_n] + \mathbf{E} [(M_{n+1} - M_n) \mathbf{1}_{\{T_{a,b} > n\}} \mid \mathcal{F}_n].$$

The first term obviously satisfies

$$\begin{split} \mathbf{E} \left[(M_{n+1} - M_n) \mathbf{1}_{\{T_{a,b} \le n\}} \mid \mathcal{F}_n \right] \\ &= \mathbf{E} \left[(f(a + S(T_{a,b}), T_{a,b} + b) - f(a + S(T_{a,b}), T_{a,b} + b)) \, \mathbf{1}_{T_{a,b} \le n} \mid \mathcal{F}_n \right] = 0. \end{split}$$

For the other term, we obtain from Lemma 12 with $\gamma = c(p) - c(p_1) > 0$

$$\begin{aligned} \mathbf{E} \left[(M_{n+1} - M_n) \mathbf{1}_{\{T_{a,b} > n\}} \mid \mathcal{F}_n \right] \\ &= \mathbf{E} \left[h(a + S(n+1), n+1+b) - h(a + S(n), n+b) \mid \mathcal{F}_n \right] \mathbf{1}_{\{T_{a,b} > n\}} \\ &= \mathbf{E} \left[h(a + S(n) + X, n+1+b) - h(a + S(n), n+b) \mid \mathcal{F}_n \right] \mathbf{1}_{\{T_{a,b} > n\}} \le 0. \end{aligned}$$

This proves the claim.

We now turn to the case p < 1. It turns out that the function h(x,t) is not appropriate for this case and that one has to use a different type of correction for $V_p(x,t)$. We define

$$G(x,t) := \overline{V}_p(x+R,t+R) - C(t+R)^{(p-\delta)/2}$$
 and $g(t,x) := (G(x,t))^+$.

If $x > (c(p) + \gamma)\sqrt{t}$ for some $\gamma > 0$ then

$$\overline{V}_p(x+R,t+R) = (t+R)^{p/2}\psi_p\left(\frac{R+x}{\sqrt{R+t}}\right) \ge (t+R)^{p/2}\psi_p(c(p)+\gamma)$$

for all sufficiently large R. Therefore, there exists R(C) such that, for every $R \ge R(C)$,

$$g(x,t) > 0$$
 for all $x \ge (c(p) + \gamma)\sqrt{t}$. (46)

Lemma 14. Assume that p < 1. For every $\gamma > 0$ there exist constants C and R such that, for all $t \ge 0$ and $x \ge (c(p) + \gamma)\sqrt{t}$,

$$\mathbf{E}[g(x+X,t+1) - g(x,t)] \le 0.$$

Proof. It follows from (46) that

$$\begin{split} \mathbf{E}[g(x+X,t+1) - g(x,t)] &= \mathbf{E}\left[\overline{V}_p(x+R+X,t+R+1) - \overline{V}_p(x+R,t+R)\right] \\ &- C\left[(t+R+1)^{(p-\delta)/2} - (t+R)^{(p-\delta)/2}\right] \\ &- \mathbf{E}\left[G(x+X,t+1);G(x+X,t+1) < 0\right]. \end{split}$$

For every $x > (c(p) + \gamma)\sqrt{t}$ we have by Lemma 9

$$\begin{aligned} \mathbf{E} \left[\overline{V}_p(R+x+X,R+t+1) \right] &- \overline{V}_p(R+x,R+t) \\ &\leq C_1 (1+R+x-c(p)\sqrt{t+R})^{p-2-\delta} \leq C_2 (t+R)^{p/2-\delta/2-1}. \end{aligned}$$

Furthermore,

$$\left[(R+t+1)^{(p-\delta)/2} - (R+t)^{(p-\delta)/2} \right] \ge \frac{p-\delta}{2} (R+t+1)^{(p-\delta)/2-1}.$$

Finally, for all sufficiently large R, $\{G(x+X,t+1)<0\} \subset \{X<-\frac{\gamma}{2}\sqrt{R+t}\}$ and, consequently,

$$-\mathbf{E} \left[G(x+X,t+1); G(x+X,t+1) < 0 \right] \\\leq C(t+R)^{(p-\delta)/2} \mathbf{P}(G(x+X,t+1) < 0) \\\leq C(t+R)^{(p-\delta)/2} \mathbf{P}(X < -\frac{\gamma}{2}\sqrt{R+t}) \\< C(t+R)^{(p-\delta)/2-1-\delta/2} \mathbf{E}|X|^{2+\delta}.$$

Combining these estimates and increasing if necessary R, we get the desired property. \Box

Lemma 15. If $p_1 then there exist constants C and R such that$

$$\widehat{M}_n := \left(V_{p_1}(R + a + S(T_{a,b} \wedge n), R + b + T_{a,b} \wedge n) - C(R + b + T_{a,b} \wedge n)^{(p_1 - \delta)/2} \right)^+$$

is a supermartingale.

The proof of this lemma repeats the proof of Lemma 13 and we omit it.

4. Upper bounds for $\mathbf{P}(T_{a,b} > n)$ and for $\mathbf{P}(S(n) \in (x, x+h], T_{a,b} > n)$

4.1. Estimates for the tail of $T_{a,b}$.

Lemma 16. For every c > 0 and every $1 < p_1 < p(c)$, there exist constants C and R such that

$$\mathbf{E}\left[\overline{V}_{p_1}(R+a+S(n),n+b);T_{a,b}>n\right] \le C(1+a^{p_1}).$$

For every c < 0 and every $0 < p_1 < p(c) < 1$, there exist constants C and R such that

$$\mathbf{E}\left[\overline{V}_{p_1}(R+a+S(n),n+b);T_{a,b}>n\right] \le C(1+|a|^{p_1}+b^{p_1/2})+Cn^{(p_1-\delta)/2}\mathbf{P}(T_{a,b}>n)$$

Proof. Consider first p(c) > 1 and let p_1 be such that $1 < p_1 < p(c)$. Notice that $T_{a,b} \wedge n$ is a bounded stopping time. Hence, we can apply the optional stopping theorem to the supermartingale M_n from Lemma 13 to obtain that there is a constant C' such that

$$\mathbf{E} \left[h(a+S(n),b+n); T_{a,b} > n\right] \\
\leq \mathbf{E} \left[h(a+S(n),b+n); T_{a,b} > n\right] + \mathbf{E} \left[h(a+S(T_{a,b}),b+T_{a,b}); T_{a,b} \le n\right] \\
= \mathbf{E} \left[h(a+S(T_{a,b} \land n),b+T_{a,b} \land n)\right] = \mathbf{E} \left[M_{n}\right] \\
\leq \mathbf{E} \left[M_{0}\right] = \left(\overline{V}_{p_{1}}(a+R,b) - C \cdot (a+R)^{p_{1}-\delta}\right)^{+} \\
\leq (C'(a+R)^{p_{1}} - C \cdot (a+R)^{p_{1}-\delta}) \le C'(a+R)^{p_{1}}.$$
(47)

Notice that on the event $\{T_{a,b} > n\}$, for some sufficiently small $\gamma_0 > 0$

$$c(p_1)\sqrt{n+b} < (c(p) - \gamma_0)\sqrt{n+b} \le \frac{c(p) - \gamma_0}{c(p)}c(p)\sqrt{n+b} < (1 - \frac{\gamma_0}{c(p)})(a+S(n))$$

so in other words, for $\gamma = \frac{c(p_1)\gamma_0}{c(p)} \frac{1}{1 - \frac{\gamma_0}{c(p)}} > 0$

$$(a+S(n)) > (c(p_1)+\gamma)\sqrt{n+b}$$

and $a + S(n) - c(p_1)\sqrt{n+b} > \gamma\sqrt{n+b}$. It follows from Lemma 5 that

$$\overline{V}_{p_1}(R+a+S(n),b+n) - C(R+a+S(n))^{p_1-\delta} \\
\geq \frac{1}{A(p_1)}(R+a+S(n)-c(p_1)\sqrt{n+b}) \cdot (R+a+S(n))^{p_1-1} - C(R+a+S(n))^{p_1-\delta} \\
= \frac{(R+a+S(n))^{p_1-1}}{A(p_1)} \left[R+a+S(n)-c(p_1)\sqrt{n+b} - CA(p_1)(R+a+S(n))^{1-\delta} \right] \\
\geq \frac{\overline{V}_{p_1}(R+a+S(n),b+n)}{A^2(p_1)} \left[1 - CA(p_1)\frac{(R+a+S(n))^{1-\delta}}{R+a+S(n)-c(p_1)\sqrt{n+b}} \right].$$
(48)

Using the fact that

$$R + a + S(n) - c(p_1)\sqrt{n+b}$$

= $R + \left(1 - \frac{c(p_1)}{c}\right)(a+S(n)) + \frac{c(p_1)}{c}(a+S(n) - c\sqrt{n+b})$
 $\ge (1 - \frac{c(p_1)}{c})(R + a + S(n))$

we obtain from (48)

$$\overline{V}_{p_1}(R+a+S(n),b+n) - C(R+a+S(n))^{p_1-\delta} \\ \ge \frac{\overline{V}_{p_1}(R+a+S(n),b+n)}{A^2(p_1)} \left(1 - \frac{CA(p_1)}{R^{\delta}(1 - \frac{c(p_1)}{c})}\right)$$

Let $K := \frac{1}{A^2(p_1)} \left(1 - \frac{CA(p_1)}{R^{\delta}(1 - \frac{c(p_1)}{c})} \right)$, which is positive for sufficiently large R. Then

$$\begin{split} \mathbf{E} \left[\overline{V}_{p_1}(R+a+S(n),n+b); T_{a,b} > n \right] \\ &\leq \frac{1}{K} \mathbf{E} \left[\overline{V}_{p_1}(R+a+S(n),n+b) - C | R+a+S(n)|^{p_1-\delta}; T_{a,b} > n \right] \\ &\leq \frac{1}{K} \mathbf{E} \left[\left(\overline{V}_{p_1}(R+a+S(n),n+b) - C \cdot | R+a+S(n)|^{p_1-\delta} \right)^+; T_{a,b} > n \right] \\ &= \frac{1}{K} \mathbf{E} \left[h(a+S(n),n+b); T_{a,b} > n \right] \leq \frac{C}{K} (a+R)^{p_1} \end{split}$$

by (47) and the claim follows.

For $0 < p_1 < p(c) < 1,$ we can apply optional stopping theorem to the supermartingale \widehat{M}_n from Lemma 15 to obtain

$$\begin{split} \mathbf{E} \left[\overline{V}_{p_1}(R+a+S(n),R+b+n); T_{a,b} > n \right] \\ &= \mathbf{E} \left[\overline{V}_{p_1}(R+a+S(T_{a,b} \wedge n),R+b+T_{a,b} \wedge n); T_{a,b} > n \right] \\ &= \mathbf{E} \left[\widehat{M}_n; T_{a,b} > n \right] + C(R+b+n)^{(p_1-\delta)/2} \mathbf{P}(T_{a,b} > n) \\ &\leq \mathbf{E} \left[\widehat{M}_n \right] + C(R+b+n)^{(p_1-\delta)/2} \mathbf{P}(T_{a,b} > n) \\ &\leq \mathbf{E} \left[\widehat{M}_0 \right] + C(R+b+n)^{(p_1-\delta)/2} \mathbf{P}(T_{a,b} > n) \\ &\leq \overline{V}_{p_1}(R+a,R+b) + Cn^{(p_1-\delta)/2} \mathbf{P}(T_{a,b} > n). \end{split}$$

Using Lemma 7, we can bound the first term $\overline{V}_{p_1}(R+a, R+b) \leq A(p_1)(R+|a|+\sqrt{b})^{p_1}$, so the claim holds.

Lemma 17. For $x \ge 0$, $z \ge n$, $R \in \mathbb{R}_+$ and p > 0 we have

$$\mathbf{P}(\overline{V}_p(S(n)+R,z) > x, T_{a,b} > n) \ge \mathbf{P}(\overline{V}_p(S(n)+R,z) > x) \cdot \mathbf{P}(T_{a,b} > n).$$

Proof. Since $\overline{V}_p(x, y)$ is increasing in x,

$$\{\overline{V}_p(a+R+S(n),b+n) > x\} = \{S(n) > g_n\},\$$

where g_n is the unique solution to $\overline{V}_p(a+R+g_n,b+n)=x$. By Lemma 24 in [5],

$$\mathbf{P}(S(n) > g_n, T_{a,b} > n) \ge \mathbf{P}(S(n) > g_n)\mathbf{P}(T_{a,b} > n).$$

This gives the desired estimate.

Lemma 18. If $\mathbf{E}|X|^p < \infty$, then

$$\mathbf{P}(T_{a,b} > n) \le \frac{\mathbf{E}\left[\overline{V}_p(a+R+S(n),b+n); T_{a,b} > n\right]}{\mathbf{E}\left[\overline{V}_p(a+R+S(n),b+n)\right]}.$$

Proof. By Lemma 17,

$$\begin{split} \mathbf{E}\left[\overline{V}_{p}(a+R+S(n),b+n);T_{a,b}>n\right] \\ &= \int_{0}^{\infty} \mathbf{P}\left(\overline{V}_{p}(a+R+S(n),b+n)>x,T_{a,b}>n\right)dx \\ &\geq \mathbf{P}(T_{a,b}>n)\int_{0}^{\infty} \mathbf{P}\left(\overline{V}_{p}(a+R+S(n),b+n)>x\right)dx \\ &= \mathbf{P}(T_{a,b}>n)\mathbf{E}\left[\overline{V}_{p}(a+R+S(n),b+n)\right]. \end{split}$$

Lemma 19. For every $1 < p_1 < p(c)$, there exists a finite constant C such that for all a, b,

$$\mathbf{P}(T_{a,b} > n) \le C \frac{1 + |a|^{p_1}}{n^{p_1/2}}.$$

If $0 < p_1 < p(c) < 1$ then

$$\mathbf{P}(T_{a,b} > n) \le C \frac{1 + |a|^{p_1} + b^{p_1/2}}{n^{p_1/2}}.$$

Proof. Assume first that p(c) > 1. We know from Lemma 16 that

$$\mathbf{E}\left[\overline{V}_{p_1}(a+R+S(n),b+n);T_{a,b}>n\right] \le C(1+a^{p_1})$$

for all *a* and *b*. Thus, by Lemma 18, it remains to estimate $\mathbf{E}\left[\overline{V}_{p_1}(a+R+S(n),b+n)\right]$. By the definition of \overline{V}_{p_1} ,

$$\begin{split} \mathbf{E}\left[\overline{V}_{p_1}(a+R+S(n),b+n)\right] \\ \geq (b+n)^{p_1/2} \mathbf{E}\left[\psi_{p_1}\left(\frac{a+R+S(n)}{\sqrt{n+b}}\right); a+S(n) > c\sqrt{b+n}\right]. \end{split}$$

Applying the central limit theorem, we conclude that the expectation on the right hand side converges to a positive constant, which depends only on p. This gives the desired bound.

In the case $p_1 < p(c) < 1$ we use the same estimation for the denominator in Lemma 18. For the enumerator, we find by Lemma 16 that there are constants C_1 and C_2 such that

$$\begin{split} \mathbf{E}\left[\overline{V}_{p_1}(R+a+S(n),n+b);T_{a,b}>n\right] \\ &\leq C_1(1+|a|^{p_1}+b^{p_1/2})+C_2n^{(p_1-\delta)/2}\mathbf{P}(T_{a,b}>n). \end{split}$$

Overall, we thus obtain

$$P(T_{a,b} > n) \le C_1 \frac{1 + |a|^{p_1} + b^{p_1/2}}{n^{p_1/2}} + C_2 \frac{n^{(p_1 - \delta)/2} \mathbf{P}(T_{a,b} > n)}{n^{p_1/2}}.$$

Rearranging gives some constant C_2 such that

$$\mathbf{P}(T_{a,b} > n) \le \left(1 - \frac{C_2}{n^{\delta/2}}\right)^{-1} C_1 \frac{1 + |a|^{p_1} + b^{p_1/2}}{n^{p_1/2}}.$$

For $n \ge (2C_2)^{2/\delta}$ we have the desired inequality with $C = 2C_1$. This completes the proof of the lemma.

Lemma 20. For every $0 < p_1 < p(c)$, there exists a constant $C < \infty$ such that

$$\mathbf{E}\left[\left(\max_{k \le T_{a,b}} |S(k)|\right)^{p_1}\right] \le C(1+|a|^{p_1}+b^{p_1/2}\mathbf{1}\{p(c)<1\})$$

and

$$\mathbf{E}(T_{a,b})^{p_1/2} \le C(1+|a|^{p_1}+b^{p_1/2}\mathbf{1}\{p(c)<1\}).$$

Proof. By the Burkholder-Davis-Gundy inequality,

$$\mathbf{E}\left[\left(\max_{k\leq T_{a,b}}|S(k)|\right)^{p_1}\right]\leq C\mathbf{E}\left[T_{a,b}^{p_1/2}\right].$$

For every $k \ge 0$, one has

$$\begin{split} \mathbf{E}\left[T_{a,b}^{p_{1}/2}\right] &= \frac{p_{1}}{2} \int_{0}^{\infty} u^{p_{1}/2-1} \mathbf{P}(T_{a,b} > u) du \\ &\leq k^{p_{1}/2} + \frac{p_{1}}{2} \int_{k}^{\infty} u^{p_{1}/2-1} \mathbf{P}(T_{a,b} > u) du. \end{split}$$

Applying Lemma 19 with some $p_2 \in (p_1, p)$, we get

$$\frac{p_1}{2} \int_k^\infty u^{p_1/2-1} \mathbf{P}(T_{a,b} > u) du \le \frac{p_1}{2} \int_k^\infty u^{p_1/2-1} \frac{1+a^{p_2}}{u^{p_2/2}} du$$
$$= \frac{p_1}{2} (1+a^{p_2}) \int_k^\infty u^{(p_1-p_2)/2-1} du \le \widehat{C}(1+a^{p_2}) k^{(p_1-p_2)/2}$$

Taking $k = (1+a)^2$, we get

$$\mathbf{E}\left[T_{a,b}^{p_1/2}\right] \le (1+a)^{p_1} \left(1 + \widehat{C}\frac{1+a^{p_2}}{(1+a)^{p_2}}\right) \le (1+\widehat{C})(1+a)^{p_1}.$$

This finishes the proof.

4.2. Estimates for local probabilities.

Lemma 21. For every $p_1 < p(c)$ there exists a constant C such that

$$\mathbf{P}(S(n) \in (x, x+1], T_{a,b} > n) \le C \frac{1+|a|^{p_1}}{n^{(p_1+1)/2}}.$$

Proof. Set $m = \lfloor \frac{n}{2} \rfloor$. It follows from the local central limit theorem that

$$\mathbf{P}(S(n-m) \in (z, z+1]) \le C_1(n-m)^{-1/2} \le C_2 n^{-1/2}$$

uniformly in z. Using this bound we obtain

$$\begin{aligned} \mathbf{P}(S(n) &\in (x, x+1], T_{a,b} > n) \\ &\leq \int \mathbf{P}(S(m) \in dz, T_{a,b} > m) \mathbf{P}(S(n-m) \in (x-z, x-z+1]) \\ &\leq C_2 n^{-1/2} \mathbf{P}(T_{a,b} > m)). \end{aligned}$$

Applying now Lemma 19, we get the desired inequality.

Lemma 22. If c > 0 then for every $1 < p_1 < p(c)$ there exists a constant C such that, for all $y \leq \sqrt{n}/2$,

$$\mathbf{P}(a+S(n) \in (c\sqrt{b+n}+y, c\sqrt{b+n}+y+1], T_{a,b} > n) \le C(1+y)^{1/2} \frac{1+|a|^{p_1}}{n^{p_1/2+3/4}}.$$
 (49)

If c < 0 then for every $0 < p_1 < p(c)$ there exists a constant C such that, for all $y \le \sqrt{n}/2$,

$$\mathbf{P}(a+S(n)\in(c\sqrt{b+n}+y,c\sqrt{b+n}+y+1], T_{a,b}>n)\leq C(1+y)\frac{1+|a|^{p_1}+b^{p_1/2}}{n^{p_1/2+1}}.$$
(50)

Proof. Set

$$\Delta(y) = (c\sqrt{b+n} + y, c\sqrt{b+n} + y + 1].$$

Then, by the Markov property at time n - m < n,

$$\mathbf{P}(a+S(n)\in\Delta(y), T_{a,b}>n)$$

=
$$\int_{c\sqrt{b+n-m}}^{\infty} \mathbf{P}(a+S(n-m)\in dz, T_{a,b}>n-m)\mathbf{P}(z+S(m)\in\Delta(y), T_{z,b+n-m}>m)$$

Using the duality lemma for random walks, we get

$$\begin{aligned} \mathbf{P}(z+S(m)\in\Delta(y), T_{z,b+n-m} > m) \\ &\leq \mathbf{P}\left(z+S(m)\in\Delta(y), \min_{j\leq m}(z+S(j)) > \min_{j\leq m}c\sqrt{b+n-m+j}\right) \\ &\leq \mathbf{P}\left(y+1-S(m)\in[z-c\sqrt{b+n}, z+1-c\sqrt{b+n}), \\ &\min_{j\leq m}(y+1-S(j)) > -c\sqrt{b+n} + \min_{j\leq m}c\sqrt{b+n-m+j}\right). \end{aligned}$$

It follows from Lemma 21 that, for $m \leq n/2,$

$$\mathbf{P}(a + S(n - m) \in (c\sqrt{b + n - m} + k, c\sqrt{b + n - m} + k + 1], T_{a,b} > n - m)$$

$$\leq C \frac{1 + |a_1|^{p_1}}{n^{(p_1 + 1)/2}}$$

uniformly in $k \ge 0$. Therefore,

$$\begin{aligned} \mathbf{P}(a+S(n) \in \Delta(y), T_{a,b} > n) \\ &\leq C \frac{1+|a_1|^{p_1}}{n^{(p_1+1)/2}} \sum_{k=0}^{\infty} \mathbf{P}\Big(y+1-S(m) \in \Delta'(k), \\ &\qquad \min_{j \leq m} (y+1-S(j)) > -c\sqrt{b+n} + \min_{j \leq m} c\sqrt{b+n-m+j} \Big) \\ &\leq C \frac{1+|a_1|^{p_1}}{n^{(p_1+1)/2}} \mathbf{P}\left(\min_{j \leq m} (y+1-S(j)) > -c\sqrt{b+n} + \min_{j \leq m} c\sqrt{b+n-m+j} \right), \end{aligned}$$

where

$$\Delta'(k) = [c\sqrt{b+n-m} - c\sqrt{b+n} + k, c\sqrt{b+n-m} - c\sqrt{b+n} + k + 1].$$

We next notice that

$$c\sqrt{b+n} - \min_{j \le m} c\sqrt{b+n-m+j} = c^+(\sqrt{b+n} - \sqrt{b+n-m}).$$

Therefore,

$$\mathbf{P}(a+S(n) \in \Delta(y), T_{a,b} > n) \\ \leq C \frac{1+|a_1|^{p_1}}{n^{(p_1+1)/2}} \mathbf{P}\left(\min_{j \le m} (y+1-S(j)) > -c^+(\sqrt{b+n}-\sqrt{b+n-m})\right).$$
(51)

If $c>0~(c^+=c)$ then we take $m=\lfloor (y+1)\sqrt{n}+1 \rfloor.$ In this case

$$\sqrt{b+n} - \sqrt{b+n-m} = \frac{m}{\sqrt{b+n} + \sqrt{b+n-m}} \ge \frac{(y+1)\sqrt{n}}{\sqrt{n}} = y+1.$$

Then, using Lemma 3 from [8], we get

$$\mathbf{P}(\min_{j \le m}(y+1-S(j)) > -c^{+}(\sqrt{b+n} - \sqrt{b+n-m}))$$

$$\leq \mathbf{P}(\min_{j \le m}(-S(j)) > -(1+c)(1+y)) \le C\frac{1+y}{\sqrt{m}} \le C\frac{(1+y)^{1/2}}{n^{1/4}}.$$
 (52)

Combining (51) and (52), we get (49).

If $c^+ = 0$, i.e. $c \le 0$, then we take $m = \lfloor n/2 \rfloor$. Using once again Lemma 3 from [8], we have

$$\mathbf{P}(\min_{j \le m}(y+1-S(j)) > -c^{+}(\sqrt{b+n} - \sqrt{b+n-m}))$$

$$\leq \mathbf{P}(\min_{j \le m}(-S(j)) > -(1+y)) \le C\frac{1+y}{\sqrt{m}} \le C\frac{(1+y)}{n^{1/2}}.$$
 (53)

Combining (51) and (53), we get (50).

5. Construction of the harmonic function: proof of Theorem 1

Now let

$$f(x,y) := \mathbf{E}\left[\overline{V}_{p(c)}(x+X,y+1)\right] - \overline{V}_{p(c)}(x,y).$$

Then the sequence

$$M_n := \overline{V}_{p(c)}(a + S(n), b + n) - \sum_{k=0}^{n-1} f(a + S(k), b + k)$$

is a martingale. Therefore, using optional stopping, we obtain

$$\begin{split} \overline{V}_{p(c)}(a,b) &= M_0 = \mathbf{E}\left[M_{T_{a,b}\wedge n}\right] = \mathbf{E}\left[M_n; T_{a,b} > n\right] + \mathbf{E}\left[M_{T_{a,b}}; T_{a,b} \le n\right] \\ &= \mathbf{E}\left[\overline{V}_{p(c)}(a + S(n), b + n); T_{a,b} > n\right] + \mathbf{E}\left[\sum_{k=0}^{n-1} f_k; T_{a,b} > n\right] \\ &+ \mathbf{E}\left[\overline{V}_{p(c)}(a + S(T_{a,b}), b + T_{a,b}); T_{a,b} \le n\right] - \mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} f_k; T_{a,b} \le n\right], \end{split}$$

where $f_k := f(a + S(k), b + k)$. The third expectation vanishes since $V_{p(c)}(a + S(T_{a,b}), b + T_{a,b}) = 0$. Rearranging this, we obtain

$$\mathbf{E}\left[\overline{V}_{p(c)}(a+S(n),b+n);T_{a,b}>n\right] \\
= \overline{V}_{p(c)}(a,b) - \mathbf{E}\left[\sum_{k=0}^{n-1}f_k;T_{a,b}>n\right] - \mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1}f_k;T_{a,b}\leq n\right] \\
= \overline{V}_{p(c)}(a,b) - \mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1}f_k\right] + \mathbf{E}\left[\sum_{k=n}^{T_{a,b}-1}f_k;T_{a,b}>n\right].$$
(54)

A priori, existence of the limit of the expectation on the left hand side is not clear: **Proposition 23.** The limit $\lim_{n\to\infty} \mathbf{E} \left[V_{p(c)}(a + S(n), b + n); T_{a,b} > n \right]$ is finite. *Proof.* We first notice that

 $\mathbf{E}\left[V_{p(c)}(a+S(n),b+n);T_{a,b}>n\right] = \mathbf{E}\left[\overline{V}_{p(c)}(a+S(n),b+n);T_{a,b}>n\right].$

It is then immediate from the representation (54) that the claim will be proven if we show that the random variable $\sum_{k=0}^{T_{a,b}-1} |f_k|$ has finite expectation. Indeed, this will imply that the second term in (54) is well-defined and that the third term will vanish as $n \to \infty$.

For $p(c) \geq 3$, we notice that by Lemma 8

$$|f(x,y)| \le C(1+x)^{p(c)-3} + C\frac{(1+x)^{p(c)-1}}{(1+x-c\sqrt{y})^{p(c)-1}}$$

for some suitable positive constant C. Therefore,

$$\sum_{k=0}^{T_{a,b}-1} |f_k| \le C \sum_{k=0}^{T_{a,b}-1} (a+1+S(k))^{p(c)-3} + C \sum_{k=0}^{T_{a,b}-1} \frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}.$$
 (55)

For the first sum we have

$$\sum_{k=0}^{T_{a,b}-1} (a+1+S(k))^{p(c)-3} \le C_1 \left[(a+1)^{p(c)-3} T_{a,b} + \sum_{k=0}^{T_{a,b}-1} (S(k))^{p(c)-3} \right]$$
$$\le C_1 \left[(a+1)^{p(c)-3} + (\max_{k \le T_{a,b}} |S(k)|)^{p(c)-3} \right] T_{a,b}$$

for some positive constant C_1 . We define $\bar{S}_T := \max_{k \leq T_{a,b}} |S(k)|$. Taking the expectation and applying the Cauchy-Schwarz inequality, we get

$$\mathbf{E} \left[\sum_{k=0}^{T_{a,b}-1} (a+1+S(k))^{p(c)-3} \right] \\
\leq C_1 \left[(a+1)^{p(c)-3} \mathbf{E} T_{a,b} + \mathbf{E} \left[(\bar{S}_T)^{p(c)-3} T_{a,b} \right] \right] \\
\leq C_1 \left((a+1)^{p(c)-3} \mathbf{E} T_{a,b} + \mathbf{E} \left[(\bar{S}_T)^{(p(c)-3)r} \right]^{1/r} \mathbf{E} \left[T_{a,b}^s \right]^{1/s} \right),$$

where we choose $s = p_1/2 < p(c)/2$ and $r = p_1/(p_1-2)$ which satisfies $(p(c)-3)r < p_1$ provided that $p(c) - 1 < p_1 < p(c)$. Applying now Lemma 20, we conclude that

$$\mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} (a+1+S(k))^{p(c)-3}\right] \le C(1+a)^{p(c)-1}.$$
(56)

We now turn to expectation of the second sum in (55). We start by the following splitting:

$$\begin{split} \mathbf{E} \left[\sum_{k=0}^{T_{a,b}-1} \frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p-1}} \right] \\ &= \sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}; T_{a,b} > k \right] \\ &= \sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}; a+S(k) > \left(c+\frac{1}{2}\right)\sqrt{k+b}, T_{a,b} > k \right] \\ &+ \sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-\sqrt{k+b})^{p(c)-1}}; a+S(k) \le \left(c+\frac{1}{2}\right)\sqrt{k+b}, T_{a,b} > k \right]. \end{split}$$

We notice that if $a + S(k) > (c + 1/2)\sqrt{k + b}$, then

$$\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}} \le C.$$

Therefore, applying Lemma 20, we obtain

$$\sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}; a+S(k) > \left(c+\frac{1}{2}\right)\sqrt{k+b}, T_{a,b} > k \right]$$

$$\leq C \sum_{k=0}^{\infty} \mathbf{P}(T_{a,b} > k) = C \mathbf{E} T_{a,b} \leq C(1+a)^2.$$
(57)

We now split

$$\mathbf{E}\left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}};a+S(k)\leq (c+1)\sqrt{k+b},T_{a,b}>k\right]$$
$$\leq C\sqrt{k+b}^{p(c)-1}\sum_{j=1}^{\frac{1}{2}\sqrt{k+b}}j^{-p(c)+1}\mathbf{P}\left(a+S(k)\in\Delta(j),T_{a,b}>k\right),$$

where

$$\Delta(j) = \left(c\sqrt{k+b} + j - 1, c\sqrt{k+b} + j\right].$$

Using now Lemma 22, we get

$$\sum_{j=1}^{\frac{1}{2}\sqrt{k+b}} j^{-p(c)+1} \mathbf{P} \left(a + S(k) \in \Delta(j), T_{a,b} > k \right)$$
$$\leq \frac{C(1+a)^{p_1}}{k^{p_1/2+3/4}} \sum_{j=1}^{\frac{1}{2}\sqrt{k+b}} j^{-p(c)+1} (1+j)^{1/2}.$$
(58)

Since $p(c) \ge 3$ the sum on the right hand side of (58) is bounded and, consequently,

$$\sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}; a+S(k) \le (c+\frac{1}{2})\sqrt{k+b}, T_{a,b} > k \right]$$

$$\le C(1+a)^{p(c)-1} + C(1+a)^{p_1} \sum_{k=1}^{\infty} k^{(p(c)-1)/2-p_1/2-3/4}.$$
(59)

Therefore, for every $p_1 \in (p(c) - 1/2, p(c))$ we have

$$\sum_{k=0}^{\infty} \mathbf{E}\left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}; a+S(k) \le (c+\frac{1}{2})\sqrt{k+b}, T_{a,b} > k\right] \le C(1+a)^{p_1}.$$
(60)

Combining (58) and (60), we finally get

$$\mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} \frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}\right] \le C(1+a)^{p(c)_1}.$$
(61)

This finishes the proof in the case $p(c) \ge 3$.

In the case $p(c) \in (2,3)$ we have from Lemma 8

$$|f_k| \le C \frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p(c)-1}}.$$

Thus, we need to prove an analogue of (61) for $p(c) \in (2,3)$. Clearly, (57) holds for all p(c) > 2. Furthermore, it is easy to see that if p(c) > 2.5 then the sum in (58) is still bounded and, consequently, (60) remains valid for p(c) > 2.5. Assume now that $p(c) \leq 2.5$. In this case

$$\sum_{j=1}^{\frac{1}{2}\sqrt{k+b}} j^{-p(c)+1} (1+j)^{1/2} \le Ck^{5/4-p(c)/2}$$

and, consequently, instead of (59) we have

$$\sum_{k=0}^{\infty} \mathbf{E} \left[\frac{(1+a+S(k))^{p(c)-1}}{(1+a+S(k)-c\sqrt{k+b})^{p-1}}; a+S(k) \le (c+\frac{1}{2})\sqrt{k+b}, T_{a,b} > k \right]$$
$$\le C(1+a)^{p(c)-1} + C(1+a)^{p_1} \sum_{k=1}^{\infty} k^{(p(c)-1)/2-p_1/2-3/4+5/4-p(c)/2}$$
$$\le C(1+a)^{p(c)-1} + C(1+a)^{p_1} \sum_{k=1}^{\infty} k^{-p_1/2}.$$

Now we can infer that (61) remains valid in the case $p(c) \in (2,3)$ with $p_1 \in (2, p(c))$.

The case $p(c) \in (1, 2]$ is rather similar to $p(c) \in (2, 3)$ and we omit its proof. We now consider the remaining case p(c) < 1. According to Lemma 9,

$$|f_k| \le C(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}.$$

Therefore,

$$\mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} |f_k|\right] \le \mathbf{E}\left[C\sum_{k=0}^{T_{a,b}-1} (1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}\right] \\ = C\sum_{k=0}^{\infty} \mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}; T_{a,b} > k\right].$$

Here we again split every expectation into two parts:

$$\begin{split} & \mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}; T_{a,b} > k\right] \\ &= \mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}; a+S(k) \le (c+1/2)\sqrt{b+k}, T_{a,b} > k\right] \\ &+ \mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}; a+S(k) > (c+1/2)\sqrt{b+k}, T_{a,b} > k\right]. \end{split}$$

The second expectation is bounded by $k^{p(c)/2-1-\delta/2}\mathbf{P}(T_{a,b} > k)$. This implies that

$$\sum_{k=0}^{\infty} \mathbf{E} \left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta}; a+S(k) > (c+1/2)\sqrt{b+k}, T_{a,b} > k \right]$$

$$\leq C \sum_{k=0}^{\infty} k^{p(c)/2-1-\delta/2} \mathbf{P}(T_{a,b} > k)$$

$$\leq C_1 \mathbf{E} T_{a,b}^{p(c)/2-\delta/2} \leq C_2 (1+|a|^{p(c)-\delta}+b^{p(c)/2-\delta/2}).$$
(62)

In the last step we have used Lemma 20. For every k we also have

$$\mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta};a+S(k) \le (c+1/2)\sqrt{b+k}, T_{a,b} > k\right]$$

$$\le \sum_{j=1}^{\sqrt{b+k}/2} j^{p(c)-2-\delta} \mathbf{P}(a+S(k) \in \Delta(j), T_{a,b} > k).$$

Applying now (50), we have

$$\mathbf{E}\left[(1+a+S(k)-c\sqrt{b+k})^{p(c)-2-\delta};a+S(k)\leq (c+1/2)\sqrt{b+k},T_{a,b}>k\right] \\
\leq C\frac{(1+|a|^{p_1}+b^{p_1/2})}{k^{p_1/2+1}}\sum_{j=1}^{\sqrt{b+k}/2}j^{p(c)-1-\delta}\leq C\frac{(1+|a|^{p_1}+b^{p_1/2})}{k^{p_1/2+1}}k^{p(c)/2-\delta/2}.$$

The sequence on the right hand side is summable for any $p_1 \in (p(c) - \delta, p(c))$. Thus,

$$\sum_{k=0}^{\infty} \mathbf{E} \left[(1+a+S(k)-c\sqrt{b+k})^{p-2-\delta}; a+S(k) \le (c+1/2)\sqrt{b+k}, T_{a,b} > k \right]$$

$$\le C(1+|a|^{p_1}+b^{p_1/2}).$$

Combining this with (62) completes the proof of the proposition.

As the expectation exists, we can use dominated convergence to take the limit $n \to \infty.$ Thus, we obtain

$$\lim_{n \to \infty} \mathbf{E} \left[V_{p(c)}(a + S(n), b + n); T_{a,b} > n \right] = V_{p(c)}(a, b) - \mathbf{E} \left[\sum_{k=0}^{T_{a,b}-1} f_k \right] =: W(a, b)$$

Lemma 24. Under the assumptions of Theorem 1,

$$\mathbf{E}[W(a+X, b+1); T_{a,b} > 1] = W(a, b).$$

Proof. Let $S^*(k) := S(k+1) - X_1$. Clearly, S(k) and X_1 are independent, and S(k) and $S^*(k)$ are identically distributed. Let $T^*_{a,b} := \inf\{k \ge 0 : S^*(k) + a < c\sqrt{k+b}\}$. We notice first that

$$\begin{aligned} \{T_{a,b} > n+1\} \\ &= \{a+S(k) > c\sqrt{k+b} \text{ for all } k \le n+1\} \\ &= \{a+S(1) > c\sqrt{1+b}\} \cap \{X_1 + a + (S(k) - X_1) > c\sqrt{k+b} \text{ for all } 2 \le k \le n+1\} \\ &= \{T_{a,b>1}\} \cap \{X_1 + a + S^*(k) > c\sqrt{k+1+b} \text{ for all } k \le n\} \\ &= \{a+S(1) > c\sqrt{1+b}\} \cap \{T^*_{a+X_1,b+1} > n\}.\end{aligned}$$

Therefore, using law of total expectation to condition on the first step of the random walk,

$$\begin{split} \mathbf{E} \left[V_{p(c)}(a + S(n+1), b + n + 1); T_{a,b} > n + 1 \right] \\ &= \mathbf{E} \left[\mathbf{E} \left[V_{p(c)}(a + S(n+1), b + n + 1); T_{a,b} > n + 1 | X_1 \right] \right] \\ &= \mathbf{E} \left[\mathbf{E} \left[V_{p(c)}(a + X_1 + S^*(n), b + 1 + n); T_{a+X_1, b+1}^* > n | X_1 \right]; T_{a,b} > 1 \right] \\ &= \int \mathbf{E} \left[V_{p(c)}(a + x + S(n), b + 1 + n); T_{a+x, b+1} > n \right] \mathbf{P}(X_1 \in dx, T_{a,b} > 1) \\ &= \int \mathbf{E} \left[V_{p(c)}(y + S(n), b + 1 + n); T_{y, b+1} > n \right] \mathbf{P}(a + X_1 \in dy, T_{a,b} > 1) \end{split}$$

We have seen in the proof of Lemma 23 that $\mathbf{E}\left[V_{p(c)}(y+S(n),b+1+n);T_{y,b+1}>n\right]$ is bounded by $C(1+|y|^p)$. Then, due to the finiteness of $\mathbf{E}|X|^p$, we may apply dominated convergence theorem:

$$\begin{split} W(a,b) &= \lim_{n \to \infty} \mathbf{E} \left[V_{p(c)}(a + S(n+1), b + n + 1); T_{a,b} > n + 1 \right] \\ &= \lim_{n \to \infty} \int \mathbf{E} \left[V_{p(c)}(y + S(n), b + 1 + n); T_{y,b+1} > n \right] \mathbf{P}(a + X_1 \in dy, T_{a,b} > 1) \\ &= \int W(y, b + 1) \mathbf{P}(a + X_1 \in dy, T_{a,b} > 1) \\ &= \mathbf{E} \left[W(a + X_1, b + 1); T_{a,b} > 1 \right]. \end{split}$$

Lemma 25. The function W(a,b) is monotone increasing in a. Furthermore, if $b \to \infty$ with $a > (c(p) + \gamma)\sqrt{b}$ for some $\gamma > 0$, then

$$W(a,b) \sim V_{p(c)}(a,b).$$
 (63)

Moreover, we can pick $\gamma_b \downarrow 0$ sufficiently slow so that (63) still holds for $a > (c(p) + \gamma_b)\sqrt{b}$.

Proof. Monotonicity of $a \mapsto W(a, b)$ is rather obvious. It is immediate from the observation that $V_{p(c)}(a, b)$ and $T_{a,b}$ are increasing in a. If $a > (c(p) + \gamma)\sqrt{b}$, then $V_{p(c)}(a, b) \ge C(a - c\sqrt{b})^{p(c)}$ for some suitable constant

If $a > (c(p) + \gamma)\sqrt{b}$, then $V_{p(c)}(a, b) \ge C(a - c\sqrt{b})^{p(c)}$ for some suitable constant C, C'. It suffices thus that in every case, $\mathbf{E}\begin{bmatrix}T_{a,b}^{-1}\\\sum_{k=0}^{T_{a,b}-1}|f_k|\end{bmatrix}$ is of lower order. This statement follows from the proof of Proposition 23: there we have shown

This statement follows from the proof of Proposition 23: there we have shown that if p(c) > 1 then $\mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} |f_k|\right] = O((1+|a|)^{p_1})$ for an appropriately chosen $\begin{bmatrix} T_{a,b}-1 \end{bmatrix}$

 $p_1 < p(c)$. And for p(c) < 1 we have proven that $\mathbf{E}\left[\sum_{k=0}^{T_{a,b}-1} |f_k|\right] = O((1+|a|^{p_1}+b^{p_1/2}))$ for an appropriate $p_1 < p(c)$. In both cases this term is of smaller order

 $(a - c\sqrt{b})^{p(c)}$. Thus, the proof is complete.

Proof of Corollary 2. Recalling that $a \mapsto W(a, b)$ is increasing and using Lemma 17, we obtain

$$\mathbf{P}(T_{a,b} > n) \le \frac{\mathbf{E}[W(a + S(n), b + n); T_{a,b} > n]}{\mathbf{E}[W(a + S(n), b + n); a + S(n) > c\sqrt{b + n}]}.$$

Then, by the harmonicity of W,

$$\mathbf{P}(T_{a,b} > n) \le \frac{W(a,b)}{\mathbf{E}[W(a+S(n),b+n);a+S(n) > c\sqrt{b+n}]}$$

Thus, it remains to derive a lower bound for the expectation in the denominator.

We know from Theorem 1 that $W(a, b) \sim V_{p(c)}(a, b)$ as $a > (c+1)\sqrt{b}$ and $a \to \infty$. Combining this with Lemma 4, we conclude that

$$\mathbf{E}[W(a+S(n),b+n);a+S(n) > c\sqrt{b+n}]$$

$$\geq A(b+n)^{p(c)/2}\mathbf{P}(a+S(n) > (c+1)\sqrt{b+n})$$

for some constant A, which does not depend on a and b. Since $a > c\sqrt{b}$ and $n \ge b$,

$$\mathbf{P}(a+S(n) > (c+1)\sqrt{b+n}) \ge \mathbf{P}(S(n) > (c+1)\sqrt{b+n} - c\sqrt{b})$$
$$\ge \mathbf{P}(S(n) > (|c|+1)\sqrt{2n}).$$

The desired bound follows now from the central limit theorem.

6. Asymptotics for $\mathbf{P}(T_{a,b} > n)$: proof of Theorem 3.

For all k < n we define

$$Q_{k,n}(y) := \mathbf{P}(y + S(j) > g(k+j) \text{ for all } 0 \le j \le n-k)$$
$$= \mathbf{P}(y + S(j) > c\sqrt{b+k+j} \text{ for all } 0 \le j \le n-k),$$
$$= \mathbf{P}(T_{a+y,b+k} > n-k).$$

Applying Corollary 2 we conclude that

$$Q_{k,n}(y) \le M \frac{W(a+y,b+k)}{(n-k)^{p(c)/2}}, \quad k \le \frac{n-b}{2}.$$

Therefore, for every $k \leq \frac{n-b}{2}$,

$$\left| (n-k)^{p(c)/2} Q_{k,n}(y) - \varkappa W(a+y,b+k) \right| \le (M+\varkappa) W(a+y,b+k).$$
(64)

Let γ_n be a sequence, which converges to zero sufficiently slow. Choose now a further sequence μ_n such that $\mu_n \to 0$ and $\frac{\mu_n}{\gamma_n} \to \infty$. If $\gamma_n \to 0$ sufficiently slow then, using the functional central limit theorem, we get, uniformly in $(y,k) \in I_n$, where $I_n := \{(y,k) : k \leq (n-b)/2, a+y-c\sqrt{b+k} \in (\gamma_n\sqrt{n}, \mu_n\sqrt{n})\}$,

$$\mathbf{P}(T_{a+y,b+k} > n-k) \sim \mathbf{P}(T_{a+y,b+k}^{(bm)} > n-k).$$

Then, using Uchiyama [15],

$$\left|\mathbf{P}(T_{a+y,b+k} > n-k) - \varkappa \frac{V_{p(c)(a+y,b+k)}}{(n-k)^{p(c)/2}}\right| \le \varepsilon_n \frac{V_p(a+y,b+k)}{(n-k)^{p(c)/2}}$$

uniformly in $(y, k) \in I_n$ for some $\varepsilon_n \downarrow 0$. We know from Lemma 25 that

$$V_{p(c)}(a+y,b+k) \sim W(a+y,b+k)$$

uniformly in $(y, k) \in I_n$. Consequently,

$$\left| (n-k)^{p(c)/2} Q_{k,n}(y) - \varkappa W(a+y,b+k) \right| \le 2\widetilde{\varepsilon}_n W(a+y,b+k)$$
(65)

uniformly (y, k) such that $k \leq n/2$ and $y \in I_n$.

We conclude then

$$\left| (n-k)^{p(c)/2} Q_{k,n}(y) - \varkappa W(a+y,b+k) \right|$$

$$\leq 2\widetilde{\varepsilon}_n W(a+y,b+k) + (M+\varkappa) W(a+y,b+k) \mathbb{1}\{(y,k) \notin I_n\}.$$
(66)

For every m < n we define

$$\nu_m := \inf\{k \ge 1 : S(k) - g_{a,b}(k) > \sqrt{m}\} \land m.$$

By the strong Markov property at time ν_m ,

$$\mathbf{P}(T_{a,b} > n) = \sum_{k=1}^{m} \int_{\mathbb{R}} \mathbf{P}(S_k \in dy, \nu_m = k < T_{a,b}) Q_{k,n}(y).$$

Applying (66), we have

$$\begin{aligned} \left| n^{p(c)/2} \mathbf{P}(T_{a,b} > n) - \varkappa \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m] \right| \\ + 2\widetilde{\varepsilon}_n \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m] \\ + C \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, (S(\nu_m), \nu_m) \notin I_n]. \end{aligned}$$

Since ν_m is a bounded stopping time,

$$\mathbf{E}[W(a+S(\nu_m),b+\nu_m);T_{a,b}>\nu_m]=W(a,b)$$

and, consequently,

$$\left| n^{p(c)/2} \mathbf{P}(T_{a,b} > n) - (\varkappa + o(1)) W(a,b) \right| \\
\leq C \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, (S(\nu_m), \nu_m) \notin I_n]. \quad (67)$$

We are left to show that the right-hand side converges to 0 for an appropriately chosen sequence m = m(n).

Lemma 26. There exists a constant C such that

$$\mathbf{P}(T_{a,b} > \nu_m) \le C \frac{W(a,b)}{m^{\frac{p(c)}{2}}}, \quad for \ all \ m \ge b.$$

Proof. It is immediate from the definition of ν_m that

$$\mathbf{P}(T_{a,b} > \nu_m) = \mathbf{P}(T_{a,b} > \nu_m = m) + \mathbf{P}(T_{a,b} > \nu_m, \nu_m < m)$$

$$\leq \mathbf{P}(T_{a,b} > m) + \mathbf{P}(T_{a,b} > \nu_m, S(\nu_m) > g_{a,b}(\nu_m) + \sqrt{m}).$$
(68)

According to Corollary 2,

$$\mathbf{P}(T_{a,b} > m) \le M \frac{W(a,b)}{m^{p(c)/2}},$$

for some constant M.

If $y > g_{a,b}(k) + \sqrt{m}$ and $k \le m$ then

$$\begin{split} W(a+y,b+k) &\geq W(a+g_{a,b}(k),b+k) = W(c\sqrt{b+k}+\sqrt{m},b+k) \\ &\geq CV_{p(c)}(c\sqrt{b+k}+\sqrt{m}) \geq Cm^{p(c)/2} \end{split}$$

for some positive constant C. Using this fact and applying the Markov inequality, we have

$$\mathbf{P}(T_{a,b} > \nu_m, S(\nu_m) > g_{a,b}(\nu_m) + \sqrt{m}) \le C \frac{\mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m]}{m^{p(c)/2}}.$$

Since W is harmonic we further obtain that

$$\mathbf{P}(T_{a,b} > \nu_m, S(\nu_m) > g_{a,b}(\nu_m) + \sqrt{m}) \le C \frac{\mathbf{E}[W(a,b)]}{m^{p(c)/2}}.$$

Lemma 27. If $A_m \to \infty$ as $m \to \infty$ then

$$\mathbf{E}[W(a+S(\nu_m),b+\nu_m);T_{a,b}>\nu_m,S(\nu_m)>A_m\sqrt{m}]=o(1).$$
(69)

Proof. Combining Lemma 4 and Lemma 25 we have

$$W(x,t) \le C|x|^{p(c)}$$
 for all $x > (c+1)\sqrt{t}$.

Moreover, by the definition of ν_m ,

$$|S(\nu_m)| \le c\sqrt{\nu_{m-1} + b} - a + \sqrt{m} + X_{\nu_m} \le (|c| + 2)\sqrt{m} + X_{\nu_m}$$

for all sufficiently large values of m. This implies that there exists m_0 such that $\{S(\nu_m) > A_m\sqrt{m}\} \subset \{X_{\nu_m} > \frac{A_m}{2}\sqrt{m}\}$ and $W(a + S(\nu_m), b + \nu_m) \leq C(X_{\nu_m})^{p(c)}$ for all $n \geq n_0$. Combining these estimates with the strong Markov property, we obtain

$$\mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, S(\nu_m) > A_m \sqrt{m}]$$

$$\leq \sum_{k=1}^n \mathbf{P}(T_{a,b} > k-1) \mathbf{E}[X^{p(c)}; X > A_m \sqrt{m}/2].$$

When p(c) > 2 then by Corollary 2 expectation $\mathbf{E}[T_{a,b}]$ is finite and (69) follows then from the Markov inequality.

Assume now that $p(c) \leq 2$. Since in this case we additionally assume that $\mathbf{E}|X|^{2+\delta}$ is finite it follows that $\mathbf{E}[X^{p(c)}; X > A_m \sqrt{m}/2] = o(m^{p(c)/2-1-\delta/2})$. Using once again Corollary 2 we see that

$$\sum_{k=1}^{m} \mathbf{P}(T_{a,b} > k-1) \le Cm^{1-\frac{p(c)}{2}}.$$

Combining these two estimates we see that (69) holds for $p(c) \leq 2$ as well.

We can now complete the proof of Theorem 3 by showing that

$$E_n := \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, (S(\nu_m), \nu_m) \notin I_n] = o(1).$$

We have, $E_n \leq E_{n,1} + E_{n,2}$, where

$$E_{n,1} = \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, a + S(\nu_m) \ge \nu_n \sqrt{n} - |c|\sqrt{b + m}],$$

$$E_{n,2} = \mathbf{E}[W(a + S(\nu_m), b + \nu_m); T_{a,b} > \nu_m, a + S(\nu_m) - c\sqrt{b + \nu_m} \le \gamma_n \sqrt{n}].$$

When $\frac{\mu_n \sqrt{n}}{\sqrt{m}} \to \infty$ the first expectation $E_{n,1} = o(1)$ by Lemma 27. Since W(y,t) is increasing in y,

$$E_{n,2} \leq \mathbf{E}[W(\gamma_n \sqrt{n} + c\sqrt{b + \nu_m}, b + \nu_m); T_{a,b} > \nu_m].$$

Using Lemma 25 and $V_{p(c)}(x,t) \leq C(x-c\sqrt{t})^{p(c)}$ we obtain

$$E_{n,2} \leq C(\gamma_n \sqrt{n})^{p(c)} \mathbf{P}(T_{a,b} > \nu_m)$$

Applying now Lemma 26

$$E_{n,2} \le C \left(\frac{\gamma_n \sqrt{n}}{\sqrt{m}}\right)^{p(c)} = o(1)$$

for m = m(n) such that $\frac{m(n)}{\gamma_n^2 n} \to \infty$. Thus, $E_n \to 0$ if m(n) is such that $\frac{m(n)}{\gamma_n^2 n} \to \infty$ and $\frac{m(n)}{\mu_n^2 n} \to 0$. This completes the proof of Theorem 1.

References

- Blackwell, D. and Freedman, D. A remark on the coin-tossing game. Ann. Math. Statist., 35: 1345–1347, 1964.
- [2] Breiman, L. First exit times from a square root boundary. Proc. Fifth Berkeley Symp. Math. Stat. Prob., 2:1236-1249, 1965.
- [3] Denisov D., Hinrichs G., Sakhanenko A.I. and V.Wachtel. Crossing an asymptotically square root boundary by the Brownian motion. *Proceedings of the Steklov Institute of Mathematics*, **316**: 105-120, 2022.
- [4] Denisov D., Sakhanenko A.I., Terveer S. and Wachtel V. Crossing an asymptotically square root boundary by a random walk. *In progress*, 2025.
- [5] Denisov, D., Sakhanenko, A. and Wachtel, V. First-passage times for random walks with non-identically distributed increments. Ann. Probab., 46(6), 3313–3350, 2018.
- [6] Denisov, D., Sakhanenko, A. and Wachtel, V. First-passage times over moving boundaries for asymtotically stable walks. *Theory Probab. Appl.*, 63, 613–633, 2019.
- [7] Denisov, D. and Wachtel, V. Random walks in cones. Ann. Probab., 43: 992-1044, 2015.
- [8] Denisov, D. and Wachtel, V. Exact asymptotics for the moment of crossing a curved boundary by an asymptotically stable random walk. *Theory of Prob. Appl.* 60 (3): 481–500, 2016.
- [9] Greenwood, P. and Perkins, E. A conditioned limit theorem for random walk and Brownian local time on square root boundaries. Ann. Probab., 11: 227–261, 1983.
- [10] Harvey, N. J. A., Liaw, C., Perkins, E, and Randhawa, S. Optimal anytime regret with two experts. *Math. Stat. Learn.* 6:87–142, 2023.
- [11] McConnell, T.R. Exit times of N-dimensional random walks. Z. Wahrscheinlichkeitstheorie verw. Gebiete, 67:213–233, 1984.
- [12] Novikov, A.A. On stopping times for the Wiener process. *Theory Probab. Appl.*, 16:449-456, 1971.
- [13] Novikov, A.A. Martingale approach to first passage problems of nonlinear boundaries. Proc. Steklov Inst. Math., 158:130-152.
- [14] Oldham, K., Myland, J., and Spanier, J. An atlas of functions. 2nd ed., Springer, New York, 2009.
- [15] Uchiyama, K. Brownian first exit from and sojourn over one-sided moving boundary and application. Z. Wahrsch. Verw. Gebiete, 54:75-116, 1980.

Department of Mathematics, University of Manchester, Oxford Road, Manchester M13 9PL, UK

Email address: denis.denisov@manchester.ac.uk

SOBOLEV INSTITUTE OF MATHEMATICS, RUSSIA Email address: aisakh@mail.ru

Mathematisches Institut, Ludwig-Maximilians-Universität München, Theresienstrasse 39, 80333 München, Germany

Email address: terveer@math.lmu.de

FACULTY OF MATHEMATICS, BIELEFELD UNIVERSITY, GERMANY *Email address*: wachtel@math.uni-bielefeld.de