Flavor-changing Lorentz and CPT violation in muonic atoms

V. Alan Kostelecký,¹ W.P. McNulty,¹ E. Passemar,^{1,2} and N. Sherrill^{3,4}

¹Physics Department, Indiana University, Bloomington, Indiana 47405, USA ²Departament de Física Teòrica, IFIC,

> Universitat de València, E-46071 València, Spain ³Department of Physics and Astronomy,

University of Sussex, Brighton BN1 9QH, UK and

⁴Institute of Theoretical Physics, Leibniz Universität Hannover,

Appelstr. 2, 30167 Hannover, Germany

(Dated: January 2025)

Abstract

Flavor-changing signatures of Lorentz and CPT violation involving muon-electron conversions in muonic atoms are studied using effective field theory. Constraints on coefficients for Lorentz violation at parts in 10^{-12} GeV⁻¹ for flavor-changing electromagnetic muon decays and parts in 10^{-13} GeV⁻² for flavor-changing 4-point quark-lepton interactions are extracted using existing data from the SINDRUM II experiment at the Paul Scherrer Institute. Estimates are provided for sensitivities attainable in the forthcoming experiments Mu2e at Fermilab and COMET at the Japan Proton Accelerator Complex. Flavor-changing processes in the minimal Standard Model (SM) arise from the weak interactions, which at tree level can convert charged leptons to neutrinos or mix quark flavors. The existence of flavor-changing physics beyond the SM is revealed by neutrino oscillations [1, 2], which establish lepton-number violation. This implies that flavor-changing conversions between charged leptons must also occur in nature, but only via suppressed oneloop processes with branching ratios $\leq 10^{-54}$ [3]. Experimental searches for charged-lepton flavor-changing (CLFC) processes therefore offer particularly clean probes of additional new physics beyond the SM [4–6].

One prospective type of CLFC physics beyond the SM is minuscule violations of Lorentz invariance and its associated CPT symmetry, which can emerge at low energies from an underlying theory unifying gravity and quantum physics such as strings [7]. Numerous experimental searches for flavor-conserving Lorentz violation (LV) have been performed [8], but studies of flavor-changing LV have largely been relegated to flavor mixing of species during propagation. However, the general model-independent framework for LV based on effective field theory [9–12] naturally incorporates terms describing CLFC interactions, of which only a subset contributing to muon decays has been explored to date [9, 13–22]. Here, we use existing data to revisit bounds from electromagnetic decays and to obtain first constraints on CLFC 4-point quark-lepton interactions.

A golden channel for experimental studies of CLFC processes, which currently is the subject of a worldwide experimental effort [23], is the coherent conversion $\mu + N \rightarrow e + N$ of a muon μ into an electron e in the presence of a nucleus N. The experimental procedure entails the capture of a muon in the ground state of a target atom, followed by its direct conversion into an electron with a characteristic monoenergetic spectrum. In model-independent effective field theory for LV searches, the leading contributions to this channel appear already at tree level instead of suppressed loop processes. No observable CLFC effects in muon decay involve LV terms of mass dimension d = 3 or 4 [22], so the dominant contributions arise via electromagnetic interactions with d = 5 and 4-point quark-lepton interactions with d = 6rather than from standard LV propagator terms [24]. Since the 4-point LV quark-lepton interactions involve the quarks in the nucleus, the corresponding CLFC effects are uniquely accessed in the conversion $\mu + N \rightarrow e + N$. No experimental constraints on these effects have been reported to date. Note that terms with d = 5 or 6 appear naturally in certain models such as noncommutative quantum field theories [25]. Their magnitudes are theoretically undetermined but are expected either to be suppressed by the scale of unification or by countershading effects [26].

In the experiments of interest here, a proton beam hits a production target, generating pions and other hadrons that decay to muons and other products. The muons with negative charge are collected and directed as a secondary beam onto a stopping target, where some are captured to form muonic atoms. The muonic atoms quickly decay to the 1S ground state. The signal for a neutrinoless muon-electron conversion is a monoenergetic electron ejected from a muonic atom with energy $E_e^{\text{conv}} = m_{\mu} - E_{\text{bind}} - E_{\text{recoil}}$, where m_{μ} is the muon mass, E_{bind} is the muon binding energy, and E_{recoil} is the nuclear recoil energy. This process can be characterized experimentally by the dimensionless ratio $R_{\mu e}$ of the conversion rate ω_{conv} to the capture rate ω_{capt} . The dominant contribution to ω_{conv} is the coherent conversion $\mu^- + \frac{A}{Z}N \rightarrow e^- + \frac{A}{Z}N$, where the nucleus $\frac{A}{Z}N$ of the muonic atom has atomic number A and charge Z. The ratio $R_{\mu e}$ can then be written as

$$R_{\mu e} = \frac{\omega_{\text{conv}}}{\omega_{\text{capt}}} = \frac{\Gamma[\mu^- + \frac{A}{Z}N \to e^- + \frac{A}{Z}N]}{\Gamma[(\mu^- + \frac{A}{Z}N \to \nu_\mu + \frac{A}{Z-1}N)]}.$$
(1)

To date, the experiment SINDRUM II [27] at the Paul Scherrer Institute has achieved the most stringent constraint, $R_{\mu e} < 7 \times 10^{-13}$ at the 90% confidence level (CL), using ${}^{197}_{79}$ Au as the target. Upcoming searches using an ${}^{27}_{13}$ Al target include the Muon-to-Electron Conversion (Mu2e) experiment at Fermilab [28], which anticipates achieving $R_{\mu e} \simeq 6.2 \times 10^{-16}$, and the Coherent Muon to Electron Transition (COMET) experiment [29] at the Japan Proton Accelerator Complex (J-PARC), which expects to reach $R_{\mu e} \simeq 7 \times 10^{-15}$. The forthcoming DeeMe experiment at J-PARC [30], using graphite and with a different configuration combining the production and stopping target, expects sensitivity at the 10^{-14} level. Planned searches for the more distant future include phase two of Mu2e at Fermilab [31], and phase two of COMET [32] and other experiments [33] at J-PARC, which are expected to attain $R_{\mu e}$ values of order 10^{-18} .

Here, we consider the experiments SINDRUM II, Mu2e, and COMET, which are performed with muonic gold or aluminum atoms. For ¹⁹⁷₇₉Au the capture rate is $\omega_{\text{capt}} \simeq$ 13.07 MHz, while for ²⁷₁₃Al the capture rate is $\omega_{\text{capt}} \simeq 0.7054$ MHz [34]. The capture rate typically increases with Z due to a greater overlap between the muon and nucleus wavefunctions, but separating the leptonic and nuclear physics becomes more challenging for heavier nuclei. The conversion rate $\omega_{\text{conv}} = |\mathcal{M}|^2$ is the appropriate spin-averaged squared modulus of the transition amplitude \mathcal{M} for the decay. We focus here on the coherent contribution where the nucleus remains in the ground state, which is experimentally the cleanest channel and is enhanced relative to incoherent conversion [35]. It has recently been shown that additional sensitivities to CLFC physics can be accessed from studies of the near-endpoint spectrum [36], and a corresponding analysis in the LV context would be a worthwhile topic for future investigation. Neglecting corrections below parts in 10³, the ground-state nuclear Coulomb field $\rho(r)$ can be taken as approximately static and spherically symmetric [37], and the nucleus can be treated as infinitely massive without nuclear recoil and reduced-mass corrections. The electron energy is then $E_e^{\text{conv}} \approx m_{\mu} - E_{\text{bind}}$ where $E_{\text{bind}} \ll m_{\mu}$.

The leptonic wavefunctions can be written in the form

$$\psi_{\kappa,s}(r,\theta,\phi) = \begin{pmatrix} g(r)\chi_{\kappa,s}(\theta,\phi)\\ if(r)\chi_{-\kappa,s}(\theta,\phi) \end{pmatrix},\tag{2}$$

where $\kappa = \mp (J+1/2)$ and the eigenspinors $\chi_{\kappa,s}$ satisfy $(\boldsymbol{\sigma} \cdot \boldsymbol{l}+I)\chi_{\kappa,s} = -\kappa\chi_{\kappa,s}$ and $J_z\chi_{\kappa,s} = s\chi_{\kappa,s}$, with normalization $\int d\Omega \ \chi^{\dagger}_{\kappa',s'}\chi_{\kappa,s} = \delta_{\mu'\mu}\delta_{\kappa'\kappa}$. Note that the muon wavefunction in the 1S orbital has $\kappa = -1$, and also that that $g^+ = if^-$ and $f^+ = -ig^-$ for a massless electron. The radial wavefunctions are obtained by numerically solving the Dirac equation

$$\frac{d}{dr} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix} = \begin{pmatrix} -\kappa/r & W - V(r) + m \\ -(W - V(r) - m) & \kappa/r \end{pmatrix} \begin{pmatrix} u_1 \\ u_2 \end{pmatrix}$$
(3)

for $u_1(r) = rg(r)$ and $u_2(r) = rf(r)$. Here, W is the lepton energy and V(r) is the nuclear electric potential. The total muon wavefunction is normalized to unity and so the muon field has mass dimension 3/2, while the electron wavefunction is normalized as a plane wave and hence the electron field has mass dimension 1. The potential V(r) is obtained from $\rho(r)$, which is normalized as $\int_0^\infty 4\pi\rho(r)r^2 dr = Z$ for a nucleus of atomic number Z. The functional form of $\rho(r)$ is taken as a two-parameter Fermi model for ¹⁹⁷₇₉Au and as a Fourier-Bessel expansion for ²⁷₁₃Al [38, 39].

When the muon-electron conversion is mediated via LV electromagnetic interactions with d = 5, the conversion rate takes the form

$$\omega_{\rm conv} = \frac{1}{2} \sum_{s=\pm\frac{1}{2}} \sum_{\kappa=\pm1} \sum_{s'=\pm\frac{1}{2}} \left| \int d^3 x F_{\alpha\beta} \overline{\psi}^{(e)}_{\kappa,s'} \mathcal{O}^{\alpha\beta} \psi^{(\mu)}_s \right|^2.$$
(4)

Here, $F_{\alpha\beta}$ is the electromagnetic field strength tensor, $\psi_{\kappa,s'}^{(e)}$ is a continuum electron wavefunction of energy W, and $\psi_s^{(\mu)}$ is the bound-state muon wavefunction in the 1S state. The sums implement the effects of the spins and angular momenta, with the factor of one half arising in averaging the muon spin. The matrix $\mathcal{O}^{\alpha\beta}$ contains the d = 5 coefficients for LV and appropriate Dirac matrices, which form the set [11, 22] { $(m_F^{(5)})_{\mu e}^{\alpha\beta}, i(m_{5F}^{(5)})_{\mu e}^{\alpha\beta}\gamma_5,$ $(a_F^{(5)})_{\mu e}^{\lambda\alpha\beta}\gamma_\lambda, (b_F^{(5)})_{\mu e}^{\lambda\alpha\beta}\gamma_5\gamma_\lambda, \frac{1}{2}(H_F^{(5)})_{\mu e}^{\kappa\lambda\alpha\beta}\sigma_{\kappa\lambda}$ }. These modify interactions rather than propagators, so standard perturbative quantization methods hold [40] and the conversion rate (4) can be derived using tree-level Feynman rules. Also, in contrast to conventional Lorentzinvariant treatments, neutrino flavor-changing LV effects [41] can be neglected here because no loop diagrams are involved.

In an inertial frame in the vicinity of the Earth, the cartesian coefficients for LV appearing in $\mathcal{O}^{\alpha\beta}$ can be taken to be independent of time and location [10]. The coefficients carry spacetime indices and so can change under observer Lorentz transformations, which implies that experimental measurements of their values must be provided in a specified inertial frame. The standard choice in the literature is the Sun-centered frame (SCF) with right-handed cartesian coordinates (T, X, Y, Z), where the time origin T = 0 is defined as the 2000 vernal equinox, the X axis is chosen to point from the Earth to the Sun at the time zero, and the Z axis parallels the rotation axis of the Earth [42]. We therefore must determine the transformation from the SCF to the detector frame (DF). For this purpose, it is convenient to fix also a standard Earth-based laboratory frame (LF), with right-handed cartesian coordinates (x, y, z) having x axis pointing to local south and y axis to local east.

Any laboratory frame distant from the poles is noninertial due to the rotation of the Earth, which has sidereal frequency $\omega_{\oplus} \simeq 2\pi/(23 \text{ h} 56 \text{ min})$. The boost of the Earth in the SCF is small, so at leading order the transformation from the SCF to the LF is a rotation $\mathcal{R}(\chi, \omega_{\oplus} T_{\oplus})$ that depends on the laboratory colatitude χ and is harmonic in the laboratory sidereal time $T_{\oplus} \equiv T - T_0$. The time T_{\oplus} in the LF is shifted by an experiment-dependent amount T_0 relative to the time T in the SCF, arising from the laboratory longitude λ and other effects [43]. The explicit form of the rotation $\mathcal{R}(\chi, \omega_{\oplus} T_{\oplus})$ is given as Eq. (7) of Ref. [22], and it can be used to transform coefficient components from the SCF to the LF. The DF typically differs from the standard LF, so we must perform an additional transformation $\mathcal{R}_{\text{detector}}(\psi)$ given in Eq. (6) of Ref. [22] that involves the angle ψ of the laboratory z-axis measured north of east along the direction of the beamline. The quantities χ , λ , T_0 , and ψ for the experiments considered here are listed in Table I.

The above considerations show that the net transformation \mathcal{R}_{total} from the SCF to the

Experiment	χ	λ	T_0	ψ	с	\widetilde{I}_1	\widetilde{I}_2	\widetilde{I}_3	\widetilde{I}_4
SINDRUM II	42.5°	8.2°	$3.86~\mathrm{h}$	242°	0.44	-0.053	0.070	0.040	0.18
COMET	53.6°	140.6°	-4.94 h	188°	0.34	-0.012	0.012	7.6×10^{-4}	0.041
Mu2e	48.2°	-88.2°	$10.27~\mathrm{h}$	122°	0.50	-0.012	0.012	7.6×10^{-4}	0.041

TABLE I. Detector geometric factors, parameters, and radial integrals.

DF is given by the combination

$$\mathcal{R}_{\text{total}} = \mathcal{R}_{\text{detector}}(\psi) \mathcal{R}(\chi, \omega_{\oplus} T_{\oplus}).$$
(5)

The components of the coefficients for LV expressed in the DF therefore depend on the detector orientation and location, and they can oscillate at harmonics of the sidereal frequency [44]. This reveals that the conversion rate and the ratio $R_{\mu e}$ have corresponding dependences, $\omega_{\text{conv}} = \omega_{\text{conv}}(\psi, \chi, \omega_{\oplus} T_{\oplus})$ and $R_{\mu e} = R_{\mu e}(\psi, \chi, \omega_{\oplus} T_{\oplus})$. An experiment recording data with time stamps can measure the amplitudes and phases of the harmonics by binning the data in sidereal time and hence can detect or constrain the various components of the coefficients for LV in the SCF.

The spherical symmetry of the nuclear charge distribution makes it convenient for some calculations to adopt DF spherical-polar coordinates (r, θ, ϕ) with z axis directed along the muon beam direction. Results must then be matched to DF cartesian coordinates to apply the map (5) from the SCF to the DF. The spherical symmetry also implies that the nonzero contributions to the rate (4) from the field-strength tensor arise only from the components $F_{tr} = -F_{rt} = E_r(r)$, where the index t represents the time T_{\oplus} . Like $F_{\alpha\beta}$, the matrix $\mathcal{O}^{\alpha\beta}$ is antisymmetric in $\alpha\beta$, so it suffices to consider the case $\alpha = t$, $\beta = r$. The matrix component \mathcal{O}^{tr} is a spatial vector, so the match between polar and cartesian coordinates in the DF can be accomplished by $\mathcal{O}^{tr} = \mathcal{O}^{tx} \sin \theta \cos \phi + \mathcal{O}^{ty} \sin \theta \sin \phi + \mathcal{O}^{tz} \cos \theta$, as usual. These additional angular factors must be taken into account when performing the integration in Eq. (4) to obtain the conversion rate ω_{conv} .

Working in DF spherical-polar coordinates, the integral (4) can be taken over the full azimuthal range $\phi \in [0, 2\pi)$. However, the angle θ is limited by the detector acceptance to a range $|\cos \theta| \leq c$. For the experiments considered here, approximate values of the acceptances c are listed in Table I. For the angular part of the integral (4), the choices $\kappa = \pm 1$, $s = \pm 1/2$, and $s' = \pm 1/2$ yield eight possible cases in principle. However, for any given coefficient for LV, only two nonzero cases occur in practice. They depend on the acceptance c for the given experiment, via two geometrical factors $w_{xy} = -c^3/6 + c/2$ and $w_z = c^3/3$. For the radial part two integrals of mass dimension 3/2 occur,

$$I_{1} = \int dr \ r^{2} E(r) \left(f_{e}^{-} g_{\mu}^{-} + g_{e}^{-} f_{\mu}^{-} \right),$$

$$I_{2} = \int dr \ r^{2} E(r) \left(g_{e}^{-} f_{\mu}^{-} - f_{e}^{-} g_{\mu}^{-} \right).$$
(6)

These integrals also take unique values for each experiment. Table I displays numerical values of the quantities $\tilde{I}_1 = I_1/m_{\mu}^{3/2}$ and $\tilde{I}_2 = I_2/m_{\mu}^{3/2}$ for the various experiments we analyze here. Combining these results and converting to cartesian coordinates gives the desired equations for the transition amplitudes \mathcal{M} and hence for the conversion rate $\omega_{\rm conv}$ expressed using coefficients for LV in the DF. Applying the transformation (5) then yields $\omega_{\rm conv}$ in terms of coefficients for LV in the SCF. Explicit forms for these results are presented in the Appendix.

The SINDRUM II bound was obtained using a primary dataset taken over a period of 81 days [27]. It can therefore be intepreted as a limit on the time-averaged conversion rate $\overline{\omega}_{\text{conv}} = \overline{\omega}_{\text{conv}}(\psi, \chi)$ for the corresponding ψ and χ values given in Table I. Adopting the standard methodology in the literature [8], we can translate this experimental limit into constraints on the coefficients for LV in the SCF, taken one at a time. The key expressions involved in this procedure are presented in the Appendix, and the ensuing constraints are tabulated in Table II. In the table, each entry is a bound at 90% CL on a coefficient component in the SCF, where the indices J and $K \neq J$ are X or Y. The table reveals that SINDRUM II achieved sensitivities down to a few parts in 10¹² to 96 real components of the coefficients for LV at d = 5. Note that the constraints involving the coefficients $(a_F^{(5)})_{\mu e}^{\lambda \alpha \beta}$ and $(b_F^{(5)})_{\mu e}^{\lambda \alpha \beta}$ also represent limits on CPT violation. The same methodology can be applied to the anticipated reaches of the Mu2e and COMET experiments, yielding the estimated attainable sensitivities displayed in Table II. Together with the enhanced resolving power arising via the separation of harmonics in sidereal time, these estimates suggest improvements of one to two orders of magnitude over the SINDRUM II results are feasible within the near future.

If instead the muon-electron conversion is mediated through 4-point interactions coupling the nuclear quarks to the leptons with d = 6, the conversion rate is

$$\omega_{\rm conv} = \frac{1}{2} \sum_{s,s',\kappa} \left| \int d^3 x \left(\alpha \overline{\psi}_{\kappa,s'}^{(e)} \mathcal{K} \psi_s^{(\mu)} + \beta \overline{\psi}_{\kappa,s'}^{(e)} \mathcal{K}_0 \psi_s^{(\mu)} \right) \right|^2, \tag{7}$$

Coefficients	SINDRUM I	I COMET	Mu2e
$ (m_F^{(5)})_{\mu e}^{TJ} , (m_{5F}^{(5)})_{\mu e}^{TJ} $	< 8	< 1	< 0.2
$ (m_F^{(5)})_{\mu e}^{TZ} , (m_{5F}^{(5)})_{\mu e}^{TZ} $	< 8	< 0.9	< 0.2
$ (a_F^{(5)})_{\mu e}^{TTJ} , (b_F^{(5)})_{\mu e}^{TTJ} $	< 6	< 1	< 0.2
$ (a_F^{(5)})_{\mu e}^{TTZ} , (b_F^{(5)})_{\mu e}^{TTZ} $	< 6	< 0.9	< 0.2
$ (a_F^{(5)})_{\mu e}^{JTJ} , (b_F^{(5)})_{\mu e}^{JTJ} $	< 6	< 1	< 0.2
$ (a_F^{(5)})_{\mu e}^{JTK} , (b_F^{(5)})_{\mu e}^{JTK} $	< 8	< 1	< 0.2
$ (a_F^{(5)})_{\mu e}^{JTZ} , (b_F^{(5)})_{\mu e}^{JTZ} $	< 8	< 0.9	< 0.2
$ (a_F^{(5)})_{\mu e}^{ZTJ} , (b_F^{(5)})_{\mu e}^{ZTJ} $	< 8	< 1	< 0.2
$ (a_F^{(5)})_{\mu e}^{ZTZ} , (b_F^{(5)})_{\mu e}^{ZTZ} $	< 7	< 0.9	< 0.2
$ (H_F^{(5)})_{\mu e}^{TJTJ} , (H_F^{(5)})_{\mu e}^{JZTK} $	< 7	< 1	< 0.2
$ (H_F^{(5)})_{\mu e}^{TJTK} , (H_F^{(5)})_{\mu e}^{JZTJ} $	< 6	< 1	< 0.2
$ (H_F^{(5)})_{\mu e}^{TJTZ} , (H_F^{(5)})_{\mu e}^{JZTZ} $	< 6	< 0.9	< 0.2
$ (H_F^{(5)})_{\mu e}^{TZTJ} , (H_F^{(5)})_{\mu e}^{XYTJ} $	< 6	< 1	< 0.2
$ (H_F^{(5)})_{\mu e}^{TZTZ} , (H_F^{(5)})_{\mu e}^{XYTZ} $	< 7	< 0.9	< 0.2

TABLE II. Constraints on d = 5 coefficients for LV in units of 10^{-12} GeV⁻¹. The results for SIN-DRUM II are constraints at 90% CL. The results for Mu2e and COMET are projected constraints based on expected rate sensitivities.

where α and β are conventional nuclear matrix elements given by $\alpha = \langle N | \overline{\psi}^{(q)} \psi^{(q)} | N \rangle$ with q summed over the quark flavors q = u, d, s and $\beta = \langle N | \overline{\psi}^{(q)} \gamma_0 \psi^{(q)} | N \rangle$ with q summed over q = u, d. Other possible nuclear matrix elements vanish in coherent conversion [39]. The operators \mathcal{K} and \mathcal{K}_0 incorporate the d = 6 coefficients for LV and appropriate Dirac matrices [11], with \mathcal{K} drawn from the set $\{(k_{SV}^{(6)})_{qqe\mu}^{\lambda}\gamma_{\lambda}, (k_{SA}^{(6)})_{qqe\mu}^{\lambda}\gamma_{5}\gamma_{\lambda}, \frac{1}{2}(k_{ST}^{(6)})_{qqe\mu}^{\kappa\lambda} \sigma_{\kappa\lambda}\}$ for q = u, d. These arise from $\{(k_{VS}^{(6)})_{qqe\mu}^{t}, i(k_{VP}^{(6)})_{qqe\mu}^{t}\gamma_{5}, (k_{VV}^{(6)})_{qqe\mu}^{t\lambda}\gamma_{\lambda}, (k_{VA}^{(6)})_{qqe\mu}^{t\lambda}\gamma_{5}\gamma_{\lambda}, \frac{1}{2}(k_{VT}^{(6)})_{qqe\mu}^{t\kappa\lambda} \sigma_{\kappa\lambda}\}$ for q = u, d. These arise from 4-point quark-lepton interactions in the underlying effective field theory, for which the standard perturbative quantization yields the expression (7) arising from vertex contributions to the Feynman rules.

In DF spherical-polar coordinates, the angular part of the integral (7) involves eight possible cases $\kappa = \pm 1$, $s = \pm 1/2$, and $s' = \pm 1/2$. Their values depend on the acceptance cfor the given experiment. The radial part involves two integrals of mass dimension 5/2,

$$I_3 = \int dr \ r^2 \rho^{(p)}(r) f_e^- f_\mu^-, \quad I_4 = \int dr \ r^2 \rho^{(p)}(r) g_e^- g_\mu^-.$$
(8)

For the experiments considered here, numerical values of $\tilde{I}_3 = I_3/m_{\mu}^{5/2}$ and $\tilde{I}_4 = I_4/m_{\mu}^{5/2}$ are

Coefficients	SINDRUM II	COMET	Mu2e
$ (k_{SV}^{(6)})_{uue\mu}^T , (k_{SA}^{(6)})_{uue\mu}^T $	< 6	< 1	< 0.2
$ (k_{SV}^{(6)})_{uue\mu}^{J} , (k_{SA}^{(6)})_{uue\mu}^{J} $	< 7	< 1	< 0.2
$ (k_{SV}^{(6)})_{uue\mu}^Z , (k_{SA}^{(6)})_{uue\mu}^Z $	< 7	< 1	< 0.2
$ (k_{SV}^{(6)})_{dde\mu}^{T} , (k_{SA}^{(6)})_{dde\mu}^{T} $	< 6	< 1	< 0.2
$ (k_{SV}^{(6)})_{dde\mu}^{J} , (k_{SA}^{(6)})_{dde\mu}^{J} $	< 7	< 1	< 0.2
$ (k_{SV}^{(6)})_{dde\mu}^{Z} , (k_{SA}^{(6)})_{dde\mu}^{Z} $	< 7	< 1	< 0.2
$ (k_{SV}^{(6)})_{sse\mu}^T , (k_{SA}^{(6)})_{sse\mu}^T $	< 10	< 2	< 0.4
$ (k_{SV}^{(6)})_{sse\mu}^{J} , (k_{SA}^{(6)})_{sse\mu}^{J} $	< 15	< 2	< 0.4
$ (k_{SV}^{(6)})_{sse\mu}^Z , (k_{SA}^{(6)})_{sse\mu}^Z $	< 15	< 2	< 0.4
$ (k_{VS}^{(6)})_{uue\mu}^T , (k_{VP}^{(6)})_{uue\mu}^T $	< 30	< 4	< 0.8
$ (k_{VS}^{(6)})_{dde\mu}^T , (k_{VP}^{(6)})_{dde\mu}^T $	< 30	< 4	< 0.7
$ (k_{ST}^{(6)})_{uue\mu}^{TJ} , (k_{ST}^{(6)})_{uue\mu}^{JZ} $	< 7	< 1	< 0.2
$ (k_{ST}^{(6)})_{uue\mu}^{TZ} , (k_{ST}^{(6)})_{uue\mu}^{XY} $	< 7	< 1	< 0.2
$ (k_{ST}^{(6)})_{dde\mu}^{TJ} , (k_{ST}^{(6)})_{dde\mu}^{JZ} $	< 7	< 1	< 0.2
$ (k_{ST}^{(6)})_{dde\mu}^{TZ} , (k_{ST}^{(6)})_{dde\mu}^{XY} $	< 7	< 1	< 0.2
$ (k_{ST}^{(6)})_{sse\mu}^{TJ} , (k_{ST}^{(6)})_{sse\mu}^{JZ} $	< 15	< 2	< 0.4
$ (k_{ST}^{(6)})_{sse\mu}^{TZ} , (k_{ST}^{(6)})_{sse\mu}^{XY} $	< 15	< 2	< 0.4
$ (k_{VV}^{(6)})_{uue\mu}^{TT} , (k_{VA}^{(6)})_{uue\mu}^{TT} $	< 20	< 4	< 0.7
$ (k_{VV}^{(6)})_{uue\mu}^{TJ} , (k_{VA}^{(6)})_{uue\mu}^{TJ} $	< 25	< 4	< 0.7
$ (k_{VV}^{(6)})_{uue\mu}^{TZ} , (k_{VA}^{(6)})_{uue\mu}^{TZ} $	< 25	< 4	< 0.7
$ (k_{VV}^{(6)})_{dde\mu}^{TT} , (k_{VA}^{(6)})_{dde\mu}^{TT} $	< 20	< 4	< 0.7
$ (k_{VV}^{(6)})_{dde\mu}^{TJ} , (k_{VA}^{(6)})_{dde\mu}^{TJ} $	< 20	< 4	< 0.7
$ (k_{VV}^{(6)})_{dde\mu}^{TZ} , (k_{VA}^{(6)})_{dde\mu}^{TZ} $	< 20	< 4	< 0.7
$ (k_{VT}^{(6)})_{uue\mu}^{TTJ} , (k_{VT}^{(6)})_{uue\mu}^{TJZ} $	< 25	< 4	< 0.7
$ (k_{VT}^{(6)})_{uue\mu}^{TTZ} , (k_{VT}^{(6)})_{uue\mu}^{TXY} $	< 25	< 4	< 0.7
$ (k_{VT}^{(6)})_{dde\mu}^{TTJ} , (k_{VT}^{(6)})_{dde\mu}^{TJZ} $	< 20	< 4	< 0.7
$ (k_{VT}^{(6)})_{dde\mu}^{TTZ} , (k_{VT}^{(6)})_{dde\mu}^{TXY} $	< 20	< 4	< 0.7

TABLE III. Constraints on d = 6 coefficients for LV in units of 10^{-13} GeV⁻². The results for SIN-DRUM II are constraints at 90% CL. The results for Mu2e and COMET are projected constraints based on expected rate sensitivities.

given in Table I. These equations permit evaluation of the transition amplitudes \mathcal{M} and hence of ω_{conv} expressed using coefficients for LV in the DF. The result for ω_{conv} in terms of coefficients for LV in the SCF can then be obtained by applying the transformation (5). Some details of this procedure are provided in the Appendix. Interpreting the limit from SINDRUM II as a bound on the time-averaged conversion rate $\overline{\omega}_{conv}$, we can establish first constraints in the SCF on each component of coefficients for d = 6 LV taken in turn. These constraints are presented in Table III, where J is X or Y. The results show that SINDRUM II attained sensitivities down to parts in 10¹³ to CLFC effects from 148 real components of coefficients for LV arising from 4-point quarklepton interactions. Note that bounds on coefficients with one or three spacetime indices are also constraints on CPT violation. Table III also displays projected sensitivities in the Mu2e and COMET experiments. Allowing also for analyses of the harmonic siderealtime variations, these results suggest that impressive enhancements in sensitivities and a corresponding significant discovery potential lie within future reach.

This work is supported in part by the U.S. Department of Energy under grants DE-SC0010120 and DE-AC05-06OR23177, by the U.S. National Science Foundation under grant PHY-2013184, by the Generalitat Valenciana (Spain) through the plan GenT program CIDEGENT/2021/037, by the Spanish Government Agencia Estatal de Investigación MCIN/AEI/10.13039/501100011033 under grants PID2020–114473GB-I00 and PID2023-146220NB-I00, by the Agencia Estatal de Investigación MCIU/AEI (Spain) under grant IFIC Centro de Excelencia Severo Ochoa CEX2023-001292-S, by the U.K. Science and Technology Facilities Council under grants ST/T006048/1 and ST/Y004418/1, by the Deutsche Forschungsgemeinschaft under the Heinz Maier Leibnitz Prize BeyondSM HML-537662082, and by the Indiana University Center for Spacetime Symmetries. Appendix: conversion rates. The conversion rate via d = 5 LV electromagnetic interactions is given by Eq. (4) and can be evaluated in the DF using spherical polar coordinates. Combining the geometrical factors w_{xy} and w_z with the radial integrals (6) for the allowed values of the muon spin $s = \pm 1/2$, electron spin $s' = \pm 1/2$, and electron angular momentum $\kappa = \pm 1$ yields expressions for the transition amplitudes $\mathcal{M}_{s',s}^{\kappa}$ in cartesian coordinates in the DF,

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{+1} = \mp i w_z (m_F^{(5)})_{\mu e}^{tz} I_1 \pm i w_z (a_F^{(5)})_{\mu e}^{ttz} I_2$$

$$\pm w_{xy} ((b_F^{(5)})_{\mu e}^{xty} - (b_F^{(5)})_{\mu e}^{ytx}) I_1$$

$$- i [w_{xy} ((b_F^{(5)})_{\mu e}^{xtx} + (b_F^{(5)})_{\mu e}^{yty}) + w_z (b_F^{(5)})_{\mu e}^{ztz}] I_2$$

$$\pm w_{xy} ((H_F^{(5)})_{\mu e}^{xztx} + (H_F^{(5)})_{\mu e}^{yzty}) I_2$$

$$+ i [w_{xy} ((H_F^{(5)})_{\mu e}^{xzty} - (H_F^{(5)})_{\mu e}^{yztx})$$

$$- w_z (H_F^{(5)})_{\mu e}^{xytz}] I_1, \qquad (9)$$

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{\pm1} = -iw_{xy}((m_F^{(5)})_{\mu e}^{tx} \pm i(m_F^{(5)})_{\mu e}^{ty})I_1 + iw_{xy}((a_F^{(5)})_{\mu e}^{ttx} \pm i(a_F^{(5)})_{\mu e}^{tty})I_2 + [(w_z(b_F^{(5)})_{\mu e}^{ytz} - w_{xy}(b_F^{(5)})_{\mu e}^{zty}) \pm i(w_z(b_F^{(5)})_{\mu e}^{xtz} - w_{xy}(b_F^{(5)})_{\mu e}^{ztx})]I_1 - [(w_{xy}(H_F^{(5)})_{\mu e}^{xyty} + w_z(H_F^{(5)})_{\mu e}^{xztz}) \mp i(w_{xy}(H_F^{(5)})_{\mu e}^{xytx} - w_z(H_F^{(5)})_{\mu e}^{yztz})]I_2,$$
(10)

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{-1} = \pm w_{z}(m_{5F}^{(5)})_{\mu e}^{tz}I_{1}$$

$$\pm w_{xy}((a_{F}^{(5)})_{\mu e}^{xty} - (a_{F}^{(5)})_{\mu e}^{ytx})I_{1}$$

$$- i[w_{xy}((a_{F}^{(5)})_{\mu e}^{xtx} + (a_{F}^{(5)})_{\mu e}^{yty})$$

$$+ w_{z}(a_{F}^{(5)})_{\mu e}^{ztz}]I_{2} \pm iw_{z}(b_{F}^{(5)})_{\mu e}^{ttz}I_{2}$$

$$\pm iw_{xy}((H_{F}^{(5)})_{\mu e}^{txty} - (H_{F}^{(5)})_{\mu e}^{tytx})I_{2}$$

$$+ [w_{xy}((H_{F}^{(5)})_{\mu e}^{txtx} + (H_{F}^{(5)})_{\mu e}^{tyty})$$

$$+ w_{z}(H_{F}^{(5)})_{\mu e}^{tztz}]I_{1}, \qquad (11)$$

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{-1} = w_{xy}((m_{5F}^{(5)})_{\mu e}^{tx} \pm i(m_{5F}^{(5)})_{\mu e}^{ty})I_{1} + [(w_{z}(a_{F}^{(5)})_{\mu e}^{ytz} - w_{xy}(a_{F}^{(5)})_{\mu e}^{zty}) \pm i(w_{z}(a_{F}^{(5)})_{\mu e}^{xtz} - w_{xy}(a_{F}^{(5)})_{\mu e}^{ztx})]I_{1} + iw_{xy}((b_{F}^{(5)})_{\mu e}^{tx} \pm i(b_{F}^{(5)})_{\mu e}^{ty})I_{2} \pm [(w_{z}(H_{F}^{(5)})_{\mu e}^{txtz} - w_{xy}(H_{F}^{(5)})_{\mu e}^{tztx}) + i(w_{z}(H_{F}^{(5)})_{\mu e}^{tytz} - w_{xy}(H_{F}^{(5)})_{\mu e}^{tzty})]I_{2}.$$
(12)

The conversion rate for any desired combination of coefficients in the DF can be obtained from these equations as $\omega_{\text{conv}} = \overline{|\mathcal{M}|^2}$.

To determine ω_{conv} in terms of coefficients for LV in the SCF, where the coefficients are constants and useful constraints can thus be established, we apply the map (5). This produces a lengthy expression containing harmonics of $\omega_{\oplus}T_{\oplus}$. For applications to datasets taken over extended time intevals, time averaging can be performed, which leaves only the time-independent contribution $\overline{\omega}_{\text{conv}}$. An experimental measurement of $\overline{\omega}_{\text{conv}}$ can thereby be translated into a limit on the components of the coefficients for LV taken one at a time. The constraints from SINDRUM II and the estimated sensitivities of Mu2e and COMET resulting from this procedure are presented in Table II.

As an illustration, consider the coefficient $(m_F^{(5)})_{\mu e}^{\alpha \beta}$. In the DF, we find

$$\omega_{\rm conv} = \left(w_{xy}^2 \sum_j |(m_F^{(5)})_{\mu e}^{tj}|^2 + w_z^2 |(m_F^{(5)})_{\mu e}^{tz}|^2 \right) I_1^2, \tag{13}$$

where j = x, y. Converting to the SCF and taking the time average yields $\overline{\omega}_{\text{conv}} = \zeta_{TJ}^{(0)} |(m_F^{(5)})_{\mu e}^{TJ}|^2$, where J = X, Y and $\zeta_{TJ}^{(0)}$ is an experiment-dependent quantity that depends on the geometric factors in Table I. For SINDRUM II, for example, we find $\zeta_{TJ}^{(0)} \simeq 9.5 \times 10^{-8}$ GeV³. This result provides the constraints on the coefficient components $(m_F^{(5)})_{\mu e}^{TJ}$ listed in the first row of Table II.

For completeness, we provide here the terms in $\omega_{\rm conv}$ involving the squared moduli of

each coefficient for LV expressed in the SCF,

$$\begin{aligned}
\omega_{\rm conv} \supset \left(\zeta_{TJ}^{(0)} + \zeta_{TJ}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + \zeta_{TJ}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}}\right) \\
\times \left(|(m_{F}^{(5)})_{\mu e}^{TJ}|^{2} + |(m_{5F}^{(5)})_{\mu e}^{TJ}|^{2}\right) \\
+ \left(\zeta_{\alpha TJ}^{(0)} + \zeta_{\alpha TJ}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + \zeta_{\alpha TJ}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}} \\
+ \zeta_{\alpha TJ}^{(4c)} c_{4\omega_{\oplus}T_{\oplus}} + \zeta_{\alpha TJ}^{(4s)} s_{4\omega_{\oplus}T_{\oplus}}\right) \\
\times \left(|(a_{F}^{(5)})_{\mu e}^{\alpha TJ}|^{2} + |(b_{F}^{(5)})_{\mu e}^{\alpha TJ}|^{2}\right) \\
+ \left(\zeta_{\alpha \beta TJ}^{(0)} + \zeta_{\alpha \beta TJ}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + \zeta_{\alpha \beta TJ}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}} \\
+ \zeta_{\alpha \beta TJ}^{(4c)} c_{4\omega_{\oplus}T_{\oplus}} + \zeta_{\alpha \beta TJ}^{(4s)} s_{4\omega_{\oplus}T_{\oplus}}\right)|(H_{F}^{(5)})_{\mu e}^{\alpha \beta TJ}|^{2},
\end{aligned}$$
(14)

where $c_{n\omega_{\oplus}T_{\oplus}} = \cos n\omega_{\oplus}T_{\oplus}$ and $s_{n\omega_{\oplus}T_{\oplus}} = \sin n\omega_{\oplus}T_{\oplus}$ for n = 2, 4. Terms in the full expression that are omitted above involve products of two different coefficients for LV, which are irrelevant for constraints obtained with one coefficient component taken nonzero at a time.

If instead the conversion is mediated by d = 6 LV operators governing 4-point quarklepton interactions, the conversion rate is given by Eq. (7). The values of the conventional nuclear matrix elements α and β can be expressed in terms of the nuclear proton density $\rho^{(p)}$, the neutron density $\rho^{(n)}$, A, Z, and numerical parameters $G_S^{(q,p)}$, $G_S^{(q,n)}$ [39]. The latter take the values $G_S^{(q,p)} = \{5.1, 4.3, 2.5\}$ and $G_S^{(q,n)} = \{4.3, 5.1, 2.5\}$ for $q = \{u, d, s\}$, respectively [45]. We take $\rho^{(p)}$ to be normalized to Z and $\rho^{(n)}$ to follow $\rho^{(p)}$ but to be normalized to A - Z, and we define $\tilde{G}_S^q \equiv [ZG_S^{(q,p)} + (A - Z)G_S^{(q,n)}]/Z$, $\tilde{G}_V^u \equiv (Z + A)/Z$, and $\tilde{G}_V^d \equiv (2A - Z)/Z$. Combining these results with geometric factors and the integrals (8) permits expressing the transition amplitude \mathcal{M} in terms of the combinations $I_{S\pm}^q = \pm c \tilde{G}_S^q (I_3 \pm I_4), I_{S1}^q = c \tilde{G}_S^q (\frac{1}{3}c^2I_3 + I_4)$, and $I_{S2}^q = c \tilde{G}_S^q [(1 - \frac{2}{3}c^2)I_3 + I_4]$ for q = u, d, s, along with $I_{V\pm}^q = \pm c \tilde{G}_V^q (I_3 \pm I_4), I_{V1}^q = c \tilde{G}_V^q (\frac{1}{3}c^2I_3 + I_4)$, and $I_{V2}^q = c \tilde{G}_V^q [(1 - \frac{2}{3}c^2)I_3 + I_4]$ for q = u, d only. The contributions to $\mathcal{M}_{s',s}^\kappa$ arising from the coefficients for LV at d = 6given in cartesian coordinates in the DF are

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{+1} = \pm I_{S2}^{q} (k_{SV}^{(6)})_{qqe\mu}^{z} + I_{S+}^{q} (k_{SA}^{(6)})_{qqe\mu}^{t} + i I_{V-}^{q} (k_{VP}^{(6)})_{qqe\mu}^{t} \pm i I_{S2}^{q} (k_{ST}^{(6)})_{qqe\mu}^{tz} \pm I_{V2}^{q} (k_{VV}^{(6)})_{qqe\mu}^{tz} + I_{V+}^{q} (k_{VA}^{(6)})_{qqe\mu}^{tt} \pm i I_{V2}^{q} (k_{VT}^{(6)})_{qqe\mu}^{ttz},$$
(15)

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{+1} = I_{S1}^{q}[(k_{SV}^{(6)})_{qqe\mu}^{x} \pm i(k_{SV}^{(6)})_{qqe\mu}^{y}]$$

$$\mp I_{S1}^{q}[(k_{ST}^{(6)})_{qqe\mu}^{ty} \mp i(k_{ST}^{(6)})_{qqe\mu}^{tx}]$$

$$+ I_{V1}^{q}[(k_{VV}^{(6)})_{qqe\mu}^{tx} \pm i(k_{VV}^{(6)})_{qqe\mu}^{ty}]$$

$$\mp I_{V1}^{q}[(k_{VT}^{(6)})_{qqe\mu}^{tty} \mp i(k_{VT}^{(6)})_{qqe\mu}^{ttx}], \qquad (16)$$

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{-1} = I_{S+}^{q} (k_{SV}^{(6)})_{qqe\mu}^{t} \pm I_{S2}^{q} (k_{SA}^{(6)})_{qqe\mu}^{z} + I_{V-}^{q} (k_{VS}^{(6)})_{qqe\mu}^{t} \pm i I_{S2}^{q} (k_{ST}^{(6)})_{qqe\mu}^{xy} + I_{V+}^{q} (k_{VV}^{(6)})_{qqe\mu}^{tt} \pm I_{V2}^{q} (k_{VA}^{(6)})_{qqe\mu}^{tz} \pm i I_{V2}^{q} (k_{VT}^{(6)})_{qqe\mu}^{txy},$$
(17)

$$\mathcal{M}_{\pm\frac{1}{2},\pm\frac{1}{2}}^{-1} = I_{S1}^{q}[(k_{SA}^{(6)})_{qqe\mu}^{x} \pm i(k_{SA}^{(6)})_{qqe\mu}^{y}] \\ + I_{S1}^{q}[(k_{ST}^{(6)})_{qqe\mu}^{yz} \mp i(k_{ST}^{(6)})_{qqe\mu}^{xz}] \\ + I_{V1}^{q}[(k_{VA}^{(6)})_{qqe\mu}^{tx} \pm i(k_{VA}^{(6)})_{qqe\mu}^{ty}] \\ + I_{V1}^{q}[(k_{VT}^{(6)})_{qqe\mu}^{tyz} \mp i(k_{VT}^{(6)})_{qqe\mu}^{txz}].$$
(18)

The conversion rate $\omega_{\text{conv}} = \overline{|\mathcal{M}|^2}$ corresponding to any given choice of coefficients in the DF can be derived directly from these equations.

.

In parallel with the d = 5 case, applying the map (5) transforms the coefficients for LV in ω_{conv} from the DF to the SCF, producing an expression involving constant coefficients for LV in the SCF and time-dependent harmonics of the sidereal frequency ω_{\oplus} . Averaging over many sidereal days eliminates all but the constant term $\overline{\omega}_{\text{conv}}$, which can be used to extract constraints on components of the coefficients for LV in turn. The ensuing constraints from SINDRUM II and the estimated sensitivities of Mu2e and COMET are displayed in Table III.

In the SCF, the contributions to $\omega_{\rm conv}$ proportional to the squared moduli of each coeffi-

cient for LV are

$$\begin{split} \omega_{\text{conv}} &\supset \left((\zeta_q)_{\alpha}^{(0)} + (\zeta_q)_{\alpha}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + (\zeta_q)_{\alpha}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}} \right) \\ &\times \left(| (k_{SV}^{(6)})_{qqe\mu}^{\alpha} |^2 + | (k_{SA}^{(6)})_{qqe\mu}^{\alpha} |^2 \right) \\ &+ (\zeta_q^0)_T^{(0)} (| (k_{VS}^{(6)})_{qqe\mu}^T |^2 + | (k_{VP}^{(6)})_{qqe\mu}^T |^2) \\ &+ \left[((\zeta_q)_{TJ}^{(0)} + (\zeta_q)_{TJ}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + (\zeta_q)_{TJ}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}} \right) \\ &\times | (k_{ST}^{(6)})_{qqe\mu}^T |^2 + (TJ \to JZ)] \\ &+ \left[(\zeta_q)_{TZ}^{(0)} | (k_{ST}^{(6)})_{qqe\mu}^T |^2 + (TZ \to JK) \right] \\ &+ ((\zeta_q^0)_{T\alpha}^{(0)} + (\zeta_q^0)_{T\alpha}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + (\zeta_q^0)_{T\alpha}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}}) \\ &\times (| (k_{VV}^{(6)})_{qqe\mu}^T |^2 + | (k_{VA}^{(6)})_{qqe\mu}^T |^2) \\ &+ \left[((\zeta_q^0)_{TTJ}^{(0)} + (\zeta_q^0)_{TTJ}^{(2c)} c_{2\omega_{\oplus}T_{\oplus}} + (\zeta_q^0)_{TTJ}^{(2s)} s_{2\omega_{\oplus}T_{\oplus}}) \right] \\ &\times | (k_{VT}^{(6)})_{qqe\mu}^T |^2 + (TTJ \to TJZ)] \\ &+ \left[(\zeta_q^0)_{TTZ}^{(0)} | (k_{VT}^{(6)})_{qqe\mu}^T |^2 + (TTZ \to TJK) \right], \end{split}$$

where a sum over the relevant quark flavors q is understood. Terms containing products of distinct coefficients for LV are omitted because they play no role when all coefficient components vanish but one.

- R. Davis, Jr., D.S. Harmer, and K.C. Hoffman, Phys. Rev. Lett. 20, 1205 (1968); Q.R. Ahmad et al., Phys. Rev. Lett. 87, 071301 (2001).
- [2] Y. Fukuda *et al.*, Phys. Rev. Lett. **81**, 1562 (1998).
- [3] W.J. Marciano and A.I. Sanda, Phys. Lett. B 67, 303 (1977); S.M. Bilenky, S.T. Petcov, and
 B. Pontecorvo, Phys. Lett. B 67, 309 (1977); B.W. Lee, S. Pakvasa, R.E. Schrock, and H. Sugawara, Phys. Rev. Lett. 38, 937 (1977).
- [4] R. Barbieri and L.J. Hall, Phys. Lett. B **338**, 212 (1994).
- [5] L. Calibbi and G. Signorelli, Riv. Nuovo Cim. 41, 71 (2018).
- [6] R.H. Bernstein and P.S. Cooper, Phys. Rept. 532, 27 (2013).
- [7] V.A. Kostelecký and S. Samuel, Phys. Rev. D 39, 683 (1989); V.A. Kostelecký and R. Potting,
 Nucl. Phys. B 359, 545 (1991); Phys. Rev. D 51, 3923 (1995).
- [8] V.A. Kostelecký and N. Russell, Data Tables for Lorentz and CPT Violation, Rev. Mod. Phys.

83, 11 (2011), arXiv:0801.0287v17 (2024).

- [9] D. Colladay and V.A. Kostelecký, Phys. Rev. D 55, 6760 (1997); Phys. Rev. D 58, 116002 (1998).
- [10] V.A. Kostelecký, Phys. Rev. D 69, 105009 (2004).
- [11] V.A. Kostelecký and Z. Li, Phys. Rev. D 99, 056016 (2019); Phys. Rev. D 103, 024059 (2021).
- [12] For reviews see, for example, Q.G. Bailey, in C. Bambi and A. Cardenas-Avendano, eds., *Recent Progress on Gravity Tests*, Springer, Singapore, 2024; Á. Roberts, Galaxies 9, 47 (2021); J.D. Tasson, Rep. Prog. Phys. 77, 062901 (2014); R. Bluhm, Lect. Notes Phys. 702, 191 (2006).
- [13] S.R. Coleman and S.L. Glashow, Phys. Rev. D 59, 116008 (1998).
- [14] E.O. Iltan, JHEP **06**, 016 (2003).
- [15] Y.M.P. Gomes and J.A. Helayel-Neto, Eur. Phys. J. C 80, 287 (2020).
- [16] A. Crivellin, F. Kirk and M. Schreck, JHEP 04, 082 (2021).
- [17] S. Hollenberg and P.B. Pal, Phys. Lett. B **701**, 89 (2011).
- [18] J.P. Noordmans, C.J.G. Onderwater, H.W. Wilschut, and R.G.E. Timmermans, Phys. Rev. D 93, 116001 (2016).
- [19] M.A. López-Osorio, E. Martínez-Pascual, and J.J. Toscano, J. Phys. G 43, 025003 (2016).
- [20] V.E. Mouchrek-Santos and M.M. Ferreira, Jr., Phys. Rev. D 95, 071701 (2017).
- [21] Y.M.P. Gomes, P.C. Malta, and M.J. Neves, Phys. Rev. D 101, 075001 (2020).
- [22] V.A. Kostelecký, E. Passemar and N. Sherrill, Phys. Rev. D 106, 076016 (2022).
- [23] For a review see, for example, A.K. Perrevoort, PoS FPCP2023, 015 (2023).
- [24] A.H. Gomes, V.A. Kostelecký, and A.J. Vargas, Phys. Rev. D 90, 076009 (2014); R. Bluhm,
 V.A. Kostelecký, and C.D. Lane, Phys. RevLett. 84, 1098 (2000).
- [25] S.M. Carroll, J.A. Harvey, V.A. Kostelecký, C.D. Lane and T. Okamoto, Phys. Rev. Lett. 87, 141601 (2001).
- [26] V.A. Kostelecký and J.D. Tasson, Phys. Rev. Lett. **102**, 010402 (2009).
- [27] W.H. Bertl *et al.*, Eur. Phys. J. C 47, 337 (2006); A. van der Schaaf, SciPost Phys. Proc. 5, 008 (2021).
- [28] L. Bartoszek et al., arXiv:1501.05241; K. Byrum et al., Universe 2023, 54 (2023).
- [29] R. Abramishvili *et al.*, PTEP **2020**, 033C01 (2020).
- [30] H. Natori, Nucl. Phys. B Proc. Suppl. 248, 52 (2014); M.J. Lee, Front. Phys. 6, 133 (2018);

K. Yamamoto, Phys. Sci. Forum 8, 39 (2023).

- [31] K. Byrum *et al.*, arXiv:2203.07569.
- [32] A. Kurup, PoS NUFACT2014, 101 (2014); B.E. Krikler, Ph.D. thesis, Imperial College, London, England (2017).
- [33] M. Aoki *et al.*, Letter of Intent P20-LoI, First PAC Meeting at J-PARC, June 2006; Y. Kuno, Nucl. Phys. B Proc. Suppl. **149**, 376 (2005).
- [34] T. Suzuki, D.F. Measday, and J.P. Roalsvig, Phys. Rev. C 35, 2212 (1987).
- [35] A. Czarnecki, W.J. Marciano, and K. Meinikov, hep-ph/9801218; H.C. Chiang, E. Oset, T.S. Kosmas, A. Faessler, and J.D. Vergados, Nucl. Phys. A 559, 526 (1993).
- [36] W.C. Haxton and E. Rule, Phys. Rev. Lett. **133**, 261801 (2024).
- [37] P. Möller, A.J. Sierk, T. Ichikawa, and H. Sagawa, Atom. Data and Nucl. Data Tables 109, 1 (2016); M. Haouas, F. Taulelle, and C. Martineau, Prog. Nucl. Mag. Res. Spectr. 94, 11 (2016); B. Bally, G. Giacalone, and M. Bender, Eur. Phys. J. A 59, 58 (2023).
- [38] H. de Vries, C.W. de Jager, and C. de Vries, At. Data Nucl. Data Tables 36, 495 (1987);
- [39] R. Kitano, M. Koike, and Y. Okada, Phys. Rev. D 66, 096002 (2002).
- [40] V.A. Kostelecký and R. Lehnert, Phys. Rev. D 63, 065008 (2001); V.A. Kostelecký, R. Lehnert, M. Schreck, and B. Seradjeh, arXiv:2412.18034.
- [41] V.A. Kostelecký and M. Mewes, Phys. Rev. D 85, 096005 (2012).
- [42] R. Bluhm, V.A. Kostelecký, C.D. Lane, and N. Russell, Phys. Rev. D 68, 125008 (2003);
 Phys. Rev. Lett. 88, 090801 (2002); V.A. Kostelecký and M. Mewes, Phys. Rev. D 66, 056005 (2002).
- [43] V.A. Kostelecký, A.C. Melissinos, and M. Mewes, Phys. Lett. B 761, 1 (2016); Y. Ding and
 V.A. Kostelecký, Phys. Rev. D 94, 056008 (2016).
- [44] V.A. Kostelecký, Phys. Rev. Lett. 80, 1818 (1998).
- [45] T.S. Kosmas, S. Kovalenko, and I. Schmidt, Phys. Lett. B 511, 203 (2001).