Self-dual pp-wave solutions in chiral higher-spin gravity

Tung Tran 1

¹Asia Pacific Center for Theoretical Physics, POSTECH, $\gamma\gamma$ Cheongamro, Nam-gu, Pohang-si, Gyeongsangbuk-do, 37673, Korea

E-mail: tran.tung@apctp.org

ABSTRACT: We show that chiral higher-spin gravity with a vanishing cosmological constant admits a class of exact self-dual pp-wave solutions derived from harmonic scalar functions and two principal spinors. These solutions satisfy both the linear and non-linear equations of motion, as they annihilate all higher-order vertices, leading to the equations of motion for free fields on a self-dual background sourced by a positive-helicity spin-2 field. Our method employs a simple light-cone ansatz for positive-helicity chiral higher-spin fields, along with a modified Kerr-Schild ansatz adapted for the self-dual gravity framework.

Contents

1 Introduction

By extending the usual GR with massless higher-spin fields, one can enlarge the symmetries of gravity by those associated with higher-spin fields, see e.g. [\[1\]](#page-25-0) for an overview. This led to various higher-spin gravities (HSGRA)s, which serve as simple toy models toward a theory of quantum gravity, one that is presumably UV-finite and free of black hole-like singularities.

Intriguingly, to construct theories of interacting massless higher-spin fields, one often needs to relax some of the usual assumptions underlying field theories such as unitarity or parity invariance, which, to some extent, is not a surprise fact (see discussion in $[2-8]$ $[2-8]$).

At the moment, QFT-compatible HSGRAs with propagating degrees of freedom are either higher-spin extension of (self-dual) Weyl gravity $[9-14]$ $[9-14]$, or chiral higher-spin theories, see e.g. $[15-22]$. There are also in 3d HSGRA $[23-31]$ $[23-31]$ and a higher-spin extension of Jackiw-Teitelboim gravity in 2d [\[32\]](#page-27-1). Another interesting example is the higher-spin gauge theory induced by the IKKT matrix model on an FLRW cosmological background with a truncated higher-spin spectrum [\[33,](#page-27-2) [34\]](#page-27-3).^{[1](#page-2-0)}

In this work, we focus on chiral higher-spin gravity and explore the space of its exact solutions to examine how much they differ from GR.[2](#page-2-1)

One of the key features of chiral HSGRA is that its gravitational background is selfdual [\[39\]](#page-27-4), i.e. background with Euclidean or split signature where half of the component of the Weyl tensor (the anti self-dual part) is zero. Of course, one can choose to work with a complexified setting, which provides more flexibility in doing computation. In fact, this philosophy, which has deep roots in twistor theory $[40]$, is what we want to employ in this paper in finding exact solutions of chiral HSGRA using its equations of motion [\[41,](#page-27-6) [42\]](#page-27-7).

The rest of the paper is organized as follows:

- In Section [2,](#page-3-0) we present crucial information, including the spectrum and explicit forms of local vertices, to study chiral higher-spin gravity (HSGRA) through its equations of motion derived from certain minimal model with local L_{∞} -structures of a certain Q-manifold [\[43,](#page-27-8) [44\]](#page-27-9).
- Section [3](#page-8-0) studies the exact solutions of chiral HSGRA sourced by a set of harmonic scalar functions $\{H_s\}$ and the principal spinors (λ, λ) à la Newman-Penrose. Since all local vertices of chiral HSGRA are smooth in the cosmological constant Λ , as shown in [\[42\]](#page-27-7), it seems reasonable to first explore the case $\Lambda = 0$. This is exactly what we do in this paper.

The spin-2 sector of the theory needs some special attention, since it generates the proposed self-dual pp-wave background away from a flat background. In particular, we will adapt the Cartan structure equations to the context of self-dual GR as discussed in [\[45\]](#page-27-10), which admits a smooth deformation from flat space to $(A)dS$. By considering an appropriate harmonic ansatz for fluctuation, akin to those in $[46, 47]$ $[46, 47]$, we show that the spin-2 sector of chiral HSGRA indeed admit a class of exact solutions, which solve the full non-linear EoMs, for any harmonic profile function H_2 . From here, the construction for exact higher-spin solutions is completely analogous. Note that our solution has a simple form, and it makes most of the structure maps of the underlying L_{∞} -algebra (for $\Lambda = 0$) vanish, see earlier discussion in [\[41\]](#page-27-6). Moreover, it is reasonable to speculate that once the space of solutions is found, it should be 'portable' to other nearby local higher-spin theories [\[48,](#page-27-13) [49\]](#page-28-0) as well.

¹See e.g. [\[35](#page-27-14)[–37\]](#page-27-15) and references therein for the recent development of this model.

²See [\[38\]](#page-27-16) for the lower-spin BPST solution in chiral HSGRA.

In the concluding part of Section [3,](#page-8-0) we also derive the effective action corresponding to our exact solutions and show that this action indeed reduces to the standard kinetic term for free spinning fields propagating on the proposed self-dual pp-wave background. That is, despite having non-trivial interactions, the theory appears to be free when the scattering data are those obtained exact solutions.

- We end with a discussion in Section [4.](#page-22-0) There are also appendices which the reader can refer to in due time.

Notation. We use lowercase Roman letters $a, b = 1, 2, 3, 4$ to denote tangent indices, while uppercase letters $A, B = 1, 2, 3, 4$ will be used for twistor or $\mathfrak{sp}(4)$ -indices. The Greek letters $\alpha, \beta = 0, 1$ and $\dot{\alpha}, \dot{\beta} = \dot{0}, \dot{1}$ are used to for spinorial indices. Symmetrized indices are denoted by the same letters, e.g. $A_a B_a = \frac{1}{2}$ $\frac{1}{2}(A_{a_1}B_{a_2}+A_{a_2}B_{a_1})$ while fully symmetric rank-s tensor will be denoted by $T_{a(s)} = T_{a_1...a_s}$.

2 Review

This section provides the mathematical background used to construct explicit representations of chiral HSGRA with vanishing cosmological constant. We shall be brief and refer the reader to $[41, 42]$ $[41, 42]$ for the details and to $[50, 51]$ $[50, 51]$ for a recent development.

2.1 Free differential algebra

Let M be a 4d spacetime manifold and $C^{\infty}(T[1]\mathcal{M}) \equiv \Omega^{\bullet}(\mathcal{M})$ be its associated graded commutative algebra of differential forms endowed with the exterior differential d, which squares to zero. Then, the pair $(T[1]\mathcal{M}, Q_{\mathcal{M}} \equiv d)$ present the simplest example of a Q manifold.

Now, consider a map $\Phi^A(x, dx) : T[1]\mathcal{M} \to \mathcal{N}$ from the source $T[1]\mathcal{M}$ to another target Q-manifold $(\mathcal{N}, Q_{\mathcal{N}})$ with coordinates Φ^A preserving the differential. Then, around a stationary point $p \in \mathcal{N}$, the associated set of pde. for Φ^A takes the form

$$
d\Phi^A = Q^A_N(\Phi). \tag{2.1}
$$

This is also known as free differential algebra (FDA) [\[43,](#page-27-8) [52–](#page-28-3)[54\]](#page-28-4). The nilpotency of $Q_N \equiv Q$ is equivalent to the statement:

$$
Q^B \frac{\partial}{\partial \Phi^B} Q^A = 0 \qquad \Leftrightarrow \qquad \sum \pm Q^B_{M_1 \dots M_n} Q^A_{N_1 \dots B \dots N_k} = 0. \tag{2.2}
$$

This gives us an L_{∞} -algebra [\[44,](#page-27-9) [55\]](#page-28-5) whose L_{∞} -structure maps are the Taylor's coefficients $Q_{M_1...N_n}^B$.

In the simplest setting, which works for generic gauge theories and gravity, we can choose $\Phi(x) = (\omega(x), \mathsf{C}(x))$ to be the graded coordinates of N. Here, ω is a gauge field of degree one taking values in some Lie algebra g, and C is a 'matter' field of degree zero taking values in some representation $\rho(\mathfrak{g})$ of \mathfrak{g} . Then, the equations of motion resulting from [\(2.1\)](#page-3-2) with the above field content, in general, are

$$
d\omega = \mathcal{V}(\omega, \omega) + \mathcal{V}(\omega, \omega, \mathsf{C}) + \mathcal{V}(\omega, \omega, \mathsf{C}, \mathsf{C}) + \dots,
$$
\n(2.3a)

$$
d\mathsf{C} = \mathcal{U}(\omega, \mathsf{C}) + \mathcal{U}(\omega, \mathsf{C}, \mathsf{C}) + \mathcal{U}(\omega, \mathsf{C}, \mathsf{C}, \mathsf{C}) + \dots
$$
\n(2.3b)

where the notations (V, U) are obvious. To sum things up, we can define a theory whenever we can specify $\{(\omega, C)\}\$ and $\{\mathcal{V}_n(\omega, \omega, C, \ldots), \mathcal{U}_n(\omega, C, \ldots)\}$.^{[3](#page-4-2)}

Note that [\(2.3\)](#page-4-3) can be referred to as a deformation of the following differential graded Lie algebra (dgLa)

$$
d\omega = \frac{1}{2}[\omega, \omega], \qquad d\mathsf{C} = \rho(\omega)\mathsf{C}.
$$
 (2.4)

The first equation is simply the definition of a flat connection associated to the one-form ω , while the second expresses the covariant constancy of the 0-form C wrt. the flat connection.

2.2 Chiral HSGRA

2.2.1 Chiral FDA data

The local isomorphism $SO(4,\mathbb{C}) \simeq SL(2,\mathbb{C}) \times SL(2,\mathbb{C})$ in 4d complexified spacetime allows us to write any tensorial fields in terms of spin-tensor fields $T_{\alpha(m)\dot{\alpha}(n)}$.

In the case of chiral HSGRA, (ω, C) can be written in terms of generating functions as

1-form :
$$
\omega = \sum_{s} \frac{1}{(2s)!} \omega_{A(2s)} Y^{A(2s)} = \sum_{m,n \ge 0} \frac{1}{m!n!} \omega_{\alpha(m)\dot{\alpha}(n)} y^{\alpha(m)} \tilde{y}^{\dot{\alpha}(n)}, \qquad m+n \in 2\mathbb{N},
$$
 (2.5a)

$$
0 \text{-form}: \ \mathsf{C} = \sum_{s} \frac{1}{(2s)!} \mathsf{C}_{A(2s)} Y^{A(2s)} = \sum_{m,n \ge 0} \frac{1}{m!n!} \mathsf{C}_{\alpha(m)\dot{\alpha}(n)} y^{\alpha(m)} \tilde{y}^{\dot{\alpha}(n)}, \qquad m+n \in 2\mathbb{N}. \tag{2.5b}
$$

where $Y^A = (y^\alpha, \tilde{y}^{\dot{\alpha}})$. Here, $y^\alpha, \tilde{y}^{\dot{\alpha}}$ are commutative spinorial variables, and $y^{\alpha(n)} = y^{\alpha_1} \dots y^{\alpha_n}$ etc. is our standard abbreviation. The dynamical bosonic fields, which give us free action, are singled out as

helicity +s:
$$
\mathcal{A} := \sum_{s \in \mathbb{N}} \frac{1}{(2s)!} \mathcal{A}_{\alpha(2s)} y^{\alpha(2s)} = \omega(y, \tilde{y} = 0), \qquad (2.6a)
$$

helicity
$$
-s
$$
:
$$
\mathcal{B} := \sum_{s \in \mathbb{N}} \frac{1}{(2s)!} B_{\alpha(2s)} y^{\alpha(2s)} = \mathsf{C}(y, \tilde{y} = 0), \qquad (2.6b)
$$

helicity
$$
0:
$$
 $\varphi := C(y = 0, \tilde{y} = 0)$. (2.6c)

³The subscripts of V and U denote the number of arguments that enter the corresponding vertices.

Using $B^{\alpha(2s)}$ and $\mathcal{A}_{\alpha(2s-2)}$ one can write down the free action for chiral spinning fields as [\[39\]](#page-27-4)

$$
S = \int B^{\alpha(2s)} \wedge H_{\alpha\alpha} \nabla \mathcal{A}_{\alpha(2s-2)} , \qquad H_{\alpha\alpha} := h_{\alpha\dot{\gamma}} \wedge h_{\alpha}^{\dot{\gamma}} . \tag{2.7}
$$

In the above, $h_{\alpha\dot{\alpha}}$ is the background vierbein of any SD spacetime where the ASD component $C_{\alpha(4)}$ of the Weyl tensor vanish, and ∇ is the associated self-dual connection. This action is invariant under $\delta A_{\alpha(2s-2)} = \nabla \xi_{\alpha(2s-2)} + h_{\alpha}{}^{\dot{\beta}} \eta_{\alpha(2s-3)\dot{\beta}}$, which can be checked immediately by noting that $\nabla^2 \xi_{\alpha(2s-2)} = H_\alpha{}^\beta \xi_{\alpha(2s-3)\beta}$ and $H_{\alpha\alpha} \wedge h_\alpha{}^\beta = 0$.

As explained in [\[39\]](#page-27-4), the physical one-form $\mathcal{A}_{\alpha(2s-2)} \equiv \omega_{\alpha(2s-2)}$ can be written as

$$
\omega_{\alpha(2s-2)} = h^{\beta\dot{\beta}} A_{\alpha(2s-2)\beta\dot{\beta}} + h_{\alpha}{}^{\dot{\alpha}} \vartheta_{\alpha(2s-3)\dot{\alpha}}.
$$
\n(2.8)

Since we can use the gauge transformation $\delta_{\eta}\omega_{\alpha(2s-2)} = h_{\alpha}{}^{\dot{\alpha}}\eta_{\alpha(2s-3)\dot{\alpha}}$ to gauge away the second component ϑ in $\omega^{\alpha(2s-2)}$, we will simply consider for most of the time

$$
\omega_{\alpha(2s-2)} = h^{\beta\dot{\beta}} A_{\alpha(2s-2)\beta\dot{\beta}}.\tag{2.9}
$$

Now, at the free level the FDA for chiral HSGRA is simple

$$
\nabla \omega_{\alpha(2s-2-k)\,\dot{\alpha}(k)} = h_{\alpha}{}^{\dot{\beta}} \omega_{\alpha(2s-3-k)\,\dot{\alpha}(k)\dot{\beta}}, \qquad 0 \le k \le 2s-2, \qquad (2.10a)
$$

$$
\nabla \omega_{\dot{\alpha}(2s-2)} = \tilde{H}^{\dot{\beta}\dot{\beta}} C_{\dot{\alpha}(2s-2)\dot{\beta}\dot{\beta}}, \qquad \qquad \tilde{H}^{\dot{\alpha}\dot{\alpha}} := h_{\gamma}^{\dot{\alpha}} \wedge h^{\gamma\dot{\alpha}}, \qquad (2.10b)
$$

$$
\nabla C_{\alpha(k)\dot{\alpha}(2s+k)} = h^{\beta\dot{\beta}} C_{\alpha(k)\beta\dot{\alpha}(2s+k)\dot{\beta}},
$$
\n(2.10c)

$$
\nabla B^{\alpha(2s+k)\dot{\alpha}(k)} = h_{\beta\dot{\beta}} B^{\alpha(2s+k)\beta\dot{\alpha}(k)\dot{\beta}},
$$
\n(2.10d)

where $C_{\alpha(k)\dot{\alpha}(2s+k)}$ are the ∇ -generated auxiliary fields of $C_{\dot{\alpha}(2s)}$, and $B^{\alpha(2s+k)\dot{\alpha}(k)}$ are the ∇ -generated auxiliary fields of $B^{\alpha(2s)}$, cf. [\(2.6\)](#page-4-4).

Since the next (auxiliary) fields can be generated by the previous (auxiliary) fields by acting ∇ on the physical fields again and again, the whole spectrum of fields can be obtained starting from the *chiral fields* $A_{\alpha(2s-2)}$ and $B^{\alpha(2s)}$ and the scalar field φ (see Fig. [1\)](#page-6-0).

2.2.2 Vertices

The next ingredients we need are the structure maps/vertices (V, U) in [\(2.3\)](#page-4-3). Conveniently, we can write all vertices of chiral HSGRA as poly-differential operators:

$$
\mathsf{F}(\Phi_1,\ldots,\Phi_n) := \mathsf{F}(y,\tilde{y},\partial_{i\alpha},\tilde{\partial}_{i\dot{\alpha}})\Phi_1(y_1,\tilde{y}_1)\ldots\Phi_n(y_n,\tilde{y}_n)\Big|_{y_i,\tilde{y}_i=0}, \quad \Phi = (\omega,\mathsf{C})\,. \tag{2.11}
$$

For conciseness, we will suppress the notation $|_{y_i=0, \tilde{y}_i=0}$ at the end of each expression. Here, $C = (C, B)$ is the master 0-form field, which contains both negative-helicity B's and positivehelicity C's fields and the scalar field φ .

Below, we present some of the key features of these structure maps. The technical details in deriving these maps can be found in [\[41,](#page-27-6) [42\]](#page-27-7) (see also Appendix [B](#page-23-1) for a recap).

Figure 1. The horizontal/vertices axes represent the number of dotted and un-dotted spinorial indices that a tensorial field has. Here, [red] are fields with positive helicity, and [blue] are fields with negative helicity. The arrows indicate the directions in which auxiliary fields will be generated by acting ∇ on the previous ones. All fields generated this way are referred to as chiral FDA data.

 \star -product The first structure map in the minimal model of interest is the binary structure map F_2 . It is given by a graded associative star-product:

$$
\Phi_1 \star \Phi_2 = \exp \left(Y^A \frac{\partial}{\partial Y_1^A} + Y^A \frac{\partial}{\partial Y_2^A} - I^{AB} \frac{\partial}{\partial Y_1^A} \frac{\partial}{\partial Y_2^B} \right) \Phi_1(Y_1) \Phi_2(Y_2)
$$

= $\exp \left(\langle y \partial_1 \rangle + \langle y \partial_2 \rangle + \Lambda \langle \partial_1 \partial_2 \rangle + [y \partial_1] + [y \partial_2] + [\partial_1 \partial_2] \right) \Phi_1 \Phi_2,$ (2.12)

where Λ is the cosmological constant, I^{AB} is the infinity twistor [\[40\]](#page-27-5) obeying

$$
I^{AB} = \begin{pmatrix} \Lambda \epsilon^{\alpha \beta} & 0 \\ 0 & \epsilon^{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \qquad I_{AB} = \begin{pmatrix} \epsilon_{\alpha\beta} & 0 \\ 0 & \Lambda \epsilon_{\dot{\alpha}\dot{\beta}} \end{pmatrix}, \qquad I_{AC}I^{BC} = \Lambda \delta_A{}^B. \tag{2.13}
$$

Note that

$$
\langle y \,\partial_i \rangle = y^\alpha \partial_{i\alpha} \,, \quad \langle \partial_i \,\partial_j \rangle = \partial_i^\alpha \partial_{j\alpha} \,;\qquad [y \,\partial_i] = \tilde{y}^{\dot{\alpha}} \tilde{\partial}_{i\dot{\alpha}} \,, \quad [\partial_i \,\partial_j] = \tilde{\partial}_i^{\dot{\alpha}} \tilde{\partial}_{j\dot{\alpha}} \,, \tag{2.14}
$$

where $\partial_{i\alpha} = \frac{\partial}{\partial y_i^{\alpha}}, \ \tilde{\partial}_{i\dot{\alpha}} = \frac{\partial}{\partial \tilde{y}}$ $\frac{\partial}{\partial \tilde{y}_i^{\alpha}}$. Here, $e^{\langle y|\partial i\rangle}$ may be referred to as a shift operator with the property $e^{\langle y \partial_i \rangle} \Phi(y_i) = \Phi(y_i + y)$. We will use the following convention to raise and lower spinorial indices

$$
v^{\alpha} = v_{\beta} \epsilon^{\alpha \beta}, \quad v_{\alpha} = v^{\beta} \epsilon_{\beta \alpha}, \quad v^{\dot{\alpha}} = v_{\dot{\beta}} \epsilon^{\dot{\alpha} \dot{\beta}}, \quad v_{\dot{\alpha}} = v^{\dot{\beta}} \epsilon_{\dot{\beta} \dot{\alpha}}.
$$
 (2.15)

The Weyl algebra A_2 of polynomial functions in $Y^A = (y^{\alpha}, \tilde{y}^{\dot{\alpha}})$ defined in terms of the \star -product cf. [\(2.12\)](#page-6-1) admits the decomposition $A_2 = A_1(y) \otimes A_1(\tilde{y})$ where the subscript denotes the number of canonical pairs. We then define

$$
\mathfrak{hs}_{\Lambda} := \mathsf{A}_1 \otimes \big(\mathsf{A}_1 \otimes \text{Mat}(n, \mathbb{K}) \big) , \tag{2.16}
$$

to be the underlying higher-spin algebra of the system. Note that when $\Lambda = 0$, the Poisson structure defined by the infinity twistor is degenerate, and A_1 becomes a commutative algebra while $A_1 \otimes \text{Mat}(n, \mathbb{K})$ remains to be an non-commutative (NC) algebra. This is the setting that we consider in this work.

Higher-order vertices. The details of deriving higher-order vertices for chiral FDA, cf. (2.3) , can be found in e.g. $[51]$. Here, we shall emphasize some of the main points, which are relevant for our analysis in Section [3.](#page-8-0) Henceforth, we focus on the case where $\Lambda = 0$.

 $\diamond \mathcal{V}_{n\geq3}(\omega,\omega,\mathsf{C},\ldots,\mathsf{C})$ vertices. The $\mathcal{V}_n(\omega,\omega,\mathsf{C},\ldots)$ maps consist of $\binom{n}{2}$ $\binom{n}{2} = n(n-1)/2$ sub-vertices:

$$
\mathcal{V}_n(\omega, \omega, \mathsf{C}, \dots, \mathsf{C}) = V_{\omega, \omega, \mathsf{C}, \dots, \mathsf{C}}^{(n)} + V_{\omega, \mathsf{C}, \omega, \mathsf{C}, \dots, \mathsf{C}}^{(n)} + \dots
$$
(2.17)

We shall denote i and j to be the positions of ω 's in the sub-vertices V's (assuming $i < j$) since they can be useful references for describing the sub-vertices V 's. It is worth noting that \mathcal{V}_n are L_{∞} structure maps obtained by suitable graded symmetrization of a certain A_{∞} structure maps/sub-vertices $V^{(n)}$ (see e.g. Appendix [B\)](#page-23-1).

For instance, a vertex of the form $V_{\mathcal{C}}^{(n)}$ $C_{\text{c},\dots,\omega_i,\text{c},\dots,\omega_j,\dots,\text{c}}^{(n)}$ has the following expressions [\[50\]](#page-28-1):

$$
V_{\mathsf{C},\ldots,\omega_{i},\mathsf{C},\ldots,\omega_{j},\ldots,\mathsf{C}}^{(n)} = (-)^{i+j-1} \sigma_{G_{ij}} \, \mathbb{U}_{n} \langle \partial_{i} \, \partial_{j} \rangle^{n-2} \int_{\Delta_{n-2}} \exp \operatorname{tr} \left(\mathsf{P}_{G_{ij}} \mathsf{Q}^{T} \right), \tag{2.18}
$$

where

$$
Q = \begin{pmatrix} u_1 & u_2 & \dots & u_{n-2} & 1 & -\sum u_i \\ v_1 & v_2 & \dots & v_{n-2} & 1 & -\sum v_i \end{pmatrix},
$$
\n(2.19a)

$$
\mathsf{P}_{G_{ij}} = \begin{pmatrix} \langle \partial_i \, \partial_1 \rangle & \dots & -\langle \partial_i \, \partial_j \rangle & \dots & 0 & \dots & \langle \partial_i \, \partial_{n-2} \rangle & -\langle y \, \partial_i \rangle \\ \langle \partial_j \, \partial_1 \rangle & \dots & 0 & \dots & -\langle \partial_j \, \partial_i \rangle & \dots & \langle \partial_j \, \partial_{n-2} \rangle & -\langle y \, \partial_j \rangle \end{pmatrix} . \tag{2.19b}
$$

The integration domain $\Delta_n \in \mathbb{R}^{2n}$ in [\(2.18\)](#page-7-0) is a compact convex *n*-gon and the integrating variables u_k and v_k obey

$$
\frac{u_1}{v_1} \le \frac{u_2}{v_2} \le \dots \le \frac{u_n}{v_n} \le \frac{1-U}{1-V}, \qquad U = \sum u_i, \quad V = \sum v_i.
$$

Furthermore, G_{ij} denotes the corresponding tree/graph in homological perturbation theory from which V's come from. Additionally, $\sigma_{G_{ij}}$ represents the permutation that arranges fields into 'natural' order of the associated graph. Lastly,

$$
\mathcal{U}_n := \exp\left(\sum_{i=1}^n [y\,\partial_i] + \sum_{i < j \le n} [\partial_i\,\partial_j]\right),\tag{2.20}
$$

is a poly-differential operator acting solely on the \tilde{y} -sector by virtue of the \star -product [\(2.12\)](#page-6-1).

Canonically, we can take $G_{12} = (\omega_1, \omega_2, \mathsf{C}_3, \ldots, \mathsf{C}_{n-2})$ to be the reference graph where all C's are on the right of ω 's. For concreteness, the explicit form of the sub-vertex $V_{\omega,\omega}^{(n)}$ $\omega, \omega, C, \ldots, C$ associated with G_{12} has the following form

$$
V_{\omega_1,\omega_2,\mathbf{C},\dots,\mathbf{C}}^{(n)}(y,\tilde{y};\partial_i,\tilde{\partial}_i) = \mathbb{U}_n \langle \partial_1 \partial_2 \rangle^{n-2} \times \times \int_{\Delta_{n-2}} \exp \left[\left(1 - \sum_{k=1}^{n-2} u_k \right) \langle y \partial_1 \rangle + \left(1 - \sum_{k=1}^{n-2} v_k \right) \langle y \partial_2 \rangle + \sum_{m \neq 1,2} u_m \langle \partial_1 \partial_m \rangle + \sum_{m \neq 1,2} v_m \langle \partial_2 \partial_m \rangle \right].
$$
\n(2.21)

Then, any other graph can be obtained from G_{12} by moving some number of C's in between $ω$'s and cyclicing all arguments to a specific position e.g. $V_{\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\omega,\mathsf{C$ This allows us to constructs the remaining vertices/graphs from the reference ones.

 \Diamond $\mathcal{U}(\omega, \mathsf{C}, \ldots, \mathsf{C})$ vertices. It turns out that $(\mathcal{V}, \mathcal{U})$ vertices can be related to each others through a \mathbb{C} -linear pairing inherited from a degree 2 cyclic A_{∞} -algebra:

$$
\langle -|-\rangle : \mathcal{N} \otimes \mathcal{N}^{\vee}[-1] \to \mathbb{C}[-2], \text{ where } \langle a|b\rangle = (-)^{(|a|+1)(|b|+1)} \langle b|a\rangle.
$$

The natural pairings in the y-sector used to construct all vertices $[41, 42, 51]$ $[41, 42, 51]$ $[41, 42, 51]$ $[41, 42, 51]$ are as follows:

$$
\langle \omega | \mathsf{C} \rangle = -\langle \mathsf{C} | \omega \rangle = \exp[\langle \partial_1 \partial_2 \rangle] \omega(y_1) \mathsf{C}(y_2)
$$

$$
\langle V(\omega, \omega, \mathsf{C}, \dots, \mathsf{C}) | C \rangle = +\langle \omega | U(\omega, \mathsf{C}, \dots, \mathsf{C}) \rangle
$$

$$
V(\partial_1, \dots, \partial_n | y) \mapsto + U(y, \partial_1, \dots, \partial_{n-1} | - \partial_n)
$$

$$
\langle \mathsf{C} | V(\mathsf{C}, \dots, \omega, \mathsf{C}, \dots, \mathsf{C}, \omega) \rangle = -\langle U(\mathsf{C}, \mathsf{C}, \dots, \omega, \mathsf{C}, \dots, \mathsf{C}) | \omega \rangle
$$

$$
V(y | \partial_1, \dots, \partial_n) \mapsto -U(-\partial_1 | \partial_2, \dots, \partial_n, y)
$$

Table 1. Some natural pairings in chiral-FDA.

For instance, to obtain $U_{\mathsf{C},\dots,\mathsf{C},\omega,\dots,\mathsf{C}} \equiv U(\mathsf{C},\dots,\mathsf{C},\omega,\dots,\mathsf{C})$ from $V_{\omega,\dots,\mathsf{C},\omega,\dots,\mathsf{C}}$, we replace $y \mapsto -\partial_n, \partial_1 \mapsto y$ and $\partial_i \mapsto \partial_{i-1}$ (for $i \neq n$), while to obtain $U_{\mathsf{C},...,\omega,\dots,\mathsf{C}}$ from $V_{\mathsf{C},...,\omega,\mathsf{C},...,\mathsf{C},\omega} \equiv$ $V(\mathsf{C},\ldots,\omega,\mathsf{C},\ldots,\mathsf{C},\omega)$, we replace $y \mapsto -\partial_1$, $\partial_i \mapsto \partial_{i+1}$ (for $i \leq n$) and $\partial_n \mapsto y$ with an overall minus sign. Since there are $\binom{n}{2}$ $\binom{n}{2} = \frac{n(n-2)}{2}$ $\frac{n-2}{2}$ number of V's sub-vertices compared to n number of U 's sub-vertices at a given order n , this replacement rule always works. Therefore, as a matter of convenience, one only needs to construct all A_{∞} structure maps $V^{(n)}$, and everything will follow.

Note that the pairing $\langle -|-\rangle$ not only facilitates the cyclic reordering of fields and vertices, but is also defined to eliminate all auxiliary γ and $\tilde{\gamma}$ variables. This ensures that all possible contractions between the stand-alone field ω or C with the multilinear maps or sub-vertices V 's and U 's are fully captured.

3 Light-cone ansatz and exact solutions

Armed with the above chiral FDA data, we now propose a family of explicit representations of (ω, C) and show that they form exact harmonic solutions in chiral HSGRA.

Recall that since the positive-helicity fields C's can be generated from ω via linear chiral FDA, cf. [\(2.10\)](#page-5-1), we can construct all positive-helicity data starting from the physical fields $\mathcal{A}_{\alpha(2s-2)}$. Furthermore, we show below that $\mathcal{A}_{\alpha(2)}$ is the field that generate a selfdual background. It is also worth stressing before running our analysis that $B^{\alpha(2s)}$ and their ∇ -generated thereof are *not* derived objects from the positive helicity fields $\mathcal{A}_{\alpha(2s-2)}$. Rather, they should be regarded simply as linearized fields that propagate on the self-dual background generated by $\mathcal{A}_{\alpha(2)} \equiv \omega_{\alpha(2)}$.

3.1 Cartan's structure equations for empty self-dual background

One of the nice features of \mathfrak{hs}_Λ is that its $\mathfrak{so}(2,3)$ subalgebra can be defined via the generators:

$$
L_{\alpha\alpha} = -\frac{1}{2} y_{\alpha} y_{\alpha} , \qquad P_{\alpha\dot{\alpha}} = -\frac{1}{2} y_{\alpha} \tilde{y}_{\dot{\alpha}} , \qquad \tilde{L}_{\dot{\alpha}\dot{\alpha}} = -\frac{1}{2} \tilde{y}_{\dot{\alpha}} \tilde{y}_{\dot{\alpha}} . \tag{3.1}
$$

As usual, $P_{\alpha\dot{\alpha}}$ is translation generator, and $(L_{\alpha\alpha}, L_{\dot{\alpha}\dot{\alpha}})$ are Lorentz's. The algebra which describes AdS_4 is [\[45\]](#page-27-10)

$$
[L_{\alpha\beta}, L_{\gamma\delta}]_{\star} = \Lambda \big(\epsilon_{\beta\gamma} L_{\alpha\gamma} + \epsilon_{\alpha\gamma} L_{\beta\delta} + \epsilon_{\beta\delta} L_{\alpha\gamma} + \epsilon_{\alpha\gamma} L_{\beta\delta}\big),\tag{3.2a}
$$

$$
[\widetilde{L}_{\dot{\alpha}\dot{\beta}}, \widetilde{L}_{\dot{\gamma}\dot{\delta}}]_{\star} = (\epsilon_{\dot{\beta}\dot{\gamma}} L_{\dot{\alpha}\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\gamma}} L_{\dot{\beta}\dot{\delta}} + \epsilon_{\dot{\beta}\dot{\delta}} L_{\dot{\alpha}\dot{\gamma}} + \epsilon_{\dot{\alpha}\dot{\gamma}} L_{\dot{\beta}\dot{\delta}}),
$$
\n(3.2b)

$$
[L_{\alpha\beta}, P_{\gamma\dot{\gamma}}]_{\star} = \Lambda \Big(\epsilon_{\beta\gamma} P_{\alpha\dot{\gamma}} + \epsilon_{\alpha\gamma} P_{\beta\dot{\gamma}}\Big), \qquad (3.2c)
$$

$$
[\widetilde{L}_{\dot{\alpha}\dot{\beta}}, P_{\dot{\gamma}\dot{\gamma}}]_{\star} = (\epsilon_{\dot{\beta}\dot{\gamma}} P_{\gamma\dot{\alpha}} + \epsilon_{\dot{\alpha}\dot{\gamma}} P_{\gamma\dot{\beta}}),
$$
\n(3.2d)

$$
[P_{\alpha\dot{\alpha}}, P_{\beta\dot{\beta}}]_{\star} = (\epsilon_{\dot{\alpha}\dot{\beta}}L_{\alpha\beta} + \Lambda \epsilon_{\alpha\beta}\dot{L}_{\dot{\alpha}\dot{\beta}}). \tag{3.2e}
$$

It is easy to notice that $L_{\alpha\beta}$ becomes central of the algebra [\(3.2\)](#page-9-1) in the flat limit $\Lambda \to 0$. That is

$$
[P_{\alpha\dot{\alpha}}, P_{\beta\dot{\beta}}]_{\star}|_{\Lambda=0} = \epsilon_{\dot{\alpha}\dot{\beta}} L_{\alpha\beta} , \qquad (3.3)
$$

instead of being trivial. For this reason, the algebra generated by [\(3.1\)](#page-9-2) in the flat limit is referred to as a *deformed* chiral Poincaré algebra $[56]$.

The Cartan structure equations associated to empty flat spacetime read:

$$
d\Omega - \frac{1}{2}[\Omega, \Omega] = 0, \qquad \Omega = \frac{1}{2}\varpi_{\alpha\beta}L^{\alpha\beta} + h_{\alpha\dot{\alpha}}P^{\alpha\dot{\alpha}} + \frac{1}{2}\widetilde{\varpi}_{\dot{\alpha}\dot{\beta}}\widetilde{L}^{\dot{\alpha}\dot{\beta}}.
$$
 (3.4)

where $h_{\alpha\dot{\alpha}}$ is the background vierbein and $\varpi_{\alpha\beta}$, $\widetilde{\varpi}_{\dot{\alpha}\dot{\beta}}$ are the (A)SD components of the spin connection. In terms of components (2.4) unfolds into connection. In terms of components, [\(3.4\)](#page-9-3) unfolds into

$$
R_{\alpha\beta} := d\varpi_{\alpha\alpha} + H_{\alpha\alpha} = 0, \qquad (3.5a)
$$

$$
T_{\alpha\dot{\alpha}} := dh_{\alpha\dot{\alpha}} + \tilde{\varpi}_{\dot{\alpha}\dot{\gamma}} \wedge h_{\alpha}^{\ \dot{\gamma}} \equiv \nabla h_{\alpha\dot{\alpha}} = 0, \qquad (3.5b)
$$

$$
R_{\dot{\alpha}\dot{\beta}} := d\tilde{\varpi}_{\dot{\alpha}\dot{\alpha}} + \tilde{\varpi}_{\dot{\alpha}\dot{\gamma}} \wedge \tilde{\varpi}_{\dot{\alpha}}^{\dot{\gamma}} = 0.
$$
 (3.5c)

The reader may notice the difference with the Cartan structure equations associated with the conventional Poincaré algebra where $[P, P] = 0$. In particular, for SD background, the ASD component of the spin connection does *not* participate in the torsion constraint $(3.5b)$. Furthermore, the Riemann two-form [\(3.5a\)](#page-9-5) reveals an intriguing feature for SD background. That is, the spin connection $\varpi_{\alpha\alpha}$ is the main player of this game since the vierbein and the SD component of the spin connection can be solved for once we know $\varpi_{\alpha\alpha}$.

3.2 Deformed self-dual pp-wave background

This subsection proposes a class of SD pp-wave backgrounds in which chiral HSGRA is welldefined. This class of backgrounds is generated by a fluctuation of the ASD component of the spin connection, which is characterized by a function $H_2(v, z, \tilde{z})$ that is harmonic in the complex 2-plane (z, \tilde{z}) , and two principal spinors $(\lambda, \tilde{\lambda})$.^{[4](#page-10-1)}

Deformed Cartan's structure equation. Suppose we want to construct a non-trivial vacuum solution which does not describe an empty flat space. Then, the new vacuum solution whose geometrical information is encoded by the traceless part of the Riemann tensor, i.e. the Weyl tensor, can be regarded as a deformation away from empty flat space. In this case, we shall modify the Cartan's structure equations [\(3.5\)](#page-9-6) to

$$
\mathbf{R}_{\alpha\alpha} : = d\boldsymbol{\varpi}_{\alpha\alpha} + \boldsymbol{H}_{\alpha\alpha} = 0, \qquad (3.7a)
$$

$$
De_{\alpha\dot{\alpha}} = de_{\alpha\dot{\alpha}} + \widetilde{\varpi}_{\dot{\alpha}\dot{\beta}} \wedge e_{\alpha}{}^{\dot{\beta}} = 0, \qquad (3.7b)
$$

$$
\widetilde{\boldsymbol{R}}_{\dot{\alpha}\dot{\alpha}} := d\widetilde{\boldsymbol{\varpi}}_{\dot{\alpha}\dot{\beta}} + \widetilde{\boldsymbol{\varpi}}_{\dot{\alpha}\dot{\gamma}} \wedge \widetilde{\boldsymbol{\varpi}}_{\dot{\beta}}^{\dot{\gamma}} = \widetilde{\boldsymbol{H}}^{\dot{\beta}\dot{\beta}} C_{\dot{\beta}\dot{\beta}\dot{\alpha}\dot{\alpha}},
$$
\n(3.7c)

where

$$
\boldsymbol{\varpi}_{\alpha\beta} = \varpi_{\alpha\alpha} + \omega_{\alpha\alpha}, \quad \boldsymbol{e}_{\alpha\dot{\alpha}} = h_{\alpha\dot{\alpha}} + \omega_{\alpha\dot{\alpha}}, \quad \boldsymbol{\widetilde{\varpi}}_{\dot{\alpha}\dot{\beta}} = \boldsymbol{\widetilde{\varpi}}_{\dot{\alpha}\dot{\beta}} + \omega_{\dot{\alpha}\dot{\beta}}, \tag{3.8}
$$

are the deformed vierbein and spin connections with $\omega_{\alpha\alpha}, \omega_{\alpha\dot{\alpha}}, \omega_{\dot{\alpha}\dot{\alpha}}$ being the associated fluctuations.^{[5](#page-10-2)} Here, D stands for the connection of the deformed SD background, and (3.7) contains only the SD component $C_{\dot{\alpha}(4)}$ of the Weyl tensor due to SD condition, cf. [\[39\]](#page-27-4).

Tetrad and dyad bases. In 4d, we can associate to each point $x \in \mathcal{M}$ a tetrad of complex null vectors $\zeta^a = (\ell^a, n^a, p^a, \tilde{p}^a)$ (see e.g. [\[57\]](#page-28-7)) where

$$
\ell_a n^a = -1 = p_a \tilde{p}^a , \qquad \qquad \ell_a p^a = \ell_a \tilde{p}^a = n_a p^a = n_a \tilde{p}^a = 0 . \qquad (3.9)
$$

$$
S_{\alpha_1...\alpha_s} = \tau_{(\alpha_1}...\zeta_{\alpha_s)},\tag{3.6}
$$

where each spinor $\{\tau, \ldots, \zeta\}$ is referred to as *principal spinor* of $S_{\alpha(s)}$. Note that these principal spinors can be used to classify the Weyl tensors in classical GR, see e.g. [\[57,](#page-28-7) [58\]](#page-28-8).

⁵All the *bold* symbols are geometrical datum of the deformed background.

⁴As is well-known, any symmetric tensor of the form $S_{\alpha(s)} \equiv S_{\alpha_1...\alpha_s}$ can be factorized into a symmetrized product of one-index spinors [\[57\]](#page-28-7), i.e.

Then, a vector V^a may be written in the tetrad basis as

$$
V^a = u\ell^a + v n^a + z p^a + \tilde{z}\tilde{p}^a. \tag{3.10}
$$

Here, we may refer to (u, v, z, \tilde{z}) as spacetime coordinates.

Since the local Lorentz group of a 4d complexified \mathcal{M} is $SL(2,\mathbb{C}) \times SL(2,\mathbb{C})$, we can also associate to each point $x \in \mathcal{M}$ the dyad bases $(\lambda^{\alpha}, \tilde{\lambda}^{\dot{\alpha}})$ and $(\mu^{\alpha}, \tilde{\mu}^{\dot{\alpha}})$, cf. [\[57\]](#page-28-7), where $(\mu, \tilde{\mu})$ is another pair of principal spinors such that

$$
\langle \mu \lambda \rangle = 1 = [\tilde{\mu} \tilde{\lambda}]. \tag{3.11}
$$

Using the same normalization in [\(3.9\)](#page-10-4), we can fix unambiguously

$$
\ell_a = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}}, \qquad n_a = \mu_\alpha \tilde{\mu}_{\dot{\alpha}}, \qquad p_a = \lambda_\alpha \tilde{\mu}_{\dot{\alpha}}, \qquad \tilde{p}_a = \mu_\alpha \tilde{\lambda}_{\dot{\alpha}}.
$$
 (3.12)

In constructing the proposed harmonic solutions for chiral HSGRA, we will use $(\lambda, \tilde{\lambda})$ to characterize positive-helicity fields and $(\mu, \tilde{\mu})$ to describe the negative-helicity fields as they are related in the sense of [\(2.7\)](#page-5-2).

Note that we will define the spinors $(\lambda, \tilde{\lambda})$ as well as $(\mu, \tilde{\mu})$ to be covariantly constant wrt. the connection associated to empty flat background, i.e.

$$
\nabla \lambda_{\alpha} = 0, \qquad \nabla \tilde{\lambda}_{\dot{\alpha}} = 0, \qquad \nabla \mu_{\alpha} = 0, \qquad \nabla \tilde{\mu}_{\dot{\alpha}} = 0. \tag{3.13}
$$

Light-cone ansatz. Typically, in standard GR, one can simplify the process of finding exact solutions for Einstein equations by considering the Kerr-Schild ansatz [\[59\]](#page-28-9)

$$
\mathbf{g}_{ab} = g_{ab} + \mathrm{H}\ell_a \ell_b , \qquad \qquad \mathbf{g}^{ab} = g^{ab} - \mathrm{H}\ell^a \ell^b \tag{3.14}
$$

where g_{ab} is the metric of the background and H is a scalar profile function whose role is to generate a deformed spacetime (M, g) . However, in SDGR, the situation is slightly different. In particular, as the ASD component of the spin connection is the main object where all other geometrical data are derived from, cf. [\[60\]](#page-28-10), one must consider a modified version of the Kerr-Schild ansatz, adapted to accommodate the ASD component of the spin connection. For our purpose of constructing exact solutions in SD setting, we can consider

$$
\boldsymbol{\varpi}_{\alpha\alpha} = \boldsymbol{\varpi}_{\alpha\alpha} + \mathbf{H}_2 \langle \lambda | h | \lambda \rangle \lambda_\alpha \lambda_\alpha , \qquad \langle \lambda | h | \lambda \rangle := \lambda^\gamma h_{\gamma\gamma} \tilde{\lambda}^\gamma . \tag{3.15}
$$

Note that the above ansatz is inspired from the recent work [\[46,](#page-27-11) [47\]](#page-27-12). For the sake of generality, we have introduced a subscript "2" to the profile function H to indicate that it corresponds to the ansatz for the spin-2 sector. For spin-s sector, we may denote the corresponding profile function as H_s (see below).

Harmonicity of H_2 **.** There is a few properties that we want to impose on the profile function H_2 . Namely, H_2 must be *u*-independent and harmonic in (z, \tilde{z}) -plane spanned by (p^a, \tilde{p}^a) [\[61\]](#page-28-11), i.e.

$$
\ell^a \nabla_a \mathbf{H}_2 = 0, \qquad p^a \tilde{p}^b \nabla_a \nabla_b \mathbf{H}_2 = 0.
$$
 (3.16)

This allows us to write $H_2(v, z, \tilde{z})$ as the sum of two functions

$$
H_2(v, z, \tilde{z}) = H_2(v, z) + \tilde{H}_2(v, \tilde{z}), \qquad (3.17)
$$

so that H_2 effectively behaves as a 2d off-shell scalar field. Nevertheless, as shown below, the self-duality condition will set one of them, specifically $H_2(v, \tilde{z})$, to zero.

The harmonicity of H_2 can be recast into the following pde:

$$
\nabla \mathbf{H}_2 = h^{\alpha \dot{\alpha}} (\lambda_\alpha \tilde{\mathbf{H}}_{2\dot{\alpha}} + \tilde{\lambda}_{\dot{\alpha}} \mathbf{H}_{2\alpha}), \qquad (3.18)
$$

where $H_{2\alpha}$ and $\tilde{H}_{2\dot{\alpha}}$ are first-order derivatives of H_2 in the y- and \tilde{y} -sector, respectively. Furthermore, all derivatives of H_2 , i.e. $\nabla^{\alpha\dot{\alpha}} \dots \nabla^{\alpha\dot{\alpha}} H_2 \equiv H_2^{\alpha(k)\dot{\alpha}(k)}$ $a^{(\kappa)}\alpha^{(\kappa)}$ can be factorized as

$$
\mathbf{H}^{\alpha(k)\,\dot{\alpha}(k)} = \lambda^{\alpha(k)}\widetilde{\mathbf{H}}_2^{\dot{\alpha}(k)} + \widetilde{\lambda}^{\dot{\alpha}(k)}\mathbf{H}_2^{\alpha(k)}.
$$
\n(3.19)

This special property of H_2 can be used to generate the linear FDA for the spin-2 sector. However, let us first specify the deformed data in [\(3.7\)](#page-10-3).

Solving deformed Cartan's structure equations. Assuming $C_{\dot{\alpha}(4)} = \phi_{(0,4)}\tilde{\lambda}_{\dot{\alpha}(4)}$, where $\phi_{(0,4)}$ is a scalar function, cf. [\(3.20e\)](#page-12-0), and plugging the ansatz [\(3.15\)](#page-11-0) to [\(3.7\)](#page-10-3), we obtain

$$
[\lambda \mathbf{H}_2] = 0, \tag{3.20a}
$$

$$
\boldsymbol{\varpi}_{\alpha\alpha} = \boldsymbol{\varpi}_{\alpha\alpha} + \mathbf{H}_2 \langle \lambda | h | \lambda \rangle \lambda_\alpha \lambda_\alpha , \qquad (3.20b)
$$

$$
\boldsymbol{e}_{\alpha\dot{\alpha}} = h_{\alpha\dot{\alpha}} + \langle \lambda \mathbf{H}_2 \rangle \langle \lambda | h | \lambda \rangle \lambda_{\alpha} \tilde{\lambda}_{\dot{\alpha}}, \qquad (3.20c)
$$

$$
\widetilde{\boldsymbol{\varpi}}_{\dot{\alpha}\dot{\beta}} = \widetilde{\boldsymbol{\varpi}}_{\dot{\alpha}\dot{\beta}} + \langle \lambda^{\gamma(2)} \mathbf{H}_{2\gamma(2)} \rangle \langle \lambda | h | \lambda] \widetilde{\lambda}_{\dot{\alpha}} \widetilde{\lambda}_{\dot{\alpha}},
$$
\n(3.20d)

$$
C_{\dot{\alpha}(4)} = \phi_{(0,4)}\tilde{\lambda}_{\dot{\alpha}(4)} , \qquad \phi_{(0,4)} = \frac{1}{2} \langle \lambda^{\gamma(3)} \mathcal{H}_{2\gamma(3)} \rangle , \qquad (3.20e)
$$

where

$$
[\lambda \mathrm{H}_2] := \tilde{\lambda}^{\dot{\alpha}} \mathrm{H}_{2\dot{\alpha}}, \qquad \langle \lambda^{\alpha(n)} \mathrm{H}_{2,\alpha(n)} \rangle := \lambda^{\alpha(n)} \mathrm{H}_{2\alpha(n)}, \qquad \lambda^{\alpha(n)} \equiv \lambda^{\alpha_1} \dots \lambda^{\alpha_n}.
$$
 (3.21)

In deriving the above results, the following relations are useful:

$$
h_{\alpha\dot{\alpha}}\langle\lambda|h|\lambda] = -\frac{1}{2}\Big(H_{\alpha\beta}\lambda^{\beta}\tilde{\lambda}_{\dot{\alpha}} + \tilde{H}_{\dot{\alpha}\dot{\beta}}\tilde{\lambda}^{\dot{\beta}}\lambda_{\alpha}\Big),\tag{3.22a}
$$

$$
d\langle \lambda | h | \lambda] H_2 = + \frac{1}{2} \Big(\mathcal{H}[\lambda H_2] + \widetilde{\mathcal{H}} \langle \lambda H_2 \rangle \Big) , \qquad (3.22b)
$$

$$
d\langle\lambda|h|\lambda]\langle\lambda^{\alpha(k)}\mathbf{H}_{2\alpha(k)}\rangle = +\frac{1}{2}\widetilde{\mathcal{H}}\langle\lambda^{\alpha(k+1)}\mathbf{H}_{2\alpha(k+1)}\rangle, \qquad (3.22c)
$$

where

$$
\mathcal{H} := H_{\alpha(2)} \lambda^{\alpha(2)}, \quad \widetilde{\mathcal{H}} := \widetilde{H}_{\dot{\alpha}(2)} \widetilde{\lambda}^{\dot{\alpha}(2)}.
$$
\n(3.23)

Observe that [\(3.20a\)](#page-12-1) implies the vanishing of all higher-order derivatives of H_2 in the \tilde{y} -sector. Thus, our analysis will focus solely on the derivative of H_2 in the y-sector.

From [\(3.20\)](#page-12-2), it is easy to see that the self-dual connection takes the form

$$
D = \nabla - \frac{1}{2} \mathbf{H}_2 \langle \lambda | h | \lambda \rangle \lambda_\alpha \lambda_\alpha L^{\alpha \alpha} - \frac{1}{2} \langle \lambda^{\gamma(2)} \mathbf{H}_2 \rangle_{\gamma(2)} \langle \lambda | h | \lambda \rangle \tilde{\lambda}_\alpha \tilde{\lambda}_\alpha \tilde{L}^{\dot{\alpha} \dot{\alpha}}.
$$
 (3.24)

Remarkably, the principle spinors λ_{α} , $\tilde{\lambda}_{\dot{\alpha}}$ are again covariantly constant wrt. the deformed connection D, i.e.

$$
D\lambda_{\alpha} = 0, \qquad D\tilde{\lambda}_{\dot{\alpha}} = 0. \tag{3.25}
$$

Therefore, D only acts non-trivially on the profile function H_2 and its derivative thereof, while "effortlessly" passing through (λ, λ) . This fact is essential for us to obtain a simple exact solution for chiral HSGRA with the above setting.

Note, however, that

$$
D\mu^{\alpha} = -\mathbf{H}_2 \langle \lambda | h | \lambda \rangle \lambda^{\alpha} , \qquad D^2 \mu^{\alpha} = 0 . \tag{3.26}
$$

Thus, unlike $(\lambda, \tilde{\lambda})$, the principal spinors $(\mu, \tilde{\mu})$ are not covariantly constant wrt. D.

Metric of self-dual pp-wave background. Remarkably, the SD background metric generated by $\omega_{\alpha\alpha}$ only slightly deviates from the conventional solutions in GR, cf. [\[61–](#page-28-11)[65\]](#page-28-12), by being self-dual and well-defined in Euclidean or split signature.

− Recall that the line element for Brinkmann metric in the conventional GR is

$$
ds2 = -du dv + H(v, z, \bar{z})dv2 + dz d\bar{z}.
$$
 (3.27)

Here, H is the profile function, which is harmonic in (z, \bar{z}) -plane, that generates a generic Petrov type-N spacetime given that $\ell_a = \lambda_\alpha \tilde{\lambda}_{\dot{\alpha}}$. As $H(v, z, \bar{z})$ is harmonic in the (z, \bar{z}) plane, we can write

$$
H(v, z, \bar{z}) = f_1(v, z) + f_2(v, \bar{z}), \qquad (3.28)
$$

for generic functions $f_1(v, z)$, $f_2(v, \overline{z})$. For real solutions, then it is necessary that $(f_2)^* = f_1$.

− In SD setting, one can deduce from the deformed vierbein [\(3.20c\)](#page-12-3) that

$$
g_{\mu\nu}^{SD} = \eta_{\mu\nu} + \langle \lambda \, \mathbf{H}_2 \rangle \ell_\mu \ell_\nu \,. \tag{3.29}
$$

That is, the metric is now generated by the first-derivative of H_2 in the z-direction.^{[6](#page-14-2)} In such a setting, we have

$$
ds_{SD}^2 = -du dv + \langle \lambda \, \mathbf{H}_2 \rangle dv^2 + dz d\tilde{z} \,, \qquad \langle \lambda \, \mathbf{H}_2 \rangle = j(v, z) \,, \tag{3.30a}
$$

$$
\phi_{(0,4)} = \langle \lambda^{\alpha(3)} \mathcal{H}_{2\alpha(3)} \rangle \sim \partial_z^3 j(v, z). \tag{3.30b}
$$

That is, the Brinkmann-like profile function is now dependent solely on (v, z) , consistent with the self-duality of the solution.

Intriguingly, even in the SD case, it is possible to place multiple pp-waves at different locations v_i along the v-direction so that the total profile function for the self-dual pp-wave metric can be written as $\langle \lambda H_2 \rangle = \sum_i \langle \lambda H_2^i(v_i) \rangle$.

3.3 Linear chiral-FDA

This subsection studies the linearized FDA for chiral HSGRA. Let us begin by stating our light-cone ansatz proposal for all positive-helicity spinning fields.

3.3.1 Light-cone ansatz for all spin

Following the discussion around [\(2.9\)](#page-5-3), it is reasonable to consider the following ansatz for all spins

$$
\omega_{\alpha(2s-2)} := \langle \lambda | h | \lambda \rangle \lambda_{\alpha(2s-2)} \mathbf{H}_s \,, \tag{3.31a}
$$

$$
B^{\alpha(2s)} := \psi_s \mu^{\alpha(2s)}, \qquad \psi_s := \lambda_{\alpha(2s)} B^{\alpha(2s)}, \quad \langle \mu \lambda \rangle = 1, \tag{3.31b}
$$

where H_s is a harmonic scalar function associated with a positive-helicity massless spin-s field, i.e.

$$
\ell^a \nabla_a \mathbf{H}_s = 0, \qquad p^a \tilde{p}^b \nabla_a \nabla_b \mathbf{H}_s = 0,
$$
\n(3.32)

as in the case of H_2 , cf. [\(3.16\)](#page-12-4). Furthermore, for the negative-helicity fields $B^{\alpha(2s)}$, we will assume that

$$
\nabla \mu^{\alpha} = 0, \qquad \ell^{a} \nabla_{a} \psi_{s} = 0, \qquad p^{a} \tilde{p}^{b} \nabla_{a} \nabla_{b} \psi_{s} = 0.
$$
 (3.33)

This guarantees that the free action of the model is [\(2.7\)](#page-5-2). Note that on the proposed SD pp-wave background

$$
D_{\beta}{}^{\dot{\alpha}}B^{\beta\alpha(2s-1)} = 0, \qquad (3.34)
$$

if $p^a \nabla_a \psi_s = 0$, or $\psi = \psi(v, \tilde{z})$. This is in accordance with the fact that ψ_s are harmonic, cf. [\(3.33\)](#page-14-3), and $B^{\alpha(2s)}$ are linear fluctuations on the proposed self-dual background. Thus, [\(3.33\)](#page-14-3) can serve as a suitable ansatz for the negative-helicity fields.

⁶The derivative is represented by the operator $p^a \nabla_a$.

3.3.2 H-FDA

Let us now cast the system (3.18) into generating function form. Since we have assumed H_s to be harmonic, cf. [\(3.32\)](#page-14-4), all $H_s^{\alpha(k) \dot{\alpha}(k)}$ will admit the same factorization form as in [\(3.19\)](#page-12-6). Then, we can write the factorization condition for $H_s^{\alpha(k) \dot{\alpha}(k)}$ using some generating functions $(H_s(y), \widetilde{H}_s(\tilde{y}))$ as

$$
\nabla \mathbf{H}_s(y) = \theta^\alpha \partial_\alpha \mathbf{H}_s(y) , \qquad \theta^\alpha := \tilde{\lambda}_{\dot{\beta}} h^{\alpha \dot{\beta}} , \qquad (3.35a)
$$

$$
\nabla \widetilde{\mathbf{H}}_s(y) = \widetilde{\theta}^{\dot{\alpha}} \partial_{\dot{\alpha}} \widetilde{\mathbf{H}}_s(\widetilde{y}), \qquad \widetilde{\theta}^{\dot{\alpha}} := \lambda_\beta h^{\beta \dot{\alpha}}.
$$
 (3.35b)

One can check that the above system satisfies the Bianchi identities

$$
\nabla^2 \mathbf{H}_s(y) = 0, \qquad \nabla^2 \widetilde{\mathbf{H}}_s(\widetilde{y}) = 0.
$$
 (3.36)

On the deformed background sourced by $\omega_{\alpha\alpha}$, cf. [\(3.15\)](#page-11-0), it can be shown that

$$
D\mathbf{H}_s^{\alpha(k)} = \theta_\beta \mathbf{H}_s^{\beta\alpha(k)} + k \mathbf{H}_2 \langle \lambda | h | \lambda \rangle \lambda^\alpha \lambda_\gamma \mathbf{H}_s^{\gamma\alpha(k-1)}, \tag{3.37a}
$$

$$
D\widetilde{\mathbf{H}}_{s}^{\dot{\alpha}(k)} = \widetilde{\theta}_{\dot{\beta}}\widetilde{\mathbf{H}}_{s}^{\dot{\beta}\dot{\alpha}(k)} + k\mathbf{H}_{2}\langle\lambda^{\gamma(2)}\mathbf{H}_{2\gamma(2)}\rangle\langle\lambda|h|\lambda]\widetilde{\lambda}^{\dot{\alpha}}\widetilde{\lambda}_{\dot{\gamma}}\widetilde{\mathbf{H}}_{s}^{\dot{\gamma}\dot{\alpha}(k-1)},\tag{3.37b}
$$

or in terms of generating function

$$
D\mathbf{H}_s(y) = \theta^{\alpha} (\partial_{\alpha} + \lambda_{\alpha} \langle \lambda y \rangle \langle \lambda \partial \rangle) \mathbf{H}_s(y),
$$

\n
$$
D\widetilde{\mathbf{H}}_s(\widetilde{y}) = \widetilde{\theta}^{\dot{\alpha}} (\widetilde{\partial}_{\dot{\alpha}} + \widetilde{\lambda}_{\dot{\alpha}} \langle \lambda^{\gamma(2)} \mathbf{H}_{2\gamma(2)} \rangle [\lambda y] [\lambda \partial]) \widetilde{\mathbf{H}}_s(\widetilde{y}).
$$
\n(3.38a)

Note that

$$
D^{2}H_{s}(y) = 0, \qquad D^{2}\tilde{H}_{s}(y) = 0.
$$
 (3.39)

Thus, as long as H_s is harmonic, cf. (3.32) , the above system can generate all auxiliary scalars which are derivative of H_s .

3.3.3 Linear FDA for spin-2 sector

Let us now consider the FDA for the harmonic spin-2 solution. As observed earlier in $(3.20a)$, where $[\lambda H_2] = 0$ due to self-duality constraint, it is natural to impose

$$
\widetilde{\mathrm{H}}_{2}^{\dot{\alpha}(k\geq 1)} = 0\,,\tag{3.40}
$$

which turns out to be a consistent condition for constructing a SD pp-wave background.

In the spin-2 sector, all data we need to generate to complete a linear-FDA are

$$
\{\omega_{\alpha\alpha}, \omega_{\alpha\dot{\alpha}}, \omega_{\dot{\alpha}\dot{\alpha}}\}\bigcup_{k=0}^{\infty} \{C^{\alpha(k)\dot{\alpha}(k+4)}\}\bigcup_{m=0}^{\infty} \{B^{\alpha(m+4)\dot{\alpha}(m)}\}.
$$
 (3.41)

Using [\(3.37\)](#page-15-2), and covariant constancy of $(\lambda, \tilde{\lambda})$ wrt. D, cf. [\(3.25\)](#page-13-0), we obtain

$$
C^{\alpha(k)\dot{\alpha}(4+k)} = \frac{1}{2} \langle \mathbf{H}_2^{\alpha(3+k)} \lambda_{\alpha(3)} \rangle \tilde{\lambda}^{\dot{\alpha}(4+k)}
$$
(3.42)

from solving iteratively $\nabla C_{\alpha(k)} \dot{\alpha}_{(4+k)} = h^{\beta \dot{\beta}} C_{\alpha(k) \beta \dot{\alpha}_{(4+k)} \dot{\beta}}$. The solution can be written in terms of generating function as

$$
C_4(y) = \frac{1}{(4+N_y)!} [y\lambda]^{4+N_y} \langle \lambda \partial' \rangle^3 \mathbf{H}_2(y') \Big|_{y'=y}, \qquad [y\lambda] = \tilde{y}^{\dot{\alpha}} \tilde{\lambda}_{\dot{\alpha}}, \quad N_y = y^{\alpha} \partial_{\alpha}.
$$
 (3.43)

For the negative-helicity auxiliary fields $B^{\alpha(m+4)\dot{\alpha}(m)}$, we observe that $B^{\alpha(4+m)\dot{\alpha}(m)}$ for $m \geq 1$ is non-linear in H₂ and their derivatives even for the case $\psi_2 = const.$ This is due to the fact that $D\mu^{\alpha} \neq 0$ when the deformation sourced by $\omega_{\alpha\alpha}$ occurs. Although we cannot find a closed form for $B^{\alpha(m+4)\dot{\alpha}(m)}$, we can still prove that the set of linearized data [\(3.41\)](#page-15-3) can solve the non-linear EOMs for chiral HSGRA, cf. [\(2.3\)](#page-4-3). Therefore, they provide us exact solutions for the spin-2 sector.

3.3.4 Linear FDA for all spins

Proceeding analogously with the case of spin-2, we will assume that

$$
\widetilde{\mathbf{H}}_s^{\dot{\alpha}(\bar{k}\geq 1)} = 0\,,\tag{3.44}
$$

and find the following linear data

$$
\bigcup_{s} \left\{ \bigcup_{j=0}^{2s-2} \{ \omega_{\alpha(2s-2-j)\,\dot{\alpha}(j)} \} \bigcup_{k=0}^{\infty} \{ C^{\alpha(k)\,\dot{\alpha}(2s+k)} \} \bigcup_{m=0}^{\infty} \{ B^{\alpha(2s+m)\,\dot{\alpha}(m)} \} \right\}.
$$
 (3.45)

one-form sector. In analogy with the spin-2 sector, we start with the light-cone ansatz [\(3.31a\)](#page-14-5), solve iteratively the FDA equation $\nabla \omega_{\alpha(2s-2-k)\dot{\alpha}(k)} = h_{\alpha}{}^{\beta} \omega_{\alpha(2s-3-k)\dot{\alpha}(k)\dot{\beta}}$ and obtain

$$
\omega_{\alpha(2s-2-k)\,\dot{\alpha}(k)} = \langle \lambda^{\gamma(k)} \mathcal{H}_{s\,\gamma(k)} \rangle \langle \lambda | h | \lambda] \lambda_{\alpha(2s-2-k)} \tilde{\lambda}_{\dot{\alpha}(k)} , \qquad 0 \le k \le 2s-2 . \tag{3.46}
$$

This can be written in terms of generating function as

$$
\omega_s = \frac{\langle \lambda | h | \lambda]}{\Gamma[2s - 1]} \Big(\langle y \lambda \rangle + [y \lambda] \langle \lambda \partial' \rangle \Big)^{2s - 2} \mathbf{H}_s(y') \Big|_{y' = 0} . \tag{3.47}
$$

Here, we keep $H_{s>3} \neq H_2$ for generality. However, it is intriguing noting that when $\{H_s\} = H_2$ for $\forall s$, we can subsume the full generating function ω of the one-form sector as

$$
\omega\big|_{\mathcal{H}_s = \mathcal{H}_2} = \sum_s \omega_{s \ge 1} \big|_{\mathcal{H}_s = \mathcal{H}_2} = \cosh\left(\langle y \,\lambda \rangle + [y \,\lambda] \langle \lambda \,\partial' \rangle\right) \mathcal{H}_2(y')\big|_{y'=0}.
$$
 (3.48)

zero-form sector. Similarly to the spin-2 case, we obtain the generating function for a spin-s 0-form field C as

$$
C^{\alpha(k)\dot{\alpha}(2s+k)} = \frac{1}{2} \langle \mathbf{H}_2^{\alpha(2s-1+k)} \lambda_{\alpha(2s-1)} \rangle \tilde{\lambda}^{\dot{\alpha}(2s+k)}
$$
(3.49)

or

$$
C_s(y) = \frac{1}{(2s + N_y)!} [y \lambda]^{2s + N_y} \langle \lambda \partial \lambda \rangle^{2s - 1} \mathcal{H}_s(y') \Big|_{y' = y}, \qquad [y \lambda] = \tilde{y}^{\dot{\alpha}} \tilde{\lambda}_{\dot{\alpha}}, \quad N_y = y^{\alpha} \partial_{\alpha}.
$$
 (3.50)

As before, $B^{\alpha(2s+m)\dot{\alpha}(m)}$ for $m \geq 1$ is non-linear in H_2 and their derivatives. However, we observe in general that

$$
B_s = B_s(\langle y \lambda \rangle, [y \lambda], \langle y \mu \rangle, \mathbf{H}_2, \psi_s). \tag{3.51}
$$

This fact is sufficient to prove that the set in [\(3.45\)](#page-16-1), whose 1-form subset consists of ω_s generating functions (cf. (3.47)), and the 0-form subset consists of $C_s = (C_s, B_s)$ fields, indeed solves the full non-linear equations of motion of chiral HSGRA.

Plane-wave solutions. In the case where [\[66\]](#page-29-0)

$$
H(y) = \exp\left(\pm x^{\alpha \dot{\alpha}} \kappa_{\alpha} \tilde{\lambda}_{\dot{\alpha}} + \langle y \kappa \rangle\right), \qquad \langle \lambda \kappa \rangle = 1, \qquad (3.52)
$$

where κ is a spinor associated with the momentum of a plane-wave. Then, it can be computed from [\(3.50\)](#page-17-1) and [\(3.47\)](#page-16-2) that

$$
C \sim \exp\left(\pm x^{\alpha \dot{\alpha}} \kappa_{\alpha} \tilde{\lambda}_{\dot{\alpha}} + [y \lambda]\right),\tag{3.53a}
$$

$$
\omega \sim \langle \lambda | h | \lambda \rangle \exp \left(\pm x^{\alpha \dot{\alpha}} \kappa_{\alpha} \tilde{\lambda}_{\dot{\alpha}} + \langle y \lambda \rangle + [y \lambda] \right). \tag{3.53b}
$$

These represent a family of plane-wave solutions for chiral HSGRA.

3.4 Non-linear chiral-FDA

Typically, when finding solutions for the non-linear equations of motion [\(2.3\)](#page-4-3), one often consider the following expansion of fields:

$$
\omega = \omega^{(1)} + \omega^{(2)} + \omega^{(3)} + \ldots = \Omega + \underline{\omega} + \omega^{(3)} + \ldots,
$$
\n(3.54a)

$$
C = C(1) + C(2) + C(3) + ... = C + C(2) + C(3) + ... ,
$$
\n(3.54b)

where $\Omega = \frac{1}{2}\boldsymbol{\varpi}_{\alpha\beta}L^{\alpha\beta} + \boldsymbol{e}_{\alpha\dot{\alpha}}P^{\alpha\dot{\alpha}} + \frac{1}{2}\boldsymbol{\widetilde{\varpi}}_{\dot{\alpha}\dot{\beta}}\tilde{L}^{\dot{\alpha}\dot{\beta}}$, and solve [\(2.3\)](#page-4-3) order by order as in e.g. [\[67,](#page-29-1) [68\]](#page-29-2). However, it is not always the case one can solve for (ω, C) to all orders and sum them up, since the computations quickly become quite involved.

Remarkably, it turns out that the 0-form $C = (C, B)$ in [\(3.50\)](#page-17-1) and the 1-form ω in [\(3.48\)](#page-16-3) solve the full non-linear equations of motion of chiral HSGRA. Thus, they form a family of exact solutions in chiral HSGRA. Note that we have denoted $\omega^{(2)} \equiv \omega = (3.47)$ $\omega^{(2)} \equiv \omega = (3.47)$ as well as $C^{(1)} = \underline{C} = (\underline{C}, \underline{B}), \text{ cf. } (3.50).$ $C^{(1)} = \underline{C} = (\underline{C}, \underline{B}), \text{ cf. } (3.50).$ $C^{(1)} = \underline{C} = (\underline{C}, \underline{B}), \text{ cf. } (3.50).$

Free equations of motion. As a consistency check, let us first derive free EOMs from [\(2.3\)](#page-4-3). Observe that for any generating function $f(y, \tilde{y})$, we have

$$
f \star \Omega \Big|_{\Lambda=0} = (1 + [\partial_1 \partial_2]) e^{\langle y \partial_1 \rangle + \langle y \partial_2 \rangle + [y \partial_1] + [y \partial_2]} f(y_1, \tilde{y}_1) \Omega(y_2, \tilde{y}_2) \Big|_{\substack{y_i = 0 \\ \tilde{y}_i = 0}}.
$$
 (3.55)

Thus, at linear order in fluctuation, we have

$$
d\underline{\omega} = \mathcal{V}(\Omega, \underline{\omega}) + \mathcal{V}(\Omega, \Omega, \underline{\mathsf{C}}) , \qquad (3.56a)
$$

$$
d\underline{\mathsf{C}} = \mathcal{U}(\Omega, \underline{\mathsf{C}})\,,\tag{3.56b}
$$

where $\mathcal{U}(\omega, \underline{C}) = U_{\omega, C} + U_{C, \omega}$ and

$$
U_{\omega,\underline{C}} = +\frac{1}{2} \mathcal{O}_2 \exp\left(\langle y \,\partial_2 \rangle + \langle \partial_1 \,\partial_2 \rangle\right) \omega_1(y_1,\tilde{y}_1) \underline{\mathsf{C}}_2(y_2,\tilde{y}_2) ,\tag{3.57a}
$$

$$
U_{\underline{\mathsf{C}},\omega} = -\frac{1}{2} \mathbf{U}_2 \exp\left(\langle y \, \partial_1 \rangle - \langle \partial_1 \, \partial_2 \rangle \right) \underline{\mathsf{C}}_1(y_1, \tilde{y}_1) \omega_2(y_2, \tilde{y}_2) ,\tag{3.57b}
$$

can be obtained from the natural pairings,^{[7](#page-18-0)} cf. Table [1.](#page-8-1) It can be then checked that (3.56) reproduces [\(2.10\)](#page-5-1).

 (ω, C) are exact solutions. It is useful noticing that as V's and U's vertices are constructed from the set of $\{\langle y \partial_i \rangle, \langle \partial_i \partial_j \rangle, [y \partial_i], [\partial_i \partial_j]\}$ operators, the building blocks for fields will not change, i.e. fields are always functions of $\{\langle y \lambda \rangle, [y \lambda], \langle y \mu \rangle, \langle \lambda \partial \prime \rangle, H_s, \psi_s\}.$

Proposition 3.1. Let

$$
\Omega \quad and \quad \underline{\mathsf{C}} = \underline{\mathsf{C}}(\langle y \lambda \rangle, \langle y \mu \rangle, [y \lambda], N_y, \langle \lambda \partial \rangle, \mathsf{H}_2, \psi_2)
$$

be the field content of the spin-2 sector. Then, all higher-order vertices $V_{n\geq 4}(\Omega,\Omega,\underline{C},\ldots,\underline{C})$ and $\mathcal{U}_{n>3}(\Omega,\underline{C},\ldots,\underline{C})$ vanish. Thus, the spin-2 sector characterized by (Ω,\underline{C}) forms a family of exact solutions in chiral HSGRA.

Proof. Let us first consider $V_{n\geq 4}(\Omega, \Omega, \underline{C}, \ldots, \underline{C})$. From [\(2.18\)](#page-7-0), it is easy to notice that there are forced contractions between the background connection 1-form Ω 's in the y-sector via the operations $\langle \partial_i \partial_j \rangle^{n-2}$ with n being the number of fields entering the sub-vertices V. It is easy to notice that $\langle \partial_i \partial_j \rangle^{n-2} \Omega_i \Omega_j = 0$ whenever $n > 4$. For the case $n = 4$, the contraction $\langle \partial_i \partial_j \rangle^2$ give^{[8](#page-18-2)}

$$
\boldsymbol{\varpi}^{\alpha\alpha}\boldsymbol{\varpi}_{\alpha\alpha} = (\varpi + \omega)^{\alpha\alpha}(\varpi + \omega)_{\alpha\alpha} = 0.
$$
\n(3.58)

Therefore, $V_{n\geq 4}(\Omega,\Omega,\underline{\mathsf{C}},\ldots,\underline{\mathsf{C}})=0$. Note that the cubic vertex $\mathcal{V}(\Omega,\Omega,\underline{\mathsf{C}})$ is needed for consistency as it produces the SD components of the Weyl tensor cf. [\(3.7c\)](#page-10-5).

⁷ recall that \mathcal{V}_2 is defined in (2.20) .

⁸The ∧-product is suppressed temporally for simplicity.

Next, we turn our focus to $\mathcal{U}_{n\geq3}(\Omega,\underline{C},\ldots,\underline{C})$ vertices, which can be obtained via the $U-V$ duality maps, cf. [\(B.9\)](#page-25-4). For $n > 5$, each sub-vertex U consists of a forced contraction of type $\langle y \partial_i \rangle^{n-2}$ where i is the position of Ω in $U_{\mathsf{C},\dots,\mathsf{C},\mathsf{\Omega}_i,\dots,\mathsf{C}}$. These sub-vertices clearly vanish for $n \geq 5$. Therefore, we only need to pay special attention to the cases where $n = 3, 4$.

• For $n = 3$, we have $\mathcal{U}(\Omega, \underline{C}, \underline{C}) = U_{\Omega, \underline{C}, \underline{C}} + U_{\underline{C}, \Omega, \underline{C}} + U_{\underline{C}, \underline{C}, \Omega}$ where [\[51\]](#page-28-2)

$$
U_{\Omega,\underline{C},\underline{C}} = +\mathcal{O}_3\langle y\,\partial_1\rangle \int e^{+(1-u)\langle y\,\partial_3\rangle + (1-v)\langle\partial_1\,\partial_3\rangle + u\langle y\,\partial_2\rangle + v\langle\partial_1\,\partial_2\rangle} ,\tag{3.59a}
$$

$$
U_{\mathbf{\underline{C}},\mathbf{\Omega},\mathbf{\underline{C}}} = -\mathcal{U}_3 \langle y \, \partial_2 \rangle \int e^{-(1-u)\langle \partial_1 \partial_2 \rangle + (1-v)\langle y \, \partial_1 \rangle + u\langle \partial_2 \partial_3 \rangle + v\langle y \, \partial_3 \rangle} + (v \leftrightarrow u) \,, \tag{3.59b}
$$

$$
U_{\mathsf{C},\mathsf{C},\mathbf{\Omega}} = +\mathcal{O}_3\langle y\,\partial_3\rangle \int e^{-(1-v)\langle\partial_1\,\partial_3\rangle + (1-u)\langle y\,\partial_1\rangle - v\langle\partial_2\,\partial_3\rangle + u\langle y\,\partial_2\rangle}.
$$
 (3.59c)

Here, $\mathcal{O}_3 = \exp\left([y \partial_1] + [y \partial_2] + [y \partial_3] + [\partial_1 \partial_2] + [\partial_1 \partial_3] + [\partial_2 \partial_3] \right)$ (cf., [\(2.20\)](#page-7-1)). We notice that there will be no contraction between \mathbb{C} 's in the \tilde{y} sector because the \star -product [\(2.12\)](#page-6-1) acts trivially on \underline{C}_i due to contractions between the principal spinors λ 's.^{[9](#page-19-0)} Note that the prefactors $\langle y \partial_i \rangle$ in each of the U sub-vertices act on Ω , so only $\varpi_{\alpha\alpha}$ and $e_{\alpha\dot{\alpha}}$ will give non-trivial contributions.

 $-\underline{\boldsymbol{\varpi}}$'s contributions. In this case, \mathcal{U}_3 acts trivially. Thus, all U's vertices reduce to

$$
U_{\nabla,\underline{C},\underline{C}} = + \int e^{+(1-u)\langle y\partial_3\rangle + (1-v)\langle \partial_1 \partial_3\rangle + u\langle y\partial_2\rangle + v\langle \partial_1 \partial_2\rangle} (\nabla_{1\alpha\alpha} y^{\alpha} y_1^{\alpha}, \underline{C}_2(y_2, \tilde{y}), \underline{C}_3(y_3, \tilde{y})) ,
$$

\n
$$
U_{\underline{C},\nabla,\underline{C}} = - \int e^{-(1-u)\langle \partial_1 \partial_2\rangle + (1-v)\langle y\partial_1\rangle + u\langle \partial_2 \partial_3\rangle + v\langle y\partial_3\rangle} (\underline{C}_1(y_1, \tilde{y}), \nabla_{2\alpha\alpha} y^{\alpha} y_2^{\alpha}, \underline{C}_3(y_3, \tilde{y}))
$$

\n
$$
- \int e^{-(1-v)\langle \partial_1 \partial_2\rangle + (1-u)\langle y\partial_1\rangle + v\langle \partial_2 \partial_3\rangle + u\langle y\partial_3\rangle} (\underline{C}_1(y_1, \tilde{y}), \nabla_{2\alpha\alpha} y^{\alpha} y_2^{\alpha}, \underline{C}_3(y_3, \tilde{y})) ,
$$

\n
$$
U_{\underline{C},\underline{C},\nabla} = + \int e^{-(1-v)\langle \partial_1 \partial_3\rangle + (1-u)\langle y\partial_1\rangle - v\langle \partial_2 \partial_3\rangle + u\langle y\partial_2\rangle} (\underline{C}_1(y_1, \tilde{y}), \underline{C}_2(y_2, \tilde{y}), \nabla_{3\alpha\alpha} y^{\alpha} y_3^{\alpha}).
$$

To see the pattern of cancellation, we can bring ϖ_i in to the first position in U. Since C's are 0-forms, the act of swapping positions of fields only provides minus signs in the exponential. Let the canonical ordering of fields in $\mathcal U$ be $\varpi, \underline{C}, \underline{C}$, we get

$$
\mathcal{U}(\boldsymbol{\varpi}, \underline{C}, \underline{C}) = + \int e^{(1-u)\langle y \partial_3 \rangle + u \langle y \partial_2 \rangle} \Big[2 + (1-v) \langle \partial_1 \partial_3 \rangle + v \langle \partial_1 \partial_2 \rangle \Big] (\boldsymbol{\varpi}_1, \underline{C}_2, \underline{C}_3)
$$

-
$$
\int e^{(1-v)\langle y \partial_1 \rangle + v \langle y \partial_3 \rangle} \Big[2 + (1-u) \langle \partial_2 \partial_1 \rangle + u \langle \partial_2 \partial_3 \rangle \Big] (\boldsymbol{\varpi}_2, \underline{C}_1, \underline{C}_3)
$$

-
$$
\int e^{(1-u)\langle y \partial_1 \rangle + u \langle y \partial_3 \rangle} \Big[2 + (1-v) \langle \partial_2 \partial_1 \rangle + v \langle \partial_2 \partial_3 \rangle \Big] (\boldsymbol{\varpi}_2, \underline{C}_1, \underline{C}_3)
$$

+
$$
\int e^{(1-u)\langle y \partial_1 \rangle + u \langle y \partial_2 \rangle} \Big[2 + (1-v) \langle \partial_3 \partial_1 \rangle + v \langle \partial_3 \partial_2 \rangle \Big] (\boldsymbol{\varpi}_3, \underline{C}_1, \underline{C}_2).
$$

⁹This is how one imposes locality into the formally consistent FDA (2.3) making it becomes a non-trivial system that describes chiral HSGRA, cf. [\[41,](#page-27-6) [42,](#page-27-7) [51\]](#page-28-2).

By simple change in variables, say $u \leftrightarrow v$ in the first line of [\(3.62a\)](#page-20-0), we see that all subvertices of U add up to zero since they cancel each other pairwise. We conclude that $U(\varpi, \underline{C}, \underline{C}) = 0$, which is a beautiful consequence of chiral HSGRA being a Lorentz-invariant theory.

− <u>e contributions.</u> In the case where $\mathcal{U}(\Omega, \underline{C}, \underline{C}) \sim \mathcal{U}(e, \underline{C}, \underline{C})$, we obtain

$$
U_{\mathbf{e},\underline{C},\underline{C}} = +\Xi_3 \big(2 + [\partial_1 \partial_2] + [\partial_1 \partial_3]\big) \int e^{+(1-u)\langle y \partial_3 \rangle + u\langle y \partial_2 \rangle} (e_{1\alpha\dot{\alpha}} y^{\alpha} \tilde{y}_1^{\dot{\alpha}}, \underline{C}_2, \underline{C}_3) ,
$$

\n
$$
U_{\underline{C},\mathbf{e},\underline{C}} = -\Xi_3 \big(2 - [\partial_2 \partial_1] + [\partial_2 \partial_3]\big) \int e^{+(1-v)\langle y \partial_1 \rangle + v\langle y \partial_3 \rangle} (e_{2\alpha\dot{\alpha}} y^{\alpha} \tilde{y}_2^{\dot{\alpha}}, \underline{C}_1, \underline{C}_3) + (v \leftrightarrow u) ,
$$

\n
$$
U_{\underline{C},\underline{C},\mathbf{e}} = +\Xi_3 \big(2 - [\partial_3 \partial_1] - [\partial_3 \partial_2]\big) \int e^{+(1-u)\langle y \partial_1 \rangle + u\langle y \partial_2 \rangle} (e_{3\alpha\dot{\alpha}} y^{\alpha} \tilde{y}_3^{\dot{\alpha}}, \underline{C}_1, \underline{C}_2) .
$$

where $\Xi_3 = \exp((y \partial_1) + (y \partial_2) + (y \partial_3))$. Observe that all $U_{e,\underline{C},\underline{C}}$ vertices cancel pairwise.

• For $n = 4$ case, where

$$
\mathcal{U}(\Omega,\underline{C},\underline{C},\underline{C})=U_{\varpi,\underline{C},\underline{C},\underline{C}}+U_{\underline{C},\varpi,\underline{C},\underline{C}}+U_{\underline{C},\underline{C},\varpi,\underline{C}}+U_{\underline{C},\underline{C},\underline{C},\varpi},
$$

and [\[51\]](#page-28-2)

$$
U_{\varpi,\underline{C},\underline{C},\underline{C}} = + \langle y\partial_1 \rangle^2 \int e^{(1-u_1-u_2)\langle y\partial_4 \rangle + (1-v_1-v_2)\langle \partial_1 \partial_4 \rangle + u_1 \langle y\partial_2 \rangle + u_2 \langle y\partial_3 \rangle + v_1 \langle \partial_1 \partial_2 \rangle + v_2 \langle \partial_1 \partial_3 \rangle} \qquad (3.62a)
$$

\n
$$
U_{\underline{C},\varpi,\underline{C},\underline{C}} = -\langle y\partial_2 \rangle^2 \int e^{(1-u_1-u_2)\langle y\partial_4 \rangle + (1-v_1-v_2)\langle \partial_2 \partial_4 \rangle + u_2 \langle y\partial_1 \rangle + u_1 \langle y\partial_3 \rangle - v_2 \langle \partial_1 \partial_2 \rangle + v_1 \langle \partial_2 \partial_3 \rangle} \qquad (3.62a)
$$

\n
$$
- \langle y\partial_2 \rangle^2 \int e^{(1-u_1-u_2)\langle y\partial_4 \rangle + (1-v_1-v_2)\langle \partial_2 \partial_4 \rangle + u_1 \langle y\partial_1 \rangle + u_2 \langle y\partial_3 \rangle - v_1 \langle \partial_1 \partial_2 \rangle + v_2 \langle \partial_2 \partial_3 \rangle} \qquad (3.62b)
$$

\n
$$
- \langle y\partial_2 \rangle^2 \int e^{(1-u_1-u_2)\langle y\partial_4 \rangle + (1-v_1-v_2)\langle \partial_3 \partial_4 \rangle + v^L \langle y\partial_1 \rangle + u^R \langle y\partial_3 \rangle - u^L \langle \partial_1 \partial_2 \rangle + v^R \langle \partial_2 \partial_3 \rangle} \qquad (3.62b)
$$

\n
$$
U_{\underline{C},\underline{C},\varpi,\underline{C}} = + \langle y\partial_3 \rangle^2 \int e^{(1-u_1-u_2)\langle y\partial_4 \rangle + (1-v_1-v_2)\langle \partial_3 \partial_4 \rangle + u_2 \langle y\partial_1 \rangle + u_1 \langle y\partial_2 \rangle - v_2 \langle \partial_1 \partial_3 \rangle - v_1 \langle \partial_2 \partial_3 \rangle} \qquad (3.62b)
$$

\n
$$
+ \langle y\partial_3 \rangle^2 \int e^{(1-v_1-v_2)\langle
$$

with $(u_i, v_i, u^L, u^R, v^L, v^R)$ are certain coefficients that are not crucial in the followings.

Again, let the canonical ordering of the quartic in the spin-2 sector be $(\varpi, \underline{C}, \underline{C}, \underline{C})$. We can ignore the pre-factors $\langle y \partial_i \rangle^2$ of each U's sub-vertices since they only produce $\varpi_{\alpha\alpha} y^{\alpha(2)}$. It can then be shown in a simple manner that all U sub-vertices cancel pairwise after some repositioning.

Therefore,
$$
V_{n\geq 4}(\Omega, \Omega, C, \dots, C) = 0
$$
 and $U_{n\geq 3}(\Omega, C, \dots, C) = 0$ as claimed.

Let us now state our main results.

Theorem 3.2. Given $\underline{\omega} = \underline{\omega}([y \lambda], \langle y \lambda \rangle, \langle \lambda \partial \theta \rangle, \mathbb{H})$, then $\mathcal{V}_{n \geq 3}(\underline{\omega}, \underline{\omega}, \underline{\mathsf{C}}, \dots, \underline{\mathsf{C}}) = 0$.

Proof. Recall that $V_n(\underline{\omega}, \underline{\omega}, \underline{\mathsf{C}}, \dots, \underline{\mathsf{C}})$ vertices have $\langle \partial_i \partial_j \rangle^{n-2}$ as prefactors where *i*th and *j*th are the positions of $\underline{\omega}_i, \underline{\omega}_j$ in the sub-vertices V's. As $\underline{\omega}_i$ are functions of $\langle y \lambda_i \rangle$, it is clear that $\mathcal{V}_n = 0$ for $n \geq 3$. \Box

Theorem 3.3. For $(\underline{\omega}, \underline{\mathsf{C}})$ defined as above, all vertices $\mathcal{U}_{n\geq 2}(\underline{\omega}, \underline{\mathsf{C}}, \ldots, \underline{\mathsf{C}}) = 0$.

Proof. Consider $\mathcal{U}_{n\geq 2}(\underline{\omega}, \underline{\mathsf{C}}, \ldots, C)$ whose sub-vertices U's carry $\langle y \partial_i \rangle^{n-2}$ as pre-factors. Then, all U's will have the same pre-factors which are proportional to $(\langle y \lambda \rangle^{n-2} \times ...)$. At this stage, we can repeat the proof of Proposition [3.1](#page-18-3) where we take the canonical ordering of fields in each U's sub-vertices to be $(\underline{\omega}, \underline{\mathsf{C}}, \ldots, \underline{\mathsf{C}})$. The claim is that all U's vertices will cancel pairwise based on how vertices of chiral HSGRA were designed [\[41,](#page-27-6) [42,](#page-27-7) [51\]](#page-28-2). This statement can be checked explicitly with $\mathcal{U}(\underline{\omega}, \underline{\mathsf{C}}) = U_{\omega, \mathsf{C}} + U_{\mathsf{C}, \omega}$, cf. [\(3.57\)](#page-18-4). It is analogous for the cubic [\(3.59\)](#page-19-1) (where we replace Ω by ω), and the quartic [\(3.62\)](#page-20-1). \Box

With the above results, it is clear that the complete non-linear FDA is solved by (ω, C) . Thus, the non-linear FDA solved by (ω, C) truncates to [\(3.56\)](#page-18-1) as it is not necessary to solve for higher-order fluctuations.

3.5 Effective action for chiral HSGRA on SD pp-wave background

We now show that the effective action associated to the above exact solutions is precisely the usual kinetic action for chiral higher-spin fields on any SD background.

In what follows, we denote

$$
\mathcal{F}(\mathcal{A}) := \left(d\omega - \mathcal{V}(\Omega, \omega) \right) \Big|_{\tilde{y}=0} = d\mathcal{A},\tag{3.63}
$$

to be the 'effective curvature tensor' for chiral HSGRA, since we have shown all higher-order vertices vanish on the solutions. Then, the effective action associated to the above field strength reads

$$
S_{\chi HSGRA}^{SD\,pp-wave} = \int_{\mathcal{M}} \langle \mathsf{C} \, | \, \mathcal{H} \wedge d\mathcal{A} \rangle_{y} \, , \qquad \mathcal{H} := \mathbf{e}_{\alpha\dot{\alpha}} \mathbf{e}_{\alpha}{}^{\dot{\alpha}} y^{\alpha} y^{\alpha} \, . \tag{3.64}
$$

where $\langle \ \rangle_y$ stands for the operation of integrating out y to produce appropriate contractions, cf. [\[39\]](#page-27-4). We obtain

$$
S[B,A] = \frac{1}{2} \sum_{s=1}^{\infty} \int \langle \mathcal{B}; \mathcal{H} \wedge d\mathcal{A} \rangle_{y} = \frac{1}{2} \sum_{s=1}^{\infty} \int \sqrt{g} B^{\alpha(2s)} D_{\alpha}{}^{\dot{\alpha}} A_{\alpha(2s-1)\dot{\alpha}}.
$$
 (3.65)

Thus, the proposed action provides the kinetic terms for all chiral higher-spin fields on a generic SD background, as promised.

4 Discussion

In this paper, we obtain a class of exact self-dual pp-wave solutions, in chiral HSGRA by considering a SD modification of the Kerr-Schild ansatz, cf. [\(3.15\)](#page-11-0). We then generalize our analysis to the case of higher spins. For completeness, we show that the spacetime action of the obtained solutions reduces to the standard kinetic action for free fields on any SD background.

Given that there are other self-dual solutions such as self-dual Taub-NUT black hole (see e.g. $[69–71]$ $[69–71]$), BPST instantons $[72]$, or ADHM instantons $[73]$, we expect that there will be more exact solutions in chiral HSGRA, which are higher-spin generalizations of the aforementioned self-dual solutions.

The reason we believe there should be more classes of exact solutions in chiral HSGRA is that this theory is integrable in the sense of Bardeen [\[74\]](#page-29-7), see e.g. [\[48\]](#page-27-13) and [\[49\]](#page-28-0) for relevant discussion. Therefore, chiral HSGRA should be solvable. This speculation is backed by the trivialization of the scattering amplitudes of chiral HSGRA in flat space, cf. [\[75–](#page-29-8)[77\]](#page-29-9), and some recent twistor construction of the theory or some closed subsectors of it, see e.g. [\[78](#page-29-10)[–81\]](#page-29-11).

It will be interesting to check whether all self-dual/chiral higher-spin theories possess this property. Note that, even if some of these theories may have quantum anomalies, as discussed in $[49]$, it is still possible to make them quantum integrable through a Green-Schwarz-like mechanism, as shown in [\[82,](#page-29-12) [83\]](#page-29-13). In particular, by introducing suitable axionic interactions in twistor space [\[84,](#page-29-14) [85\]](#page-29-15), one should be able to cure any quantum anomalies that arise in self-dual/chiral theories. A beautiful discussion related to this story in spacetime can also be found, e.g., in [\[86\]](#page-30-0).

While the analysis of chiral HSGRA is relatively straightforward in $\Lambda = 0$ case, it is more intricate in the case of (A)dS, where higher-order vertices will arise, as observed in [\[18,](#page-26-4) [19,](#page-26-5) [42\]](#page-27-7). (See also [\[87\]](#page-30-1) for a higher-dimensional and [\[88\]](#page-30-2) for a double-copy generalization of the theory). This fact makes the task of construction a covariant action for chiral HSGRA difficult. Nevertheless, one may hope that the light-cone ansatz introduced in [\[46,](#page-27-11) [47\]](#page-27-12) will provide further insight in guessing the correct action for chiral HSGRA due to the resulting simple form of higher-order vertices after substituting them in. See [\[89,](#page-30-3) [90\]](#page-30-4) for some work along this line.

Acknowledgement

Many discussions with Yasha Neiman and Zhenya Skvortsov are gratefully acknowledged. This work is supported by the Young Scientist Training (YST) program at the Asia Pacific Center of Theoretical Physics (APCTP) through the Science and Technology Promotion Fund and Lottery Fund of the Korean Government, and also the Korean Local Governments – Gyeongsangbuk-do Province and Pohang City. The author also appreciates the support from the Fonds de la Recherche Scientifique under grant number F.4503.20 (HighSpinSymm), grant number F.4544.21 (HigherSpinGraWave) and funding from the European Research Council (ERC) under grant number 101002551, when part of this work took place in Mons, Belgium.

A Convention and useful identities

In the main text, we have raised and lowered spinorial indices with the following convention:

 $h^{\mu}_{\alpha\dot{\alpha}}h_{\mu\beta\dot{\beta}}=2\epsilon_{\alpha\beta}\epsilon_{\dot{\alpha}\dot{\beta}}\,,\quad v_{\alpha}=v^{\beta}\epsilon_{\beta\alpha}\,,\quad v^{\alpha}=v_{\beta}\epsilon^{\alpha\dot{\beta}}\,,\quad v_{\dot{\alpha}}=v^{\dot{\beta}}\epsilon_{\dot{\beta}\dot{\alpha}}\,,\quad v^{\dot{\alpha}}=v_{\dot{\beta}}\epsilon^{\dot{\alpha}\dot{\beta}}\,$ $(A.1)$ Here, $\epsilon^{\alpha\beta}$ is the $\mathfrak{sp}(2)$ -invariant tensor with the properties $\epsilon^{\alpha\beta} = \epsilon_{\alpha\beta} = -\epsilon^{\beta\alpha}$ and $\epsilon^{01} = 1$. Furthermore, the following identities are useful

$$
h^{\alpha\dot{\alpha}} \wedge h^{\beta\dot{\beta}} = +\frac{1}{2} H^{\alpha\beta} \epsilon^{\dot{\alpha}\dot{\beta}} + \frac{1}{2} \widetilde{H}^{\dot{\alpha}\dot{\beta}} \epsilon^{\alpha\beta} , \qquad H^{\alpha\alpha} \wedge \widetilde{H}^{\dot{\alpha}\dot{\alpha}} = 0 , \qquad (A.2a)
$$

$$
H_{\alpha\alpha} \wedge H^{\alpha\alpha} = -\widetilde{H}_{\dot{\alpha}\dot{\alpha}} \wedge \widetilde{H}^{\dot{\alpha}\dot{\alpha}} = \text{Vol}\,,\tag{A.2b}
$$

$$
h^{\alpha\dot{\alpha}} \wedge H^{\beta\beta} = -\frac{1}{3} \epsilon^{\alpha\beta} \mathcal{H}_3^{\beta\dot{\alpha}}, \qquad h^{\alpha\dot{\alpha}} \wedge \widetilde{H}^{\dot{\beta}\dot{\beta}} = +\frac{1}{3} \epsilon^{\dot{\alpha}\dot{\beta}} \mathcal{H}_3^{\alpha\dot{\beta}}, \qquad (A.2c)
$$

where

$$
h^{\alpha}_{\dot{\gamma}} \wedge h^{\alpha \dot{\gamma}} = H^{\alpha \alpha} , \qquad h_{\gamma}^{\dot{\alpha}} \wedge h^{\gamma \dot{\alpha}} = \widetilde{H}^{\dot{\alpha} \dot{\alpha}} , \qquad h^{\alpha}_{\dot{\alpha}} \wedge H^{\dot{\beta} \dot{\alpha}} = \mathcal{H}_{3}^{\alpha \dot{\beta}} . \qquad (A.3)
$$

B Strong homotopy algebras

This appendix provides some basic facts about homotopy algebras adapted to the notations used in the main text.

Recall that the Q-manifold (N, Q) is a Z-graded target space $\mathcal{N} = \bigoplus_{n \in \mathbb{Z}} \mathcal{N}_n$ whose homogeneous elements a_n has degree $|a_n| = n$. Have we chosen to work with a degree k-shifted target space $\mathcal N$, the corresponding subspaces would be

$$
\mathcal{N}[k]_n := \mathcal{N}_{n+k} \,. \tag{B.1}
$$

Hereinafter, we use Koszul sign convention where $ab = (-)^{|a||b|}ba$ for two graded objects (a, b) of degrees |a| and |b|, respectively.

 A_{∞}/L_{∞} **-algebras.** Denote $C^{\bullet}(\mathcal{N})$ as the space of homomorphisms $\text{Hom}_{\mathbb{C}}(T\mathcal{N}, \mathcal{N})$, where $T\mathcal{N} := \bigoplus_{n=1}^{\infty} \mathcal{N}^{\otimes n}$ is the tensor algebra associated with \mathcal{N} , whose elements $\mathsf{F}_m \in C^m(\mathcal{N})$ are F_m maps with degree 1. We define the Gerstenhaber \circ -product as

$$
\mathsf{F}_n \circ \mathsf{F}_m := \sum_{k=0}^{n-1} (-)^{|\mathsf{F}_m|(|\Phi_1| + \ldots + |\Phi_k|)} \mathsf{F}_n(\Phi_1, \ldots, \Phi_k, \mathsf{F}_m(\Phi_{k+1}, \ldots, \Phi_{k+m}), \ldots, \Phi_{m+n-1}) \tag{B.2}
$$

where Φ 's are elements of \mathcal{N}_0 or \mathcal{N}_1 .

Definition B.1. The pair $(\mathcal{N}, \mathsf{F})$ where $\mathsf{F} = \sum_{i=1}^{\infty} \mathsf{F}_i$ is called A_{∞} -algebra if

$$
\mathsf{F} \circ \mathsf{F} = 0. \tag{B.3}
$$

Following from the above definition of A_{∞} -algebra, it is clear that $\mathsf{F}_1^2 = 0$, and therefore F_1 defines a differential. The next structure map F_2 is a Leibniz-compatible binary operation. Then, the homotopy given by F_3 measures the failure of the associativity of F_2 , and so on. For more detail, see e.g. [\[91\]](#page-30-5).

Note that an A_{∞} -algebra can induce an L_{∞} -algebra defined by a pair (\mathcal{N}, L) , where $\mathsf{L} = \sum \mathsf{L}_i$ with $|\mathsf{L}| = 1$ called L_{∞} -structure maps with suitable symmetrization among the arguments, i.e.

$$
\mathsf{L}_n(\Phi_1,\ldots,\Phi_n) = \sum_{\sigma \in S_n} (-)^K \mathsf{F}_n(\Phi_{\sigma(1)},\ldots,\Phi_{\sigma(n)})\,. \tag{B.4}
$$

Here, $(-)^K$ is the Koszul sign resulting from σ permutation under the symmetric group S_n . For instance,

$$
\mathsf{L}_n(\Phi_1,\ldots,\Phi_i,\Phi_{i+1},\ldots,\Phi_n) = (-)^{|\Phi_i||\Phi_{i+1}|} \mathsf{L}_n(\Phi_1,\ldots,\Phi_{i+1},\Phi_i,\ldots,\Phi_n).
$$
 (B.5)

As the reader may notice, L can be identified with the homological vector field Q on \mathcal{N} . Furthermore, following from [\(B.2\)](#page-23-2) and Definition [B.1,](#page-24-0) L's must obey:

$$
\sum_{m+n-1=\text{const}}\sum_{\sigma\in S_n}(-)^K \mathsf{L}_m(\mathsf{L}_n(\Phi_{\sigma(1)},\ldots,\Phi_{\sigma(n)}),\Phi_{\sigma(n+1)},\ldots,\Phi_{\sigma(n+m-1)})=0.
$$
 (B.6)

As usual, $L_1 \equiv d$ is the defining differential on $C^{\bullet}(\mathcal{N})$. Note that an L_{∞} -algebra with $L_{n>3} = 0$ is referred to as a dgLa.

The L_{∞} -structure maps $(\mathcal{V}, \mathcal{U})$ for chiral HSGRA can be obtained by restricting the super coordinates Φ 's to their appropriate components, i.e. to ω 's or C's. In particular, when there are strictly two ω arguments in L, then $L(\omega, \omega, C, \ldots, C) \equiv \mathcal{V}(\omega, \omega, C, \ldots, C)$, and when there is only one ω we have the identification $L(\omega, C, \ldots, C) \equiv \mathcal{U}(\omega, C, \ldots, C)$.

Although V's and U's are L_{∞} -structure maps, it will be more convenient to work with their sub-vertices which originate from a cyclic A_{∞} -algebra where the signs and permutations can be kept in track in a simple manner, cf. [\[41\]](#page-27-6). This can be done by letting ω and C take values in some matrix algebra $Mat(n, \mathbb{C})$ as alluded to in the main text. Furthermore, since V's and U's originated from the same L_{∞} -algebra, we can relate them together through the following nature pairing.

Definition B.2. A degree-2 cyclic L_{∞} -algebra is an L_{∞} -algebra endowed with a nondegenerate $\mathbb{C}\text{-linear pairing }\langle -|-\rangle:\mathcal{N}\oplus\mathcal{N}^{\vee}[-1]\to\mathbb{C}[-2]$ such that

$$
\langle a|b\rangle = (-)^{(|a|+1)(|b|+1)} \langle b|a\rangle. \tag{B.7}
$$

From the above definition, we have for instance

$$
\langle L_n(\Phi_1,\ldots,\Phi_n),\Phi_{n+1}\rangle = (-)^{|\Phi_1| \left(\sum_{k=2}^{n+1} |\Phi_k|\right) + |\Phi_1| + |\Phi_{n+2}|} \langle L_n(\Phi_2,\ldots,\Phi_{n+2}),\Phi_1\rangle, \quad (B.8a)
$$

$$
\langle L_n(\Phi_1,\ldots,\Phi_n)|\Phi_{n+1}\rangle = (-)^{\sum_{k=1}^n |\Phi_k|} \langle \Phi_1|L_n(\Phi_2,\ldots,\Phi_{n+1})\rangle ,\qquad (B.8b)
$$

Upon restricting Φ 's to ω and C, we obtain

$$
\langle V(\omega, \mathsf{C}, \dots, \omega, \mathsf{C}, \dots, \mathsf{C}) | C \rangle = + \langle \omega | U(\mathsf{C}, \dots, \omega, \mathsf{C}, \dots, \mathsf{C}) \rangle, \tag{B.9a}
$$

$$
\langle C|V(C,\ldots,\omega,C,\ldots,C,\omega)\rangle = -\langle U(C,\ldots,\omega,C,\ldots,C)|\omega\rangle, \qquad (B.9b)
$$

where V's and U's are sub-vertices originated from A_{∞} -algebra of V and U. The maps [\(B.9\)](#page-25-4) are called the V-U duality maps [\[41,](#page-27-6) [42,](#page-27-7) [51\]](#page-28-2), which can be used to extract U's vertices from \mathcal{V} 's (at a fixed order *n*), and vice versa.

References

- [1] X. Bekaert, N. Boulanger, A. Campoleoni, M. Chiodaroli, D. Francia, M. Grigoriev, E. Sezgin and E. Skvortsov, Snowmass White Paper: Higher Spin Gravity and Higher Spin Symmetry, [2205.01567](http://arXiv.org/abs/2205.01567).
- [2] S. Weinberg, Photons and Gravitons in S-Matrix Theory: Derivation of Charge Conservation and Equality of Gravitational and Inertial Mass, Phys. Rev. 135 (1964) B1049–B1056.
- [3] S. R. Coleman and J. Mandula, All Possible Symmetries of the S Matrix, Phys. Rev. 159 (1967) 1251–1256.
- [4] J. Maldacena and A. Zhiboedov, Constraining Conformal Field Theories with A Higher Spin Symmetry, J. Phys. A 46 (2013) 214011 [[1112.1016](http://arXiv.org/abs/1112.1016)].
- [5] X. Bekaert, J. Erdmenger, D. Ponomarev and C. Sleight, Quartic AdS Interactions in Higher-Spin Gravity from Conformal Field Theory, JHEP 11 (2015) 149 [[1508.04292](http://arXiv.org/abs/1508.04292)].
- [6] C. Sleight and M. Taronna, *Higher-Spin Gauge Theories and Bulk Locality*, *Phys. Rev. Lett.* 121 (2018), no. 17 171604 [[1704.07859](http://arXiv.org/abs/1704.07859)].
- [7] T. Tran, Constraining higher-spin S-matrices, JHEP 02 (2023) 001 [[2212.02540](http://arXiv.org/abs/2212.02540)].
- [8] Y. Neiman, Quartic locality of higher-spin gravity in de Sitter and Euclidean anti-de Sitter space, Phys. Lett. B 843 (2023) 138048 [[2302.00852](http://arXiv.org/abs/2302.00852)].
- [9] A. Y. Segal, Conformal higher spin theory, Nucl. Phys. B 664 (2003) 59–130 [[hep-th/0207212](http://arXiv.org/abs/hep-th/0207212)].
- [10] A. A. Tseytlin, On limits of superstring in $AdS(5)$ x S^{**} 5, Theor. Math. Phys. 133 (2002) 1376–1389 [[hep-th/0201112](http://arXiv.org/abs/hep-th/0201112)].
- [11] X. Bekaert, E. Joung and J. Mourad, Effective action in a higher-spin background, JHEP 02 (2011) 048 [[1012.2103](http://arXiv.org/abs/1012.2103)].
- [12] P. Hähnel and T. McLoughlin, *Conformal higher spin theory and twistor space actions*, *J.* Phys. A 50 (2017), no. 48 485401 [[1604.08209](http://arXiv.org/abs/1604.08209)].
- [13] T. Adamo, P. Hähnel and T. McLoughlin, Conformal higher spin scattering amplitudes from twistor space, JHEP 04 (2017) 021 [[1611.06200](http://arXiv.org/abs/1611.06200)].
- [14] T. Basile, M. Grigoriev and E. Skvortsov, Covariant action for conformal higher spin gravity, J. Phys. A 56 (2023), no. 38 385402 [[2212.10336](http://arXiv.org/abs/2212.10336)].
- [15] R. R. Metsaev, Poincare invariant dynamics of massless higher spins: Fourth order analysis on mass shell, Mod. Phys. Lett. A 6 (1991) 359–367.
- [16] R. R. Metsaev, S matrix approach to massless higher spins theory. 2: The Case of internal symmetry, Mod. Phys. Lett. A 6 (1991) 2411–2421.
- [17] D. Ponomarev and E. D. Skvortsov, Light-Front Higher-Spin Theories in Flat Space, J. Phys. A 50 (2017), no. 9 095401 [[1609.04655](http://arXiv.org/abs/1609.04655)].
- [18] R. R. Metsaev, Light-cone gauge cubic interaction vertices for massless fields in $AdS(4)$, Nucl. Phys. B 936 (2018) 320–351 [[1807.07542](http://arXiv.org/abs/1807.07542)].
- [19] E. Skvortsov, Light-Front Bootstrap for Chern-Simons Matter Theories, JHEP 06 (2019) 058 [[1811.12333](http://arXiv.org/abs/1811.12333)].
- [20] R. R. Metsaev, *Cubic interaction vertices for N=1 arbitrary spin massless supermultiplets in* flat space, JHEP 08 (2019) 130 [[1905.11357](http://arXiv.org/abs/1905.11357)].
- [21] R. R. Metsaev, Cubic interactions for arbitrary spin $\mathcal N$ -extended massless supermultiplets in 4d flat space, JHEP 11 (2019) 084 [[1909.05241](http://arXiv.org/abs/1909.05241)].
- [22] M. Tsulaia and D. Weissman, Supersymmetric quantum chiral higher spin gravity, JHEP 12 (2022) 002 [[2209.13907](http://arXiv.org/abs/2209.13907)].
- [23] M. P. Blencowe, A Consistent Interacting Massless Higher Spin Field Theory in $D = (2+1)$, Class. Quant. Grav. 6 (1989) 443.
- [24] E. Bergshoeff, M. P. Blencowe and K. S. Stelle, Area Preserving Diffeomorphisms and Higher Spin Algebra, Commun. Math. Phys. 128 (1990) 213.
- [25] A. Campoleoni, S. Fredenhagen, S. Pfenninger and S. Theisen, Asymptotic symmetries of three-dimensional gravity coupled to higher-spin fields, JHEP 11 (2010) 007 [[1008.4744](http://arXiv.org/abs/1008.4744)].
- [26] M. Henneaux and S.-J. Rey, *Nonlinear* $W_{infinity}$ as Asymptotic Symmetry of Three-Dimensional Higher Spin Anti-de Sitter Gravity, JHEP 12 (2010) 007 [[1008.4579](http://arXiv.org/abs/1008.4579)].
- [27] C. N. Pope and P. K. Townsend, Conformal Higher Spin in (2+1)-dimensions, Phys. Lett. B 225 (1989) 245–250.
- [28] E. S. Fradkin and V. Y. Linetsky, A Superconformal Theory of Massless Higher Spin Fields in $D = (2+1)$, Mod. Phys. Lett. A 4 (1989) 731.
- [29] M. Grigoriev, I. Lovrekovic and E. Skvortsov, New Conformal Higher Spin Gravities in 3d, JHEP 01 (2020) 059 [[1909.13305](http://arXiv.org/abs/1909.13305)].
- [30] M. Grigoriev, K. Mkrtchyan and E. Skvortsov, Matter-free higher spin gravities in 3D: Partially-massless fields and general structure, Phys. Rev. D 102 (2020), no. 6 066003 [[2005.05931](http://arXiv.org/abs/2005.05931)].
- [31] A. Sharapov, E. Skvortsov and A. Sukhanov, Matter-coupled higher spin gravities in 3d: noand yes-go results, [2409.12830](http://arXiv.org/abs/2409.12830).
- [32] K. Alkalaev and X. Bekaert, *Towards higher-spin AdS₂/CFT*₁ holography, JHEP 04 (2020) 206 [[1911.13212](http://arXiv.org/abs/1911.13212)].
- [33] M. Sperling and H. C. Steinacker, Covariant cosmological quantum space-time, higher-spin and gravity in the IKKT matrix model, JHEP 07 (2019) 010 [[1901.03522](http://arXiv.org/abs/1901.03522)].
- [34] H. C. Steinacker, Gravity as a quantum effect on quantum space-time, Phys. Lett. B 827 (2022) 136946 [[2110.03936](http://arXiv.org/abs/2110.03936)].
- [35] H. C. Steinacker and T. Tran, Soft limit of higher-spin interactions in the IKKT model, [2311.14163](http://arXiv.org/abs/2311.14163).
- [36] H. C. Steinacker and T. Tran, Quantum hs-Yang-Mills from the IKKT matrix model, Nucl. Phys. B 1005 (2024) 116608 [[2405.09804](http://arXiv.org/abs/2405.09804)].
- [37] A. Manta, H. C. Steinacker and T. Tran, hs-extended gravity from the IKKT matrix model, [2411.02598](http://arXiv.org/abs/2411.02598).
- [38] E. Skvortsov and Y. Yin, Low spin solutions of higher spin gravity: BPST instanton, JHEP 07 (2024) 032 [[2403.17148](http://arXiv.org/abs/2403.17148)].
- [39] K. Krasnov, E. Skvortsov and T. Tran, Actions for self-dual Higher Spin Gravities, JHEP 08 (2021) 076 [[2105.12782](http://arXiv.org/abs/2105.12782)].
- [40] R. Penrose, Twistor algebra, J. Math. Phys. 8 (1967) 345.
- [41] A. Sharapov, E. Skvortsov, A. Sukhanov and R. Van Dongen, Minimal model of Chiral Higher Spin Gravity, [2205.07794](http://arXiv.org/abs/2205.07794).
- [42] A. Sharapov and E. Skvortsov, *Chiral Higher Spin Gravity in* $(A)dS₄$ *and secrets of* Chern-Simons Matter Theories, [2205.15293](http://arXiv.org/abs/2205.15293).
- [43] D. Sullivan, *Infinitesimal computations in topology*, *Publications Mathématiques de l'IHÉS* 47 (1977) 269–331.
- [44] M. Alexandrov, A. Schwarz, O. Zaboronsky and M. Kontsevich, The Geometry of the master equation and topological quantum field theory, Int. J. Mod. Phys. A 12 (1997) 1405–1429 [[hep-th/9502010](http://arXiv.org/abs/hep-th/9502010)].
- [45] K. Krasnov and E. Skvortsov, Flat Self-dual Gravity, [2106.01397](http://arXiv.org/abs/2106.01397).
- [46] T. Adamo and T. Tran, *Higher-spin Yang-Mills, amplitudes and self-duality*, [2210.07130](http://arXiv.org/abs/2210.07130).
- [47] Y. Neiman, Self-dual gravity in de Sitter space: Light-cone ansatz and static-patch scattering, Phys. Rev. D 109 (2024), no. 2 024039 [[2303.17866](http://arXiv.org/abs/2303.17866)].
- [48] D. Ponomarev, Chiral Higher Spin Theories and Self-Duality, JHEP 12 (2017) 141

[[1710.00270](http://arXiv.org/abs/1710.00270)].

- [49] R. Monteiro, From Moyal deformations to chiral higher-spin theories and to celestial algebras, JHEP 03 (2023) 062 [[2212.11266](http://arXiv.org/abs/2212.11266)].
- [50] A. Sharapov, E. Skvortsov and R. Van Dongen, Chiral Higher Spin Gravity and Convex Geometry, [2209.01796](http://arXiv.org/abs/2209.01796).
- [51] A. Sharapov, E. Skvortsov, A. Sukhanov and R. Van Dongen, More on Chiral Higher Spin Gravity and convex geometry, Nucl. Phys. B 990 (2023) 116152 [[2209.15441](http://arXiv.org/abs/2209.15441)].
- [52] P. Van Nieuwenhuizen, Free graded differential superalgebras, in Group Theoretical Methods in Physics: Proceedings of the XIth International Colloquium Held at Boğaziçi University, Istanbul, Turkey August 23–28, 1982, pp. 228–247, Springer, 2005.
- [53] R. D'Auria, P. Fre and T. Regge, Graded Lie Algebra Cohomology and Supergravity, Riv. Nuovo Cim. 3N12 (1980) 1.
- [54] M. A. Vasiliev, Consistent Equations for Interacting Massless Fields of All Spins in the First Order in Curvatures, Annals Phys. 190 (1989) 59–106.
- [55] J. Stasheff, Homotopy associativity, H-spaces from a Homotopy Point of View (2006) 27–30.
- [56] D. Ponomarev, Towards higher-spin holography in flat space, JHEP 01 (2023) 084 [[2210.04035](http://arXiv.org/abs/2210.04035)].
- [57] R. Penrose and W. Rindler, SPINORS AND SPACE-TIME. VOL. 2: SPINOR AND TWISTOR METHODS IN SPACE-TIME GEOMETRY. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 4, 1988.
- [58] E. Newman and R. Penrose, An Approach to gravitational radiation by a method of spin coefficients, J. Math. Phys. 3 (1962) 566–578.
- [59] R. P. Kerr and A. Schild, Republication of: A new class of vacuum solutions of the einstein field equations, General Relativity and Gravitation 41 (2009), no. 10 2485–2499.
- [60] K. Krasnov, Self-Dual Gravity, Class. Quant. Grav. 34 (2017), no. 9 095001 [[1610.01457](http://arXiv.org/abs/1610.01457)].
- [61] J. M. Stewart, Advanced general relativity. Cambridge Monographs on Mathematical Physics. Cambridge University Press, 4, 1994.
- [62] H. Brinkmann, Einstein spaces which are mapped conformally on each other, Mathematische Annalen 94 (1925), no. 1 119–145.
- [63] P. C. Aichelburg and R. U. Sexl, On the Gravitational field of a massless particle, Gen. Rel. Grav. 2 (1971) 303–312.
- [64] A. Bilge and M. Gürses, *Generalized kerr-schild transformation*, in *Group Theoretical* Methods in Physics, pp. 252–255. Springer, 1983.
- [65] H. Stephani, D. Kramer, M. A. H. MacCallum, C. Hoenselaers and E. Herlt, Exact solutions of Einstein's field equations. Cambridge Monographs on Mathematical Physics. Cambridge Univ. Press, Cambridge, 2003.
- [66] E. Skvortsov and R. Van Dongen, Minimal models of field theories: Chiral Higher Spin Gravity, [2204.10285](http://arXiv.org/abs/2204.10285).
- [67] V. E. Didenko and A. V. Korybut, Planar solutions of higher-spin theory. Part I. Free field level, JHEP 08 (2021) 144 [[2105.09021](http://arXiv.org/abs/2105.09021)].
- [68] V. E. Didenko and A. V. Korybut, Planar solutions of higher-spin theory. Nonlinear corrections, JHEP 01 (2022) 125 [[2110.02256](http://arXiv.org/abs/2110.02256)].
- [69] S. W. Hawking, Gravitational Instantons, Phys. Lett. A 60 (1977) 81.
- [70] C. P. Boyer and J. D. Finley, III, Killing Vectors in Selfdual, Euclidean Einstein Spaces, J. Math. Phys. 23 (1982) 1126.
- [71] T. Adamo, G. Bogna, L. Mason and A. Sharma, Scattering on self-dual Taub-NUT, Class. Quant. Grav. 41 (2024), no. 1 015030 [[2309.03834](http://arXiv.org/abs/2309.03834)].
- [72] A. A. Belavin, A. M. Polyakov, A. S. Schwartz and Y. S. Tyupkin, Pseudoparticle Solutions of the Yang-Mills Equations, Phys. Lett. B 59 (1975) 85–87.
- [73] M. F. Atiyah, N. J. Hitchin, V. G. Drinfeld and Y. I. Manin, Construction of Instantons, Phys. Lett. A 65 (1978) 185–187.
- [74] W. A. Bardeen, Selfdual Yang-Mills theory, integrability and multiparton amplitudes, Prog. *Theor. Phys. Suppl.* 123 (1996) 1–8.
- [75] E. D. Skvortsov, T. Tran and M. Tsulaia, Quantum Chiral Higher Spin Gravity, Phys. Rev. Lett. 121 (2018), no. 3 031601 [[1805.00048](http://arXiv.org/abs/1805.00048)].
- [76] E. Skvortsov, T. Tran and M. Tsulaia, More on Quantum Chiral Higher Spin Gravity, Phys. Rev. D 101 (2020), no. 10 106001 [[2002.08487](http://arXiv.org/abs/2002.08487)].
- [77] E. Skvortsov and T. Tran, One-loop Finiteness of Chiral Higher Spin Gravity, JHEP 07 (2020) 021 [[2004.10797](http://arXiv.org/abs/2004.10797)].
- [78] T. Tran, Twistor constructions for higher-spin extensions of (self-dual) Yang-Mills, JHEP 11 (2021) 117 [[2107.04500](http://arXiv.org/abs/2107.04500)].
- [79] T. Tran, Toward a twistor action for chiral higher-spin gravity, [2209.00925](http://arXiv.org/abs/2209.00925).
- [80] Y. Herfray, K. Krasnov and E. Skvortsov, Higher-spin self-dual Yang-Mills and gravity from the twistor space, JHEP 01 (2023) 158 [[2210.06209](http://arXiv.org/abs/2210.06209)].
- [81] Y. Neiman, Higher-spin self-dual General Relativity: 6d and 4d pictures, covariant vs. lightcone, JHEP 07 (2024) 178 [[2404.18589](http://arXiv.org/abs/2404.18589)].
- [82] K. Costello and S. Li, Quantization of open-closed BCOV theory, I, [1505.06703](http://arXiv.org/abs/1505.06703).
- [83] K. Costello and S. Li, Anomaly cancellation in the topological string, Adv. Theor. Math. Phys. 24 (2020), no. 7 1723-1771 [[1905.09269](http://arXiv.org/abs/1905.09269)].
- [84] K. J. Costello, *Quantizing local holomorphic field theories on twistor space*, [2111.08879](http://arXiv.org/abs/2111.08879).
- [85] R. Bittleston, D. Skinner and A. Sharma, Quantizing the Non-linear Graviton, Commun.

Math. Phys. 403 (2023), no. 3 1543-1609 [[2208.12701](http://arXiv.org/abs/2208.12701)].

- [86] R. Monteiro, R. Stark-Muchão and S. Wikeley, Anomaly and double copy in quantum self-dual Yang-Mills and gravity, JHEP 09 (2023) 030 [[2211.12407](http://arXiv.org/abs/2211.12407)].
- [87] T. Basile, Massless chiral fields in six dimensions, [2409.12800](http://arXiv.org/abs/2409.12800).
- [88] D. Ponomarev, Chiral higher-spin double copy, [2409.19449](http://arXiv.org/abs/2409.19449).
- [89] C. Chowdhury, G. Doran, A. Lipstein, R. Monteiro, S. Nagy and K. Singh, Light-cone actions and correlators of self-dual theories in AdS4, [2411.04172](http://arXiv.org/abs/2411.04172).
- [90] K. Krasnov and A. Lipstein, Pure connection formalism and Plebanski's second heavenly equation, [2412.15779](http://arXiv.org/abs/2412.15779).
- [91] H. Kajiura, Noncommutative homotopy algebras associated with open strings, Rev. Math. Phys. 19 (2007) 1-99 [[math/0306332](http://arXiv.org/abs/math/0306332)].