

Pathwise guessing in categorical time series with unbounded alphabets

J.-R. Chazottes ^{*1}, S. Gallo ^{†2}, and D. Y. Takahashi ^{‡3}

¹Centre de Physique Théorique, CNRS, Ecole Polytechnique,
Institut Polytechnique de Paris, France

²Departamento de Estatística, Universidade Federal de São Carlos,
São Paulo, Brazil

³Instituto do Cérebro, Universidade Federal do Rio Grande do
Norte, Natal, Brazil

January 15, 2025

Abstract

The following learning problem arises naturally in various applications: Given a finite sample from a categorical or count time series, can we learn a function of the sample that (nearly) maximizes the probability of correctly guessing the values of a given portion of the data using the values from the remaining parts? Unlike the classical task of estimating conditional probabilities in a stochastic process, our approach avoids explicitly estimating these probabilities.

We propose a non-parametric guessing function with a learning rate that is independent of the alphabet size. Our analysis focuses on a broad class of time series models that encompasses finite-order Markov chains, some hidden Markov chains, Poisson regression for count process, and one-dimensional Gibbs measures.

Additionally, we establish a minimax lower bound for the rate of convergence of the risk associated with our guessing problem. This lower bound matches the upper bound achieved by our estimator up to a logarithmic factor, demonstrating its near-optimality.

Key-words: stochastic chain of unbounded memory, countable alphabets, Dvoretzky-Kiefer-Wolfowitz type inequality, Markov chains, autoregressive models, hidden Markov chains, one-dimensional Gibbs measures, Poisson regression for count time series.

*Email: jeanrene@cpt.polytechnique.fr

†Email: sandro.gallo@ufscar.br

‡Email: takahashiyd@gmail.com

1 Introduction

Prediction and interpolation, *i.e.*, *guessing*, are foundational problems in science with wide-ranging applications, particularly in the context of categorical time series. Prediction involves estimating future values of a sequence based on prior observations, while interpolation focuses on inferring missing values within a sequence of known data. Both tasks aim to construct a function from observed data that can accurately predict or interpolate the missing components. These problems have garnered renewed interest due to the rise of large language models, which can be viewed as categorical time series models with large alphabets [12]. In this new context, both the alphabet size and the dependence order are typically high. Our work specifically addresses models with unbounded dependence and alphabet sizes, making them suitable for modern applications.

A common approach to these problems relies on estimating conditional probabilities. Classical prediction involves pointwise estimation of all transition probabilities, as discussed in [19]. Other studies, such as [25] and [14], use PAC learning frameworks to investigate optimal rates for learning conditional probabilities across various metrics, with [13] addressing k -step Markov chains with large alphabets using minimax Kullback-Leibler risk. This last article introduces alternative loss functions to mitigate sensitivity to rare transitions, showing that optimal prediction can be achieved with an alphabet size scaling as $\mathcal{O}(\sqrt{n})$. Additionally, parametric regression models, known for their flexibility and algorithmic efficiency [9], are another common approach to time series prediction. However, such models often impose restrictive assumptions on the underlying processes and may not efficiently solve the guessing problem in general. In all these approaches, the guessing becomes challenging when the alphabet size is large or when transition probabilities are very small. Accurately estimating rare events requires a substantial amount of data, which can be impractical.

A more practical approach is to concentrate on the events that are most likely to happen. This means focusing our efforts on predicting the most probable outcomes and giving less weight to rare, unlikely events. This approach is especially useful when dealing with sequences that have a huge or even infinite number of possible symbols. Even with large sets of symbols, this method can still make accurate predictions, unlike traditional methods that might struggle to estimate the probabilities of all possible transitions.

In this paper, we present a probabilistic guessing scheme that addresses these challenges. In guessing, we want to learn from the sample a function that correctly guesses the values on a given portion of the time series, given the remaining parts. Our non-parametric estimator applies to a broad class of categorical time series [16], accommodating arbitrary alphabet sizes and relaxing constraints on memory or order. A key feature of our approach is that the learning rate of the associated risk is independent of the alphabet size, and we establish that this rate is essentially sharp. The proof of the upper bound is based on a Dvoretzky-Kiefer-Wolfowitz-type inequality, which we previously established in a slightly different form [1]. To prove the lower bound, we use classical Le Cam's type of argument [26].

This paper is organized as follows: Section 2 introduces the basic notation and formalizes the probabilistic guessing scheme. Section 3 details our assumptions and presents the main theoretical results. Examples of time series models that satisfy our assumptions are discussed in Section 4, and the proofs of the main results are provided in Section 5.

2 The probabilistic guessing problem

We will now explain the formulation of what we call the probabilistic guessing problem. Before that let us introduce the notation.

Notation. Let \mathcal{A} be a countable set with cardinality at least 2 (the set of categories, or the alphabet) endowed with the discrete topology and denote by $\mathcal{A}^{\mathbb{Z}}$ the set of bi-infinite sequences drawn from \mathcal{A} . We then put the product topology on $\mathcal{A}^{\mathbb{Z}}$. It is generated by the cylinder sets $[a_{-n+1}, \dots, a_{n-1}] = \{x \in \mathcal{A}^{\mathbb{Z}} : x_i = a_i, |i| \leq n-1\}$, $a_i \in \mathcal{A}$, $n \in \mathbb{N}$, and comprises all Borel sets of $\mathcal{A}^{\mathbb{Z}}$.

For $D \subset \mathbb{Z}$, $D_k := D \cap [-k, k]$ and for $b \in \mathcal{A}^D$, we define the truncation $b_k \in \mathcal{A}^{D_k}$ by $(b_k)_j = b_j$ for all $j \in D_k$. We use the notation \Subset to mean “finite subset of”. Given $H \Subset \mathbb{Z}$, the diameter of H is defined as $\text{diam}(H) := \max H - \min H + 1$. For integers $m < n$, let $\llbracket m, n \rrbracket := [m, n] \cap \mathbb{Z}$.

For any pair $x, y \in \mathcal{A}^{\mathbb{Z}}$, and any $\Lambda \Subset \mathbb{Z}$, the point $z = x_{\Lambda} y_{\Lambda^c}$ is defined by $z_i = x_i$ for $i \in \Lambda$, and $z_i = y_i$ for $i \notin \Lambda$. We define the shift operator θ acting on subsets of \mathbb{Z} . For $H \subset \mathbb{Z}$ and $i \in \mathbb{Z}$, let $\theta^i H = H + i$, where $H + i = \{i + j : j \in H\}$. Similarly, the shift operator T on $\mathcal{A}^{\mathbb{Z}}$ is defined by $(T^i x)_n = x_{n+i}$, for $i, n \in \mathbb{Z}$.

Given a \mathcal{A} -valued process $(X_i)_{i \in \mathbb{Z}}$, we write $X_m^n = (X_m, X_{m+1}, \dots, X_n)$ ($m < n$), and more generally, for $G \subset \mathbb{Z}$, $X_G = (X_i)_{i \in G}$. We denote by $\sigma(X_G)$ the σ -algebra generated by $(X_i)_{i \in G}$.

The probabilistic guessing problem. We want to find an estimator in which the probability to correctly guess a given subset of symbols is close (with a precision that we can choose) to the best possible guess. We will formulate this problem as follows. Let $(X_j)_{j \in \mathbb{Z}}$ and $(Y_j)_{j \in \mathbb{Z}}$ be two independent copies of the same process with distribution \mathbb{P} . We want to use the information of the *data set* D to predict the value on the *guess set* G . An estimator is a function

$$\begin{aligned} \hat{f}_{D,G}^n : \mathcal{A}^n \times \mathcal{A}^D &\rightarrow \mathcal{A}^G \\ \hat{f}_{D,G}^n[x_1^n](y_D) &= y_G, \end{aligned}$$

which we usually think as a random function on the second coordinate

$$\hat{f}_{D,G}^n[X_1^n](y_D) := \hat{f}_{D,G}^n(y_D),$$

and it naturally depends on n . To measure the performance of our estimator, we define the *excess risk* as

$$R(\hat{f}_{D,G}^n) \tag{1} \\ := \sup_{b \in \mathcal{A}^D} \left(\tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G \mid Y_D = b) - \inf_{a \in \mathcal{A}^G} \tilde{\mathbb{P}}(a \neq Y_G \mid Y_D = b) \right) \tilde{\mathbb{P}}(Y_D = b),$$

where $\tilde{\mathbb{P}} = \mathbb{P} \otimes \mathbb{P}$ (product measure). Note that $R(\hat{f}_{D,G}^n)$ depends \mathbb{P} , although we do not explicitly indicate this in the notation as long as \mathbb{P} is fixed.

Assume that we observe a finite sample X_1, \dots, X_n . We want to obtain an estimator $\hat{f}_{D,G}^n$ such that $R(\hat{f}_{D,G}^n)$ is small. Two natural cases are the “prediction” problem, which corresponds to the case $\max D < \min G$, and the “interpolation” problem, which corresponds to the case $G \subset \llbracket \min D, \max D \rrbracket$.

Our choice of the risk in (1) warrants some explanation. An alternative risk could be defined as the “unweighted” risk:

$$\sup_{b \in \mathcal{A}^D} \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G \mid Y_D = b) - \inf_{a \in \mathcal{A}^G} \tilde{\mathbb{P}}(a \neq Y_G \mid Y_D = b).$$

The drawback of this alternative is that it requires controlling errors even when $\tilde{\mathbb{P}}(Y_D = b)$ is small. This typically demands larger sample sizes, with minimal improvement to the overall probability of guessing.

Another natural risk can be defined using the function $f : \mathcal{A}^D \rightarrow \mathcal{A}^G$, where

$$f(b) = \arg \min_{a \in \mathcal{A}^G} \mathbb{P}(a \neq Y_G \mid Y_D = b).$$

The minimum is guaranteed to exist since \mathbb{P} is a probability measure. A seemingly equivalent risk to (1) is then given by

$$\sup_{b \in \mathcal{A}^D} \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq f(Y_D), Y_D = b).$$

However, this choice has its own caveat. When several symbols have probabilities close to the minimum, estimating f becomes more challenging. Despite this, these symbols yield similar guessing errors, making it unnecessary to distinguish between them.

Consider a simple example: a biased die where the side labeled with the symbol “one” has the highest probability of appearing. If each roll is independent, the optimal prediction is to always guess “one”. In this scenario, it suffices to correctly identify that “one” has the highest probability; there is no need to precisely estimate the probabilities of the other symbols. Furthermore, the total number of symbols on the die is irrelevant to the accuracy of this prediction.

Now suppose the probabilities for all symbols on the die are nearly identical. In this case, any symbol can be chosen, and the probability of a correct guess will still be close to the theoretical optimum. Here, even though it might be challenging to identify the symbol with the highest probability, the exact identification might be less relevant, as the prediction error is negligible. Such situations are common when the number of categories is large.

Finally, consider a die with a Markovian dynamics, where the outcome of each roll depends on the previous one. In this case, the optimal guess depends on the past observations, with the accuracy of the guess tied to the probabilities associated with prior outcomes. When a past outcome has a low probability, it becomes less critical to make an accurate prediction because the event requiring a guess will occur rarely. Moreover, even when the rare event happens, the chance of correctly predicting it remains slim. Our proposed formulation explicitly captures and formalizes this practical consideration.

To conclude with our formulation of the guessing problem, observe that it shares similarities with non-parametric multi-class classification for dependent processes. However, a key distinction is that we do not aim to classify all labels accurately. Instead, we focus only on those labels with a high probability of occurrence, enabling us to extend the method to cases involving countably infinite alphabets.

The estimator. We will use the following estimator.

DEFINITION 2.1. Given $n \geq 1$, X_1^n , $G, D \in \mathbb{Z}$, such that $\min(G \cup D) = 1$, and $b \in \mathcal{A}^D$, let

$$\hat{f}_{D,G}^n[X_1^n](b) := \arg \max_{a \in \mathcal{A}^G} \frac{N_{D,G}^n[X_1^n](b, a)}{N_{D,G}^n[X_1^n](b)} = \arg \max_{a \in \mathcal{A}^G} \frac{N_{D,G}^n[X_1^n](b, a)}{n}, \quad (2)$$

where

$$N_{D,G}^n[X_1^n](b, a) := \sum_{i=0}^{n-1} \mathbb{1}\{X_{\theta^i D} = b, X_{\theta^i G} = a\},$$

and

$$N_{D,G}^n[X_1^n](b) := \sum_{a \in \mathcal{A}^G} N_{D,G}^n[X_1^n](b, a)$$

are the counting functions, with the convention that the indicator function is zero if $\theta^i D$ or $\theta^i G$ is not included in $\llbracket 1, n \rrbracket$. We will often omit the reference to the sample on which we are counting occurrences, writing $\hat{f}_{D,G}^n(b)$ and $N_{D,G}^n(b, a)$. The key observation for our estimator is that the maximizing argument for the conditioned empirical probability is equal to the maximizing argument for the unconditioned empirical probability.

3 Assumption and statement of the results

Our main assumption about the categorical process concerns the degree of dependence it exhibits on its “past”. Let us denote by p a regular version of the left conditional expectations of the process $(X_i)_{i \in \mathbb{Z}}$ with distribution \mathbb{P} , that is, $p : \mathcal{A} \times \mathcal{A}^{\mathbb{Z}^-} \rightarrow [0, 1]$ is such that

$$\mathbb{E}(\mathbb{1}_{\{X_0=a\}} \mid \sigma(X_{-\infty}^{-1})) (x) = p(a|x) \quad \mathbb{P}\text{-a.s.}, \quad (3)$$

where $\mathbb{Z}_- := \{-1, -2, \dots\}$ is the set of negative integers. We say that, for \mathbb{P} -almost all “pasts” $x \in \mathcal{A}^{\mathbb{Z}_-}$, p specifies the transition probabilities to any symbol of \mathcal{A} . Let us also define the *variation of p* , $(\text{Var}_n(p))_{n \geq 0}$, as

$$\text{Var}_n(p) := \sup \left\{ \frac{1}{2} \sum_{a \in \mathcal{A}} |p(a|x) - p(a|y)| : x, y \in \mathcal{A}^{\mathbb{Z}_-}, x_i = y_i, i \geq -n \right\}, n \geq 0,$$

with the convention that for $n = 0$ the supremum is taken over any pair $x, y \in \mathcal{A}^{\mathbb{Z}_-}$ without restriction. Our main regularity assumption concerning p is that $\text{Var}_n(p)$ goes to 0 sufficiently fast when n diverges.

ASSUMPTION A. Let $(X_i)_{i \in \mathbb{Z}}$ be a stationary process. We say that $(X_i)_{i \in \mathbb{Z}}$ satisfy Assumption A if its left conditional expectations satisfies

$$\Gamma(p) := \prod_{j=0}^{\infty} (1 - \text{Var}_j(p)) > 0.$$

Observe that Assumption A is equivalent to having

$$\text{Var}_0(p) < 1 \quad \text{and} \quad \sum_{j \geq 1} \text{Var}_j(p) < \infty.$$

Stationary processes defined by (3) are commonly referred to in the literature as chains of infinite order, chains with complete connections, or stochastic chains with unbounded memory, or g -measures (in the context of ergodic theory); see *e.g.* [8]. An insightful reference that examines these objects in the context of categorical time series is [10]. Assumption A aligns with classical conditions from the literature, providing a framework to ensure that conditional expectations are not overly influenced by distant past events. This assumption captures a wide range of well-known models, including Markov chains and generalized linear models for categorical and count time series, one-dimensional Gibbs measures, as detailed in Section 4.

We are now ready to state our results.

THEOREM 3.1. Let $G, D \in \mathbb{Z}$ and put $L := \text{diam}(G \cup D)$. Let $(X_j)_{j \in \mathbb{Z}}$ be a stationary process with kernel p satisfying Assumption A. Let $\varepsilon > 0$. If

$$n \geq \frac{16}{\varepsilon^2} \left(\frac{(|G| + |D|)^2}{\Gamma(p)^2} \log \left(\frac{1}{\varepsilon} \right) + \frac{4(|G| + |D|)(1 - \Gamma(p))}{\Gamma(p)} + 2 \right) + L - 2, \quad (4)$$

then

$$R(\hat{f}_{D,G}^n) \leq \varepsilon. \quad (5)$$

A direct consequence is that taking $\varepsilon = C \sqrt{\log n/n}$ for some constant $C > 0$ in Theorem 3.1, C can be taken large enough such that (4) is satisfied. This proves the following corollary.

COROLLARY 3.1. Let $G, D \in \mathbb{Z}$. If $(X_j)_{j \in \mathbb{Z}}$ is a stationary process with kernel p and measure \mathbb{P} that satisfies Assumption A, then there exist a positive constant $C = C(p, G, D)$ such that, for all $n \geq 2$,

$$R(\hat{f}_{D,G}^n) \leq C \sqrt{\frac{\log n}{n}}. \quad (6)$$

A natural question then arises: is the learning rate achieved in this corollary optimal? The result below establishes that, up to a logarithmic factor, our estimator attains the minimax optimal rate of convergence for the risk.

To state the result, we let \mathcal{P} denote the set of distributions of all stationary stochastic processes on alphabet \mathcal{A} whose left conditional expectation satisfies Assumption A. We will now explicitly write the dependence on \mathbb{P} of the excess risk (see (1)).

THEOREM 3.2. Fix $G, D \in \mathbb{Z}$. For every positive integer n , let Ψ_n be the set of estimators $\psi_n : \mathcal{A}^n \times \mathcal{A}^D \rightarrow \mathcal{A}^G$. There exists a positive constant $C = C(|G|, |D|)$ such that for all $n \geq 1$,

$$\inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} R(\psi_n; \mathbb{P}) \geq \frac{C}{\sqrt{n}}. \quad (7)$$

4 Examples satisfying Assumption A

As previously mentioned, Assumption A is quite general. We now motivate this claim listing some classical classes of categorical times series and some examples that fit this assumption.

Independent random variables. For independent random variables, the function p defined in (3) is independent of $x \in \mathcal{A}^{\mathbb{Z}^-}$. Consequently, $\text{Var}_n(p) = 0$ for all $n \geq 0$, which implies $\Gamma(p) = 1$. Even in this simple case, our result seems to be novel, as it provides a learning rate that does not depend on the alphabet size.

Markov chains. Let us consider a Markov chain with transition matrix $Q = (Q(a, b))_{a, b \in \mathcal{A}}$. In this case $p(a|x) = Q(x_{-1}, a)$ for any $x \in \mathcal{A}^{\mathbb{Z}^-}$, thus $\text{Var}_n(p) = 0$ for any $n \geq 1$, and condition (A) becomes equivalent to

$$\text{Var}_0(p) = d(Q) := \sup_{a, b \in \mathcal{A}} \|Q(a, \cdot) - Q(b, \cdot)\|_{\text{TV}} < 1,$$

and $\Gamma(p)$ can be substituted by $1 - d(Q)$. The quantity $d(Q)$ is called the Dobrushin ergodicity coefficient (see for instance [6] and [7, Section 18.2]).

More generally, a k -steps Markov chain will satisfy $\text{Var}_n(p) = 0$ for any $n \geq k$, and therefore our results apply as long as $\text{Var}_0(p) < 1$ as well.

Autoregressive models. Consider a function $\Upsilon : \mathbb{R} \rightarrow (0, 1)$ such that $\Upsilon(r) + \Upsilon(-r) = 1$ and an absolutely summable sequence of real numbers $(\xi_j)_{j \geq 0}$. Consider then the binary process $(X_j)_{j \in \mathbb{Z}}$ with transition probabilities given by

$$p(a|x) = \Upsilon\left(a \sum_{j=1}^{\infty} \xi_j x_{-j} + a\xi_0\right).$$

The process generated by this kernel is called a binary auto-regressive process [16]. A simple and yet classical situation is when $\Upsilon(u) = (1 + e^{-2u})^{-1}$, $\xi_i \geq \xi_j \geq 0$ for all $j > i \geq 1$, and $\xi_0 = \sum_{k=1}^{\infty} \xi_k$. In this case we have

$$\frac{2e^{2\xi_1}}{(1 + e^{2\xi_1})^2} \sum_{k>j} \xi_k \leq \text{Var}_j(g) \leq \sum_{k>j} \xi_k,$$

indicating that if $\sum_{j>1} \sum_{k>j} \xi_k < \infty$ to get $\Gamma(p) > 0$, the autoregressive process satisfies Assumption A.

Poisson regression for count time series. Let $\mathcal{A} = \mathbb{N}$ and $(\xi_j)_{j \geq 0}$ be a sequence of non-positive absolutely summable real numbers, and a constant $c > 0$. For all $x \in \mathcal{A}^{\mathbb{Z}^-}$, let

$$v(x) = \exp\left(\sum_{j=1}^{\infty} \xi_j \min\{x_{-j}, c\}\right).$$

For all $a \in \mathbb{N}$ and $x \in \mathcal{A}^{\mathbb{Z}^-}$, the kernel of a Poisson regression model (see [16]) is defined as

$$p(a|x) = \frac{e^{-v(x)} v(x)^a}{a!}.$$

Applying the mean value theorem to $r \mapsto e^{-e^r} e^{ra} / a!$, and maximizing on $r \in (-\infty, 0]$ for each $a \in \mathbb{N}$, we obtain that $\Gamma(p)$ is finite if $\sum_{j>1} \sum_{k>j} \xi_k < \infty$.

Hidden Markov chains. Let $(X_j)_{j \in \mathbb{Z}}$ be a Markov chain with transition matrix P on a finite alphabet \mathcal{A} and a function $f : \mathcal{A} \rightarrow \mathcal{B}$ where \mathcal{B} is another alphabet with $|\mathcal{B}| < |\mathcal{A}|$. Specifically, this means that certain symbols in \mathcal{A} are indistinguishable or merged, which can, for instance, be attributed to a measurement error. The process $(Y_j)_{j \in \mathbb{Z}}$ defined by $Y_i = f(X_i)$ is a hidden Markov process, and it is usually not Markov of any order. The works of [2, Theorem 3.1], [5] and [20] for instance, give quantitative information on the variation of $(Y_j)_{j \in \mathbb{Z}}$, allowing us to compute $\Gamma(p)$. In particular, when $P > 0$ and \mathcal{A} is finite, the variation of the projected process decays exponentially fast. If, in turn, certain symbols of \mathcal{B} become indistinguishable or are merged, the variation of the resulting process will continue to decay exponentially [20].

Convex mixture of Markov chains. Let $(\lambda_j)_{j \geq 1}$ be a sequence of non-negative real numbers such that $\sum_{j=1}^{\infty} \lambda_j = 1$. Define a family of finite-order Markov kernels $p^{[k]} : \mathcal{A} \times \mathcal{A}^{\llbracket -k, -1 \rrbracket} \rightarrow [0, 1]$, $k \geq 1$. The kernel for mixture of Markov chains is defined, for all $a \in \mathcal{A}$ and $x \in \mathcal{A}^{\mathbb{Z}^-}$, as

$$p(a|x) = \sum_{j=1}^{\infty} \lambda_j p^{[j]}(a|x_{-j}^{-1}).$$

We have $\Gamma(p) \geq \prod_{k \geq 1} (1 - \sum_{j \geq k} \lambda_j)$ which is > 0 whenever $\sum_{j=1}^{\infty} j \lambda_j < \infty$ and $\lambda_0 > 0$. This is a quite general class of kernels since all kernels p such that $\lim_{j \rightarrow \infty} \text{Var}_j(p) = 0$ can be represented as a convex mixture of Markov chains [15, 4].

Gibbs measures. Here we assume that $|\mathcal{A}| < \infty$. Gibbs measures, or Gibbs random fields, originally introduced in the context of statistical physics, form a natural and extensively studied class of examples in probability theory. These measures are particularly relevant to our guessing problem because they are defined via two-sided conditional probabilities (as formalized in (9)). Their inherent structure and rich theoretical framework make them well-suited for addressing such probabilistic inference challenges. Here we only consider Gibbs measures on the one-dimensional lattice \mathbb{Z} .

Formally, consider a collection $\Phi := (\Phi_{\Lambda})_{\Lambda \in \mathbb{Z}}$ of real-valued functions on $\mathcal{A}^{\mathbb{Z}}$ such that, for any x , $\Phi_{\Lambda}(x)$ depends only on x_{Λ} . This is called an *interaction potential*. We assume translation invariance, that is, $\Phi_{\theta\Lambda} \circ T = \Phi_{\Lambda}$, and absolutely summability, that is,

$$\sum_{\substack{\Lambda \in \mathbb{Z} \\ 0 \in \Lambda}} \|\Phi_{\Lambda}\|_{\infty} < \infty. \quad (8)$$

The *energy* of a state x in the “box” Λ is $H_{\Lambda}(x) := \sum_{\Lambda' \in \mathbb{Z}: \Lambda' \cap \Lambda \neq \emptyset} \Phi_{\Lambda'}(x)$.

A Gibbs measure with a potential Φ is given by any measure that satisfies

$$\mathbb{P}(X_{\Lambda} = x_{\Lambda} \mid X_{\Lambda^c} = y_{\Lambda^c}) \stackrel{\mathbb{P}\text{-a.s.}}{=} \frac{\exp(-\beta H_{\Lambda}^{\Phi}(x_{\Lambda} y_{\Lambda^c}))}{Z_{\Lambda, \beta}^{\Phi}(y)}, \quad \Lambda \in \mathbb{Z}, \quad (9)$$

where $Z_{\Lambda}^{\Phi}(y)$ is the normalizing factor and $\beta \geq 0$ is a constant known as “inverse temperature”. In other words, they are specified by inside *vs* outside (finite boxes Λ) conditioning instead of the present *vs* past conditioning of (3). For finite range potentials, the corresponding Gibbs measure is a k -steps Markov chain for some k and with $\text{Var}_0(p) < 1$ (see [11, Theorem 3.5]). For Gibbs measure with unbounded range potential Φ , we prove the following proposition.

PROPOSITION 4.1. Let \mathbb{P} be a Gibbs measure with potential Φ . Define $\Delta(\Phi_{\Lambda}) := \max \Phi_{\Lambda} - \min \Phi_{\Lambda}$. If Φ is such that

$$\sum_{k \geq 1} \sum_{i \geq k} \sum_{\substack{\min \Lambda = 0 \\ \max \Lambda \geq i}} \Delta(\Phi_{\Lambda}) < \infty, \quad (10)$$

then the corresponding left conditional expectation satisfies Assumption (A).

As an example, consider the 1D long range Ising model, a classical example of non-Markovian Gibbs measure. Let $\mathcal{A} = \{-1, +1\}$. Fix $\alpha > 1$ and consider the potential $\Phi_\Lambda(x) = |i - j|^{-\alpha} x_i x_j$ if $\Lambda = \{i, j\}$ with $i \neq j$, and $\Phi_\Lambda(x) = 0$ otherwise. For $\alpha > 2$, the process $(X_j)_{j \in \mathbb{Z}}$ is uniquely specified by (9) (see for instance [11, (8.41)]). We have that $\sum_{\min \Lambda=0, \max \Lambda \geq i} \Delta(\Phi_\Lambda) = \sum_{\ell \geq i} \Delta(\Phi_{\{0, \ell\}})$ which is $\mathcal{O}(i^{-\alpha+1})$. Thus, as a consequence of Proposition 4.1, Assumption (A) is satisfied when $\alpha > 3$.

In the framework of Gibbs measures, there exists a natural extension of Markov chains where conditional independence is a two-sided property. This leads to the exploration of concepts analogous to those found in hidden Markov chains. It was shown in [22] that the resulting process is a Gibbs measure with an interaction potential that decays exponentially. Starting from an exponentially decaying interaction potential, the transformed potential retains this property. In the specific case of the one-dimensional long-range Ising model with $\alpha > 2$, the transformation yields a potential of the same type but with $\alpha' = \alpha - 1$. Consequently, to ensure that the transformed process satisfies Assumption (A), it is necessary to begin with $\alpha > 4$.

5 Proofs

5.1 Proof of Theorem 3.1

Let us define

$$\gamma_{D,G}(b, \varepsilon) = \left\{ c \in \mathcal{A}^G : \mathbb{P}(X_G \neq c, X_D = b) - \inf_{a \in \mathcal{A}^G} \mathbb{P}(X_G \neq a, X_D = b) \leq \varepsilon \right\}.$$

We have

$$\begin{aligned} & \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G, Y_D = b) \\ &= \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G, Y_D = b, \hat{f}_{D,G}^n(Y_D) \in \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})) \\ & \quad + \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G, Y_D = b, \hat{f}_{D,G}^n(Y_D) \notin \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})). \end{aligned}$$

From the definition of $\gamma_{D,G}$, we have

$$\begin{aligned} & \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G, Y_D = b, \hat{f}_{D,G}^n(Y_D) \in \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})) \\ &= \sum_{c \in \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})} \tilde{\mathbb{P}}(c \neq Y_G, Y_D = b, \hat{f}_{D,G}^n(b) = c) \\ &= \sum_{c \in \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})} \mathbb{P}(c \neq Y_G, Y_D = b) \mathbb{P}(\hat{f}_{D,G}^n(b) = c) \\ &\leq \inf_{a \in \mathcal{A}^G} \mathbb{P}(a \neq Y_G, Y_D = b) + \frac{\varepsilon}{2}. \end{aligned}$$

We also have that, for all $b \in A^D$,

$$\begin{aligned} & \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(Y_D) \neq Y_G, Y_D = b, \hat{f}_{D,G}^n(Y_D) \notin \gamma_{D,G}(Y_D, \frac{\varepsilon}{2})) \\ & \leq \tilde{\mathbb{P}}(\hat{f}_{D,G}^n(b) \notin \gamma_{D,G}(b, \frac{\varepsilon}{2})). \end{aligned}$$

Observe that

$$\gamma_{D,G}(b, \varepsilon) = \left\{ c \in \mathcal{A}^G : \sup_{a \in \mathcal{A}^G} \mathbb{P}(X_G = a, X_D = b) - \mathbb{P}(X_G = c, X_D = b) \leq \varepsilon \right\}.$$

Hence, if

$$\sup_{a \in \mathcal{A}^G, b \in \mathcal{A}^D} \left| \frac{N_{G,D}^n(b, a)}{n} - \mathbb{P}(X_G = a, X_D = b) \right| \leq \frac{\varepsilon}{4}, \quad (11)$$

then, for all $b \in A^D$,

$$\hat{f}_{D,G}^n(b) \in \gamma_{D,G}(b, \frac{\varepsilon}{2}).$$

It remains to bound from above the probability that (11) is not satisfied. For this, we use Theorem A.1 which gives us the following Dvoretzky-Kiefer-Wolfowitz inequality: for any $u > 0$ and $n \geq L := \text{diam}(G \cup D)$,

$$\begin{aligned} & \mathbb{P} \left(\max_{a \in \mathcal{A}^G, b \in \mathcal{A}^D} \left| \frac{N_{D,G}^n(b, a)}{n} - \mathbb{P}(X_G = a, X_D = b) \right| > u + \sqrt{\frac{2M(1-\Gamma) + \Gamma}{\Gamma(n-L+2)}} \right) \\ & \leq \exp \left(-2(n-L+2)\Gamma^2 M^{-2} u^2 \right), \end{aligned} \quad (12)$$

where we used $M := |D| + |G|$, and wrote Γ for $\Gamma(p)$ to alleviate notation. Equating the upper bound of (12) to ε , we obtain

$$u = \frac{M}{\Gamma} \sqrt{\frac{\log(\frac{1}{\varepsilon})}{2(n-L+2)}}. \quad (13)$$

If n satisfies (4), we obtain

$$\frac{\varepsilon^2}{16} \geq 2 \left(u^2 + \frac{2M(1-\Gamma) + \Gamma}{\Gamma(n-L+2)} \right).$$

Now, using the elementary inequality $(\sqrt{u} + \sqrt{v})^2 \leq 2(u+v)$ for $u, v \geq 0$, we conclude that

$$\frac{\varepsilon}{4} \geq u + \sqrt{\frac{2M(1-\Gamma) + \Gamma}{\Gamma(n-L+2)}},$$

which finishes the proof.

5.2 Proof of Theorem 3.2

Our proof follows a classical procedure to obtain minimax lower bounds for estimators. Let us first write the minimax problem. Let $n \geq 1$. Writing $\tilde{\mathbb{P}} = \mathbb{P} \otimes \mathbb{P}$, we have by definition

$$\begin{aligned} & \inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} R(\psi_n; \mathbb{P}) \\ &= \inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} \sup_{b \in \mathcal{A}^D} \left(\tilde{\mathbb{P}}(\psi_n(X_1^n, b) \neq Y_G, Y_D = b) - \inf_{a \in \mathcal{A}^G} \tilde{\mathbb{P}}(a \neq Y_G, Y_D = b) \right). \end{aligned}$$

Now, we want to obtain a lower bound for the left hand side of the above equation. Fix an element $b \in \mathcal{A}^D$. We will later choose a specific b . Let $a_{\mathbb{P}} = \arg \min_{a \in \mathcal{A}^G} \tilde{\mathbb{P}}(a \neq Y_G, Y_D = b)$. A minimizer exists, because the probability sums to one. If there is more than one minimizer, we can just choose one of them. To simplify the notation, let us abbreviate $\psi_n(X_1^n, b)$ as $\psi_n(b)$ in the next estimations. We have

$$\begin{aligned} & \tilde{\mathbb{P}}(\psi_n(b) \neq Y_G, Y_D = b) - \tilde{\mathbb{P}}(a_{\mathbb{P}} \neq Y_G, Y_D = b) \\ &= \sum_{c \neq a_{\mathbb{P}}} \tilde{\mathbb{P}}(\psi_n \neq c, Y_G = c, Y_D = b) - \sum_{c \neq a_{\mathbb{P}}} \tilde{\mathbb{P}}(Y_G = c, Y_D = b) \\ & \quad + \tilde{\mathbb{P}}(\psi_n(b) \neq a_{\mathbb{P}}, Y_G = a_{\mathbb{P}}, Y_D = b) \\ &= \tilde{\mathbb{P}}(\psi_n(b) \neq a_{\mathbb{P}}, Y_G = a_{\mathbb{P}}, Y_D = b) - \sum_{c \neq a_{\mathbb{P}}} \tilde{\mathbb{P}}(\psi_n(b) = c, Y_G = c, Y_D = b) \\ &= \sum_{c \neq a_{\mathbb{P}}} \tilde{\mathbb{P}}(\psi_n(b) = c, Y_G = a_{\mathbb{P}}, Y_D = b) - \sum_{c \neq a_{\mathbb{P}}} \tilde{\mathbb{P}}(\psi_n(b) = c, Y_G = c, Y_D = b) \\ &= \sum_{c \neq a_{\mathbb{P}}} \mathbb{P}(\psi_n(b) = c) \left(\mathbb{P}(Y_G = a_{\mathbb{P}}, Y_D = b) - \mathbb{P}(Y_G = c, Y_D = b) \right). \end{aligned}$$

Let define the gap $\delta_{\mathbb{P}} := \inf \{ \mathbb{P}(Y_G = a_{\mathbb{P}}, Y_D = b) - \mathbb{P}(Y_G = c, Y_D = b) : c \neq a_{\mathbb{P}} \}$. Let \mathbb{P}_0 and \mathbb{P}_1 be any two probability distributions in \mathcal{P} . We have

$$\begin{aligned} & \inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} R(\psi_n) \tag{14} \\ & \geq \inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} \delta_{\mathbb{P}} \mathbb{P}(\psi_n(X_1^n, b) \neq a_{\mathbb{P}}) \\ & \geq \inf_{\psi_n \in \Psi_n} \sup_{i \in \{0,1\}} \delta_{\mathbb{P}_i} \mathbb{P}_i(\psi_n(X_1^n, b) \neq a_{\mathbb{P}_i}) \\ & \geq \left(\inf_{i \in \{0,1\}} \delta_{\mathbb{P}_i} \right) \inf_{\psi_n \in \Psi_n} \sup_{i \in \{0,1\}} \mathbb{P}_i(\psi_n(X_1^n, b) \neq a_{\mathbb{P}_i}) \\ & \geq \left(\inf_{i \in \{0,1\}} \delta_{\mathbb{P}_i} \right) \inf_{\psi_n \in \Psi_n} \frac{\mathbb{P}_0(\psi_n(X_1^n, b) \neq a_{\mathbb{P}_0}) + \mathbb{P}_1(\psi_n(X_1^n, b) \neq a_{\mathbb{P}_1})}{2}. \tag{15} \end{aligned}$$

For $x \in \mathcal{A}^n$, let $\phi_*(x, b) = a_{\mathbb{P}_0}$ if $\mathbb{P}_0(X_1^n = x) \geq \mathbb{P}_1(X_1^n = x)$ and $\phi_*(x, b) = a_{\mathbb{P}_1}$ otherwise. It is straightforward to verify that the function ϕ_* minimizes the

righthand side of the inequality (15). Let $\mathbb{P}_0^{(n)}$ and $\mathbb{P}_1^{(n)}$ be the projection of \mathbb{P}_0 and \mathbb{P}_1 on $\{1, \dots, n\}$. From the definition of total variation distance d_{TV} , we have that

$$\frac{\mathbb{P}_0(\phi_*(X_1^n, b) \neq a_{\mathbb{P}_0}) + \mathbb{P}_1(\phi_*(X_1^n, b) \neq a_{\mathbb{P}_1})}{2} = 1 - d_{\text{TV}}(\mathbb{P}_0^{(n)}, \mathbb{P}_1^{(n)}).$$

To get a lower bound of the right hand side we use Bretagnolle-Huber inequality [23, Lemma 2.6, p. 89], that is,

$$1 - d_{\text{TV}}(\mathbb{P}_0^{(n)}, \mathbb{P}_1^{(n)}) \geq e^{-\text{KL}(\mathbb{P}_0^{(n)} \parallel \mathbb{P}_1^{(n)})},$$

where $\text{KL}(\mathbb{P}_0^{(n)} \parallel \mathbb{P}_1^{(n)})$ is the Kullback-Leibler divergence, that is,

$$\begin{aligned} \text{KL}(\mathbb{P}_0^{(n)} \parallel \mathbb{P}_1^{(n)}) &:= \sum_{x_1^n \in \mathcal{A}^n} \mathbb{P}_0^{(n)}(X_1^n = x_1^n) \log \frac{\mathbb{P}_0^{(n)}(X_1^n = x_1^n)}{\mathbb{P}_1^{(n)}(X_1^n = x_1^n)} \\ &= \sum_{x_1^n \in \mathcal{A}^n} \mathbb{P}_0^{(n)}(X_1^n = x_1^n) \sum_{k=0}^{n-1} \log \frac{\mathbb{P}_0^{(n)}(X_{n-k} = x_{n-k} \mid X_1^{n-k-1} = x_1^{n-k-1})}{\mathbb{P}_1^{(n)}(X_{n-k} = x_{n-k} \mid X_1^{n-k-1} = x_1^{n-k-1})} \\ &= \sum_{k=0}^{n-1} \mathbb{E}_{\mathbb{P}_0^{(n)}} \left[\sum_{a \in \mathcal{A}} \mathbb{P}_0^{(n)}(X_{n-k} = a \mid X_1^{n-k-1}) \log \frac{\mathbb{P}_0^{(n)}(X_{n-k} = a \mid X_1^{n-k-1})}{\mathbb{P}_1^{(n)}(X_{n-k} = a \mid X_1^{n-k-1})} \right]. \end{aligned}$$

Using Lemma C.1 (see Appendix C), we obtain

$$\text{KL}(\mathbb{P}_0^{(n)} \parallel \mathbb{P}_1^{(n)}) \leq \sum_{k=0}^{n-1} \sup_{x, y, z \in \mathcal{A}^{z-}} \sum_{a \in \mathcal{A}} \frac{(p_{\mathbb{P}_0}(a \mid x_{-k}^{-1} y) - p_{\mathbb{P}_1}(a \mid x_{-k}^{-1} z))^2}{p_{\mathbb{P}_1}(a \mid x_{-k}^{-1} z)}.$$

Putting everything together, we obtain

$$\begin{aligned} &\inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} R(\psi_n; \mathbb{P}) \\ &\geq \left(\inf_{i \in \{0, 1\}} \delta_{\mathbb{P}_i} \right) \exp \left(- \sum_{k=0}^{n-1} \sup_{x, y, z \in \mathcal{A}^{z-}} \sum_{a \in \mathcal{A}} \frac{(p_{\mathbb{P}_0}(a \mid x_{-k}^{-1} y) - p_{\mathbb{P}_1}(a \mid x_{-k}^{-1} z))^2}{p_{\mathbb{P}_1}(a \mid x_{-k}^{-1} z)} \right). \end{aligned}$$

Let index the elements of the alphabet as $\mathcal{A} = \{a_0, a_1, a_2, \dots, a_{|\mathcal{A}|}\}$ when the alphabet is finite, otherwise write $\mathcal{A} = \{a_0, a_1, a_2, \dots\}$. Choose two distinct elements $a_0, a_1 \in \mathcal{A}$. Consider that \mathbb{P}_0 and \mathbb{P}_1 are product measures with $\mathbb{P}_0(X_1 = a_0) = 4^{-1} + 1/(8\sqrt{n}) + 2^{-|\mathcal{A}|}$, $\mathbb{P}_0(X_1 = a_1) = 4^{-1} - 1/(8\sqrt{n}) + 2^{-|\mathcal{A}|}$ and, for $i \geq 2$, $\mathbb{P}_0(X_1 = a_i) = 2^{-i}$. Similarly $\mathbb{P}_1(X_1 = a_1) = 4^{-1} + 1/(8\sqrt{n}) + 2^{-|\mathcal{A}|}$, $\mathbb{P}_1(X_1 = a_0) = 4^{-1} - 1/(8\sqrt{n}) + 2^{-|\mathcal{A}|}$ and for $i \geq 2$, $\mathbb{P}_1(X_1 = a_i) = 2^{-i}$. When $|\mathcal{A}| = \infty$, we have $2^{-|\mathcal{A}|} = 0$.

With these choices, the supremum in the argument of the exponential simplifies to

$$\sum_{a \in \mathcal{A}} \frac{(p_{\mathbb{P}_0}(a) - p_{\mathbb{P}_1}(a))^2}{p_{\mathbb{P}_1}(a)} \leq \frac{1}{n},$$

and

$$\delta_{\mathbb{P}_0} = \delta_{\mathbb{P}_1} = \inf_{\substack{c \in \mathcal{A} \\ c \neq a}} [\mathbb{P}_0(Y_G = (a_0, a_0, \dots, a_0)) - \mathbb{P}_0(Y_G = (c, a_0, \dots, a_0))] \mathbb{P}_1(Y_D = b).$$

Choosing $b = (a_0, \dots, a_0) \in A^D$, we obtain

$$\delta_{\mathbb{P}_0} \geq \frac{1}{\sqrt{n}} \left(\frac{1}{2} + \frac{1}{2\sqrt{n}} \right)^{|D|+|G|-1} \geq \frac{1}{\sqrt{n}} \left(\frac{1}{2} \right)^{|D|+|G|-1},$$

and we conclude that

$$\inf_{\psi_n \in \Psi_n} \sup_{\mathbb{P} \in \mathcal{P}} R(\psi_n; \mathbb{P}) \geq \frac{1}{\sqrt{n}} \left(\frac{1}{2} \right)^{|D|+|G|-1} e^{-1},$$

which ends the proof of the theorem.

A Dvoretzky-Kiefer-Wolfowitz type inequality

The following theorem is a restatement of results of [1, Theorem 4.1] to better fit the need for this paper. The proof is included for the convenience of the reader. This result is a variant of what is commonly referred to in the literature as the Dvoretzky-Kiefer-Wolfowitz inequality, specifically for the case of independent random variables.

Let $S \subseteq \mathbb{Z}$ and $\sigma \in \mathcal{A}^S$ with $\text{diam}(S) := \sup S - \inf S = k (< \infty)$. For any $n \geq k$, let us denote the frequency of occurrences of the string σ in X_1^n by

$$\hat{r}_S^n(\sigma) := \frac{N_S^n(\sigma)}{n - k + 2} := \frac{\sum_{i=0}^{n-k+1} \mathbb{1}\{X_{S+i} = \sigma\}}{n - k + 2}.$$

THEOREM A.1. Let $(X_j)_{j \in \mathbb{Z}}$ be a stationary process such that (A) holds. Then, for all $u > 0$, for all $n > 0$, and $0 < k \leq n$, we have

$$\begin{aligned} & \mu \left(\sup_{a \in \mathcal{A}^S} |\hat{r}_S^n(a) - \mathbb{P}(X_S = a)| > u + \sqrt{\frac{2|S|(1 - \Gamma(p)) + \Gamma(p)}{\Gamma(p)(n - k + 2)}} \right) \\ & \leq \exp \left(-\frac{2(n - k + 2)u^2}{|S|^2 \Gamma(p)^{-2}} \right). \end{aligned}$$

PROOF. Define the statistic $f_S = \|\hat{r}_S^n(\cdot) - \mathbb{P}(X_S \in \cdot)\|_\infty$. Without loss of generality, we assume $\inf S = 1$. By the Gaussian concentration bound in [1, Theorem 3.2], we have

$$\mathbb{P}(f_S - \mathbb{E}[f_S] > u) \leq \exp(-2|S|^{-2}(n - k + 2)\Gamma(p)^2 u^2). \quad (16)$$

Therefore, to prove Theorem A.1, it remains to find a good upper bound for $\mathbb{E}[f_S]$. Here, we follow the argument used in [17]. By Jensen's inequality, and since $\mathbb{E}[\hat{r}_S^n(\sigma)] = \mathbb{P}(X_S = \sigma)$, we have

$$\begin{aligned} (\mathbb{E}[f_S])^2 &\leq \mathbb{E}[f_S^2] \leq \mathbb{E} \left[\sum_{\sigma \in \mathcal{A}^S} (\hat{r}_S^n(\sigma) - \mathbb{P}(X_S = \sigma))^2 \right] \\ &\leq \sum_{\sigma \in \mathcal{A}^S} \left(\mathbb{E}[\hat{r}_S^n(\sigma)^2] - \mathbb{P}(X_S = \sigma)^2 \right). \end{aligned} \quad (17)$$

For all $\sigma \in \mathcal{A}^S$, we have

$$\begin{aligned} &\mathbb{E}[\hat{r}_S^n(\sigma)^2] \\ &= \frac{1}{(n-k+2)^2} \mathbb{E} \left[\left(\sum_{i=0}^{n-k+1} \mathbb{1}\{X_{S+i} = \sigma\} \right)^2 \right] \\ &= \frac{1}{(n-k+2)^2} \mathbb{E} \left[\sum_{i=0}^{n-k+1} \mathbb{1}\{X_{S+i} = \sigma\} \right] \\ &\quad + \frac{2}{(n-k+2)^2} \mathbb{E} \left[\sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} \mathbb{1}\{X_{S+i} = \sigma\} \mathbb{1}\{X_{S+j} = \sigma\} \right]. \end{aligned}$$

Hence by stationarity we get

$$\begin{aligned} &\mathbb{E}[\hat{r}_S^n(\sigma)^2] \\ &= \frac{\mathbb{P}(X_S = \sigma)}{n-k+2} + \frac{2}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} \mathbb{P}(X_{S+i} = \sigma, X_{S+j} = \sigma) \\ &= \frac{\mathbb{P}(X_S = \sigma)}{n-k+2} + \frac{2}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} \mathbb{P}(X_S = \sigma) \mathbb{P}(X_{S+j} = \sigma \mid X_{S+i} = \sigma) \\ &\leq \frac{\mathbb{P}(X_S = \sigma)}{n-k+2} + \frac{2}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} \left[\mathbb{P}(X_S = \sigma) \times \right. \\ &\quad \left. (\mathbb{P}(X_S = \sigma) + |\mathbb{P}(X_{S+j} = \sigma \mid X_{S+i} = \sigma) - \mathbb{P}(X_S = \sigma)|) \right]. \end{aligned}$$

Now, for all $\sigma \in \mathcal{A}^S$, we have

$$\begin{aligned}
& \left| \mathbb{P}(X_{S+j} = \sigma \mid X_{S+i} = \sigma) - \mathbb{P}(X_S = \sigma) \right| \\
& \leq \sup_{\tilde{\sigma} \in \mathcal{A}^S} \left| \mathbb{P}(X_{S+j} = \sigma \mid X_{S+i} = \sigma) - \mathbb{P}(X_{S+j} = \sigma \mid X_{S+i} = \tilde{\sigma}) \right| \\
& = \sup_{\tilde{\sigma} \in \mathcal{A}^S} \left| \mathbb{P}(X_{S+j-i} = \sigma \mid X_S = \sigma) - \mathbb{P}(X_{S+j-i} = \sigma \mid X_S = \tilde{\sigma}) \right| \\
& \leq \sup_{x, y \in \mathcal{A}^{\mathbb{Z}^-}} \left| \mathbb{P}^x(X_{S+j-i} = \sigma) - \mathbb{P}^y(X_{S+j-i} = \sigma) \right| \\
& \leq \sup_{x, y \in \mathcal{A}^{\mathbb{Z}^-}} \sum_{\ell \in S+j-i} \mathbb{P}^{x, y}(\eta_\ell \neq \omega_\ell) =: q_{j-i}(S),
\end{aligned}$$

where $\mathbb{P}^{x, y}$ is the one-step maximal coupling between \mathbb{P}^x and \mathbb{P}^y , which are the laws of the process $(X_i)_{i \in \mathbb{Z}}$ conditioned on starting with pasts x and y , respectively. Coming back to the estimation of $\mathbb{E}[\hat{r}_S^n(\sigma)^2]$, we have

$$\begin{aligned}
& \mathbb{E}[\hat{r}_S^n(\sigma)^2] - \frac{\mathbb{P}(X_S = \sigma)}{n-k+2} \\
& \leq \frac{2}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} \mathbb{P}(X_S = \sigma) (\mathbb{P}(X_S = \sigma) + q_{j-i}(S)) \\
& \leq \mathbb{P}(X_S = \sigma)^2 + \frac{2\mathbb{P}(X_S = \sigma)}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i=0}^{j-1} q_{j-i}(S) \\
& \leq \mathbb{P}(X_S = \sigma)^2 + \frac{2\mathbb{P}(X_S = \sigma)}{(n-k+2)^2} \sum_{j=1}^{n-k+1} \sum_{i \geq 1} q_i(S) \\
& \leq \mathbb{P}(X_S = \sigma)^2 + \frac{2\mathbb{P}(X_S = \sigma)}{n-k+2} \sum_{i \geq 1} q_i(S).
\end{aligned}$$

On the other hand,

$$\sum_{i \geq 1} q_i(S) = \sum_{i \geq 1} \sup_{x, y \in \mathcal{A}^{\mathbb{Z}^-}} \sum_{\ell \in S+i} \mathbb{P}^{x, y}(\eta_\ell \neq \omega_\ell) \leq |S| \sum_{i \geq 1} \sup_{x, y \in \mathcal{A}^{\mathbb{Z}^-}} \mathbb{P}^{x, y}(\eta_\ell \neq \omega_\ell).$$

So we only need now an upper bound for the coupling errors. Proposition 5.2 in [1] gives us that

$$\sum_{i=1}^{\infty} \sup_{x, y \in \mathcal{A}^{\mathbb{Z}^-}} \mathbb{P}^{x, y}(\eta_i \neq \omega_i) \leq \frac{1 - \Gamma(p)}{\Gamma(p)}.$$

Hence, we have that

$$\mathbb{E}[\hat{r}_S^n(\sigma)^2] - \mathbb{P}(X_S = \sigma)^2 \leq \frac{2|S|(1 - \Gamma(p))\Gamma(p)^{-1} + 1}{n-k+2} \mathbb{P}(X_S = \sigma). \quad (18)$$

Using (17) we obtain

$$\mathbb{E}[\|\hat{r}_S^n - \mathbb{P}\|_\infty] \leq \sqrt{\frac{2|S|(1 - \Gamma(p))\Gamma(p)^{-1} + 1}{n-k+2}},$$

which is the desired bound. \square

B One-dimensional Gibbs measures

The goal of this section is to establish Theorem B.1 below, from which Proposition 4.1 follows as a direct corollary. To this end, we return to the framework introduced in the example of Gibbs measures in Section 4. Let $(X_j)_{j \in \mathbb{Z}}$ be a Gibbs measure associated with the potential Φ , and denote by p_Φ the regular version of its left conditional expectations, as defined in (3).

Observe that by (8), the conditional probabilities (9) are uniformly bounded away from zero, that is

$$h_\Lambda := \inf_{x,y} \mathbb{P}(X_\Lambda = x_\Lambda | X_{\Lambda^c} = y_{\Lambda^c}) > 0, \quad (19)$$

uniformly in Λ , and in particular, $h_{\{0\}} > 0$.

The next theorem B.1 establishes a relationship between the variation of p_Φ and $\Delta(\Phi_\Lambda) := \max \Phi_\Lambda - \min \Phi_\Lambda$ (oscillation of Φ_Λ).

THEOREM B.1. We have $\text{Var}_0(p_\Phi) \leq 1 - h_{\{0\}}$ and

$$\text{Var}_k(p_\Phi) \leq \frac{|\mathcal{A}|}{2} \sum_{i \geq k} \sum_{\substack{\min \Lambda=0 \\ \max \Lambda \geq \frac{i}{2}}} \Delta(\Phi_\Lambda), \quad k \geq 1. \quad (20)$$

Proposition 4.1 follows directly as a corollary of this result, as $\text{Var}_k(p_\Phi)$ is a non-increasing sequence. Moreover, for a $(0, 1)$ -valued sequence of real numbers $(a_n)_{n \in \mathbb{N}}$, it holds that $\sum_n a_n < \infty \iff \prod_n (1 - a_n) > 0$. Furthermore, defining

$$n_\star := \inf \left\{ n \geq 0 : \frac{|\mathcal{A}|}{2} \sum_{i \geq n} \sum_{\substack{\min \Lambda=0 \\ \max \Lambda \geq \frac{i}{2}}} \Delta(\Phi_\Lambda) < 1 \right\},$$

we also see that

$$\Gamma(p_\Phi) \geq \prod_{n \geq n_\star} \left(1 - \frac{|\mathcal{A}|}{2} \sum_{i \geq n} \sum_{\substack{\min \Lambda=0 \\ \max \Lambda \geq \frac{i}{2}}} \Delta(\Phi_\Lambda) \right), \quad (21)$$

which, in principle, makes it possible to derive explicit bounds on $\Gamma(p_\Phi)$.

Before delving into the proof of this theorem, let us offer a brief prelude. For a given function ϕ , defined either on $\mathcal{A}^{\mathbb{Z}}$ or $\mathcal{A}^{\mathbb{Z}^-} \times \mathcal{A}$, an *equilibrium state* for ϕ is a measure μ_ϕ on $\mathcal{A}^{\mathbb{Z}}$, shift-invariant (that is $\mu_\phi \circ T^{-1} = \mu_\phi$ where T is the left shift) which maximizes $\int \phi d\mu + h_\mu(T)$ where $h_\mu(T)$ denotes the entropy of μ . The function ϕ is called two-sided (*resp.* one-sided) potential function if it is

defined on $\mathcal{A}^{\mathbb{Z}}$ (*resp.* on $\mathcal{A}^{\mathbb{Z}^-} \times \mathcal{A}$). The regularity of a potential is measured by its “variation” (modulus of continuity with respect to the usual distance). For $n \geq 0$, let

$$\text{var}_n(\phi) := \sup \{ |\phi(x) - \phi(y)| : x, y \in \mathcal{A}^{\mathbb{Z}}, x_i = y_i, |i| \leq n \},$$

for a two-sided potential ϕ , and

$$\text{var}_n(\varphi) := \sup \{ |\varphi(xa) - \varphi(ya)| : x, y \in \mathcal{A}^{\mathbb{Z}^-}, a \in \mathcal{A}, x_i = y_i, i \geq -n \},$$

for a one-sided potential φ .

Theorem B.1 establishes a connection between the variation $\Delta(\Phi)$ of the potential in a Gibbs (two-sided) specification and the variation $\text{Var}_n(p)$ of the one-sided conditional expectations. In essence, it links two distinct methods of specifying stationary processes: the two-sided conditioning used in Gibbs measures and the one-sided conditioning employed in stochastic processes. The proof strategy leverages equilibrium states from dynamical systems, offering a third perspective on defining stationary processes and serving as a bridge between the two conditioning frameworks.

PROOF OF THEOREM B.1. Denote by μ the law of the process $(X_j)_{j \in \mathbb{Z}}$, that is, the measure which is defined through $\mu(C) = \mathbb{P}(X_{-\infty}^{+\infty} \in C)$ for any measurable set C . Since $(X_j)_{j \in \mathbb{Z}}$ is stationary, μ is invariant under the shift ($\mu(C) = \mu(T(C))$ for any measurable set C). We know by assumption that $(X_j)_{j \in \mathbb{Z}}$ (or equivalently μ) satisfies (9). It is proven in [3, Section 3] that a two-sided potential ϕ can be constructed from Φ , which is such that μ is an equilibrium state for ϕ and moreover

$$\text{var}_n(\phi) \leq \sum_{\substack{\min \Lambda = 0 \\ \max \Lambda \geq n}} \Delta(\Phi_\Lambda). \quad (22)$$

We next use [21, Section 5] which allows us to construct a one-sided potential φ out of the two-sided potential ϕ , in such a way that μ is an equilibrium state for φ and moreover

$$\text{var}_n(\varphi) \leq \text{var}_{\frac{n}{2}}(\phi). \quad (23)$$

We now invoke [18, Section IV] (see also [24, Proof of Theorem 3.3]), which constructs a one-sided normalized function g (normalized means that it satisfies $\sum_{a \in \mathcal{A}} g(x_{-\infty}^{-1} a) = 1$, for any $x_{-\infty}^{-1}$) such that $\inf g > 0$, and μ is an equilibrium state for $\log g$. Moreover

$$\text{var}_n(\log g) \leq \sum_{k \geq n} \text{var}_k(\varphi). \quad (24)$$

Now, by the mean value theorem applied to the exponential function, we have for all n , $x, y \in \mathcal{A}^{\mathbb{Z}^-}$, $a \in \mathcal{A}$, such that $x_i = y_i, i \geq -n$,

$$|g(xa) - g(ya)| = |e^{\log g(xa)} - e^{\log g(ya)}| \leq |\log g(xa) - \log g(ya)|,$$

where we used that $0 < g < 1$. Thus

$$\text{var}_n(g) \leq \text{var}_n(\log g), \quad n \geq 1. \quad (25)$$

To conclude, [18] also tells us that μ being an equilibrium state for $\log g$ is equivalent to saying that the stationary process of the corresponding equilibrium state satisfies (3) with $g(xa)$ in place of $p(a|x)$. But by the a.s. uniqueness of the regular version, we conclude that $g(xa) = p_\Phi(a|x)$ \mathbb{P} -a.s., and thus $\text{var}_n(g) = \text{var}_n(p_\Phi)$.

Combining (22), (23), (24), and (25), and noting that $\text{Var}_n(p) \leq \frac{|A|}{2} \text{var}_n(p)$, we thus proved (20).

It only remains to prove that $\text{Var}_0(p_\Phi) \leq 1 - h_{\{0\}}$. We claim that $1 - \text{Var}_0(p_\Phi) = \inf_{x,y \in \mathcal{A}^{\mathbb{Z}^-}} \sum_{a \in \mathcal{A}} p_\Phi(a|x) \wedge p_\Phi(a|y) \geq \inf_{a \in \mathcal{A}, x \in \mathcal{A}^{\mathbb{Z}^-}} p_\Phi(a|x) \geq h_{\{0\}}$. Indeed, for any $x \in \mathcal{A}^{\mathbb{Z}^-}$,

$$\begin{aligned} \inf_{a \in \mathcal{A}, x \in \mathcal{A}^{\mathbb{Z}^-}} p_\Phi(a|x) &= \inf_{a \in \mathcal{A}, x \in \mathcal{A}^{\mathbb{Z}^-}} \int_{\mathcal{A}^{\mathbb{N}}} \frac{\exp\left(-H_{\{0\}}^\Phi(xay)\right)}{Z_\Lambda^\Phi((xay)_{\mathbb{Z} \setminus \{0\}})} d\mathbb{P}(y) \\ &\geq \int_{\mathcal{A}^{\mathbb{N}}} \inf_{a \in \mathcal{A}, x \in \mathcal{A}^{\mathbb{Z}^-}} \frac{\exp\left(-H_{\{0\}}^\Phi(xay)\right)}{Z_\Lambda^\Phi((xay)_{\mathbb{Z} \setminus \{0\}})} d\mathbb{P}(y) \\ &\geq \int_{\mathcal{A}^{\mathbb{N}}} \inf_{xay \in \mathcal{A}^{\mathbb{Z}}} \frac{\exp\left(-H_{\{0\}}^\Phi(xay)\right)}{Z_\Lambda^\Phi((xay)_{\mathbb{Z} \setminus \{0\}})} d\mathbb{P}(y) \\ &\geq \inf_{xay \in \mathcal{A}^{\mathbb{Z}}} \frac{\exp\left(-H_{\{0\}}^\Phi(xay)\right)}{Z_\Lambda^\Phi(y)} = h_{\{0\}}. \end{aligned}$$

The theorem is proved. \square

C A lemma

The following lemma is a classical upper bound for the Kullback-Leibler divergence using chi-square distance.

LEMMA C.1. Let \mathbb{P} and \mathbb{Q} be two probability measures on a countable alphabet \mathcal{A} . Then

$$\sum_{a \in \mathcal{A}} \mathbb{P}(X = a) \log \frac{\mathbb{P}(X = a)}{\mathbb{Q}(X = a)} \leq \sum_{a \in \mathcal{A}} \frac{(\mathbb{P}(X = a) - \mathbb{Q}(X = a))^2}{\mathbb{Q}(X = a)},$$

where we define $\log 0/0 = 0$ for the left hand side and $0/0 = 0$ for the right hand side.

PROOF. Using the concavity of the logarithm, then adding and subtracting $\mathbb{Q}(X = a)$ and expanding the square, and finally using the inequality $\log(1+u) \leq$

u for $u \geq 0$, we get

$$\begin{aligned} & \sum_{a \in \mathcal{A}} \mathbb{P}(X = a) \log \frac{\mathbb{P}(X = a)}{\mathbb{Q}(X = a)} \leq \log \left(\sum_{a \in \mathcal{A}} \frac{\mathbb{P}(X = a)^2}{\mathbb{Q}(X = a)} \right) \\ & = \log \left(1 + \sum_{a \in \mathcal{A}} \frac{(\mathbb{P}(X = a) - \mathbb{Q}(X = a))^2}{\mathbb{Q}(X = a)} \right) \\ & \leq \sum_{a \in \mathcal{A}} \frac{(\mathbb{P}(X = a) - \mathbb{Q}(X = a))^2}{\mathbb{Q}(X = a)}, \end{aligned}$$

which proves the lemma. \square

Acknowledgements. The authors acknowledge FAPESP (Regular Research Grants 2019/23439-4) for support. D. Y. T. and S. G. gratefully acknowledge École Polytechnique for supporting their visits to CPHT, funding a one-month stay in 2022 and another in 2024.

References

- [1] J.-R. Chazottes, S. Gallo, and D. Y. Takahashi. Gaussian concentration bounds for stochastic chains of unbounded memory. *The Annals of Applied Probability*, 33(5):3321–3350, 2023.
- [2] J.-R. Chazottes and E. Ugalde. On the preservation of gibbsianness under symbol amalgamation. In Brian Marcus, Karl Petersen, and Tsachy Weissman, editors, *Entropy of Hidden Markov Processes and Connections to Dynamical Systems: Papers from the Banff International Research Station Workshop*, London Mathematical Society Lecture Note Series, pages 72–97. Cambridge University Press, 2011.
- [3] Z. Coelho and A. Quas. Criteria for \bar{d} -continuity. *Transactions of the American Mathematical Society*, 350(8):3257–3268, 1998.
- [4] F. Comets, R. Fernández, and P. A. Ferrari. Processes with long memory: regenerative construction and perfect simulation. *The Annals of Applied Probability*, 12(3):921–943, 2002.
- [5] W. De Carvalho, S. Gallo, and N. L. Garcia. Continuity properties of a factor of Markov chains. *Journal of Applied Probability*, 53(1):216–230, 2016.
- [6] R. L. Dobrushin. Central limit theorem for nonstationary Markov chains. I. *Theory of Probability & Its Applications*, 1(1):65–80, 1956.
- [7] R. Douc, E. Moulines, P. Priouret, and Ph. Soulier. *Markov chains*. Springer, 2018.

- [8] R. Fernández and G. Maillard. Chains with complete connections: general theory, uniqueness, loss of memory and mixing properties. *Journal of Statistical Physics*, 118(3-4):555–588, 2005.
- [9] K. Fokianos. *Categorical time series: Prediction and control*. University of Maryland, College Park, 1996.
- [10] K. Fokianos and L. Truquet. On categorical time series models with covariates. *Stochastic processes and their applications*, 129(9):3446–3462, 2019.
- [11] H.-O. Georgii. *Gibbs measures and phase transitions*, volume 9. Walter de Gruyter, 2011.
- [12] N. Gruver, M. Finzi, S. Qiu, and A. G. Wilson. Large language models are zero-shot time series forecasters. *Advances in Neural Information Processing Systems*, 36, 2024.
- [13] Y. Han, S. Jana, and Y. Wu. Optimal prediction of Markov chains with and without spectral gap. *IEEE Transactions on Information Theory*, 69(6):3920–3959, 2023.
- [14] Y. Hao, A. Orłitsky, and V. Pichapati. On learning Markov chains. *Advances in Neural Information Processing Systems*, 31, 2018.
- [15] S. Kalikow. Random Markov processes and uniform martingales. *Israel Journal of Mathematics*, 71(1):33–54, 1990.
- [16] B. Kedem and K. Fokianos. *Regression models for time series analysis*, volume 488. John Wiley & Sons, 2005.
- [17] A. Kontorovich and R. Weiss. Uniform Chernoff and Dvoretzky-Kiefer-Wolfowitz-type inequalities for Markov chains and related processes. *Journal of Applied Probability*, 51(4):1100–1113, 2014.
- [18] F. Ledrappier. Principe variationnel et systèmes dynamiques symboliques. *Probability Theory and Related Fields*, 30(3):185–202, 1974.
- [19] G. Morvai and B. Weiss. On universal algorithms for classifying and predicting stationary processes. *Probability Surveys*, 18:77 – 131, 2021.
- [20] M. Piraino. Single site factors of Gibbs measures. *Nonlinearity*, 33(2):742–761, 2020.
- [21] M. Pollicott. Rates of mixing for potentials of summable variation. *Transactions of the American Mathematical Society*, 352(2):843–853, 2000.
- [22] F. Redig and F. Wang. Transformations of one-dimensional gibbs measures with infinite range interaction. *Markov Processes and Related Fields*, 16(1):737–752, 2010.

- [23] A.B. Tsybakov. *Introduction to Nonparametric Estimation*. Springer Series in Statistics. Springer New York, 2009.
- [24] P. Walters. Ruelle’s operator theorem and g -measures. *Transactions of the American Mathematical Society*, 214:375–387, 1975.
- [25] G. Wolfer and A. Kontorovich. Minimax learning of ergodic Markov chains. In *Algorithmic Learning Theory*, pages 904–930. PMLR, 2019.
- [26] Bin Yu and Fano Assouad. Le cam. *Festschrift for Lucien Le Cam: Research Papers in Probability and Statistics*, Springer-Verlag, New York, pages 423–435, 1997.