

PT-Symmetric $SU(2)$ - like Random Matrix Ensembles: Invariant Distributions and Spectral Fluctuations

Stalin Abraham^{1,2†}, A. Bhagwat^{1,2*†}, Sudhir Ranjan Jain^{1†}

¹*School of Physical Sciences, UM-DAE Centre for Excellence in Basic Sciences, University of Mumbai, Vidyanagari Campus, Santacruz East, Mumbai, 400098, Maharashtra, India.

²Centre for Excellence in Theoretical and Computational Sciences, University of Mumbai, Vidyanagari Campus, Santacruz East, Mumbai, 400098, Maharashtra, India.

*Corresponding author(s). E-mail(s): ameeya@cbs.ac.in;

Contributing authors: stalin.abraham@cbs.ac.in; srjain@cbs.ac.in;

[†]These authors contributed equally to this work.

Abstract

We consider an ensemble of 2×2 normal matrices with complex entries representing operators in the quantum mechanics of 2 - level parity-time reversal (PT) symmetric systems. The randomness of the ensemble is endowed by obtaining probability distributions based on symmetry and statistical independence. The probability densities turn out to be power law with exponents that depend on the boundedness of the domain. For small spacings, σ , the probability density varies as σ^ν , $\nu \geq 2$. The degree of level repulsion is a parameter of great interest as it makes a connection to quantum chaos; the lower bound of ν for our ensemble coincides with the Gaussian Unitary Ensemble. We believe that the systematic development presented here paves the way for further generalizations in the field of random matrix theory for PT-symmetric quantum systems.

Keywords: Random Matrix Theory, PT-Symmetric Matrices, Probability Distribution Functions, Level Repulsion

MSC Classification: 15B52 , 15B57 , 60B20

1 Introduction

Symmetries guide mathematical classification of physical systems [1]. Among them, discrete symmetries like time reversal and parity play a special role. One of these appears in the statistical treatment of quantum systems with complex spectra, often mathematically treated by random matrix theory [2, 3]. For systems where parity and time-reversal are broken (PT -symmetric), random matrix theory was introduced [4, 5]. These works introduced the concept of pseudo-unitary symmetry in the context of random matrices and proved the existence of a new universality class with novel level repulsion. The physical situations related to these random matrix models are diverse and most interesting. Historically, these models appeared in the context of quantum field theory with an indefinite metric [6] and in the electromagnetic theory of transmission lines [7]. By introducing an indefinite metric, a general criterion for a set of non-Hermitian operators was established [8], consistent with the conventional interpretation of quantum mechanics [9]. The momentum distribution of an ideal gas of identical particles in two dimensions is an open problem. One of the ways to obtain this was shown by employing the eigenstate ensemble exploiting quantum chaos [10, 11] where the second virial coefficient for such gas was expressed in terms of a counting problem in the theory of Braid groups. Such systems are related on the one hand with Nambu's [12] proposal of quark confinement in the medium of monopoles and with Aharonov-Bohm type quantum billiards which are pseudo-integrable [13]. The latter work paved the way towards a new universality class of random matrices for explaining the spectral fluctuations of pseudo-integrable and almost integrable quantum billiards [14]. Subsequently, these results have been generalized for $N \times N$ random cyclic matrices [15, 16]. Yet another fascinating connection is made with the exactly solvable models of many-particle systems. The joint probability distribution function of the eigenvalues of several random matrix ensembles coincides with the joint probability distribution function obtained from the many-body ground state [17, 18]. A similar connection for random cyclic matrices exists [19] where the connection was made with a screened harmonic oscillator. By a further transformation employing the normal coordinates, eigenfunctions of integrable [18] and chaotic billiards [20] have been obtained.

Here we re-visit the random matrix theory for non-Hermitian matrices. There has been a lot of work on random matrices with complex elements in the last sixty years [21–23]. In all these works, the real and imaginary parts of the matrix elements were drawn from a Gaussian distribution. In the celebrated work pioneered by Wigner, Mehta, Gaudin, Dyson and others, this distribution was derived for the matrix ensembles based on time-reversal invariance, rotational symmetry, and statistical independence [2]. However, for the case of non-Hermitian matrices, the distribution has been assumed to be Gaussian. As one of the main results, we present here a symmetry-based argument for deriving a probability distribution for the ensemble of non-Hermitian matrices considered here. Subsequently, we calculate the fluctuation properties of the eigenvalues and obtain new results. This leads to a degree of level repulsion, which is shown to be greater than or equal to two. It may be recalled that the degree of level repulsion is qualitatively associated with quantum chaos. The degree of level repulsion measures the rate at which the two eigenvalues go away from

each other in the limit of small spacing. For the invariant Gaussian ensembles, the degree of repulsion is respectively 1, 2, and 4 for Orthogonal, Unitary, and Symplectic ensembles [2]. It was found to be stronger for certain pseudo-Hermitian ensembles some twenty years ago [4, 5].

Specifically, in this work, we focus on the 2×2 parity and time-reversal symmetric (PT-symmetric) complex matrices that are non-Hermitian but invertible and normal. It is worth pointing to a recent review [24] of the subject of PT-symmetric quantum mechanics, our effort falls very much into the physics and mathematics of the theme. These are interesting due to the fact that they can have real or complex eigenvalues and are used to understand, non - dissipative and dissipative systems, respectively (see, for example, [25–30]).

2 Random matrix ensemble

We consider an ensemble of normal 2×2 matrices with complex entries expressed as

$$H = \begin{pmatrix} z_1 & z_2 \\ -\bar{z}_2 & \bar{z}_1 \end{pmatrix} \quad (1)$$

with z_1 and z_2 are complex numbers, $z_1 = (x_1, y_1)$, and $z_2 = (x_2, y_2)$. These matrices are complex normal matrices and hence are unitarily diagonalisable. Further, they satisfy $H^\dagger H = HH^\dagger = \det(H)I_2$ where, I_2 is an identity matrix. Notice that $\det(H) = z_1\bar{z}_1 + z_2\bar{z}_2 \neq 0$, unless $z_1 = z_2 = 0$. If $\det(H) = 1$, these would have been $SU(2)$ matrices. With this motivation, the matrices considered in this analysis are called $SU(2)$ -like matrices.

Suppose that $\epsilon \in \mathbb{R}$ is a very small number. We demand invariance of probability distribution under similarity transformations. Let us write an infinitesimal transformation,

$$Q = \begin{pmatrix} 1 & -\epsilon \\ \epsilon & 1 \end{pmatrix} \quad (2)$$

under which, H transforms as

$$Q^{-1}HQ = \begin{pmatrix} z_1 + \epsilon(z_2 - \bar{z}_2) & z_2 + \epsilon(\bar{z}_1 - z_1) \\ -z_2 + \epsilon(\bar{z}_1 - z_1) & \bar{z}_1 + \epsilon(\bar{z}_2 - z_2) \end{pmatrix} \quad (3)$$

2.1 Construction of Probability Distribution Function

Let $P(z)$ denote the probability distribution function of the parameters defining the matrix elements. The invariance of $P(z)$ under similarity transformation implies that

$$P(z_1 + \epsilon(z_2 - \bar{z}_2))P(z_2 + \epsilon(\bar{z}_1 - z_1)) = P(z_1)P(z_2) \quad (4)$$

We now make two assumptions: 1) $P(z) = f(x^2 + y^2) = f_1(x^2)f_2(y^2)$ with $z \in \mathbb{C}$ and $z = (x, y)$ and f_1, f_2 sufficiently smooth, and 2) $P(z_1z_2) = P(z_1)P(z_2)$. With

these assumptions, we get:

$$P(z_1 + \epsilon(z_2 - \bar{z}_2)) = P(x_1 + iy_1 + i2\epsilon y_2) = f_1(x_1^2 + y_1^2 + 4\epsilon y_1 y_2) \quad (5)$$

$$P(z_2 + \epsilon(\bar{z}_1 - z_1)) = P(x_2 + iy_2 - i2\epsilon y_1) = f_2(x_2^2 + y_2^2 - 4\epsilon y_1 y_2) \quad (6)$$

with

$$f_1(x_1^2 + y_1^2 + 4\epsilon y_1 y_2) = f_1(x_1^2 + y_1^2) + 8\epsilon y_1 y_2 \left(x_1 \frac{\partial f_1}{\partial x_1} + y_1 \frac{\partial f_1}{\partial y_1} \right) + o(\epsilon^2) \quad (7)$$

$$f_2(x_2^2 + y_2^2 - 4\epsilon y_1 y_2) = f_2(x_2^2 + y_2^2) - 8\epsilon y_1 y_2 \left(x_2 \frac{\partial f_2}{\partial x_2} + y_2 \frac{\partial f_2}{\partial y_2} \right) + o(\epsilon^2) \quad (8)$$

Invariance of probability distribution function under similarity transformation requires that:

$$f_1(x_1^2 + y_1^2 + 4\epsilon y_1 y_2) f_2(x_2^2 + y_2^2 - 4\epsilon y_1 y_2) = f_1(x_1^2 + y_1^2) f_2(x_2^2 + y_2^2) \quad (9)$$

giving us

$$f_2(x_2^2 + y_2^2) \left(x_1 \frac{\partial f_1}{\partial x_1} + y_1 \frac{\partial f_1}{\partial y_1} \right) = f_1(x_1^2 + y_1^2) \left(x_2 \frac{\partial f_2}{\partial x_2} + y_2 \frac{\partial f_2}{\partial y_2} \right). \quad (10)$$

Let us separate $f_{1,2}$ in x - and y - parts:

$$f_1(x_1^2 + y_1^2) = F_{11}(x_1^2) F_{12}(y_1^2) \quad (11)$$

$$f_2(x_2^2 + y_2^2) = F_{21}(x_2^2) F_{22}(y_2^2) \quad (12)$$

Combining Eqs. (10,12), we get:

$$F_{21} F_{22} \left[x_1 \frac{dF_{11}}{dx_1} F_{12} + y_1 \frac{dF_{12}}{dy_1} F_{11} \right] = F_{11} F_{12} \left[x_2 \frac{dF_{21}}{dx_2} F_{22} + y_2 \frac{dF_{22}}{dy_2} F_{21} \right] \quad (13)$$

This leads to:

$$\frac{x_1}{F_{11}} \frac{dF_{11}}{dx_1} + \frac{y_1}{F_{12}} \frac{dF_{12}}{dy_1} = \frac{x_2}{F_{21}} \frac{dF_{21}}{dx_2} + \frac{y_2}{F_{22}} \frac{dF_{22}}{dy_2} \quad (14)$$

Using the separation of variables,

$$\frac{x_1}{F_{11}} \frac{dF_{11}}{dx_1} + \frac{y_1}{F_{12}} \frac{dF_{12}}{dy_1} - \frac{x_2}{F_{21}} \frac{dF_{21}}{dx_2} = \frac{y_2}{F_{22}} \frac{dF_{22}}{dy_2} = \lambda_0 \in \mathbb{R} \quad (15)$$

where λ_0 is a separation constant. This leads directly to

$$F_{22}(y_2) = y_2^{\lambda_0} \quad (16)$$

Further,

$$\frac{x_1}{F_{11}} \frac{dF_{11}}{dx_1} + \frac{y_1}{F_{12}} \frac{dF_{12}}{dy_1} = \lambda_0 + \frac{x_2}{F_{21}} \frac{dF_{21}}{dx_2} = \lambda_2 \in \mathbb{R} \quad (17)$$

leading to:

$$F_{21}(x_2) = x_2^{\lambda_2 - \lambda_0}. \quad (18)$$

Also,

$$\frac{x_1}{F_{11}} \frac{dF_{11}}{dx_1} = \lambda_2 - \frac{y_1}{F_{12}} \frac{dF_{12}}{dy_1} = \lambda_3 \in \mathbb{R}. \quad (19)$$

Finally,

$$F_{11}(x_1) = x_1^{\lambda_3} \quad \text{and} \quad F_{12}(y_1) = y_1^{\lambda_2 - \lambda_3} \quad (20)$$

In summary, we get the probability distribution function:

$$P(z_1, z_2) = P(x_1, y_1, x_2, y_2) = Ax_1^{\lambda_3} y_1^{\lambda_2 - \lambda_3} x_2^{\lambda_2 - \lambda_0} y_2^{\lambda_0} \quad (21)$$

2.2 Determination of Normalisation Constant

Let $(x_1, y_1, x_2, y_2) \in \mathbb{R}^4$. Take the weight function $w : \mathbb{C} \times \mathbb{C} \rightarrow \mathbb{R}$ which is given by

$$w(z_1, z_2) = w(x_1, y_1, x_2, y_2) = \exp \left\{ -\alpha \left(|z_1|^2 + |z_2|^2 \right) \right\} \quad (22)$$

$$= \exp \left\{ -\alpha \left(x_1^2 + y_1^2 + x_2^2 + y_2^2 \right) \right\} \quad (23)$$

with $\alpha > 0$. We demand that

$$\int_{\mathbb{R}^4} P(x_1, y_1, x_2, y_2) w(x_1, y_1, x_2, y_2) d^4x = 1 \quad (24)$$

where $d^4x := dx_1 dx_2 dy_1 dy_2$. Explicitly,

$$A \int_{\mathbb{R}} x_1^{\lambda_3} e^{-\alpha x_1^2} dx_1 \int_{\mathbb{R}} y_1^{\lambda_2 - \lambda_3} e^{-\alpha y_1^2} dy_1 \int_{\mathbb{R}} x_2^{\lambda_2 - \lambda_0} e^{-\alpha x_2^2} dx_2 \int_{\mathbb{R}} y_2^{\lambda_0} e^{-\alpha y_2^2} dy_2 = 1 \quad (25)$$

It is well known that

$$\int_{\mathbb{R}} x^a e^{-bx^2} dx = \frac{1}{2} [1 + (-1)^a] b^{-\frac{a+1}{2}} \Gamma\left(\frac{a+1}{2}\right) \quad \text{with } \Re a > -1 \text{ and } \Re b > 0 \quad (26)$$

where Γ is the usual gamma function. Now, by our assumption, $a, b \in \mathbb{R}$. Thus, the above integrals are convergent when $a > -1$ and $b > 0$ in all the cases. Explicitly, we

have:

$$\int_{\mathbb{R}} x_1^{\lambda_3} e^{-\alpha x_1^2} dx_1 = \left[1 + (-1)^{\lambda_3}\right] \alpha^{-(\lambda_3+1)/2} \Gamma\left(\frac{\lambda_3+1}{2}\right) \quad (27)$$

$$\int_{\mathbb{R}} x_2^{\lambda_2-\lambda_0} e^{-\alpha x_2^2} dx_2 = \left[1 + (-1)^{\lambda_2-\lambda_0}\right] \alpha^{-(\lambda_2-\lambda_0+1)/2} \Gamma\left(\frac{\lambda_2-\lambda_0+1}{2}\right) \quad (28)$$

$$\int_{\mathbb{R}} y_1^{\lambda_2-\lambda_3} e^{-\alpha y_1^2} dy_1 = \left[1 + (-1)^{\lambda_2-\lambda_3}\right] \alpha^{-(\lambda_2-\lambda_3+1)/2} \Gamma\left(\frac{\lambda_2-\lambda_3+1}{2}\right) \quad (29)$$

$$\int_{\mathbb{R}} y_2^{\lambda_0} e^{-\alpha y_2^2} dy_2 = \left[1 + (-1)^{\lambda_0}\right] \alpha^{(\lambda_0+1)/2} \Gamma\left(\frac{\lambda_0+1}{2}\right) \quad (30)$$

In order that the above integrals are convergent, we need to demand that $\lambda_3 > -1$, $\lambda_2 - \lambda_0 > -1$, $\lambda_2 - \lambda_3 > -1$ and $\lambda_0 > -1$. Further, in order that the integrals are non-zero, we need

$$(-1)^{\lambda_3} \neq -1 \Rightarrow \lambda_3 \neq 2n + 1 \quad \forall n \in \mathbb{Z} \quad (31)$$

$$(-1)^{\lambda_2-\lambda_0} \neq -1 \Rightarrow \lambda_2 - \lambda_0 \neq 2n + 1 \quad \forall n \in \mathbb{Z} \quad (32)$$

$$(-1)^{\lambda_2-\lambda_3} \neq -1 \Rightarrow \lambda_2 - \lambda_3 \neq 2n + 1 \quad \forall n \in \mathbb{Z} \quad (33)$$

$$(-1)^{\lambda_0} \neq -1 \Rightarrow \lambda_0 \neq 2n + 1 \quad \forall n \in \mathbb{Z} \quad (34)$$

Thus, in conclusion, the above integrals exist if $\lambda_0 > -1$, $\lambda_2 > -2$ and $\lambda_3 > -1$ with an additional constraint that none of these should be odd integers. Notice further that $(-1)^\beta \in \mathbb{C}$ unless $\beta \in \mathbb{Z}$. Thus, λ_3 and λ_0 are integers. However, these cannot be odd integers, hence λ_0 and λ_3 are even integers. Let $\lambda_0 = 2n$ and $\lambda_3 = 2l$ such that $n, l \in \mathbb{N} \cup \{0\}$. For the sake of brevity, we will denote $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. Along the same lines, it follows that $\lambda_2 - \lambda_0$ should be an integer, giving us, λ_2 is an integer, in fact, an even integer. Further, given that $\lambda_2 - \lambda_0 > -1$, if $\lambda_2 = 2m$, $m \in \mathbb{Z}$ then $2m - 2n > -1 \Rightarrow m \geq n$. Similarly, $\lambda_2 - \lambda_3 > -1$ gives $m \geq l$. Thus, it follows that $m \geq \max\{l, n\}$. In conclusion, $\lambda_0 = 2n$, $\lambda_2 = 2l$ and $\lambda_3 = 2m$ with $l, m, n \in \mathbb{N}_0$ and $m \geq \max\{l, n\}$.

With all these simplifications, and by using properties of Gamma function [31], we get:

$$A_{lnm} = \frac{2^{2m} \alpha^{2m+2}}{\pi^2 \xi_{lnm}} \quad (35)$$

with

$$\xi_{lnm} = (2l-1)!! (2n-1)!! (2m-2n-1)!! (2m-2l-1)!! \quad (36)$$

where the symbols have their usual meanings. This finally yields,

$$P(x_1, y_1, x_2, y_2) = A_{lnm} x_1^{2l} y_1^{2(m-l)} x_2^{2(m-n)} y_2^{2n} \quad (37)$$

such that $l, m, n \in \mathbb{N}_0$ and $m \geq \max\{l, n\}$.

2.3 Level Repulsion

The matrix H (Eq. (1)) has two eigenvalues,

$$E_{\pm} = x_1 \pm i\sqrt{y_1^2 + y_2^2 + x_2^2}. \quad (38)$$

The level repulsion can be computed using the joint probability distribution:

$$\mathcal{P}(\sigma) = \int_{\mathbb{R}^4} \delta(|E_+ - E_-| - \sigma) P(x_1, y_1, x_2, y_2) w(x_1, y_1, x_2, y_2) d^4x \quad (39)$$

where the function w is the weight function defined above and δ is the usual Dirac delta measure (see, for example, [32]). Explicitly,

$$\begin{aligned} \mathcal{P}(\sigma) = \frac{1}{2} A_{lnm} \int_{\mathbb{R}^4} \delta\left(\sqrt{y_1^2 + y_2^2 + x_2^2} - \sigma/2\right) & x_1^{2l} y_1^{2(m-l)} x_2^{2(m-n)} y_2^{2n} \\ & \times e^{-\alpha(y_1^2 + y_2^2 + x_2^2 + x_1^2)} d^4x \end{aligned} \quad (40)$$

A close inspection of the above integral reveals that transformation to spherical polar coordinates could be useful here. Specifically, let $R^2 = y_1^2 + y_2^2 + x_2^2$ such that $y_1 = R \sin \theta \cos \phi$, $y_2 = R \sin \theta \sin \phi$, and $x_2 = R \cos \theta$. Using this transformation, the above integral becomes

$$\begin{aligned} \mathcal{P}(\sigma) = \frac{1}{2} A_{lnm} \int_{\mathbb{R}} x_1^{2l} e^{-\alpha x_1^2} dx_1 \int_0^{\infty} R^{4m-2l+2} e^{-\alpha R^2} \delta(R - \sigma/2) dR \\ \times \int_0^{2\pi} (\cos \phi)^{2m-2l} (\sin \phi)^{2n} d\phi \int_0^{\pi} (\sin \theta)^{2m+2n-2l+1} (\cos \theta)^{2m-2n} d\theta \end{aligned} \quad (41)$$

This integral can be evaluated analytically using the properties of beta integral, and the integral representation of the Gamma function (see, for example, [31]), leading to

$$\mathcal{P}(\sigma) = \frac{1}{2} \frac{\alpha^{2m-l+3/2}}{\sqrt{\pi} (2m-l+1/2)} \left(\frac{\sigma^2}{2}\right)^{2m-l+1} e^{-\alpha\sigma^2/4} \quad (42)$$

which is the desired result. The degree of level repulsion is given by the exponent on σ as $\sigma \rightarrow 0$, in this case it is $\sigma^{2(2m-l+1)}$. Recalling the admissible values of m, l , we see that the minimum value of m is l ; this gives the asymptotic behaviour as $\sigma^{2(l+1)}$. Thus, for the minimum value of $l = 0$, the degree of level repulsion coincides with that of the Gaussian Unitary Ensemble [2]. For any other value, the degree of level repulsion is stronger than 2. This is an algebraic power, in contrast to the case of pseudo-Hermitian matrices where, for one case, it was shown to vary as $\sigma \log(1/\sigma)$ [4].

2.4 Probability Distribution Function and Level Repulsion in a bounded Region of \mathbb{R}^4

We begin with the expression for the probability distribution (up to the normalisation factor) obtained above:

$$P(x_1, y_1, x_2, y_2) = Ax_1^{\lambda_3} y_1^{\lambda_2 - \lambda_3} x_2^{\lambda_2 - \lambda_0} y_2^{\lambda_0} \quad (43)$$

For the sake of convenience, we re-label some variables here. Out of the four variables, we will re-label y_1 to x_3 and y_2 to x_4 . This leads to:

$$P(x_1, x_2, x_3, x_4) = Ax_1^{\lambda_3} x_3^{\lambda_2 - \lambda_3} x_2^{\lambda_2 - \lambda_0} x_4^{\lambda_0} \quad (44)$$

Let $V \subset \mathbb{R}^4$ be defined by $x_i \geq 0$ for $i = 1, 2, 3, 4$, such that

$$\sum_{i=1}^4 x_i^2 \leq 1 \quad (45)$$

We first state a result due to Liouville that essentially generalises the beta integral to dimensions higher than 2 (see [31] for details):

Theorem 1. *If V is a region defined by $x_i \geq 0$, $i = 1, 2, \dots, n$ and $\sum x_i \leq 1$, then for $\Re \alpha_i > 0$,*

$$\int_V \prod_{i=1}^n x_i^{\alpha_i - 1} dx_1 \dots dx_n = \frac{\prod_{i=1}^n \Gamma(\alpha_i)}{\Gamma(1 + \sum_{i=1}^n \alpha_i)} \quad (46)$$

As an immediate corollary to this theorem, we have [31]:

Corollary 1. *If V is the region enclosed by $x_i \geq 0$ and $\sum (x_i/a_i)^{p_i} \leq 1$ then for $\Re \alpha_i > 0$,*

$$\int_V \prod_{i=1}^n x_i^{\alpha_i - 1} dx_1 \dots dx_n = \frac{\prod_{i=1}^n (a_i^{\alpha_i/p_i}) \Gamma(\alpha_i/p_i)}{\Gamma(1 + \sum_{i=1}^n \alpha_i/p_i)} \quad (47)$$

The normalisation constant in the present case can be obtained readily from this corollary by setting $p_i = 2$ and $a_i = 1$. Explicitly,

$$A \int_V \prod_{i=1}^4 x_i^{\alpha_i - 1} d^4x = \frac{1}{16} \frac{\prod_{i=1}^4 \Gamma(\alpha_i/2)}{\Gamma(1 + \sum_{i=1}^4 \alpha_i/2)} \quad (48)$$

giving us,

$$A = \frac{16 \Gamma(1 + \sum_{i=1}^4 \alpha_i/2)}{\prod_{i=1}^4 \Gamma(\alpha_i/2)} \quad (49)$$

where, $\alpha_1 = \lambda_3 + 1$, $\alpha_2 = \lambda_2 - \lambda_0 + 1$, $\alpha_3 = \lambda_2 - \lambda_3 + 1$ and $\alpha_4 = \lambda_0 + 1$ and $d^4x := dx_1 dx_2 dx_3 dx_4$. Given the condition that $\Re\alpha_i > 0$, in our case, the above integral exists if and only if $\lambda_3 + 1 > 0$, $\lambda_2 - \lambda_3 + 1 > 0$, $\lambda_2 - \lambda_0 + 1 > 0$, and $\lambda_0 + 1 > 0$. Thus, it follows that the integral and hence the normalisation constant exists provided that $\lambda_3 > -1$, $\lambda_0 > -1$ and $\lambda_2 > \max\{\lambda_0, \lambda_3\}$. Further notice that for the domain over which λ_k are defined, the normalisation constant is always non-zero, as it should be. There is no further restriction on the values that the separation constants can take, which is in contrast with the case of full \mathbb{R}^4 , where the separation constants were restricted to positive even integral values.

We now compute the nearest-neighbour level spacing distribution in this case. Specifically, we need to evaluate the following integral:

$$\mathcal{P}(\sigma) = \frac{1}{2}A \int_V \prod_{i=1}^4 x_i^{\alpha_i-1} \delta\left(\sqrt{x_2^2 + x_3^2 + x_4^2} - \sigma/2\right) d^4x \quad (50)$$

It is convenient to use hyper-spherical coordinates here [33]. Specifically, we set

$$x_1 = r \cos \phi_1 \quad (51)$$

$$x_2 = r \cos \phi_2 \sin \phi_1 \quad (52)$$

$$x_3 = r \sin \theta \sin \phi_1 \sin \phi_2 \quad (53)$$

$$x_4 = r \cos \theta \sin \phi_1 \sin \phi_2 \quad (54)$$

such that $r^2 = \sum_{i=1}^4 x_i^2$ ($0 \leq r \leq 1$), $\phi_1 \in [0, \pi]$, $\phi_2 \in [0, \pi]$, and $\theta \in [0, 2\pi]$. In the present case, since we are restricting ourselves to non-negative values of x_i , it follows that $\phi_1, \phi_2, \theta \in [0, \pi/2]$. Further, the determinant of the Jacobian matrix is $r^3 (\sin \phi_1)^2 \sin \phi_2$, and $x_2^2 + x_3^2 + x_4^2 = r^2 (\sin \phi_1)^2$, which leads to

$$\mathcal{P}(\sigma) = \frac{1}{2}A \int_0^1 \int_{\xi}^{\pi/2} \int_0^{\pi/2} \int_0^{\pi/2} \delta(r \sin \phi_1 - \sigma/2) r^{\sum_i \alpha_i - 1} (\sin \theta)^{\alpha_3 - 1} (\cos \theta)^{\alpha_4 - 1} \times (\cos \phi_2)^{\alpha_2 - 1} (\sin \phi_2)^{\alpha_3 + \alpha_4 - 1} (\cos \phi_1)^{\alpha_1 - 1} (\sin \phi_1)^{\alpha_2 + \alpha_3 + \alpha_4 - 1} dr d\phi_1 d\phi_2 d\theta \quad (55)$$

Notice that the domain of integral over ϕ_1 has been cut to $[\xi, \pi/2]$ with $\xi > 0$. This is due to the constraint implied by the Dirac Delta measure, as will be explained below.

We will first carry out integral over r . In order to do that, we first write:

$$\mathcal{P}(\sigma) = \frac{1}{2}A \int_0^1 \int_{\xi}^{\pi/2} \int_0^{\pi/2} \int_0^{\pi/2} \delta(r - \sigma/(2 \sin \phi_1)) r^{\sum_i \alpha_i - 1} (\sin \theta)^{\alpha_3 - 1} (\cos \theta)^{\alpha_4 - 1} \times (\cos \phi_2)^{\alpha_2 - 1} (\sin \phi_2)^{\alpha_3 + \alpha_4 - 1} (\cos \phi_1)^{\alpha_1 - 1} (\sin \phi_1)^{\alpha_2 + \alpha_3 + \alpha_4 - 2} dr d\phi_1 d\phi_2 d\theta \quad (56)$$

leading to:

$$\begin{aligned}
\mathcal{P}(\sigma) &= \frac{1}{2} A \left(\frac{\sigma}{2} \right)^{\sum_i \alpha_i - 1} \int_{\xi}^{\pi/2} (\cos \phi_1)^{\alpha_1 - 1} (\sin \phi_1)^{-\alpha_1 - 1} d\phi_1 \\
&\quad \times \int_0^{\pi/2} (\sin \theta)^{\alpha_3 - 1} (\cos \theta)^{\alpha_4 - 1} d\theta \int_0^{\pi/2} (\cos \phi_2)^{\alpha_2 - 1} (\sin \phi_2)^{\alpha_3 + \alpha_4 - 1} d\phi_2
\end{aligned} \tag{57}$$

Out of these, the second and the third are the standard beta integrals [31], and are given by

$$\int_0^{\pi/2} (\sin \theta)^{\alpha_3 - 1} (\cos \theta)^{\alpha_4 - 1} d\theta = \frac{\Gamma(\alpha_3/2) \Gamma(\alpha_4/2)}{2\Gamma((\alpha_3 + \alpha_4)/2)} \tag{58}$$

and

$$\int_0^{\pi/2} (\cos \phi_2)^{\alpha_2 - 1} (\sin \phi_2)^{\alpha_3 + \alpha_4 - 1} d\phi_2 = \frac{\Gamma((\alpha_3 + \alpha_4)/2) \Gamma(\alpha_2/2)}{2\Gamma((\alpha_2 + \alpha_3 + \alpha_4)/2)} \tag{59}$$

In order to evaluate the first integral, we need to estimate ξ first. We know that

$$0 \leq \frac{\sigma^2}{4(\sin \phi_1)^2} \leq 1 \tag{60}$$

The lower limit of the integral, ξ , is obtained by demanding that it is the one that gives the smallest possible value of ϕ_1 , leading to $\sin \xi = \sigma/2$, giving us $\xi = \arcsin(\sigma/2)$. The first integral can now be evaluated:

$$\int_{\xi}^{\pi/2} (\cos \phi_1)^{\alpha_1 - 1} (\sin \phi_1)^{-\alpha_1 - 1} d\phi_1 = \frac{1}{\alpha_1} (\cot \xi)^2 = \frac{4 - \sigma^2}{\sigma^2} \tag{61}$$

Upon combining all these factors and inserting the value of normalisation constant, we finally get

$$\mathcal{P}(\sigma) = \frac{4\Gamma\left(1 + \sum_{i=1}^4 \alpha_i/2\right)}{\Gamma((\alpha_2 + \alpha_3 + \alpha_4)/2)} \frac{4 - \sigma^2}{\alpha_1 \sigma^2} \left(\frac{\sigma}{2}\right)^{\sum_i \alpha_i - 1} \tag{62}$$

Finally, note that $\sum_i \alpha_i = 2\lambda_2 + 4$. Thus,

$$\mathcal{P}(\sigma) = \frac{4\Gamma(\lambda_2 + 3)}{\Gamma(\lambda_2 - \lambda_3/2 + 3/2)} \frac{4 - \sigma^2}{(\lambda_3 + 1)} \left(\frac{\sigma}{2}\right)^{2\lambda_2 + 1} \tag{63}$$

3 Concluding remarks

As emphasized in the main text, one of the significant developments reported here is to obtain the distribution function of the elements of an ensemble of non-Hermitian $SU(2)$ -like matrices on the basis of symmetry and statistical independence. Hitherto, it was always assumed to be a Gaussian distribution. For at least the ensemble considered here, this assumption has been lifted. This has led us to the consideration of certain cases dictated by the domain of the matrix elements. Employing the new-found probability distribution, we have calculated the nearest-neighbour level spacing distribution. The degree of level repulsion is tunable as there appear two parameters; this result is potentially significant in modelling physical situations where interaction strength varies (see, e.g. [13]) and also in connection with number theory (see [34]). However, at the lower bound of the level repulsion parameter of our ensemble is the degree for the Gaussian Unitary ensembles (GUE). It is worth recalling that GUE corresponds to physical systems which violate time-reversal invariance. In line with earlier results [4, 5, 15, 16], we are led to conjecture that an additional breakdown of parity makes the degree of level repulsion stronger; thus, we conjecture:

$$\lim_{\sigma \rightarrow 0} P(\sigma) \sim \sigma^\nu, \nu \geq 2.$$

The connections between random matrix theory and exactly solvable models are profound [17, 20, 35–37]. Upon appropriate generalization, we believe that this work opens up the possibility of connections with new, exactly solvable models also.

Acknowledgments. SA acknowledges financial support through the Cyrus Guzder Fellowship. Valuable comments from the Doctoral Advisory Committee of UM-DAE Centre for Excellence in Basic Sciences are gratefully acknowledged.

Declarations

- Funding: Partial financial support to Stalin Abraham was received through Cyrus Guzder Fellowship.
- Conflict of interest/Competing interests: The authors have no competing interests to declare that are relevant to the content of this article.
- Ethics approval and consent to participate: Not applicable.
- Consent for publication: The authors give the publisher the due consent for publication of this article.
- Data availability: Not applicable.
- Materials availability: Not applicable.
- Code availability: Not applicable.
- Author contribution: The authors declare equal contribution to this article.

References

- [1] E. P. Wigner, Statistical Properties of Real Symmetric Matrices with Many Dimensions, Proceedings of the Fourth Canadian Mathematical Congress (University of Toronto Press, Toronto, Canada, 1959) 174-184.
- [2] M. L. Mehta, Random matrices, 3rd edn. (Academic Press, 2004).
- [3] S. R. Jain and P. Gaspard, *Dynamical response and time correlation functions in random quantum systems*, arXiv:2408.09419v1 [cond-mat.stat-mech]. <https://doi.org/10.48550/arXiv.2408.09419>
- [4] Z. Ahmed and S. R. Jain, Phys. Rev. E 67, 045106 (2003). <https://doi.org/10.1103/PhysRevE.67.045106>
- [5] Z. Ahmed and S. R. Jain, J. Phys. A: Math. Gen. 36, 3349 (2003). <https://doi.org/10.1088/0305-4470/36/37/309>
- [6] E. C. G. Sudarshan, Phys. Rev. 123, 2183 (1961). <https://doi.org/10.1103/PhysRev.123.2183>
- [7] M. C. Pease III, Methods of Matrix Algebra (Academic Press, New York, 1965).
- [8] F. G. Scholtz, H. B. Geyer, and F. Hahne, Ann. Phys. 213, 74 (1992). [https://doi.org/10.1016/0003-4916\(92\)90284-S](https://doi.org/10.1016/0003-4916(92)90284-S)
- [9] S. R. Jain, Pramana-J. Phys. 73, 251 (2009). <https://doi.org/10.1007/s12043-009-0116-6>
- [10] D. Alonso and S. R. Jain, Phys. Lett. B 387, 812 (1996). [https://doi.org/10.1016/0370-2693\(96\)01104-5](https://doi.org/10.1016/0370-2693(96)01104-5)
- [11] S. R. Jain and D. Alonso, J. Phys. A: Math. Gen. 30, 4993 (1997). <https://doi.org/10.1088/0305-4470/36/12/327>
- [12] Y. Nambu, in Preludes in theoretical physics, ed. By A. de Shalit, H. Feshbach and L. van Hove (North Holland, Amsterdam, 1966).
- [13] G. Date, S. R. Jain, and M. V. N. Murthy, Phys. Rev. E 51, 198 (1995). <https://doi.org/10.1103/PhysRevE.51.198>
- [14] B. Grémaud and S. R. Jain, J. Phys. A: Math. Gen. 31, L637 (1998). <https://doi.org/10.1088/0305-4470/31/37/003>
- [15] S. R. Jain and S. C. L. Srivastava, Phys. Rev. E 78, 036213 (2008). <https://doi.org/10.1103/PhysRevE.78.036213>
- [16] S. R. Jain and S. C. L. Srivastava, Pramana-J. Phys. 73, 989 (2009).

- <https://doi.org/10.1007/s12043-009-0174-9>
- [17] S. R. Jain, Czech J. Phys. 56, 1021 (2006).
<https://doi.org/10.1007/s10582-006-0397-7>
- [18] S. Rey and Ph. Choquard, Eur. J. Phys. 18, 94 (1997).
<https://doi.org/10.1088/0143-0807/18/2/007>
- [19] S. C. L. Srivastava and S. R. Jain, Phys. Rev. E 85, 041143 (2012).
<https://doi.org/10.1103/PhysRevE.85.041143>
- [20] S. R. Jain, B. Grémaud, and A. Khare, Phys. Rev. 66, 016216 (2002).
<https://doi.org/10.1103/PhysRevE.66.016216>
- [21] J. Ginibre, J. Math. Phys. 6, 440 (1965). <https://doi.org/10.1063/1.1704292>
- [22] F. Haake, Quantum signatures of chaos, (Springer, Heidelberg, 1991).
- [23] P. J. Forrester, Log-gases and random matrices, (Princeton University Press, Princeton, 2010).
- [24] C. M. Bender and D. W. Hook, Rev. Mod. Phys. 96, 045002 (2024).
<https://doi.org/10.1103/RevModPhys.96.045002>
- [25] Carl M. Bender and Stefan Boettcher, Phys. Rev. Lett. 80, 5243 (1998).
<https://doi.org/10.1103/PhysRevLett.80.5243>
- [26] Carl M. Bender, Stefan Boettcher, and Peter Meisinger, J. Math. Phys. 40, 2201 (1999). <https://doi.org/10.1063/1.532860>
- [27] Carl M. Bender, Peter N. Meisinger, and Qinghai Wang, J. Phys. A 36, 6791 (2003). <https://doi.org/10.1088/0305-4470/36/25/312>
- [28] Ali Mostafazadeh, J. Math. Phys. 43, 205 (2002).
<https://doi.org/10.1063/1.1418246>
- [29] Ali Mostafazadeh, J. Math. Phys. 43, 2814 (2002).
<https://doi.org/10.1063/1.1461427>
- [30] Stalin Abraham and A. Bhagwat, *On Classification and Geometric Characterizations of Ensembled 2×2 Pseudo Hermitian and PT -Symmetric Matrices*, arXiv:2410.18530v1 [math-ph]
<https://doi.org/10.48550/arXiv.2410.18530>
- [31] G. E. Andrews, R. Askey, R. Roy, Special functions (Cambridge University Press, Cambridge, 1999).

- [32] I. M. Gel'fand and G. E. Shilov, Generalized Functions, Vol. 1, Properties and Operations (AMS Chelsea Publishing, 2016).
- [33] L. E. Blumenson, The American Mathematical Monthly 67, 63 (1960).
<https://doi.org/10.2307/2308932>
- [34] Z. Ahmed and S. R. Jain, Mod. Phys. Lett. A 21, 331 (2006).
<https://doi.org/10.1142/S0217732306018792>
- [35] G. Auberson, S. R. Jain and A. Khare, J. Phys. A: Math. Gen. 34, 695 (2001).
<https://doi.org/10.1088/0305-4470/34/4/302>
- [36] F. Calogero, J. Math. Phys. 10, 2191 (1969). <https://doi.org/10.1063/1.1664820>
- [37] B. Sutherland, Beautiful models, (World Scientific, Singapore, 2005).