

High-order Accurate Inference on Manifolds

Chengzhu Huang* and Anru R. Zhang†

(January 14, 2025)

Abstract

We present a new framework for statistical inference on Riemannian manifolds that achieves high-order accuracy, addressing the challenges posed by non-Euclidean parameter spaces frequently encountered in modern data science. Our approach leverages a novel and computationally efficient procedure to reach higher-order asymptotic precision. In particular, we develop a bootstrap algorithm on Riemannian manifolds that is both computationally efficient and accurate for hypothesis testing and confidence region construction. Although locational hypothesis testing can be reformulated as a standard Euclidean problem, constructing high-order accurate confidence regions necessitates careful treatment of manifold geometry. To this end, we establish high-order asymptotics under a fixed normal chart centered at the true parameter, thereby enabling precise expansions that incorporate curvature effects. We demonstrate the versatility of this framework across various manifold settings—including spheres, the Stiefel manifold, fixed-rank matrices manifolds, and rank-one tensor manifolds—and, for Euclidean submanifolds, introduce a class of projection-like coordinate charts with strong consistency properties. Finally, numerical studies confirm the practical merits of the proposed procedure.

1 Introduction

This paper focuses on the statistical inference for parameters defined on a Riemannian manifold. Let $\mathcal{X}_n = \{X_1, \dots, X_n\} \subset \mathbb{R}^m$ be observations independently sampled from a distribution \mathbb{P}_{θ_0} , where θ_0 resides on a p -dimensional Riemannian manifold \mathcal{M} . Our aim is to make inference on θ based on \mathcal{X}_n . Given an objective function $L(\theta, x) : \mathcal{M} \times \mathbb{R}^m \rightarrow \mathbb{R}$ and its sample average analog $L_n(\theta, \mathcal{X}_n) := \sum_{i \in [n]} L(\theta, X_i)/n$, an M-estimator, an important instance of extremum estimator, can be defined as

$$\tilde{\theta}_n := \arg \min_{\theta \in \mathcal{M}} L_n(\theta, \mathcal{X}_n). \quad (1)$$

For brevity, we often write $L_n(\theta, \mathcal{X}_n)$ in place of $L_n(\theta)$.

A natural approach to assess the location of $\tilde{\theta}_n$ is to exploit a suitable coordinate representation. Under the assumption that the likelihood function is geodesically convex but possibly non-smooth, Brunel (2023) explored the first-order asymptotics of $\tilde{\theta}_n$ and proved that the coordinate of $\tilde{\theta}_n$ under the logarithmic mapping centered at θ_0 and a basis $\{e_i\}$ of $T_{\theta_0}\mathcal{M}$ is asymptotically distributed as

$$\sqrt{n} \cdot \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\tilde{\theta}_n) \Rightarrow \mathcal{N}(\mathbf{0}, \Sigma),$$

*Department of Statistics, Columbia University

†Department of Biostatistics & Bioinformatics and Department of Computer Science, Duke University

where $\pi_{\{e_i\}}$ denotes a canonical mapping from $T_{\theta_0}\mathcal{M}$ to \mathbb{R}^p induced by $\{e_i\}$, Log_{θ_0} represents the logarithmic mapping, and

$$\begin{aligned} \Sigma := & \left(\left(\mathbb{E} [\nabla^2 L(\theta_0, X_1)[e_i, e_j]] \right)_{i,j \in [p]} \right)^{-1} \left(\mathbb{E} [e_i L(\theta_0, X_1) e_j L(\theta_0, X_1)^\top] \right)_{i,j \in [p]} \\ & \cdot \left(\left(\mathbb{E} [\nabla^2 L(\theta_0, X_1)[e_i, e_j]] \right)_{i,j \in [p]} \right)^{-1}. \end{aligned} \quad (2)$$

The primary goal of this paper is to go beyond this asymptotic normality of the exact extremum estimators and establish computationally feasible inference methods that achieve high-order accuracy in manifold settings. Specifically, we aim to design a hypothesis test that ensures a high-order accurate control of type-I error. Additionally, for a confidence level $1 - \alpha$, our goal is to construct a confidence region $I \subseteq \mathcal{M}$ that satisfies

$$|\mathbb{P}_{\mathcal{X}_n}[\theta_0 \in I] - (1 - \alpha)| = o(n^{-1/2}).$$

To achieve this goal, we employ a resampling procedure for inference on θ , so as to mimic the distribution of the targeted quantity as well as to construct the region I . Toward the theoretical guarantees for the inference tasks, we focus on demonstrating the distributional closeness between the approximate solution and the resampled approximate solution after appropriate studentification, in line with the Edgeworth expansions of (approximate) M-estimators (Hall and Horowitz, 1996; Andrews, 2002).

Challenges in Manifold Settings. We describe a smooth manifold, without delving into concepts of differential geometry, as a space that locally behaves like Euclidean space. A smooth manifold endowed with a metric structure is a Riemannian manifold. The primary challenge in dealing with such an abstract structure lies in the locality of the coordinates; specifically, we must account for the different representations of a point under various charts. The core idea of this paper is to consider exponential mapping, where the relationship between different charts is determined by the Riemannian connection defined on the manifold. Leveraging results from differential geometry (Gavrilov, 2007), we are able to upper bound the curvature’s effect on the coordinates and explicitly interpret the distribution of the proposed estimator. Furthermore, exponential mappings are often computationally infeasible in many applications, prompting us to explore alternative methods, such as retraction mappings, to achieve second-order accuracy.

1.1 Extremum Estimators on Manifolds

An extremum estimator is obtained by optimizing an objective function that depends on observed data (Newey and McFadden, 1994; Fisher, 1922; Wald, 1949). Maximum likelihood estimation (MLE) is a prominent example and has been widely used for inference in parametric models. Notably, the framework of extremum estimators naturally lends itself to a manifold-based formulation in several important statistical analysis scenarios.

Constrained Parameter Space. The asymptotic properties of statistical models with a fixed-dimensional parameter in Euclidean space have been extensively studied (Van der Vaart, 2000). However, there are many statistical problems that do not exhibit the desirable properties associated with Euclidean spaces. For example, the behavior of the MLE and the associated information metric in curved exponential families exhibit notable complexity (Efron, 1975, 1978). The nonlinearity constraints in such problems offer intriguing interpretability while also posing challenges in terms

of coordinate representation, naturally falling into the differential geometric viewpoint. Given a parametric space $\Theta \in \mathbb{R}^p = \{\theta : f_i(\theta) = 0, i \in [r]\}$ ($r < p$) where $\{f_i\}_{i \in [r]}$ are smooth functions whose derivatives are linearly independent, the constant-rank level set theorem (Lee, 2013) shows Θ is an embedded submanifold with dimension $(p - r)$. However, the approach to enable high-order accurate inference in these settings remains blank.

Nonconvex Optimization. Nonconvex optimization problems often arise in modern machine learning and signal processing, often accompanied by complex geometric structures that cannot be fully captured using standard Euclidean tools. In particular, the matrix/tensor-valued parameter estimation problems with low-rank constraints naturally lead to formulations on specialized manifolds such as the fixed-rank matrices manifold, Stiefel manifold, and the Grassmannian manifold. Recent developments have shown the significance of such manifold-based approaches, either theoretically or algorithmically (e.g., Boumal et al. (2019); Li et al. (2021); Chen et al. (2020); Huang et al. (2016)).

Examples We present several key Riemannian manifolds below:

- The Stiefel manifold $\text{St}(p, r) := \{\mathbf{X} \in \mathbb{R}^{p \times r} : \mathbf{X}^\top \mathbf{X} = \mathbf{I}_r\}$ consists of all $p \times r$ orthonormal matrices. This manifold generalizes the spheres to higher dimensions and is pivotal in applications requiring orthonormal bases or dimensionality reduction, such as in factor analysis (Plumbley, 2005).
- The fixed-rank matrices manifold, denoted as $\mathcal{R}_{r, p_1, p_2}$, represents the set of all $p_1 \times p_2$ matrices with rank r and possesses a smooth manifold structure. This manifold is commonly explored in the study of low-rank matrix completion and approximation problems (Vandereycken, 2013; Mishra et al., 2014). For example, when the observed data consists of noisy measurements of an underlying element in $\mathcal{R}_{r, p_1, p_2}$, a critical challenge lies in recovering the true underlying parameter and performing uncertainty quantification.

In short, viewing estimation and inference problems through the lens of extremum estimators on manifolds can reveal new insights, particularly in scenarios where traditional Euclidean frameworks fall short.

1.2 Bootstrap and Studentization on Manifolds

As the sample size increases in a fixed model, it is widely recognized that the distribution of the extremum estimator exhibits certain asymptotic properties. However, a significant challenge lies in the lack of direct access to the actual asymptotic distribution. To address this, a notable idea is to *studentize* the target statistic while resampling the data, namely, to divide the (resampled) statistic by a sample-based estimate of its standard deviation (e.g., *t*-statistic) to achieve a more accurate approximation of the statistic’s distribution. Over the past decades, computationally efficient bootstrap procedures have been explored extensively in works such as Davidson and MacKinnon (1999); Andrews (2002). However, these multi-step estimators are not readily applicable to manifold settings, where unique geometrical structures introduce additional complexities. This limitation motivates our investigation into bootstrap procedures specifically tailored for manifold-based problems.

In this paper, we propose a fine-grained inference procedure to address the locality challenges in manifold settings, enabling the derivation of high-order accurate confidence regions through appropriate studentization techniques. To ensure high-order accuracy in the distributional approximation

obtained via bootstrap, we carefully control each resampled term, incorporating the effects of the Riemannian curvature of the underlying manifold.

1.3 Our Contributions

The contributions of this paper include the following three folds.

First, for a class of well-formalized M-estimators on Riemannian manifolds, we develop a bootstrapping algorithm that stands out for its computational efficiency and high-order accuracy in hypothesis testing and confidence region construction.

Secondly, we discover that, while performing a locational hypothesis testing on a Riemannian manifold can be naturally transformed into a well-studied testing problem on a Euclidean space, constructing a high-order accurate confidence region poses additional challenges due to a manifold’s inherent geometrical structure. To ensure both the computational efficiency and the coverage accuracy of the proposed confidence interval construction methods, we establish high-order asymptotics of the specific statistics in our algorithms. To address the ambiguity of the coordinate expression, we adopt a fixed normal chart centered at the true parameter in our analysis. This approach allows us to expand the functions of interest through a Euclidean lens while accounting for the impact of manifold curvature when transitioning across different charts.

Finally, our methodology and theoretical results extend to a wide range of statistical settings. We specifically study the inference on spheres, the Stiefel manifold, fixed-rank matrices manifolds, and rank-one tensors manifolds using our developed framework. Notably, for Euclidean submanifolds, we introduce a class of projection-like coordinate charts equipped with the necessary consistency properties, providing a robust foundation for further applications.

1.4 Related Literature

This work intersects with several key areas of research, including Riemannian optimization, high-order asymptotics, and manifold inference.

Firstly, the extension of classic optimization methods (Smith, 1993, 2014; Huper and Trumpf, 2004; Helmke and Hüper, 2000; Dennis Jr and Schnabel, 1996) to Riemannian manifold settings (Absil et al., 2009) has attracted increasing attention, driven by the emergence of large-scale matrix data (Luo et al., 2024a; Luo and Trillos, 2022) and tensor data (Luo and Zhang, 2023).

Next, the study of high-order asymptotics dates back to the efforts in refining the central limit theorem for fixed-dimensional cases. Foundational results such as Bhattacharya et al. (1978); Babu and Singh (1984); Bhattacharya (1987); Hall and Horowitz (1996); Hall (2013) established the Edgeworth expansion and its bootstrap adaptations for empirical averages and functions of weighted sums. Additionally, Andrews (2002) introduced computationally efficient algorithms combined with studentized statistics to achieve high-order asymptotics for M-estimation problems. These developments serve as a foundation for extending such methods to manifold settings. There is another series of research focusing on the Edgeworth expansion of U-statistics (Callaert et al., 1980; Bickel et al., 1986; Lai and Wang, 1993), especially on network moment (Zhang and Xia, 2022; Shao et al., 2022).

Moreover, research in manifold inference covers diverse geometrical approaches, with a focus on Barycenter problems (Bhattacharya and Patrangenaru, 2003, 2005; Eltzner and Huckemann, 2019). Recent advancements have addressed uncertainty quantification for low-rank models, offering perspectives tailored to high-dimensional settings (Chen et al., 2019; Xia and Yuan, 2021; Xia et al., 2022; Xie and Zhang, 2024).

1.5 Preliminaries on Riemannian Manifolds

To ensure this paper is self-contained, we provide a concise introduction to Riemannian manifolds, particularly in the context of numerical optimization (Absil et al., 2009).

Smooth Structure. A smooth manifold \mathcal{M} of dimension p is a connected paracompact Hausdorff space for which every point has a neighborhood U that is homeomorphic to an open subset Ω of \mathbb{R}^p . Such a homeomorphism $\phi : U \rightarrow \Omega$ is called a (coordinate) chart. A tangent vector X to the manifold \mathcal{M} at x is a mapping from the set of real-valued functions defined on \mathcal{M} (i.e., $C^\infty(\mathcal{M})$) to \mathbb{R} such that there exists a smooth curve $\gamma(t)$ on \mathcal{M} with $\gamma(0) = x$ and $X : f \mapsto \left. \frac{d}{dt} f \circ \gamma(t) \right|_{t=0} =: Xf$ for all $f \in C^\infty(\mathcal{M})$. The tangent space $T_x\mathcal{M}$ of a smooth manifold \mathcal{M} is the linear space of all tangent vectors at x . The disjoint union of $T_x\mathcal{M}$ over $x \in \mathcal{M}$ is defined as the tangent bundle $T\mathcal{M} := \cup_{x \in \mathcal{M}} T_x\mathcal{M}$. A vector field on \mathcal{M} is a smooth function that assigns a tangent vector $v \in T_\theta\mathcal{M}$ to each point $\theta \in \mathcal{M}$. $\Gamma(\mathcal{M})$ denotes the collection of possible vector fields on \mathcal{M} . Naturally, given a vector field $X \in \Gamma(\mathcal{M})$ and a smooth function $f \in C^\infty(\mathcal{M})$ on \mathcal{M} , Xf is again a smooth function on \mathcal{M} . For further use, we define the mapping $XY : C^\infty(\mathcal{M}) \rightarrow \mathbb{R}$ for $X, Y \in \Gamma(\mathcal{M})$ such that $XYf \mapsto X(Yf)$ for $f \in C^\infty(\mathcal{M})$. Given two smooth manifolds \mathcal{M}, \mathcal{N} and a smooth function $f : \mathcal{N} \rightarrow \mathcal{M}$, we define a mapping called the differential of f at $x \in \mathcal{N}$ as a linear mapping $T_x\mathcal{N} \rightarrow T_{f(x)}\mathcal{M}$ such that $df(X)g = X(g \circ f)$ for every $X \in T_x\mathcal{N}$ and every $g \in C^\infty(\mathcal{M})$.

Riemannian Manifold. Further, a Riemannian manifold \mathcal{M} is a smooth manifold equipped with a smooth positive-definite inner product $\langle \cdot, \cdot \rangle$ on the tangent space $T_p\mathcal{M}$ at each point p . With an inner product at each point, the length of a piece-wise smooth curve $\gamma(t), t \in [0, 1]$ from x to x' can be defined as $L(\gamma) = \int_0^1 \langle \dot{\gamma}(t), \dot{\gamma}(t) \rangle^{\frac{1}{2}} dt$. We also denote the distance of two points x, x' on manifold \mathcal{M} by $\text{dist}(x, x')$, which is defined by the infimum of $L(\gamma)$ over all possible piece-wise smooth curves from x to x' . Further, given two vector fields on \mathcal{M} , we note that $\langle X, Y \rangle$ is a smooth function on \mathcal{M} by the smoothness of $\langle \cdot, \cdot \rangle$.

Levi-Civita Connection. To enable the comparison of tangent vectors across disparate tangent spaces, we introduce the notion of connection $\nabla : \Gamma(\mathcal{M}) \times \Gamma(\mathcal{M}) \rightarrow \Gamma(\mathcal{M})$, as a generalization of the directional derivative on a vector field to Riemannian manifolds, satisfying that (i) $\nabla_{f_1X_1+f_2X_2}Y = f_1\nabla_{X_1}Y + f_2\nabla_{X_2}Y$, (ii) $\nabla_X(\alpha_1Y_1 + \alpha_2Y_2) = \alpha_1\nabla_XY_1 + \alpha_2\nabla_XY_2$, (iii) $\nabla_X(fY) = (Xf)Y + f\nabla_XY$ with $X, Y, X_1, X_2, Y_1, Y_2 \in \Gamma(\mathcal{M})$, $f_1, f_2, f \in C^\infty(\mathcal{M})$, and $\alpha_1, \alpha_2 \in \mathbb{R}$. Among all possible choices of affine connections, the Levi-Civita connection is the unique one that preserves the metric and satisfies the torsion-free-ness; to be specific, an affine connection is called the Levi-Civita connection if (i) $X \langle Y, Z \rangle = \langle \nabla_XY, Z \rangle + \langle Y, \nabla_XZ \rangle$ holds for any vector fields X, Y, Z ,¹ (ii) $\nabla_XY - \nabla_YX - (XY - YX) = 0$ holds for any vector fields X, Y, Z (see the details of the Levi-Civita connection's interpretation in Jost (2008) and Lee (2018)). We only consider the Levi-Civita connections throughout this paper.

Parallel Transport. Parallel transport is a local realization of a connection that intrinsically defines an isometry between tangent spaces. Letting $\mu \in \Gamma(\mathcal{M})$ be a vector field on \mathcal{M} and $\gamma(t)$ be a smooth curve, $\mu(\gamma(t))$ is called the parallel transport when $\nabla_{\dot{\gamma}(t)}(\mu(\gamma(t))) = 0$ holds for arbitrary t . If the shortest curve connecting x with x' is unique, we denote the tangent vector at x' from the

¹Since $\langle Y, Z \rangle$ is a smooth function on \mathcal{M} , $X \langle Y, Z \rangle := X(\langle Y, Z \rangle)$ is also a smooth function given an additional tensor field X on \mathcal{M} .

parallel transport of $v \in T_x\mathcal{M}$ in the tangent space of x along the shortest path (also a geodesic by (Lee, 2018, Theorem 6.4)) by $T_{x \rightarrow x'}(v)$; in fact, Theorem 6.17 in Lee (2018) ensures the uniqueness of the geodesic connecting two arbitrary points in a small neighborhood. Such a local viewpoint will frequently be employed in the subsequent discussion.

Retraction. The notion of retraction mapping is a crucial tool in developing efficient and practical optimization algorithms on manifolds. In much of the previous literature, a retraction mapping R refers to a smooth mapping from $T\mathcal{M}$ to \mathcal{M} . However, since in this paper, we are mainly interested in quantities related to specific coordinate representations, we define a retraction $R := \{R_x\}_{x \in \mathcal{M}}$ to be a collection of retraction mappings such that each retraction mapping R_x centered at x is a diffeomorphism from an open neighborhood $U \subset \mathbb{R}^p$ of $\mathbf{0}$ to \mathcal{M} satisfying $R_x(\mathbf{0}) = x$, $\{dR_x \delta_i\}$ is an orthogonal basis of $T_x\mathcal{M}$ for each $x \in \mathcal{M}$, and $R_x \circ dR_x^{-1}(y)$ with respect to x and y is a smooth mapping from an open neighborhood of the zero section of $T\mathcal{M}$ to \mathcal{M} . The Euclidean realization of a retraction presented here also leads to the subsequent discussion on the necessity of smoothness in coordinate bases across different charts. To impose regularity conditions on retractions, we further introduce the notion of a second-order retraction, defined as follows.

Definition 1. A retraction R is second-order if it holds for every $x \in \mathcal{M}$ and \mathbf{y} in an open neighborhood of \mathbb{R}^p that

$$\left\| \text{Log}_x(R_x(\mathbf{y})) - \pi_{\{dR_x e_i\}}^{-1}(\mathbf{y}) \right\| = O(\|\mathbf{y}\|_2^3).$$

Derivatives. Given a Riemannian manifold \mathcal{M} equipped with the Levi-Civita connection ∇ and a smooth function f on \mathcal{M} , the k -th derivative $\nabla^k \subset \Gamma(\otimes_{i=1}^k T\mathcal{M}^*)^2$ of f ($k \in \mathbb{N}^+$) is defined recursively as follows

$$\begin{aligned} \nabla f[X_1] &= \nabla^1 f[X_1] = X_1 f, \\ \nabla^2 f[X_1, X_2] &= X_1(\nabla^1 f[X_2]) - \nabla^1 f[\nabla_{X_1} X_2], \\ \nabla^k f[X_1, \dots, X_k] &= X_1 \nabla^{k-1} f[X_2, \dots, X_k] - \nabla^{k-1} f[\nabla_{X_1} X_2, \dots, X_k] \\ &\quad - \nabla^{k-1} f[X_2, \nabla_{X_1} X_3, \dots, X_k] - \dots - \nabla^{k-1} f[X_2, X_3, \dots, \nabla_{X_1} X_k]. \end{aligned}$$

Note that a retraction $R = \{R_x\}_{x \in \mathcal{M}}$ is a second-order retraction if and only if $\nabla^2 R_x = \mathbf{0}$ for every $x \in \mathcal{M}$.

Remark 1. In a way, a retraction typically suggests the topological correspondence between the derivative of mapping and the tangent space of the manifold but does not involve any metric structure defined on the manifold. However, a second-order retraction goes beyond the topological structure; namely, it takes the metric structure of a manifold into account. As a consequence, different metric structures on a manifold usually lead to different second-order retractions.

1.6 Notation

For any positive integer n , we denote $[n] = \{1, \dots, n\}$. We also denote the i -th canonical basis of \mathbb{R}^p by δ_i . Given a matrix $\mathbf{A} \in \mathbb{R}^{N \times N}$, we denote its Moore–Penrose inverse by \mathbf{A}^\dagger . The ℓ_2 -norm of a vector \mathbf{v} is denoted by $\|\mathbf{v}\|_2$, and the Frobenius norm of a matrix $\mathbf{A} \in \mathbb{R}^{n_1 \times n_2}$ is denoted by

² $\Gamma(\otimes_{i=1}^k T\mathcal{M}^*)$ denotes the collection of smooth functions that assign an element of $\underbrace{T_x\mathcal{M}^* \otimes \dots \otimes T_x\mathcal{M}^*}_{k \text{ times}}$ to each x in \mathcal{M} where $T_x\mathcal{M}^*$ denotes the dual space of $T_x\mathcal{M}$.

$\|\mathbf{A}\|_F$. Given a vector space \mathcal{T} with norm $\|\cdot\|$, we denote a sphere $\{y \in \mathcal{T} : \|y - x\| \leq \rho\}$ centered at $x \in \mathcal{T}$ with radius ρ by $B(x, \rho)$.

Consider a Riemannian manifold \mathcal{M} of dimension p . The norm of a tangent vector $v \in \text{T}\mathcal{M}$ is written as $\|v\| := \langle v, v \rangle^{\frac{1}{2}}$. We denote by $\text{Exp}_x(\cdot)$ and $\text{Log}_x(\cdot)$ the exponential mapping and the logarithmic mapping at $x \in \mathcal{M}$. Given a basis $\{e_i\}$ of $\text{T}\mathcal{M}$, we define the canonical mapping $\pi_{\{e_i\}} : \text{T}_x\mathcal{M} \rightarrow \mathbb{R}^p$ at x by $\pi_{\{e_i\}}(y) = \mathbf{v}$ with $y = \sum_{i=1}^p v_i e_i$ and $\mathbf{v} = (v_1, \dots, v_p) \in \mathbb{R}^p$. For a function $f : \mathcal{M} \rightarrow \mathbb{R}$ defined on \mathcal{M} , we denote its composition with R_θ by $f^{\text{Retr}(\theta)}$.

Throughout this paper, coordinate expressions and abstract tensor field expressions will be used interchangeably. To avoid confusion, we will use the notation $\nabla^k f$ for $f : \mathcal{M} \rightarrow \mathbb{R}$ to denote the k -th derivative of f in the abstract tensor field expression, while we use $\bar{\nabla}^\nu f(\mathbf{x}) := \frac{\partial^{\nu_1}}{\partial x_1^{\nu_1}} \cdots \frac{\partial^{\nu_p}}{\partial x_p^{\nu_p}} f(\mathbf{x})$ for $f : \mathbb{R}^p \rightarrow \mathbb{R}$ to denote the partial derivative of f related to $\nu \in \mathbb{N}^p$ in the coordinate expression. Moreover, given a sufficiently smooth function $f : \mathbb{R}^p \rightarrow \mathbb{R}$, we denote the gradient column vector by $\bar{\nabla} f(\mathbf{x}) = (\frac{\partial}{\partial x_j} f(\mathbf{x}))_{j \in [p]}$ and the Hessian matrix by $\bar{\nabla}^2 f(\mathbf{x}) = (\frac{\partial^2}{\partial x_i \partial x_j} f(\mathbf{x}))_{i, j \in [p]} \in \mathbb{R}^{p \times p}$. Further, given a tangent basis $\{e_i\}$ of $\text{T}_\theta\mathcal{M}$ at $\theta \in \mathcal{M}$ and a multi-index $\nu \in \mathbb{N}^p$, we denote

$$\nabla^{|\nu|} f(\theta) [\underbrace{e_1, \dots, e_1}_{\nu^{(1)}}, \underbrace{e_2, \dots, e_2}_{\nu^{(2)}}, \dots, \underbrace{e_p, \dots, e_p}_{\nu^{(p)}}]$$

by $\nabla^\nu f(\theta)$, referred to as the ν -derivative of f under the basis $\{e_i\}$; by the knowledge of differential geometry, this is also equivalent to $\frac{\partial^{\nu^{(1)}}}{\partial x_1^{\nu^{(1)}}} \cdots \frac{\partial^{\nu^{(p)}}}{\partial x_p^{\nu^{(p)}}} f \circ \text{Exp} \circ \pi_{\{e_i\}}^{-1}$.

For an objective function $L(\theta, x) : \mathcal{M} \times \mathbb{R}^m \rightarrow \mathbb{R}$ or a composition $L(R_\theta(\boldsymbol{\eta}), x)$ with a mapping $R_\theta(\boldsymbol{\eta})$ from \mathbb{R}^p to \mathcal{M} , we slightly abuse the notation by performing all differentiation operations with respect to the first argument, without explicitly indicating it.

2 High-order Accurate Inference Using Extremum Estimators

Inference with high-order accuracy typically utilizes the Edgeworth expansions for both the statistics of interest and their resampled counterparts. We first define the resampled objective function and the resampled M-estimator as

$$L_n^*(\theta) := \sum_{i \in [n]} L(\theta, X_i^*)/n, \quad \tilde{\theta}_n^* = \arg \min_{\theta \in \mathcal{M}} L_n^*(\theta), \quad (3)$$

where $\mathcal{X}_n^* = \{X_i^*\}_{i \in [n]}$ is the resampled collection with replacement from \mathcal{X}_n .

Despite the validity of the high-order asymptotics related to the exact solution, the explicit solution is often infeasible. To address this, we employ the Riemannian Newton method to proceed with estimation and inference. Intuitively, given a reasonable initialization, Newton iterates usually converge to the local stationary point of the objective function. We thereby turn to look at the property of solutions to first-order conditions as a starting point. We consider relaxed solutions to (1) and (3), satisfying the first-order conditions:

$$\nabla L_n(\tilde{\theta}_n) = 0, \quad (4)$$

$$\nabla L_n^*(\tilde{\theta}_n^*) = 0. \quad (5)$$

In cases where the solutions are not unique, they are defined as the ones closest to θ_0 in their respective cases.

2.1 Riemannian Newton Iteration

The classic Newton’s method (Dennis Jr and Schnabel, 1996) is an iterative procedure that incorporates second-order information to seek a critical point of the cost function. It is well-established that in deterministic settings, Newton’s method achieves a quadratic rate of convergence if an initial point is close enough to a minimizer with a nonsingular second derivative in Euclidean settings (Dennis Jr and Schnabel, 1996) as well as in general Riemannian manifold settings (Absil et al., 2009). In the former case, letting $\boldsymbol{\theta}_{k-1}$ be the iterative point at the $(k-1)$ st step and f be an objective function, Newton’s update is written as

$$\boldsymbol{\theta}_k := \boldsymbol{\theta}_{k-1} - [\bar{\nabla}^2 f(\boldsymbol{\theta}_{k-1})]^{-1} \bar{\nabla} f(\boldsymbol{\theta}_{k-1}) \in \mathbb{R}^p.$$

Given a Riemannian manifold \mathcal{M} , Newton’s update³ takes form

$$\boldsymbol{\theta}_k := R_{\boldsymbol{\theta}_{k-1}} \left(-[\bar{\nabla}^2 f \circ R_{\boldsymbol{\theta}_{k-1}}(\mathbf{0})]^\dagger \bar{\nabla} f \circ R_{\boldsymbol{\theta}_{k-1}}(\mathbf{0}) \right) \in \mathcal{M},$$

where R is a retraction defined in Definition 1 and $f : \mathcal{M} \rightarrow \mathbb{R}$ is an objective function.

In what follows, we consider an objective function L composed with a random data set \mathcal{Y} and introduce a Riemannian Newton iteration procedure as described in Algorithm 1⁴, primarily for constructing confidence regions in subsequent steps. A key difference between optimization on a Riemannian manifold and in Euclidean space is that there is no fixed coordinate representation (or coordinate chart) for this manifold. Instead, at each iteration of Algorithm 1, we operate on the local chart $R_{\hat{\boldsymbol{\theta}}_{s-1}}^{-1}(\cdot)$, centered at the current estimate $\hat{\boldsymbol{\theta}}_{s-1}$.

Algorithm 1: Riemannian Newton Iteration on a Manifold

Input: Initial point θ^{initial} and data collection \mathcal{Y} , iteration times t

Output: $\hat{\boldsymbol{\theta}}_t$

1 Let $\theta^{(0)} = \theta^{\text{initial}}$ **for** $s = 1, \dots, t$ **do**

2 Update:

$$\boldsymbol{\theta}^{(s)} = R_{\boldsymbol{\theta}^{(s-1)}} \left(-[\bar{\nabla}^2 \left(\sum_{X \in \mathcal{Y}} L(R_{\boldsymbol{\theta}^{(s-1)}}(\cdot), X) \right) |_{\mathbf{0}} / |\mathcal{Y}|]^\dagger [\bar{\nabla} \left(\sum_{X \in \mathcal{Y}} L(R_{\boldsymbol{\theta}^{(s-1)}}(\cdot), X) \right) |_{\mathbf{0}} / |\mathcal{Y}|] \right).$$

3 **end**

Remark 2. As discussed in Adler et al. (2002); Absil et al. (2009), local quadratic convergence is maintained even if Algorithm 1 employs only a first-order retraction. However, as we will demonstrate in Section 2.3.2, a second-order retraction is essential to explicitly incorporate the high-order asymptotics of the constructed statistics, thereby enabling high-order accurate inference.

Since the empirical solution to (4) is likely to stay close to the resampled solution to (5) with high probability, it naturally serves as an ideal initialization. We therefore propose to integrate the estimation with the resampling procedures, allowing the stochastic optimization process to produce a sequence of replicates based on a similar underlying mechanism as the original estimate. Our procedure generates the estimate $\hat{\boldsymbol{\theta}}_n$ and the resampled estimates $\{\boldsymbol{\theta}_{n,i}^*\}_{i \in [b]}$, as summarized in Algorithm 2.

³The Moore–Penrose inverse is introduced to address the potential singularity issue of the Hessian matrix.

⁴One may notice that the domain of R_θ is an open neighborhood of $\mathbf{0}$, but not necessarily \mathbb{R}^p . To ensure the well-definiteness, we simply let $R_\theta(\mathbf{y})$ be θ for any \mathbf{y} outside the domain of R_θ .

Algorithm 2: Resampled Riemannian Newton Iteration on a Manifold

Input: Initial estimate $\widehat{\theta}_n^{(0)}$, number of burnings `burn`, number of resamplings b

Output: Approximate extremum estimate $\widehat{\theta}_n$ and the resampled approximate extremum estimate series $\{\widehat{\theta}_n^{*[i]}\}_{i=1}^b$

- 1 Implement Algorithm 1 with initial point $\theta^{\text{initial}} = \widehat{\theta}_n^{(0)}$, $t = \text{burn}$, and the dataset \mathcal{X}_n to obtain $\widehat{\theta}_n$.
 - 2 **for** $i = 1, \dots, b$ **do**
 - 3 Resample the dataset \mathcal{X}_n to obtain $\mathcal{X}_n^{[i]}$ (i.e., draw n samples with replacement from \mathcal{X}_n to form $\mathcal{X}_n^{[i]}$), and implement Algorithm 1 with the initial point $\widehat{\theta}_n$, $t = 2$, and the dataset $\mathcal{X}_n^{[i]}$ to obtain $\widehat{\theta}_n^{*[i]}$.
 - 4 **end**
-

In the bootstrapping stage of Algorithm 2, we perform a two-step Riemannian Newton initialized from the point estimate $\widehat{\theta}_n$, provided the resampled dataset $\mathcal{X}_n^{[i]}$. Heuristically speaking, both $\widehat{\theta}_n$ and $\widetilde{\theta}_n$ are likely to lie within a radius- $O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$ spherical neighborhood around θ_0 with high probability, and $\widehat{\theta}_n$ usually serves as a nice approximation of $\widetilde{\theta}_n$. Provided the quadratic convergence rate of Newton's method, the two-step update is expected to provide a good approximation of the solution to the first-order condition given the dataset $\mathcal{X}_n^{[i]}$, which is further confirmed by our high-order asymptotic analysis in Section 3.

2.2 Hypothesis Testing on Manifold

We consider a hypothesis testing problem on a Riemannian manifold, specifically focusing on the following locational testing problem:

$$H_0 : \theta_0 = \theta_1 \in \mathcal{M} \quad \text{versus} \quad H_1 : \theta_0 \neq \theta_1, \quad \text{with significance level } \alpha.$$

We focus on controlling the test rejection probability when the null hypothesis holds. To achieve this, we note that by restricting attention to a neighborhood $U \subseteq \mathcal{M}$ of θ_1 , where (ϕ, U) is a smooth chart with $\phi(\theta_1) = \mathbf{0} \in \mathbb{R}^p$, the testing problem reduces to

$$H_0 : \phi(\theta_0) = \mathbf{0} \quad \text{versus} \quad H_1 : \phi(\theta_0) \neq \mathbf{0}, \quad \text{with significance level } \alpha.$$

In light of this, a natural approach is to employ an alternative loss function $\sum_{i \in [n]} L(\phi^{-1}(\cdot), X_i)/n : \phi(U) \rightarrow \mathbb{R}$. Moreover, we define the t -statistics $T_{\phi, j}$, $j \in [p]$, the Wald statistic W_ϕ , and their bootstrap alternatives $T_{\phi, j}^*$, W_ϕ^* as follows⁵:

$$T_{\phi, j} = \frac{\phi(\widehat{\theta}_n)_j}{(\boldsymbol{\Sigma}_\phi)_{j, j}^{1/2}}, \quad T_{\phi, j}^* = \frac{\phi(\widehat{\theta}_n^*)_j - \phi(\widehat{\theta}_n)_j}{(\boldsymbol{\Sigma}_\phi^*)_{j, j}^{1/2}}, \quad \text{for } j \in [p],$$

$$W_\phi = \phi(\widehat{\theta}_n)^\top \boldsymbol{\Sigma}_\phi^{-1} \phi(\widehat{\theta}_n), \quad W_\phi^* = (\phi(\widehat{\theta}_n^*) - \phi(\widehat{\theta}_n))^\top \boldsymbol{\Sigma}_\phi^{*\dagger} (\phi(\widehat{\theta}_n^*) - \phi(\widehat{\theta}_n)).$$

⁵If the denominator $(\boldsymbol{\Sigma}_\phi^*)_{j, j}^{1/2}$ is equal to zero, we simply let $T_{\phi, j}^* = 0$.

Here $\widehat{\theta}_n^*$ denotes a generic $\widehat{\theta}_{n,1}^*$ in Algorithm 2 and $\mathcal{X}^* = \{X_i^*\}_{i \in [n]}$ denotes a generic resampled dataset $\mathcal{X}_n^{[1]}$. The matrices Σ_ϕ and Σ_ϕ^* are defined as

$$\begin{aligned}\Sigma_\phi &:= \left(\bar{\nabla}^2 L_n(\phi^{-1}(\cdot))|_{\phi(\widehat{\theta}_n)} \right)^{-1} \\ &\quad \left(\frac{1}{n} \sum_{i \in [n]} \bar{\nabla} L(\phi^{-1}(\cdot), X_i)|_{\phi(\widehat{\theta}_n)} \bar{\nabla} L(\phi^{-1}(\cdot), X_i)|_{\phi(\widehat{\theta}_n)}^\top \right) \left(\bar{\nabla}^2 L_n(\phi^{-1}(\cdot))|_{\phi(\widehat{\theta}_n)} \right)^{-1}, \\ \Sigma_\phi^* &:= \left(\bar{\nabla}^2 L_n^*(\phi^{-1}(\cdot))|_{\phi(\widehat{\theta}_n^*)} \right)^{-1} \\ &\quad \left(\frac{1}{n} \sum_{i \in [n]} \bar{\nabla} L(\phi^{-1}(\cdot), X_i^*)|_{\phi(\widehat{\theta}_n^*)} \bar{\nabla} L(\phi^{-1}(\cdot), X_i^*)|_{\phi(\widehat{\theta}_n^*)}^\top \right) \left(\bar{\nabla}^2 L_n^*(\phi^{-1}(\cdot))|_{\phi(\widehat{\theta}_n^*)} \right)^{-1}.\end{aligned}$$

Note that the coordinate derivative operations, $\bar{\nabla}$ and $\bar{\nabla}^2$, refer specifically to the first argument of L , i.e., the coordinates under the retraction mapping ϕ^{-1} .

Intuitively, $\phi(\widehat{\theta}_n)$ and $\phi(\widehat{\theta}_n^*)$ are close to the exact solutions $\phi(\widetilde{\theta}_n)$ and $\phi(\widetilde{\theta}_n^*)$, where $\widetilde{\theta}_n$ and $\widetilde{\theta}_n^*$ are defined in (4) and (5). In view of the delta method (Lemma 4), it thus suffices to focus on the high-order asymptotics of the latter ones. Moreover, a useful observation is that $\phi(\widetilde{\theta}_n)$ and $\phi(\widetilde{\theta}_n^*)$ are also the solutions to

$$\bar{\nabla} L_n(\phi^{-1}(\mathbf{x})) = 0 \quad \text{and} \quad \bar{\nabla} L_n^*(\phi^{-1}(\mathbf{x})) = 0,$$

respectively, where $L_n(\phi^{-1}(\cdot))$ could be viewed as an empirical loss function defined on a neighborhood of \mathbb{R}^p . Since well-established high-order asymptotic results exist for M-estimation (Hall and Horowitz, 1996; Andrews, 2002), these can be applied in our context. We have the following formal result.

Proposition 1. *Suppose the following assumptions hold:*

1. *The zero vector is the unique solution in a sufficiently small spherical neighborhood U of θ_0 to $\mathbb{E}[\bar{\nabla} L(\phi^{-1}(\cdot), X_1)] = \mathbf{0}$. There exists a function $C_g(\mathbf{x})$ such that $\|\bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}_1), \mathbf{x}) - \bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}_2), \mathbf{x})\|_2 \leq C_g(\mathbf{x}) \|\boldsymbol{\eta}_1 - \boldsymbol{\eta}_2\|_2$ for all \mathbf{x} in the support of X_1 and all $\boldsymbol{\eta}_1, \boldsymbol{\eta}_2 \in \phi(U)$. $\mathbb{E}[C_g^{q_1}(X_1)] < \infty$ and $\mathbb{E}\|\bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}), X_1)\|_2^{q_1} < \infty$ for all $0 < q_1 < \infty$.*
2. *The p -by- p Hessian matrix $\mathbb{E}[\bar{\nabla}^2 L(\phi^{-1}(\mathbf{0}), X_1)]$ is of full rank p . $\bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}), \mathbf{x})$ is nine-times differentiable with respect to $\boldsymbol{\eta}$ on $\phi(U)$ for all \mathbf{x} in the support of X_1 . Let $f(\boldsymbol{\eta}, \mathbf{x})$ be a vector containing $\bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}), \mathbf{x})$, $\bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}), \mathbf{x}) \bar{\nabla} L(\phi^{-1}(\boldsymbol{\eta}), \mathbf{x})^\top$, and their derivatives through order 5. There is a function $C_{\partial f}(\mathbf{x})$ such that $\|\bar{\nabla}^\nu f(\mathbf{0}, \mathbf{x}) - \bar{\nabla}^\nu f(\boldsymbol{\eta}, \mathbf{x})\|_2 \leq C_{\partial f}(\mathbf{x}) \|\boldsymbol{\eta}\|_2$ for all $\boldsymbol{\eta} \in \phi(U)$, all $\boldsymbol{\nu} \in \mathbb{N}^p$ with $\sum_{i \in [p]} \nu_i \leq 4$, and all \mathbf{x} in the support of X_1 . $\mathbb{E}[C_{\partial f}(X_1)^{q_2}] < \infty$ and $\mathbb{E}[(\bar{\nabla}^\nu f(\mathbf{0}, X_1))^{q_2}] < \infty$ for all $q_2 > 0$ and all $\boldsymbol{\nu}$ with $\sum_{i \in [p]} \nu_i \leq 4$. The function $f(\mathbf{0}, \mathbf{x})$ is once differentiable with respect to \mathbf{x} with uniformly continuous first derivative.*
3. *The random vector $f(\mathbf{0}, X_1)$ satisfies Cramér's condition, that is,*

$$\limsup_{\|\mathbf{t}\| \rightarrow \infty} |\mathbb{E}[e^{i\mathbf{t}^\top f(\mathbf{0}, X_1)}]| < 1.$$

If $\theta_0 = \theta_1$, then it holds for every $\xi \in (0, \frac{1}{2}]$ that

$$\begin{aligned}\mathbb{P}_{\mathcal{X}_n} [T_{\phi, j} > z_{T_{\phi, j}, 2, \alpha}^*] &= \alpha + o(n^{-1+\xi}), \\ \mathbb{P}_{\mathcal{X}_n} [T_{\phi, j} < z_{T_{\phi, j}, 2, 1-\alpha}^*] &= \alpha + o(n^{-1+\xi}), \\ \mathbb{P}_{\mathcal{X}_n} [W_\phi > z_{W_\phi, 2, \alpha}^*] &= \alpha + o(n^{-\frac{3}{2}+\xi}),\end{aligned}$$

where $z_{T_{\phi,j},k,\alpha}^*$ is defined as a value that minimizes $|\mathbb{P}[T_{\phi,j}^* \leq z | \mathcal{X}_n] - (1 - \alpha)|$ over $z \in \mathbb{R}$ and $z_{W_{\phi,2,\alpha}}^*$ is defined analogously.

Proposition 1 indicates that by repeatedly resampling the data, we can obtain a consistent estimate of $z_{T_{\phi,j},2,\alpha}^*$ or $z_{W_{\phi,2,\alpha}}^*$ to serve as a critical value. According to the Glivenko–Cantelli theorem (Durrett, 2019, Theorem 2.4.9), this approach yields a more accurate approximation of the distribution of $T_{\phi,j}$ or W_{ϕ} , ensuring that the actual rejection probability closely aligns with the intended significance level α .

2.3 Confidence Region Constructions on Manifold

In this section, we delve into the construction of confidence regions on Riemannian manifolds. A natural approach, drawing on classic ideas from statistics, would be to convert the hypothesis test introduced in Section 2.2 into a confidence region. However, this approach may be computationally infeasible since we no longer have access to the presumed location information of θ_0 , which previously allowed for a convenient reduction from the manifold setting to a Euclidean one. Additionally, the choice of coordinate representations introduces further randomness, making the construction of confidence regions notably distinct from that of hypothesis testing.

2.3.1 Algorithms for Confidence Region Constructions

Selecting appropriate coordinate representations is essential for performing inference on Riemannian manifolds. For hypothesis testing, it is natural to adopt a fixed coordinate system at θ_0 . However, this approach no longer applies to confidence interval construction, necessitating transitions between different charts. These coordinate transformations are inherently nonlinear and influenced by the curvature of the local region. Consequently, when resampling the data and constructing studentized statistics, curvature effects may cause the distribution function of the studentized statistics for $\widehat{\theta}_n^{*[i]}$ to deviate from being second-order consistent with that related to $\widehat{\theta}_n$.

Algorithm for Wald-Type Statistic. The Wald-type statistic is often advantageous for constructing an elliptically shaped confidence region within a fixed chart or directly on the parameter manifold \mathcal{M} . By leveraging the information gathered during the resampling process, we can approximate the distribution of the following Wald-type statistic:

$$W := R_{\widehat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^\dagger R_{\widehat{\theta}_n}^{-1}(\theta_0), \quad (6)$$

where

$$\widehat{\Sigma} := (\overline{\nabla}^2 L_n(R_{\widehat{\theta}_n}(\cdot))|_{\mathbf{0}})^{-1} \left(\sum_{j \in [n]} \overline{\nabla} L(R_{\widehat{\theta}_n}(\cdot), X_j)|_{\mathbf{0}} \overline{\nabla} L(R_{\widehat{\theta}_n}(\cdot), X_j)|_{\mathbf{0}}^\top / n \right) (\overline{\nabla}^2 L_n(R_{\widehat{\theta}_n}(\cdot))|_{\mathbf{0}})^{-1}. \quad (7)$$

Since $-R_{\theta_0}^{-1}(\widehat{\theta}_n)$ is approximately equal to $R_{\widehat{\theta}_n}^{-1}(\theta_0)$ up to a rotation as noted in (6), a natural approximation of $-R_{\theta_0}^{-1}(\widehat{\theta}_n)$ is $R_{\theta_0}^{-1}(\widehat{\theta}_n) - R_{\theta_0}^{-1}(\widehat{\theta}_n^{*[i]})$, which is closely related to $R_{\widehat{\theta}_n^{*[i]}}^{-1}(\widehat{\theta}_n)$. Therefore, when working with the i -th resampled dataset $\mathcal{X}_n^{[i]} = \{X_j^{[i]}\}_{j \in [n]}$ in Algorithm 2, replacing $R_{\widehat{\theta}_n}(\theta_0)$ and $\widehat{\Sigma}$ with $R_{\widehat{\theta}_n^{*[i]}}(\widehat{\theta}_n)$ and $\widehat{\Sigma}^{[i]}$, respectively, yields the counterpart resampled statistic $W^{*[i]}$ to W as follows:

$$W^{*[i]} := R_{\widehat{\theta}_n^{*[i]}}^{-1}(\widehat{\theta}_n)^\top \widehat{\Sigma}^{[i]\dagger} R_{\widehat{\theta}_n^{*[i]}}^{-1}(\widehat{\theta}_n),$$

where $\tilde{\Sigma}^{[i]}$ is defined as

$$\tilde{\Sigma}^{[i]} := (\bar{\nabla}^2 L_n^{[i]}(R_{\hat{\theta}_n^{*[i]}}(\cdot))|_{\mathbf{0}})^{-1} \left(\sum_{j \in [n]} \bar{\nabla} L(R_{\hat{\theta}_n^{*[i]}}(\cdot), X_j^{[i]})|_{\mathbf{0}} \bar{\nabla} L(R_{\hat{\theta}_n^{*[i]}}(\cdot), X_j^{[i]})|_{\mathbf{0}}^\top / n \right) (\bar{\nabla}^2 L_n^{[i]}(R_{\hat{\theta}_n^{*[i]}}(\cdot))|_{\mathbf{0}})^{-1} \quad (8)$$

with the shorthand notation $L_n^{[i]}$ for $\sum_{j \in [n]} L(\cdot, X_j^{[i]})/n$ provided the i -th resampled dataset $\mathcal{X}_n^{[i]} = \{X_j^{[i]}\}_{j \in [n]}$.

The algorithm is summarized in Algorithm 3⁶. Heuristically, since the statistic of interest W and its bootstrapped counterpart are largely unaffected by the choice of chart, as discussed in Section 2.3.2, the conditional distribution of $W^{*[i]}$ given \mathcal{X}_n is expected to closely approximate the distribution of W with high probability. This will be further justified in Section 3.

Algorithm 3: Approximate Resampled M-Estimates on a Manifold with Wald-based Statistic

Input: Approximate extremum estimate $\hat{\theta}_n$ and the resampled approximated extremum estimate series $\{\hat{\theta}_n^{*[i]}\}_{i=1}^b$ derived from Algorithm 2

Output: confidence region for θ_0 on the chart $R_{\hat{\theta}_n}$

- 1 Compute the α -level quantile $w_\alpha := \arg \min_{x \in \mathbb{R}} |\sum_{i \in [b]} 1\{W^{*[i]} \leq x\}/b - (1 - \alpha)|$ with

$$W^{*[i]} = R_{\hat{\theta}_n^{*[i]}}^{-1}(\hat{\theta}_n)^\top \tilde{\Sigma}^{[i]\dagger} R_{\hat{\theta}_n^{*[i]}}^{-1}(\hat{\theta}_n);$$

- 2 Construct the $(1 - \alpha)$ -level confidence region on the region of $R_{\hat{\theta}_n}$:

$$I_{1-\alpha}^{\text{Wald}} = \{\mathbf{x} : \|\mathbf{x}\|_2 \leq \rho, \mathbf{x}^\top \hat{\Sigma}^\dagger \mathbf{x} \leq w_\alpha\}$$

with $\hat{\Sigma}$ defined in (7).

Algorithm for Intrinsic t -statistics. We now move on to investigate how to quantify the uncertainty given a single coordinate of θ_0 under the chart $R_{\hat{\theta}_n}$. Before presenting our algorithm for constructing confidence intervals on a single coordinate, here we would like to discuss further the insight into why inference on a single coordinate of the true parameter is an even more difficult task beyond establishing certain asymptotics in Euclidean cases.

Intractability of Local Charts on Riemannian Geometry. A natural approach to inference is to establish specific asymptotics of the targeted quantities under a specific coordinate chart \tilde{R} , namely, the limiting distribution of $\tilde{R}(\theta_0)$ where \tilde{R} represents a coordinate chart centered at some location θ' that could depend on the data. However, this local Euclidean viewpoint introduces some extra intractability, as the basis $\{\tilde{e}_i\}$ induced by the chart \tilde{R} we use is usually related to the samples themselves, which cooperate some complicated dependency with the asymptotic normality of $\pi_{\{T_{\theta_0 \rightarrow \theta'} e_i\}} \circ \text{Log}_{\theta'}(\theta_0)$, and consequently deter the expected limit distribution. Notably, the cooperated dependency varies across different manifolds, making it challenging to derive a general inference procedure for a single coordinate.

Consider, for instance, the fixed-rank matrices manifold and the SVD-based retraction in Example 3, which is elaborated on later. Suppose that all the nonzero singular values of the ground

⁶Similar to the handling of the locality of R_θ in Algorithm 2, if $\hat{\theta}_n^{*[i]}$ is outside the domain of $R_{\hat{\theta}_n}^{-1}$, we set the corresponding quantity $W^{*[i]}$ to be zero. Similar rules apply to the quantity $T^{*[i]}$ in Algorithm 4.

Hist. of intr. t-stat (n = 1000, Fx.-Rk. Mat.)




Figure 1: A Counterexample: Irregular Empirical Distribution Using an Irregular Chart. We consider the fixed-rank matrices manifold with the sample size $n = 1000$ and the ground truth parameter $\theta_0 = \text{diag}(1, 1, 0, 0)$. The histogram represents the studentized version of $(R_{\hat{\theta}_n}^{-1}(\theta_0))_1$ while the blue curve depicts the probability density function of the standard normal distribution.

truth θ_0 are identical. In this scenario, the vector field induced by the first coordinate⁷ of the charts is not even differentiable. This non-differentiability results in the limiting distribution of the studentized version of $(R_{\hat{\theta}_n}^{-1}(\theta_0))_1$ deviating significantly from the standard Gaussian distribution, even with a large sample size, as shown in Figure 1.

A Differentiable Chart Helps. However, if specific smoothness conditions, for instance, the additional assumption in Theorem 3.2, on charts locally around θ_0 are imposed, we are able to construct confidence intervals with respect to a linear combination of coordinates on the chart centered at $\hat{\theta}_n$. In particular, we aim to approximate the distribution of the studentized t -statistic

$$T = \frac{\mathbf{a}^\top R_{\hat{\theta}_n}^{-1}(\theta_0)}{(\mathbf{a}^\top \hat{\Sigma} \mathbf{a})^{\frac{1}{2}}}.$$

Similar to the construction of the resampled Wald statistic, we consider the counterpart resampled statistic $T^{*[i]}$:

$$T^{*[i]} = \frac{\mathbf{a}^\top R_{\hat{\theta}_n^{*[i]}}^{-1}(\hat{\theta}_n)}{(\mathbf{a}^\top \check{\Sigma}^{[i]} \mathbf{a})^{\frac{1}{2}}}.$$

The algorithm is outlined in Algorithm 4.

Algorithm for Statistics Related to Extrinsic Coordinates. For problems involving a submanifold embedded in the Euclidean space, our framework can also accommodate a broad range

⁷The first coordinate refers to the top left entry of \mathbf{A} for a tangent vector $\mathbf{Z} = \mathbf{U} \begin{bmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{B} & \mathbf{0} \end{bmatrix} \mathbf{V}^\top$.

Algorithm 4: Approximate Resampled M-Estimates on a Riemannian Manifold with Intrinsic t -statistic

Input: Direction \mathbf{a} , the approximate extremum estimate $\hat{\theta}_n$, and the resampled approximate extremum estimate series $\{\hat{\theta}_n^{*[i]}\}_{i=1}^b$ derived from Algorithm 2

Output: Confidence region for θ_0 on the chart $R_{\hat{\theta}_n}$ along the direction \mathbf{a}

- 1 Compute the $(1 - \alpha)$ -quantile $w_{1-\alpha} := \arg \min_{x \in \mathbb{R}} \left| \sum_{i \in [b]} 1\{T^{*[i]} \leq x\} / b - \alpha \right|$ with

$$T^{*[i]} = \frac{\mathbf{a}^\top R_{\hat{\theta}_n^{*[i]}}^{-1}(\hat{\theta}_n)}{(\mathbf{a}^\top \hat{\Sigma}^{[i]} \mathbf{a})^{1/2}};$$

- 2 Construct the $(1 - \alpha)$ -level confidence region for $\mathbf{a}^\top R_{\hat{\theta}_n}^{-1}(\theta_0)$:

$$I_{1-\alpha}^{\text{Intr.t}} = \left(-\infty, -w_{1-\alpha} \cdot (\mathbf{a}^\top \hat{\Sigma} \mathbf{a})^{\frac{1}{2}} \right]$$

with $\hat{\Sigma}$ defined in (7).

of extrinsic representations of the true parameter. Specifically, let \mathcal{M} be a submanifold embedded in \mathbb{R}^d . Suppose that the true parameter is $\theta_0 \in \mathcal{M} \subset \mathbb{R}^d$ and $f : \mathbb{R}^d \rightarrow \mathbb{R}$ is a smooth function with $\nabla f \circ R_{\theta_0} \neq \mathbf{0}$. Under these conditions, Algorithm 5 yields a second-order accurate confidence interval for $f(\theta_0)$.

Algorithm 5: Approximate Resampled M-Estimates on a Riemannian Manifold with Extrinsic t -Statistic

Input: Smooth function f , the approximate extremum estimate $\hat{\theta}_n$, and the resampled approximate extremum estimate series $\{\hat{\theta}_n^{*[i]}\}_{i=1}^b$ obtained from Algorithm 2

Output: Confidence interval for $f(\theta_0)$

- 1 Compute the $(1 - \alpha)$ -quantile $w_{1-\alpha} := \arg \min_{x \in \mathbb{R}} \left| \sum_{i \in [b]} 1\{T^{*[i]} \leq x\} / b - (1 - \alpha) \right|$ with

$$T^{*[i]} = (f(\hat{\theta}_n^{*[i]}) - f(\hat{\theta}_n)) / \sqrt{\nabla f(\hat{\theta}_n^{*[i]})^\top \hat{\Sigma}^{[i]} \nabla f(\hat{\theta}_n^{*[i]})};$$

- 2 Construct the $(1 - \alpha)$ -level confidence interval for $f(\theta_0)$:

$$I_{1-\alpha}^{\text{Extr.t}} = \left(-\infty, f(\hat{\theta}_n) - w_\alpha \cdot (\nabla f(\hat{\theta}_n)^\top \hat{\Sigma} \nabla f(\hat{\theta}_n))^{\frac{1}{2}} \right]$$

with $\hat{\Sigma}$ defined in (7).

2.3.2 Confidence Regions Construction: An Insight into Coordinate Representations

To clarify the importance of employing a second-order retraction and selecting appropriate coordinate representations across different charts in Algorithm 3 and Algorithm 4, we begin by examining the geometry of a two-dimensional submanifold \mathcal{M} embedded in \mathbb{R}^3 . Recall that θ_0 represents the true parameter, $\hat{\theta}_n$ denotes the outputs of Algorithm 2, and $\hat{\theta}_n^*$ refers to a generic version of $\hat{\theta}_{n,b}^*$.

One appealing but impractical choice to represent $\theta_0, \hat{\theta}_n, \hat{\theta}_n^*$ is to utilize the chart centered at θ_0 so that the problem is reduced to the well-studied Euclidean M-estimation problem by considering the objective function $L(R_{\theta_0}(\cdot), \cdot) : \mathbb{R}^p \times \mathbb{R}^m \rightarrow \mathbb{R}$. Although in practice the mapping R_{θ_0} is inaccessible without the knowledge of θ_0 , it enlightens us to bridge the actual coordinate representations with the oracle coordinates under R_{θ_0} ; indeed, the roadmap starts with relating

$R_{\hat{\theta}_n}^{-1}(\theta_0)$ and $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$ with $R_{\theta_0}^{-1}(\hat{\theta}_n)$ and $R_{\theta_0}^{-1}(\hat{\theta}_n^*) - R_{\theta_0}^{-1}(\hat{\theta}_n)$, respectively. Consider the case where $\hat{\theta}_n$ and $\hat{\theta}_n^*$ are contained in a sufficiently small neighborhood of θ_0 , which can be justified under some regularity conditions as n goes to infinity.




Figure 2: Tangent spaces and coordinates of a dimension-2 submanifold \mathcal{M} in \mathbb{R}^3

We note that the Edgeworth expansions of the studentized versions of $R_{\theta_0}^{-1}(\hat{\theta}_n)$ and $R_{\theta_0}^{-1}(\hat{\theta}_n^*)$ are readily accessible, owing to the previous results (e.g., [Bhattacharya et al. \(1978\)](#); [Andrews \(2002\)](#)) on Edgeworth expansions for (approximate) M-estimations. Consequently, it is natural to pursue equating $R_{\theta_0}^{-1}(\hat{\theta}_n)$ with $-R_{\hat{\theta}_n}^{-1}(\theta_0)$ and also $R_{\theta_0}^{-1}(\hat{\theta}_n^*) - R_{\theta_0}^{-1}(\hat{\theta}_n)$ with $-R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$, as illustrated in [Figure 2](#). This analysis is conducted in two distinct parts:

- **Identify $R_{\hat{\theta}_n}^{-1}(\theta_0)$ with $R_{\theta_0}^{-1}(\hat{\theta}_n)$.** We relate the coordinate of $\hat{\theta}_n$ under R_{θ_0} to the exponential mapping by letting

$$R_{\theta_0}^{-1}(\hat{\theta}_n) = \pi_{\{dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n) + \Delta_1.$$

Similarly, for $R_{\hat{\theta}_n}(\theta_0)$ one has

$$R_{\hat{\theta}_n}^{-1}(\theta_0) = \pi_{\{dR_{\hat{\theta}_n}\delta_i\}} \circ \text{Log}_{\hat{\theta}_n}(\theta_0) + \Delta'_1.$$

Moreover, the invariance property of parallel transport yields that

$$\pi_{\{dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n) = -\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n} dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\hat{\theta}_n}(\theta_0).$$

Combining the above pieces together implies that

$$\left\| R_{\theta_0}^{-1}(\hat{\theta}_n) + \pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n} dR_{\theta_0}\delta_i\}} \circ \pi_{\{dR_{\hat{\theta}_n}\delta_i\}}^{-1} \circ R_{\hat{\theta}_n}^{-1}(\theta_0) \right\| \leq \|\Delta_1\| + \|\Delta'_1\|. \quad (9)$$

- **Identify $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$ with $R_{\theta_0}^{-1}(\hat{\theta}_n^*) - R_{\theta_0}^{-1}(\hat{\theta}_n)$.** We denote that

$$R_{\theta_0}^{-1}(\hat{\theta}_n^*) = \pi_{\{dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*) + \Delta_2,$$

$$R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) = \pi_{\{dR_{\hat{\theta}_n^*}\delta_i\}} \circ \text{Log}_{\hat{\theta}_n^*}(\hat{\theta}_n) + \Delta_3.$$

Again, leveraging parallel transports, we derive that

$$\pi_{\{dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n^*) = -\pi_{\{T_{\theta_0 \rightarrow \widehat{\theta}_n^*} dR_{\theta_0}\delta_i\}} \circ \text{Log}_{\widehat{\theta}_n^*}(\theta_0).$$

Then the difference between $-\pi_{\{T_{\theta_0 \rightarrow \widehat{\theta}_n^*} dR_{\theta_0}\delta_i\}} \circ \pi_{\{dR_{\theta_0}\delta_i\}}^{-1} \circ R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)$ and $R_{\theta_0}^{-1}(\widehat{\theta}_n^*) - R_{\theta_0}^{-1}(\widehat{\theta}_n)$ could be written as

$$\left\| R_{\theta_0}^{-1}(\widehat{\theta}_n^*) - R_{\theta_0}^{-1}(\widehat{\theta}_n) + \pi_{\{T_{\theta_0 \rightarrow \widehat{\theta}_n^*} dR_{\theta_0}\delta_i\}} \circ \pi_{\{dR_{\theta_0}\delta_i\}}^{-1} \circ R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n) \right\| \leq \Delta_1 + \Delta_2 + \Delta_3 + \widetilde{\Delta}, \quad (10)$$

where $\widetilde{\Delta}$ is defined as

$$\widetilde{\Delta} := \left\| \text{Log}_{\theta_0}(\widehat{\theta}_n^*) - \text{Log}_{\theta_0}(\widehat{\theta}_n) + T_{\widehat{\theta}_n^* \rightarrow \theta_0} \text{Log}_{\widehat{\theta}_n^*}(\widehat{\theta}_n) \right\|.$$

Informally speaking, since the magnitudes of $\text{dist}(\widehat{\theta}_n, \theta_0)$ and $\text{dist}(\widehat{\theta}_n^*, \theta_0)$ are both $O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$, to prove that the distributions of $-\pi_{\{T_{\theta_0 \rightarrow \widehat{\theta}_n} dR_{\theta_0}e_i\}} \circ \pi_{\{dR_{\widehat{\theta}_n}\delta_i\}}^{-1} \circ R_{\widehat{\theta}_n}^{-1}(\theta_0)$ and $-\pi_{\{T_{\theta_0 \rightarrow \widehat{\theta}_n^*} dR_{\theta_0}\delta_i\}} \circ \pi_{\{dR_{\theta_0}\delta_i\}}^{-1} \circ R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)$ align with those of $R_{\theta_0}^{-1}(\widehat{\theta}_n)$ and $R_{\theta_0}^{-1}(\widehat{\theta}_n^*) - R_{\theta_0}^{-1}(\widehat{\theta}_n)$ in the sense of high order, we need to show that the magnitudes of $\Delta_1, \Delta_1', \Delta_2, \Delta_3, \widetilde{\Delta}$ are small enough, to be specific, of order $O(n^{-1} \log n)$.

By the definition of second-order retractions, one can show that $\Delta_1, \Delta_1', \Delta_2$, and Δ_3 satisfy the above requirement. What remains to be shown is the upper bound for $\widetilde{\Delta}$ which relies on the so-called double exponential mapping (Brewin, 2009; Gavrilov, 2006, 2007).

Double Exponential Mapping. Given $x \in \mathcal{M}$, the double exponential mapping is defined as $\text{Log}_x(\text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w)))$ with $v, w \in \text{T}_x \mathcal{M}$. Intuitively, the difference $\text{Log}_x(\text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w))) - v - w \in \text{T}_x \mathcal{M}$ represents a measure of how curvature affects the triangle relation among v , w , and $v + w$. This difference captures the deviation from the expected vector space addition $v + w$ when w is transported from x to $\text{Exp}_x(v)$. Informally, it could be shown that

$$\left\| \text{Log}_x(\text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w))) - v - w \right\| \leq C_1 \|v\| \|w\| + C_2 \|v\| \|w\|^2,$$

where the constants C_1 and C_2 depend on the local curvature around x . A formal result is stated in Lemma 1.

Back to the problem above, we let x be θ_0 , v be $\text{Log}_{\theta_0}(\widehat{\theta}_n^*)$ and w be $T_{\widehat{\theta}_n^* \rightarrow \theta_0} \text{Log}_{\widehat{\theta}_n^*}(\widehat{\theta}_n)$ so that $\text{Log}_x(\text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w)))$ becomes $\text{Log}_{\theta_0}(\widehat{\theta}_n)$, as depicted in Figure 3. Then we arrive at

$$\widetilde{\Delta} = O(n^{-3/2}(\log n)^{\frac{3}{2}}),$$

which satisfies the requirement.

Consistency after Studentization. Studentizing the differences is crucial for achieving high-order consistency between the target distribution and its resampling counterpart. However, the studentization term based on the chart $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\cdot)$ in our theoretical analysis does not directly appear in the algorithmic implementation; in practice, we naturally use the studentization terms centered at $\widehat{\theta}_n$ or $\widehat{\theta}_n^*$. These terms, originating from different perspectives, can be reconciled via Lemma 2, which establishes that the approximation error of a Hessian operator depends on both the distance between the centers of the two charts and the magnitude of the gradient.




Figure 3: Double exponential mapping

Rotation Issue. Finally, the approximations in (9) and (10) depend on certain orthogonal transformations, specifically $\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n} dR_{\theta_0} \delta_i\}} \circ \pi_{\{dR_{\hat{\theta}_n} \delta_i\}}^{-1}$ and $\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n^*} dR_{\theta_0} \delta_i\}} \circ \pi_{\{dR_{\theta_0} \delta_i\}}^{-1}$, which are typically intractable. To address this issue, we adopted the following two approaches.

- **Rotation-Insensitivity of Wald Statistics.** When computing the Wald-type statistic related to $R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^{-1} R_{\hat{\theta}_n}^{-1}(\theta_0)$ and $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \hat{\Sigma}^{*\dagger} R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$, the orthogonal mappings are canceled out regardless of the choice of retraction. Consequently, these statistics are insensitive to potential rotations.
- **Imposing Differentiability on Retraction Family.** For t -statistics, Figure 1 has informed us that the non-differentiability of a chart may lead to irregularities. To mitigate this, we impose certain differentiability conditions and focus on proving that the proposed bootstrap procedure can effectively capture the deviation tendency of a chart.

In summary, the key distinction between Euclidean spaces and general Riemannian manifolds lies in the locality of coordinate expressions, particularly when transiting across different charts. To address this, we bridge gaps between idealized and realizable coordinate systems. The imposition of differentiability conditions and the adoption of rotation-insensitive techniques further address irregularities arising from rotation issues. These efforts culminate in a rigorous framework that achieves high-order consistency between the target distribution and its resampling counterpart, providing a foundation for high-order accurate inference on Riemannian manifolds.

3 Theoretical Distributional Guarantees

In this section, we show that the resampling procedures implemented by the aforementioned algorithms yield high-order accurate approximations of the distributions for the constructed statistic.

3.1 Assumptions

Our approach requires the following regularity conditions to perform Taylor's expansion of the objective function and to ensure the validity of Edgeworth expansion, part of which is inherited from [Bhattacharya et al. \(1978\)](#); [Hall and Horowitz \(1996\)](#) from the Euclidean viewpoint, while

further modifications are introduced to adapt the method to manifold settings. We make the following assumption:

Assumption 1. We let $\{e_i\}$ be an orthogonal basis of $T_{\theta_0}\mathcal{M}$.

A There is an open subset U of \mathbb{R}^m such that for each $\theta \in \mathcal{M}$ one has $\mathbb{P}_\theta[X_1 \in U] = 1$ and $L(\theta, \mathbf{x})$ is 4-times differentiable with respect to θ for $(\theta, \mathbf{x}) \in U \times \mathcal{M}$.

B There exists a compact subset K of \mathcal{M} such that θ_0 is an interior point in K . For each ν with $1 \leq |\nu| \leq 4$, $\mathbb{E}_{\theta_0}[|\nabla^\nu L(\theta_0, X_1)|^{32}] < +\infty$ holds. For every $\theta_1 \in \mathcal{M}$, there exists a function $C(\mathbf{x})$ such that $|\nabla^\nu L(\theta_0, \mathbf{x}) - \bar{\nabla}^\nu L \circ \text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\pi_{\{e_i\}} \circ \text{Log}(\theta_1), \mathbf{x})| \leq C(\mathbf{x}) \cdot \text{dist}(\theta_0, \theta_1)$ for $x \in U$, $\nu \in \mathbb{N}^p$ with $1 \leq |\nu| \leq 4$ and $\mathbb{E}_{\theta_0}[C(X_1)^{16}] \leq \infty$.

C $\mathbb{E}_{\theta_0}[\nabla_{e_i} L(\theta_0, X_1)] = 0$ holds for $i \in [p]$, and $(\mathbb{E}_{\theta_0}[\nabla_{e_i} L(\theta_0, X_1) \nabla_{e_j} L(\theta_0, X_1)])_{i,j \in [p]}$ is non-singular.

D We let \mathbf{Z}_i be a vector that collects $\nabla^\nu L(\theta_0, X_i)$, $\nabla^{\nu_1} L(\theta_0, X_i) \nabla^{\nu_2} L(\theta_0, X_i)$ for $1 \leq |\nu| \leq 3, 1 \leq |\nu_1| + |\nu_2| \leq 2$. The random vector \mathbf{Z}_1 satisfies the Crámer condition, that is,

$$\limsup_{\|\mathbf{t}\|_2 \rightarrow \infty} |\mathbb{E}[e^{i\mathbf{t}^\top \mathbf{Z}_1}]| < 1.$$

3.2 Point Estimation Convergence Guarantees

We first showcase our convergence guarantees for the output sequence of Algorithm 2 based on Assumption 1.

Theorem 1 (Convergence guarantees: Algorithm 1 with full data). *If Assumption 1 holds and the initial estimate $\text{dist}(\theta^{\text{initial}}, \theta_0) \leq c_0$ for some sufficiently small c_0 , then with $\text{burn} = c \log \log n$, where c is a constant determined by c_0 , the objective function L , and $\mathcal{Y} = \mathcal{X}_n$, the output of Algorithm 1 satisfies*

$$\text{dist}(\hat{\theta}_n, \tilde{\theta}_n) = O(n^{-2})$$

with probability at least $1 - O(n^{-1}(\log n)^{-2})$.

Remark 3. *Given a deterministic objective function, the quadratic convergence of Newton's algorithms on Riemannian manifolds has been studied in a sequence of previous work (Smith, 1993, 2014; Absil et al., 2009). Theorem 1 extends the convergence guarantees to the stochastic setting with some additional analysis of the fluctuation of derivatives.*

Remark 4. *As demonstrated in previous works (Absil et al., 2009), quadratic convergence does not depend on the second-order property of the retraction being used. Nonetheless, the second-order retraction is primarily for the sake of high-order asymptotics upon the coordinate calculation.*

Theorem 2 (Convergence guarantees: Algorithm 1 with bootstrapped data). *Suppose that Assumption 1 holds. Then given an initial estimate θ^{initial} satisfying $\text{dist}(\theta^{\text{initial}}, \tilde{\theta}_n) = O(n^{-2})$ and $i \in [b]$, the output of Algorithm 1 with $t = 2$ and $\mathcal{Y} = \mathcal{X}_n^{[i]}$, $\tilde{\theta}_n^{*[i]}$, satisfies*

$$\text{dist}(\hat{\theta}_n^{*[i]}, \tilde{\theta}_n^{[i]}) = O(n^{-2}(\log n)^2)$$

with probability $1 - O(n^{-1}(\log n)^{-2})$ conditional on \mathcal{X}_n , where $\tilde{\theta}_n^{[i]}$ represents a solution to (5) given the resampled dataset $\mathcal{X}_n^{[i]}$.

3.3 High-order Accurate Distributional Guarantees on Riemannian Manifolds

This subsection introduces our main theorem on the distributional guarantees for the approximate M-estimators on Riemannian manifolds.

From the statement of Algorithm 2, it is clear that, conditioning on the original dataset \mathcal{X}_n , $\{\hat{\theta}_n^{*[i]}\}_{i \in [b]}$ are identically distributed and conditionally independent of each other. Therefore, we omit the superscript $[i]$ to denote by $\hat{\theta}_n^*$ the generic resampled output. For the intrinsic t -statistic and the intrinsic Wald-statistic, the main result is as follows.

Theorem 3.⁸ *Suppose Assumption 1 holds.*

1. For every $0 < \xi < \frac{1}{2}$, it holds that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{-\infty < x < \infty} \left| \mathbb{P} \left[R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) \leq x \mid \mathcal{X}_n \right] - \mathbb{P}_{\mathcal{X}_n} \left[R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0) \leq x \right] \right| > n^{-1+\xi} \right] < +\infty, \quad (11)$$

where $\hat{\Sigma}$ and $\check{\Sigma}$ are defined in (7) and (8), respectively.

2. If the retraction mappings R_θ for θ 's in a neighborhood of θ_0 satisfy that the function $dR_\theta^{-1}(T_{\theta_0 \rightarrow \theta}(dR_{\theta_0}(\delta_i)))$ is twice differentiable with respect to θ for every $i \in [p]$, then for an arbitrary $\mathbf{a} \in \mathbb{R}^p$ and every $0 < \xi < \frac{1}{2}$ it holds that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{-\infty < x < \infty} \left| \mathbb{P} \left[\frac{\mathbf{a}^\top R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)}{(\mathbf{a}^\top \hat{\Sigma} \mathbf{a})^{\frac{1}{2}}} \leq x \mid \mathcal{X}_n \right] - \mathbb{P}_{\mathcal{X}_n} \left[\frac{\mathbf{a}^\top R_{\hat{\theta}_n}^{-1}(\theta_0)}{(\mathbf{a}^\top \hat{\Sigma} \mathbf{a})^{\frac{1}{2}}} \leq x \right] \right| > n^{-1+\xi} \right] < +\infty. \quad (12)$$

Moreover, for the extrinsic t -statistic, we establish the following distributional guarantee, which can be viewed as a byproduct of the Euclidean Edgeworth expansion theory:

Proposition 2. *If \mathcal{M} is a submanifold embedded in \mathbb{R}^d and Assumption 1 holds, then for every smooth and Lipschitz-continuous function $f : \mathbb{R}^d \rightarrow \mathbb{R}$ with $\nabla f \circ \text{Exp} \neq \mathbf{0}$ it holds that*

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{-\infty < x < \infty} \left| \mathbb{P} \left[\frac{f(\hat{\theta}_n^*) - f(\hat{\theta}_n)}{(\nabla f(\hat{\theta}_n^*)^\top \hat{\Sigma} \nabla f(\hat{\theta}_n^*))^{\frac{1}{2}}} \leq x \mid \mathcal{X}_n \right] - \mathbb{P}_{\mathcal{X}_n} \left[\frac{f(\hat{\theta}_n) - f(\theta_0)}{(\nabla f(\hat{\theta}_n)^\top \hat{\Sigma} \nabla f(\hat{\theta}_n))^{\frac{1}{2}}} \leq x \right] \right| > n^{-1+\xi} \right] < \infty.$$

Armed with the above distribution consistency results, we are able to control the coverage probability for the confidence regions constructed in Algorithms 3, 4, 5 under $b = \infty$.

Corollary 1. *Suppose Assumption 1 holds.*

1. For every $0 < \xi < \frac{1}{2}$, it holds for Algorithm 3 that

$$\limsup_{b \rightarrow \infty} \left| \mathbb{P}_{\mathcal{X}_n} [\theta_0 \in I_{1-\alpha}^{\text{Wald}}] - (1 - \alpha) \right| \leq O(n^{-1+\xi}).$$

⁸Consistent with the convention in Algorithm 3 and Algorithm 4, we let $R_x^{-1}(y)$ be $\mathbf{0}$ for any y outside the range of R .

2. If the retraction mappings R_θ for θ 's in a neighborhood of θ_0 satisfy that the function $dR_\theta^{-1}(T_{\theta_0 \rightarrow \theta}(dR_{\theta_0}(\delta_i)))$ is twice differentiable with respect to θ for every $i \in [p]$, then for the confidence region in Algorithm 4, an arbitrary $\mathbf{a} \in \mathbb{R}^p$, and every $0 < \xi < \frac{1}{2}$, it holds that

$$\limsup_{b \rightarrow \infty} \left| \mathbb{P}_{\mathcal{X}_n} [\theta_0 \in I_{1-\alpha}^{\text{Intr.t}}] - (1 - \alpha) \right| \leq O(n^{-1+\xi}).$$

3. It holds for Algorithm 5 and every $0 < \xi < \frac{1}{2}$ that

$$\limsup_{b \rightarrow \infty} \left| \mathbb{P}_{\mathcal{X}_n} [\theta_0 \in I_{1-\alpha}^{\text{Extr.t}}] - (1 - \alpha) \right| \leq O(n^{-1+\xi}).$$

In practice, however, resampling the data infinitely many times is not feasible. Nevertheless, by leveraging the Glivenko-Cantelli theorem, we can approximate the ideal case with arbitrary precision by increasing resampling times.

4 Specific Applications

In this section, we showcase our methodology on several commonly encountered Riemannian manifolds that are crucial in various statistical applications. We will demonstrate the implementation of the proposed algorithms across different Riemannian manifolds under a range of statistical scenarios. Section 4.1 provides detailed expressions for Riemannian metrics, gradients, Hessian operators, retractions, and inverse retractions. Section 4.2 specifies the loss functions used in the Gaussian location estimation and Barycenter problems.

Before proceeding, we first introduce a simple method for constructing a smooth mapping $f^{\text{ortho}, \mathbf{A}_0}$ over a neighborhood of a rank- p_2 matrix $\mathbf{A}_0 \in \mathbb{R}^{p_1 \times p_2}$. This mapping maps a rank- p_2 matrix $\mathbf{A} \in \mathbb{R}^{p_1 \times p_2}$ with $p_1 > p_2 > 0$ to an orthonormal matrix $f^{\text{ortho}, \mathbf{A}_0}(\mathbf{A}) \in \text{St}(p_1, p_1 - p_2)$ such that $\mathbf{A}^\top f^{\text{ortho}, \mathbf{A}_0}(\mathbf{A}) = \mathbf{0}_{p_2 \times (p_1 - p_2)}$. The procedure is outlined as follows:

1. Compute the top- p_2 left singular vectors $\mathbf{U}_0 = (\mathbf{u}_{\mathbf{A}_0, 1}, \dots, \mathbf{u}_{\mathbf{A}_0, p_2})$ of \mathbf{A}_0 . Fix an orthonormal matrix $\mathbf{U}_{0, \perp} \in \mathbb{R}^{p_1 \times (p_1 - p_2)}$ such that $\mathbf{A}_0^\top \mathbf{U}_{0, \perp} = \mathbf{0}_{p_2 \times (p_1 - p_2)}$.
2. For any \mathbf{A} in a sufficiently small neighborhood of \mathbf{A}_0 , compute the top- p_2 left singular vectors $\mathbf{U}_{\mathbf{A}} = (\mathbf{u}_{\mathbf{A}, 1}, \dots, \mathbf{u}_{\mathbf{A}, p_2}) \in \mathbb{R}^{p_1 \times p_2}$ of \mathbf{A} .
3. Define $f^{\text{ortho}, \mathbf{A}_0}(\mathbf{A})$ as the top- $(p_2 - p_1)$ left singular vectors of $(\mathbf{I}_{p_1} - \mathbf{U}_{\mathbf{A}} \mathbf{U}_{\mathbf{A}}^\top)(\mathbf{U}_0, \mathbf{U}_{0, \perp}) \text{diag}(p_1, p_1 - 1, \dots, 1)$.

This mapping will be used in the following examples to satisfy the differentiability requirements necessary for constructing intrinsic t -statistics as discussed in Theorem 3.

Furthermore, as one may deduce from the proofs of the results in Section 3, the coordinate calculation does not necessarily require the inverse of a mapping in the adopted retraction; in fact, those results hold if we replace $\{R_x^{-1}\}$ with an atlas⁹ $\{\phi_x\}$ of \mathcal{M} , such that for every x ,

$$\|\phi_x(y) - \pi_{\{\text{d}\phi^{-1}\delta_i\}} \circ \text{Log}_x(y)\|_2 = O(\text{dist}(x, y)^3) \quad (13)$$

for every y in the domain of ϕ_x . In the subsequent discussion, the term "inverse retraction" may refer to either the inverse of a given retraction or a chart in an atlas.

⁹An atlas refers to a collection of charts which covers \mathcal{M} .

4.1 Riemannian Manifold Examples

We begin with a straightforward yet fundamental example.

Example 1 (Sphere Manifold). *Consider the unit sphere \mathbb{S}^{p-1} endowed with the Euclidean metric in \mathbb{R}^p . This induced metric is identical to the quotient metric. For every $\mathbf{x} \in \mathbb{S}^{p-1}$, there exists a canonical isometry between the tangent space $\mathbb{T}_{\mathbf{x}}\mathbb{S}^{p-1}$ at \mathbf{x} and the subspace $\{\mathbf{z} \in \mathbb{R}^p : \mathbf{z}^\top \mathbf{x} = 0\}$ of \mathbb{R}^p , therefore we usually represent the tangent vectors in terms of coordinates in \mathbb{R}^p with a bit abuse of notations hereinafter. For an arbitrary smooth function $f : \mathbb{S}^{p-1} \rightarrow \mathbb{R}$, the gradient ∇f at $\mathbf{x} \in \mathbb{S}^{p-1}$ is written as*

$$\nabla f(\mathbf{x}) = \left(\mathbf{I}_p - \mathbf{x}\mathbf{x}^\top \right) \nabla \bar{f}(\mathbf{x}) \in \{\mathbf{z} \in \mathbb{R}^p : \mathbf{z}^\top \mathbf{x} = 0\} \cong \mathbb{T}_{\mathbf{x}}\mathbb{S}^{p-1},$$

where \bar{f} is a smooth function on a neighborhood U of \mathbf{x} in \mathbb{R}^p whose restriction on $\mathbb{S}^{p-1} \cap U$ is f ¹⁰. The result in [Absil et al. \(2013\)](#) yields the Hessian operator as

$$\nabla^2 f(\mathbf{x})[\boldsymbol{\xi}] = \left(\mathbf{I}_p - \mathbf{x}\mathbf{x}^\top \right) \nabla^2 \bar{f}(\mathbf{x})[\boldsymbol{\xi}] \in \{\mathbf{z} \in \mathbb{R}^p : \mathbf{z}^\top \mathbf{x} = 0\} \cong \mathbb{T}_{\mathbf{x}}\mathbb{S}^{p-1},$$

where we again employ the canonical identification between the tangent space and a subspace in \mathbb{R}^p . Finally, we employ the projection retraction $R_{\mathbf{x}} : \mathbb{R}^{p-1} \rightarrow \mathbb{S}^{p-1}$, defined as

$$R_{\mathbf{x}}(\mathbf{v}) = \frac{\mathbf{x} + f^{\text{ortho}, \hat{\theta}_n}(\mathbf{x})\mathbf{v}}{\left\| \mathbf{x} + f^{\text{ortho}, \hat{\theta}_n}(\mathbf{x})\mathbf{v} \right\|_2}, \quad (14)$$

which has been proved to be a second-order retraction in [Gawlik and Leok \(2018\)](#). The inverse of $R_{\mathbf{x}}$ is given by

$$R_{\mathbf{x}}^{-1}(\mathbf{y}) = f^{\text{ortho}, \hat{\theta}_n}(\mathbf{x})^\top \left(\frac{\mathbf{y}}{\mathbf{y}^\top \mathbf{x}} - \mathbf{x} \right). \quad (15)$$

The role of $\hat{\theta}_n$ here is to ensure the smoothness of the retraction in the neighborhood of the true parameter, leveraging the previously established convergence rate of $\hat{\theta}_n$. By substituting this retraction and its inverse into our framework, we derive the specific algorithms.

A generalization of a sphere manifold is the Stiefel manifold. We consider the canonical metric induced by a quotient formulation ([Tagare, 2011](#); [Gawlik and Leok, 2018](#)).

Example 2 (Stiefel Manifold). *The Stiefel manifold is defined by $\text{St}(p, r) = \{\mathbf{U} \in \mathbb{R}^{p \times r} : \mathbf{U}^\top \mathbf{U} = \mathbf{I}_r\}$ with $p \geq r$. For an arbitrary $\mathbf{U} \in \text{St}(p, r)$, we can fix a perpendicular matrix $\mathbf{U}_\perp \in \mathbb{R}^{p \times (p-r)}$ such that $(\mathbf{U}, \mathbf{U}_\perp)^\top (\mathbf{U}, \mathbf{U}_\perp) = \mathbf{I}_p$ and identify the tangent space at \mathbf{U} to the subspace*

$$\mathbb{T}_{\mathbf{U}}\text{St}(p, r) = \{\mathbf{U}\mathbf{A} + \mathbf{U}_\perp\mathbf{B} : \mathbf{A} \in \mathbb{R}^{r \times r}, \mathbf{A} + \mathbf{A}^\top = \mathbf{0}_{r \times r}, \mathbf{B} \in \mathbb{R}^{(p-r) \times r}\}.$$

We define a metric at \mathbf{U} as $\langle \boldsymbol{\xi}_1, \boldsymbol{\xi}_2 \rangle = \frac{1}{2} \text{tr}(\mathbf{A}_1^\top \mathbf{A}_2) + \text{tr}(\mathbf{B}_1^\top \mathbf{B}_2)$ with $\boldsymbol{\xi}_1 = \mathbf{U}\mathbf{A}_1 + \mathbf{U}_\perp\mathbf{B}_1$, $\boldsymbol{\xi}_2 = \mathbf{U}\mathbf{A}_2 + \mathbf{U}_\perp\mathbf{B}_2 \in \mathbb{T}_{\mathbf{U}}\text{St}(p, r)$. Given a smooth function $f : \text{St}(p, r) \rightarrow \mathbb{R}$, the gradient at \mathbf{U} is written as

$$\nabla f_{\mathbf{U}} = \nabla \bar{f}_{\mathbf{U}} - \mathbf{U} \nabla \bar{f}_{\mathbf{U}}^\top \mathbf{U},$$

where \bar{f} is a smooth extension of f to a local neighborhood of \mathbf{U} in $\mathbb{R}^{p \times r}$ and $\nabla \bar{f}$ is written in a corresponding p -by- r matrix form. According to [Edelman et al. \(1998\)](#), the Hessian operator is expressed as

$$\nabla^2 f_{\mathbf{U}}[\boldsymbol{\xi}] = \nabla^2 f_{\mathbf{U}}[\boldsymbol{\xi}] + \frac{1}{2} (\mathbf{U}\boldsymbol{\xi}^\top \nabla \bar{f}_{\mathbf{U}} + \nabla \bar{f}_{\mathbf{U}} \boldsymbol{\xi}^\top \mathbf{U}) - \frac{1}{2} (\mathbf{I}_p - \mathbf{U}\mathbf{U}^\top) \boldsymbol{\xi} (\nabla \bar{f}_{\mathbf{U}}^\top \mathbf{U} + \mathbf{U}^\top \nabla \bar{f}_{\mathbf{U}}) \in \mathbb{T}_{\mathbf{U}}\text{St}(p, r)$$

¹⁰The existence of such an extension is always guaranteed by Lemma 5.34 in [Lee \(2013\)](#), for example.

for $\xi \in \mathbb{T}_U \text{St}(p, r)$.

To define retraction mappings, we introduce a second-order retraction on $\text{St}(p, r)$ based on Theorem 5 from [Gawlik and Leok \(2018\)](#). We first introduce a mapping $\text{skew}(\mathbf{v})$ that maps a vector $\mathbf{v} \in \mathbb{R}^{r(r-1)/2}$ to a skew-symmetric matrix with its upper-triangle part being \mathbf{v} . Additionally, we let $\text{mat}_{p_1 \times p_2}(\mathbf{v}) \in \mathbb{R}^{p_1 \times p_2}$ be the matricization of a vector $\mathbf{v} \in \mathbb{R}^{p_1 p_2}$. Given $\mathbf{U} \in \text{St}(p, r)$, the second-order retraction R_U is expressed as:

$$R_U(\mathbf{v}) = \mathcal{P} \left(\mathbf{U} - \frac{1}{3} \mathbf{H}^\top \mathbf{H} - \frac{1}{2} \mathbf{U}^\top \mathbf{H} \mathbf{U} \mathbf{H} + \mathbf{H} \left(\mathbf{I} + \frac{1}{2} \mathbf{U}^\top \mathbf{H} \right) \right).$$

for $\mathbf{v} = (\mathbf{v}_1^\top, \mathbf{v}_2^\top)^\top \in \mathbb{R}^{r(r-1)/2 + (p-r)r}$ with $\mathbf{v}_1 \in \mathbb{R}^{r(r-1)/2}$ and $\mathbf{v}_2 \in \mathbb{R}^{(p-r)r}$. Here, the matrix \mathbf{H} is defined as

$$\mathbf{H} = \mathbf{U} \text{skew}(\mathbf{v}_1) + f^{\text{ortho}, \hat{\theta}_n}(\mathbf{U}) \text{mat}_{r \times (p-r)}(\mathbf{v}_2) \quad (16)$$

and the unifying operator $\mathcal{P} : \mathbb{R}^{p \times r} \rightarrow \text{St}(p, r)$ is defined as $\mathcal{P} : \mathbf{X} \mapsto \mathbf{L} \mathbf{R}^\top$ where the singular value decomposition of \mathbf{X} is $\mathbf{X} = \mathbf{L} \mathbf{D} \mathbf{R}^\top$. For the inverse retraction, we recommend using the logarithmic mapping approximation proposed in [Zimmermann \(2017\)](#), in conjunction with the expression in (16).

It is worth noting that, while computationally tractable retractions exist for the examples discussed above, such explicit forms may not be always available for general Riemannian manifolds. Nonetheless, in what follows we will focus on Euclidean submanifolds, as a convenient framework that offers the (inverse) retraction mappings we require.

Euclidean Submanifold. A Euclidean submanifold refers to a submanifold of a Euclidean space, whose importance could be revealed from the following two distinct perspectives:

- **Level Sets of Constant Rank Function.** Suppose that the parameter space \mathcal{M} is determined by $\mathcal{M} = \{\mathbf{x} \in \mathbb{R}^d : g_i(\mathbf{x}) = 0, i = 1, \dots, d - p\}$ with $d > p$. If the mappings g_i are smooth and $\mathbf{g} = (g_1, \dots, g_{d-p})$ has a constant rank $d - p$, by the constant rank level set theorem (Theorem 5.12 in [Lee \(2013\)](#)) we deduce that \mathcal{M} is an embedded submanifold of \mathbb{R}^d .
- **Whitney's Embedding Theorem.** Even though we are working with a smooth manifold with an abstract and complicated differential structure, the Whitney embedding theorem shows that an arbitrary smooth manifold of dimension n always admits a proper embedding into \mathbb{R}^{2n+1} . In view of uncertainty quantification concerning parameter location, it thus suffices to establish inference procedures under an embedding mapping and its Euclidean metric.

For ease of presentation, we use the canonical representation to identify a tangent vector in the tangent space of a submanifold with a vector in \mathbb{R}^d . This identification similarly extends to normal vectors. We consider a Riemannian manifold of dimension p embedded in \mathbb{R}^d . By Theorem 5.8 in [Lee \(2013\)](#), for every $x \in M$ there exists a chart (U, ϕ) for \mathbb{R}^d centered at x such that $\phi(U \cap \mathcal{M}) = \{(x_1, \dots, x_p, 0, \dots, 0) \in U\}$. We denote by $\tilde{\phi}$ the last $d - p$ coordinates of ϕ and then define the projection operator as

$$\mathcal{P}_{\mathbf{x}} := \mathbf{I}_d - \nabla \tilde{\phi} \left(\nabla \tilde{\phi}^\top \nabla \tilde{\phi} \right)^{-1} \nabla \tilde{\phi}^\top,$$

where $\nabla \tilde{\phi} := \left(\frac{\partial}{\partial x_i} \tilde{\phi}_j \right)_{ij} \in \mathbb{R}^{d \times (d-p)}$ with a slight abuse of notation. Given a smooth function \bar{f} on \mathbb{R}^d , we denote the function whose domain is constrained on \mathcal{M} by f . The gradient of f over \mathcal{M} is given by $\nabla f = \mathcal{P}_{\mathbf{x}} \nabla \bar{f}$ where $\nabla \bar{f}$ denotes the gradient of \bar{f} over \mathbb{R}^d . The Hessian operator is given by $\nabla^2 f(\mathbf{x})[\mathbf{v}] = \mathcal{P}_{\mathbf{x}}(\nabla^2 \bar{f}(\mathbf{x})[\mathbf{v}]) + \mathcal{P}_{\mathbf{x}}((\nabla_{\mathbf{v}} \mathcal{P}_{\mathbf{x}})[\nabla \bar{f}])$.

Second-Order inverse retraction of Euclidean Submanifold. Now we are in a position to acquire a viable method to approximate the logarithmic mapping so as to obtain a coordinate representation for Euclidean submanifolds. We refer to the inverse retractions in an atlas that satisfy (13) as the second-order inverse retractions. Inspired by the projection-like retraction proposed in Absil and Malick (2012), we introduce projection-like inverse retractions as a means of approximating logarithmic mappings for general Euclidean submanifolds.

We first introduce the concept of inverse retractor. Recall that two subspaces of dimension p and $(d - p)$ in \mathbb{R}^d are transverse if their direct sum is \mathbb{R}^d .

Definition 2 (Inverse Retractor). *Let \mathcal{M} be a p -dimensional smooth submanifold of a d -dimensional Euclidean space. An inverse retractor on \mathcal{M} is a smooth mapping from a neighborhood $U \times U \in \mathcal{M} \times \mathcal{M}$ to the Grassmannian manifold $\text{Gr}(d-p, \mathbb{R}^d)$, which consist of all $(d-p)$ -dimensional linear subspaces of \mathbb{R}^d , such that the subspace $D(x, x)$ is transverse to $\text{T}_x\mathcal{M}$ for all $x \in U$.*

An inverse retraction L_x at x is called second-order if it satisfies any of the following equivalent conditions:

- $\frac{d^2}{dt^2} L_x(\text{Exp}_x(tv)) = 0$ for every $v \in \text{T}_x\mathcal{M}$;
- $\text{Log}_x(y) = L_x(y) + O(\text{dist}(x, y)^3)$ as $\text{dist}(x, y) \rightarrow 0$.

Given an inverse retractor D and $x_0 \in \mathcal{M}$ with a neighborhood U_{x_0} , we define an inverse retraction centered at x_0 as follows:

$$L_{x_0} : U_{x_0} \rightarrow \text{T}_{x_0}\mathcal{M}; \quad L_{x_0}(x) = v \text{ such that } x_0 + v - x \in D(x_0, x).$$

Building on the work Absil and Malick (2012), which established the second-order consistency of a retraction induced by a retractor (Absil and Malick, 2012, Definition 14), we justify the second-order consistency of L_{x_0} induced by an easy-to-compute inverse retractor in the following proposition, which can be viewed as an inverse counterpart to the projection-like retraction in Absil and Malick (2012).

Proposition 3. *Let L_{x_0} be a mapping defined above with an inverse retractor D satisfying that $D(x_0, y) = \text{N}_{x_0}\mathcal{M}$ for all $x, y \in U$, where $\text{N}_{x_0}\mathcal{M}$ denotes the normal space of \mathcal{M} at x_0 . Then L_{x_0} is a second-order inverse retraction.*

By combining the elements established above, we can proceed to calculate the necessary derivative and coordinate expressions across a broad spectrum of Euclidean submanifolds.

Example 3 (Example: Manifold of Fixed-Rank Matrices). *The fixed-rank matrices manifold is a well-studied matrix manifold (Mishra et al., 2012; Luo et al., 2024b), which could be viewed as a submanifold of the Euclidean space. We consider the matrix manifold $\mathcal{R}_{r, p_1, p_2}$ composed of all rank- r matrices in $\mathbb{R}^{p_1 \times p_2}$, whose dimension is $(p_1 r + p_2 r - r^2)$. For a matrix $\mathbf{X} \in \mathcal{R}_{r, p_1, p_2}$ with the singular value decomposition $\mathbf{X} = \mathbf{U} \begin{bmatrix} \boldsymbol{\Sigma} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix} \mathbf{V}^\top$, every element \mathbf{Z} of $\text{T}_{\mathbf{X}}\mathcal{R}_{r, p_1, p_2}$ could be written as*

$$\mathbf{Z} = \mathbf{U} \begin{bmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{B} & \mathbf{0} \end{bmatrix} \mathbf{V}^\top, \quad \mathbf{U} = [\mathbf{U}_1, \mathbf{U}_2], \quad \mathbf{V} = [\mathbf{V}_1, \mathbf{V}_2],$$

where $\mathbf{U}^\top \mathbf{U} = \mathbf{I}_{p_1}$ and $\mathbf{V}^\top \mathbf{V} = \mathbf{I}_{p_2}$. Note that, if identical nonzero singular values exist in the above decomposition, the choice of $\mathbf{U}_1, \mathbf{V}_1$ becomes non-unique. Moreover, the decomposition of the matrices in the neighborhood of such a matrix is not smooth, potentially leading to the irregular

asymptotic distribution shown in Figure 1. Therefore, we assume the nonzero singular values of \mathbf{X} are distinct hereinafter.

Given a smooth function $f \in \mathcal{R}_{r,p_1,p_2}$ with an extension \bar{f} over $\mathbb{R}^{p_1 \times p_2}$, the gradient of f at \mathbf{X} is given by

$$\nabla f_{\mathbf{X}} = \nabla \bar{f}(\mathbf{X}) - \mathbf{U}_2 \mathbf{U}_2^\top \nabla \bar{f}(\mathbf{X}) \mathbf{V}_2 \mathbf{V}_2^\top.$$

By Section 4.5 in Absil et al. (2013), the Hessian operator of f at \mathbf{X} is given by

$$\nabla^2 f_{\mathbf{X}}[\xi] = \nabla^2 \bar{f}_{\mathbf{X}}[\xi] - \mathbf{U}_2 \mathbf{U}_2^\top \nabla^2 \bar{f}_{\mathbf{X}}[\xi] \mathbf{V}_2 \mathbf{V}_2^\top + \nabla \bar{f}_{\mathbf{X}} \xi^\top (\mathbf{X}^\dagger)^\top + (\mathbf{X}^\dagger)^\top \xi^\top \nabla \bar{f}.$$

By Proposition 4.11 in Absil and Malick (2012), the mapping $R_{\mathbf{X}}$ centered at \mathbf{X} given by

$$R_{\mathbf{X}}(\mathbf{v}) = \mathbf{U} \begin{bmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{B} & \mathbf{B}(\boldsymbol{\Sigma} + \mathbf{A})^{-1} \mathbf{C} \end{bmatrix} \mathbf{V}^\top$$

is a second-order retraction at \mathbf{X} . Here, $\mathbf{v} = (\mathbf{v}_1^\top, \mathbf{v}_2^\top, \mathbf{v}_3^\top)^\top \in \mathbb{R}^{p_1 p_2 - (p_1 - r)(p_2 - r)}$ is transformed into $\mathbf{A} = \text{mat}_{r \times r}(\mathbf{v}_1)$, $\mathbf{B} = \text{mat}_{(p_1 - r) \times r}(\mathbf{v}_2)$, and $\mathbf{C} = \text{mat}_{r \times (p_2 - r)}(\mathbf{v}_3)$. And the matrices \mathbf{U}_2 and \mathbf{V}_2 in the definitions of \mathbf{U} , \mathbf{V} are specified to be $\mathbf{U}_2 = f^{\text{ortho}, \mathbf{U}_{\hat{\theta}_n}}(\mathbf{U}_1)$ and $\mathbf{V}_2 = f^{\text{ortho}, \mathbf{V}_{\hat{\theta}_n}}(\mathbf{V}_1)$, where $\mathbf{U}_{\hat{\theta}_n}$ and $\mathbf{V}_{\hat{\theta}_n}$ refer to the left and right top- r singular vectors of $\hat{\theta}_n \in \mathcal{R}_{r,p_1,p_2}$, respectively.

On the other hand, for any $\mathbf{Y} \in \mathcal{R}_{r,p_1,p_2}$ close to \mathbf{X} enough, the inverse of $R_{\mathbf{X}}$ is

$$R_{\mathbf{X}}^{-1}(\mathbf{Y}) = (\text{vec}(\mathbf{A})^\top, \text{vec}(\mathbf{B})^\top, \text{vec}(\mathbf{C})^\top)^\top,$$

given $\mathbf{Y} = \mathbf{U} \begin{bmatrix} \mathbf{A} & \mathbf{C} \\ \mathbf{B} & \mathbf{D} \end{bmatrix} \mathbf{V}^\top$, where vec denotes the vectorization operator. By Proposition 3, $R_{\mathbf{X}}^{-1}$ is a second-order inverse retraction.

Example 4 (Example: Manifold of Rank-one Tensors). The set $\mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)}$ of Tucker tensors Kasai and Mishra (2016) of fixed rank $\mathbf{r} = (1, \dots, 1)$ in $\mathbb{R}^{p_1 \times p_2 \times \dots \times p_k}$ also form a Riemannian submanifold in the Euclidean space. The tangent space of $\mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)}$ at $\mathbf{X} = x_0 \times_1 \mathbf{u}_1 \times_2 \mathbf{u}_2 \times_3 \dots \times_k \mathbf{u}_k$ can be expressed as

$$\mathbf{T}_{\mathbf{X}} \mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)} = \left\{ \mathbf{v} \mid \mathbf{v} = a_0 \times_{i \in [k]} \mathbf{u}_i + \sum_{i \in [k]} \sum_{j \neq i} 1 \times_{j \neq i} \mathbf{u}_j \times_i \mathbf{u}'_i, \quad \text{with } \mathbf{u}'_i{}^\top \mathbf{u}_i = 0, a_0 \in \mathbb{R} \right\}.$$

Provided with a smooth function f and $\xi \in \mathbf{T}_{\mathbf{X}} \mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)}$, the gradient and the Hessian operator at \mathbf{X} can be expressed as follows (Heidel and Schulz, 2018)

$$\begin{aligned} \nabla f_{\mathbf{X}} &= \sum_{i=1}^k \nabla \bar{f}_{\mathbf{X}} \times_{i \in [k]} \mathbf{u}_i \mathbf{u}_i^\top + \sum_{i \in [k]} \nabla \bar{f}_{\mathbf{X}} \times_{j \neq i} \mathbf{u}_j \mathbf{u}_j^\top \times_i \mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top \\ \nabla^2 f_{\mathbf{X}}[\xi] &= \sum_{i=1}^k \nabla^2 \bar{f}_{\mathbf{X}}[\xi] \times_{i \in [k]} \mathbf{u}_i \mathbf{u}_i^\top + \sum_{i \in [k]} \nabla^2 \bar{f}_{\mathbf{X}}[\xi] \times_{j \neq i} \mathbf{u}_j \mathbf{u}_j^\top \times_i \mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top \\ &\quad + \sum_{i \in [k]} a_0 \times_i \left(\mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top \sum_{k \neq j \neq i} (\nabla \bar{f}_{\mathbf{X}} \times_k \mathbf{u}'_k \times_{k \neq j \neq i} \mathbf{u}_j) \right) \times_{j \neq i} \mathbf{u}_j \in \mathbf{T}_{\mathbf{X}} \mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)}. \end{aligned}$$

We then consider a projective-like retraction, inspired by the estimator proposed in Zhang (2019), defined as follows:

$$R_{\mathbf{X}}(\mathbf{v}) = (x_0 + a_0) \times_{i \in [k]} \left(\mathbf{u}_i + \frac{x_0}{x_0 + a_0} \mathbf{u}'_i \right),$$

where $\mathbf{v} = (a_0, \mathbf{v}_1^\top, \dots, \mathbf{v}_k^\top)^\top$, $\mathbf{u}_i = \mathbf{U}_{i,\perp} \mathbf{v}_i$, and $\mathbf{U}_{i,\perp}$, $i \in [k]$ are specified as $\mathbf{U}_{i,\perp} = f^{\text{ortho}, \mathbf{u}_i, \hat{\theta}_n}(\mathbf{u}_i)$, $i \in [k]$, given $\hat{\theta}_n = a_{0, \hat{\theta}_n} \times_{i \in [k]} \mathbf{u}_{i, \hat{\theta}_n}$. In view of Theorem 22 in Absil and Malick (2012), this retraction is a second-order retraction. We then have the following second-order inverse retraction by Proposition 3

$$R_{\mathbf{X}}^{-1}(\mathbf{Y}) = (a_0, \mathbf{v}_1^\top, \dots, \mathbf{v}_k^\top)^\top,$$

where

$$a_0 = \mathbf{Y} \times_{i \in [k]} \mathbf{u}_i^\top, \quad \mathbf{v}_i = \frac{1}{x_0} \mathbf{U}_{i,\perp}^\top \mathbf{Y} \times_{j \neq i} \mathbf{u}_j^\top.$$

4.2 Statistical Applications

We now turn to two classic statistical problems that naturally align with manifold formulations. Note that all these manifolds can be written as submanifolds in \mathbb{R}^d 's, with the d 's being self-evident from the preceding discussion.

4.2.1 Maximum Likelihood Estimation under Gaussian Noise

In this section, we consider the problem of estimating the location parameter in additive noise models, using the Riemannian manifold parameter spaces discussed earlier. We specifically consider the observation $\mathbf{X}_i = \boldsymbol{\theta}_0 + \boldsymbol{\varepsilon}_i$ for unknown locational parameter $\boldsymbol{\theta}_0 \in \mathcal{M} \subseteq \mathbb{R}^d$ and noise $\boldsymbol{\varepsilon}_i \sim N(0, \mathbf{I}_d)$. The resulting objective function, derived from the maximum likelihood estimation, is given by

$$L(\boldsymbol{\theta}, \mathbf{X}) := \frac{1}{2} \|\mathbf{X} - \boldsymbol{\theta}\|_F^2 \quad \text{with} \quad \boldsymbol{\theta} \in \mathcal{M}.$$

The sample average version L_n is written as

$$L_n(\boldsymbol{\theta}) = \frac{1}{2n} \sum_{i \in [n]} \|\mathbf{X}_i - \boldsymbol{\theta}\|_F^2 \quad \text{with} \quad \boldsymbol{\theta} \in \mathcal{M}.$$

Extending L_n to the whole Euclidean coordinate space, for instance \mathbb{R}^d , we naturally obtain that $\bar{L}_n(\boldsymbol{\theta}) = \frac{1}{2n} \sum_{i \in [n]} \|\mathbf{X}_i - \boldsymbol{\theta}\|_F^2$ with $\boldsymbol{\theta} \in \mathbb{R}^d$.

With the formulations of gradients and Hessian operators in place of Section 4.1, we substitute \bar{L}_n into those expressions and collectively list the expressions in Table 1, where the notations follow the conventions in Section 4.1.

Parameter Space	Gradient $\nabla L_n(\boldsymbol{\theta})$	Hessian $\nabla^2 L_n(\boldsymbol{\theta})[\boldsymbol{\xi}]$
Sphere Manifold \mathbb{S}^{p-1}	$(\mathbf{I}_p - \mathbf{x}\mathbf{x}^\top)(\boldsymbol{\theta} - \bar{\mathbf{X}}_n)$	$(\mathbf{I}_p - \mathbf{x}\mathbf{x}^\top)\boldsymbol{\xi}$
Stiefel Manifold $\text{St}(p, r)$	$(\boldsymbol{\theta} - \bar{\mathbf{X}}_n) - \mathbf{U}(\boldsymbol{\theta} - \bar{\mathbf{X}}_n)^\top \mathbf{U}$	$\boldsymbol{\xi} + \frac{1}{2}(\mathbf{U}\boldsymbol{\xi}^\top(\boldsymbol{\theta} - \bar{\mathbf{X}}_n) + (\boldsymbol{\theta} - \bar{\mathbf{X}}_n)\boldsymbol{\xi}^\top \mathbf{U}) - \frac{1}{2}(\mathbf{I}_p - \mathbf{U}\mathbf{U}^\top)\boldsymbol{\xi}((\boldsymbol{\theta} - \bar{\mathbf{X}}_n)^\top \mathbf{U} + \mathbf{U}^\top(\boldsymbol{\theta} - \bar{\mathbf{X}}_n))$
fixed-rank matrices manifold $\mathcal{R}_{r, p_1, p_2}$	$(\boldsymbol{\theta} - \bar{\mathbf{X}}_n) - \mathbf{U}_2 \mathbf{U}_2^\top (\boldsymbol{\theta} - \bar{\mathbf{X}}_n) \mathbf{V}_2 \mathbf{V}_2^\top$	$\boldsymbol{\xi} - \mathbf{U}_2 \mathbf{U}_2^\top \boldsymbol{\xi} \mathbf{V}_2 \mathbf{V}_2^\top + (\boldsymbol{\theta} - \bar{\mathbf{X}}_n) \boldsymbol{\xi}^\top (\mathbf{X}^\dagger)^\top + (\mathbf{X}^\dagger)^\top \boldsymbol{\xi}^\top (\boldsymbol{\theta} - \bar{\mathbf{X}}_n)$
Rank-1 Tensors Manifold $\mathcal{M}_{p_1, p_2, \dots, p_k}^{(1)}$	$\sum_{i=1}^k (\boldsymbol{\theta} - \bar{\mathbf{X}}_n) \times_{i \in [k]} \mathbf{u}_i \mathbf{u}_i^\top + \sum_{i \in [k]} (\boldsymbol{\theta} - \bar{\mathbf{X}}_n) \times_{j \neq i} \mathbf{u}_j \mathbf{u}_j^\top \times_i \mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top$	$\sum_{i=1}^k \boldsymbol{\xi} \times_{i \in [k]} \mathbf{u}_i \mathbf{u}_i^\top + \sum_{i \in [k]} \boldsymbol{\xi} \times_{j \neq i} \mathbf{u}_j \mathbf{u}_j^\top \times_i \mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top + \sum_{i \in [k]} a_0 \times_i (\mathbf{U}_{i,\perp} \mathbf{U}_{i,\perp}^\top \sum_{k \neq i} ((\boldsymbol{\theta} - \bar{\mathbf{X}}_n) \times_k \mathbf{u}'_k \times_{k \neq j \neq i} \mathbf{u}_j)) \times_{j \neq i} \mathbf{u}_j$

Table 1: Gradient and Hessian Expressions for Different Manifolds

4.2.2 Barycenter Problem

Barycenter problems, serving as a specific case of M-estimations on manifolds, attracted much attention in the past decades (Fréchet, 1948; Karcher, 1977; Bhattacharya and Patrangenaru, 2003, 2005; Pennec, 2019). To be specific, given n manifold-valued samples $\{x_i\}$ independently drawn from a distribution on a Riemannian manifold \mathcal{M} , the barycenter is defined as the minimizer $x \in \mathcal{M}$ of

$$L_n(x) = \frac{1}{2n} \sum_{i \in [n]} \text{dist}^2(x_i, x).$$

While previous studies have primarily examined the geometrical and statistical properties of the barycenter, we revisit the barycenter problem from an optimization perspective. Algorithm 2 together with the algorithms in Section 2.3.1 yields a computationally feasible approach to construct confidence regions for the Barycenter with high accuracy.

Next, we focus on the sphere case $\mathcal{M} = \mathbb{S}^{p-1}$. According to Pennec (2018), the expressions of gradient and Hessian operator at \mathbf{x} are given by

$$\begin{aligned} \nabla L_n(\mathbf{x}) &= \frac{-\sum_{i \in [n]} \text{Log}_{\mathbf{x}} \mathbf{x}_i}{n} = -\frac{1}{n} \sum_{i \in [n]} \frac{\arccos(\mathbf{x}^\top \mathbf{x}_i)}{\sqrt{1 - (\mathbf{x}^\top \mathbf{x}_i)^2}} (\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top) \mathbf{x}_i, \\ \nabla^2 L_n(\mathbf{x})[\mathbf{v}] &= \frac{1}{n} \sum_{i \in [n]} \frac{(\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top) \mathbf{x}_i \mathbf{x}_i^\top (\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top) \mathbf{v}}{(1 - (\mathbf{x}^\top \mathbf{x}_i)^2)} \\ &+ \frac{1}{n} \sum_{i \in [n]} \frac{\arccos(\mathbf{x}_i^\top \mathbf{x})}{\sqrt{1 - (\mathbf{x}^\top \mathbf{x}_i)^2}} (\mathbf{x}_i^\top \mathbf{x}) \left(\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top - \frac{(\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top) \mathbf{x}_i \mathbf{x}_i^\top (\mathbf{I}_p - \mathbf{x} \mathbf{x}^\top)}{(1 - (\mathbf{x}^\top \mathbf{x}_i)^2)} \right) \mathbf{v} \end{aligned}$$

for $\mathbf{v} \in \text{T}_{\mathbf{x}} \mathbb{S}^{p-1}$. As outlined in Example 1, we employ the second-order retraction (14) in the optimization procedure and use its inverse (15) to calculate the coordinate statistics.

5 Simulation Studies

In this section, we validate our methods and theoretical results through a range of numerical simulations.

Simulation Settings Overview We set out by providing an overview of all simulation settings. The first four simulations proceeded in an additive Gaussian noise framework as discussed in Section 4.2.1 and the last simulation investigates the application of our methods to the Barycenter problem on a sphere manifold. The precise parameter setups are as follows:

- **Setting 1:** \mathbb{S}^2 . We consider the sphere manifold \mathbb{S}^2 embedded in \mathbb{R}^3 and set the ground-truth parameter as $\boldsymbol{\theta}_0 = (0, 1, 0)^\top$.

- **Setting 2:** $\text{St}(4, 2)$. We set the ground-truth parameter to be $\boldsymbol{\theta}_0 = \begin{pmatrix} 1/\sqrt{2} & 1/\sqrt{2} \\ 1/\sqrt{2} & -1/\sqrt{2} \\ 0 & 0 \\ 0 & 0 \end{pmatrix} \in \text{St}(4, 2)$.

- **Setting 3:** $\mathcal{R}_{2,4,4}$. We consider the fixed-rank matrices manifold to be the collection of 4-by-4 matrices with rank 2 and let the ground truth $\boldsymbol{\Theta}_0$ to be the matrix with all zero entries except for the first two diagonal entries, which are set to 5 and 1, respectively.

- **Setting 4:** $\mathcal{M}_{3,3,3}^{(1)}$. We consider the rank-one tensors manifold embedded in $\mathbb{R}^{3 \times 3 \times 3}$. The ground truth parameter is set to be $\Theta = \mathbf{u}_1 \otimes \mathbf{u}_1 \otimes \mathbf{u}_1$ with $\mathbf{u}_1 = (1, 0, 0)^\top$.

Using the gradient and Hessian formulas in Section 4.2, we implemented the estimating and resampling procedures in Algorithm 2. We validated the distributional guarantees for these procedures as detailed in Section 2.2 and Algorithms 3, 4, 5. We repeated the bootstrapping for the resampled estimators b times, where b is taken to be $1000 \times n$, with n being the corresponding sample size. Increasing the number of bootstrap iterations with n helps reduce the noise in the empirical cumulative distribution function (CDF), thereby better revealing differences from the true limiting distribution. We conduct four simulations as follows:

1. **Type-I Error of Tests.** To evaluate the type-I error control as discussed in Section 2.2, we conduct an illustrative simulation under Setting 1; the samples are independently drawn from $\mathcal{N}(\theta_0, 2\mathbf{I}_3)$ and we let $\theta_1 = \theta_0 = (0, 1, 0)^\top$. The chart ϕ in Section 2.2 is defined as $R_{\theta_1}^{-1}(\mathbf{y}) = \frac{\mathbf{U}_\perp \mathbf{y}}{(\mathbf{U}_\perp \mathbf{y})^\top \theta_1} - \theta_1$, where $\mathbf{U}_\perp \in \mathbb{R}^{3 \times 2}$ is an orthonormal matrix whose columns are perpendicular to θ_1 . For sample sizes $n = \{40, 80, 160\}$, we performed 30 repeated experiments and summarize the observed Type-I error in Table 2 for the t -statistic based on $\phi(\theta)_1$ and the Wald statistic. The results in Table 2 demonstrate that our method provides stable approximations of the targeted Type-I error as the sample size increases.
2. **Convergence Rate of Intrinsic Statistics.** This experiment is designed to validate our theoretical findings on the statistics that are intrinsic to the corresponding Riemannian manifolds, namely, the Wald statistic and the intrinsic t -statistic, by comparing them with benchmark statistics. To assess the approximation accuracy to the cumulant distribution function of a targeted quantity \hat{T}_n denoted by $F_{\hat{T}_n}$, we consider the following measure:

$$\text{Error}(\hat{F}_{\hat{T}_n}) = \sup_{x \in S} \left| \hat{F}_{\hat{T}_n}(x) - F_{\hat{T}_n}(x) \right|,$$

where $\hat{F}_{\hat{T}_n}$ is an approximation to the CDF $F_{\hat{T}_n}$ of \hat{T}_n . For the Wald statistic, we let $\hat{T}_n = R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^{-1} R_{\hat{\theta}_n}^{-1}(\theta_0)$ and $S = [8]$, whereas for the intrinsic t statistic, we let $\hat{T}_n = R_{\hat{\theta}_n}^{-1}(\theta_0)_1 / (\hat{\Sigma}_{1,1})^{\frac{1}{2}}$ and $S = \{0.2 \cdot s : s \in \{-10, -9, \dots, 9, 10\}\}$. For the choice of $\hat{F}_{\hat{T}_n}$, we propose using the Monte-Carlo approximation of the CDF of $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^{-1} R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$ for the Wald statistic, and $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_1 / (\check{\Sigma}_{1,1})^{\frac{1}{2}}$ for the intrinsic t statistic. For comparison purposes, we also consider alternative options for $\hat{F}_{\hat{T}_n}$ including the empirical CDF of the non-studentized version $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \hat{\Sigma}^{-1} R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$ and $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_1 / (\hat{\Sigma}_{1,1})^{\frac{1}{2}}$, as well as the CDFs of the normal/chi-squared distributions. Then, the $\text{Error}(\hat{F}_{\hat{T}_n})$ is scaled by the logarithm function and the square root of n , respectively. In this simulation, we are interested in the distributional consistency of Wald statistics and the intrinsic t -statistics, in comparison with the other benchmark methods. Figure 4 presents the average, maximum, and minimum scaled approximation errors using different methods over 30 epochs. The first and third rows of Figure 4 indicate the (almost) linear relation between $\text{Log}(\text{Error})$ and the sample size, with our proposed statistics often exhibiting the steepest slope. Meanwhile, the second and fourth rows of Figure 4 corroborate our theoretical expectation from Theorem 3 that $n^{\frac{1}{2}} \cdot \text{Error}$ decreases with increasing sample size, with our methods consistently outperforming or closely matching the benchmarks.

Hypothesis	$H_0 : \boldsymbol{\theta}_1 = \boldsymbol{\theta}_0$ v.s. $H_1 : \boldsymbol{\theta}_1 \neq \boldsymbol{\theta}_0$					
Statistic	Test Based on T_1			Test Based on W		
1–Signif. Level	0.9	0.95	0.975	0.9	0.95	0.975
$n = 40$	0.9231075	0.9705825	0.9883875	0.9607000	0.9864775	0.9955950
$n = 80$	0.9075400	0.9581725	0.9818088	0.9404200	0.9781425	0.9923438
$n = 160$	0.9057363	0.9564100	0.9807069	0.9210481	0.9667800	0.9867062

Table 2: Table: 1 - (Type-I Error) of Hypothesis Testing on Sphere.

- Convergence Rate of Extrinsic t Statistics.** Moreover, we conducted further simulations to examine the extrinsic t -statistic for the fixed-rank matrices manifold. The simulations follow the same setup and measurement approach as described earlier for the fixed-rank matrices manifold. Our focus is on a single coordinate of the fixed-rank matrix, specifically on \widehat{T}_n , which refers to the studentized version of $((\boldsymbol{\theta}_0)_{1,1} - (\widehat{\boldsymbol{\theta}}_n)_{1,1})$. The resulting errors using Algorithm 5 are shown in Figure 5. From the log-scaled plot and the $n^{\frac{1}{2}}$ -scaled plot, it shows that the distributional deviation $\text{Error}(\widehat{F})$ of the extrinsic t -statistic is of the order $o(n^{-\frac{1}{2}})$ employing Algorithm 5, thereby supporting our theoretical results.
- Barycenter Problem.** Lastly, we aim to demonstrate the performance of our methods on Barycenter problems discussed in Section 4.2.2. We consider the sphere \mathbb{S}^2 restricted to the northern hemisphere ($x_2 \geq 0$). We independently generate n samples $\{X_i\}_{i=1}^n = \{\cos \theta_i \sin \phi_i, \cos \phi_i, \sin \theta_i \sin \phi_i\}$, where θ_i is independently drawn from $\text{Unif}([0, 2\pi])$ and ϕ_i is drawn from the beta distribution $\text{Beta}(2, 2)$. We then compute the barycenter using Algorithm 2. Given that the population barycenter in this case is $(0, 1, 0)$, we analyze the distributional consistency of the corresponding statistics over 30 repetitions, with results summarized in Figure 6. These results reveal that the Wald statistic, intrinsic t -statistic, and extrinsic t -statistic all achieve a convergence rate of $o(n^{-\frac{1}{2}})$ in terms of the CDF, aligning with our theoretical predictions.

In summary, the simulations validate the effectiveness and theoretical soundness of our proposed methods across diverse statistical challenges, demonstrating superior performance and consistency compared to benchmark approaches.

6 Conclusion and Discussions

In this study, we presented a novel framework for high-order accurate inference on Riemannian manifolds, addressing the challenges faced in statistical inference when the parameter spaces are not Euclidean. This framework encompasses the development of computationally friendly bootstrapping algorithms, high-order asymptotics for specific statistics under curvature effect, and the exploration of a comprehensive treatment across various modern statistical settings, including spheres, the Stiefel manifold, Euclidean submanifolds, and specific applications like fixed-rank matrices and rank-one tensors manifolds. Compared with existing works on high-order asymptotics of specific statistics, this work elucidates some essential ingredients for treating nonconvex problems in the scope of manifold language, especially shedding light on the coordinate transformation issue.

Looking ahead, multiple avenues for future research emerge. Firstly, regarding high-order asymptotics, the symmetric case achieves higher accuracy than the asymmetric case. However, this advantage diminishes in the manifold setting due to the triangle error introduced by the dou-




Figure 4: Scaled Cumulative Distribution Function Difference. column 1: Stiefel Manifold; column 2: fixed-rank matrices manifold; column 3: Rank-One Tensor Manifold; row 1: log-scaled Wald Statistic; row 2: \sqrt{n} -scaled Wald Statistic; row 3: log-scaled intrinsic t -Statistic; row 4: \sqrt{n} -scaled intrinsic t -Statistic; **Black**: resampled statistic (our method); **Red**: non-studentized resampled statistic; **Blue**: chi-square / standard normal distribution.




Figure 5: Scaled Cumulative Distribution Function Difference for Extrinsic t -statistic for fixed-rank matrices manifold. Black: the empirical distribution of the resampled statistic; Red: the empirical distribution of the non-studentized resampled statistic; Blue: the distribution of the standard normal distribution.




Figure 6: Cumulative Distribution Function Values for Sphere Barycenter Problem. row 1: Log-Scaled Error; row 2: \sqrt{n} -Scaled Error; Column 1: Intrinsic t ; column 2: Extrinsic t ; column 3: Wald.

ble exponential mapping. Therefore, a correction accounting for local curvature might be needed to achieve the same accuracy as in Euclidean spaces.

Our current focus is on the fixed manifold case, where the sample size n tends to infinity but dimension p remains fixed. When p increases along with n , the necessary conditions on p and the loss function L become significantly more complex and stringent, and the bootstrap may fail as indicated in [El Karoui and Purdom \(2018\)](#). Therefore, this study concentrates on the fixed-dimensional case with $n \rightarrow \infty$. Given the recent advances in high-dimensional statistical theory, it would be of broad interest to investigate the curvature effect when the dimensions of a sequence of manifolds also grow with the sample size.

Moreover, our study highlights an additional application of the double exponential mapping, previously employed to investigate curvature effects in the barycenter problem ([Pennec, 2019](#)). Our technique may offer insights to various phenomena in Riemannian optimization with higher precision.

References

- Absil, P.-A., Mahony, R., and Sepulchre, R. (2009). [Optimization algorithms on matrix manifolds](#). Princeton University Press.
- Absil, P.-A., Mahony, R., and Trunpf, J. (2013). An extrinsic look at the riemannian hessian. In [International conference on geometric science of information](#), pages 361–368. Springer.
- Absil, P.-A. and Malick, J. (2012). Projection-like retractions on matrix manifolds. [SIAM Journal on Optimization](#), 22(1):135–158.
- Adler, R. L., Dedieu, J.-P., Margulies, J. Y., Martens, M., and Shub, M. (2002). Newton’s method on riemannian manifolds and a geometric model for the human spine. [IMA Journal of Numerical Analysis](#), 22(3):359–390.
- Andrews, D. W. (2002). Higher-order improvements of a computationally attractive k-step bootstrap for extremum estimators. [Econometrica](#), 70(1):119–162.
- Babu, G. J. and Singh, K. (1984). On one term edgeworth correction by efron’s bootstrap. [Sankhyā: The Indian Journal of Statistics, Series A](#), pages 219–232.
- Bhattacharya, R. (1987). Some aspects of edgeworth expansions in probability and statistics. [New Perspectives in Theoretical and Applied Statistics](#). New York: Wiley, pages 157–171.
- Bhattacharya, R. and Patrangenaru, V. (2003). Large sample theory of intrinsic and extrinsic sample means on manifolds. [The Annals of Statistics](#), 31(1):1–29.
- Bhattacharya, R. and Patrangenaru, V. (2005). Large sample theory of intrinsic and extrinsic sample means on manifolds—ii. [The Annals of Statistics](#), 33(3):1225–1259.
- Bhattacharya, R. N., Ghosh, J. K., et al. (1978). On the validity of the formal edgeworth expansion. [Ann. Statist](#), 6(2):434–451.
- Bickel, P., Götze, F., and Van Zwet, W. (1986). The edgeworth expansion for u-statistics of degree two. [The Annals of Statistics](#), pages 1463–1484.
- Boumal, N., Absil, P.-A., and Cartis, C. (2019). Global rates of convergence for nonconvex optimization on manifolds. [IMA Journal of Numerical Analysis](#), 39(1):1–33.

- Brewin, L. (2009). Riemann normal coordinate expansions using cadabra. Classical and Quantum Gravity, 26(17):175017.
- Brunel, V.-E. (2023). Geodesically convex m -estimation in metric spaces. In The Thirty Sixth Annual Conference on Learning Theory, pages 2188–2210. PMLR.
- Callaert, H., Janssen, P., and Veraverbeke, N. (1980). An edgeworth expansion for u -statistics. The Annals of Statistics, pages 299–312.
- Chandra, T. K. and Ghosh, J. (1979). Valid asymptotic expansions for the likelihood ratio statistic and other perturbed chi-square variables. Sankhyā: The Indian Journal of Statistics, Series A, pages 22–47.
- Chen, S., Ma, S., Man-Cho So, A., and Zhang, T. (2020). Proximal gradient method for nonsmooth optimization over the stiefel manifold. SIAM Journal on Optimization, 30(1):210–239.
- Chen, Y., Fan, J., Ma, C., and Yan, Y. (2019). Inference and uncertainty quantification for noisy matrix completion. Proceedings of the National Academy of Sciences, 116(46):22931–22937.
- Davidson, R. and MacKinnon, J. G. (1999). Bootstrap testing in nonlinear models. International Economic Review, 40(2):487–508.
- Denker, M. and Bhattacharya, R. (1990). Asymptotic Statistics, volume 14. Springer Science & Business Media.
- Dennis Jr, J. E. and Schnabel, R. B. (1996). Numerical methods for unconstrained optimization and nonlinear equations. SIAM.
- Durrett, R. (2019). Probability: theory and examples. Cambridge university press.
- Edelman, A., Arias, T. A., and Smith, S. T. (1998). The geometry of algorithms with orthogonality constraints. SIAM journal on Matrix Analysis and Applications, 20(2):303–353.
- Efron, B. (1975). Defining the curvature of a statistical problem (with applications to second order efficiency). The Annals of Statistics, pages 1189–1242.
- Efron, B. (1978). The geometry of exponential families. The Annals of Statistics, pages 362–376.
- El Karoui, N. and Purdom, E. (2018). Can we trust the bootstrap in high-dimensions? the case of linear models. Journal of Machine Learning Research, 19(5):1–66.
- Eltzner, B. and Huckemann, S. F. (2019). A smeary central limit theorem for manifolds with application to high-dimensional spheres. The Annals of Statistics, 47(6):3360–3381.
- Fisher, R. A. (1922). On the mathematical foundations of theoretical statistics. Philosophical transactions of the Royal Society of London. Series A, containing papers of a mathematical or physical character, 222(594-604):309–368.
- Fréchet, M. (1948). Les éléments aléatoires de nature quelconque dans un espace distancié. In Annales de l’institut Henri Poincaré, volume 10, pages 215–310.
- Gavrilov, A. V. (2006). Algebraic properties of covariant derivative and composition of exponential maps. Matematicheskie Trudy, 9(1):3–20.

- Gavrilov, A. V. (2007). The double exponential map and covariant derivation. Siberian Mathematical Journal, 48:56–61.
- Gawlik, E. S. and Leok, M. (2018). High-order retractions on matrix manifolds using projected polynomials. SIAM Journal on Matrix Analysis and Applications, 39(2):801–828.
- Hall, P. (2013). The bootstrap and Edgeworth expansion. Springer Science & Business Media.
- Hall, P. and Heyde, C. C. (2014). Martingale limit theory and its application. Academic press.
- Hall, P. and Horowitz, J. L. (1996). Bootstrap critical values for tests based on generalized-method-of-moments estimators. Econometrica: Journal of the Econometric Society, pages 891–916.
- Heidel, G. and Schulz, V. (2018). A riemannian trust-region method for low-rank tensor completion. Numerical Linear Algebra with Applications, 25(6):e2175.
- Helmke, U. and Hüper, K. (2000). A jacobi-type method for computing balanced realizations. Systems & Control Letters, 39(1):19–30.
- Huang, W., Absil, P.-A., and Gallivan, K. A. (2016). A riemannian bfgs method for nonconvex optimization problems. In Numerical Mathematics and Advanced Applications ENUMATH 2015, pages 627–634. Springer.
- Huper, K. and Trumpf, J. (2004). Newton-like methods for numerical optimization on manifolds. In Conference Record of the Thirty-Eighth Asilomar Conference on Signals, Systems and Computers, 2004., volume 1, pages 136–139. IEEE.
- James, G. (1958). On moments and cumulants of systems of statistics. Sankhyā: The Indian Journal of Statistics, pages 1–30.
- Jost, J. (2008). Riemannian geometry and geometric analysis, volume 42005. Springer.
- Karcher, H. (1977). Riemannian center of mass and mollifier smoothing. Communications on pure and applied mathematics, 30(5):509–541.
- Kasai, H. and Mishra, B. (2016). Low-rank tensor completion: a riemannian manifold preconditioning approach. In International conference on machine learning, pages 1012–1021. PMLR.
- Lai, T. L. and Wang, J. Q. (1993). Edgeworth expansions for symmetric statistics with applications to bootstrap methods. Statistica Sinica, pages 517–542.
- Lee, J. M. (2013). Introduction to Smooth Manifolds. Springer.
- Lee, J. M. (2018). Introduction to Riemannian manifolds, volume 176. Springer.
- Li, X., Chen, S., Deng, Z., Qu, Q., Zhu, Z., and Man-Cho So, A. (2021). Weakly convex optimization over stiefel manifold using riemannian subgradient-type methods. SIAM Journal on Optimization, 31(3):1605–1634.
- Luo, Y., Huang, W., Li, X., and Zhang, A. (2024a). Recursive importance sketching for rank constrained least squares: Algorithms and high-order convergence. Operations Research, 72(1):237–256.

- Luo, Y., Li, X., and Zhang, A. R. (2024b). On geometric connections of embedded and quotient geometries in riemannian fixed-rank matrix optimization. Mathematics of Operations Research, 49(2):782–825.
- Luo, Y. and Trillos, N. G. (2022). Nonconvex matrix factorization is geodesically convex: Global landscape analysis for fixed-rank matrix optimization from a riemannian perspective. arXiv preprint arXiv:2209.15130.
- Luo, Y. and Zhang, A. R. (2023). Low-rank tensor estimation via riemannian gauss-newton: Statistical optimality and second-order convergence. The Journal of Machine Learning Research, 24(1):18274–18321.
- Mishra, B., Apuroop, K. A., and Sepulchre, R. (2012). A Riemannian geometry for low-rank matrix completion. arXiv preprint arXiv:1211.1550.
- Mishra, B., Meyer, G., Bonnabel, S., and Sepulchre, R. (2014). Fixed-rank matrix factorizations and Riemannian low-rank optimization. Computational Statistics, 29(3-4):591–621.
- Newey, W. K. and McFadden, D. (1994). Large sample estimation and hypothesis testing. Handbook of econometrics, 4:2111–2245.
- Pennec, X. (2018). Barycentric subspace analysis on manifolds. The Annals of Statistics, 46(6A):2711 – 2746.
- Pennec, X. (2019). Curvature effects on the empirical mean in riemannian and affine manifolds: a non-asymptotic high concentration expansion in the small-sample regime. arXiv preprint arXiv: Arxiv-1906.07418.
- Plumbley, M. D. (2005). Geometrical methods for non-negative ica: Manifolds, lie groups and toral subalgebras. Neurocomputing, 67:161–197.
- Shao, M., Xia, D., Zhang, Y., Wu, Q., and Chen, S. (2022). Higher-order accurate two-sample network inference and network hashing. arXiv preprint arXiv: 2208.07573.
- Smith, S. T. (1993). Geometric optimization methods for adaptive filtering. Harvard University.
- Smith, S. T. (2014). Optimization techniques on riemannian manifolds. arXiv preprint arXiv:1407.5965.
- Tagare, H. D. (2011). Notes on optimization on stiefel manifolds. Yale University, New Haven.
- Van der Vaart, A. W. (2000). Asymptotic statistics, volume 3. Cambridge university press.
- Vandereycken, B. (2013). Low-rank matrix completion by Riemannian optimization. SIAM Journal on Optimization, 23(2):1214–1236.
- von Bahr, B. (1967). On the central limit theorem in r k. Arkiv för matematik, 7(1):61–69.
- Wald, A. (1949). Note on the consistency of the maximum likelihood estimate. The Annals of Mathematical Statistics, 20(4):595–601.
- Xia, D. and Yuan, M. (2021). Statistical inferences of linear forms for noisy matrix completion. Journal of the Royal Statistical Society Series B: Statistical Methodology, 83(1):58–77.

- Xia, D., Zhang, A. R., and Zhou, Y. (2022). Inference for low-rank tensors—no need to debias. The Annals of Statistics, 50(2):1220–1245.
- Xie, F. and Zhang, Y. (2024). Higher-order entrywise eigenvectors analysis of low-rank random matrices: Bias correction, edgeworth expansion, and bootstrap. arXiv preprint arXiv: 2401.15033.
- Zhang, A. (2019). Cross. The Annals of Statistics, 47(2):936–964.
- Zhang, Y. and Xia, D. (2022). Edgeworth expansions for network moments. The Annals of Statistics, 50(2):726–753.
- Zimmermann, R. (2017). A matrix-algebraic algorithm for the riemannian logarithm on the stiefel manifold under the canonical metric. SIAM Journal on Matrix Analysis and Applications, 38(2):322–342.

Appendices

A Proof of Main Result

A.1 Proof of Theorem 3

Instead of getting entangled in the complexities across various charts, our strategy throughout the paper is to concentrate on a fixed normal coordinate chart centered at θ_0 to simplify the discussions. Our primary analytical tool is the Taylor expansion on the coordinate chart $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\cdot)$, where $\{e_i\}$ is a normal basis of $\text{T}_{\theta_0}\mathcal{M}$ to be specified later, along with the Edgeworth expansion results developed in the Euclidean case.

Firstly, to ensure that the geodesics and exponential mappings are well-defined, we restrict the region to a geodesically complete subset of \mathcal{M} . By Theorem 6.17 in Lee (2018), there exists a positive ρ such that for arbitrary $\theta \in \text{Exp}_{\theta_0}(B(0, \rho))$, Log_{θ} defined on $\text{Exp}_{\theta_0}(B(0, \rho))$ is a diffeomorphism onto its codomain in $\text{T}_{\theta}\mathcal{M}$. We denote this region by N in the following context.

In Algorithm 2, each resampling process, including Newton's step on resampled datasets, is performed independently. Consequently, we treat $\hat{\theta}_n^*$ as a random variable, and Algorithm 2 generates b independent replications of $\hat{\theta}_n^*$ conditioned on the dataset \mathcal{X}_n . We denote the generic resampled dataset by $\mathcal{X}_n^* := \{X_i^*\}_{i \in [n]}$ and the corresponding objective function by $L_n^*(\cdot) := \sum_{i \in [n]} L(\cdot, X_i^*)/n$.

To simplify notations, we define the orthonormal basis induced by R_{θ_0} at θ_0 as $\{e_i\}_{i \in [p]} \subset \text{T}_{\theta_0}\mathcal{M}$, where $e_i = \text{d}R_{\theta_0}\delta_i$. We use $L^{\text{Retr},\theta}(\cdot, x)$ to denote the function $L(R_{\theta}(\cdot), x)$. Further, given $\theta \in \mathcal{M}$ and a basis $\{e_i\}$ of $\text{T}_{\theta}\mathcal{M}$, we denote the function $L(\text{Exp}_{\theta} \circ \pi_{\{e_i\}}^{-1}(\cdot), x)$ by $L^{\text{Exp},\theta,\{e_i\}}(\cdot, x)$ for every $\theta \in \mathcal{M}$. Similarly, we denote by $L_n^{\text{Exp},\theta,\{e_i\}}(\cdot)$, $L_n^{*\text{Exp},\theta,\{e_i\}}(\cdot)$, $L_n^{\text{Retr},\theta}(\cdot)$, $L_n^{*\text{Retr},\theta}$ the composite functions $L_n(\text{Exp}_{\theta} \circ \pi_{\{e_i\}}^{-1}(\cdot))$, $L_n^*(\text{Exp}_{\theta} \circ \pi_{\{e_i\}}^{-1}(\cdot))$, $L_n(R_{\theta}(\cdot))$, $L_n^*(R_{\theta}(\cdot))$, respectively.

We use the following coordinated forms to represent the first-order condition's solutions and the approximate solutions returned by Algorithm 2:

$$\begin{aligned} \tilde{\eta}_n &\text{ is the solution to } \bar{\nabla} L_n^{\text{Exp},\theta_0,\{e_i\}}(\eta) = \mathbf{0}, \\ \tilde{\eta}_n^* &\text{ is the solution to } \bar{\nabla} L_n^{*\text{Exp},\theta_0,\{e_i\}}(\eta) = \mathbf{0}, \\ \hat{\eta}_n &= \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), \quad \hat{\eta}_n^* = \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), \end{aligned}$$

where $\hat{\theta}_n$ is produced by Algorithm 2 and $\hat{\theta}_n^*$ is a generic version of $\hat{\theta}_n^{*[i]}$ based on the dataset \mathcal{X}_n^* .

Following the spirit of expressing quantities under a fixed coordinate system, we also define an analog of Σ in (2) as follows:

$$\begin{aligned} \Omega &:= \left(\mathbb{E}[\bar{\nabla}^2 L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_1)] \right)^{-1} \mathbb{E}[\bar{\nabla} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_1) \bar{\nabla} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_1)^\top] \\ &\quad \cdot \left(\mathbb{E}[\bar{\nabla}^2 L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_1)] \right)^{-1}. \end{aligned}$$

One subtle yet crucial point is that the previously defined exact solutions $\tilde{\theta}_n$ and $\tilde{\theta}_n^*$ may not be unique, and the approximations $\hat{\theta}_n$ and $\hat{\theta}_n^*$ may be associated with a singular covariance estimate. On the computational side, we address potential singularity issues by employing the generalized matrix inverse; on the theoretical side, we will demonstrate that the solutions are well-defined and the covariance estimates are consistent with high probability, so that, in the exceptional event, the approximate solutions can be assigned arbitrary values without affecting the overall conclusion with the help of the delta method.

A.1.1 Implicit Function of Approximate Solutions

We first follow the route of [Bhattacharya et al. \(1978\)](#) to show in the section that $\widehat{\boldsymbol{\eta}}_n$ can be approximated by a smooth function of the sample-averaged derivatives of L .

The first step is to prove the existence of a solution to (4). To begin with, we utilize the Taylor expansion of $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ to derive that for each $i \in [p]$,

$$\begin{aligned} 0 &= \frac{\partial}{\partial x_i} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}) \\ &= e_i L_n(\theta_0) + \sum_j \bar{\nabla}^{\delta_i + \delta_j} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \eta_j + \sum_{\boldsymbol{\nu}, |\boldsymbol{\nu}|=2} \frac{\bar{\nabla}^{\boldsymbol{\nu} + \delta_i} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \boldsymbol{\eta}^{\boldsymbol{\nu}}}{\boldsymbol{\nu}!} + \text{Res}_i(\boldsymbol{\eta}), \end{aligned} \quad (17)$$

where $\text{Res}_i(\boldsymbol{\eta}) := \sum_{\boldsymbol{\nu}: |\boldsymbol{\nu}|=3} \frac{\bar{\nabla}^{\boldsymbol{\nu} + \delta_i} L_n^{\text{Exp}, \theta_0, \{e_i\}}(t\boldsymbol{\eta}) \boldsymbol{\eta}^{\boldsymbol{\nu}}}{\boldsymbol{\nu}!}$ with some $t \in [0, 1]$ represents the residual term and $\boldsymbol{\nu}!$ denotes the factorial product $\prod_{i=1}^p \nu_i!$.

Following the convention in the previous context, we denote that

$$\bar{\mathbf{Z}}_n := \sum_{i \in [n]} \mathbf{Z}_i / n, \quad (18)$$

where \mathbf{Z}_i is defined in Assumption 1.D.

We first set out to control the fluctuation of $\bar{\mathbf{Z}}_n$. By the finite moment conditions (Assumption 1.B) and the moderate deviation result (Lemma 8), we have

$$\mathbb{P}_{\mathcal{X}_n} \left[\left\| n^{\frac{1}{2}} (\bar{\mathbf{Z}}_n - \boldsymbol{\mu}) \right\|_2 \geq (2\Lambda \log n)^{\frac{1}{2}} \right] \leq d_1 n^{-1} (\log n)^{-2}, \quad (19)$$

where Λ is the largest eigenvalue of the covariance matrix of \mathbf{Z}_1 and d_1 is some constant that depends on Λ . Similarly, applying Lemma 8 to $C(x)$ defined in Assumption 1.B yields that

$$\mathbb{P}_{\mathcal{X}_n} \left[\left| \frac{\sum_{i \in [n]} C(X_i)}{n} - \mathbb{E}[C(X_1)] \right| \geq d_2 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \right] \leq d_2 n^{-1} (\log n)^{-2} \quad (20)$$

for some sufficiently large constant d_2 .

Looking into each entry of $\bar{\mathbf{Z}}_n$ separately, (19) implies that

$$\begin{aligned} \mathbb{P}_{\mathcal{X}_n} \left[|e_i L_n(\theta_0)| \geq d_3 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \right] &\leq d_3 n^{-1} (\log n)^{-2}, \quad 1 \leq i \leq p, \\ \mathbb{P}_{\mathcal{X}_n} \left[\left| \bar{\nabla}^{\boldsymbol{\nu} + \delta_i} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \mathbb{E}[\bar{\nabla}^{\boldsymbol{\nu}} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_1)] \right| \geq d_3 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \right] &\leq d_3 n^{-1} (\log n)^{-2}, \\ \mathbb{P}_{\mathcal{X}_n} \left[\left| \sum_{i \in [n]} \bar{\nabla}^{\boldsymbol{\nu}_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i) \bar{\nabla}^{\boldsymbol{\nu}_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i) / n \right. \right. \\ &\quad \left. \left. - \mathbb{E}[\bar{\nabla}^{\boldsymbol{\nu}_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_1) \bar{\nabla}^{\boldsymbol{\nu}_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_1)] \right| \geq d_3 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \right] \leq d_3 n^{-1} (\log n)^{-2}, \end{aligned} \quad (21)$$

simultaneously hold for $1 \leq |\boldsymbol{\nu}| \leq 4$, $1 \leq |\boldsymbol{\nu}_1|, |\boldsymbol{\nu}_2| \leq 3$, and some sufficiently large constant d_3 .

Moreover, invoking the definition of $C(\mathbf{x})$ in Assumption 1.B, it follows from (20) that

$$\begin{aligned} \mathbb{P}_{\mathcal{X}_n} \left[\max_{\boldsymbol{\eta} \in \text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}} \left| \text{Res}_i(\boldsymbol{\eta}) \right| - \left(\left(\sum_{\boldsymbol{\nu}: |\boldsymbol{\nu}|=3} \left| \mathbb{E}_{\mathcal{X}_n} \left[\frac{1}{\boldsymbol{\nu}!} \bar{\nabla}^{\boldsymbol{\nu} + \delta_i} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right] \right| \right) \right. \right. \\ \left. \left. + d_4 n^{-1} (\log n)^{-2} \boldsymbol{\eta}^{\boldsymbol{\nu}} + d_5 \rho (\mathbb{E}[C(X_1)] + d_2 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}) \|\boldsymbol{\eta}\|_2^3 \right) \right] \geq 0 \end{aligned} \quad (22)$$

$$\begin{aligned}
&\leq \mathbb{P} \mathcal{X}_n \left[\left(\sum_{\nu:|\nu|=3} \frac{1}{\nu!} \bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) - \sum_{\nu:|\nu|=3} \left| \mathbb{E} \left[\frac{1}{\nu!} \bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \right] \right| \right) \geq d_4 n^{-1} (\log n)^{-2} \right] \\
&+ \mathbb{P} \mathcal{X}_n \left[\max_{\boldsymbol{\eta}: \text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\boldsymbol{\eta}) \in N} \left(\sum_{\nu:|\nu|=3} \frac{1}{\nu!} \max_{t \in [0,1]} \bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(t\boldsymbol{\eta}) \boldsymbol{\eta}^\nu - \sum_{\nu:|\nu|=3} \frac{1}{\nu!} \bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \boldsymbol{\eta}^\nu \right. \right. \\
&\left. \left. - d_5 \rho (\mathbb{E}[C(X_1)] + d_2 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}) \|\boldsymbol{\eta}\|_2^3 \right) \geq 0 \right] \\
&\leq \mathbb{P} \mathcal{X}_n \left[\left(\sum_{\nu:|\nu|=3} \frac{\bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0})}{\nu!} - \mathbb{E} \left[\sum_{\nu:|\nu|=3} \frac{\bar{\nabla}^{\nu+\delta_i} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0})}{\nu!} \right] \right) \geq d_4 n^{-1} (\log n)^{-2} \right] \\
&+ \mathbb{P} \mathcal{X}_n \left[\max_{\boldsymbol{\eta}: \text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\boldsymbol{\eta}) \in N} \left(\rho \sum_{\nu:|\nu|=3} \frac{\boldsymbol{\eta}^\nu \sum_{i \in [n]} C(X_i)}{\nu! n} - d_5 \rho (\mathbb{E}[C(X_1)] + d_2 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}) \|\boldsymbol{\eta}\|_2^3 \right) \geq 0 \right] \\
&\leq d_6 n^{-1} (\log n)^{-2},
\end{aligned}$$

holds for some constants d_4, d_5, d_6 . Therefore, we can find an event \mathcal{F}_n to be

$$\mathcal{F}_n := \left\{ \|\bar{\mathbf{Z}}_n\|_\infty < d_7 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}, \left| \max_{\boldsymbol{\eta}: \text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\boldsymbol{\eta}) \in N} \text{Res}_i(\boldsymbol{\eta}) \right| < d_7 \|\boldsymbol{\eta}\|_2^3, \text{ for } i, j \in [p] \right\} \quad (23)$$

for some sufficiently large constant d_7 satisfying

$$\mathbb{P}(\mathcal{F}_n^c) \leq d_7 n^{-1} (\log n)^{-2}. \quad (24)$$

Under the event \mathcal{F}_n in (23), multiplying by $(\bar{\nabla}^2 L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}))^{-1}$ both sides of (17) and rearranging the terms yields that

$$\boldsymbol{\eta} = (\mathbb{E}[\bar{\nabla}^2 L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0})] + \boldsymbol{\zeta}_n)^{-1} \left[\boldsymbol{\delta}_n + \sum_{\nu:|\nu|=2} \frac{1}{\nu!} \bar{\nabla}^{\nu+1} L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \boldsymbol{\eta}^\nu + \|\boldsymbol{\eta}\|_2^3 \boldsymbol{\epsilon}_n \right] \quad (25)$$

where $\boldsymbol{\zeta}_n$ is a random matrix and $\boldsymbol{\delta}_n, \boldsymbol{\epsilon}_n$ are random vectors, each with a norm ($\|\cdot\|_2$ or $\|\cdot\|_F$) less than $d_7 p n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}$.

Thus, by the Brouwer Fixed Point Theorem, at least one solution $\tilde{\boldsymbol{\eta}}_n$ exists solving (25) (also (17)) under the event \mathcal{F}_n , within the ball $B_n := B(0, d_8 n^{-1} (\log n)^{-2}) \subset B(\theta_0, \frac{\ell}{2}) \subseteq N$ with a constant d_8 for some sufficiently large n .

The next step is to leverage the implicit function theorem to obtain an approximation. We consider the following equation

$$\mathbf{0} = \mathbf{z}^{\delta_i} + \sum_{\nu:|\nu|=1,2} \frac{1}{\nu!} \mathbf{z}^{(\delta_i+\nu)} \mathbf{u}^\nu =: P(\mathbf{u}, \mathbf{z}).$$

Obviously, the equation has a solution at $\mathbf{u} = \mathbf{0}, \mathbf{z} = \boldsymbol{\mu}$ where $\boldsymbol{\mu} = \mathbb{E} \mathbf{Z}_1$ by Assumption 1.C. By the implicit function theorem and the smoothness of the function $P(\mathbf{u}, \mathbf{z})$ with respect to \mathbf{u} and \mathbf{z} , we conclude that there uniquely exists an injective and infinitely differentiable function \mathbf{H} defined on a neighborhood of $\boldsymbol{\mu}$ such that

$$\mathbf{u} = \mathbf{H}(\mathbf{z}) = (H_1(\mathbf{z}), \dots, H_p(\mathbf{z})) \quad (26)$$

with $\mathbf{H}(\boldsymbol{\mu}) = \mathbf{0}$. We assume that N is contained in the image of \mathbf{H} by shrinking N if necessary. This in turn implies that the first-order solution $\tilde{\theta}_n$ is unique on N with $\text{dist}(\tilde{\theta}_n, \theta_0) = O(n^{-1} \log n)$ under the event \mathcal{F}_n . Therefore it is straightforward by the existence of $\tilde{\boldsymbol{\eta}}_n$ that

$$\tilde{\boldsymbol{\eta}}_n = \mathbf{H}(\bar{\mathbf{Z}}'_n) = \mathbf{H}(\bar{\mathbf{Z}}_n) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \quad (27)$$

solves (17) and also implies that $\text{Exp} \circ \pi_{\{e_i\}}^{-1}(\tilde{\boldsymbol{\eta}}_n)$ is a solution of (4), where $\bar{\mathbf{Z}}'_n$ is a modified version of $\bar{\mathbf{Z}}_n$, by replacing $e_i L_n(\theta_0)$ with $e_i L_n(\theta_0) + \text{Res}_i(\tilde{\boldsymbol{\eta}}_n)$.

Recall that the actual output $\hat{\theta}_n$ of Algorithm 2 is an approximate solution to (4) according to Theorem 1, and $\hat{\boldsymbol{\eta}}_n = \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n)$. Replacing $\tilde{\boldsymbol{\eta}}_n$ with $\hat{\boldsymbol{\eta}}_n$ together with Theorem 1, (23) yields that

$$\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n}(e_i)\}} \circ \text{Log}_{\hat{\theta}_n}(\theta_0) = -(\mathbf{H}(\bar{\mathbf{Z}}_n)) + d_9 n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}} \quad (28)$$

under the event \mathcal{F}_n for some constant d_9 .

A.1.2 Implicit Function of Approximate Resampled Solutions

We now move on to prove an empirical version of the implicit approximation. Given the extremum estimator $\hat{\theta}_n$ based on the dataset $\mathcal{X}_n = \{X_i\}$, we generate a resampled dataset $\mathcal{X}_n^* = \{X_i^*\}$ by drawing n samples with replacement from \mathcal{X}_n , leading to the corresponding resampled extremum estimator $\hat{\theta}_n^*$ in Algorithm 2.

Similar to (17), by performing a Taylor expansion of $L_n^{*\text{Exp}, \theta_0, \{e_i\}}$ at $\tilde{\boldsymbol{\eta}}_n$, we are interested in the solution to

$$\begin{aligned} 0 &= \frac{\partial}{\partial x_i} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}) \\ &= \frac{\partial}{\partial x_i} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) + \sum_j \bar{\nabla}^{\delta_i, \delta_j} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) (\eta_j - \tilde{\eta}_{n,j}) \\ &\quad + \frac{1}{\nu!} \sum_{\nu, |\nu|=2} \bar{\nabla}^{\nu + \delta_i} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) (\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n)^\nu + \text{Res}_i^*(\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n) \end{aligned}$$

for $1 \leq i \leq p$.

Given the samples \mathcal{X}_n , we naturally define a resampled version \mathbf{Z}_n^* of the derivative collection \mathbf{Z}_n as in Assumption 1.D by replacing \mathcal{X}_n with \mathcal{X}_n^* . Conditional on \mathcal{X}_n , we shall control each entry of $\bar{\mathbf{Z}}_n^*$ by decomposing it into two parts:

$$\begin{aligned} &\left| \bar{\nabla}^\nu L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) - \bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) \right| \\ &\leq \underbrace{\left| \bar{\nabla}^\nu L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right|}_{\text{an entry of } \bar{\mathbf{Z}}_n^*} + \underbrace{\left| \bar{\nabla}^\nu L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) - \bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right|}_{\text{an entry of } \bar{\mathbf{Z}}_n} \\ &+ \left| \bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) - \bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right|, \quad (29) \end{aligned}$$

for $1 \leq |\nu| \leq 4$,

$$\begin{aligned}
& \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i^*) / n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i) / n \right| \\
& \leq \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) / n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) / n \right| \\
& \quad + \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i^*) / n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) / n \right| \\
& \quad + \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\eta}_n, X_i) / n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) / n \right|, \quad \text{for } 1 \leq |\nu_1|, |\nu_2| \leq 3. \quad (30)
\end{aligned}$$

For an arbitrary entry $(\mathbf{Z}_1^*)_k$ of \mathbf{Z}_1^* , we notice that

$$\begin{aligned}
& \mathbb{E} \left[|(\mathbf{Z}_1^*)_k - (\bar{\mathbf{Z}}_n)_k|^4 \mid \mathcal{X}_n \right] \\
& \leq 2^4 \max_{0 \leq r \leq 4} \mathbb{E} [(\mathbf{Z}_1^*)_k^r \mid \mathcal{X}_n] (\bar{\mathbf{Z}}_n)_k^{4-r} \\
& \stackrel{\text{by Jensen's inequality}}{\leq} 2^4 \max_{0 \leq r \leq 4} \mathbb{E} [(\mathbf{Z}_1^*)_k^4 \mid \mathcal{X}_n]^{\frac{r}{4}} (\bar{\mathbf{Z}}_n)_k^{4-r} \\
& \leq 2^4 \mathbb{E}[(\mathbf{Z}^*)_k^4] + 2^4 |(\bar{\mathbf{Z}}_n)_k|^4.
\end{aligned}$$

Hence, we have

$$\begin{aligned}
& \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{E} [|(\mathbf{Z}_1^*)_k - (\bar{\mathbf{Z}}_n)_k|^4 \mid \mathcal{X}_n] \geq 2^4 (\mathbb{E} [|(\mathbf{Z}_1)_k|^4] + |\mathbb{E} [|(\bar{\mathbf{Z}}_n)_k|^4] + 2^5) \right] \\
& \leq \mathbb{P}_{\mathcal{X}_n} \left[\left| \sum_{i=1}^n (|(\mathbf{Z}_i)_k|^4 - \mathbb{E} [|(\mathbf{Z}_1)_k|^4]) \right| + n \left| |(\bar{\mathbf{Z}}_n)_k|^4 - |\mathbb{E} [|(\bar{\mathbf{Z}}_n)_k|^4]| \right| \geq 2n \right] \\
& \leq \mathbb{P}_{\mathcal{X}_n} \left[\left| \sum_{i=1}^n (|(\mathbf{Z}_i)_k|^4 - \mathbb{E} [|(\mathbf{Z}_1)_k|^4]) \right| \geq n \right] + \mathbb{P}_{\mathcal{X}_n} \left[\left| |(\bar{\mathbf{Z}}_n)_k|^4 - |\mathbb{E} [|(\bar{\mathbf{Z}}_n)_k|^4]| \right| \geq 1 \right] \quad (31) \\
& \leq n^{-2} \mathbb{E}_{\mathcal{X}_n} \left[\sum_{i=1}^n (|(\mathbf{Z}_i)_k|^4 - \mathbb{E}_{\mathcal{X}_n} [|(\mathbf{Z}_i)_k|^4])^2 \right] + \mathbb{E}_{\mathcal{X}_n} [|(\bar{\mathbf{Z}}_n)_k|^4] \\
& \leq c_{\nu} n^{-1} \mathbb{E}_{\mathcal{X}_n} [|(\mathbf{Z}_1)_k|^8] + c_k n^{-2} \mathbb{E}_{\mathcal{X}_n} [|(\mathbf{Z}_1)_k|^4] \\
& = O(n^{-1})
\end{aligned}$$

for some constant c_k , where the finite moment condition in Assumption 1.B comes into play. Here the penultimate line follows from the observation that, by interpreting $S_t := \sum_{i=1}^t (\mathbf{Z}_i)_k$ for $t = 1, \dots, n$ as a martingale with its natural filtration, and applying Lemma 9 along with Jensen's inequality, we obtain:

$$\mathbb{E}_{\mathcal{X}_n} [|(\bar{\mathbf{Z}}_n)_k|^4] = n^{-4} \mathbb{E}_{\mathcal{X}_n} [S_n^4] \leq c_k n^{-4} \mathbb{E}_{\mathcal{X}_n} \left[\left| \sum_{i \in [n]} (\mathbf{Z}_i)_k^2 \right|^2 \right] \leq c_k n^{-2} \mathbb{E}_{\mathcal{X}_n} [|(\mathbf{Z}_1)_k|^4].$$

Analysis of Eq. (29). For the term $\left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \right|$ with $1 \leq |\nu| \leq 4$, Assumption 1.B implies that

$$\left| \bar{\nabla}^\nu L^{\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n, x) - \bar{\nabla}^\nu L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, x) \right| \leq C(x) \text{dist}(\tilde{\theta}_n, \theta_0),$$

which leads to

$$\left| \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \right| \leq \frac{\sum_{i \in [n]} C(X_i) \text{dist}(\tilde{\theta}_n^*, \theta_0)}{n}, \quad 1 \leq |\nu| \leq 4, \quad (32)$$

$$\left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \right| \leq \frac{\sum_{i \in [n]} C(X_i^*) \text{dist}(\tilde{\theta}_n, \theta_0)}{n}, \quad 1 \leq |\nu| \leq 4. \quad (33)$$

Analogous to the bound in (31), we establish that

$$\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{E} \left[\left| C(X_1^*) - \sum_{i \in [n]} C(X_i)/n \right|^4 \middle| \mathcal{X}_n \right] \geq 2^3 (\mathbb{E}[|C(X)|^4] + |\mathbb{E}[C(X)]|^4) + 2^5 \right] = O(n^{-1}).$$

This together with Lemma 8 leads to the fact that

$$\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\frac{\sum_{i \in [n]} C(X_i^*)}{n} \geq f_1 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} + \mathbb{E}[C(X)] \middle| \mathcal{X}_n \right] \geq f_1 n^{-1} (\log n)^{-2} \right] = O(n^{-1}). \quad (34)$$

for some sufficiently large constant f_1 .

By (29), we can bound the difference $\left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) \right|$ via

$$\begin{aligned} & \left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) \right| \\ & \leq \left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) - \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}) \right| + \frac{C \sum_{i \in [n]} C(X_i) \|\tilde{\eta}_n\|_2}{n} + \frac{C \sum_{i \in [n]} C(X_i^*) \|\tilde{\eta}_n\|_2}{n}. \end{aligned} \quad (35)$$

The decomposition (35) together with (20), (21), (31), and (34) gives that

$$\begin{aligned} & \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left| \bar{\nabla}^\nu L_n^{*\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) - \bar{\nabla}^\nu L_n^{\text{Exp},\theta_0,\{e_i\}}(\tilde{\eta}_n) \right| \geq f_2 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \middle| \mathcal{X}_n \right] \leq f_2 n^{-1} (\log n)^{-2} \right] \\ & \geq 1 - f_2 n^{-1} \end{aligned}$$

for some sufficiently large constant f_2 .

Analysis of Eq. (30). In terms of the counterparts to the quadratic terms in $\bar{\mathbf{Z}}_n$ as decomposed in (30), we begin by bounding each term on the right-hand side of (30).

- Note that (31) has already implied the uniform boundedness of the fourth moment for every entry of \mathbf{Z}_1 , except on an event with probability $O(n^{-1})$. Invoking Lemma 8 again, one has

$$\begin{aligned} & \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_i^*) / n \right. \right. \right. \\ & \quad \left. \left. \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp},\theta_0,\{e_i\}}(\mathbf{0}, X_i) / n \right| \geq f_3 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \middle| \mathcal{X}_n \right] \leq f_3 n^{-1} (\log n)^{-2} \right] \\ & \geq 1 - f_3 n^{-1} \end{aligned} \quad (36)$$

for some constant f_3 .

- Further, by Assumption 1.B, the triangle inequality, and the Cauchy inequality, the second term in (30) can be bounded by

$$\begin{aligned}
& \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*)/n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*)/n \right| \\
& \leq \sum_{i \in [n]} C(X_i^*) \text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0) (|\bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*)| + |\bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*)|)/n \\
& \quad + \sum_{i \in [n]} C(X_i^*)^2 \text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0)^2/n \\
& \leq \text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0) \left(\sum_{i \in [n]} C(X_i^*)^2/n \right) \\
& \quad \cdot \left(2 \left(\sum_{i \in [n]} (\bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*))^2/n \right)^{\frac{1}{2}} + 2 \left(\sum_{i \in [n]} (\bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*))^2/n \right)^{\frac{1}{2}} + 1 \right),
\end{aligned}$$

where we use $\text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0)^2 \leq \text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0)$ in a small enough N to obtain the last inequality.

Following the route toward proving (31) exploiting Lemma 8, we derive that $\sum_{i \in [n]} C(X_i^*)^2/n$ is upper bounded by a constant with probability at least $1 - f_C n^{-1}$ with some constant f_C , where we make use of the finite 16-th moment condition on $C(X_1)$. This together with (36) and the concentration on $\text{dist}(\tilde{\boldsymbol{\theta}}_n, \boldsymbol{\theta}_0)$ under \mathcal{F}_n yields that

$$\begin{aligned}
& \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*)/n \right. \right. \right. \\
& \quad \left. \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i^*)/n \right| \geq f_4 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} |\mathcal{X}_n| \leq f_4 n^{-1} (\log n)^{-2} \right] \\
& \geq 1 - f_4 n^{-1}
\end{aligned} \tag{37}$$

for some constant f_4 .

- Similar to the proof of (37), for the third term in (30) we have

$$\begin{aligned}
& \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i)/n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)/n \right| \\
& \leq f_5 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}
\end{aligned} \tag{38}$$

with probability at least $1 - f_5 n^{-1} (\log n)^{-2}$.

Combining (36), (37), and (38) together with (30) gives that

$$\begin{aligned} & \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*)/n \right. \right. \right. \\ & \quad \left. \left. \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i)/n \right| \geq f_6 n^{-1} (\log n)^{-2} | \mathcal{X}_n \right] \leq f_6 n^{-1} (\log n)^{-2} \right] \\ & \geq 1 - f_6 n^{-1} \end{aligned}$$

holds for some constant f_6 .

Bound for Residual Term. Finally, by a similar argument to (22), (34), and (35), we can also prove that

$$\begin{aligned} & \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\max_{\eta \in N} |\text{Res}_i^*(\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n)| - f_7 \|\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n\|_2^s \left(1 + n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \right) \geq 0 | \mathcal{X}_n \right] \leq f_7 n^{-1} (\log n)^{-2} \right] \\ & \geq 1 - f_7 n^{-1} \end{aligned}$$

with sufficiently large constant f_7 .

Putting Pieces Together. Bringing together the above concentration inequalities, we denote by \mathcal{F}_n^* the event

$$\begin{aligned} \mathcal{F}_n^* := & \left\{ \max_{1 \leq |\boldsymbol{\nu}| \leq 4} \left\{ |\bar{\nabla}^{\boldsymbol{\nu}} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}) - \bar{\nabla}^{\boldsymbol{\nu}} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta})|, 1 \leq |\boldsymbol{\nu}| \leq 4 \right\} < f_8 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}, \right. \\ & \max_{1 \leq |\boldsymbol{\nu}_1|, |\boldsymbol{\nu}_2| \leq 3} \left\{ \left| \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i^*)/n \right. \right. \\ & \quad \left. \left. - \sum_{i \in [n]} \bar{\nabla}^{\nu_1} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i) \bar{\nabla}^{\nu_2} L^{\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n, X_i)/n \right| \right\} < f_8 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}, \quad (39) \\ & \frac{\sum_{i \in [n]} C(X_i^*)}{n} < f_8 n^{-1} (\log n)^{-2} + \mathbb{E}[C(X)], \|\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n\|_\infty < f_8 n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}, \\ & \left. |\text{Res}^*(\boldsymbol{\eta}^* - \tilde{\boldsymbol{\eta}}_n)| < f_8 \|\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n\|_2^3 \right\}, \end{aligned}$$

and by \mathcal{F}'_n the event

$$\mathcal{F}'_n := \left\{ \mathbb{P}[\mathcal{F}_n^* | \mathcal{X}_n] \leq f_8 n^{-1} (\log n)^{-2} \right\} \cap \mathcal{F}_n \quad (40)$$

with a sufficiently large f_8 , where its complementary event is controlled by

$$\mathbb{P}_{\mathcal{X}_n} \left[\mathcal{F}'_n^c \right] \leq f_8 n^{-1} \quad (41)$$

according to the above analysis.

Again, using a similar argument to the first part and the Brouwer fixed point theorem but under the event $\mathcal{F}_n^* \cap \mathcal{F}_n$, we can prove the existence of a solution solving

$$\begin{aligned} \boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n = & (\bar{\nabla}^2 L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) + \boldsymbol{\zeta}_n^*)^{-1} \\ & \cdot \left[\boldsymbol{\delta}_n^* + \sum_{|\boldsymbol{\nu}|=2} \frac{1}{\nu!} \bar{\nabla}^{\boldsymbol{\nu}+1} L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\tilde{\boldsymbol{\eta}}_n) (\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n)^\nu + \|\boldsymbol{\eta} - \tilde{\boldsymbol{\eta}}_n\|_2^3 \boldsymbol{\epsilon}_n^* \right], \quad (42) \end{aligned}$$

where ϵ_n^* is a random vector related to $\bar{\nabla}^\nu L_n^{\text{Exp}, \theta_0, \{e_i\}}(t(\tilde{\eta}_n) + (1-t)\eta), |\nu| = 3, t \in [0, 1]$. We denote the solution to (42) by $\tilde{\eta}_n^*$.

Recall the smooth function defined in (26). Applying the implicit function theorem yields that

$$\tilde{\eta}_n^* - \tilde{\eta}_n = \mathbf{H}(\bar{\mathbf{Y}}'_n)$$

under the event $\mathcal{F}_n^* \cap \mathcal{F}_n$ for a sufficiently large n , where $\bar{\mathbf{Y}}'_n$ corresponds to $\bar{\mathbf{Z}}'_n$ in (27) with \mathcal{X}_n replaced by \mathcal{X}_n^* , while the derivatives are evaluated at $\tilde{\eta}_n$ instead of $\mathbf{0}$.

Since the implicit function \mathbf{H} is deterministic and smooth, replacing $\bar{\mathbf{Y}}'_n$ with $\bar{\mathbf{Y}}_n$ implies that

$$\tilde{\eta}_n^* = \mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n) + f_9 n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}, \quad (43)$$

with some constant f_9 , where we invoke (27) to represent $\tilde{\eta}_n$, and $\bar{\mathbf{Y}}_n$ is the counterpart to $\bar{\mathbf{Z}}_n$ with \mathcal{X}_n replaced by \mathcal{X}_n^* with the derivatives are taken at $\tilde{\eta}_n$ instead of $\mathbf{0}$.

Invoking Theorem 2, (43) turns out to be

$$\hat{\eta}_n^* = \mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}})$$

under $\mathcal{F}_n^* \cap \mathcal{F}_n$.

Finally, we relate the coordinate difference between $\hat{\theta}_n^*$ and $\hat{\theta}_n$ under the chart $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\cdot)$ to the coordinate of $\hat{\theta}_n$ under the chart $\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n^*}(e_i)\}} \circ \text{Log}_{\hat{\theta}_n^*}(\cdot)$. Specifically, by Lemma 1, it holds

$$\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n^*}(e_i)\}} \circ \text{Log}_{\hat{\theta}_n^*}(\hat{\theta}_n) = -\mathbf{H}(\bar{\mathbf{Y}}_n) + f_{10} n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}} \quad (44)$$

under the event $\mathcal{F}_n^* \cap \mathcal{F}_n$, provided a sufficiently large constant f_{10} .

A.1.3 High-Order Asymptotics of Approximate Solutions under Second-Order Retractions

Now we are in a position to combine the above pieces together so as to establish the high-order asymptotics of the iteration series generated by Algorithm 3 and Algorithm 4. We first introduce two covariance matrices associated with the implicit functions in the previous steps:

$$\begin{aligned} \hat{\Omega} &:= \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i)^\top / n \right) \\ &\quad \cdot \left(\bar{\nabla}^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1}, \\ \hat{\Omega}^* &:= \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) \right)^{-1} \\ &\quad \cdot \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n), X_i^*)^\top / n \right) \\ &\quad \cdot \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) \right)^{-1}. \end{aligned}$$

In terms of the involved matrix inversion, we will shortly elaborate on the well-definedness of these quantities under \mathcal{F}_n or \mathcal{F}_n^* .

The essential idea to demonstrate the high-order asymptotics is to exploit the delta method (Lemma 4) to prove the equivalence between the approximate solutions and the implicit function after coordinate transformations.

Proof of (11) (Wald Statistic). For the quantity $R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0)^\top$, we once again note that the generalized-inverse is designed to prevent the algorithm from failure after hundred thousands of resampling in practice; in the theoretical analysis, the generalized inverse can be replaced by the matrix inverse under the event \mathcal{F}_n defined in (23).

Next, we are going to express $R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^{-1} R_{\hat{\theta}_n}^{-1}(\theta_0)$ and $R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^{-1} R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)$ in the coordinate system of $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\cdot)$.

The core of this analysis is that the difference across various charts can be eliminated in the Wald-type statistic regardless of the bases across various charts. Formally, we let $\widehat{\mathbf{Q}}$ denote the orthonormal matrix whose i -th row corresponds to the coordinate of $dR_{\hat{\theta}_n}^{-1}[T_{\theta_0 \rightarrow \hat{\theta}_n} e_i]$. Similarly, let $\widehat{\mathbf{Q}}^*$ denote an analogous orthogonal matrix, with the i -th row being $dR_{\hat{\theta}_n^*}^{-1}[T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_i]$. Then, under the event \mathcal{F}_n and the event $\mathcal{F}_n^* \cap \mathcal{F}_n$, respectively, applying the property of second-order retractions gives that

$$\begin{aligned} \widehat{\mathbf{Q}} R_{\hat{\theta}_n}^{-1}(\theta_0) &= \pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}} \circ \text{Log}_{\hat{\theta}_n}(\theta_0) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) = -\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}), \\ \widehat{\mathbf{Q}}^* R_{\hat{\theta}_n^*}^{-1}(\theta_0) &= \pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_i\}} \circ \text{Log}_{\hat{\theta}_n^*}(\theta_0) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) = -\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}). \end{aligned} \quad (45)$$

Invoking the property of a second-order retraction, it holds for every $\mathbf{x} \in U$ that

$$\begin{aligned} \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(\mathbf{0}, \mathbf{x}) &= \widehat{\mathbf{Q}} \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n}(\mathbf{0}, \mathbf{x}), \\ \bar{\nabla}^2 L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(\mathbf{0}, \mathbf{x}) &= \widehat{\mathbf{Q}} \bar{\nabla}^2 L^{\text{Retr}, \hat{\theta}_n}(\mathbf{0}, \mathbf{x}) \widehat{\mathbf{Q}}^\top. \end{aligned}$$

Then we make the following claim to establish the consistency of studentized terms across different charts whose proof is postponed to Section A.1.5.

Claim 1. *Under the event \mathcal{F}_n , it holds that*

$$\left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) - \widehat{\mathbf{Q}} \bar{\nabla}^2 L_n^{\text{Retr}, \hat{\theta}_n} \widehat{\mathbf{Q}}^\top \right\|_F = O(n^{-1} \log n), \quad (46)$$

$$\begin{aligned} &\left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i)^\top / n \right. \\ &\left. - \widehat{\mathbf{Q}} \sum_{i=1}^n \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n}(\mathbf{0}, X_i)^\top \widehat{\mathbf{Q}}^\top / n \right\|_F = O(n^{-1} \log n). \end{aligned} \quad (47)$$

Similarly, under the event \mathcal{F}_n^* it has

$$\left\| \bar{\nabla}^2 L_n^{*\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) - \widehat{\mathbf{Q}}^* \bar{\nabla}^2 L_n^{*\text{Retr}, \hat{\theta}_n^*} \widehat{\mathbf{Q}}^{*\top} \right\|_F = O(n^{-1} \log n), \quad (48)$$

$$\begin{aligned} &\left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n), X_i^*)^\top / n \right. \\ &\left. - \widehat{\mathbf{Q}}^* \sum_{i=1}^n \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n^*}(\mathbf{0}, X_i^*) \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n^*}(\mathbf{0}, X_i^*)^\top \widehat{\mathbf{Q}}^{*\top} / n \right\|_F = O(n^{-1} \log n). \end{aligned} \quad (49)$$

As a consequence of the above claim, we have proved that, given a sufficiently large n , the involved matrix inverse operations in the definitions of $\widehat{\Sigma}$ and $\check{\Sigma}$ are always feasible, under \mathcal{F}_n and

$\mathcal{F}_n^* \cap \mathcal{F}_n$, respectively, in light of (46) and (48). Moreover, by the definitions of $\widehat{\Omega}$ and $\widehat{\Sigma}_n$, Claim 1 immediately implies that

$$\begin{aligned} \left\| \widehat{Q} \widehat{\Sigma} \widehat{Q}^\top - \widehat{\Omega} \right\| &= O(n^{-1} \log n), \\ \left\| \widehat{Q}^* \widehat{\Sigma} \widehat{Q}^{*\top} - \widehat{\Omega}^* \right\| &= O(n^{-1} \log n). \end{aligned} \quad (50)$$

As a consequence of (45) and (50), we have

$$\begin{aligned} \left| R_{\widehat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^{-1} R_{\widehat{\theta}_n}^{-1}(\theta_0)^\top - \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n)^\top \widehat{\Omega}^{-1} \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n) \right| &= O(n^{-2}(\log n)^2), \\ &\text{under } \mathcal{F}_n, \\ \left| R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)^\top \check{\Sigma}^{-1} R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)^\top - (\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n^*) - \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n))^\top \right. \\ &\quad \left. \widehat{\Omega}^{*-1} (\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n^*) - \pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n)) \right| = O(n^{-2}(\log n)^2) \\ &\text{under } \mathcal{F}_n^*. \end{aligned}$$

Here we are going to utilize the smooth implicit function approximation obtained in Section A.1.1 and A.1.2. Substitution of $H(\bar{\mathbf{Z}}_n)$ and $H(\bar{\mathbf{Z}}_n^*)$ into $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n)$ and $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\widehat{\theta}_n^*)$ respectively together with (28) and (44) gives that

$$\begin{aligned} \left| R_{\widehat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^{-1} R_{\widehat{\theta}_n}^{-1}(\theta_0) - \mathbf{H}(\bar{\mathbf{Z}}_n)^\top \widehat{\Omega} \mathbf{H}(\bar{\mathbf{Z}}_n) \right| &= O(n^{-2}(\log n)^2) \\ \left| R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)^\top \check{\Sigma}^{-1} R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n) - \mathbf{H}(\bar{\mathbf{Y}}_n)^\top \widehat{\Omega}^* \mathbf{H}(\bar{\mathbf{Y}}_n) \right| &= O(n^{-2}(\log n)^2). \end{aligned} \quad (51)$$

Now we are left with establishing the Edgeworth expansion of $\mathbf{H}(\bar{\mathbf{Z}}_n)^\top \widehat{\Omega}^{-1} \mathbf{H}(\bar{\mathbf{Z}}_n)$ as well as $\mathbf{H}(\bar{\mathbf{Y}}_n)^\top \widehat{\Omega}^{*-1} \mathbf{H}(\bar{\mathbf{Y}}_n)$. To see the distributional consistency of quadratic terms, we first establish the consistency of the multivariate studentized terms, namely, $\widehat{\Omega}^{-\frac{1}{2}} \mathbf{H}(\bar{\mathbf{Z}}_n)$ and $\widehat{\Omega}^{*- \frac{1}{2}} \mathbf{H}(\bar{\mathbf{Y}}_n)$. We first note that under the event \mathcal{F}_n with every sufficiently large n ,

$$\begin{aligned} \widehat{\Omega}^{-\frac{1}{2}} &= \Omega^{-\frac{1}{2}} (\mathbf{I} + (\widehat{\Omega}^{\frac{1}{2}} - \Omega^{\frac{1}{2}}) \Omega^{-\frac{1}{2}})^{-1} = \Omega^{-\frac{1}{2}} \sum_{k=0}^{\infty} (-1)^k ((\widehat{\Omega}^{\frac{1}{2}} - \Omega^{\frac{1}{2}}) \Omega^{-\frac{1}{2}})^k \\ &= \Omega^{-\frac{1}{2}} - \Omega^{-\frac{1}{2}} (\widehat{\Omega}^{\frac{1}{2}} - \Omega^{\frac{1}{2}}) \Omega^{-\frac{1}{2}} + O(n^{-1} \log n), \end{aligned} \quad (52)$$

where the validity of the expansion is ensured by the conditions in \mathcal{F}_n .

Invoking Proposition 4 first yields that

$$\sup_{B \in \mathcal{B}} \left| \mathbb{P} \left[\sqrt{n} (\Omega^{-\frac{1}{2}} - \Omega^{-\frac{1}{2}} (\widehat{\Omega}^{\frac{1}{2}} - \Omega^{\frac{1}{2}}) \Omega^{-\frac{1}{2}}) \mathbf{H}(\bar{\mathbf{Z}}_n) \in B \right] - \int_B \psi_{4,n}^{(\mathbf{H})} d\mathbf{x} \right| = o(n^{-1})$$

holds for a class \mathcal{B} of Borel set satisfying, for some $a > 0$,

$$\Phi_{\Sigma}((\partial B)^\epsilon) = O(\epsilon^a),$$

where Φ_{Σ} represents the multivariate Gaussian measure with covariance Σ . Here $\psi_{4,n}^{(\mathbf{H})}$ denotes the formal Edgeworth expansion of $(\Omega^{-\frac{1}{2}} - \Omega^{-\frac{1}{2}} (\widehat{\Omega}^{\frac{1}{2}} - \Omega^{\frac{1}{2}}) \Omega^{-\frac{1}{2}}) \mathbf{H}(\bar{\mathbf{Z}}_n)$ as detailed in Section D.

Then applying Lemma 4 yields that

$$\sup_{B \in \mathcal{B}} \left| \mathbb{P}[\sqrt{n}\widehat{\Omega}^{-\frac{1}{2}}\mathbf{H}(\bar{\mathbf{Z}}_n) \in B] - \int_B \psi_{4,n}^{(\mathbf{H})} d\mathbf{x} \right| = O(n^{-1+\xi})$$

holds for every $0 < \xi \leq \frac{1}{2}$, where we make use of (52) and (24).

On the other hand, for its resample counterpart $\mathbf{H}(\bar{\mathbf{Y}}_n)$, a similar derivation to (52) gives that

$$\widehat{\Omega}^{*-\frac{1}{2}} = \widehat{\Omega}^{-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}}(\widehat{\Omega}^{*-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}})\widehat{\Omega}^{-\frac{1}{2}} + O(n^{-1} \log n) \quad (53)$$

under the event \mathcal{F}_n^* .

Applying Proposition 5, and Proposition 6 to $(\widehat{\Omega}^{-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}}(\widehat{\Omega}^{*-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}})\widehat{\Omega}^{-\frac{1}{2}})\mathbf{H}(\bar{\mathbf{Y}}_n)$, we arrive at

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}} \left| \mathbb{P}[(\widehat{\Omega}^{-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}}(\widehat{\Omega}^{*-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}})\widehat{\Omega}^{-\frac{1}{2}})\mathbf{H}(\bar{\mathbf{Y}}_n) \in B | \mathcal{X}_n] - \int_B \psi_{4,n}^{*(\mathbf{H})} d\mathbf{x} \right| > n^{-1+\xi} \right] < \infty \quad (54)$$

for every $0 < \xi \leq \frac{1}{2}$, where we invoke Lemma 4 together with (41) and $\psi_{4,n}^{*(\mathbf{H})}$ denotes the empirical Edgeworth expansion of the random vector $(\widehat{\Omega}^{-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}}(\widehat{\Omega}^{*-\frac{1}{2}} - \widehat{\Omega}^{-\frac{1}{2}})\widehat{\Omega}^{-\frac{1}{2}})\mathbf{H}(\bar{\mathbf{Y}}_n)$ conditional on \mathcal{X}_n .

Combining (54) with (53), Lemma 4 yields that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}} \left| \mathbb{P}[\widehat{\Omega}^{*-\frac{1}{2}}\mathbf{H}(\bar{\mathbf{Y}}_n) \in B | \mathcal{X}_n] - \int_B \psi_{4,n}^{*(\mathbf{H})} d\mathbf{x} \right| > n^{-1+\xi} \right] < \infty.$$

Then, as stated in Remark 2.2, Chandra and Ghosh (1979), given a random vector $\mathbf{X} \in \mathbb{R}^d$, the validity of Edgeworth expansion for $\|\mathbf{X}\|_2^2$ can be demonstrated by (Chandra and Ghosh, 1979, Theorem 1) once we derive a valid Edgeworth expansion for the distribution of \mathbf{X} . To be specific, one has

$$\begin{aligned} & \sup_{t \in \mathbb{R}^+} \left| \mathbb{P} \left[n\mathbf{H}(\bar{\mathbf{Z}}_n)^\top \widehat{\Omega}^{-1} \mathbf{H}(\bar{\mathbf{Z}}_n) \leq t \right] - \int_{x \leq t} \psi_{4,n}^{(\mathbf{H}^2)}(x) dx \right| \leq O(n^{-1+\xi}), \\ & n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{t \in \mathbb{R}^+} \left| \mathbb{P} \left[n\mathbf{H}(\bar{\mathbf{Y}}_n)^\top \widehat{\Omega}^{*-1} \mathbf{H}(\bar{\mathbf{Y}}_n) \leq t | \mathcal{X}_n \right] - \int_{x \leq t} \psi_{4,n}^{*(\mathbf{H}^2)}(x) dx \right| > n^{-1+\xi} \right] < \infty. \end{aligned}$$

Here the function $\psi_{4,n}^{(\mathbf{H}^2)}(x)$ is given by $f_{\chi_p^2}(x) \cdot (1 + n^{-1}\pi_1^{(\mathbf{H}^2)}(x))$, where $f_{\chi_p^2}$ denotes the density function of a chi-squared distribution of freedom p and $\pi_1^{(\mathbf{H}^2)}(x)$ is a polynomial whose coefficients are determined by those of $\psi_{4,n}^{(\mathbf{H})}(\mathbf{x})$. Similarly, the function $\psi_{4,n}^{*(\mathbf{H}^2)}(x)$ obeys the same form, with its coefficients determined by the ones of $\psi_{4,n}^{*(\mathbf{H})}(\mathbf{x})$.

Leveraging (51), we apply the delta method (Lemma 4) to the above approximations and derive that

$$\sup_{t \in \mathbb{R}^+} \left| \mathbb{P} \left[R_{\widehat{\theta}_n}^{-1}(\theta_0)^\top \widehat{\Sigma}^\dagger R_{\widehat{\theta}_n}^{-1}(\theta_0) \leq t \right] - \int_{x \leq t} \psi_{4,n}^{(\mathbf{H}^2)}(x) dx \right| \leq O(n^{-1+\xi}), \quad (55)$$

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{t \in \mathbb{R}^+} \left| \mathbb{P} \left[nR_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n)^\top \widehat{\Sigma}^\dagger R_{\widehat{\theta}_n^*}^{-1}(\widehat{\theta}_n) \leq t | \mathcal{X}_n \right] - \int_{x \leq t} \psi_{4,n}^{*(\mathbf{H}^2)}(x) dx \right| > n^{-1+\xi} \right] < \infty. \quad (56)$$

Finally, the desired result follows by combining (56) with (55):

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{x \in \mathbb{R}^+} \left| \mathbb{P} \left[R_{\hat{\theta}_n}^{-1}(\hat{\theta}_n)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\hat{\theta}_n) \leq x \mid \mathcal{X}_n \right] - \mathbb{P} \left[R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0) \leq x \right] \right| > n^{-1+\xi} \right] < +o(1) \quad (57)$$

where we make use of the fact that the coefficients of the n^{-1} term in the polynomial of $\psi_{4,n}^{*(\mathbf{H}^2)}(x)$ converge to those of $\psi_{4,n}^{(\mathbf{H}^2)}(x)$ at a rate of $O(n^{-\xi})$ for every $0 < \xi < \frac{1}{2}$, since:

1. The coefficients are determined by the first four moments of \mathbf{Z}_1 , and their empirical counterpart, respectively, as well as the derivatives of the functions $(\mathbf{\Omega}^{-\frac{1}{2}} - \mathbf{\Omega}^{-\frac{1}{2}}(\hat{\mathbf{\Omega}}^{\frac{1}{2}} - \mathbf{\Omega}^{\frac{1}{2}})\mathbf{\Omega}^{-\frac{1}{2}})\mathbf{H}(\bar{\mathbf{Z}}_n)$ and $(\hat{\mathbf{\Omega}}^{-\frac{1}{2}} - \hat{\mathbf{\Omega}}^{-\frac{1}{2}}(\hat{\mathbf{\Omega}}^{*\frac{1}{2}} - \hat{\mathbf{\Omega}}^{-\frac{1}{2}})\hat{\mathbf{\Omega}}^{-\frac{1}{2}})\mathbf{H}(\bar{\mathbf{Y}}_n)$, respectively.
2. By the finite moment condition Assumption 1.B, Lemma 8 yields that the empirical moments converge to the population ones at a rate of $O(n^{-\xi})$ with probability $1 - O(n^{-1})$, for every $0 < \xi < \frac{1}{2}$.
3. By the convergence rate of $\bar{\mathbf{Z}}_n$ in (19) and Proposition 6, we can prove that the involved derivatives also converge to their counterparts at a rate of $O(n^{-\xi})$ with probability $1 - O(n^{-1})$, for every $0 < \xi < \frac{1}{2}$.
4. The function $f_{\mathcal{X}_p^2}(x)$, when multiplied by any polynomial is always integrable.

Proof of (12) (t -Statistic). In contrast to the Wald-type statistic, where orthogonal matrices corresponding to basis transformations are canceled out, analyzing the intrinsic t -type statistic is more complicated because of the existence of basis transformation. Without loss of generality, we let \mathbf{a} be δ_i for an arbitrary $i \in [p]$. In what follows, we claim the existence of a precise enough approximation, provided the differentiability of the retraction:

Claim 2. *Under the event \mathcal{F}_n^* , for each $i \in [p]$, there exists a function \tilde{H}_i such that*

$$\left| \frac{\tilde{H}_i(\bar{\mathbf{Z}}_n, \mathbb{E}[\mathbf{Z}_1])}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} - \frac{R_{\hat{\theta}_n}(\theta_0)_i}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} \right| = O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}), \quad (58)$$

$$\left| \frac{\tilde{H}_i(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n)}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} - \frac{R_{\hat{\theta}_n}(\hat{\theta}_n)_i}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} \right| = O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}). \quad (59)$$

Moreover, the function \tilde{H}_i satisfies that

1. $\tilde{H}_i(\cdot, \mathbf{x}')$ is three-times continuously differentiable on O for every $\mathbf{x}' \in O$ where O is an open neighborhood of $\mathbb{E}[\mathbf{Z}_1]$.
2. The first three-times derivatives of $\tilde{H}_i(\mathbf{x}_1, \mathbf{x}_2)$ with respect to the first variable $\mathbf{x}_1 \in O$ are continuously differentiable with respect to $\mathbf{x}_2 \in O$.

The lengthy proof of Claim 2 is postponed to Section A.1.6. For the function \tilde{H}_i , it follows from Proposition 4 that

$$\sup_{B \in \mathcal{B}} \left| \mathbb{P} \left[n^{\frac{1}{2}} \tilde{H}_i(\bar{\mathbf{Z}}_n, \mathbb{E}[\mathbf{Z}_1]) \in B \right] - \int_B \Psi_{4,n}^{(\tilde{H}_i)}(v) dv \right| = o(n^{-1}), \quad (60)$$

holds for all class $\mathcal{B} \in \mathcal{B}^p$ (the collection of Borel sets in \mathbb{R}^p) such that, for some $a > 0$,

$$\sup_{B \in \mathcal{B}} \Phi_{\Sigma}((\partial B)^\epsilon) = O(\epsilon^a), \quad \text{as } \epsilon \downarrow 0.$$

Here the explicit form of $\Psi_{4,n}^{(\tilde{H}_i)}(x)$ is introduced in Remark 5.

Combining the fact that $\mathbb{P}(\mathcal{F}_n^{\mathcal{L}}) \leq Cn^{-1}(\log n)^{-2}$, (58), and Lemma 4 with (60) yields that

$$\sup_{B \in \mathcal{B}} \left| \mathbb{P}_{\mathcal{X}_n} \left[n^{\frac{1}{2}} \frac{R_{\hat{\theta}_n}^{-1}(\theta_0)_i}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} \in B \right] - \int_B \Psi_{4,n}^{(\tilde{H}_i)}(v) dv \right| = O(n^{-1+\xi}) \quad (61)$$

for every $0 < \xi \leq \frac{1}{2}$.

Similarly, Proposition 5 and Proposition 6 implies that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}} \left| \mathbb{P} \left[n^{\frac{1}{2}} \tilde{H}_i(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n) \in B | \mathcal{X} \right] - \int_B \Psi_{4,n}^{*(\tilde{H}_i)}(v) dv \right| > n^{-1} \right] < \infty \quad (62)$$

for every class \mathcal{B} of Borel set satisfying, for some $a > 0$,

$$\Phi_{\Sigma}((\partial B)^\epsilon) = O(\epsilon^a),$$

where $\Psi_{4,n}^{*(\tilde{H}_i)}$ is an empirical analog to $\Psi_{4,n}^{(\tilde{H}_i)}$.

Again, invoking (58) and the delta method (Lemma 4), one has

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}'} \left| \mathbb{P} \left[n^{\frac{1}{2}} \frac{R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_i}{(\check{\Sigma})_{i,i}^{\frac{1}{2}}} \in B | \mathcal{X}_n \right] - \int_B \Psi_{4,n}^{*(\tilde{H}_i)}(v) dv \right| > n^{-1+\xi} \right] < \infty$$

for every $0 < \xi \leq \frac{1}{2}$. Here, the coefficients of $n^{-\frac{1}{2}}$ in $\Psi_{4,n}^{(\tilde{H}_i)}$ and $\Psi_{4,n}^{*(\tilde{H}_i)}$ coincides and the coefficient of n^{-1} in $\Psi_{4,n}^{*(\tilde{H}_i)}$ converges to that in $\Psi_{4,n}^{(\tilde{H}_i)}$ at a rate of $O(n^{-\xi})$ with probability $1 - O(n^{-1})$ for $0 < \xi < \frac{1}{2}$, following similar arguments to those below (57).

Consequently, combining (61) with (62) gives that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{x \in \mathbb{R}} \left| \mathbb{P}_{\mathcal{X}_n} \left[n^{\frac{1}{2}} \frac{R_{\hat{\theta}_n}^{-1}(\theta_0)_i}{(\hat{\Sigma})_{i,i}^{\frac{1}{2}}} \leq x \right] - \mathbb{P} \left[n^{\frac{1}{2}} \frac{R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_i}{(\check{\Sigma})_{i,i}^{\frac{1}{2}}} \leq x | \mathcal{X}_n \right] \right| > n^{-1+\xi} \right] < \infty$$

for every $0 < \xi \leq \frac{1}{2}$. □

A.1.4 Proof of Corollary 1

The coverage probability is a consequence of the uniform controls in the proof of Theorem 3. For clarity, we provide the proof for the Wald statistic for instance.

We define w_α^* as the exact solution t to $\int_{x \leq t} \psi_{4,n}^{*(\mathbf{H}^2)}(x) dx = \alpha$, whose existence is ensured by the continuity of the function. We then recall the definition of the approximate quantile in Algorithm 3 with $b = \infty$, expressed as:

$$w_\alpha = \arg \min_{t \in \mathbb{R}^+} \left| \mathbb{P} [W^* \leq t | \mathcal{X}_n] - (1 - \alpha) \right|.$$

From this, we deduce the following:

$$\begin{aligned}
& \left| \mathbb{P}_{\mathcal{X}_n} [R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0) \leq w_\alpha] - (1 - \alpha) \right| \\
& \lesssim \left| \mathbb{P}_{\mathcal{X}_n} [R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0) \leq w_\alpha] - \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} [R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^\dagger R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) \leq w_\alpha | \mathcal{X}_n] \right] \right| \\
& \quad + \left| \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} [R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^\dagger R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) \leq w_\alpha | \mathcal{X}_n] \right] - (1 - \alpha) \right| \\
& \leq \left| \mathbb{P}_{\mathcal{X}_n} [R_{\hat{\theta}_n}^{-1}(\theta_0)^\top \hat{\Sigma}^\dagger R_{\hat{\theta}_n}^{-1}(\theta_0) \leq w_\alpha] - \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} [R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^\dagger R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) \leq w_\alpha | \mathcal{X}_n] \right] \right| \\
& \quad + \left| \mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} [R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)^\top \check{\Sigma}^\dagger R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n) \leq w_\alpha^* | \mathcal{X}_n] \right] - (1 - \alpha) \right| \\
& \leq O(n^{-1+\xi}),
\end{aligned}$$

for every $0 < \xi < \frac{1}{2}$, where, in the penultimate line, we use (11) to upper bound the first term and (56) to upper bound the second term. Finally, we leverage the Glivenko-Cantelli theorem to conclude.

The coverage guarantees for the intrinsic t -statistic and the extrinsic t -statistic can be derived using analogous arguments, which we omit for brevity.

A.1.5 Proof of Claim 1

The basic idea is to pull the gradient and the Hessian back to the coordinate chart induced by $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}$ by controlling the accumulated error in parallel transports. With regard to the gradient term $\bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(X_i)$, it follows by Lemma 7 and the definition of \mathcal{F}'_n that

$$\begin{aligned}
& \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(X_i) \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(X_i)^\top / n \right. \\
& \quad \left. - \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i)^\top / n \right\|_F \\
& \leq 2\sqrt{p} \cdot \max_{j \in [p]} \left\| \text{dExp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}[\delta'_j] - T_{\theta_0 \rightarrow \hat{\theta}_n} e_j \right\| \\
& \quad \cdot \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i)^\top / n \right\|_F \\
& \quad + p \max_{j \in [p]} \left\| \text{dExp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}[\delta'_j] - T_{\theta_0 \rightarrow \hat{\theta}_n} e_j \right\|^2 \\
& \quad \cdot \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i)^\top / n \right\|_F \\
& = O(n^{-1} \log n), \tag{63}
\end{aligned}$$

where δ'_j denotes the j -th canonical tangent vector of \mathbb{R}^p at $\pi_{\{e_i\}}^{-1} \circ \text{Log}_{\theta_0}(\hat{\theta}_n)$ with a slight abuse of notation. Here the last line arises since the term

$$\left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n), X_i)^\top / n \right\|_F$$

is bounded by a constant as a consequence of (20) and $\max_{j \in [p]} \left\| \text{dExp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}[\delta'_j] - T_{\theta_0 \rightarrow \hat{\theta}_n} e_j \right\| \leq O(\text{dist}(\theta_0, \hat{\theta}_n)^2)$ by Lemma 7.

Following similar arguments, we can also prove that the inequality (49) holds as a result of Lemma 7 and (34) under the event \mathcal{F}'_n

$$\begin{aligned}
& \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n^*, \{T_{\theta_0 \rightarrow \hat{\theta}_n^*}\}}(X_i^*) \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n^*, \{T_{\theta_0 \rightarrow \hat{\theta}_n^*}\}}(X_i^*)^\top / n \right. \\
& - \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*)^\top / n \left. \right\|_F \\
& \leq 2\sqrt{p} \cdot \max_{j \in [p]} \left\| \text{dExp}_{\theta_0} \cdot \pi_{\{e_i\}}^{-1} \delta_j'' - T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_j \right\| \\
& \quad \cdot \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*)^\top / n \right\|_F \\
& \quad + p \max_{j \in [p]} \left\| \text{dExp}_{\theta_0} \cdot \pi_{\{e_i\}}^{-1} \delta_j'' - T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_j \right\|^2 \\
& \quad \cdot \left\| \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*), X_i^*)^\top / n \right\|_F \\
& = O(n^{-1} \log n).
\end{aligned}$$

Finally, invoking the relations (28) and (43) as well as the uniform bound yields the desired results (47) and (49).

Regarding the Hessian terms, Lemma 2 demonstrates that the difference is not only related to the distances among θ_0 , $\hat{\theta}_n$, and $\hat{\theta}_n^*$ but also to the magnitude of averaged gradients, that is,

$$\begin{aligned}
& \left\| \hat{\mathbf{Q}} \bar{\nabla}^2 L_n^{\text{Retr}, \hat{\theta}_n}(\mathbf{0}) \hat{\mathbf{Q}}^\top - \bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n)) \right\| \\
& \leq O(n^{-1} \log n) + O(n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}) \left\| \nabla L_n(\hat{\theta}_n) \right\| \\
& = O(n^{-1} \log n), \tag{64} \\
& \left\| \hat{\mathbf{Q}}^* \bar{\nabla}^2 L_n^{\text{Retr}, \hat{\theta}_n^*}(\mathbf{0}) \hat{\mathbf{Q}}^{*\top} - \bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}_n^*, \{T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*)) \right\| \\
& \leq O(n^{-1} \log n) + O(n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}}) \left\| \nabla L_n^*(\hat{\theta}_n^*) \right\| \\
& = O(n^{-1} \log n),
\end{aligned}$$

where $\left\| \nabla L_n(\hat{\theta}_n) \right\|$ and $\left\| \nabla L_n^*(\hat{\theta}_n^*) \right\|$ are of the order $o(n^{-\frac{1}{2}} (\log n)^{\frac{1}{2}})$ by Theorem 1, Theorem 2, (32), and (33).

Analogously, replacing $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n)$ and $\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n^*)$ with $\mathbf{H}(\bar{\mathbf{Z}}_n)$ and $\mathbf{H}(\bar{\mathbf{Y}}_n)$, respectively, implies (46) and (48). \square

A.1.6 Proof of Claim 2

Step 1: Basis Transformation Analysis. The goal of this part is to establish explicit approximations for the rotated coordinates as functions of $\bar{\mathbf{Z}}_n$ and $\bar{\mathbf{Y}}_n$ respectively. Before continuing, we remind that in Section A.1.1 and Section A.1.2, we established that $\mathbf{H}(\bar{\mathbf{Z}}_n)$, $\mathbf{H}(\bar{\mathbf{Y}}_n)$ serve as nice approximations to $-\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}} \circ \text{Log}_{\hat{\theta}_n}(\theta_0)$ and $\pi_{\{T_{\theta_0 \rightarrow \hat{\theta}_n^*} e_i\}} \circ \text{Log}_{\hat{\theta}_n^*}(\theta_0)$, respectively. The remaining task is to address how to reconcile the bases induced by parallel transports and those arising from the adopted retraction $\{R_\theta\}$.

The subsequent analysis is a natural exploitation of the differentiability of $dR_\theta^{-1}(T_{\theta_0 \rightarrow \theta}(dR_{\theta_0}\delta_i))$. For ease of notations, we denote the composite function $dR_{\text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}}^{-1}(T_{\theta_0 \rightarrow \text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}}(\mathbf{x})(e_j))$ by $\mathbf{t}_j(\mathbf{x}) = (t_{i,k}(\mathbf{x}))_{k \in [p]}$ for every $j \in [p]$, which is twice-differentiable as assumed in the second part of Theorem 3.

In the context of this section and the next section, we proceed with our analysis under the event $\mathcal{F}_n^* \cap \mathcal{F}_n$ for every n , which heuristically means that, $\hat{\theta}_n, \hat{\theta}_n^*, \tilde{\theta}_n$, and $\tilde{\theta}_n^*$ are always contained in a sequence of shrinking regions as illustrated in Section A.1.1 and Section A.1.2, and the deviations of derivatives are appropriately bounded.

We begin with the Taylor expansion of $\mathbf{t}_i(\mathbf{x}) = (t_{i,k}(\mathbf{x}))_{k \in [p]}$ with respect to \mathbf{x} . One has

$$t_{i,k}(\mathbf{x}) = t_{i,k}(\mathbf{0}) + \sum_{j \in [p]} \frac{\partial t_{i,k}}{\partial x_j}(\mathbf{0})x_j + O(\|\mathbf{x}\|_2^2). \quad (65)$$

Recall that the i -th canonical basis of \mathbb{R}^p is denoted by δ_i for $i \in [p]$. For the quantity $R_{\hat{\theta}_n}^{-1}(\theta)_i$ with $\theta \in \mathcal{M}$, it then holds that

$$\begin{aligned} (R_{\hat{\theta}_n}^{-1}(\theta))_i &= \pi_{\{dR_{\hat{\theta}_n}\delta_i\}} \circ \text{Log}_{\hat{\theta}_n}(\theta) + O(\text{dist}(\theta, \hat{\theta}_n)^3) \\ &= \sum_{j \in [p]} (\mathbf{t}_j(\hat{\boldsymbol{\eta}}_n))_i \left(\pi_{\{T_{\theta_0 \rightarrow \text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}}(\hat{\boldsymbol{\eta}}_n)(e_i)\}} \circ \text{Log}_{\hat{\theta}_n}(\theta) \right)_j + O(\text{dist}(\theta, \hat{\theta}_n)^3) \\ &= \sum_{j \in [p]} (\mathbf{t}_j(\hat{\boldsymbol{\eta}}_n))_i \left(\pi_{\{T_{\theta_0 \rightarrow \text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}}(\hat{\boldsymbol{\eta}}_n)(e_i)\}} \circ \text{Log}_{\hat{\theta}_n}(\theta) \right)_j + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\ &= \sum_{j \in [p]} ((\mathbf{t}_j(\mathbf{H}(\bar{\mathbf{Z}}_n)))_i + O(n^{-1} \log n)) \left(\pi_{\{T_{\theta_0 \rightarrow \text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}}(\hat{\boldsymbol{\eta}}_n)(e_i)\}} \circ \text{Log}_{\hat{\theta}_n}(\theta) \right)_j + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}), \end{aligned}$$

where the first equality holds by the property of second-order retractions, the second one follows by the definition of $\mathbf{t}_i(\mathbf{x})$, and the last equality arises by (28) and the fact that \mathbf{t}_j is continuously differentiable.

Taking $\theta = \theta_0$ and invoking the coordinate invariance under parallel transports give that

$$\begin{aligned} - (R_{\hat{\theta}_n}^{-1}(\theta_0))_i &= \sum_{j \in [p]} -(\mathbf{t}_j(\mathbf{H}(\bar{\mathbf{Z}}_n)))_i (\hat{\boldsymbol{\eta}}_n)_j + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\ &= \sum_{j \in [p]} t_{j,i}(\mathbf{H}(\bar{\mathbf{Z}}_n))H(\bar{\mathbf{Z}}_n)_j + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\ &= H(\bar{\mathbf{Z}}_n)_i + \sum_{j,k,l \in [p]} \frac{\partial t_{j,i}}{\partial x_k}(\mathbf{0})H(\bar{\mathbf{Z}}_n)_kH(\bar{\mathbf{Z}}_n)_j + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}), \end{aligned} \quad (66)$$

where the last line holds by the fact that $t_{i,j}(\mathbf{0}) = \mathbb{1}\{i = j\}$ and (65).

We now move on to the empirical counterpart $R_{\hat{\theta}_n}(\hat{\theta}_n^*)$. Again, by jointly applying the property of second-order retractions, the property of the double exponential mapping (Lemma 1), and a similar expansion of \mathbf{t}_j as above, we find:

$$\begin{aligned} - R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_i &= \pi_{\{dR_{\hat{\theta}_n^*}e_i\}} \circ \text{Log}_{\hat{\theta}_n^*}(\hat{\theta}_n) + O(\text{dist}(\hat{\theta}_n, \hat{\theta}_n^*)^3) \\ &= \left(\pi_{\{dR_{\hat{\theta}_n^*}e_i\}} \circ T_{\theta_0 \rightarrow \hat{\theta}_n^*} \left(\text{Log}_{\theta_0}(\hat{\theta}_n) - \text{Log}_{\theta_0}(\hat{\theta}_n^*) \right) \right)_i + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\ &= \sum_{j \in [p]} t_{j,i}(\mathbf{H}(\bar{\mathbf{Y}}_n))H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k,l \in [p]} \frac{\partial t_{j,i}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n))H(\bar{\mathbf{Y}}_n)_kH(\bar{\mathbf{Y}}_n)_l + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}). \end{aligned}$$

Step 2: Studentization Analysis. Now we are placed to prove Claim 2 associated with the studentization quantities in Algorithm 4. We aim to show that the numerators and the denominators of the studentized statistics can be replaced by clearer forms related to $\bar{\mathbf{Z}}_n$ and $\bar{\mathbf{Y}}_n$. For clarity, we herein prove (59) for $i = 1$, as the other cases similarly follow in a similar manner.

We start with the analysis of the covariance terms. We collect the coordinate-transformation functions in a matrix $\mathbf{T}(\mathbf{x}) = (\mathbf{t}_1(\mathbf{x}), \dots, \mathbf{t}_p(\mathbf{x})) \in \mathbb{R}^{p \times p}$. By the orthogonality assumption in the definition of a retraction, it follows that $\mathbf{T}(\mathbf{x})$ is an orthogonal matrix for every $\mathbf{x} \in \mathbb{R}^p$. By replacing $\mathbf{T}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\hat{\theta}_n))$ with the rotation matrix $\hat{\mathbf{T}} := \mathbf{T}(\mathbf{H}(\bar{\mathbf{Z}}_n))$ at $\mathbf{H}(\bar{\mathbf{Z}}_n)$ and invoking (28), $(\hat{\Sigma})_{1,1}$ can be approximated as follows:

$$\begin{aligned}
(\hat{\Sigma})_{1,1} &= \left((\bar{\nabla}^2 L_n^{\text{Retr}, \hat{\theta}_n})^{-1} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n}(X_i) \bar{\nabla} L^{\text{Retr}, \hat{\theta}_n}(X_i)^\top / n \right) (\bar{\nabla}^2 L_n^{\text{Retr}, \hat{\theta}_n})^{-1} \right)_{1,1} \\
&= \left((\hat{\mathbf{T}} \bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}} \hat{\mathbf{T}}^\top)^{-1} \right. \\
&\quad \cdot \left(\hat{\mathbf{T}} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(X_i) \bar{\nabla} L^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}}(X_i)^\top / n \right) \hat{\mathbf{T}}^\top \right) \\
&\quad \left. \cdot (\hat{\mathbf{T}} \bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}_n, \{T_{\theta_0 \rightarrow \hat{\theta}_n} e_i\}} \hat{\mathbf{T}}^\top)^{-1} \right)_{1,1} + O(n^{-1} \log n). \tag{67}
\end{aligned}$$

In what follows, we shall switch back to the chart centered at θ_0 using intermediate results established in the proof of Claim 1. Invoking the error controls (63) and (64) on the sum of squared gradient terms and on the sample Hessian matrix, (67) turns into the form that

$$\begin{aligned}
(\hat{\Sigma})_{1,1} &= \left((\hat{\mathbf{T}} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \hat{\mathbf{T}}^\top)^{-1} \right. \\
&\quad \cdot \left(\sum_{i \in [n]} \hat{\mathbf{T}} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i)^\top \hat{\mathbf{T}}^\top / n \right) \\
&\quad \left. \cdot (\hat{\mathbf{T}} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \hat{\mathbf{T}}^\top)^{-1} \right)_{1,1} + O(n^{-1} \log n) \\
&= \left(\hat{\mathbf{T}} (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)))^{-1} \right. \\
&\quad \cdot \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i)^\top / n \right) \\
&\quad \left. \cdot (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)))^{-1} \hat{\mathbf{T}}^\top \right)_{1,1} + O(n^{-1} \log n), \tag{68}
\end{aligned}$$

where the last line holds since $\hat{\mathbf{T}}$ is always an orthogonal matrix.

We proceed to parse the difference $(\Sigma)_{1,1} - (\hat{\Sigma})_{1,1}$ by applying Taylor's expansion to the second term in (68). To facilitate this, we define a function $\phi_{\mathbf{A}}(\mathbf{X}, \mathbf{Z}) : \mathbb{R}^{p \times p} \times \mathbb{R}^{p \times p} \rightarrow \mathbb{R}^{p \times p}$ as

$$\phi_{\mathbf{A}}(\mathbf{X}, \mathbf{Z}) := \mathbf{Z} \mathbf{X}^{-1} \mathbf{A} \mathbf{X}^{-1} \mathbf{Z}^\top,$$

where \mathbf{A} is a fixed p -by- p matrix.

We assume that the smallest singular value of \mathbf{X}_0 , is bounded away from 0 and the largest singular values of \mathbf{X}, \mathbf{Z} are bounded by a constant c_σ . By the Taylor expansion, for every $\Delta_1, \Delta_2 \in \mathbb{R}^{p \times p}$ with $\|\Delta_i\|_F \leq c_\Delta \leq c_\sigma, i = 1, 2$, we have

$$\begin{aligned}
&\phi_{\mathbf{A}}(\mathbf{X}_0 + \Delta_1, \mathbf{Z}_0 + \Delta_2) \\
&= \phi_{\mathbf{A}}(\mathbf{X}_0, \mathbf{Z}_0) - \mathbf{Z}_0 \mathbf{X}_0^{-1} \Delta_1 \mathbf{X}_0^{-1} \mathbf{A} \mathbf{X}_0^{-1} \mathbf{Z}_0^\top - \mathbf{Z}_0 \mathbf{X}_0^{-1} \mathbf{A} \mathbf{X}_0^{-1} \Delta_2 \mathbf{X}_0^{-1} \mathbf{Z}_0^\top \\
&\quad + \Delta_2 \mathbf{X}_0^{-1} \mathbf{A} \mathbf{X}_0^{-1} \mathbf{Z}_0^\top + \mathbf{Z}_0 \mathbf{X}_0^{-1} \mathbf{A} \mathbf{X}_0^{-1} \Delta_1^\top + c_\delta \max\{\|\Delta_1\|_F^2, \|\Delta_2\|_F^2\}, \tag{69}
\end{aligned}$$

where the constant c_δ is determined by c_σ , c_Δ , and $\|\mathbf{A}\|_F$.

Moreover, we introduce the following function:

$$\mathbf{G}(\boldsymbol{\eta}, \mathcal{X}_n) := \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}, X_i)^\top / n, \quad (70)$$

and denote the matrix $\mathbf{G}(\mathbf{H}(\bar{\mathbf{Z}}_n), \mathcal{X}_n)$ by $\widehat{\mathbf{G}}_n$. Applying the above fact (69) to (68) yields that

$$\begin{aligned} & \left(\widehat{\mathbf{T}} \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{T}}^\top \right)_{1,1} \\ &= \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right)_{1,1} \\ &+ \left((\widehat{\mathbf{T}} - \mathbf{I}) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right)_{1,1} \\ &+ \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} (\widehat{\mathbf{T}} - \mathbf{I})^\top \right)_{1,1} \\ &+ \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right) \right. \\ &\quad \cdot \left. \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right)_{1,1} \\ &+ \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right. \\ &\quad \cdot \left. \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right)_{1,1} \\ &+ c' \max \left\{ \left\| \widehat{\mathbf{T}} - \mathbf{I} \right\|_F^2, \left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right\|_F^2 \right\} \end{aligned} \quad (71)$$

holds for some constant c' since $\|\widehat{\mathbf{G}}_n\|$ is upper bounded by some constant under the event \mathcal{F}'_n .

Furthermore, applying a Taylor expansion again to $\widehat{\mathbf{T}} - \mathbf{I}$ and $\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n))$ provides us with a linear approximation in terms of $\mathbf{H}(\bar{\mathbf{Z}}_n)$ as follows

$$\widehat{\mathbf{T}} - \mathbf{I} = \sum_{j \in [p]} \frac{\partial}{\partial x_j} \mathbf{T}(\mathbf{0})(\mathbf{H}(\bar{\mathbf{Z}}_n))_j + c_T \|\mathbf{H}(\bar{\mathbf{Z}}_n)\|_2^2, \quad (72)$$

$$\begin{aligned} & \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \\ &= - \sum_{j \in [p]} (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}} + c_h \|\mathbf{H}(\bar{\mathbf{Z}}_n)\|_2^2 \end{aligned} \quad (73)$$

for some constants c_T and c_h since the fourth-time derivatives of $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ is locally bounded by a constant under the event \mathcal{F}'_n .

What remains to be justified is the quantity $\widehat{\mathbf{G}}_n$. In view of Taylor's expansion of \mathbf{G}_n , $\widehat{\mathbf{G}}_n$ can

be written as

$$\begin{aligned}
\widehat{\mathbf{G}}_n &= \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \\
&+ \sum_{j \in [p]} \left(\sum_{i \in [n]} \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \\
&+ \sum_{j \in [p]} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \\
&+ \sum_{j \in [p]} \sum_{k \in [p]} \int_0^1 (1-t) \frac{\partial^2}{\partial x_i \partial x_j} \mathbf{G}_n(t\mathbf{H}(\bar{\mathbf{Z}}_n)) dt (\mathbf{H}(\bar{\mathbf{Z}}_n))_i (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \\
&= \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \\
&+ \sum_{j \in [p]} \left(\sum_{i \in [n]} \frac{\partial}{\partial x_j} \bar{\nabla}^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \\
&+ \sum_{j \in [p]} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \\
&+ c_G \|\mathbf{H}(\bar{\mathbf{Z}}_n)\|_2^2
\end{aligned} \tag{74}$$

for some constant c_G where the last equality holds since $\frac{\partial^2}{\partial x_i \partial x_j} \mathbf{G}_n(t\mathbf{H}(\bar{\mathbf{Z}}_n))$ for $t \in [0, 1]$ is uniformly upper bounded by some constant c_G under the event \mathcal{F}_n .

Armed with the above approximations, we substitute (71), (72), (73), and (74) into (68), and derive that

$$\begin{aligned}
&(\widehat{\boldsymbol{\Sigma}})_{1,1} \\
&= \left(\widehat{\mathbf{T}} \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{G}}_n \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{T}}^\top \right)_{1,1} + O(n^{-1} \log n) \\
&= (\boldsymbol{\Sigma})_{1,1} + \widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu}) + O(n^{-1} \log n),
\end{aligned} \tag{75}$$

where $\widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu})$ is a quantity determined by $\bar{\mathbf{Z}}_n$ and can be expressed as

$$\begin{aligned}
\widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu}) &:= \left(\mathbf{T}(\mathbf{H}(\boldsymbol{\mu})) \widehat{\boldsymbol{\xi}}(\bar{\mathbf{Z}}_n) \mathbf{T}(\mathbf{H}(\boldsymbol{\mu}))^\top \right)_{1,1} \\
&+ \left(\left(\sum_{j \in [p]} \frac{\partial}{\partial x_j} \mathbf{T}(\mathbf{H}(\boldsymbol{\mu})) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \right) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \mathbf{G}(\mathbf{0}, \mathcal{X}_n) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \right)_{1,1} \\
&+ \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \mathbf{G}(\mathbf{0}, \mathcal{X}_n) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right)^{-1} \left(\sum_{j \in [p]} \frac{\partial}{\partial x_j} \mathbf{T}(\mathbf{H}(\boldsymbol{\mu})) (\mathbf{H}(\bar{\mathbf{Z}}_n))_j \right)^\top \right)_{1,1},
\end{aligned}$$

with

$$\begin{aligned}
\widehat{\boldsymbol{\xi}}(\bar{\mathbf{Z}}_n) &:= \left((\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \mathbf{G}(\mathbf{0}, \mathcal{X}_n) (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} - \boldsymbol{\Sigma} \right)_{1,1} \\
&+ \left((\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \left(\sum_{j \in [p]} \left(\sum_{i \in [n]} \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) H(\bar{\mathbf{Z}}_n)_j \right) \right. \\
&\quad \cdot (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \Big)_{1,1} \\
&+ \left((\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \left(\sum_{j \in [p]} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i) \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_i)^\top / n \right) H(\bar{\mathbf{Z}}_n)_j \right) \right. \\
&\quad \cdot (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \Big)_{1,1} \\
&- \left((\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \left(\sum_{j \in [p]} H(\bar{\mathbf{Z}}_n)_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}} \right) \right. \\
&\quad \cdot (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \mathbf{G}(\mathbf{0}, \mathcal{X}_n) (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \Big)_{1,1} \\
&- \left((\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \mathbf{G}(\mathbf{0}, \mathcal{X}_n) (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \right. \\
&\quad \cdot \left. \left(\sum_{j \in [p]} H(\bar{\mathbf{Z}}_n)_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}} \right) (\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}))^{-1} \right)_{1,1}.
\end{aligned}$$

Here we make use of the fact that $\|\mathbf{H}(\bar{\mathbf{Z}}_n)\|_2$ is of the order $O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$ under the event \mathcal{F}_n . We hereby make note of the quantity $\widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu})$ that all involved derivatives of $L^{\text{Exp}, \theta_0, \{e_i\}}$ or $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ can be viewed as the components of $\bar{\mathbf{Z}}_n$. Thus $\widehat{\boldsymbol{\xi}}_R : \mathbb{R}^{\dim(\bar{\mathbf{Z}}_n)} \times \mathbb{R}^{\dim(\bar{\mathbf{Z}}_n)} \rightarrow \mathbb{R}^{p \times p}$ is a function independently of $L^{\text{Exp}, \theta_0, \{e_i\}}$ or $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ and will be utilized again in the analysis of the resampled quantities.

An immediate consequence of (75) is that

$$|(\boldsymbol{\Sigma})_{1,1} - (\widehat{\boldsymbol{\Sigma}})_{1,1}| = O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$$

under the event \mathcal{F}_n .

Now we are ready to decompose the studentized statistic $-\frac{R_{\widehat{\theta}_n}^{-1}(\theta_0)_i}{(\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}}}$. Combining (66) and the fact that $\|\mathbf{H}(\bar{\mathbf{Z}}_n)\|_2$ is of the order $O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$ under the event \mathcal{F}_n gives that

$$\begin{aligned}
&-\frac{R_{\widehat{\theta}_n}^{-1}(\theta_0)_1}{(\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}}} \\
&= \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}}} + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\
&= \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}}
\end{aligned}$$

$$\begin{aligned}
& + \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}} \left((\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}} - (\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}} \right) \\
& + \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1} (\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}}} \left((\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}} - (\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}} \right)^2 + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}) \\
& = \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& + \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}} \left((\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}} - (\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}} \right) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}) \\
& = \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& + \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{2(\boldsymbol{\Sigma})_{1,1}^{\frac{3}{2}}} \left((\boldsymbol{\Sigma})_{1,1} - (\widehat{\boldsymbol{\Sigma}})_{1,1} \right) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}). \quad (76)
\end{aligned}$$

Substituting (75) into (76) establishes the first part of the claim:

$$\begin{aligned}
& - \frac{R_{\widehat{\theta}_n}(\theta_0)_1}{(\widehat{\boldsymbol{\Sigma}})_{1,1}^{\frac{1}{2}}} \\
& = \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& - \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{2(\boldsymbol{\Sigma})_{1,1}^{\frac{3}{2}}} \widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu}) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}) \\
& = \frac{1}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}} \left(H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j \right. \\
& \quad \left. - \frac{H(\bar{\mathbf{Z}}_n)_1 + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{0}) H(\bar{\mathbf{Z}}_n)_k H(\bar{\mathbf{Z}}_n)_j}{2(\boldsymbol{\Sigma})_{1,1}} \widehat{\boldsymbol{\xi}}_R(\bar{\mathbf{Z}}_n, \boldsymbol{\mu}) \right) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}) \\
& = \frac{\widetilde{H}_1(\bar{\mathbf{Z}}_n, \boldsymbol{\mu})}{(\boldsymbol{\Sigma})_{1,1}^{\frac{1}{2}}} + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}),
\end{aligned}$$

where we define the differentiable function $\widetilde{H}_i(\mathbf{x}, \mathbf{y})$ appearing in (58) as

$$\begin{aligned}
\widetilde{H}_1(\mathbf{x}, \mathbf{y}) & := \sum_{j \in [p]} t_{j,1}(\mathbf{H}(\mathbf{y})) H(\mathbf{x})_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\mathbf{y})) H(\mathbf{x})_k H(\mathbf{x})_j \\
& - \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\mathbf{y})) H(\mathbf{x})_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\mathbf{y})) H(\mathbf{x})_k H(\mathbf{x})_j}{2(\boldsymbol{\Sigma})_{1,1}} \widehat{\boldsymbol{\xi}}_R(\mathbf{x}, \mathbf{y}).
\end{aligned}$$

The proof of (59) closely follows the arguments presented earlier, but proceeds under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$. Below, we provide a brief outline of the main steps.

Recalling the function \mathbf{G} defined in (70), we introduce the quantity:

$$\mathbf{G}(\boldsymbol{\eta}, \mathcal{X}_n^*) := \sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}, X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\boldsymbol{\eta}, X_i^*)^\top / n.$$

We also denote that $\widehat{\mathbf{T}}^* := \mathbf{T}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n))$ and $\widehat{\mathbf{G}}_n^* := \mathbf{G}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n), \mathcal{X}_n^*)$.

Notice that a similar fact to (72) and (73) holds under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$ that

$$\begin{aligned} \widehat{\mathbf{T}}^* - \widehat{\mathbf{T}} &= \sum_{j \in [p]} \frac{\partial}{\partial x_j} T(\bar{\mathbf{Z}}_n) H(\bar{\mathbf{Y}}_n)_j + c'_T \|\mathbf{H}(\bar{\mathbf{Y}}_n)\|_2^2, \\ \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) \\ &= - \sum_{j \in [p]} H(\bar{\mathbf{Y}}_n)_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) + c'_h \|\mathbf{H}(\bar{\mathbf{Y}}_n)\|_2^2. \end{aligned}$$

With the idea of Taylor's expansion in mind together with Claim 1, we can deduce that under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$

$$\begin{aligned} & (\tilde{\boldsymbol{\Sigma}})_{1,1} \\ &= \left(\widehat{\mathbf{T}}^* \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) \right)^{-1} \widehat{\mathbf{G}}_n^* \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n) + \mathbf{H}(\bar{\mathbf{Y}}_n)) \right)^{-1} \widehat{\mathbf{T}}^{*\top} \right)_{1,1} \\ & \quad + O(n^{-1} \log n) \\ &= (\widehat{\boldsymbol{\Sigma}})_{1,1} + \left(\widehat{\mathbf{T}} \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{G}}_n^* \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \widehat{\mathbf{T}}^\top - (\widehat{\boldsymbol{\Sigma}})_{1,1} \right)_{1,1} \\ & \quad + \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right. \\ & \quad \cdot \left(\sum_{j \in [p]} \left(\sum_{i \in [n]} \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i^*) \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i^*)^\top / n \right) H(\bar{\mathbf{Y}}_n)_j \right) \\ & \quad \cdot \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \Big)_{1,1} \\ & \quad + \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right. \\ & \quad \cdot \left(\sum_{j \in [p]} \left(\sum_{i \in [n]} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i^*) \frac{\partial}{\partial x_j} \bar{\nabla} L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n), X_i^*)^\top / n \right) H(\bar{\mathbf{Y}}_n)_j \right) \\ & \quad \cdot \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \Big)_{1,1} \\ & \quad + \left(\left(\sum_{i \in [p]} \frac{\partial}{\partial x_i} \mathbf{T}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H_i(\bar{\mathbf{Y}}_n) \right) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \mathbf{G}_n^*(\mathbf{H}(\bar{\mathbf{Z}}_n)) \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right)_{1,1} \\ & \quad + \left(\left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \mathbf{G}_n^*(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right. \\ & \quad \cdot \left. \left(\bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \left(\sum_{i \in [p]} \frac{\partial}{\partial x_i} \mathbf{T}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H_i(\bar{\mathbf{Y}}_n) \right)^\top \right)_{1,1} \end{aligned}$$

$$\begin{aligned}
& - \left(\left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \left(\sum_{j \in [p]} H(\bar{\mathbf{Y}}_n)_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right) \right. \\
& \cdot \left. \left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \mathbf{G}_n^*(\mathbf{H}(\bar{\mathbf{Z}}_n)) \left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right)_{1,1} \\
& - \left(\left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \mathbf{G}_n^*(\mathbf{H}(\bar{\mathbf{Z}}_n)) \left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right. \\
& \cdot \left. \left(\sum_{j \in [p]} H(\bar{\mathbf{Y}}_n)_j \frac{\partial}{\partial x_j} \bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right) \left(\bar{\nabla}^2 L_n^{* \text{Exp}, \theta_0, \{e_i\}}(\mathbf{H}(\bar{\mathbf{Z}}_n)) \right)^{-1} \right)_{1,1} + O(n^{-1} \log n) \\
& = (\hat{\Sigma})_{1,1} + \hat{\xi}_R(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n) + O(n^{-1} \log n). \tag{77}
\end{aligned}$$

Again, for the resampled studentized term $-\frac{R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_1}{(\check{\Sigma})_{1,1}^{\frac{1}{2}}}$ under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$, we can decompose it as follows

$$\begin{aligned}
& - \frac{R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_1}{(\check{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& = \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{(\check{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& + \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{(\hat{\Sigma})_{1,1}} \left((\hat{\Sigma})_{1,1}^{\frac{1}{2}} - (\check{\Sigma})_{1,1}^{\frac{1}{2}} \right) \\
& + \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,i}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{(\hat{\Sigma})_{1,1} (\check{\Sigma})_{1,1}^{\frac{1}{2}}} \left((\hat{\Sigma})_{1,1}^{\frac{1}{2}} - (\check{\Sigma})_{1,1}^{\frac{1}{2}} \right)^2 + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}) \\
& = \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{(\hat{\Sigma})_{1,1}} \\
& + \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{2(\hat{\Sigma})_{1,1}^{\frac{3}{2}}} \left((\hat{\Sigma})_{1,1} - (\check{\Sigma})_{1,1} \right) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}}), \tag{78}
\end{aligned}$$

where we leverage the concentration conditions under $\mathcal{F}_n \cap \mathcal{F}_n^*$ again.

Plugging (77) into (78) deduces that under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$

$$\begin{aligned}
& - \frac{R_{\hat{\theta}_n^*}^{-1}(\hat{\theta}_n)_1}{(\check{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& = \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{(\hat{\Sigma})_{1,1}^{\frac{1}{2}}} \\
& - \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Y}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_l}{2(\hat{\Sigma})_{1,1}^{\frac{3}{2}}} \hat{\xi}_R(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n) + O(n^{-\frac{3}{2}} (\log n)^{\frac{3}{2}})
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{(\widehat{\Sigma})_{1,1}^{\frac{1}{2}}} \left(\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j \right. \\
&\quad \left. - \frac{\sum_{j \in [p]} t_{j,1}(\mathbf{H}(\bar{\mathbf{Y}}_n)) H(\bar{\mathbf{Y}}_n)_j + \sum_{j,k \in [p]} \frac{\partial t_{j,1}}{\partial x_k}(\mathbf{H}(\bar{\mathbf{Z}}_n)) H(\bar{\mathbf{Y}}_n)_k H(\bar{\mathbf{Y}}_n)_j}{2\widehat{\Sigma}_{j,j}} \widehat{\xi}_R(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n) \right) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}) \\
&=: \frac{\widetilde{H}_1(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n)}{(\widehat{\Sigma})_{1,1}^{\frac{1}{2}}} + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}),
\end{aligned}$$

which completes the proof. \square

A.2 Proof of Proposition 2

The proof of this part inherits the notations from the proof of Theorem 3. Compared with Theorem 3, the extrinsic t -statistic returns to the concise representation under a fixed coordinate system. The condition that $f \circ \text{Exp} \circ \pi_{\{e_i\}}^{-1}$ is smooth in a neighborhood of $\mathbf{0}$, together with (27), (28), Theorem 1, and Theorem 2, reveals that

$$f(\widehat{\theta}_n^*) - f(\widehat{\theta}_n) = f(\text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\mathbf{H}(\bar{\mathbf{Z}}_n))) - f(\text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\mathbf{H}(\bar{\mathbf{Z}}_n^*))) + O(n^{-\frac{3}{2}}(\log n)^{\frac{3}{2}}). \quad (79)$$

Note that $f(\text{Exp}_{\theta_0} \circ \pi_{\{e_i\}}^{-1}(\mathbf{H}(\mathbf{x})))$ is a smooth function. Combining the Edgeworth expansions with respect to $f(\text{Exp} \circ \pi_{\{e_i\}}^{-1}(\mathbf{H}(\bar{\mathbf{Z}}_n)))$ and $f(\text{Exp} \circ \pi_{\{e_i\}}^{-1}(\mathbf{H}(\bar{\mathbf{Z}}_n^*)))$ with (79), it follows that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{-\infty < x < \infty} \left| \mathbb{P} \left[\frac{f(\widehat{\theta}_n^*) - f(\widehat{\theta}_n)}{(\nabla f(\widehat{\theta}_n^*)^\top \widehat{\Sigma}_i \nabla f(\widehat{\theta}_n^*))^{\frac{1}{2}}} \leq x \mid \mathcal{X}_n \right] - \mathbb{P}_{\mathcal{X}_n} \left[\frac{f(\widehat{\theta}_n) - f(\theta_0)}{(\nabla f(\widehat{\theta}_n)^\top \widehat{\Sigma}_i \nabla f(\widehat{\theta}_n))^{\frac{1}{2}}} \leq x \right] \right| \geq n^{-1+\xi} \right] \leq \infty,$$

again, by Proposition 4, Proposition 5, and Lemma 4. \square

A.3 Proof of Proposition 1

The assumptions we make in Proposition 1 is an i.i.d. version of the assumptions in (Andrews, 2002, Theorem 2.(a) and Theorem 2.(b)). More precisely, in the context of the GMM estimator discussion in Andrews (2002) we let the function $g(X_i, \boldsymbol{\eta})$ be $\nabla L(\phi^{-1}(\boldsymbol{\eta}), X_i)$ and Ω be an identity matrix. To apply Theorem 2.(a) and Theorem 2.(b) in Andrews (2002) to the t -statistic and the Wald statistic, respectively, we let d_1 and d_2 be 5 and 4, respectively. Then those results immediately imply that

$$\begin{aligned}
\mathbb{P}[T_{\phi,j} > z_{T_{\phi,j},2,\alpha}^*] &= \alpha + o(n^{-1+\xi}), \\
\mathbb{P}[T_{\phi,j} < z_{T_{\phi,j},1-\alpha}^*] &= \alpha + o(n^{-1+\xi}), \\
\mathbb{P}[W_\phi > z_{W_\phi,\alpha}^*] &= \alpha + o(n^{-\frac{3}{2}+\xi}),
\end{aligned}$$

where $z_{T_{\phi,j},2,\alpha}^*$, $z_{T_{\phi,j},1-\alpha}^*$, and $z_{W_\phi,\alpha}^*$ corresponds to the quantile analogs of the ‘‘exact-version’’ statistics by replacing $\widehat{\theta}_n$ and $\widehat{\theta}_n^*$ with $\widetilde{\theta}_n$ and $\widetilde{\theta}_n^*$, respectively. It remains to prove the distributional proximity between the ‘‘exact’’ versions and the ‘‘two-step-Newton’’ version, which was originally provided by Theorem 1 in Andrews (2002) for the Euclidean Newton updates. Nonetheless, their arguments solely rely on the convergence rate of the approximation instead of in which way it is generated, which can also apply to $\widehat{\theta}_n$ generated in Algorithm 2. This is because the assumptions we make in Proposition 1 can imply the assumptions in Theorem 1 and Theorem 2, and the convergence

rate provided in these theorems is sufficient for proving the distributional proximity. Consequently, by Lemma 4, we conclude the claimed results by a similar argument to the one in Theorem 2 of Andrews (2002). \square

B Normal Coordinate Transformation

B.1 Double Exponential Mapping

In this part, we are going to present one of the essential tools for coordinate analysis, which was first studied in Riemannian geometry and general relativity theory and further developed in Gavrilov (2006, 2007) in the sense of high-order expansion. Here we introduce the formula derived from the result in Gavrilov (2007), which fulfills our third-order requirements.

Lemma 1. *Let (\mathcal{M}, ∇) be a smooth manifold with a smooth, symmetric connection. Given $x \in \mathcal{M}$, we let $U \subseteq \mathcal{M}$ be a geodesically complete neighborhood of x . Then for $v, w \in T_x \mathcal{M}$ with $\|v\|, \|w\| \leq \epsilon$, it has*

$$\begin{aligned} & \text{Log}_x(\text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w))) \\ &= v + w + \frac{1}{6}R(w, v)v + \frac{1}{3}R(w, v)w + \frac{1}{12}\nabla_v R(w, v)v + \frac{1}{24}\nabla_w R(w, v)v \\ &+ \frac{5}{24}\nabla_v R(w, v)w + \frac{1}{12}\nabla_w R(w, v)w + C\epsilon^5 \end{aligned}$$

for a sufficiently small ϵ and a sufficiently large constant C .

Proof. By Equation (3) and the main theorem in Gavrilov (2007), it can be inferred that the terms on the right hand are exactly the derivatives up to the fourth order. Noticing that v, w are parameters in the system of the second-order ODE, the Picard-Lindelöf theorem yields that $h(v, w) := \text{Exp}_{\text{Exp}_x(v)}(T_{x \rightarrow \text{Exp}_x(v)}(w))$ is a smooth function with respect to v, w . By the integral form of the remainder and the boundedness of the fifth-order derivative of $h(v, w)$ around x , the term $r(v, w)$ can be replaced by $C\epsilon^5$ with some constant C . \square

B.2 Consistency of Derivative Operators

This part is aimed at proving the desired consistency between $\nabla^k f(\hat{\theta}_n)$ and $\bar{\nabla}^k f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}(\hat{\theta}_n))$ with $k = 2, 3$. Invoking that $\nabla^k f$ is a $(0, 2)$ tensor, we consider their matrix/tensor forms.

Lemma 2. *Let \mathcal{M} be a dimension- p Riemannian manifold and $\{e_i\}$ is a basis of $T_x \mathcal{M}$. Given $\theta \in \mathcal{M}$ and a third-order differentiable function $f : \mathcal{M} \rightarrow \mathbb{R}^p$, it holds in a neighborhood of θ that*

$$\begin{aligned} & \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{T_{\theta \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^2 f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_\theta(\theta')) \right\|_F \\ & \leq C \left(\left\| \bar{\nabla}^2 f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_\theta(\theta')) \right\|_F \text{dist}(\theta, \theta')^2 + \|\bar{\nabla} f(\theta')\|_2 \text{dist}(\theta, \theta') \right) \end{aligned}$$

for some positive constants C .

Proof. Given $\{e_i\} \in T_\theta \mathcal{M}$ and θ' in a sufficient small neighborhood of θ , we denote the vector bundle induced by the function $\text{Exp} \circ \pi_{\{e_i\}}^{-1}$ at θ' by $\{e_i(\theta')\}$. To begin with, noting that the

exponential mapping is a second-order retraction, Lemma 7 allows us to relate $T_{\theta \rightarrow \theta'} e_j$ and $e_j(\theta')$ with a second-order error:

$$\|e_j(\theta') - T_{\theta \rightarrow \theta'} e_j\| = \left\| \frac{d}{dx_j} \text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}(\theta') - T_{\theta \rightarrow \theta'} e_j \right\| \leq C_1 \text{dist}(\theta, \theta')^2 \quad (80)$$

with some constant C_1 for $j = 1, \dots, p$. Thus, for the Hessian matrices given two different bases, the tensor property of the Hessian operator yields that

$$\begin{aligned} & \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{T_{\theta \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \\ & \leq 2\sqrt{p} C_1 \text{dist}(x, \theta')^2 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \\ & \quad + p C_1^2 \text{dist}(x, \theta')^4 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \\ & \leq C_2 \text{dist}(x, \theta')^2 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \end{aligned} \quad (81)$$

for some constant C_2 .

On the other hand, the compatibility of a Riemannian connection yields that

$$\begin{aligned} & \frac{d}{dx_{j_1}} \frac{d}{dx_{j_2}} f(\text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(y)) = \frac{d}{dx_{j_1}} \langle \nabla f, e_{j_2}(y) \rangle \\ & = \langle \nabla_{e_{j_1}(y)} \nabla f, e_{j_2}(y) \rangle + \langle \nabla f(\text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(y)), \nabla_{e_{j_1}(y)} e_{j_2}(y) \rangle \\ & = \nabla^2 f(e_{j_1}(y), e_{j_2}(y)) + \langle \nabla f(\text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(y)), \nabla_{e_{j_1}(y)} e_{j_2}(y) \rangle. \end{aligned} \quad (82)$$

By the property of geodesics, we have $\nabla_{e_i(\theta)} e_j(\theta) = 0$ and thus $\|\nabla_{e_i(\theta')} e_j(\theta')\| = O(\text{dist}(\theta, \theta'))$ by the smoothness of the connection. Rearranging (82) gives

$$\left| \frac{d}{dx_{j_1}} \frac{d}{dx_{j_2}} f(\theta') - \bar{\nabla}^2 f(e_i(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')), e_j(\pi_{\{e_i\}} \circ \text{Log}_x(\theta'))) \right| \leq C_3 \|\nabla f(\theta')\|_2 \text{dist}(\theta, \theta').$$

for some constant C_3 . Therefore, controlling the entrywise difference of the Hessian matrices yields a bound on the Frobenius norm of the difference:

$$\left\| \bar{\nabla}^2 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) - \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_{j_1}} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \leq C_4 \|\nabla f(\theta')\|_2 \text{dist}(\theta, \theta') \quad (83)$$

holds for some constant C_4 .

Then combining (80) and (82) yields that

$$\begin{aligned} & \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{T_{x \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^2 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \\ & \leq \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_i} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) - \bar{\nabla}^2 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \\ & \quad + C_2 \text{dist}(x, \theta')^2 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F \\ & \leq C_2 \text{dist}(x, \theta')^2 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_j} \text{Exp}_x \circ \pi_{\{e_i\}}^{-1}(\theta')\}}(\mathbf{0}) \right\|_F + C_4 \|\nabla f(\theta')\|_2 \text{dist}(\theta, \theta') \\ & \leq C \left(\left\| \bar{\nabla}^2 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \text{dist}(x, \theta')^2 + \|\nabla f(\theta')\|_2 \text{dist}(\theta, \theta') \right) \end{aligned}$$

holds for some constant C where we make use of (83) again. □

Lemma 3. *Let \mathcal{M} be a Riemannian manifold. Given $\theta \in \mathcal{M}$, a basis $\{e_i\}$ of $\mathbb{T}_\theta\mathcal{M}$, and a third-order differentiable function $f : \mathcal{M} \rightarrow \mathbb{R}^p$, it holds in a neighborhood of θ that*

$$\begin{aligned} & \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta', \{T_{x \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^3 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \\ & \leq C \left(\left\| \bar{\nabla}^3 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \text{dist}(x, \theta')^2 \right. \\ & \quad \left. + \left\| \bar{\nabla}^2 f^{\text{Exp}, x, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_x(\theta')) \right\|_F \text{dist}(x, \theta') + \|\nabla f(\theta')\| \right) \end{aligned}$$

for some constant C .

Proof. Similar to (80) and (81), we have

$$\begin{aligned} & \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta', \{T_{\theta \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^3 f^{\text{Exp}, \theta', \{\frac{d}{dx_i} \text{Exp}_{\theta \circ \pi_{\{e_i\}}^{-1}}(\theta')\}}(\mathbf{0}) \right\|_F \\ & \leq C_1 \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta', \{\frac{d}{dx_i} \text{Exp}_{\theta \circ \pi_{\{e_i\}}^{-1}}(\theta')\}}(\mathbf{0}) \right\|_F \text{dist}(\theta, \theta')^2 \end{aligned}$$

for some constant $C_1 > 0$.

Recall the definition of $e_k(\theta')$ in the proof of Lemma 2. By the definition of $\nabla^3 f$, we have

$$\begin{aligned} & \nabla^3 f[e_i(\theta'), e_j(\theta'), e_k(\theta')] \\ & = e_i(\theta') \langle \nabla^2 f(e_j(\theta')), e_k(\theta') \rangle - \langle \nabla^2 f(e_j(\theta')), \nabla_{e_i(\theta')} e_k(\theta') \rangle - \langle \nabla^2 f(\nabla_{e_i(\theta')} e_j(\theta')), e_k(\theta') \rangle \\ & = e_i(\theta') \left(\frac{d}{dx_j} \frac{d}{dx_k} f(\text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}(y)) - \langle \nabla f, \nabla_{e_j(\theta')} e_k(\theta') \rangle \right) - \langle \nabla^2 f(e_j(\theta')), \nabla_{e_i(\theta')} e_k(\theta') \rangle \\ & \quad - \langle \nabla^2 f(e_k(\theta')), \nabla_{e_i(\theta')} e_j(\theta') \rangle \\ & = \frac{d}{dx_i} \frac{d}{dx_j} \frac{d}{dx_k} f(\text{Exp}_\theta \circ \pi_{\{e_i\}}^{-1}(y)) - \langle \nabla f^2(e_i(\theta')), \nabla_{e_j(\theta')} e_k(\theta') \rangle - \langle \nabla^2 f(e_j(\theta')), \nabla_{e_i(\theta')} e_k(\theta') \rangle \\ & \quad - \langle \nabla^2 f(e_k(\theta')), \nabla_{e_i(\theta')} e_j(\theta') \rangle - \langle \nabla f, \nabla_{e_i(\theta')} \nabla_{e_j(\theta')} e_k(\theta') \rangle. \end{aligned}$$

Again, for $i, j \in [p]$, we have $\|\nabla_{e_i(\theta')} e_j(\theta')\| = O(\text{dist}(\theta, \theta'))$ and $\|\nabla_{e_i(\theta')} \nabla_{e_j(\theta')} e_k(\theta')\|$ is bounded away from ∞ by smoothness.

Therefore, invoking Lemma 2, we arrive at the conclusion that

$$\begin{aligned} & \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta', \{T_{\theta \rightarrow \theta'}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^3 f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_\theta(\theta')) \right\|_F \\ & \leq C_1 \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta', \{\frac{d}{dx_i} \text{Exp}_{\theta \circ \pi_{\{e_i\}}^{-1}}(\theta')\}}(\mathbf{0}) \right\|_F \text{dist}(\theta, \theta')^2 \\ & \quad + C_2 \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta', \{\frac{d}{dx_i} \text{Exp}_{\theta \circ \pi_{\{e_i\}}^{-1}}(\theta')\}}(\mathbf{0}) \right\|_F \text{dist}(\theta, \theta') + C_3 \|\nabla f(\theta')\| \\ & \leq C \left(\left\| \bar{\nabla}^3 f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_\theta(\theta')) \right\|_F \text{dist}(\theta, \theta')^2 \right. \\ & \quad \left. + \left\| \bar{\nabla}^2 f^{\text{Exp}, \theta, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_\theta(\theta')) \right\|_F \text{dist}(\theta, \theta') + \|\nabla f(\theta')\| \right) \end{aligned}$$

for some constants $C_2, C_3, C > 0$. □

C Convergence Rates of (Resampled) Newton Iterations

In this section, we will work on the convergence rates of Newton's iteration (Algorithm 1) with the original data \mathcal{X}_n and the resampled data $\mathcal{X}_n^{[i]}$, $i = 1, \dots, b$, respectively. We also remind that the desired quadratic convergence indeed doesn't rely on the second-order property of retractions; this condition is necessary only for constructing the confidence regions.

Proof of Theorem 1. We begin by focusing on a neighborhood $N := \text{Exp}_{\theta_0}(B(0, \rho))$, ensuring that N is geodesically convex, i.e., for every p, q in N there is a unique minimizing geodesic from p to q lying in N , by Theorem 6.17 in Lee (2018). We also assume that $\theta^{\text{initial}} \in \text{Exp}_{\theta_0}(B(0, c_\rho \rho))$ for some $0 < c_\rho < 1$. We carry out the following analysis under the event \mathcal{F}_n defined in (23) with $\mathbb{P}[\mathcal{F}_n^c] = O(n^{-1}(\log n)^{-2})$. Given the conditions under \mathcal{F}_n and the relation (27), it follows that $\tilde{\theta}_n$ shrinks to θ_0 and thus is contained in $\text{Exp}_{\theta_0}(B(0, c_\rho \rho))$ for sufficiently large n .

Recap that $\hat{\theta}^{(0)}$ represents the initial estimate θ^{initial} and $\theta^{(k)}$, $k = 1, \dots, t$ be the updated point after k steps of Newton's update in Algorithm 1. We aim to demonstrate the targeted convergence rate by inductively establishing that, for $k = 0, \dots, b-1$,

$$\text{dist}(\theta^{(k+1)}, \tilde{\theta}_n) \leq c_t \text{dist}(\theta^{(k+1)}, \tilde{\theta}_n)^2.$$

holds, given $\theta^{(k+1)} \in \text{Exp}_{\theta_0}(B(0, c_\rho \rho))$.

Recall that $\tilde{\theta}_n$ denotes the exact solution of $\nabla L_n(\theta) = 0$ which lies closest to θ_0 . Applying Lemma 6 to ∇L_n at $\tilde{\theta}_n$ gives that

$$\begin{aligned} 0 &= T_{\tilde{\theta}_n \rightarrow \theta^{(k)}} \nabla L_n(\tilde{\theta}_n) = \nabla L_n(\theta^{(k)}) + \nabla^2 L_n(\theta^{(k)}) [\text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)] \\ &+ \int_0^1 (1-t) T_{\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)) \rightarrow \theta^{(k)}} \left(\nabla^3 L_n(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))) [\text{Log}_{\theta^{(k)}}(\tilde{\theta}_n), \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)] \right) dt. \end{aligned} \quad (84)$$

Each term on the right-hand side of equation (84) requires a delicate analysis.

For the Hessian term, a basic fact is that

$$\nabla^2 L_n(\theta^{(k+1)}) [T_{\theta_0 \rightarrow \theta} e_i, T_{\theta_0 \rightarrow \theta} e_j] = \left(\bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}^{(k)}, \{T_{\theta_0 \rightarrow \theta^{(k+1)}}(e_i)\}}(\mathbf{0}) \right)_{i,j}. \quad (85)$$

Further, by Lemma 2, we replace the right hand side of (85) with the Hessian matrix of the function $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ at $\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})$ to obtain that

$$\begin{aligned} &\left\| \bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}^{(k)}, \{T_{\theta_0 \rightarrow \theta^{(k+1)}}(e_i)\}}(\mathbf{0}) - \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) \right\|_F \\ &\leq C \rho^2 \left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) \right\|_F + C \rho \left\| \nabla L_n(\theta^{(k+1)}) \right\|. \end{aligned} \quad (86)$$

for some constant C .

In order to control $\left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) \right\|_F$, we combine (20) and (21) to obtain

that

$$\begin{aligned}
& \left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) \right\|_F \\
& \leq \left(\left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{\pi_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) - \nabla^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right\|_F + \left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) \right\|_F \right) \\
& \leq C_p \rho (\mathbb{E}[C(X_1)] + \left| \sum_{i \in [n]} C(X_i)/n - \mathbb{E}[C(X_1)] \right|) \\
& \quad + \left(\left\| \bar{\nabla}^2 L_n^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}) - \mathbb{E}[\bar{\nabla}^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_1)] \right\|_F + \left\| \mathbb{E}[\bar{\nabla}^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_1)] \right\|_F \right) \\
& \leq C'
\end{aligned} \tag{87}$$

for some constant C_p, C' under the event \mathcal{F}_n .

Analogously, for the gradient term $\nabla L_n(\theta^{(k+1)})$ we have

$$\begin{aligned}
& \left\| \nabla L_n(\theta^{(k+1)}) \right\| \\
& = \left\| \bar{\nabla} L_n^{\text{Exp}, \theta^{(k+1)}, \{T_{\theta_0 \rightarrow \theta^{(k+1)}} e_i\}}(\mathbf{0}) \right\|_2 \\
& \leq 2 \left\| \bar{\nabla} L_n^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\theta^{(k+1)})) \right\|_2 \\
& \leq c_p \rho (\mathbb{E}[C(X_1)] + \left| \sum_{i \in [n]} C(X_i)/n - \mathbb{E}[C(X_1)] \right|) + cn^{-\frac{1}{2}} (\log n)^{\frac{1}{2}} \leq c',
\end{aligned} \tag{88}$$

for some constants c, c', c_p under the event \mathcal{F}_n , where the first inequality follows by Lemma 7.

With an appropriate choice of ρ , this ensures that the minimum singular value of $\bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}^{(k)}, \{T_{\theta_0 \rightarrow \theta^{(k+1)}}(e_i)\}}(\mathbf{0})$ exceeds half of the smallest eigenvalue of $\mathbb{E}[\nabla^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_1)]$, making it invertible with probability at least $1 - O(n^{-1}(\log n)^{-2})$.

For the last term in (84), we decompose it using Lemma 3 as follows:

$$\begin{aligned}
& \sup_{t \in [0, 1]} \sup_{u_1, u_2, u_3 \in \mathbb{T}_{\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))} \mathcal{M}} \left| \nabla^3 L_n(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)))[u_1, u_2, u_3] \right| \\
& \leq \sup_{t \in [0, 1]} \left\| \bar{\nabla}^3 L_n^{\text{Exp}, \text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)), \{T_{\theta_0 \rightarrow \text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))}(e_i)\}}(\mathbf{0}) \right. \\
& \quad \left. - \bar{\nabla}^3 f^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)))) \right\|_F \\
& \quad + \sup_{t \in [0, 1]} \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)))) \right\|_F \\
& \stackrel{(i)}{\leq} C_1 \left(\sup_{t \in [0, 1]} \left\| \bar{\nabla}^3 f^{\text{Exp}, \theta_0, \{e_i\}}(\pi_{\{e_i\}} \circ \text{Log}_{\theta_0}(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)))) \right\|_F + 1 \right) \\
& \stackrel{(ii)}{\leq} C_2
\end{aligned} \tag{89}$$

for some constants C_1 and C_2 under the event \mathcal{F}_n , where (i) holds by Lemma 3, (87), and (88), and (ii) holds by arguments similar to (87).

To the end, we are required to ensure the well-definedness of the geodesics between $\theta^{(k+1)}$ and

the other elements in N . Notice that under the event \mathcal{F}_n ,

$$\begin{aligned}
& \left\| \nabla^2 L_n(\theta^{(k)})^{-1} \nabla L_n(\theta^{(k)}) \right\| \\
& \stackrel{\text{by (84)}}{\leq} \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\| \\
& \quad + \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\|^2 \sup_{t \in [0,1]} \left\| \bar{\nabla}^3 L_n^{\text{Exp,Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)), \{T_{\theta_0 \rightarrow \text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))}^{(e_i)}\}}(\mathbf{0}) \right\|_F \\
& \quad \cdot \left\| (\bar{\nabla}^2 L_n^{\text{Exp,Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)), \{T_{\theta_0 \rightarrow \text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))}^{(e_i)}\}}(\mathbf{0}))^{-1} \right\| \\
& \leq C_3 \left\| \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n) \right\| \left(1 + C_3 \left\| \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n) \right\| \right) \leq C_4 \text{dist}(\theta^{(k)}, \tilde{\theta}_n), \\
& \quad \text{dist}(\hat{\theta}^{(k)}, \theta^{(k+1)}) = \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\| \stackrel{\text{by Definition 1}}{\leq} c'_R \left\| \nabla^2 L_n(\theta^{(k)})^{-1} \nabla L_n(\theta^{(k)}) \right\|
\end{aligned}$$

hold for some constants C_3 , C_4 , and c'_R . Therefore, a proper choice of c_ρ and ρ ensures that the distance $\text{dist}(\theta_0, \theta^{(k+1)}) \leq \text{dist}(\theta_0, \tilde{\theta}_n) + \text{dist}(\tilde{\theta}_n, \theta^{(k+1)}) + \text{dist}(\theta^{(k+1)}, \theta^{(k+1)}) < \rho$. By the definition of a geodesically convex neighborhood, a geodesic that lies in N and connects $\theta^{(k+1)}$ with a point in N is always unique.

Now we are prepared to prove the quadratic convergence rate of Newton's iteration. By Lemma 1 and the well-definedness of logarithmic mapping related to $\theta^{(k+1)}$, there exist some constants c_t, c_R such that

$$\begin{aligned}
& \left\| \text{Log}_{\tilde{\theta}_n}(\theta^{(k+1)}) \right\| = \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) - \text{Log}_{\theta^{(k+1)}}(\theta^{(k+1)}) \right\| \\
& \quad + c_t \left(\left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\|^2 \left\| \text{Log}_{\theta^{(k+1)}}(\theta^{(k+1)}) \right\| + \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\| \left\| \text{Log}_{\theta^{(k+1)}}(\theta^{(k+1)}) \right\|^2 \right) \\
& \stackrel{(i)}{\leq} \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) + \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\| \\
& \quad + c_t \left(\left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\|^2 \left\| \text{Log}_{\theta^{(k+1)}}(\theta^{(k+1)}) \right\| + \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n) \right\| \left\| \text{Log}_{\theta^{(k+1)}}(\theta^{(k+1)}) \right\|^2 \right) \\
& \quad + c_R \left\| \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\|^3
\end{aligned}$$

where (i) holds by the property of a second-order retraction and (84).

Since the operator $\nabla^2 L_n(\theta^{(k)})$ is invertible with probability at least $1 - O(n^{-1}(\log n)^{-2})$, rearranging (84) and plugging the above inequalities gives the desired quadratic convergence rate:

$$\begin{aligned}
& \text{dist}(\tilde{\theta}_n, \theta^{(k+1)}) = \left\| \text{Log}_{\tilde{\theta}_n}(\theta^{(k+1)}) \right\| \\
& = \left\| \int_0^1 (1-t) T_{\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)) \rightarrow \theta^{(k)}} \left(\nabla^3 L_n(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n))) (\text{Log}_{\theta^{(k)}}(\tilde{\theta}_n), \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n)) \right) dt \right\| \\
& \quad + c_t \left(\left\| \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n) \right\|^2 \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\| + \left\| \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n) \right\| \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\|^2 \right) \\
& \quad + c_R \left\| \nabla^2 L_n(\theta^{(k)})^{-1} \nabla L_n(\theta^{(k)}) \right\|^3 \\
& \leq c_l \text{dist}(\tilde{\theta}_n, \theta^{(k)})^2 \leq c_\rho \rho
\end{aligned} \tag{90}$$

holds for some constant c_l and sufficiently large n , where we make use of the fact that $\left\| \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n) \right\| = \text{dist}(\theta^{(k)}, \tilde{\theta}_n)$.

Finally, the conclusion is a direct consequence of the quadratic convergence rate (90) that we have proved. \square

Proof of Theorem 2. Since this proof closely follows the structure of Theorem 1, we provide only a sketch of the argument below. Let \mathcal{X}_n^* be a generic data collection obtained from resampling \mathcal{X}_n with replacement. Throughout the proof, we focus on the event $\mathcal{F}_n \cap \mathcal{F}_n^*$ defined in (23) and (39). Our goal is to investigate the distance $\text{dist}(\theta^{(k+1)}, \tilde{\theta}_n^*)$, provided $\theta^{(k)} \in \text{Exp}_{\theta_0}(B(\theta_0, c_\rho\rho))$ for $k = 0, 1$ and the constants ρ and c_ρ in the proof of Theorem 1.

A similar application of Lemma 6 yields that

$$0 = T_{\tilde{\theta}_n^* \rightarrow \theta^{(k)}} \nabla L_n(\tilde{\theta}_n^*) = \nabla L_n(\theta^{(k)}) + \nabla^2 L_n(\theta^{(k)}) (\text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*)) \\ + \int_0^1 (1-t) T_{\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*)) \rightarrow \theta^{(k)}} \left(\nabla^3 L_n(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*))) (\text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*), \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*)) \right) dt.$$

In terms of the control of $\nabla^2 L_n(\theta^{(k)})$ and $\nabla^3 L_n(\theta^{(k)})$, similar treatments to (86), (87), (88), (89) but replacing the concentrations based on (20) and (21) with applying (34) and the conditions in (40) gives that

1. the minimum singular value of $\bar{\nabla}^2 L_n^{\text{Exp}, \hat{\theta}^{(k)}, \{T_{\theta_0 \rightarrow \theta^{(k+1)}}(e_i)\}}$ ($\mathbf{0}$) is greater than half of the smallest singular value of $\mathbb{E}[\nabla^2 L^{\text{Exp}, \theta_0, \{e_i\}}(\mathbf{0}, X_1)]$,
2. $\sup_{t \in [0,1]} \sup_{u_1, u_2, u_3 \in \mathbb{T}_{\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*))} \mathcal{M}} \left| \nabla^3 L_n(\text{Exp}_{\theta^{(k)}}(t \text{Log}_{\theta^{(k)}}(\tilde{\theta}_n^*))) [u_1, u_2, u_3] \right|$ is upper bounded by some constant.
3. $\hat{\theta}^{(k+1)} \in \text{Exp}_{\theta_0}(B(0, c_\rho\rho))$.

Thus $\left\| \text{Log}_{\tilde{\theta}_n^*}(\theta^{(k+1)}) \right\|$ is upper bounded by

$$\text{dist}(\tilde{\theta}_n^*, \theta^{(k+1)}) = \left\| \text{Log}_{\tilde{\theta}_n^*}(\theta^{(k+1)}) \right\| \\ \leq \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n^*) + \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\| \\ + c_t \left(\left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n^*) \right\|^2 \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\| + \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n^*) \right\| \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\|^2 \right) \\ + c_R \left\| \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\|^3 \\ \leq c_l^* \text{dist}(\tilde{\theta}_n^*, \theta^{(k+1)})^2$$

for some constant c_l^* , where we use the facts:

$$\left\| \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\| \leq \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n^*) \right\| (1 + C_2 \left\| \text{Log}_{\theta^{(k+1)}}(\tilde{\theta}_n^*) \right\|) \leq C_4^* \text{dist}(\theta^{(k)}, \tilde{\theta}_n^*), \\ \left\| \text{Log}_{\hat{\theta}^{(k)}}(\hat{\theta}^{(k+1)}) \right\| \leq c_R^* \left\| \nabla^2 L_n(\theta^{(k+1)})^{-1} \nabla L_n(\theta^{(k+1)}) \right\|$$

for some constants C_4^* and c_R^* .

Finally, the conclusion that $\text{dist}(\theta^{(2)}, \tilde{\theta}_n^*) = O(n^{-2}(\log n)^2)$ follows from the quadratic convergence under the event $\mathcal{F}_n \cap \mathcal{F}_n^*$ since $\text{dist}(\tilde{\theta}_n^*, \theta^{\text{initial}}) = O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$. \square

D Classic Edgeworth Expansion Results

In this section, we restate the settings considered in [Bhattacharya et al. \(1978\)](#); [Bhattacharya \(1987\)](#); [Denker and Bhattacharya \(1990\)](#) and provide a version with some slight but necessary

modifications. Consider p -dimensional i.i.d. observations \mathbf{X}_j , $j \in [n]$ with common distribution G . Write

$$\begin{aligned}\boldsymbol{\mu} &:= \mathbb{E}\mathbf{X}_j, \\ \mathbf{V} &= (v_{rr'})_{r,r' \in [p]} := \text{Cov}(\mathbf{X}_1).\end{aligned}$$

Let $\mathbf{H}(\mathbf{x}) = (H^{(1)}(\mathbf{x}), H^{(2)}(\mathbf{x}), \dots, H^{(k)}(\mathbf{x}))$ be a measurable function defined on \mathbb{R}^k . We consider the asymptotic distribution of $H(\bar{\mathbf{X}})$, where

$$\bar{\mathbf{X}}_n := (\mathbf{X}_1 + \dots + \mathbf{X}_n)/n.$$

The assumptions are as follows:

- (A₁) $\mathbb{E}\|\mathbf{X}_1\|_2^s < \infty$ for an integer $s \geq 3$.
- (A₂) Derivatives of $H^{(i)}(\mathbf{x})$, $i \in [p]$ of order up to and including $s - 1$ exist and are continuous in a neighborhood of $\boldsymbol{\mu}$.
- (A₃) $\bar{\nabla}H^{(i)}(\boldsymbol{\mu}) := (\partial/\partial x^{(1)}H^{(i)}(\mathbf{x}), \partial/\partial x^{(2)}H^{(i)}(\mathbf{x}), \dots, \partial/\partial x^{(k)}H^{(i)}(\mathbf{x}))|_{\mathbf{x}=\boldsymbol{\mu}}$, $1 \leq i \leq p$, are linearly independent elements of \mathbb{R}^p .
- (A₄) The distribution of \mathbf{X}_1 satisfies Cramèr's condition, namely, $\limsup_{\|t\|_2 \rightarrow \infty} |\mathbb{E}[e^{it^\top \mathbf{X}_1}]| < 1$.

In view of the population, the following result from [Denker and Bhattacharya \(1990\)](#) provides us with the consistency guarantee of the population Edgeworth expansion.

Proposition 4 (Theorem 2.1 in [Denker and Bhattacharya \(1990\)](#)). *Under the assumptions (A₁) - (A₄), one has*

$$\sup_{B \in \mathcal{B}} \left| \mathbb{P}[\sqrt{n}(\mathbf{H}(\bar{\mathbf{X}}_n) - \mathbf{H}(\boldsymbol{\mu})) \in B] - \int_B \psi_{s,n}(\mathbf{x}) d\mathbf{x} \right| = o(n^{-\frac{(s-2)}{2}}),$$

for all class $\mathcal{B} \in \mathcal{B}^p$ (the collection of Borel sets in \mathbb{R}^p) such that, for some $a > 0$,

$$\sup_{B \in \mathcal{B}} \Phi_{\Sigma}((\partial B)^\epsilon) = O(\epsilon^a), \quad \text{as } \epsilon \downarrow 0.$$

Remark 5. Note that $\psi_{s,n}(\mathbf{x})$ takes the form of $e^{-\frac{\mathbf{x}^\top \Sigma^{-1} \mathbf{x}}{2}}$ multiplied by a polynomial of \mathbf{x} where the coefficients in the polynomial are determined by the moments of \mathbf{X} and the derivatives of H as discussed in [Bhattacharya et al. \(1978\)](#). For the sake of completeness, we restate the details as follows:

We first reduce the target function \mathbf{H} to its $(s - 1)$ -order Taylor expansion \mathbf{H}_{s-1} at $\boldsymbol{\mu}$ by the fact that $\sqrt{n}(\mathbf{H}(\bar{\mathbf{X}}_n) - \mathbf{H}_{s-1}(\bar{\mathbf{X}}_n)) = o_{\mathbb{P}}(n^{-\frac{s-2}{2}})$ and the delta method ([Lemma 4](#)). We denote the ν -th cumulant of $\sqrt{n}(\mathbf{H}_{s-1}(\bar{\mathbf{X}}_n) - \mathbf{H}_{s-1}(\boldsymbol{\mu}))$ by $\kappa_{\nu,n}$ ¹¹, and denote the approximate ν -th cumulant with respect to $\kappa_{\nu,n}$ by

$$\tilde{\kappa}_{\nu,n} = \begin{cases} \Sigma_{i,j} + \sum_{l=1}^{s-2} a_{\nu,l} n^{-\frac{l}{2}} & \text{if } \nu = \{i, j\}; \\ \sum_{l=1}^{s-2} a_{\nu,l} n^{-\frac{l}{2}} & \text{otherwise} \end{cases},$$

¹¹Since H_k is a polynomial function, it is straightforward to see that $\kappa_{\nu,n}$ is of the form $\kappa_{\nu,n} = \sum_{k=0}^{(s-2)|\nu|} a_{\nu,k} n^{-\frac{k}{2}}$.

after removing the terms related to $n^{-\frac{k}{2}}$ for $l \geq s-1$ where the coefficients $\{a_{\nu,l}\}$ depend on the cumulants of \mathbf{X}_1 and the coefficients of \mathbf{H}_k ; in fact, by [James \(1958\)](#) we have $\tilde{\kappa}_{\nu,n} = O(n^{-\frac{|\nu|-2}{2}})$ for $|\nu| \geq 3$.

Expanding the characteristic function of $\sqrt{n}(\mathbf{H}(\bar{\mathbf{X}}_n) - \mathbf{H}(\boldsymbol{\mu}))$ gives that for each \mathbf{t} one has

$$\begin{aligned} & \mathbb{E}[\exp(\sqrt{n}i\mathbf{t}^\top(\mathbf{H}(\bar{\mathbf{X}}_n) - \mathbf{H}(\boldsymbol{\mu})))] \\ &= \exp\left(-\frac{\mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t}}{2}\right) \exp\left(i \sum_{i \in [k]} t_i \tilde{\kappa}_{\{i\},n} + \frac{-1}{2} \sum_{i,j \in [k]} (\tilde{\kappa}_{\{i,j\},n} - \Sigma_{i,j}) t_i t_j + \sum_{|\nu| \leq s} \frac{i^{|\nu|}}{\nu!} \tilde{\kappa}_{\nu,n} \mathbf{t}^\nu\right) \\ &= o(n^{-\frac{s-2}{2}}), \end{aligned}$$

where $\boldsymbol{\Sigma} = (\Sigma_{i,j})_{i,j \in [p]}$. Then expanding the exponential term $\exp\left(i \sum_{i \in [k]} t_i \tilde{\kappa}_{\{i\},n} + \frac{-1}{2} \sum_{i,j \in [k]} (\tilde{\kappa}_{\{i,j\},n} - \Sigma_{i,j}) t_i t_j + \sum_{|\nu| \leq s} \frac{1}{\nu!} \tilde{\kappa}_{\nu,n} \mathbf{t}^\nu\right)$ implies the existence of polynomials $\{\pi_i\}_{i=1}^{s-2}$ whose coefficients do not depend on n but depend on the coefficients $\{a_{\nu,l}\}$ in $\{\tilde{\kappa}_{\nu,n}\}$ such that

$$\mathbb{E}[\exp(\sqrt{n}i\mathbf{t}^\top(\mathbf{H}(\bar{\mathbf{X}}_n) - \mathbf{H}(\boldsymbol{\mu})))] = \exp\left(-\frac{\mathbf{t}^\top \boldsymbol{\Sigma} \mathbf{t}}{2}\right) \left(1 + \sum_{i=1}^{s-2} n^{-\frac{i}{2}} \pi_i(i\mathbf{t})\right) + o(n^{-\frac{s-2}{2}}).$$

In the end, we define the formal edgeworth expansion $\psi_{s,n}$ as

$$\psi_{s,n}(x) = \left(1 + \sum_{i=1}^{s-2} n^{-\frac{i}{2}} \pi_i\left(-\frac{d}{dx}\right)\right) \exp\left(-\frac{\mathbf{x}^\top \boldsymbol{\Sigma}^{-1} \mathbf{x}}{2}\right).$$

Regarding the bootstrap procedure, we shall provide a modified version of Theorem 5.3 in [Bhattacharya \(1987\)](#) to accommodate the impact of coordinate chart adjustment. We first consider a continuous function $\tilde{\mathbf{H}} : \mathbb{R}^k \times \mathbb{R}^k \rightarrow \mathbb{R}^p$ satisfying that

1. $\tilde{\mathbf{H}}(\mathbf{x}, \boldsymbol{\mu}) = \mathbf{H}(\mathbf{x})$, and $\tilde{\mathbf{H}}(\cdot, \mathbf{x}')$ is $(s-1)$ -times continuously differentiable on O for every $\mathbf{x}' \in O$ where O is an open neighborhood of $\boldsymbol{\mu}$.
2. The first $(s-1)$ -times derivatives of $\tilde{\mathbf{H}}(\mathbf{x}_1, \mathbf{x}_2)$ with respect to the first variable $\mathbf{x}_1 \in O$ are continuous with respect to $\mathbf{x}_2 \in O$.

Proposition 5 (A Modified Version of Theorem 5.3 in [Bhattacharya \(1987\)](#)). *If Assumption (A₁) is strengthened to $\mathbb{E}\|\mathbf{X}_1\|_2^{2s} < \infty$ for some integer $s \geq 3$, and the assumptions (A₂) - (A₄) are satisfied, then one has*

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}} \left| \mathbb{P}[\sqrt{n}(\tilde{\mathbf{H}}(\bar{\mathbf{X}}_n^*, \bar{\mathbf{X}}_n) - \tilde{\mathbf{H}}(\bar{\mathbf{X}}_n, \bar{\mathbf{X}}_n)) \in B | \mathcal{X}_n] - \int_B \tilde{\psi}_{s,n}(\mathbf{x}) d\mathbf{x} \right| \geq cn^{-\frac{(s-1)}{2}} \right] < \infty \quad (91)$$

for every constant $c > 0$ as $n \rightarrow \infty$, for every class \mathcal{B} of Borel sets satisfying, for some $a > 0$,

$$\sup_{B \in \mathcal{B}} \Psi_{\boldsymbol{\Sigma}}((\partial B)^\epsilon) = O(\epsilon^a), \quad \text{as } \epsilon \downarrow 0.$$

Here the coefficients of the polynomial in $\tilde{\psi}_{s,n}$ depends on the empirical moments of $\{\mathbf{X}_i\}_{i \in [n]}$ as well as the derivatives up to $s-1$ times of $\tilde{\mathbf{H}}(\cdot, \bar{\mathbf{X}}_n)$ evaluated at $\bar{\mathbf{X}}_n$. Moreover, the coefficients of $\tilde{\psi}_{s,n}$ converge to their population counterparts at a rate of $O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$ with probability at least $1 - O(n^{-\frac{s-2}{2}}(\log n)^{-\frac{s}{2}})$.

Proof. Following the proof of Theorem 5.2 in [Bhattacharya \(1987\)](#) together with moment control similar to (31), we have

$$n\mathbb{P}_{\mathcal{X}_n} \left[\sup_{B \in \mathcal{B}} \left| \mathbb{P}[\sqrt{n}(\bar{\mathbf{X}}_n^* - \bar{\mathbf{X}}_n) \in B | \mathcal{X}_n] - \int_B \left[1 + \sum_{r=1}^{s-2} n^{-\frac{r}{2}} \widehat{\psi}_r(\mathbf{x}) \right] \phi_{\bar{\mathbf{V}}}(\mathbf{x}) d\mathbf{x} \right| \geq cn^{-\frac{s-1}{2}} \right] < \infty \quad (92)$$

for every constant $c > 0$ as $n \rightarrow \infty$, for every class \mathcal{B} satisfying, for some $a > 0$, $\sup_{B \in \mathcal{B}} \Psi_{\mathbf{V}}((\partial B)^\epsilon) = O(\epsilon^a)$, as $\epsilon \downarrow 0$, where $\{\widehat{\phi}_r\}$ are polynomials that do not depend on n but depend on the moments of \mathbf{X}_1^* , and $\bar{\mathbf{V}}$ denotes the sample covariance matrix of $\{\mathbf{X}_i\}_{i=1}^n$. Then the reduction step from (92) to (91) follows by the same arguments as the one in [Bhattacharya et al. \(1978\)](#), with an observation from Remark 5 that the coefficients in the formal Edgeworth expansion only depends on the moments of \mathbf{X}_1^* conditional on \mathcal{X}_n and the derivatives up to $s - 1$ times of $\widetilde{\mathbf{H}}(\cdot, \bar{\mathbf{X}}_n)|_{\bar{\mathbf{X}}_n}$. The second part follows from the continuity of $\widetilde{\mathbf{H}}$ and the moderate deviation (Lemma 8). \square

Now we turn back to the preceding analysis in Section A.1.1 and Section A.1.2. One subtle aspect of this analysis involves approximating $\widetilde{\boldsymbol{\eta}}_n^* - \widetilde{\boldsymbol{\eta}}$ by a smooth function of the derivatives of $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ evaluated at $\widetilde{\boldsymbol{\eta}}_n$. To leverage the assumptions we made on the derivatives of L at θ_0 , we shall utilize the next proposition to facilitate the replacement of derivatives.

Proposition 6. *Instate the assumptions in Theorem 3 and consider $\bar{\mathbf{Z}}_n$ in (18), its resampled counterpart $\bar{\mathbf{Z}}_n^*$, and $\bar{\mathbf{Y}}_n$ defined in (43). Given a function $\widetilde{\mathbf{H}}$ as introduced in Proposition 5, it holds almost surely that*

$$\begin{aligned} n\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left\| \left(\widetilde{\mathbf{H}}(\bar{\mathbf{Z}}_n^*, \bar{\mathbf{Z}}_n) - \widetilde{\mathbf{H}}(\bar{\mathbf{Z}}_n, \bar{\mathbf{Z}}_n) \right) - \left(\widetilde{\mathbf{H}}(\bar{\mathbf{Y}}_n, \bar{\mathbf{Z}}_n) - \widetilde{\mathbf{H}}(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n], \bar{\mathbf{Z}}_n) \right) \right\|_2 \leq cn^{-1} \log n | \mathcal{X}_n \right] \right. \\ \left. \leq 1 - cn^{-1}(\log n)^{-2} \right] < \infty \end{aligned}$$

for some constant c .

Proof. We shall discuss a univariate \widetilde{H} since the proof for a multivariate $\widetilde{\mathbf{H}}$ simply follows a similar argument. We denote the (random) function $H(\cdot, \bar{\mathbf{Z}}_n)$ by $g(\cdot)$ herein. Expanding the function g at $\bar{\mathbf{Z}}_n$ and at $\bar{\mathbf{Y}}_n$ respectively yields that

$$\begin{aligned} & (g(\bar{\mathbf{Z}}_n^*) - g(\bar{\mathbf{Z}}_n)) - (g(\bar{\mathbf{Y}}_n) - g(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n])) \\ &= (\bar{\nabla} g(\bar{\mathbf{Z}}_n))^\top (\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n) - (\bar{\nabla} g(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]))^\top (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) + \Xi_1 - \Xi_2 \\ &= (\bar{\nabla} g(\bar{\mathbf{Z}}_n))^\top \left((\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n) - (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \right) + \Xi_1 - \Xi_2 \\ & \quad + (\bar{\nabla} g(\bar{\mathbf{Z}}_n) - \bar{\nabla} g(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n])) (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \end{aligned} \quad (93)$$

where Ξ_1, Ξ_2 are the residual terms of the Taylor expansions. By (41) and similar reasoning in its proof, it holds that

$$\begin{aligned} n\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left\| \bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n] \right\|_2 \leq c_1 (\log n)^{\frac{1}{2}} n^{-\frac{1}{2}}, \quad \left\| \bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n \right\|_2 \leq c_1 (\log n)^{\frac{1}{2}} n^{-\frac{1}{2}}, \right. \right. \\ \left. \left| \Xi_1 \right| \leq c_1 (\log n) n^{-1}, \quad \left| \Xi_2 \right| \leq c_1 (\log n) n^{-1}, \right. \\ \left. (\bar{\nabla} g(\bar{\mathbf{Z}}_n) - \bar{\nabla} g(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n])) (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \leq c_1 (\log n) n^{-1} | \mathcal{X}_n \right] \leq 1 - c_1 n^{-1} (\log n)^{-2} \right] < +\infty \end{aligned}$$

for some constant c_1 .

Then we look into the term $(\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n) - (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n])$. By definition and Taylor's expansion, we have

$$\begin{aligned} & (\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n) - (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \\ &= (\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Y}}_n) - (\bar{\mathbf{Z}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \\ &= -(\mathbf{\Pi}_1 - \mathbf{\Pi}_2)^\top \tilde{\boldsymbol{\eta}}_n + \boldsymbol{\Xi}_3 + \boldsymbol{\Xi}_4, \end{aligned}$$

where $\mathbf{\Pi}_i, i = 1, 2$ are the vectors that collect the derivatives of $L_n^{\text{Exp}, \theta_0, \{e_i\}}$ and $L_n^{* \text{Exp}, \theta_0, \{e_i\}}$ up to the fifth time evaluated at $\mathbf{0}$, respectively, and $\boldsymbol{\Xi}_3$ and $\boldsymbol{\Xi}_4$ are the residual terms. By a similar moderate deviation argument invoking Lemma 8, we arrive at

$$\begin{aligned} n\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\|\mathbf{\Pi}_1 - \mathbf{\Pi}_2\|_2 \leq c_2(\log n)^{\frac{1}{2}} n^{-\frac{1}{2}}, \right. \right. \\ \left. \left. \|\boldsymbol{\Xi}_1\|_2 \leq c_2(\log n)^{\frac{1}{2}} n^{-\frac{1}{2}}, \|\boldsymbol{\Xi}_2\|_2 \leq c_2(\log n)^{\frac{1}{2}} n^{-\frac{1}{2}} | \mathcal{X}_n \right] \leq 1 - c_2 n^{-1} (\log n)^{-2} \right] < \infty \end{aligned}$$

for some constant c_2 . Together with the fact that $\|\tilde{\boldsymbol{\eta}}_n\|_2 = O(n^{-\frac{1}{2}}(\log n)^{\frac{1}{2}})$ under \mathcal{F}_n , we derive that

$$n\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left\| (\bar{\mathbf{Z}}_n^* - \bar{\mathbf{Z}}_n) - (\bar{\mathbf{Y}}_n - \mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n]) \right\|_2 \leq c_9(\log n)n^{-1} | \mathcal{X}_n \right] \leq 1 - c_3 n^{-1} (\log n)^{-2} \right] < \infty.$$

Plugging these bounds into (93) concludes that

$$\begin{aligned} n\mathbb{P}_{\mathcal{X}_n} \left[\mathbb{P} \left[\left| (g(\bar{\mathbf{Z}}_n^*) - g(\bar{\mathbf{Z}}_n)) - (g(\bar{\mathbf{Y}}_n) - g(\mathbb{E}[\bar{\mathbf{Y}}_n | \mathcal{X}_n])) \right| \leq c(\log n)n^{-1} | \mathcal{X}_n \right] \right. \\ \left. \leq 1 - cn^{-1}(\log n)^{-2} \right] < \infty \end{aligned}$$

holds for some constant c . □

E Auxiliary Lemmas

An elementary tool in this paper is called the delta method in the distribution theory, which allows us to neglect small terms in expansions. For the sake of completeness, we restate Lemma 5 in Andrews (2002) in the following.

Lemma 4 (Lemma 5 in Andrews (2002)). *1. Let $\{A_n\} \in \mathbb{R}^{L_A}$ be a sequence of random vectors with an Edgeworth expansion with coefficients of order $O(1)$ satisfying that*

$$\lim_{n \rightarrow \infty} n^a \sup_{\mathbf{x} \in \mathbb{R}^{L_A}} \left| \mathbb{P}[(A_n)_i \leq x_i, i \in [L_A]] - \int_{z_i \leq x_i} \left[1 + \sum_{i=1}^{2a} n^{-\frac{i}{2}} \pi_i(\partial/\partial z) \right] \phi_{\boldsymbol{\Sigma}_n}(\mathbf{z}) d\mathbf{z} \right| = 0$$

for some a with $2a$ being a positive integer, where $\phi_{\boldsymbol{\Sigma}_n}$ denotes the density function of $\mathcal{N}(\mathbf{0}, \boldsymbol{\Sigma}_n)$ and the eigenvalues of $\boldsymbol{\Sigma}_n$'s are bounded away from 0 and ∞ . Let $\{\xi_N\} \in \mathbb{R}^{L_A}$ be a sequence of random vectors with $\mathbb{P}[\|\xi_n\|_2 > \varrho_n] = o(n^{-b})$ for $\varrho_n = o(n^{-b})$ and some $0 < b \leq a$. Then

$$\lim_{n \rightarrow \infty} \sup_{x \in \mathbb{R}} n^b |\mathbb{P}[A_n + \xi_n \leq z] - \mathbb{P}[A_n \leq z]| = 0.$$

2. Let $\{A_n^*, n \geq 1\} \in \mathbb{R}^{L_A}$ be a sequence of bootstrap random vectors that possesses

$$\lim_{n \rightarrow \infty} n^a \mathbb{P}_{\mathcal{X}_n} \left[n^a |\mathbb{P}[A_n^* \leq z | \mathcal{X}_n] - [1 + \sum_{i=1}^{2a} n^{-\frac{i}{2}} \pi_i^*(\partial/\partial z)] \Phi_{\Sigma_n^*}(z)| \right] = 0$$

for some a with $2a$ being a positive integer, where the coefficients of polynomials $\pi_i^*(\partial/\partial z)$ are of order $O(1)$ and the smallest eigenvalue of Σ_n^* is bounded away from 0 and ∞ , except in a sequence of sets with probability $o(N^{-a})$. Let $\{\xi_n^*, n \geq 1\} \in \mathbb{R}^{L_A}$ be a sequence of random vectors with $\lim_{n \rightarrow \infty} n^b \mathbb{P}[\mathbb{P}^*[\|\xi_n\|_2 > \varrho_n] > n^{-b}] = 0$ for some sequence $\varrho_n = o(n^{-b})$ with $b \leq a$. Then

$$\lim_{n \rightarrow \infty} n^b \mathbb{P}_{\mathcal{X}_n} \left[\sup_{z \in \mathbb{R}^{L_A}} |\mathbb{P}[A_n^* + \xi_n^* \leq z | \mathcal{X}_n] - \mathbb{P}[A_n^* \leq z | \mathcal{X}_n]| > n^{-b} \right] = 0.$$

Lemma 5 (Taylor Expansion of a Real-valued Function). *Let \mathcal{M} be a dimension- p Riemannian manifold and $f : \mathcal{M} \rightarrow \mathbb{R}$ is a k -times differentiable function on \mathcal{M} . Then for $\theta \in \mathcal{M}$, there exists a positive ρ such that for arbitrary $\theta \in \text{Exp}_\theta(B(0, \rho))$, Log_θ defined on $\text{Exp}_{\theta_0}(B(0, \rho))$ is a diffeomorphism onto its codomain in $\mathbb{T}_\theta \mathcal{M}$. Moreover, given a fixed normal basis of $\mathbb{T}_\theta \mathcal{M}$ and $\theta' \in B(\theta, \rho)$ it holds that*

$$f(\theta') - f(\theta) = \sum_{i=1}^{k-1} \sum_{\nu: |\nu|=i} \frac{1}{\nu!} \nabla^\nu f|_{x=\theta} [h^\nu] + \frac{1}{k!} \frac{d}{dt} f \circ \text{Exp}(t \text{Log}_\theta(\theta')) \Big|_{t=t_0} \|\text{Log}_\theta(\theta')\|^k,$$

where $t_0 \in [0, \|\text{Log}_\theta(\theta')\|]$.

Proof. The first part is a well-known conclusion in Riemannian geometry. And the Taylor expansion is an immediate consequence of the Taylor expansion of $f \circ \text{Exp} \circ \pi_{\{e_i\}}^{-1}$ on \mathbb{R}^p . \square

Lemma 6 (Taylor Expansion of a $(0, 1)$ Tensor). *Let V be a $0-1$ tensor on a Riemannian manifold \mathcal{M} and $c : [a, b] \rightarrow \mathcal{M}$ be a geodesic curve connecting $c(a)$ and $c(b)$. Then it holds that*

$$T_{c(b) \rightarrow c(a)} V(c(b)) = V(c(a)) + \nabla V(c(a))(\dot{c}(t)) + \int_a^b (1-t) T_{c(b) \rightarrow c(a)}(\nabla^2 T(c(t))[\dot{c}(t), \dot{c}(t)]) dt.$$

Proof. The proof is a direct consequence of the Taylor expansion of $V \circ c$ at a and the property of parallel transport. \square

Lemma 7. *Let \mathcal{M} be a Riemannian manifold of dimension p , R be a second-order retraction which is defined on $U \subset \mathbb{R}^p$ centered at x , and x' be a point satisfying $x' \in R_x(U)$ with $\text{dist}(x, x') = \epsilon$. Then given the canonical basis $\{\delta_i\}$ of \mathbb{R}^p , it holds that*

$$\|\text{d}R_x(R_x^{-1}(x'))\delta_i - T_{x \rightarrow x'}(\text{d}R_x(0)\delta_i)\| \leq O(\epsilon^2), \quad \forall i \in [p].$$

Proof. Given a sufficiently small $\epsilon > 0$, we assume that $\gamma(t), t \in [0, 1]$ is a unique shortest constant-speed geodesic between x and x' and $\{\tilde{e}_i(t)\} \subset \mathbb{T}_{\gamma(t)} \mathcal{M}$ is the basis derived from the parallel transport of $\{\text{d}R(0)\delta_i\}$ along γ . We decompose $\text{d}R_a(\gamma(t))\delta_i$ as $\text{d}R_x(\gamma(t))\delta_i = \sum_{j=1}^p a_{i,j}(t)\tilde{e}_j$. Then we have

$$\|\text{d}R_x(R_x^{-1}(x'))\delta_i - T_{x \rightarrow x'}(\text{d}R_x(0)\delta_i)\|^2 \leq (|a_{i,i}(1) - 1|^2 + \sum_{j \neq i}^p |a_{i,j}(1)|^2).$$

On the other hand, for $v := \text{Log}_x(x') \in \text{T}_x\mathcal{M}$, the property of second-order retractions yields that

$$\nabla_v(\text{d}R_x[\delta_i])(0) = 0 = \sum_{j=1}^p (va_j)\tilde{e}_j(0)$$

which leads to the fact that $va_i = 0$ for $i = 1, \dots, p$.

Therefore, it follows by the Taylor expansion and the smoothness of R that

$$\begin{aligned} |a_1 - 1| &\leq \frac{c\epsilon^2}{2} \\ |a_i| &\leq \frac{c\epsilon^2}{2}, \quad \forall i = 2, \dots, p \end{aligned}$$

with a constant c in a local neighborhood of x which leads to the conclusion. \square

For completeness, we present the classical results on moderate deviation and moment control in the following.

Lemma 8 (Theorem 1 in [von Bahr \(1967\)](#)). *Suppose that we have n i.i.d. random vector $\{\mathbf{X}_i\}_{i \in [n]}$ of dimension k . If $\mathbb{E}[\|\mathbf{X}_1\|_2^r]$ holds for an integer $r \geq 3$, and m is the largest singular value of the covariance matrix of \mathbf{X}_1 , then*

$$\left| \mathbb{P}\left[\left\|\sum_{i \in [n]} \mathbf{X}_i / \sqrt{n}\right\|_2 \leq a\right] - \int_{\|\mathbf{x}\|_2 \leq a} \text{d}\Phi(\mathbf{x}) \right| \leq C \cdot a^{-r} \cdot n^{-\frac{r-2}{2}}$$

holds for some constant C and every $a \geq (\frac{5}{4}m(r-2)\log n)^{\frac{1}{2}}$, where Φ denotes the standard k -dimensional normal distribution.

Lemma 9 (Theorem 2.10 in [Hall and Heyde \(2014\)](#)). *If $\{S_i, \mathcal{F}_i, 1 \leq i \leq n\}$ is a martingale and $1 < p < \infty$, $X_1 = S_1$, and $X_i = S_i - S_{i-1}$, $2 \leq i \leq n$, then there exist constants C_1 and C_2 depending only on p such that*

$$C_1 \mathbb{E} \left| \sum_{i \in [n]} X_i^2 \right|^{\frac{p}{2}} \leq \mathbb{E} |S_n|^p \leq C_2 \mathbb{E} \left| \sum_{i \in [n]} X_i^2 \right|^{\frac{p}{2}}.$$

Lastly, we provide a proof of [Proposition 3](#).

Proof of Proposition 3 This proposition is a consequence of [Lemma 20](#) and [Theorem 22](#) in [Absil and Malick \(2012\)](#). To be specific, we consider a function $R : \text{T}\mathcal{M} \rightarrow \text{M}$ that maps $(x_0, u) \in \text{T}\mathcal{M}$ to $x_0 + u + v$, where v is a vector in $\text{N}_{x_0}\mathcal{M}$ of the shortest length ensuring $x_0 + u + v \in \mathcal{M}$. By [Lemma 20](#) in [Absil and Malick \(2012\)](#), the existence and uniqueness of v in a sufficiently small neighborhood of $(x_0, 0_{x_0}) \in \text{T}\mathcal{M}$ is guaranteed, and $\text{d}R(x_0, \cdot)|_{0_{x_0}} = \text{id}$. By choosing a sufficiently small neighborhood $U \in \text{T}_{x_0}\mathcal{M}$, $R(x_0, \cdot)$ is a diffeomorphism from U to $R(x_0, U)$. By the construction of L_{x_0} , it immediately follows that $L_{x_0}(\cdot) = (R(x_0, \cdot))^{-1}$ on $R(x_0, U)$. Notice that R is proved to satisfy $\mathcal{P}_{x_0} \frac{\text{d}^2}{\text{d}t^2} R(x_0, tv)|_{t=0} = 0$, or equivalently, $\frac{\text{d}^2}{\text{d}t^2} \text{Log}_{x_0}(R(x_0, tv))|_{t=0} = 0$, for every $v \in \text{T}_{x_0}\mathcal{M}$, by [Theorem 22](#) in [Absil and Malick \(2012\)](#). We thus conclude that L_{x_0} is a second-order inverse retraction as it satisfies $\frac{\text{d}^2}{\text{d}t^2} L_x(\text{Exp}_x(tv)) = 0$ for every $v \in \text{T}_x\mathcal{M}$. \square