

Universal Constraints for Conformal Line Defects

Barak Gabai,¹ Amit Sever,² and De-liang Zhong (钟德亮)³

¹*Laboratory for Theoretical Fundamental Physics, EPFL, Rte de la Sorge, CH-1015, Lausanne*

²*School of Physics and Astronomy, Tel Aviv University, Ramat Aviv 69978, Israel*

³*Theoretical Physics Group, The Blackett Laboratory, Imperial College London, Prince Consort Road London, SW7 2AZ, UK*

We present a novel framework for deriving integral constraints for correlators on conformal line defects. These constraints emerge from the non-linearly realized ambient-space conformal symmetry. To validate our approach, we examine several examples and compare them against existing data for the four-point function of the displacement operator. Additionally, we provide a few new predictions that extend the current understanding of these correlators.

I. INTRODUCTION

The conformal bootstrap is a powerful technique for carving out the space of conformal field theories (CFTs). It is based on the conformal and crossing symmetries of CFT correlation functions; see [1–7] and references therein. This approach extends straightforwardly to conformal defects [8, 9]. In this paper, we focus on one-dimensional line defects. We argue that the correlation functions on these extended operators are subject to a new infinite set of bootstrap constraints, which go beyond those that follow from the conformal and crossing symmetries on a straight line.

The origin of these constraints is simple. Conformal line operators do not exhibit a conformal anomaly, allowing them to be placed along arbitrary smooth paths without breaking conformal symmetry.¹ When expanded around a straight line, the conformal symmetry of the deformed line translates into a set of constraints for integrated correlators.

More concretely, a straight conformal line operator is invariant under the $SL(2, \mathbb{R}) \times SO(d-1)$ subgroup of the ambient conformal symmetry. Operators localized on the line are classified by this symmetry, which also constrains their correlators. This classification extends to smooth lines, as they are locally straight. Small deformations of the straight line can be parameterized in terms of correlators of operators living on the line. At linear order in the deformation, these considerations imply the existence of the so-called *displacement operator*, \mathbb{D} , a primary under $SL(2, \mathbb{R})$ with dimension two and transverse spin one. At higher orders in the deformation, the deformed line can be expressed in terms of integrated correlators of the displacement operators. Contributions from non-protected operators are excluded due to their anomalous dimensions, making their appearance incompatible with conformal symmetry. Applying a conformal transformation to a deformed line and expanding around the straight line yields an infinite tower of constraints on these integrated correlators.

It is interesting to explore how restrictive these constraints are in determining the expectation value of the line defect. In this paper, we take a first step in this direction by spelling some of these explicitly and checking them against known data at weak and strong coupling.

II. SETUP

We consider a smooth deformation of the straight line

$$x_{\text{straight}}(\tau) \mapsto x(\tau) = x_{\text{straight}}(\tau) + v(\tau), \quad (1)$$

where $x(\tau)$ is some parametrization of the line and $v(\tau)$ is the deformation vector, transverse to the line. At leading order in v , the change in the line operator \mathcal{W} is given by

$$\delta\mathcal{W} = \int d\tau |\dot{x}_\tau| v_\tau^i \mathcal{P}[\mathbb{D}_i(x_\tau)\mathcal{W}], \quad (2)$$

where $\dot{x}_\tau = \partial_\tau x(\tau)$ and $i = 1, \dots, d-1$ is a transverse index. Here, \mathbb{D}_i is the displacement operator, whose two-point function is fixed by the conformal symmetry up to an overall constant, Λ , as

$$\langle\langle \mathbb{D}_i(x)\mathbb{D}_j(0) \rangle\rangle = \frac{\Lambda \delta_{ij}}{|x|^4}. \quad (3)$$

Here, $\langle\langle \dots \rangle\rangle$ denotes correlation functions of operators inserted along the infinite straight line, normalized such that $\langle\langle \mathbb{1} \rangle\rangle = 1$.

At higher order in the deformation v , multiple displacement operators can be integrated along the line. The correlation of any odd number of these spinning operators is zero. Hence, the next non-zero correlator is the four-point function

$$\langle\langle \mathbb{D}_i(x_1)\mathbb{D}_j(x_2)\mathbb{D}_k(x_3)\mathbb{D}_l(x_4) \rangle\rangle = \frac{\widehat{\mathcal{F}}_{ijkl}(t)}{|x_{14}x_{23}|^4}, \quad (4)$$

where $x_{ij} = x_i - x_j$ and $x_1 < x_2 < x_3 < x_4$ are four ordered points on the line. Here,

$$t = \frac{x_{12}x_{34}}{x_{14}x_{23}} \geq 0, \quad (5)$$

is the $SL(2, \mathbb{R})$ invariant cross-ratio and $\widehat{\mathcal{F}}_{ijkl}$ can be written as a sum of tensor structures.

¹ In other words, their expectation value and correlation functions are conformal invariant functionals of their shape.

The correlator (4) is invariant under cyclic permutation of the four points, which translates into the crossing relation

$$\widehat{\mathcal{F}}_{ijkl}(t) = t^{-4} \widehat{\mathcal{F}}_{jkli}(1/t). \quad (6)$$

We can start equally with the two-point function of a $SL(2, \mathbb{R})$ primary operator \mathbb{O} on the straight line and then smoothly deform it. At the second order of the deformation v , we have the four-point function

$$\frac{\langle\langle \mathbb{O}_I(x_1) \mathbb{D}_i(x_2) \mathbb{D}_j(x_3) \mathbb{O}_J(x_4) \rangle\rangle}{\langle\langle \mathbb{D}_1(x_2) \mathbb{D}_1(x_3) \rangle\rangle \langle\langle \mathbb{O}_1(x_1) \mathbb{O}_1(x_4) \rangle\rangle} = \widehat{\mathcal{F}}_{IijJ}^{\text{ODDDO}}(t). \quad (7)$$

where $\Delta_{\mathbb{O}}$ is the $SL(2, \mathbb{R})$ conformal dimension of the line operator \mathbb{O} . If \mathbb{O} transforms under internal or space-time symmetries, then I and J represent the corresponding indices.

The constraints that we consider are most naturally written in terms of the *subtracted four-point function*, obtained from $\widehat{\mathcal{F}}$ by subtracting the generalized free field contribution $\mathcal{F} \equiv \widehat{\mathcal{F}} - \mathcal{F}_{\text{GFF}}$, where

$$\begin{aligned} (\mathcal{F}_{\text{GFF}})_{IijJ}^{\text{ODDDO}} &= \delta_{ij} \delta_{IJ}, \\ (\mathcal{F}_{\text{GFF}})_{ijkl} &= \delta_{il} \delta_{jk} + \delta_{ij} \delta_{kl} t^{-4} + \delta_{ik} \delta_{jl} (1+t)^{-4}. \end{aligned} \quad (8)$$

III. CONSTRAINTS FOR 4-POINT FUNCTIONS

A four-point function first appears at the fourth-order variation. Instead, we find it simpler to start with a two-point function of a $SL(2, \mathbb{R})$ primary operator \mathbb{O}_I on the line and apply a second-order variation to it. If $\mathbb{O} = \mathbb{D}$, the resulting four-point function corresponds to that of four displacement operators, as given in (4). Otherwise, it takes the form (7).

We start with the two-point function $\langle\langle \mathbb{O}_I \mathbb{O}_J \rangle\rangle$ and apply an arbitrary smooth variation to it, imposing that the result transforms covariantly under conformal transformations. This condition involves integrations of the form (2), see Appendix A for details. By applying a Mellin transform, one of the integrations can be performed straightforwardly. The remaining integral still depends on an arbitrary deformation vector. However, using the fact that an infinitesimal conformal deformation of the line corresponds to a quadratic polynomial, only a limited number of moments of this vector contribute. In this way, we arrive at two types of integrated constraints.

The first type is homogeneous and takes the form

$$\begin{aligned} \int_0^\infty dt \left[t^2 (\mathcal{F}_{JijI}^{\text{ODDDO}} + \mathcal{F}_{iIjJ}^{\text{DODO}}) + (1+t)^2 \mathcal{F}_{JjiI}^{\text{ODDDO}} \right] &= 0, \\ \int_0^\infty dt t \left[(1+t) (\mathcal{F}_{JijI}^{\text{ODDDO}} + \mathcal{F}_{JjiI}^{\text{ODDDO}}) - \mathcal{F}_{iIjJ}^{\text{DODO}} \right] &= 0, \\ \int_0^\infty dt \left[(1+t)^2 \mathcal{F}_{JijI}^{\text{ODDDO}} + t^2 \mathcal{F}_{JjiI}^{\text{ODDDO}} + \mathcal{F}_{iIjJ}^{\text{DODO}} \right] &= 0, \end{aligned} \quad (9)$$

where as before $\mathcal{F} = \widehat{\mathcal{F}} - \mathcal{F}_{\text{GFF}}$ and ²

$$\widehat{\mathcal{F}}_{iIjJ}^{\text{DODO}}(t) \equiv \frac{1}{(1+t)^4} \frac{\langle\langle \mathbb{D}_i(x_1) \mathbb{O}_I(x_2) \mathbb{D}_j(x_3) \mathbb{O}_J(x_4) \rangle\rangle}{\langle\langle \mathbb{D}_1(x_1) \mathbb{D}_1(x_3) \rangle\rangle \langle\langle \mathbb{O}_1(x_2) \mathbb{O}_1(x_4) \rangle\rangle}. \quad (10)$$

These types of constraints also apply in the case where \mathbb{O}_I and \mathbb{O}_J are replaced by any two defect operators that are not related by symmetry. The second type of constraint is inhomogeneous and takes the form ³

$$\begin{aligned} \Lambda \int_0^\infty dt \left[\left(t^2 [\mathcal{F}_{IjiJ}^{\text{ODDDO}}(t) + \mathcal{F}_{JijI}^{\text{ODDDO}}(t)] - (t+1)^2 [\mathcal{F}_{IijJ}^{\text{ODDDO}}(t) + \mathcal{F}_{JjiI}^{\text{ODDDO}}(t)] \right) \log \frac{t}{t+1} \right. \\ \left. + \left(t^2 \mathcal{F}_{jIiJ}^{\text{DODO}}(t) - \mathcal{F}_{iIjJ}^{\text{DODO}}(t) \right) \log t \right] = 2 (\delta_{ij} \delta_{IJ} \Delta_{\mathbb{O}} + [M_{ij}]_{IJ}), \end{aligned} \quad (11)$$

and

$$\begin{aligned} \Lambda \int_0^\infty dt t \left[(t+1) \left(\mathcal{F}_{IjiJ}^{\text{ODDDO}}(t) - \mathcal{F}_{IijJ}^{\text{ODDDO}}(t) + \mathcal{F}_{JijI}^{\text{ODDDO}}(t) - \mathcal{F}_{JjiI}^{\text{ODDDO}}(t) \right) \log \frac{t}{t+1} \right. \\ \left. + \left(\mathcal{F}_{iIjJ}^{\text{DODO}}(t) - \mathcal{F}_{jIiJ}^{\text{DODO}}(t) \right) \log t \right] = [M_{ij}]_{IJ}, \end{aligned} \quad (12)$$

² The prefactors in (10) are introduced so that when $\mathbb{O} = \mathbb{D}$ all the \mathcal{F} 's are equal and the constraints can be further simplified, see Appendix A 6.

³ When operators of noninteger dimensions lower than 4 appear in one of the OPE channels, fractional power divergences should be subtracted.

where M_{ij} is the rotation generator in the $i - j$ transverse plane, acting in the representation of the operator \mathbb{O} . For example, in the vector representation, $[M_{ij}]_{IJ} = \delta_{iJ}\delta_{jI} - \delta_{iI}\delta_{jJ}$. The source on the right-hand side of these constraints originates from the conformal transformation properties of this operator.

Once the defect CFT is specified, the integrated constraints can also be expressed as sum rules on the conformal data. These take the schematic form

$$\sum_j \left[A(\tilde{\Delta}_j) C_{\mathbb{D}\mathbb{D}\tilde{\mathbb{O}}_j} C_{\tilde{\mathbb{O}}_j\mathbb{O}\mathbb{O}} + B(\tilde{\Delta}_j) C_{\mathbb{D}\mathbb{O}\tilde{\mathbb{O}}_j}^2 \right] = \frac{F}{\Lambda}, \quad (13)$$

where the sum runs over all DCFT operators and A, B, F are universal functions, see Appendix B for more details.

IV. CHECKS

We have performed several checks of the constraints (9), (11), (12) against available data in the literature at weak, strong, and finite coupling. In all cases, we have found them to be satisfied. The examples we considered include

- The 1/2-BPS Wilson line in $\mathcal{N} = 4$ SYM theory in the 't Hooft large N limit. This defect preserves a part of the superconformal symmetry, which relates \mathbb{D} to other members of a short multiplet. We find that the constraints are consistent with these SUSY relations. At weak 't Hooft coupling, perturbative data is available up to two loops, [10, 11]. In the strong coupling limit, the perturbative data (in $1/g$) is available up to four orders [12–15]. Finally, the leading OPE behavior of the four-point function is known at finite coupling [14, 15].
- The 1/2-BPS Wilson line in ABJM theory in the 't Hooft large N limit [16]. At strong coupling, the displacement four-point function is known at the tree level. It is identical to the result for the 1/2-BPS Wilson line in $\mathcal{N} = 4$ SYM theory, up to the identification of the string tension with the coupling constant in ABJM theory. Consequently, the checks performed there apply directly to this case.
- The 1/2-BPS defect in the CFT dual to string theory on $AdS_3 \times S^3 \times T^4$ with mixed Ramond-Ramond (RR) and Neveu-Schwarz-Neveu-Schwarz (NS-NS) three-form fluxes [17]. A one-parameter family of such backgrounds, interpolating between the pure NS-NS and pure R-R cases, was studied at strong coupling in [17]. The two-point function and the connected four-point functions of the displacement operator have been computed at leading order. In Appendix E, we demonstrate that in this case the integrated constraints can be employed to fix an undetermined free parameter in the analytic conformal bootstrap.

- Localized magnetic field line defect in the $O(N)$ Wilson-Fisher fixed point. The two-point function and the connected four-point functions involving at least two displacement operators were calculated in [18, 19] at the leading order in the ε -expansion.

In the following subsection, we present the check for the localized magnetic field line defect at the $O(N)$ Wilson-Fisher fixed point. This relatively brief example highlights the general features of the weak coupling expansion and allows us to make new predictions.

A. Localized Magnetic Field Line Defect in the $O(N)$ Model

The $O(N)$ Wilson-Fisher CFT has a simple line defect that was studied in [18, 19] in the ε -expansion, as we now review. The model consists of N scalars $\vec{\phi} = (\phi_1, \dots, \phi_N)$ in $4 - \varepsilon$ dimensions, with action

$$S = \int d^d x \left[\frac{1}{2} (\partial \vec{\phi})^2 + \frac{\lambda}{4!} (\vec{\phi}^2)^2 \right], \quad (14)$$

where, at the fixed point, the coupling constant λ takes the value

$$\frac{\lambda_*}{(4\pi)^2} = \frac{3\varepsilon}{N+8} + \frac{9(3N+14)\varepsilon^2}{(N+8)^3} + \mathcal{O}(\varepsilon^3). \quad (15)$$

A localized magnetic field defect along a smooth line \mathcal{C} is introduced by deforming the action as

$$S \rightarrow S + h \int_{\mathcal{C}} \phi_1. \quad (16)$$

It breaks the $O(N)$ symmetry down to $O(N-1)$, and induces an RG flow on the defect. The magnetic field h reaches a fixed point at

$$h_*^2 = (N+8) + \varepsilon \frac{4N^2 + 45N + 170}{2N+16} + \mathcal{O}(\varepsilon^2). \quad (17)$$

The corresponding DCFT contains the following line operators

ϕ_1 : An $O(N-1)$ singlet of dimension $\Delta_{\phi_1} = 1 + \varepsilon + \mathcal{O}(\varepsilon^2)$.

ϕ_I : A $O(N-1)$ vector of dimension one. These are protected *tilt operators* associated with the broken $O(N)$ symmetry.

\mathbb{D} : The displacement operator (2). It takes the form $\mathbb{D}_j = h_* \partial_j \phi_1$, where j is a transverse vector index.

The relevant conformal data for our computation are

$$\begin{aligned} \Delta_{\phi_1} &= 1 + \varepsilon - \frac{3N^2 + 49N + 194}{2(N+8)^2} \varepsilon^2 + \mathcal{O}(\varepsilon^3), \\ C_{\mathbb{D}\mathbb{D}\phi_1} &= \frac{\pi}{\sqrt{N+8}} \varepsilon + C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)} \varepsilon^2 + \mathcal{O}(\varepsilon^3), \\ C_{\phi_1\phi_1\phi_1} &= \frac{3\pi}{\sqrt{N+8}} \varepsilon + C_{\phi_1\phi_1\phi_1}^{(2)} \varepsilon^2 + \mathcal{O}(\varepsilon^3), \\ \Lambda &= \frac{N+8}{2\pi^2} + \frac{5N^2 + 23N + 62}{12\pi^2(N+8)} \varepsilon + \mathcal{O}(\varepsilon^2), \end{aligned} \quad (18)$$

where $C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}$ and $C_{\phi_1\phi_1\phi_1}^{(2)}$ are unknown.

In the following, we will focus on the four-point function that involves two displacement operators and two tilt operators or two $O(N-1)$ singlets. These correlators take the form (7), (10), with $\mathbb{O}_I = \phi_I$ or ϕ_1 . The tree-level contribution $\mathcal{O}(\varepsilon^0)$ is a factorized product of two free propagators. In our normalization, this disconnected contribution to \mathcal{F} does not receive higher-order corrections.

The leading connected contributions $\mathcal{O}(\varepsilon)$ take the form

$$\mathcal{F}_{ij;(1)}^{\phi_1\mathbb{D}\mathbb{D}\phi_1}(t) = -\frac{\varepsilon\delta_{ij}\mathcal{I}_1(t)}{N+8}, \quad \mathcal{F}_{ij;(1)}^{\mathbb{D}\phi_1\mathbb{D}\phi_1}(t) = -\frac{\varepsilon\delta_{ij}\mathcal{I}_2(t)}{N+8}, \quad (19)$$

where the subscript (1) indicates the order in the ε -expansion. The two functions of t are given by [19]

$$\begin{aligned} \mathcal{I}_1(t) &= \frac{2t-1}{t^2} \log(1+t) + \frac{1}{t(t+1)} - \frac{2t+3}{(t+1)^2} \log t, \\ \mathcal{I}_2(t) &= \frac{\log(1+t)}{t^2} - \frac{1}{t(t+1)^2} - \frac{t+3}{(t+1)^3} \log t. \end{aligned} \quad (20)$$

The four-point functions with the tilt operators are related to these by an overall factor of $\delta_{IJ}/3$. The details of checking the constraints are very similar for that case. Hence, we only present the constraint check with the singlet (19).

As noted in [11], it is dangerous to substitute perturbative results, such as (20), directly into finite-coupling integrals. This is because, at some order in the ε -expansion, the corresponding integral may diverge. In such cases, the ε -expansion does not commute with the integration.

For example, consider an operator $\tilde{\mathbb{O}}$ of dimension $\tilde{\Delta}$ that appears in the $[\mathbb{D} \times \mathbb{D}] \rightarrow [\mathbb{O} \times \mathbb{O}]$ OPE channel of $\mathcal{F}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}$ in (7), (which could be either a primary or a descendant). Its contribution to the four-point function is of the form

$$\mathcal{F}_{ij}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}(t) = \delta_{ij} \mathcal{N} \times t^{-\tilde{\Delta}} (1 + \mathcal{O}(1/t)), \quad (21)$$

where $\mathcal{N} = C_{\mathbb{D}\mathbb{D}\tilde{\mathbb{O}}} C_{\tilde{\mathbb{O}}\mathbb{O}\mathbb{O}}$ is the product of two structure constants. This behavior results in a contribution to the constraints of the form

$$\int_u^\infty dt t^n \mathcal{F}_{11}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}(t) = \frac{\mathcal{N} u^{n+1-\tilde{\Delta}}}{\tilde{\Delta} - n - 1} + \dots \quad (22)$$

where $n < \tilde{\Delta} - 1$ is an integer and u is a large value for which the OPE expansion (21) is still valid. Suppose that $\tilde{\Delta} = (n+1) + \gamma_1 \varepsilon + \mathcal{O}(\varepsilon^2)$, and $\mathcal{N} = \mathcal{N}_1 \varepsilon + \mathcal{N}_2 \varepsilon^2 + \mathcal{O}(\varepsilon^3)$. In perturbation theory, we have

$$\frac{\mathcal{N} u^{n+1-\tilde{\Delta}}}{\tilde{\Delta} - n - 1} = \frac{\mathcal{N}_1}{\gamma_1} + \varepsilon \frac{\mathcal{N}_2}{\gamma_1} - \varepsilon \mathcal{N}_1 \log u + \mathcal{O}(\varepsilon^2). \quad (23)$$

If we naively expand (21) in ε before performing the integration, the $\mathcal{O}(\varepsilon^0)$ term would be missed, and at $\mathcal{O}(\varepsilon^1)$,

the integral (22) would diverge. This shows that the integral enhances certain contributions in the ε -expansion. Hence, when plugging a perturbative result into the constraints, we must add all the enhanced contributions and regulate the divergences by an analytic continuation in the dimension $\tilde{\Delta}$.

Note that, when $\tilde{\Delta} = m + \mathcal{O}(\varepsilon)$ with $m < n + 1$, the integral diverges also for small finite ε . Such cases are regulated by an analytical continuation in $\tilde{\Delta}$, starting from $\tilde{\Delta} > n + 1$.

a. The homogeneous constraints For the relevant case $i = j$ and $I = J$ in (9), the first and third lines are equivalent. The integrals over \mathcal{I}_1 have power divergences. They originate from the $[\phi_1 \times \phi_1] \rightarrow [\mathbb{D} \times \mathbb{D}]$ OPE limit, where

$$\mathcal{F}_{ij;(1)}^{\phi_1\mathbb{D}\mathbb{D}\phi_1}(t) = -\frac{3\varepsilon\delta_{ij}}{8+N} (t^{-2} - t^{-3}) + \mathcal{O}(t^{-4}). \quad (24)$$

These first two orders match the expansion of the $SL(2, \mathbb{R})$ conformal block of a dimension two operator, see (B2). There are two such line operators, labeled by their leading-order form as ϕ_1^2 and $\phi^I \phi_I$. Because $C_{\mathbb{D}\mathbb{D}\phi_1^2} C_{\phi_1^2\phi_1\phi_1} = \mathcal{O}(\varepsilon)$ and $C_{\mathbb{D}\mathbb{D}(\phi^I\phi_I)} C_{(\phi^I\phi_I)\phi_1\phi_1} = \mathcal{O}(\varepsilon^2)$, at order $\mathcal{O}(\varepsilon)$ only ϕ_1^2 is exchanged. Hence, before integrating \mathcal{I}_1 it is sufficient to replace the two leading terms in (24) with the finite coupling behavior

$$t^{-2} (1 - t^{-1}) \rightarrow t^{-\Delta_{\phi_1^2}} \left(1 - \frac{\Delta_{\phi_1^2}}{2} t^{-1}\right). \quad (25)$$

After this substitution the integrals in (9) converge for $\Delta_{\phi_1^2} > 3$, and we re-expand the result in ε . The corresponding enhanced contributions cancel out.

We find that the enhanced contributions for the first homogeneous constraint cancel among themselves. For the second homogeneous constraint, a single enhanced contribution remains. It originates from the exchange of the first two descendants of ϕ_1 in the $[\mathbb{D} \times \mathbb{D}] \rightarrow [\phi_1 \times \phi_1]$ OPE channel and is of the form (23) with $\mathcal{N}_1 = 0$. Collecting all the contributions, we find that the two independent homogeneous constraints in (9) are satisfied.

b. The inhomogeneous constraint For $I = J$, the second inhomogeneous equation in (12) becomes trivial, so we only have to check the first. The source on the right-hand side of this equation is of order $\mathcal{O}(\varepsilon^0)$, see (18). On the left-hand side, the connected four-point function is of order $\mathcal{O}(\varepsilon^1)$, which means that all contributions at order $\mathcal{O}(\varepsilon^0)$ are enhanced ones. They originate from the \mathbb{D} exchange in the $[\mathbb{D} \times \phi_1] \rightarrow [\mathbb{D} \times \phi_1]$ and $[\phi_1 \times \mathbb{D}] \rightarrow [\mathbb{D} \times \phi_1]$ OPE channels. Due to a version of (22) with logarithms in the kernels, they include enhancements from $\mathcal{O}(\varepsilon^2)$ to down to $\mathcal{O}(\varepsilon^0)$. These enhanced contributions sum up to exactly reproduce the right-hand side of the constraint equation.

At next order $\mathcal{O}(\varepsilon^1)$, the inhomogeneous constraint receives both regular and enhanced contributions. In this case the enhanced ones arise from the ϕ_1 -exchange in the $[\mathbb{D} \times \mathbb{D}] \rightarrow [\phi_1 \times \phi_1]$ OPE channel and the \mathbb{D} -exchange

in the $[\mathbb{D} \times \phi_1] \rightarrow [\mathbb{D} \times \phi_1]$, $[\phi_1 \times \mathbb{D}] \rightarrow [\mathbb{D} \times \phi_1]$ OPE channels. They combine to an answer that depends on the unknown structure constant $C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}$. As a result, the inhomogeneous constraint yields a new prediction

$$C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)} = -\frac{\pi}{12} \frac{29N^2 + 413N + 1610}{(N+8)^{5/2}}. \quad (26)$$

Using a similar analysis, predictions can be extracted for structure constants involving the tilt operator ϕ_I and displacements. For example, we find

$$C_{\mathbb{D}\phi_I\partial_\perp\phi_I}^{(2)} = \pi \frac{22 + N(3-N)}{12(N+8)^{5/2}} - \frac{\gamma_{\partial_\perp\phi_I}^{(2)}}{\sqrt{N+8}}. \quad (27)$$

Further constraints can be formulated using the $\mathcal{O}(\varepsilon^2)$ perturbative data alone. For example, by taking a linear combination of the $\mathcal{O}(\varepsilon^2)$ homogeneous constraints such that the enhanced contribution (proportional to $C_{\phi_1\phi_1\phi_1}^{(2)}$) drops out, one obtains an integrated constraint on the four-point function at this order that does not involve any other unknown conformal data. Details can be found in Appendix C.

V. DISCUSSION

The expectation value of a critical line defect along an arbitrary smooth path is a conformally invariant functional of its shape. When expanded around a straight path, this invariance imposes an infinite set of integrated constraints on correlation functions involving the displacement operator.

In this Letter, we have established and studied the first non-trivial constraints in this set; see (9), (11) and (12). A comprehensive review of available defect CFT data from the literature revealed that these constraints are universally satisfied in all studied examples. Furthermore, in certain cases, these constraints enabled us to make novel predictions.

The constraints we have considered are only the tip of the iceberg, with many future directions to pursue.

Below, we outline several of them: I) The constraints studied in this Letter can be naturally extended by considering higher-order variations. II) In scenarios where the line defect breaks internal symmetry or supersymmetry, protected “tilt” operators emerge [20–23]. These constraints can then be extended to integrated correlators involving both displacement and tilt operator insertions.⁴ III) A novel avenue for future exploration is understanding how restrictive these constraints are. For example, in CS-matter theories at large N , it has been shown that minimal CFT data, together with these constraints, uniquely determine the line defect CFT and its expectation values [25, 26]. IV) The same ideas may also be implemented for higher-dimensional defects. A new ingredient that appears when an even-dimensional defect is placed along a curved sub-manifold is a gravitational anomaly. Its presence is expected to yield an extra source term in the integrated constraints. V) It would be advantageous to combine these constraints with existing numerical and analytical bootstrap methods to improve our understanding and bounds on defect CFTs.⁵ VI) Finally, it would be interesting if similar constraints could also be derived for the correlation functions of the stress-energy tensor of the CFT itself, [52].

Note: We have been informed that similar ideas are pursued in [53].

ACKNOWLEDGMENTS

We thank Victor Gorbenko, Shota Komatsu, Ziwen Kong, Slava Rychkov, Philine van Vliet, Miguel Paulos, Michelangelo Preti, and Xi Yin for the discussions. We are grateful to Nikolay Gromov and Zohar Komargodski for insightful discussions and comments on the manuscript. AS is supported by the Israel Science Foundation (grant number 1197/20 and 1099/24). BG is supported by the Simons Collaboration Grant on Confinement and QCD Strings. DIZ is supported in part by the UK Engineering and Physical Sciences Research Council grant number EP/Z000106/1, and the Royal Society under the grant URF\R1\221310.

[1] S. Rychkov, *EPFL Lectures on Conformal Field Theory in D = 2, 3 Dimensions*, SpringerBriefs in Physics (2016) arXiv:1601.05000 [hep-th].

[2] D. Simmons-Duffin, The Conformal Bootstrap, in *Theoretical Advanced Study Institute in Elementary Particle Physics* (2017) pp. 1–74, arXiv:1602.07982 [hep-th].

[3] D. Poland, S. Rychkov, and A. Vichi, The Conformal Bootstrap: Theory, Numerical Techniques, and Applications, *Rev. Mod. Phys.* **91**, 015002 (2019), arXiv:1805.04405 [hep-th].

[4] S. M. Chester, Weizmann lectures on the numerical conformal bootstrap, *Phys. Rept.* **1045**, 1 (2023), arXiv:1907.05147 [hep-th].

[5] T. Hartman, D. Mazac, D. Simmons-Duffin, and A. Zhiboedov, Snowmass White Paper: The Ana-

⁴ See [24] for integrated constraints involving the tilt operators only.

⁵ Integrated constraints have been proven to be very successful in the CFT bootstrap. One prominent example is the set of constraints derived from supersymmetric localization, [27–47]. Another example involves integrated correlators on supersymmetric 1/2-BPS Wilson line in $N = 4$ super Yang-Mills theory, [11, 48–51].

- lytic Conformal Bootstrap, in *Snowmass 2021* (2022) [arXiv:2202.11012 \[hep-th\]](#).
- [6] A. Bissi, A. Sinha, and X. Zhou, Selected topics in analytic conformal bootstrap: A guided journey, *Phys. Rept.* **991**, 1 (2022), [arXiv:2202.08475 \[hep-th\]](#).
- [7] S. Rychkov and N. Su, New developments in the numerical conformal bootstrap, *Rev. Mod. Phys.* **96**, 045004 (2024), [arXiv:2311.15844 \[hep-th\]](#).
- [8] M. Billò, V. Gonçalves, E. Lauria, and M. Meineri, Defects in conformal field theory, *JHEP* **04**, 091, [arXiv:1601.02883 \[hep-th\]](#).
- [9] A. Gadde, Conformal constraints on defects, *JHEP* **01**, 038, [arXiv:1602.06354 \[hep-th\]](#).
- [10] N. Kiryu and S. Komatsu, Correlation Functions on the Half-BPS Wilson Loop: Perturbation and Hexagonalization, *JHEP* **02**, 090, [arXiv:1812.04593 \[hep-th\]](#).
- [11] A. Cavaglià, N. Gromov, J. Julius, and M. Preti, Bootstrability in defect CFT: integrated correlators and sharper bounds, *JHEP* **05**, 164, [arXiv:2203.09556 \[hep-th\]](#).
- [12] S. Giombi, R. Roiban, and A. A. Tseytlin, Half-BPS Wilson loop and $\text{AdS}_2/\text{CFT}_1$, *Nucl. Phys. B* **922**, 499 (2017), [arXiv:1706.00756 \[hep-th\]](#).
- [13] P. Liendo, C. Meneghelli, and V. Mitev, Bootstrapping the half-BPS line defect, *JHEP* **10**, 077, [arXiv:1806.01862 \[hep-th\]](#).
- [14] P. Ferrero and C. Meneghelli, Unmixing the Wilson line defect CFT. Part II. Analytic bootstrap, *JHEP* **06**, 010, [arXiv:2312.12551 \[hep-th\]](#).
- [15] P. Ferrero and C. Meneghelli, Unmixing the Wilson line defect CFT. Part I. Spectrum and kinematics, *JHEP* **05**, 090, [arXiv:2312.12550 \[hep-th\]](#).
- [16] L. Bianchi, G. Bliard, V. Forini, L. Griguolo, and D. Seminara, Analytic bootstrap and Witten diagrams for the ABJM Wilson line as defect CFT₁, *JHEP* **08**, 143, [arXiv:2004.07849 \[hep-th\]](#).
- [17] G. Bliard, D. H. Correa, M. Lagares, and I. Salazar Landea, Bootstrapping line defects in $\text{AdS}_3/\text{CFT}_2$, (2024), [arXiv:2410.02685 \[hep-th\]](#).
- [18] G. Cuomo, Z. Komargodski, and M. Mezei, Localized magnetic field in the $\text{O}(N)$ model, *JHEP* **02**, 134, [arXiv:2112.10634 \[hep-th\]](#).
- [19] A. Gimenez-Grau, E. Lauria, P. Liendo, and P. van Vliet, Bootstrapping line defects with $\text{O}(2)$ global symmetry, *JHEP* **11**, 018, [arXiv:2208.11715 \[hep-th\]](#).
- [20] A. J. Bray and M. A. Moore, Critical behaviour of semi-infinite systems, *J. Phys. A* **10**, 1927 (1977).
- [21] G. Cuomo, M. Mezei, and A. Raviv-Moshe, Boundary conformal field theory at large charge, *JHEP* **10**, 143, [arXiv:2108.06579 \[hep-th\]](#).
- [22] J. Padayasi, A. Krishnan, M. A. Metlitski, I. A. Gruzberg, and M. Meineri, The extraordinary boundary transition in the 3d $\text{O}(N)$ model via conformal bootstrap, *SciPost Phys.* **12**, 190 (2022), [arXiv:2111.03071 \[cond-mat.stat-mech\]](#).
- [23] C. P. Herzog and K.-W. Huang, Boundary Conformal Field Theory and a Boundary Central Charge, *JHEP* **10**, 189, [arXiv:1707.06224 \[hep-th\]](#).
- [24] N. Drukker, Z. Kong, and G. Sakkas, Broken Global Symmetries and Defect Conformal Manifolds, *Phys. Rev. Lett.* **129**, 201603 (2022), [arXiv:2203.17157 \[hep-th\]](#).
- [25] B. Gabai, A. Sever, and D.-l. Zhong, Line operators in Chern-Simons-Matter theories and Bosonization in Three Dimensions II: Perturbative analysis and all-loop resummation, *JHEP* **04**, 070, [arXiv:2212.02518 \[hep-th\]](#).
- [26] B. Gabai, A. Sever, and D.-l. Zhong, Bootstrapping smooth conformal defects in Chern-Simons-matter theories, *JHEP* **03**, 055, [arXiv:2312.17132 \[hep-th\]](#).
- [27] N. B. Agmon, S. M. Chester, and S. S. Pufu, Solving M-theory with the Conformal Bootstrap, *JHEP* **06**, 159, [arXiv:1711.07343 \[hep-th\]](#).
- [28] D. J. Binder, S. M. Chester, S. S. Pufu, and Y. Wang, $\mathcal{N} = 4$ Super-Yang-Mills correlators at strong coupling from string theory and localization, *JHEP* **12**, 119, [arXiv:1902.06263 \[hep-th\]](#).
- [29] N. B. Agmon, S. M. Chester, and S. S. Pufu, The M-theory Archipelago, *JHEP* **02**, 010, [arXiv:1907.13222 \[hep-th\]](#).
- [30] S. M. Chester, R. R. Kalloor, and A. Sharon, Squashing, Mass, and Holography for 3d Sphere Free Energy, *JHEP* **04**, 244, [arXiv:2102.05643 \[hep-th\]](#).
- [31] L. F. Alday, S. M. Chester, and H. Raj, ABJM at strong coupling from M-theory, localization, and Lorentzian inversion, *JHEP* **02**, 005, [arXiv:2107.10274 \[hep-th\]](#).
- [32] S. M. Chester, Bootstrapping 4d $\mathcal{N} = 2$ gauge theories: the case of SQCD, *JHEP* **01**, 107, [arXiv:2205.12978 \[hep-th\]](#).
- [33] S. M. Chester, R. Dempsey, and S. S. Pufu, Bootstrapping $\mathcal{N} = 4$ super-Yang-Mills on the conformal manifold, *JHEP* **01**, 038, [arXiv:2111.07989 \[hep-th\]](#).
- [34] S. M. Chester, R. Dempsey, and S. S. Pufu, Level repulsion in $\mathcal{N} = 4$ super-Yang-Mills via integrability, holography, and the bootstrap, *JHEP* **07**, 059, [arXiv:2312.12576 \[hep-th\]](#).
- [35] S. M. Chester, S. S. Pufu, Y. Wang, and X. Yin, Bootstrapping M-theory orbifolds, *JHEP* **06**, 001, [arXiv:2312.13112 \[hep-th\]](#).
- [36] S. Caron-Huot, F. Coronado, and Z. Zahraee, Bootstrapping $\mathcal{N} = 4$ sYM correlators using Integrability and Localization, (2024), [arXiv:2412.00249 \[hep-th\]](#).
- [37] S. Caron-Huot, F. Coronado, A.-K. Trinh, and Z. Zahraee, Bootstrapping $\mathcal{N} = 4$ sYM correlators using integrability, *JHEP* **02**, 083, [arXiv:2207.01615 \[hep-th\]](#).
- [38] C. Wen and S.-Q. Zhang, Integrated correlators in $\mathcal{N} = 4$ super Yang-Mills and periods, *JHEP* **05**, 126, [arXiv:2203.01890 \[hep-th\]](#).
- [39] D. Dorigoni, M. B. Green, and C. Wen, Exact results for duality-covariant integrated correlators in $\mathcal{N} = 4$ SYM with general classical gauge groups, *SciPost Phys.* **13**, 092 (2022), [arXiv:2202.05784 \[hep-th\]](#).
- [40] D. Dorigoni, M. B. Green, and C. Wen, Novel Representation of an Integrated Correlator in $\mathcal{N} = 4$ Supersymmetric Yang-Mills Theory, *Phys. Rev. Lett.* **126**, 161601 (2021), [arXiv:2102.08305 \[hep-th\]](#).
- [41] S. M. Chester and S. S. Pufu, Far beyond the planar limit in strongly-coupled $\mathcal{N} = 4$ SYM, *JHEP* **01**, 103, [arXiv:2003.08412 \[hep-th\]](#).
- [42] A. Brown, P. Heslop, C. Wen, and H. Xie, Integrated correlators in $\mathcal{N} = 4$ SYM beyond localisation, *Phys. Rev. Lett.* **132**, 101602 (2024), [arXiv:2308.07219 \[hep-th\]](#).
- [43] S. M. Chester, M. B. Green, S. S. Pufu, Y. Wang, and C. Wen, New modular invariants in $\mathcal{N} = 4$ Super-Yang-

- Mills theory, *JHEP* **04**, 212, [arXiv:2008.02713 \[hep-th\]](#).
- [44] L. F. Alday, S. M. Chester, D. Dorigoni, M. B. Green, and C. Wen, Relations between integrated correlators in $\mathcal{N} = 4$ supersymmetric Yang-Mills theory, *JHEP* **05**, 044, [arXiv:2310.12322 \[hep-th\]](#).
- [45] D. Dorigoni, M. B. Green, C. Wen, and H. Xie, Modular-invariant large- N completion of an integrated correlator in $\mathcal{N} = 4$ supersymmetric Yang-Mills theory, *JHEP* **04**, 114, [arXiv:2210.14038 \[hep-th\]](#).
- [46] H. Paul, E. Perlmutter, and H. Raj, Integrated correlators in $\mathcal{N} = 4$ SYM via $SL(2, \mathbb{Z})$ spectral theory, *JHEP* **01**, 149, [arXiv:2209.06639 \[hep-th\]](#).
- [47] D. Dorigoni, Note on 't Hooft-line defect integrated correlators in $N=4$ supersymmetric Yang-Mills theory, *Phys. Rev. D* **110**, L121702 (2024), [arXiv:2410.02377 \[hep-th\]](#).
- [48] A. Cavaglià, N. Gromov, J. Julius, and M. Preti, Integrability and conformal bootstrap: One dimensional defect conformal field theory, *Phys. Rev. D* **105**, L021902 (2022), [arXiv:2107.08510 \[hep-th\]](#).
- [49] A. Cavaglià, N. Gromov, J. Julius, and M. Preti, Integrated correlators from integrability: Maldacena-Wilson line in $\mathcal{N} = 4$ SYM, *JHEP* **04**, 026, [arXiv:2211.03203 \[hep-th\]](#).
- [50] A. Cavaglià, N. Gromov, and M. Preti, Computing Four-Point Functions with Integrability, Bootstrap and Parity Symmetry, (2023), [arXiv:2312.11604 \[hep-th\]](#).
- [51] A. Cavaglià, N. Gromov, J. Julius, M. Preti, and N. S. Sokolova, Probing Line Defect CFT with Mixed-Correlator Bootstrability, (2024), [arXiv:2412.07624 \[hep-th\]](#).
- [52] B. Gabai, A. Sever, and D.-l. Zhong, Work in progress, .
- [53] N. Drukker, P. Kravchuk, and Z. Kong, To appear, .
- [54] B. Gabai, A. Sever, and D.-l. Zhong, Line Operators in Chern-Simons-Matter Theories and Bosonization in Three Dimensions, *Phys. Rev. Lett.* **129**, 121604 (2022), [arXiv:2204.05262 \[hep-th\]](#).
- [55] B. Gabai, A. Sever, and D.-l. Zhong, Line operators in chern-simons-matter theories and bosonization in three dimensions, *Physical Review Letters* **129**, 10.1103/physrevlett.129.121604 (2022).
- [56] B. Gabai, A. Sever, and D.-l. Zhong, Bootstrapping smooth conformal defects in chern-simons-matter theories, *Journal of High Energy Physics* **2024**, 10.1007/jhep03(2024)055 (2024).
- [57] D. Pappadopulo, S. Rychkov, J. Espin, and R. Rattazzi, OPE Convergence in Conformal Field Theory, *Phys. Rev. D* **86**, 105043 (2012), [arXiv:1208.6449 \[hep-th\]](#).
- [58] N. B. Agmon and Y. Wang, Classifying Superconformal Defects in Diverse Dimensions Part I: Superconformal Lines, (2020), [arXiv:2009.06650 \[hep-th\]](#).
- [59] M. Gunaydin and R. J. Scalise, Unitary Lowest Weight Representations of the Noncompact Supergroup $Osp(2m^*/2n)$, *J. Math. Phys.* **32**, 599 (1991).
- [60] N. Dorey and A. Singleton, An Index for Superconformal Quantum Mechanics, *JHEP* [arXiv:1812.11816 \[hep-th\]](#).
- [61] L. Eberhardt, Summing over Geometries in String Theory, *JHEP* **05**, 233, [arXiv:2102.12355 \[hep-th\]](#).
- [62] F. A. Dolan and H. Osborn, Conformal four point functions and the operator product expansion, *Nucl. Phys. B* **599**, 459 (2001), [arXiv:hep-th/0011040](#).
- [63] F. A. Dolan and H. Osborn, Conformal partial waves and the operator product expansion, *Nucl. Phys. B* **678**, 491 (2004), [arXiv:hep-th/0309180](#).
- [64] R. Jackiw and S. Y. Pi, Conformal Blocks for the 4-Point Function in Conformal Quantum Mechanics, *Phys. Rev. D* **86**, 045017 (2012), [Erratum: *Phys. Rev. D* **86**, 089905 (2012)], [arXiv:1205.0443 \[hep-th\]](#).
- [65] K. Ghosh, A. Kaviraj, and M. F. Paulos, Polyakov blocks for the 1D conformal field theory mixed-correlator bootstrap, *Phys. Rev. D* **109**, L061703 (2024), [arXiv:2307.01257 \[hep-th\]](#).

SUPPLEMENTAL MATERIAL

Appendix A: Derivation of the Constraints

The key idea that leads to the integrated constraint is to compute the conformal transformation of a non-straight path in two different ways and then expand the results on the straight line. One way is to use the conformal covariance of smooth lines with insertions (or invariance where there are no insertions). The other way is by operator insertions on the line. This bootstrap method was first developed to study defects in Chern-Simons matter theory [25, 26, 54], but is generally applicable.

Here, we start from a two-point function on a straight defect. Without loss of generality, we place the two line operators at the origin and at infinity, $\langle\langle \mathbb{O}(0)\mathbb{O}(\infty) \rangle\rangle$. We then smoothly deform the path as in (1). For the order in which we work in this paper, it is sufficient to focus on the second order in the deformation parameter, v . The terms at this order can be classified according to the number of operators that have been integrated on the line

$$\delta^2 \langle\langle \mathbb{O}(0)\mathbb{O}(\infty) \rangle\rangle = [\text{int}^2] + [\text{int}] + [B_0^2] + [B_\infty^2] + [B_0 B_\infty]. \quad (\text{A1})$$

Here, $[\text{int}^2]$ is a double integral over two displacement operators, $[\text{int}]$ is a single integral, and $[B^2]$, $[B_\infty^2]$, $[B_0 B_\infty]$ are boundary terms, without integrations. They consist of all the operators that are allowed by the symmetries of the straight line and are listed in the following sections.

We then decompose the transverse deformation vector in (1) as $v = v_a + v_c$, where the transverse deformation vector v_c corresponds to a conformal deformation and v_a is an arbitrary smooth deformation that is not associated with a conformal one. These two are independent at the second order that we are working at.

By equating (A1) with the conformal transformation of the two operators at orders $\mathcal{O}(v_c^2)$ and $\mathcal{O}(v_a v_c)$ we can fix all coefficients in front of the operators appearing in (A1). We now describe each of the terms in detail and then derive the constraints from them.

1. Double Integration

The double integration term consists of two ordered insertions of displacement operators along the line. They come in three possible orderings

$$\begin{aligned} [\text{int}^2] &= \int_{\tau_1 < 0 < \tau_2} d\tau_1 d\tau_2 v^i(\tau_1) v^j(\tau_2) \langle\langle \mathbb{D}_i(\tau_1) \mathbb{O}_I(0) \mathbb{D}_j(\tau_2) \mathbb{O}_J(\infty) \rangle\rangle \\ &+ \int_{0 < \tau_1 < \tau_2} d\tau_1 d\tau_2 v^i(\tau_1) v^j(\tau_2) \langle\langle \mathbb{O}_I(0) \mathbb{D}_i(\tau_1) \mathbb{D}_j(\tau_2) \mathbb{O}_J(\infty) \rangle\rangle \\ &+ \int_{\tau_2 < \tau_1 < 0} d\tau_1 d\tau_2 v^i(\tau_1) v^j(\tau_2) \langle\langle \mathbb{D}_j(\tau_2) \mathbb{D}_i(\tau_1) \mathbb{O}_I(0) \mathbb{O}_J(\infty) \rangle\rangle, \end{aligned} \quad (\text{A2})$$

where we are using a "proper time" parametrization of the straight line in which $x(\tau) = \tau$, so $|\partial_\tau x(\tau)| = 1$ in (2). These integrals are generically divergent. We regularize them using point splitting in a conformal frame where the two \mathbb{O} operators are at $\tau = 0$ and $\tau = 1$. After using the conformal transformation $\tau \rightarrow \tau/(1 - \tau)$ to map this frame to the one where the operators are inserted at $\tau = 0$ and $\tau = \infty$, this regularization scheme reads

$$\epsilon < |\tau_j| < 1/\epsilon, \quad |\tau_2 - \tau_1| > \epsilon|(1 + \tau_1)(1 + \tau_2)|. \quad (\text{A3})$$

Using the normalization where $\langle\langle \mathbb{O}_I(0)\mathbb{O}_J(x) \rangle\rangle = \delta_{IJ}/|x|^{2\Delta}$, the correlators in $[\text{int}^2]$ are related to the ones in (7) and (10) as

$$\begin{aligned} \langle\langle \mathbb{D}_i(x_1) \mathbb{O}_I(x_2) \mathbb{D}_j(x_3) \mathbb{O}_J(\infty) \rangle\rangle &= \frac{\Lambda}{x_{23}^4} \times \widehat{\mathcal{F}}_{iIjJ}^{\text{DODO}}(t), \\ \langle\langle \mathbb{O}_I(x_1) \mathbb{D}_i(x_2) \mathbb{D}_j(x_3) \mathbb{O}_J(\infty) \rangle\rangle &= \frac{\Lambda}{x_{23}^4} \times \widehat{\mathcal{F}}_{IijJ}^{\text{ODOO}}(t), \\ \langle\langle \mathbb{D}_j(x_1) \mathbb{D}_i(x_2) \mathbb{O}_I(x_3) \mathbb{O}_J(\infty) \rangle\rangle &= \frac{\Lambda}{x_{12}^4} \times \widehat{\mathcal{F}}_{JjiI}^{\text{ODOO}}(1/t). \end{aligned} \quad (\text{A4})$$

These are defined so that for $\mathbb{O} = \mathbb{D}$ they are all equal. In terms of these $\widehat{\mathcal{F}}$'s, (A2) takes the form

$$\begin{aligned} [\text{int}^2]/\Lambda &= \int_0^\infty \frac{d\tau_1 d\tau_2}{\tau_2^4} \widehat{\mathcal{F}}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(\tau_1/\tau_2) v^i(-\tau_1) v^j(\tau_2) \\ &+ \int_{0 < \tau_1 < \tau_2} \frac{d\tau_1 d\tau_2}{(\tau_2 - \tau_1)^4} \left[\widehat{\mathcal{F}}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}\left(\frac{\tau_1}{\tau_2 - \tau_1}\right) v^i(\tau_1) v^j(\tau_2) + \widehat{\mathcal{F}}_{JjI}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}\left(\frac{\tau_1}{\tau_2 - \tau_1}\right) v^i(-\tau_1) v^j(-\tau_2) \right], \end{aligned} \quad (\text{A5})$$

where we leave the point-splitting regularization implicit. Here, we have made some changes of the integration variables that is inferred from the argument of the v 's.

Next, we rewrite the double integral (A5) in terms of the subtracted correlators $\mathcal{F} = \widehat{\mathcal{F}} - \mathcal{F}_{\text{GFF}}$, where \mathcal{F}_{GFF} is given in (8). We find that $[\text{int}^2]$ takes the same form in terms of the subtracted correlators. To see that, we plug \mathcal{F}_{GFF} into (A5) and take the deformation vector v to be a combination of the deformations that we will use in the following to derive the constraints (see (A25) and (A31)). We then perform the integrations using the regularization scheme (A3). After expanding the result in ϵ , we find that there are no finite contributions remaining.

The integrals in (A5) have a convolution structure. Hence, it is natural to express them in Mellin space,

$$\tilde{f}(s) = \int_0^\infty d\tau \tau^{s-1} f(\tau), \quad f(\tau) = \int_{c-i\infty}^{c+i\infty} \frac{ds}{2\pi i} \tau^{-s} \tilde{f}(s), \quad (\text{A6})$$

where the real part of the Mellin contour c is determined by the small and large τ asymptotics of the function $f(\tau)$.

For pure Mellin modes $v^i(\tau_1) = \tau_1^{-s_1}$ and $v^j(\tau_2) = \tau_2^{-s_2}$, the first line in $[\text{int}^2]$, (A5) can be written in terms of the Mellin transformed correlator as

$$\begin{aligned} &\int_\epsilon^{1/\epsilon} \frac{d\tau_1 d\tau_2}{\tau_2^4} \tau_1^{-s_1} \tau_2^{-s_2} \mathcal{F}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(\tau_1/\tau_2) = \int_\epsilon^{1/\epsilon} d\tau_2 \int_{\epsilon/\tau_2}^{1/(\tau_2\epsilon)} dt \tau_2^{-s_2-s_1-3} t^{-s_1} \mathcal{F}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(t) \\ &= \int_\epsilon^{1/\epsilon} d\tau_2 \int_0^\infty dt \tau_2^{-s_2-s_1-3} t^{-s_1} \mathcal{F}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(t) - \int_\epsilon^{1/\epsilon} d\tau_2 \left[\int_0^{\epsilon/\tau_2} + \int_{1/(\epsilon\tau_2)}^\infty \right] dt \tau_2^{-s_2-s_1-3} t^{-s_1} \mathcal{F}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(t) \\ &= 2\pi i \delta_\epsilon(s_1 + s_2 + 2) \widetilde{\mathcal{F}}_{iIjJ}^{\mathbb{D}\mathbb{O}\mathbb{D}\mathbb{O}}(1 - s_1) + [\text{boundary-contributions}]. \end{aligned} \quad (\text{A7})$$

where $[\text{boundary-contributions}]$ stands for terms in which at least one of the integrations is localized near the boundary. Such terms are absorbed in the boundary and single integration terms in (A1) that are analyzed in the next subsections. Here, δ_ϵ is the regularized delta function

$$\delta_\epsilon(x) \equiv \int_\epsilon^{1/\epsilon} \frac{d\tau}{\tau} \tau^x. \quad (\text{A8})$$

Similarly, the integrals on the second line of (A5) can be evaluated using

$$\begin{aligned} &\int_{0 < \tau_1 < \tau_2} \frac{d\tau_1 d\tau_2}{(\tau_2 - \tau_1)^4} \tau_1^{-s_1} \tau_2^{-s_2} \mathcal{F}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}\left(\frac{\tau_1}{\tau_2 - \tau_1}\right) = \int_{2\epsilon}^{1/\epsilon} d\tau_2 \tau_2^{-s_1-s_2-3} \int_{\epsilon/\tau_2}^{1-\frac{\epsilon}{\tau_2(1+\tau_2)^2}} dx \frac{x^{-s_1}}{(x-1)^4} \mathcal{F}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}\left(\frac{x}{1-x}\right) \\ &= 2\pi i \delta_\epsilon(s_1 + s_2 + 2) \widetilde{\mathcal{H}}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}(1 - s_1) + [\text{boundary-contributions}], \end{aligned} \quad (\text{A9})$$

where

$$\widetilde{\mathcal{H}}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}(s) \equiv \int_0^\infty dt (1+t)^2 \left(\frac{t}{t+1}\right)^{s-1} \mathcal{F}_{iIjJ}^{\mathbb{O}\mathbb{D}\mathbb{D}\mathbb{O}}(t). \quad (\text{A10})$$

By introducing the inverse Mellin integrals

$$\tilde{v}_\pm(s) \equiv \int_0^\infty d\tau \tau^{s-1} v(\pm\tau), \quad v(\pm|\tau|) = \int_{c-i\infty}^{c+i\infty} \frac{ds}{2\pi i} \tau^{-s} \tilde{v}_\pm(s), \quad (\text{A11})$$

we arrive at

$$[\text{int}^2] = \Lambda \int \frac{ds_1 ds_2}{2\pi i} \delta_\epsilon(2 + s_1 + s_2) \quad (\text{A12})$$

$$\times \left[\tilde{\mathcal{F}}_{iIjJ}^{\text{DDDD}}(1 - s_1) \tilde{v}_-^i(s_1) \tilde{v}_+^j(s_2) + \tilde{\mathcal{H}}_{iIjJ}^{\text{ODDD}}(1 - s_1) \tilde{v}_+^i(s_1) \tilde{v}_+^j(s_2) + \tilde{\mathcal{H}}_{JjiI}^{\text{ODDD}}(1 - s_1) \tilde{v}_-^i(s_1) \tilde{v}_-^j(s_2) \right].$$

2. Single Integration

For a single integral, the two deformations can be either integrated together or one is integrated and the other is at the location of one of the \mathbb{O} operators. We would like to enumerate all such terms that are consistent with the $SL(2, \mathbb{R} \times SO(d-1))$ symmetry of the straight line. This symmetry imposes two conditions. First, the boundary terms must retain the same $SL(2, \mathbb{R})$ dimension as \mathbb{O} prior to the deformation; second, the integrand on the defect must have dimension one, ensuring that the integral remains dimensionless. We assume that the CFT is “generic”. That is, we assume that

1. The only scalar operator that is a singlet under all internal symmetries and has integer dimension $\Delta < 4$ is the identity.
2. There is no operator of dimension $\Delta_{\mathbb{O}} + 1$ that otherwise has the same charges as \mathbb{O} and a spin larger by one unit.

Under this assumption, we find that for a generic spacetime dimension, the only terms with a single bulk integration and a non-zero expectation value are total derivatives. Such terms can be absorbed in the boundary terms in (A1). For example, we have

$$\int_a^b d\tau v(\tau) v'''(\tau) \times \mathbf{1} = \left[v(\tau) v''(\tau) - \frac{1}{2} (v'(\tau))^2 \right] \Big|_a^b. \quad (\text{A13})$$

In two, three, and four spacetime dimensions an additional operator on the line is allowed. Specifically, in four spacetime dimensions the following single integral is consistent with the straight-line symmetries

$$\int d\tau \epsilon^{ijk} v^i(\tau) v'^j(\tau) \langle\langle \mathbb{O}_I(0) \mathbb{D}^k(\tau) \mathbb{O}_J(\infty) \rangle\rangle \in [\text{int}]_{4d}, \quad (\text{A14})$$

where the i -index runs over the three transverse directions. We have found, however, that this term is ruled out by our bootstrap (although we do not include the detailed analysis in this note). Hence, we will not include it in the following.

Similarly, in three spacetime dimensions the following term is allowed

$$\int d\tau \epsilon^{ij} v^i(\tau) v'^j(\tau) \langle\langle \mathbb{O}_I(0) \mathbb{O}_J(\infty) \rangle\rangle \in [\text{int}]_{3d}. \quad (\text{A15})$$

While this term is not a total derivative, its addition has the effect of shifting the spin source on the right-hand side of the inhomogeneous constraints (11), (12). In our earlier work on Chern-Simons matter theories [55, 56], this term was excluded at third order, $O(v^3)$, of the bootstrap. This fact implies that also for any other three-dimensional theory, the coefficient of this term is fixed at third order. We expected, but did not prove, that it would be fixed to zero. In what follows we assume that this is indeed the case and therefore the constraints are not altered as we go down to three spacetime dimensions.

In two-dimensional CFTs in which there is no reflection symmetry in the direction perpendicular to the line the integral

$$\int d\tau v(\tau) dv(\tau) \langle\langle \mathbb{O}_I(0) \mathbb{D}(\tau) \mathbb{O}_J(\infty) \rangle\rangle \in [\text{int}]_{2d}, \quad (\text{A16})$$

is allowed. Similarly, there are also new boundary terms that can be added. In this case, we have repeated the derivation of the constraints for the four-point function of the displacement operator and found that they are not modified. This fact will be relevant for us when checking the constraints in the two-dimensional example studied in appendix E.

3. Boundary Terms

We now list all the boundary terms that are allowed by the $SL(2, \mathbb{R}) \times SO(d-1)$ symmetry of the straight line. In total, such terms have zero $SO(d-1)$ transverse spin and zero $SL(2, \mathbb{R})$ conformal dimension. Since the two v 's together have dimension minus two and spin zero, these terms consist of the operator \mathbb{O} , its first and second descendants.

We take the straight line to be placed along the third direction, x_3 . For simplicity, we restrict our discussion to the case where v is in a two-dimensional transverse space that is spanned by $x_{\pm} = (x_1 \pm ix_2)/\sqrt{2}$. This restricted kinematical configuration is sufficient for the derivation of the constraints. The boundary terms that are allowed in any CFT are

$$\begin{aligned} [B_0 B_0] &= \langle\langle [\gamma_0 v_0^+ v_0^- d^2 \mathbb{O}_I(0) + \gamma_1 (v_0^+ dv_0^- + v_0^- dv_0^+) d\mathbb{O}_I(0) + \gamma_4 (v_0^+ dv_0^- - v_0^- dv_0^+) d\mathbb{O}_I(0) \\ &\quad + (\gamma_3 dv_0^+ dv_0^- + \gamma_2 (v_0^+ d dv_0^- + v_0^- d dv_0^+) + \gamma_5 (v_0^+ d dv_0^- - v_0^- d dv_0^+)) \mathbb{O}_I(0)] \mathbb{O}_J(\infty) \rangle\rangle, \quad (\text{A17}) \\ [B_{\infty} B_{\infty}] &= \langle\langle \mathbb{O}_I(0) [\tilde{\gamma}_0 v_1^+ v_1^- d^2 \mathbb{O}_J(\infty) + \tilde{\gamma}_1 (v_{\infty}^+ dv_{\infty}^- + v_{\infty}^- dv_{\infty}^+) d\mathbb{O}_J(\infty) - \tilde{\gamma}_4 (v_{\infty}^+ dv_{\infty}^- - v_{\infty}^- dv_{\infty}^+) d\mathbb{O}_J(\infty) \\ &\quad + (\tilde{\gamma}_3 dv_{\infty}^+ dv_{\infty}^- + \tilde{\gamma}_2 (v_{\infty}^+ d dv_{\infty}^- + v_{\infty}^- d dv_{\infty}^+) - \tilde{\gamma}_5 (v_{\infty}^+ d dv_{\infty}^- - v_{\infty}^- d dv_{\infty}^+)) \mathbb{O}_J(\infty)] \rangle\rangle, \end{aligned}$$

where $\gamma_i, \tilde{\gamma}_i$ are unknown coefficients. Here, dv and $d\mathbb{O}$ stand for a derivative of v and \mathbb{O} along the straight line. As for descendant operators, the form of these derivatives depends on the conformal frame in which they are defined, which is tailored to the scheme in which we regulate the line integrals. Recall that we use the point-splitting regularization scheme in the conformal frame where the two operators are placed at $\tau = 0$ and $\tau = 1$. Hence, in this frame $dv = \partial_{\tau} v$ and $d\mathbb{O} = \partial_{\tau} \mathbb{O}$, where $x^3(\tau) = \tau$ with $\tau \in [0, 1]$. Using the conformal transformation $w \rightarrow w/(1-w)$ employed in (A3), this definition translates to

$$d^n A(\tau) \equiv \lim_{\sigma \rightarrow \tau} ((1+\sigma)^2 \partial_{\sigma})^n \frac{A(\sigma)}{(1+\sigma)^2}, \quad (\text{A18})$$

the $[0, \infty]$ frame.

Additionally, there are terms that are allowed when the defect contains operators of special dimension and spin. For example

- If the theory has an operator $\tilde{\mathbb{O}}$ of dimensions $\Delta_{\tilde{\mathbb{O}}} = \Delta_{\mathbb{O}} - m$ where m is a positive integer, then there may be additional allowed boundary terms. For example, if $\mathbb{O} = \mathbb{D}$ and $\tilde{\mathbb{O}} = \mathbf{1}$ then the term

$$dddv_0^i dddv_{\infty}^j \in \delta^2 \langle\langle \mathbb{D}^i(0) \mathbb{D}^j(\infty) \rangle\rangle, \quad (\text{A19})$$

is allowed and is part of $[B_0 B_{\infty}]$ in the decomposition (A1). This term, however, does not contribute when at least one of the v 's corresponds to a conformal transformation, which is the case relevant to our discussion.

- If \mathbb{O} has transverse spin 1/2 then terms such as

$$dv_0^{\pm} dv_{\infty}^{\mp} \langle\langle \mathbb{O}_{\pm \frac{1}{2}} \mathbb{O}_{\mp \frac{1}{2}} \rangle\rangle \in \delta^2 \langle\langle \mathbb{O}_{\mp \frac{1}{2}}(0) \mathbb{O}_{\pm \frac{1}{2}} \rangle\rangle, \quad (\text{A20})$$

are allowed and are part of $[B_0 B_{\infty}]$ in the decomposition (A1).

- If \mathbb{O} of spin one, then terms such as

$$v_0^+ dv_0^+ \langle\langle d\mathbb{O}_+(0) \mathbb{O}_-(\infty) \rangle\rangle \in \delta^2 \langle\langle \mathbb{O}_-(0) \mathbb{O}_-(\infty) \rangle\rangle, \quad (\text{A21})$$

are allowed. In this case, the two v 's together have transverse spin 2.

In these cases, we have verified that the presence of the new terms does not affect the constraints. Since these terms are nongeneric, in the following we will assume that they are not present.

4. Source Term

We now choose the deformation vector in (1) that will lead to the constraints eluded in the main text. We start with a non-straight path parameterized by $v(\tau) = v_a(\tau)$. At the second order, applying to it a conformal transformation

amounts to simply adding to it the infinitesimal conformal variation vector, $v_a(\tau) \rightarrow v_a(\tau) + v_c(\tau)$.⁶ Consequently, the change in the expectation value of the deformed line receives contributions at orders $\mathcal{O}(v_c^2)$ and $\mathcal{O}(v_c v_a)$. The source term of $\delta^2 \langle \mathbb{O}(0) \mathbb{O}(\infty) \rangle$ arises from the conformal covariance of the two operators. It is composed of two contributions. The first accounts for the scaling dimension of the operators. At the second order under consideration, this term takes the form,

$$[\text{conformal factor}] = -\Delta \delta_{IJ} (\vec{v}_\infty - \vec{v}_0)^2. \quad (\text{A22})$$

The second contribution accounts for the spin of the operators and takes the form

$$[\text{spin source}]_{\mathcal{O}(v_c^2)} = \frac{1}{2} \left[[M_{ij}]_{IJ} v_c^i \text{d}v_c^j / 2 \right] \Big|_0^\infty. \quad (\text{A23})$$

$$[\text{spin source}]_{\mathcal{O}(v_c v_a)} = \left[[M_{ij}]_{IJ} (v_a^i \text{d}v_c^j + \text{d}v_a^i \text{d}v_c^j / 2) \right] \Big|_0^\infty. \quad (\text{A24})$$

Finally, an infinitesimal conformal transformation is a degree two polynomial with arbitrary vector coefficients that we denote by \mathbf{c} ,

$$v_c^i(\tau) = \mathbf{c}_0^i + \mathbf{c}_1^i \tau + \mathbf{c}_2^i \tau^2. \quad (\text{A25})$$

The Mellin transform (A11) of $v_c(\tau)$ is

$$\tilde{v}_{c,\pm}^i(s) = 2\pi i \left[\mathbf{c}_0^i \delta(s) \pm \mathbf{c}_1^i \delta(1+s) + \mathbf{c}_2^i \delta(2+s) \right]. \quad (\text{A26})$$

5. Constraints

The constraints arise from equating $\delta^2 \langle \mathbb{O}(0) \mathbb{O}(\infty) \rangle$ in (A1) with the source term at orders $\mathcal{O}(v_c^2)$ and $\mathcal{O}(v_a v_c)$. We consider each of these orders in turn and separate between the cases where $\tilde{v}_a(s)$ is regular or has a pole at the support of the delta function in $[\text{int}^2]$, (A12).

a. $\mathcal{O}(v_a v_c)$ Constraints from Regular v_a

We first consider profiles $v_a(\tau)$ that vanish sufficiently fast at the boundaries, such that the boundary terms $[B_0 B_0]$ and $[B_\infty B_\infty]$ in (A17) vanish. For such a choice, the source terms in (A22)-(A24) also vanish. Moreover, for $\mathbb{O} = \mathbb{D}$, the term in (A19) vanishes because v_c is a degree 2 polynomial. Hence, the only contribution with which we remain is the double integral $[\text{int}^2]$ and the corresponding constraint is homogeneous.

With this choice, the Mellin transformed $\tilde{v}_a^i(s)$ is regular at the support of the delta function in (A12), $s = 0, -1, -2$, so we can safely take the $\epsilon \rightarrow 0$ limit there. Plugging in \tilde{v}_c from (A26) leads to the expression

$$\begin{aligned} [\text{int}^2] \Big|_{\mathcal{O}(v_a v_c)} &= \sum_{\ell=0}^2 \mathbf{c}_\ell^i \left(\tilde{v}_{a,-}^j(\ell-2) \left[\tilde{\mathcal{F}}_{jIiJ}^{\text{DODO}}(3-\ell) + (-1)^\ell \tilde{\mathcal{H}}_{JijI}^{\text{ODDO}}(3-\ell) + (-1)^\ell \tilde{H}_{JjiI}^{\text{ODDO}}(\ell+1) \right] \right. \\ &\quad \left. + \tilde{v}_{a,+}^j(\ell-2) \left[(-1)^\ell \tilde{\mathcal{F}}_{iIjJ}^{\text{DODO}}(\ell+1) + \tilde{\mathcal{H}}_{IjiJ}^{\text{ODDO}}(3-\ell) + \tilde{H}_{IijJ}^{\text{ODDO}}(\ell+1) \right] \right). \end{aligned} \quad (\text{A27})$$

Since $\tilde{v}_{a,\pm}(\ell-2)$ are free parameters, the equation imposes the following homogeneous constraints

$$\begin{aligned} (-1)^\ell \tilde{\mathcal{F}}_{jIiJ}^{\text{DODO}}(3-\ell) + \tilde{\mathcal{H}}_{JijI}^{\text{ODDO}}(3-\ell) + \tilde{H}_{JjiI}^{\text{ODDO}}(1+\ell) &= 0 \\ (-1)^\ell \tilde{\mathcal{F}}_{iIjJ}^{\text{DODO}}(1+\ell) + \tilde{\mathcal{H}}_{IjiJ}^{\text{ODDO}}(3-\ell) + \tilde{H}_{IijJ}^{\text{ODDO}}(1+\ell) &= 0 \end{aligned}, \quad \ell = 0, 1, 2. \quad (\text{A28})$$

These constraints apply for any choice of transverse directions i and j . Only three out of these six constraints are independent. Specifically, the first and the second lines are interchanged under $i \leftrightarrow j$, $I \leftrightarrow J$, and $\ell \leftrightarrow 2 - \ell$. After plugging in the definitions of the Mellin transforms $\tilde{\mathcal{F}}$ (A11) and $\tilde{\mathcal{H}}$ (A10), we arrive at the homogeneous constraints (9) in the main text.

⁶ At higher orders v_c depends on v_a .

b. $O(v_c^2)$ Constraints

Consider first the double integral at order $O(v_c^2)$. By plugging \tilde{v}_c in (A26) into (A12) we arrive at

$$[\text{int}^2] \Big|_{O(v_c^2)} = 2\pi i \Lambda \sum_{\ell_1, \ell_2=0}^2 \mathbf{c}_{\ell_1}^i \mathbf{c}_{\ell_2}^j \delta_\epsilon(2-\ell_1-\ell_2) \left[(-1)^{\ell_1} \tilde{\mathcal{F}}_{iIjJ}^{\text{DODO}}(1+\ell_1) + \tilde{\mathcal{H}}_{iIjJ}^{\text{ODDO}}(1-\ell_1) + (-1)^{\ell_1+\ell_2} \tilde{\mathcal{H}}_{JjiI}^{\text{ODDO}}(1-\ell_1) \right]. \quad (\text{A29})$$

The combination in the parentheses vanishes separately for each value of ℓ_1 and ℓ_2 by (A28).

We remained with the boundary and source terms. By comparing them for \mathbf{c}_i 's in a two-dimensional transverse plane we arrive at the following equations for the boundary coefficients

$$\begin{aligned} \tilde{\gamma}_0 = -\gamma_0 &= \frac{1-2\gamma_1}{1+2\Delta}, & \gamma_1 = 1 - \tilde{\gamma}_1 &= \frac{\gamma_2 + \gamma_3}{\Delta}, & \tilde{\gamma}_2 + \gamma_2 &= \Delta, \\ \tilde{\gamma}_3 = \tilde{\gamma}_3, & \gamma_4 = \tilde{\gamma}_4, & \Delta\gamma_4 - \gamma_5 &= \mathfrak{s}_0/2, & \Delta\tilde{\gamma}_4 + \tilde{\gamma}_5 &= \mathfrak{s}_\infty/2, \end{aligned} \quad (\text{A30})$$

where \mathfrak{s}_0 and \mathfrak{s}_∞ are the clockwise $U(1)$ spins of the two operators in the x_1 - x_2 transverse plane that we use here. This implies that γ_5 and $\tilde{\gamma}_5$ depend on the polarizations of the two \mathbb{O} operators.

c. $O(v_a v_c)$ Constraints from More General v_a

We consider v_a profiles of the form

$$v_a^i \in \left\{ \frac{\tau}{\tau^2+1}, \frac{\tau^2}{\tau^2+1}, \frac{\tau^3}{\tau^2+1} \right\}. \quad (\text{A31})$$

Their Mellin transform, $\tilde{v}_a^i(s)$, has a simple pole at the support of the delta function in (A12). That is, at $s = 0$, $s = -1$ or $s = -2$. However, these poles multiply expressions that vanish by the homophonous constraint (A28). As a result, the corresponding integrand in $[\text{int}^2]$ is regular at $\epsilon \rightarrow 0$. Yet, to obtain the finite terms in $[\text{int}^2]$ correctly, it is crucial to introduce the ϵ regulator in (A12) and take it to zero at the very end.

The source term at order $O(v_a v_c)$ that corresponds to the profiles (A31) is nontrivial. When combined with the results of the previous subsection (A30), these lead to the constraints

$$\Lambda \left[\tilde{\mathcal{F}}_{jIiJ}^{\text{DODO}}(3) - \tilde{\mathcal{F}}_{iIjJ}^{\text{DODO}}(1) + \tilde{\mathcal{H}}_{IjiJ}^{\text{ODDO}}(3) - \tilde{H}'_{iIjJ}^{\text{ODDO}}(1) + \tilde{\mathcal{H}}_{JjiI}^{\text{ODDO}}(3) - \tilde{H}'_{JjiI}^{\text{ODDO}}(1) \right] = 2\delta_{ij}\delta_{IJ}\Delta + 2[M_{ij}]_{IJ}, \quad (\text{A32})$$

and

$$\Lambda \left[-\tilde{\mathcal{F}}_{jIiJ}^{\text{DODO}}(2) + \tilde{\mathcal{F}}_{iIjJ}^{\text{DODO}}(2) + \tilde{\mathcal{H}}_{IjiJ}^{\text{ODDO}}(2) - \tilde{H}'_{iIjJ}^{\text{ODDO}}(-2) + \tilde{\mathcal{H}}_{JjiI}^{\text{ODDO}}(2) - \tilde{H}'_{JjiI}^{\text{ODDO}}(2) \right] = [M_{ij}]_{IJ}. \quad (\text{A33})$$

After plugging in the definitions of the Mellin transforms $\tilde{\mathcal{F}}$ (A11) and $\tilde{\mathcal{H}}$ (A10), we arrive at the non-homogeneous constraint (11) and (12) in the main text.

6. Simplified Constraints for $\mathbb{O} = \mathbb{D}$

In the case where the operator \mathbb{O} in (7) is the displacement operator and $I = J = i = j$, we have a single four-point function to consider and we can use crossing symmetry to map all the integration regions to $t \in [0, 1]$. The homogeneous and inhomogeneous constraints each reduce to a single independent nontrivial constraint. These take the form

$$\int_0^1 dt \mathcal{K}_{\text{Homo}}(t) \mathcal{F}_{1111}^{\text{DDDD}}(t) = 0, \quad \Lambda \int_0^1 dt \mathcal{K}_{\text{Inhomo}}(t) \mathcal{F}_{1111}^{\text{DDDD}}(t) = 4, \quad (\text{A34})$$

where

$$\mathcal{K}_{\text{Homo}}(t) = 1 + t + t^2, \quad \mathcal{K}_{\text{Inhomo}}(t) = (t^2 - 2t - 2) \log t + (t^2 + 4t + 1) \log(1 + t). \quad (\text{A35})$$

Appendix B: OPE Sum Rules

The integrated constraints can be rephrased as OPE sum rules. In this appendix, we demonstrate how to convert between the two for the four-point function $\mathcal{F}_{1111}^{\mathbb{D}\mathbb{D}\mathbb{D}\mathbb{D}}(t)$. The homogeneous and inhomogeneous integral constraints (A34) are of the form

$$\int_0^1 dt \mathcal{K}(t) \mathcal{F}_{1111}^{\mathbb{D}\mathbb{D}\mathbb{D}\mathbb{D}}(t) = \frac{F}{\Lambda}, \quad (\text{B1})$$

where \mathcal{K} is one of the kernels in (A35) and F is either zero or four.

The OPE expansion of $\mathcal{F}_{1111}^{\mathbb{D}\mathbb{D}\mathbb{D}\mathbb{D}}(t)$ in the $t \rightarrow 0$ channel takes the form

$$\mathcal{F}_{1111}^{\mathbb{D}\mathbb{D}\mathbb{D}\mathbb{D}}(t) = \sum_{\mathbb{O}} C_{\mathbb{D}\mathbb{D}\mathbb{O}}^2 \text{Block1D}[\Delta_{\mathbb{O}}, t], \quad \text{with} \quad \text{Block1D}[\Delta, t] = \left(\frac{t}{1+t}\right)^\Delta {}_2F_1\left(\Delta, \Delta, 2\Delta, \frac{t}{1+t}\right). \quad (\text{B2})$$

This expansion converges for $t \in [0, 1]$, [57]. Hence, after plugging it into the constraint (B1) we can exchange the order of integration and summation. In this way, we arrive at

$$\sum_{\mathbb{O}} C_{\mathbb{D}\mathbb{D}\mathbb{O}}^2 \widetilde{\text{Block}}(\Delta_{\mathbb{O}}) = \Lambda F, \quad \text{where} \quad \widetilde{\text{Block}}(\Delta) = \int_0^1 dt \mathcal{K}(t) \text{Block1D}[\Delta, t]. \quad (\text{B3})$$

For small Δ the integral may diverge. In such cases, it is evaluated by analytic continuation from larger Δ .

In the generic case where $\mathbb{O} \neq \mathbb{D}$ the four-point functions are integrated over the range $t \in [0, \infty]$. This integration domain can be split into the regions $t \in [0, 1]$ and $t \in [1, \infty]$. In the first region, the $t \rightarrow 0$ OPE expansion converges, and in the latter, the $t \rightarrow \infty$ OPE expansion converges. Using the convergent expansion in each region, we can exchange the order of summation and integration freely. In this way, we arrive at the sum rule of the form (13).

Appendix C: A new prediction for $O(N)$ defect at order $\mathcal{O}(\varepsilon^2)$

In the main text, we have presented a prediction for $C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}$, see (26). In this appendix, we use this result to obtain an integral constraint for the $\mathcal{O}(\varepsilon^2)$ correction to the four-point function that does not involve any additional unfixed conformal data.

We start by substituting $C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}$ from (26) into the homogeneous constraints (9) and consider the result at order $\mathcal{O}(\varepsilon^2)$. There are three sources of divergences, for the exchange of ϕ_1 , from the exchange of ϕ_1^2 , and from the exchange of ϕ_I^2 . The last two operators are degenerate at tree level and mix with each other at higher orders. We denote the two operators with a good anomalous dimension that result from this mixing by \mathbb{S}_+ and \mathbb{S}_- . Following the enhancement mechanism, we subtract from the four-point function the divergent contributions and separately add their finite coupling expressions before integration. We denote the subtracted four-point functions by \mathcal{F}_{reg} . Since the singular behavior of the constraints depends on the form of the integration kernels, the subtraction differs between the first and second homogeneous constraints in (9), and we treat them separately. For the first homogeneous constraint, we define the subtracted four-point function as

$$\mathcal{F}_{\text{reg},1}^{\phi_1\mathbb{D}\mathbb{D}\phi_1} = \begin{cases} \mathcal{F}^{\phi_1\mathbb{D}\mathbb{D}\phi_1} - C_{\mathbb{D}\mathbb{D}\phi_1}^2 \frac{1}{t^{\Delta_{\phi_1}}} & \text{for } 0 < t < 1 \\ \mathcal{F}^{\phi_1\mathbb{D}\mathbb{D}\phi_1} - C_{\mathbb{D}\mathbb{D}\phi_1} C_{\phi_1\phi_1\phi_1} \frac{1}{t^{\Delta_{\phi_1}}} \left(1 - \frac{\Delta_{\phi_1}}{2t} + \frac{(\Delta_{\phi_1}^2 + 2\Delta_{\phi_1} + 1)\Delta_{\phi_1}}{4(2\Delta_{\phi_1} + 1)t^2}\right) & \text{for } t > 1 \\ - \sum_{a=\pm} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a} C_{\phi_1\phi_1\mathbb{S}_a} \frac{1}{t^{\Delta_{\mathbb{S}_a}}} \left(1 - \frac{\Delta_{\mathbb{S}_a}}{2t}\right) & \end{cases}, \quad (\text{C1})$$

and

$$\mathcal{F}_{\text{reg},1}^{\mathbb{D}\phi_1\mathbb{D}\phi_1} = \begin{cases} \mathcal{F}^{\mathbb{D}\phi_1\mathbb{D}\phi_1} & \text{for } 0 < t < 1 \\ \mathcal{F}^{\mathbb{D}\phi_1\mathbb{D}\phi_1} - C_{\mathbb{D}\mathbb{D}\phi_1}^2 t^{\Delta_{\phi_1}-4} & \text{for } t > 1 \end{cases}, \quad (\text{C2})$$

where the subtracted finite coupling terms are understood to be expanded at small ε up to the same order as \mathcal{F} before integration.

Substituting those definitions into the first homogeneous constraint and separately adding back the enhanced contributions from the $\mathcal{O}(\varepsilon^3)$, we arrive at the constraint⁷

$$\begin{aligned} & \int_0^\infty dt \left[t^2 (\mathcal{F}_{\text{reg},1}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} + \mathcal{F}_{\text{reg},1}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1}) + (1+t)^2 \mathcal{F}_{\text{reg},1}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} \right]_{\mathcal{O}(\varepsilon^2)} \\ &= \frac{2\pi C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}}{\sqrt{N+8}} - \frac{2\pi C_{\phi_1 \phi_1 \phi_1}^{(2)}}{3\sqrt{N+8}} + \frac{7\pi^2}{3(N+8)} + \sum_{a=\pm} \left[C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} \left(2\gamma_{\mathbb{S}_a}^{(1)} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} + \frac{9}{2} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)} \right) + \frac{9}{2} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(1)} \right]. \end{aligned} \quad (\text{C3})$$

Similarly, for the second homogeneous constraint, we define the subtracted four-point function as

$$\mathcal{F}_{\text{reg},2}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} = \begin{cases} \mathcal{F}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1}, & \text{for } 0 < t < 1 \\ \mathcal{F}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} - C_{\mathbb{D}\mathbb{D}\phi_1} C_{\phi_1 \phi_1 \phi_1} \frac{1}{t^{\Delta_{\phi_1}}} \left(1 - \frac{\Delta_{\phi_1}}{2t} + \frac{(\Delta_{\phi_1}^2 + 2\Delta_{\phi_1} + 1)\Delta_{\phi_1}}{4(2\Delta_{\phi_1} + 1)t^2} \right) \\ \quad - \sum_{a=\pm} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a} C_{\phi_1 \phi_1 \mathbb{S}_a} \frac{1}{t^{\Delta_{\mathbb{S}_a}}} \left(1 - \frac{\Delta_{\mathbb{S}_a}}{2t} \right) & \text{for } t > 1 \end{cases}, \quad (\text{C4})$$

and $\mathcal{F}_{\text{reg},2}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1} = \mathcal{F}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1}$ with no subtractions. Substituting those definitions into the second homogeneous constraint and adding the enhanced contributions from the $\mathcal{O}(\varepsilon^3)$, we arrive at

$$\begin{aligned} & \int_0^\infty dt \left[2t t(1+t) \left(\mathcal{F}_{\text{reg},2}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} + \mathcal{F}_{\text{reg},2}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1} \right) - 2t \mathcal{F}_{\text{reg},2}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1} \right]_{\mathcal{O}(\varepsilon^2)} \\ &= \frac{\pi C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}}{\sqrt{N+8}} \frac{\pi C_{\phi_1 \phi_1 \phi_1}^{(2)}}{3\sqrt{N+8}} - \frac{\pi^2 \gamma_{\phi_1}^{(2)}}{N+8} + \frac{13\pi^2}{3(N+8)} + \sum_{a=\pm} \left[C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} (\gamma_{\mathbb{S}_a}^{(1)} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} + 5C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)}) + 5C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(1)} \right]. \end{aligned} \quad (\text{C5})$$

By taking a linear combination of (C3) and (C5) so that unknown structure constant at this order, $C_{\phi_1 \phi_1 \phi_1}^{(2)}$, drops out we arrive at the constraint

$$\begin{aligned} & \int_0^\infty dt \left[t^2 (\mathcal{F}_{\text{reg},1}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} + \mathcal{F}_{\text{reg},1}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1}) + (1+t)^2 \mathcal{F}_{\text{reg},1}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} + 2t t(1+t) \left(\mathcal{F}_{\text{reg},2}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} + \mathcal{F}_{\text{reg},2}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1} \right) - 2t \mathcal{F}_{\text{reg},2}^{\mathbb{D}\phi_1 \mathbb{D}\phi_1} \right]_{\mathcal{O}(\varepsilon^2)} \\ &= \frac{\pi^2 (2\gamma_{\phi_1}^{(2)} - 11) + 4\pi\sqrt{N+8} C_{\mathbb{D}\mathbb{D}\phi_1}^{(2)}}{N+8} + \sum_{a=\pm} \left[4\gamma_{\mathbb{S}_a}^{(1)} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} + \frac{29}{2} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} + \frac{29}{2} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(1)} \right]. \end{aligned} \quad (\text{C6})$$

Note that the OPE coefficient $C_{\phi_1 \phi_1 \mathbb{S}_a}$ up to $\mathcal{O}(\varepsilon)$ can be found in [19], while for the OPE coefficient $C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}$ can be found in the expansion of the four-point function of $\mathcal{F}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1}$. Therefore, the RHS of the equation consists of known CFT data and data that can be extracted from the four-point function at this order.⁸

In more detail, the part subtracted in the $t \in [1, \infty)$ region takes the form

$$\begin{aligned} t \in [1, \infty) : \quad \mathcal{F}_{\text{reg},1 \text{ or } 2}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} \Big|_{\mathcal{O}(\varepsilon^2)} &= \mathcal{F}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1} \Big|_{\mathcal{O}(\varepsilon^2)} - \varepsilon^2 \frac{(6t^2 - 3t + 2)}{6t^3} \left[C_{\phi_1 \phi_1 \phi_1}^{(1)} C_{\mathbb{D}\mathbb{D}\phi_1}^{(1)} \right] \\ &\quad - \varepsilon^2 \frac{1}{t^2} \sum_{a=\pm} \left[C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} + C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(1)} - \log t \gamma_{\mathbb{S}_a}^{(1)} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} \right] \\ &\quad - \varepsilon^2 \frac{1}{t^3} \sum_{a=\pm} \left[\frac{1}{2} \gamma_{\mathbb{S}_a}^{(1)} (2 \log t - 1) C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} - C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(0)} - C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1 \phi_1 \mathbb{S}_a}^{(1)} \right]. \end{aligned} \quad (\text{C7})$$

The first subtracted term corresponds to the ϕ_1 exchange and depends only on the known OPE data from (18). After subtracting this term, the combinations appearing on the right-hand side of (C6) can be extracted from the remaining $\mathcal{O}(1/t^2)$ terms:

⁷ Note that $C_{\phi_1 \phi_1 \mathbb{S}_a} = \mathcal{O}(1)$ while $C_{\mathbb{D}\mathbb{D}\mathbb{S}_a} = \mathcal{O}(\varepsilon)$.

⁸ Note that $C_{\phi_1 \phi_1 \phi_1}^{(2)}$ cannot be extracted from $\mathcal{F}_{\text{reg},2}^{\phi_1 \mathbb{D}\mathbb{D}\phi_1}$ because it appears multiplied by $C_{\mathbb{D}\mathbb{D}\phi_1}$ which starts at order $\mathcal{O}(\varepsilon)$.

- The coefficient of $\log t$ gives us the average $\sum_{a\pm} \gamma_{\mathbb{S}_a}^{(1)} C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1\phi_1\mathbb{S}_a}^{(0)}$.
- The $\log t$ -independent coefficient gives the average OPE coefficient $\sum_{a=\pm} \left[C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(2)} C_{\phi_1\phi_1\mathbb{S}_a}^{(0)} + C_{\mathbb{D}\mathbb{D}\mathbb{S}_a}^{(1)} C_{\phi_1\phi_1\mathbb{S}_a}^{(1)} \right]$.

Those data can also be extracted from the $\mathcal{O}(1/t^3)$ term.

Appendix D: $\mathcal{N} = 4$ Displacement OPE and Supercorrelator

1. Preliminary

The 1/2 BPS Wilson loop in $\mathcal{N} = 4$ SYM theory preserves an $\mathfrak{osp}(4^*|4)$ superconformal subgroup. Consequently, defect operators are labeled by their conformal dimension Δ , their representation under the $\mathfrak{so}(3)$ transverse rotation group, and an $\mathfrak{sp}(4) \simeq \mathfrak{so}(5)$ R -symmetry Dynkin label $[a, b]$. In the following, we will label them by $[a, b]_s^\Delta$.

The displacement operator \mathbb{D}_i , with $i = 1, 2, 3$ is the top component of the 1/2 BPS \mathcal{D}_1 supermultiplet, which consists of three states

$$\mathcal{D}_1 : [0, 1]_{s=0}^{\Delta=1} \rightarrow [1, 0]_{s=1}^{\Delta=3/2} \rightarrow [0, 0]_{s=2}^{\Delta=2}. \quad (\text{D1})$$

The OPE of two \mathcal{D}_1 supermultiplets contains the identity \mathcal{I} , the 1/2 BPS \mathcal{D}_2 supermultiplet, and an infinite set of unprotected *scalar* long multiplets $\mathcal{L}_{s=0, [0, 0]}^\Delta$

$$\mathcal{D}_1 \times \mathcal{D}_1 = \mathcal{I} + \mathcal{D}_2 + \sum_{\Delta} \mathcal{L}_{s=0, [0, 0]}^\Delta. \quad (\text{D2})$$

Here, the \mathcal{D}_2 multiplet consists of the states

$$\mathcal{D}_2 : [0, 2]_{s=0}^{\Delta=2} \rightarrow [1, 1]_{s=1}^{\Delta=5/2} \rightarrow [0, 1]_{s=2}^{\Delta=3} \oplus [2, 0]_{s=0}^{\Delta=3} \rightarrow [1, 0]_{s=1}^{\Delta=7/2} \rightarrow [0, 0]_{s=0}^{\Delta=4}. \quad (\text{D3})$$

Details of those multiplets and their properties can be found in [14, 15, 58–60].

We turn to the structure of the four-point function $\langle\langle \mathcal{D}_1 \mathcal{D}_1 \mathcal{D}_1 \mathcal{D}_1 \rangle\rangle$. Superconformal symmetry fixes this correlator up to an unknown function of the cross-ratio called *the reduced correlator* $f(\chi)$

$$\frac{\langle\langle \mathcal{D}_1 \mathcal{D}_1 \mathcal{D}_1 \mathcal{D}_1 \rangle\rangle}{\langle\langle \mathcal{D}_1 \mathcal{D}_1 \rangle\rangle \langle\langle \mathcal{D}_1 \mathcal{D}_1 \rangle\rangle} = \mathbf{F} \mathfrak{X} + \hat{\mathbf{D}} f(\chi), \quad (\text{D4})$$

where \mathbf{F} is a constant, \mathfrak{X} is the super cross-ratio, and $\hat{\mathbf{D}}$ is a differential operator. We refer the reader to [13] for their definition. In particular, in terms of the reduced correlator, crossing symmetry takes the form

$$\chi^2 f(\chi) = (1 - \chi)^2 f(1 - \chi). \quad (\text{D5})$$

To check the constraints, we consider the four-point function of the displacement operator. It takes the form⁹

$$\langle\langle \mathbb{D}_i(x_1) \mathbb{D}_j(x_2) \mathbb{D}_k(x_3) \mathbb{D}_l(x_4) \rangle\rangle = \frac{1}{x_{12}^4 x_{34}^4} \left[\delta_{ij} \delta_{kl} \mathcal{F}_S(\chi) + (\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk}) \mathcal{F}_A(\chi) + \left(\delta_{ik} \delta_{jl} + \delta_{il} \delta_{jk} - \frac{2}{3} \delta_{ij} \delta_{kl} \right) \mathcal{F}_T(\chi) \right], \quad (\text{D6})$$

⁹ We thank Michelangelo Preti for kindly sharing a `Mathematica` code for the superspace computations.

where the cross ration χ is related to our cross ratio t as $\chi = \frac{t}{1+t}$, and

$$\begin{aligned}
\mathcal{F}_S(\chi) &= -\frac{1}{108}(\chi-1)^2((\chi-3)\chi+3)\chi^4 f^{(5)}(\chi) - \frac{1}{108}(\chi-1)(\chi(\chi(16\chi-33)+3)+18)\chi^3 f^{(4)}(\chi) \\
&\quad - \frac{1}{54}(\chi(\chi(\chi(37\chi-54)+9)-36)+45)\chi^2 f^{(3)}(\chi) + \frac{1}{3}(-3\chi^4+\chi^3-6\chi+9)\chi f''(\chi) \\
&\quad + \left(-\frac{\chi^4}{3}+4\chi-7\right) f'(\chi) + \left(\frac{8}{\chi}-4\right) f(\chi) + \frac{\mathbb{F}\chi^4}{3}, \\
\mathcal{F}_A(\chi) &= -\frac{1}{72}(\chi-2)(\chi-1)^2\chi^5 f^{(5)}(\chi) - \frac{1}{36}(\chi-1)(\chi(8\chi-13)+3)\chi^4 f^{(4)}(\chi) \\
&\quad - \frac{1}{36}(\chi(\chi(37\chi-53)+9)+6)\chi^3 f^{(3)}(\chi) + \frac{1}{4}(-6\chi^5+3\chi^4+2\chi^2) f''(\chi) \\
&\quad - \frac{1}{2}(\chi^3+2)\chi f'(\chi) + f(\chi) + \frac{\mathbb{F}\chi^4}{2}, \\
\mathcal{F}_T(\chi) &= -\frac{1}{72}(\chi-1)^2\chi^6 f^{(5)}(\chi) - \frac{1}{18}(\chi-1)(4\chi-3)\chi^5 f^{(4)}(\chi) \\
&\quad - \frac{1}{36}(\chi(37\chi-51)+15)\chi^4 f^{(3)}(\chi) + \frac{1}{4}(5-6\chi)\chi^4 f''(\chi) - \frac{1}{2}\chi^4 f'(\chi) + \frac{\mathbb{F}\chi^4}{2}.
\end{aligned} \tag{D7}$$

The $\mathcal{D}_1\mathcal{D}_1$ OPE shown in (D2) implies that the reduced correlator can be expanded as:

$$f(\chi) = f_{\mathbf{1}} + C_{\mathcal{D}_1\mathcal{D}_1\mathcal{D}_2}^2 f_{\mathcal{D}_2} + \sum_{\mathbb{O}} C_{\mathcal{D}_1\mathcal{D}_1\mathbb{O}}^2 f_{\mathbb{O}}, \tag{D8}$$

where the terms $f_{\mathbf{1}}$, $f_{\mathcal{D}_2}$, and $f_{\mathbb{O}}$ are referred to as *reduced blocks*, serving as direct analogs of the standard conformal blocks. As shown in [13–15], supersymmetry fixes their form to be

$$\begin{aligned}
f_{\mathbf{1}} &= \chi, \\
f_{\mathcal{D}_2} &= \chi - \chi {}_2F_1(1, 2; 4; \chi) = \mathcal{O}(\chi^2), \\
f_{\mathbb{O}} &= \frac{\chi^{\Delta_{\mathbb{O}}+1} {}_2F_1(\Delta_{\mathbb{O}}+1, \Delta_{\mathbb{O}}+2; 2\Delta_{\mathbb{O}}+4; \chi)}{1-\Delta_{\mathbb{O}}} = \mathcal{O}(\chi^{\Delta_{\mathbb{O}}+1}).
\end{aligned} \tag{D9}$$

At a generic value of the YM coupling, there are no $\mathfrak{sp}(4)$ singlets operators with integer dimension $\Delta < 4$. This fact is sufficient for our constraints to hold, see section A 2. We test them below using the CFT data equated above.

2. Simplifying the Homogeneous Constraint

Here, we will show that the homogeneous constraints fix the leading OPE behavior of the reduced correlator. After subtracting \mathcal{F}_{GFF} , we plug the form of the correlator (D7) into the homogeneous constraints (9) and perform multiple integrations by parts to remove the derivatives from the reduced correlator $f(\chi)$. After this, we find that the integrations over the reduced correlator reduce to a set of boundary terms. We are left with the integration of a topological term, $\mathbb{F}\mathfrak{X}$ in (D4), and the homogeneous constraint reduces to

$$\int_0^1 d\chi \mathbb{F} (\chi^\ell \delta_{ij} \delta_{IJ} + \chi^{2-\ell} \delta_{iI} \delta_{jJ} + (\chi-1)^\ell \chi^{2-\ell} \delta_{iJ} \delta_{jI}) = [\text{boundary terms}], \tag{D10}$$

where $\ell = 0, 1, 2$ corresponds to the three homogeneous constraints in (9). We have verified that (D10) is satisfied, provided that the reduced correlator has the following asymptotics behavior close to $\chi = 0$ and $\chi = 1$

$$f(\chi) = -\mathbb{F}\chi^2/2 + \mathcal{O}(\chi^3), \quad f(\chi) = \mathbb{F}/2 + \mathcal{O}((1-\chi)^2). \tag{D11}$$

These two behaviors are related by crossing. They are the result of the exchange of the \mathcal{D}_2 multiplet, see (D9). The other operators in the OPE (D2) are scalar long multiplets. They give contributions of order $\mathcal{O}(\chi^{\Delta_{\mathbb{O}}+1})$. These are subleading due to the unitarity bound, $\Delta_{\mathbb{O}} > 1$. This asymptotic behavior fixes the value of the protected OPE coefficient, defined in (D8), as $C_{\mathcal{D}_1\mathcal{D}_1\mathcal{D}_2}^2 = \mathbb{F}$. It agrees with the finite-coupling asymptotics in [13], where this structure constant is fixed by supersymmetric localization.

3. Simplifying the Inhomogeneous Constraint

As before, we subtract from (D4) \mathcal{F}_{GFF} and plug the result into the inhomogeneous constraints (11) and (12). We then perform integrations by parts to remove the derivatives from the reduced correlator. We first consider the constraint (11) and contract it with $\delta_{I_i}\delta_{J_j}$. Since there may be potential divergences at the integration boundaries, we introduce a cut-off ϵ_1 near $\chi = 1$ and ϵ_0 near $\chi = 0$. After several integrations by parts, the equation reduces to

$$\int_{\epsilon_0}^{1-\epsilon_1} d\chi \left[2f(\chi) \left(\frac{1}{\chi} - \frac{1}{\chi^3} + \frac{1}{3\chi^2} + \frac{1}{3(1-\chi)} \right) \right] + [\text{boundary terms}] = \frac{4}{\Lambda}. \quad (\text{D12})$$

Using (D11) from the homogeneous constraint and the explicit form of the $\text{F}\mathfrak{X}$ term, the boundary terms take the form

$$[\text{boundary terms}] = -\frac{4\text{F}}{3} + \frac{\text{F}}{3} \log(\epsilon_1) + \text{F} \log(\epsilon_0) + \mathcal{O}(\epsilon_0, \epsilon_1). \quad (\text{D13})$$

Here, the last two cutoff-dependent terms can be written as

$$\frac{\text{F}}{3} \log(\epsilon_1) + \text{F} \log(\epsilon_0) = -\frac{\text{F}}{3} \int_{\epsilon_0}^{1-\epsilon_1} \frac{d\chi}{1-\chi} - \text{F} \int_{\epsilon_0}^{1-\epsilon_1} \frac{d\chi}{\chi} + \mathcal{O}(\epsilon_0, \epsilon_1). \quad (\text{D14})$$

Adding them back to the bulk integration, we observe that it becomes *finite* due to the boundary asymptotics given in (D11). As a result, the equation reduces to

$$\int_0^1 d\chi \left[2f(\chi) \left(\frac{1}{\chi} - \frac{1}{\chi^3} + \frac{1}{3\chi^2} + \frac{1}{3(1-\chi)} \right) - \frac{\text{F}}{3} \frac{1}{1-\chi} - \frac{\text{F}}{\chi} - \frac{4\text{F}}{3} \right] = \frac{\Lambda}{4}. \quad (\text{D15})$$

To further simplify the result, we notice that the crossing (D5) implies that

$$\int_0^1 d\chi f(\chi) = \int_0^1 d\chi \frac{f(\chi)}{\chi}, \quad \int_0^1 d\chi \frac{f(\chi)}{\chi^2} = 0. \quad (\text{D16})$$

Thus we can rewrite (D15) as

$$\frac{4}{3} \int_0^1 d\chi \left[f(\chi) \left(1 - \frac{2}{\chi^3} \right) - \left(1 + \frac{1}{\chi} \right) \text{F} \right] = \frac{4}{\Lambda}, \quad (\text{D17})$$

where we have simultaneously performed the change of variable $\chi \rightarrow 1-\chi$ for the terms with $1-\chi$ in the denominator. This form of the equation is finite due to (D11). This constraint (D17) is in agreement with a relation that was found and tested in [24, 49].

The analysis of the other inhomogeneous equations follows a similar procedure. We find that they reduce to multiples of (D17). The only exception is the $\delta_{iI}\delta_{jJ}$ term in the first inhomogeneous constraint, which, in addition to involving multiples of (D17), also depends on two subleading Taylor coefficients of the reduced correlator near $\chi = 0$. Given (D17), these new contributions must cancel among themselves. This leads to the condition

$$6a_3 - 4a_4 + \text{F} = 0, \quad \text{where} \quad f(\chi) = -\frac{\text{F}}{2}\chi^2 + a_3\chi^3 + a_4\chi^4 + \mathcal{O}(\chi^5). \quad (\text{D18})$$

At large coupling the dimension of long operators is large. Therefore, they do not contribute to the small χ OPE expansion up to order $\mathcal{O}(\chi^4)$. Since $f(\chi)$ is expected to be analytic in the 't Hooft coupling, they do not contribute to (D18) at finite coupling. As a result, the coefficients a_3 and a_4 receive contributions only from the protected multiplet \mathcal{D}_2 . We find that the expansion of the reduced block, $f_{\mathcal{D}_2}$ in (D9), indeed satisfies this relation.

In conclusion, for the 1/2 BPS line in $\mathcal{N} = 4$ SYM theory, the constraints agree with the available CFT data.

Appendix E: 1/2 BPS defects in $AdS_3 \times S^3 \times T^4$

Type IIB string theory in $AdS_3 \times S^3 \times T^4$ is dual to a 2 dimensional CFT. The geometric background $AdS_3 \times S^3 \times T^4$ can support both Ramond-Ramond (RR) and Neveu-Schwarz–Neveu-Schwarz (NS-NS) three-form fluxes. For our purposes, it suffices to know that the mixed fluxes are parameterized by a real angle $\lambda \in [0, \pi/2]$ that extrapolates between the pure RR ($\lambda = 0$) and the pure NS-NS ($\lambda = \pi/2$) cases. This parameter represents a true moduli of the dual CFT. Additionally, the theory depends on the string coupling g_s which is related to an integer parameter, N , in the CFT¹⁰ and the ratio between the string scale and the AdS radius that is related to a sort of 't Hooft coupling in the dual CFT, $g = R_{\text{AdS}}^2/(2\pi l_s^2)$.

In [17], the authors have studied a classical string configuration in this background that is holographically dual to a $\frac{1}{2}$ -BPS line defect in the 't Hooft limit where the CFT is strongly coupled. We will test the constraints to leading nontrivial order at strong coupling, $g \rightarrow \infty$.

At the leading order in $1/g$, the four-point function of the displacement operator is given by \mathcal{F}_{GFF} (8), which we subtract. The next order, $\mathcal{O}(1/g)$, has been computed in two different ways. One is based on tree-level diagrams in the effective theory on the classical AdS_2 string. The other is done using analytical bootstrap methods. The analytical bootstrap result depends on two undetermined parameters, b_1 and b_2 , and takes the following form

$$\begin{aligned}
 g\mathcal{F}_{1111} = & -\frac{2(t(t+1)(t(t(t(2t+9)+16)+14)+6)+2)(b_1-2b_2)\log(t)}{3(t+1)^5} \\
 & -\frac{4b_1(t^2+t+1)(t(t+1)(t(3(t+2)t^2+t-2)+9)+3)}{9(t+1)^4t^4} \\
 & +\frac{2b_2(t^2+t+1)(t(t+1)(t(12(t+2)t^2+t-11)+36)+12)}{9(t+1)^4t^4} \\
 & +\frac{2(2t^6+t^5+t+2)(b_1-2b_2)\log(t+1)}{3t^5},
 \end{aligned} \tag{E1}$$

where g enters the bootstrap through

$$\Lambda = \frac{6g}{\pi} + \mathcal{O}(g^0). \tag{E2}$$

The effective string result, on the other hand, depends on only one parameter, λ . By comparison of the two results, [17] determine,

$$b_1 + 4b_2 + 6/\pi = 0. \tag{E3}$$

In the following, we find that our integrated constraints can also be used to fix this relation without using the result of the holographic computation. Note that without additional input, any bootstrap computation cannot fix a moduli of a CFT such as λ . Its relation to these parameters is found to be $b_1 = -2\sin^2(\lambda)/\pi$.

A new feature that appears in this case is that the OPE of two displacement operators includes the displacement operator itself. That is because the NS-NS flux breaks parity and thus reflection in the transverse direction is not a symmetry of the straight defect. This fact is reflected in the small- t expansion of (E1)

$$g\mathcal{F}_{1111}(t) = -\frac{b_1}{3}(t^{-2} - t^{-1} + \mathcal{O}(t^0)), \tag{E4}$$

where the first two terms match with the expansion of the conformal block of a dimension two operator.

We have repeated the procedure described in the main text for the exchange of an operator of dimension Δ and send $\Delta \rightarrow 2$ at the end. We find that the homophonous constraints are satisfied while the inhomogeneous constraints reduce to the condition (E3). Hence, the constraints are compatible with both the string theory and the bootstrap computations.

¹⁰ To be more precise, the string theory is dual to a grand canonical ensemble of CFTs with different N , [61].

Appendix F: Conformal Block in One Dimension

The generic scalar four-point function can be expanded as [62, 63]

$$\langle\langle \mathbb{O}_{\Delta_1}(x_1)\mathbb{O}_{\Delta_2}(x_2)\mathbb{O}_{\Delta_3}(x_3)\mathbb{O}_{\Delta_4}(x_4) \rangle\rangle = \left(\frac{x_{24}}{x_{14}}\right)^{\Delta_2-\Delta_1} \left(\frac{x_{14}}{x_{13}}\right)^{\Delta_3-\Delta_4} \frac{\widehat{\mathcal{F}}(\chi)}{x_{12}^{\Delta_1+\Delta_2} x_{34}^{\Delta_3+\Delta_4}}, \quad (\text{F1})$$

where

$$\chi = \frac{x_{12}x_{34}}{x_{13}x_{24}} = \frac{t}{1+t}, \quad (\text{F2})$$

is the cross-ratio. It is convenient to work with this cross-ratio in this section, as it simplifies the differential equation of the conformal block.

The function $\widehat{\mathcal{F}}(\chi)$ can be expanded in conformal blocks. In the S -channel, the expansion is given by:

$$\widehat{\mathcal{F}}(\chi) = \sum_{\tilde{\mathbb{O}}} C_{\mathbb{O}_{\Delta_1}\mathbb{O}_{\Delta_2}\tilde{\mathbb{O}}} C_{\tilde{\mathbb{O}}\mathbb{O}_{\Delta_3}\mathbb{O}_{\Delta_4}} \text{Block1D}_{\Delta_{\tilde{\mathbb{O}}}}(\chi), \quad (\text{F3})$$

where the one-dimensional conformal block can be expressed in terms of the hypergeometric ${}_2F_1$ function as [64, 65]

$$\text{Block1D}_{\Delta}(\chi) = \chi^{\Delta} {}_2F_1(\Delta - \Delta_1 + \Delta_2, \Delta + \Delta_3 - \Delta_4; 2\Delta; \chi). \quad (\text{F4})$$

For our purposes, it is sufficient to consider the conformal block for a scalar intermediate operator.

Since our normalization of the four-point function differs from that used in (F1), we must account for this change. Specifically, we have:

$$\text{Block1D}_{\Delta}^{\text{ODDO}}(\chi) = \text{Block1D}_{\Delta}^{\text{DODO}}(\chi) = (1-\chi)^4 \chi^{\Delta-\Delta_0-2} {}_2F_1(\Delta - \Delta_0 + 2, \Delta - \Delta_0 + 2; 2\Delta; \chi). \quad (\text{F5})$$

These are equal because we have rescaled $\mathcal{F}^{\text{DODO}}$ as defined in (10).

In the main text, we will also consider the expansion in the T -channel where we take $x_1 \rightarrow x_4$ and $x_2 \rightarrow x_3$,

$$\widehat{\mathcal{F}}(\chi) = \sum_{\tilde{\mathbb{O}}} C_{\mathbb{O}_{\Delta_2}\mathbb{O}_{\Delta_3}\tilde{\mathbb{O}}} C_{\tilde{\mathbb{O}}\mathbb{O}_{\Delta_4}\mathbb{O}_{\Delta_1}} \overline{\text{Block1D}}_{\Delta_{\tilde{\mathbb{O}}}}(\chi), \quad (\text{F6})$$

the conformal blocks $\overline{\text{Block1D}}$ can be obtained from Block1D by crossing, which takes

$$x_i \rightarrow x_{(i+1) \bmod 4}, \quad \chi \rightarrow 1-\chi, \quad \Delta_i \rightarrow \Delta_{(i+1) \bmod 4}. \quad (\text{F7})$$

Taking into account the change in the prefactor from (F1), the crossed conformal block is given by:

$$\overline{\text{Block1D}}_{\Delta}(\chi) = \chi^{\Delta_3+\Delta_4} (1-\chi)^{-\Delta_2-\Delta_3+\Delta} {}_2F_1(\Delta - \Delta_2 + \Delta_3, \Delta - \Delta_1 + \Delta_4; 2\Delta; 1-\chi). \quad (\text{F8})$$

For the four-point function of interest, the corresponding conformal blocks in the crossed channel, after accounting for the change in prefactor from (7) and (10), read:

$$\begin{aligned} \overline{\text{Block1D}}_{\Delta}^{\text{ODDO}}(\chi) &= (1-\chi)^{\Delta} {}_2F_1(\Delta, \Delta; 2\Delta; 1-\chi), \\ \overline{\text{Block1D}}_{\Delta}^{\text{DODO}}(\chi) &= (1-\chi)^{\Delta-\Delta_0+2} {}_2F_1(\Delta - \Delta_0 + 2, \Delta + \Delta_0 - 2; 2\Delta; 1-\chi). \end{aligned} \quad (\text{F9})$$