Solar Particle Acceleration

Donald V. Reames (https://orcid.org/0000-0001-9048-822X)
Institute for Physical Science and Technology, University of Maryland, College Park, MD, USA 20742

Abstract High-energy particles may be accelerated widely in stellar coronae; probably by the same processes we find in the Sun. Here, we have learned of two physical mechanisms that dominate the acceleration of solar energetic particles (SEPs). The highest energies and intensities are produced in "gradual" events at shock waves driven from the Sun by fast, wide coronal mass ejections (CMEs). Smaller, but more numerous, "impulsive" events with unusual particle composition are produced during magnetic reconnection in solar jets and flares. Jets provide open magnetic field lines where SEPs escape; closed magnetic loops contain this energy to produce bright, hot flares, perhaps even contributing to heating the low corona in profuse nanoflares. Streaming protons amplify Alfvén waves upstream of the shocks. These waves scatter and trap SEPs and, in large events, modify the element abundances and flatten the low-energy spectra upstream. Shocks also reaccelerate residual ions from earlier impulsive events, when available, that characteristically dominate the energetic heavy-ion abundances. The large CME-driven shock waves develop an extremely wide longitude span, filling much of the inner heliosphere with energetic particles.

Keywords Solar cosmic rays · Solar energetic particles · Shock waves · Coronal mass ejections · Solar jets · Solar flares · Solar system abundances

1 Introduction

High-energy particles are likely accelerated in the coronae of most stars, but we can only study them directly near the Sun. Energetic particle radiation constitutes a recurring hazard to modern astronauts and missions in space and may have influenced the evolution of life here and on exoplanets around other stars.

Particle acceleration can only occur where the density is low enough that particles do not immediately dissipate their energy in Coulomb collisions. We have found that the most powerful physical processes for acceleration of solar energetic particles (SEPs) involve shock waves driven out from the Sun by fast, wide coronal mass ejections (CMEs) [1, 2], but regions of magnetic reconnection and turbulence in solar jets also produce electron-rich bursts of SEPs with unusual and distinctive element abundances although at lower average energies and intensities.

Energetic particles can be accelerated in and above the corona where the plasma densities fall rapidly below about 10° atoms cm⁻³; they rapidly lose energy in Coulomb scattering and stop at the higher densities of the low corona and chromosphere. The magnetic energy of reconnection can spawn energetic particles in the corona or generate heat below. In solar flares, accelerated particles trapped on closed magnetic loops soon scatter into the loss cone and plunge into denser plasma below to produce heat, X-rays, gamma-rays, and thermal EUV emission. Particles accelerated on open field lines in solar jets, called *impulsive* SEPs, escape along field lines where the streaming electrons generate type III radio emission. A signature of these impulsive SEPs is their unusual abundances including 10,000-fold enhancements of ³He/⁴He and a systematic enhancement of increasingly heavy elements continuing as high as Pb [3 – 14]. In contrast, fast (>>500 km s⁻¹) CMEs drive shock waves that accelerate particles in *gradual* SEP events [1] that include the widespread, intense, energetic, and long-lasting SEP events [15 – 20].

2 Magnetic Reconnection

Since the Sun is composed of ionized plasma, differences in forces on ions and electrons, especially in differential rotation, cause currents that generate magnetic fields. Magnetic fields begin deep in the Sun but are deformed at the *tachocline*, the boundary between the inner radiative zone, which rotates uniformly, and the convective zone, which rotates faster at the equator than at the poles. As the plasma swirls, magnetically active regions or sunspots are visible at the radiatively-cooled photosphere [21]. When the fields become twisted and tangled, oppositely directed fields are brought together to cancel and reconnect, releasing energy. The magnetic energy that vanishes during reconnection drives all the activity on the Sun that we see as flares, coronal mass ejections (CMEs), solar energetic particles (SEPs), X-rays, gamma-rays, and radio emission. Energy trapped in closed field configurations is emitted eventually as photons radiated from flares or can help drive out CMEs.

Patterns of field reconnection can be complicated, but Figure 1 shows two examples of topology that results when reconnection occurs either between open and closed lines, which allows SEPs to escape, or among completely closed lines, where they cannot. In Figure 1a, emerging loops have opposite polarity from the overlying field lines, with the energy in islands of reconnection driving SEP acceleration in the

2

collapsing current sheet [22 – 24], thus producing a solar jet. In Figure 1a, newly opened field lines on the right are matched by newly closed fields on the left which capture some SEPs. These will eventually plunge into the denser corona to produce a hot solar flare. Thus, jets may always have an accompanying flare, but the SEPs we see do not come from this flaring region. This is why SEP properties were historically found to correlate poorly with associated X-ray intensities, for example [25, 26].

Figure 1b shows a topology where shear in the footpoints of an arcade of loops produces reconnection where all the field lines remain closed. All accelerated particles are trapped and eventually plunge into the footpoints of the loops to produce a flare with no SEPs in space [28]. However, the liberated flux rope shown in the figure may rise as part of a CME which can be fast enough to form a shock wave and produce a gradual SEP event. Note that the right-most cartoon in Figure 1 is often viewed end-on and the flux rope is shown as a completely open region (e.g. the Kopp and Pneuman model [27]). Other cartoons show it as a closed circle, disconnected from the Sun. In the likely topology, all particles accelerated in this reconnecting region plunge into the Sun to produce flaring.




Figure 1 Examples of the topology of magnetic reconnection are shown: (a) In a solar jet, initially open field lines of one polarity (*black*) reconnect with newly emerging loops (*blue*) in the *red hashed* region where accelerated SEPs can escape easily, however, some of the SEPs are captured in a newly closed region on the left, where they produce a hot flare. (b) A sheared arcade of field lined can reconnect in the *red hatched* regions to produce a semi-detached flux rope that can rise to form a CME. Here, all SEPs are trapped on closed field lines and eventually plunge into the denser corona to produce a flare. SEPs are accelerated in collisionless plasma; flaring is produced in collisional plasma [28].

Figure 1 does not cover all possibilities. If the reconnection region is too deep in the corona, there are no SEPs and the magnetic energy is dissipated as heat. Modifications of the topology in Figure 1a may also produce a CME and that CME may drive a shock wave that can reaccelerate the SEPs from the reconnection region. We will consider this possibility below. Of course, jets can be more complicated than the one shown [29-31].

3 Impulsive Events

Impulsive events were first associated with the streaming electrons that produce type III radio bursts, as distinct from the shock-accelerated electrons producing type II bursts [32, 33]. Later, some small SEP events were found to have 10,000-fold enhancements of ${}^{3}\text{He}/{}^{4}\text{He} > 1$ [34], and these electron- and ${}^{3}\text{He}$ -rich events were subsequently found to be the same [35, 36]. It became clear that ${}^{3}\text{He}$ enhancements were a resonance phenomenon produced by waves near the ${}^{3}\text{He}$ gyrofrequency = $Q\omega_{p}/A$, where ω_{p} is the proton gyrofrequency, and the charge-to-mass ratio Q/A = 2/3 for ${}^{3}\text{He}$. The wave spectrum could be damped at the gyrofrequencies of the dominant ions, H at Q/A = 1, and ${}^{4}\text{He}$ at Q/A = 2/4, but the low abundance of plasma ${}^{3}\text{He}$ (${}^{3}\text{He}/{}^{4}\text{He} \approx 5 \times 10^{-4}$) is unable to damp

the resonant waves at that gyrofrequency. Many wave modes were suggested for the selective preheating of ³He [37-44], however, Temerin and Roth [45] suggested a more complete model with electromagnetic ion-cyclotron waves being generated by the streaming electrons that could be resonantly absorbed by mirroring ³He ions to accelerate them. As acceleration increases, available ³He can become depleted completely [46], which contributes to spectral differences between ³He and ⁴He [47 – 49].

Subsequent observations also showed heavy-ion enhancements up to Fe [50, 10] that seemed to increase as A/Q [8]. Later observations extended this pattern across the periodic table of elements, seen up to about Pb [4, 9, 11, 12]. This behavior was unlike the sharp resonance for 3 He and was first associated with cascading waves [51 – 53] and later associated with a shock-like acceleration as ions are scattered back and forth from the ends of collapsing islands of magnetic reconnection [22].

4 Shock Acceleration

The dominant mechanism for producing high-energy particles is acceleration by shock waves [54] driven out by the Sun by fast, wide coronal mass ejections (CMEs) with speeds of $\sim 800 - 3000$ km s⁻¹. When the magnetic field is quasi-parallel to the shock normal, particles can be scattered back and forth across the shock by resonant scattering against waves upstream and downstream [55, 56, 15, 16, 57]. As the particles stream away from the shock at each higher energy, they increase the scattering by amplifying resonant waves [58, 59] of wave number k with

$$k \approx B/\mu P \tag{1}$$

where P in the magnetic rigidity, i.e. momentum per unit charge, of the particle and μ is its pitch angle with respect to the magnetic field of vector \mathbf{B} . On each transit of the shock, particles gain an increment in velocity related to the plasma-wave velocity difference across the shock. The growth of self-amplified waves is an essential aspect of shock acceleration. For quasi-parallel regions of the shock, where the magnetic field lies nearly in the shock plane, particles are also accelerated in the $\mathbf{V}_{shock} \times \mathbf{B}$ electric field [60]. Some early authors [15] assumed $\mu \approx 1$, so each rigidity had its own unique resonant wave number, a convenient simplification.

For even greater simplicity, many authors assume that particle scattering does not vary at all with time. More generally, Bell [56, 57] assumed that equilibrium is maintained between particles and resonant waves and this assumption was applied to SEP shocks by Lee [15, 16]. Later Ng and Reames [57] allowed full time evolution of both particles and waves, allowing estimates of the timescale of shock acceleration.

Element abundances accelerated by shock waves tend to reflect the abundances in the corona at 2-3 solar radii [61-63] where the ions are first sampled. The coronal abundances differ from abundances in the photosphere as a function of the first ionization potential (FIP) because of ion-neutral fractionation as the particles cross the chromosphere to the corona [64, 65] where they all become ionized. Averaged SEP abundances can be a good measure of coronal abundances [66, 67] although their FIP pattern differs from that seen in the solar wind [68-71]. SEPs are *not* just accelerated solar wind.

For impulsive SEP events, we were able to neglect scattering of the ions during transport since impulsive events have low intensities, the particles tend to propagate scatter free [72], and they are focused by the diverging magnetic field. However, wave

scattering is a basis of shock acceleration since the particles streaming away grow waves resonant with higher- and higher-energy ions during acceleration, and these waves extend farther and farther from the shock as intensities increase [16, 57]. The wave intensity and thus particle scattering is a strong function of space and time. For different ion species of a given velocity, the rigidity dependence of scattering tends to trap low-rigidity (low-A/Q) ions near the shock while the higher-rigidity (higher-A/Q) ions leak away. This power-law dependence upon A/Q was first directly observed by Breneman and Stone [73]. Thus, early in an event, Fe/O will be first observed to be enhanced, for example, although later, when the shock approaches, Fe/O will always be depressed from its coronal value. With the proper averaging we should be able to recover the source coronal abundance. In small gradual events the depletion in Fe/O, from Fe loss by preferential leakage, will be most noticeable throughout. In large gradual events, preferential trapping of O near the shock can cause elevated Fe/O to persist for a day or so at moderate energies until the shock actually passes over the observer.

During the early acceleration, as intensities of protons of a given rigidity rise at a shock, they first stream away, causing more and more resonant waves to grow. Wave scattering reduces this streaming, until an equilibrium limit is reached where increasing intensities at the shock just grow enough additional waves to stop the streaming out to observers away from the shock. This has been called the "streaming limit" that causes an intensity plateau [74-76], or the "flat spectra" rediscovered recently [77]. Figure 2a shows superposed time profiles for several large SEP events where the early proton intensities near 1 AU are bounded at just over 200 protons cm⁻² sr⁻¹ s⁻¹ MeV⁻¹ although in some events intensities rise to much higher values later, near the shock [74]. If source intensities increase, the protons streaming outward simply grow more resonant waves so as to restrict the streaming, trapping ions near the source. Figure 2b shows flattening of plateau H and O spectra, typical of all species, for 5 SEP events [76]. Figure 2c shows the importance of high intensities at ~ 10 MeV for flattening the intensities at energies below [78-80]. Ng [81] discusses the way the limiting proton intensities vary with radius and with plasma and shock parameters.

The streaming limit is a common feature in all large gradual SEP events with sufficiently high intensities. For stronger shocks, the flattening extends to higher energies. It is a transport-generated limit involving equilibrium between streaming protons and resonant waves. When there is no streaming, e.g. at the shock, there is no intensity limit. Also, when intensities rise extremely rapidly, they can briefly exceed the limit, until enough waves grow to establish it. This situation occurs in the Jan 20 2005 event, as shown in [80], and is well described by theories that follow the time evolution of both particles and waves [79].




Figure 2 (a) Superposed time profiles of 3-6 MeV protons show limited plateau intensities early in large SEP events that often rise later when the shock approaches [74]. (b) H and O plateau spectra for several events show flattening at low energies [76]. (c) Theoretical fits are compared, using the Ng et al. [79] theory, for the flattened plateau proton spectra of the large 2003 October 28 event and the un-flattened, power-law spectrum in the smaller event of 1998 May 2 [78]. The ~100-fold intensity difference above ~10 MeV is emphasized by the gray arrow; the higher intensities amplify more waves that restrict outward flows, trapping ions and flattening spectra.

4.1 Spatial Distributions

Spatial distributions of SEPs driven by shock acceleration can be quite extensive. The earliest particles are released at \sim 2 solar radii [61, 62] and particles from these earliest spectra are effectively the dominant seed population for all additional acceleration. As the shock moves outward wave turbulence tends to trap particles with the shock, but B decreases on average, so the existing waves tend to resonate with ions of lower rigidity (Eq. 1), allowing the highest rigidity ions tend to leak away first.

The earliest particles that escape the shock when it is near the Sun usually contain energies of greatest radiation hazard. Figure 2b shows early proton spectra up to 700 MeV and some are flat out to \sim 50 MeV. A recent measurement near the Sun at 0.35 AU [23] also observed the flat spectra but only below \sim 1 MeV. Figure 3 shows measurements by spacecraft conveniently located to show the radial evolution of the central region in a gradual SEP event. *Helios 1* is well connected to the center of the shock near the Sun, *Helios 2* and IMP 8 when it is near 1 AU, and *Voyager 2* when it is near 2 AU, as shown in Figure 3b. Proton intensities at 6-11 MeV are compared in Figure 3a and more complete energy coverage is shown in Figures 3c-3e. Protons of 100-200 MeV peak early, are still visible at 1 AU, but have dissipated by 2 AU. In contrast 20-40 MeV protons still peak at the shock near 2 AU in this event.




Figure 3 (a) The radial evolution of 6-11 MeV proton intensities is captured by four spacecraft as the accelerating shock wave moves out to ~ 2 AU in the 1978 Jan 1 SEP event. (b) The spatial distribution of the spacecraft is shown and four shock positions S1 – S4 correlate with times in (a). Full energy coverage vs. time from $\sim 3-200$ MeV is shown for (c) *Helios 1*, (d) IMP 8, and (e) *Voyager 2* [18, 82].

While the *Helios*-IMP8-*Voyager* period provided several revealing coincidences including a case of successive strikes of field lines by both flanks of a wide shock in September 1978 [18, 82], the STEREO period showed the wide distribution of SEPs and of the shock waves that produce them. The example in Figure 4 shows intensities of 20 MeV protons at the widely-spaced spacecraft distribution shown in Figure 4b. The shock itself is seen at all three spacecraft at similar times, suggesting a nearly spherical shape.




Figure 4 (a) Intensities of 20 MeV protons vs. time are shown for the three spacecraft located as in (b). Similar shock arrival times at the spacecraft indicated in (a) suggest a nearly spherical shock [82]. The pink field lines labeled 1 and 2 in (b) will rotate to intercept *Wind* when the shock reaches 1 or 2 AU, respectively, bearing some of the SEPs acquired at their initial longitudes.

4.2 Element Abundances and Seed Particles

Abundances of elements, accelerated by shock waves to a given velocity, or energy per nucleon, tend to reflect the abundances of the seed particles initially injected into the shock. In most gradual SEP events this is the ambient abundances of the solar corona near the active region where the shock begins, but shocks will accelerate any (and all) seed particles that are available. The question is: which will dominate?

Mason, Mazur, and Dwyer [83] first noticed that shocks could reaccelerate ³He and other impulsive ions provided by earlier impulsive SEP events. It was recognized [84 – 87] that quasi-perpendicular shocks might selectively prefer the pre-accelerated seed population of ions that had sufficient speed to overtake the shock from behind, i.e. residual suprathermal particles from impulsive events. It turns out that residual ions from many persistent impulsive flares can accumulate above active regions, providing a seed population, in addition to the ambient corona plasma, thus complicating the abundance pattern of elements in SEP events. Reames [88] identified four processes leading to different SEP abundance patterns:

- SEP1: "Pure" impulsive SEPs with enhanced heavy elements from reconnection in solar jets, but no fast shocks.
- SEP2: Impulsive SEP1 ions reaccelerated as seed particles of a fast CME-driven shock from the same jet [89 91] dominate high Z.
- SEP3: Enhanced abundances accelerated by a fast CME-driven shock traversing an active region with residual seed ions from many small impulsive events that will dominate high *Z*.
- SEP4: Particles accelerated by a fast, wide, CME-driven shock with only ambient coronal abundances available as the dominant seed particles.

Since shock waves will accelerate any ions available, SEP2 and SEP3 events tend to reflect both types of seed particles, the heavy-ion-rich impulsive seeds dominate the high-Z SEPs, and the ambient abundances are left to dominate protons. These patterns

are compared in Figure 5. Fits to the abundance enhancements vs. A/Q involve varying plasma temperatures to determine ion A/Q values from their well-determined temperature dependence [92, 93]. Best-fit temperatures are shown in Figure 4b and fit lines in Figure 4c, for three time-intervals. The events in the left panel in Figure 4c show rising enhancements vs. A/Q for shockless SEP1 impulsive SEP events, the event in the right panel shows SEP4 falling enhancements vs. A/Q where higher-A/Q elements from the ambient-coronal seeds have preferentially leaked away, and the middle panel shows a SEP2 or SEP3 event with components of both rising and falling seed populations.




Figure 5 In (a) intensities of H, 4 He, O, and Fe in the listed intervals of MeV amu $^{-1}$ are shown for three time intervals. (b) Temperatures derived from best-fits of enhancement vs. A/Q show with power-law fits in the same color in (c) above [88, 94, 95].

In the impulsive SEP1 events, protons either lie on the fit line, or perhaps, theory allows the increase to begin above He [22] so H and ⁴He would both be equally unenhanced. For SEP4 events protons again lie on the fit line since all ions belong to a single seed-particle population. For SEP2 and SEP3 events, proton enhancements signal the presence of two seed populations. Sequences of SEP3 events, dominated by impulsive seed ions, can come from a single active region, [96]. These help distinguish SEP3 events from smaller, isolated SEP2 events.

Very intense SEP4 events do have rising initial A/Q dependence when only the higher-Z ions can penetrate out to the observer. For these SEP4 events, the A/Q-

dependence suddenly declines as the shock passes; for SEP3 events, the enhancements persist across the shock since they are a fundamental property of the impulsive seeds, not of transport. Element abundances and their time dependence is a powerful tool in the study of SEP acceleration and transport.

Using the assumption of power-law abundance enhancements to derive source plasma temperatures may seem uncertain, but Bučík et al. [97] have now directly confirmed similar temperatures of ~2.5 MK in the source jets that produce impulsive SEP events. Also, nearly all of the SEP abundance studies have used Maxwellian electron distributions to determine A/Q vs. temperature. However, using kappa distributions for the thermal electrons, Lee et al. [98] have shown that the Maxwellian distributions used to derive A/Q vs. temperature [92, 93] did not produce large errors. They also suggested that since the SEP and source temperatures are similar, the SEPs must leave the acceleration region on a time scale shorter than their ionization time. After acceleration these ions evidently pass through enough matter to strip them to an equilibrium ionization state that varies with velocity [99] and leaves $Q_{\rm Fe} \approx 20$ near 1 MeV amu⁻¹. This stripping suggests [99] that impulsive SEP events occur at about 1.5 solar radii. These ions from SEP1 events must retain their higher "stripped" charge states if they get reaccelerated by shock waves to produce SEP2 or SEP3 events, even though the shock acceleration may occur at the much lower densities at 2-3 solar radii. Thus, the events with impulsive seeds have enhanced Fe/O and charge states appropriate to their stripping energy after the SEP1 phase, but their abundance-derived temperatures and O-values are appropriate to their earlier state before and during the SEP1 phase. In contrast, the SEP4 events have ambient coronal charge states (at ~ 1 MK) with coronal source abundances modified only by transport.

5 Discussion

Recently, Mason et al. [100] suggested that the enhancement of Fe/O could come from the suppression of O which resonates with waves damped by 4 He at A/O = 2. However, this fails to explain the strong continuing enhancement of heavier elements well above Fe, fairly well approximated as a power law in A/O observed at all energies [9, 11,12]. This power-law increase does not exclude the possibility that additional resonant wave absorption can produce selective peaks in Si and S in some small events [101], presumable near $A/Q \sim 3.0$ as the second harmonic of ³He at A/Q = 1.5. Turbulent reconnection in jets may well produce both power-law reconnection [22] and resonant [102] processes, especially in the small SEP events with steep spectra that have only been seen below ~1 MeV amu⁻¹. Jets that produce larger impulsive SEP events with O and Fe spectra measurable out to 10 MeV amu-1 or more are likely to include CMEs that can drive shock waves to 500 km s⁻¹ or more, reaccelerate the SEPs, and perhaps even smooth the A/O dependence by including extensive or multiple reconnection regions that average differing characteristics. In the list of Fe-rich events at 3 - 10 MeV amu⁻¹ presented by Reames et al. [9], about 40% of the events have CMEs with measured speeds above 500 km s⁻¹; these are surely SEP2 events and many of them are probably SEP3 events that average extensive sources of seeds. Generally, they are fit with power-laws in A/Qdetermined at temperatures of 2.5 – 3.2 MK [6, 94]. Despite the dominance of powerlaw dependence, resonant wave absorption processes may still contribute the occasional

10

additional enhancements seen in Ne or Mg may suggest a small resonant component, perhaps injected at lower temperature so that $A/Q \sim 3$.

For gradual SEP events, we have come a long way since Gosling's "Solar flare myth" [103, 104] pointed out the dominance of CMEs and shock waves, rather than flares, in interplanetary phenomena, including SEPs. The STEREO mission has shown how shock waves, and the SEPs they accelerate, can envelope the Sun (e.g. Figure 5) [82], while gamma-ray-line measurements suggest that the particles that are accelerated and magnetically trapped in flares are ³He-rich [105] and Fe-rich [106, 107] like those of the impulsive SEPs we now know to come from jets [14]. Thus impulsive SEPs are logically connected to flares, but do not actually come from flares; they come out to us from jets. For gradual events we must consider the hierarchy of seed particles, when available, and their reacceleration by shocks.

In most large gradual events, heavy ion abundances seem to be dominated by either impulsive (SEP3) or ambient-coronal (SEP4) seed ions across the whole longitude span of the shock [108, 96]. However, one might expect to find SEP events where the shock has sampled different ions at different longitudes. Recently, Xu et al. [109] have found just such an event, where spacecraft separated by $\sim 60^{\circ}$ longitude clearly see Ferich material at one spacecraft and Fe-poor material at the other over the full energy range of 0.1-10 MeV amu⁻¹.

Resonant wave-particle interactions are an essential factor in SEP acceleration. They selectively enhance ³He in impulsive SEP events [39, 45], they enable the dominant acceleration as they scatter ions back and forth across a shock [15, 16, 57], and, as intensities increase, they produce rigidity-dependent scattering that can flatten spectra and enhance or suppress element abundances [110, 111, 79, 75, 76]. These abundances and spectra tell us about the underlying physical processes.

Parker [112] proposed the release of magnetic energy in current sheets of many small *nanoflares* as a mechanism for heating the entire solar corona. If this is the case, it is easy to imagine the acceleration of energetic particles in magnetic loops that carry that energy downward and deposit it in the footpoints of the loops. Electrons emit much of their energy as X-ray bremstralung but stopping ions largely heat the plasma. Thus SEP acceleration may actually be a significant agent in the heating of solar and stellar coronae.

For stars, other than the Sun, there is information on flares but not on CMEs or SEPs. Kahler's [113] paper on "big flare syndrome" provided the first caution that a correlation between SEPs and flare-related parameters, for example, does not mean that flares *cause* SEPs, but rather suggests that increasingly large dissipation of magnetic energy can result in many increasing phenomena. Nevertheless, big flare syndrome does suggest that flare stars may have more intense SEPs, so the underlying correlation is explored further by Kahler and Ling [114]. Other studies like Hu et al. [115] apply CME models to superflares, while Fu et al. [116] study the dependence of SEP properties on stellar rotation speeds.

Conflict of Interests

The author declares that this research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Funding

No institutional funding was provided for this work.

Reference

- Kahler, S.W., Sheeley, N.R., Jr., Howard, R.A., Koomen, M.J., Michels, D.J., McGuire R.E., von Rosenvinge, T.T., Reames, D.V., Associations between coronal mass ejections and solar energetic proton events, *J. Geophys. Res.* 1984 89, 9683 https://doi.org/10.1029/JA089iA11p09683
- 2 Reames D.V., Solar Energetic Particles, (Second Edition) Lec. Notes Phys. 978 Springer Nature, Cham, Switzerland, (2021) open access, https://doi.org/10.1007/978-3-030-66402-2
- 3 Reames, D.V., Bimodal abundances in the energetic particles of solar and interplanetary origin, Astrophys. J. Lett. 1988 330, L71 doi: 10.1086/185207
- Reames, D.V., Abundances of trans-iron elements in solar energetic particle events, Astrophys. J. Lett. 2000 540, L111 doi: 10.1086/312886
- 5 Reames, D.V., The two sources of solar energetic particles, Space Sci. Rev. 2013 175, 53 https://doi.org/10.1007/s11214-013-9958-9
- Reames, D. V., Abundances, ionization states, temperatures, and FIP in solar energetic particles. Space Sci. Rev. 2018 214, 61. https://doi.org/10.1007/s11214-018-0495-4
- 7 Reames, D. V., Fifty Years of ³He-Rich Events. Front. Astron. Space Sci. 2021 8,164. https://doi.org/10.3389/fspas.2021.760261
- 8 Reames, D.V., Meyer, J.P., von Rosenvinge, T.T., Energetic-particle abundances in impulsive solar flare events, Astrophys. J. Suppl. 1994 90, 649 https://doi.org/10.1086/191887
- Reames, D.V., Cliver, E.W., Kahler, S.W., Abundance enhancements in impulsive solar energetic-particle events with associated coronal mass ejections, *Solar Phys.* 2014 289, 3817 https://doi.org/10.1007/s11207-014-0547-1
- 10 Mason, G.M., Reames, D.V., Klecker, B., Hovestadt, D., von Rosenvinge, T.T., The heavy-ion compositional signature in He-3-rich solar particle events, Astrophys. J. 1986 303, 849 doi: 10.1086/164133
- 11 Mason, G.M., Mazur, J.E., Dwyer, J.R., Jokippi, J.R., Gold, R.E., Krimigis, S.M., Abundances of heavy and ultraheavy ions in ³He-rich solar flares, Astrophys. J. 2004 606, 555 doi: 10.1086/382864
- 12 Reames, D.V., Ng, C.K., Heavy-element abundances in solar energetic particle events, Astrophys. J. 2004 610, 510 https://doi.org/10.1088/0004-637X/723/2/1286
- 13 Mason, G.M. ³He-rich solar energetic particle events. Space Sci. Rev. **2007** 130, 231 https://doi.org/10.1007/s11214-007-9156-8
- 14 Bučík, R., ³He-rich solar energetic particles: solar sources, Space Sci. Rev. 2020 216 24 doi: 10.1007/s11214-020-00650-5
- 15 Lee, M.A., Coupled hydromagnetic wave excitation and ion acceleration at interplanetary traveling shocks, J. Geophys. Res. 1983 88, 6109. https://doi.org/10.1029/JA088iA08p06109
- 16 Lee, M.A., Coupled hydromagnetic wave excitation and ion acceleration at an evolving coronal/interplanetary shock, Astrophys. J. Suppl., 2005 158, 38 https://doi.org/10.1086/428753
- 17 Reames, D.V., Particle acceleration at the Sun and in the heliosphere, Space Sci. Rev. 1999 90, 413 https://doi.org/10.1023/A:1005105831781
- 18 D.V. Reames, How do shock waves define the space-time structure of gradual solar energetic-particle events? Space Sci. Rev. 2023 219, 14. https://doi.org/10.1007/s11214-023-00959-x
- 19 Zank, G.P., Rice, W.K.M., Wu, C.C., Particle acceleration and coronal mass ejection driven shocks: A theoretical model, J. Geophys. Res. 2000 105, 25079 https://doi.org/10.1029/1999JA000455
- 20 Desai, M.I., Giacalone, J., Large gradual solar energetic particle events, Living Reviews of Solar Physics 2016 13 3 https://doi.org/10.1007/s41116-016-0002-5

2 Reconn

- 21 Parker, E.N., The formation of sunspots from the solar toroidal field, Astrophys. J. 1955 121 491 doi: 10.1086/146010
- 22 Drake, J.F., Cassak, P.A., Shay, M.A., Swisdak, M., Quataert, E., A magnetic reconnection mechanism for ion acceleration and abundance enhancements in impulsive flares, *Astrophys. J. Lett.* 2009 700, L16 doi: https://doi.org/10.1088/0004-637X/700/1/L16
- 23 Kramoliš, D., Bárta, M., Varady, M., Bučík, R., Preferential acceleration of heavy ions in a spontaneously fragmenting flare current sheet. Astrophys. J. 2022 927 (2) 177 https://doi.org/10.3847/1538-4357/ac4fc9
- 24 Fitzmaurice, A., Drake, J.F., Swisdak, M., Wave generation by flare-accelerated ions and implications for 3He acceleration. Astrophys. J. 2024 964, 97 https://doi.org/10.3847/1538-4357/ad217f
- 25 Reames, D.V., Dennis, B.R., Stone, R.G., Lin, R.P., X-ray and radio properties of solar ³He-rich events Astrophys. J. 1988 327, 998 doi: 10.1086/166257
- 26 Reames, D.V., Temperature dependence of the abundances of elements in solar ³He-rich events, Astrophys. J. Lett. 1988 325 L53. doi: 10.1086/185109

- 27 Kopp, R.A., Pneuman, G. W. Magnetic reconnection in the corona and the loop prominence phenomena, Solar Physics 1976 50 85 doi: 10.1007/BF00206193
- 28 Reames, D. V., Magnetic topology of impulsive and gradual solar energetic particle events. Astrophys. J. Lett. 2002 571 L63. https://doi.org/doi:10.1086/341149
- 29 Archontis, V., Hood, A.W., A numerical model of standard to blowout jets, *Astrophys. J. Lett.* 2013, 769 L21 doi: 10.1088/2041-8205/769/2/L21 [Erratum: *Astrophys. J. Lett.* 2013, 770, L41].
- 30 Lee, E.J., Archontis, V., Hood, A.W., Plasma jets and eruptions in solar coronal holes: a three-dimensional flux emergence experiment, Astrophys. J. Lett. 2015, 798 L10 doi: 10.1088/2041-8205/798/1/L10
- 31 Pariat, E., Dalmasse, K., DeVore, C.R., Antiochos, S.K., Karpen, J.T., Model for straight and helical solar jets. I. Parametric studies of the magnetic field geometry, Astron. Astrophys. 2015 573 A130 doi: 10.1051/0004-6361/201424209

3 Impul

- 32 Wild, J.P., Smerd, S.F., Weiss, A.A., Solar Bursts, Annu. Rev. Astron. Astrophys. 1963 1, 291 doi: 10.1146/annurev.aa.01.090163.001451
- 33 Lin, R.P., The emission and propagation of 40 keV solar flare electrons. I: The relationship of 40 keV electron to energetic proton and relativistic electron emission by the sun, Sol. Phys. 1970 12, 266 doi: 10.1007/BF00227122
- 34 Serlemitsos, A.T., Balasubrahmanyan, V.K., Solar particle events with anomalously large relative abundance of ³He, Astrophys. J. 1975 198, 195 doi: 10.1086/153592
- 35 Reames, D.V., von Rosenvinge, T.T., Lin, R.P., Solar ³He-rich events and nonrelativistic electron events A new association, *Astrophys. J.* **1985** 292, 716 doi: 10.1086/163203
- 36 Reames, D.V., Stone, R.G., The identification of solar ³He-rich events and the study of particle acceleration at the sun, *Astrophys. J.* **1986**, 308, 902 https://doi.org/10.1086/164560
- 37 Ibragimov, I. A., Kocharov, G. E., Possible mechanism for enrichment of solar cosmic rays by helium-three and heavy nuclei, 15th Int. Conf. on Cosmic Rays (Plovdiv: Bulgarian Academy of Sciences), 11, 340 (1977)
- 38 Kocharov, G. E., Kocharov, L. G., Present state of experimental and theoretical investigations of solar events enriched by helium-3, in *Proc. 10th Leningrad Sympos. on Cosmic Rays* (Leningrad: A. F. Yoffe Phys.-Tech. Inst.), p. 37 (1978)
- 39 Fisk, L.A., ³He-rich flares a possible explanation, *Astrophys. J.* **1978** 224, 1048 doi: 10.1086/156456
- 40 Varvoglis, H., Papadopoulis, K., Selective nonresonant acceleration of He-3(2+) and heavy ions by H(+) cyclotron waves Astrophys. J. Lett. 1983 270, L95 doi: 10.1086/184077
- 41 Kocharov, G. E., Kocharov, L. G., ³He-rich solar flares, Space Science Rev. 1984 38, 89 doi: 10.1007/BF00180337
- 42 Weatherall, J., Turbulent heating in solar cosmic-ray theory, Astrophys. J. 1984 281, 468 doi: 10.1086/162119
- 43 Winglee, R.M., Heating and acceleration of heavy ions during solar flares, Astrophys. J. 1989 343, 511 doi: 10.1086/167726
- 44 Riyopoulos, S., Subthreshold stochastic diffusion with application to selective acceleration of ³He in solar flares, *Astrophys. J.* **1991** 381 578 doi: 10.1086/170682
- 45 Temerin, M., Roth, I., The production of ³He and heavy ion enrichment in ³He-rich flares by electromagnetic hydrogen cyclotron waves, *Astrophys. J. Lett.* 1992 391, L105 https://doi.org/10.1086/186408
- 46 Ho, G.C., Roelof, E.C., Mason, G.M., The upper limit on ³He fluence in solar energetic particle events, *Atrophys. J. Lett.* **2005** 621, L141 doi: 10.1086/429251
- 47 Liu, S., Petrosian, V., Mason, G.M.: Stochastic acceleration of ³He and ⁴He in solar flares by parallel-propagating plasma waves, *Astrophys. J. Lett.* 2004 613, L13 doi: 10.1086/425070
- 48 Liu, S., Petrosian, V., Mason, G.M.: Stochastic acceleration of ³He and ⁴He in solar flares by parallel-propagating plasma waves: general results. *Astrophys. J.* **2006** 636, 462 doi: 10.1086/497883
- 49 Petrosian, V., Jiang, Y. W., Liu, S., Ho, G. C., and Mason, G. M., Relative distributions of fluences of ³He and ⁴He in solar energetic particles. *Astrophys. J.* 2009 701 1 https://doi.org/10.1088/0004-637X/701/1/1
- 50 Mogro-Campero, A., and Simpson, J. A. Enrichment of very heavy nuclei in the composition of solar accelerated particles. Astrophys. J. Lett. 1972 171, L5 https://doi.org/10.1086/180856
- 51 Eichler, D. Particle acceleration in solar flares by cyclotron damping of cascading turbulence Astrophys. J. 1979 229 413 doi: 10.1086/156969
- 52 Miller, J. A., and Reames, D.V. Heavy ion acceleration by cascading Alfvén waves in impulsive solar flares, AIP Conf. Proc. 1996 374, 450 https://doi.org/10.1063/1.50981
- 53 Eichler, D. Ultraheavy element enrichment in impulsive solar flares. Astrophys. J. 2014 794(1): 6, https://doi.org/10.1088/0004-637X/794/1/6

4 Shock Accel.

- 54 Jones, F.C., Ellison, D.E., The plasma physics of shock acceleration, Space Sci. Rev. 1991 58, 259 doi: 10.1007/BF01206003
- 55 Bell, A. R. The acceleration of cosmic rays in shock fronts. I. Mon. Not. Roy. Astron. Soc. 1978 182, 147–156. https://doi.org/10.1093/mnras/182.2.147
- Bell, A. R. The acceleration of cosmic rays in shock fronts. II. Mon. Not. Roy. Astron. Soc. 1978 182, 443–455. https://doi.org/10.1093/mnras/182.3.443

4.1

- 57 Ng, C.K., Reames, D.V., Shock acceleration of solar energetic protons: the first 10 minutes, Astrophys. J. Lett. 2008 686, L123 doi: 10.1086/592996
- 58 Stix, T.H., Waves in Plasmas (New York: AIP) (1992)
- 59 Melrose, D. B. *Plasma astrophysics*. New York: Gordon & Breach (1980).
- 60 Decker, R.B.: Formation of shock-spike events at quasi-perpendicular shocks. J. Geophys. Res. 1983 88 9959 https://doi.org/10.1029/JA088iA12p09959
- 61 Reames, D. V., Solar release times of energetic particles in ground-level events, Astrophys. J. 2009 693, 812 http://doi.org/10.1088/0004-637X/693/1/812
- 62 Reames, D. V., Solar energetic-particle release times in historic ground-level events, Astrophys. J. 2009 706, 844 http://doi.org/10.1088/0004-637X/706/1/844
- 63 Cliver, E.W., Kahler, S.W., and Reames, D.V., Coronal shocks and solar energetic proton events, *Astrophys. J.* 2004 605, 902 http://doi.org/10.1086/382651
- 64 Meyer, J.P., The baseline composition of solar energetic particles, Astrophys. J. Suppl. 1985 57, 151 http://doi.org/10.1086/191000
- 65 Laming, J.M., The FIP and inverse FIP effects in solar and stellar coronae, Living Reviews in Solar Physics 2015 12, 2 doi: 10.1007/Irsp-2015-2
- 66 Reames, D.V., Coronal Abundances determined from energetic particles, Adv. Space Res. 1995 15 (7), 41
- 67 Reames, D.V., Element abundances in solar energetic particles and the solar corona, Solar Phys. 2014 289, 977 doi: 10.1007/s11207-013-0350-4
- 68 Reames, D. V., Richardson, I. G., and Barbier, L. M. On the differences in element abundances of energetic ions from corotating events and from large solar events. *Astrophys. J. Lett.* **1991** 382, L43. http://doi.org/10.1086/186209
- 69 Mewaldt, R.A., Cohen, C.M.S., Leske, R.A., Christian, E.R., Cummings, A.C., Stone, E.C., von Rosenvinge, T.T. and Wiedenbeck, M. E., Fractionation of solar energetic particles and solar wind according to first ionization potential, *Advan. Space Res.* 2002 30, 79 doi: 10.1016/S0273-1177(02)00263-6
- 70 Reames, D.V., The "FIP effect" and the origins of solar energetic particles and of the solar wind, Solar Phys. 2018 293 47 https://doi.org/10.1007/s11207-018-1267-8 (arXiv 1801.05840)
- 71 Laming, J.M., Vourlidas, A., Korendyke, C., et al., Element abundances: a new diagnostic for the solar wind, Astrophys. J. 2019 879 124 doi: 10.3847/1538-4357/ab23f1
- 72 Mason, G.M., Ng, C.K., Klecker, B., Green, G., Impulsive acceleration and scatter-free transport of about 1 MeV per nucleon ions in ³He-rich solar particle events, *Astrophys. J.* 1989 339, 529 doi: 10.1086/167315
- 73 Breneman, H.H., Stone, E.C., Solar coronal and photospheric abundances from solar energetic particle measurements, Astrophys. J. Lett. 1985 299, L57 http://doi.org/10.1086/184580
- 74 Reames, D.V., Acceleration of energetic particles by shock waves from large solar flares, Astrophys. J. Lett. 1990 358, L63 doi: 10.1086/185780
- 75 Reames, D.V. Ng, C.K., Streaming-limited intensities of solar energetic particles, Astrophys. J. 1998 504, 1002 doi: 10.1086/306124
- 76 Reames, D.V., Ng, C.K., Streaming-limited intensities of solar energetic particles on the intensity plateau, Astrophys. J. 2010 723 1286 doi: 10.1088/0004-637X/723/2/1286
- 77 Giacalone, J. et al. Analyses of ~0.05–2 MeV ions associated with the 2022 February 16 energetic storm particle event observed by Parker Solar Probe, *Astrophys, J.* 2023 958 144 https://doi.org/10.3847/1538-4357/acfb86
- 78 Ng, C.K., Reames, D.V., Tylka, A.J., Solar energetic particles: shock acceleration and transport through self-amplified waves, AIP Conf. Proc. 2012 1436, 212 doi: 10.1063/1.4723610
- 79 Ng, C.K., Reames, D.V., Tylka, A.J., Modeling shock-accelerated solar energetic particles coupled to interplanetary Alfvén waves, Astrophys. J. 2003 591, 461 https://doi.org/10.1086/375293
- 80 Reames, D.V., and Ng, C.K. The streaming limit of solar energetic-particle intensities, *Living with a star workshop on extreme space weather events*. Boulder, Co. June 9-11 (2014) https://arxiv.org/abs/1412.2279
- 81 Ng, C.K., Effect of solar-wind velocity, magnetic field and density on solar energetic particle transport, ASP Conf. Ser. 2014 484 156 Qiang Hu and Gary P. Zank, eds.

82 Reames, D.V. Review and outlook of solar-energetic-particle measurements on multispacecraft missions, Front. Astron. Space Sci. 2023 10 https://doi.org/10.3389/fspas.2023.1254266

- 4.2
 83 Mason, G.M., Mazur, J.E., Dwyer, J.R., ³He enhancements in large solar energetic particle events, *Astrophys. J.* Lett. 1999 525, L133 https://doi.org/10.1086/312349
- 84 Desai, M.I., Mason, G.M., Dwyer, J.R., Mazur, J.E., Gold, R.E., Krimigis, S.M., Smith, C.W., Skoug, R.M., Evidence for a suprathermal seed population of heavy ions accelerated by interplanetary shocks near 1 AU, Astrophys. J. 2003, 588, 1149 https://doi.org/10.1086/374310
- 85 Tylka, A.J., Cohen, C.M.S., Dietrich, W.F., Lee, M.A., Maclennan, C.G., Mewaldt, R.A., Ng, C.K., Reames, D.V., Shock geometry, seed populations, and the origin of variable elemental composition at high energies in large gradual solar particle events, *Astrophys. J.* 2005 625, 474 https://doi.org/10.1086/429384

- 7 Tylka, A.J., Lee, M.A., Spectral and compositional characteristics of gradual and impulsive solar energetic particle events, *Astrophys. J.* **2006** 646, 1319 https://doi.org/10.1086/505106
- 87 Sandroos, A., Vainio, R., Simulation results for heavy ion spectral variability in large gradual solar energetic particle events, *Astrophys. J.* **2007** 662, L127 doi: 10.1086/519378
- 88 Reames, D.V., Four distinct pathways to the element abundances in solar energetic particles, *Space Sci. Rev.* **2020** 216 20 doi: 10.1007/s11214-020-0643-5
- 89 Kahler, S.W., Reames, D.V., Sheeley, N.R., Jr., Coronal mass ejections associated with impulsive solar energetic particle events, *Astrophys. J.* **2001** 562, 558 https://doi.org/10.1086/323847
- 90 Nitta, N.V., Reames, D.V., DeRosa, M.L., Yashiro, S., Gopalswamy, N., Solar sources of impulsive solar energetic particle events and their magnetic field connection to the earth, Astrophys. J. 2006 650, 438 doi: 10.1086/507442
- Wang, Y.-M., Pick, M., Mason, G.M., Coronal holes, jets, and the origin of ³He-rich particle events, *Astrophys. J.* **2006** 639, 495 doi: 10.1086/499355
- 92 Mazzotta, P., Mazzitelli, G., Colafrancesco, S., Vittorio, N. Ionization balance for optically thin plasmas: rate coefficients for all atoms and ions of the elements H to Ni. Astron. Astrophys Suppl. 1988 133, 403 https://doi.org/10.1051/aas:1998330
- 93 Post, D.E., Jensen, R.V., Tarter, C.B., Grasberger, W.H., Lokke, W.A. Steady-state radiative cooling rates for low-density, high temperature plasmas. At. Data Nucl. Data Tables. 1977 20, 397 https://doi.org/10.1016/0092-640X(77)90026-2
- 94 Reames, D.V., Cliver, E.W., Kahler, S.W., Variations in abundance enhancements in impulsive solar energetic-particle events and related CMEs and flares, Solar Phys. 2014 289, 4675 http://doi.org/10.1007/s11207-014-0589-4
- 95 Reames, D.V., Temperature of the source plasma in gradual solar energetic particle events, *Solar Phys.* **2016**, 291 911 http://doi.org/10.1007/s11207-016-0854-9, arXiv: 1509.08948
- 96 Reames, D.V. Seeds and sequences of element abundances in solar energetic particle events, Eur. Phys. J. Plus 2024 139 710. http://doi.org/10.3389/fspas.2024.1368043
- 97 Bučík, R., Mulay, S.M., Mason, G.M., Nitta, N.V., Desai, M.I., Dayeh, M.A., Temperature in solar sources of ³Herich solar energetic particles and relation to ion abundances, *Astrophys. J.* 2021 908 243 doi: 10.3847/1538-4357/abd62d
- 28 Lee, J. Y., Kahler, S., Raymond, J. C., and Ko, Y. K. Solar energetic particle charge states and abundances with nonthermal electrons. *Astrophys. J.* 2024 963 70 doi 10.3847/1538-4357/ad1ab6
- 99 DiFabio, R., Guo, Z., Möbius, E., Klecker, B., Kucharek, H., Mason, G.M., Popecki, M., Energy-dependent charge states and their connection with ion abundances in impulsive solar energetic particle events, *Astrophys. J.* 2008 687, 623 doi: 10.1086/591833

5 Discussion

- 100 Mason, G.M., Roth, I., Nitta, N.V., Bučík, R., Lario, D., Ho, G.C. Allen, R.C., Kouloumvakos, A., Wimmer-Schweingruber, R.F., Rodriguez-Pacheco, J., Heavy-ion acceleration in ³He-rich solar energetic particle events: new insights from Solar Orbiter. Astrophys. J. 2023 957, 112. http://doi.org/10.3847/1538-4357/acf31b
- 101 Mason, G. M., Nitta, N. V., Wiedenbeck, M. E., and Innes, D. E. Evidence for a common acceleration mechanism for enrichments of ³He and heavy ions in impulsive EP events. Astrophys. J. 2016, 138. http://doi.org/10.3847/0004-637X/823/2/138
- 102 Roth, I., and Temerin, M. Enrichment of ³He and heavy ions in impulsive solar flares. Astrophys. J. 1997 477, 940–957. http://doi.org/10.1086/303731
- 103 Gosling, J.T., The solar flare myth. J. Geophys. Res. 1993 98, 18937 http://doi.org/10.1029/93JA01896
- 104 Gosling, J.T., Corrections to "The solar flare myth." J. Geophys. Res. 1994 99, 4259 doi: 10.1029/94JA00015
- 105 Mandzhavidze, N., Ramaty, R., Kozlovsky, B., Determination of the abundances of subcoronal ⁴He and of solar flare-accelerated ³He and ⁴He from gamma-ray spectroscopy, Astrophys. J. 1999 518, 918 doi: 10.1086/307321
- 106 Murphy, R.J., Kozlovsky, B., Share, G.H.,: Evidence for enhanced ³He in flare-accelerated particles based on new calculations of the gamma-ray line spectrum, Astrophys. J. 2016 833, 166 doi: 10.3847/1538-4357/833/2/196
- 107 Murphy, R.J., Ramaty, R., Kozlovsky, B., Reames, D.V., Solar abundances from gamma-ray spectroscopy: Comparisons with energetic particle, photospheric, and coronal abundances, *Astrophys. J.* 1991 371, 793 doi: 10.1086/169944
- 108 Cohen, C. M. S., Mason, G. M., and Mewaldt, R. A. Characteristics of solar energetic ions as a function of longitude. Astrophys. J. 2017 132. http://doi.org/10.3847/1538-4357/aa7513
- 109 Xu, Z.G., Cohen, C.M.S., Leske, R.A., et al. Composition variation of the May 16 2023 Solar Energetic Particle Event observed by SolO and PSP Astrophys. J. Lett. 2024 976 L3 doi: 10.3847/2041-8213/ad8b18
- 110 Ng, C.K., Reames, D.V., Focused interplanetary transport of approximately 1 MeV solar energetic protons through self-generated Alfven waves, Astrophys. J. 1994 424, 1032 doi: 10.1086/173954
- 111 Ng, C.K., Reames, D.V., Tylka, A.J., Effect of proton-amplified waves on the evolution of solar energetic particle composition in gradual events, *Geophys. Res. Lett.* 1999 26, 2145 doi: 10.1029/1999GL900459
- 112 Parker, E.N.: Nanoflares and the solar X-ray corona. Astrophys. J. 1988 330, 474 https://doi.org/10.1086/166485
- 113 Kahler, S.W. The role of the big flare syndrome in correlations of solar energetic proton fluxes and associated microwave burst parameters. J. Geophys. Res. 1982 87 3439 https://doi.org/10.1029/JA087iA05p03439
- 114 Kahler, S.W., Ling, A.G., Forecasting solar energetic particle (SEP) events with flare X-ray peak ratios, Space Weather Space Clim. 2018, 8, A47 https://doi.org/10.1051/swsc/2018033
- 115 Hu, J., Airapetian, V., Li, G., Zank, G., Jin, M., Extreme energetic particle events by superflare-associated CMEs from solar-like stars, Sci. Adv. 2022 8, eabi9743 https://doi.org/10.1126/sciadv.abi9743

116 Fu, S., Jiang, Y., Airapetian, V., Hu, J., Li, G., Zank, G., Effect of star rotation rates on the characteristics of energetic particle events *Astrophys. J. Lett.* 2019 878, L36 doi: 10.3847/2041-8213/ab271d